

Free actions on products of spheres at high dimensions

OSMAN BERAT OKUTAN
ERGÜN YALÇIN

A classical conjecture in transformation group theory states that if $G = (\mathbb{Z}/p)^r$ acts freely on a product of k spheres $S^{n_1} \times \cdots \times S^{n_k}$, then $r \leq k$. We prove this conjecture in the case where the dimensions $\{n_i\}$ are high compared to all the differences $|n_i - n_j|$ between the dimensions.

57S25; 20J06

1 Introduction

Let G be a finite group. The rank of G , denoted by $\text{rk}(G)$, is defined as the largest integer s such that $(\mathbb{Z}/p)^s \leq G$ for some prime p . It is known that G acts freely and cellularly on a finite complex homotopy equivalent to a sphere S^n if and only if $\text{rk}(G) = 1$. This follows from the results due to P A Smith [12] and R Swan [13]. As a generalization of this, it has been conjectured by Benson and Carlson [3] that $\text{rk}(G) = \text{hrk}(G)$ where $\text{hrk}(G)$ is defined as the smallest integer k such that G acts freely and cellularly on a finite CW-complex homotopy equivalent to a product of k spheres. This conjecture is often referred to as the rank conjecture. Note that one direction of the Benson–Carlson conjecture is the following statement:

Conjecture 1.1 *Let p be a prime. If $G = (\mathbb{Z}/p)^r$ acts freely and cellularly on a finite CW-complex X homotopy equivalent to $S^{n_1} \times \cdots \times S^{n_k}$, then $r \leq k$.*

This conjecture is a classical conjecture that was studied intensely through the 80's and it has been proven that the conjecture is true under some additional assumptions. For example it is known that when the dimensions of the spheres are all equal, ie, $n = n_1 = \cdots = n_k$, then the conjecture is true for all primes p and for all positive integers n except when $p = 2$ and $n = 3, 7$. This was proved by G Carlsson [6] in the case where the G -action on the homology of X is trivial and the general case is due to Adem and Browder [2]. The $p = 2$ and $n = 1$ case was proven later by Yalçin [14]. More recently, B Hanke [10] proved that [Conjecture 1.1](#) is true in the case where

$p \geq 3 \dim X$, ie, when the prime p is large compared to the dimension of the space. In this paper, we prove [Conjecture 1.1](#) for the other extreme, ie, when the dimensions of the spheres are high compared to all the differences between the dimensions.

Theorem 1.2 *Suppose $G = (\mathbb{Z}/p)^r$ for a prime p and k, l are positive integers. Then there is an integer N that depends only on k, l and G such that if G acts freely and cellularly on a finite dimensional CW-complex X homotopy equivalent to $S^{n_1} \times \dots \times S^{n_k}$ where $n_i \geq N$ and $|n_i - n_j| \leq l$ for all i, j , then $r \leq k$.*

The proof follows from a theorem of Browder [\[4\]](#) that gives a restriction on the order of groups acting freely on a finite dimensional CW-complex in terms of homology groups of the complex. We also use a method of gluing homology groups at different dimensions, which we first saw in a paper by Habegger [\[9\]](#), and a crucial result on the exponents of cohomology groups of elementary abelian p -groups, which is due to Pakianathan [\[11\]](#).

At the end of the paper we also prove a generalization of [Theorem 1.2](#) to non-free actions, which was suggested to us by A Adem.

The paper is organized as follows: In [Section 2](#), we list some well-known results about hypercohomology and in [Section 3](#), we introduce Habegger’s theorem on gluing homology at different dimensions. In [Section 4](#), we discuss the exponents of Tate cohomology groups and in [Section 5](#), we prove [Theorem 1.2](#), which is our main theorem.

2 Tate hypercohomology

Let G be a finite group and M be a $\mathbb{Z}G$ -module. The Tate cohomology of G with coefficients in M is defined as follows:

$$\hat{H}^i(G, M) := H^i(\text{Hom}_G(F_*, M))$$

for all $i \in \mathbb{Z}$, where F_* is a complete $\mathbb{Z}G$ -resolution of \mathbb{Z} (see Brown [\[5\]](#), page 134). We can generalize this and define Tate hypercohomology of G with coefficients in a chain complex C_* of $\mathbb{Z}G$ -modules. To do this, we need to extend the contravariant functor $\text{Hom}_G(-, M)$ to $\mathcal{H}om_G(-, C_*)$. We will define it as in Brown [\[5\]](#), page 5], but instead of defining it as a chain complex, we consider it as a cochain complex.

Suppose C_* and D_* are chain complexes over $\mathbb{Z}G$ with differentials ∂^C and ∂^D , respectively. For all $n \in \mathbb{Z}$, let $\mathcal{H}om_G(C_*, D_*)^n$ denote the set of graded G -module homomorphisms of degree $-n$ and define the boundary map δ^n by

$$\delta^n(f) = f \partial^C - (-1)^n \partial^D f.$$

Note that $\mathcal{H}om_G(-, C_*)$ (resp. $\mathcal{H}om_G(C_*, -)$) becomes a covariant (resp. contravariant) functor from the category of chain complexes of $\mathbb{Z}G$ -modules to the category of cochain complexes of abelian groups. Also, if C_* is a chain complex concentrated at 0 with $C_0 = M$, then $\mathcal{H}om_G(-, C_*)$ is naturally equivalent to the functor $\text{Hom}_G(-, M)$.

Now, we define the Tate hypercohomology of a finite group G with coefficients in C_* as

$$\hat{H}^i(G, C_*) := H^i(\mathcal{H}om_G(F_*, C_*))$$

for all $i \in \mathbb{Z}$, where F_* is a complete $\mathbb{Z}G$ -resolution of \mathbb{Z} . We immediately have $\hat{H}^i(G, \Sigma C_*) \cong \hat{H}^{i+1}(G, C_*)$, where $(\Sigma C_*)_i = C_{i-1}$ for all i . Therefore, if C_* is a chain complex concentrated at n , then $\hat{H}^i(G, C_*) \cong \hat{H}^{i+n}(G, C_n)$. Also note that given a short exact sequence of chain complexes

$$0 \rightarrow C_* \rightarrow D_* \rightarrow E_* \rightarrow 0$$

of $\mathbb{Z}G$ -modules, there is a long exact sequence of the following form

$$\dots \rightarrow \hat{H}^i(G, C_*) \rightarrow \hat{H}^i(G, D_*) \rightarrow \hat{H}^i(G, E_*) \rightarrow \hat{H}^{i+1}(G, C_*) \rightarrow \dots$$

An important property of $\mathcal{H}om$ functor is that if P_* is a chain complex of projective $\mathbb{Z}G$ -modules and $f_*: C_* \rightarrow D_*$ a weak equivalence of nonnegative chain complexes of $\mathbb{Z}G$ -modules, then $f_*: \mathcal{H}om_G(P_*, C_*) \rightarrow \mathcal{H}om_G(P_*, D_*)$ is also a weak equivalence (see Brown [5, page 29]). Actually, Brown proves this result by assuming P_* is nonnegative and C_* and D_* are arbitrary, but the same proof remains true if we assume P_* is arbitrary and C_* and D_* are nonnegative. Using this, we obtain the following proposition:

Proposition 2.1 *If C_* is a nonnegative chain complex of $\mathbb{Z}G$ -modules with homology concentrated at dimension n and $H_n(C_*) = M$, then $\hat{H}^i(G, C_*) \cong \hat{H}^{i+n}(G, M)$.*

Proof Let Z_n denote the group of n -cycles in C_* . We have the following weak equivalences:

$$\begin{array}{ccccccc} C_*: & \dots & \longrightarrow & C_{n+1} & \longrightarrow & C_n & \longrightarrow & C_{n-1} & \longrightarrow & \dots \\ & & & \uparrow & & \uparrow & & \uparrow & & \\ & & & \text{id} & & & & & & \\ D_*: & \dots & \longrightarrow & C_{n+1} & \longrightarrow & Z_n & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & \downarrow & & \downarrow & & \downarrow & & \\ E_*: & \dots & \longrightarrow & 0 & \longrightarrow & M & \longrightarrow & 0 & \longrightarrow & \dots \end{array}$$

Therefore, $\hat{H}^i(G, C_*) \cong \hat{H}^i(G, D_*) \cong \hat{H}^i(G, E_*) \cong \hat{H}^{i+n}(G, M)$. □

An exact sequence

$$K \xrightarrow{f} L \xrightarrow{g} M$$

of $\mathbb{Z}G$ -modules is called *admissible* if the inclusion map $\text{im}(g) \hookrightarrow M$ is \mathbb{Z} -split (see Brown [5, page 129]). A $\mathbb{Z}G$ -module M is called relatively injective if $\text{Hom}_G(-, M)$ takes an admissible exact sequence to an exact sequence of abelian groups. Projective $\mathbb{Z}G$ -modules are relatively injective (see Brown [5, page 130]). Since a complete $\mathbb{Z}G$ -resolution F_* of \mathbb{Z} is an exact sequence of free $\mathbb{Z}G$ -modules, the sequence $F_{i+1} \rightarrow F_i \rightarrow F_{i-1}$ is admissible for all i . Hence if P is a projective $\mathbb{Z}G$ -module, then the Tate cohomology group $\hat{H}^i(G, P) = 0$ for all i . This result generalizes to hypercohomology.

Proposition 2.2 *If P_* is a chain complex of projective $\mathbb{Z}G$ -modules which has finite length, then $\hat{H}^i(G, P_*) = 0$ for all i .*

Proof Recall that we say a chain complex C_* has finite length if there are integers n and m such that $C_i = 0$ for all $i > n$ and $i < m$. By shifting P_* if necessary, we can assume that P_* is a finite dimensional nonnegative chain complex and prove the proposition by an easy induction on the dimension of P_* . □

We say that two chain complexes C_* and D_* are *freely equivalent* if there is a sequence of chain complexes $C_* = E_*^0, \dots, E_*^n = D_*$ such that either E_*^i is an extension of E_*^{i-1} or E_*^{i-1} is an extension of E_*^i by a finite length chain complex of free modules. Note that we say a chain complex D_* is an extension of C_* by a finite length chain complex of free modules if there is short exact sequence of chain complexes either of the form $0 \rightarrow C_* \rightarrow D_* \rightarrow F_* \rightarrow 0$ or of the form $0 \rightarrow F_* \rightarrow D_* \rightarrow C_* \rightarrow 0$, where F_* is a finite length chain complex of free modules. As a corollary of Proposition 2.2, we have the following.

Corollary 2.3 *If two chain complexes C_* and D_* are freely equivalent, then*

$$\hat{H}^i(G, C_*) \cong \hat{H}^i(G, D_*)$$

for all i .

Before we conclude this section, we would like to note that there is a hypercohomology spectral sequence that converges to the Tate hypercohomology $\hat{H}^*(G, C_*)$ for a given chain complex C_* of $\mathbb{Z}G$ -modules. One way to obtain this spectral sequence is to consider the double complex $D^{p,q} = \text{Hom}_G(F_p, C_{-q})$ where the vertical and horizontal

differentials are given by $\delta_0 = \text{Hom}_G(-, \partial)$ and $\delta_1 = \text{Hom}_G(\partial, -)$. Note that the total complex $\text{Tot } D^{*,*}$ with

$$\text{Tot}^n D^{*,*} = \bigoplus_{p+q=n} D^{p,q}$$

and $\delta^n = \delta_0 - (-1)^n \delta_1$ is a cochain complex homotopy equivalent to the cochain complex $\mathcal{H}om_G(F_*, C_*)$. Filtering this double complex with respect to the index p and then with respect to the index q , we obtain two spectral sequences

$$\begin{aligned} \text{I } E_2^{p,q} &= \widehat{H}^p(G, H_{-q}(C_*)) \Rightarrow \widehat{H}^{p+q}(G, C_*), \\ \text{II } E_1^{p,q} &= \widehat{H}^q(G, C_{-p}) \Rightarrow \widehat{H}^{p+q}(G, C_*). \end{aligned}$$

Note that using these two spectral sequences it is possible to give alternative proofs for Propositions 2.1 and 2.2.

3 Habegger’s Theorem

In [9, pages 433–434], Habegger uses a technique to “glue” homology groups of a chain complex at different dimensions. This technique will be crucial in the proof of Theorem 1.2, so we give a proof for it here. Before we state Habegger’s Theorem, we recall the definition of syzygies of modules.

For every positive integer n , the n^{th} syzygy of a $\mathbb{Z}G$ -module M is defined as the kernel of ∂_{n-1} in a partial resolution of the form

$$P_{n-1} \xrightarrow{\partial_{n-1}} \cdots \rightarrow P_1 \xrightarrow{\partial_1} P_0 \rightarrow M \rightarrow 0$$

where P_0, \dots, P_{n-1} are projective $\mathbb{Z}G$ -modules. We denote the n^{th} syzygy of M by $\Omega^n M$ and by convention we take $\Omega^0 M = M$.

The n^{th} syzygy of a module M is well-defined only up to stable equivalence. Recall that two $\mathbb{Z}G$ -modules M and N are called stably equivalent if there are projective $\mathbb{Z}G$ -modules P and Q such that $M \oplus P \cong N \oplus Q$. Well-definedness of syzygies up to stable equivalence follows from a generalization of Schanuel’s Lemma (see Brown [5, page 193]). Since for any two stably equivalent modules M and N , we have $\widehat{H}^i(G, M) \cong \widehat{H}^i(G, N)$ for all i , we will ignore the fact that syzygies are well-defined only up to stable equivalence and treat $\Omega^n M$ as a unique module depending only on M and n . Alternatively, one can fix a resolution for every $\mathbb{Z}G$ -module M and define $\Omega^n M$ as the kernel of ∂_{n-1} in this unique resolution.

Theorem 3.1 (Habegger [9]) *Let C_* be a chain complex of $\mathbb{Z}G$ -modules and n, m are integers such that $m < n$. If $H_k(C_*) = 0$ for all k with $m < k < n$, then C_* is freely equivalent to a chain complex D_* such that:*

- (i) $H_i(D_*) = H_i(C_*)$ for every $i \neq n, m$,
- (ii) $H_m(D_*) = 0$, and
- (iii) there is an exact sequence of $\mathbb{Z}G$ -modules

$$0 \rightarrow H_n(C_*) \rightarrow H_n(D_*) \rightarrow \Omega^{n-m} H_m(C_*) \rightarrow 0.$$

Proof Let $F_{n-1} \rightarrow \dots \rightarrow F_m \rightarrow H_m(C_*) \rightarrow 0$ be an exact sequence where the F_i are free $\mathbb{Z}G$ -modules. Consider the following diagram

$$\begin{array}{cccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & F_{n-1} & \longrightarrow & \dots & \longrightarrow & F_m & \longrightarrow & H_m(C_*) & \longrightarrow & 0 & \longrightarrow & \dots \\ & & & & & & & & & & \text{id} \downarrow & & \downarrow & & \\ \dots & \longrightarrow & C_n & \longrightarrow & C_{n-1} & \longrightarrow & \dots & \longrightarrow & Z_m & \longrightarrow & H_m(C_*) & \longrightarrow & 0 & \longrightarrow & \dots \end{array}$$

where Z_m denotes the group of m -cycles in C_* . Since all F_i are projective and the bottom row has no homology below dimension n , the identity map extends to a chain map between rows. Notice that this chain map gives a chain map $f_*: F_* \rightarrow C_*$ as follows

$$\begin{array}{cccccccccccc} \dots & \longrightarrow & 0 & \longrightarrow & F_{n-1} & \longrightarrow & \dots & \longrightarrow & F_m & \longrightarrow & 0 & \longrightarrow & \dots \\ & & \downarrow & & f_{n-1} \downarrow & & & & f_m \downarrow & & \downarrow & & \\ \dots & \longrightarrow & C_n & \longrightarrow & C_{n-1} & \longrightarrow & \dots & \longrightarrow & C_m & \longrightarrow & C_{m-1} & \longrightarrow & \dots \end{array}$$

where the maps $f_i: F_i \rightarrow C_i$ for $i > m$ are the same as the maps in the first diagram above. The map $f_m: F_m \rightarrow C_m$ is defined as the composition

$$F_m \xrightarrow{f'_m} Z_m \hookrightarrow C_m$$

where $f'_m: F_m \rightarrow Z_m$ is the map defined as the lifting of the identity map in the first diagram.

Now, let D_* be the mapping cone of f_* . We have the following short exact sequence of the form

$$0 \rightarrow C_* \rightarrow D_* \rightarrow \Sigma F_* \rightarrow 0$$

so C_* is freely equivalent to D_* . The corresponding long exact sequence of homology groups is

$$\dots \rightarrow H_i(F_*) \xrightarrow{f_*} H_i(C_*) \rightarrow H_i(D_*) \rightarrow H_{i-1}(F_*) \rightarrow \dots$$

Assume first that $n > m + 1$. Then F_* has at least two terms and its homology is nonzero only at two dimensions $n - 1$ and m . So, $H_i(C_*) \cong H_i(D_*)$ for all i such that $i \neq m, m + 1, n - 1, n$. At dimension m , the map $f_*: H_m(F_*) \rightarrow H_m(C_*)$ is an

isomorphism, so we get $H_m(D_*) = H_{m+1}(D_*) = 0$. At dimension $n - 1$, we have $H_{n-1}(C_*) = 0$, so we get $H_{n-1}(D_*) = 0$. We also have a short exact sequence of the form

$$0 \rightarrow H_n(C_*) \rightarrow H_n(D_*) \rightarrow H_{n-1}(F_*) \rightarrow 0.$$

Since $H_{n-1}(F_*) \cong \Omega^{n-m}(H_m(C_*))$, this gives the desired result.

If $n = m + 1$, then F_* has a single term F_m , so we have a sequence of the form

$$0 \rightarrow H_n(C_*) \rightarrow H_n(D_*) \rightarrow F_m \xrightarrow{f_*} H_m(C_*) \rightarrow H_m(D_*) \rightarrow 0.$$

Since f_* is surjective by construction, we conclude that $H_m(D_*) = 0$ and there is a short exact sequence of the form

$$0 \rightarrow H_n(C_*) \rightarrow H_n(D_*) \rightarrow \Omega^1(H_m(C_*)) \rightarrow 0$$

as desired. □

4 Exponents of Tate cohomology groups

To prove the main theorem, we need some results about the exponents of Tate cohomology groups. We first recall some definitions. The exponent of a finite abelian group A is defined as the smallest positive integer n such that $na = 0$ for all $a \in A$. We denote the exponent of A by $\text{exp } A$. Note that if $A \rightarrow B \rightarrow C$ is an exact sequence of finite abelian groups, then $\text{exp } B$ divides $\text{exp } A \cdot \text{exp } C$. In this situation we sometimes write $\text{exp } B / \text{exp } A$ divides $\text{exp } C$ to refer to the same fact even though $\text{exp } B / \text{exp } A$ may not be an integer in general.

The first result we prove is a proposition on the exponent of Tate cohomology group with coefficients in a filtered module. First let us explain the terminology that we will be using throughout the paper. Let M be a $\mathbb{Z}G$ -module and A_1, A_2, \dots, A_n be a sequence of $\mathbb{Z}G$ -modules. If M has a filtration $0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_n = M$ such that $M_j / M_{j-1} \cong A_j$ for all j , then we say M has a filtration with sections $A_1 - A_2 - \dots - A_n$.

Proposition 4.1 *Let M be a $\mathbb{Z}G$ -module which has a filtration with sections $A_1 - A_2 - \dots - A_n$. Then, $\text{exp } \hat{H}^i(G, M)$ divides $\prod_{j=1}^n \text{exp } \hat{H}^i(G, A_j)$.*

Proof Let $0 = M_0 \subseteq M_1 \subseteq \dots \subseteq M_n = M$ be the filtration of M with the sections as above. Then for every j , we have an exact sequence of $\mathbb{Z}G$ -modules

$$0 \rightarrow M_{j-1} \rightarrow M_j \rightarrow A_j \rightarrow 0,$$

which gives a long exact Tate cohomology sequence of the following form:

$$\dots \rightarrow \widehat{H}^i(G, M_{j-1}) \rightarrow \widehat{H}^i(G, M_j) \rightarrow \widehat{H}^i(G, A_j) \rightarrow \dots .$$

From this we observe that

$$\exp \widehat{H}^i(G, M_j) / \exp \widehat{H}^i(G, M_{j-1})$$

divides the exponent of $\widehat{H}^i(G, A_j)$. Multiplying these relations through all $j = 1, \dots, n$, we get $\exp \widehat{H}^i(G, M)$ divides $\prod_{j=1}^n \exp \widehat{H}^i(G, A_j)$. □

In [4], Browder proves a theorem that gives an upper bound on the order of a finite group G in terms of the exponents of cohomology groups with coefficients in homology groups of a CW-complex on which G acts freely. Since we use this theorem in the proof of our main theorem, we state it below and give a proof for it. The proof we give here is slightly different than the original proof. It uses [Theorem 3.1](#) and [Proposition 4.1](#).

Theorem 4.2 (Browder [4]) *Let C_* be a nonnegative, free, connected chain complex of dimension n . Then $|G|$ divides $\prod_{j=1}^n \exp H^{j+1}(G, H_j(C_*))$.*

Proof Let us take $C_*^{(0)} = C_*$ and for $j = 1$ to n , define $C_*^{(j)}$ to be the chain complex obtained from $C_*^{(j-1)}$ by applying the method in [Theorem 3.1](#) for the dimensions $n - j$ and n . Since C_* is a finite dimensional chain complex of free $\mathbb{Z}G$ -modules, by [Proposition 2.2](#), $\widehat{H}^i(G, C_*) = 0$ for all i . Hence by [Corollary 2.3](#) and [Theorem 3.1](#), we have $\widehat{H}^i(G, C_*^{(j)}) = 0$ for all i, j . Notice that $C_*^{(n)}$ is a chain complex with homology concentrated at n . Let us denote the homology of $C_*^{(n)}$ at n by M . Hence, by [Proposition 2.1](#), we have $\widehat{H}^i(G, M) = 0$ for all i . By [Theorem 3.1](#), M has a filtration

$$0 \subseteq H_n(C_*^{(0)}) \subseteq \dots \subseteq H_n(C_*^{(n-1)}) \subseteq H_n(C_*^{(n)}) = M$$

with sections $H_n(C_*) - \Omega^1 H_{n-1}(C_*) - \dots - \Omega^{n-1} H_1(C_*) - \Omega^n H_0(C_*)$. If we let $M' := H_n(C_*^{(n-1)})$, then M' has a filtration with sections

$$H_n(C_*) - \Omega^1 H_{n-1}(C_*) - \dots - \Omega^{n-1} H_1(C_*)$$

and there is a short exact sequence of the form

$$0 \rightarrow M' \rightarrow M \xrightarrow{\pi} \Omega^n H_0(C_*) \rightarrow 0.$$

Note that $H_0(C_*) \cong \mathbb{Z}$, so we obtain an exact sequence of the form

$$\dots \rightarrow \widehat{H}^n(G, M) \xrightarrow{\pi_*} \widehat{H}^n(G, \Omega^n \mathbb{Z}) \rightarrow \widehat{H}^{n+1}(G, M') \rightarrow \widehat{H}^{n+1}(G, M) \rightarrow \dots .$$

Since $\widehat{H}^i(G, M) = 0$ for all i , we obtain

$$\widehat{H}^{n+1}(G, M') \cong \widehat{H}^n(G, \Omega^n \mathbb{Z}) \cong \widehat{H}^0(G, \mathbb{Z}) \cong \mathbb{Z}/|G|.$$

Hence by Proposition 4.1, we get $|G| = \exp \widehat{H}^{n+1}(G, M')$ divides the product

$$\prod_{j=1}^n \exp \widehat{H}^{n+1}(G, \Omega^{n-j} H_j(C_*)).$$

Since $\widehat{H}^{n+1}(G, \Omega^{n-j} H_j(C_*)) \cong H^{j+1}(G, H_j(C_*))$, this gives the desired result. \square

As a corollary of Theorem 4.2, Browder gives a proof for a theorem of G Carlsson [6] that says that if $G = (\mathbb{Z}/p)^r$ acts freely on a finite dimensional CW-complex $X \simeq (S^n)^k$ with trivial action on homology, then $r \leq k$. The main observation is that when $G = (\mathbb{Z}/p)^r$ and M is a trivial $\mathbb{Z}G$ -module, the exponent of $H^i(G, M)$ divides p for all $i \geq 1$. This follows easily by induction on r using properties of the transfer map in group cohomology. So, from the relation given in Theorem 4.2, one obtains that if G acts freely on a finite dimensional CW-complex $X \simeq (S^n)^k$ with trivial action on homology, then $|G| = p^r$ divides p^k , which gives $r \leq k$.

Note that the assumption that G acts trivially on the homology of X is crucial in the above argument since for an arbitrary $\mathbb{Z}G$ -module, the exponent of $H^i(G, M)$ can be as large as the order of $|G|$. In fact, if we take $M = \Omega^i(\mathbb{Z})$ for some positive integer i , then we have $H^i(G, M) \cong \mathbb{Z}/|G|$, so the exponent of $H^i(G, M)$ is equal to $|G|$ in this case. Taking the direct sum of all such modules over all i , one can obtain a $\mathbb{Z}G$ -module M such that the exponent of $H^i(G, M)$ is equal to $|G|$ for every $i \geq 0$. The following theorem says that when M is finitely generated this situation cannot happen and that the exponent of $H^i(G, M)$ eventually becomes small at high dimensions.

Theorem 4.3 (Pakianathan [11]) *Let $G = (\mathbb{Z}/p)^r$ and M be a finitely generated $\mathbb{Z}G$ -module. Then, there is an integer N such that the exponent of $H^i(G, M)$ divides p for all $i \geq N$.*

Proof By Theorem 7.4.1 in Evens [8, page 87], $H^*(G, M)$ is a finitely generated module over the ring $H^*(G, \mathbb{Z})$. Let u_1, \dots, u_k be homogeneous elements generating $H^*(G, M)$ as an $H^*(G, \mathbb{Z})$ -module and let $N = 1 + \max_j \{\deg u_j\}$. If $u \in H^i(G, M)$ such that $i \geq N$, then we can write $u = \sum_{j=1}^k \alpha_j u_j$ for some homogeneous elements α_j in $H^*(G, \mathbb{Z})$ with $\deg \alpha_j \geq 1$ for all j . Since $\exp H^i(G, \mathbb{Z})$ divides p for all $i \geq 1$, we have $p\alpha_j = 0$ for all j . Hence we obtain $pu = \sum_{j=1}^k p\alpha_j u_j = 0$ as desired. \square

5 Proof of the main theorem

Let $G = (\mathbb{Z}/p)^r$ and k, l be positive integers. We will show that there is an integer N such that if G acts freely and cellularly on a CW-complex X homotopy equivalent to $S^{n_1} \times \cdots \times S^{n_k}$ where $|n_i - n_j| \leq l$ and $n_i \geq N$ for all i, j , then $r \leq k$.

Suppose that G acts freely and cellularly on some CW-complex X homotopy equivalent to $S^{n_1} \times \cdots \times S^{n_k}$ where $|n_i - n_j| \leq l$ for all i, j . Let $n = \max\{n_i \mid i = 1, \dots, k\}$ and let $a_i = n - n_i$ for all i . Consider the cellular chain complex $C_*(X)$ of the CW-complex X . The complex $C_*(X)$ is a nonnegative, connected, and finite-dimensional chain complex of free $\mathbb{Z}G$ -modules and has nonzero homology only at the following dimensions other than dimension zero:

- (1) $n - a_1, n - a_2, \dots, n - a_k$
- (2) $2n - a_1 - a_2, 2n - a_1 - a_3, \dots, 2n - a_{k-1} - a_k$
- \vdots
- (j) $jn - (a_1 + \cdots + a_j), \dots, jn - (a_{k-j+1} + \cdots + a_k)$
- \vdots
- (k) $kn - (a_1 + a_2 + \cdots + a_k)$

If $n > lk$, then we have $n > a_1 + \cdots + a_k$, which implies that for all j , the dimensions listed on the j^{th} row are strictly larger than the dimensions listed on the $(j - 1)^{\text{st}}$ row. Since this fact is crucial for our argument, we will assume that the integer N in the statement of the theorem satisfies $N > lk$ to guarantee that this condition holds.

Now we can apply Habegger’s argument given in [Theorem 3.1](#) to glue all the homology groups at the dimensions listed on the j^{th} row above to the homology at dimension jn for all $j = 1, \dots, k$. The resulting complex D_* is a connected, finite-dimensional chain complex of free $\mathbb{Z}G$ -modules that has homology only at dimensions $0, n, 2n, \dots, kn$. Let $M_j := H_{jn}(D_*)$ for all $j = 1, \dots, k$. Note that by construction M_j is a finitely generated $\mathbb{Z}G$ -module for all j since syzygies of finitely generated $\mathbb{Z}G$ -modules are finitely generated when G is a finite group.

Now we can apply [Theorem 4.3](#) to find an integer N_j for each j such that if $i \geq N_j$, then $\exp H^i(G, M_j)$ divides p . Suppose that for a fixed $G = (\mathbb{Z}/p)^r$, k and l , there are only finitely many possibilities for $\mathbb{Z}G$ -modules M_j up to stable equivalence. Then by taking the maximum of the N_j over all possible M_j , we can find an integer N_j^{\max} for each j such that if $i \geq N_j^{\max}$, then $\exp H^i(G, M_j)$ divides p for all possible M_j that may occur. Then we can take $N = \max_j N_j^{\max}$ and complete the proof in the

following way. By [Theorem 4.2](#), we have $|G| = p^r$ divides

$$\prod_{j=1}^k H^{jn+1}(G, H_{jn}(D_*)) = \prod_{j=1}^k H^{jn+1}(G, M_j).$$

So, if $n \geq N$, then p^r divides p^k , which gives $r \leq k$ as desired.

Hence to complete the proof, it only remains to show that for fixed $G = (\mathbb{Z}/p)^r$, k and l , there are only finitely many possibilities for $\mathbb{Z}G$ -modules M_j up to stable equivalence. To show this, first note that for a fixed l , there are finitely many k -tuples (a_1, \dots, a_k) with the property that $0 \leq a_i \leq l$ for all i . So we can assume that we have a fixed k -tuple (a_1, \dots, a_k) . Let us also fix an integer j and show there are only finitely many possibilities for $M_j = H_{jn}(D_*)$.

Let $s_1 < \dots < s_m$ be a sequence of integers such that $\{jn - s_1, \dots, jn - s_m\}$ is the set of all distinct dimensions on the j^{th} row of the above diagram. Note that the complex D_* is constructed with the repeated usage of [Theorem 3.1](#), so the module $M_j = H_{jn}(D_*)$ has a filtration

$$0 = K_0 \subseteq K_1 \subseteq \dots \subseteq K_m = M_j$$

such that $K_i/K_{i-1} \cong \Omega^{s_i}(A_i)$ where $A_i = H_{jn-s_i}(X)$. For all i , the module A_i is a \mathbb{Z} -free $\mathbb{Z}G$ -module with \mathbb{Z} -rank less than or equal to $\binom{k}{j}$, so by the Jordan-Zassenhaus Theorem (see Corollary 79.12 in Curtis and Reiner [\[7, page 563\]](#)), there are only finitely many possibilities for A_i up to isomorphism.

We will inductively show that there exist only finitely many possibilities for K_i up to stable equivalence. For $i = 1$, we have $K_1 = \Omega^{s_1}(A_1)$ so this follows from the fact that there are only finitely many possibilities for A_1 and that syzygies are well-defined up to stable equivalence. For $i > 1$, consider the following short exact sequence:

$$0 \rightarrow K_{i-1} \rightarrow K_i \rightarrow \Omega^{s_i} A_i \rightarrow 0$$

By induction we know that there are only a finite number of possibilities for K_{i-1} up to stable equivalence. By a similar argument as above, the same is true for $\Omega^{s_i}(A_i)$. The extensions like the ones above are classified by the ext-group $\text{Ext}_{\mathbb{Z}G}^1(\Omega^{s_i}(A_i), K_{i-1})$ and since both modules are \mathbb{Z} -free, these ext-groups are well-defined up to stable equivalence. So, it remains to show that

$$\text{Ext}_{\mathbb{Z}G}^1(\Omega^{s_i}(A_i), K_{i-1}) = \text{Ext}_{\mathbb{Z}G}^{s_i+1}(A_i, K_{i-1})$$

is a finite group. Note that since both A_i and K_{i-1} are finitely generated,

$$\text{Ext}_{\mathbb{Z}G}^{s_i+1}(A_i, K_{i-1})$$

is a finitely generated abelian group. Moreover, since A_i is \mathbb{Z} -free, it has an exponent divisible by $|G|$. So, $\text{Ext}_{\mathbb{Z}G}^{s_i+1}(A_i, K_{i-1})$ is a finite group. This completes the proof of [Theorem 1.2](#).

We conclude this section with a generalization of [Theorem 1.2](#) to non-free actions. The exact statement is as follows.

Theorem 5.1 *Suppose $G = (\mathbb{Z}/p)^r$ for a prime p and k, l are positive integers. Then there is an integer N that depends only on k, l and G such that if G acts cellularly on a finite dimensional CW-complex X homotopy equivalent to $S^{n_1} \times \dots \times S^{n_k}$ where $n_i \geq N$ and $|n_i - n_j| \leq l$ for all i, j , then $r - s \leq k$ where s is the largest integer such that $|G_x| = p^s$ for some $x \in X$.*

Proof Let $C_* := C_*(X)$ denote the cellular chain complex of X and let $\varepsilon: C_* \rightarrow \mathbb{Z}$ be the map induced by the constant map $X \rightarrow pt$. The arguments in the proof of [Theorem 1.2](#) can be repeated to prove that there is an integer N such that if $n_i \geq N$ and $|n_i - n_j| \leq l$ for all i, j , then

$$p^k \hat{H}^0(G, \mathbb{Z}) \subseteq \text{im}\{\varepsilon_*: \hat{H}^0(G, C_*) \rightarrow \hat{H}^0(G, \mathbb{Z})\}.$$

This can be seen by a spectral sequence argument or by the filtration argument given in the proof of [Theorem 4.2](#). To see it using the filtration argument, observe that the map ε_* can be written as a composition

$$\varepsilon_*: \hat{H}^0(G, C_*) \cong \hat{H}^0(G, C_*^{(n)}) \cong H^n(G, M) \xrightarrow{\pi_*} \hat{H}^n(G, \Omega^n \mathbb{Z}) \cong \hat{H}^0(G, \mathbb{Z})$$

where the module M and the map π_* are as given in the proof of [Theorem 4.2](#). Repeating the arguments in the proof of [Theorem 1.2](#), we can show that there is an integer N such that if $n_i \geq N$ and $|n_i - n_j| \leq l$ for all i, j , then $\exp \hat{H}^{n+1}(G, M')$ divides p^k where M' is as in the proof of [Theorem 4.2](#). Using the long exact sequence given in the proof of [Theorem 4.2](#), we can conclude that $p^k \hat{H}^0(G, \mathbb{Z}) \subseteq \text{im } \varepsilon_*$.

The inclusion given above implies that $|G| = p^r$ divides $p^k \cdot \exp \hat{H}^0(G, C_*)$. Hence the proof will be complete if we can show that $\exp \hat{H}^0(G, C_*)$ divides p^s where s is the largest integer such that $|G_x| = p^s$ for some $x \in X$. However this is already known to be true as proven by A Adem [[1](#), Theorems 3.1 and 3.2]. So the proof is complete. □

Acknowledgements

We thank Jonathan Pakianathan and Alejandro Adem for many helpful conversations on the paper and the referee for a careful reading of the paper and for his/her corrections and helpful comments. The second author is partially supported by TÜBİTAK-TBAG/110T712.

References

- [1] **A Adem**, *Torsion in equivariant cohomology*, Comment. Math. Helv. 64 (1989) 401–411 [MR998857](#)
- [2] **A Adem, W Browder**, *The free rank of symmetry of $(S^n)^k$* , Invent. Math. 92 (1988) 431–440 [MR936091](#)
- [3] **D J Benson, J F Carlson**, *Complexity and multiple complexes*, Math. Z. 195 (1987) 221–238 [MR892053](#)
- [4] **W Browder**, *Cohomology and group actions*, Invent. Math. 71 (1983) 599–607 [MR695909](#)
- [5] **K S Brown**, *Cohomology of groups*, Graduate Texts in Mathematics 87, Springer, New York (1982) [MR672956](#)
- [6] **G Carlsson**, *On the rank of abelian groups acting freely on $(S^n)^k$* , Invent. Math. 69 (1982) 393–400 [MR679764](#)
- [7] **C W Curtis, I Reiner**, *Representation theory of finite groups and associative algebras*, Pure and Applied Mathematics 11, Interscience Publishers, a division of John Wiley & Sons, New York (1962) [MR0144979](#)
- [8] **L Evens**, *The cohomology of groups*, Oxford Mathematical Monographs, The Clarendon Press, New York (1991) [MR1144017](#)
- [9] **N Habegger**, *Hypercohomology varieties for complexes of modules, the realizability criterion, and equivalent formulations of a conjecture of Carlsson*, from: “The Arcata conference on representations of finite groups”, (P Fong, editor), Proc. Sympos. Pure Math. 47, Amer. Math. Soc. (1987) 431–437 [MR933380](#)
- [10] **B Hanke**, *The stable free rank of symmetry of products of spheres*, Invent. Math. 178 (2009) 265–298 [MR2545682](#)
- [11] **J Pakianathan**, *private communication* (2012)
- [12] **P A Smith**, *Permutable periodic transformations*, Proc. Nat. Acad. Sci. U. S. A. 30 (1944) 105–108 [MR0010278](#)
- [13] **R G Swan**, *Periodic resolutions for finite groups*, Ann. of Math. 72 (1960) 267–291 [MR0124895](#)
- [14] **E Yalçın**, *Group actions and group extensions*, Trans. Amer. Math. Soc. 352 (2000) 2689–2700 [MR1661282](#)

Department of Mathematics, The Ohio State University, Columbus, OH 43210-1174, USA

Department of Mathematics, Bilkent University, 06800 Ankara, Turkey

okutan.1@osu.edu, yalcine@fen.bilkent.edu.tr

<http://www.math.osu.edu/people/okutan.1>,

<http://www.fen.bilkent.edu.tr/~yalcine/>

Received: 17 October 2012 Revised: 13 February 2013