

Betti numbers and stability for configuration spaces via factorization homology

BEN KNUDSEN

Using factorization homology, we realize the rational homology of the unordered configuration spaces of an arbitrary manifold M , possibly with boundary, as the homology of a Lie algebra constructed from the compactly supported cohomology of M . By locating the homology of each configuration space within the Chevalley–Eilenberg complex of this Lie algebra, we extend theorems of Bödiger, Cohen and Taylor and of Félix and Thomas, and give a new, combinatorial proof of the homological stability results of Church and Randal-Williams. Our method lends itself to explicit calculations, examples of which we include.

57R19; 17B56, 55R80

1 Introduction

We study the configuration space $B_k(M)$ of k unordered points in a manifold M , defined as

$$B_k(M) = \text{Conf}_k(M)_{\Sigma_k} := \{(x_1, \dots, x_k) \in M^k \mid x_i \neq x_j \text{ for } i \neq j\} / \Sigma_k,$$

where the permutation group Σ_k acts by permuting the x_i . Our main theorem concerns the homology of these spaces.

Theorem 1.1 *Let M be an n -manifold. There is an isomorphism of bigraded vector spaces*

$$\bigoplus_{k \geq 0} H_*(B_k(M); \mathbb{Q}) \cong H^{\mathcal{L}}(H_c^{-*}(M; \mathcal{L}(\mathbb{Q}^w[n-1]))).$$

Here H_c^{-*} denotes compactly supported cohomology, \mathbb{Q}^w is the orientation sheaf of M , $H^{\mathcal{L}}$ denotes Lie algebra homology and \mathcal{L} is the free graded Lie algebra functor. The auxiliary grading on the left is by cardinality of the configuration and on the right by powers of the Lie generator.

Our methods apply equally to the calculation of the twisted homology of configuration spaces and of the homology of certain relative configuration spaces defined for manifolds

with boundary; precise statements may be found in Theorem 4.5 and Theorem 4.9, respectively. All results and arguments herein are valid over an arbitrary field of characteristic zero.

The study of configuration spaces is classical. To name some highlights, the space $B_k(\mathbb{R}^2)$ is a classifying space for the braid group on k strands (see Artin [2]); the space $\text{Conf}_k(\mathbb{R}^n)$ has the homotopy type of the space of k -ary operations of the little n -cubes operad and so plays a central role in the theory of n -fold loop spaces (see eg Cohen, Lada and May [17], May [43] and Segal [51]); certain spaces of labeled configurations provide models for more general types of mapping spaces (see Bödigheimer [8], McDuff [44], Salvatore [48], Segal [51]); and, according to a striking theorem of Longoni and Salvatore [39], the homotopy type of $B_k(M)$ is not an invariant of the homotopy type of M .

As this last fact indicates, configuration spaces depend in subtle ways on the structure of the background manifold. On the other hand, the homology of these spaces has often been shown to be surprisingly simple, provided one is willing to work over a field of characteristic zero. Indeed, Bödigheimer, Cohen and Taylor [10] show that the Betti numbers of $B_k(M)$ are determined by those of M when M is of odd dimension, and Félix and Thomas [24] show that, in the even-dimensional case, the Betti numbers of $B_k(M)$ are determined by the rational cohomology ring of M , as long as M is closed, orientable and nilpotent. We recover extensions of these results as immediate consequences of Theorem 1.1.

Corollary 1.2 *The groups $H_*(B_k(M); \mathbb{Q})$ depend only on n and*

- *the graded abelian group $H_*(M; \mathbb{Q})$ if n is odd, or*
- *the cup product $H_c^{-*}(M; \mathbb{Q}^w)^{\otimes 2} \rightarrow H_c^{-*}(M; \mathbb{Q})$ if n is even.*

The computational power of Theorem 1.1 lies in the bigrading, which permits one to isolate the homology of a single configuration space within the Chevalley–Eilenberg complex computing the appropriate Lie homology. Employing this strategy, we show that the chain complexes computing $H_*(B_k(M); \mathbb{Q})$ exhibited in [10] and [24] are isomorphic to subcomplexes of the Chevalley–Eilenberg complex; precise statements appear in Section 4.3. Better yet, in dealing with the entire Chevalley–Eilenberg complex at once, one is able to perform computations for all k simultaneously; see Section 6.

Another important aspect of the study of configuration spaces is the phenomenon of *homological stability*. As k tends to infinity, the Betti numbers of $B_k(M)$ are eventually constant, despite the absence of a map of spaces $B_k(M) \rightarrow B_{k+1}(M)$ in

general; see Church [14], Church, Eilenburg and Farb [15], Randall-Williams [46] and Cantero and Palmer [13]. Here too, characteristic zero is special.

Regarding stability, we prove the following.

Theorem 1.3 *Let M be a connected n -manifold with $n > 1$. The cap product with the unit in $H^0(M; \mathbb{Q})$ induces a map*

$$H_*(B_{k+1}(M); \mathbb{Q}) \rightarrow H_*(B_k(M); \mathbb{Q})$$

that is

- an isomorphism for $* < k$ and a surjection for $* = k$ when M is an orientable surface, and
- an isomorphism for $* \leq k$ and a surjection for $* = k + 1$ in all other cases.

The sense in which the homology of configuration spaces forms a coalgebra, so that the cap product is defined, will be explained in Section 5. We lack a conceptual explanation for the exceptional behavior in dimension 2, as it emerges from our argument solely as a numerical/combinatorial coincidence.

This result improves on the stable range of Church [14] and very slightly on that of Randal-Williams [46]. As in the former work, our stable range can be further improved if the low-degree Betti numbers of M vanish. As the example of the Klein bottle shows, the bound $* \leq k$ is sharp in the sense that no better stable range holds for all manifolds that are not orientable surfaces. When M is open, the surjectivity statement is proven in [46]; to the author's knowledge, the result is new for compact manifolds.

Conceptually, we think of Theorem 1.1 as providing an explanation and organizing principle for the behavior of configuration spaces in characteristic zero. The germ of our approach, and the source of the connection to Lie algebras, is the calculation, due to Arnol'd and Cohen, of the homology of the ordered configuration spaces of \mathbb{R}^n , which is the fundamental result of the subject; see Arnol'd [1] and Cohen, Lada and May [17]. Specifically, for $n \geq 2$, the homology groups of the spaces $\text{Conf}_k(\mathbb{R}^n)$ form a shifted version of the operad governing Poisson algebras, with the shifted Lie bracket given by the fundamental class of $\text{Conf}_2(\mathbb{R}^n) \simeq S^{n-1}$; see Sinha [52] for a beautiful geometric discussion of this identification. Locally, then, configuration spaces enjoy a rich algebraic structure; factorization homology, our primary tool in this work, provides a means of assembling this structure across coordinate patches of a general manifold, globalizing the calculation of Arnol'd and Cohen. Theorem 1.1 is the natural output of this procedure.

At a more formal level, we rely on the fact that the factorization homology of M , with coefficients taken in a certain free algebra, can be computed in two different ways.

On the one hand, according to Proposition 3.1, it has an expression in terms of the configuration spaces of M . On the other hand, the free algebra may be thought of as a kind of enveloping algebra, and a calculation of the author's in [35] identifies the same invariant as Lie algebra homology. On the face of it, these calculations only coincide for framed manifolds; we show that they agree in general in characteristic zero.

In keeping with our metamathematical goal of making the case for factorization homology as a computational tool, we do not focus on the technical underpinnings of the theory. The interested reader may find these in Ayala and Francis [5; 4; 3], Ayala, Francis and Tanaka [7; 6], Francis [25] and Lurie [42].

The paper is split into seven sections. In Sections 2–3, we review the basics of factorization homology and discuss calculations thereof in several cases of interest. Theorem 1.1 and its variants are proved in Section 4 assuming several deferred results, and the classical results alluded to above follow. In Section 5, we discuss coalgebraic phenomena arising from configuration spaces, which lead us to the proof of Theorem 1.3 and one of the missing ingredients in the main theorem. Finally, Section 6 is concerned with explicit computations, and Section 7 supplies the remaining missing ingredients.

Conventions (1) In accordance with the bulk of the literature on factorization homology, we work in an ∞ -categorical context, where for us an ∞ -category will always mean a quasicategory. The standard references here are Lurie [40; 42], but we will need to ask only very little of the vast theory developed therein, and the reader may obtain a sense of the arguments and results by substituting “homotopy colimit” for “colimit” everywhere, for example.

(2) Every manifold is smooth and may be embedded as the interior of a compact manifold with boundary (such an embedding is not part of the data). We view manifolds as objects of the ∞ -category \mathcal{Mfld}_n , the topological nerve of the topological category of n -manifolds and smooth embeddings, which is symmetric monoidal under disjoint union.

(3) Our homology theories are valued in $\mathcal{Ch}_{\mathbb{Q}}$, the underlying ∞ -category of the category of \mathbb{Q} -chain complexes equipped with the standard model structure. With the single exception of Theorem 2.1, $\mathcal{Ch}_{\mathbb{Q}}$ is understood to be symmetric monoidal under tensor product.

(4) The homology of a chain complex V is written $H(V)$, while the homology of a space X is written $H_*(X)$. Hence $H_*(X) = H(C_*(X))$. If \mathfrak{g} is a differential graded Lie algebra, then $H(\mathfrak{g})$ is a graded Lie algebra.

(5) Chain complexes are homologically graded. If V is a chain complex, $V[k]$ is the chain complex with $(V[k])_n = V_{n-k}$, and, for $x \in V$, the corresponding element

in $V[k]$ is denoted $\sigma^k x$. Cohomology is concentrated in negative degrees; to reinforce this point, we write $H^{-*}(X)$ for the graded vector space whose degree $-k$ part is the k^{th} cohomology group of X ; for example,

$$H^{-*}(S^n; \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & \text{if } * \in \{-n, 0\}, \\ 0 & \text{otherwise.} \end{cases}$$

(6) If X is a space and V is a chain complex, the *tensor of X with V* is the chain complex

$$X \otimes V := C_*(X) \otimes V.$$

(7) If (X, A) is a pair of spaces, the quotient of X by A is the pointed space X/A defined as the pushout in the following diagram:

$$\begin{array}{ccc} A & \longrightarrow & X \\ \downarrow & & \downarrow \\ \text{pt} & \longrightarrow & X/A \end{array}$$

In particular, we have $X/\emptyset = X_+$.

(8) If X is an object of the ∞ -category \mathcal{C} with an action of the group G , then X_G and X^G denote the G -coinvariants and G -invariants of X , respectively, which are objects of \mathcal{C} . When \mathcal{C} is topological spaces or chain complexes, this object coincides in the homotopy category with *homotopy* coinvariants.

Acknowledgments I am grateful to John Francis for suggesting this project, and for his help and patience.

I would like to thank Lee Cohn, Elden Elmanto, Boris Hanin, Sander Kupers, Jeremy Miller, Martin Palmer and Dylan Wilson for their comments on earlier versions of this paper, Joel Specter and Eric Wofsey for suggested examples, Bruno Vallette for an enlightening lecture, and Jordan Ellenberg and Aaron Mazel-Gee for crucial conversations leading to significant improvements of the results.

After writing the paper, I learned that some of the same results are accessible through the work of Ezra Getzler by first passing to symmetric group invariants in the complex constructed in Getzler [28; 29], and then invoking Proposition 7.6 below. I am grateful to Dan Petersen for bringing these papers to my attention.

This paper is derived from my PhD thesis [36]. Revision was undertaken during visits to the Mathematisches Forschungsinstitut Oberwolfach and the Hausdorff Research Institute for Mathematics in Bonn.

Finally, I would like to thank the referees for their comments.

2 Factorization homology

2.1 Homology theories

In this section, we review the basic notions of factorization homology, also known as topological chiral homology. The primary reference is [5]. As there, our point of view is that factorization homology is a natural theory of homology for manifolds. To illustrate in what sense this is so, we first recall the classical characterization of ordinary homology, phrased in a way that invites generalization.

Theorem 2.1 (Eilenberg–Steenrod axioms) *Let V be a chain complex. There is a symmetric monoidal functor $C_*(-; V)$ from spaces with disjoint union to chain complexes with direct sum, called **singular homology with coefficients in V** , which is characterized up to natural equivalence by the following properties:*

- (1) $C_*(\text{pt}; V) \simeq V$;
- (2) *the natural map*

$$C_*(X_1; V) \bigoplus_{C_*(X_0; V)} C_*(X_2; V) \rightarrow C_*(X; V)$$

is an equivalence, where X is the pushout of the diagram of cofibrations

$$X_1 \hookrightarrow X_0 \hookrightarrow X_2.$$

Property (2), a local-to-global principle equivalent to the usual excision axiom, is the reason that homology is computable and hence useful.

Of course, ordinary homology is a homotopy invariant. In the study of manifolds, the equivalence relation of interest is often finer than homotopy equivalence, and one could hope for a theory better suited to such geometric investigations. To discover what form this theory might take, let us contemplate a generic symmetric monoidal functor $(\text{Mfld}_n, \sqcup) \rightarrow (\text{Ch}_{\mathbb{Q}}, \otimes)$. By analogy with Theorem 2.1, we ask that this functor be determined by its value on \mathbb{R}^n , the basic building block in the construction of n -manifolds. Unlike a point, however, Euclidean space has interesting internal structure.

Definition 2.2 An n -disk algebra in $\text{Ch}_{\mathbb{Q}}$ is a symmetric monoidal functor

$$A: (\text{Disk}_n, \sqcup) \rightarrow (\text{Ch}_{\mathbb{Q}}, \otimes),$$

where $\text{Disk}_n \subseteq \text{Mfld}_n$ is the full subcategory spanned by manifolds diffeomorphic to $\bigsqcup_k \mathbb{R}^n$ for some $k \in \mathbb{Z}_{\geq 0}$.

In other words, Disk_n is the (nerve of the) category of operations associated to the endomorphism operad of the manifold \mathbb{R}^n , and an n -disk algebra is an algebra over this operad. In contrast, the endomorphism operad of a point in topological spaces is the commutative operad, and every chain complex is canonically and essentially uniquely a commutative algebra in $(\text{Ch}_{\mathbb{Q}}, \oplus)$.

Taking the extra structure of \mathbb{R}^n into account, [5, Theorem 3.24] provides an analogous classification theorem.

Theorem 2.3 (Ayala and Francis) *Let A be an n -disk algebra. There is a symmetric monoidal functor $\int_{(-)} A$ from n -manifolds with disjoint union to chain complexes with tensor product, called **factorization homology with coefficients in A** , which is characterized up to natural equivalence by the following properties:*

- (1) $\int_{\mathbb{R}^n} A \simeq A$ as n -disk algebras;
- (2) the natural map

$$\int_{M_1} A \otimes_{\int_{M_0 \times \mathbb{R}} A} \int_{M_2} A \rightarrow \int_M A$$

is an equivalence, where M is obtained as the collar-gluing of the diagram of embeddings $M_1 \hookrightarrow M_0 \times \mathbb{R} \hookrightarrow M_2$.

Just as the functor of singular chains is but one model for ordinary homology, factorization homology may be constructed in several equivalent ways. The construction that we will favor is as follows.

Let $A: \text{Disk}_n \rightarrow \text{Ch}_{\mathbb{Q}}$ be an n -disk algebra. Then factorization homology with coefficients in A is the left Kan extension in the following diagram of ∞ -categories:

$$\begin{array}{ccc} \text{Disk}_n & \xrightarrow{A} & \text{Ch}_{\mathbb{Q}} \\ \downarrow & \nearrow \int_{(-)} A & \\ \text{Mfld}_n & & \end{array}$$

Explicitly, it may be calculated as the colimit

$$\int_M A \simeq \text{colim}(\text{Disk}_n/M \rightarrow \text{Disk}_n \xrightarrow{A} \text{Ch}_{\mathbb{Q}}).$$

Remark 2.4 Since $\text{Ch}_{\mathbb{Q}}$ admits sifted colimits and \otimes distributes over them, Theorem 3.2.3 of [4] guarantees that the left Kan extension and the symmetric monoidal left Kan extension exist and coincide.

2.2 Variant: framed manifolds

The category Disk_n is closely related to the classical operad E_n of little n -cubes. To make this connection, we recall that a *framing* of an n -manifold M is a nullhomotopy of its tangent classifier

$$M \xrightarrow{TM \simeq *} BO(n).$$

With the corresponding notion of *framed embedding* between framed manifolds in hand, one obtains an ∞ -category $\mathcal{M}\text{fld}_n^{\text{fr}}$ of framed n -manifolds; see [5, Definition 2.7].

Definition 2.5 A *framed n -disk algebra* in $\text{Ch}_{\mathbb{Q}}$ is a symmetric monoidal functor $A: (\text{Disk}_n^{\text{fr}}, \sqcup) \rightarrow (\text{Ch}_{\mathbb{Q}}, \otimes)$, where $\text{Disk}_n^{\text{fr}} \subseteq \mathcal{M}\text{fld}_n^{\text{fr}}$ is the full subcategory spanned by framed manifolds diffeomorphic to $\bigsqcup_k \mathbb{R}^n$ for some $k \in \mathbb{Z}_{\geq 0}$.

As before, the factorization homology of a framed n -manifold with coefficients in a framed n -disk algebra is defined as the left Kan extension from $\text{Disk}_n^{\text{fr}}$. Indeed, the whole theory carries over into the context of topological manifolds equipped with a microtangential B -structure arising from a map $B \rightarrow B\text{Top}(n)$. In this paper, we will only make use of the cases $B = BO(n)$, corresponding to smooth manifolds (see [5, Example 2.11 and Remark 3.29]), and $B = *$, corresponding to framed manifolds.

Now, the topological operad E_n has an associated ∞ -operad (see [42, Section 2.1]), and [6, Example 2.11] asserts an equivalence

$$\text{Alg}_{\text{Disk}_n^{\text{fr}}}(\mathcal{C}) \xrightarrow{\sim} \text{Alg}_{E_n}(\mathcal{C})$$

for any symmetric monoidal ∞ -category \mathcal{C} . Moreover, this equivalence induces a further equivalence

$$\text{Alg}_{\text{Disk}_n}(\mathcal{C}) \xrightarrow{\sim} \text{Alg}_{E_n}(\mathcal{C})^{O(n)}.$$

Informally, an n -disk algebra is an E_n -algebra with an action of $O(n)$ compatible with the action on E_n given by rotating disks. In the language of [49], n -disk algebras are algebras for the *semidirect product* $E_n \rtimes O(n)$.

Remark 2.6 The reader is cautioned not to confuse the framed n -disk algebras employed here with the “framed E_n -algebras” that occur elsewhere in the literature. These algebras carry an action of $SO(n)$ and yield homology theories for *oriented* manifolds.

2.3 Free algebras


We introduce several functors that will be important for us in what follows. The reference here is [3].

Within the ∞ -category Disk_n there is a Kan complex with a single vertex, the object \mathbb{R}^n , whose endomorphisms are $\text{Emb}(\mathbb{R}^n, \mathbb{R}^n) \simeq O(n)$, so that we may identify this Kan complex with $BO(n)$. Restricting to this subcategory defines a forgetful functor

$$\text{Alg}_{\text{Disk}_n}(\text{Ch}_{\mathbb{Q}}) \rightarrow \text{Fun}(BO(n), \text{Ch}_{\mathbb{Q}}) \xrightarrow{\sim} \text{Mod}_{O(n)}(\text{Ch}_{\mathbb{Q}}).$$

The latter symbol denotes the ∞ -category of chain complexes equipped with an action of $C_*(O(n); \mathbb{Q})$, which we refer to simply as $O(n)$ -modules. This functor admits a left adjoint \mathbb{F}_n , the free n -disk algebra generated by an $O(n)$ -module.

Evaluation on \mathbb{R}^n defines a still more forgetful functor, which we think of as associating to an algebra its underlying chain complex. The situation is summarized in the following commuting diagram of adjunctions, in which the straight arrows are right and the bent arrows left adjoints:



In particular, for a chain complex V , the free n -disk algebra on V is naturally equivalent to $\mathbb{F}_n(O(n) \otimes V)$. More generally, there is the following description.

Proposition 2.7 *There is a natural equivalence*

$$\mathbb{F}_n(K) \xrightarrow{\sim} \bigoplus_{k \geq 0} \left(\text{Emb}(\bigsqcup_k \mathbb{R}^n, -) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \right),$$

where K is an $O(n)$ -module.

Proof The map is supplied by the universal property of the free algebra. In the case $K = O(n) \otimes V$, it is an equivalence, since $\mathbb{F}_n(K)$ is now the free n -disk algebra on the chain complex V , so that

$$\begin{aligned} \mathbb{F}_n(K) &\simeq \bigoplus_{k \geq 0} \left(\text{Emb}(\bigsqcup_k \mathbb{R}^n, -) \otimes_{\Sigma_k} V^{\otimes k} \right) \\ &\cong \bigoplus_{k \geq 0} \left(\text{Emb}(\bigsqcup_k \mathbb{R}^n, -) \otimes_{\Sigma_k \times O(n)^k} (O(n)^k \otimes V^{\otimes k}) \right) \\ &\cong \bigoplus_{k \geq 0} \left(\text{Emb}(\bigsqcup_k \mathbb{R}^n, -) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \right). \end{aligned}$$

Since a general $O(n)$ -module may be expressed as a split geometric realization of free $O(n)$ -modules, and since \mathbb{F}_n , as a left adjoint, preserves geometric realizations, it suffices to show that the right-hand side shares this property. But both $\text{Mod}_{O(n)}(\text{Ch}_{\mathbb{Q}})$ and $\text{Alg}_{\mathcal{D}\text{isk}_n}(\text{Ch}_{\mathbb{Q}})$ are monadic over $\text{Ch}_{\mathbb{Q}}$, so on both sides the geometric realization is computed in $\text{Ch}_{\mathbb{Q}}$, and the right-hand side clearly preserves colimits in chain complexes. \square

In the framed case, $\text{Emb}^{\text{fr}}(\mathbb{R}^n, \mathbb{R}^n)$ is contractible, so there is only the one forgetful functor

$$\text{Alg}_{\mathcal{D}\text{isk}_n^{\text{fr}}}(\text{Ch}_{\mathbb{Q}}) \rightarrow \text{Ch}_{\mathbb{Q}},$$

whose left adjoint, the free framed n -disk algebra functor, is denoted \mathbb{F}_n^{fr} .

By restriction along the natural inclusion $\mathcal{D}\text{isk}_n^{\text{fr}} \rightarrow \mathcal{D}\text{isk}_n$, any n -disk algebra is in particular a framed n -disk algebra, and there is an equivalence of $\mathcal{D}\text{isk}_n^{\text{fr}}$ -algebras

$$\mathbb{F}_n(V) \simeq \mathbb{F}_n^{\text{fr}}(V),$$

where V is a chain complex considered as a trivial $O(n)$ -module.

3 Calculations

3.1 Frame bundles

The object of this section is twofold. First, we compute the factorization homology of the free n -disk algebra generated by an $O(n)$ -module K . Second, for suitable K , we interpret this calculation in terms of the homology of configuration spaces.

For a manifold M , let $\text{Fr}_M \rightarrow M$ denote the corresponding principal $O(n)$ -bundle. Since $\text{Conf}_k(M)$ is an open submanifold of M^k , its structure group is canonically reducible to $O(n)^k$, and we denote the corresponding principal $O(n)^k$ -bundle by $\text{Conf}_k^{\text{fr}}(M)$.

Proposition 3.1 *There is a natural equivalence*

$$\int_M \mathbb{F}_n(K) \xrightarrow{\sim} \bigoplus_{k \geq 0} (\text{Conf}_k^{\text{fr}}(M) \otimes_{\Sigma_k \ltimes O(n)^k} K^{\otimes k}),$$

where K is an $O(n)$ -module.

Proof The natural map

$$\text{colim}_{\mathcal{D}\text{isk}_n/M} (\text{Emb}(\bigsqcup_k \mathbb{R}^n, -)) \xrightarrow{\sim} \text{Emb}(\bigsqcup_k \mathbb{R}^n, M)$$

is an equivalence by [42, page 726], so we have

$$\begin{aligned} \int_M \mathbb{F}_n(K) &\simeq \operatorname{colim}_{\operatorname{Disk}_n/M} \left(\bigoplus_{k \geq 0} \left(\operatorname{Emb}(\bigsqcup_k \mathbb{R}^n, -) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \right) \right) \\ &\simeq \bigoplus_{k \geq 0} \left(\operatorname{colim}_{\operatorname{Disk}_n/M} \left(\operatorname{Emb}(\bigsqcup_k \mathbb{R}^n, -) \right) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \right) \\ &\simeq \bigoplus_{k \geq 0} \left(\operatorname{Emb}(\bigsqcup_k \mathbb{R}^n, M) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \right). \end{aligned}$$

To conclude, we note that evaluation at the origin defines a projection

$$\operatorname{Emb}(\bigsqcup_k \mathbb{R}^n, M) \rightarrow \operatorname{Conf}_k(M),$$

and the natural derivative map $\operatorname{Emb}(\bigsqcup_k \mathbb{R}^n, M) \rightarrow \operatorname{Conf}_k^{\text{fr}}(M)$ covering the identity is an equivalence of $O(n)^k$ -spaces over $\operatorname{Conf}_k(M)$. \square

Remark 3.2 This proposition is a special case of a calculation carried out in the more general context of *zero-pointed manifolds* in [3, Theorem 2.4.1]. We have included this simplified argument for the reader’s convenience.

It will be important in what follows to be able to identify the summand of this object corresponding to a particular choice of k .

Definition 3.3 The *cardinality grading* of the functor $\int_M \mathbb{F}_n(K)$ is the grading corresponding to the direct sum decomposition of Proposition 3.1.

Note that this grading corresponds to the grading induced on the colimit by the cardinality grading of the functor $\mathbb{F}_n(K)$.

We will be most interested in this calculation for particularly simple choices of $O(n)$ -module K .

Corollary 3.4 *There is a natural equivalence*

$$\int_M \mathbb{F}_n(\mathbb{Q}) \xrightarrow{\sim} \bigoplus_{k \geq 0} C_*(B_k(M); \mathbb{Q}).$$

Proposition 3.1 can also be used to study the twisted homology of $B_k(M)$. To pursue this direction, we must first identify the orientation cover $\widetilde{B}_k(M)$ of $B_k(M)$. For this we note that the orientation cover

$$\widetilde{\operatorname{Conf}_k(M)} \rightarrow \operatorname{Conf}_k(M) \rightarrow B_k(M)$$

has structure group $\Sigma_k \times C_2$ when considered as a bundle over $B_k(M)$; that the automorphism corresponding to $-1 \in C_2$ reverses orientation; and that the automorphism corresponding to $\tau \in \Sigma_k$ reverses orientation if $\text{sgn}(\tau) = -1$ and n is odd and preserves orientation otherwise. Therefore, the action of the subgroup

$$H := \{(\tau, \text{sgn}(\tau)^n) \mid \tau \in \Sigma_k\} < \Sigma_k \times C_2$$

is orientation-preserving, and we deduce the following proposition.

Proposition 3.5 $\widehat{B_k(M)} \cong \widehat{\text{Conf}_k(M)}_H$ as covers of $B_k(M)$.

For a chain complex V , let V^{sgn} denote the sign representation of C_2 on V , and V^{det} the $O(n)$ -module obtained from the latter by restriction along the determinant $O(n) \rightarrow C_2$. Recall that, for an n -manifold N , the homology of N twisted by the orientation character may be computed as the homology of the complex

$$C_*(N; \mathbb{Q}^w) := \widetilde{N} \otimes_{C_2} \mathbb{Q}^{\text{sgn}} \cong \text{Fr}_N \otimes_{O(n)} \mathbb{Q}^{\text{det}}.$$

Proposition 3.6 Let M be an n -manifold.

(1) If n is even, there is a natural equivalence

$$\int_M \mathbb{F}_n(\mathbb{Q}^{\text{det}}) \xrightarrow{\sim} \bigoplus_{k \geq 0} C_*(B_k(M); \mathbb{Q}^w).$$

(2) If n is odd, there is a natural equivalence

$$\int_M \mathbb{F}_n(\mathbb{Q}^{\text{det}}[1]) \xrightarrow{\sim} \bigoplus_{k \geq 0} C_*(B_k(M); \mathbb{Q}^w)[k].$$

Proof (1) We have that

$$\begin{aligned} \text{Conf}_k^{\text{fr}}(M) \otimes_{\Sigma_k \times O(n)^k} (\mathbb{Q}^{\text{det}})^{\otimes k} &\cong \text{Conf}_k^{\text{fr}}(M) \otimes_{\Sigma_k \times O(nk)} \mathbb{Q}^{\text{det}} \\ &\cong \widehat{\text{Conf}_k(M)} \otimes_{\Sigma_k \times C_2} \mathbb{Q}^{\text{sgn}} \\ &\cong \widehat{\text{Conf}_k(M)}_{\Sigma_k} \otimes_{C_2} \mathbb{Q}^{\text{sgn}} \\ &\cong \widehat{B_k(M)} \otimes_{C_2} \mathbb{Q}^{\text{sgn}}, \end{aligned}$$

where we used the commutativity of the diagram

$$\begin{array}{ccc} O(n)^k & \longrightarrow & O(nk) \\ \det^k \downarrow & & \downarrow \det \\ C_2^k & \xrightarrow{\text{multiply}} & C_2 \end{array}$$

and the fact that $H = \Sigma_k \times \{1\}$ when n is even. The claim follows after summing over k and applying Proposition 3.1.

(2) Similarly, we have that

$$\begin{aligned} \text{Conf}_k^{\text{fr}}(M) \otimes_{\Sigma_k \times O(n)^k} (\mathbb{Q}^{\det}[1])^{\otimes k} &\cong \text{Conf}_k^{\text{fr}}(M) \otimes_{\Sigma_k \times O(nk)} (\mathbb{Q}^{\det} \otimes \mathbb{Q}[1]^{\otimes k}) \\ &\cong \widetilde{\text{Conf}}_k(M) \otimes_{\Sigma_k \times C_2} (\mathbb{Q}^{\text{sgn}} \otimes \mathbb{Q}[1]^{\otimes k}) \\ &\cong \widetilde{\text{Conf}}_k(M)_H \otimes_{C_2} \mathbb{Q}^{\text{sgn}}[k] \\ &\cong \widetilde{B}_k(M) \otimes_{C_2} \mathbb{Q}^{\text{sgn}}[k], \end{aligned}$$

where we used that $\mathbb{Q}^{\text{sgn}} \otimes \mathbb{Q}[1]^{\otimes k}$ is a trivial H -module and $[\Sigma_k \times C_2 : H] = 2$. \square

3.2 Commutative algebras

We now consider a calculation of factorization homology in a certain degenerate case, which is a slight generalization of that considered in [5, Proposition 5.1]. We will make use of this calculation in the next section.

Restriction of embeddings defines a map $\text{Emb}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n) \rightarrow \prod_k \text{Emb}(\mathbb{R}^n, \mathbb{R}^n) \simeq \prod_k O(n)$, which assemble to form a symmetric monoidal functor

$$\pi: \text{Disk}_n \rightarrow BO(n)^\sqcup,$$

where $BO(n)^\sqcup$ is the ∞ -category obtained as the nerve of the topological category with objects the natural numbers and morphism spaces given by

$$\text{Map}_{BO(n)^\sqcup}(r, s) = \bigsqcup_{f: \langle r \rangle \rightarrow \langle s \rangle} \prod_{i=1}^s O(n)^{f^{-1}(i)},$$

which is symmetric monoidal under addition. For more on this and related ∞ -categories, the reader may consult [42, Section 2.4.3]. For us, the relevance of this object is the following consequence of [42, Theorem 2.4.3.18].

Theorem 3.7 (Lurie) *There is an equivalence*

$$\text{Fun}^\otimes(BO(n)^\sqcup, \text{Ch}_\mathbb{Q}) \xrightarrow{\sim} \text{Mod}_{O(n)}(\text{Alg}_{\text{Com}}(\text{Ch}_\mathbb{Q})).$$

This result motivates our next definition.

Definition 3.8 *A commutative refinement of an n -disk algebra A is a factorization*

$$\begin{array}{ccc}
 \mathcal{D}isk_n & \xrightarrow{A} & \mathcal{C}h_{\mathbb{Q}} \\
 \pi \downarrow & \nearrow & \\
 BO(n)^{\sqcup} & &
 \end{array}$$

through a symmetric monoidal functor $BO(n)^{\sqcup} \rightarrow \mathcal{C}h_{\mathbb{Q}}$.

By the previous theorem, a commutative refinement endows the underlying object of A with the structure of a commutative algebra for which the n -disk algebra structure maps are homomorphisms. More formally, we obtain a factorization

$$\begin{array}{ccc}
 \mathcal{D}isk_n & \xrightarrow{A} & \mathcal{C}h_{\mathbb{Q}} \\
 A_{\text{Com}} \downarrow & \nearrow & \\
 \text{Alg}_{\text{Com}}(\mathcal{C}h_{\mathbb{Q}}) & &
 \end{array}$$

of A through the forgetful functor.

Example 3.9 By the Künneth theorem, the functor $H: \mathcal{C}h_{\mathbb{Q}} \rightarrow \mathcal{C}h_{\mathbb{Q}}$ is symmetric monoidal, whence the homology of an n -disk algebra is canonically an n -disk algebra. Since H factors through the discrete ∞ -category of graded vector spaces, we have a symmetric monoidal factorization

$$\begin{array}{ccc}
 \mathcal{D}isk_n & \xrightarrow{H(A)} & \mathcal{C}h_{\mathbb{Q}} \\
 \pi \downarrow & \nearrow & \\
 BO(n)^{\sqcup} & & \\
 \downarrow & \nearrow & \\
 BC_2^{\sqcup} & &
 \end{array}$$

π_0 (curved arrow from $\mathcal{D}isk_n$ to BC_2^{\sqcup})

through the homotopy category of $\mathcal{D}isk_n$, so that $H(A)$ is canonically commutative.

Definition 3.10 Let X be a topological space and B a commutative algebra. The tensor of X and B is the colimit

$$X \otimes B = \text{colim}(X \rightarrow \text{pt} \xrightarrow{B} \text{Alg}_{\text{Com}}(\mathcal{C}h_{\mathbb{Q}}))$$

of the constant functor from X , viewed as an ∞ -groupoid, with value B .

Remark 3.11 When $X = S^1$, this construction has the homotopy type of the Hochschild chains of A . In general, one recovers Pirashvili’s higher Hochschild homology.

Let $\mathcal{D}isk_{n/M}^1$ denote the full subcategory of $\mathcal{D}isk_{n/M}$ spanned by those arrows $\bigsqcup_k \mathbb{R}^n \rightarrow M$ with $k = 1$.

Proposition 3.12 *Suppose that A admits a commutative refinement. There is a natural equivalence*

$$\text{Fr}_M \otimes_{O(n)} A \simeq \text{colim}(\mathcal{D}isk_{n/M}^1 \rightarrow \mathcal{D}isk_n \xrightarrow{A_{\text{Com}}} \text{Alg}_{\text{Com}}(\text{Ch}_{\mathbb{Q}})).$$

Proof Since the colimit is the left Kan extension to a point, and since Kan extensions compose, we may write

$$\text{colim}_{\mathcal{D}isk_{n/M}^1} A_{\text{Com}} \simeq \text{colim}_{BO(n)} \pi_! A_{\text{Com}} \simeq (\pi_! A_{\text{Com}})_{O(n)},$$

so that it suffices to identify $\pi_! A_{\text{Com}}$.

Since the projection $\mathcal{D}isk_{n/M} \rightarrow \mathcal{D}isk_n$ is a left fibration, so is $\pi: \mathcal{D}isk_{n/M}^1 \rightarrow BO(n)$; in particular, this functor is a co-Cartesian fibration, which implies that the inclusion $\pi^{-1}(\text{pt}) \rightarrow \pi_{/\text{pt}}$ of the fiber over the basepoint into the overcategory is a right adjoint and hence final. Therefore, we have

$$\pi_! A_{\text{Com}} = \text{colim}_{\pi_{/\text{pt}}} A_{\text{Com}} \simeq \text{colim}_{\pi^{-1}(\text{pt})} A_{\text{Com}} = \pi^{-1}(\text{pt}) \otimes A$$

According to [5, Corollary 2.13], the ∞ -category $\mathcal{D}isk_{n/M}^1$ is equivalent to the Kan complex M , and the map $\pi: \mathcal{D}isk_{n/M}^1 \rightarrow BO(n)$ coincides under this identification with the classifying map for the tangent bundle of M . In particular, the fiber of this map is $O(n)$ -equivalent to Fr_M , which completes the proof. \square

Proposition 3.13 *Suppose that A admits a commutative refinement. There is a natural equivalence*

$$\int_M A \simeq \text{Fr}_M \otimes_{O(n)} A.$$

Proof By the previous proposition, it suffices to show that the inclusion $\mathcal{D}isk_{n/M}^1 \rightarrow \mathcal{D}isk_{n/M}$ and the forgetful functor $\text{Alg}_{\text{Com}}(\text{Ch}_{\mathbb{Q}}) \rightarrow \text{Ch}_{\mathbb{Q}}$ induce equivalences

$$\text{colim}_{\mathcal{D}isk_{n/M}^1} A_{\text{Com}} \xrightarrow{\sim} \text{colim}_{\mathcal{D}isk_{n/M}} A_{\text{Com}} \xrightarrow{\sim} \text{colim}_{\mathcal{D}isk_{n/M}} A$$

when A is commutative.

Since $\text{Ch}_{\mathbb{Q}}$ is \otimes -presentable (see [5, Definition 3.4]), the second equivalence follows from [5, Corollary 3.22], which asserts that $\mathcal{D}isk_{n/M}$ is sifted, and [42, Corollary 3.2.3.2], which implies that the forgetful functor from commutative algebras preserves sifted colimits.

The first equivalence holds whenever M is framed by [5, Proposition 5.1], since in this case the diagram

$$\begin{array}{ccc}
 \mathcal{D}isk_{n/M}^{fr} & \xrightarrow{\sim} & \mathcal{D}isk_{n/M} \\
 \downarrow & & \downarrow \\
 \mathcal{D}isk_n^{fr} & \longrightarrow & \mathcal{D}isk_n \\
 \downarrow & & \downarrow \\
 \mathit{Fin} = \{e\}^\sqcup & \longrightarrow & BO(n)^\sqcup
 \end{array}$$

commutes. In particular, the equivalence holds for $M = \bigsqcup_k \mathbb{R}^n$, and we conclude that

$$A \simeq \mathit{Fr}_{(-)} \otimes_{O(n)} A$$

as n -disk algebras. Therefore, the claim will be established once we are assured that the expression on the right satisfies condition (2) of Theorem 2.3. For this, we note that the functor $\mathit{Fr}_{(-)}$ takes a collar-gluing of manifolds to a pushout of $O(n)$ -spaces, and that the functor $- \otimes_{O(n)} A$ preserves colimits of $O(n)$ -spaces. \square

3.3 A spectral sequence

We employ a certain “commutative-to-noncommutative” spectral sequence in the proof of Theorem 1.1. For technical reasons, it will be convenient to restrict our attention to n -disk algebras valued in $\mathit{Ch}_{\mathbb{Q}}^{\geq 0}$, the full subcategory of chain complexes concentrated in nonnegative homological degree. This restriction is not essential.

Proposition 3.14 *Let M be an n -manifold and A an n -disk algebra in $\mathit{Ch}_{\mathbb{Q}}^{\geq 0}$. There is a natural first-quadrant spectral sequence*

$$E_{p,q}^2 \cong H_{p,q}(\mathit{Fr}_M \otimes_{O(n)} H(A)) \implies H_{p+q} \left(\int_M A \right),$$

with differential d^r of bidegree $(-r, r - 1)$.

The nature of the bigrading will become clear in the proof.

To construct this spectral sequence, we employ a rigidified version of the overcategory $\mathcal{D}isk_{n/M}$, denoted $\mathit{Disj}(M)$ following [42, Chapter 5], which is the poset of those open subsets of M diffeomorphic to $\bigsqcup_k \mathbb{R}^n$ for some k . We refer the reader to [42, Proposition 5.5.2.13] for the proof of the following result.

Proposition 3.15 *There is a final functor $N(\text{Disj}(M)) \rightarrow \text{Disk}_n/M$.*

Thus, by [40, Proposition 4.1.1.8], the factorization homology of M may be computed as a colimit over the nerve of the ordinary category $\text{Disj}(M)$. Having achieved this simplification, we proceed as follows. Using the fact that $\text{Ch}_{\mathbb{Q}}^{\geq 0}$ arises from a combinatorial simplicial model category, [40, Proposition 4.2.4.4] implies that any functor $N(\text{Disj}(M)) \rightarrow \text{Ch}_{\mathbb{Q}}^{\geq 0}$ of ∞ -categories is equivalent in the ∞ -category of functors to one coming from a functor of ordinary categories. Having chosen such a “straightening” of A , which we abusively denote by A , [40, Theorem 4.2.4.1] now guarantees that the homotopy colimit of A coincides with the ∞ -categorical colimit.

Proof of Proposition 3.14 From the discussion of the previous paragraph and [47, Corollary 5.1.3], we have equivalences

$$\int_M A \simeq \text{hocolim}_{\text{Disj}(M)} A \simeq B(\text{pt}, \text{Disj}(M), A),$$

where $B(\text{pt}, \text{Disj}(M), A)$ denotes the realization of the simplicial chain complex given in simplicial degree p by

$$B_p(\text{pt}, \text{Disj}(M), A) = \bigoplus_{U_p \rightarrow \dots \rightarrow U_0 \rightarrow M} A(U_p)$$

(here we use for a second time the fact that the model structure on nonnegatively graded chain complexes is simplicial). Filtering by skeleta in the usual way, we obtain a spectral sequence

$$E_{p,q}^1 = \bigoplus_{U_p \rightarrow \dots \rightarrow U_0 \rightarrow M} H_q(A(U_p)) \implies H_{p+q} \left(\int_M A \right),$$

with the differential d^1 given by the alternating sum of the face maps (see [50, Proposition 5.1], for example, which treats the case of a simplicial space). In other words, the E^1 page is the (graded) chain complex associated to the (graded) simplicial chain complex $B_{\bullet}(\text{pt}, \text{Disj}(M), H(A))$ via the Dold–Kan correspondence, so that, invoking Proposition 3.13, we have natural isomorphisms

$$E_{p,q}^2 \cong H_{p,q}(B(\text{pt}, \text{Disj}(M), H(A))) \cong H_{p,q} \left(\int_M H(A) \right) \cong H_{p,q}(\text{Fr}_M \otimes_{O(n)} H(A)). \quad \square$$

Remark 3.16 Horel discusses a version of this spectral sequence in [33, Section 5].

3.4 Enveloping algebras

In this section, we outline the place of Lie algebras in the theory of factorization homology, the general reference for which is [35].

It has long been known that configuration spaces are intimately related to Lie algebras; see [17; 18; 16], for example. To see the connection, suppose that A is a $\text{Disk}_n^{\text{fr}}$ -algebra in chain complexes, with $n \geq 2$. Part of the structure of such an object is a multiplication map

$$m: \text{Emb}^{\text{fr}}(\bigsqcup_2 \mathbb{R}^n, \mathbb{R}^n) \otimes A^{\otimes 2} \rightarrow A,$$

and since the homology of $\text{Emb}^{\text{fr}}(\bigsqcup_2 \mathbb{R}^n, \mathbb{R}^n) \simeq \text{Conf}_2(\mathbb{R}^n) \simeq S^{n-1}$ is concentrated in degrees 0 and $n - 1$, this multiplication encodes two maps

$$m_0: A^{\otimes 2} \rightarrow A \quad \text{and} \quad m_{n-1}: A^{\otimes 2} \rightarrow A[1-n]$$

defining a commutative multiplication on A and a Lie bracket on $A[n - 1]$, again up to homotopy. The Jacobi identity for m_{n-1} follows from the three-term or Yang–Baxter relations in $H_*(\text{Conf}_3(\mathbb{R}^n))$ (see [20]), and $O(n)$, acting on S^{n-1} by degree ± 1 maps, interchanges it with the opposite bracket.

The fact that this discussion illustrates is the existence of a forgetful functor from $\text{Disk}_n^{\text{fr}}$ -algebras to Lie algebras at the level of ∞ -categories. Indeed, according to [35], there is the following commuting diagram of adjunctions:

$$\begin{array}{ccc}
 & \xleftarrow{U_n} & \\
 \text{Alg}_{\text{Disk}_n^{\text{fr}}}(\text{Ch}\mathbb{Q}) & \xrightarrow{\quad} & \text{Alg}_{\mathcal{L}}(\text{Ch}\mathbb{Q}) \\
 \uparrow \mathbb{F}_n^{\text{fr}} & & \downarrow \mathcal{L} \\
 \text{Ch}\mathbb{Q} & \xrightarrow{[n-1]} & \text{Ch}\mathbb{Q} \\
 & \xleftarrow{[1-n]} &
 \end{array}$$

Here \mathcal{L} denotes the free Lie algebra functor.

The $\text{Disk}_n^{\text{fr}}$ -algebra $U_n(\mathfrak{g})$ is known as the n -enveloping algebra of \mathfrak{g} ; see [31, Section 4.6] for a discussion of the identification between U_1 and the usual universal enveloping algebra. The factorization homology of these algebras is computed in [35].

Theorem 3.17 (Knudsen) *There is a natural equivalence*

$$\int_M U_n(\mathfrak{g}) \xrightarrow{\sim} C^{\mathcal{L}}(\mathfrak{g}^{M^+}).$$

We pause briefly to explain the terms of the theorem.

- (1) The ∞ -category of differential graded Lie algebras has limits and is therefore cotensored over pointed spaces; we denote by \mathfrak{g}^X the cotensor of the pointed space X with the Lie algebra \mathfrak{g} . A model for this object is provided by [32, Lemma 4.8.3].

Proposition 3.18 *Let X be a pointed finite CW complex. There is a natural equivalence*

$$\mathfrak{g}^X \simeq A_{\text{PL}}(X) \otimes \mathfrak{g}.$$

Here A_{PL} denotes the functor of reduced piecewise-linear de Rham forms (see [22, Section 10(c)], for example), and the right-hand side carries the canonical Lie bracket on the tensor product of a nonunital commutative algebra and a Lie algebra, which is defined by the formula

$$[a \otimes v, b \otimes w] = (-1)^{|v||b|} ab \otimes [v, w].$$

(2) The symbol $C^{\mathcal{L}}$ denotes the functor of Lie algebra chains. This coaugmented cocommutative coalgebra is defined abstractly via the monadic bar construction against the free Lie algebra monad, but it has a concrete incarnation as the *Chevalley–Eilenberg complex*

$$\text{CE}(\mathfrak{g}) = (\text{Sym}(\mathfrak{g}[1]), d_{\mathfrak{g}} + D),$$

where D is defined as a coderivation by specifying that

$$D(\sigma x \wedge \sigma y) = (-1)^{|x|} \sigma[x, y].$$

See [27, Section 6] for a discussion of the comparison between the monadic bar construction and the Chevalley–Eilenberg complex. We remark that $\text{CE}(\mathfrak{g})$ is a coaugmented cocommutative differential graded coalgebra, and the resulting coproduct on $H^{\mathcal{L}}(\mathfrak{g})$ coincides with the one inherited from the monadic bar construction; indeed, both are induced by the diagonal $\mathfrak{g} \rightarrow \mathfrak{g} \oplus \mathfrak{g}$, which is a map of Lie algebras.

The equivalence of Theorem 3.17 specializes to a natural equivalence

$$U_n(\mathfrak{g}) \simeq C^{\mathcal{L}}(\mathfrak{g}^{(\mathbb{R}^n)^+})$$

of $\text{Disk}_n^{\text{fr}}$ -algebras. In this way, Theorem 3.17 can be thought of as identifying an n -disk algebra refinement of the $\text{Disk}_n^{\text{fr}}$ -algebra $U_n(\mathfrak{g})$, so that the expression $\int_M U_n(\mathfrak{g})$ is sensible for manifolds M that are not necessarily framed.

Returning to the discussion that began this section, if A is now an n -disk algebra rather than merely a $\text{Disk}_n^{\text{fr}}$ -algebra, then A determines a shifted Lie algebra in $O(n)$ -modules, but now with $O(n)$ acting on the suspension coordinates. A full discussion of this phenomenon and the corresponding enveloping algebra is beyond the scope of this paper. Since the analogue of Theorem 3.17 is true in that context, we will content ourselves with making it our definition.

As a matter of notation, if X is a pointed $O(n)$ -space and \mathfrak{g} a Lie algebra in $O(n)$ -modules, we denote the $O(n)$ -invariants of \mathfrak{g}^X by $\text{Map}^{O(n)}(X, \mathfrak{g})$.

Definition 3.19 Let \mathfrak{g} be a Lie algebra in $O(n)$ -modules. The n -enveloping algebra of \mathfrak{g} is the n -disk algebra

$$U_n(\mathfrak{g}) = C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{(\mathbb{R}^n)^+}, \mathfrak{g})).$$

Here we take the frame bundle of the one-point compactification to be the cofiber

$$\text{Fr}_{M^+} = \text{cofib}(\text{Fr}_{\overline{M}}|_{\partial\overline{M}} \rightarrow \text{Fr}_{\overline{M}})$$

of $O(n)$ -spaces, where \overline{M} is a compact n -manifold with boundary whose interior is M ; see [4, Definition 4.5.1] for a more invariant interpretation of this object.

A choice of framing of \mathbb{R}^n trivializes $\text{Fr}_{(\mathbb{R}^n)^+}$, inducing an equivalence

$$\text{Map}^{O(n)}(\text{Fr}_{(\mathbb{R}^n)^+}, \mathfrak{g}) \simeq \mathfrak{g}^{(\mathbb{R}^n)^+},$$

which is even equivariant for the diagonal action of $O(n)$ on the target, so this definition specializes via Theorem 3.17 to our earlier one when \mathfrak{g} is an ordinary Lie algebra.

The corresponding factorization homology calculation is the following.

Proposition 3.20 *There is a natural equivalence*

$$\int_M U_n(\mathfrak{g}) \xrightarrow{\sim} C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathfrak{g}))$$

for M an n -manifold and \mathfrak{g} a Lie algebra in $O(n)$ -modules.

Proof Since $C^{\mathcal{L}}$, as a left adjoint, preserves colimits, it suffices to exhibit an equivalence of Lie algebras

$$\int_M \mathfrak{g}^{(\mathbb{R}^n)^+} \xrightarrow{\sim} \text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathfrak{g}),$$

which is supplied by the argument of [5, Proposition 5.13], since sifted colimits of Lie algebras are computed in $\text{Ch}_{\mathbb{Q}}$ by [41, Proposition 2.1.16]. □

We close this section with a definition of a grading that will play an important role in what follows. Let \mathfrak{g} be a differential graded Lie algebra with a *weight decomposition* as a direct sum of complexes $\mathfrak{g} = \bigoplus_k \mathfrak{g}(k)$ with the property that $[v, w] \in \mathfrak{g}(r + s)$ when $v \in \mathfrak{g}(r)$ and $w \in \mathfrak{g}(s)$.

Example 3.21 A free Lie algebra has a canonical weight decomposition

$$\mathcal{L}(V) = \bigoplus_{k \geq 0} \mathcal{L}(k) \otimes_{\Sigma_k} V^{\otimes k}.$$

Example 3.22 If \mathfrak{g} has a weight decomposition, then $A_{\text{PL}}(X) \otimes \mathfrak{g}$ carries a canonical weight decomposition for any space X .

Such a decomposition induces a *weight grading* on the underlying graded vector space of $\text{Sym}(\mathfrak{g}[1])$ of the Chevalley–Eilenberg complex. In fact, since we have assumed that the bracket and differential of \mathfrak{g} each respect the weight decomposition, the Chevalley–Eilenberg differential applied to a monomial of pure weight k again has pure weight k , so that $\text{CE}(\mathfrak{g})$ is a bicomplex. In this way, a weight decomposition of \mathfrak{g} induces a weight grading on $H(U_n(\mathfrak{g}))$.

4 Configuration spaces

4.1 The main result

In this section, we prove Theorem 1.1 assuming the validity of several results, discussion of which is postponed for the sake of continuity, as the proofs involve different techniques from those used thus far.

As a preliminary step, we have the following basic pair of observations.

Proposition 4.1 (1) *Let K be an $O(n)$ -module and \underline{K} its underlying chain complex. There is a natural equivalence of framed n -disk algebras*

$$\mathbb{F}_n^{\text{fr}}(\underline{K}) \simeq \mathbb{F}_n(K).$$

(2) *Let \mathfrak{g} be a Lie algebra in $O(n)$ -modules and $\underline{\mathfrak{g}}$ its underlying Lie algebra. There is a natural equivalence of framed n -disk algebras*

$$U_n(\underline{\mathfrak{g}}) \simeq U_n(\mathfrak{g}).$$

Proof A choice of framing for \mathbb{R}^n induces an $O(n)$ -equivariant homotopy equivalence

$$\text{Emb}^{\text{fr}}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n) \times O(n)^k \xrightarrow{\sim} \text{Emb}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n),$$

whence from Proposition 2.7 we have

$$\begin{aligned} \mathbb{F}_n(K) &\cong \bigoplus_{k \geq 0} \left(\text{Emb}^{\text{fr}}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n) \times O(n)^k \right) \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k} \\ &\cong \bigoplus_{k \geq 0} \left(\left(\text{Emb}^{\text{fr}}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n) \times O(n)^k \right) \otimes_{O(n)^k} K^{\otimes k} \right)_{\Sigma_k} \\ &\cong \bigoplus_{k \geq 0} \text{Emb}^{\text{fr}}(\bigsqcup_k \mathbb{R}^n, \mathbb{R}^n) \otimes_{\Sigma_k} K^{\otimes k} \\ &\cong \mathbb{F}_n^{\text{fr}}(\underline{K}). \end{aligned}$$

This proves (1), and (2) is immediate from Definition 3.19. □

Remark 4.2 Thinking topologically, the generic example of an n -disk algebra in spaces is an n -fold loop space on an $O(n)$ -space X ; see [49] or [54]. In this context, the statement is that, as an n -fold loop space, the homotopy type of $\Omega^n X$ does not depend on the action of $O(n)$ on X .

Connecting (1) and (2) is the following formal observation, which amounts to the statement that left adjoints compose.

Proposition 4.3 *Let V be a chain complex. There is a natural equivalence*

$$\mathbb{F}_n^{\text{fr}}(V) \xrightarrow{\sim} U_n(\mathcal{L}(V[n-1]))$$

of framed n -disk algebras, where \mathcal{L} is the free Lie algebra functor.

This observation is a generalization of the familiar fact that the universal enveloping algebra of the free Lie algebra on a set of generators S is free on S as an associative algebra; however, equipped with the involution given by its Hopf algebra antipode, the universal enveloping algebra of the free Lie algebra on S is *not* the free algebra-with-involution on S . This classical fact illustrates the $n = 1$ case of the general phenomenon that the free n -disk algebra on the trivial $O(n)$ -module V is *not* the n -enveloping algebra of the free Lie algebra on V . As the following proposition shows, the $O(n)$ -action must be twisted to restore the equivalence.

Proposition 4.4 *Let K be an $O(n)$ -module. There is a natural equivalence*

$$\mathbb{F}_n(K) \xrightarrow{\sim} U_n(\mathcal{L}((\mathbb{R}^n)^+ \otimes K[-1]))$$

of n -disk algebras, where $(\mathbb{R}^n)^+ \otimes K$ carries the diagonal $O(n)$ -action.

Proof First, we note that the unit

$$K \rightarrow ((\mathbb{R}^n)^+ \otimes K)^{(\mathbb{R}^n)^+}$$

of the tensor/cotensor adjunction is an equivalence of $O(n)$ -modules. Indeed, it suffices to verify this in the case $K = \mathbb{Q}$, in which case the map induces the isomorphism $\mathbb{Q} \cong (\mathbb{Q}^{\det})^{\otimes 2}$ in homology.

Now, composing this unit map with the natural inclusions

$$((\mathbb{R}^n)^+ \otimes K)^{(\mathbb{R}^n)^+} \rightarrow \mathcal{L}((\mathbb{R}^n)^+ \otimes K[-1])^{(\mathbb{R}^n)^+}[1] \rightarrow C^{\mathcal{L}}(\mathcal{L}((\mathbb{R}^n)^+ \otimes K[-1])^{(\mathbb{R}^n)^+})$$

of $O(n)$ -modules, we obtain a map of n -disk algebras

$$\mathbb{F}_n(K) \rightarrow U_n(\mathcal{L}((\mathbb{R}^n)^+ \otimes K[-1]))$$

from the universal property of the free algebra. It will suffice to show that this map is an equivalence upon passing to underlying $\text{Disk}_n^{\text{fr}}$ -algebras, which follows from the previous two propositions and the (nonequivariant) equivalence $\mathbb{Q}[n] \simeq \tilde{C}_*((\mathbb{R}^n)^+; \mathbb{Q})$. □

Proof of Theorem 1.1 An equivalence of n -disk algebras induces an equivalence on passing to factorization homology. Using the indicated results, we obtain equivalences

$$\bigoplus_{k \geq 0} C_*(B_k(M); \mathbb{Q}) \simeq \int_M \mathbb{F}_n(\mathbb{Q}) \tag{3.4}$$

$$\simeq \int_M U_n(\mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+)[-1])) \tag{4.4}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+)[-1]))) \tag{3.20}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathcal{L}(\mathbb{Q}^{\det}[n-1]))) \tag{7.1}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{C_2}(\tilde{M}^+, \mathcal{L}(\mathbb{Q}^{\text{sgn}}[n-1])))$$

$$\simeq C^{\mathcal{L}}(H_c^{-*}(M, \mathcal{L}(\mathbb{Q}^w[n-1]))) \tag{7.5}$$

Applying Proposition 3.14 to this equivalence of algebras, we obtain an isomorphism of spectral sequences. The weight and cardinality gradings of the two algebras pass to factorization homology, so that these spectral sequences are each trigraded. According to Proposition 5.4, the isomorphism preserves the extra grading on E^2 and hence on E^∞ . □

4.2 Variations

In this section, we discuss the corresponding results for twisted homology and manifolds with boundary.

Theorem 4.5 *Let M be an n -manifold.*

- (1) *If n is even, there is an isomorphism of bigraded vector spaces*

$$\bigoplus_{k \geq 0} H_*(B_k(M); \mathbb{Q}^w) \cong H^{\mathcal{L}}(H_c^{-*}(M; \mathcal{L}(\mathbb{Q}[n-1])))$$

- (2) *If n is odd, there is an isomorphism of bigraded vector spaces*

$$\bigoplus_{k \geq 0} H_*(B_k(M); \mathbb{Q}^w)[k] \cong H^{\mathcal{L}}(H_c^{-*}(M; \mathcal{L}(\mathbb{Q}[n])))$$

Proof We imitate the proof of Theorem 1.1. In the even case, we have

$$\bigoplus_{k \geq 0} C_*(B_k(M); \mathbb{Q}^w) \simeq \int_M \mathbb{F}_n(\mathbb{Q}^{\det}) \tag{3.6}$$

$$\simeq \int_M U_n(\mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+) \otimes \mathbb{Q}^{\det}[-1])) \tag{4.4}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+) \otimes \mathbb{Q}^{\det}[-1]))) \tag{3.20}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathcal{L}((\mathbb{Q}^{\det})^{\otimes 2}[n-1]))) \tag{7.1}$$

$$\simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M^+}, \mathcal{L}(\mathbb{Q}[n-1])))$$

$$\simeq C^{\mathcal{L}}(\mathcal{L}(\mathbb{Q}[n-1])^{M^+})$$

$$\simeq C^{\mathcal{L}}(H_c^{-*}(M, \mathcal{L}(\mathbb{Q}[n-1])),) \tag{7.5}$$

and the odd case is essentially identical. The same argument as in the proof of Theorem 1.1 shows that the resulting isomorphism is bigraded. \square

Now, if M is a manifold with boundary, then $B_k(M) \simeq B_k(\overset{\circ}{M})$, since configuration spaces are isotopy functors. A more interesting configuration space in this context is the *relative configuration space*

$$B_k(M, \partial M) := \frac{B_k(M)}{\{(x_1, \dots, x_k) \mid x_i \in \partial M \text{ for some } i\}}.$$

From the point of view of factorization homology, the natural setting in which to study these spaces is that of the *zero-pointed manifolds* of [4], a class of pointed spaces that are manifolds away from the basepoint. Indeed, if M is a manifold with boundary, then $M/\partial M$ is naturally a zero-pointed manifold.

The algebraic counterpart of a basepoint is an *augmentation*.

Definition 4.6 An *augmented n -disk algebra* is an n -disk algebra A together with a map of n -disk algebras $\epsilon: A \rightarrow \mathbb{Q}$.

Example 4.7 The free n -disk algebra $\mathbb{F}_n(K)$ is naturally augmented via the unique map of $O(n)$ -modules $K \rightarrow 0$.

Example 4.8 The n -enveloping algebra $U_n(\mathfrak{g})$ is naturally augmented via the unique map of Lie algebras $\mathfrak{g} \rightarrow 0$.

The theory of factorization homology for zero-pointed n -manifolds with coefficients in augmented n -disk algebras is expounded at length in [4] and [3]. For us, what is

important is that, if M is a manifold with boundary, then the factorization homology of $M/\partial M$ is defined for any choice of augmented n -disk algebra; moreover, if $\partial M = \emptyset$, then $M/\partial M = M_+$, and the factorization homology of the zero-pointed manifold M_+ with coefficients in $\epsilon: A \rightarrow \mathbb{Q}$ is equivalent to the factorization homology of M with coefficients in A defined previously.

Our arguments go through in this more general context.

Theorem 4.9 *Let M be an n -manifold with boundary. There is an isomorphism of bigraded vector spaces*

$$\bigoplus_{k \geq 0} \tilde{H}_*(B_k(M, \partial M); \mathbb{Q}) \cong H^{\mathcal{L}}(H_c^{-*}(M; \mathcal{L}(\mathbb{Q}^w[n-1]))).$$

Proof We explain the adjustments necessary in the proof of Theorem 1.1. First, [3, Theorem 2.4.1] guarantees the equivalence

$$\bigoplus_{k \geq 0} \tilde{C}_*(B_k(M, \partial M); \mathbb{Q}) \simeq \int_{M/\partial M} \mathbb{F}_n(\mathbb{Q}).$$

Second, it is immediate from its definition that the map of Proposition 4.4 is a map of augmented n -disk algebras, so that

$$\int_{M/\partial M} \mathbb{F}_n(\mathbb{Q}) \simeq \int_{M/\partial M} U_n(\mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+)[-1])).$$

The proof of Proposition 3.20 translates verbatim into the zero-pointed context, so that we have the further equivalence

$$\int_{M/\partial M} U_n(\mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+)[-1])) \simeq C^{\mathcal{L}}(\text{Map}^{O(n)}(\text{Fr}_{M_+}, \mathcal{L}(\tilde{C}_*((\mathbb{R}^n)^+)[-1]))).$$

The remainder of the proof goes through unchanged. □

Remark 4.10 When M has boundary, there are two obvious candidates for the orientation sheaf of M , namely the ordinary and the exceptional pushforwards of the orientation sheaf of the interior of M . We intend the former here.

4.3 Formulas

In this section, we use Theorem 1.1 and the Chevalley–Eilenberg complex to reproduce and extend the classical results on the rational homology of configuration spaces alluded to in the introduction.

We remind the reader that the free Lie algebra on $\mathbb{Q}^w[r]$ is given as a graded vector space by

$$\mathcal{L}(\mathbb{Q}^w[r]) \cong \begin{cases} \mathbb{Q}^w[r] \oplus \mathbb{Q}[2r] & \text{for } r \text{ odd,} \\ \mathbb{Q}^w[r] & \text{for } r \text{ even.} \end{cases}$$

When r is odd, the only nonvanishing bracket is the isomorphism $(\mathbb{Q}^w[r])^{\otimes 2} \cong \mathbb{Q}[2r]$.

Corollary 4.11 *If n is odd, there is an isomorphism*

$$H_*(B_k(M); \mathbb{Q}) \cong \text{Sym}^k(H_*(M; \mathbb{Q})).$$

Proof Since n is odd, the Lie algebra in question is abelian, so that the Chevalley–Eilenberg complex has no differential, and the weight grading coincides with the usual grading of the symmetric algebra. The claim follows after replacing shifted, twisted, compactly supported cohomology with homology using Poincaré duality. \square

This result is [10, Theorem C] as formulated in dual form in [23, Theorem 4], in which the isomorphism on cohomology is shown to be an isomorphism of algebras.

Corollary 4.12 *If n is even, $H_*(B_k(M); \mathbb{Q})$ is isomorphic to the homology of the complex*

$$\left(\bigoplus_{i=0}^{\lfloor k/2 \rfloor} \text{Sym}^{k-2i}(H_c^{-*}(M; \mathbb{Q}^w)[n]) \otimes \text{Sym}^i(H_c^{-*}(M; \mathbb{Q})[2n-1]), D \right),$$

where the differential D is defined as a coderivation by the equation

$$D(\sigma^n \alpha \wedge \sigma^n \beta) = (-1)^{(n-1)|\beta|} \sigma^{2n-1}(\alpha \smile \beta).$$

Proof It suffices by Theorem 1.1 to identify the complex in question with the weight- k part of the Chevalley–Eilenberg complex for $\mathfrak{g} = H_c^{-*}(M; \mathcal{L}(\mathbb{Q}^w[n-1]))$, which as a graded vector space is given by

$$\text{Sym}(\mathfrak{g}[1]) \cong \text{Sym}(H_c^{-*}(M; \mathbb{Q}^w)[n]) \otimes \text{Sym}(H_c^{-*}(M; \mathbb{Q})[2n-1]),$$

with differential determined as a coderivation by the bracket of \mathfrak{g} , which is none other than the shifted cup product shown above, with the sign determined by the usual Koszul rule of signs. Since the cogenerators of the first tensor factor have weight 1 and those of the second tensor factor weight 2, the subcomplex of total weight k is exactly the sum shown above. \square

When M is closed, orientable and nilpotent, we recover the linear and Poincaré dual of [24, Theorem A], as formulated in [23, Theorem 1]. When M is a once-punctured surface, we recover [9, Theorem C].

Remark 4.13 The proofs of Theorem 1 and the even-dimensional half of Theorem 3 of [23] rely crucially on the results of [24] and thereby on the hypotheses of compactness, orientability and nilpotence. At the time of writing, these hypotheses do not appear in the statements of the theorems.

It follows from our results, however, that these theorems are true at the stated level of generality. Indeed, by [23, Theorem 6], the Σ_k -invariants of the E_1 page of the Cohen–Taylor–Totaro spectral sequence (see [18] and [53]) coincide with the linear dual of the complex exhibited in Corollary 4.12.

An analogous spectral sequence in the nonorientable case, possibly with twisted coefficients, is available due to [28] and [29]; see also [45].

We leave it to the reader to formulate the analogous results on twisted homology and those concerning the homology of the relative configuration spaces $B_k(M, \partial M)$, which follow in the same way from Theorems 4.5 and 4.9, respectively. To the author's knowledge, the computation in the twisted case is new in all cases except when M is orientable and n is even, so that $B_k(M)$ is orientable, and the computation in the relative case is new in all cases except when $\partial M = \emptyset$.

5 Coalgebraic structure

5.1 Primitives and weight

Our present goal is to supply the first of the missing ingredients in the proof of the main theorem, namely the identification of the cardinality and weight gradings at the level of homology (see Definition 3.3 and the end of Section 3.4 for definitions of these gradings). We make this identification locally on M in this section and globalize in the following section using a spectral sequence argument.

Let K be an $O(n)$ -module. We define the following maps:

- (1) $\iota: K \rightarrow \mathbb{F}_n(K)$ is the map of $O(n)$ -modules given by the unit of the free/forgetful adjunction;
- (2) $\eta: \mathbb{Q} \rightarrow \mathbb{F}_n(K)$ is the unit of $\mathbb{F}_n(K)$ as an n -disk algebra;

- (3) $\delta: \mathbb{F}_n(K) \rightarrow \mathbb{F}_n(K) \otimes \mathbb{F}_n(K)$ is the map of n -disk algebras induced by the composite

$$K \xrightarrow{\Delta} K \oplus K \xrightarrow{\eta \otimes \iota + \iota \otimes \eta} \mathbb{F}_n(\mathbb{Q}) \otimes \mathbb{F}_n(\mathbb{Q}),$$

where Δ is the diagonal and we have tacitly employed the canonical identifications $K \otimes \mathbb{Q} \cong K \cong \mathbb{Q} \otimes K$;

- (4) δ_M and η_M are the maps on factorization homology induced by δ and η , respectively.


Note that we have suppressed the choice of K from the notation.

Although we will only use the case $M = \mathbb{R}^n$ here, we record the following result for its inherent interest.

Proposition 5.1 *The maps $H(\delta_M)$ and $H(\eta_M)$ endow $H(\int_M \mathbb{F}_n(K))$ with the structure of a coaugmented cocommutative coalgebra.*

Proof The functor \int_M is symmetric monoidal in the algebra variable by [42, Theorem 5.5.3.2], so it suffices to verify the claim in the case $M = \mathbb{R}^n$. The required axioms all follow from the universal property of the free algebra; we spell out the argument for coassociativity, leaving the remainder to the reader.

Consider the following cubical diagram:



It will suffice to show that the square diagram given by the front face of the cube commutes in the ∞ -category of n -disk algebras, since this square witnesses coassociativity after applying factorization homology and passing to the homotopy category of chain complexes. Applying the universal property of the free algebra, the required commutativity is equivalent to commutativity as a diagram of $O(n)$ -modules after precomposing with η . By a standard diagram chase, it suffices to verify that the remaining five faces each commute:

- The left and top face commute by the definition of δ .
- The back face commutes by the universal property of the direct sum, considered as the ∞ -categorical product.
- The right and bottom face commute by the definition of δ and the universal property of the direct sum, considered as the ∞ -categorical coproduct. \square

Although we have defined this coalgebra structure in abstract terms, it has an appealing geometric interpretation, which we discuss in Section 5.2 below.

When $M = \mathbb{R}^n$, the same homology is also an algebra, and even commutative for $n \geq 2$. Since δ is a map of n -disk algebras, $H(\mathbb{F}_n(\mathbb{Q}))$ inherits the structure of a *bialgebra*, and in fact a Hopf algebra, although we will not make use of the antipode.

For the duration of this section, we make the abbreviation

$$\mathfrak{g}(K) := \mathcal{L}((\mathbb{R}^n)^+ \otimes K[-1])^{(\mathbb{R}^n)^+}.$$


Proposition 5.2 *The isomorphism on homology induced by the equivalence of Proposition 4.4 is an isomorphism of bialgebras.*

Proof Denote by φ the equivalence

$$\mathbb{F}_n(K) \xrightarrow{\sim} C^{\mathcal{L}}(\mathfrak{g}(K))$$

of Proposition 4.4. Since φ is a map of n -disk algebras, the induced map on homology is a map of algebras; therefore, it will suffice to show that this map is also a map of coalgebras.

Consider the cubical diagram



where γ denotes the comultiplication on Lie algebra chains. As before, we wish to show that the front face commutes in the ∞ -category of n -disk algebras, and, as before, this reduces to checking the commutativity of the remaining five faces in the ∞ -category of $O(n)$ -modules:

- The left face commutes by the definition of δ .
- The back face commutes by functoriality of the diagonal.
- The top face commutes by the definition of φ .
- The bottom face commutes by the definition of φ and the universal property of the direct sum, considered as the categorical coproduct.
- The right face commutes because the functor $C^{\mathcal{L}}$ is Cartesian monoidal. \square

This bialgebra is a familiar one, and the various components of its structure interact predictably with the bigradings.

Proposition 5.3 (1) *There are isomorphisms*

$$H^{\mathcal{L}}(\mathfrak{g}(K)) \cong \text{Sym}(H(\mathfrak{g}(K))[1]) \cong H(\mathbb{F}_n(K))$$

of graded bialgebras, where Sym is equipped with the standard product and coproduct.

- (2) *The product in $H(\mathbb{F}_n(K))$ preserves the cardinality grading.*
- (3) *The coproduct in $H(\mathbb{F}_n(K))$ preserves the cardinality grading.*
- (4) *The product in $H^{\mathcal{L}}(\mathfrak{g}(K))$ preserves the weight grading.*
- (5) *The coproduct in $H^{\mathcal{L}}(\mathfrak{g}(K))$ preserves the weight grading.*

Proof (1) We note that $\mathfrak{g}(K)$ is a formal Lie algebra, since the pointed space $(\mathbb{R}^n)^+$ is formal; moreover, since $H(\mathfrak{g}(K))$ is abelian, there is no differential in the Chevalley–Eilenberg complex, so we have isomorphisms of coaugmented coalgebras

$$H^{\mathcal{L}}(\mathfrak{g}(K)) \cong H^{\mathcal{L}}(H(\mathfrak{g}(K))) \cong \text{Sym}(H(\mathfrak{g}(K))[1]).$$

From the discussion of Section 3.4, the product of $H^{\mathcal{L}}(\mathfrak{g}(K))$ is the map induced on Lie algebra homology by the n –disk algebra structure map of $\mathfrak{g}(K)$ corresponding to any embedding $\mathbb{R}^n \sqcup \mathbb{R}^n \rightarrow \mathbb{R}^n$, and any such structure map induces the fold map

$$H(\mathfrak{g}(K)) \oplus H(\mathfrak{g}(K)) \xrightarrow{+} H(\mathfrak{g}(K))$$

at the level of homology. Likewise, the coproduct is induced by the diagonal, and we recognize the standard bialgebra structure on Sym. The second isomorphism now follows by Proposition 5.2.

- (2) The cardinality grading is natural, and the product is the map induced on homology by the n –disk algebra structure map corresponding to any embedding $\mathbb{R}^n \sqcup \mathbb{R}^n \rightarrow \mathbb{R}^n$.

- (3) Since the coproduct preserves the cardinality grading on generators by definition, the claim follows from (2) and the fact that $H(\mathbb{F}_n(K))$ is a bialgebra.
- (4) The claim is immediate from the definition of the weight grading.
- (5) Since the coproduct preserves the cardinality grading on generators by definition, the claim follows from (4) and the fact that $H^{\mathcal{L}}(\mathfrak{g}(K))$ is a bialgebra. \square

The desired identification of bigradings now follows easily.

Proposition 5.4 *In the case $M = \mathbb{R}^n$, the isomorphisms of Theorems 1.1, 4.5 and 4.9 are isomorphisms of bigraded vector spaces.*

Proof We present the argument for Theorem 1.1, the others being essentially identical.

We follow the convention that a subscript indicates homological degree, a generator decorated with a tilde has weight 2 and an unadorned generator has weight 1. There is an isomorphism of bialgebras $H(\mathbb{F}_n(\mathbb{Q})) \cong \text{Sym}(V_n)$, where

$$V_n = \begin{cases} \mathbb{Q}\langle x_0 \rangle & \text{for } n \text{ odd,} \\ \mathbb{Q}\langle x_0, \tilde{y}_{n-1} \rangle & \text{for } n \text{ even.} \end{cases}$$

Identifying both sides of the isomorphism of Theorem 1.1 with $\text{Sym}(V_n)$, Proposition 5.2 permits us to view this isomorphism as an automorphism f of this graded bialgebra. Now, as a morphism of graded coalgebras, f takes primitives to primitives, so that there is an induced map $f|_{V_n}: V_n \rightarrow V_n$ of graded vector spaces, which we claim is a bigraded isomorphism. In the case of odd n , the claim is implied by the injectivity of $f|_{V_n}$, while in the even case we note that, for degree reasons, $f(x)$ is a scalar multiple of x and $f(\tilde{y})$ is a scalar multiple of \tilde{y} . By injectivity, this scalar is nonzero, and we conclude that $f|_{V_n}$ is a bigraded isomorphism.

Now, since f is also a map of algebras, we have $f(x_1 \cdots x_r) = f(x_1) \cdots f(x_r)$, which, together with the previous paragraph, shows that f preserves weight on monomials. Since monomials form a bihomogeneous basis and f is linear, the proof is complete. \square

5.2 Interlude: splitting configurations

Configuration spaces of different cardinalities are interrelated by splitting and forgetting maps inherited from the Cartesian product via the embedding $\text{Conf}_k(M) \rightarrow M^k$. This rich structure invites an inductive way of thinking that appears in one form or another in essentially every classical approach to these spaces; see [1] and [21] for the origins of this approach and [14] for a modern implementation.

In the setting of factorization homology, the importance of these splitting maps is that they assemble to form a coproduct, a shadow of which we have seen in the previous section, endowing $\mathbb{F}_n(K)$ with the structure of an n -disk algebra *in cocommutative coalgebras*. We will not need the full force of this statement, nor will we need the geometric interpretation of this coalgebra structure; nevertheless, we devote the remainder of this section to elucidating this interpretation, both for its general interest and for the motivation it provides for our proof of homological stability.

Remark 5.5 The constructions of this section are valid in more general stable settings than chain complexes, including the symmetric monoidal ∞ -category of spectra with smash product. We intend to return to this setting in future work.

The basic ingredient is the collection of natural transformations

$$s_{i,j}: \text{Conf}_k^{\text{fr}} \rightarrow \text{Conf}_i^{\text{fr}} \times \text{Conf}_j^{\text{fr}},$$

defined whenever $i + j = k$, which make the diagram

$$\begin{array}{ccc} \text{Conf}_k^{\text{fr}}(M) & \xrightarrow{(s_{i,j})_M} & \text{Conf}_i^{\text{fr}}(M) \times \text{Conf}_j^{\text{fr}}(M) \\ \downarrow & & \downarrow \\ \prod_k \text{Fr}_M & \xrightarrow{\cong} & \prod_i \text{Fr}_M \times \prod_j \text{Fr}_M \end{array}$$

commute; in other words,

$$(s_{i,j})_M(x_1, \dots, x_k) = ((x_1, \dots, x_i), (x_{i+1}, \dots, x_k)).$$

Given an $O(n)$ -module K , we have maps

$$s_{i,j}^K: \text{Conf}_k^{\text{fr}} \otimes K^{\otimes k} \xrightarrow{\delta_{i,j} \otimes 1} (\text{Conf}_i^{\text{fr}} \times \text{Conf}_j^{\text{fr}}) \otimes K^{\otimes k} \xrightarrow{\simeq} \text{Conf}_i^{\text{fr}} \otimes K^{\otimes i} \otimes \text{Conf}_j^{\text{fr}} \otimes K^{\otimes j},$$

which are $(\Sigma_i \times \Sigma_j) \times O(n)^k$ -equivariant. Taking $O(n)^k$ -coinvariants and using that induction is right adjoint to restriction for the inclusion $\Sigma_i \times \Sigma_j \rightarrow \Sigma_k$, we obtain by adjunction a Σ_k -equivariant map

$$\tilde{s}_{i,j}^K: \text{Conf}_k^{\text{fr}} \otimes_{O(n)^k} K^{\otimes k} \rightarrow \text{Ind}_{\Sigma_i \times \Sigma_j}^{\Sigma_k} (\text{Conf}_i^{\text{fr}} \otimes_{O(n)^i} K^{\otimes i} \otimes \text{Conf}_j^{\text{fr}} \otimes_{O(n)^j} K^{\otimes j}).$$

Finally, taking Σ_k -coinvariants and summing over k, i and j , we obtain a map

$$\begin{aligned} s^K: \bigoplus_{k \geq 0} (\text{Conf}_k^{\text{fr}} \otimes_{\Sigma_k \times O(n)^k} K^{\otimes k}) & \\ & \rightarrow \bigoplus_k \bigoplus_{i+j=k} (\text{Conf}_i^{\text{fr}} \otimes_{\Sigma_i \times O(n)^i} K^{\otimes i} \otimes \text{Conf}_j^{\text{fr}} \otimes_{\Sigma_j \times O(n)^j} K^{\otimes j}). \end{aligned}$$

Collecting terms and restricting to Disk_n , we recognize this as a monoidal natural transformation

$$s^K: \mathbb{F}_n(K) \rightarrow \mathbb{F}_n(K) \otimes \mathbb{F}_n(K).$$

The proof of homological stability given in the next section is completely internal to the Chevalley–Eilenberg complex, but the motivation behind it comes from thinking of the symmetric coproduct, given by splitting monomials in all possible ways, as corresponding to this geometric coproduct, given by splitting configurations in all possible ways. To see the connection, we recall that, in the approach of [14], stability is induced by the transfer maps

$$\begin{array}{ccc} H_*(\text{Conf}_{k+1}(M); \mathbb{Q}) & \xrightarrow{\sum_i (p_i)_*} & H_*(\text{Conf}_k(M); \mathbb{Q}) \\ \downarrow & & \downarrow \\ H_*(B_{k+1}(M); \mathbb{Q}) & \dashrightarrow^{\text{tr}} & H_*(B_k(M); \mathbb{Q}) \end{array}$$

where p_i denotes the projection that forgets x_i . In terms of our splitting maps, we have a factorization

$$\begin{array}{ccc} \text{Conf}_{k+1}(M) & \xrightarrow{p_i} & \text{Conf}_k(M) \\ \sigma_i \downarrow & & \uparrow \\ \text{Conf}_{k+1}(M) & \xrightarrow{s_{1,k}} & M \times \text{Conf}_k(M) \end{array}$$

where σ_i denotes the permutation that moves x_i to the first position while maintaining the relative order of the remaining points, and the unmarked arrow is the projection. The composite $s_{1,k}\sigma_i$ is a component of the coproduct defined above, and the projection away from the M factor corresponds at the level of homology to evaluating against the unit in $H^0(M; \mathbb{Q})$. Together, these observations suggest that homological stability should be induced taking a *cap product*. We realize this idea in the next section.

5.3 Stability

This section assembles the proof of Theorem 1.3. Throughout, unless otherwise noted, M will be connected, without boundary and of dimension $n > 1$. For the sake of brevity, we make the abbreviation

$$\mathfrak{g}_M = H_c^{-*}(M; \mathcal{L}(\mathbb{Q}^w[n-1])).$$

Let $\lambda \in H^0(M)$ denote the multiplicative unit. We view this cohomology class as a functional on $H_0(M) \cong H_c^0(M; \mathbb{Q}^w)[n]$ and hence, extending by zero, on $\text{CE}(\mathfrak{g}_M)$, since the former is canonically a summand of the underlying bigraded vector space of the

latter. Thus we may contemplate the *cap product* with this element, denoted $\lambda \frown (-)$, which is defined as the composite

$$\text{CE}(\mathfrak{g}_M) \cong \mathbb{Q} \otimes \text{CE}(\mathfrak{g}_M) \xrightarrow{\lambda \otimes \gamma} \text{CE}(\mathfrak{g}_M)^\vee \otimes \text{CE}(\mathfrak{g}_M) \otimes \text{CE}(\mathfrak{g}_M) \xrightarrow{\langle -, - \rangle \otimes \text{id}} \text{CE}(\mathfrak{g}_M).$$

Denote by $p \in H_c^n(M; \mathbb{Q}^w) \subset \text{CE}(\mathfrak{g}_M)$ the Poincaré dual of a point in M , which is well-defined since M is connected. Extend the set $\{1, p\}$ once and for all to a bihomogeneous basis \mathcal{B} for $\mathfrak{g}_M[1]$. Then the set of nonzero monomials in elements of \mathcal{B} form a bihomogeneous basis for $\text{CE}(\mathfrak{g}_M)$, and, under the resulting identification of this vector space with its dual, λ is identified with the dual functional to p . Since p is closed of degree 0 and weight 1, we conclude the following:

Proposition 5.6 $\lambda \frown (-)$ is a chain map of degree 0 and weight -1 .

There is a simple formula describing this map. Here and throughout, when we speak of divisibility, multiplication and differentiation in the Chevalley–Eilenberg complex, we refer only to the formal manipulation of bigraded polynomials; in particular, $\text{CE}(\mathfrak{g}_M)$ is *not* in general a differential graded algebra.

Proposition 5.7 *The formula*

$$\lambda \frown x = \frac{dx}{dp}$$

holds for all $x \in \text{CE}(\mathfrak{g}_M)$.

Proof Both sides are linear, so the claim is equivalent to the equality

$$\lambda \frown p^r y = r p^{r-1} y$$

whenever $r \geq 0$ and y is a monomial in elements of $\mathcal{B} \setminus \{p\}$. There are now two cases.

The first case is when y is a scalar, in which case we may assume by linearity that $y = 1$, so that $x = p^r$, and

$$\begin{aligned} \gamma(x) &= \gamma(p)^r \\ &= (p \otimes 1 + 1 \otimes p)^r \\ &= \sum_{i=0}^r \binom{r}{i} p^i \otimes p^{r-i}, \end{aligned}$$

so that

$$\lambda \frown x = \sum_{i=0}^r \binom{r}{i} \langle \lambda, p^i \rangle p^{r-i} = r p^{r-1}.$$

The second is when y is a monomial in $\mathcal{B} \setminus \{1, p\}$, in which case we may write

$$\gamma(y) = y \otimes 1 + 1 \otimes y + \sum_j y_j \otimes y'_j$$

with y_j and y'_j monomials in $\mathcal{B} \setminus \{1, p\}$. Then we have

$$\begin{aligned} \gamma(p^r y) &= \gamma(p)^r \gamma(y) \\ &= (p \otimes 1 + 1 \otimes p)^r \left(y \otimes 1 + 1 \otimes y + \sum_j y_j \otimes y'_j \right) \\ &= \sum_{i=0}^r \binom{r}{i} \left(p^i y \otimes p^{r-i} + p^i \otimes p^{r-i} y + \sum_j p^i y_j \otimes p^{r-i} y'_j \right), \end{aligned}$$

whence

$$\lambda \frown p^r y = \sum_{i=0}^r \binom{r}{i} \left(\langle \lambda, p^i y \rangle p^{r-i} + \langle \lambda, p^i \rangle p^{r-i} y + \sum_j \langle \lambda, p^i y_j \rangle p^{r-i} y'_j \right) = r p^{r-1} y,$$

since $p^i y$ is not a scalar multiple of p for any i , nor is $p^i y_j$ a scalar multiple of p for any (i, j) . □

Corollary 5.8 *The chain map $\lambda \frown (-)$ is surjective.*

Proof It suffices to show that a general monomial in elements of \mathcal{B} lies in the image. Such a monomial may be written as $p^r y$ with $r \geq 0$ and y a monomial in elements of $\mathcal{B} \setminus \{p\}$. We then have

$$\frac{d}{dp} \left(\frac{1}{r+1} p^{r+1} y \right) = p^r y. \quad \square$$

The central observation behind our approach to stability is the following.

Proposition 5.9 *Let x be a nonzero monomial in $\text{CE}(\mathfrak{g}_M)$. Then x is divisible by p provided either*

- $\text{wt}(x) > |x| + 1$ and M is an orientable surface, or
- $\text{wt}(x) > |x|$ and M is not an orientable surface.

Proof Suppose $\text{wt}(x) > |x|$, and write $x = x_1 \cdots x_r$ with $x_i \in \mathcal{B}$. Then $\text{wt}(x_j) > |x_j|$ for some j . Since $x_j \in \mathfrak{g}_M[1]$, the weight of this element is either 1 or 2.

In the first case, $x_j \in H_c^{-*}(M; \mathbb{Q}^w)[n]$, and we have

$$|x_j| < \text{wt}(x_j) = 1 \quad \text{implies} \quad |x_j| = 0,$$

since $H_c^{-*}(M; \mathbb{Q}^w)[n]$ is concentrated in degrees $0 \leq * \leq n$. But $H_c^n(M; \mathbb{Q}^w)$ is one-dimensional on the class p , so x_j is a scalar multiple of p .

In the second case, $x_j \in H_c^{-*}(M; \mathbb{Q})[2n - 1]$, and we have $|x_j| < 2$. Because $H_c^{-*}(M; \mathbb{Q})[2n - 1]$ is concentrated in degrees $n - 1 \leq * \leq 2n - 1$, we conclude that $x_j = 0$ provided $n \neq 2$ (recall that we have already assumed $n > 1$). Thus $x = 0$, which is a contradiction. This proves the claim when M is not a surface.

If M is a nonorientable surface, then $H_c^2(M; \mathbb{Q}) \cong H_0(M; \mathbb{Q}^w) = 0$, and therefore $H_c^{-*}(M; \mathbb{Q})[2n - 1]$ is concentrated in degrees 2 and 3. Thus, in this case as well, we have a contradiction.

Assume now that M is an orientable surface and $\text{wt}(x) > |x| + 1$. As before, write $x = x_1 \cdots x_r$ and choose x_j with $\text{wt}(x_j) > |x_j|$, and assume that x_j is not a scalar multiple of p . Then by the argument above, $\text{wt}(x_j) = 2$, so $|x_j| = 1$, since $H_c^{-*}(M; \mathbb{Q})[3]$ is concentrated in degrees $1 \leq * \leq 3$.

Now, the monomial $x' = x_1 \cdots \hat{x}_j \cdots x_r$ has the property that

$$\text{wt}(x') = \text{wt}(x) - 2 > |x| - 1 = |x'|,$$

so there is some x_i with $i \neq j$ and $\text{wt}(x_i) > |x_i|$. If x_i is a scalar multiple of p , we are finished; otherwise, repeating the same argument shows that x_i has degree 1. But $H_c^2(M; \mathbb{Q}) \cong H_0(M; \mathbb{Q})$ is one-dimensional, so that x_i is a scalar multiple of x_j , and x is divisible by x_j^2 . Since x_j is of odd degree, this implies that $x = 0$, which is a contradiction. □

We are now equipped to prove Theorem 1.3. Denote by $C(k)$ the subcomplex of the Chevalley–Eilenberg complex spanned by the weight- k monomials. Taking the cap product with 1 restricts to a map $\Phi_k: C(k + 1) \rightarrow C(k)$, and we aim to show that this map induces an isomorphism in homology in the specified range.

Recall that the r^{th} brutal truncation of a chain complex V is the chain complex $\tau_{\leq r} V$ whose underlying graded vector space is

$$(\tau_{\leq r} V)_i = \begin{cases} V_i & \text{if } i \leq r, \\ 0 & \text{otherwise.} \end{cases}$$

and whose differential is the restriction of the differential of V . Truncation is a functor on chain complexes in the obvious way.

We make use of the following elementary fact.

Proposition 5.10 *Let $f: V \rightarrow W$ be a surjective chain map such that $\tau_{\leq r} f$ is a chain isomorphism. Then f is a homology isomorphism through degree r and a homology surjection in degree $r + 1$.*

Proof From the definition of the brutal truncation, it is immediate that f is a homology isomorphism through degree $r - 1$. Moreover, f induces a bijection on r -cycles and an injection on r -boundaries.

To show that f is a homology isomorphism in degree r , it suffices to show that $f^{-1}(w)$ is a boundary if $w \in W_r$ is a boundary. Write $w = du$; then, by surjectivity, there is some $\tilde{u} \in V_r$ such that $f(\tilde{u}) = u$, and

$$f(d\tilde{u}) = df(\tilde{u}) = du = w \quad \text{implies} \quad f^{-1}(w) = d\tilde{u},$$

as desired.

To show that f is a homology surjection in degree $r + 1$, let $v \in W_{r+1}$ be a cycle. By surjectivity, $v = f(\tilde{v})$, and it will suffice to show that \tilde{v} is a cycle, for which we have

$$f(d\tilde{v}) = df(\tilde{v}) = dv = 0 \quad \text{implies} \quad d\tilde{v} = f^{-1}(0) = 0. \quad \square$$

Proof of Theorem 1.3 Assume first that M is not an orientable surface. By the previous proposition and Corollary 5.8, we are reduced to showing that $\tau_{\leq k} \Phi_k$ is a chain isomorphism. To see this, let $x \in \tau_{\leq k} C(k + 1)$ be a monomial. Then $\text{wt}(x) > |x|$, so that $x = p^r y$ with $r > 0$ and y a monomial in $\mathcal{B} \setminus \{p\}$ by Proposition 5.9. By Proposition 5.7, $\Phi_k(x) = rp^{r-1}y$, so $\tau_{\leq k} \Phi_k$ maps distinct elements of our preferred basis for $C(k + 1)$ to nonzero scalar multiples of distinct elements of our preferred basis for $C(k)$, which implies that $\tau_{\leq k} \Phi_k$ is injective. But Φ and hence Φ_k are surjective by Corollary 5.8, so $\tau_{\leq k} \Phi_k$ is as well.

Assume now that M is an orientable surface. For the same reason, we are reduced to showing that $\tau_{k-1} \Phi_k$ is a chain isomorphism, which is accomplished by the same argument, using the other half of Proposition 5.9. □

Remark 5.11 Let \mathbb{K} denote the Klein bottle. As shown in Section 6,

$$\dim H_*(B_k(\mathbb{K}); \mathbb{Q}) = \begin{cases} 1 & i \in \{0, 1, 2, k + 1\}, \\ 2 & 3 \leq i \leq k, \\ 0 & \text{else.} \end{cases}$$

In particular, $H_{k+1}(B_{k+1}(\mathbb{K}); \mathbb{Q}) \not\cong H_{k+1}(B_k(\mathbb{K}); \mathbb{Q})$, so our bound is sharp in the sense that no better stable range holds for all manifolds that are not orientable surfaces.

Remark 5.12 If M is orientable and $H_*(M; \mathbb{Q}) = 0$ for $1 \leq * \leq r - 1$, then $H_c^{-*}(M; \mathbb{Q}) = 0$ for $n - r + 1 \leq -* \leq n - 1$, and the argument of Proposition 5.9 shows that a monomial x is divisible by p provided its weight is greater than $\frac{|x|}{r} + 1$. This improved estimate leads to an improved stable range, as in [14, Proposition 4.1].

Remark 5.13 In [37], factorization homology is used to obtain homological stability results for various constructions on open manifolds. The approach there is through certain “partial algebras” and appears unrelated to ours.

6 Examples

We now present a selection of computations illustrating the following general procedure for determining the rational homology of the configuration space of k points in an n -manifold M :

- (1) compute the compactly supported cohomology of M , twisted if necessary;
- (2) compute the Lie algebra homology of $H_c^{-*}(M; \mathcal{L}(\mathbb{Q}^w[n-1]))$;
- (3) count basis elements of weight k .

It is worth noting that the Chevalley–Eilenberg complex allows one to obtain answers simultaneously for all k , reducing an infinite sequence of computations to one.

The computations of this section are all relatively elementary, and one can do better with more effort. In [19], this approach is used to determine the Betti numbers of $B_k(\Sigma)$ for every surface Σ .

Convention In the following examples, a variable decorated with a tilde has weight 2, while an unadorned variable has weight 1.

6.1 Punctured euclidean space

As a warm-up and base case, we recover the classical computation of $H_*(B_k(\mathbb{R}^n); \mathbb{Q})$. Since there are no cup products in the compactly supported cohomology of \mathbb{R}^n , there are no differentials in the corresponding Chevalley–Eilenberg complex. Thus $H_*(B_k(\mathbb{R}^n); \mathbb{Q})$ is identified with the subspace of $\mathbb{Q}[x]$ spanned by x^k when n is odd, while for n even, the identification is with the subspace of

$$\mathbb{Q}[x] \otimes \Lambda[\tilde{x}], \quad |x| = 0, |\tilde{x}| = n - 1,$$

spanned by elements of weight k , a basis for which is given by $\{x^k, x^{k-2}\tilde{x}\}$. We conclude, for all $k > 1$, that

$$H_*(B_k(\mathbb{R}^n); \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & \text{for } n \text{ odd,} \\ \mathbb{Q} \oplus \mathbb{Q}[n-1] & \text{for } n \text{ even.} \end{cases}$$

Now, choose $\bar{p} = \{p_1, \dots, p_m\} \in \mathbb{R}^n$. There is a homotopy equivalence $(\mathbb{R}^n \setminus \bar{p})^+ \simeq S^n \vee (S^1)^{\vee m}$, so that $H_c^{-*}(\mathbb{R}^n \setminus \bar{p}; \mathbb{Q}) \cong \mathbb{Q}^m[-1] \oplus \mathbb{Q}[-n]$. There are no cup products, so there can be no differentials.

If n is odd, Theorem 1.1 identifies $H_*(\mathbb{R}^n \setminus \bar{p}; \mathbb{Q})$ with the weight- k part of

$$\mathbb{Q}[x, y_1, \dots, y_m], \quad |x| = 0, |y_i| = n - 1,$$

and an easy induction now shows that

$$\dim H_*(B_k(\mathbb{R}^n \setminus \bar{p}); \mathbb{Q}) = \begin{cases} \binom{m+i-1}{i} & \text{for } * = i(n-1), 0 \leq i \leq k, \\ 0 & \text{otherwise.} \end{cases}$$

(It is helpful to recall that $\binom{m+i-1}{i}$ is the number of ways to choose i not-necessarily-distinct elements from a set of m elements.)

If n is even, then the corresponding vector space is the weight- k part of

$$\mathbb{Q}[x, \tilde{y}_1, \dots, \tilde{y}_m] \otimes \Lambda[\tilde{x}, y_1, \dots, y_m], \quad |x| = 0, |y_i| = |\tilde{x}| = n - 1, |\tilde{y}_i| = 2n - 2.$$

Counting inductively in terms of less punctured Euclidean spaces, one finds that

$$H_*(B_k(\mathbb{R}^n \setminus \bar{p}); \mathbb{Q}) \cong \bigoplus_{l=0}^k \bigoplus_{j_1 + \dots + j_m = l} H_{*-l(n-1)}(B_{k-l}(\mathbb{R}^n); \mathbb{Q}),$$

from which it follows easily that

$$\dim H_*(B_k(\mathbb{R}^n \setminus \bar{p}); \mathbb{Q}) = \begin{cases} \binom{m+i-1}{m-1} + \binom{m+i-2}{m-1} & \text{for } * = i(n-1), 0 \leq i < k, \\ \binom{m+k-1}{m-1} & \text{for } * = k(n-1), \\ 0 & \text{otherwise.} \end{cases}$$

(It is helpful to recall that $\binom{m+i-1}{m-1}$ is the number of ways to write i as the sum of m nonnegative integers.)

It should be clear from this example that Theorem 1.1 reduces calculations to counting problems whenever n is odd or the relevant compactly supported cohomology has no cup products.

6.2 Punctured torus

Since $H_c^{-*}(T^2 \setminus \text{pt}; \mathbb{Q}) \cong \tilde{H}^{-*}(T^2; \mathbb{Q})$, the relevant Lie algebra is isomorphic to

$$\mathfrak{h} \oplus \mathbb{Q}\langle \tilde{a}, \tilde{b}, c \rangle,$$

where $\mathfrak{h} = \mathbb{Q}\langle a, b, \tilde{c} \rangle$ as a vector space,

$$|a| = |b| = |\tilde{c}| = 0, \quad |\tilde{a}| = |\tilde{b}| = 1, \quad |c| = -1,$$

and the bracket is defined by the equation

$$[a, b] = \tilde{c}.$$

The Lie homology of \mathfrak{h} is calculated by the complex

$$(\Lambda[x, y, \bar{z}], d(xy) = \bar{z}),$$

(where for ease of notation we have set $x = \sigma a$ and so on), a basis for the homology of which is easily seen to be given by the image in homology of the set $\{1, x, y, x\bar{z}, y\bar{z}, xy\bar{z}\}$. Thus we have an identification of $H_*(B_k(T^2 \setminus \text{pt}); \mathbb{Q})$ with the weight- k part of

$$\mathbb{Q}\langle 1, x, y, x\bar{z}, y\bar{z}, xy\bar{z} \rangle \otimes \mathbb{Q}[\tilde{x}, \tilde{y}, z], \quad |z| = 0, |x| = |y| = |\bar{z}| = 1, |\tilde{x}| = |\tilde{y}| = 2.$$

Counting, we find that

$$\dim H_*(B_k(T^2 \setminus \text{pt}); \mathbb{Q}) = \begin{cases} \frac{3i-1}{2} + 1 & \text{for } * = 2i + 1 < k, \\ \frac{3i}{2} + 1 & \text{for } * = 2i < k, \\ k + 1 & \text{for } * = k \text{ odd,} \\ \frac{k}{2} + 1 & \text{for } * = k \text{ even,} \\ 0 & \text{otherwise.} \end{cases}$$

An amusing comparison can be seen by taking $k = 2$ in the above formula, which yields

$$H_*(B_2(T^2 \setminus \text{pt}); \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}^2[1] \oplus \mathbb{Q}^2[2].$$

On the other hand, from the preceding example, one calculates that

$$H_*(B_2(\mathbb{R}^2 \setminus \{p_1, p_2\}); \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}^3[1] \oplus \mathbb{Q}^3[2].$$

Thus, despite the fact that the punctured torus and the twice-punctured plane are homotopy equivalent, having $S^1 \vee S^1$ as a common deformation retract, their configuration spaces are not homotopy equivalent.

6.3 Real projective space

Let n be even, so that $\mathbb{R}P^n$ is nonorientable. Then, as a ring, $H_c^{-*}(\mathbb{R}P^n; \mathbb{Q}) \cong \mathbb{Q}$, and the Lie homology of interest is $H^{\mathcal{L}}(\mathcal{L}(\mathbb{Q}[n-1])) \cong \mathbb{Q} \oplus \mathbb{Q}[n]$, whence, for $k > 1$,

$$H_*(B_k(\mathbb{R}P^n); \mathbb{Q}^w) = 0.$$

As for the untwisted homology, we note that $H_c^{-*}(\mathbb{R}P^n; \mathbb{Q}^w) \cong \mathbb{Q}[-n]$ by Poincaré duality, so that the cup product map $H_c^{-*}(\mathbb{R}P^n; \mathbb{Q}^w)^{\otimes 2} \rightarrow H_c^{-*}(\mathbb{R}P^n; \mathbb{Q})$ is trivial for degree reasons. Thus

$$H_c^{-*}(\mathbb{R}P^n; \mathcal{L}(\mathbb{Q}^w[n-1])) \cong \mathbb{Q}[-1] \oplus \mathbb{Q}[2n-2]$$

is abelian, so that $H_*(B_k(\mathbb{R}P^n); \mathbb{Q})$ is isomorphic to the weight- k part of

$$\mathbb{Q}[x] \otimes \Lambda[\tilde{y}], \quad |x| = 0, |\tilde{y}| = 2n - 1.$$

Hence for all $k > 1$,

$$H_*(B_k(\mathbb{R}P^n); \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}[2n - 1].$$

See [55] for an alternate method of computation in the case $n = 2$.

6.4 Klein bottle, twisted

Let \mathbb{K} denote the Klein bottle. Then $H_c^{-*}(\mathbb{K}; \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}[-1]$, with the generator in degree zero acting as a unit for the multiplication. As a vector space, the Lie algebra in question is $\mathfrak{g} := \mathbb{Q}\langle a, \tilde{a}, b, \tilde{b} \rangle$, where $|b| = 0$, $|a| = |\tilde{b}| = 1$ and $|\tilde{a}| = 2$, and the bracket is defined by the equations

$$[a, a] = \tilde{a}, \quad [a, b] = -\tilde{b}.$$

The subspace spanned by $\{b, \tilde{b}\}$ is an ideal realizing \mathfrak{g} as an extension

$$0 \rightarrow \mathbb{Q}\langle b, \tilde{b} \rangle \rightarrow \mathfrak{g} \rightarrow \mathcal{L}(\mathbb{Q}\langle a \rangle) \rightarrow 0,$$

so that we may avail ourselves of the Lyndon–Hochschild–Serre spectral sequence

$$E_{p,q}^2 \cong H_p^{\mathcal{L}}(\mathcal{L}(\mathbb{Q}\langle a \rangle); H_q^{\mathcal{L}}(\mathbb{Q}\langle b, \tilde{b} \rangle)) \implies H_{p+q}^{\mathcal{L}}(\mathfrak{g}).$$

There are no differentials for degree reasons, and the E^2 page is computed as the homology of the complex

$$0 \rightarrow \mathbb{Q}\langle a \rangle[1] \otimes \text{Sym}(\mathbb{Q}\langle b, \tilde{b} \rangle[1]) \rightarrow \text{Sym}(\mathbb{Q}\langle b, \tilde{b} \rangle[1]) \rightarrow 0,$$

where the differential is the action of a . It follows that a basis for $H^{\mathcal{L}}(\mathfrak{g})$ is given by $\{\sigma a \otimes (\sigma \tilde{b})^i, \sigma b \otimes (\sigma \tilde{b})^j \mid i, j \geq 0\}$. Counting monomials of weight k , we find that

$$H_*(B_k(\mathbb{K}); \mathbb{Q}^w) \cong \begin{cases} \mathbb{Q}[k] \oplus \mathbb{Q}[k + 1] & \text{for } k \text{ odd,} \\ 0 & \text{for } k \text{ even.} \end{cases}$$

6.5 Nonorientable surfaces

Let $N_h = (\mathbb{R}P^2)^{\#h}$. Using the method of the previous example, one could proceed to obtain a general formula for the twisted homology of $B_k(N_h)$. Here we will determine the corresponding untwisted homology. We have

$$H_c^{-*}(N_h; \mathbb{Q}) \cong \mathbb{Q} \oplus \mathbb{Q}[-1]^{h-1}, \quad H_c^{-*}(N_h; \mathbb{Q}^w) \cong \mathbb{Q}[-1]^{h-1} \oplus \mathbb{Q}[-2],$$

so that there can be no cup products. Thus $H_*(B_k(N_h); \mathbb{Q})$ is the weight- k part of

$$\mathbb{Q}[x, \tilde{y}_1, \dots, \tilde{y}_{h-1}] \otimes \Lambda[\tilde{z}, w_1, \dots, w_{h-1}], \quad |x| = 0, |w_i| = 1, |\tilde{y}_i| = 2, |\tilde{z}| = 3.$$

Counting inductively as in the example of punctured Euclidean space, we find that

$$H_*(B_k(N_h); \mathbb{Q}) \cong \bigoplus_{l=0}^k \bigoplus_{j_1+\dots+j_{h-1}=l} H_{*-l}(B_{k-l}(\mathbb{R}\mathbb{P}^2); \mathbb{Q}),$$

from which it follows that

$$\dim H_*(B_k(N_h); \mathbb{Q}) = \begin{cases} \binom{h+*-2}{h-2} + \binom{h+*-5}{h-2} & \text{for } * \leq k, \\ \binom{h+*-5}{h-2} & \text{for } * = k + 1, \\ 0 & \text{otherwise.} \end{cases}$$

6.6 Open and closed Möbius band

Let \mathbb{M} denote the closed Möbius band. Then since \mathbb{M} has the same compactly supported cohomology ring as the Klein bottle, our earlier calculation shows that

$$\tilde{H}_*(B_k(\mathbb{M}, \partial\mathbb{M}); \mathbb{Q}^w) \cong \begin{cases} \mathbb{Q}[k] \oplus \mathbb{Q}[k + 1] & \text{for } k \text{ odd,} \\ 0 & \text{for } k \text{ even.} \end{cases}$$

On the other hand, by Poincaré duality, we have $H_c^{-*}(\mathbb{M}; \mathbb{Q}^w) = 0$, and hence $H_c^{-*}(\mathbb{M}; \mathcal{L}(\mathbb{Q}^w[1])) \cong H^{-*}(\mathbb{M}; \mathbb{Q})[2]$ is abelian, and $\tilde{H}_*(B_k(\mathbb{M}, \partial\mathbb{M}); \mathbb{Q})$ is the weight- k part of

$$\mathbb{Q}[\tilde{x}] \otimes \Lambda[\tilde{y}], \quad |\tilde{x}| = 2, |\tilde{y}| = 3,$$

so

$$\tilde{H}_*(B_k(\mathbb{M}, \partial\mathbb{M}); \mathbb{Q}) \cong \begin{cases} 0 & \text{for } k \text{ odd,} \\ \mathbb{Q}[k] \oplus \mathbb{Q}[k + 1] & \text{for } k \text{ even.} \end{cases}$$

The situation with the corresponding open manifold is quite different. We have $H_c^{-*}(\mathring{\mathbb{M}}; \mathbb{Q}) = 0$ since $(\mathring{\mathbb{M}})^+ \cong \mathbb{R}\mathbb{P}^2$, so

$$H_*(B_k(\mathring{\mathbb{M}}); \mathbb{Q}^w) = 0$$

for all $k > 1$. On the other hand, $H_c^{-*}(\mathring{\mathbb{M}}; \mathbb{Q}^w) \cong \mathbb{Q}[-1] \oplus \mathbb{Q}[-2]$ by Poincaré duality, so that $H_*(B_k(\mathring{\mathbb{M}}); \mathbb{Q})$ is the weight- k part of

$$\mathbb{Q}[x] \otimes \Lambda[y], \quad |x| = 0, |y| = 1,$$

whence

$$H_*(B_k(\mathring{\mathbb{M}}); \mathbb{Q}) \cong \mathbb{Q}[0] \oplus \mathbb{Q}[1]$$

for all $k \geq 1$.

7 Two formality results

In this final section, we supply the remaining two ingredients in the proof of Theorem 1.1. Although unrelated to each other, these formality statements may be of independent interest.

7.1 The $O(n)$ -equivariant sphere

Since the reduced homology of S^n is one-dimensional, any choice of representative of a homology generator defines a quasi-isomorphism

$$C_*(S^n) \simeq \mathbb{Z} \oplus \mathbb{Z}[n].$$

The goal of this section is to prove that, rationally, this equivalence can be made $O(n)$ -equivariant.

Theorem 7.1 *There is an equivalence of $O(n)$ -modules*

$$C_*(S^n; \mathbb{Q}) \simeq \mathbb{Q} \oplus \mathbb{Q}^{\det}[n].$$

The proof has three main ingredients, the first of which is rational homotopy theory. We consider the Borel construction

$$\hat{\xi}: ESO(n) \times_{SO(n)} S^n \rightarrow BSO(n)$$

where $SO(n)$ acts on $S^n \cong (\mathbb{R}^n)^+$ by extension of its canonical action on \mathbb{R}^n . In other words, $\hat{\xi}$ is the fiberwise one-point compactification of the universal oriented n -plane bundle ξ . We denote by $E(\hat{\xi})$ the total space of this sphere bundle.

Sphere bundles over simply connected spaces admit particularly simple rational descriptions. According to [22, Sections 15(a)–(b)], we have the following commutative diagram, whose terms we will explain presently:

$$\begin{array}{ccccc}
 (\text{Sym}(W_n), d_1) & \xrightarrow{\sim} & A_{\text{PL}}(S^n) & \xrightarrow{\sim} & C^{-*}(S^n; \mathbb{Q}) \\
 \uparrow & & \uparrow & & \uparrow \\
 (S \otimes \text{Sym}(W_n), d_1 + d_2) & \xrightarrow{\sim} & A_{\text{PL}}(E(\hat{\xi})) & \xrightarrow{\sim} & C^{-*}(E(\hat{\xi}); \mathbb{Q}) \\
 \uparrow & & \uparrow & & \uparrow \\
 S & \xrightarrow{\sim} & A_{\text{PL}}(BSO(n)) & \xrightarrow{\sim} & C^{-*}(BSO(n); \mathbb{Q})
 \end{array}$$

In this diagram,

- (1) $S := H^{-*}(BSO(n); \mathbb{Q})$ is a polynomial algebra,
- (2) A_{PL} denotes the functor of PL de Rham forms,

- (3) the horizontal arrows in the right-hand column are components of the natural quasi-isomorphism $\phi: A_{PL} \rightarrow C^{-*}$ given by integrating forms over simplices,
- (4) each term appearing in the leftmost column is a Sullivan model for the corresponding space, and
- (5) W_n denotes the graded vector space

$$W_n = \begin{cases} \mathbb{Q}\langle x_{-n} \rangle & \text{for } n \text{ odd,} \\ \mathbb{Q}\langle x_{-n}, y_{-2n+1} \rangle & \text{for } n \text{ even,} \end{cases}$$

the differential d_1 is defined by the equation $d_1(y) = x^2$ and the differential d_2 is specified by its value on y , which is an element of P determined by the bundle $\hat{\xi}$.

We direct the reader to [11] for more on (1), and to [22, Sections 10(c), 10(e), 12, 15(b)], respectively, for more on (2)–(5). The reader is advised that, although we have maintained our convention of homological grading, the prevailing convention in rational homotopy theory is cohomological.

The second ingredient is the theory of A_∞ -algebras and their modules, for which we refer the reader to [34]. The relevance here is that, according to [12, Section 3.1], the integration map ϕ extends to a map of A_∞ -algebras (referred to in [12] as “strongly homotopic differential algebras”), so that $C^{-*}(E(\hat{\xi}); \mathbb{Q})$ becomes an A_∞ - S -module via the bottom composite in the above diagram.

Proposition 7.2 *There is a quasi-isomorphism of A_∞ - S -modules*

$$S \oplus S[-n] \xrightarrow{\sim} C^{-*}(E(\hat{\xi}); \mathbb{Q}).$$

Proof The fiberwise basepoint furnishes $\hat{\xi}$ with a section, and the Gysin sequence now implies that the top map in the commuting diagram

$$\begin{array}{ccc} S \oplus S[-n] & \longrightarrow & (S \otimes \text{Sym}(W_n), d_1 + d_2) \\ \uparrow & & \uparrow \\ S & \xlongequal{\quad} & S \end{array}$$

is a quasi-isomorphism. Combining this diagram with the previous yields the result. \square

The third ingredient is the Koszul duality between modules for the symmetric algebra S and modules for the exterior algebra Λ on the same generators with degrees shifted by 1. According to [11], there is a Hopf algebra isomorphism $\Lambda \cong H_*(\text{SO}(n); \mathbb{Q})$, where the latter carries the Pontryagin product induced by the group structure of $\text{SO}(n)$.

Koszul duality is the algebraic avatar of the correspondence between $SO(n)$ -spaces and spaces fibered over $BSO(n)$ witnessed by the Borel construction. There are many variations on this theme; the relevant facts for our purposes are the following, which are extracted from [26, Theorem 1.2 and Proposition 3.1]; see also [30]. Our notation differs slightly from that in [26], and we maintain the terminology of A_∞ -modules rather than “weak modules”.

Theorem 7.3 (Franz; Goresky, Kottwitz and MacPherson) *There is a functor h from A_∞ - S -modules to A_∞ - Λ -modules with the following properties:*

- (1) *Let $\pi: X \rightarrow BSO(n)$ be a space over $BSO(n)$. Then the A_∞ - Λ -modules $h(C^{-*}(X))$ and $C^{-*}(\text{hofiber}(\pi))$ are connected by a zig-zag of natural quasi-isomorphisms.*
- (2) *Let V be a graded vector space. Then $h(S \otimes V) \cong V$, where V is regarded as a trivial A_∞ - Λ -module.*

Proposition 7.4 *There is an equivalence of $SO(n)$ -modules*

$$C_*(S^n; \mathbb{Q}) \simeq \mathbb{Q} \oplus \mathbb{Q}[n],$$

where the latter is regarded as a trivial $SO(n)$ -module.

Proof Both of the $SO(n)$ -modules in question are dualizable objects of $\text{Ch}_{\mathbb{Q}}$, so it suffices to exhibit an $SO(n)$ -equivalence $C^{-*}(SO(n); \mathbb{Q}) \simeq \mathbb{Q} \oplus \mathbb{Q}[-n]$ between the duals. By [42, Theorem 4.3.3.17], the homotopy category of the ∞ -category of $SO(n)$ -modules coincides with the homotopy category obtained from the model category of $C_*(SO(n); \mathbb{Q})$ -modules equipped with the usual model structure on modules over a differential graded algebra. By [34, Section 4.3], this homotopy category in turn coincides with the full subcategory of the homotopy category of A_∞ - $C_*(SO(n); \mathbb{Q})$ -modules spanned by the “homologically unital modules”, so that, since the modules in question are homologically unital, it will suffice to produce to an isomorphism in the homotopy category of A_∞ -modules. By [34, Section 6.2], it suffices to produce an isomorphism in the homotopy category of A_∞ - Λ -modules after restricting along the A_∞ -quasi-isomorphism $\Lambda \rightarrow C_*(SO(n); \mathbb{Q})$ of [26]. For this, we apply the Koszul duality of Theorem 7.3 to the A_∞ -quasi-isomorphism of Proposition 7.2, yielding the zig-zag of A_∞ -quasi-isomorphisms

$$\mathbb{Q} \oplus \mathbb{Q}[-n] \simeq h(S \oplus S[-n]) \rightarrow h(C^{-*}(E(\hat{\xi})); \mathbb{Q}) \simeq C^{-*}(S^n; \mathbb{Q}). \quad \square$$

Proof of Theorem 7.1 We explain the following diagram of $O(n)$ -modules:

$$\mathbb{Q} \oplus \mathbb{Q}^{\det}[n] \rightarrow \mathbb{Q}[C_2] \oplus \mathbb{Q}[C_2][n] \simeq \mathbb{Q}[C_2] \otimes C_*(S^n; \mathbb{Q}) \rightarrow C_*(S^n; \mathbb{Q}).$$

- (1) Let e and σ denote the basis elements of $\mathbb{Q}[C_2]$ corresponding to the identity and generator, respectively. The left-hand map sends $1 \in \mathbb{Q}$ to $\frac{e+\sigma}{2}$ and $1 \in \mathbb{Q}^{\det}$ to $\frac{e-\sigma}{2}$. This is a map of C_2 -modules and therefore of $O(n)$ -modules, since $O(n)$ acts on both domain and codomain by restriction along the determinant.
- (2) Fixing a choice of isomorphism $O(n) \cong C_2 \times \text{SO}(n)$, we obtain an isomorphism $C_*(O(n); \mathbb{Q}) \cong \mathbb{Q}[C_2] \otimes C_*(\text{SO}(n); \mathbb{Q})$ of $O(n)$ -modules. The middle equivalence is now obtained by applying the functor of induction from $\text{SO}(n)$ to $O(n)$ to the equivalence of Proposition 7.4.
- (3) The right-hand arrow is the counit of the induction-restriction adjunction.

Applying homology yields an isomorphism, completing the proof. □

7.2 Two-step nilpotent Lie algebras

In this section, we prove that the Lie algebras of interest to us are formal.

Proposition 7.5 *Let K be either \mathbb{Q} or \mathbb{Q}^{sgn} . For any $r \in \mathbb{Z}$ and any manifold M , the Lie algebra $\text{Map}^{C^2}(\tilde{M}^+, \mathcal{L}(K[r]))$ is formal.*

The proof will rely on the following technical result.

Proposition 7.6 *Let*

$$0 \rightarrow \mathfrak{h} \rightarrow \mathfrak{e} \rightarrow \mathfrak{g} \rightarrow 0$$

be an exact sequence of Lie algebras in $\text{Ch}_{\mathbb{Q}}$ with \mathfrak{g} and \mathfrak{h} abelian. Assume that \mathfrak{g} acts trivially on \mathfrak{h} and that the underlying sequence of chain complexes splits. Then \mathfrak{e} is formal.

Proof The hypotheses imply that the bracket on $\mathfrak{e} \cong \mathfrak{g} \oplus \mathfrak{h}$ is given by

$$[(g_1, h_1), (g_2, h_2)] = f(g_1, g_2)$$

for some (not uniquely defined) map $f: \text{Sym}^2(\mathfrak{g}[1])[-2] \rightarrow \mathfrak{h}$, and the bracket on $H(\mathfrak{e}) \cong H(\mathfrak{g}) \oplus H(\mathfrak{h})$ is determined in the same way by f_* .

Choose quasi-isomorphisms $\varphi: \mathfrak{g} \rightarrow H(\mathfrak{g})$ and $\psi: \mathfrak{h} \rightarrow H(\mathfrak{h})$. Without loss of generality, we may assume that both maps induce the identity on homology. Let $\bar{\psi}$ be a quasi-inverse to ψ . Then $(\bar{\psi} \circ f_* \circ \varphi^{\wedge 2})_* = f_*$, so

$$\bar{\psi} \circ f_* \circ \varphi^{\wedge 2} - f = d_{\mathfrak{h}}G + Gd_{\text{Sym}}$$

for some homotopy operator $G: \text{Sym}^2(\mathfrak{g}[1])[-2] \rightarrow \mathfrak{h}[-1]$.

Now, since \mathfrak{g} is abelian and acts trivially, this equation may be written as

$$D(G) = \bar{\psi} \circ f_* \circ \varphi^{\wedge 2} - f,$$

where D denotes the differential in the Chevalley–Eilenberg cochain complex computing $H_{\mathcal{L}}^*(\mathfrak{g}, \mathfrak{h})$. Since extensions of \mathfrak{g} by the module \mathfrak{h} are classified by $H_{\mathcal{L}}^2(\mathfrak{g}, \mathfrak{h})$, it follows that f and $\bar{\psi} \circ f_* \circ \varphi^{\wedge 2}$ determine isomorphic extensions, so that we may take $f = \bar{\psi} \circ f_* \circ \varphi^{\wedge 2}$ after choosing a different splitting. But then $\psi \circ f = f_* \circ \varphi^{\wedge 2}$, so that the composite

$$\mathfrak{e} \xrightarrow{\cong} \mathfrak{g} \oplus \mathfrak{h} \xrightarrow{(\varphi, \psi)} H(\mathfrak{g}) \oplus (\mathfrak{h}) \xrightarrow{\cong} H(\mathfrak{e})$$

is a map of Lie algebras. Since it is also a quasi-isomorphism of chain complexes, the proof is complete. \square

Proof of Proposition 7.5 The exact sequence

$$0 \rightarrow \mathfrak{h} \rightarrow \mathcal{L}(K[r]) \rightarrow K[r] \rightarrow 0$$

satisfies the hypotheses of Proposition 7.6, where

$$\mathfrak{h} = \begin{cases} K^{\otimes 2}[2r] & \text{for } r \text{ odd,} \\ 0 & \text{for } r \text{ even.} \end{cases}$$

By Proposition 3.18, we have

$$\text{Map}^{C_2}(\tilde{M}^+, \mathcal{L}(K[r])) \simeq (A_{\text{PL}}(\tilde{M}^+) \otimes \mathcal{L}(K[r]))^{C_2}.$$

Since the operations of tensoring with the commutative algebra $A_{\text{PL}}(\tilde{M}^+)$ and taking C_2 fixed points preserve the hypotheses of Proposition 7.6, the claim follows. \square

Remark 7.7 Proposition 7.5 asserts that $A_{\text{PL}}(M) \otimes \mathcal{L}(Q[r])$ is formal whenever M is compact and orientable. When r is odd, this fact may be surprising at first glance, since M is not assumed to be formal.

A conceptual understanding of this phenomenon is afforded by the homotopy transfer theorem; see [38, Section 10.3], for example. Indeed, let A be any nonunital differential graded commutative algebra and \mathfrak{g} any two-step nilpotent graded Lie algebra. Fixing an additive homotopy equivalence between A and $H(A)$, we obtain a transferred L_{∞} -algebra structure on $H(A) \otimes \mathfrak{g}$. The higher brackets of the transferred structure combine information about the Massey products of A and the Lie bracket of \mathfrak{g} .

In our case, using the fact that \mathfrak{g} has no nontrivial iterated brackets, the explicit formulas of the homotopy transfer theorem show that these higher brackets all vanish, which implies that $A \otimes \mathfrak{g}$ is formal. In other words, although the Massey products in $H(A)$ may be nontrivial, they are damped out by the nilpotence of \mathfrak{g} .

References

- [1] **VI Arnol'd**, *The cohomology ring of the colored braid group*, Math. Notes 5 (1969) 138–140 MR
- [2] **E Artin**, *Theory of braids*, Ann. of Math. 48 (1947) 101–126 MR
- [3] **D Ayala, J Francis**, *Poincaré/Koszul duality*, preprint (2014) arXiv
- [4] **D Ayala, J Francis**, *Zero-pointed manifolds*, preprint (2014) arXiv
- [5] **D Ayala, J Francis**, *Factorization homology of topological manifolds*, J. Topol. 8 (2015) 1045–1084 MR
- [6] **D Ayala, J Francis, H L Tanaka**, *Factorization homology of stratified spaces*, Selecta Math. 23 (2017) 293–362 MR
- [7] **D Ayala, J Francis, H L Tanaka**, *Local structures on stratified spaces*, Adv. Math. 307 (2017) 903–1028 MR
- [8] **C-F Bökigheimer**, *Stable splittings of mapping spaces*, from “Algebraic topology” (HR Miller, DC Ravenel, editors), Lecture Notes in Math. 1286, Springer (1987) 174–187 MR
- [9] **C-F Bökigheimer, F R Cohen**, *Rational cohomology of configuration spaces of surfaces*, from “Algebraic topology and transformation groups” (T tom Dieck, editor), Lecture Notes in Math. 1361, Springer (1988) 7–13 MR
- [10] **C-F Bökigheimer, F Cohen, L Taylor**, *On the homology of configuration spaces*, Topology 28 (1989) 111–123 MR
- [11] **A Borel**, *Sur l’homologie et la cohomologie des groupes de Lie compacts connexes*, Amer. J. Math. 76 (1954) 273–342 MR
- [12] **A K Bousfield, V K A M Gugenheim**, *On PL de Rham theory and rational homotopy type*, Mem. Amer. Math. Soc. 179, Amer. Math. Soc. (1976) MR
- [13] **F Cantero, M Palmer**, *On homological stability for configuration spaces on closed background manifolds*, preprint (2014) arXiv
- [14] **T Church**, *Homological stability for configuration spaces of manifolds*, Invent. Math. 188 (2012) 465–504 MR
- [15] **T Church, JS Ellenberg, B Farb**, *FI-modules and stability for representations of symmetric groups* (2012) arXiv
- [16] **F R Cohen**, *On configuration spaces, their homology, and Lie algebras*, J. Pure Appl. Algebra 100 (1995) 19–42 MR
- [17] **F R Cohen, T J Lada, J P May**, *The homology of iterated loop spaces*, Lecture Notes in Math. 533, Springer (1976) MR

- [18] **F R Cohen, L R Taylor**, *Computations of Gel'fand–Fuks cohomology, the cohomology of function spaces, and the cohomology of configuration spaces*, from “Geometric applications of homotopy theory, I” (HR Miller, DC Ravenel, editors), Lecture Notes in Math. 657, Springer (1978) 106–143 MR
- [19] **G Drummond-Cole, B Knudsen**, *Betti numbers of configuration spaces of surfaces*, J. London Math. Soc. (online publication July 2017)
- [20] **E R Fadell, S Y Husseini**, *Geometry and topology of configuration spaces*, Springer (2001) MR
- [21] **E Fadell, L Neuwirth**, *Configuration spaces*, Math. Scand. 10 (1962) 111–118 MR
- [22] **Y Félix, S Halperin, J-C Thomas**, *Rational homotopy theory*, Graduate Texts in Math. 205, Springer (2001) MR
- [23] **Y Félix, D Tanré**, *The cohomology algebra of unordered configuration spaces*, J. London Math. Soc. 72 (2005) 525–544 MR
- [24] **Y Félix, J-C Thomas**, *Rational Betti numbers of configuration spaces*, Topology Appl. 102 (2000) 139–149 MR
- [25] **J Francis**, *The tangent complex and Hochschild cohomology of \mathcal{E}_n -rings*, Compos. Math. 149 (2013) 430–480 MR
- [26] **M Franz**, *Koszul duality and equivariant cohomology*, Doc. Math. 11 (2006) 243–259 MR
- [27] **B Fresse**, *Koszul duality of operads and homology of partition posets*, from “Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K -theory” (P Goerss, S Priddy, editors), Contemp. Math. 346, Amer. Math. Soc. (2004) 115–215 MR
- [28] **E Getzler**, *The homology groups of some two-step nilpotent Lie algebras associated to symplectic vector spaces*, preprint (1999) arXiv
- [29] **E Getzler**, *Resolving mixed Hodge modules on configuration spaces*, Duke Math. J. 96 (1999) 175–203 MR
- [30] **M Goresky, R Kottwitz, R MacPherson**, *Equivariant cohomology, Koszul duality, and the localization theorem*, Invent. Math. 131 (1998) 25–83 MR
- [31] **O Gwilliam**, *Factorization algebras and free field theories*, PhD thesis, Northwestern University (2012) MR Available at <https://search.proquest.com/docview/1018399857>
- [32] **V Hinich**, *Homological algebra of homotopy algebras*, Comm. Algebra 25 (1997) 3291–3323 MR
- [33] **G Horel**, *Higher Hochschild cohomology of the Lubin–Tate ring spectrum*, Algebr. Geom. Topol. 15 (2015) 3215–3252 MR

- [34] **B Keller**, *Introduction to A -infinity algebras and modules*, Homology Homotopy Appl. 3 (2001) 1–35 MR
- [35] **B Knudsen**, *Higher enveloping algebras*, preprint (2016) arXiv
- [36] **B Knudsen**, *Higher enveloping algebras and configuration spaces of manifolds*, PhD thesis, Northwestern University (2016) MR Available at <https://search.proquest.com/docview/1799681505?pq-origsite=gscholar>
- [37] **A Kupers, J Miller**, *Homological stability for topological chiral homology of completions*, Adv. Math. 292 (2016) 755–827 MR
- [38] **J-L Loday, B Vallette**, *Algebraic operads*, Grundlehrer Math. Wissen. 346, Springer (2012) MR
- [39] **R Longoni, P Salvatore**, *Configuration spaces are not homotopy invariant*, Topology 44 (2005) 375–380 MR
- [40] **J Lurie**, *Higher topos theory*, Annals of Mathematics Studies 170, Princeton Univ. Press (2009) MR
- [41] **J Lurie**, *Derived algebraic geometry, X: Formal moduli problems*, preprint (2011) Available at <http://www.math.harvard.edu/~lurie/papers/DAG-X.pdf>
- [42] **J Lurie**, *Higher algebra* (2014) Available at <http://www.math.harvard.edu/~lurie/papers/HA.pdf>
- [43] **J P May**, *The geometry of iterated loop spaces*, Lectures Notes in Math. 271, Springer (1972) MR
- [44] **D McDuff**, *Configuration spaces of positive and negative particles*, Topology 14 (1975) 91–107 MR
- [45] **D Petersen**, *A spectral sequence for stratified spaces and configuration spaces of points*, Geom. Topol. 21 (2017) 2527–2555 MR
- [46] **O Randal-Williams**, *Homological stability for unordered configuration spaces*, Q. J. Math. 64 (2013) 303–326 MR
- [47] **E Riehl**, *Categorical homotopy theory*, New Mathematical Monographs 24, Cambridge Univ. Press (2014) MR
- [48] **P Salvatore**, *Configuration spaces with summable labels*, from “Cohomological methods in homotopy theory” (J Aguadé, C Broto, C Casacuberta, editors), Progr. Math. 196, Birkhäuser, Basel (2001) 375–395 MR
- [49] **P Salvatore, N Wahl**, *Framed discs operads and Batalin–Vilkovisky algebras*, Q. J. Math. 54 (2003) 213–231 MR
- [50] **G Segal**, *Classifying spaces and spectral sequences*, Inst. Hautes Études Sci. Publ. Math. 34 (1968) 105–112 MR
- [51] **G Segal**, *Configuration-spaces and iterated loop-spaces*, Invent. Math. 21 (1973) 213–221 MR

- [52] **DP Sinha**, *The (non-equivariant) homology of the little disks operad*, from “Operads 2009” (J-L Loday, B Vallette, editors), Sémin. Congr. 26, Soc. Math. France, Paris (2013) 253–279 MR
- [53] **B Totaro**, *Configuration spaces of algebraic varieties*, *Topology* 35 (1996) 1057–1067 MR
- [54] **N Wahl**, *Ribbon braids and related operads*, PhD thesis, Oxford (2001) Available at <http://eprints.maths.ox.ac.uk/43/1/wahl.pdf>
- [55] **JH Wang**, *On the braid groups for \mathbf{RP}^2* , *J. Pure Appl. Algebra* 166 (2002) 203–227 MR

*Department of Mathematics, Harvard University
Cambridge, MA, United States*

`knudsen@math.harvard.edu`

<http://scholar.harvard.edu/knudsen/>

Received: 8 December 2016 Revised: 9 December 2016

