

Taut branched surfaces from veering triangulations

MICHAEL LANDRY

Let M be a closed hyperbolic 3–manifold with a fibered face σ of the unit ball of the Thurston norm on $H_2(M)$. If M satisfies a certain condition related to Agol’s veering triangulations, we construct a taut branched surface in M spanning σ . This partially answers a 1986 question of Oertel, and extends an earlier partial answer due to Mosher.

57M99

1 Introduction

Let M be a closed, irreducible, atoroidal 3–manifold. In [16], Oertel shows that each closed face σ of the Thurston norm ball of M possesses a finite collection of taut oriented branched surfaces which together carry representatives of all integral classes in $\text{cone}(\sigma)$. We say that these branched surfaces *span* σ . One question he asked and left open was whether this collection could have size 1. That is, when is it possible to find a single taut oriented branched surface which spans σ ? Our main result is a partial answer to this question when M is hyperbolic and σ is fibered.

Main Theorem (Theorem 3.9) *If every boundary torus of \mathring{M} witnesses at most two ladderpole vertex classes of $H_2(M)$, then M has a taut homology branched surface spanning σ .*

The definition of \mathring{M} , and that of *ladderpole vertex class*, which involves Agol’s veering triangulation, can be found in Sections 2 and 3.1, respectively.

To quickly summarize the previous study of Oertel’s question: Sterba-Boatwright [19] shows by a counterexample that the answer is “not always”. However, Mosher [15] proves that in the case when M is hyperbolic and σ is fibered, there exists such a branched surface if the vertices of σ have positive intersection with the singular orbits of the Fried suspension flow φ corresponding to σ .

The main theorem, Theorem 3.9, strengthens Mosher’s result because ladderpole classes in particular have intersection 0 with some singular orbits of the Fried flow φ . While

the condition in the statement of the main theorem may seem mysterious, we prove in [Theorem 3.11](#) that it holds, in particular, when $H_2(M)$ has rank at most 3. This gives us the following corollary.

Corollary 3.12 *If $b_2(M) \leq 3$, then M has a taut homology branched surface spanning σ .*

The technique we employ to improve Mosher's result uses Agol's veering triangulation of a pseudo-Anosov mapping torus M' which is missing the singular orbits of Fried's flow φ . One nice property of this veering triangulation is that its 2-skeleton has the structure of a taut oriented branched surface which spans a fibered face σ' of M' .

We work in the cusped manifold M' and make use of the veering triangulation before moving back to M by Dehn filling. By understanding how the veering triangulation sits in M , we can construct a face-spanning taut oriented branched surface in M as long as our condition regarding ladderpole vertex classes is met.

Veering triangulations were introduced in Agol [1], where they were used to provide an alternative proof of the theorem in Farb, Leininger and Margalit [4], stating that the mapping tori of small-dilatation pseudo-Anosovs come from Dehn filling on finitely many cusped hyperbolic manifolds. Hodgson, Rubinstein, Segerman and Tillmann [12] show that veering triangulations admit positive angle structures, and Futer and Guéritaud [7] give a lower bound on the smallest angle in such a triangulation in terms of combinatorial data coming from the veering triangulation. In light of these results, one could hope that veering triangulations might be realized geometrically. However, in Hodgson, Issa and Segerman [11] there is an explicit example of a nongeometric veering triangulation. Guéritaud [10] proved that the veering triangulation controls the combinatorics of the Cannon–Thurston map associated to the fully punctured manifold M' . Most recently, Minsky and Taylor found connections between subsurface projections and the veering triangulation in [14].

Acknowledgements I would like to thank the anonymous referee for her or his helpful comments and suggestions. Thanks are also due to Siddhi Krishna, who identified an expositional error in an earlier version of the paper. Finally, I would like to thank my advisor Yair Minsky for many helpful conversations during this research.

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No DGE-1122492, and by the National Science Foundation under Grant No DMS-1610827 (PI Yair Minsky). Any

opinions, findings, and conclusions or recommendations expressed in this material are those of the author and do not necessarily reflect the views of the National Science Foundation.

2 Background

2.1 Notation

Throughout this document, certain notation will remain fixed. All homology groups are assumed to have coefficients in \mathbb{R} unless otherwise stated. We consider a closed pseudo-Anosov mapping torus M , and a closed face σ of the unit ball of the Thurston norm x on $H_2(M)$. There is a pseudo-Anosov flow φ on M which Fried showed in [5] is associated to σ in a natural way. We denote the union of the singular orbits of φ by c and define $\mathring{M} := M \setminus U$, where U is an open tubular neighborhood of c . The boundary $\partial\mathring{M}$ of \mathring{M} is a union of tori. There is a natural embedding $P: H_2(M) \rightarrow H_2(\mathring{M}, \partial\mathring{M})$, which comes from the exact sequence $0 = H_2(U) \rightarrow H_2(M) \rightarrow H_2(M, U) \rightarrow \cdots$ and the excision isomorphism $H_2(\mathring{M}, \partial\mathring{M}) \cong H_2(M, U)$. This is induced at the level of chains by sending a chain S to $S \setminus U$. We use the notation $P(\alpha) = \mathring{\alpha}$. More exposition of some of these ideas will be provided in the following subsections.

2.2 Thurston norm

We begin by reviewing the definition of, and some salient facts about, the Thurston norm. For a more detailed treatment, see Thurston's original paper [20] or Candel and Conlon's textbook [3].

Let Z be an orientable three-manifold. The Thurston norm is a seminorm on $H_2(Z, \partial Z)$ defined as follows. If S is a connected surface embedded in Z , define $\chi_-(S) := \max\{0, -\chi(S)\}$, where χ denotes Euler characteristic. If S has multiple connected components, then $\chi_-(S) := \sum_i \chi_-(S_i)$, where the sum is over the components S_i of S .

If $\alpha \in H_2(Z, \partial Z)$ is an integral class, we can always find an embedded surface representing α . We define

$$x(\alpha) = \min_{[S]=\alpha} \chi_-(S),$$

where the minimum is taken over embedded representatives of α , and call $x(\alpha)$ the *Thurston norm* of α .

The Thurston norm is homogeneous and satisfies the triangle inequality for integral classes. It can be extended to all rational classes in $H_2(Z, \partial Z)$ by homogeneity, whereafter it extends uniquely to a seminorm on all of $H_2(Z, \partial Z)$. If Z is incompressible, atoroidal, anannular and boundary-incompressible (ie has no essential surfaces of nonnegative Euler characteristic), then x is a bona fide norm on $H_2(Z, \partial Z)$, and not just a seminorm. In this paper we will deal only with pseudo-Anosov mapping tori, so we will assume x is a norm from this point forward.

From the fact that x takes integer values on the integral lattice, one can show that the unit sphere of x is a finite-sided convex polyhedron. Moreover, this polyhedron encodes information about how Z fibers over S^1 . If F is a fiber of some fiber bundle $F \rightarrow Z \rightarrow S^1$, then $[F]$ lies in the interior of the positive cone over a top-dimensional face. If α is any integral class in the interior of the same cone, then α is represented by a fiber of some fiber bundle over the circle. The closed top-dimensional face associated to this cone is called a *fibred face*.

In the setting of this paper, a closed hyperbolic manifold M with fibred face σ , David Fried proved in [5] that we can associate to σ a flow φ with very nice properties. Namely, any primitive integral class $\alpha \in \text{int cone}(\sigma)$, where int denotes interior, is represented by a cross-section S to φ , and the first return map of S is pseudo-Anosov. Hence we can think of φ as the simultaneous suspension flow of all monodromies of fibers corresponding to σ , and the singular orbits of φ as the suspensions of the singular points of those monodromies.

2.3 Taut branched surfaces

A *branched surface* B is a smooth codimension-1 object in a 3-manifold, analogous to a train track in a surface, which organizes the data of various embedded surfaces. Locally, a branched surface looks like a stack of disks D_1, \dots, D_n such that D_i is glued to D_{i+1} for $i < n$ along the closure of one component of the complement of a smooth arc through D_i . The union of the images of smooth curves is called the *branching locus*. The smooth structure of B is such that the inclusion of each D_i is smooth [17]. The *sectors* of B are the connected components of the complement of the branching locus; these are analogous to the edges of a train track.

A regular neighborhood $N(B)$ of a branched surface B can be foliated by line segments transverse to B . This foliation is called the *vertical foliation* of $N(B)$. If it is possible to consistently orient the leaves of the vertical foliation, then B is called an *oriented branched surface*.

We say that B carries S if S is embedded in $N(B)$ transverse to the vertical foliation; this is analogous to a train track carrying a curve. In the same way that a train track inherits nonnegative integer edge weights from a carried curve, a surface S carried by B assigns a weight to each sector of B . If these weights are all positive, we say B fully carries S .

Conversely, a collection of nonnegative integral weights on sectors of B which satisfy the linear equations determined by the branching determines an isotopy class of surface carried by B . In fact, any real weights satisfying the branching equations naturally determine a homology class; we say that the homology classes corresponding to nonnegative weights are carried by B .

If we allow negative weights, there is a natural vector space whose elements are collections of real weights satisfying the branching equations. In this vector space, the integral points in the cone of nonnegative weights correspond to surfaces carried by B .

Branched surfaces are interesting in part because of the surfaces they carry, so it is natural to distinguish types of branched surfaces which carry surfaces with nice properties. If a branched surface carries only surfaces realizing the minimal χ_- in their homology class, we say it is *almost taut*. Following Oertel, we say B is *taut* if it carries only *incompressible* surfaces which attain the minimal χ_- in their homology class. We say B is a *homology branched surface* if B is oriented and for each point $p \in B$, there exists a closed oriented transversal through p , ie an oriented loop through p whose intersection with $N(B)$ consists of leaves of the vertical foliation with the correct orientation. Since any surface carried by a homology branched surface has nonzero algebraic intersection number with a closed curve, homology branched surfaces carry only homologically nontrivial surfaces.

The following lemma, which we will use later, is probably known to many. We record a proof here for convenience.

Lemma 2.1 *Let Z be a compact 3-manifold, and $B \subset Z$ a homology branched surface. Then the cone of classes carried by B is closed.*

Proof Consider the natural linear map L from the vector space V of weights satisfying the branching equations on B to $H_2(Z, \partial Z)$. Let $A \subset V$ be the closed cone of nonnegative weights on B ; our goal is to show that $L(A)$ is closed.

Choose a norm $\|\cdot\|$ on V , and let $A^1 = \{v \in A \mid \|v\| = 1\}$. Then A^1 is compact, and hence $L(A^1)$ is compact. Moreover, $0 \notin L(A^1)$ because B is a homology branched surface.

Note that $L(A) = \text{cone}(L(A^1))$. Since the cone over a compact set not containing 0 is closed, we are done. \square

Note that an almost taut branched surface is not necessarily taut. For example, if B carries the boundary of a solid torus, B is not taut. However, since this torus realizes the Thurston norm of the homology class 0, carrying the torus does not preclude B from being almost taut.

If an almost taut homology branched surface B lies in a pseudo-Anosov mapping torus, it is taut. In general, we have the following.

Lemma 2.2 *Let N be a manifold such that the Thurston norm x is a norm (ie not just a seminorm) on $H_2(N, \partial N)$, and let $B \subset N$ be an almost taut branched surface. If B is also a homology branched surface, then B is taut.*

Proof Suppose S is a compressible surface carried by B , and let S' be the surface obtained by compressing S along a compression disk. Since B is almost taut, $\chi_-(S) = \chi_-(S')$, and since $\chi(S') = \chi(S) + 2$, S must be a torus, annulus, disk or sphere. Since B is a homology branched surface, S is homologically nontrivial, a contradiction. \square

In the course of proving Theorem 4 in [16], Oertel proves the following useful criterion for almost tautness. Since he doesn't state the result explicitly, we record a proof here for the reader's convenience.

Lemma 2.3 [16] *Let N be as above. Suppose $B \subset M$ is an oriented branched surface which fully carries a minimal- χ_- representative Σ of a single class in $H_2(N, \partial N)$. If B does not carry any spheres or disks, then B is almost taut.*

Proof Suppose for a contradiction that B carries a surface S which is homologous to S' with $\chi_-(S) > \chi_-(S')$. Since Σ is fully carried, there exists $n \in \mathbb{N}$ such that $n\Sigma$ (ie n parallel copies of Σ) is carried with greater weights than S on each sector of B . We have

$$n\Sigma = (n\Sigma - S) + S,$$

where $n\Sigma - S$ denotes the surface arising from subtracting the weights corresponding to S from the weights corresponding to $n\Sigma$, and $+$ denotes oriented sum in a regular neighborhood of B . Let $F := n\Sigma - S$. Now

$$\begin{aligned}\chi_-(n\Sigma) &= \chi_-(F) + \chi_-(S) \quad (\text{because } B \text{ does not carry disks or spheres}) \\ &> \chi_-(F) + \chi_-(S') \\ &\geq x([F]) + x([S']) \\ &\geq x([n\Sigma]).\end{aligned}$$

The last inequality is the triangle inequality for x . This is a contradiction since $n\Sigma$ realizes the minimal χ_- in $[n\Sigma]$. \square

2.4 Branched surfaces and faces of the Thurston norm ball

If two homologically nontrivial surfaces S and T are carried by a taut oriented branched surface B , we can perform an oriented cut-and-paste along the branching locus of B to form a surface $S + T$ representing $[S] + [T]$ and carried by B . Since B is taut, $S + T$ is norm-minimizing. Also, $\chi_-(S + T) = \chi_-(S) + \chi_-(T)$, as none of S , T or $S + T$ has any sphere or disk components. This is because they are all carried by a taut branched surface. Thus,

$$x([S] + [T]) = x([S + T]) = \chi_-(S + T) = \chi_-(S) + \chi_-(T) = x([S]) + x([T]).$$

The faces of the Thurston norm unit ball are projectivizations of the maximal cones on which x is linear, so we conclude that $[S]$ and $[T]$ lie in $\text{cone}(\sigma)$ for some face σ of the Thurston norm ball. Oertel observed this and asked the following question.

Question 2.4 [16] Let M be a simple (compact, irreducible, atoroidal) 3-manifold. For each face of the unit ball of the Thurston norm on $H_2(M, \partial M)$, is it possible to find a taut oriented branched surface which carries a norm-minimizing representative of every projective homology class in $\text{cone}(\sigma)$?

In [19], a closed 3-manifold is constructed for which the answer to [Question 2.4](#) is no. More specifically, Sterba-Boatwright produces a face of this manifold's Thurston norm ball with the following property: any branched surface carrying norm-minimizing representatives of all classes in that face also carries a compressible torus, so it cannot be taut. Hence the answer to Oertel's question is not an unqualified yes. However, in [15], [Question 2.4](#) is answered in the affirmative for a fibered face of a closed

pseudo-Anosov mapping torus in the case that each integral class in the cone over the face has positive intersection number with the singular orbits of the suspension flow for that face.

In this paper we extend Mosher's result by using the relatively new technology of veering triangulations, which we describe now.

2.5 Veering triangulations

A *taut ideal tetrahedron* is an ideal tetrahedron with the following extra data: each edge is labeled 0 or π so that the sum of the labels of edges incident to any ideal vertex is π , each face is cooriented so that two faces point out and two faces point in, and face coorientations change only along edges labeled 0. Note that the word "taut" is used here in a different sense than when it modifies "branched surface". A *taut ideal triangulation* of a 3-manifold is an ideal triangulation by taut tetrahedra such that face coorientations agree, and for each edge e , the sum of e 's labels over all incident tetrahedra is 2π . The 2-skeleton of such a triangulation Δ can be pinched along each edge to give a branched surface B_Δ in the manifold, as seen in [Figure 3](#). In this way we think of the edge labels of a taut tetrahedron as angles. Taut ideal triangulations were introduced in [\[13\]](#).

The condition that makes a triangulation veering is simple to draw but more complicated to define.

Consider a taut ideal triangulation of an oriented 3-manifold L . Up to combinatorial equivalence there is only one taut ideal tetrahedron. However, if we distinguish a 0-edge, there are 2 types embedded in L up to oriented equivalence, as you can see in [Figure 1](#). The taut ideal triangulation is *veering* if, for each edge e ,

- (i) if we circle around e and read off edge angles, no two angles of π are circularly adjacent, and
- (ii) each tetrahedron for which e is a 0-edge is of the same type when e is distinguished.

This situation is shown in [Figure 2](#). If L is not orientable, a taut ideal triangulation of L is veering if its lift to the orientation cover of L is veering.

There is an alternative definition of veering, due to Guéritaud [\[10\]](#), which says that a taut triangulation is veering if its 1-skeleton possesses a certain type of two-coloring. That definition is equivalent to the one given above, which we use throughout the paper.




Figure 1: The two types of taut tetrahedron with distinguished (bold) 0-edge. In this drawing the coorientation is pointing out of the paper, the vertical and horizontal edges are 0-edges, and the diagonal edges are π -edges.

Recall that c denotes the union of singular orbits of the Fried suspension flow in our pseudo-Anosov mapping torus M . In [1], Ian Agol shows how to construct a canonical veering triangulation of $M \setminus c$. He builds this triangulation using a sequence of ideal triangulations of the punctured surface which are dual to a periodic train track splitting sequence. The triangulations are related by Whitehead moves which determine the incidences of taut tetrahedra. Taut ideal triangulations obtained from Whitehead moves in this way, a construction due to Lackenby [13], are called *layered triangulations*.

Agol has proven the following theorem, which shows that his veering triangulation of $M \setminus c$ is canonically associated to a fibered face of M . Guéritaud provided an alternative proof, which is exposted in [14], based on his alternative construction of the canonical veering triangulation in [10].




Figure 2: The bold edge satisfies the veering condition.




Figure 3

Theorem 2.5 (Agol) *The ideal layered veering triangulation τ of $M \setminus c$ coming from a fibration associated to $\text{int cone}(\sigma)$ is constant over $\text{int cone}(\sigma)$. The 2-skeleton of this triangulation is a branched surface B_τ such that if S is a fiber with $[S] \in \text{int cone}(\sigma)$, some multiple of $S \setminus c$ is fully carried by B_τ .*

We remark that B_τ is taut. Indeed, B_τ is transverse to φ , the flow associated to σ , and a generic orbit of φ is dense in M ; this allows us to find a closed transversal through every point of B_τ , so B_τ is a homology branched surface. Since $M \setminus c$ is irreducible and boundary-irreducible, this implies B_τ carries no disks or spheres. Indeed, any carried disk or sphere would be homologically nontrivial because B_τ is a homology branched surface, but this is ruled out by the irreducibility and boundary-irreducibility of $M \setminus c$.

The fact that B_τ fully carries a fiber (in fact, many fibers) of $M \setminus c$ means that B_τ is almost taut by [Lemma 2.3](#) since fibers are norm-minimizing; see [20]. By [Lemma 2.2](#), B_τ is taut.

Since we want to use the veering triangulation τ to extract information about the Dehn filling M of $M \setminus c$, it will be useful to consider the restriction of τ to $\mathring{M} = M \setminus U$ (recall that U is a small tubular neighborhood of the collection c of singular orbits of φ). That way we will have room to work in the solid tori of U . Homologically this changes nothing, and we will use the notation $\mathring{\tau}$ and $B_{\mathring{\tau}}$ to denote $\tau \setminus U$ and $B_\tau \setminus U$, respectively. A taut ideal triangulation of \mathring{M} will mean a taut ideal triangulation of $M \setminus c$ restricted to \mathring{M} .



Figure 4: On the left and right, respectively, are upward and downward flat triangles. The coorientation of each edge points upward.

2.6 On the boundary of \mathring{M}

Let Δ be a taut ideal triangulation of \mathring{M} . Then we may assume its 2–skeleton B_Δ intersects $\partial\mathring{M}$ transversely in a cooriented train track as in Figure 3. This train track divides each component of $\partial\mathring{M}$ into bigons, which we will think of as triangles with one vertex whose interior angle is π (corresponding to a smooth path through a switch) and two cuspidal vertices whose angles are 0.

A triangle as above, with two vertices of interior angle 0 and one with angle π , is called *flat*. The edge between the two 0–vertices is called a 0–0 edge and the other two are called 0– π edges. We say the triangle is *upward* or *downward* if the train track’s coorientation points out of or into a flat triangle at its π –vertex, respectively. See Figure 4.

We draw the train track with the convention that we view $\partial\mathring{M}$ from inside \mathring{M} and the coorientation points upward. Having made this convention, we define the *triangles to the left* and *to the right* of a switch p to be the triangles with interior angle 0 at p which lie to the left and right of p .

Lemma 2.6 (veering condition on boundary) *Let Δ be a taut ideal triangulation of \mathring{M} . Then Δ is veering if and only if the train track on $\partial\mathring{M}$ has the property that, for every switch p , the triangles to the left of p are all upward or all downward, and the triangles to the right of p are all the opposite.*




Figure 5: The edge of the taut ideal triangulation corresponding to the central switch satisfies the veering condition.

Proof Each switch p in the train track corresponds to an edge e in the triangulation, and it is clear that, with respect to a single component of $\partial\overset{\circ}{M}$, the intersections with tetrahedra to the right of the switch are flat triangles of the same type if and only if the corresponding tetrahedra are of the same type when e is distinguished. The triangles to the left of p will be of the opposite type if and only if their corresponding tetrahedra are of the same type as those to the right of p . \square

An example of a switch satisfying the condition of [Lemma 2.6](#) is shown in [Figure 5](#). In [\[7\]](#), Futer and Guéritaud observed the following structure in the intersection of a veering triangulation with $\partial\overset{\circ}{M}$.

Lemma 2.7 Fix a component T_i of $\partial\overset{\circ}{M}$. Let $u, d \subset T_i$ be the closed regions consisting of upward and downward triangles, respectively. Then u and d are collections of essential annuli. If t is a triangle, then all three vertices of t lie in $\partial u = \partial d$.

For example, if t is upward then each edge of t either lies entirely in $\partial u = \partial d$ or traverses u with its endpoints on the boundary.

Proof The reader can check that a violation of the lemma contradicts [Lemma 2.6](#). \square




Figure 6: An example of a train track coming from a veering triangulation, endowed with its inherited orientation, lifted to the plane. This particular example comes from the veering triangulation associated to a minimal dilation 4-strand braid, discussed in [\[1\]](#). This train track is associated to the cusp coming from the suspension of the monodromy's lone singular point.

We will call the annulus connected components of u and d *upward* and *downward bands*, respectively. Part of the content of [Lemma 2.7](#) is that each band is only one edge across. Guéritaud calls the edges forming the boundaries of bands *ladderpole edges*, and the edges which traverse the bands *rungs*. For an example of one of these triangulations lifted to \mathbb{R}^2 , see [Figure 6](#).

3 Moving forward with Oertel's question

3.1 More on the structure of $\partial \overset{\circ}{\mathcal{T}}$

Define $\gamma := \partial \overset{\circ}{\mathcal{T}}$, ie the intersection of the veering triangulation with $\partial \overset{\circ}{\mathcal{M}}$. Fix a component T_i of $\partial \overset{\circ}{\mathcal{M}}$, and set $\gamma_i = \gamma \cap T_i$. We say the slope s_i corresponding to the union of all ladderpole edges in γ_i is the *ladderpole slope* for T_i . It will be convenient to think of γ_i as oriented in addition to being cooriented, meaning that each edge of γ_i has a preferred direction, and the preferred directions are compatible at switches. Our choice of orientation is the one induced by the boundary of any surface carried by $B_{\mathcal{Q}}$. In particular, the orientation on all rungs of γ_i is from right to left (as indicated in [Figure 6](#)). We will say γ_i *positively carries* an oriented curve a if the orientation of a agrees with the orientations of the edges in γ_i . Define an *upward* (resp. *downward*) *ladderpole* to be the right (resp. left) boundary component of an upward band in T_i , endowed with the orientation it inherits from γ_i . In our pictures, the upward ladderpoles are oriented upwards.

Lemma 3.1 *Let s_i and T_i be as above. Then*

- (1) γ_i *positively carries* curves s_i^+ and s_i^- with slope s_i such that $[s_i^+] + [s_i^-] = 0$ in $H_1(T_i)$, and s_i is the unique such slope on T_i ;
- (2) *any curve carried by γ_i with slope s_i traverses only ladderpole edges; and*
- (3) γ_i *positively carries a representative of every integral class in a halfspace of $H_1(T_i)$ bounded by $\mathbb{R} \cdot [s_i^+]$.*

Proof The union of all edges in a positive ladderpole forms an oriented curve s_i^+ which is positively carried by γ_i , and similarly a downward ladderpole gives an oriented positively carried curve s_i^- . It is clear that $[s_i^+]$ and $[s_i^-]$ sum to 0 in homology.

Let α be a curve positively carried by γ_i which traverses a rung of γ_i , and recall that all rungs of γ_i are oriented from right to left. As we trace along α , we must traverse a rung

of every band in T_i from right to left, as otherwise α could not close up. Therefore, α intersects s_i^+ , and after a perturbation we can assume that the intersections are all positive, so $i([s_i^+], [\alpha]) > 0$. Hence, any curve carried positively or negatively by γ_i which traverses a rung has nonzero intersection number with s_i^+ and cannot have slope s_i . This proves claim (2).

If β is a curve carried positively by γ_i with slope $\neq s_i$, then it traverses rungs of γ_i and has positive intersection with s_i^+ as above. This means γ_i cannot positively carry a representative of $-\beta$, completing the proof of claim (1).

Convex combinations of classes with positively carried representatives can be represented by oriented cut-and-paste sums carried by γ_i . Also, if γ_i positively carries a curve representing ka for $a \in H_1(T_i)$ and $k \in \mathbb{Z}_{>1}$, then γ_i positively carries a curve representing a . Indeed, if ρ is such a curve, we can perform oriented surgeries to eliminate self-intersections that do not change the homology class of ρ , yielding k parallel curves carried by γ_i which represent a . In other words, to show that a homology class is represented by a positively carried curve, it is enough to show some multiple of the homology class is represented by a positively carried curve.

Therefore to prove claim (3), it suffices by (1) and (2) to show that γ_i positively carries a curve traversing a rung of γ_i . It is easy to see that there is such a curve: you can draw one by starting at a switch and tracing along rungs of γ_i until you reach the ladderpole containing your path's initial point. Then the path can be closed up along the ladderpole. □

Lemma 3.1 concerned all curves carried by γ . Now we consider the boundaries of surfaces carried by $B_{\tilde{\tau}}$. If an oriented surface is carried by $B_{\tilde{\tau}}$, its boundary traces out an oriented collection of curves on $\partial \mathring{M}$ that is positively carried by γ . Our understanding of γ allows us to deduce information about the surfaces carried by $B_{\tilde{\tau}}$, and thus surfaces representing classes in σ .

For example, we observe that every surface carried by $B_{\tilde{\tau}}$ has a nonladderpole boundary component.

Lemma 3.2 *There is no surface carried by $B_{\tilde{\tau}}$ whose boundary components are all ladderpoles.*

Proof Suppose S is a surface carried by $B_{\tilde{\tau}}$. An edge of γ traversed by ∂S corresponds to a truncated ideal triangle of $\overset{\circ}{\tau}$ on which S has positive weight. Let t be

such a triangle. Choose a truncated ideal tetrahedron Y of which t is a face. The truncated vertices of t correspond to 3 edges of the truncated vertices of Y , which are flat triangles lying in $\partial\overset{\circ}{M}$. Exactly one of these edges, which we will call e , is a $0-0$ edge with respect to a truncated vertex of Y .

Each edge in γ is incident to two flat triangles. Our knowledge of γ gives us that a ladderpole edge is $0-\pi$ with respect to both of its incident triangles, while a rung is $0-0$ with respect to one incident triangle and $0-\pi$ with respect to the other. It follows that e is a rung, so ∂S cannot consist of all ladderpole edges. \square

Recall from Section 2.1 the injective puncturing map $P: H_2(M) \rightarrow H_2(\overset{\circ}{M}, \partial\overset{\circ}{M})$, induced at the level of chains by sending S to $S \setminus U$. As a reminder, we will write $P(\alpha) =: \overset{\circ}{\alpha}$ when convenient.

We can classify the possible boundaries of all surfaces carried by $B_{\frac{\pi}{2}}$ coming from M , ie those representing classes that lie in the image of P .

Lemma 3.3 *Let $\alpha \in \text{cone}(\sigma)$ be an integral class, and consider a surface $Y \subset \overset{\circ}{M}$ carried by $B_{\frac{\pi}{2}}$ and representing $\overset{\circ}{\alpha}$. Then, for each component T_i of $\partial\overset{\circ}{M}$, $\partial Y \cap T_i$ is positively carried by γ_i , and is either*

- (1) empty,
- (2) a collection of ladderpoles which is nullhomologous in T_i , or
- (3) a collection of meridians.

Further, if $\alpha \in \text{int cone}(\sigma)$, then $\partial Y \cap T_i$ is a nonempty collection of meridians.

Proof Because the coorientation of Y agrees with that of $B_{\frac{\pi}{2}}$, the orientation of $\partial Y \cap T_i$ agrees with that of γ_i . Hence $\partial Y \cap T_i$ is positively carried.

Let Q be the map $H_2(M) \rightarrow H_2(\text{cl}(U), \partial\overset{\circ}{M})$ defined similarly to the puncturing map P by excising $M \setminus \text{cl}(U)$. Then we have the commutative diagram

$$(3.4) \quad \begin{array}{ccc} H_2(M) & \xleftarrow{P} & H_2(\overset{\circ}{M}, \partial\overset{\circ}{M}) \\ \downarrow Q & & \downarrow \partial \\ H_2(\text{cl}(U), \partial\overset{\circ}{M}) & \xrightarrow{-\partial} & H_1(\partial\overset{\circ}{M}) \end{array}$$

(A quick remark: the minus sign on the bottom boundary map of (3.4) reflects the fact that $\partial: H_2(\text{cl}(U), \partial\overset{\circ}{M}) \rightarrow H_1(\partial\overset{\circ}{M})$ is induced by the boundary map on 2-chains inside U , while the vertical boundary map is induced by the map on 2-chains outside U .)

Fix a component U_i of U with boundary T_i and meridional disk D_i and set $m_i = -\partial D_i$. Since the images of $\partial \circ P$ and $-\partial \circ Q$ are equal, and $H_2(\text{cl}(U_i), T_i) = \langle [D_i] \rangle$, the boundary of Y on T_i is homologous to km_i , where $k \in \mathbb{Z}$. Moreover, ∂Y is embedded and carried by γ . In particular this means the curves of $\partial Y \cap T_i$ are parallel.

If $\partial Y \cap T_i$ is nonempty and nullhomologous in T_i , then it must consist of an even number of ladderpoles, by Lemma 3.1.

Otherwise $k \neq 0$, in which case $\partial Y \cap T_i$ must be a collection of k meridians. If $\alpha \in \text{int cone}(\sigma)$, then by Fried’s results in [6], α has positive intersection with each singular orbit of φ . Thus $k > 0$ in the above analysis, completing the proof. \square

A vertex class of σ is a primitive integral homology class $v \in H_2(M)$ projecting to a vertex of σ .

Let T_i be a torus boundary component of \mathring{M} . An embedded surface $Y \subset \mathring{M}$ carried by B_{φ} is ladderpole at T_i if $Y \cap T_i$ is a collection of ladderpoles. The homology class $\alpha \in H_2(M)$ is ladderpole at T_i if $\mathring{\alpha}$ has an embedded representative carried by B_{φ} which is ladderpole at T_i .

Note that by Lemma 3.3, ladderpole classes must lie in $\partial \text{cone}(\sigma)$.

3.2 Our approach to Question 2.4

In [15], Mosher gives an example of how to construct a branched surface spanning any fibered face F of a 3-manifold N . He simply takes embedded minimal- χ -representatives S_i of each vertex class v_i for the face and perturbs them to intersect transversely. Then there is an isotopy in a neighborhood of the intersection locus, shown in Figure 7, which gives the union of the surfaces the structure of a branched surface B_F carrying each vertex class. We will refer to this operation on transverse cooriented surfaces as *branched sum*. Since the vertex classes span F , the surface B_F spans F . The problem with this method, as Mosher notes, is that there is no guarantee that all surfaces carried by the B_F are incompressible.

However, B_F is frequently almost taut. Indeed, consider the surface $\sum_i S_i$, ie the oriented cut-and-paste sum of the S_i . We have

$$\chi_-\left(\sum_i S_i\right) = \sum_i \chi_-(S_i) = \sum_i x(v_i) = x\left(\sum_i v_i\right),$$




Figure 7: The branched sum operation. A generic intersection, shown on the left, can be smoothed in a unique way, shown on the right, such that it preserves coorientations. Here the coorientations are such that they all point into the octant facing the reader.

where the last equality follows from the fact that x is linear on $\text{cone}(F)$. Therefore $\sum_i S_i$ realizes the minimal χ_- in the homology class $\sum_i v_i$. Moreover, $\sum_i S_i$ is fully carried by B_F , so almost-tautness follows from Lemma 2.3 as long as B_F carries no disks or spheres. Slightly more generally, we have proved the following lemma.

Lemma 3.5 *Let F be a closed face of the Thurston norm ball in $H_2(N, \partial N)$. If $\{S_i\}$ is a finite collection of norm-minimizing surfaces embedded in M such that $\{[S_i]\} \subset \text{cone}(F)$, then the branched sum of the S_i is almost taut provided it carries no disks or spheres.*

If B_F is indeed almost taut and carries no disks or spheres, it follows that the only way B_F can fail to be an answer to Oertel’s Question 2.4 is if it carries a compressible torus.

The method we use to address Oertel’s Question 2.4 is similar to Mosher’s example above. The difference is that rather than take any embedded representatives of vertex classes, we take representatives of their punctured images in $H_2(\overset{\circ}{M}, \overset{\circ}{\partial M})$ lying in a regular neighborhood of τ . This is possible because, by Theorem 2.5, B_τ carries a representative of $\overset{\circ}{\alpha}$ for every integral class in $\text{int cone}(\sigma)$. By Lemma 2.1, the same can be said for representatives of classes $\overset{\circ}{\beta}$ for $\beta \in \partial \text{cone}(\sigma)$. With an extra hypothesis, we show how to extend the surfaces over the Dehn filling so that their branched sum B_σ carries no tori. Rather than show directly that no tori are carried, our method of proof is to demonstrate that B_σ is a homology branched surface, which is enough to imply tautness by Lemmas 2.3 and 2.2.

3.3 A lemma concerning x and Dehn filling

We are particularly interested in the restriction of the puncturing map $P: H_2(M) \rightarrow H_2(\overset{\circ}{M}, \partial\overset{\circ}{M})$ to $\text{cone}(\sigma)$, and it will be useful to have the following lemma concerning the relationship of P to the norm x . The subject of how the Thurston norm behaves under Dehn filling has been studied in [8; 9; 18], and more recently in [2].

Recall that c denotes the union of the singular orbits of φ , which is the flow associated to σ .

Lemma 3.6 *Let $\alpha \in \text{cone}(\sigma)$ be an integral homology class. Then*

$$x(\alpha) = x(\overset{\circ}{\alpha}) - i(\alpha, c),$$

where $i(\cdot, \cdot)$ is the algebraic intersection number.

Before proving Lemma 3.6 we state some results from [15] that require some definitions.

We say a flow φ' is a *dynamic blowup* of φ if it is obtained by the following procedure: We replace a singular orbit θ by the suspension of a homeomorphism f of a finite tree T . This homeomorphism's first return map on each edge of T should fix the endpoints and act without fixed points on the interior. Thus each edge of T can be given an orientation according to the direction points are moved by its first return map, and around each vertex these orientations should alternate between outward and inward. The suspended tree forms a complex of annuli K which is invariant under φ' . The new flow φ' is semiconjugate to φ by a map which collapses K to θ , and is one-to-one in the complement of K . The vector fields generating φ and φ' differ only inside a small neighborhood of θ .

We say that a surface S embedded in M is *almost transverse* to φ if there is a dynamic blowup $\varphi^\#$ of φ such that S is transverse to $\varphi^\#$, and the sum of tangent spaces $TS \oplus T\varphi^\#$ is positively oriented in M . Here TS and $T\varphi^\#$ denote the tangent spaces to the surface S and flow $\varphi^\#$, respectively.

Lemma 3.7 [15] *Let $\alpha \in H_2(M)$ be an integral class. Then α can be represented by a surface almost transverse to φ if and only if $\alpha \in \text{cone } \sigma$ (recall that φ depends on σ). More specifically, there exists a way to dynamically blow up φ to a flow $\varphi^\#$ along only singular orbits θ of φ with $i(\alpha, \theta) = 0$ such that α is represented by a surface S transverse to $\varphi^\#$.*

Singular orbits θ with $i(\alpha, \theta) = 0$ are called α -null and, if $i(\alpha, \theta) > 0$, we will say that θ is α -positive. It will also be useful to know:

Lemma 3.8 [15] *Let S be a surface almost transverse to a pseudo-Anosov flow on M . Then S is norm-minimizing.*

Proof of Lemma 3.6 First we will show that $x(\alpha) \leq x(\mathring{\alpha}) - i(\alpha, c)$.

Let Y be a representative of $\mathring{\alpha}$ carried by $B_{\mathring{c}}$, and let $T_i = \partial U_i$ be one of the torus boundary components of \mathring{M} . By Lemma 3.3, $\partial Y \cap T_i$ is either a collection of meridians or a collection of ladderpoles that represents 0 in $H_2(T_i)$. Such a nullhomologous collection can be realized as the boundary of a family of closed annuli whose interiors are embedded in U_i .

Observe that $x(\alpha) \leq x(\mathring{\alpha}) - n(Y, \partial \mathring{M})$, where n is the number of meridians of $\partial \mathring{M}$ in ∂Y . Indeed, we can glue a disk to each meridian boundary component, and cap off all other boundary components with annuli. Because $B_{\mathring{c}}$ is taut and thus Y is norm-minimizing, we obtain a representative S of α with $\chi_-(S) = x(\mathring{\alpha}) - n(Y, \partial \mathring{M})$.

Now we claim that $n(Y, \partial \mathring{M}) = i(\alpha, c)$.

First, note that by the proof of Lemma 3.3, the images of the boundary maps

$$\partial|_{\text{image}(P)}: H_2(\mathring{M}, \partial \mathring{M}) \rightarrow H_1(\partial \mathring{M}) \quad \text{and} \quad \partial: H_2(\text{cl}(U), \partial \mathring{M}) \rightarrow H_1(\partial \mathring{M})$$

are contained in the subgroup $\bigoplus_i \langle m_i \rangle$ of $H_1(\partial \mathring{M})$. Here m_i is the homology class of a meridional curve on T_i , oriented so that it is positively carried by γ_i . Let $\pi_i: \langle m_i \rangle \rightarrow \mathbb{Z}$ be the map $km_i \mapsto k$. Now, Y is carried by $B_{\mathring{c}}$, so $n(Y, \partial \mathring{M}) = \sum_i \pi_i(\partial(\mathring{\alpha}))$, and it thus makes sense to write $n(\mathring{\alpha}, \partial \mathring{M})$.

We can package this information into an updated version of the diagram (3.4):

$$\begin{array}{ccc} H_2(M) & \xrightarrow{P} & H_2(\mathring{M}, \partial \mathring{M}) \\ \downarrow Q & & \downarrow \partial \\ H_2(\text{cl}(U), \partial \mathring{M}) & \xrightarrow{-\partial} & \bigoplus_i \langle m_i \rangle \\ & \swarrow i(-, c) & \nwarrow \sum \pi_i \\ & \mathbb{Z} & \end{array}$$

where the dotted arrows are defined on the integral lattice of their domains. Since $i(\alpha, c) = i(Q(\alpha), c)$, the claim that $n(\mathring{\alpha}, \partial \mathring{M}) = i(\alpha, c)$ reduces to the claim that the

above diagram is commutative, which is true since the square and triangle commute. Therefore $x(\alpha) \leq x(\hat{\alpha}) - i(\alpha, c)$.

Next we will show that $x(\hat{\alpha}) \leq x(\alpha) + i(\alpha, c)$. The idea of this direction is to take a norm-minimizing surface transverse to c representing α and delete its intersections with U . This will give a representative of $\hat{\alpha}$ with $\chi_- = x(\alpha) + i(\alpha, c)$ as long as the orientations of all intersections with c agree. Hence it suffices to show that there exists such a norm-minimizing surface representing α , and this is where we use Lemmas 3.7 and 3.8.

Let S be a representative of α which is almost transverse to φ with blown-up flow $\varphi^\#$, let θ be an α -null orbit of φ , let K_θ be the $\varphi^\#$ -invariant annulus complex blowing up θ , and let $U_\theta^\#$ be a solid torus containing K_θ . Since S is transverse to $\varphi^\#$ and $TS \oplus T\varphi^\#$ is positively oriented, we have in particular that each intersection point of S with any α -positive singular orbit of $\varphi^\#$ is positively oriented. This is close to the property we want, but we need it to hold for φ and not just $\varphi^\#$. Thus we will show that S intersects the solid torus $U_\theta^\#$ in a collection of disjoint annuli, and can be isotoped outside $U_\theta^\#$. Then the positivity of S with respect to the α -positive singular orbits of φ will be preserved by the semiconjugacy collapsing K_θ to θ . Arguing in this way for each α -null orbit, we will obtain a norm-minimizing surface representing α that has only positively oriented intersection points with c .

Observe that $[S]$ maps to 0 under the map $H_2(M) \rightarrow H_2(\text{cl}(U_\theta^\#), \partial U_\theta^\#)$ because θ is α -null. After applying the boundary map $H_2(\text{cl}(U_\theta^\#), \partial U_\theta^\#) \rightarrow H_1(\partial U_\theta^\#)$ we see that $S \cap \partial U_\theta^\#$ represents 0 in $H_1(\partial U_\theta^\#)$. It follows that $S \cap \partial U_\theta^\#$, after perturbation for transversality, is a possibly empty nullhomologous embedded collection of curves. If it is empty, there is nothing for us to prove.

If $S \cap \partial U_\theta^\#$ contains a curve a which is inessential in $\partial U_\theta^\#$, then a bounds a disk in $\partial U_\theta^\#$. If a is essential in S , this contradicts the incompressibility of S . If a is inessential in S , then we can perform an isotopy of S which removes a from $\partial U_\theta^\#$. Therefore we can assume $S \cap \partial U_\theta^\#$ is a collection of curves essential in $\partial U_\theta^\#$. Since the collection is embedded and nullhomologous, there must be $2n$ curves of the same slope, half having one orientation and half the other.

If this slope is the meridional slope for $U_\theta^\#$, the norm-minimality of S implies that $S \cap \text{cl}(U_\theta^\#)$ is a collection of $2n$ disks. However, since the orientations of these disks must match up with those of $S \cap \partial U_\theta^\#$, n of the disks must intersect flow lines of $\varphi^\#$ negatively, contradicting the definition of almost-transversality.

The last possibility is that these curves do not have meridional slope, ie do not bound disks in $U_\theta^\#$. The norm-minimality of S then implies that $S \cap \text{cl}(U_\theta^\#)$ is a collection of annuli. As we noted before, this means S_α can be isotoped outside of $U_\theta^\#$, completing the proof. \square

3.4 Main theorem

Theorem 3.9 *If every boundary torus of $\overset{\circ}{M}$ has at most two ladderpole vertex classes, then M has a taut homology branched surface spanning σ .*

Proof Let v_1, \dots, v_n be the vertex classes of σ . Take embedded representatives $\overset{\circ}{S}_1, \dots, \overset{\circ}{S}_n$ of $\overset{\circ}{v}_1, \dots, \overset{\circ}{v}_n$ which lie in a regular neighborhood of $B_{\frac{\pi}{2}}$, transverse to its vertical foliation. After performing the Dehn filling at each boundary torus, these surfaces with boundary are embedded in M . By capping off their boundaries as follows, we can extend them over the Dehn filling so that they represent v_1, \dots, v_n .

For each boundary torus T_i which $\overset{\circ}{S}_j$ meets in a collection of meridians, we glue an embedded family of disks $D_{i,j} \subset \text{cl}(U_i)$ to $\overset{\circ}{S}_j \cap T_i$.

If $\overset{\circ}{S}_j$ is ladderpole at T_i for some i , then $\overset{\circ}{S}_j \cap T_i$ is an even-sized collection of cooriented ladderpoles which sums to 0 in the first homology of T_i . Thus we may glue in a disjoint collection of annuli embedded in U_i whose coorientation matches that of $\overset{\circ}{S}_j \cap T_i$. Call this collection $A_{i,j}$.

By assumption, at most one other vertex class is ladderpole at T_i ; if $\overset{\circ}{S}_k$ is also ladderpole at T_i , we take another such collection of annuli, $A_{i,k}$. We isotope the two families rel T_i so that they intersect essentially. These families of annuli, $A_{i,j}$ and $A_{i,k}$, have ladderpole boundaries by the construction, and in particular they cut the solid torus U_i into smaller solid subtori. By possibly choosing a new $A_{i,j}$ and $A_{i,k}$, we assume that they cut U_i into a minimal number of solid subtori among all possible choices.

The boundaries of these solid subtori are partitioned into subannuli which are subsets of either $A_{i,j}$, $A_{i,k}$ or ∂U_i ; the subannuli coming from $A_{i,j}$ and $A_{i,k}$ are cooriented.

We now describe configurations of annuli which we call *sources* and *sinks*, and we will then explain why the $A_{i,j}$ and $A_{i,k}$ do not form these configurations. A source is a solid subtorus of U_i having a boundary composed of subannuli from $A_{i,j}$ and $A_{i,k}$ whose coorientations point out of the subtorus. Similarly, a sink has cooriented




Figure 8: Cross-sectional views of (from left to right) a source, surgered source, sink and surgered sink. The solid lines are portions of the intersections of a meridional disk of U_i with an embedded family of annuli, and the dashed lines correspond to a separate family of annuli. By examining the coorientations in the leftmost picture, we see that none of the regions incident to A correspond to sources, so that all of the incident regions correspond to subtori distinct from the source corresponding to A . Thus the surgery merges distinct subtori, reducing the total number of subtori; the argument is symmetric for sinks. In the general case, sources and sinks may have any even number of boundary subannuli.

boundary subannuli all pointing inward. Note that because $A_{i,j}$ is embedded, no two subannuli from $A_{i,j}$ are adjacent on the boundary of a source or sink; the same holds for $A_{i,k}$. Given a source or sink, we can perform an oriented cut-and-paste surgery on $A_{i,j}$ and $A_{i,k}$ as shown in Figure 8. As explained in the caption of Figure 8, this surgery reduces the number of solid subtori of U_i . Because we chose $A_{i,j}$ and $A_{i,k}$ to minimize the number of subtori, we conclude there are no sources or sinks in U_i .

Now define $S_\ell = \mathring{S}_\ell \cup (\bigcup_i A_{i,\ell}) \cup (\bigcup_i D_{i,\ell})$. This gives us a collection of closed surfaces S_1, \dots, S_n embedded in M . We see from the definition of the puncturing map P that $P([S_\ell]) = \mathring{v}_\ell$, and since P is injective, it follows that $[S_\ell] = v_\ell$.

Each disk of $\bigcup_i D_{i,\ell}$ intersects c positively because \mathring{S}_ℓ is carried by $\mathring{\tau}$. Therefore, $\#(\bigcup_i D_{i,\ell}) = i(v_\ell, c)$. Each annulus of $\bigcup_i A_{i,\ell}$ contributes 0 to $\chi_-(S_\ell)$. Therefore, we have

$$(3.10) \quad \chi_-(S_\ell) = \chi_-(\mathring{S}_\ell) - i(v_\ell, c) = x(\mathring{v}_\ell) - i(v_\ell, c) = x(v_\ell),$$

where the second equality follows from the fact that \mathring{S}_ℓ is carried by $\mathring{\tau}$ and is thus norm-minimizing, and the third comes from Lemma 3.6.

Let B_σ be the branched sum of S_1, \dots, S_n . As the branched sum of norm-minimizing surfaces, it will be almost taut by Lemma 3.5 provided it carries no spheres (we need not consider disks since B_σ has no boundary). It spans σ because it carries a representative of each vertex.

Let us briefly review the structure of B_σ . Inside $\overset{\circ}{M}$, B_σ is a branched surface lying in a regular neighborhood of B_τ . Its boundary is a train track lying in a regular neighborhood of γ . Inside each U_i , B_σ is a branched sum of meridional disks with at most two embedded collections of annuli whose boundaries have ladderpole slope on T_i . We will use the notations $A_{i,\ell}$ and $D_{i,\ell}$ to denote the images of the $A_{i,\ell}$ and $D_{i,\ell}$ under the branched sum isotopy.

We now show that B_σ is a homology branched surface, which will show that B_σ carries no spheres, whence it is almost taut by [Lemma 2.3](#). Since an almost taut homology branched surface in a pseudo-Anosov mapping torus is taut by [Lemma 2.2](#), this will complete the proof.

If p is any point in B_σ outside of U , there is a closed positive transversal through p because B_τ is a homology branched surface. Now suppose $p \in B_\sigma$ lies inside U_i for some i . We construct a path f from p to the interior of $\overset{\circ}{M}$ which is a positive transversal to B_σ .

Begin the path f at p by traveling from B_σ into $U \setminus B_\sigma$ in the direction of the coorientation of B_σ . The endpoint of f lies in a component C of $\text{cl}(U_i) \setminus (\bigcup_\ell D_{i,\ell})$ that is homeomorphic to a solid cylinder $\{x \in \mathbb{R}^2 \mid \|x\| \leq 1\} \times (0, 1)$. The annuli $\bigcup_\ell A_{i,\ell}$ cut C into smaller subcylinders whose sides are either cooriented portions of $\bigcup_\ell A_{i,\ell}$ or subsets of ∂U_i .

Because there are no sinks in U_i , this subcylinder is either adjacent to ∂U_i or possesses an outwardly cooriented wall. If the subcylinder is adjacent to ∂U_i , extend f to a point q outside of U_i . Otherwise, extend f through an outwardly cooriented wall to enter a new subcylinder of C , and iterate this procedure. Each component of $(\bigcup_\ell A_{i,\ell}^1) \cap C$ is a cooriented rectangle separating C into two components. Each time f passes through one of these rectangles, f is blocked from passing through a second time because of the rectangle's coorientation. It follows that f never returns to the same subcylinder of C . Thus the procedure eventually terminates, and f may be extended to $q \notin U_i$.

A symmetric construction using the fact that there are no sources in U_i shows that there is a negative transversal h^{-1} from p to a point r exterior to U_i . Using the fact that B_τ is transverse to the pseudo-Anosov suspension flow in $M \setminus c$, we can find a positive transversal g from q to r . The concatenation fgh is then a closed positive transversal through p . \square

We remark that [Theorem 3.9](#) extends [Theorem 1.5](#) in [\[15\]](#), which states that if every vertex of the fibered face σ has positive intersection with each singular orbit of φ (and in particular is nonladderpole), then M has a taut oriented branched surface.

Theorem 3.11 *If $b_2(M) \leq 3$, then each boundary torus of $\overset{\circ}{M}$ witnesses at most two ladderpole vertex classes.*

Proof The oriented sum of two surfaces which are ladderpole at a component T_i of T is again ladderpole at T_i . Therefore the same is true for homology classes, and in particular the sum of two ladderpole classes lies in the boundary of σ by [Lemma 3.3](#). We conclude that all vertex classes which are ladderpole at the same boundary component of $\overset{\circ}{M}$ lie in the same facet of σ . The dimension of σ is at most 2 by assumption, so this facet has dimension at most 1. Since a 1-cell has two boundary points, at most two vertex classes can be ladderpole at the same T_i . \square

Corollary 3.12 *If $b_2(M) \leq 3$, then M has a taut homology branched surface spanning σ .*

The following corollary was known to Mosher in [\[15\]](#), but we include it here because it follows very easily from our results.

Corollary 3.13 *If φ has only one singular orbit, then M has a taut homology branched surface spanning σ .*

Proof In this case there are no ladderpole classes in $H_2(M)$. Indeed, if $\alpha \in H_2(M)$, then a representative of $\overset{\circ}{\alpha}$ carried by $B_{\frac{1}{2}}$ cannot have all ladderpole boundary components by [Lemma 3.2](#). Since $\overset{\circ}{M}$ has only one boundary component, such a representative of $\overset{\circ}{\alpha}$ cannot have any ladderpole boundary components. The result follows from [Theorem 3.9](#). \square

References

- [1] **I Agol**, *Ideal triangulations of pseudo-Anosov mapping tori*, from “Topology and geometry in dimension three” (W Li, L Bartolini, J Johnson, F Luo, R Myers, JH Rubinstein, editors), *Contemp. Math.* 560, Amer. Math. Soc., Providence, RI (2011) 1–17 [MR](#)
- [2] **KL Baker, SA Taylor**, *Dehn filling and the Thurston norm*, preprint (2016) [arXiv](#)

- [3] **A Candel, L Conlon**, *Foliations, II*, Graduate Studies in Mathematics 60, Amer. Math. Soc., Providence, RI (2003) [MR](#)
- [4] **B Farb, C J Leininger, D Margalit**, *Small dilatation pseudo-Anosov homeomorphisms and 3-manifolds*, Adv. Math. 228 (2011) 1466–1502 [MR](#)
- [5] **D Fried**, *Fibrations over S^1 with pseudo-Anosov monodromy*, from “Travaux de Thurston sur les surfaces”, Astérisque 66, Soc. Math. France, Paris (1979) Exposé 14, 251–266 [MR](#) Reprinted in *Thurston’s work on surfaces*, Mathematical Notes 48, Princeton Univ. Press (2012) 281–301
- [6] **D Fried**, *The geometry of cross sections to flows*, Topology 21 (1982) 353–371 [MR](#)
- [7] **D Futer, F Guéritaud**, *Explicit angle structures for veering triangulations*, Algebr. Geom. Topol. 13 (2013) 205–235 [MR](#)
- [8] **D Gabai**, *Foliations and the topology of 3-manifolds, II*, J. Differential Geom. 26 (1987) 461–478 [MR](#)
- [9] **D Gabai**, *Foliations and the topology of 3-manifolds, III*, J. Differential Geom. 26 (1987) 479–536 [MR](#)
- [10] **F Guéritaud**, *Veering triangulations and Cannon–Thurston maps*, J. Topol. 9 (2016) 957–983 [MR](#)
- [11] **C D Hodgson, A Issa, H Segerman**, *Non-geometric veering triangulations*, Exp. Math. 25 (2016) 17–45 [MR](#)
- [12] **C D Hodgson, J H Rubinstein, H Segerman, S Tillmann**, *Veering triangulations admit strict angle structures*, Geom. Topol. 15 (2011) 2073–2089 [MR](#)
- [13] **M Lackenby**, *Taut ideal triangulations of 3-manifolds*, Geom. Topol. 4 (2000) 369–395 [MR](#)
- [14] **Y N Minsky, S J Taylor**, *Fibered faces, veering triangulations, and the arc complex*, preprint (2016) [arXiv](#)
- [15] **L Mosher**, *Surfaces and branched surfaces transverse to pseudo-Anosov flows on 3-manifolds*, J. Differential Geom. 34 (1991) 1–36 [MR](#)
- [16] **U Oertel**, *Homology branched surfaces: Thurston’s norm on $H_2(M^3)$* , from “Low-dimensional topology and Kleinian groups” (D B A Epstein, editor), London Math. Soc. Lecture Note Ser. 112, Cambridge Univ. Press (1986) 253–272 [MR](#)
- [17] **U Oertel**, *Measured laminations in 3-manifolds*, Trans. Amer. Math. Soc. 305 (1988) 531–573 [MR](#)
- [18] **Z Sela**, *Dehn fillings that reduce Thurston norm*, Israel J. Math. 69 (1990) 371–378 [MR](#)
- [19] **B D Sterba-Boatwright**, *Thurston norm and taut branched surfaces*, Proc. Amer. Math. Soc. 102 (1988) 1052–1056 [MR](#)

- [20] **W P Thurston**, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. 339, Amer. Math. Soc., Providence, RI (1986) 99–130 [MR](#)

Mathematics Department, Yale University
New Haven, CT, United States

michael.landry@yale.edu

Received: 2 May 2017 Revised: 21 September 2017