

Dimension functions for spherical fibrations

CIHAN OKAY
ERGÜN YALÇIN

Given a spherical fibration ξ over the classifying space BG of a finite group G we define a dimension function for the m -fold fiber join of ξ , where m is some large positive integer. We show that the dimension functions satisfy the Borel–Smith conditions when m is large enough. As an application we prove that there exists no spherical fibration over the classifying space of $\mathrm{Qd}(p) = (\mathbb{Z}/p)^2 \rtimes \mathrm{SL}_2(\mathbb{Z}/p)$ with p -effective Euler class, generalizing a result of Ünlü (2004) about group actions on finite complexes homotopy equivalent to a sphere. We have been informed that this result will also appear in upcoming work of Alejandro Adem and Jesper Grodal as a corollary of a previously announced program on homotopy group actions due to Grodal.

55M35; 55S10, 55S37

1 Introduction

This paper is motivated by a conjecture about group actions on products of spheres due to Benson and Carlson [5]. The conjecture states that the maximal rank of an elementary abelian p -group contained in a finite group is at most k if and only if there exists a finite free G -CW-complex X homotopy equivalent to a product of spheres $\mathbb{S}^{n_1} \times \mathbb{S}^{n_2} \times \cdots \times \mathbb{S}^{n_k}$. For $k = 1$ this conjecture is proved by Swan [22]. The next case $k = 2$ is proved by Adem and Smith [2] for finite groups that do not involve $\mathrm{Qd}(p) = (\mathbb{Z}/p)^2 \rtimes \mathrm{SL}_2(\mathbb{Z}/p)$ for any odd prime p .

An important technique developed in [2] for constructing free actions starts with a spherical fibration over BG whose Euler class is p -effective and uses fiber joins to construct a free action on a finite complex homotopy equivalent to a product of two spheres. One source of such a spherical fibration is a finite G -CW-complex $X \simeq \mathbb{S}^n$ with rank-one isotropy. Ünlü [23] proved that for $G = \mathrm{Qd}(p)$ there exists no such finite G -CW-complex. The main goal of this paper is to extend this result by showing that there exists no spherical fibration over BG with p -effective Euler class when G is $\mathrm{Qd}(p)$. We also show that $\mathrm{Qd}(p)$ cannot act freely on a finite complex homotopy

equivalent to $\mathbb{S}^n \times \mathbb{S}^n$. However, the case of the Benson–Carlson conjecture where the dimensions of the spheres are different remains open.

Given a spherical fibration $\xi: E \rightarrow BG$ over BG with fibers \mathbb{S}^n , there is an infinite-dimensional free G -space X_ξ , defined as the pullback of ξ along the universal fibration $EG \rightarrow BG$, such that the Borel construction $EG \times_G X_\xi \rightarrow BG$ is fiber homotopy equivalent to ξ . Two G -spaces X and Y are said to be hG -equivalent if there is a zig-zag sequence of G -maps between X and Y that are weak equivalences (nonequivariantly). The fiber homotopy classes of n -dimensional spherical fibrations over BG are in one-to-one correspondence with hG -equivalence classes of G -spaces that are homotopy equivalent to \mathbb{S}^n (see Section A.3 for details). We will use this correspondence throughout the paper without further explanations.

Let G be a p -group and X be a finite-dimensional G -CW-complex. We write $H(-)$ for mod- p cohomology. Classical Smith theory says that if $H(X) \cong H(\mathbb{S}^n)$ for some n , then the fixed point space X^G also has the mod- p cohomology of a sphere. A systematic way of studying fixed point subspaces is to define dimension functions n_X by setting

$$H(X^K) \cong H(\mathbb{S}^{n_X(K)-1})$$

for a subgroup $K \leq G$. It is a fundamental fact that n_X satisfies certain properties called the Borel–Smith conditions. Smith theory fails for infinite-dimensional complexes. The problem is that, up to homotopy, every action can be made free by taking a product with the universal contractible free G -space EG . One way around this problem is to consider homotopy fixed points $X^{hK} = \text{Map}(EK, X)^K$ instead of ordinary fixed points. An important algebraic tool for studying cohomology of homotopy fixed points is Lannes’ T -functor and its variant the Fix functor. Here a technical point is that X needs to be replaced by its Bousfield–Kan p -completion X_p^\wedge , and the theory only works for elementary abelian p -groups. Then a theorem of Lannes relates the mod- p cohomology of homotopy fixed points $(X_p^\wedge)^{hV}$ to the algebraically defined object $\text{Fix}_V(H(X_{hV}))$, where V is an elementary abelian p -subgroup of G , and $X_{hV} = EV \times_V X$.

Lannes’ theory can be applied under certain conditions. We show that these conditions can be satisfied by replacing a given G -space $X \simeq \mathbb{S}^n$ with the p -completion of its m -fold join

$$X[m] = \underbrace{(X * \cdots * X)}_m^\wedge.$$

For large m we prove that classical Smith theory holds for infinite-dimensional complexes, where the role of fixed points is played by homotopy fixed points.

Theorem 1.1 [1] *Let P be a finite p -group and $X \simeq (\mathbb{S}^n)_p^\wedge$ be a P -space. Then there exists a positive integer m such that $(X[m])^{hP} \simeq (\mathbb{S}^r)_p^\wedge$ for some r .*

We have been informed that this result is going to appear in Adem and Grodal [1] and is part of a program on homotopy group actions due to Grodal which was announced previously. Using this result we can define dimension functions for mod- p spherical fibrations. A mod- p spherical fibration is a fibration whose fiber has the homotopy type of a p -completed sphere. Given a mod- p spherical fibration $\xi: E \rightarrow BG$ and a p -subgroup $Q \leq G$, we can restrict the fibration ξ to a fibration $\xi|_{BQ}: E_Q \rightarrow BQ$ by taking the pullback along the inclusion map $BQ \rightarrow BG$. This corresponds to restricting the G -action on X_ξ to a Q -action via the inclusion map. We define the integer $n_{\xi[m]}(Q)$ via the weak equivalence

$$(X_\xi[m])^{hQ} \simeq (\mathbb{S}^{n_{\xi[m]}(Q)-1})_p^\wedge$$

which is a consequence of Theorem 1.1. It turns out that for m large enough, $n_{\xi[m]}$ satisfies the Borel–Smith conditions when regarded as an integer-valued function on the set of p -subgroups of G (see Theorem 4.6). The dimension function can be made independent of m by considering a rational-valued dimension function defined by

$$\text{Dim}_\xi(Q) = \frac{1}{m} n_{\xi[m]}(Q) \quad (m \gg 0)$$

for every p -subgroup $Q \leq G$.

The Euler class of a fibration is said to be p -effective if its restriction to elementary abelian p -subgroups of maximal rank is not nilpotent. This is a condition on the Euler class of a spherical fibration that is required to obtain a free action of a rank-two group on a product of two spheres using the Adem–Smith method. As an application of the dimension function that we defined, we obtain the following.

Theorem 1.2 [1] *Assume that $p > 2$. There exists no mod- p spherical fibration $\xi: E \rightarrow BQd(p)$ with a p -effective Euler class.*

We have been informed that this result is also going to appear in [1] and was previously announced as a theorem by Grodal. As a consequence of Theorem 1.2, we obtain that the Adem–Smith method of constructing free actions on finite complexes homotopy equivalent to a product of spheres does not work for $Qd(p)$.

Another method for constructing free actions on a product of two spheres $\mathbb{S}^{n_1} \times \mathbb{S}^{n_2}$ is given by Hambleton and Ünlü [14]. This method applies only to the equidimensional case ($n_1 = n_2$). The following theorem shows that this method cannot be used for $\text{Qd}(p)$ either.

Theorem 1.3 *Let $G = \text{Qd}(p)$. Then for any $n \geq 0$, there is no finite free G -CW-complex X homotopy equivalent to $\mathbb{S}^n \times \mathbb{S}^n$.*

Therefore, if the Benson–Carlson conjecture holds, then in the construction of a complex $X \simeq \mathbb{S}^{n_1} \times \mathbb{S}^{n_2}$ with free $\text{Qd}(p)$ -action the possibilities are narrowed down to distinct-dimensional spheres with a more exotic action.

The general theory of homotopy group actions has been considered by Adem and Grodal. They have informed us that Theorems 1.1 and 1.2 will also appear in their paper [1] under preparation. The idea of using dimension functions for studying mod- p spherical fibrations goes back to Grodal and Smith’s unpublished earlier work, although an outline of their ideas can be found in the extended abstract [13]. Theorems 1.1 and 1.2 can also be thought of as corollaries of a program on homotopy group actions due to Grodal. We are grateful to Adem and Grodal for sharing their ideas with us on the subject, and we are looking forward to reading their complete account on the subject. Here we offer our proofs for Theorems 1.1 and 1.2 for completeness and to cover a gap in the existing literature. We should also mention that a result stated by Assadi [4, Corollary 4] also implies Theorems 1.1 and 1.2. Unfortunately, no proofs were provided for this result either.

The organization of the paper is as follows. In Section 2 we compute $\text{Fix}(HE)$ for a fibration $\xi: E \rightarrow B\mathbb{Z}/p$ whose fiber has the cohomology of a sphere. Our main result Theorem 1.1 (Theorem 3.11) is proved in Section 3, where we study the space of sections of a mod- p spherical fibration over the classifying space of a p -group. The dimension function for an m -fold join of a mod- p spherical fibration is defined in Section 4. We prove the nonexistence result Theorem 1.2 (Corollary 4.9) in that section. In Section 5 we prove Theorem 1.3 (Theorem 5.1). We collect some results about mapping spaces, homotopy fixed points, and fiber joins in the appendix.

Acknowledgements We thank Alejandro Adem, Matthew Gelvin, and Jesper Grodal for their comments on the first version of this paper. Yalçın is supported by a Tübitak 1001 project (grant no. 116F194).

2 Spherical fibrations and Lannes' T -functor

In this section we compute $\text{Fix}(HE)$ for a mod- p spherical fibration $E \rightarrow B\mathbb{Z}/p$. More generally, we work with fibrations where the mod- p cohomology of the fiber is isomorphic to the cohomology of a sphere. We modify the argument of [9, Chapter 3, Section 4] for the classical case, which works for group actions on finite-dimensional complexes, and use the connection between Lannes' T -functor and localization, established in [10].

2.1 Lannes' T -functor

Let \mathbf{U} (resp. \mathbf{K}) denote the category of unstable modules (resp. unstable algebras) over the mod- p Steenrod algebra \mathcal{A}_p . Let V denote an elementary abelian p -group and HV the mod- p cohomology ring of V . The tensor product functor $HV \otimes -: \mathbf{U} \rightarrow \mathbf{U}$ has a left adjoint $T^V: \mathbf{U} \rightarrow \mathbf{U}$, which is called the Lannes T -functor. Let $\mathbf{U}(HV)$ denote the category of unstable modules M with an HV -module structure such that the multiplication map $HV \otimes M \rightarrow M$ satisfies the Cartan formula. Let $f: HW \rightarrow HV$ denote the map induced by a subgroup inclusion $V \subset W$. Its adjoint $\hat{f}: T^VHW \rightarrow \mathbb{F}_p$ is determined by a ring homomorphism $\hat{f}_0: (T^VHW)^0 \rightarrow \mathbb{F}_p$ in degree zero. We define

$$T_f^V(M) = \mathbb{F}_p \otimes_{(T^VHW)^0} T^VM,$$

where the $(T^VHW)^0$ -module structure on \mathbb{F}_p is the one determined by \hat{f}_0 . Let S_f denote the multiplicatively closed subset of HV generated by the images of the Bocksteins of one-dimensional classes in HW that map nontrivially under f . The following is the main theorem of [10]. $\text{Un}(-)$ denotes the largest unstable \mathcal{A}_p -submodule.

Theorem 2.1 (Dwyer–Wilkerson [10]) *Let W be an elementary abelian p -group, V a subgroup of W , and $f: HW \rightarrow HV$ the map induced by subgroup inclusion. Suppose that M is an object of $\mathbf{U}(HV)$ that is finitely generated as a module over HV . Then there is a natural isomorphism*

$$T_f^V(M) \cong \text{Un } S_f^{-1}(M).$$

For an object M in $\mathbf{U}(HV)$ the Fix functor is defined by

$$\text{Fix}_V(M) = \mathbb{F}_p \otimes_{T^VHV} T^VM,$$

where \mathbb{F}_p is regarded as a T^VHV -module via the adjoint $\hat{\varphi}: T^VHV \rightarrow \mathbb{F}_p$ of the identity map $\varphi: HV \rightarrow HV$; see [16, Section 4.4.3] for details. We record the following properties.

Proposition 2.2 *Let M be an object in $U(HV)$. Then:*

- (1) *The natural map $T_\varphi^V M \rightarrow HV \otimes \text{Fix}_V M$ is an isomorphism in $U(HV)$.*
- (2) *If M is a finitely generated HV -module, then the localization of the natural map $M \rightarrow T_\varphi^V M$ with respect to S_φ is an isomorphism.*

Proof The first result is proved in [16, Proposition 4.5]. For the second result, the natural map is obtained as follows: Let $M \rightarrow HV \otimes T^V M$ denote the adjoint of the identity map $T^V M \rightarrow T^V M$. Composing this map with the unique algebra map $HV \rightarrow \mathbb{F}_p$ gives a map $M \rightarrow T^V M$. The desired map is obtained by applying the natural projection $T^V M \rightarrow T_\varphi^V M$ to the second factor. The fact that the resulting map is an isomorphism can be found in [10, Lemma 4.3, Section 5]. □

2.2 Spherical fibrations over $B\mathbb{Z}/p$

We will study fibrations $\xi: E \rightarrow B\mathbb{Z}/p$ where the cohomology HF of the fiber F is isomorphic to $H(\mathbb{S}^n)$ for some $n \geq 0$, and show that $\text{Fix}(HE) \cong H(\mathbb{S}^r)$ for some $-1 \leq r \leq n$. Note that mod- p spherical fibrations satisfy this condition.

We start with recalling the mod- p cohomology ring of \mathbb{Z}/p . If $p = 2$, the cohomology ring $H(\mathbb{Z}/2)$ is a polynomial algebra $\mathbb{F}_2[t]$, where t is of degree one. When $p > 2$ we have $H(\mathbb{Z}/p) = \mathbb{F}_p[t] \otimes \Lambda[s]$, where s is of degree one and $t = \beta s$. Here β is the Bockstein map. The set S_φ corresponding to the identity map $\varphi: H\mathbb{Z}/p \rightarrow H\mathbb{Z}/p$ is generated by the Bockstein of the one-dimensional class in each case. If $S = \{1, t, t^2, \dots\}$, then localization with respect to S is the same as localization with respect to S_φ . For simplicity of notation, when $V = \mathbb{Z}/p$ we will write $T = T^V$, $T_\varphi = T_\varphi^V$, and $\text{Fix} = \text{Fix}_V$.

Lemma 2.3 *For an arbitrary fibration $\xi: E \rightarrow BV$ we have $(\text{Fix}_V HE)^0 = 0$ if and only if $\xi^*: HV \rightarrow HE$ does not split in \mathbf{K} .*

Proof $(\text{Fix}_V HE)^0$ is isomorphic to $(T_\varphi^V HE)^0$, which has an \mathbb{F}_p -basis Z_φ given by the set of \mathbf{K} -maps $\alpha: HE \rightarrow HV$ such that $\alpha\xi^*$ is the identity map on HV ; see [20, Theorem 3.8.6]. □

Now we are ready to prove our main result in this section.

Theorem 2.4 *Let $\xi: E \rightarrow B\mathbb{Z}/p$ be a fibration such that $HF \cong H(\mathbb{S}^n)$. Then $\text{Fix}(HE) \cong H(\mathbb{S}^r)$ for some $-1 \leq r \leq n$.*

Proof The Serre spectral sequence of the fibration ξ has E_2 -page given by

$$H(\mathbb{Z}/p) \otimes HF \Rightarrow HE,$$

which is nonzero only in two rows since $HF \cong H(\mathbb{S}^n)$. The spectral sequence is determined by the differential $d_{n+1}: E_2^{0,n} \rightarrow E_2^{n+1,0}$, whose image lies in the polynomial part of $H\mathbb{Z}/p$ [3, page 137]. First we assume that d_{n+1} is nonzero, ie ξ^* does not split. In this case t is nilpotent in HE . Hence the localization vanishes: $S^{-1}HE = 0$. By Theorem 2.1 we have $T_\varphi HE = 0$ and the first part of Proposition 2.2 implies that $\text{Fix } HE = 0$. Next assume that $d_{n+1} = 0$, so that ξ^* splits. Localizing the natural map $HE \rightarrow T_\varphi HE$ with respect to S gives a diagram

$$(2.2.1) \quad \begin{array}{ccc} HE & \longrightarrow & T_\varphi HE \\ \downarrow & & \downarrow \\ S^{-1}HE & \xrightarrow{\cong} & S^{-1}(T_\varphi HE) \end{array}$$

Here the fact that the bottom map is an isomorphism is a consequence of the second part of Proposition 2.2. The right vertical monomorphism maps onto the unstable part of $S^{-1}(T_\varphi HE)$ as a consequence of Theorem 2.1 and the commutativity of the diagram. Since $HE \cong H(\mathbb{Z}/p) \otimes HF$ is a free $H(\mathbb{Z}/p)$ -module generated by an element of degree n , the localization map $HE \rightarrow S^{-1}HE$ is a monomorphism. Note that in the spectral sequence multiplication by t is an isomorphism. After localizing the spectral sequence the two rows extend to negative degrees. Therefore comparing the spectral sequences we see that the localization map is an isomorphism in degrees $i \geq n$. Hence from the diagram (2.2.1) it follows that the natural map $H^i E \rightarrow (T_\varphi HE)^i \cong (H(\mathbb{Z}/p) \otimes \text{Fix}(HE))^i$ is an isomorphism for $i \geq n$. Therefore

$$(2.2.2) \quad \bigoplus_{k=0}^i \text{Fix}(HE)^k \cong \bigoplus_{k=0}^i H^k(F) = \mathbb{F}_p \oplus \mathbb{F}_p \quad \text{for } i \geq n.$$

One of the factors corresponds to a generator of $\text{Fix}(HE)^0$, which is nonzero by Lemma 2.3. The other one corresponds to a generator of $\text{Fix}(HE)$ in degree $r \leq n$. Therefore $\text{Fix}(HE)$ is isomorphic to $H(\mathbb{S}^r)$, where $0 \leq r \leq n$. □

3 Cohomology of homotopy fixed points

In this section we study the homotopy fixed point space or equivalently the space of sections of a fibration by applying Lannes' results. For the relationship between the

T -functor and mapping spaces, our main references are [16] and [20]. See also the appendix for preliminaries on homotopy fixed points and space of sections.

3.1 Lannes' theorems

Lannes' T -functor gives an approximation to the cohomology of the mapping space $\text{Map}(BV, Y)$, where Y is an arbitrary space. The evaluation map

$$\text{ev}: BV \times \text{Map}(BV, Y) \rightarrow Y$$

induces a map in cohomology $HY \rightarrow HV \otimes H(\text{Map}(BV, Y))$ whose adjoint is $T^VHY \rightarrow H(\text{Map}(BV, Y))$. The adjoint map factors through

$$(3.1.1) \quad \widehat{\text{ev}}: T^VHY \rightarrow H(\text{Map}(BV, Y_p^\wedge)).$$

In degree zero it is induced by the isomorphism $[BV, Y_p^\wedge] \rightarrow \mathbf{K}(HY, HV)$ defined by applying the cohomology functor [20, page 187]. It is convenient to work on a connected component associated to the homotopy class of a map $\alpha: BV \rightarrow Y_p^\wedge$. Let $\alpha^*: HY \rightarrow HV$ denote the homomorphism induced in cohomology. The component of T^VHY at α^* is defined by

$$T^V(HY, \alpha^*) = \mathbb{F}_p \otimes_{(T^VHY)^0} T^VHY,$$

where the module structure on \mathbb{F}_p is given by the adjoint $T^VHY \rightarrow \mathbb{F}_p$ of α^* . Then $\widehat{\text{ev}}$ in (3.1.1) is the product of the maps

$$\widehat{\text{ev}}_\alpha: T^V(HY, \alpha^*) \rightarrow H(\text{Map}(BV, Y_p^\wedge)_\alpha),$$

where α runs over the homotopy classes of maps $BV \rightarrow Y_p^\wedge$.

We need the notion of freeness for the next theorem due to Lannes. Let G denote the left adjoint of the forgetful functor $\mathbf{K} \rightarrow \mathbf{E}$, where \mathbf{E} denotes the category of graded vector spaces over \mathbb{F}_p . For an object $K \in \mathbf{K}$ let ΣK^1 denote the graded vector space isomorphic to K^1 in degree one and zero in other degrees. There is an inclusion of graded vector spaces $\Sigma K^1 \rightarrow K$. Applying G to this map and composing with the counit $GK \rightarrow K$ of the adjunction gives a canonical map

$$(3.1.2) \quad \chi: G(\Sigma K^1) \rightarrow K.$$

An unstable algebra K is said to be *free in degrees ≤ 2* if χ is an isomorphism in degrees < 2 and a monomorphism in degree 2. For a more explicit definition, see [16, page 25].

Theorem 3.1 (Lannes [16, Theorem 3.2.4]) *Assume that HY and T^VHY are of finite type. If T^VHY is free in degrees ≤ 2 , then*

$$\widehat{v}_\alpha: T^V(HY, \alpha^*) \rightarrow H(\text{Map}(BV, Y_p^\wedge)_\alpha)$$

is an isomorphism of unstable algebras.

In the next section we will apply this theorem to mod- p spherical fibrations.

3.2 Mod- p spherical fibrations

A fibration whose fiber is homotopy equivalent to a p -completed sphere is called a mod- p spherical fibration. A source for such fibrations is the fiberwise completion of spherical fibrations. Let $\xi: E \rightarrow BV$ be a mod- p spherical fibration with connected fiber. There is a map of fibrations

$$\begin{array}{ccc} E & \longrightarrow & E_p^\wedge \\ \downarrow \xi & & \downarrow \xi_p^\wedge \\ BV & \longrightarrow & BV_p^\wedge \end{array}$$

where the horizontal maps are weak equivalences. In particular E is p -complete. Moreover, the diagram is a homotopy pullback diagram. This implies that there is a weak equivalence $\text{Sec}(\xi) \rightarrow \text{Sec}(\xi_p^\wedge)$ between the space of sections of these fibrations induced by the p -completion map. Therefore in applying Lannes' theory we can ignore the p -completions up to weak equivalence. We are interested in the cohomology of the space of sections $\text{Sec}(\xi)$. As explained in Appendices A.1 and A.2, the space of sections $\text{Sec}(\xi)$ is weakly equivalent to the homotopy fixed point space X_ξ^{hV} , where X_ξ is the V -space defined as the pullback of ξ along the universal bundle $EV \rightarrow BV$. The space of homotopy sections $\text{hSec}(\xi)$ is isomorphic to $BV \times \text{Sec}(\xi)$ as simplicial sets (see Section A.1). We will use Lannes' theory to study the cohomology of space of homotopy sections. Consider the diagram

$$\begin{array}{ccc} [BV, E] & \xrightarrow{\cong} & \mathbf{K}(HE, HV) \\ \downarrow & & \downarrow \\ [BV, BV] & \xrightarrow{\cong} & \mathbf{K}(HV, HV) \end{array}$$

induced by ξ , where the horizontal maps are bijections. Let Z_φ denote the subset of maps in $\mathbf{K}(HE, HV)$ which split ξ^* induced in cohomology. The subset of maps

in $[BV, E]$ that split ξ up to homotopy is in one-to-one correspondence with Z_φ . Then

$$T_\varphi^V HE = \prod_{\alpha^* \in Z_\varphi} T^V(HE, \alpha^*),$$

and the product of the evaluation maps \widehat{ev}_α gives a map

$$(3.2.1) \quad T_\varphi^V HE \rightarrow H(\text{hSec}(\xi)).$$

By Theorem 3.1 this map is an isomorphism of unstable algebras if $T_\varphi^V HE$ is free in degrees ≤ 2 . Note that the conditions that HE and $T^V HE$ are of finite type are satisfied in this case because of the spectral sequence calculation and by Theorem 2.4.

Theorem 3.2 *Let $\xi: E \rightarrow BV$ be a mod- p spherical fibration and X_ξ denote the pullback of ξ along the universal fibration $EV \rightarrow BV$. Assume $\text{Fix}_V(HE) \cong H(S^r)$ for some $-1 \leq r \leq n$. If $r \neq 1$, then*

$$H(X_\xi^{hV}) \cong H(S^r).$$

Proof By the results in Section A.2, $X_\xi^{hV} \simeq \text{Sec}(\xi)$. Since $\text{hSec}(\xi) \simeq BV \times \text{Sec}(\xi)$, it is enough to show that the map in (3.2.1) is an isomorphism. When $r = -1$, the result follows from Lemma 2.3. For the cases $r = 0$ and $r > 1$ we check the freeness condition. In (3.1.2) it turns out that the object $G(\Sigma K^1)$ is isomorphic to HW , where W is the \mathbb{F}_p -dual of K^1 . Therefore χ is an isomorphism in degrees ≤ 2 if and only if $H^2W \rightarrow K^2$ is a monomorphism. We claim that $T_\varphi^V HE$ is free in degrees ≤ 2 when $r = 0$ and $r > 1$. If $r = 0$, then the set Z_φ contains two maps α_0 and α_1 , and $T_\varphi^V HE = T^V(HE, \alpha_0) \oplus T^V(HE, \alpha_1)$, where each component is isomorphic to HV . Note that for $K = HV$ the map χ is an isomorphism. Hence the freeness condition is satisfied. For $r > 1$ we have $T_\varphi^V HE = T^V(HE, \alpha)$ for a unique homotopy class of a map α . The freeness property holds since $(T_\varphi^V HE)^1 = H^1V$ and $H^2V \subset (T_\varphi^V HE)^2$. \square

Remark 3.3 Note that in general $T_\varphi^V HE$ is not free in degrees ≤ 2 when $r = 1$. Hence in this case we cannot apply the theorem of Lannes (Theorem 3.1) to calculate the cohomology of the homotopy fixed point space. See also [16, Theorem 4.9.3].

We turn to another theorem of Lannes to study the homotopy type of the homotopy fixed point space.

Theorem 3.4 [16, Corollary 3.4.3] *Assume $HY, T^V HY$, and $H(\text{Map}(BV, Y)_\alpha)$ are of finite type. Then $T^V(HY, \alpha^*) \rightarrow H(\text{Map}(BV, Y)_\alpha)$ is an isomorphism of unstable algebras if and only if $(\text{Map}(BV, Y)_\alpha)_p^\wedge \rightarrow \text{Map}(BV, Y_p^\wedge)_\alpha$ is a homotopy equivalence.*

Using this theorem, the Fix calculation of Theorem 2.4, and Theorem 3.2 we can determine the homotopy type of the space of sections of a mod- p spherical fibration over $B\mathbb{Z}/p$.

Theorem 3.5 *Let $\xi: E \rightarrow B\mathbb{Z}/p$ be a mod- p spherical fibration such that $HF \cong H(S^n)$ and $\text{Fix}(HE) \cong H(S^r)$, where $r \neq 1$. Let X_ξ denote the pullback of ξ along $E\mathbb{Z}/p \rightarrow B\mathbb{Z}/p$. Then*

$$X_\xi^{h\mathbb{Z}/p} \simeq (S^r)_p^\wedge,$$

where $-1 \leq r \leq n$.

Proof In Theorem 2.4 we showed that $\text{Fix}(HE) \cong H(S^r)$ for some $-1 \leq r \leq n$. If $r \neq 1$, then Theorem 3.2 implies that $H(\text{Sec}(\xi)) \cong H(S^r)$. The section space is the product of mapping spaces $\text{Map}(BV, E)_\alpha$, where α is a representative of a homotopy class such that α^* lies in Z_φ . Applying Theorem 3.4 to each component we obtain a homotopy equivalence

$$\text{Sec}(\xi)_p^\wedge \rightarrow \text{Sec}(\xi)$$

after identifying $\text{Sec}(\xi) \simeq \text{Sec}(\xi_p^\wedge)$ up to homotopy. Therefore $\text{Sec}(\xi) \simeq X_\xi^{h\mathbb{Z}/p}$ is a p -complete space that has the cohomology of a sphere. \square

3.3 Fiber joins and the Fix functor

Next we look at the relationship between the Fix functor and fiber joins to be able to go around the problem in Theorem 3.5 when $r = 1$. Let $\xi_1: E_1 \rightarrow B\mathbb{Z}/p$ and $\xi_2: E_2 \rightarrow B\mathbb{Z}/p$ be two fibrations with fibers F_1 and F_2 , respectively. We assume that $HF_i \cong H(S^{n_i})$ and $\xi_1^*: H\mathbb{Z}/p \rightarrow HE_1$ splits.

Lemma 3.6 *The natural map*

$$\text{Fix}(HE_1) \otimes \text{Fix}(HE_2) \rightarrow \text{Fix } H(E_1 \times_{B\mathbb{Z}/p} E_2)$$

is an isomorphism.

Proof Consider the pullback diagram of fibrations

$$\begin{array}{ccccc} & & F_1 & \xlongequal{\quad} & F_1 \\ & & \downarrow & & \downarrow \\ F_2 & \longrightarrow & E_1 \times_{B\mathbb{Z}/p} E_2 & \xrightarrow{p_1} & E_1 \\ \parallel & & \downarrow p_2 & & \downarrow \xi_1 \\ F_2 & \longrightarrow & E_2 & \xrightarrow{\xi_2} & B\mathbb{Z}/p \end{array}$$

We have $HE_1 = H\mathbb{Z}/p \otimes HF_1$. The differential d_{n+1} in the spectral sequence of ξ_2 is either zero or t^α . By comparing the spectral sequences we see that $H(E_1 \times_{B\mathbb{Z}/p} E_2)$ is either isomorphic to $HE_1 \otimes HF_2$ or $HE_1/(t^\alpha)$. Consider the natural map

$$\theta: HE_1 \otimes_{H\mathbb{Z}/p} HE_2 \rightarrow H(E_1 \times_{B\mathbb{Z}/p} E_2)$$

of unstable modules induced by p_1 and p_2 . If $d_{n+1} = 0$, then the tensor product is isomorphic to $HE_1 \otimes HF_2$. If the differential is given by t^α , then it becomes $HE_1 \otimes_{H\mathbb{Z}/p} (H\mathbb{Z}/p)/(t^\alpha) \cong HE_1/(t^\alpha)$. Therefore in both cases θ is an isomorphism of unstable modules. In fact it is a morphism in $U(H\mathbb{Z}/p)$. Then the result follows from the isomorphism $\text{Fix}(M_1 \otimes_{H\mathbb{Z}/p} M_2) \cong \text{Fix}(M_1) \otimes \text{Fix}(M_2)$, which is valid for $U(H\mathbb{Z}/p)$ -modules [16, Theorem 4.6.2.1]. □

Proposition 3.7 *Assume that $\text{Fix } H(E_i) \cong H(S^{r_i})$ for some r_i . Then there is an isomorphism*

$$\text{Fix}(H(E_1 *_{B\mathbb{Z}/p} E_2)) \cong H(S^{r_1+r_2+1}).$$

Proof Consider the homotopy pushout square

$$(3.3.1) \quad \begin{array}{ccc} E_1 \times_{B\mathbb{Z}/p} E_2 & \longrightarrow & E_1 \\ \downarrow & & \downarrow \\ E_2 & \longrightarrow & E_1 *_{B\mathbb{Z}/p} E_2 \end{array}$$

We can assume that $E_i = (X_i)_{h\mathbb{Z}/p}$ for some \mathbb{Z}/p -space X_i . The assignment $X \mapsto \text{Fix } H(X_{h\mathbb{Z}/p})$ defines an equivariant cohomology theory on the category of \mathbb{Z}/p -spaces [16, Section 4.7]. Then associated to the pushout diagram there is a Mayer–Vietoris sequence, which breaks into short exact sequences

$$0 \rightarrow \text{Fix } H(E_1)^q \oplus \text{Fix } H(E_2)^q \rightarrow (\text{Fix } H(E_1) \otimes \text{Fix } H(E_2))^q \rightarrow \text{Fix } H(E_1 *_{B\mathbb{Z}/p} E_2)^{q+1} \rightarrow 0,$$

where we used Lemma 3.6 for the middle term. In degree zero we need to consider the reduced cohomology groups. Compare this sequence to the Mayer–Vietoris sequence of the homotopy pushout

$$\begin{array}{ccc} S^{r_1} \times S^{r_2} & \longrightarrow & S^{r_1} \\ \downarrow & & \downarrow \\ S^{r_2} & \longrightarrow & S^{r_1} * S^{r_2} \end{array}$$

The result follows from $S^{r_1} * S^{r_2} \cong S^{r_1+r_2+1}$. □

Let $\xi: E \rightarrow B$ be a fibration with fiber F . The fiberwise p -completion (Section A.4) of ξ is a fibration $\xi_{p/B}^\wedge: E_{p/B}^\wedge \rightarrow B$ whose fiber is F_p^\wedge . We use the following notation (Section A.5)

$$X[m] = \underbrace{(X * \cdots * X)}_m^\wedge_p$$

and for a fibration $\xi: E \rightarrow B$ we define

$$E_{/B}[m] = \underbrace{(E *_B \cdots *_B E)}_m^\wedge_{p/B}$$

and denote the associated fibration by $\xi[m]: E_{/B}[m] \rightarrow B$.

Corollary 3.8 *Let $X = X_\xi$ and r be defined as in Theorem 3.5. Then for all $m > 2$,*

$$(X[m])^{h\mathbb{Z}/p} \simeq \mathbb{S}^r[m].$$

Proof By Corollary A.3 we have a fiber homotopy equivalence $(X[m])_{h\mathbb{Z}/p} \simeq E_{/B\mathbb{Z}/p}[m]$. Since $X[m]$ is homotopy equivalent to a p -completed sphere, using Proposition 3.7 we obtain

$$\text{Fix } H(E_{/B\mathbb{Z}/p}[m]) \cong \text{Fix } H(E *_B\mathbb{Z}/p \cdots *_B\mathbb{Z}/p E) \cong H(\mathbb{S}^r[m]).$$

Therefore we can apply Theorem 3.5 to $\xi[m]$. □

Remark 3.9 According to Theorem 3.5, as long as $r \neq 1$ the statement of Corollary 3.8 holds with $m = 1$. The problem we faced for $r = 1$ can be handled by taking joins. When $r = 1$ it suffices to take $m = 2$ to obtain

$$(X[2])^{h\mathbb{Z}/p} \simeq (\mathbb{S}^3)_p^\wedge.$$

3.4 Finite p -groups

Next we extend our results to p -groups. Let P be a finite p -group and $Z \cong \mathbb{Z}/p$ be a subgroup of P contained in the center. Consider a mod- p spherical fibration $\xi: E \rightarrow BP$, and let $X = X_\xi$. We are interested in computing the homotopy type of the homotopy fixed point space X^{hP} . By transitivity of homotopy fixed points (Appendix A.2) we have

$$X^{hP} \simeq Y^{hP/Z},$$

where $Y = \text{Map}(EP, X)^Z \simeq X^{hZ}$. By replacing X with $X[k]$ for some k and using Corollary 3.8, we can ensure that X^{hZ} is homotopy equivalent to a p -completed

sphere. Now we can consider the P/Z -space Y . But to be able to determine the homotopy type of $Y^{hP/Z}$ we may need to replace Y with $Y[l]$. At this step we need the following lemma.

Lemma 3.10 *Let X_1 and X_2 be P -spaces such that for $i = 1, 2$, we have that $X_i \simeq (\mathbb{S}^{n_i})_p^\wedge$ and $X_i^{hZ} \simeq (\mathbb{S}^{r_i})_p^\wedge$ for some $r_i > 0$. There is a weak equivalence*

$$\alpha: (X_1^{hZ} * X_2^{hZ})_p^\wedge \rightarrow ((X_1 * X_2)_p^\wedge)^{hZ},$$

which is induced by a map of P/Z -spaces when the homotopy fixed point spaces are interpreted as mapping spaces.

Proof We describe the map α . There is a natural map of P/Z -spaces

$$\alpha': \text{Map}(EP, X_1)^Z * \text{Map}(EP, X_2)^Z \rightarrow \text{Map}(EP, X_1 * X_2)^Z$$

defined by $\alpha'[f, g, t](z) = [f(z), g(z), t]$. Note that $\text{Map}(EP, X)^Z$ is weakly equivalent to X^{hZ} via the natural map $EZ \rightarrow EP$. Hence we obtain

$$X_1^{hZ} * X_2^{hZ} \rightarrow (X_1 * X_2)^{hZ}.$$

Composing this with the natural map $(X_1 * X_2)^{hZ} \rightarrow ((X_1 * X_2)_p^\wedge)^{hZ}$, we obtain a map

$$X_1^{hZ} * X_2^{hZ} \rightarrow ((X_1 * X_2)_p^\wedge)^{hZ}.$$

Completion of this map at p gives the map α . Note that since $r_1 + r_2 + 1 > 1$ we can apply Theorem 3.5 to conclude that $((X_1 * X_2)_p^\wedge)^{hZ}$ is p -complete. To see that α is a weak equivalence it suffices to show that the map induced in mod- p cohomology is an isomorphism. A Mayer-Vietoris type of argument shows that

$$H((X_1^{hZ} * X_2^{hZ})_p^\wedge) \cong H(\mathbb{S}^{r_1+r_2+1}).$$

On the other hand, Proposition 3.7 implies that $\text{Fix } H(X_1 * X_2)_{hZ} \cong H(\mathbb{S}^{r_1+r_2+1})$ and by Theorem 3.5 $((X_1 * X_2)_p^\wedge)^{hZ}$ is weakly equivalent to the p -completion of $\mathbb{S}^{r_1+r_2+1}$. □

Now we are ready to prove the main theorem of this section.

Theorem 3.11 *Let P be a finite p -group and $\xi: E \rightarrow BP$ be a mod- p spherical fibration. Then there exists a positive integer m such that $X[m]^{hP} \simeq (\mathbb{S}^r)_p^\wedge$, where $X = X_\xi$.*

Proof We will use the transitivity property (Appendix A.2) of homotopy fixed points,

$$(\text{Map}(EP, X)^Z)^{hP/Z} \simeq X^{hP},$$

where $\text{Map}(EP, X)^Z \simeq X^{hZ}$, and we will do induction on the order of P . We can assume X^{hZ} is nonempty, otherwise the result holds trivially. Using Corollary 3.8 we can replace X by $X[k]$ to ensure that $Y = \text{Map}(EP, X[k])^Z \simeq X[k]^{hZ}$ has the homotopy type of a p -completed simply connected sphere. We regard Y as a P/Z -space. Since the order of P/Z is less than the order of P there exists, by the induction hypothesis, some l such that the homotopy fixed points $(Y[l])^{hP/Z}$ is weakly equivalent to a p -completed sphere. We claim that the homotopy fixed points of $X[kl]$ under the action of P is a p -completed sphere. To see this let $A = X[k]$. There is a weak equivalence

$$(A^{hZ})[l] \rightarrow (A[l])^{hZ}$$

that is induced by a map of P/Z -spaces when regarded as a map between the associated mapping spaces. This can be shown by using Lemma 3.10 and doing induction on l . Now consider the natural P -map

$$X[kl] \rightarrow (X[k])[l],$$

which is also a weak equivalence. Using these two maps we obtain a zig-zag of weak equivalences

$$(A^{hZ})[l] \rightarrow (A[l])^{hZ} \leftarrow X[kl]^{hZ}$$

through P/Z -maps. Thus we have

$$X[kl]^{hP} \simeq (Y[l])^{hP/Z}$$

and the result follows by induction. □

4 Dimension functions

In this section we will define dimension functions for spherical fibrations and show that they satisfy the Borel–Smith conditions after taking fiber joins.

4.1 Dimension functions

Let $C(G)$ denote the ring of integer-valued functions defined on the set of all subgroups of G that are constant on G -conjugacy classes. Let $H \leq G$ be a subgroup. Given a

finite G -CW-complex Y with $H(Y) \cong H(\mathbb{S}^n)$, the dimension function n_Y of Y is defined by $H(Y^H) \cong H(\mathbb{S}^{n_Y(H)-1})$. Thus Y gives rise to an element n_Y of $C(G)$.

We extend this definition to our situation. Let $\xi: E \rightarrow BG$ be a mod- p spherical fibration. For a p -subgroup $Q \leq G$ let X_Q denote the pullback $\xi|_{BQ}$ along the universal fibration $EQ \rightarrow BQ$. By Theorem 3.11 there exists an m such that $(X_Q[m])^{hQ}$ has the homotopy type of $(\mathbb{S}^{r_Q})_p^\wedge$ for some r_Q for all p -subgroups $Q \leq G$. Note that by standard properties of homotopy fixed points and Theorem 3.5, we have $r_{Q'} \leq r_Q$ if $Q \leq Q'$ and $r_Q = r_{Q'}$ if Q is conjugate to Q' in G . Let $\mathcal{S}_p(G)$ denote the set of all p -subgroups of G . We define a function

$$n_{\xi[m]}: \mathcal{S}_p(G) \rightarrow \mathbb{Z} \quad \text{by } n_{\xi[m]}(Q) = r_Q + 1,$$

which is constant on G -conjugacy classes and call it the dimension function associated to the fibration $\xi[m]$. Given a spherical fibration we can consider the dimension function associated to its fiberwise p -completion.

Remark 4.1 To associate a dimension function to a mod- p spherical fibration independent of m we can define a rational-valued dimension function

$$\text{Dim}_\xi(Q) = \frac{1}{m} n_{\xi[m]}(Q)$$

for all p -subgroups $Q \leq G$, where m is a positive integer large enough that Theorem 3.11 holds.

4.2 Dimensions and subgroups

We will prove an important relation satisfied by the dimension functions. Let V be an elementary abelian p -group of rank two. Let $\xi: E \rightarrow BV$ be a mod- p spherical fibration and $X = X_\xi$. Assume that n_ξ is defined. (This can be achieved by replacing ξ with $\xi[m]$.) This means that the homotopy fixed points of X under the action of a subgroup of V is a p -completed sphere.

By the Thom isomorphism theorem for $W \leq V$ the reduced cohomology ring of the Thom space $\text{Th}(\xi_W)$ of the fibration $\xi_W: (X^{hW})_{hV} \rightarrow BV$ is a free HV -module on a single generator $t(\xi_W)$. There is a map $X^{hV} \rightarrow X^{hW}$ defined as the composition

$$\text{Map}(EV, X)^V \rightarrow \text{Map}(EV, X)^W \rightarrow \text{Map}(EW, X)^W$$

of the natural inclusion of the fixed points, and the map induced by $EW \rightarrow EV$. This map induces a diagram

$$(4.2.1) \quad \begin{array}{ccccc} (X^{hV})_{hV} & \longrightarrow & (X^{hW})_{hV} & & \\ \downarrow \xi_V & & \downarrow \xi_W & & \\ BV & \xlongequal{\quad} & BV & & \\ \downarrow & & \downarrow & & \\ \text{Th}(\xi_V) & \xrightarrow{\alpha} & \text{Th}(\xi_W) & \xrightarrow{\beta} & \Sigma(X_{W,V})_{hV} \end{array}$$

where $X_{W,V}$ is the cofiber of $X^{hV} \rightarrow X^{hW}$. The bottom row is a cofibration sequence. In the long exact sequence of cohomology groups

$$(4.2.2) \quad \dots \rightarrow \tilde{H}^i(\text{Th}(\xi_W)) \xrightarrow{\alpha^*} \tilde{H}^i(\text{Th}(\xi_V)) \xrightarrow{\beta^*} H^i((X_{W,V})_{hV}) \rightarrow \dots$$

we have $\alpha^*(t(\xi_W)) = e_{W,V}t(\xi_V)$ for some element $e_{W,V}$ in HV . Let us set $S_W = S_f$, where $f: HV \rightarrow HW$ is the map induced by a subgroup inclusion $W \leq V$.

Lemma 4.2 *Let $W \subset V$ be a subspace of codimension one. Then there is an isomorphism $H((X_{W,V})_{hV}) \cong HV/(e_{W,V})$.*

Proof Let Y denote the space of homotopy fixed points $\text{Map}(EV, X)^W \simeq X^{hW}$. Let $V = W \times L$ be a splitting. Then $Y^{hL} \simeq X^{hV}$, and $(X^{hV})_{hV} \rightarrow (X^{hW})_{hV}$ induces a map in cohomology $H((X^{hW})_{hV}) \rightarrow H((X^{hV})_{hV})$. Note that $(X^{hW})_{hV} \simeq BW \times Y_{hL}$ and $(X^{hV})_{hV} \simeq BW \times (Y^{hL})_{hL}$. Therefore we obtain

$$HW \otimes H(Y_{hL}) \rightarrow HW \otimes H((Y^{hL})_{hL}),$$

which becomes an isomorphism after localizing with respect to S_L . This is a consequence of the isomorphism $T_\varphi^L H(Y_{hL}) \cong H((Y^{hL})_{hL})$ implied by Theorem 3.2 and the second part of Proposition 2.2 applied to $M = H(Y_{hL})$. Therefore the localization of $H((X^{hW})_{hV}) \rightarrow H((X^{hV})_{hV})$ with respect to S_V is an isomorphism. From the map between the cofiber sequences in (4.2.1) we see that the map between the cohomology rings of Thom spaces becomes an isomorphism after localizing with respect to S_V . Thus there is a diagram

$$\begin{array}{ccc} \tilde{H}(\text{Th}(\xi_W)) & \xrightarrow{\alpha^*} & \tilde{H}(\text{Th}(\xi_V)) \\ \downarrow & & \downarrow \\ S_V^{-1} \tilde{H}(\text{Th}(\xi_W)) & \xrightarrow{\cong} & S_V^{-1} \tilde{H}(\text{Th}(\xi_V)) \end{array}$$

where the vertical arrows are injective since $\tilde{H}(\text{Th}(\xi_W))$ and $\tilde{H}(\text{Th}(\xi_V))$ are HV -free. Therefore in (4.2.2) we have that α^* is injective and β^* is surjective. Then $H((X_{W,V})_{hV})$ is the quotient of the map

$$\tilde{H}(\text{Th}(\xi_W)) \rightarrow \tilde{H}(\text{Th}(\xi_V))$$

induced by $t(\xi_W) \mapsto e_{W,V}t(\xi_V)$. □

Let us simply denote $e_{W,V}$ by e_V when W is the trivial group. Let t_L denote the generator of the polynomial part of HL . We regard t_L as an element of HV via the isomorphism $HV \cong HL \otimes HW$.

Lemma 4.3 *Assume that e_V belongs to the polynomial part of HV . We have $n_X(W) > n_X(V)$ if and only if t_L divides e_V . Moreover,*

$$e_V = u \prod_W e_{W,V},$$

where $u \in \mathbb{F}_p$ is a unit and W runs over subspaces of codimension one in V such that $n_X(W) > n_X(V)$.

Proof Note that $n_X(W) > n_X(V)$ if and only if $e_{W,V} = at_L^\alpha$ for some $\alpha > 0$ and nonzero $a \in \mathbb{F}_p$. Let β be the maximal natural number such that t_L^β divides e_V . Consider the last cofiber sequences in the diagram (4.2.1) for the pair of subgroup inclusions given by $W \subset V$ and $1 \subset V$. There is a map between the cofiber sequences

$$\begin{array}{ccccc} \text{Th}(\xi_V) & \longrightarrow & \text{Th}(\xi_W) & \longrightarrow & \Sigma(X_{W,V})_{hV} \\ \parallel & & \downarrow & & \downarrow \\ \text{Th}(\xi_V) & \longrightarrow & \text{Th}(\xi_1) & \longrightarrow & \Sigma(X_{1,V})_{hV} \end{array}$$

We claim that the map $S_W^{-1}H(X_{hV}) \rightarrow S_W^{-1}H((X^{hW})_{hV})$ is an isomorphism. This follows from the transitivity of the Borel construction. The map $((X^{hW})_{hW})_{hL} \rightarrow (X_{hW})_{hL}$ between Borel constructions with respect to the action of L induces a map between the E_2 -pages

$$H(L, S_W^{-1}H(X_{hW})) \rightarrow H(L, S_W^{-1}H((X^{hW})_{hW}))$$

of the associated spectral sequences. Then the claim follows from the isomorphism

$$S_W^{-1}H(X_{hW}) \cong S_W^{-1}H((X^{hW})_{hW}).$$

Now comparing the diagrams for $W \subset V$ and $1 \subset V$ we see that localization of $H((X_{1,V})_{hV}) \rightarrow H((X_{W,V})_{hV})$ with respect to S_W is an isomorphism. Thus by Lemma 4.2 there is an isomorphism $S_W^{-1}HV/(e_V) \cong S_W^{-1}HV/(e_{W,V})$. This forces $\alpha = \beta$. □

Proposition 4.4 *Assume e_V belongs to the polynomial part of HV . If W_1, \dots, W_s denote the subspaces of codimension one in V , then*

$$n_X(1) - n_X(V) = \sum_{i=1}^s (n_X(W_i) - n_X(V)).$$

Proof By Lemma 4.3 a codimension one subspace W contributes to the sum on the right-hand side if and only if t_L divides e_V . Thus the result follows from comparing the degrees. Note that $|e_V| = n_X(1) - n_X(V)$ and $|e_{W,V}| = n_X(W) - n_X(V)$. □

Remark 4.5 In view of Corollary A.6, by choosing m large enough in $\xi[m]$ we can achieve the property that e_V belongs to the polynomial part of HV .

4.3 Borel–Smith functions

An element τ of $C(G)$ is called a Borel–Smith function [9, Definition 5.1] if it satisfies the following conditions:

- (i) If $H \triangleleft K \leq G$ such that $K/H \cong \mathbb{Z}/p \times \mathbb{Z}/p$, and H_i/H denotes the cyclic subgroups, then

$$\tau(H) - \tau(K) = \sum_{i=0}^p (\tau(H_i) - \tau(K)).$$

- (ii) If $H \triangleleft K \leq G$ such that $K/H \cong \mathbb{Z}/p$, where $p > 2$, then $\tau(H) - \tau(K)$ is even.
- (iii) If $H \triangleleft L \triangleleft K \leq G$ such that $L/H \cong \mathbb{Z}/2$, then $\tau(H) - \tau(L)$ is even if $K/H \cong \mathbb{Z}/4$, and $\tau(H) - \tau(L)$ is divisible by 4 if K/H is a generalized quaternion group of order $\geq 2^3$.

Let $C_b(G)$ denote the additive subgroup of Borel–Smith functions in $C(G)$. We also say a function $S_p(G) \rightarrow \mathbb{Z}$ constant on the G -conjugacy classes satisfies Borel–Smith conditions if it satisfies (i), (ii), and (iii) on p -subgroups.

Theorem 4.6 *There exists a positive integer m such that $n_{\xi[m]}: S_p(G) \rightarrow \mathbb{Z}$ satisfies the Borel–Smith conditions.*

Proof Monotonicity is a consequence of Theorem 3.5. Condition (i) is proved in Proposition 4.4, where the hypothesis that e_V belongs to the polynomial part of HV holds by choosing m large enough. This is a consequence of Corollary A.6. Conditions (ii) and (iii) can be achieved by taking m large. \square

When G is a finite nilpotent group, Borel–Smith functions can always be realized as dimension functions of virtual representations. Let $\text{RO}(G)$ denote the Grothendieck group of real representations. There is an additive morphism $\dim: \text{RO}(G) \rightarrow C(G)$ which sends a real representation ρ to the function which sends a subgroup H to the dimension of the fixed subspace ρ^H . Let $C_{\text{rep}(G)}$ denote the image of this homomorphism. A key fact is that if G is a finite nilpotent group, then $C_b(G) = C_{\text{rep}(G)}$. In the case of p -groups we can use honest representations when the Borel–Smith function is also monotone.

Theorem 4.7 [9, Theorem 5.13] *If P is a p -group and τ is a monotone Borel–Smith function, then there is a real representation ρ such that $\tau = \dim \rho$.*

Up to fiber joins the dimension function of a mod- p spherical fibration can be realized by the dimension function of a real representation.

Corollary 4.8 *Given a finite p -group P and a mod- p spherical fibration $\xi: E \rightarrow BP$, there is a positive integer m and a real representation ρ such that $n_{\xi[m]} = \dim \rho$.*

4.4 Proof of Theorem 1.2

A cohomology class in $H(G)$ is called p -effective if its restriction to maximal elementary abelian p -subgroups is not nilpotent. Let $\text{Qd}(p)$ denote the semidirect product $(\mathbb{Z}/p)^2 \rtimes \text{SL}_2(\mathbb{Z}/p)$, where the special linear group acts in the obvious way.

Corollary 4.9 *Let $p > 2$. There exists no mod- p spherical fibration $\xi: E \rightarrow B\text{Qd}(p)$ with a p -effective Euler class.*

Proof The idea of the proof follows [23, Theorem 3.3] but here we use dimension functions for mod- p spherical fibrations. Assume that there is a fibration ξ with an effective Euler class. Consider the dimension function $n_{\xi[m]}$ for some large m . Its Euler class is still effective by Corollary A.6. Let P be a Sylow p -subgroup of $G = \text{Qd}(p)$. The center $Z(P)$ is a cyclic group of order p . By Theorem 4.6 we can

choose m large enough that the dimension function of the restricted bundle $\xi[m]|_{BP}$ belongs to $C_b(P)$. Then Theorem 4.7 implies that there is a real representation ρ which realizes this dimension function. Since the Euler class is p -effective, the dimensions of subspaces of ρ fixed under a cyclic subgroup $C \leq P_i$ have the property that $\dim \rho^C = 0$ if and only if $C = Z(P)$ (see [23, Lemma 3.4]). Therefore on cyclic p -subgroups of P the dimension function $n_{\xi[m]}$ is zero only at $Z(P)$ but in G the center $Z(P)$ is conjugate to a noncentral cyclic p -subgroup C of P . Then $n_{\xi[m]}(Z(P)) = n_{\xi[m]}(C)$, but this gives a contradiction. \square

5 Qd(p)-action on $S^n \times S^n$

In this section we prove the following theorem.

Theorem 5.1 *Let $G = \text{Qd}(p)$. Then for any $n \geq 0$, there is no finite free G -CW-complex X homotopy equivalent to $S^n \times S^n$.*

If $p = 2$, then $G = \text{Qd}(2)$ is isomorphic to the symmetric group S_4 , which includes A_4 as a subgroup. In this case the theorem follows from a result of Oliver [19] which says that the group A_4 does not act freely on a finite complex X homotopy equivalent to a product of two equal-dimensional spheres. Also note that for $n = 0$, the statement holds for obvious reasons. Hence it is enough to prove the theorem when p is an odd prime and $n \geq 1$.

Lemma 5.2 *Let G be a finite group generated by elements of odd order. Let X be a finite free G -CW-complex homotopy equivalent to $S^n \times S^n$ for some $n \geq 1$. Then n is odd, and the induced G -action on $H^*(X; \mathbb{Z})$ is trivial.*

Proof It is enough to prove this for the case $G = \mathbb{Z}/p^k$, where p is an odd prime. By induction we can assume that the action of the maximal subgroup $H \leq G$ on cohomology is trivial. Consider the $G/H \cong \mathbb{Z}/p$ action on $H^*(X; \mathbb{Z})$. The only indecomposable \mathbb{Z} -free $\mathbb{Z}[\mathbb{Z}/p]$ -modules are either 1-dimensional, $(p-1)$ -dimensional, or p -dimensional [15, Theorem 2.6]. This gives that for $p > 3$, the G -action on $H^*(X; \mathbb{Z})$ is trivial. For $p = 3$, the only nontrivial module can occur in dimension n , and in this case G/H acts on $H^n(X; \mathbb{Z})$ with the action $x \rightarrow -y$ and $y \rightarrow x - y$, where x and y are generators of $H^n(X; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}$. Note that the trace of this action is -1 , so by the Lefschetz trace formula, $L(f) = 2 - (-1) = 3$ when n is odd, and $L(f) = 0 + (-1) = -1$ when n is even. In either case $L(f) \neq 0$, hence G cannot

admit a free action on X if the G/H action on homology is nontrivial. If the action is trivial, then again by the Lefschetz trace formula, n must be odd. \square

The group $SL_2(p)$ is generated by elements of order p . For example, we can take

$$A = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$$

as a set of generators. Since $Qd(p) = (\mathbb{Z}/p)^2 \rtimes SL_2(\mathbb{Z}/p)$ is a semidirect product of $(\mathbb{Z}/p)^2$ with $SL_2(p)$, it is also generated by elements of order p . Hence we conclude the following.

Proposition 5.3 *Let $G = Qd(p)$, where p is an odd prime. Suppose that there exists a finite free G -CW-complex X homotopy equivalent to $\mathbb{S}^n \times \mathbb{S}^n$ for some $n \geq 1$. Then n is odd and G acts trivially on $H^*(X; \mathbb{Z})$.*

To complete the proof of Theorem 5.1, we use the Borel construction. Let $G = Qd(p)$ with p odd, and let X be a finite free G -CW-complex homotopy equivalent to $\mathbb{S}^n \times \mathbb{S}^n$ for some integer $n \geq 1$. By Proposition 5.3, the induced action of G on X is trivial and $n = 2k - 1$ for some $k \geq 1$. Consider the Borel fibration $X_{hG} \rightarrow BG$, where $X_{hG} = EG \times_G X$. There is an associated spectral sequence with E_2 -term

$$E_2^{i,j} = H^i(G; H^j(X))$$

that converges to $H^{i+j}(X_{hG})$.

Note that since G acts freely on X we have $X_{hG} \simeq X/G$. From this one obtains that the cohomology ring $H^*(X_{hG})$ is finite-dimensional in each degree and vanishes above some degree. The first nonzero differential in the above spectral sequence takes the generators of $H^{2k-1}(X) = \mathbb{F}_p \oplus \mathbb{F}_p$ to the cohomology classes μ_1 and μ_2 in $H^{2k}(G)$. These classes are called the k -invariants of the G -space X .

For any subgroup $H \leq G$, we can restrict the above spectral sequence to the one for the action of H on X . This follows from the fact that the Borel construction is functorial. The k -invariants of this restricted action will be $\text{res}_H^G \mu_1$ and $\text{res}_H^G \mu_2$, where

$$\text{res}_H^G: H^*(G) \rightarrow H^*(H)$$

denotes the homomorphism induced by inclusion of H into G .

Let V denote the (unique) normal elementary abelian subgroup $\mathbb{Z}/p \times \mathbb{Z}/p$ in G . Let θ_1 and θ_2 denote the k -invariants of the restricted V -action on X . Note that the classes θ_i are restrictions of cohomology classes μ_1 and μ_2 in $H^{2k}(G)$. By the

Cartan–Eilenberg stable element theorem, the classes θ_i lie in the invariant subring $H^*(V)^{\text{SL}_2(p)}$. This invariant ring is described in detail in [17, Proposition 1.4.1 and Claim 1.4.2]. If we write

$$H^*(V) = \mathbb{F}_p[x, y] \otimes \wedge(u, v),$$

then $H^{\text{ev}}(V)^{\text{SL}_2(p)} = \mathbb{F}_p[x, y]^{\text{SL}_2(p)} \otimes \wedge(vu)$, where $\mathbb{F}_p[x, y]^{\text{SL}_2(p)} = \mathbb{F}_p[\xi, \zeta]$ is a polynomial subalgebra generated by

$$\xi = \sum_{i=0}^p (x^{p-i} y^i)^{p-1} \quad \text{and} \quad \zeta = xy^p - yx^p.$$

For $i = 1, 2$, let $\theta_i = f_i(\xi, \zeta) + uv g_i(\xi, \zeta)$ for some polynomials f_i and g_i . Since θ_1 and θ_2 are integral classes, ie they are in the image of the map $H^*(V, \mathbb{Z}) \rightarrow H^*(V, \mathbb{F}_p)$ induced by mod- p reduction, we have $g_i = 0$ for $i = 1, 2$. This can be seen easily by applying the Bockstein operator $\beta: H^*(V) \rightarrow H^{*+1}(V)$ to the classes θ_i . Since $\beta(u) = x$ and $\beta(v) = y$, we obtain

$$0 = \beta(\theta_i) = \beta(uv)g_i = (xv - uy)g_i.$$

This gives $g_i = 0$. Hence the k -invariants θ_1 and θ_2 lie in the polynomial subalgebra $\mathbb{F}_p[\xi, \zeta]$.

Let $I = (\theta_1, \theta_2)$ be the ideal in $H^*(V)$ generated by θ_1 and θ_2 . By a theorem of Carlsson [8, Corollary 7], the cohomology ring $H^*(X_{hV}) \cong H^*(X/V)$ is isomorphic to $H^*(V)/I$. In fact it is proved that $\{\theta_1, \theta_2\}$ is a homogenous system of parameters, hence it gives a regular sequence (in any order it is taken). This makes the spectral sequence collapse at the E_{n+2} -page, and gives that $H^*(X_{hV}) \cong H^*(V)/I$, which is finite-dimensional as a vector space.

For every $j \geq 0$, let $P^j: H^r(V) \rightarrow H^{r+2(p-1)j}(V)$ denote the Steenrod operation. From the isomorphism above, we obtain that the ideal I is closed under Steenrod operations, meaning that for every $j \geq 0$, we have $P^j(u) \in I$ for every $u \in I$. A slightly stronger Steenrod closeness condition also holds:

Lemma 5.4 *Let M denote the \mathbb{F}_p -vector space generated by θ_1 and θ_2 . For every $m \in M$, and for $j \geq 0$, there exist α_1 and α_2 in the invariant subring $\mathbb{F}_p[\xi, \zeta]$ such that $P^j(m) = \sum_i \alpha_i \theta_i$.*

Proof Note that it is enough to show that the elements α_i can be chosen from $H^*(V)^{\text{SL}_2(p)}$. Since both $P^j(m)$ and θ_i belongs to the polynomial part of the invariant

subring, this will imply α_i also belongs to $\mathbb{F}_p[\xi, \zeta]$. For this we will show that α_i can be taken as $\alpha_i = \text{res}_V^G \lambda_i$ for some $\lambda_i \in H^*(G)$. Hence it is enough to show that the ideal $J \subset H^*(G)$ generated by μ_1 and μ_2 is closed under Steenrod operations, where μ_1 and μ_2 are the k -invariants of the G -action on $S^n \times S^n$. For this, first note that $\{\mu_1, \mu_2\}$ is a homogeneous system of parameters because its restriction to elementary abelian subgroups is a homogeneous system of parameters. If Z is the center of a Sylow p -subgroup of G , then $\text{res}_Z^G \mu_i \neq 0$ for some i . Assume that $\text{res}_Z^G \mu_1 \neq 0$, then by [7, Theorem 12.3.3], μ_1 is a nonzero divisor. This gives that the spectral sequence $E_2^{i,j} = H^i(G; H^j(X))$ described above collapses at E_{n+2} -page, in particular, $d_{2n+1} = 0$. Hence $H^*(G)/J$ is a module over Steenrod algebra, giving that J is closed under Steenrod operations. \square

If $M \subset H^{2k}(V)$ is a subspace which lies inside the subring $\mathbb{F}_p[\xi, \zeta]$ and has the Steenrod closeness property given in the above lemma, then we say the subspace M is closed under Steenrod operations in the invariant subring.

Proposition 5.5 *Let $M \subset H^{2k}(V)$ be a nonzero subspace that lies inside the invariant subring $\mathbb{F}_p[\xi, \zeta]$. If M is closed under the Steenrod operations in the invariant subring, then M lies in the principal ideal generated by ζ .*

Proof We will use an argument similar to the argument given by Oliver [19] for A_4 -actions on $S^n \times S^n$. A nonzero homogeneous polynomial $f \in \mathbb{F}_p[\xi, \zeta]$ can be written as a sum

$$f = (f_0(\xi) + f_1(\xi)\zeta + \dots + f_t(\xi)\zeta^t)\zeta^s$$

for some homogeneous polynomials $f_i(\xi) = a_i \xi^{d_i}$, where f_0 and f_t are nonzero and $s \geq 0$. Note that the degree of ξ is $2p(p-1)$, the degree of ζ is $2(p+1)$, hence for dimension reasons $f_i = 0$ when p does not divide i . So we can conclude that every homogenous element f in $\mathbb{F}_p[\xi, \zeta]$ is of the form

$$f = \left(\sum_{i=0}^{t'} f_i(\xi)\zeta^{ip} \right) \zeta^s$$

for some homogenous polynomials $f_i(\xi) = a_i \xi^{d_i}$ with $a_i \in \mathbb{F}_p$ and $f_0(\xi) \neq 0$. By direct calculation, it is easy to see $P^1(\zeta) = 0$ and $P^1(\xi) = \zeta^{p-1}$. So if $P^1(f) = 0$, then

$$\left(\sum_{i=0}^{t'} P^1(f_i)\zeta^{ip} \right) \zeta^s = 0,$$

which gives $0 = P^1(f_i) = a_i d_i \xi^{d_i-1} \zeta^{p-1}$ for all i . Hence for every i , we have $d_i = p d'_i$ for some d'_i . From this we conclude that f is of the form $(f')^p \zeta^s$ for some f' when $P^1(f) = 0$. Furthermore, if the degree of f is divisible by p , then $s = p s'$ for some s' , and hence in this case we have $f = (f'')^p$ for some homogeneous polynomial f'' in $\mathbb{F}_p[\xi, \zeta]$. We will use these observations in the calculations below.

Now let $M \subset H^{2k}(V)$ be a subspace as in the proposition. If p does not divide k , then each $m \in M$ can be written as $m = \zeta m'$ for some $m' \in \mathbb{F}_p[\xi, \zeta]$. In this case we have $M \subset (\zeta)$ as desired. So let us assume that p divides k . Then every $m \in M$ is of the form

$$m = \left(\sum_{i=0}^{t'} f_i(\xi) \zeta^{ip} \right) \zeta^{s'p}$$

for some homogeneous polynomials $f_i(\xi) = a_i \xi^{d_i}$ with $a_i \in \mathbb{F}_p$ and $f_0(\xi) \neq 0$. Since M is closed under Steenrod operations in the invariant subring, $P^1(m) = \sum_i \alpha_i m_i$ for some $m_i \in M$ and $\alpha_i \in \mathbb{F}_p[\xi, \zeta]$. However, there are no $2(p-1)$ -dimensional classes in the invariant subring, hence we must have $P^1(m) = 0$ for every $m \in M$. By the above observation this implies that every $m \in M$ is equal to $(m')^p$ for some (unique) $m' \in H^{2k/p}(V)$. Let $M' \subset H^{2k/p}(V)$ be the subspace formed by elements m' such that $(m')^p \in M$. Note that M' also lies in the invariant subring $\mathbb{F}_p[\xi, \zeta]$.

Let $r \geq 1$ be the largest integer such that there exists a subspace $M_r \subset H^{2k/p^r}(V)$ such that M_r lies in the invariant subring $\mathbb{F}_p[\xi, \zeta]$ and every $m \in M$ is of the form $m = u^{p^r}$ for some $u \in M_r$. For every $j \geq 0$, we have

$$P^j(f^{p^b}) = \begin{cases} (P^a(f))^{p^b} & \text{if } j = ap^b, \\ 0 & \text{otherwise.} \end{cases}$$

So we have $P^{p^r}(u^{p^r}) = (P^1(u))^{p^r}$. Since M is closed under Steenrod operations in the invariant subring, we also have $P^{p^r}(m) = \sum_i \alpha_i m_i$ for some α_i in $\mathbb{F}_p[\xi, \zeta]$. Hence we have

$$(P^1(u))^{p^r} = \sum_i \alpha_i u_i^{p^r}$$

for some α_i in the invariant subring. We claim that the coefficients α_i are of the form $\alpha_i = (\alpha'_i)^{p^r}$ for some α'_i .

To see this first note that P^{p^r} increases the dimension by $2p^r(p-1)$, so the dimension of α_i is $2p^r(p-1)$. If we apply P^1 to the above equation we get $0 = \sum_i P^1(\alpha_i) u_i^{p^r}$. Each element $P^1(\alpha_i)$ has degree $2(p^r+1)(p-1)$ and each element $u_i^{p^r}$ has degree at least $2(p+1)p^r$, so by dimension reasons we have $P^1(\alpha_i) = 0$ for all i . Hence

we have $\alpha_i = (\alpha'_i)^p$ for some α'_i . Plugging this into the above equation and taking the p^{th} root we get

$$(P^1(u))^{p^{r-1}} = \sum_i \alpha'_i u_i^{p^{r-1}}.$$

Repeating the same argument r times, we get $\alpha_i = (\alpha'_i)^{p^r}$ for some α'_i that lies in the invariant subring. This gives $P^1(u) = \sum_i \alpha'_i u_i$ for some α'_i . Since the degree of α'_i is $2(p-1)$ and there is no invariant element in that dimension, we get $\alpha'_i = 0$. This gives $P^1(u) = 0$. From this we conclude that for all u we have $u = (u')^p$ for some u' . This contradicts the assumption that r is the largest integer such that every $m \in M$ is of the form $m = u^{p^r}$ for some $u \in \mathbb{F}_p[\xi, \zeta]$. □

We are now ready to complete the proof of Theorem 5.1.

Proof of Theorem 5.1 Let $I = (\theta_1, \theta_2)$ denote the ideal generated by the k -invariants of the V -action on $X \simeq \mathbb{S}^n \times \mathbb{S}^n$. By [8, Corollary 7], there is an isomorphism

$$H^*(X/V) \cong H^*(V)/I,$$

hence $H^*(V)/I$ is a finite-dimensional vector space. Moreover the ideal I is closed under the Steenrod operations and it is of the form $M \cdot H^*(V)$, where $M \subset \mathbb{F}_p[x, y]^{\text{SL}_2(p)}$ is the subspace generated by θ_1 and θ_2 . By Lemma 5.4, M is closed under Steenrod operations in the invariant subring. Hence by Proposition 5.5 the ideal I is included in the principal ideal (ζ) . This gives a contradiction because the fact that $H^*(V)/I$ is finite-dimensional implies that I cannot be included in a principal ideal generated by $\zeta = xy(x+y) \cdots (x+(p-1)y)$ by standard results in commutative algebra (see [8, Proposition 3]). □

Appendix

By a space we mean either a topological space or a simplicial set. The relation between the two is given by the singular simplicial set functor and the geometric realization functor. The category of simplicial sets is a model category with Quillen model structure with the usual weak equivalences and Kan fibrations. The geometric realization functor carries a Kan fibration to a Serre fibration.

A.1 Mapping spaces

Let X and Y be simplicial sets. The mapping space $\text{Map}(X, Y)$ is the simplicial set whose set of n -simplices is given by $\mathcal{S}(\Delta[n] \times X, Y)$, and the simplicial structure

is induced by the ordinal maps $\Delta[n] \rightarrow \Delta[m]$. We will denote by $\text{Map}_B(X, Y)_f$ the connected component of a map $f: X \rightarrow Y$, in other words the space of maps which are homotopic to f . Let \mathcal{S}_B denote the overcategory whose objects are maps $X \rightarrow B$ and whose morphisms are commutative triangles over B . Let $\text{Map}_B(X, Y)$ denote the mapping space for the overcategory, which is defined to be the simplicial set with n -simplices given by the set $\mathcal{S}_B(\Delta[n] \times X, Y)$ and with the simplicial structure defined similarly.

Let $\xi: E \rightarrow B$ be a Kan fibration of simplicial sets. Then ξ induces a fibration of mapping spaces

$$\xi_*: \text{Map}(B, E) \rightarrow \text{Map}(B, B)$$

whose fiber over the identity map $\text{id}: B \rightarrow B$ is the mapping space $\text{Map}_B(B, E)$. We will also denote the fiber by $\text{Sec}(\xi)$ and call it the space of sections of ξ . There is a pullback diagram

$$\begin{array}{ccc} \text{hSec}(\xi) & \longrightarrow & \text{Map}(B, E) \\ \downarrow & & \downarrow \xi_* \\ \text{Map}(B, B)_{\text{id}} & \longrightarrow & \text{Map}(B, B) \end{array}$$

where $\text{hSec}(\xi)$ is the space of maps $B \rightarrow E$ such that the triangle

$$\begin{array}{ccc} B & \longrightarrow & E \\ \parallel & \searrow \xi & \\ B & & \end{array}$$

commutes up to homotopy.

In Lannes' theory we will consider fibrations over the classifying space of an elementary abelian p -group V . The classifying space BV is a simplicial group with product $BV \times BV \rightarrow BV$ induced by the product on V . The mapping space $\text{Map}(BV, BV)$ is isomorphic to $\text{Hom}(V, V) \times BV$ as a simplicial set [18, Proposition 25.2], and the adjoint $BV \rightarrow \text{Map}(BV, BV)$ of the product map identifies BV with the identity component of the mapping space. For any X the mapping space $\text{Map}(BV, X)$ has an induced action of BV . Also the simplicial monoid $\text{Map}(BV, BV)$ acts by precomposition on the mapping space. The actions are equivariant with respect to the isomorphism $BV \rightarrow \text{Map}(BV, BV)_{\text{id}}$ of simplicial abelian groups [18, Proposition 25.3]. Given a

fibration $\xi: E \rightarrow BV$ in the diagram of fibrations

$$\begin{array}{ccc} BV \times \text{Sec}(\xi) & \longrightarrow & \text{hSec}(\xi) \\ \downarrow & & \downarrow \\ BV & \longrightarrow & \text{Map}(BV, BV)_{\text{id}} \end{array}$$

the horizontal arrows are isomorphisms.

A.2 Homotopy fixed points

Let G be a discrete group and X be a simplicial set with G -action, also called a G -space. The homotopy orbit space X_{hG} is the quotient $(EG \times X)/G$ under the diagonal action. The homotopy fixed point space X^{hG} is the simplicial subset $\text{Map}(EG, X)^G$ of G -equivariant simplicial set maps in $\text{Map}(EG, X)$. Let $f: X \rightarrow Y$ be a map of G -spaces which is also a weak equivalence. Then the induced maps $X_{hG} \rightarrow Y_{hG}$ and $X^{hG} \rightarrow Y^{hG}$ are weak equivalences.

Next we describe a transitivity property of homotopy fixed points [11, Lemma 10.5]. Let H be a normal subgroup of G . There is a natural action of G/H on the mapping space $\text{Map}(EG, X)^H$, and there is a weak equivalence

$$X^{hG} \simeq (\text{Map}(EG, X)^H)^{hG/H},$$

where $\text{Map}(EG, X)^H \simeq X^{hH}$.

Let $\xi: E \rightarrow BG$ be a fibration with fiber F . Consider the pullback diagram

$$\begin{array}{ccc} X_\xi & \longrightarrow & E \\ \downarrow & & \downarrow \xi \\ EG & \longrightarrow & BG \end{array}$$

along the universal principal G -fibration. Since X_ξ is a free G -space, the natural map $h: (X_\xi)_{hG} \rightarrow (X_\xi)/G = E$ is a weak equivalence, in fact a trivial fibration. There is a map of fibrations

$$\begin{array}{ccc} X_\xi & \longrightarrow & F \\ \downarrow & & \downarrow \\ (X_\xi)_{hG} & \xrightarrow{h} & E \\ \downarrow \tilde{\xi} & & \downarrow \xi \\ BG & \xlongequal{\quad} & BG \end{array}$$

where $\tilde{\xi}$ is the composition $\xi \circ h$. We will usually switch from an arbitrary fibration to the natural projection $(X_\xi)_{hG} \rightarrow BG$. As a consequence we have the identifications

$$(X_\xi)^{hG} = \text{Map}(EG, X_\xi)^G = \text{Sec}(\tilde{\xi}) \simeq \text{Sec}(\xi).$$

The equivalence $\text{Sec}(\tilde{\xi}) \simeq \text{Sec}(\xi)$ is a consequence of the fact that the map

$$\text{Map}(BG, Y) \rightarrow \text{Map}(BG, Y')$$

induced by a trivial fibration $Y \rightarrow Y'$ is also a trivial fibration [12]. Note that both spaces are fibrations over $\text{Map}(BG, BG)$. Pulling back along the subspace $\text{Map}(BG, BG)_{\text{id}} \rightarrow \text{Map}(BG, BG)$ induces the required weak equivalence between the spaces of sections.

A.3 hG -equivalence

Let $\xi_i: E_i \rightarrow BG$ for $i = 1, 2$ be two fibrations. A map $E_1 \rightarrow E_2$ of fibrations over BG is called a fiber homotopy equivalence if there is a homotopy inverse over BG . Let X_{ξ_i} denote the associated G -spaces. Then f induces a map $X_{\xi_1} \rightarrow X_{\xi_2}$ which is a G -homotopy equivalence. Conversely one can start with two G -spaces X_1 and X_2 and compare the fibrations associated to their Borel constructions. In this case a weaker notion of equivalence is enough. We say X_1 and X_2 are hG -equivalent if there is a zig-zag of G -maps that are also weak equivalences. Given hG -equivalent G -spaces X_1 and X_2 , the products $EG \times X_1$ and $EG \times X_2$ are G -equivalent. Therefore $(X_1)_{hG}$ is fiber homotopy equivalent to $(X_2)_{hG}$. This implies that there is a one-to-one correspondence between fiber homotopy classes of fibrations over BG and hG -equivalence classes of G -spaces.

A.4 Completion at a prime

Let X_p^\wedge denote the Bousfield–Kan completion of X at a prime p as defined in [6]. It comes with a natural map $X \rightarrow X_p^\wedge$. A space is called p -complete if this map is a weak equivalence. For example, the classifying space BP of a p -group is p -complete. A map $f: X \rightarrow Y$ induces an isomorphism $\tilde{H}_*(f, \mathbb{F}_p)$ if and only if its p -completion $f_p^\wedge: X_p^\wedge \rightarrow Y_p^\wedge$ is a weak equivalence. Note that any weak equivalence between p -completed spaces is a homotopy equivalence since the p -completion of a space is a fibrant simplicial set, ie a Kan complex. Let $\xi: E \rightarrow B$ be a fibration with fiber F . The p -completion $\xi_p^\wedge: E_p^\wedge \rightarrow B_p^\wedge$ is still a fibration. The fiber lemma [6, Chapter II, 5.1] implies that if $\pi_1 B$ is a p -group and F is connected, then the fiber of ξ_p^\wedge is the p -completion of F . There is also a relative version of the completion construction

which applies to a fibration $\xi: E \rightarrow B$, called the fiberwise completion at a prime p . We will denote the fiberwise p -completion by $\xi_{p/B}^\wedge: E_{p/B}^\wedge \rightarrow B$. This is a fibration whose fiber is given by F_p^\wedge . If ξ is a fibration over BP , then up to homotopy ξ_p^\wedge can be identified with $\xi_{p/B}^\wedge$. More explicitly, there is a map of fibrations

$$\begin{array}{ccc} E_{p/BP}^\wedge & \longrightarrow & E_p^\wedge \\ \downarrow & & \downarrow \\ BP & \longrightarrow & BP_p^\wedge \end{array}$$

where the horizontal maps are weak equivalences.

Proposition A.1 *Let $\xi: E \rightarrow BP$ be a fibration with connected fiber and let $X = (X_\xi)_p^\wedge$ denote the p -completion of the P -space associated to ξ . Then there is a fiber homotopy equivalence*

$$\begin{array}{ccc} X_{hP} & \xrightarrow{\sim} & E_p^\wedge \\ \downarrow & & \downarrow \\ BP & \xlongequal{\quad} & BP \end{array}$$

Proof This result is proved in [16, Proposition 4.3.1] for an elementary abelian p -group. For a general p -group we proceed as follows. Let X_ξ denote the pullback of ξ along $EP \rightarrow BP$. It fits into a fibration sequence $F \rightarrow X_\xi \rightarrow EP$. After completion the sequence $F_p^\wedge \rightarrow (X_\xi)_p^\wedge \rightarrow EP_p^\wedge$ is a fiber sequence since EP_p^\wedge is still contractible. Then the diagram

$$\begin{array}{ccc} (X_\xi)_p^\wedge & \longrightarrow & EP_p^\wedge \\ \downarrow & & \downarrow \\ E_p^\wedge & \longrightarrow & BP_p^\wedge \end{array}$$

is a homotopy pullback diagram since $E_p^\wedge \rightarrow BP_p^\wedge$ is a fibration with fiber F_p^\wedge by the fiber lemma. This implies that $X = (X_\xi)_p^\wedge$ is a P -covering of E_p^\wedge . Thus X is a free P -space and the desired map is given by the projection $X_{hP} \rightarrow X/P$. □

A.5 Fiber joins

Next we will discuss the fiber join construction. First we look at the behavior with respect to the Borel construction and p -completion, and then we study the Euler class

of fiber joins. The fiber join of two fibrations $\xi_1: E_1 \rightarrow B$ and $\xi_2: E_2 \rightarrow B$ is defined to be the homotopy pushout of

$$(A.5.1) \quad \begin{array}{ccc} E_1 \times_B E_2 & \longrightarrow & E_1 \\ \downarrow & & \downarrow \\ E_2 & \longrightarrow & E_1 *_B E_2 \end{array}$$

where $E_1 \times_B E_2$ is the pullback of the maps ξ_1 and ξ_2 over B . When B is a point this definition specializes to the join construction, and we simply write $E_1 * E_2$.

Proposition A.2 *Let X and Y be G -spaces. There is a fiber homotopy equivalence*

$$\begin{array}{ccc} X_{hG} *_B Y_{hG} & \xrightarrow{\sim} & (X * Y)_{hG} \\ \downarrow & & \downarrow \\ BG & \xlongequal{\quad} & BG \end{array}$$

Proof The map is induced by the Borel construction of the natural maps

$$X \rightarrow X * Y \leftarrow Y$$

and restricts to the identity map between the fibers. □

Given two spaces X and Y the join $X_p^\wedge * Y_p^\wedge$ is in general not p -complete [21, page 107]. For example, for spheres the p -completion of $(\mathbb{S}^n)_p^\wedge * (\mathbb{S}^m)_p^\wedge$ is given by $(\mathbb{S}^n * \mathbb{S}^m)_p^\wedge$. We use the notation

$$X[m] = \underbrace{(X * \cdots * X)}_m^\wedge_p.$$

A fiberwise version of this definition is as follows: given a fibration $\xi: E \rightarrow B$ we define

$$E_{/B}[m] = \underbrace{(E *_B \cdots *_B E)}_m^\wedge_{p/B}$$

and denote the associated fibration by $\xi[m]: E_{/B}[m] \rightarrow B$.

Corollary A.3 *Let P be a p -group. If ξ is the fibration $E \rightarrow BP$ associated to the Borel construction of a P -space X , then $\xi[m]$ is fiber homotopy equivalent to $(X[m])_{hP}$.*

Proof Using Proposition A.2 and induction we see that there is a fiber homotopy equivalence

$$*_{BP}^n E \rightarrow (*^n X)_{hP}$$

over BP . Let Y denote the pullback of the fiber join $*_{BP}^n E$ along $EP \rightarrow BP$. Then Y is G -equivalent to $*^n X$. Applying Proposition A.1 to $*^n X$ we obtain a fiber homotopy equivalence

$$((^n X)_p^\wedge)_{hP} \rightarrow (*_{BP}^n E)_p^\wedge. \quad \square$$

Remark A.4 The natural map $X[mn] \rightarrow (X[m])[n]$ is a homotopy equivalence. Similarly, $\xi[mn]$ is fiber homotopy equivalent to $(\xi[m])[n]$.

A.6 Euler class

We will study the Euler class of fiber joins of mod- p spherical fibrations. Note that the fiber join construction does not result in a mod- p spherical fibration until we fiberwise complete it at p . But the resulting fibration has a fiber whose mod- p cohomology is the mod- p cohomology of a sphere. Therefore for our purposes we consider a larger class of fibrations. Let $\xi: E \rightarrow B$ be a fibration whose fiber F satisfies $HF \cong H(\mathbb{S}^d)$ for some d . The Thom space $\text{Th}(\xi)$ of ξ is defined to be the cofiber of the map ξ . Consider the diagram of cofibrations

$$\begin{array}{ccc} F & \longrightarrow & E \\ \downarrow & & \downarrow \xi \\ * & \longrightarrow & B \\ \downarrow & & \downarrow \\ \Sigma F & \xrightarrow{\mu} & \text{Th}(\xi) \end{array}$$

The reduced cohomology ring $\tilde{H}(\text{Th}(\xi))$ is a free HB -module generated on a generator $t(\xi)$ of degree $d + 1$ called the Thom class of ξ . We can take $t(\xi)$ to be the dual of the image of a chosen generator under the map $H_{d+1}(\Sigma F) \rightarrow H_{d+1}(\text{Th}(\xi))$ induced by μ . The image of the Thom class under the natural map $H^{d+1}(\text{Th}(\xi)) \rightarrow H^{d+1}(B)$ is called the Euler class $e(\xi)$. Alternatively, $e(\xi)$ is the image of the transgression of the generator of $H^d(F)$ in the Serre spectral sequence of ξ .

We will need an alternative description of fiber join construction. Given fibrations $\xi_i: E_i \rightarrow B$ with fiber F_i , where $i = 1, 2$, let us define a quotient space

$$E_1 \hat{*} E_2 = E_1 \times E_2 \times [0, 1] / \sim$$

by the relations $(e_1, e_2, 0) \sim (e_1, e'_2, 0)$ if $\xi_2(e_2) = \xi_2(e'_2)$ and $(e_1, e_2, 1) \sim (e'_1, e_2, 1)$ if $\xi_1(e_1) = \xi_1(e'_1)$. Projecting onto each factor induces a fibration $\xi_1 \hat{*} \xi_2: E_1 \hat{*} E_2 \rightarrow B \times B$ with fiber $F_1 * F_2$. The pullback along the diagonal inclusion $\Delta: B \rightarrow B \times B$ is exactly the fiber join construction

$$\begin{array}{ccc}
 F_1 * F_2 & \xlongequal{\quad} & F_1 * F_2 \\
 \downarrow & & \downarrow \\
 E_1 *_B E_2 & \longrightarrow & E_1 \hat{*} E_2 \\
 \downarrow & & \downarrow \xi_1 \hat{*} \xi_2 \\
 B & \xrightarrow{\quad \Delta \quad} & B \times B
 \end{array}$$

In more detail, the diagram (A.5.1) maps to the pullback and induces a weak equivalence between the spaces given in the two definitions.

Proposition A.5 *Let ξ_1 and ξ_2 be mod- p spherical fibrations over B . In the commutative diagram*

$$\begin{array}{ccc}
 \Sigma(F_1 * F_2) & \xrightarrow{\sim} & \Sigma F_1 \wedge \Sigma F_2 \\
 \downarrow & & \downarrow \Sigma\mu_1 \wedge \Sigma\mu_2 \\
 \text{Th}(\xi_1 \hat{*} \xi_2) & \xrightarrow{\sim} & \text{Th}(\xi_1) \wedge \text{Th}(\xi_2)
 \end{array}$$

the horizontal maps are weak equivalences.

Proof Let $D(\xi)$ denote the fiber join of the identity map $B \rightarrow B$ (regarded as a fibration) with a mod- p spherical fibration $\xi: E \rightarrow B$. The Thom space $\text{Th}(\xi)$ can be described as the quotient $D(\xi)/B$. We have the identifications

$$\begin{aligned}
 \text{Th}(\xi_1) \wedge \text{Th}(\xi_2) &= \frac{\text{Th}(\xi_1) \times \text{Th}(\xi_2)}{\text{Th}(\xi_1) \vee \text{Th}(\xi_2)} \\
 &= \frac{D(\xi_1) \times D(\xi_2)}{(E \times D(\xi_2)) \cap (D(\xi_1) \times E)} \\
 &\simeq \frac{D(\xi_1 \hat{*} \xi_2)}{\xi_1 \hat{*} \xi_2} \\
 &= T(\xi \hat{*} \xi),
 \end{aligned}$$

which are compatible with the equivalence $\Sigma F_1 \wedge \Sigma F_2 \simeq \Sigma(F_1 * F_2)$. □

An almost immediate consequence of Proposition A.5 is that the Euler class of the fiber join of two mod- p spherical fibrations is the cup product of Euler classes of the individual fibrations.

Corollary A.6 *If ξ_1 and ξ_2 are mod- p spherical fibrations, then*

$$e(\xi_1 *_B \xi_2) = e(\xi_1)e(\xi_2).$$

Proof Proposition A.5 implies that the Thom class of $\xi_1 \hat{*} \xi_2$ is the cross product of the Thom classes of ξ_1 and ξ_2 . Looking at the corresponding diagram of cohomology groups associated to the diagram

$$\begin{array}{ccc} B & \xrightarrow{\Delta} & B \times B \\ \downarrow & & \downarrow \\ \text{Th}(\xi_1 *_B \xi_2) & \longrightarrow & \text{Th}(\xi_1) \wedge \text{Th}(\xi_2) \end{array}$$

we see that the Euler class of $\xi_1 *_B \xi_2$ is the cup product of the Euler classes of ξ_1 and ξ_2 . \square

References

- [1] **A Adem, J Grodal**, *Homotopical group actions and periodicity*, in preparation
- [2] **A Adem, J H Smith**, *Periodic complexes and group actions*, Ann. of Math. 154 (2001) 407–435 MR
- [3] **C Allday, V Puppe**, *Cohomological methods in transformation groups*, Cambridge Studies in Advanced Mathematics 32, Cambridge Univ. Press (1993) MR
- [4] **A H Assadi**, *Some local-global results in finite transformation groups*, Bull. Amer. Math. Soc. 19 (1988) 455–458 MR
- [5] **D J Benson, J F Carlson**, *Complexity and multiple complexes*, Math. Z. 195 (1987) 221–238 MR
- [6] **A K Bousfield, D M Kan**, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics 304, Springer (1972) MR
- [7] **J F Carlson, L Townsley, L Valeri-Elizondo, M Zhang**, *Cohomology rings of finite groups*, Algebra and Applications 3, Kluwer Academic, Dordrecht (2003) MR
- [8] **G Carlsson**, *On the rank of abelian groups acting freely on $(S^n)^k$* , Invent. Math. 69 (1982) 393–400 MR
- [9] **T tom Dieck**, *Transformation groups*, De Gruyter Studies in Mathematics 8, de Gruyter, Berlin (1987) MR
- [10] **W G Dwyer, C W Wilkerson**, *Smith theory and the functor T* , Comment. Math. Helv. 66 (1991) 1–17 MR
- [11] **W G Dwyer, C W Wilkerson**, *Homotopy fixed-point methods for Lie groups and finite loop spaces*, Ann. of Math. 139 (1994) 395–442 MR

- [12] **P G Goerss, J F Jardine**, *Simplicial homotopy theory*, Progress in Mathematics 174, Birkhäuser, Basel (1999) MR
- [13] **J Grodal, J H Smith**, *Algebraic models for finite G -spaces*, Oberwolfach Rep. 4 (2007) 2690–2692 MR
- [14] **I Hambleton, O Ünlü**, *Examples of free actions on products of spheres*, Q. J. Math. 60 (2009) 461–474 MR
- [15] **A Heller, I Reiner**, *Representations of cyclic groups in rings of integers, I*, Ann. of Math. 76 (1962) 73–92 MR
- [16] **J Lannes**, *Sur les espaces fonctionnels dont la source est le classifiant d'un p -groupe abélien élémentaire*, Inst. Hautes Études Sci. Publ. Math. 75 (1992) 135–244 MR
- [17] **J H Long**, *The cohomology rings of the special affine group of \mathbb{F}_p^2 and of $\mathrm{PSL}(3, p)$* , PhD thesis, University of Maryland (2008) MR <https://search.proquest.com/docview/304560975>
- [18] **J P May**, *Simplicial objects in algebraic topology*, Van Nostrand Mathematical Studies 11, Van Nostrand, Princeton, NJ (1967) MR
- [19] **R Oliver**, *Free compact group actions on products of spheres*, from “Algebraic topology” (J L Dupont, I H Madsen, editors), Lecture Notes in Math. 763, Springer (1979) 539–548 MR
- [20] **L Schwartz**, *Unstable modules over the Steenrod algebra and Sullivan’s fixed point set conjecture*, Univ. Chicago Press (1994) MR
- [21] **D P Sullivan**, *Geometric topology: localization, periodicity and Galois symmetry*, K -Monographs in Mathematics 8, Springer (2005) MR
- [22] **R G Swan**, *Periodic resolutions for finite groups*, Ann. of Math. 72 (1960) 267–291 MR
- [23] **O Ünlü**, *Constructions of free group actions on products of spheres*, PhD thesis, University of Wisconsin (2004) MR <https://search.proquest.com/docview/305110769>

*Department of Mathematics, Middlesex College, Western University
London, ON, Canada*

*Department of Mathematics, Bilkent University
Ankara, Turkey*

`okay@math.ubc.ca, yalcine@fen.bilkent.edu.tr`

Received: 30 October 2017 Revised: 1 June 2018

