

AG  
T

*Algebraic & Geometric  
Topology*

Volume 23 (2023)

**$G_\infty$ -ring spectra and Moore spectra for  $\beta$ -rings**

MICHAEL STAHLHAUER





# $G_\infty$ -ring spectra and Moore spectra for $\beta$ -rings

MICHAEL STAHLHAUER

We introduce the notion of  $G_\infty$ -ring spectra. These are globally equivariant homotopy types with a structured multiplication, giving rise to power operations on their equivariant homotopy and cohomology groups. We illustrate this structure by analyzing when a Moore spectrum can be endowed with a  $G_\infty$ -ring structure. Such  $G_\infty$ -structures correspond to power operations on the underlying ring, indexed by the Burnside ring. We exhibit a close relation between these globally equivariant power operations and the structure of a  $\beta$ -ring, thus providing a new perspective on the theory of  $\beta$ -rings.

55P43, 55P91; 18C15, 19A22, 55S91

Introduction	87
1. Power operations on ultracommutative ring spectra	92
2. $G_\infty$ -ring spectra and their properties	98
3. Power operations on Moore spectra and $\beta$ -rings	119
Appendix. Transferring monads under lax functors	142
References	151

## Introduction

Our aim is the introduction of the new notion of  $G_\infty$ -ring spectra. These support power operations on their equivariant homotopy groups and cohomology with coefficients in such spectra. We moreover provide an algebraic description of this notion on Moore spectra, linking  $G_\infty$ -ring structures on a Moore spectrum to  $\beta$ -ring structures on the represented ring.

Algebraic invariants are more useful the more structure they are endowed with. One example of this slogan is power operations on cohomology. The Steenrod operations on mod- $p$  cohomology, the Adams operations on  $K$ -theory and power operations on

complex bordism and stable cohomotopy all carry a lot of additional information and have seen extensive use in classical homotopy theory. More recently, Hill, Hopkins and Ravenel used equivariant power operations in the guise of norm maps to prove the nonexistence of elements of Kervaire invariant one in [22]. This work renewed interest in both multiplicative aspects of homotopy theory and equivariant techniques.

Classical power operations in cohomology arise from an  $H_\infty$ -ring structure on the representing spectrum, as defined by Bruner, May, McClure and Steinberger in [13]. We generalize the notion of an  $H_\infty$ -ring spectrum to a globally equivariant context in order to represent equivariant power operations. Here, globally equivariant means that we encode compatible actions by all compact Lie groups, using the framework provided by Schwede in [35]. Thus, a  $G_\infty$ -ring spectrum encodes power operations on equivariant cohomology groups for all compact Lie groups. Hence, the notion of a  $G_\infty$ -ring spectrum relates to the stricter notion of an ultracommutative ring spectrum as an  $H_\infty$ -ring structure relates to an  $E_\infty$ -ring spectrum. This is visualized in the following diagram, which exhibits forgetful functors between the corresponding homotopy categories:

$$\begin{array}{ccc} \text{ultracommutative ring spectra} & \longrightarrow & G_\infty\text{-ring spectra} \\ \downarrow & & \downarrow \\ E_\infty\text{-ring spectra} & \longrightarrow & H_\infty\text{-ring spectra} \end{array}$$

Algebraically, power operations can be packaged in different ways. The Adams operations on  $K$ -theory endow it with the structure of a  $\lambda$ -ring, and the power operations on stable cohomotopy give it the structure of a  $\beta$ -ring. Among these,  $\lambda$ -rings are better behaved and are widely studied in algebraic topology and representation theory, eg by Hoffman [23], Atiyah and Tall [5], and Knutson [26]. On the other hand, the theory of  $\beta$ -rings is still largely mysterious, with different definitions and many subtleties not present in the study of  $\lambda$ -rings. For an overview on classical literature on  $\beta$ -rings, see Rymer [34], Ochoa [32] and Vallejo [46]. In this paper, we present a different approach to the notion of  $\beta$ -rings, coming from a well-structured theory of global power operations, where the question of scalar extensions of the Burnside ring global power functor naturally leads to considering  $\beta$ -rings. In this way,  $G_\infty$ -ring structures on Moore spectra, which yield scalar extensions of the global power operations on the sphere spectrum, are intimately tied to  $\beta$ -ring structures. Moreover, we can also obtain the  $\beta$ -ring structure on stable cohomotopy  $\pi^0(X)$  in this way. We believe that this new approach to  $\beta$ -rings clarifies the structure of power operations indexed by the Burnside

ring, and underlines that the notion of a global power functor is more fundamental than the notion of  $\beta$ -rings.

**Results** In the first part of this work, we introduce the notion of a  $G_\infty$ -ring spectrum. This is a derived version of a structured ring spectrum. In contrast to, for example, an  $E_\infty$ -ring spectrum, the definition is at the level of the homotopy category. Concretely, we take the free commutative algebra monad  $\mathbb{P} : Sp \rightarrow Sp$  at the level of spectra. This functor is left derivable for the positive global model structure and thus induces a monad on the global homotopy category  $\mathcal{GH}$ .

**Definition 2.3** A  $G_\infty$ -ring spectrum is an algebra over the monad  $\mathbb{G} = L\mathbb{P}$ .

We then study properties of  $G_\infty$ -ring spectra. As mentioned above, the main property of a  $G_\infty$ -ring spectrum is that it supports power operations on its equivariant homotopy groups:

**Proposition 2.13** Let  $E$  be a  $G_\infty$ -ring spectrum. The structure map  $\mathbb{G}E \rightarrow E$  defines the structure of a global power functor on  $\pi_0(E)$ .

To prove this, we construct external power operations  $\pi_0^G(E) \rightarrow \pi_0^{\Sigma_m^G}(\mathbb{G}E)$ , which exist for any global spectrum  $E$ . In the presence of a  $G_\infty$ -structure, the power operations on the homotopy groups are then obtained by postcomposition with the multiplication.

Any ultracommutative ring spectrum induces a  $G_\infty$ -ring structure on its homotopy type. However, there are also examples of  $G_\infty$ -ring spectra which are not induced from any strictly commutative multiplication. These are constructed by means of an adjunction

$$G_\infty\text{-Rings} \begin{array}{c} \xrightarrow{U} \\ \xleftarrow{R} \end{array} H_\infty\text{-Rings},$$

where  $U$  is the forgetful functor from  $G_\infty$ -rings to  $H_\infty$ -rings and  $R$  does not change the underlying  $H_\infty$ -ring spectrum. Applying the right adjoint  $R$  to an  $H_\infty$ -ring spectrum  $E$  which cannot be rigidified to a strict commutative ring spectrum, we see that  $RE$  is not the homotopy type of an ultracommutative ring spectrum.

In the second part of this paper, we analyze the structure of a  $G_\infty$ -multiplication on Moore spectra. We obtain the following result, which characterizes the  $G_\infty$ -ring structures on global Moore spectra via purely algebraic data:

**Theorem 3.14** *The functor*

$$\pi_0 : G_\infty\text{-Moore}^{\text{torsion-free}} \rightarrow \mathcal{GIPow}_{\text{left}}^{\text{torsion-free}}$$

is an equivalence of categories between the homotopy category of  $G_\infty$ -Moore spectra for countable torsion-free rings and the category of countable torsion-free left-induced global power functors.

Here, a left-induced global power functor is a global power functor where the multiplication map  $\mathbb{A} \otimes R(e) \rightarrow R$  is an isomorphism of global Green functors, where  $\mathbb{A}$  denotes the Burnside ring global power functor. The restriction to torsion-free rings is necessary, since the existence of multiplications on Moore spectra in the presence of torsion is already a subtle question. Countability is a mild technical assumption we use in order to construct such Moore spectra. The above theorem shows that whenever multiplications on Moore spectra are tractable, then also the power operations are completely determined by algebraic power operations on the represented ring. This relation can be used to prove that neither the Moore spectra  $\mathbb{S}(\mathbb{Z}/p)$  nor  $\mathbb{S}(\mathbb{Z}[i])$  can be endowed with a  $G_\infty$ -ring structure. This is an analogue of the classical results that these spectra do not support an  $A_\infty$ - or  $E_\infty$ -structure, respectively. However, the  $G_\infty$ -result can be obtained by elementary calculations of the power operations in the Burnside ring.

Moreover, the theory of left-induced global power functors is closely linked to the theory of  $\beta$ -rings. In fact, we prove the following theorem, which allows to induce  $\beta$ -ring structures from global power operations:

**Theorem 3.38** *The assignment  $(G, R) \mapsto R(G)$  extends to a functor*

$$\text{ev} : \text{Rep}^{\text{op}} \times \mathcal{GIPow}_{\mathbb{A}\text{-defl}} \rightarrow \beta\text{-Rings},$$

which sends a conjugacy class of a morphism of compact Lie groups to the corresponding restriction.

Here,  $\text{Rep}$  is the category of compact Lie groups and conjugacy classes of continuous homomorphisms between them, and  $\mathcal{GIPow}_{\mathbb{A}\text{-defl}}$  denotes the category of global power functors equipped with deflation maps  $R(K \times G) \otimes \mathbb{A}(K) \rightarrow R(G)$ . This includes all left-induced global power functors. In particular, this shows that for a Moore spectrum  $\mathbb{S}B$  that supports a  $G_\infty$ -ring structure, all equivariant homotopy groups  $\pi_0^G(\mathbb{S}B) \cong \mathbb{A}(G) \otimes B$  come endowed with the structure of a  $\beta$ -ring. Moreover, this

condition also includes (equivariant) stable cohomotopy  $\pi_G^0(X)$  for a based space  $X$ , hence we reprove the fact that stable cohomotopy can be equipped with the structure of a  $\beta$ -ring from Guillot [20].

The above theorem shows that the notion of a global power functor with  $\mathbb{A}$ -deflations captures the structure of a ring supporting  $\beta$ -ring structures at all compact Lie groups at once. Hence, it provides a unifying point of view to the theory of  $\beta$ -rings, which has proven rather hard to understand.

**Structure** In Section 1, we recall orthogonal spectra as a model for global homotopy theory, and the multiplicative aspects leading to power operations. Moreover, we give a construction of the external power operations on the equivariant homotopy groups of any spectrum.

In Section 2, we define  $G_\infty$ -ring spectra and see that they support power operations on their homotopy groups. We also compare this new notion to classical  $H_\infty$ -ring spectra by means of an adjunction featuring the forgetful functor, and provide a homotopical comparison of the derived symmetric algebra monad  $\mathbb{G} = L\mathbb{P}$  and extended symmetric powers  $\Sigma_+^\infty(E_{\text{gl}}\Sigma_m) \wedge_{\Sigma_m} X^{\wedge m}$ .

In Section 3, we analyze  $G_\infty$ -ring Moore spectra. On the topological side, we prove that power operations on  $\mathbb{A} \otimes B$  are equivalent to a  $G_\infty$ -ring structure on  $\mathbb{S}B$  for torsion-free  $B$ . On the algebraic side, we prove that such power operations give rise to  $\beta$ -ring structures on  $\mathbb{A}(G) \otimes B$  for all compact Lie groups  $G$ . This comparison also yields  $\beta$ -operations on stable cohomotopy groups  $\pi^0(X)$ .

In the appendix, we collect results from the theory of monads used throughout our work. In particular, we study under which 2-functors and double functors monads and monad functors are preserved. We utilize these results when constructing the adjunction between  $G_\infty$ - and  $H_\infty$ -ring spectra.

**Acknowledgements** This paper is a revised version of a master’s thesis written at the University of Bonn. I would like to thank my supervisor Stefan Schwede for suggesting the topic of  $G_\infty$ -ring spectra and  $\beta$ -rings to me, and for his constant support and encouragement.

I would also like to thank Markus Hausmann for helpful remarks, Irakli Patchkoria for answering questions on Moore spectra and Jack Davies for interesting conversations on global power functors. I also thank the anonymous referee for helpful comments, leading to several improvements of the exposition.

The research in this paper was supported by the DFG Schwerpunktprogramm 1786 Homotopy Theory and Algebraic Geometry (GZ SCHW 860/1-1) and the Max Planck Institut für Mathematik Bonn. The author is an associate member of the Hausdorff Center for Mathematics at the University of Bonn (DFG GZ 2047/1, project id 390685813).

## 1 Power operations on ultracommutative ring spectra

In this section we give an introduction to power operations on the global homotopy groups of an ultracommutative ring spectrum, and construct external power operations for any orthogonal spectrum. We work throughout this article in the context of global homotopy theory, where the adjective “global” indicates that we study equivariant spectra for all compact Lie groups at once. As a model, we use the model category of global orthogonal spectra provided by Stefan Schwede in [35], and we use this work as a general reference point for the foundations of global homotopy theory.

The category of orthogonal spectra is endowed with a symmetric monoidal structure, and commutative monoids are called ultracommutative ring spectra. These spectra support power operations on their global homotopy groups. Such power operations are important additional structure, as emphasized most prominently in the work of Hill, Hopkins and Ravenel on the nonexistence of elements of Kervaire invariant one in [22]. We recall the multiplicative aspects of global homotopy theory and the formalism of power operations.

Moreover, we introduce external power operations in Construction 1.13. These are defined on the homotopy groups of any orthogonal spectrum  $X$ , but only take values in the homotopy groups of the symmetric powers  $\mathbb{P}^m X$ . For an ultracommutative ring spectrum, the structure morphism  $\mathbb{P} X \rightarrow X$  then recovers the usual power operations. These external power operations are used in Section 2.1 to define power operations on the homotopy groups of any  $G_\infty$ -ring spectrum.

**Orthogonal spectra and homotopy groups** We assume familiarity with the context of global orthogonal spectra and ultracommutative ring spectra, as established in [35, Chapters 3–5]. We quickly collect the relevant notions.

We denote by  $\mathcal{Sp}$  the category of orthogonal spectra; see [30]. For any compact Lie group  $G$  and integer  $k$ , we can associate to an orthogonal spectrum  $X$  its  $k^{\text{th}}$   $G$ -equivariant homotopy group  $\pi_k^G(X)$  [35, (3.1.11)]. A morphism  $f: X \rightarrow Y$  of orthogonal spectra



is called a *global equivalence* if it induces isomorphisms  $f_*: \pi_k^G(X) \rightarrow \pi_k^G(Y)$  for all compact Lie groups  $G$  and all  $k$ . The resulting homotopy category obtained by inverting global equivalences is called the *global homotopy category*  $\mathcal{GH}$ .

The collection of homotopy groups  $\underline{\pi}_0(X) = \{\pi_0^G(X)\}_G$  for any orthogonal spectrum  $X$  comes equipped with *restriction maps*  $\alpha^*: \pi_0^K(X) \rightarrow \pi_0^G(X)$  for any continuous homomorphism  $\alpha: G \rightarrow K$  of compact Lie groups [35, Construction 3.1.15], and with *transfer maps*  $\text{tr}_H^G: \pi_0^H(X) \rightarrow \pi_0^G(X)$  for any closed subgroup  $H \subset G$  [35, Construction 3.2.22], which are trivial if the Weyl group  $W_G H$  is infinite. These morphisms endow  $\underline{\pi}_0(X)$  with the structure of a *global functor* [35, Definition 4.2.2], and the category of global functors is denoted by  $\mathcal{GF}$ . It is defined as the category of additive functors from the global Burnside category  $\mathbf{A}$  [35, Construction 4.2.1] to abelian groups. Here the Burnside category has as objects the compact Lie groups and the morphisms are generated by restrictions and transfers [35, Proposition 4.2.5]. Composition inside  $\mathbf{A}$  contains the information about the compositions of transfers and restrictions, such as the *double coset formula* [35, Theorem 3.4.9].

Note that the notion of a global functor is a global version of a Mackey functor for a fixed compact Lie group  $G$ . The adjective “global” refers to the fact that we allow restrictions along arbitrary group homomorphisms, not just inclusions. There are various different global versions of Mackey functors, and we refer to [35, Remark 4.2.16] for a discussion of the different definitions.

**Remark 1.1** The global homotopy groups for a fixed compact Lie group  $G$  can be considered as a functor

$$\mathcal{GH} \rightarrow \text{Sets} \quad \text{given by } X \mapsto \pi_0^G(X).$$

This functor is representable by the suspension spectrum of the global classifying space  $B_{\text{gl}}G$  by [35, Theorem 4.4.3(i)]. The orthogonal space  $B_{\text{gl}}G$  is a generalization of the classical space  $BG$ , and we recall its construction now.

For two inner product spaces  $V$  and  $W$ , we denote by  $\mathbf{L}(V, W)$  the space of linear isometric embeddings from  $V$  into  $W$ . Let  $G$  be a compact Lie group. We define, for any  $G$ -representation  $V$ , the orthogonal  $G$ -space  $\mathbf{L}_V = \mathbf{L}(V, \_)$  with the right  $G$ -action by precomposition with the  $G$ -action on  $V$ , and the orthogonal space  $\mathbf{L}_{G,V} = \mathbf{L}(V, \_)/G$ . By [35, Proposition 1.1.26], the global homotopy types of  $\mathbf{L}_V$  and  $\mathbf{L}_{G,V}$  are independent of the choice of the  $G$ -representation  $V$  as long as  $V$  is faithful. We then denote, for any faithful  $V$ , these orthogonal  $G$ -spaces by  $E_{\text{gl}}G = \mathbf{L}_V$  and  $B_{\text{gl}}G = \mathbf{L}_{G,V}$ .

We have a stable tautological class  $e_G \in \pi_0^G(\Sigma_+^\infty B_{\text{gl}}G)$  as defined in [35, (4.1.12)], and the pair  $(\Sigma_+^\infty B_{\text{gl}}G, e_G)$  represents  $\pi_0^G$  in the sense that

$$(1.2) \quad \mathcal{GH}(\Sigma_+^\infty B_{\text{gl}}G, X) \rightarrow \pi_0^G(X) \quad \text{given by } [f] \mapsto f_*(e_G)$$

is a bijection for every orthogonal spectrum  $X$ .

We can moreover describe the suspension spectrum of  $B_{\text{gl}}G$  as follows (see [35, (3.1.2)]). Choose a faithful  $G$ -representation  $V$ , such that  $L_{G,V}$  represents  $B_{\text{gl}}G$ . Then, for any inner product space  $W$ , we define a homeomorphism

$$(1.3) \quad S^W \wedge \mathbf{L}(V, W)_+ \cong \mathbf{O}(V, W) \wedge S^V,$$

called the untwisting isomorphism, using that we can trivialize the orthogonal complement bundle over  $\mathbf{L}(V, W)$  with an additional copy of  $V$ . These homeomorphisms descend to  $G$ -orbits and assemble for varying  $W$  into an isomorphism

$$(1.4) \quad \Sigma_+^\infty B_{\text{gl}}G \rightarrow \mathbf{O}(V, -) \wedge_G S^V =: F_{G,V}S^V.$$

**Global Green and power functors** We now give more details about the multiplicative structure on orthogonal spectra and homotopy groups.

The category of orthogonal spectra has a symmetric monoidal structure using the smash product as defined in [35, Definition 3.5.1]. Using this symmetric monoidal structure, we can define a symmetric algebra monad  $\mathbb{P}X = \bigvee_{m \geq 0} \mathbb{P}^m X$  with  $\mathbb{P}^m X = X^{\wedge m} / \Sigma_m$  for orthogonal spectra  $X$ , and a notion of commutative monoids in this category.

**Definition 1.5** An ultracommutative ring spectrum is a commutative monoid in the category  $\mathcal{Sp}$  of orthogonal spectra. We write  $\text{ucom}$  for the category of ultracommutative ring spectra and multiplicative maps.

Note that the categories of  $\mathbb{P}$ -algebras and of ultracommutative ring spectra are isomorphic.

This strict multiplication induces a rich structure on the homotopy groups of ultracommutative ring spectra. Besides endowing the homotopy groups with the structure of a commutative monoid in the category of global functors, it also induces power operations. We now give a short recollection of how these multiplications and power operations are constructed.

The category of global functors has a symmetric monoidal structure, called the box product, arising as a Day convolution product; compare [35, Construction 4.2.27].

**Definition 1.6** A global Green functor is a commutative monoid in the category  $\mathcal{GF}$  endowed with the symmetric monoidal structure provided by the box product. A morphism of global Green functors is a morphism of global functors compatible with the multiplication.

Explicitly, the multiplication map  $R \square R \rightarrow R$  of a global Green functor is equivalent both to a family of multiplication maps

$$\times: R(G) \times R(K) \rightarrow R(G \times K)$$

for all compact Lie groups  $G$  and  $K$ , and to a family of diagonal products

$$(1.7) \quad \cdot: R(G) \times R(G) \rightarrow R(G)$$

for all compact Lie groups  $G$ . These multiplications have to satisfy the properties explained after [35, Definition 5.1.3]. The relationship between these formulations via restrictions along diagonal and projections is elaborated upon in [35, Remark 4.2.20].

Classically, it is well known that the equivariant homotopy groups of a homotopy commutative  $G$ -ring spectrum support the structure of a Green functor. The same is true here.

**Proposition 1.8** *There is an external multiplication map*

$$(1.9) \quad \boxtimes: \pi_0^G(X) \times \pi_0^K(Y) \rightarrow \pi_0^{G \times K}(X \wedge Y),$$

defined for any two orthogonal spectra  $X$  and  $Y$ . Upon composition with the induced map of the multiplication  $E \wedge E \rightarrow E$  for a homotopy commutative ring spectrum  $E$ , this defines the structure of a global Green functor on  $\pi_0(E)$ .

**Proof** This statement is already contained in the discussion of multiplications on global functors from [35]. The definition of the external multiplication map is given in [35, Construction 4.1.20], and its properties are listed in [35, Theorem 4.1.22]. These, together with the properties of the multiplication on  $E$ , imply that  $\pi_0(E)$  with this multiplication is indeed a global Green functor.  $\square$

**Remark 1.10** To define the power operations induced by an ultracommutative ring spectrum, we recall the wreath product  $\Sigma_m \wr G$  of the symmetric group  $\Sigma_m$  on  $m$  letters with a group  $G$ . We refer to [35, Construction 2.2.3] for details. The wreath product is defined as the semidirect product  $\Sigma_m \wr G = \Sigma_m \ltimes G^m$  with respect to the permutation

action of the symmetric group on the factors of  $G^m$ . This has a natural action on the  $m^{\text{th}}$  power  $A^m$  of a  $G$ -set  $A$ , given by the  $G$ -action on each factor and the permutation action of  $\Sigma_m$  on the factors. In particular, if  $V$  is a  $G$ -representation, then  $V^m$  is a  $\Sigma_m \wr G$ -representation which is faithful if  $V$  is a nonzero faithful  $G$ -representation.

Now we can define the power operations. Let  $E$  be an ultracommutative ring spectrum, then define for  $f: S^V \rightarrow E(V)$  in  $\pi_0^G(E)$ , where  $V$  is some  $G$ -representation, the  $(\Sigma_m \wr G)$ -map

$$(1.11) \quad P^m(f): S^{V^m} \cong (S^V)^{\wedge m} \xrightarrow{f^m} E(V)^{\wedge m} \xrightarrow{\mu_{V, \dots, V}} E(V^m),$$

which represents an element in  $\pi_0^{\Sigma_m \wr G}(E)$ . The map  $\mu_{V, \dots, V}$  is the value of the  $m$ -fold multiplication map  $E^{\wedge m} \rightarrow E$  on the inner product space  $V^m$ . This assignment defines a morphism

$$P^m: \pi_0^G(E) \rightarrow \pi_0^{\Sigma_m \wr G}(E)$$

for all  $m \geq 1$ .

These power operations satisfy equivariant versions of the properties of powers in a ring, and compatibility results with the restriction and transfer maps present on the global homotopy groups. These properties are condensed into the notion of a global power functor, defined in [35, Definition 5.1.6].

**Theorem 1.12** *For an ultracommutative ring spectrum  $E$ , the global homotopy groups  $\pi_0(E)$  together with the operations*

$$P^m: \pi_0^G(E) \rightarrow \pi_0^{\Sigma_m \wr G}(E),$$

*as defined in (1.11), form a global power functor.*

For the proof we refer to [35, Theorem 5.1.11].

**External power operations** We see that a strict multiplication on an orthogonal spectrum gives rise to power operations on the global homotopy groups. We now ask the question whether we need the full structure of an ultracommutative ring spectrum to obtain these power operations. In classical homotopy theory we have the notion of  $H_\infty$ -ring spectra from [13], which is only defined in the stable homotopy category. Such  $H_\infty$ -ring spectra also define power operations on their homotopy and cohomology groups. We define a global analogue. To facilitate the proof that the global homotopy

groups of a  $G_\infty$ -ring spectrum support power operations, we introduce external power operations.

The multiplication on the homotopy groups of a homotopy commutative ring spectrum can be constructed by defining an external multiplication as described in (1.9), which exists for any two orthogonal spectra. Upon composition with the homotopy multiplication, this gives the structure of a global Green functor on the homotopy groups. We now define a similar external power operation for any orthogonal spectrum.

**Construction 1.13** Let  $X$  be an orthogonal spectrum. We set  $\mathbb{P}^m X = X^{\wedge m} / \Sigma_m$  and define maps

$$\hat{P}^m : \pi_0^G(X) \rightarrow \pi_0^{\Sigma_m \wr G}(\mathbb{P}^m X)$$

for every compact Lie group  $G$  and  $m \geq 1$  as follows. For a  $G$ -representation  $V$  and a  $G$ -equivariant map  $f : S^V \rightarrow X(V)$  representing an element in  $\pi_0^G(X)$ , we set

$$\hat{P}^m(f) : S^{V^m} \cong (S^V)^{\wedge m} \xrightarrow{f^{\wedge m}} X(V)^{\wedge m} \xrightarrow{i_{V, \dots, V}} X^{\wedge m}(V^m) \xrightarrow{\text{pr}(V^m)} (\mathbb{P}^m X)(V^m),$$

where the morphism  $i_{V, \dots, V}$  is the iteration of the universal bimorphism from the definition of the smash product (see [35, Definition 3.5.1]), and  $\text{pr} : X^{\wedge m} \rightarrow X^{\wedge m} / \Sigma_m$  is the projection.

We claim that this map is  $\Sigma_m \wr G$ -equivariant. To see this, let

$$(\sigma; g_*) = (\sigma; g_1, \dots, g_m) \in \Sigma_m \wr G$$

be an element of the wreath product. Then we consider the following diagram, where  $\sigma \cdot (\_)$  signifies the  $\Sigma_m$ -action by permutation:

$$\begin{array}{ccccccc} (S^V)^{\wedge m} & \xrightarrow{f^{\wedge m}} & X(V)^{\wedge m} & \xrightarrow{i_{V, \dots, V}} & X^{\wedge m}(V^m) & \longrightarrow & (\mathbb{P}^m X)(V^m) \\ \downarrow g_* & & \downarrow g_* & & \downarrow g_* & & \downarrow g_* \\ (\sigma; g_*) \left( (S^V)^{\wedge m} \xrightarrow{f^{\wedge m}} X(V)^{\wedge m} \xrightarrow{i_{V, \dots, V}} X^{\wedge m}(V^m) \longrightarrow (\mathbb{P}^m X)(V^m) \right) (\sigma; g_*) \\ \downarrow \sigma & & \downarrow \sigma & & \downarrow \sigma & & \downarrow \sigma \\ S^{\sigma \cdot V^m} = \sigma \cdot (S^V)^{\wedge m} & \xrightarrow{f^{\wedge m}} & \sigma \cdot X(V)^{\wedge m} & \xrightarrow{i_{V, \dots, V}} & (\sigma \cdot X^{\wedge m})(\sigma \cdot V^m) & \longrightarrow & (\mathbb{P}^m X)(\sigma \cdot V^m) \end{array}$$

In this diagram, the upper left square commutes by equivariance of  $f$ , the other squares on the top commute as the horizontal map  $i_{V, \dots, V}$  is an  $m$ -morphism of spectra, and  $\text{pr}$  also is a morphism of spectra. The squares on the bottom row commute by the symmetry property of both the direct sum of inner product spaces and the smash product of spectra, and as we exactly quotient out the permutation action on  $X^{\wedge m}$  in the passage

to  $\mathbb{P}^m X$ . Thus this diagram is commutative and proves that the morphism  $\widehat{P}^m(f)$  is  $\Sigma_m \wr G$ -equivariant, hence it defines an element in  $\pi_0^{\Sigma_m \wr G}(\mathbb{P}^m X)$ .

These maps  $\widehat{P}^m$  are called external power operations. The quotient  $X^{\wedge m} \rightarrow \mathbb{P}^m(X)$  is necessary for this definition, since on  $X^{\wedge m}$  we have to consider the  $\Sigma_m$ -action by permuting the factors.

These external operations fit into a commutative diagram

$$(1.14) \quad \begin{array}{ccccc} \pi_0^G(X) & \xrightarrow{(\text{incl}_1)_*} & \pi_0^G(\mathbb{P} X) & \xrightarrow{(\text{pr}_1)_*} & \pi_0^G(X) \\ \widehat{P}^m \downarrow & & P^m \downarrow & & \downarrow \widehat{P}^m \\ \pi_0^{\Sigma_m \wr G}(\mathbb{P}^m X) & \xrightarrow{(\text{incl}_m)_*} & \pi_0^{\Sigma_m \wr G}(\mathbb{P} X) & \xrightarrow{(\text{pr}_m)_*} & \pi_0^{\Sigma_m \wr G}(\mathbb{P}^m X) \end{array}$$

where  $\text{incl}_m: \mathbb{P}^m X \rightarrow \mathbb{P} X$  is the inclusion as the wedge summand indexed by  $m$ , and  $\text{pr}_m$  is the projection onto the wedge summand indexed by  $m$ . Moreover, we use that  $\mathbb{P} X$  as an ultracommutative ring spectrum has power operations on its homotopy groups. This diagram exhibits  $\widehat{P}^m$  as a retract of  $P^m$ .

Note that by definition we obtain the power operation of an ultracommutative ring spectrum  $E$  by composing with the map induced by the multiplication  $\mathbb{P}^m E \rightarrow E$ . However, we also can consider a weaker type of structure which also allows us to internalize these external power operations. This leads to the notion of a  $G_\infty$ -ring spectrum.

## 2 $G_\infty$ -ring spectra and their properties

In this section we give the definition of  $G_\infty$ -ring spectra in Definition 2.3. This notion is a homotopical version of structured ring spectra, with structure morphisms only defined in the global homotopy category. This structured multiplication allows us to construct power operations on the equivariant homotopy groups of a  $G_\infty$ -ring spectrum in Construction 2.12.

The notion of  $G_\infty$ -ring spectra is a global generalization of the nonequivariant notion of an  $H_\infty$ -ring spectrum from [13, Definition I.3.1]. In Section 2.2 we construct an adjunction between  $G_\infty$ - and  $H_\infty$ -ring spectra. The left adjoint is a forgetful functor from the globally equivariant  $G_\infty$ -ring spectrum to the nonequivariant  $H_\infty$ -ring spectrum. The right adjoint exhibits a way to obtain a  $G_\infty$ -ring spectrum from an  $H_\infty$ -ring spectrum, thought of as a “global Borel construction”. This also gives a way

to generate examples of  $G_\infty$ -ring spectra which do not come as the homotopy types of ultracommutative ring spectra; see Remark 2.32. For this, we use the nonequivariant examples of Noel in [31] and Lawson in [27] of  $H_\infty$ -ring spectra which do not rigidify to commutative ring spectra.

In Section 2.3, we compare the derived symmetric powers to a global version of the extended powers  $D_m X = (E\Sigma_m)_+ \wedge_{\Sigma_m} X^{\wedge m}$  in Theorem 2.37. This can be used to give an alternative description of  $G_\infty$ -ring spectra which is closer to the original definition from [13].

### 2.1 Definition of $G_\infty$ -ring spectra and their power operations

Recall that the multiplication maps  $\mathbb{P}^m E \rightarrow E$  of an ultracommutative ring spectrum can be used to define internal power operations on the homotopy groups of  $E$  from the external power operations defined in Construction 1.13. But all that is really needed are such maps on the homotopy groups, hence we define the corresponding structure on the level of the global homotopy category  $\mathcal{GH}$ . To do so, we make use of the positive global model structures on  $Sp$  and  $ucom$ , which are constructed in [35, Proposition 4.3.33 and Theorem 5.4.3].

**Lemma 2.1** *The functors  $Sp \xrightleftharpoons[U]{\mathbb{P}}$  form a Quillen adjoint functor pair with respect to the positive global model structures on both sides.*

**Proof** It is clear that these functors are adjoint to one another. To prove that this is a Quillen adjunction, it suffices to show that the right adjoint  $U$  preserves both fibrations and acyclic fibrations. This is directly evident from the characterization of global equivalences and positive global fibrations of ultracommutative ring spectra by their underlying maps. □

In fact, the model structure on  $ucom$  is transferred from the positive model structure on  $Sp$  along this adjunction.

Now, every Quillen adjunction defines an adjunction on the homotopy categories, see [24, Lemma 1.3.10], hence we get an adjunction

$$\mathcal{GH} \begin{array}{c} \xrightarrow{L_{gl}\mathbb{P}} \\ \xleftarrow{\text{Ho}(U)} \end{array} \text{Ho}(ucom).$$

Note that  $U$  is already homotopical, so it can be derived without a fibrant replacement. From this adjunction, we obtain the monad

(2.2) 
$$\mathbb{G} = \text{Ho}(U) \circ L_{gl}\mathbb{P} : \mathcal{GH} \rightarrow \mathcal{GH}.$$

**Definition 2.3** A  $G_\infty$ -ring spectrum is an algebra over the monad  $\mathbb{G}$ .

**Example 2.4** As we obtain the notion of  $G_\infty$ -ring spectra as algebras over a derived monad  $\mathbb{G} = L_{\text{gl}}\mathbb{P}$ , we see that algebras over the point-set monad  $\mathbb{P}$ , ie ultracommutative ring spectra, also induce a  $G_\infty$ -ring structure on their global homotopy type. This already gives a broad class of examples, which encompasses the sphere spectrum  $\mathbb{S}$ , Eilenberg–Mac Lane spectra  $HR$  for a global power functor  $R$  as constructed in [35, Theorem 5.4.14], and the global versions of Thom and  $K$ -theory spectra from [35, Chapter 6].

These examples that arise as homotopy types of ultracommutative ring spectra, however, do not provide all  $G_\infty$ -ring spectra. In Theorems 2.33 and 2.34 we provide examples of  $G_\infty$ -ring spectra which are not the homotopy type of an ultracommutative ring spectrum such that the  $G_\infty$ -ring structure is induced by the ultracommutative multiplication.

**Remark 2.5** We also note that the definition of  $G_\infty$ -ring spectra is not the same as that of a homotopy commutative ring spectrum in  $\mathcal{GH}$ . This can be seen from the fact that a homotopy commutative ring spectrum does not support power operations on its homotopy groups, whereas Proposition 2.13 proves the existence of equivariant power operations for  $G_\infty$ -ring spectra. For the same reason,  $G_\infty$ -ring spectra posses more structure than  $H_\infty$ -rings internal to the global homotopy category  $\mathcal{GH}$ . The difference lies in the fact that the derived symmetric power  $\mathbb{G}^m X$  can be represented by the global extended power  $\Sigma_+^\infty E_{\text{gl}}\Sigma_m \wedge_{\Sigma_m} X^{\wedge m}$  as shown in Theorem 2.37. In contrast, for an  $H_\infty$ -ring spectrum, the nonequivariant extended powers  $\Sigma_+^\infty E\Sigma_m \wedge_{\Sigma_m} X^{\wedge m}$  would be used.

Our aim is to define power operations on the homotopy groups of a  $G_\infty$ -ring spectrum. Since the definition of  $G_\infty$ -ring spectra is internal to the global homotopy category, we also rephrase the external power operations in terms of the representability of the homotopy groups in  $\mathcal{GH}$ . To do this, we derive the levels of  $\mathbb{P}$  separately with respect to the positive global model structures from [35, Proposition 4.3.33 and Theorem 5.4.3].

**Lemma 2.6** *Let  $f : X \rightarrow Y$  be a global equivalence between positively cofibrant spectra, and let  $m \geq 0$ . Then  $\mathbb{P}^m f : \mathbb{P}^m X \rightarrow \mathbb{P}^m Y$  is a global equivalence in  $Sp$ .*

**Proof** We follow the argument given at the end of the proof of [35, Theorem 5.4.12]. By Lemma 2.1, we know that the functor  $\mathbb{P} : Sp \rightarrow \text{ucom}$  is left Quillen. By Ken Brown’s



lemma [24, Lemma 1.1.12],  $\mathbb{P}f: \mathbb{P}X \rightarrow \mathbb{P}Y$  is a global equivalence of ultracommutative ring spectra, hence by definition a global equivalence of the underlying spectra. But the transformations  $\mathbb{P}^m \xrightarrow{\text{incl}_m} \mathbb{P} = \bigvee_{m \geq 0} \mathbb{P}^m \xrightarrow{\text{pr}_m} \mathbb{P}^m$  exhibit  $\mathbb{P}^m f$  as a retract of  $\mathbb{P}f$ , thus also this morphism is a global equivalence.  $\square$

**Remark 2.7** This lemma is enough to conclude that  $\mathbb{P}^m: Sp \rightarrow Sp$  admits a left derived functor

$$\mathbb{G}^m: \mathcal{GH} \rightarrow \mathcal{GH}.$$

However, one can indeed show more: for any  $m > 0$ , the functor  $\mathbb{P}^m: Sp \rightarrow Sp$  preserves positive cofibrations and acyclic positive cofibrations between positively cofibrant spectra. This uses [19, Theorem 22], that the positive cofibrations are symmetrizable [35, Theorem 5.4.1] and that smashing with positively cofibrant spectra preserves weak equivalences [35, Theorem 4.3.27].

We now calculate the value of the functor  $\mathbb{G}$  on the sphere spectrum and on the representing spectra  $\Sigma_+^\infty B_{\text{gl}}G$  for the global homotopy groups. We use this to define the external power operations intrinsically in  $\mathcal{GH}$ .

**Example 2.8** Since  $\mathbb{S}$  is not positively cofibrant, we need to positively replace it. For this, consider the map from [35, (4.1.28)]:

$$\lambda_{\Sigma_1, \mathbb{R}, 0}: F_{\Sigma_1, \mathbb{R}} S^1 \rightarrow F_{\Sigma_1, 0} = \mathbb{S}.$$

This map is a global equivalence by [35, Theorem 4.1.29], as 0 is a faithful representation of the trivial group  $\Sigma_1$ . Moreover, the spectrum  $F_{\Sigma_1, \mathbb{R}} S^1 = \mathbf{O}(\mathbb{R}, \_) \wedge S^1$  is positively cofibrant, hence  $\lambda_{\Sigma_1, \mathbb{R}, 0}$  can be chosen as a positively cofibrant replacement. Then, we have that  $\mathbb{G}^m \mathbb{S}$  is represented by

$$\mathbb{P}^m(F_{\Sigma_1, \mathbb{R}} S^1) \cong \mathbf{O}(\mathbb{R}^m, \_) \wedge_{\Sigma_m} S^m = F_{\Sigma_m, \mathbb{R}^m} S^m.$$

Hence, by the description of the global classifying spaces via semifree orthogonal spectra in (1.4) and since  $\mathbb{R}^m$  is a faithful  $\Sigma_m$ -representation, we see that

$$\mathbb{G}^m \mathbb{S} \cong \Sigma_+^\infty B_{\text{gl}} \Sigma_m.$$

More generally, we can calculate  $\mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}}G)$  for any compact Lie group, with the previous calculation a special case for  $G = e$ , using the identification  $\Sigma_+^\infty B_{\text{gl}}e \cong \mathbb{S}$ . For this calculation, choose a nonzero faithful  $G$ -representation  $V$ . Then we can write

$$\Sigma_+^\infty B_{\text{gl}}G \cong F_{G, V} S^V$$

as in (1.4). Now, the spectrum  $F_{G,V}S^V$  is positively cofibrant and we calculate

$$\mathbb{P}^m(F_{G,V}S^V) \cong F_{G^m,V^m}S^{V^m} / \Sigma_m \cong F_{\Sigma_m \wr G, V^m}S^{V^m},$$

where for the last identification we used that the permutation action of  $\Sigma_m$  and the action of  $G^m$  on  $V^m$  assemble into the natural action of  $\Sigma_m \wr G$ . Then  $V^m$  is a faithful  $\Sigma_m \wr G$ -representation, hence we see that

$$(2.9) \quad \mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}}G) \cong \Sigma_+^\infty B_{\text{gl}}(\Sigma_m \wr G).$$

**Construction 2.10** We now give another description of the external power operation, using our calculation of  $\mathbb{G}^m$  on the representing spectra for  $\pi_0^G$ .

Let  $f \in \pi_0^G(X)$  be an element of the homotopy groups of a global homotopy type  $X$ . By the representability result (1.2), we can represent  $f$  by a map  $f: \Sigma_+^\infty B_{\text{gl}}G \rightarrow X$  in the global homotopy category. Then, we define

$$\mathbb{G}^m(f): \Sigma_+^\infty B_{\text{gl}}(\Sigma_m \wr G) \cong \mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}}G) \xrightarrow{\mathbb{G}^m f} \mathbb{G}^m X$$

for  $m \geq 1$ , and this morphism represents an element in  $\pi_0^{\Sigma_m \wr G}(\mathbb{G}^m X)$ . Thus we define the external operations as the effect of the functor  $\mathbb{G}^m$  on the homotopy groups  $\pi_0^G(X) \cong \mathcal{GH}(\Sigma_+^\infty B_{\text{gl}}G, X)$ , and obtain maps

$$\mathbb{G}^m: \pi_0^G(X) \rightarrow \pi_0^{\Sigma_m \wr G}(\mathbb{G}^m X).$$

**Lemma 2.11** For a positively cofibrant spectrum  $X$ , the two external operations  $\hat{P}^m$  and  $\mathbb{G}^m: \pi_0^G(X) \rightarrow \pi_0^{\Sigma_m \wr G}(\mathbb{P}^m X)$  agree.

**Proof** Let  $f \in \mathcal{GH}(\Sigma_+^\infty B_{\text{gl}}G, X)$ . Then by (1.2) the corresponding class in  $\pi_0^G(X)$  is  $f_*(e_G)$ . Concretely, let  $f: S^V \wedge (B_{\text{gl}}G)_+(V) \rightarrow X(V)$  for a nonzero faithful  $G$ -representation  $V$ . We can always represent  $f$  on a faithful  $G$ -representation, since we can embed any  $G$ -representation into a faithful one. Then, the tautological class in  $\pi_0^G(\Sigma_+^\infty B_{\text{gl}}G)$  is

$$e_G: S^V \xrightarrow{-\wedge \text{Id}_V} S^V \wedge (B_{\text{gl}}G)_+(V) = S^V \wedge \mathbf{L}(V, V)_+/G,$$

and the tautological class for  $\Sigma_m \wr G$  is

$$e_{\Sigma_m \wr G}: S^{V^m} \xrightarrow{-\wedge \text{Id}_{V^m}} S^{V^m} \wedge \mathbf{L}(V^m, V^m)_+(\Sigma_m \wr G).$$

Note that this element agrees with

$$\hat{P}^m e_G: S^{V^m} \xrightarrow{(-\wedge \text{Id}_V)^m} S^{V^m} \wedge (\mathbf{L}(V, V)_+/G)^m \xrightarrow{\text{pr}} S^{V^m} \wedge \mathbf{L}(V^m, V^m)_+(\Sigma_m \wr G).$$

We now compare  $\widehat{P}^m(f_*(e_G))$  and  $(\mathbb{G}^m f)_*(e_{\Sigma_m \wr G})$ . Since  $X$  is positively cofibrant, we can write  $\mathbb{P}^m$  instead of  $\mathbb{G}^m$ . Then by naturality of the external power operations, we obtain

$$\widehat{P}^m(f_*(e_G)) = \mathbb{P}^m(f)_*(\widehat{P}^m(e_G)) = \mathbb{P}^m(f)_*(e_{\Sigma_m \wr G}).$$

Hence, the two operations agree. □

We now prove that a  $G_\infty$ -ring spectrum structure indeed gives rise to power operations on the homotopy groups.

**Construction 2.12** Let  $E$  be a  $G_\infty$ -ring spectrum. We consider the multiplication map  $\mathbb{G}E \rightarrow E$  in the homotopy category. Since we can derive  $\mathbb{P}$  levelwise, this decomposes into maps  $\zeta_m: \mathbb{G}^m E \rightarrow E$ . Thus, we define the power operations as

$$P^m = \underline{\pi}_0(\zeta_m) \circ \mathbb{G}^m: \pi_0^G(E) \rightarrow \pi_0^{\Sigma_m \wr G}(\mathbb{G}^m E) \rightarrow \pi_0^{\Sigma_m \wr G}(E).$$

**Proposition 2.13** Let  $E$  be a  $G_\infty$ -ring spectrum with structure map  $\zeta: \mathbb{G}E \rightarrow E$ . Then the operations  $P^m$  defined in Construction 2.12, together with the multiplication given by  $\zeta_2$ , define the structure of a global power functor on  $\underline{\pi}_0(E)$ .

**Proof** The map  $\zeta_2: \mathbb{G}^2 E \rightarrow E$  together with the unit map makes  $E$  into an homotopy commutative ring spectrum, hence  $\underline{\pi}_0(E)$  is a global Green functor by Proposition 1.8. Moreover, we need to check the relations for the global power operations as listed in [35, Definition 5.1.6]. For this, we use the same naturality as in the proof of Lemma 2.11 and check on the representing spectra  $\Sigma_+^\infty B_{\text{gl}}G$ . Since the arguments are all similar, we focus on one property here.

Let  $i, j \geq 0$ , and  $x \in \pi_0^G(E)$ . We need to check that  $\Phi_{i,j}^*(P^{i+j}(x)) = P^i(x) \times P^j(x)$  holds in  $\pi_0^{\Sigma_i \wr G \times \Sigma_j \wr G}(E)$ . Here, the map  $\Phi_{i,j}: \Sigma_i \wr G \times \Sigma_j \wr G \rightarrow \Sigma_{i+j} \wr G$  is given by juxtaposition of permutations. Let  $x$  be represented by a map  $f: \Sigma_+^\infty B_{\text{gl}}G \rightarrow E$  in  $\mathcal{GH}$ . Then the power operations on  $x$  are given as  $P^m(x) = \underline{\pi}_0(\zeta_m)(\mathbb{G}^m f)_*(e_{\Sigma_m \wr G})$ .

We now consider the maps  $\varphi_{i,j}: \mathbb{P}^i X \wedge \mathbb{P}^j X \rightarrow \mathbb{P}^{i+j} X$ , given for any  $X$  by forming orbits along  $\Phi_{i,j}$  in  $X^{\wedge i+j} / (\Sigma_i \times \Sigma_j) \rightarrow X^{\wedge i+j} / (\Sigma_{i+j})$ . For  $X = \Sigma_+^\infty B_{\text{gl}}G$ , this map represents the restriction

$$\Phi_{i,j}^*: \pi_0^{\Sigma_{i+j} \wr G} \rightarrow \pi_0^{\Sigma_i \wr G \times \Sigma_j \wr G},$$

meaning that  $(\varphi_{i,j})_*(e_{\Sigma_i \wr G} \times e_{\Sigma_j \wr G}) = \Phi_{i,j}^*(e_{\Sigma_{i+j} \wr G})$ . Moreover,  $\varphi_{i,j}: \mathbb{P}^i \wedge \mathbb{P}^j \rightarrow \mathbb{P}^{i+j}$  features in the monad structure of the functor  $\mathbb{P}$ , hence also in the derived monad

structure for  $\mathbb{G}$ . The fact that  $\zeta$  defines a  $G_\infty$ -structure hence shows that we have the commutative square

$$(2.14) \quad \begin{array}{ccc} \mathbb{G}^i E \wedge^L \mathbb{G}^j E & \xrightarrow{\zeta_i \wedge \zeta_j} & E \wedge^L E \\ \varphi_{i,j} \downarrow & & \downarrow \mu \\ \mathbb{G}^{i+j} E & \xrightarrow{\zeta_{i+j}} & E \end{array}$$

Here,  $\mu$  is the homotopy multiplication induced by  $\zeta_2$ . This square is also used in the original definition of  $H_\infty$ -ring spectra in [13, Definition I.3.1]. In total, we calculate

$$\begin{aligned} \underline{\pi}_0(\zeta_{i+j})(\mathbb{G}^{i+j} f) \Phi_{i,j}^*(e_{\Sigma_{i+j} \wr G}) &= \underline{\pi}_0(\zeta_{i+j})(\mathbb{G}^{i+j} f)(\varphi_{i,j})_*(e_{\Sigma_i \wr G} \times e_{\Sigma_j \wr G}) \\ &= \underline{\pi}_0(\zeta_{i+j})(\varphi_{i,j})_*(\mathbb{G}^i f \wedge \mathbb{G}^j f)_*(e_{\Sigma_i \wr G} \times e_{\Sigma_j \wr G}) \\ &= \underline{\pi}_0(\zeta_i)(\mathbb{G}^i f)_*(e_{\Sigma_i \wr G}) \times \underline{\pi}_0(\zeta_j)(\mathbb{G}^j f)_*(e_{\Sigma_j \wr G}). \end{aligned}$$

This proves one of the relations of a global power functor, the other ones may be proved similarly by considering  $\mathbb{G}^k \mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}} G)$ ,  $\mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}} G \wedge \Sigma_+^\infty B_{\text{gl}} K)$  and  $\mathbb{G}^m(\Sigma_+^\infty B_{\text{gl}} G \vee \Sigma_+^\infty B_{\text{gl}} G)$ . □

**Remark 2.15** As an application of this description of the external power operations, we also define external cohomology operations, and show that a  $G_\infty$ -structure can be used to internalize these operations. These internal cohomology operations are also constructed for an ultracommutative ring spectrum in [35, Remark 5.1.14].

Let  $X$  be an orthogonal spectrum and  $A$  be a cofibrant based  $G$ -space. We define an orthogonal space  $\mathbf{L}_{G,V} A = \mathbf{L}(V, \_) \wedge_G A$  for any  $G$ -representation  $V$ , similar to the construction in Remark 1.1. Then we define the  $G$ -equivariant  $X$ -cohomology of  $A$  as

$$X_G^0(A) = [\Sigma_+^\infty \mathbf{L}_{G,V} A, X],$$

where  $[\_, \_]$  denotes the morphisms in  $\mathcal{GH}$ , and  $V$  is any faithful  $G$ -representation. Then, external power operations on this  $X$ -cohomology are defined by

$$\begin{aligned} \hat{P}^m : X_G^0(A) &= [\Sigma_+^\infty \mathbf{L}_{G,V} A, X] \xrightarrow{\mathbb{G}^m} [\mathbb{G}^m \Sigma_+^\infty \mathbf{L}_{G,V} A, \mathbb{G}^m X] \\ &= [\Sigma_+^\infty \mathbf{L}_{\Sigma_m \wr G, V^m} A^m, \mathbb{G}^m X] = (\mathbb{G}^m X)_{\Sigma_m \wr G}^0(A^m). \end{aligned}$$

Here, we used a relative version of the calculations in Example 2.8 to calculate

$$\mathbb{G}^m \Sigma_+^\infty \mathbf{L}_{G,V} A \cong \Sigma_+^\infty \mathbf{L}_{\Sigma_m \wr G, V^m} A^m.$$

Using a  $G_\infty$ -ring structure on  $X$ , given by morphisms  $\zeta_m: \mathbb{G}^m X \rightarrow X$ , we can internalize these operations to

$$P^m: X_G^0(A) \xrightarrow{\widehat{P}^m} (\mathbb{G}^m X)_{\Sigma_m \wr G}^0(A^m) \xrightarrow{(\zeta_m)_*} X_{\Sigma_m \wr G}^0(A^m).$$

In [35, Remark 5.1.14], it is shown that these power operations forget to the classical power operations  $X^0(A) \rightarrow X^0(B\Sigma_m \times A)$  on the nonequivariant  $X$ -cohomology of  $A$  upon postcomposition with the diagonal on  $A$ . These are the power operations induced by an  $H_\infty$ -structure on  $X$  in [13, Definition I.4.1].

### 2.2 An adjunction between $G_\infty$ - and $H_\infty$ -ring spectra

In this section, we compare the notion of  $G_\infty$ -ring spectra to the classical notion of  $H_\infty$ -ring spectra. This is accomplished by lifting the adjunction

$$\mathcal{GH} \begin{array}{c} \xrightarrow{U} \\ \xleftarrow{R} \end{array} \mathcal{SH}$$

to structured ring spectra, where  $U$  is the forgetful functor and  $R$  its right adjoint. This adjunction is exhibited in [35, Theorem 4.5.1].

**Remark 2.16** In this section we use for the stable homotopy category the positive stable model structure defined by Stolz in [41, Section 1.3]. There, it is called the  $\mathbb{S}$ -model structure. This model structure has the following two desirable properties, which we use in order to analyze the derivability of the symmetric product functor  $\mathbb{P}$  in Lemma 2.19:

(i) The cofibrations and positive acyclic cofibrations are symmetrizable [19, Definition 3]. If  $f: X \rightarrow Y$  is a cofibration, then for all  $n \geq 1$  the iterated pushout product map

$$f^{\square n} / \Sigma_n: Q^n(f) / \Sigma_n \rightarrow \mathbb{P}^n(Y)$$

is a cofibration. Here,  $Q^n(f)$  is the colimit over the punctured cube diagram

$$\begin{aligned} \{0 \rightarrow 1\}^n \setminus \{1, \dots, 1\} &\rightarrow Sp, \\ (i_1, \dots, i_n) &\mapsto Z_{i_1} \wedge \dots \wedge Z_{i_n}, \\ \text{id} \wedge \dots \wedge (0 \rightarrow 1) \wedge \dots \wedge \text{id} &\mapsto \text{id} \wedge \dots \wedge f \wedge \dots \wedge \text{id}. \end{aligned}$$

In this definition, we set

$$Z_i = \begin{cases} X & \text{if } i = 0, \\ Y & \text{if } i = 1. \end{cases}$$

The corresponding property also holds for the positive acyclic cofibrations. That these properties hold for the stable model structure constructed by Stolz follows from observing that the cofibrations agree with those of the global model structure constructed by Schwede in [35, Section 4.3], where symmetrizability is verified in [35, Theorem 5.4.1]. For the acyclic cofibrations, a similar calculation can be carried out.

(ii) Cofibrant objects are flat. If  $X$  is cofibrant in the stable model structure, then  $X \wedge_-$  preserves stable equivalences [41, Proposition 1.3.11].

These properties are required in order to apply the results from [19], in particular Theorem 25, which states that the functor  $\mathbb{P}^n$  preserves weak equivalences between positively cofibrant objects.

**2.2.1 Lifting the forgetful functor  $\mathcal{GH} \rightarrow \mathcal{SH}$  to structured ring spectra** We first recall the classical definition of  $H_\infty$ -ring spectra. As defined in [13, I, Definition 3.1], an  $H_\infty$ -ring spectrum  $X$  is defined by maps

$$\xi_m: D_m X \rightarrow X,$$

where  $D_m X = (E\Sigma_m)_+ \wedge_{\Sigma_m} X^{\wedge m}$ . These maps are required to satisfy compatibility conditions such as the one used in (2.14). Note that this formulation uses the modern smash product, which was not yet available in the original definition. Unravelling the definitions in [13] however gives this formulation. In contrast, our definition of  $G_\infty$ -ring spectra uses a modern point-set category of spectra to obtain the monad  $\mathbb{G}$ , and defines  $G_\infty$ -ring spectra as algebras over this monad. As this definition is more conceptual and allows us to use the results from the appendix, we also formulate the notion of  $H_\infty$ -ring spectra in this way.

For this, note that the adjunction

$$Sp \begin{array}{c} \xrightarrow{\mathbb{P}} \\ \xleftarrow{U_{\text{Com}}} \end{array} \text{Com}$$

is also a Quillen adjunction with respect to the positive stable model structures defined by Stolz in [41, Proposition 1.3.10 and Theorem 1.3.28]. Thus we obtain a derived adjunction

$$\mathcal{SH} \begin{array}{c} \xrightarrow{L_{\text{st}}\mathbb{P}} \\ \xleftarrow{\text{Ho}(U_{\text{Com}})} \end{array} \text{Ho}^{\text{st}}(\text{Com}).$$

**Definition 2.17** An  $H_\infty$ -ring spectrum is an algebra over the monad

$$\mathbb{H} = \text{Ho}(U_{\text{Com}}) \circ L_{\text{st}}\mathbb{P}.$$

By abuse of notation, we will also denote  $\mathbb{H}$  by  $L_{\text{st}}\mathbb{P}$ , since it is the left derived functor of  $\mathbb{P} : Sp \rightarrow Sp$ . In the same way, we denote  $\mathbb{G}$  by  $L_{\text{gl}}\mathbb{P}$ .

That our definition using  $L_{\text{st}}\mathbb{P}$  agrees with the original definition follows from the following statement, after the necessary translations regarding the different models for spectra:

**Lemma 2.18** *Let  $X$  be a positive stably cofibrant orthogonal spectrum. Then the map*

$$p : D_m X = (E\Sigma_m)_+ \wedge_{\Sigma_m} X^{\wedge m} \rightarrow X^{\wedge m} / \Sigma_m = \mathbb{P}^m X$$

*that collapses  $E\Sigma_m$  is a stable weak equivalence.*

**Proof** Since we work in the stable model structure constructed by Stolz, this is [41, Lemma 1.3.17], where a cellular induction along the lines of [13, pages 36–37] is carried out. The analogous statement for the more commonly used projective model structure by Mandell, May, Schwede and Shipley is [30, Lemma 15.5].  $\square$

Using this definition of  $H_\infty$ -ring spectra, we show that the underlying stable homotopy type of a  $G_\infty$ -ring spectrum is an  $H_\infty$ -spectrum. To do this, we show that the derived functor  $U : \mathcal{GH} \rightarrow \mathcal{SH}$  is a monad functor in the sense of Definition A.4. We deduce this formally from a variant of the fact that taking the homotopy category of a model category is a pseudo-2-functor [24, Definition 1.4.2]:

We consider the 2-category (Model, left) of model categories and left Quillen functors. Then, [24, Theorem 1.4.3] shows that taking homotopy categories and left derived functors is a pseudo-2-functor  $L : (\text{Model, left}) \rightarrow \text{Cat}$ . Hence, by Corollary A.6 the functor  $L$  preserves monads and monad morphisms.

However, the functor  $\mathbb{P} : Sp \rightarrow Sp$  is not left Quillen, but merely left derivable, ie it sends weak equivalences between cofibrant objects to weak equivalences, in both the stable and the global positive model structures. Moreover, all compositions  $\mathbb{P}^{\circ k} : Sp \rightarrow Sp$  can be derived:

**Lemma 2.19** *Let  $X$  be a positively cofibrant spectrum in either the stable or global model structure, and  $A = \bigvee_I \mathbb{S}$  be a wedge of sphere spectra. Then  $\mathbb{P}(A \vee X) \cong B \vee Y$ , where  $B = \bigvee_J \mathbb{S}$  is a wedge of spheres which only depends on  $A$ , and where  $Y$  is a positively cofibrant spectrum. Moreover, if  $f : X \rightarrow X'$  is a weak equivalence between positively cofibrant spectra, then  $\mathbb{P}(\text{id} \vee f)$  is also a weak equivalence of the form  $\text{id} \vee g : B \vee Y \rightarrow B \vee Y'$ .*

In particular, for any  $k \geq 1$ , the functor  $\mathbb{P}^{\circ k}: Sp \rightarrow Sp$  sends weak equivalences between positively cofibrant spectra to weak equivalences.

**Proof** We write

$$\mathbb{P}(A \vee X) \cong \mathbb{P}(A) \wedge \mathbb{P}(X) \cong \mathbb{P}(A) \wedge (\mathbb{S} \vee \mathbb{P}_{>0}(X)) \cong \mathbb{P}(A) \vee (\mathbb{P}(A) \wedge \mathbb{P}_{>0}(X)).$$

Now, we see that

$$\mathbb{P}(A) = \mathbb{P}(\bigvee_I \mathbb{S}) \cong \bigwedge_I (\mathbb{P}\mathbb{S}) \cong \bigwedge_I (\bigvee_{i \geq 0} \mathbb{S}) \cong \bigvee_{\mathbb{N}^I} (\bigwedge_I \mathbb{S}) \cong \bigvee_{\mathbb{N}^I} \mathbb{S}$$

is a wedge of spheres. Moreover, the spectrum  $\mathbb{P}_{>0}X$  is positively cofibrant by applying [19, Corollary 10] to the positive model structures, and hence  $\mathbb{P}(A) \wedge \mathbb{P}_{>0}X$  is also positively cofibrant. This proves the first assertion, putting  $B = \mathbb{P}(A)$  and  $Y = \mathbb{P}(A) \wedge \mathbb{P}_{>0}X$ . If  $f: X \rightarrow X'$  is a weak equivalence between positively cofibrant spectra, so are  $\mathbb{P}_{>0}(f)$  and  $\mathbb{P}(A) \wedge \mathbb{P}_{>0}(f)$  by the observations in Remark 2.16. This proves the second part of the lemma, since  $\mathbb{P}(\text{id} \vee f) = \text{id}_{\mathbb{P}(A)} \vee (\mathbb{P}(A) \wedge \mathbb{P}_{>0}(f))$ .

In total, this proves the conclusion that  $\mathbb{P}^{\circ k}$  preserves weak equivalences between positively cofibrant spectra by induction. □

Now we generalize the statement of [24, Theorem 1.4.3] to encompass all left derivable functors. There are two problems: the class of left derivable functors is not closed under composition, and if  $F$  and  $G$  are composable left derivable functors such that  $GF$  also is left derivable, the natural transformation  $LG \circ LF \rightarrow L(GF)$  might not be invertible. However, we obtain:

**Proposition 2.20** *Let  $(\text{Model}, \text{all})$  be the 2–category of model categories and all functors and natural transformations, and let  $\mathcal{L}Der_1$  denote the class of all left derivable functors and  $\mathcal{L}Der_2$  the class of all natural transformations between left derivable functors. Then the assignment*

$$\begin{aligned} L: (\text{Model}, \mathcal{L}Der_1, \mathcal{L}Der_2) &\rightarrow \text{Cat}, \\ \mathcal{C} &\mapsto \text{Ho}(\mathcal{C}), \quad F \mapsto LF, \quad \eta \mapsto L\eta, \end{aligned}$$

comes equipped with the following structure:

- (i) a unitality isomorphism  $\alpha_{\mathcal{C}}: \text{id}_{\text{Ho}(\mathcal{C})} \rightarrow L(\text{id}_{\mathcal{C}})$  for any model category  $\mathcal{C}$ ,
- (ii) a natural transformation  $\mu_{G,F}: LG \circ LF \rightarrow L(GF)$  for any pair of left derivable functors  $F: \mathcal{C} \rightarrow \mathcal{D}$  and  $G: \mathcal{D} \rightarrow \mathcal{E}$  such that  $GF$  is also left derivable.

These satisfy the properties of a lax 2–functor from Definition A.2 where defined.



Moreover, if  $F: \mathcal{C} \rightarrow \mathcal{D}$  is left derivable and  $U: \mathcal{D} \rightarrow \mathcal{E}$  is homotopical, then  $UF$  is left derivable and  $\mu_{U,F}$  is invertible.

**Proof** The proof is the same as that of [24, Theorem 1.4.3], where we weaken the requirement of being left Quillen to sending weak equivalences between cofibrant objects to weak equivalences. □

We consider the commutative diagram

$$(2.21) \quad \begin{array}{ccc} Sp^{gl} & \xrightarrow{U} & Sp^{st} \\ \mathbb{P}_{gl} \downarrow & & \downarrow \mathbb{P}_{st} \\ Sp^{gl} & \xrightarrow{U} & Sp^{st} \end{array}$$

which exhibits the functor  $U$  as a monad functor between  $(\mathbb{P}_{gl}, \mu_{gl}, \eta_{gl})$  and  $(\mathbb{P}_{st}, \mu_{st}, \eta_{st})$ . Also, since  $U$  is homotopical, it guarantees that all composites  $\mathbb{P}^{oi} \circ U \circ \mathbb{P}^{oj} = U \circ \mathbb{P}^{oi+j}$  are left derivable. Proposition 2.20 allows us to conclude that taking homotopy categories and left derived functors preserves the monads  $\mathbb{P}$  on  $Sp^{gl}$  and  $Sp^{st}$  as well as the functor  $U$  between them.

**Proposition 2.22** *The left derived functors  $L\mathbb{P}_{st}$  and  $L\mathbb{P}_{gl}$  have the structure of monads via the natural transformations*

$$L\mu_{gl} \circ \mu_{\mathbb{P}_{gl}, \mathbb{P}_{gl}} \quad \text{and} \quad L\eta_{gl} \circ \alpha_{Sp^{gl}},$$

and the analogous transformations for  $L\mathbb{P}_{st}$ .

Moreover, the derived functor  $\text{Ho}(U): \text{Ho}(Sp^{gl}) \rightarrow \text{Ho}(Sp^{st})$  has the structure of a monad functor between  $L\mathbb{P}_{gl}$  and  $L\mathbb{P}_{st}$  via the transformation  $\mu_{U, \mathbb{P}_{gl}}^{-1} \circ \mu_{\mathbb{P}_{st}, U}$ .

**Proof** This is the statement of Corollary A.6. To apply this corollary as stated, we would need to have a lax 2-functor  $L: \text{Model} \rightarrow \text{Cat}$  encompassing all left derivable functors. However, it suffices that we have the required structure morphisms for all composites  $\mathbb{P}_{st}^{oi} \circ U \circ \mathbb{P}_{gl}^{oj}$  with  $i + j \leq 3$ . This is the case by Lemma 2.19 and the commutative square (2.21). □

Since monad functors lift to functors on the categories of algebras, we have proven:

**Proposition 2.23** *The functor  $U$  lifts to a functor from the category of  $G_\infty$ -ring spectra to the category of  $H_\infty$ -ring spectra.*

Explicitly, let  $X$  be a  $G_\infty$ -ring spectrum with structure map  $h: L\mathbb{P}_{\text{gl}}X \rightarrow X$ . Then the map  $h \circ \mu_{U, \mathbb{P}_{\text{gl}}}^{-1} \circ \mu_{\mathbb{P}_{\text{st}}, U}: (L_{\text{st}}\mathbb{P})UX \rightarrow UX$  defines an  $H_\infty$ -ring structure on the stable homotopy type  $UX$ . Moreover, for a  $G_\infty$ -ring morphism  $f: X \rightarrow Y$  in  $\mathcal{GH}$  between two  $G_\infty$ -ring spectra, the map  $U(f) \in \mathcal{SH}(UX, UY)$  is an  $H_\infty$ -ring map.

Using this result, we get the commutative diagram

$$(2.24) \quad \begin{array}{ccc} \text{Ho}^{\text{gl}}(\text{ucom}) & \longrightarrow & (G_\infty\text{-ring spectra}) \\ \downarrow & & \downarrow \\ \text{Ho}^{\text{st}}(\text{Com}) & \longrightarrow & (H_\infty\text{-ring spectra}) \end{array}$$

of homotopy categories of structured ring spectra, where all functors are forgetful ones.

**2.2.2 Lifting the right adjoint  $\mathcal{SH} \rightarrow \mathcal{GH}$  to structured ring spectra** In this section, we study whether the forgetful functor  $U$  from the category of  $G_\infty$ -ring spectra to  $H_\infty$ -ring spectra from (2.24) has adjoints. The corresponding question for the homotopy categories  $\mathcal{SH}$  and  $\mathcal{GH}$  is investigated in [35, Section 4.5], and we use these results to obtain a right adjoint to the forgetful functor. This gives us a way to define  $G_\infty$ -ring spectra from  $H_\infty$ -ring spectra, and we use this to give examples of  $G_\infty$ -ring spectra which do not come from ultracommutative ring spectra.

We first recall the right adjoint  $R: \mathcal{SH} \rightarrow \mathcal{GH}$  to the forgetful functor from [35, Construction 4.5.21].

**Construction 2.25** We define the functor

$$b: Sp \rightarrow Sp$$

as a ‘‘global Borel construction’’: for an orthogonal spectrum  $X$  and an inner product space  $V$ , we set

$$(bX)(V) = \text{map}(\mathbf{L}(V, \mathbb{R}^\infty), X(V)),$$

with structure morphisms defined as in [35, Construction 4.5.21].

We also define a natural transformation  $i: \text{Id} \rightarrow b$  via the map

$$i_X(V): X(V) \rightarrow \text{map}(\mathbf{L}(V, \mathbb{R}^\infty), X(V)) \quad \text{given by } x \mapsto \text{const}_x.$$

The morphism  $i_X(V)$  is a nonequivariant homotopy equivalence for all inner product spaces  $V$ , since the space  $\mathbf{L}(V, \mathbb{R}^\infty)$  is contractible. Hence the induced morphism  $i_X: X \rightarrow bX$  is invertible in the stable homotopy category.

Moreover,  $b$  comes equipped with a lax symmetric monoidal structure for the smash product of orthogonal spectra, so that  $i: \text{Id} \rightarrow b$  is a monoidal transformation. Thus:

**Corollary 2.26** *The functor  $b : Sp \rightarrow Sp$  defines a monad endofunctor in the sense of Definition A.4 of the symmetric algebra monad  $\mathbb{P}$  on  $Sp$ .*

*Moreover, the transformation  $i : \text{Id} \rightarrow b$  is a monadic transformation.*

By [35, Proposition 4.5.22], the functor  $b$  represents the right adjoint to the forgetful functor  $U : \mathcal{GH} \rightarrow \mathcal{SH}$  on stable  $\Omega$ -spectra. We modify this statement to hold on positive  $\Omega$ -spectra, since we need positive model structures for the study of commutative ring spectra. Recall that for obtaining the stable homotopy category, we use the stable  $\mathbb{S}$ -model structure constructed by Stolz in [41, Proposition 1.3.10], in order to achieve derivability of the symmetric powers in Lemma 2.19. Note that this model structure has fewer fibrant objects than the projective positive stable model structure from [30, Theorem 14.2], so all fibrant objects are in particular positive  $\Omega$ -spectra.

**Proposition 2.27** *Let  $X$  be a positive orthogonal  $\Omega$ -spectrum. Then:*

- (i)  *$bX$  is a positive global  $\Omega$ -spectrum whose homotopy type lies in the image of the right adjoint  $R$ .*
- (ii) *For every orthogonal spectrum  $A$ , the two homomorphisms*

$$\mathcal{GH}(A, bX) \xrightarrow{U} \mathcal{SH}(A, bX) \xrightarrow{(i_X)_*^{-1}} \mathcal{SH}(A, X)$$

*are isomorphisms. In particular, the counit of the adjunction between  $U$  and  $R$  is given by  $i_X^{-1} : bX \rightarrow X \in \mathcal{SH}(bX, X)$ .*

**Proof** The proof is completely analogous to the proof of [35, Proposition 4.5.22]. The cited proof that  $bX$  is a global  $\Omega$ -spectrum works levelwise, hence if  $X$  is only a positive  $\Omega$ -spectrum,  $bX$  is a positive global  $\Omega$ -spectrum. That  $bX$  is right induced from the stable homotopy category can be seen by replacing  $bX$  by the globally equivalent  $\Omega \text{ sh}(bX)$  and using that the shift of a positive  $\Omega$ -spectrum is a  $\Omega$ -spectrum.

Moreover, (ii) is a formal consequence of (i), as indicated in the proof of [35, Proposition 4.5.22]. □

We also check that  $b$  is right derivable:

**Lemma 2.28** *Let  $f : X \rightarrow Y$  be a stable equivalence between (positive)  $\Omega$ -spectra. Then  $b(f)$  is a global equivalence.*

**Proof** Since  $f$  is a stable equivalence between (positive)  $\Omega$ -spectra, it is a (positive) level equivalence. Since the  $G$ -space  $\mathbf{L}(V, \mathbb{R}^\infty)$  is  $G$ -cofibrant and free for any faithful  $G$ -representation  $V$ , mapping out of it takes weak equivalences to  $G$ -weak

equivalences. Thus,  $b$  takes (positive) level equivalences to (positive) level equivalences and thus to global equivalences. Hence,  $b(f)$  is a global equivalence.  $\square$

Using this, we describe the unit of the adjunction  $\mathcal{GH} \xrightleftharpoons[R]{U} \mathcal{SH}$  in a similar way to the results for the counit in Proposition 2.27(ii).

**Lemma 2.29** *Let  $X$  and  $Y$  be (positive)  $\Omega$ -spectra, and assume that  $X$  is (positively) cofibrant. Then, the composition*

$$\mathcal{SH}(X, Y) \xrightarrow{b} \mathcal{GH}(bX, bY) \xrightarrow{(i_X)^*} \mathcal{GH}(X, bY)$$

is a bijection inverse to

$$\mathcal{GH}(X, bY) \xrightarrow{U} \mathcal{SH}(X, bY) \xrightarrow{(i_Y)_*^{-1}} \mathcal{SH}(X, Y).$$

**Proof** We consider the diagram

$$\begin{array}{ccc} \mathcal{GH}(X, bY) & \xleftarrow{(i_X)^*} & \mathcal{GH}(bX, bY) \\ U \downarrow & & \uparrow b \\ \mathcal{SH}(X, bY) & \xleftarrow{(i_Y)_*} & \mathcal{SH}(X, Y) \end{array}$$

As both  $(i_Y)_*^{-1}$  and  $U$  are bijective, it suffices to show that this diagram is commutative, ie that  $(i_Y)_* = U \circ (i_X)^* \circ b$ . This is a consequence of the fact that  $i : \text{Id} \rightarrow b$  is natural.  $\square$

Now, we set up the double categorical context we use to prove that the above adjunction lifts to  $G_\infty$ - and  $H_\infty$ -ring spectra.

We first consider the double category **Model** of model categories, left Quillen functors as vertical morphisms, right Quillen functors as horizontal morphisms and all natural transformations as 2-cells. Then [39, Theorem 7.6] shows that taking the homotopy category and derived functors defines a double pseudofunctor into the double category **Sq**(Cat) of categories, functors as horizontal and vertical morphisms and natural transformations as 2-cells.

In our context, however, neither the symmetric algebra monads  $\mathbb{P}_{\text{gl}}$  and  $\mathbb{P}_{\text{st}}$  nor  $b$  are Quillen functors, but merely derivable. Hence, as in Proposition 2.20, we restrict to the classes of left and right derivable functors, respectively, and obtain:

**Proposition 2.30** *Let (Model, all, all) denote the double category of model categories, all functors as horizontal and vertical morphisms and natural transformations as 2-cells. Let  $\mathcal{LDer}$  denote the class of left derivable functors,  $\mathcal{RDer}$  denote the class of right*

derivable functors and  $Der_2$  denote the class of natural transformations of the form  $FG \rightarrow KH$  with  $G$  and  $K$  right derivable and  $F$  and  $H$  left derivable. Then the assignment

$$\begin{aligned} \text{Ho} : (\mathbf{Model}, \mathcal{L}Der, \mathcal{R}Der, Der_2) &\rightarrow \mathbf{Sq}(\text{Cat}), \\ \mathcal{C} \mapsto \text{Ho}(\mathcal{C}), \quad F \mapsto LF, \quad G \mapsto RG, \quad \eta \mapsto \text{Ho}(\eta), \end{aligned}$$

comes equipped with the following structure:

- (i) unitality isomorphisms  $\alpha_C^v : \text{id}_{\text{Ho}(\mathcal{C})} \rightarrow L(\text{id}_C)$  and  $\alpha_C^h : R(\text{id}_C) \rightarrow \text{id}_{\text{Ho}(\mathcal{C})}$  for any model category  $\mathcal{C}$ ,
- (ii) a natural transformation  $\mu_{G,F}^v : LG \circ LF \rightarrow L(GF)$  for any pair of left derivable functors  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{E}$  such that  $GF$  is also left derivable,
- (iii) a natural transformation  $\mu_{G,F}^h : R(GF) \rightarrow RG \circ RF$  for any pair of right derivable functors  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{E}$  such that  $GF$  is also right derivable.

These satisfy the properties of a lax-oplax double functor from Definition A.7 where defined.

Moreover, if  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{E}$  are right derivable and either  $F$  is right Quillen or  $G$  is homotopical, then  $GF$  is right derivable and  $\mu_{G,F}^h$  is invertible.

**Proof** The proof is the same as for [39, Theorem 7.6], where we weaken the requirements from being Quillen to being derivable. The last statement about invertibility of  $\mu_{G,F}^h$  follows from the description of this transformation as  $(GF)P \xrightarrow{G_{PFP}} GPF P$ , where  $p : \text{id} \rightarrow P$  denotes a functorial fibrant replacement. If  $G$  is homotopical, it sends the weak equivalence  $p_{FP}$  to a weak equivalence. If  $F$  is right Quillen,  $FP$  is fibrant and thus  $p_{FP}$  is a weak equivalence between fibrant objects. Since  $G$  is right derivable, it then sends  $p_{FP}$  to a weak equivalence. □

Now we have all the ingredients to prove that we have an adjunction between  $G_\infty$ -ring spectra and  $H_\infty$ -ring spectra.

**Theorem 2.31** *Let  $R : \mathcal{SH} \rightarrow \mathcal{GH}$  denote the right adjoint to  $U : \mathcal{GH} \rightarrow \mathcal{SH}$ .*

- (i) *The functor  $R : \mathcal{SH} \rightarrow \mathcal{GH}$  induces a functor  $\widehat{R}$  from the category of  $H_\infty$ -ring spectra to the category of  $G_\infty$ -ring spectra.*
- (ii) *The functor  $\widehat{R}$  is right adjoint to the forgetful functor  $U$ , with adjunction unit lifted from  $I : \text{id}_{\mathcal{GH}} \rightarrow RU$  and adjunction counit lifted from  $J^{-1} : UR \rightarrow \text{id}_{\mathcal{SH}}$ , where both  $I$  and  $J$  are obtained from deriving  $i : \text{id} \rightarrow b$ .*

**Proof** (i) We have seen that  $\mathbb{P}_{\text{gl}}^{\circ i}$  and  $\mathbb{P}_{\text{st}}^{\circ j}$  are left derivable for all  $i, j \geq 0$  and that  $b$  is right derivable, and moreover that  $b$  has the structure of a monad morphism between  $\mathbb{P}_{\text{st}}$  and  $\mathbb{P}_{\text{gl}}$  by Corollary 2.26. Hence Proposition 2.30 suffices to invoke Proposition A.11 to conclude that the right derived functor  $R = Rb: \mathcal{SH} \rightarrow \mathcal{GH}$  is a monad functor between  $L_{\text{st}}\mathbb{P}$  and  $L_{\text{gl}}\mathbb{P}$ . Hence, it lifts to a functor  $\hat{R}: (H_{\infty}\text{-Rings}) \rightarrow (G_{\infty}\text{-Rings})$ .

(ii) We know that both  $R = R(b)$  and  $U = R(u)$  for the forgetful functor  $u: Sp_{\text{gl}} \rightarrow Sp_{\text{st}}$  are monad functors. Hence both compositions  $RU$  and  $UR$  are monad functors.

We note that both compositions  $ub$  and  $bu$  are right derivable, since  $u$  is homotopical and sends global  $\Omega$ -spectra to nonequivariant  $\Omega$ -spectra, on which  $b$  is homotopical by Lemma 2.28. Moreover, they are monad functors as composites of monad functors, and hence so are the derived functors  $R(ub)$  and  $R(bu)$ . We start by constructing  $I$ .

We define

$$I = \mu_{b,u}^h \circ \text{Ho}(i) \circ (\alpha_{Sp_{\text{gl}}}^h)^{-1} : \text{id}_{\mathcal{GH}} \rightarrow R(\text{id}_{Sp_{\text{gl}}}) \rightarrow R(bu) \rightarrow Rb \circ Ru = RU.$$

Since  $i$  is a monadic transformation by Corollary 2.26 and any lax-oplax double functor preserves these by Proposition A.11, we see that  $\text{Ho}(i)$  is monadic. Moreover, by Lemma A.12, both  $\mu_{b,u}^h$  and  $\alpha_{Sp_{\text{gl}}}^h$  are monadic, and thus so is  $(\alpha_{Sp_{\text{gl}}}^h)^{-1}$ . In total,  $I$  is a monadic transformation as a composition of such.

Analogously, we define  $J = \mu_{u,b}^h \circ \text{Ho}(i) \circ (\alpha_{Sp_{\text{st}}}^h)^{-1}$ . This also is a monadic transformation by the same arguments. Moreover,  $J$  is invertible, since  $\mu_{u,b}^h$  is by Proposition 2.30,  $(\alpha_{Sp_{\text{st}}}^h)^{-1}$  is by definition and  $\text{Ho}(i)$  is invertible in the stable homotopy category since  $i$  is a stable equivalence. Thus  $J^{-1}: UR \rightarrow \text{id}_{\mathcal{SH}}$  is also a monadic transformation.

Hence, the two transformations  $I: \text{id}_{\mathcal{GH}} \rightarrow RU$  and  $J^{-1}: UR \rightarrow \text{id}_{\mathcal{SH}}$  lift to the categories of algebras. Moreover, Lemma 2.29 shows that for any homotopy types  $X \in \mathcal{GH}$  and  $Y \in \mathcal{SH}$ , the morphisms

$$\mathcal{SH}(UX, Y) \xrightarrow{R} \mathcal{GH}(RUX, RY) \xrightarrow{I^*} \mathcal{GH}(X, RY)$$

and

$$\mathcal{GH}(X, RY) \xrightarrow{U} \mathcal{SH}(UX, URY) \xrightarrow{J_*^{-1}} \mathcal{SH}(UX, Y)$$

are inverse isomorphisms, and thus  $I$  and  $J^{-1}$  are unit and counit of an adjunction. As the forgetful functor from the categories of algebras to the base category is faithful, this property lifts to prove that the lifts of  $I$  and  $J^{-1}$  are unit and counit of an adjunction

$$G_{\infty}\text{-Rings} \begin{array}{c} \xrightarrow{U} \\ \xleftarrow{\hat{R}} \end{array} H_{\infty}\text{-Rings}. \quad \square$$

As an application of this result, we use the right adjoint to give examples of  $G_\infty$ -ring spectra which are not obtained as the homotopy type of an ultracommutative ring spectrum.

**Remark 2.32** Let  $X \in \mathcal{SH}$  be an  $H_\infty$ -ring spectrum. Then we consider the induced  $G_\infty$ -ring structure on the global homotopy type  $RX$ . This induces an  $H_\infty$ -ring structure on  $U(RX)$ . The map  $J_X: X \rightarrow U(RX)$ , defined in the proof of Theorem 2.31, is a stable equivalence and, by monadicity of the transformation, an isomorphism of  $H_\infty$ -ring spectra.

Assume now that there is an ultracommutative ring spectrum  $Y$  such that the homotopy type of  $Y$  is  $RX$ , and such that the  $G_\infty$ -ring structure is induced from the structure map  $\mathbb{P}Y \rightarrow Y$  of  $Y$ . Then the  $H_\infty$ -ring structure on  $U(RX)$  is induced by the commutative multiplication on  $UY$ . But the  $H_\infty$ -ring spectrum  $U(RX)$  is equivalent to  $X$ . Hence, if we find an ultracommutative representative  $Y$  for the  $G_\infty$ -ring spectrum  $RX$ , then  $UY$  is a commutative ring spectrum which induces the  $H_\infty$ -ring structure on  $X$ .

Thus, in order to provide examples of  $G_\infty$ -ring spectra that are not induced by ultracommutative ring spectra, it is enough to consider this question nonequivariantly, where counterexamples are already exhibited in [31] and [27].

**Theorem 2.33** [31, Theorem 1.2] *Let  $s_k \in H^{2k}(BU; \mathbb{Z}_{(2)})$  be a primitive generator. Define the space  $KL_k$  as the homotopy fibre*

$$KL_k \xrightarrow{i_k} BU_{(2)} \xrightarrow{4s_k} K(\mathbb{Z}_{(2)}, 2k)$$

and consider the suspension spectra

$$\Sigma_+^\infty KL_k \xrightarrow{\Sigma_+^\infty i_k} \Sigma_+^\infty BU_{(2)} \xrightarrow{\Sigma_+^\infty 4s_k} \Sigma_+^\infty K(\mathbb{Z}_{(2)}, 2k).$$

Then, for any  $k$ , the spectrum  $\Sigma_+^\infty KL_k$  admits the structure of an  $H_\infty$ -ring spectrum, and  $\Sigma_+^\infty i_k$  is an  $H_\infty$ -ring map. Moreover, for  $k = 15$ , the  $H_\infty$ -ring structure of  $\Sigma_+^\infty KL_{15}$  is not induced by an  $E_\infty$ -ring spectrum.

As the homotopy category of  $E_\infty$ -ring spectra is equivalent to the homotopy category of commutative ring spectra, for example by [17, Sections II.3 and II.4], this indeed gives rise to an example of a  $G_\infty$ -ring spectrum whose structure is not induced from an ultracommutative ring spectrum.

**Theorem 2.34** [27, Theorem 1] *Let  $R_k$  be a wedge of Eilenberg–Mac Lane spectra such that  $\pi_* R_k$  is isomorphic to the graded ring  $\mathbb{F}_2[x]/(x^3)$ , where  $|x| = -2^k$ . Then  $R_k$  has an  $H_\infty$ –ring structure, and for  $k > 3$  these structures are not induced from commutative ring spectra.*

For more details on these examples, the reader is referred to the cited articles.

### 2.3 Homotopical analysis of the extended powers

In this section we generalize the analysis of the symmetric powers  $\mathbb{P}^m X$  classically given by Lemma 2.18, comparing them to the extended powers  $(E\Sigma_m)_+ \wedge_{\Sigma_m} X^{\wedge m}$ , to the global context. Thus, we connect our definition of  $G_\infty$ –ring spectra using the derived monad  $\mathbb{G}$  to the original definition of  $H_\infty$ –ring spectra using the extended powers. As in the  $G$ –equivariant version of Lemma 2.18 given in [22, Proposition B.117], we need to replace the universal space  $E\Sigma_m$  with an appropriate global object. The correct analogue is the global universal space  $E_{\text{gl}}\Sigma_m$  defined in Remark 1.1.

Recall that the global universal space  $E_{\text{gl}}\Sigma_m$  is constructed as  $\mathbf{L}_V$ , where  $V$  is a faithful  $\Sigma_m$ –representation. Then its suspension spectrum can be described by the untwisting isomorphism

$$(2.35) \quad \Sigma_+^\infty \mathbf{L}_V \rightarrow \mathbf{O}(V, -) \wedge S^V = F_V S^V,$$

as defined in (1.3). This isomorphism  $\Sigma_+^\infty \mathbf{L}_V \rightarrow F_V S^V$  induces the isomorphism  $\Sigma_+^\infty \mathbf{L}_{G,V} \cong F_{G,V} S^V$  from (1.4) on  $G$ –orbits.

We also consider the morphism

$$(2.36) \quad \lambda_{G,V,W} : F_{G,V \oplus W} S^V \rightarrow F_{G,W}$$

defined in [35, (4.1.28)] for any compact Lie group  $G$  and  $G$ –representations  $V$  and  $W$  with  $W$  faithful. At an inner product space  $U$ , the map  $\lambda_{G,V,W}$  is represented as

$$\begin{aligned} \mathbf{O}(V \oplus W, U) \wedge_G S^V &\rightarrow \mathbf{O}(W, U)/G, \\ [(u, \varphi), t] &\mapsto [u + \varphi(t), \varphi \circ i_2], \end{aligned}$$

where  $i_2 : W \rightarrow V \oplus W$  is the inclusion as the second factor. The map  $\lambda_{G,V,W}$  is a global equivalence by [35, Theorem 4.1.29].

**Theorem 2.37** *Let  $X$  be a positively cofibrant orthogonal spectrum, and  $n \geq 1$ . Then the map*

$$q = q_n^X : \Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n} \wedge_{\Sigma_n} X^{\wedge n} \rightarrow X^{\wedge n} / \Sigma_n = \mathbb{P}^n X$$

*that collapses  $\Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n}$  to  $\mathbb{S} = \Sigma_+^\infty *$  is a global equivalence.*



**Proof** For the proof, we use a  $\Sigma_n$ -equivariant decomposition

$$\Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n} \cong F_{\mathbb{R}^n} S^n \xleftarrow{j} (F_{\mathbb{R}} S^1)^{\wedge n}.$$

The isomorphism  $j$  arises from the homeomorphism  $(S^1)^{\wedge n} \cong S^n$  and the isomorphism  $(F_{\mathbb{R}})^{\wedge n} \cong F_{\mathbb{R}^n}$  from [35, Remark C.11]. This decomposition is  $\Sigma_n$ -equivariant, since both of the involved maps are symmetric. Explicitly, this isomorphism is given at inner product spaces  $U_1, \dots, U_n$  as

$$(F_{\mathbb{R}} S^1)(U_1) \wedge \cdots \wedge (F_{\mathbb{R}} S^1)(U_n) \xrightarrow{j_{U_1, \dots, U_n}} (F_{\mathbb{R}^n} S^n)(U_1 \oplus \cdots \oplus U_n),$$

$$[(u_1, \varphi_1), t_1] \wedge \cdots \wedge [(u_n, \varphi_n), t_n] \mapsto [(u_1 \oplus \cdots \oplus u_n, \varphi_1 \oplus \cdots \oplus \varphi_n), t_1 \wedge \cdots \wedge t_n].$$

Using this decomposition, we can rewrite the domain of the morphism  $q$ :

$$\Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n} \wedge_{\Sigma_n} X^{\wedge n} \cong F_{\mathbb{R}^n} S^n \wedge_{\Sigma_n} X^{\wedge n} \cong (F_{\mathbb{R}} S^1)^{\wedge n} \wedge_{\Sigma_n} X^{\wedge n} \cong (F_{\mathbb{R}} S^1 \wedge X)^{\wedge n} / \Sigma_n.$$

We claim that under this translation, the morphism  $q$  corresponds to the morphism

$$(\lambda_{\Sigma_1, \mathbb{R}, 0} \wedge X)^{\wedge n} / \Sigma_n : (F_{\Sigma_1, \mathbb{R}} S^1 \wedge X)^{\wedge n} / \Sigma_n \rightarrow (F_{\Sigma_1, 0} \wedge X)^{\wedge n} / \Sigma_n$$

from (2.36). Note that  $\Sigma_1 = e$  is the trivial group, and hence  $0$  is a faithful  $\Sigma_1$ -representation. Moreover,  $F_{\Sigma_1, 0} = \mathbf{O}(0, \_)/\Sigma_1 \cong \mathbb{S}$  is the sphere spectrum. To prove this claim, we consider the diagram

$$(2.38) \quad \begin{array}{ccc} (F_{\mathbb{R}} S^1)^{\wedge n} \wedge_{\Sigma_n} X^{\wedge n} & \longrightarrow & (F_{\mathbb{R}} S^1 \wedge X)^{\wedge n} / \Sigma_n \\ j \wedge_{\Sigma_n} X^{\wedge n} \downarrow & & \downarrow (\lambda_{\Sigma_1, \mathbb{R}, 0} \wedge X)^{\wedge n} / \Sigma_n \\ F_{\mathbb{R}^n} S^n \wedge_{\Sigma_n} X^{\wedge n} & \xrightarrow{\lambda_{\Sigma_1, \mathbb{R}^n, 0} \wedge X^{\wedge n}} & \\ \text{untwisting} \downarrow & \searrow & \downarrow \\ \Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n} \wedge_{\Sigma_n} X^{\wedge n} & \xrightarrow{q} & X^{\wedge n} / \Sigma_n \end{array}$$

We first consider the upper diagram. Let  $U_1, \dots, U_n$  be inner product spaces, and consider the diagram

$$(F_{\mathbb{R}} S^1)(U_1) \wedge \cdots \wedge (F_{\mathbb{R}} S^1)(U_n) \xrightarrow{\lambda_{\mathbb{R}, 0}(U_1) \wedge \cdots \wedge \lambda_{\mathbb{R}, 0}(U_n)} \mathbf{O}(0, U_1) \wedge \cdots \wedge \mathbf{O}(0, U_n)$$

$$\begin{array}{ccc} j_{U_1, \dots, U_n} \downarrow & & \oplus \downarrow \mathbb{R} \\ (F_{\mathbb{R}^n} S^n)(U_1 \oplus \cdots \oplus U_n) & \xrightarrow{\lambda_{\mathbb{R}^n, 0}(U_1 \oplus \cdots \oplus U_n)} & \mathbf{O}(0, U_1 \oplus \cdots \oplus U_n) \cong S^{U_1 \oplus \cdots \oplus U_n} \end{array}$$

Evaluating this on an element yields

$$\begin{array}{ccc}
 [(u_1, \varphi_1), t_1] \wedge \cdots \wedge [(u_n, \varphi_n), t_n] & \longmapsto & [u_1 + \varphi_1(t_1)] \wedge \cdots \wedge [u_n + \varphi_n(t_n)] \\
 \downarrow & & \downarrow \\
 [(u_1 \oplus \cdots \oplus u_n, \varphi_1 \oplus \cdots \oplus \varphi_n), t_1 \wedge \cdots \wedge t_n] & \longmapsto & [(u_1 \oplus \cdots \oplus u_n) + (\varphi_1(t_1) \oplus \cdots \oplus \varphi_n(t_n))]
 \end{array}$$

By applying  $(\_) \wedge_{\Sigma_n} X^{\wedge n}$  to this diagram, we see that the upper half of (2.38) commutes. For the second half of (2.38), let  $U$  be an inner product space. We need to consider the left diagram of

$$\begin{array}{ccc}
 \mathbf{O}(\mathbb{R}^n, U) \wedge S^n & & [(u, \varphi), t] \\
 \text{untwisting} \downarrow & \searrow \lambda_{\Sigma_1, \mathbb{R}^n, 0}(U) & \downarrow \\
 S^U \wedge \mathbf{L}(\mathbb{R}^n, U) & \xrightarrow{q(U)} & \mathbf{O}(0, U) = S^U & \begin{array}{ccc} & & \searrow \\ & & [u + \varphi(t)] \\ & & \longmapsto \\ [u + \varphi(t), \varphi] & & [u + \varphi(t)] \end{array}
 \end{array}$$

On elements, this takes the form of the right diagram.

Thus, this part of (2.38) also commutes. Hence, we have translated the statement of the theorem into the claim that the map

$$\mathbb{P}^n(\lambda_{\Sigma_1, \mathbb{R}, 0} \wedge X) = (\lambda_{\Sigma_1, \mathbb{R}, 0} \wedge X)^{\wedge n} / \Sigma_n : \mathbb{P}^n(F_{\Sigma_1, \mathbb{R}} S^1 \wedge X) \rightarrow \mathbb{P}^n X$$

is a global equivalence. But by [35, Theorem 4.1.29], the morphism  $\lambda_{\Sigma_1, \mathbb{R}, 0}$  is a global equivalence. As the spectrum  $X$  is positively cofibrant, smashing with  $X$  preserves global equivalences by [35, Theorem 4.3.27]. Moreover, we know by Lemma 2.6 that  $\mathbb{P}^n$  sends global equivalences between positively cofibrant spectra to global equivalences. As both  $X$  and  $F_{\Sigma_1, \mathbb{R}} S^1 = F_{\mathbb{R}} S^1$  are positively cofibrant,  $\mathbb{P}^n(\lambda_{\Sigma_1, \mathbb{R}, 0} \wedge X)$  is a global equivalence, and hence  $q$  is as well.  $\square$

**Remark 2.39** Here we used a decomposition of the global classifying space  $\Sigma_+^\infty \mathbf{L}_{\mathbb{R}^n}$  and the fact that we already know that  $\mathbb{P}^n$  preserves global equivalences by Lemma 2.6 to give an easy proof of the theorem. Our proof thus relies on the fact that we already have a model structure on commutative ring spectra, following the approach of White in [48] via analyzing the symmetric powers  $\mathbb{P}^n$ . More classically, in [17; 30; 41], the theorems analogous to Theorem 2.37 are used to provide the model structure on commutative ring spectra. In these sources, the proof of the above theorem is done by cellular induction; see for example the proof of [17, Theorem III.5.1]. A similar proof can also be done in our context, using the above calculations for the induction start and then using [19, Theorem 22] for the induction over the cell attachments.

### 3 Power operations on Moore spectra and $\beta$ -rings

In this part of the article we study  $G_\infty$ -structures on global Moore spectra  $\mathbb{S}B$  for torsion-free commutative rings  $B$ . We show that  $G_\infty$ -ring structures on  $\mathbb{S}B$  provide  $\beta$ -ring structures on all equivariant homotopy groups. The reason that we study Moore spectra is first of all that they are (almost) completely determined by the underlying algebra of the ring  $B$ , so we can translate the topological structure of being a  $G_\infty$ -ring spectrum into an algebraic structure on  $B$ . Moreover, a Moore spectrum can be thought of as an extension of coefficients of the sphere spectrum, and hence has relevance to talking about cohomology theories with coefficients in a ring  $B$ . Hence, providing power operations in the Moore spectrum  $\mathbb{S}B$  is a first step to providing power operations on these extended cohomology theories.

It is known classically that torsion in the ring  $B$  obstructs the existence of a highly structured multiplication on the Moore spectrum  $\mathbb{S}B$ . We can also show that these obstructions occur in the algebra of global power functors. Hence, we restrict our analysis to the class of torsion-free rings, where these phenomena are not visible.

In Section 3.1 we study the topological side of the situation and construct from a global power structure on the homotopy groups of a Moore spectrum a  $G_\infty$ -ring structure. In Theorem 3.14 we arrive at an equivalence between the homotopy category of Moore spectra for torsion-free rings equipped with a  $G_\infty$ -structure and the category of corresponding global power functors. Using one direction of this relationship, which does not need the torsion-freeness assumption, we also obtain easy arguments that the Moore spectra for  $\mathbb{Z}/n$  and  $\mathbb{Z}[i]$  cannot support  $G_\infty$ -ring structures. This is an analogue of the classical facts that these spectra cannot support an  $A_\infty$ - or  $E_\infty$ -structure, respectively.

Then, in an algebraic Section 3.2, we study for which rings there can be global power operations on the homotopy groups of  $\mathbb{S}B$ . For this, we link the power operations on global functors of the form  $\mathbb{A} \otimes B$  to  $\beta$ -ring structures on  $\mathbb{A}(G) \otimes B$ . In particular, we give a new perspective on these objects, using a well-structured theory of global power operations to obtain  $\beta$ -rings. This approach is also used to obtain the  $\beta$ -ring structure of stable cohomotopy.

A similar analysis has been done by Julia Singer in her PhD thesis [40, Section 2.4] in the case of  $H_\infty$ -structures. There, the representation ring functor  $R$  takes the role of the Burnside ring. Our treatment generalizes the results to a global context.

In this section all rings  $B$  are commutative and unital.

### 3.1 $G_\infty$ -structure on Moore spectra for global power functors

In this section we construct  $G_\infty$ -ring structures on Moore spectra from power operations on their homotopy groups. It is well known that the existence of a multiplication on Moore spectra is a subtle question, which traces back to the fact that the cone of a map is only well defined in the homotopy category up to noncanonical isomorphism. This distinguishes them notably from Eilenberg–MacLane spectra, which are unique in a more rigorous way and hence are better behaved for algebraic manipulations. However, there are classes of rings which work better than others, in particular, torsion-free rings support multiplications on their Moore spectra. We thus restrict our attention to the subcategory of torsion-free rings in the following section. The main examples of Moore spectra for torsion rings that do not support commutative multiplications are  $\mathbb{S}(\mathbb{Z}/2)$ , which does not admit a unital multiplication, and  $\mathbb{S}(\mathbb{Z}/3)$ , where Massey products obstruct the associativity of the multiplication. A remark about the latter phenomenon can be found in [3, Example 3.3], and details for the first can be found in [4, Theorem 1.1]. In fact, no Moore spectrum  $\mathbb{S}(\mathbb{Z}/n)$  can have an  $A_n$ -ring structure. That no Moore spectrum  $\mathbb{S}(\mathbb{Z}/n)$  can be endowed with a  $G_\infty$ -structure can be seen in Corollary 3.5, where we show that  $\mathbb{A} \otimes \mathbb{Z}/n$  does not admit the structure of a global power functor. In a similar way, we also observe that the Moore spectrum for the Gaussian integers  $\mathbb{S}(\mathbb{Z}[i])$  does not support a  $G_\infty$ -ring structure. This is an analogue of the result that this spectrum also cannot be endowed with an  $E_\infty$ -multiplication. Note, however, that the nonexistence results for  $G_\infty$ -ring structures for  $\mathbb{S}(\mathbb{Z}/p)$  and  $\mathbb{S}(\mathbb{Z}[i])$  are obtained by easy algebraic calculations rather than by higher obstructions. In this way, the global viewpoint simplifies the calculations.

For countable torsion-free rings  $B$  however, these problems vanish, and we are able to describe  $G_\infty$ -ring structures on Moore spectra completely algebraically. Theorem 3.14 provides an equivalence of  $G_\infty$ -ring structures on  $\mathbb{S}B$  and global power functor structures on  $\mathbb{A} \otimes B$ .

In this section we use the triangulated structure of the categories  $\mathcal{SH}$  and  $\mathcal{GH}$ . As a reference for these triangulated structures we use [35, Section 4.4].

Recall from [35, Theorem 4.5.1] that the forgetful functor  $U: \mathcal{GH} \rightarrow \mathcal{SH}$  has both adjoints, and that the left adjoint is the left derived functor of the identity functor  $\text{Id}: Sp \rightarrow Sp$ . As such, it can be represented by assigning to a nonequivariant spectrum  $X$  a nonequivariant cofibrant replacement  $QX$ , considered as a global homotopy type. By construction of the model structures, in fact any nonequivariant stable equivalence

between cofibrant spectra is a global equivalence. We call the objects in the image of the left adjoint *left-induced homotopy types*, and a spectrum *left-induced (from the trivial group)* if its homotopy type is. When we talk about Moore spectra for rings, we require that all their homology is not only concentrated in degree 0, but also determined by nonequivariant data. Hence we make the following definition:

**Definition 3.1** Let  $B$  be a ring. A (global) Moore spectrum for  $B$  is a connective spectrum  $X$ , left-induced from the trivial group, such that  $H_0^e(X) \cong \pi_0^e(X) \cong B$  and  $H_*^e(X) = 0$  for all  $* \neq 0$ . We denote a Moore spectrum for the ring  $B$  by  $\mathbb{S}B$ .

Since we study power operations on the homotopy groups of a Moore spectrum, we need to calculate  $\underline{\pi}_0(\mathbb{S}B)$  as a global functor. This can be done in terms of  $B$ , using that  $\mathbb{S}B$  is connective and left-induced.

**Proposition 3.2** Let  $X$  be a connective spectrum left-induced from the trivial group. Then the exterior product

$$\boxtimes: \pi_0^G(\mathbb{S}) \otimes \pi_0^e(X) \rightarrow \pi_0^{G \times e}(\mathbb{S} \wedge X) \cong \pi_0^G(X)$$

is an isomorphism of abelian groups. As the group  $G$  varies, these assemble into an isomorphism

$$\underline{\pi}_0(\mathbb{S}) \otimes \pi_0^e(X) \rightarrow \underline{\pi}_0(X)$$

of global functors. If  $X$  is a homotopy ring spectrum, then this is an isomorphism of global Green functors.

**Proof** We first observe that the proposition is true in the case  $X = \mathbb{S}$ , since in this case  $\pi_0^e(\mathbb{S}) \cong \mathbb{Z}$ , and the exterior product is the multiplication map

$$\pi_0^G(\mathbb{S}) \otimes \mathbb{Z} \rightarrow \pi_0^G(\mathbb{S}),$$

which is an isomorphism. Moreover, we note that the class of spectra  $X$  for which the exterior product map is an isomorphism is closed under coproducts, since  $\boxtimes$  is additive in the spectrum  $X$ . It is also closed under cones as defined on [35, page 419] by the following argument.

Suppose we have a distinguished triangle

$$X \rightarrow Y \rightarrow Z \rightarrow X[1]$$

of connective spectra in the stable homotopy category, where for  $X$  and  $Y$  the map

$$\boxtimes: \pi_0^G(\mathbb{S}) \otimes \pi_0^e(-) \rightarrow \pi_0^G(-)$$

is an isomorphism. Then the left-induced triangle is also distinguished, and we obtain a long exact sequence in homotopy groups

$$\pi_0^G(X) \rightarrow \pi_0^G(Y) \rightarrow \pi_0^G(Z) \rightarrow \pi_{-1}^G(X) = 0$$

for all compact Lie groups  $G$ . As the exterior product is natural, we obtain the commutative diagram

$$\begin{array}{ccccccc} \pi_0^G(\mathbb{S}) \otimes \pi_0^e(X) & \longrightarrow & \pi_0^G(\mathbb{S}) \otimes \pi_0^e(Y) & \longrightarrow & \pi_0^G(\mathbb{S}) \otimes \pi_0^e(Z) & \longrightarrow & 0 \\ \boxtimes \downarrow & & \boxtimes \downarrow & & \boxtimes \downarrow & & \\ \pi_0^G(X) & \longrightarrow & \pi_0^G(Y) & \longrightarrow & \pi_0^G(Z) & \longrightarrow & 0 \end{array}$$

with exact rows, where the two left vertical maps are isomorphisms. Then by the 5–lemma the exterior product is also an isomorphism for  $Z$ .

Then by [35, Proposition 4.4.13], and respectively its nonequivariant analogue, the map  $\boxtimes$  is an isomorphism for all connective left-induced spectra, since the sphere spectrum is a compact weak generator of  $\mathcal{SH}$ .

Now, by the properties of the external product [35, Theorem 4.1.22], the map  $\boxtimes$  is a morphism of global functors, and levelwise an isomorphism, hence it is an isomorphism of global functors. Moreover, if  $X$  is a homotopy ring spectrum, then  $\boxtimes$  is a map of global Green functors. This proves the proposition.  $\square$

**Remark 3.3** The above statement can also be deduced from an identification of the genuine fixed points of a left-induced spectrum. Let  $X$  be a nonequivariant spectrum and denote by  $LX$  the corresponding left-induced spectrum. Then we obtain

$$(LX)^G \simeq \mathbb{S}^G \wedge X,$$

where  $(-)^G$  denotes the genuine fixed point functor. This relation can easily be seen using an  $\infty$ –categorical approach. Here, the functor  $(-) \mapsto (L-)^G$  preserves colimits, hence we are able to calculate  $(LX)^G$  by evaluating the functor on the sphere spectrum and smashing with  $X$ . From this formula, we deduce the above proposition by observing that  $\pi_0^G(X) = \pi_0(X^G)$  and that  $\pi_0$  is strict monoidal on connective spectra.

**Corollary 3.4** *Let  $X$  be a left-induced spectrum. If  $X$  supports the structure of a  $G_\infty$ -ring spectrum, then it induces the structure of a global power functor on*

$$\pi_0(X) \cong \mathbb{A} \otimes \pi_0^e(X).$$

Thus, we take particular interest in global power functors of the form  $\mathbb{A} \otimes B$  in the following. These can also be described as global power functors  $R$ , where the multiplication map  $\mathbb{A} \otimes R(e) \rightarrow R$ , arising from the  $\mathbb{A}$ -module structure of  $R$ , is an isomorphism of global Green functors. We call such global power functors *left-induced* and call the full subcategory of such global power functors  $\mathcal{GIPow}_{\text{left}}$ . Any morphism between such left-induced global power functors is already determined by the morphism at the trivial group  $e$ .

Note that, a priori,  $\mathbb{A} \otimes B$  only has the structure of a global Green functor for a ring  $B$ . The existence of power operations is additional structure. In Section 3.2, we exhibit a relationship between global power functor structures on  $\mathbb{A} \otimes B$  and  $\beta$ -ring structures on  $\mathbb{A}(G) \otimes B$ . Moreover, we can show that for certain rings, such power operations cannot exist:

**Corollary 3.5** *The Moore spectra  $\mathbb{S}(\mathbb{Z}/n)$  for  $n \in \mathbb{N}$  and  $\mathbb{S}(\mathbb{Z}[i])$  do not support a  $G_\infty$ -ring structure.*

**Proof** We claim that the global functor  $\mathbb{A} \otimes \mathbb{Z}/n$  does not support a global power functor structure. Suppose otherwise, then we in particular obtain power operations

$$P^m : \mathbb{Z}/n \rightarrow \mathbb{A}(\Sigma_m) \otimes \mathbb{Z}/n$$

for all  $m \geq 0$ . Note that since  $\mathbb{Z}/n$  is additively cyclic, if a power operation exists, it is induced by the power operations on  $\mathbb{A}$  upon taking the quotient by  $n$ , using additivity. Hence, we can provide formulas for the power operations in terms of powers of finite sets. We claim that if  $p$  is any prime factor of  $n$ , then the power operation  $P^p : \mathbb{Z} \rightarrow \mathbb{A}(\Sigma_p)$  does not descend to  $\mathbb{Z}/n \rightarrow \mathbb{A}(\Sigma_p) \otimes \mathbb{Z}/n$ .

To check this, we calculate  $P^p(n) \in \mathbb{A}(\Sigma_p)$ . We observe by the explicit description of the power operations on the Burnside ring of a finite group in [35, Example 5.3.1] that the element  $P^p(n) \in \mathbb{A}(\Sigma_p)$  is represented by the  $\Sigma_p$ -set  $[n]^p$  with the permutation action, where we denote by  $[n]$  the set  $\{1, \dots, n\}$  with no group actions. We decompose this  $\Sigma_p$ -set as a disjoint union of  $\Sigma_p$ -orbits.

In particular, we consider the free orbits in  $[n]^p$ . Such orbits are in bijection with equivalence classes of points  $(k_1, \dots, k_p)$  with pairwise different  $k_i$ , up to reordering. There are  $\binom{n}{p}$  such classes, hence this is the coefficient of  $\Sigma_p/e$  in  $[n]^p$ . Since we have

$$\binom{n}{p} = \frac{n \cdots (n - p + 1)}{p \cdots 1},$$

and in the numerator only  $n$  is divisible by  $p$ , we see that  $\binom{n}{p}$  is divisible by  $\frac{n}{p}$ , but not by  $n$ . Thus the element  $P^p(n) \in \mathbb{A}(\Sigma_p)$  is not divisible by  $n$ . Hence we do not have power operations on the global functors  $\mathbb{A} \otimes \mathbb{Z}/n$  for any  $n$ .

For the case  $\mathbb{Z}[i]$ , we use the relation  $(P^m(i))^2 = P^m(-1)$ . Then we observe that  $P^2(-1) = t - 1$ , where we write  $t = \text{tr}_e^{\Sigma_2} \in \mathbb{A}(\Sigma_2)$ . This is not a square in  $\mathbb{A}(\Sigma_2) \otimes \mathbb{Z}[i]$ , as we can calculate

$$(at + b)^2 = 2a(a + b)t + b^2$$

for  $a, b \in \mathbb{Z}[i]$ , and 2 is not invertible in  $\mathbb{Z}[i]$ . □

Note that  $\mathbb{A} \otimes \mathbb{Z}/n$  does however support truncated power operations, namely maps  $P^k$  for all  $k < p$ , where  $p$  is the smallest prime divisor of  $n$ . This can be shown by similar computations as above for all stabilizers of points in  $[n]^p$ . Such truncated power operations are also an object of research, for example in [9], and relate to the fact that  $\mathbb{S}(\mathbb{Z}/p)$  has an  $A_{p-1}$ -multiplication. Moreover, we observe that  $P^2(-1) = t - 1$  becomes a square after inverting 2 in  $\mathbb{A} \otimes \mathbb{Z}[i]$ . In fact, there is a strict commutative model for the Moore spectrum of  $\mathbb{Z}[i, \frac{1}{2}]$ .

We now study the reverse direction in the case that  $X$  is a global Moore spectrum. We aim to define a  $G_\infty$ -ring structure on  $\mathbb{S}B$  from power operations on its homotopy groups. As already mentioned, we restrict to torsion-free rings in the following.

We first show that the ring structure of  $B$  induces the structure of a homotopy ring spectrum on  $\mathbb{S}B$ . For this, we show that we can test properties of morphisms between Moore spectra on the homotopy groups. This lemma should be well-known, but since the author is not aware of a proof in the literature, it is included here for the sake of completeness.

**Lemma 3.6** *The functor*

$$\pi_0^e : \text{Moore}^{\text{torsion-free}} \rightarrow \text{Ab}^{\text{torsion-free}}$$



between the homotopy category of Moore spectra of torsion-free abelian groups and the category of torsion-free abelian groups is fully faithful, and hence an equivalence of categories.

**Proof** Let  $A$  and  $B$  be torsion-free groups. We have to calculate the group of morphisms  $[\mathbb{S}A, \mathbb{S}B] := \mathcal{GH}(\mathbb{S}A, \mathbb{S}B)$ . Note that in fact  $\mathcal{GH}(\mathbb{S}A, \mathbb{S}B) \cong \mathcal{SH}(\mathbb{S}A, \mathbb{S}B)$ , since Moore spectra are left-induced. To calculate this group of morphisms, we consider a free resolution

$$(3.7) \quad 0 \rightarrow \mathbb{Z}^{\oplus I} \rightarrow \mathbb{Z}^{\oplus J} \rightarrow A \rightarrow 0$$

of  $A$ . Then we take as a model of the Moore spectrum for  $A$  the cone

$$\bigvee_I \mathbb{S} \rightarrow \bigvee_J \mathbb{S} \rightarrow \mathbb{S}A.$$

Using this distinguished triangle and mapping into  $\mathbb{S}B$ , we obtain an exact sequence

$$[\bigvee_J \mathbb{S}\mathbb{S}, \mathbb{S}B] \rightarrow [\bigvee_I \mathbb{S}\mathbb{S}, \mathbb{S}B] \rightarrow [\mathbb{S}A, \mathbb{S}B] \rightarrow [\bigvee_J \mathbb{S}, \mathbb{S}B] \rightarrow [\bigvee_I \mathbb{S}, \mathbb{S}B].$$

Since the homotopy classes are additive under wedges, we can write this sequence as

$$\begin{aligned} \text{Hom}(\mathbb{Z}^{\oplus J}, \pi_1(\mathbb{S}B)) &\rightarrow \text{Hom}(\mathbb{Z}^{\oplus I}, \pi_1(\mathbb{S}B)) \rightarrow [\mathbb{S}A, \mathbb{S}B] \\ &\rightarrow \text{Hom}(\mathbb{Z}^{\oplus J}, B) \rightarrow \text{Hom}(\mathbb{Z}^{\oplus I}, B). \end{aligned}$$

As  $\text{Hom}$  is left exact and the sequence (3.7) is exact, we see that the kernel of the rightmost map is  $\text{Hom}(A, B)$ . Moreover, the cokernel of the leftmost map is isomorphic to  $\text{Ext}_{\mathbb{Z}}^1(A, \pi_1(\mathbb{S}B))$ . We now calculate  $\pi_1(\mathbb{S}B)$ , using a free resolution

$$0 \rightarrow \mathbb{Z}^{\oplus I'} \rightarrow \mathbb{Z}^{\oplus J'} \rightarrow B \rightarrow 0,$$

which gives a cofibre sequence in the global homotopy category. The associated long exact sequence in homotopy groups gives

$$\bigoplus_{I'} \mathbb{Z}/2 \rightarrow \bigoplus_{J'} \mathbb{Z}/2 \rightarrow \pi_1(\mathbb{S}B) \rightarrow \bigoplus_{I'} \mathbb{Z} \rightarrow \bigoplus_{J'} \mathbb{Z},$$

using  $\pi_1(\mathbb{S}) \cong \mathbb{Z}/2$ . As the rightmost map is injective and tensor product is right-exact, we obtain  $\pi_1(\mathbb{S}B) \cong B \otimes \mathbb{Z}/2$ .

Since  $\text{Ext}_{\mathbb{Z}}^1$  is additive, we see that  $\text{Ext}_{\mathbb{Z}}^1(A, B/2)$  is 2-torsion. Moreover, the sequence  $0 \rightarrow A \xrightarrow{2} A \rightarrow A/2 \rightarrow 0$  is exact, as  $A$  is torsion-free, and the long exact sequence of  $\text{Ext}$ -functors yields that

$$\text{Ext}_{\mathbb{Z}}^1(A, B/2) \xrightarrow{2} \text{Ext}_{\mathbb{Z}}^1(A, B/2) \rightarrow \text{Ext}_{\mathbb{Z}}^2(A/2, B/2) = 0$$

is exact. Thus  $\text{Ext}_{\mathbb{Z}}^1(A, B/2)$  is also 2-divisible and hence vanishes. □

To consider ring structures on Moore spectra, we need to calculate whether  $\mathbb{S}B \wedge \mathbb{S}B$  is again a Moore spectrum:

**Proposition 3.8** *Let  $A$  and  $B$  be commutative rings whose underlying abelian group is torsion-free. Then the external product*

$$H_*(\mathbb{S}A, \mathbb{Z}) \otimes H_*(\mathbb{S}B, \mathbb{Z}) \rightarrow H_*(\mathbb{S}A \wedge \mathbb{S}B, \mathbb{Z})$$

*in homology is an isomorphism. Hence  $\mathbb{S}A \wedge \mathbb{S}B$  is a Moore spectrum for  $A \otimes B$ .*

**Proof** This follows from the Künneth theorem; see for example [1, (KT1) and Note 12]. The homology groups  $H_*(\mathbb{S}A)$  and  $H_*(\mathbb{S}B)$  are flat over  $\mathbb{Z}$ , since they are torsion-free. This proves that  $H_*(\mathbb{S}A \wedge \mathbb{S}B)$  is  $A \otimes B$  concentrated in degree 0. □

**Corollary 3.9** *The equivalence from Lemma 3.6 induces an equivalence*

$$\pi_0^e : \text{Moore}_{\text{Rings}}^{\text{torsion-free}} \rightarrow \text{Rings}^{\text{torsion-free}}$$

*between the homotopy category of commutative homotopy ring Moore spectra for torsion-free rings and the category of torsion-free commutative rings.*

We now follow the same approach in order to put a  $G_\infty$ -ring structure on  $\mathbb{S}B$ . Note that  $\mathbb{G}(\mathbb{S}B)$  is not a Moore spectrum, since it is not left-induced. However, we still can calculate the group  $[\mathbb{G}(\mathbb{S}B), \mathbb{S}B]$  in terms of the homotopy groups of  $\mathbb{S}B$ . The proof of the following lemma uses representability of the equivariant homotopy groups for the case that  $B$  is finitely generated and free. In order to reduce to this case, we assume countability of the group  $B$ .

**Lemma 3.10** *Let  $B$  be a countable torsion-free abelian group, and let  $Y$  be an orthogonal spectrum such that there is an isomorphism  $\underline{\pi}_1(Y) \cong R \otimes A$  of global functors, where  $A$  is a commutative ring and  $R$  is a global functor such that for any finite group  $G$ ,  $R(G)$  is finite. Then for either of the spectra  $X = \mathbb{G}(\mathbb{S}B)$  or  $X = \mathbb{G}(\mathbb{G}(\mathbb{S}B))$ , the morphism*

**(3.11)** 
$$\underline{\pi}_0 : \mathcal{GH}(X, Y) \rightarrow \mathcal{GF}(\underline{\pi}_0(X), \underline{\pi}_0(Y))$$

*is an isomorphism.*

**Proof** We start by considering  $X = \mathbb{G}(\mathbb{S}B)$ , and first consider the case that  $B$  is free. Then we choose a basis  $(x_i)_{i \in I}$  of  $B$ , such that  $\mathbb{S}B \cong \bigvee_{i \in I} \mathbb{S}$ . We calculate

$$\begin{aligned}
 (3.12) \quad \mathbb{G}^m(\mathbb{S}B) &= \mathbb{G}^m\left(\bigvee_{i \in I} \mathbb{S}\langle x_i \rangle\right) \\
 &= \bigvee_{(m)} \bigwedge_{i \in I, m_i \neq 0} \mathbb{G}^{m_i} \mathbb{S} = \bigvee_{(m)} \bigwedge_{i \in I, m_i \neq 0} \Sigma_+^\infty B_{\text{gl}} \Sigma_{m_i} \\
 &= \bigvee_{(m)} \Sigma_+^\infty B_{\text{gl}} \left( \bigtimes_{\substack{i \in I \\ m_i \neq 0}} \Sigma_{m_i} \right),
 \end{aligned}$$

where  $(m) = (m_i)_{i \in I}$  runs through all partitions  $m = \sum_{i \in I, m_i \neq 0} m_i$ .

By representability of the homotopy groups  $\pi_0^G$  by  $\Sigma_+^\infty B_{\text{gl}} G$  from (1.2), the map  $\underline{\pi}_0$  from (3.11) is an isomorphism for the spectra

$$X = \Sigma_+^\infty B_{\text{gl}} \left( \bigtimes_{\substack{i \in I \\ m_i \neq 0}} \Sigma_{m_i} \right).$$

As both the domain and codomain of the map (3.11) are additive under wedges, for  $\mathbb{G}(\mathbb{S}B)$  the morphism  $\underline{\pi}_0$  is also an isomorphism.

Now let  $B$  be a general torsion-free abelian group. By [28], any flat module  $M$  is isomorphic to a directed colimit of finitely generated free modules, generated on finite sets of elements in  $M$ . Since, over  $\mathbb{Z}$ , flat and torsion-free are equivalent, we can write  $B \cong \text{colim}_{i \in I} B_i$  as a directed colimit of finitely generated free  $\mathbb{Z}$ -modules. Moreover, since  $B$  is countable, we find a cofinal sequential system in the directed indexing system, so that we can write  $B$  as a sequential colimit of finitely generated free modules  $B_n$  with  $n \in \mathbb{N}$ . To see that this is the case, we consider the directed system  $(\varphi_A: F_A \rightarrow B)_{A \in \mathcal{S}(B)}$  for the directed set  $\mathcal{S}(B)$  of finite subsets of  $B$ . Here,  $F_A$  is the free module on  $A$ . Then  $\mathcal{S}(B)$  is countable. We choose a bijection  $s: \mathbb{N} \rightarrow \mathcal{S}(B)$  and iteratively define  $B_n$ . Set  $B_1 = F_{s(1)}$ . Once we have defined  $B_n = F_{A_n}$ , we take an element  $A_{n+1}$  such that both  $A_n$  and  $s(n+1)$  map to  $A_{n+1}$  in the directed set  $\mathcal{S}(B)$ , and define  $B_{n+1} = F_{A_{n+1}}$ . This clearly defines a cofinal sequential subset of  $\mathcal{S}(B)$ .

We can then lift this sequential system to a cofibrant system  $\mathbb{S}B_n$  of Moore spectra, such that the colimit  $\text{colim}_n \mathbb{S}B_n$  models the homotopy colimit. Since homology commutes with sequential homotopy colimits, we see that  $\mathbb{S}B \cong \text{colim}_n \mathbb{S}B_n$  is a model for the Moore spectrum of  $B$ . Since sequential colimits of ultracommutative ring spectra are calculated on the underlying spectra, the functor  $\mathbb{P}: \mathcal{S}p \rightarrow \mathcal{S}p$  also commutes with

sequential colimits, and thus  $\mathbb{G}(\mathbb{S}B) \cong \text{colim}_n \mathbb{G}(\mathbb{S}B_n)$ . Then we obtain the Milnor exact sequence

$$0 \rightarrow \lim_n^1 \mathcal{GH}(\mathbb{G}(\mathbb{S}B_n), \Omega Y) \rightarrow \mathcal{GH}(\mathbb{G}(\mathbb{S}B), Y) \rightarrow \lim_n \mathcal{GH}(\mathbb{G}(\mathbb{S}B_n), Y) \rightarrow 0.$$

By the above arguments for free modules  $B$ , for the right-hand object

$$\begin{aligned} \lim_n \mathcal{GH}(\mathbb{G}(\mathbb{S}B_n), Y) &\cong \lim_n \text{Hom}_{\mathcal{GF}}(\underline{\pi}_0(\mathbb{G}(\mathbb{S}B_n)), \underline{\pi}_0(Y)) \\ &\cong \text{Hom}_{\mathcal{GF}}(\underline{\pi}_0(\mathbb{G}(\mathbb{S}B)), \underline{\pi}_0(Y)). \end{aligned}$$

Similarly, for the left-hand object

$$\lim_n^1 \mathcal{GH}(\mathbb{G}(\mathbb{S}B_n), \Omega Y) \cong \lim_n^1 \text{Hom}_{\mathcal{GF}}(\underline{\pi}_0(\mathbb{G}(\mathbb{S}B_n)), \underline{\pi}_1(Y)).$$

By (3.12), this last term is isomorphic to

$$\lim_n^1 \prod_{(m^n)} \pi_1^{\Sigma(m^n)}(Y),$$

where the product is over all tuples of natural numbers  $m_i^n \geq 0$  indexed on a basis  $I_n$  of  $B_n$ , and we write  $\Sigma(m^n) = \times_{i \in I_n} \Sigma m_i^n$ . Now this  $\lim^1$ -term decomposes as the product

$$\prod_{m \geq 0} \left[ \lim_n^1 \prod_{\sum_{I_n} m_i^n = m} \pi_1^{\Sigma(m^n)}(Y) \right] \cong \prod_{m \geq 0} \left[ \lim_n^1 \left( A \otimes \prod_{\sum_{I_n} m_i^n = m} R(\Sigma(m^n)) \right) \right].$$

In each of the individual  $\lim^1$ -terms, the inverse system consists only of tensor products of finite groups with  $A$ , since  $I_n$  is finite for every  $n$  and  $R(G)$  is finite for any finite group  $G$ . Thus, these systems satisfy the Mittag-Leffler condition and thus the  $\lim^1$ -term vanishes.

This proves that the morphism  $\underline{\pi}_0$  is an isomorphism.

The arguments for  $\mathbb{G}(\mathbb{G}(\mathbb{S}B))$  are completely analogous. □

We now check that the assumptions on  $\underline{\pi}_1(Y)$  are satisfied in the case of a Moore spectrum  $\mathbb{S}B$ .

**Lemma 3.13** *Let  $B$  be an abelian group and  $\mathbb{S}B$  be a Moore spectrum for  $B$ . Then:*

- (i) *There is an isomorphism  $\underline{\pi}_1(\mathbb{S}B) \cong \underline{\pi}_1(\mathbb{S}) \otimes B$  of global functors, given by  $\boxtimes$ .*
- (ii) *For any finite group  $G$ , the homotopy group  $\pi_1^G(\mathbb{S})$  is finite.*

**Proof** Let  $0 \rightarrow \mathbb{Z}^{\oplus I} \rightarrow \mathbb{Z}^{\oplus J} \rightarrow B \rightarrow 0$  be a free resolution of  $B$ . Then we construct the Moore spectrum  $\mathbb{S}B$  as the mapping cone in

$$\bigvee_I \mathbb{S} \rightarrow \bigvee_J \mathbb{S} \rightarrow \mathbb{S}B.$$

Thus, the long exact sequence of homotopy groups becomes

$$\begin{aligned} \cdots \rightarrow \bigoplus_I \pi_1(\mathbb{S}) &\rightarrow \bigoplus_J \pi_1(\mathbb{S}) \rightarrow \pi_1(\mathbb{S}B) \\ &\rightarrow \bigoplus_I \pi_0(\mathbb{S}) \rightarrow \bigoplus_J \pi_0(\mathbb{S}) \rightarrow \pi_0(\mathbb{S}B). \end{aligned}$$

Now, by freeness of  $\pi_0^G(\mathbb{S}) \cong \mathbb{A}(G)$ , we know that the second row of this sequence is left exact, so the first row is right exact. Moreover, tensoring with  $\pi_1(\mathbb{S})$  is right exact, hence applying this to the sequence  $0 \rightarrow \mathbb{Z}^{\oplus I} \rightarrow \mathbb{Z}^{\oplus J} \rightarrow B \rightarrow 0$  proves  $\pi_1(\mathbb{S}B) \cong \pi_1(\mathbb{S}) \otimes B$ .

For the second statement, we use the tom Dieck splitting [15, Satz 2] and the Adams isomorphism [2, Theorem 5.4], which decompose the homotopy groups for any compact Lie group  $G$  as

$$\pi_1^G(\mathbb{S}) \cong \bigoplus_{(H) \subset G} \pi_1^{W_G H}(E W_G H_+ \wedge \mathbb{S}^H) \cong \bigoplus_{(H) \subset G} \pi_1(\Sigma_+^\infty B W_G H).$$

Here the sum runs over conjugacy classes of closed subgroups of  $G$ , and  $W_G H$  denotes the Weyl group of  $H$  in  $G$ . Now the based suspension spectrum of  $B W_G H$  splits stably as  $\mathbb{S} \vee \widetilde{\Sigma}_+^\infty B W_G H$ , a sum of the sphere spectrum and the reduced suspension spectrum of  $B W_G H$ . Thus, we find

$$\pi_1(\Sigma_+^\infty B W_G H) \cong \pi_1(\mathbb{S}) \oplus \pi_1^{\text{st}}(B W_G H) \cong \mathbb{Z}/2 \oplus \pi_0(W_G H)^{\text{ab}}.$$

Hence,  $\pi_1^G(\mathbb{S})$  is finite for any finite group  $G$ . □

**Theorem 3.14** *The functor*

$$\underline{\pi}_0 : G_\infty\text{-Moore}^{\text{torsion-free}} \rightarrow \mathcal{GIPow}_{\text{left}}^{\text{torsion-free}}$$

*is an equivalence of categories between the homotopy category of  $G_\infty$ -Moore spectra for countable torsion-free commutative rings and the category of countable torsion-free left-induced global power functors  $R$ .*

**Proof** We first prove that the functor  $\underline{\pi}_0$  from the theorem is essentially surjective. To do this, let  $B = R(e)$  for a torsion-free left-induced global power functor  $R \cong \mathbb{A} \otimes B$ . We have to define a  $G_\infty$ -multiplication  $\zeta : \mathbb{G}(\mathbb{S}B) \rightarrow \mathbb{S}B$  on  $\mathbb{S}B$ .

By Lemma 3.10, we know that the map

$$\pi_0: \mathcal{GH}(\mathbb{G}(\mathbb{S}B), \mathbb{S}B) \rightarrow \mathcal{GF}(\pi_0(\mathbb{G}(\mathbb{S}B)), \pi_0(\mathbb{S}B))$$

is an isomorphism. We have calculated the homotopy groups global functor of  $\mathbb{S}B$  in Proposition 3.2 to be  $\mathbb{A} \otimes B$ . Moreover, by [35, Theorem 5.4.11], we have that  $\pi_0(\mathbb{G}(\mathbb{S}B))$  is the free global power functor on the global functor  $\pi_0(\mathbb{S}B) \cong \mathbb{A} \otimes B$ . We denote this free global power functor by  $F(\mathbb{A} \otimes B)$ .

This free global power functor is part of a diagram

$$\begin{array}{ccccc}
 & & U & & \\
 & \swarrow & & \searrow & \\
 \mathcal{G}IPow & \xrightarrow{U} & \mathcal{G}IGreen & \xrightarrow{U} & \mathcal{G}\mathcal{F} \\
 & \swarrow C & & \searrow \mathbb{P} & \\
 & & F & & 
 \end{array}$$

of adjunctions, where all functors labelled  $U$  are forgetful functors and right adjoints,  $\mathbb{P}$  is the symmetric algebra functor for the box product of global functors and  $C$  is the free global power functor for a global Green functor constructed in [35, Proposition 5.2.21]. We now claim that the composite adjunction featuring  $F$  is monadic. For this, we use Beck’s monadicity theorem; see for example [29, VI.7 Theorem 1].

Let  $R \rightrightarrows S$  be a pair of parallel arrows in  $\mathcal{G}IPow$  that has a split coequalizer in  $\mathcal{G}\mathcal{F}$ . Then by monadicity of the adjunction  $\mathcal{G}IGreen \xrightleftharpoons[\mathbb{P}]{U} \mathcal{G}\mathcal{F}$ , the coequalizer in  $\mathcal{G}\mathcal{F}$  has the unique structure of a global Green functor such that it is a coequalizer of  $R \rightrightarrows S$  in  $\mathcal{G}IGreen$ . Since  $\mathcal{G}IPow$  is also comonadic over  $\mathcal{G}IGreen$  by [35, Theorem 5.2.13], colimits in  $\mathcal{G}IPow$  are created by  $U: \mathcal{G}IPow \rightarrow \mathcal{G}IGreen$ . Thus, Beck’s monadicity theorem shows that the adjunction  $\mathcal{G}IPow \xrightleftharpoons[F]{U} \mathcal{G}\mathcal{F}$  is monadic. We also denote the associated monad  $UF$  by  $F$ .

Hence, the power functor structure on  $R \cong \mathbb{A} \otimes B$  is equivalent to a morphism  $\tau: F(\mathbb{A} \otimes B) \rightarrow \mathbb{A} \otimes B$ , satisfying the compatibility conditions with the monad structure on  $F$ . Since  $\pi_0$  (3.11) is an isomorphism for  $\mathbb{G}(\mathbb{S}B)$  and  $\mathbb{S}B$ , this morphism  $\tau: F(\mathbb{A} \otimes B) \rightarrow \mathbb{A} \otimes B$  is the image of a unique morphism  $\zeta: \mathbb{G}(\mathbb{S}B) \rightarrow \mathbb{S}B$  under the functor  $\pi_0$ . We claim that  $\zeta$  endows  $\mathbb{S}B$  with the structure of a  $G_\infty$ -ring spectrum.

For this, we check that the functor  $\pi_0: \mathcal{GH}_{\geq 0} \rightarrow \mathcal{G}\mathcal{F}$  sends the monad diagrams for  $\mathbb{G}$  to those of  $F$ . Here the subscript  $\geq 0$  denotes the full subcategory on the connective

spectra. For this, we consider the diagrams

$$(3.15) \quad \begin{array}{ccc} \mathcal{GH}_{\geq 0} & \xrightarrow{L_{\text{gl}}\mathbb{P}} & \text{Ho}(\text{ucom})_{\geq 0} \\ \pi_0 \downarrow & & \downarrow \pi_0 \\ \mathcal{GF} & \xrightarrow{F} & \mathcal{GIPow} \end{array} \quad \text{and} \quad \begin{array}{ccc} \text{Ho}(\text{ucom})_{\geq 0} & \xrightarrow{U} & \mathcal{GH}_{\geq 0} \\ \pi_0 \downarrow & & \downarrow \pi_0 \\ \mathcal{GIPow} & \xrightarrow{U} & \mathcal{GF} \end{array}$$

The right diagram commutes, thus we get a natural transformation  $\rho: F \circ \pi_0 \rightarrow \pi_0 \circ L_{\text{gl}}\mathbb{P}$  in the left diagram as the mate of the right isomorphism (see [25, Proposition 2.1] for the definition of mates). This transformation is thus defined by freeness of  $F$ , and [35, Theorem 5.4.11] proves that  $\rho$  is a natural isomorphism. The pasting of the two diagrams in (3.15), using the inverse of the transformation  $\rho$ , then exhibits  $\pi_0: \mathcal{GH}_{\geq 0} \rightarrow \mathcal{GF}$  as a monad functor between the monads  $\mathbb{G}$  and  $F$ . For compatibility with the monad structure, naturality of mates is used.

Thus, we see that the  $G_\infty$ -diagrams

$$\begin{array}{ccc} \mathbb{G}(\mathbb{G}(\mathbb{S}B)) & \xrightarrow{\mathbb{G}(\zeta)} & \mathbb{G}(\mathbb{S}B) \\ \mu \downarrow & & \downarrow \zeta \\ \mathbb{G}(\mathbb{S}B) & \xrightarrow{\zeta} & \mathbb{S}B \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathbb{S}B & \xrightarrow{\eta} & \mathbb{G}(\mathbb{S}B) \\ & \searrow & \downarrow \zeta \\ & & \mathbb{S}B \end{array}$$

are sent under  $\pi_0$  to the corresponding diagrams for the monad  $F$ . Since  $\tau$  defines the structure of an  $F$ -algebra on  $\mathbb{A} \otimes B$  by assumption, the monad diagrams for  $\mathbb{A} \otimes B$  commute. Using Lemma 3.10 for morphisms out of  $\mathbb{G}(\mathbb{G}(\mathbb{S}B))$  and Lemma 3.6 for  $\mathbb{S}B$ , the  $G_\infty$ -diagrams for  $\mathbb{S}B$  also commute. This proves that

$$\pi_0: G_\infty\text{-Moore}^{\text{torsion-free}} \rightarrow \mathcal{GIPow}_{\text{left}}^{\text{torsion-free}}$$

is essentially surjective.

To check that  $\pi_0$  is also fully faithful, we only need to check that the unique induced map  $\mathbb{S}B \rightarrow \mathbb{S}C$  from Lemma 3.6 for a map  $f: \mathbb{A} \otimes B \rightarrow \mathbb{A} \otimes C$  of left-induced global power functors is a morphism of  $G_\infty$ -ring spectra. But this again follows from Lemma 3.10 by looking at the diagrams

$$\left( \begin{array}{ccc} \mathbb{G}(\mathbb{S}B) & \xrightarrow{\zeta_B} & \mathbb{S}B \\ \downarrow & & \downarrow \\ \mathbb{G}(\mathbb{S}C) & \xrightarrow{\zeta_C} & \mathbb{S}C \end{array} \right) \mapsto \left( \begin{array}{ccc} F(\mathbb{A} \otimes B) & \xrightarrow{\tau_B} & \mathbb{A} \otimes B \\ Ff \downarrow & & \downarrow f \\ F(\mathbb{A} \otimes C) & \xrightarrow{\tau_C} & \mathbb{A} \otimes C \end{array} \right).$$

Here, the right diagram commutes by assumptions on  $f$ , so the left diagram also commutes. In total, the functor

$$\underline{\pi}_0 : G_\infty\text{-Moore}^{\text{torsion-free}} \rightarrow \mathcal{GIPow}_{\text{left}}^{\text{torsion-free}}$$

is an equivalence of categories.  $\square$

### 3.2 The relation to $\beta$ -rings

We now connect the theory of global power functors to the theory of  $\beta$ -rings. For this we use the perspective of global power functors as coalgebras over the comonad  $\exp$  on the category of global Green functors; see [35, Section 5.2]. Thus, a global power functor  $R$  comes with power operations  $R(G) \rightarrow \exp(R; G) \subset \prod_{m \geq 0} R(\Sigma_m \wr G)$ . On the other hand, a  $\beta$ -ring  $A$  has power operations indexed by the Burnside rings of symmetric groups, given as maps  $A \otimes \mathbb{A}(\Sigma_m) \rightarrow A$ . In order to obtain such structure from the power operations on a global power functor, we assume that the global power functor  $R$  comes equipped with *deflation maps*  $R(K \times G) \times \mathbb{A}(K) \rightarrow R(G)$ . These allow us to dualize the power operations to obtain  $\beta$ -operations on the values  $R(G)$ .

It is a classical observation that the Burnside rings in fact support such deflations. Thus, we can apply the theory presented in this section to the left-induced global power functors  $\mathbb{A} \otimes B$ , which by Theorem 3.14 completely parametrize  $G_\infty$ -ring structures on Moore spectra  $\mathbb{S}B$ . Such structures are hence tied to  $\beta$ -ring structures on  $\mathbb{A}(G) \otimes B$ , extending the classical  $\beta$ -ring  $\mathbb{A}(G)$ .

Another example possessing the necessary deflations is given by stable cohomotopy  $\underline{\pi}^0(X)$  for any based space  $X$ . Here the deflations can be constructed by the theory of polynomial operations, as described in [20; 45]. Thus, our approach of providing  $\beta$ -operations via global power operations and deflations gives back the classical examples of  $\beta$ -rings.

Our approach is loosely based on the discussion of  $\tau$ -rings in [18, Section 4.2], which itself goes back to [23].

Note that we could also consider deflations indexed by arbitrary global power functors  $T$  instead of  $\mathbb{A}$ . For the representation ring global power functor, this yields the theory of  $\lambda$ -rings. However, there are not many other examples of global power functors supporting deflations, so we focus on the Burnside ring.



**Remark 3.16** We give a short overview over the history of the notion of  $\beta$ -rings.

The notion of a  $\beta$ -ring was first introduced by Rymer in [34], based on the question, posed by Boorman in [10], of whether there is a theory of  $\beta$ -rings which formalizes the  $\beta$ -operations on the Burnside ring defined in this work. Rymer, however, did not define his operator ring structure on  $\mathbf{B} = \bigoplus_{m \geq 0} \mathbb{A}(\Sigma_m)$  properly, a fact explained and amended by Ochoa in [32]. An explicit construction of the  $\beta$ -operations, using the language of polynomial operations, is given by Vallejo in [44], and he extended the definition of a  $\beta$ -ring by a unitality condition in [46].

Lastly, the survey article [20] of Guillot provides more details on the history of  $\beta$ -rings and their connection to  $\lambda$ -rings, as well as showing that stable cohomotopy is an example of a  $\beta$ -ring. Moreover, there an additivity condition is added to the notion of  $\beta$ -rings.

A  $\beta$ -ring encodes power operations indexed by Burnside rings. This is formalized by a certain operator ring:

**Definition 3.17** We define

$$\mathbf{B} = \bigoplus_{m \geq 0} \mathbb{A}(\Sigma_m).$$

We endow this abelian group with a commutative multiplication via

$$x \cdot y = \text{tr}_{\Sigma_k \times \Sigma_l}^{\Sigma_{k+l}}(x \times y)$$

for  $x \in \mathbb{A}(\Sigma_k)$  and  $y \in \mathbb{A}(\Sigma_l)$ . This defines a ring structure on  $\mathbf{B}$ . Moreover, we define an operation  $*$  on  $\mathbf{B}$ . Let  $x \in \mathbb{A}(\Sigma_k)$  and  $y_i \in \mathbb{A}(\Sigma_{l_i})$  for  $1 \leq i \leq n$ . Then

$$(3.18) \quad x * (y_1 + \dots + y_n) = \sum_{(k)} \text{tr}_{\times \Sigma_{k_i} \wr \Sigma_{l_i}}^{\Sigma_{(k) \cdot (l)}} \left( \prod_{i=1}^n P^{k_i}(y_i) \cdot ((\times \Sigma_{k_i} \wr p_{\Sigma_{l_i}})^* \Phi_{(k)}^*)(x) \right),$$

where we sum over all partitions  $(k) = (k_i)_{i=1, \dots, n}$  of  $k$ , we define  $(k) \cdot (l) := \sum_{i=1}^n k_i l_i$  and the transfer is along the monomorphisms  $\Psi_{k_i, l_i}$  and  $\Phi_{(k_i l_i)_i}$  [35, (2.2.5) and (2.2.6)]. In the map  $\times \Sigma_{k_i} \wr p_{\Sigma_{l_i}} : \times \Sigma_{k_i} \wr \Sigma_{l_i} \rightarrow \times \Sigma_{k_i}$ , the product runs from  $i = 1$  to  $n$  and the map  $p_{\Sigma_{l_i}} : \Sigma_{l_i} \rightarrow e$  is the unique map to the trivial group.

The operation  $*$  is additive in the first component and can be extended linearly to give a map  $* : \mathbf{B} \times \mathbf{B} \rightarrow \mathbf{B}$ .

The operation  $*$  is sometimes called plethysm, for example in [32] and in [23] for the representation ring instead of the Burnside ring. It makes  $\mathbf{B}$  into an operator ring

by [46, Theorem 1.11]. We then define a  $\beta$ -ring as an *operator module* over  $\mathbf{B}$ . The following definition is given in [46, Definition 1.12]. Note that here and in the following, we denote by  $\text{Map}$  the collection of maps of *sets*. In contrast,  $\text{Hom}$  in the following denotes ring homomorphisms.

**Definition 3.19** A  $\beta$ -ring is a commutative ring  $A$  with a map  $\vartheta: \mathbf{B} \rightarrow \text{Map}(A, A)$  such that for all  $a \in A$  and  $x, y \in \mathbf{B}$ :

- (i)  $\vartheta(x + y) = \vartheta(x) + \vartheta(y)$ .
- (ii)  $\vartheta(x \cdot y) = \vartheta(x) \cdot \vartheta(y)$ .
- (iii)  $\vartheta(x * y) = \vartheta(x) \circ \vartheta(y)$ .
- (iv)  $\vartheta(1)(a) = 1$  for the multiplicative unit  $1 \in \mathbb{A}(\Sigma_0) \subset \mathbf{B}$  on the left and  $1 \in A$  on the right.
- (v)  $\vartheta(e) = \text{id}_A$ , where  $e = 1 \in \mathbb{A}(\Sigma_1) \subset \mathbf{B}$  is a unit for the operation  $*$ .

In the first two statements, we use the pointwise ring structure on  $\text{Map}(A, A)$ .

For the construction of a  $\beta$ -ring structure on  $R(G)$  for a global power functor  $R$ , we need an additional structure in the form of a pairing with the Burnside ring.

**Definition 3.20** Let  $R$  be a global power functor. Maps

$$\langle \_, \_ \rangle_{K,G}: R(K \times G) \times \mathbb{A}(K) \rightarrow R(G),$$

defined for all compact Lie groups  $K$  and  $G$ , are called  $\mathbb{A}$ -deflations if the maps  $\langle \_, \_ \rangle_{K,G}$  are biadditive and satisfy:

- (i)  $\langle (K \times \alpha)^* r, x \rangle_{K,G} = \alpha^* \langle r, x \rangle_{K,L}$  for any continuous group homomorphism  $\alpha: G \rightarrow L$ .
- (ii)  $\langle \text{tr}_{L \times G}^{K \times G} r, x \rangle_{K,G} = \langle r, \text{res}_L^K x \rangle_{L,G}$  for any closed subgroup  $L \subset K$ , and the reversed relation also holds.
- (iii)  $\langle (\alpha \times G)^* r, \alpha^* x \rangle = \langle r, x \rangle$  for  $r \in R(K \times G)$ ,  $x \in \mathbb{A}(K)$  and a surjective group homomorphism  $\alpha: L \rightarrow K$ .
- (iv)  $\langle r, 1 \rangle = r$  for all  $r \in R(G)$  and  $1 \in \mathbb{A}(e)$ .
- (v)  $\langle r \cdot s, x \cdot y \rangle_{K,G} = \langle r, x \rangle_{K,G} \cdot \langle s, y \rangle_{K,G}$  for all  $r, s \in R(K \times G)$  and  $x, y \in \mathbb{A}(K)$ .
- (vi)  $\langle r \cdot \text{pr}_K^* y, x \rangle_{K,G} = \langle r, y \cdot x \rangle_{K,G}$  for all  $r \in R(K \times G)$  and  $x, y \in \mathbb{A}(K)$ .

(vii)  $\langle (\delta_n^G)^* P^n \langle r, y \rangle, x \rangle = \langle (\delta_n^{\Sigma_k, G})^* (P^n(r) \cdot (\Sigma_n \wr \text{pr}_{\Sigma_k})^* P^n(y)), (\Sigma_n \wr p_{\Sigma_k})^* x \rangle$   
 for all  $r \in R(\Sigma_k \times G)$ ,  $y \in \mathbb{A}(\Sigma_k)$  and  $x \in \mathbb{A}(\Sigma_n)$ . Here, we considered the diagonal inclusion

(3.21)  $\delta_n^G : \Sigma_n \times G \rightarrow \Sigma_n \wr G$  given by  $(\sigma, g) \mapsto (\sigma; g, \dots, g)$   
 and the relative version  $\delta_n^{\Sigma_k, G} : (\Sigma_n \wr \Sigma_k) \times G \rightarrow \Sigma_n \wr (\Sigma_k \times G)$ .

We denote by  $\mathcal{G}IPow_{\mathbb{A}\text{-def}}$  the category of global power functors with  $\mathbb{A}$ -deflations and morphisms of global power functors compatible with the pairing  $\langle \_ , \_ \rangle$ .

**Remark 3.22** In practice, such a deflation pairing on a global power functor often arises from actual deflations, ie maps  $\varphi_* : R(G) \rightarrow R(K)$  for surjective group homomorphisms  $\varphi : G \rightarrow K$ . These satisfy certain relations, as exhibited in [12] for finite groups, and packaged in [47, Chapter 8] in the notion of a globally defined Mackey functor. The compatibility of the deflations with the power operations also explains the compatibility condition of the above deflation pairing with the power operations.

In the following, we only use the deflation pairing on a global power functor  $R$ .

**Lemma 3.23** For the left-induced global power functor  $\mathbb{A} \otimes C$  for a ring  $C$ , the composition

$$\mathbb{A}(K \times G) \otimes C \otimes \mathbb{A}(K) \xrightarrow{\times} \mathbb{A}(K \times G \times K) \otimes C \xrightarrow{\Delta_K^*} \mathbb{A}(K \times G) \otimes C \xrightarrow{(\text{pr}_G)_*} \mathbb{A}(G) \otimes C$$

defines an  $\mathbb{A}$ -deflation. Here  $\Delta_K : K \times G \rightarrow K \times G \times K$  is the diagonal of  $K$ , and  $(\text{pr}_G)_*$  denotes the deflation along  $\text{pr}_G : K \times G \rightarrow G$  present in the Burnside rings.

We omit the calculations of the properties of this pairing. We recall however the definition of the deflations, since they are the main part of this construction:

**Construction 3.24** In the case of finite groups, the deflations in the Burnside ring global functor are described as follows. Let  $f : G \rightarrow K$  be a group homomorphism, and  $X$  be a finite  $G$ -set. Then we define  $f_*(X) = K \times_G X = (K \times X)/G$ , where we identify  $(k \cdot f(g), x)$  with  $(k, gx)$  for all  $g \in G$ , and consider  $f_*(X)$  as a  $K$ -set by the multiplication on the  $K$ -factor. This defines a map  $\mathbb{A}(G) \rightarrow \mathbb{A}(K)$ . For the general case of a morphism  $G \rightarrow K$  of compact Lie groups, this construction is generalized in [14, Proposition 20; 16, Proposition IV.2.18].

**Remark 3.25** Another main example of a global functor with  $\mathbb{A}$ -deflations is stable cohomotopy  $\pi^0(X)$  for a based space  $X$ . By Remark 2.15, we have restricted power operations on the equivariant stable cohomotopy defined by

$$\pi_G^0(X) \xrightarrow{P^m} \pi_{\Sigma_m \wr G}^0(X^m) \xrightarrow{(\delta_m^G)^*} \pi_{\Sigma_m \times G}^0(X^m) \xrightarrow{\Delta^*} \pi_{\Sigma_m \times G}^0(X).$$

As explained below, these restricted power operations, using  $\delta_m^G$ , are also used to obtain the  $\beta$ -operations. Hence, the theory developed below is also applicable to equivariant stable cohomotopy.

Moreover, we obtain a pairing  $\pi_{K \times G}^0(X) \times \mathbb{A}(K) \rightarrow \pi_G^0(X)$  by generalizing the definition for  $\mathbb{A}$  to an arbitrary base. We again restrict to finite groups. We may define operations on stable cohomotopy by defining additive (or only polynomial) operations on the monoid  $\text{Cov}^+(X)$  of isomorphism classes of (equivariant) coverings over  $X$ . Nonequivariantly this is [45, Theorem 2.4], building on the observations by Segal in [38]. The main observation for this is that stable cohomotopy can be seen as the group completion of the monoid of coverings over  $X$ , which is a consequence of the Barratt–Priddy–Quillen theorem [6; 33; 37]. The equivariant statement holds analogously, using the equivariant Barratt–Priddy–Quillen theorem established in [36; 21; 7].

Hence, let  $E \rightarrow X$  be a  $(K \times G)$ -equivariant covering of  $X$  and  $T$  be a finite  $K$ -set. Define  $\langle E \rightarrow X, T \rangle$  to be  $E \times_K T$ , which is a  $G$ -equivariant covering. One can check that this is biadditive and induces a pairing  $\pi_{K \times G}^0(X) \times \mathbb{A}(K) \rightarrow \pi_G^0(X)$  as required.

Suppose now that  $R$  is a global power functor with  $\mathbb{A}$ -deflations. We are ultimately interested in a pairing  $R(\Sigma_m \wr G) \otimes \mathbb{A}(\Sigma_m) \rightarrow R(G)$ , hence we use the morphism  $\delta_m^G: \Sigma_m \times G \rightarrow \Sigma_m \wr G$  from (3.21) to define

$$R(\Sigma_m \wr G) \otimes \mathbb{A}(\Sigma_m) \xrightarrow{(\delta_m^G)^* \otimes \text{id}} R(\Sigma_m \times G) \otimes \mathbb{A}(\Sigma_m) \xrightarrow{(\cdot, \cdot)_{\Sigma_m, G}} R(G).$$

Using these definitions, for any global power functor  $R$  with  $\mathbb{A}$ -deflations we can define a morphism

(3.26) 
$$D_G: \exp(R; G) \rightarrow \text{Map}(\mathbf{B}, R(G)),$$

$$x = (x_n)_{n \geq 0} \mapsto \left( (y_n) \mapsto \sum_{n \geq 0} \langle (\delta_n^G)^* x_n, y_n \rangle \right).$$

**Proposition 3.27** *Let  $R$  be a global power functor with  $\mathbb{A}$ -deflations and  $G$  be a compact Lie group. Then for any  $x \in \exp(R; G)$ , the morphism  $D_G(x)$  is a ring homomorphism.*

**Proof** Additivity of the morphism  $D_G(x): \mathbf{B} \rightarrow R(G)$  is clear from biadditivity of the pairing  $\langle \_, \_ \rangle$ . Moreover, let  $1 \in \mathbb{A}(\Sigma_0)$  be the multiplicative unit of  $\mathbf{B}$ . Then we calculate

$$D_G(x)(1) = \langle (\delta_0^G)^* x_0, 1 \rangle = p_G^*(x_0) = 1,$$

using that exponential sequences have as zeroth term the unit of  $R(e)$  and that  $\delta_0^G$  is the unique map  $p_G: G \rightarrow e$ .

Now, we check that the map  $D_G(x)$  is multiplicative. Recall that the product on  $\mathbf{B}$  is the transfer product from Definition 3.17. For two elements  $y, z \in \mathbf{B}$ , we have

$$\begin{aligned} D_G(x)(y \cdot z) &= \sum_{n \geq 0} \left\langle (\delta_n^G)^* x_n, \sum_{k+l=n} \text{tr}_{\Sigma_k \times \Sigma_l}^{\Sigma_{k+l}}(y_k \times z_l) \right\rangle \\ &= \sum_{k,l \geq 0} \langle \text{res}_{\Sigma_k \times \Sigma_l \times G}^{\Sigma_{k+l} \times G} (\delta_{k+l}^G)^* x_{k+l}, y_k \times z_l \rangle \\ &= \sum_{k,l \geq 0} \langle \Delta_G^* ((\delta_k^G)^* x_k \times (\delta_l^G)^* x_l), y_k \times z_l \rangle \\ &= \sum_{k,l \geq 0} \langle (\delta_k^G)^* x_k, y_k \rangle \cdot \langle (\delta_l^G)^* x_l, z_l \rangle = D_G(x)(y) \cdot D_G(x)(z). \end{aligned}$$

Here, we use that  $\delta_{k+l}^G \circ (\Phi_{k,l} \times G)$  and  $\Phi_{k,l}^G \circ (\delta_k^G \times \delta_l^G) \circ \Delta_G: \Sigma_k \times \Sigma_l \times G \rightarrow \Sigma_{k+l} \wr G$  agree and that the sequence  $x$  is exponential. □

Using this morphism, we can now study how any global power functor  $R$  with  $\mathbb{A}$ -deflations induces the structure of a  $\beta$ -ring on  $R(G)$ :

**Construction 3.28** Let  $R$  be a global power functor with  $\mathbb{A}$ -deflations and let  $G$  be a compact Lie group. Then we define

$$\bar{\vartheta}_G: R(G) \xrightarrow{P} \text{exp}(R; G) \xrightarrow{D_G} \text{Hom}_{\text{Rings}}(\mathbf{B}, R(G)),$$

where  $P$  denotes the power operation on  $R$ .

This map is adjoint to a map

$$\vartheta: \mathbf{B} \rightarrow \text{Map}(R(G), R(G)).$$

**Proposition 3.29** *The map  $\vartheta$  makes  $R(G)$  into a  $\beta$ -ring.*

**Proof** This proof is a lengthy calculation, using the properties of the global power operations and the pairing  $\langle \_, \_ \rangle$ . An essentially similar calculation in the case  $R = \mathbb{A}$

has been carried out in [34, Theorem 2], where the relations are only checked for some additive generators of  $\mathbf{B}$ , and in [46, Corollary 1.16], where these calculations are extended to the entirety of  $\mathbf{B}$  using the theory of polynomial operations. Also, a similar calculation can be found in [32].  $\square$

**Corollary 3.30** *The rings  $\mathbb{A}(G)$  and  $\pi_G^0(X)$  for any based space  $X$  carry the structure of  $\beta$ -rings.*

**Corollary 3.31** *Let  $C$  be a commutative ring. If  $\mathbb{A} \otimes C$  supports the structure of a global power functor, then  $\mathbb{A}(G) \otimes C$  inherits the structure of a  $\beta$ -ring.*

Note that the  $\beta$ -ring structures on Burnside rings and stable cohomotopy rings are already known in the classical literature. However, the above definition of the structure map  $\vartheta$  illustrates how this structure can be obtained from a more naturally arising structure, namely from a global power functor structure on  $R$  together with deflations. We hope that this allows for a more insightful picture of  $\beta$ -rings.

The definition of a  $\beta$ -ring only incorporates relations between the different operations  $\vartheta(x)$  on  $R(G)$ , indexed by  $x \in \mathbf{B}$ . It does not provide any compatibility of these operations with the ring structure on  $R(G)$ . We now add a condition of “external” additivity, due to [20]:

**Definition 3.32** We define

$$\mathbf{B}^2 = \bigoplus_{p,q \geq 0} \mathbb{A}(\Sigma_p \times \Sigma_q).$$

This has a ring structure analogous to that of  $\mathbf{B}$ , given by  $x \cdot y = \text{tr}_{\Sigma_p \times \Sigma_q \times \Sigma_r \times \Sigma_s}^{\Sigma_{p+r} \times \Sigma_{q+s}}(x \times y)$  for  $x \in \mathbb{A}(\Sigma_p \times \Sigma_q)$  and  $y \in \mathbb{A}(\Sigma_r \times \Sigma_s)$ . Moreover, we have maps

$$\Phi: \mathbf{B} \rightarrow \mathbf{B}^2 \quad \text{given by } x \mapsto \sum_{p+q=m} \Phi_{p,q}^* x \text{ for } x \in \mathbb{A}(\Sigma_m)$$

and

$$\times: \mathbf{B} \otimes \mathbf{B} \rightarrow \mathbf{B}^2 \quad \text{given by } x \otimes y \mapsto x \times y \text{ for } x \in \mathbb{A}(\Sigma_p) \text{ and } y \in \mathbb{A}(\Sigma_q).$$

Additivity is then expressed by a morphism  $\vartheta^2: \mathbf{B}^2 \rightarrow \text{Map}(A \times A, A)$  analogous to  $\vartheta$ , using the map  $\Phi$ .

**Definition 3.33** An additive  $\beta$ -ring is a commutative ring  $A$  together with maps  $\vartheta: \mathbf{B} \rightarrow \text{Map}(A, A)$  and  $\vartheta^2: \mathbf{B}^2 \rightarrow \text{Map}(A \times A, A)$  such that  $(A, \vartheta)$  is a  $\beta$ -ring and:

- (i)  $\vartheta^2(x \times y)(c, d) = \vartheta(x)(c) \cdot \vartheta(y)(d)$  for all  $x, y \in \mathbf{B}$  and all  $c, d \in A$ .
- (ii)  $\vartheta(x)(c + d) = \vartheta^2(\Phi x)(c, d)$  for all  $x \in \mathbf{B}$  and  $c, d \in A$ .

We construct such a map  $\vartheta^2$  for  $R(G)$  when  $R$  a global power functor with  $\mathbb{A}$ -deflations.

**Construction 3.34** Let  $R$  be a global power functor with  $\mathbb{A}$ -deflations. Then we define

$$D^2: \exp(R; G) \times \exp(R; G) \rightarrow \text{Hom}(\mathbf{B}^2, R(G)),$$

$$(x, y) \mapsto \left( z \mapsto \sum_{p,q \geq 0} \langle (\delta_{p,q}^G)^*(x_p \times y_q), z_{p,q} \rangle \right),$$

where  $\delta_{p,q}^G = (\delta_p^G \times \delta_q^G) \circ \Delta_G: \Sigma_p \times \Sigma_q \times G \rightarrow \Sigma_p \wr G \times \Sigma_q \wr G$  is the diagonal on  $G$ . This in fact takes values in ring homomorphisms by a similar argument to Proposition 3.27.

Moreover, we define

$$\bar{\vartheta}^2: R(G) \times R(G) \xrightarrow{P \times P} \exp(R; G) \times \exp(R; G) \xrightarrow{D^2} \text{Hom}(\mathbf{B}^2, R(G)),$$

and denote the morphism adjoint to  $\bar{\vartheta}^2$  by

$$\vartheta^2: \mathbf{B}^2 \rightarrow \text{Map}(R(G) \times R(G), R(G)).$$

**Proposition 3.35** *The morphisms  $\vartheta$  and  $\vartheta^2$  make  $R(G)$  into an additive  $\beta$ -ring.*

**Proof** We only prove part (ii) from Definition 3.33, since the first assertion is an easy calculation, using the description  $\delta_{p,q}^G = (\delta_p^G \times \delta_q^G) \circ \Delta_G$ .

We thus calculate

$$\begin{aligned} \vartheta(x)(c + d) &= \sum_{m \geq 0} \langle (\delta_m^G)^* P^m(c + d), x_m \rangle \\ &= \sum_{m \geq 0} \left\langle (\delta_m^G)^* \sum_{p+q=m} \text{tr}_{\Sigma_p \wr G \times \Sigma_q \wr G}^{\Sigma_{p+q} \wr G} (P^p(c) \times P^q(d)), x_m \right\rangle \\ &= \sum_{p,q \geq 0} \langle \text{tr}_{\Sigma_p \times \Sigma_q \times G}^{\Sigma_{p+q} \times G} (\delta_{p,q}^G)^* (P^p(c) \times P^q(d)), x_{p+q} \rangle \\ &= \sum_{p,q \geq 0} \langle (\delta_{p,q}^G)^* (P^p(c) \times P^q(d)), \Phi_{p,q}^* x_{p+q} \rangle \\ &= D^2(P(c) \times P(d))(\Phi x) = \vartheta^2(\Phi x)(c, d). \end{aligned}$$

Here, in the third line, we use the observation that there is only one double coset in

$$\Sigma_{p+q} \times G \backslash \Sigma_{p+q} \wr G / \Sigma_p \wr G \times \Sigma_q \wr G,$$

and hence the double coset formula for  $(\delta_{p+q}^G)^* \text{tr}_{\Sigma_p \wr G \times \Sigma_q \wr G}^{\Sigma_{p+q} \wr G}$  consists of a single summand. □

Finally, we consider in which sense this construction is functorial. We can study functoriality both in the global power functor  $R$  and in the compact Lie group  $G$ .

**Definition 3.36** Let  $A$  and  $A'$  be additive  $\beta$ -rings with structure morphisms  $\vartheta$  and  $\vartheta^2$ , and  $\vartheta'$  and  $\vartheta'^2$ , respectively. Then a morphism  $f: A \rightarrow A'$  of  $\beta$ -rings is a ring homomorphism  $f$  such that the relations

$$f(\vartheta(x)(a)) = \vartheta'(x)(f(a)) \quad \text{and} \quad f(\vartheta^2(y)(a_1, a_2)) = \vartheta'^2(y)(f(a_1), f(a_2))$$

hold for all  $x \in \mathbf{B}$ ,  $y \in \mathbf{B}^2$  and  $a, a_1, a_2 \in A$ .

**Proposition 3.37** (i) Let  $G$  be a compact Lie group and  $f: R \rightarrow S$  be a morphism of global power functors with  $\mathbb{A}$ -deflations. Then  $f(G)$  is a morphism between the  $\beta$ -rings  $R(G)$  and  $S(G)$ .

(ii) Let  $\varphi: K \rightarrow G$  be a homomorphism of compact Lie groups and  $R$  be a global power functor with  $\mathbb{A}$ -deflations. Then  $\varphi^*$  is a morphism between the  $\beta$ -rings  $R(G)$  and  $R(K)$ .

**Proof** We only check the compatibility with  $\vartheta$ , the calculations for  $\vartheta^2$  are similar.

For the first assertion, we calculate, for  $x \in \mathbf{B}$  and  $b \in R(G)$ ,

$$\begin{aligned} \vartheta_S(x)(f(G)(b)) &= D_G(P_S(f(G)(b)))(x) = \sum_{n \geq 0} \langle (\delta_n^G)^* P_S^n(f(G)(b)), x_n \rangle \\ &= \sum_{n \geq 0} \langle f(\Sigma_n \times G)(\delta_n^G)^* P_R^n(b), x_n \rangle \\ &= \sum_{n \geq 0} f(G) \langle (\delta_n^G)^* P_R^n(b), x_n \rangle = f(G)(\vartheta_R(x)(b)). \end{aligned}$$

For the second assertion, we calculate for  $x \in \mathbf{B}$  and  $c \in R(G)$ :

$$\begin{aligned} \vartheta_K(x)(\varphi^*(c)) &= \sum_{n \geq 0} \langle (\delta_n^K)^* P^n(\varphi^*(c)), x_n \rangle = \sum_{n \geq 0} \langle (\delta_n^K)^* (\Sigma_n \wr \varphi)^* P^n(c), x_n \rangle \\ &= \sum_{n \geq 0} \langle (\Sigma_n \times \varphi)^* (\delta_n^G)^* P^n(c), x_n \rangle \\ &= \sum_{n \geq 0} \varphi^* \langle (\delta_n^G)^* P^n(c), x_n \rangle = \varphi^* \vartheta_G(x)(c). \end{aligned}$$

The corresponding calculations with  $\vartheta^2$  work analogously. □



Thus we have proven the following result, where we denote by  $\text{Rep}$  the category of compact Lie groups and conjugacy classes of continuous group homomorphisms, and by  $\beta$ -Rings the category of additive  $\beta$ -rings. Moreover, we denote by  $\mathcal{GIPow}_{\mathbb{A}\text{-defl}}$  the category of global power functors with  $\mathbb{A}$ -deflations.

**Theorem 3.38** *The assignment  $(G, R) \mapsto R(G)$  extends to a functor*

$$\text{ev}: \text{Rep}^{\text{op}} \times \mathcal{GIPow}_{\mathbb{A}\text{-defl}} \rightarrow \beta\text{-Rings},$$

*which sends a conjugacy class of a morphism of compact Lie groups to the corresponding restriction.*

In this theorem, we only treat restrictions. In fact, transfers do not induce morphisms of  $\beta$ -rings. The reason is that transfers do not commute with the morphism  $(\delta_n^G)^*$ .

To illustrate the theory of  $\beta$ -rings, we calculate the  $\beta$ -operations in one example.

**Example 3.39** We now apply our theory to the global power functor  $\mathbb{A}$ , where  $C = \mathbb{Z} = \mathbb{A}(e)$ . Then we obtain  $\beta$ -ring structures on the Burnside rings  $\mathbb{A}(G)$  for all compact Lie groups  $G$ . The operations here are given as follows:

For the element  $x = \Sigma_n/H \in \mathbf{B} = \bigoplus_{n \geq 0} \mathbb{A}(\Sigma_n)$ , we obtain for finite  $G$  and a finite  $G$ -set  $X$  the formula

$$\vartheta_H(X) := \vartheta_{\Sigma_n/H}(X) = \langle P^n(X), \Sigma_n/H \rangle = \Sigma_n/H \times_{\Sigma_n} X^n = X^n/H,$$

where we consider the resulting set as a  $G$ -set. This formula agrees with the one from [44] and generalizes to compact Lie groups as shown in [34]. Thus, in this case, we obtain the classical  $\beta$ -ring structure on  $\mathbb{A}(G)$ , using our abstract definition. Also the iterated operations  $\vartheta^2$  used for additivity agree with those defined in [20, Example 3.2]. In fact, we have

$$\vartheta_H^2(X, Y) = (X^p \times Y^q)/H$$

for a finite group  $G$ , finite  $G$ -sets  $X$  and  $Y$  and  $H \subset \Sigma_p \times \Sigma_q$ .

The above construction of  $\beta$ -ring structures on  $R(G)$  highlights the importance of a global point of view. Theorem 3.38 shows that the notion of a global power functor with  $\mathbb{A}$ -deflations encodes compatible  $\beta$ -ring structures for all compact Lie groups at

once. In this way, we may approach the still rather mysterious theory of  $\beta$ -rings from the direction of the well-structured global power functors.

The comparison of Theorem 3.38 is not perfect, however. It remains open to what extent we can represent all  $\beta$ -rings by global power functors, for example. In general, the condition of having global power operations on a ring is stronger than admitting a  $\beta$ -ring structure. Also, we require no multiplicative behaviour of the operations  $\vartheta$ , whereas the power operations of a global power functor are multiplicative. In the face of the complications posed in the analysis of  $\beta$ -rings, starting with finding a feasible definition, it seems sensible to propose that the notion of global power functors is the more fundamental one.

### Appendix Transferring monads under lax functors

In Section 2.2, we studied two lifting theorems for functors between algebras over the derived symmetric algebra monad in the global and stable homotopy categories. To separate the homotopy theoretic properties needed to provide the liftings from the formal background in monad theory, it is convenient to use the language of 2-categories. We also treat aspects of double categories, which we use when we encounter both left and right derived functors. For the theory of 2-categories, we refer to [11, Chapter 7; 25; 43] and for the theory of double categories, we refer to [25; 39].

**Definition A.1** A 2-category is a category enriched in the category  $\text{Cat}$  of categories, and a double category is a category object in  $\text{Cat}$ .

Thus, explicitly, a 2-category consists of classes of objects, morphisms and transformations, where we have a horizontal composition  $\star$  and a vertical composition  $\circ$  of transformations, and composition of morphisms is strictly unital and associative. For horizontal and vertical composition, we use the conventions

$$(\eta: g \rightarrow g') \star (\vartheta: f \rightarrow f') = X \begin{array}{ccc} \xrightarrow{f} & & \xrightarrow{g} \\ \Downarrow \vartheta & & \Downarrow \eta \\ \xrightarrow{f'} & & \xrightarrow{g'} \end{array} Y \quad Z$$

and

$$(\eta: g \rightarrow h) \circ (\vartheta: f \rightarrow g) = X \begin{array}{ccc} & \xrightarrow{f} & \\ & \Downarrow \vartheta & \\ & \xrightarrow{g} & \\ \Downarrow \eta & & \\ & \xrightarrow{h} & \end{array} Y$$

A double category consists of a class of objects, classes of horizontal and vertical morphisms each being part of a category with common objects, and a class of transformations

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & \Downarrow & \downarrow \\ Z & \longrightarrow & W \end{array}$$

also called squares or 2-cells. Transformations can be composed both horizontally and vertically, and all possible orders of composition agree. We denote horizontal composition by  $\boxtimes$  and vertical composition by  $\boxplus$ . Note that for any double category  $\mathbf{C}$ , we obtain two 2-categories  $\mathcal{V}(\mathbf{C})$  and  $\mathcal{H}(\mathbf{C})$  by considering only vertical morphisms and 2-cells with identities as horizontal morphisms, or similarly only considering horizontal morphisms.

For these notions of higher categories, there exist various versions of functors and natural transformations between them. We need:

**Definition A.2** Let  $\mathcal{C}$  and  $\mathcal{D}$  be 2-categories. A lax 2-functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  consists of the following data:

- (i) the assignments  $X \mapsto F(X)$ ,  $(f: X \rightarrow Y) \mapsto (F(f): F(X) \rightarrow F(Y))$  and  $(\eta: f \rightarrow g) \mapsto (F(\eta): F(f) \rightarrow F(g))$  of objects, morphisms and transformations, and
- (ii) the transformations  $\alpha_X: \text{id}_{F(X)} \rightarrow F(\text{id}_X)$  and  $\mu_{g,f}: F(g) \circ F(f) \rightarrow F(gf)$  for any object  $X$  and any pair  $(g, f)$  of composable morphisms in  $\mathcal{C}$ .

These have to satisfy the compatibility conditions given in [11, Definition 7.5.1].

**Definition A.3** Let  $F, G: \mathcal{C} \rightarrow \mathcal{D}$  be two lax functors between 2-categories. A lax natural transformation  $\eta: F \rightarrow G$  between  $F$  and  $G$  consists of assignments

$$X \mapsto (\eta_X: F(X) \rightarrow G(X)) \quad \text{and} \quad (f: X \rightarrow Y) \mapsto \left( \begin{array}{ccc} F(X) & \xrightarrow{\eta_X} & G(X) \\ Ff \downarrow & \Downarrow_{\eta_f} & \downarrow Gf \\ F(Y) & \xrightarrow{\eta_Y} & G(Y) \end{array} \right)$$

such that the compatibility conditions given in [11, Definition 7.5.2] are satisfied.

For two lax transformations  $\eta: F \rightarrow G$  and  $\theta: G \rightarrow H$  between lax 2-functors, the composite is given by sending  $X$  to the morphism  $\theta_X \circ \eta_X$ , and a morphism  $f: X \rightarrow Y$  to the transformation

$$\begin{array}{ccccc}
 F(X) & \xrightarrow{\eta_X} & G(X) & \xrightarrow{\theta_X} & H(X) \\
 Ff \downarrow & \swarrow_{\eta_f} & \downarrow Gf & \swarrow_{\theta_f} & \downarrow Hf \\
 F(Y) & \xrightarrow{\eta_Y} & G(Y) & \xrightarrow{\theta_Y} & H(Y)
 \end{array}$$

We now relate these notions to the theory of monads. First note that the definition of a monad can be given in any 2-category, generalizing an endofunctor  $T: \mathcal{C} \rightarrow \mathcal{C}$  to an endomorphism  $T: X \rightarrow X$  of an object  $X$ , and the multiplication and unitality natural transformations  $\mu: TT \rightarrow T$  and  $\eta: \text{Id} \rightarrow T$  to corresponding transformations. Moreover, we can consider morphisms between such monads, compare [42, Section 1].

**Definition A.4** Let  $\mathcal{C}$  be a 2-category and  $(P, \mu, \eta)$  and  $(Q, \nu, \varepsilon)$  be monads on objects  $X$  and  $Y$  of  $\mathcal{C}$ , respectively. A (lax) monad morphism is a pair  $(F, \rho)$ , consisting of a morphism  $F: X \rightarrow Y$  and a transformation  $\rho: QF \rightarrow FP$ , such that the diagrams

$$\begin{array}{ccc}
 F \xrightarrow{\varepsilon F} QF & & QQF \xrightarrow{Q\rho} QFP \\
 \searrow F\eta & \text{and} & \downarrow \nu F \\
 & & QF \xrightarrow{\rho} FP \\
 & & \downarrow \rho P \\
 & & FFP \\
 & & \downarrow F\mu \\
 & & FP
 \end{array}$$

commute. Let  $(F, \rho), (G, \sigma): X \rightarrow Y$  be two monad functors between  $P$  and  $Q$ . Then a monadic transformation between  $F$  and  $G$  is a transformation  $\theta: F \rightarrow G$  such that  $\theta P \circ \rho = \sigma \circ Q\theta$  as transformations  $QF \rightarrow GP$ .

Note that if a 2-category  $\mathcal{C}$  admits a construction of algebras, ie a right adjoint to the inclusion of  $\mathcal{C}$  into the category of monads in  $\mathcal{C}$ , then a monad morphism  $(F, \rho): P \rightarrow Q$  induces a morphism between the corresponding objects of algebras over  $P$  and  $Q$ . In particular, a monad functor in  $\text{Cat}$  induces a functor between the categories of algebras, and a monadic transformation a transformation between the induced functors.

The following observation goes back to [8, Definition 5.4.1]:

**Lemma A.5** Let  $\mathcal{C}$  be a 2-category, and let  $1$  be the terminal 2-category with a single object  $*$ , its identity morphism and the identity natural transformation. Then,

the category of lax 2-functors  $1 \rightarrow \mathcal{C}$  and lax natural transformations, and the category of monads and lax monad morphisms in  $\mathcal{C}$  are isomorphic via the functor

$$(T : 1 \rightarrow \mathcal{C}) \mapsto (T(\text{id}_*) : T(*) \rightarrow T(*), \mu : T(\text{id}_*) \circ T(\text{id}_*) \rightarrow T(\text{id}_*), \eta : \text{id}_{T(*)} \rightarrow T(\text{id}_*)),$$

$$(\rho : S \rightarrow T) \mapsto \left( \begin{array}{ccc} T(*) & \xrightarrow{\rho_*} & T(*) \\ \rho_* : S(*) \rightarrow T(*) & \rho(\text{id}_*) : S(\text{id}_*) \downarrow & \not\cong \downarrow T(\text{id}_*) \\ S(*) & \xrightarrow{\rho_*} & T(*) \end{array} \right).$$

The proof is an easy translation of the corresponding properties.

Using this description, we see that monads are preserved under any lax 2-functor, and lax monad morphisms are preserved if we moreover assume that some of the structure maps of a lax functor are invertible.

**Corollary A.6** *Let  $\mathcal{C}$  and  $\mathcal{D}$  be 2-categories and let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a lax 2-functor. Let  $(T : X \rightarrow X, \nu : T \circ T \rightarrow T, \varepsilon : \text{id}_X \rightarrow T)$  be a monad in  $\mathcal{C}$ . Then*

$$(F(T) : F(X) \rightarrow F(X), F(\nu) \circ \mu_{T,T} : FT \circ FT \rightarrow FT, F(\varepsilon) \circ \eta_X : \text{id}_{FX} \rightarrow FT)$$

*is a monad in  $\mathcal{D}$ .*

Moreover, let  $S$  and  $T$  be two monads in  $\mathcal{C}$  on objects  $X$  and  $Y$  respectively, let  $(f : X \rightarrow Y, \rho : Tf \rightarrow fS)$  be a lax monad morphism, and assume that  $\mu_{f,S}$  is invertible. Then  $(Ff : FX \rightarrow FY, \mu_{f,S}^{-1} \circ F(\rho) \circ \mu_{T,f} : FT \circ Ff \rightarrow Ff \circ FS)$  is a lax monad morphism between  $FS$  and  $FT$ .

**Proof** The first part of this corollary follows directly from the above lemma. We can consider the monad  $T$  in  $\mathcal{C}$  as a lax 2-functor  $T : 1 \rightarrow \mathcal{C}$ . Then, the composition  $F \circ T : 1 \rightarrow \mathcal{D}$  is a lax 2-functor with coherence morphisms the composites of the coherence morphisms of  $F$  and  $T$ . Thus,  $FT$  is a monad in  $\mathcal{D}$ , and the structure of the monad is exactly given by the described morphism.

For the second part, we note that the above lemma shows that a lax monad morphism from  $S$  to  $T$  is the same as a lax natural transformation between the corresponding lax 2-functors  $1 \rightarrow \mathcal{C}$ . It is an easy argument that a lax 2-functor with invertible transformation  $\mu_{f,S}$  preserves such a transformation. □

We now consider the double categorical context. This is used in Section 2.2 in order to handle the occurrence of both right and left derived functors. These different types of functors can conveniently be handled by assigning them as vertical and horizontal morphisms of a double category, respectively.

In a double category  $\mathbf{C}$ , we use a similar formalism as for 2-categories to consider whether a corresponding notion of weak double functor preserves monads and morphisms between them. We thus first define the appropriate notion of a weak double functor.

**Definition A.7** Let  $\mathbf{C}$  and  $\mathbf{D}$  be double categories. A lax–oplax double functor  $F: \mathbf{C} \rightarrow \mathbf{D}$  consists of assignments of objects, horizontal 1–cells, vertical 1–cells and 2–cells of  $\mathbf{D}$  to those of  $\mathbf{C}$ , and the following coherence data:

- (i) invertible unitality 2–cells

$$\begin{array}{ccc}
 FX & \xrightarrow{F(\text{id}_X)} & FX \\
 \parallel & \swarrow \alpha_X^h & \parallel \\
 FX & \xrightarrow{\text{id}_{F(X)}} & FX
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 FX & \xlongequal{\quad} & FX \\
 F(\text{id}_X) \downarrow & \swarrow \alpha_X^v & \parallel \text{id}_{F(X)} \\
 FX & \xlongequal{\quad} & FX
 \end{array}$$

for any object  $X$  of  $\mathbf{C}$ ,

- (ii) composition 2–cells

$$\begin{array}{ccc}
 FX & \xrightarrow{F(gf)} & FZ \\
 \parallel & \swarrow \mu_{g,f}^h & \parallel \\
 FX & \xrightarrow{Ff} FY \xrightarrow{Fg} & FZ
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 FX & \xlongequal{\quad} & FX \\
 \downarrow F(gf) & \swarrow \mu_{g,f}^v & \downarrow Ff \\
 FZ & \xlongequal{\quad} & FZ
 \end{array}$$

for composable pairs  $X \xrightarrow{f} Y \xrightarrow{g} Z$  of horizontal and vertical morphisms, respectively.

These coherence cells need to satisfy the unitality, associativity and naturality relations as written down in [39, Definition 6.1(v) and (vi)].

**Remark A.8** Note that in [39], the direction of the vertical structure of 2–cells is reversed. In that work, all of the above are assumed to be isomorphisms, so the direction of the cells is irrelevant. In our application, the 2–cells  $\mu_{g,f}^v$  are not invertible in general, so we have to take care of the orientation. We choose the given convention

since deriving (vertical) left derivable functors comes endowed with an oplax structure, and deriving (horizontal) right derivable functors with a lax structure.

On the other hand, the unitality 2-cells  $\alpha_X^h$  and  $\alpha_X^v$  are assumed to be invertible. This allows us to obtain from a lax–oplax double functor a lax 2–functor  $\mathcal{V}(F): \mathcal{V}(\mathbf{C}) \rightarrow \mathcal{V}(\mathbf{D})$  by applying  $F$  to vertical morphisms, and by defining

$$\mathcal{V}(F) \left( \begin{array}{ccc} X & \xlongequal{\quad} & X \\ f \downarrow & \Downarrow_{\eta} & \downarrow g \\ Y & \xlongequal{\quad} & Y \end{array} \right) = \begin{array}{ccccc} FX & \xlongequal{\quad} & FX & & FX \\ \parallel & \Downarrow_{(\alpha_X^h)^{-1}} & \parallel & & \parallel \\ FX & \xrightarrow{F(\text{id}_X)} & FX & & FX \\ Ff \downarrow & \Downarrow_{F\eta} & \downarrow Fg & & \downarrow Fg \\ FY & \xrightarrow{F(\text{id}_Y)} & FY & & FY \\ \parallel & \Downarrow_{\alpha_X^h} & \parallel & & \parallel \\ FY & \xlongequal{\quad} & FY & & FY \end{array}$$

In the same way, we obtain an oplax 2–functor  $\mathcal{H}(F): \mathcal{H}(\mathbf{C}) \rightarrow \mathcal{H}(\mathbf{D})$ .

In fact, the constraint that all  $\alpha_X^h$  are invertible can be used to strictify  $F$  into a lax–oplax functor where  $\alpha_X^h = \text{id}_X$ . The main result is:

**Lemma A.9** *Let  $\mathbf{C}$  and  $\mathbf{D}$  be double categories and  $F: \mathbf{C} \rightarrow \mathbf{D}$  be a lax–oplax double functor. Let*

$$\begin{array}{ccc} X & \xlongequal{\quad} & X \\ f \downarrow & \Downarrow_{\theta} & \downarrow g \\ Y & \xlongequal{\quad} & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X & \xrightarrow{h} & X' \\ g \downarrow & \Downarrow_{\eta} & \downarrow g' \\ Y & \xrightarrow{k} & Y' \end{array}$$

be 2–cells in  $\mathbf{C}$ . Then the 2–cells

$$\begin{array}{ccc} FX & \xrightarrow{Fh} & FX' \\ Ff \downarrow & \Downarrow_{F(\theta \square \eta)} & \downarrow Fg' \\ FY & \xrightarrow{Fk} & FY' \end{array} \quad \text{and} \quad \begin{array}{ccccc} FX & \xlongequal{\quad} & FX & \xrightarrow{Fh} & FX' \\ \parallel & \Downarrow_{(\alpha_X^h)^{-1}} & \parallel & & \parallel \\ FX & \xrightarrow{F(\text{id}_X)} & FX & & FX \\ Ff \downarrow & \Downarrow_{F(\theta)} & \downarrow Fg & \Downarrow_{F(\eta)} & \downarrow Fg' \\ FY & \xrightarrow{F(\text{id}_Y)} & FY & & FY \\ \parallel & \Downarrow_{\alpha_Y^h} & \parallel & & \parallel \\ FY & \xlongequal{\quad} & FY & \xrightarrow{Fk} & FY' \end{array}$$

in  $\mathbf{D}$  agree. The analogous statement holds for  $F(\eta \square \theta)$ .

**Proof** We use the following chain of pasting diagrams:

$$\begin{array}{ccc}
 FX \xlongequal{\quad} FX \xrightarrow{Fh} FX' & & FX \xlongequal{\quad} FX \xrightarrow{Fh} FX' \\
 \parallel \swarrow_{(\alpha_X^h)^{-1}} \parallel & & \parallel \swarrow_{(\alpha_X^h)^{-1}} \parallel \swarrow_{\text{id}_{Fh}} \parallel \\
 FX \xrightarrow{F(\text{id}_X)} FX & & FX \xrightarrow{F(\text{id}_X)} FX \xrightarrow{Fh} FX' \\
 Ff \downarrow \swarrow_{F(\theta)} \downarrow Fg \swarrow_{F(\eta)} & = & Ff \downarrow \swarrow_{F(\theta)} \downarrow Fg \swarrow_{F(\eta)} \downarrow Fg' \\
 FY \xrightarrow{F(\text{id}_Y)} FY & & FY \xrightarrow{F(\text{id}_Y)} FY \xrightarrow{Fk} FY' \\
 \parallel \swarrow_{\alpha_Y^h} \parallel & & \parallel \swarrow_{\alpha_Y^h} \parallel \swarrow_{\text{id}_{Fk}} \parallel \\
 FY \xlongequal{\quad} FY \xrightarrow{Fk} FY' & & FY \xlongequal{\quad} FY \xrightarrow{Fk} FY'
 \end{array}$$
  

$$\begin{array}{ccc}
 FX \xrightarrow{Fh} FX' & & FX \xrightarrow{Fh} FX' \\
 \parallel \swarrow_{\mu_{h,\text{id}}^h} \parallel & & \parallel \\
 FX \xrightarrow{F(\text{id}_X)} FX \xrightarrow{Fh} FX' & & FX \xrightarrow{F(\text{id}_X)} FX \xrightarrow{Fh} FX' \\
 = Ff \downarrow \swarrow_{F(\theta)} \downarrow Fg \swarrow_{F(\eta)} \downarrow Fg' & & = Ff \downarrow \swarrow_{F(\theta)} \downarrow Fg \swarrow_{F(\eta)} \downarrow Fg' \\
 FY \xrightarrow{F(\text{id}_Y)} FY \xrightarrow{Fk} FY' & & FY \xrightarrow{F(\text{id}_Y)} FY \xrightarrow{Fk} FY' \\
 \parallel \swarrow_{(\mu_{k,\text{id}}^h)^{-1}} \parallel & & \parallel \\
 FY \xrightarrow{Fk} FY' & & FY \xrightarrow{Fk} FY'
 \end{array}$$
  

$$\begin{array}{ccc}
 FX \xrightarrow{Fh} FX' & & \\
 = Ff \downarrow \swarrow_{F(\theta \square \eta)} \downarrow Fg' & & \\
 FY \xrightarrow{Fk} FY' & &
 \end{array}$$

Here, we used the unitality and naturality conditions on a lax–oplax double functor in the second and third steps, respectively. Also note that the unitality condition guarantees that the transformation  $\mu_{k,\text{id}}^h$  is indeed invertible. □

Now, we define the relevant notions of monads and morphisms between them in a double category. In our application, we have a left derivable monad and a right derivable monad morphism, and this motivates the following definition. Moreover, we define monadic transformations between monad morphisms in this context.

**Definition A.10** Let  $\mathbf{C}$  be a double category. A vertical monad  $T$  in  $\mathbf{C}$  is a monad in the vertical 2–category  $\mathcal{V}(\mathbf{C})$ . A horizontal monad morphism between two vertical monads  $S$  and  $T$  on objects  $X$  and  $Y$ , respectively, is a horizontal morphism  $F: X \rightarrow Y$



together with a 2-cell

$$\begin{array}{ccc} X & \xrightarrow{F} & Y \\ S \downarrow & \Downarrow \rho & \downarrow T \\ X & \xrightarrow{F} & Y \end{array}$$

satisfying the unitality and multiplicativity conditions

$$\begin{array}{ccc} X & \xrightarrow{F} & Y \quad \equiv \quad Y \\ S \downarrow & \Downarrow \rho & \downarrow T \quad \Downarrow \eta_T \quad \parallel \\ X & \xrightarrow{F} & Y \quad \equiv \quad Y \end{array} = \begin{array}{ccc} X & \equiv & X \xrightarrow{F} Y \\ S \downarrow & \Downarrow \eta_S & \parallel \quad \Downarrow \text{id}_F \quad \parallel \\ X & \equiv & X \xrightarrow{F} Y \end{array}$$

and

$$\begin{array}{ccc} X & \equiv & X \xrightarrow{F} Y \\ S \downarrow & \Downarrow \mu_S & \downarrow S \quad \Downarrow \rho \quad \downarrow T \\ X & \equiv & X \xrightarrow{F} Y \end{array} = \begin{array}{ccc} X & \xrightarrow{F} & Y \quad \equiv \quad Y \\ S \downarrow & \Downarrow \rho & \downarrow T \quad \Downarrow \mu_T \\ X & \xrightarrow{F} & Y \quad \equiv \quad Y \end{array}$$

For two horizontal monad morphisms  $(F, \rho)$  and  $(G, \sigma)$  between  $S$  and  $T$ , a monadic transformation is a 2-cell

$$\begin{array}{ccc} X & \xrightarrow{F} & Y \\ \parallel & \Downarrow \eta & \parallel \\ X & \xrightarrow{G} & Y \end{array} \quad \text{such that} \quad \begin{array}{ccc} X & \xrightarrow{F} & Y \\ S \downarrow & \Downarrow \rho & \downarrow T \\ X & \xrightarrow{F} & Y \\ \parallel & \Downarrow \eta & \parallel \\ X & \xrightarrow{G} & Y \end{array} = \begin{array}{ccc} X & \xrightarrow{F} & Y \\ \parallel & \Downarrow \eta & \parallel \\ X & \xrightarrow{G} & Y \\ S \downarrow & \Downarrow \sigma & \downarrow T \\ X & \xrightarrow{G} & Y \end{array}$$

**Proposition A.11** *Let  $\mathbf{C}$  and  $\mathbf{D}$  be two double categories and let  $L: \mathbf{C} \rightarrow \mathbf{D}$  be a lax-oplax double functor. Let  $(S, \mu, \eta)$  be a vertical monad in  $\mathbf{C}$ . Then*

$$(\mathcal{V}(L)(S), \mathcal{V}(L)(\mu) \circ \mu_{S,S}, \mathcal{V}(L)(\eta) \circ \alpha_X^{\mathcal{V}})$$

*is a vertical monad in  $\mathbf{D}$ .*

*Moreover, let  $S$  and  $T$  be vertical monads in  $\mathbf{C}$  and let  $(F, \rho)$  be a horizontal monad morphism between them. Then  $(LF, L\rho)$  is a horizontal monad morphism between  $\mathcal{V}(L)(S)$  and  $\mathcal{V}(L)(T)$ .*

*Furthermore, for any monadic transformation  $\eta: F \rightarrow G$  between two monad morphisms, the natural transformation  $\mathcal{H}(L)(\eta)$  is a monadic transformation between  $LF$  and  $LG$ .*

**Proof** The first part is a direct consequence of A.6, since a vertical monad is a monad in the vertical 2–category  $\mathcal{V}(\mathbf{C})$  and  $L$  induces a lax 2–functor  $\mathcal{V}(L): \mathcal{V}(\mathbf{C}) \rightarrow \mathcal{V}(\mathbf{D})$ .

We now prove the second part. We check the unitality condition on  $(LF, L\rho)$ , and thus consider

$$\begin{array}{c}
 \begin{array}{ccccc}
 LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & \xlongequal{\quad} & LY \\
 \downarrow LS & & \downarrow \text{\scriptsize } \Downarrow_{\alpha_Y^{-1}} & & \downarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & & \downarrow \text{\scriptsize } \Downarrow_{\alpha_Y} \\
 & & LY & \xrightarrow{L(\text{id}_Y)} & LY & & \\
 & \swarrow \text{\scriptsize } \Downarrow_{L\rho} & \downarrow LT & \swarrow \text{\scriptsize } \Downarrow_{L(\eta_T)} & \downarrow L(\text{id}_Y) & & \\
 & & LY & \xrightarrow{L(\text{id}_Y)} & LY & & \\
 & & \downarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & & \downarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & & \\
 LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & \xlongequal{\quad} & LY
 \end{array} \\
 \\
 \begin{array}{ccc}
 LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & & LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY \\
 \downarrow LS & \swarrow \text{\scriptsize } \Downarrow_{L(\rho \square \eta_T)} & \downarrow L(\text{id}_Y) & \swarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & \parallel & = & \downarrow LS & \swarrow \text{\scriptsize } \Downarrow_{L(\eta_S \square \text{id}_F)} & \downarrow L(\text{id}_Y) & \swarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & \parallel \\
 LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & & LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY
 \end{array} \\
 \\
 \begin{array}{ccc}
 LX & \xlongequal{\quad} & LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & & LX & \xlongequal{\quad} & LX & \xlongequal{\quad} & LX & \xrightarrow{LF} & LY \\
 \parallel & \swarrow \text{\scriptsize } \Downarrow_{\alpha_X^{-1}} & \parallel & & \downarrow L(\text{id}_Y) & & \parallel & & \parallel & \swarrow \text{\scriptsize } \Downarrow_{\alpha_X^{-1}} & \parallel & & \downarrow L(\text{id}_Y) & & \parallel \\
 LX & \xrightarrow{LF} & LX & & \downarrow L(\text{id}_X) & & LX & \xrightarrow{LF} & LX & & \downarrow L(\text{id}_X) & & \downarrow L(\text{id}_Y) & & \parallel \\
 \downarrow LS & \swarrow \text{\scriptsize } \Downarrow_{L(\eta_S)} & \downarrow L(\text{id}_X) & \swarrow \text{\scriptsize } \Downarrow_{L(\text{id}_F)} & \swarrow \text{\scriptsize } \Downarrow_{\alpha_Y} & & \downarrow LS & \swarrow \text{\scriptsize } \Downarrow_{L(\eta_S)} & \downarrow L(\text{id}_X) & \swarrow \text{\scriptsize } \Downarrow_{\alpha_X} & \swarrow \text{\scriptsize } \Downarrow_{\text{id}_{LF}} & & & & \\
 LX & \xrightarrow{LF} & LX & & & & LX & \xrightarrow{LF} & LX & & \swarrow \text{\scriptsize } \Downarrow_{\alpha_X} & & \downarrow L(\text{id}_Y) & & \parallel \\
 LX & \xlongequal{\quad} & LX & \xrightarrow{LF} & LY & \xlongequal{\quad} & LY & & LX & \xlongequal{\quad} & LX & \xlongequal{\quad} & LX & \xrightarrow{LF} & LY
 \end{array}
 \end{array}$$

Here we use Lemma A.9 in the first and third step.

The multiplicativity condition is obtained by similar pasting diagrams. Thus  $(LF, L(\rho))$  is a horizontal monad morphism.

The fact that  $\mathcal{H}(L)(\eta)$  is a monadic transformation between  $LF$  and  $LG$  is proven in the same way, using a horizontal version of Lemma A.9 for the exchange relation.  $\square$

Using this proposition, we also consider how the structure transformations of a lax–oplax double functor behave for a composite of monad functors.

**Lemma A.12** *Let  $\mathbf{C}$  and  $\mathbf{D}$  be double categories,  $R: X \rightarrow X$ ,  $S: Y \rightarrow Y$  and  $T: Z \rightarrow Z$  be vertical monads in  $\mathbf{C}$  and let  $(F, \rho)$  be a horizontal monad morphism from  $R$  to  $S$  and  $(G, \sigma)$  be a horizontal monad morphism from  $S$  to  $T$ . Let  $L: \mathbf{C} \rightarrow \mathbf{D}$*

be a lax-oplax double functor. Then the structure maps  $\mu_{G,F}^h: L(GF) \rightarrow LG \circ LF$  and  $\alpha_X^h: L(\text{id}_X) \rightarrow \text{id}_{LX}$  are monadic transformations in  $\mathbf{D}$ .

The proof follows easily using the naturality constraints of a lax-oplax double functor; see [39, (6.2)].

## References

- [1] **J F Adams**, *Lectures on generalised cohomology*, from “Category Theory, Homology Theory and their Applications, III” (P J Hilton, editor), Springer (1969) 1–138 MR Zbl
- [2] **J F Adams**, *Prerequisites (on equivariant stable homotopy) for Carlsson’s lecture*, from “Algebraic topology” (I Madsen, B Oliver, editors), Lecture Notes in Math. 1051, Springer (1984) 483–532 MR Zbl
- [3] **V Angeltveit**, *Topological Hochschild homology and cohomology of  $A_\infty$  ring spectra*, *Geom. Topol.* 12 (2008) 987–1032 MR Zbl
- [4] **S Araki, H Toda**, *Multiplicative structures in mod  $q$  cohomology theories, I*, *Osaka Math. J.* 2 (1965) 71–115 MR Zbl
- [5] **M F Atiyah, D O Tall**, *Group representations,  $\lambda$ -rings and the  $J$ -homomorphism*, *Topology* 8 (1969) 253–297 MR Zbl
- [6] **M G Barratt**, *A free group functor for stable homotopy*, from “Algebraic topology” (A Liulevicius, editor), Amer. Math. Soc., Providence, RI (1971) 31–35 MR Zbl
- [7] **C Barwick**, *Spectral Mackey functors and equivariant algebraic  $K$ -theory (I)*, *Adv. Math.* 304 (2017) 646–727 MR Zbl
- [8] **J Bénabou**, *Introduction to bicategories*, from “Reports of the Midwest Category Seminar”, Springer (1967) 1–77 MR Zbl
- [9] **A J Blumberg, M A Hill**, *Incomplete Tambara functors*, *Algebr. Geom. Topol.* 18 (2018) 723–766 MR Zbl
- [10] **E H Boorman**,  *$S$ -operations in representation theory*, *Trans. Amer. Math. Soc.* 205 (1975) 127–149 MR Zbl
- [11] **F Borceux**, *Handbook of categorical algebra, I: Basic category theory*, *Encycl. Math. Appl.* 50, Cambridge Univ. Press (1994) MR Zbl
- [12] **S Bouc**, *Biset functors for finite groups*, *Lecture Notes in Math.* 1990, Springer (2010) MR Zbl
- [13] **R R Bruner, J P May, J E McClure, M Steinberger**,  *$H_\infty$  ring spectra and their applications*, *Lecture Notes in Math.* 1176, Springer (1986) MR Zbl
- [14] **T tom Dieck**, *The Burnside ring of a compact Lie group, I*, *Math. Ann.* 215 (1975) 235–250 MR Zbl
- [15] **T tom Dieck**, *Orbittypen und äquivariante Homologie, II*, *Arch. Math. (Basel)* 26 (1975) 650–662 MR Zbl

- [16] **T tom Dieck**, *Transformation groups*, De Gruyter Studies in Math. 8, de Gruyter, Berlin (1987) MR Zbl
- [17] **A D Elmendorf, I Kriz, M A Mandell, J P May**, *Rings, modules, and algebras in stable homotopy theory*, Math. Surv. and Monogr. 47, Amer. Math. Soc., Providence, RI (1997) MR Zbl
- [18] **N Ganter**, *Global Mackey functors with operations and  $n$ -special lambda rings*, preprint (2013) arXiv 1301.4616
- [19] **S Gorchinskiy, V Guletskiĭ**, *Symmetric powers in abstract homotopy categories*, Adv. Math. 292 (2016) 707–754 MR Zbl
- [20] **P Guillot**, *Adams operations in cohomotopy*, preprint (2006) arXiv math/0612327
- [21] **B J Guillou, J P May**, *Equivariant iterated loop space theory and permutative  $G$ -categories*, Algebr. Geom. Topol. 17 (2017) 3259–3339 MR Zbl
- [22] **M A Hill, M J Hopkins, D C Ravenel**, *On the nonexistence of elements of Kervaire invariant one*, Ann. of Math. 184 (2016) 1–262 MR Zbl
- [23] **P Hoffman**,  *$\tau$ -rings and wreath product representations*, Lecture Notes in Math. 746, Springer (1979) MR Zbl
- [24] **M Hovey**, *Model categories*, Math. Surv. and Monogr. 63, Amer. Math. Soc., Providence, RI (1999) MR Zbl
- [25] **G M Kelly, R Street**, *Review of the elements of 2-categories*, from “Category Seminar” (G M Kelly, editor), Lecture Notes in Math. 420, Springer (1974) 75–103 MR Zbl
- [26] **D Knutson**,  *$\lambda$ -rings and the representation theory of the symmetric group*, Lecture Notes in Math. 308, Springer (1973) MR Zbl
- [27] **T Lawson**, *A note on  $H_\infty$  structures*, Proc. Amer. Math. Soc. 143 (2015) 3177–3181 MR Zbl
- [28] **D Lazard**, *Sur les modules plats*, C. R. Acad. Sci. Paris 258 (1964) 6313–6316 MR Zbl
- [29] **S MacLane**, *Categories for the working mathematician*, Graduate Texts in Math. 5, Springer (1971) MR Zbl
- [30] **M A Mandell, J P May, S Schwede, B Shipley**, *Model categories of diagram spectra*, Proc. London Math. Soc. 82 (2001) 441–512 MR Zbl
- [31] **J Noel**,  $H_\infty \neq E_\infty$ , from “An alpine expedition through algebraic topology” (C Ausoni, K Hess, B Johnson, W Lück, J Scherer, editors), Contemp. Math. 617, Amer. Math. Soc., Providence, RI (2014) 237–240 MR Zbl
- [32] **G Ochoa**, *Outer plethysm, Burnside ring and  $\beta$ -rings*, J. Pure Appl. Algebra 55 (1988) 173–195 MR Zbl
- [33] **S B Priddy**, *On  $\Omega^\infty S^\infty$  and the infinite symmetric group*, from “Algebraic topology” (A Liulevicius, editor), Amer. Math. Soc., Providence, RI (1971) 217–220 MR Zbl

- [34] **N W Rymer**, *Power operations on the Burnside ring*, J. London Math. Soc. 15 (1977) 75–80 MR Zbl
- [35] **S Schwede**, *Global homotopy theory*, New Math. Monogr. 34, Cambridge Univ. Press (2018) MR Zbl
- [36] **G B Segal**, *Equivariant stable homotopy theory*, from “Actes du Congrès International des Mathématiciens” (M Berger, J Dieudonné, J Leray, J-L Lions, P Malliavin, J-P Serre, editors), volume 2, Gauthier-Villars, Paris (1971) 59–63 MR Zbl
- [37] **G Segal**, *Categories and cohomology theories*, Topology 13 (1974) 293–312 MR Zbl
- [38] **G Segal**, *Operations in stable homotopy theory*, from “New developments in topology” (G Segal, editor), London Math. Soc. Lecture Note Ser. 11, Cambridge Univ. Press (1974) 105–110 MR Zbl
- [39] **M Shulman**, *Comparing composites of left and right derived functors*, New York J. Math. 17 (2011) 75–125 MR Zbl
- [40] **J Singer**, *Äquivariante  $\lambda$ -Ringe und kommutative Multiplikationen auf Moore-Spektren*, PhD thesis, Universität Bonn (2007) Zbl Available at <https://hdl.handle.net/20.500.11811/3607>
- [41] **M Stolz**, *Equivariant structure on smash powers of commutative ring spectra*, PhD thesis, University of Bergen (2011)
- [42] **R Street**, *The formal theory of monads*, J. Pure Appl. Algebra 2 (1972) 149–168 MR Zbl
- [43] **R Street**, *Two constructions on lax functors*, Cahiers Topologie Géom. Différentielle 13 (1972) 217–264 MR Zbl
- [44] **E Vallejo**, *Polynomial operations from Burnside rings to representation functors*, J. Pure Appl. Algebra 65 (1990) 163–190 MR Zbl
- [45] **E Vallejo**, *Polynomial operations on stable cohomotopy*, Manuscripta Math. 67 (1990) 345–365 MR Zbl
- [46] **E Vallejo**, *The free  $\beta$ -ring on one generator*, J. Pure Appl. Algebra 86 (1993) 95–108 MR Zbl
- [47] **P Webb**, *A guide to Mackey functors*, from “Handbook of algebra, II” (M Hazewinkel, editor), North-Holland, Amsterdam (2000) 805–836 MR Zbl
- [48] **D White**, *Model structures on commutative monoids in general model categories*, J. Pure Appl. Algebra 221 (2017) 3124–3168 MR Zbl

Max Planck Institute for Mathematics  
Bonn, Germany

mstahlhauer@uni-bonn.de

<https://guests.mpim-bonn.mpg.de/stahlhauer/>

Received: 22 September 2020      Revised: 22 July 2021



# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](http://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
etnyre@math.gatech.edu  
Georgia Institute of Technology

Kathryn Hess  
kathryn.hess@epfl.ch  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

Matthew Ando	University of Illinois mando@math.uiuc.edu	Tsuyoshi Kobayashi	Nara Women's University tsuyoshi09@gmail.com
Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Joan Birman	Columbia University jb@math.columbia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Indira Chatterji	CNRS & Université Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas Flores	Universidad de Santiago de Chile andres.navas@usach.cl
Fred Cohen	University of Rochester cohf@math.rochester.edu	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris-Nord bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Luis Paris	Université de Bourgogne lparis@u-bourgogne.fr
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
David Futер	Temple University dfuter@temple.edu	Peter Scott	University of Michigan pscott@umich.edu
Soren Galatius	Stanford University galatius@math.stanford.edu	Zoltán Szabó	Princeton University szabo@math.princeton.edu
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
J. Elisenda Grigsby	Boston College grigsbyj@bc.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Daniel T. Wise	McGill University, Canada daniel.wise@mcgill.ca
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu		

---

See inside back cover or [msp.org/agt](http://msp.org/agt) for submission instructions.


The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23 Issue 1 (pages 1–508) 2023

---

Proper 2–equivalences between infinite ended finitely presented groups	1
MANUEL CÁRDENAS, FRANCISCO FERNÁNDEZ LASHERAS, ANTONIO QUINTERO and RANJA ROY	
Groups acting on CAT(0) cube complexes with uniform exponential growth	13
RADHIKA GUPTA, KASIA JANKIEWICZ and THOMAS NG	
Admissible replacements for simplicial monoidal model categories	43
HALDUN ÖZGÜR BAYINDIR and BORIS CHORNY	
The bridge number of arborescent links with many twigs	75
SEBASTIAN BAADER, RYAN BLAIR, ALEXANDRA KJUCHUKOVA and FILIP MISEV	
$G_\infty$ –ring spectra and Moore spectra for $\beta$ –rings	87
MICHAEL STAHLHAUER	
External Spanier–Whitehead duality and homology representation theorems for diagram spaces	155
MALTE LACKMANN	
On the surjectivity of the $\mathrm{tmf}$ –Hurewicz image of $A_1$	217
VIET-CUONG PHAM	
The handlebody group and the images of the second Johnson homomorphism	243
QUENTIN FAES	
Recognition of connective commutative algebra spectra through an idempotent quasiajunction	295
RENATO VASCONCELLOS VIEIRA	
$\mathrm{SL}_2$ quantum trace in quantum Teichmüller theory via writhe	339
HYUN KYU KIM, THANG T Q LÊ and MIRI SON	
Constraints on families of smooth 4–manifolds from $\mathrm{Pin}^-(2)$ –monopole	419
HOKUTO KONNO and NOBUHIRO NAKAMURA	
Suspension homotopy of 6–manifolds	439
RUIZHI HUANG	
Time-periodic solutions of Hamiltonian PDEs using pseudoholomorphic curves	461
OLIVER FABERT and NIEK LAMOREE	