

AG
T

*Algebraic & Geometric
Topology*

Volume 23 (2023)

Mod 2 power operations revisited

DYLAN WILSON



Mod 2 power operations revisited

DYLAN WILSON

In this mostly expository note we take advantage of homotopical and algebraic advances to give a modern account of power operations on the mod 2 homology of \mathbb{E}_∞ -ring spectra. The main advance is a quick proof of the Adem relations utilizing the Tate-valued Frobenius as a homotopical incarnation of the total power operation. We also give a streamlined derivation of the action of power operations on the dual Steenrod algebra.

55P43, 55S12

Introduction	2993
1. The Tate construction	2995
2. Tate powers	2999
3. Power operations	3001
4. Adem relations	3007
5. Relationship to the Steenrod algebra	3011
References	3013

Introduction

As someone who entered college at about the time that Netflix started automatically playing the next episode of a series, I cannot imagine discovering or verifying the Adem relations using the tools available to Adem [1].¹ I even find it hard to *remember* the Adem and Nishida relations.

¹It was precisely while trying and failing multiple times to prove the Adem relations in equivariant homotopy theory that, in an act of true laziness, I stumbled upon the technique explained in this note.

Luckily, there is a useful mnemonic device which utilizes the *total power operation*

$$Q(t) := \sum_{i \in \mathbb{Z}} Q^i t^i.$$

Here t is an indeterminate, and the operation $Q^i : A_* \rightarrow A_{*+i}$ acts on the homotopy of any $\mathbb{E}_\infty\text{-}\mathbb{F}_2$ -algebra A . The total power operation then produces a map

$$Q(t) : A_* \rightarrow A_*((t)).$$

We extend $Q(t)$ to a ring map

$$Q(t) : A_*((s)) \rightarrow A_*[[s, t]][s^{-1}, t^{-1}]$$

by requiring that

$$Q(t)(s) = s + s^2 t^{-1}.$$

With this convention, it is possible to restate the Adem relations, following Bullett and Macdonald [5], Steiner [16], and Bisson and Joyal [3], as:

- **Adem relations** For any $x \in A_*$, $Q(t)Q(s)x$ is symmetric in s and t .

The usual Adem relations are recovered using a trick with residues which we will review in Section 4.3. Steiner’s proof that the above identity holds is to reduce it to one of the expressions met in the proof of the Adem relations as in Steenrod [15, page 119] and May [12, 4.7(e,g,i)].

In the case of Steenrod operations acting on the cohomology of a space X , there is a more conceptual argument due to Segal [5, Section 4]. One can use the diagonal map to produce a version of the total power operation taking values in $H^*(X \times B\Sigma_2)$. Indeed, this is one of the earlier constructions of Steenrod operations [15, Chapter VII]. The iterated total square then takes values in $H^*(X \times B\Sigma_2 \times B\Sigma_2) = H^*(X)[s, t]$ but factors through the total fourth power which takes values in $H^*(X \times B\Sigma_4)$. The automorphism swapping s and t arises as an inner automorphism of Σ_4 so the formula for the iterated square must be symmetric in s and t .

Our primary goal is to explain how the Tate diagonal (Section 2.3) on spectra allows for a similar argument for general power operations. The reader could probably reconstruct the argument themselves just from the observation that the total power operation is the effect on homotopy of the (non- \mathbb{F}_2 -linear) map of spectra

$$A \xrightarrow{\Delta} (A \otimes_{\mathbb{F}_2} A)^{t\Sigma_2} \rightarrow A^{t\Sigma_2}.$$

In fact, we take this as a definition and develop all the basic properties of power operations efficiently from there. We hope that this note will give a mnemonic for the *proofs* of the standard identities for power operations in much the same way that the work of Steiner [16], Bisson and Joyal [3], and Baker [2] has provided mnemonics for their *statements*.

Outline

In Sections 1 and 2 we review the facts we need about the Tate construction and the Tate diagonal, following Nikolaus and Scholze [14]. In Section 3 we give three definitions of the operations Q^i : the classical one, one due to Lurie [10, Section 2.2], and one in terms of the Tate-valued Frobenius. We then explain how to recover the first properties of power operations.

In Section 4 we turn to the Adem relations. The key thing to prove is that having a Σ_4 -equivariant map $A^{\otimes 4} \rightarrow A$ produces a lift of the iterated total power operation through the Frobenius $A \rightarrow A^{t\Sigma_4}$. This takes a little bit of work but the reader could come up with the argument themselves if they remember to use the universal property of the Tate diagonal amongst natural transformations of exact, lax symmetric monoidal functors over and over again. Indeed, this proof is an excellent illustration of the computational utility of establishing such universal properties in the first place.

Finally, in Section 5, we show how the Bisson–Joyal and Baker formulations of the Nishida relations arise naturally from the perspective of the Tate-valued Frobenius. We end by explaining how to recover Steinberger’s formulas [4, Section III.2] for the action of power operations on the dual Steenrod algebra. This last step is mostly algebraic, and essentially due to Bisson and Joyal, but we have included it for completeness.

Acknowledgements The author is grateful to Tom Bachmann for comments on an earlier draft, and to the referee for careful reading and helpful suggestions.

1 The Tate construction

We review the Tate construction (Section 1.1) and its universal property (Section 1.3) as well as the important Warwick duality (Section 1.2) of Greenlees [8] which allows an alternative computation of the Tate construction. We end (Section 1.4) by spelling out what happens in the case $G = \Sigma_2$.

1.1 Definitions

Let G be a finite group and k an \mathbb{E}_∞ -ring, and denote by

$$\text{Mod}_k^{hG} := \text{Psh}(\text{BG}; \text{Mod}_k)$$

the ∞ -category of Borel G -modules. There is a fully faithful embedding

$$\text{Mod}_k^{hG} \rightarrow \text{Mod}_k^G$$

from Borel G -modules to modules over k in genuine G -spectra whose essential image consists of the Borel complete G -modules, ie those X such that $X \rightarrow F(\text{EG}_+, X)$ is an equivalence. Let \mathcal{F} be a collection of subgroups closed under subconjugacy, and $\text{E}\mathcal{F}$ the G -space characterized up to homotopy by the requirement

$$\text{E}\mathcal{F}^H = \begin{cases} * & \text{if } H \in \mathcal{F}, \\ \emptyset & \text{if } H \notin \mathcal{F}, \end{cases}$$

and define $\widetilde{\text{E}}\mathcal{F}$ as the cofiber of $\text{E}\mathcal{F}_+ \rightarrow S^0$. Then the \mathcal{F} -Tate spectrum of a Borel G -spectrum can be computed as [7, page 443]

$$X^{t\mathcal{F}} = (\widetilde{\text{E}}\mathcal{F} \wedge F(\text{EG}_+, X))^G,$$

where the right-hand side is computed in genuine G -spectra.

It will be more convenient for us to think of the above as a computation and not a definition. Instead, we opt to define the Tate construction by a universal property, following [14].

To that end, let

$$(\text{Mod}_k^{hG})_{\mathcal{F}\text{-ind}} \subseteq \text{Mod}_k^{hG}$$

be the smallest full, stable subcategory containing all objects which are left Kan extended from diagrams $\text{BH} \rightarrow \text{Sp}$ for some $H \in \mathcal{F}$.

Recall [14, Section I.3] that, associated to any exact functor $F: \text{Mod}_k^{hG} \rightarrow \mathcal{E}$ to a presentable stable ∞ -category \mathcal{E} , there is a natural transformation

$$F \rightarrow L_{\mathcal{F}}F$$

which is initial amongst natural transformations to exact functors which annihilate the subcategory $(\text{Mod}_k^{hG})_{\mathcal{F}\text{-ind}}$. Concretely, $L_{\mathcal{F}}F$ is specified by the formula [14, I.3.3]

$$L_{\mathcal{F}}F(X) = (\text{Mod}_k^{hG})_{\mathcal{F}\text{-ind}/X} Y \text{ colim } F(\text{cofib}(Y \rightarrow X)).$$

Definition 1.1.1 With notation as above, we define

$$(-)^{t\mathcal{F}} = L_{\mathcal{F}}((-)^{hG}): \text{Mod}_k^{hG} \rightarrow \text{Mod}_k.$$

More generally, if $G \subseteq G'$, we define

$$(-)^{t\mathcal{F}} = L_{\mathcal{F}}((-)^{hG}): \text{Mod}_k^{hG'} \rightarrow \text{Mod}_k^{hW_{G'}G}$$

where $W_{G'}G = N_{G'}G/G$ is the Weyl group of G in G' .

Example 1.1.2 When \mathcal{F} consists only of the trivial subgroup, we denote $X^{t\mathcal{F}}$ by X^{tG} . This can be computed as the cofiber of the trace map $X_{hG} \rightarrow X^{hG}$.

Example 1.1.3 Suppose $G \subseteq \Sigma_n$ is a subgroup and let $\mathcal{F} = \mathcal{T}$ be the family of subgroups of G which do *not* act transitively on $\{1, \dots, n\}$. When $G = C_n$ this coincides with the more commonly seen family of proper subgroups, and when $G = C_p$ this coincides with the family consisting of only the trivial subgroup.

1.2 Warwick duality

We can dualize the construction in the previous section and define the *opposite \mathcal{F} -Tate spectrum*² as

$$X^{t^{op}\mathcal{F}} := \text{holim}_{((\text{Mod}_k^{hG})_{\mathcal{F}\text{-ind}})_{X/\ni Y}} \text{fib}(X \rightarrow Y)_{hG}.$$

Greenlees proved [8, Section B] that this construction is not really new:

Theorem 1.2.1 (Warwick duality) *There is a canonical equivalence*

$$X^{t^{op}\mathcal{F}} \simeq \Sigma^{-1} X^{t\mathcal{F}}.$$

In particular, we obtain extra functoriality: if $\mathcal{F} \subseteq \mathcal{F}'$, then the original construction produces a canonical map $(-)^{t\mathcal{F}'} \rightarrow (-)^{t\mathcal{F}}$ while the opposite construction, composed with suspension, produces a map $(-)^{t\mathcal{F}} \rightarrow (-)^{t\mathcal{F}'}$.

1.3 Monoidal structure

We will make much use of the following excellent description of the lax symmetric monoidal structure on the Tate construction.

²We stole this name from [6].

Proposition 1.3.1 *There is a natural transformation of lax symmetric monoidal functors*

$$(-)^{hG} \rightarrow (-)^{t\mathcal{F}}$$

which is initial amongst natural transformations of lax symmetric monoidal functors with target an exact functor that annihilates $(\text{Mod}_k^{hG})_{\mathcal{F}\text{-ind}}$.

This follows from the more general result [14, I.3.6] about the relationship between Verdier quotients and lax symmetric monoidal structures.

1.4 An example

Let k be a field of characteristic 2. Then

$$\pi_* k^{h\Sigma_2} \simeq H^{-*}(\mathbb{B}\Sigma_2, k) = k[[t]],$$

where $t \in \pi_{-1} k^{h\Sigma_2}$ is the Stiefel–Whitney class of the canonical line bundle. The Tate construction has the effect of inverting t and we can compute

$$\pi_* k^{t\Sigma_2} = k((t)),$$

the algebra of Laurent series over k .

On the other side, the homotopy orbits $k_{h\Sigma_2}$ have a dual basis on homotopy

$$\pi_* k_{h\Sigma_2} = k\{e_0, e_1, \dots\},$$

where e_i is the linear dual of t^i . The trace map

$$k_{h\Sigma_2} \rightarrow k^{h\Sigma_2}$$

is zero on homotopy groups and so we have a short exact sequence

$$0 \rightarrow k[[t]] \rightarrow k((t)) \rightarrow \pi_* \Sigma k_{h\Sigma_2} \rightarrow 0$$

which identifies the last term as the quotient $k((t))/k[[t]]$. This provides another basis for the homotopy of $k_{h\Sigma_2}$, and the two are related by the correspondence

$$e_i \leftrightarrow t^{-i-1}.$$

Under this interpretation, the composite map

$$k^{t\Sigma_2} \rightarrow \Sigma k_{h\Sigma_2} \rightarrow \Sigma k$$

is given by sending a Laurent series $g(t) = \sum a_i t^i$ to the *residue* a_{-1} .

Finally, Warwick duality in this context translates to the computation [9, 16.1]

$$\Sigma^{-1}k^{t\Sigma_2} = \operatorname{holim}_n(\Sigma^{-n\tau}k)_{h\Sigma_2} = \operatorname{holim}_n k \wedge (\mathbb{R}P^\infty)^{-n\tau} = \operatorname{holim}_n k \wedge \mathbb{R}P_{-n}^\infty,$$

where τ is the sign representation.

2 Tate powers

The source of power operations is the symmetry present on $X^{\otimes n}$. In Section 2.1 we review several constructions based on this symmetry. In Section 2.2 we explain how the construction $X \mapsto (X^{\otimes n})^{t\mathcal{T}}$ arises as a Goodwillie derivative; in particular this construction is exact. In Section 2.3, following [14], we describe the spectral analog of the diagonal map we will use when defining power operations.

2.1 Variants of extended powers

Let \mathcal{C} be a symmetric monoidal ∞ -category. Then there is a natural functor

$$\mathcal{C} \rightarrow \mathcal{C}^{h\Sigma_n} = \operatorname{Fun}(\mathbf{B}\Sigma_n, \mathcal{C})$$

given as the composite

$$\mathcal{C} \xrightarrow{\delta} (\mathcal{C}^{\times n})^{h\Sigma_n} \rightarrow \mathcal{C}^{h\Sigma_n}$$

where the latter map is a choice of tensor product. In other words, for every $X \in \mathcal{C}$, the object $X^{\otimes n}$ has a Σ_n -action.

If \mathcal{C} admits homotopy limits and colimits, we can form both a “symmetric” power of an object and a “divided” power of an object. We do this more generally for a fixed subgroup $G \subseteq \Sigma_n$.

Definition 2.1.1 We define symmetric and divided power functors as

$$\operatorname{Sym}^G(X) := (X^{\otimes n})_{hG}, \quad \Gamma^G(X) := (X^{\otimes n})^{hG}.$$

Finally, if $\mathcal{C} = \operatorname{Mod}_k$ is the ∞ -category of k -modules over an \mathbb{E}_∞ -ring k , then:

Definition 2.1.2 Let $G \subseteq \Sigma_n$ be a subgroup. We define the *Tate power* of X as

$$\mathbf{T}_G(X) := (X^{\otimes n})^{t\mathcal{T}}$$

where \mathcal{T} is the family of nontransitive subgroups of G .

In each case we abbreviate G as n if $G = \Sigma_n$.

2.2 Tate powers as a Goodwillie derivative

Let \mathcal{C} and \mathcal{D} be stable, presentable ∞ -categories. Then the full subcategory

$$\text{Fun}^{\text{ex}}(\mathcal{C}, \mathcal{D}) \subseteq \text{Fun}(\mathcal{C}, \mathcal{D})$$

admits a left adjoint [11, 6.1.1.10], the 1-excisive approximation

$$P_1: \text{Fun}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Fun}^{\text{ex}}(\mathcal{C}, \mathcal{D}).$$

In the case where $F(0) = 0$, we may compute $P_1 F$ as [11, 6.1.1.23 and 6.1.1.27]

$$P_1 F(X) = \text{hocolim}_n \Omega_{\mathcal{D}}^n F(\Sigma_{\mathcal{C}}^n X).$$

I believe the following is well known but do not know a reference.

Proposition 2.2.1 *With notation as in Section 2.1, there is an equivalence*

$$P_1 \Gamma^G \simeq T_G.$$

Proof Let V denote the standard representation of Σ_n on \mathbb{R}^n and \bar{V} the reduced standard representation. By the formula above,

$$\begin{aligned} P_1 \Gamma^G(X) &= \text{hocolim}_j \Omega^j \Gamma^G(\Sigma^j X) \\ &\simeq \text{hocolim} \Omega^j (\Sigma^{jV} X^{\otimes n})^{h\Sigma_n} \\ &\simeq \text{hocolim} (\Sigma^{j\bar{V}} X^{\otimes n})^{h\Sigma_n} \\ &\simeq \text{hocolim} (S^{j\bar{V}} \wedge F(\text{EG}_+, X^{\otimes n}))^G \\ &\simeq (S^{\infty\bar{V}} \wedge F(\text{EG}_+, X^{\otimes n}))^G. \end{aligned}$$

The last identification used that genuine fixed points commute with all homotopy limits and colimits. Finally, observe that $S^{\infty\bar{V}}$ is a model for $\widetilde{\text{E}\mathcal{T}}$. □

The same argument computes the Goodwillie coderivative of Sym^G :

Proposition 2.2.2 *The Goodwillie coderivative of Sym^G is $((-)^{\otimes n})^{t^{\text{op}}\mathcal{T}} = \Sigma^{-1} T_G$.*

This last observation motivates the excellent account of stable power operations given by Glasman and Lawson [6].

2.3 The Tate diagonal

Recall the following result of Nikolaus [13, Corollary 6.9]:

Proposition 2.3.1 *The forgetful functor $U: \text{Mod}_k \rightarrow \text{Sp}$ is initial amongst exact, lax symmetric monoidal functors to spectra.*

In the previous section we identified T_G as a Goodwillie derivative. In particular, T_G is exact. It also has a lax symmetric monoidal structure, being a composite of lax symmetric monoidal functors. So we get the following:

Corollary 2.3.2 *There is an essentially unique natural transformation of lax symmetric monoidal functors $U \rightarrow UT_G$.*

We refer to this map $\Delta_G: M \rightarrow T_G(M)$ as the *Tate diagonal*.

Remark 2.3.3 This is not the same as the Tate diagonal in [14] unless $k = S^0$, since we use the tensor product in Mod_k . Of course there is an evident relationship between the two: the Tate diagonal above is just the composite

$$M \rightarrow (M^{\wedge n})^{t\mathcal{T}} \rightarrow (M^{\otimes n})^{t\mathcal{T}}.$$

Warning 2.3.4 The Tate diagonal is *not* k -linear.

3 Power operations

We now fix a field k of characteristic 2 and let Mod_k be the ∞ -category of k -module (spectra). In Section 3.1 we serve up power operations three ways, and then verify they agree in Section 3.5. In between we verify the first properties of power operations up to the Cartan formula. We emphasize that this section does not show off the utility of the approach via the Tate-valued Frobenius, but we have included the proofs since they are still pleasant.

3.1 Three definitions of operations

First we specify the objects on which power operations will act.

Definition 3.1.1 We say that $A \in \text{Mod}_k$ is *equipped with a symmetric multiplication* if we have specified a map $\text{Sym}^2(A) \rightarrow A$ of k -modules. Equivalently, if we have specified a map $A^{\otimes 2} \rightarrow A$ in $\text{Mod}_k^{h\Sigma_2}$.

Remark 3.1.2 A k -module with a symmetric multiplication is the same as an object of $\mathcal{C}(2, \infty)$ in the notation of [12].

To give the classical construction of power operations we'll need a computation.

Lemma 3.1.3 For any integer n there is a canonical equivalence

$$\text{Sym}^2(\Sigma^n k) \simeq \Sigma^{2n} k_{h\Sigma_2}.$$

Proof The object $(\Sigma^n k)^{\otimes 2} = \Sigma^{nV} k$ in $\text{Mod}_k^{h\Sigma_2}$ corresponds to a map $B\Sigma_2 \rightarrow \text{Mod}_k^{h\Sigma_2}$ which is determined by a map

$$\Sigma_2 \rightarrow \text{End}_k(\Sigma^{2n} k) \simeq \text{End}_k(k, k) \simeq k$$

of \mathbb{E}_1 -monoids. The map factors through the units k^\times , but k has characteristic 2 and hence no nontrivial square roots of unity. So the action is trivial and the result follows. □

The following construction is the current standard definition of power operations.

Construction 3.1.4 (hands-on power operations) Let A be a k -module equipped with a symmetric multiplication. Given $x \in \pi_n A$ and $i \geq n$, define $Q^i(x) \in \pi_{n+i} A$ as the composite

$$S^{n+i} \xrightarrow{\Sigma^{2n} e_{i-n}} \Sigma^{2n} k_{h\Sigma_2} \simeq \text{Sym}^2(\Sigma^n k) \xrightarrow{\text{Sym}^2(x)} \text{Sym}^2(A) \rightarrow A.$$

This has the benefit of generalizing well to power operations for other cohomology theories, but in the case of mod 2 cohomology there is a more uniform option. The author learned this next approach from [10, Section 2.2] and has not found an earlier reference, but a more recent and detailed account can be found in [6].

First we need a preliminary observation. Let $T'_2: \text{Mod}_k \rightarrow \text{Mod}_k$ denote the left Kan extension of the restriction of T_2 to the full subcategory of compact objects. This endomorphism commutes with all colimits and so—see [11, 7.1.2.4]—there is a bimodule B and an equivalence $T'_2(M) \simeq B \otimes M$. By evaluating on $M = k$ we deduce that $B = k^{t\Sigma_2}$ as a left k -module. Notice, by construction, we have a natural map $B \otimes M \rightarrow T_2(M)$.

Construction 3.1.5 (stable power operations) Let A be a k -module equipped with a symmetric multiplication. The element $t^{-i-1} \in \pi_{i+1} k^{t\Sigma_2}$ extends to a right module map $\Sigma^i k \rightarrow \Sigma^{-1} B$. We now define $Q^i: \Sigma^i A \rightarrow A$ as the (non- k -linear!) composite

$$\Sigma^i A = \Sigma^i k \otimes A \rightarrow \Sigma^{-1} B \otimes A \rightarrow \Sigma^{-1} T_2(A) \rightarrow \text{Sym}^2(A) \rightarrow A.$$

This construction emphasizes the role of $\Sigma^{-1} k^{t\Sigma_2}$ as acting on A , but we can also record this information in a kind of coaction. For that we first need a computation.

Lemma 3.1.6 For any k -module M equipped with the trivial Σ_2 -action, there is a canonical equivalence of $\pi_*k^{t\Sigma_2}$ -modules

$$\pi_*M^{t\Sigma_2} \simeq M_*((t)).$$

Proof It suffices to prove $\pi_*M^{h\Sigma_2} \simeq M_*[[t]]$. From the skeletal filtration on $B\Sigma_2$,

$$M^{h\Sigma_2} \simeq \operatorname{holim} F(\operatorname{sk}_j B\Sigma_2, k) \otimes M$$

and $\pi_*F(\operatorname{sk}_j B\Sigma_2, k) \otimes M = M_*[t]/t^{j+1}$. The transition maps are surjective so there is no \lim^1 term in the Milnor exact sequence and the result follows. \square

Construction 3.1.7 (Tate-valued Frobenius) Let A be a k -module equipped with a symmetric multiplication. Define the *total power operation* as the composite

$$Q(t) : A \xrightarrow{\Delta_2} T_2(A) = (A^{\otimes 2})^{t\Sigma_2} \rightarrow A^{t\Sigma_2}.$$

We then define $Q^i : A \rightarrow \Sigma^{-i}A$ as the composite

$$A \rightarrow A^{t\Sigma_2} \xrightarrow{t^{-i-1}} \Sigma^{-i-1}A^{t\Sigma_2} \rightarrow \Sigma^{-i}A_{h\Sigma_2} \rightarrow \Sigma^{-i}A.$$

In Section 3.5 we will verify that the two definitions of the endomorphism $Q^i : \Sigma^i A \rightarrow A$ coincide and that each induce the operation $Q^i : \pi_n A \rightarrow \pi_{n+i} A$ on homotopy. For now we will assume this compatibility.

Remark 3.1.8 (naturality of Frobenius) The Tate-valued Frobenius can be defined for any *spectrum* equipped with a symmetric multiplication, as the composite

$$A \rightarrow (A \wedge A)^{t\Sigma_2} \rightarrow A^{t\Sigma_2}.$$

Since the k -module Tate diagonal factors through the spectrum Tate diagonal, we learn that the Tate-valued Frobenius only depends on the underlying \mathbb{E}_∞ -ring. In particular, the Tate-valued Frobenius is natural for maps $A \rightarrow B$ of \mathbb{E}_∞ -rings, independent of any compatibility with k -module structures.

3.2 First properties

The first properties follow easily from the Tate-valued Frobenius description, with the exception of the squaring property, which is most readily seen through the classical definition.

Proposition 3.2.1 The operations Q^i satisfy the following properties:

- (i) **Additivity** $Q^i(x + y) = Q^i(x) + Q^i(y)$.

- (ii) **Suspension** $\Omega Q^i(x) = Q^i(\Omega x)$.
- (iii) **Squaring** $Q^{|x|}(x) = x^2$.
- (iv) **Instability** $Q^i(x) = 0$ if $i < |x|$.
- (v) **Action on cohomology** If $A = F(X, k)$, where X is a pointed space, then $Q^i(x) = 0$ for $i > 0$ and $Q^0(x) = x$.

Proof (i) **Additivity** Since Q^i is induced by a map of spectra, it is automatically additive.

(ii) **Suspension** The Tate diagonal is a natural transformation of exact functors, so $\Delta_{2, \Omega A} \simeq \Omega \Delta_2$. Exactness of T_2 then ensures that $\Omega T_2(A) \rightarrow T_2(\Omega A)$ is an equivalence, and composing with the multiplication on ΩA identifies $\Omega Q(t)$ with the total power operation for ΩA , which was to be shown.

(iii) **Squaring** Using Construction 3.1.4, observe that $Q^{|x|}(x)$ is the image of the bottom class in $\text{Sym}^2(\Sigma^n k)$, which is the left vertical arrow in the diagram

$$\begin{array}{ccc}
 \Sigma^n k \otimes \Sigma^n k & \xrightarrow{x \otimes x} & M \otimes M \\
 \downarrow & & \downarrow \\
 \text{Sym}^2(\Sigma^n k) & \longrightarrow & \text{Sym}^2(M)
 \end{array}$$

The result follows by chasing the diagram clockwise.

(iv) **Instability** By (ii) we may replace A by $\Omega^{|x|-i} A$ and thereby reduce to the case that $A = \Omega B$ and $i = |x|$. By (iii), $Q^i x = x^2$, but the multiplication on ΩB is always trivial, since $S^1 \rightarrow S^1 \wedge S^1$ is null.

(v) **Action on cohomology** By naturality we may replace X with $K(k, n)$ and x with the fundamental class. The vanishing now follows for degree reasons. To check that $Q^0(x) = x$ we may reduce, by naturality, to the case $X = S^n$ and then, by stability, to $X = S^0$. The result now follows from the equivalence $F(S^0, k) = k$. □

3.3 Cartan formula

If A and A' are equipped with symmetric multiplications then $A \otimes A'$ inherits a canonical symmetric multiplication as well. In this case we have an external Cartan formula:

Proposition 3.3.1 (Cartan formula)

$$Q(t)(x \otimes y) = Q(t)(x) \otimes Q(t)(y) \in (A \otimes A')((t)).$$

Proof The formula is equivalent to commutativity of the square

$$\begin{array}{ccccc}
 A \otimes A' & \longrightarrow & T_2(A) \otimes T_2(A') & \longrightarrow & A^{t\Sigma_2} \otimes (A')^{t\Sigma_2} \\
 \parallel & & \downarrow & & \downarrow \\
 A \otimes A' & \longrightarrow & T_2(A \otimes A') & \longrightarrow & (A \otimes A')^{t\Sigma_2}
 \end{array}$$

The left square commutes because the Tate diagonal is a transformation of lax symmetric monoidal functors. The right-hand square commutes by naturality of the lax structure map

$$(-)^{t\Sigma_2} \otimes (-)^{t\Sigma_2} \rightarrow (- \otimes -)^{t\Sigma_2}$$

applied to $(A \otimes A')^{\otimes 2} \simeq A^{\otimes 2} \otimes A'^{\otimes 2} \rightarrow A \otimes A'$. □

Corollary 3.3.2
$$Q^n(x \otimes y) = \sum_{i+j=n} Q^i(x) \otimes Q^j(y).$$

As a corollary of the proof, we see:

Corollary 3.3.3 *If $A \otimes A \rightarrow A$ is a map of objects equipped with symmetric multiplications, then $Q(t): A \rightarrow A^{t\Sigma_2}$ is also a map of objects equipped with symmetric multiplications.*

3.4 An example

We revisit our example $k^{t\Sigma_2}$, but to avoid confusion we change the name of the generator: $k_*^{t\Sigma_2} = k((s))$. From the equivalence $k^{h\Sigma_2} = F(B\Sigma_{2+}, k)$ together with properties (iii), (iv), and (v), we see that

$$Q(t)(s) = s + s^2t^{-1}.$$

The Cartan formula now determines the behavior of $Q(t)$ in general:

$$Q(t) \sum_i a_i s^i = \sum_i a_i (s + s^2t^{-1})^i.$$

3.5 Comparing the definitions

Let B denote the bimodule from Construction 3.1.5, which is equivalent to $k^{t\Sigma_2}$ as a left k -module. Let $k \rightarrow B$ extend $1 \in \pi_0 k^{t\Sigma_2}$ as a right module map.

Lemma 3.5.1 *The composite*

$$A \rightarrow B \otimes A \rightarrow T_2(A)$$

above is equivalent to the Tate diagonal Δ_2 .

Proof Indeed, first observe that by the universal property of spectra [11, 1.4.2.23],

$$\Omega^\infty : \text{Fun}^{\text{ex}}(\text{Mod}_k, \text{Sp}) \xrightarrow{\cong} \text{Fun}^{\text{lex}}(\text{Mod}_k, \text{Spaces}).$$

Now let $U : \text{Mod}_k \rightarrow \text{Sp}$ be the forgetful functor. Then $\Omega^\infty U$ is corepresented by k , so the Yoneda lemma applied to the previous observation implies that

$$\text{Map}_{\text{Fun}^{\text{ex}}(\text{Mod}_k, \text{Sp})}(U, UT_2) \simeq \Omega^\infty k^{t\Sigma_2}.$$

Since the Tate diagonal is a transformation of lax symmetric monoidal functors, the transformation $U \rightarrow UT_2$ evaluates on k to the unit $k \rightarrow k^{t\Sigma_2}$. Combining this with the previous observation we learn that the Tate diagonal is the unique transformation $U \rightarrow UT$ which corresponds to the element $1 \in \pi_0 k^{t\Sigma_2}$. \square

Thus the map

$$A \rightarrow B \otimes A \rightarrow T_2(A) \rightarrow A^{t\Sigma_2}$$

coincides with the Tate-valued Frobenius. Now observe that the last three terms are left modules over $k^{t\Sigma_2}$, so multiplication by t^{-i-1} and naturality of $(-)^{t\Sigma_2} \rightarrow \Sigma(-)_{h\Sigma_2}$ gives a commutative diagram

$$\begin{array}{ccccccc} A & \longrightarrow & B \otimes A & \longrightarrow & T_2(A) & \longrightarrow & A^{t\Sigma_2} \\ & \searrow & \downarrow & & \downarrow & & \downarrow \\ & & \Sigma^{-i-1} B \otimes A & \longrightarrow & \Sigma^{-i-1} T_2(A) & \longrightarrow & \Sigma^{-i-1} A^{t\Sigma_2} \\ & & & & \downarrow & & \downarrow \\ & & & & \Sigma^{-i} \text{Sym}^2(A) & \longrightarrow & \Sigma^{-i} A \end{array}$$

Chasing the diagram around clockwise gives the definition of Q^i in terms of the total power operation. Chasing the diagram around counterclockwise gives the definition of Q^i in terms of Construction 3.1.5. So these two constructions agree.

Now we compare with the classical construction. The equivalence $(\Sigma^n k)^{\otimes 2} \simeq \Sigma^{2n} k$ in $\text{Mod}_k^{h\Sigma_2}$ gives a commutative diagram

$$\begin{array}{ccc} \Sigma^{-1} T_2(\Sigma^n k) & \longrightarrow & \text{Sym}^2(\Sigma^n k) \\ \downarrow \simeq & & \downarrow \simeq \\ \Sigma^{2n-1} k^{t\Sigma_2} & \longrightarrow & \Sigma^{2n} k_{h\Sigma_2} \end{array}$$

Since the bottom horizontal map is surjective on homotopy, so is the top, and we see that $\Sigma^{2n} e_{i-n}$ on the lower right corresponds to $t^{-i-1}y$ on the top left, where $y \in \pi_n \Sigma^n k$ is the generator. Now let $x: S^n \rightarrow A$ be a class and form the diagram

$$\begin{array}{ccccc}
 S^{i+n} & \xrightarrow{t^{-i-1}y} & \Sigma^{-1}T_2(\Sigma^n k) & \longrightarrow & \text{Sym}^2(\Sigma^n k) \\
 & \searrow^{t^{-i-1}x} & & & \downarrow \\
 & & \Sigma^{-1}T_2(A) & \longrightarrow & \text{Sym}^2(A)
 \end{array}$$

Traversing clockwise gives $Q^i(x)$ as in Construction 3.1.4 and traversing counterclockwise gives the image of x under Q^i as in Construction 3.1.5, and this completes the argument.

4 Adem relations

The Adem relations arise from relating the iterated total power operation to a total fourth power operation. In Section 4.1 we first explain how to lift the iterated total power operation to an intermediate Tate spectrum. In Section 4.2 we show that the existence of extra symmetry on iterated multiplication allows us to factor further through a total fourth power operation. This implies a version of the Adem relations as an identity between formal Laurent series in two variables, and in Section 4.3 we essentially perform the maneuver from [5] to recover the usual Adem relations.

For notational ease we adopt the following convention in this section:

Convention 4.0.1 If $G \subseteq \Sigma_n$ is a subgroup, and \mathcal{T} denotes the family of nontransitive subgroups of G , then we denote $(-)^{t\mathcal{T}}$ by $(-)^{\tau G}$.

4.1 Iterated power operations

Suppose A is a k -module equipped with a symmetric multiplication. Iterating the multiplication gives a map

$$A^{\otimes 4} \rightarrow A$$

which need not admit an Σ_4 -equivariant structure. However, it can be made $\Sigma_2 \wr \Sigma_2$ -equivariant, so we may define a map

$$A \rightarrow T_{\Sigma_2 \wr \Sigma_2}(A) \rightarrow A^{\tau \Sigma_2 \wr \Sigma_2}.$$

Our first goal is to show that this lifts the iterated total power operation.

Proposition 4.1.1 *Let A be a k -module equipped with a symmetric multiplication. Then there is a canonical commutative diagram*

$$\begin{array}{ccc}
 & & A^{\tau\Sigma_2\wr\Sigma_2} \\
 & \nearrow & \downarrow \\
 A & \xrightarrow{Q(t)\circ Q(s)} & (A^{t\Sigma_2})^{t\Sigma_2}
 \end{array}$$

Proof First consider the diagram

$$\begin{array}{ccccc}
 T_2(A) & \longrightarrow & T_2(T_2(A)) & \longrightarrow & ((A^{\otimes 4})^{t\Sigma_2})^{t\Sigma_2} \\
 \downarrow & & \downarrow & & \downarrow \\
 A^{t\Sigma_2} & \longrightarrow & T_2(A^{t\Sigma_2}) & \longrightarrow & (A^{t\Sigma_2})^{t\Sigma_2}
 \end{array}$$

The first square commutes by naturality of the Tate diagonal applied to the map $T_2(A) \rightarrow A$. The second square commutes by naturality of the lax structure map for $(-)^{t\Sigma_2}$.

It follows that $Q(t) \circ Q(s)$ can be written as the composite

$$A \rightarrow T_2(T_2(A)) \rightarrow ((A^{\otimes 4})^{t\Sigma_2})^{t\Sigma_2} \rightarrow (A^{t\Sigma_2})^{t\Sigma_2}.$$

Now consider both $(-)^{\tau\Sigma_2\wr\Sigma_2}$ and $((-)^{t\Sigma_2})^{t\Sigma_2}$ as exact functors $\text{Mod}_k^{h\Sigma_4} \rightarrow \text{Mod}_k$. We have a natural transformation

$$(-)^{h\Sigma_2\wr\Sigma_2} \rightarrow (-)^{h\Sigma_2 \times \Sigma_2} = ((-)^{h\Sigma_2})^{h\Sigma_2} \rightarrow ((-)^{t\Sigma_2})^{t\Sigma_2},$$

where the first map is induced by the inclusion

$$\Sigma_2 \times \Sigma_2 \rightarrow (\Sigma_2 \times \Sigma_2) \rtimes \Sigma_2 = \Sigma_2 \wr \Sigma_2$$

given by the diagonal on the first factor. By the universal property of the Tate construction (Section 1.1), we get a natural transformation $(-)^{\tau\Sigma_2\wr\Sigma_2} \rightarrow ((-)^{t\Sigma_2})^{t\Sigma_2}$. In particular, applied to the multiplication map $A^{\otimes 4} \rightarrow A$, we get a commutative diagram

$$\begin{array}{ccc}
 T_{\Sigma_2\wr\Sigma_2}(A) & \longrightarrow & A^{\tau\Sigma_2\wr\Sigma_2} \\
 \downarrow & & \downarrow \\
 ((A^{\otimes 4})^{t\Sigma_2})^{t\Sigma_2} & \longrightarrow & (A^{t\Sigma_2})^{t\Sigma_2}
 \end{array}$$

Finally, the composite

$$\Gamma^{\Sigma_2\wr\Sigma_2} \rightarrow \Gamma^{\Sigma_2 \times \Sigma_2} \simeq \Gamma^2 \circ \Gamma^2 \rightarrow T_2 \circ T_2$$

yields a natural transformation $T_{\Sigma_2 \wr \Sigma_2} \rightarrow T_2 \circ T_2$ from the universal property of $T_{\Sigma_2 \wr \Sigma_2}$ as the Goodwillie derivative of $\Gamma^{\Sigma_2 \wr \Sigma_2}$. The diagram

$$\begin{array}{ccc} T_{\Sigma_2 \wr \Sigma_2} & & \\ \downarrow & \searrow & \\ T_2 \circ T_2 & \longrightarrow & (((-)\otimes^4)^{t\Sigma_2})^{t\Sigma_2} \end{array}$$

commutes by the same universal property, and the result follows. □

4.2 Adem objects

For the Adem relations to hold we need the symmetric multiplication to satisfy an extra condition.

Definition 4.2.1 We say that a k -module A equipped with a symmetric multiplication is an *Adem object* if there exists a map $\text{Sym}^4(A) \rightarrow A$ such that the diagram

$$\begin{array}{ccc} \text{Sym}^2(\text{Sym}^2(A)) & \longrightarrow & \text{Sym}^2(A) \\ \downarrow & & \downarrow \\ \text{Sym}^4(A) & \longrightarrow & A \end{array}$$

commutes up to homotopy.

Proposition 4.2.2 If A is an Adem object, then we have a commutative diagram

$$\begin{array}{ccc} & & A^{\tau\Sigma_4} \\ & \nearrow & \downarrow \\ & A^{\tau\Sigma_2 \wr \Sigma_2} & \\ & \nearrow & \downarrow \\ A & \xrightarrow{Q(t)\circ Q(s)} & (A^{t\Sigma_2})^{t\Sigma_2} \end{array}$$

Proof By Proposition 4.1.1, the bottom triangle commutes. Factor the top triangle as

$$\begin{array}{ccc} & T_4(A) & \longrightarrow & A^{\tau\Sigma_4} \\ & \nearrow & \downarrow & \downarrow \\ A & \longrightarrow & T_{\Sigma_2 \wr \Sigma_2}(A) & \longrightarrow & A^{\tau\Sigma_2 \wr \Sigma_2} \end{array}$$

The triangle commutes because each arrow is a transformation of exact, lax symmetric monoidal functors, and $U : \text{Mod}_k \rightarrow \text{Sp}$ is initial amongst such functors (Proposition 2.3.1). The square commutes by the definition of an Adem object, ie the structure of a Σ_4 -equivariant map $A^{\otimes 4} \rightarrow A$ refining the given $(\Sigma_2 \wr \Sigma_2)$ -equivariant structure. \square

Theorem 4.2.3 (Adem relations) *If A is an Adem object and $x \in \pi_* A$ is an element, then $Q(t)(Q(s)x)$ is symmetric in the variables s and t . Explicitly,*

$$\sum_{i,j} (Q^i Q^j x)(s + s^2 t^{-1})^j t^i = \sum_{i,j} (Q^i Q^j x)(t + t^2 s^{-1})^j s^i.$$

Proof By Proposition 4.2.2, the iterated total power operation factors through $A^{\tau \Sigma_4}$ and the operation which swaps s and t arises from an inner automorphism of Σ_4 which thus acts trivially on the Tate construction, whence the claim. The explicit formula follows from the basic properties of power operations, the Cartan formula, and the computation in Section 3.4. \square

4.3 Residues and relations

Now we recall how to recover the individual Adem relations using the power series identity above.

Proposition 4.3.1 *Let A be an Adem object and $x \in A_*$ a homotopy class. Then*

$$Q^i Q^j(x) = \sum_{\ell} \binom{\ell-j-1}{2\ell-i} Q^{i+j-\ell} Q^{\ell}(x).$$

Proof In the previous section we showed

$$\sum_j Q(t)(Q^j x)(s + s^2 t^{-1})^j = \sum_{k,j} (Q^k Q^j x)(t + t^2 s^{-1})^j s^k.$$

Let $u = s + s^2 t^{-1}$ and observe that this is composition invertible as a power series in s with coefficients in $k((t))$. Now,

$$Q(t)(Q^j x) = \sum_i (Q^i Q^j x) t^i$$

is the coefficient of u^j on the left-hand side, so we would like to compute the coefficient of u^j on the right-hand side. It will be convenient to reindex the right-hand side, for fixed j , as

$$\sum_{i,\ell} (Q^{i+j-\ell} Q^{\ell} x)(t + t^2 s^{-1})^{\ell} s^{i+j-\ell}.$$

Observe that $du = ds$ since $2 = 0$ in k , and hence

$$\begin{aligned} \text{res}(u^{-j-1}(Q^{i+j-\ell}Q^\ell x)(t+t^2s^{-1})^\ell s^{i+j-\ell} du) \\ = \text{res}(u^{-j-1}(Q^{i+j-\ell}Q^\ell x)(t+t^2s^{-1})^\ell s^{i+j-\ell} ds). \end{aligned}$$

Fixing i and ℓ and writing $u = st^{-1}(t+s)$ and $(t+t^2s^{-1}) = s^{-1}t(t+s)$, we have

$$u^{-j-1}(t+t^2s^{-1})^\ell s^{i+j-\ell} = t^{\ell+j+1}s^{i-2\ell-1}(t+s)^{\ell-j-1}.$$

The coefficient of s^{-1} in the previous expression is then

$$\binom{\ell-j-1}{2\ell-i} t^i$$

and the result follows. □

5 Relationship to the Steenrod algebra

In this section we restrict to the case $k = \mathbb{F}_2$ for ease of exposition. In Section 5.1 we recall the Steenrod coaction on the Tate spectrum, then in Section 5.2 we use this to give a succinct proof of the Nishida relations. Finally, in Section 5.3 we show how this determines the action of $Q(t)$ on the dual Steenrod algebra, following an idea of Bisson and Joyal.

5.1 Coaction on the Tate spectrum

The map $k = S^0 \wedge k \rightarrow k \wedge k$ gives rise to a map $k^{t\Sigma_2} \rightarrow (k \wedge k)^{t\Sigma_2}$ if we equip the source and target with trivial Σ_2 -action.

This induces a completed coaction

$$\psi_R : k((t)) \rightarrow \mathcal{A}_*((t)).$$

More generally, for any spectrum X , the composite

$$k \wedge X \simeq X \wedge k = S^0 \wedge X \wedge k \rightarrow k \wedge X \wedge k$$

gives a completed coaction $\psi_R : H_*(X)((t)) \rightarrow (H_*(X) \otimes \mathcal{A}_*)((t))$. Now recall that Milnor defined generators³ of the dual Steenrod algebra by the identity

$$\psi_R(t) = \sum \zeta_i t^{2^i}.$$

³We are following Milnor's convention and not the more recent trend of using ζ_i to denote the *conjugates* of Milnor's generators.

5.2 Nishida relations

The easier version of the Nishida relations in this context is in terms of the coaction.

Theorem 5.2.1 (Bisson–Joyal, Baker) *Let X be a spectrum equipped with an equivariant symmetric multiplication $X_{h\Sigma_2}^{\wedge 2} \rightarrow X$. Then*

$$\sum_i \psi_R(Q^i x)t^i = Q(\bar{\zeta}(t))\psi_R(x) \in (H_*X \otimes \mathcal{A}_*)((t)).$$

Proof The right coaction $k \wedge X \rightarrow (k \wedge X) \otimes_k (k \wedge k)$ is a map of *spectra* equipped with symmetric multiplications (though it is not a map of k -modules equipped with symmetric multiplications). By Remark 3.1.8 this yields a commutative diagram

$$\begin{array}{ccc} k \wedge X & \xrightarrow{\psi_R} & (k \wedge X) \otimes_k (k \wedge k) \\ \downarrow & & \downarrow \\ (k \wedge X)^{t\Sigma_2} & \xrightarrow{(\psi_R)^{t\Sigma_2}} & ((k \wedge X) \otimes_k (k \wedge k))^{t\Sigma_2} \end{array}$$

The bottom map is the completed coaction defined in the previous subsection. Thus,

$$\psi_R(Q(t)x) = Q(t)(\psi_R(x)).$$

Since ψ_R is a ring map, and $\psi_R(t) = \zeta(t)$, this becomes

$$\sum \psi_R(Q^i x)\zeta(t)^i = Q(t)(\psi_R(x)).$$

Now substitute the conjugate series $\bar{\zeta}(t)$ for t and use the relation $\zeta(\bar{\zeta}(t)) = t$. □

5.3 Action on the dual Steenrod algebra

The following description of the action of the Q^i on \mathcal{A}_* is essentially that of Bisson and Joyal [3, Section 1, Proposition 6].

Theorem 5.3.1 (Bisson–Joyal) *The total power operation on the Milnor generators ζ_i is determined implicitly by the identity*

- (1)
$$\zeta(s) + \zeta(s)^2\zeta(t)^{-1} = \sum_i (Q(t)\zeta_i)(s^{2^i} + s^{2^{i+1}}t^{-2^i}),$$
- (2)
$$t^{2^n} Q(t)\zeta_n = \left(\sum_{i \geq n+1} \zeta_i t^{2^i} \right) + \zeta(t)^{-1} \left(\sum_{i \geq n} \zeta_i^2 t^{2^{i+1}} \right).$$

Proof Write $\pi_*k^{h\Sigma_2} = k[[s]]$. Then

$$\psi_R(Q(t)s) = Q(t)\psi_R(s).$$

Now use the identities $Q(t)s = s + s^2t^{-1}$ and $\psi_R(s) = \zeta(s)$. Comparing coefficients for s^{2^n} gives a recursion for $Q(t)\zeta_n$ starting with $Q(t)\zeta_0 = Q(t)1 = 1$ and (2) solves the recursion. □

It is not difficult to extract the earlier results of Steinberger [4, Section III.2].

Corollary 5.3.2 (Steinberger) For $i \geq 2$, $Q^{2^i-2}\zeta_1 = \bar{\zeta}_i$.

Proof From Theorem 5.3.1(2) above in the case $n = 1$,

$$Q(t)\zeta_1 = t^{-1} + \zeta_1 + \zeta(t)^{-1}.$$

So, for $i \geq 2$, change of variables and a quick computation gives

$$Q^{2^i-2}\zeta_1 = \text{res}(t^{-2^i+1}\zeta(t)^{-1} dt) = \text{res}(\bar{\zeta}(u)^{-2^i+1}u^{-1} du) = \bar{\zeta}_i. \quad \square$$

Corollary 5.3.3 (Steinberger) We have $Q^{2^i}\zeta_i = \zeta_{i+1} + \zeta_i^2\zeta_1$ and $Q^{2^i}\bar{\zeta}_i = \bar{\zeta}_{i+1}$.

Proof The case $i = 0$ is evident, so assume $i \geq 1$. The coefficient of $t^0s^{2^i+1}$ on the right-hand side of Theorem 5.3.1(1) is visibly $Q^{2^i}\zeta_i + Q^0(\zeta_i) = Q^{2^i}\zeta_i$. The constant term of $\zeta(t)^{-1}$ is ζ_1 , so the coefficient of $t^0s^{2^i+1}$ on the left-hand side is $\zeta_{i+1} + \zeta_i^2\zeta_1$. The other identity follows from this one by induction and the defining relation for conjugation. □

References

- [1] **J Adem**, *The iteration of the Steenrod squares in algebraic topology*, Proc. Nat. Acad. Sci. U.S.A. 38 (1952) 720–726 MR Zbl
- [2] **A Baker**, *Power operations and coactions in highly commutative homology theories*, Publ. Res. Inst. Math. Sci. 51 (2015) 237–272 MR Zbl
- [3] **T P Bisson, A Joyal**, *Q-rings and the homology of the symmetric groups*, from “Operads: proceedings of renaissance conferences” (J-L Loday, J D Stasheff, A A Voronov, editors), Contemp. Math. 202, Amer. Math. Soc., Providence, RI (1997) 235–286 MR Zbl
- [4] **R R Bruner, J P May, J E McClure, M Steinberger**, *H_∞ ring spectra and their applications*, Lecture Notes in Math. 1176, Springer (1986) MR Zbl

- [5] **S R Bullett, I G Macdonald**, *On the Adem relations*, *Topology* 21 (1982) 329–332 MR Zbl
- [6] **S Glasman, T Lawson**, *Stable power operations*, preprint (2020) arXiv 2002.02035
- [7] **J P C Greenlees**, *Representing Tate cohomology of G -spaces*, *Proc. Edinburgh Math. Soc.* 30 (1987) 435–443 MR Zbl
- [8] **J P C Greenlees**, *Tate cohomology in commutative algebra*, *J. Pure Appl. Algebra* 94 (1994) 59–83 MR Zbl
- [9] **J P C Greenlees, J P May**, *Generalized Tate cohomology*, *Mem. Amer. Math. Soc.* 543, Amer. Math. Soc., Providence, RI (1995) MR Zbl
- [10] **J Lurie**, *Derived algebraic geometry, XIII: Rational and p -adic homotopy theory*, preprint (2011) Available at <https://www.math.ias.edu/~lurie/papers/DAG-XIII.pdf>
- [11] **J Lurie**, *Higher algebra*, book project (2017) Available at <https://url.msp.org/Lurie-HA>
- [12] **J P May**, *A general algebraic approach to Steenrod operations*, from “The Steenrod algebra and its applications” (F P Peterson, editor), *Lecture Notes in Math.* 168, Springer (1970) 153–231 MR Zbl
- [13] **T Nikolaus**, *Stable ∞ -Operads and the multiplicative Yoneda lemma*, preprint (2016) arXiv 1608.02901
- [14] **T Nikolaus, P Scholze**, *On topological cyclic homology*, *Acta Math.* 221 (2018) 203–409 MR Zbl
- [15] **N E Steenrod**, *Cohomology operations*, *Annals of Mathematics Studies* 50, Princeton Univ. Press (1962) MR Zbl
- [16] **R Steiner**, *Homology operations and power series*, *Glasgow Math. J.* 24 (1983) 161–168 MR Zbl

*Department of Mathematics, West Virginia University
Morgantown, WV, United States*

dylan.wilson2@mail.wvu.edu

Received: 26 November 2020 Revised: 5 February 2022

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23 Issue 7 (pages 2925–3415) 2023

- Differential geometric invariants for time-reversal symmetric Bloch bundles, II: The low-dimensional “quaternionic” case 2925
GIUSEPPE DE NITTIS and KIYONORI GOMI
- Detecting isomorphisms in the homotopy category 2975
KEVIN ARLIN and J DANIEL CHRISTENSEN
- Mod 2 power operations revisited 2993
DYLAN WILSON
- The Devinatz–Hopkins theorem via algebraic geometry 3015
ROK GREGORIC
- Neighboring mapping points theorem 3043
ANDREI V MALYUTIN and OLEG R MUSIN
- Stable cohomology of the universal degree d hypersurface in \mathbb{P}^n 3071
ISHAN BANERJEE
- On the wheeled PROP of stable cohomology of $\text{Aut}(F_n)$ with bivariant coefficients 3089
NARIYA KAWAZUMI and CHRISTINE VESPA
- Anchored foams and annular homology 3129
ROSTISLAV AKHMECHET and MIKHAIL KHOVANOV
- On a problem of Hopf for circle bundles over aspherical manifolds with hyperbolic fundamental groups 3205
CHRISTOFOROS NEOFYTIDIS
- The mod 2 cohomology of the infinite families of Coxeter groups of type B and D as almost-Hopf rings 3221
LORENZO GUERRA
- Operads in unstable global homotopy theory 3293
MIGUEL BARRERO
- On some p -differential graded link homologies, II 3357
YOU QI and JOSHUA SUSSAN
- Leighton’s theorem and regular cube complexes 3395
DANIEL J WOODHOUSE