

AG
T

*Algebraic & Geometric
Topology*

Volume 23 (2023)

Neighboring mapping points theorem

ANDREI V MALYUTIN

OLEG R MUSIN



Neighboring mapping points theorem

ANDREI V MALYUTIN

OLEG R MUSIN

We introduce and study a new family of theorems extending the class of Borsuk–Ulam and topological Radon type theorems. The defining idea for this new family is to replace requirements of the form “the image of a subset that is large in some sense is a singleton” with requirements of the milder form “the image of a subset that is large in some sense is a subset that is small in some sense”. This approach covers the case of mappings $S^m \rightarrow \mathbb{R}^n$ with $m < n$ and extends to wider classes of spaces.

An example of a statement from this new family is the following theorem. Let f be a continuous map of the boundary $\partial\Delta^n$ of the n -dimensional simplex Δ^n to a contractible metric space M . Then $\partial\Delta^n$ contains a subset E such that E (is “large” in the sense that it) intersects all facets of Δ^n and the image $f(E)$ (is “small” in the sense that it) is either a singleton or a subset of the boundary ∂B of a metric ball $B \subset M$ whose interior does not meet $f(\partial\Delta^n)$.

We generalize this theorem to noncontractible normal spaces via covers and deduce a series of its corollaries. Several of these corollaries are similar to the topological Radon theorem.

55M20, 55M25, 55P05

1 Introduction

We introduce and study a new family of theorems extending the class of Borsuk–Ulam- and topological Radon-type theorems (though none of our theorems is a generalization of the Borsuk–Ulam or topological Radon theorem itself). By the Borsuk–Ulam- and topological Radon-type theorems we mean those stating that a continuous map takes a “wide” set of some specific kind to a point. Let us list several of the most influential examples:

- *The Borsuk–Ulam theorem* itself says that every continuous map of a Euclidean n -sphere S^n into Euclidean n -space \mathbb{R}^n identifies two antipodes.

- *The Hopf theorem* states that, if X is a compact Riemannian n -manifold and $f: X \rightarrow \mathbb{R}^n$ is a continuous map, then, for any $\delta > 0$, there exists a geodesic $\gamma: [0, \delta] \rightarrow X$ of length δ such that $f(\gamma(0)) = f(\gamma(\delta))$.
- *The topological Radon theorem* says that if P is a convex n -polytope, then any continuous map $\partial P \rightarrow \mathbb{R}^{n-1}$ identifies two points from disjoint faces.
- *The topological Tverberg theorem* says that, if $d \geq 1$ is an integer, r is a prime power and P is a convex $(r-1)(d+1)$ -polytope, then any continuous map $\partial P \rightarrow \mathbb{R}^d$ identifies r points from r pairwise disjoint faces.

See eg Steinlein [35; 36], Matoušek [22], Karasev [17], Akopyan, Karasev and Volovikov [3], Frick [11], Bárány, Blagojević and Ziegler [5], Skopenkov [33] and Bárány and Soberón [6] for more examples, including various extensions and generalizations for \mathbb{Z}_p -spaces, maps between manifolds, matroids and colored versions. Another related family is Knaster's conjecture-type theorems (see Matschke [23]).

All of these examples involve rigid dimensional restrictions. It is a natural question whether the maps not satisfying these restrictions have any properties of the Borsuk–Ulam kind. In particular, we are interested in whether the Borsuk–Ulam theorem has any reasonable extensions to the case of mappings $\mathbb{S}^m \rightarrow \mathbb{R}^n$ with $m < n$ (a related idea appears in Adams, Bush and Frick [2]).

Extensions of this kind are found in a new class we study. This class emerges by replacing conditions of the form “*the image of a subset that is large in some sense is a singleton*” with conditions of the milder form “*the image of a subset that is wide in some sense is a subset that is restricted in some sense*”. This approach covers the case of mappings $\mathbb{S}^m \rightarrow \mathbb{R}^n$ with $m < n$ and extends to wider classes of spaces.

Here is an example for the simplest nondegenerate case $\mathbb{S}^1 \rightarrow \mathbb{R}^2$:

Proposition 1 (Malyutin [20]) *Let a, b and c be three closed arcs covering the circle \mathbb{S}^1 such that no two of them cover \mathbb{S}^1 , and let $f: \mathbb{S}^1 \rightarrow \mathbb{R}^2$ be a continuous map. Then either $f(a) \cap f(b) \cap f(c) \neq \emptyset$ or each of $f(a)$, $f(b)$ and $f(c)$ touches a closed Euclidean disk $D^2 \subset \mathbb{R}^2$ whose interior does not meet $f(\mathbb{S}^1)$.*

Proposition 1 works for plane curves and knot diagrams and has a corollary with applications in knot theory (see [20]). We formulate this corollary here. Let $\gamma: \mathbb{S}^1 \rightarrow \mathbb{R}^2$ be a regular smooth plane curve in general position (that is, its only singularities are transversal double points). By an *edge* of γ we mean the closure of a component of the set $\gamma(\mathbb{S}^1) \setminus V$, where V is the set of double points of γ . We say that two edges I and J




Figure 1: For Proposition 1. The circle ∂D^2 touches $f(a)$, $f(b)$ and $f(c)$

of γ are *neighboring edges* or *neighbors* if there exists a component Q of $\mathbb{R}^2 \setminus \gamma(\mathbb{S}^1)$ such that the boundary ∂Q contains both I and J . We say that two edges I and J of γ are *consecutive* if the union $I \cup J$ coincides with the image $\gamma(\alpha)$ of a (connected) arc α in \mathbb{S}^1 . We denote by ρ the maximal metric on the set $E(\gamma)$ of edges of γ in the class of metrics satisfying the condition “ $\rho(I, J) = 1$ whenever I and J are consecutive edges of γ ”.

Proposition 2 [20] *If the curve γ has k double points, then γ has a pair of neighboring edges I and J with $\rho(I, J) \geq \frac{2}{3}k$.*

Proposition 1 readily implies Proposition 2 if we choose the arcs a, b and c appropriately. Proposition 2 appears in [20] as an auxiliary lemma (Lemma 5.1) needed to obtain a series of statements related to knot theory. In [20], this lemma is deduced from the topological Helly theorem (see Bogatyĭ [7] and Montejano [24]). The statement of Proposition 2 was one of the starting points for our study.

Here, we generalize Proposition 1 to noncontractible normal spaces via covers. The generalizations and their corollaries will be formulated in the next sections, after definitions. Our method is based on obstruction theory and uses a variation of the concept of non-nullhomotopic covers introduced by Musin [27; 28].

Acknowledgments The authors are grateful to Florian Frick, Sergei Ivanov, Roman Karasev, Gaiane Panina and Arkadiy Skopenkov for helpful discussions and comments. Also, the authors are grateful to the referees for helpful remarks and suggestions.

Malyutin was supported by RFBR by the research project 20-01-00070. Musin is supported by the Ministry of Science and Higher Education of the Russian Federation, agreement 075-15-2019-1620 dated 08/11/2019.

2 Definitions and results

Throughout this paper we mainly consider normal topological spaces,¹ all simplicial complexes and covers will be finite, all manifolds will be both compact and PL, S^n will denote the n -dimensional sphere, Δ^n will denote the n -dimensional simplex and $\text{sk}_k(\Delta^n)$ will denote the k -skeleton of Δ^n . We shall denote the set of homotopy classes of continuous maps $V \rightarrow W$ by $[V, W]$. The nerve of a (finite) collection \mathcal{S} of sets will be denoted by $\mathcal{N}(\mathcal{S})$. When this does not cause confusion we use the same notation for an abstract simplicial complex and its underlying space (carrier).

The further exposition in this section is structured as follows: first we give a chain of successively stronger generalizations of Proposition 1 (Theorem 4 is the weakest, Theorem 36 is the strongest); then we present a family of corollaries (all but one of which follow from Theorem 4).

2.1 Spherical f -neighbors

All of the following generalizations and corollaries replace the condition “the image is a singleton” appearing in the Borsuk–Ulam-type theorems with the following milder condition of “spherical neighboring”:

Definition 3 (spherical f -neighbors) Let X be a set, let Y be a metric space and let $f: X \rightarrow Y$ be a map. We say that a subset $N \subset X$ is a *set of spherical f -neighbors* if N contains at least two points and the image $f(N)$ is either a point or a subset of the boundary ∂B of a metric ball² $B \subset Y$ whose interior does not meet $f(X)$. If a two-point set $\{p, q\}$ is a set of spherical f -neighbors, we say that p and q are *spherical f -neighbors*. (See Figure 2.)

The first extension generalizes Proposition 1 to the case of spheres of arbitrary dimension and replaces Euclidean spaces with arbitrary contractible metric spaces. (Recall that *facets* of a polytope of dimension n are its faces of dimension $n - 1$.)

¹A topological space X is *normal* if any two disjoint closed sets of X are contained in disjoint open sets of X ; see [32, page 446] for equivalent definitions via the Urysohn and shrinking lemmas.

²By a *metric ball* in a metric space (Y, d) with metric d we mean a subset of the form $\{y \in Y \mid d(y, x) \leq R\}$ with $x \in Y$ and $R \geq 0$.




Figure 2: The image of a set $\{p_1, p_2, p_3\}$ of spherical f -neighbors

Theorem 4 Let f be a continuous map of the boundary $\partial\Delta^n$ of the n -dimensional simplex Δ^n to a contractible metric space M . Then a set of spherical f -neighbors intersects all facets of Δ^n .

Proof Let d denote the metric on M . If z is a point and N is a subset in M , we write

$$d(z, N) := \inf_{p \in N} d(z, p).$$

Let $\Delta_1, \dots, \Delta_{n+1}$ be the facets of Δ^n . For each $i \in \{1, \dots, n + 1\}$, we set

$$E_i := \{z \in M \mid d(z, f(\Delta_i)) = d(z, f(\partial\Delta^n))\}.$$

Observe that E_i contains $f(\Delta_i)$ and is closed, so $\{E_1, \dots, E_{n+1}\}$ is a closed cover of M . Since M is contractible, f extends to a continuous map $F: \Delta^n \rightarrow M$. Then $\{F^{-1}(E_1), \dots, F^{-1}(E_{n+1})\}$ is a closed cover of Δ^n extending the closed cover $\{\Delta_1, \dots, \Delta_{n+1}\}$ of $\partial\Delta^n$. By the Knaster–Kuratowski–Mazurkiewicz (KKM) lemma, the elements of $\{F^{-1}(E_1), \dots, F^{-1}(E_{n+1})\}$ have a common point p . Then $x = F(p)$ lies in $E_1 \cap \dots \cap E_{n+1}$. Then either $x \in f(\partial\Delta^n)$, so that x belongs to all of $f(\Delta_i)$ by the definition of E_i ; or $r = d(x, f(\partial\Delta^n)) > 0$ and the ball $B_r(x)$ of radius r centered at x touches all of $f(\Delta_i)$ while its interior does not meet $f(\partial\Delta^n)$. \square

We generalize Theorem 4 by replacing the set of facets with a more general class of covers as in the KKM lemma.

2.2 KKM covers and spherical f -neighbors

Definition 5 (KKM covers) Let Δ^{n+1} be an $(n+1)$ -dimensional simplex with vertices labeled v_1, \dots, v_{n+2} . A closed cover $\{C_1, \dots, C_{n+2}\}$ of the n -sphere S^n is called a *KKM cover* if there exists a homeomorphism $h: S^n \rightarrow \partial\Delta^{n+1}$ such that, for each $J \subset \{1, \dots, n+2\}$, the convex hull of the vertices v_j with $j \in J$ is covered by the union $\bigcup_{j \in J} h(C_j)$.

The argument in the proof of Theorem 4 also proves the following theorem:

Theorem 6 Let \mathcal{C} be a KKM cover of the n -sphere S^n , and let $f: S^n \rightarrow M$ be a continuous map to a contractible metric space M . Then a set of spherical f -neighbors intersects all elements of \mathcal{C} .

The key role in Theorem 6 is played by the properties of the cover, and not by the fact that the underlying space is a sphere. To move on to the next generalization, we define non-nullhomotopic covers (we generalize the concept of non-nullhomotopic covers given in [27; 28]).

2.3 Non-nullhomotopic covers and spherical f -neighbors

Let X be a normal topological space and let $\mathcal{U} = \{U_1, \dots, U_n\}$ be an open cover of X . Let $\mathcal{N}(\mathcal{U})$ be the nerve of \mathcal{U} . Let $\Phi = \{\varphi_1, \dots, \varphi_n\}$ be a partition of unity subordinate to \mathcal{U} . Let v_1, \dots, v_n be the vertices of the $(n-1)$ -dimensional unit simplex Δ^{n-1} , where

$$\Delta^{n-1} := \{x \in \mathbb{R}^n \mid x_i \geq 0, x_1 + \dots + x_n = 1\}.$$

For each i , we identify the vertex of $\mathcal{N}(\mathcal{U})$ corresponding to U_i with v_i , so that $\mathcal{N}(\mathcal{U})$ becomes a subcomplex of Δ^{n-1} . We set

$$h_{\mathcal{U}, \Phi}(x) := \sum_{i=1}^n \varphi_i(x) v_i.$$

Then $h_{\mathcal{U}, \Phi}$ is a continuous map from X to $\mathcal{N}(\mathcal{U}) \subset \Delta^{n-1}$. Since the linear homotopy $\Theta(t) = (1-t)\Phi + t\Psi$ of two partitions of unity Φ and Ψ subordinate to \mathcal{U} induces a homotopy between the corresponding maps, it follows that the homotopy class $[h_{\mathcal{U}, \Phi}]$ in $[X, \mathcal{N}(\mathcal{U})]$, where by $[V, W]$ we denote the set of homotopy classes of continuous maps $V \rightarrow W$, does not depend on Φ (see [27, Lemma 1.6]). We denote this class in $[X, \mathcal{N}(\mathcal{U})]$ by $[\mathcal{U}]$.

The homotopy classes of covers are also well defined for closed sets. Indeed, in a normal space, any finite closed cover has an open extension with the same nerve (see eg [25, Theorem 1.3; 16, pages 31–33]). Furthermore, if $\mathcal{C} = \{C_1, \dots, C_n\}$ is a closed cover of a normal space X and $\mathcal{S} = \{S_1, \dots, S_n\}$ and $\mathcal{U} = \{U_1, \dots, U_n\}$ are two open covers such that $S_i \cap U_i$ contains C_i for all i that have the same nerve $\mathcal{N}(\mathcal{S}) = \mathcal{N}(\mathcal{U}) = \mathcal{N}(\mathcal{C})$, then each partition of unity subordinate to the open cover

$$\mathcal{T} := \{S_1 \cap U_1, \dots, S_n \cap U_n\}$$

is also subordinate to both \mathcal{S} and \mathcal{U} . This implies that $[S] = [\mathcal{T}] = [\mathcal{U}]$ in $[X, \mathcal{N}(\mathcal{C})]$ due to the independence of the choice of partition of unity mentioned above. Then we set

$$[\mathcal{C}] := [S] = [\mathcal{T}] = [\mathcal{U}].$$

Definition 7 (non-nullhomotopic covers) We say that an open or closed cover \mathcal{C} of a normal topological space X is *non-nullhomotopic* if the corresponding homotopy class $[\mathcal{C}]$ in $[X, \mathcal{N}(\mathcal{C})]$ contains no constant map.

Remark 8 Any non-nullhomotopic map $X \rightarrow K$ to a finite simplicial complex yields non-nullhomotopic covers on X ; to obtain an example, take the inverse images of all elements in one of the collections

- open stars of vertices of K ,
- stars of vertices of K in its first barycentric subdivision,
- maximal simplexes of K .

Definition 9 (homotopy ranks of maps) Let X be a topological space, let K be a finite simplicial complex and let $h: X \rightarrow K$ be a continuous map. Let Δ_K be the simplex spanned by the vertices of K , so K is a subcomplex of Δ_K . We define the *homotopy rank* $\text{rk}(h)$ of h to be the least nonnegative integer k such that h is nullhomotopic in $K \cup \text{sk}_k(\Delta_K)$, where sk_k stands for the k -skeleton.³ (Since Δ_K is contractible, the homotopy rank is well defined and does not exceed the dimension of Δ_K .)

Remark 10 In terms of Definition 9, h is nullhomotopic if and only if $\text{rk}(h) = 0$.

Definition 11 (ranks of covers) We define the (*homotopy*) *rank* $\text{rk}(\mathcal{C})$ of a (closed or open) finite cover \mathcal{C} of a normal space X to be the homotopy rank of maps $X \rightarrow \mathcal{N}(\mathcal{C})$ in the class $[\mathcal{C}]$ determined by \mathcal{C} .

Remark 12 A cover is non-nullhomotopic if and only if it is of nonzero rank.

³We say that $K \cup \text{sk}_k(\Delta_K)$ is the k -*exoskeleton* of K .

Remark 13 Since $\text{sk}_m(\Delta^m) = \Delta^m$ is contractible, it follows that the rank of an n -element cover does not exceed $n - 1$.

Definition 14 (principal covers) An n -element cover ($n \geq 2$) of rank $n - 1$ is said to be *principal*.

Remark 15 Since any proper nonempty subcomplex of $\partial\Delta^m$ is contractible in $\partial\Delta^m$, it follows that a cover is principal if and only if it is non-nullhomotopic and its nerve is the boundary of a simplex.

Remark 16 Remark 15 implies that no principal cover has a proper subcollection of elements with empty intersection; in particular, no principal cover has disjoint elements.

Remark 17 Any non-nullhomotopic map $X \rightarrow \mathbb{S}^k$ to the k -sphere yields a principal cover of X of rank $k + 1$ (see Remark 8 and [27, Theorem 1.5]). Thus, a space can have principal covers of distinct ranks.

Remark 18 (conditions for cover non-nullhomotopicity) If the composition of continuous maps is non-nullhomotopic, then each of them is non-nullhomotopic.

- On the one hand, this implies that any refinement of a cover of rank k has rank at least k . In particular, any refinement of a non-nullhomotopic cover is non-nullhomotopic.
- On the other hand, this implies that, if $f: X \rightarrow Y$ is a continuous map of normal spaces and $\mathcal{C} = \{C_1, \dots, C_n\}$ is a closed cover of Y such that the dimension of the nerve $\mathcal{N}(\mathcal{C})$ is less than the rank $\text{rk}(\tilde{\mathcal{C}})$ of the induced cover $\tilde{\mathcal{C}} = \{f^{-1}(C_1), \dots, f^{-1}(C_n)\}$, then $\text{rk}(\mathcal{C}) \geq \text{rk}(\tilde{\mathcal{C}})$. In particular, if the induced cover is principal and $\bigcap_{i=1}^n C_i = \emptyset$, then \mathcal{C} is principal. (See Lemma 52.)

We now have all the definitions needed to replace spheres in Theorem 6 with general “noncontractible” spaces.

Theorem 19 Let X be a compact normal space, let M be a contractible metric space and let $f: X \rightarrow M$ be a continuous map. Then, for any non-nullhomotopic cover \mathcal{C} of X , a set of spherical f -neighbors intersects at least $\text{rk}(\mathcal{C}) + 1$ elements of \mathcal{C} . In particular, for any principal cover, a set of spherical f -neighbors intersects all elements of the cover.




Figure 3: An example with disconnected $X = \mathbb{S}^1 \cup \{b', c'\}$.

Theorem 19 implies Theorem 6 because each KKM cover either is principal or all of its elements have a common point; furthermore, the maps in the homotopy class $[C]$ corresponding to each principal KKM cover C are of degree one, so $[C]$ contains a homeomorphism (see [28, Corollaries 2.1–2.3]).

Remark 20 X in Theorem 19 is not assumed to be connected. Figure 3 shows an example with $X = \mathbb{S}^1 \cup \{b', c'\}$ (compare Figure 1).

Remark 21 Combining the idea that X in Theorem 19 is not necessarily connected with switching attention to the image of the cover leads to generalizations of Helly’s theorem and the KKM lemma. See also Lemma 52 below. We do not develop this line here.

2.4 EP triples and ranks, and spherical f -neighbors

We are going to upgrade Theorem 19 to the more general Theorem 36, which covers the case of maps to not necessarily contractible spaces. In order to state and prove Theorem 36, we introduce concepts of Eilenberg–Pontryagin and Knaster–Kuratowski–Mazurkiewicz ranks.

Definition 22 (Eilenberg–Pontryagin triples and ranks) Let Z be a topological space with a subspace A , let K be a finite simplicial complex and let $[h]$ be a homotopy class in $[A, K]$. We say that $(Z, A, [h])$ is an *Eilenberg–Pontryagin triple* if no map in $[h]$ extends to a continuous map $Z \rightarrow K$.

We define the *Eilenberg–Pontryagin rank (EP rank)* $\text{rk}(Z, A, [h])$ of the triple $(Z, A, [h])$ to be the least nonnegative integer k such that there exists a continuous map $H: Z \rightarrow K \cup \text{sk}_k(\Delta_K)$ whose restriction $H|_A$ is homotopic in $K \cup \text{sk}_k(\Delta_K)$ to the maps of $[h]$, where Δ_K is the simplex spanned by the vertices of K and containing K as a subcomplex. (Since Δ_K is contractible, the EP rank is well defined and does not exceed the dimension of Δ_K .)

Remark 23 In terms of Definition 22, a triple $(Z, A, [h])$ is Eilenberg–Pontryagin if and only if it is of nonzero EP rank (because $K \cup \text{sk}_0(\Delta_K) = K$).

Remark 24 Since any constant map extends to any ambient space, it follows that in terms of Definitions 9 and 22, for any Z, A, K and h ,

$$\text{rk}(Z, A, [h]) \leq \text{rk}(h).$$

Furthermore, if A is contractible in Z , then

$$\text{rk}(Z, A, [h]) = \text{rk}(h).$$

In particular, if \mathcal{C} is a finite closed cover of A and A is contractible in Z , then

$$\text{rk}(Z, A, [\mathcal{C}]) = \text{rk}(\mathcal{C}).$$

Example 25

- If $Z = \Delta^n$, $A = K = \partial\Delta^n$ and $h = \text{id}$, then $\text{rk}(Z, A, [h]) = \text{rk}(h) = n$.
- If $Z = K = \Delta^n$, $A = \partial\Delta^n$, and $h = \text{id}$, then $\text{rk}(Z, A, [h]) = \text{rk}(h) = 0$.

Example 26 We have $\text{rk}(Z, A, [h]) = 0$ whenever A is a retract of Z .

Example 27 Let W be an orientable, compact PL m -manifold with connected nonempty boundary ∂W , and let $h: \partial W \rightarrow \partial\Delta^n$ be a continuous map. Then $\text{rk}(h) \in \{0, n\}$, $\text{rk}(W, \partial W, [h]) \in \{0, n\}$ and $\text{rk}(W, \partial W, [h]) \leq \text{rk}(h)$.

- If $m = n$, then $\text{rk}(W, \partial W, [h]) = \text{rk}(h)$ (this follows from the Hopf degree theorem; see the proof of Corollary 39 below).
- If $W = \Delta^m$, then $\text{rk}(W, \partial W, [h]) = \text{rk}(h)$ (because Δ^m is contractible; see Remark 24).
- Results of [29] imply however that, for any m and n with nontrivial $\pi_{m-1}(\mathbb{S}^{n-1})$ and for any non-nullhomotopic $h: \mathbb{S}^{m-1} \rightarrow \partial\Delta^n$, there exists an m -manifold W with $\partial W = \mathbb{S}^{m-1}$ such that h extends to a continuous map $W \rightarrow \partial\Delta^n$, so $\text{rk}(W, \partial W, [h]) = 0$ and $\text{rk}(h) = n$.

Definition 28 (Knaster–Kuratowski–Mazurkiewicz rank) Let Z be a topological space and let $\mathcal{S} = \{S_1, \dots, S_n\}$ be a collection of subsets in Z . We say that the pair (Z, \mathcal{S}) is a *Knaster–Kuratowski–Mazurkiewicz (KKM) system* if no closed cover $\{E_1, \dots, E_n\}$ of Z with $S_i \subset E_i$ for all i has the same nerve as \mathcal{S} .

We define the *KKM rank* $\text{rk}(Z, \mathcal{S})$ of the pair (Z, \mathcal{S}) to be the least integer k such that there exists a closed cover $\mathcal{E} = \{E_1, \dots, E_n\}$ of Z with $S_i \subset E_i$ for all i such that the dimension of $\mathcal{N}(\mathcal{E}) \setminus \mathcal{N}(\mathcal{S})$ is k .

Remark 29 In terms of Definition 28, a pair (Z, \mathcal{S}) is a KKM system if and only if it is of nonzero KKM rank.

Example 30 If $Z = \Delta^m$ and $\mathcal{S} = \{S_1, \dots, S_{m+1}\}$ is a KKM cover of $\partial\Delta^m$, then $\text{rk}(Z, \mathcal{S}) = m$ by the KKM lemma.

Example 31 We have $\text{rk}(Z, \mathcal{S}) = 0$ whenever \mathcal{S} is a closed cover of a retract of Z .

Example 32 We have $\text{rk}(Z, \{S_1, \dots, S_n\}) = 0$ whenever $\bigcap_{i=1}^n S_i \neq \emptyset$.

Lemma 33 Let a normal space Z contain a normal space A as a subspace, let \mathcal{C} be a closed cover of A , and let $[\mathcal{C}]$ be the corresponding homotopy class in $[A, \mathcal{N}(\mathcal{C})]$, where $\mathcal{N}(\mathcal{C})$ is the nerve. Then the EP rank of the triple $(Z, A, [\mathcal{C}])$ does not exceed the KKM rank of the system (Z, \mathcal{C}) :

$$\text{rk}(Z, A, [\mathcal{C}]) \leq \text{rk}(Z, \mathcal{C}).$$

Furthermore, if A is closed in Z , then

$$\text{rk}(Z, A, [\mathcal{C}]) = \text{rk}(Z, \mathcal{C}).$$

Lemma 33 is proved in the next section.

Example 34 (showing that the closedness requirement of A in the second part of Lemma 33 is essential) If X is a compact normal space, $\mathcal{C}' = \{C'_1, \dots, C'_n\}$ is a closed cover of X with $\bigcap_{i=1}^n C_i = \emptyset$ and each C'_i nonempty, Z is the cone over X , z_0 is the top of Z , $A = Z \setminus \{z_0\}$, C''_i is the subcone in Z over C'_i , $C_i = C''_i \setminus \{z_0\}$, and $\mathcal{C} = \{C_1, \dots, C_n\}$, then the KKM rank $\text{rk}(Z, \mathcal{C})$ is $n - 1$ and the EP rank $\text{rk}(Z, A, [\mathcal{C}])$ is one more than the dimension of the nerve $\mathcal{N}(\mathcal{C}')$. (See Figure 4 with $X = \mathbb{S}^1$.) For example, if $n > 2$ and the elements of \mathcal{C}' are pairwise disjoint, then

$$\text{rk}(Z, A, [\mathcal{C}]) = 1 < n - 1 = \text{rk}(Z, \mathcal{C}).$$




Figure 4: A disk with $\text{rk}(Z, A, [\mathcal{C}]) = 2 < 4 = \text{rk}(Z, \mathcal{C})$.

Remark 35 Lemma 33 implies (see Remark 24) that, given a compact normal space A with a finite closed cover \mathcal{C} , for any ambient normal space $Z \supset A$, $\text{rk}(Z, A, [\mathcal{C}]) = \text{rk}(Z, \mathcal{C}) \leq \text{rk}(\mathcal{C})$, while $\text{rk}(Z, A, [\mathcal{C}]) = \text{rk}(Z, \mathcal{C}) = \text{rk}(\mathcal{C})$ if A is contractible in Z . (This generalizes [27, Theorem 2.2].)

Theorem 36 *Let A be a compact normal space, let \mathcal{C} be a closed cover of A , and let $[\mathcal{C}]$ be the corresponding homotopy class in $[A, \mathcal{N}(\mathcal{C})]$, where $\mathcal{N}(\mathcal{C})$ is the nerve. Let Z be a normal space containing A as a subspace. If the triple $(Z, A, [\mathcal{C}])$ is Eilenberg–Pontryagin, with EP rank $\text{rk}(Z, A, [\mathcal{C}]) > 0$, then, for any metric space M and any continuous map $f : A \rightarrow M$ that extends to a continuous map $Z \rightarrow M$, a set of spherical f –neighbors intersects at least $\text{rk}(Z, A, [\mathcal{C}]) + 1$ elements of \mathcal{C} .*

Theorem 36 is proved in the next section.

Proof of Theorem 19 We deduce Theorem 19 from Theorem 36. Let X, M, f and \mathcal{C} be as in Theorem 19. Set $\text{Cone}(X) := (X \times [0, 1]) / (X \times \{0\})$ and identify X with $X \times \{1\} \subset \text{Cone}(X)$. (The cone is normal because X is compact and normal; see eg [30].) Definitions of ranks imply (see Remark 24) that

$$(***) \quad \text{rk}(\text{Cone}(X), X, [\mathcal{C}]) = \text{rk}(\mathcal{C}).$$

In particular, the triple $(\text{Cone}(X), X, [\mathcal{C}])$ is Eilenberg–Pontryagin since \mathcal{C} is non-nullhomotopic. Since M is contractible, it follows that f extends to a continuous map $F : \text{Cone}(X) \rightarrow M$. We apply Theorem 36 to the Eilenberg–Pontryagin triple $(\text{Cone}(X), X, [\mathcal{C}])$, with $Z = \text{Cone}(X)$ and $A = X$ in the notation of Theorem 36, and see that a set of spherical f –neighbors intersects at least $\text{rk}(\text{Cone}(X), X, [\mathcal{C}]) + 1$ elements of \mathcal{C} . Then Theorem 19 follows by (***) □

Remark 37 Theorem 36 has further refinements regarding the number of distinct sets of spherical f -neighbors intersecting the prescribed number of cover elements, but we do not develop this line here.

2.5 Corollaries

Next, we list several corollaries of Theorems 4, 6, 19 and 36. In fact, all of the following corollaries, except for Corollary 39, follow from Theorem 4.

Definition 38 A continuous map $f: A \rightarrow Y$ of an orientable, connected, closed PL manifold A to a space Y is said to be *null-cobordant* if there exists an orientable, compact PL manifold W with $\partial W = A$ and a continuous map $F: W \rightarrow Y$ such that $F|_A = f$.

Corollary 39 (see [27, Theorem 2.6]) *Let A be an orientable, connected, closed PL n -manifold and let \mathcal{C} be a non-nullhomotopic cover of A such that the nerve of \mathcal{C} is homeomorphic to the n -sphere. Then, for any metric space M and any null-cobordant map $f: A \rightarrow M$, a set of spherical f -neighbors intersects at least $n + 2$ elements of \mathcal{C} . In particular, if \mathcal{C} is principal and contains precisely $n + 2$ elements, then a set of spherical f -neighbors intersects all elements of \mathcal{C} .*

Proof If $f: A \rightarrow M$ is null-cobordant, then there is an orientable, compact PL $(n+1)$ -manifold Z with $\partial Z = A$ and a continuous map $F: Z \rightarrow M$ with $F|_A = f$. A homological argument shows that, for each continuous map $H: Z \rightarrow \mathcal{N}(\mathcal{C}) \cong \mathbb{S}^n$, the restriction $H|_A$ is of zero degree. Then the Hopf degree theorem implies that $H|_A$ is nullhomotopic. This means that the triple $(Z, A, [\mathcal{C}])$ is Eilenberg–Pontryagin and the statement follows by Theorem 36. □

Remark 40 (the dimensional restriction in Corollary 39 is essential) It is shown in [29] that any continuous map $\mathbb{S}^m \rightarrow \mathbb{S}^n$ is null-cobordant if $m > n$. Let m and n be such that $m > n$ and $\pi_m(\mathbb{S}^n)$ is nontrivial, and let $h: \mathbb{S}^m \rightarrow \partial\Delta^{n+1}$ be a non-nullhomotopic continuous map. Then there exists an orientable, compact PL $(m+1)$ -manifold W with $\partial W = \mathbb{S}^m$ and a continuous map $H: W \rightarrow \partial\Delta^{n+1}$ such that $H|_{\partial W} = h$. Let \mathcal{C} be the closed cover of ∂W composed of the inverse images of the facets of Δ^{n+1} . Then $[\mathcal{C}] = [h]$ and \mathcal{C} is principal. We embed W into a Euclidean ball B^N of large dimension and “tiny” diameter, then embed W into the product $\partial\Delta^{n+1} \times B^N$ such that the projection of this embedding to $\partial\Delta^{n+1}$ yields H , and take the induced metric on W .

Now, let $f: \partial W \rightarrow W$ be the identity map. Then f is null-cobordant but no set of spherical f -neighbors intersects all elements of \mathcal{C} if the diameter of B^N is sufficiently small.

Corollary 41 *Let M be a contractible metric space, let \mathbb{S}^n be the Euclidean unit n -sphere in Euclidean $(n+1)$ -space \mathbb{R}^{n+1} and let $f: \mathbb{S}^n \rightarrow M$ be a continuous map. Then there exists a pair $\{p, q\}$ of spherical f -neighbors such that the Euclidean distance $\|p - q\|$ is at least $\sqrt{(n+2)/n}$.*

Corollary 41 is proved in the next section.

Remark 42 In [21] we show that, if $M = \mathbb{R}^m$ with $m > n$, then the constant $\sqrt{(n+2)/n}$ in Corollary 41 (the Euclidean distance between the centers of adjacent $(n-1)$ -simplices of the regular triangulation of \mathbb{S}^n) can be replaced with $\sqrt{2(n+2)/(n+1)}$ (the Euclidean distance between vertices of the regular triangulation of \mathbb{S}^n), which is the best possible. Our proof for the Euclidean case $M = \mathbb{R}^m$ is based on the Delaunay triangulations and we do not know whether it extends to all contractible M .

Corollary 43 *Let M be a contractible metric space, let P be a convex n -polytope and let $f: \partial P \rightarrow M$ be a continuous map. Then a set of spherical f -neighbors intersects at least $n + 1$ facets of P .*

Proof via Theorem 4 Corollary 43 is an “equivalent generalization” of Theorem 4 because the $(n-2)$ -skeleton of any convex n -polytope contains the $(n-2)$ -skeleton of the n -simplex as a topological subspace (see [14]). \square

Proof via Theorem 19 Clearly, the cover \mathcal{C} of ∂P composed of the facets of P is non-nullhomotopic of rank n because \mathcal{C} is a *good* cover (that is, any intersection of elements in \mathcal{C} is contractible), so the nerve of \mathcal{C} has homotopy type of $\partial P \cong \mathbb{S}^{n-1}$ by the nerve theorem, while the maps in the class $[\mathcal{C}]$ are homotopy equivalences. Then a set of spherical f -neighbors intersects at least $n + 1$ facets of P by Theorem 19. \square

Since any collection of $n + 1$ facets of the n -cube contains a pair of antipodal facets, Corollary 43 implies the following:

Corollary 44 *Let M be a contractible metric space, let $\partial[0, 1]^m$ be the boundary of the m -dimensional cube $[0, 1]^m$ and let $f: \partial[0, 1]^m \rightarrow M$ be a continuous map. Then there is a pair of spherical f -neighbors intersecting antipodal facets of $[0, 1]^m$.*

There exists an example of continuous map $\mathbb{S}^2 \rightarrow \mathbb{R}^3$ showing that the statement of Corollary 44 about spherical f -neighbors lying on antipodal facets holds for neither regular octahedra nor regular dodecahedra nor regular icosahedra. A weaker version of Corollary 44 where “antipodal” is replaced with “disjoint” holds for many polytopes.

2.6 Radon-type theorems

Definition 45 (weak Radon polytopes) We say that an n -polytope P is *weakly Radon* if, for any continuous map $f : \partial P \rightarrow M$ into any contractible metric space M , there is a pair of spherical f -neighbors intersecting two disjoint faces of P .

We recall some standard definitions. A *flag* polytope is a convex polytope such that every collection of pairwise intersecting facets has a nonempty intersection. A (*combinatorial*) *fullerene* is a simple 3-polytope with all facets pentagons and hexagons.

A “visual” simply checked sufficient condition for weakly Radon polytopes is provided by the so-called *belts*. A k -*belt* (or a *prismatic k -circuit*) in a 3-polytope is a cyclic sequence (F_1, \dots, F_k) of $k \geq 3$ facets in which pairs of consecutive facets (including $\{F_k, F_1\}$) are adjacent, other pairs of facets do not intersect, and no three facets have a common vertex.

- Corollary 46** (1) *If the $(n-2)$ -skeleton of a convex n -polytope P contains the $(n-2)$ -skeleton of the n -cube as a topological subspace, then P is weakly Radon.*
- (2) *Each convex 3-polytope having a k -belt with $k \geq 4$ is weakly Radon.*
- (3) *Each flag 3-polytope is weakly Radon.*
- (4) *Each fullerene is weakly Radon.*
- (5) *The regular dodecahedron and the regular icosahedron are weakly Radon.*

Proof Assertion (1) follows from Corollary 44 in an obvious way. Assertions (2) and (5) directly follow from assertion (1). Assertion (3) follows from Corollary 51(2) below. Assertion (4) is a particular case of assertion (3). \square

Definition 47 (weak Radon rank) If P is an n -polytope, Y is a metric space and $f : \partial P \rightarrow M$ is a map, we say that two facets F_1 and F_2 of P are *spherical f -neighbors* (or that the pair $\{F_1, F_2\}$ is a *pair of spherical f -neighbors*) if there is a pair $\{p, q\}$ of spherical f -neighbors with $p \in F_1$ and $q \in F_2$. We say that $f : \partial P \rightarrow M$ has *weak Radon rank m* if there are exactly m distinct pairs of facets of P such that each of these pairs is a pair of disjoint spherical f -neighbors. By the *weak Radon rank*

of a polytope P we mean the least of the weak Radon ranks of continuous maps $f: \partial P \rightarrow M$ into contractible metric spaces.

Corollary 44 allows us to obtain rough lower bounds on the weak Radon rank.

Definition 48 (cubic hemisphere) Let H be a subset of the boundary ∂P of a convex n -polytope P . We say that H is a *cubic hemisphere* if there exists a homeomorphism $h: [0, 1]^n \rightarrow P$ such that the restriction of h to the $(n-2)$ -skeleton of $[0, 1]^n$ is a topological embedding to the $(n-2)$ -skeleton of P and H is the image of the union of n facets of $[0, 1]^n$ that have a common vertex.

Definition 49 (lighthouse independence number) We say that a set Z of vertices of an n -polytope is *lighthouse independent* if no two vertices in Z share a facet (equivalently, the corresponding facets of the dual polytope are pairwise disjoint). The *lighthouse independence number* $\text{lin}(P)$ of an n -polytope P is the cardinality of a largest lighthouse independent set of P .

Remark 50 The lighthouse independence number of an n -polytope equals the cardinality of a largest set of pairwise disjoint facets of the dual polytope.

Corollary 51 (1) *Let P be a convex n -polytope. If ∂P contains k cubic hemispheres with pairwise disjoint interiors, then the weak Radon rank of P is at least $\frac{1}{2}k$.*

(2) *Let P be a flag 3-polytope (eg a fullerene). Then the weak Radon rank of P is at least half the lighthouse independence number of P .*

(3) *If P is a flag simple 3-polytope with ψ facets and g is the largest number of edges in a facet of P , then the weak Radon rank of P is at least*

$$\frac{1}{2} \left\lfloor \frac{2\psi - 7}{3g - 8} \right\rfloor,$$

where $\lfloor \cdot \rfloor$ stands for the floor function.

(4) *If P is a fullerene with ψ facets, then the weak Radon rank of P is at least*

$$\frac{1}{2} \left\lfloor \frac{1}{5}(\psi - 3) \right\rfloor.$$

(5) *The weak Radon rank of the regular dodecahedron is at least 2.*

(6) *The weak Radon rank of the regular icosahedron is at least 2.*

(7) *The weak Radon rank of the cube is 1.*

Corollary 51 is proved in the next section.

3 Proofs

Proof of Lemma 33 (1) We show that $\text{rk}(Z, A, [C]) \leq \text{rk}(Z, C)$.

Let $C = \{C_1, \dots, C_n\}$, let Δ^{n-1} denote the simplex spanned by the vertices of the nerve $\mathcal{N}(C)$ of C , so $\mathcal{N}(C)$ is a subcomplex of Δ^{n-1} , and let $r := \text{rk}(Z, C)$. By the definition of the KKM rank, there exists a closed cover $\mathcal{E} = \{E_1, \dots, E_n\}$ of Z with $C_i \subset E_i$ for all i such that the dimension of $\mathcal{N}(\mathcal{E}) \setminus \mathcal{N}(C)$ is r . Therefore, the union $\mathcal{N}(C) \cup \text{sk}_r(\Delta^{n-1})$ contains $\mathcal{N}(\mathcal{E})$. Set

$$\mathcal{E}_A := \{E_1 \cap A, \dots, E_n \cap A\}.$$

Since $C_i \subset E_i$ for all i , it follows that the nerve $\mathcal{N}(\mathcal{E}_A)$ contains $\mathcal{N}(C)$. We have

$$\mathcal{N}(C) \subset \mathcal{N}(\mathcal{E}_A) \subset \mathcal{N}(\mathcal{E}) \subset \mathcal{N}(C) \cup \text{sk}_r(\Delta^{n-1}).$$

Let $[\mathcal{E}]$ be the homotopy class in $[Z, \mathcal{N}(\mathcal{E})]$ determined by \mathcal{E} and let $F: Z \rightarrow \mathcal{N}(\mathcal{E})$ be a map in $[\mathcal{E}]$. Let $[\mathcal{E}_A]$ be the homotopy class in $[A, \mathcal{N}(\mathcal{E}_A)]$ determined by \mathcal{E}_A and let $f': A \rightarrow \mathcal{N}(\mathcal{E}_A)$ be a map in $[\mathcal{E}_A]$. Let $f: A \rightarrow \mathcal{N}(C)$ be a map in $[C] \in [A, \mathcal{N}(C)]$.

Since $E_i \cap A$ contains C_i for each i , the argument preceding Definition 7 shows that f and f' are homotopic in $\mathcal{N}(\mathcal{E}_A)$. By construction, $F|_A$ and f' are homotopic in $\mathcal{N}(\mathcal{E})$. Thus, $F|_A$ and f are homotopic in $\mathcal{N}(\mathcal{E})$ and hence in $\mathcal{N}(C) \cup \text{sk}_r(\Delta^{n-1})$ as well. By the definition of the EP rank this means that $\text{rk}(Z, A, [C]) \leq r = \text{rk}(Z, C)$.

(2) We show that $\text{rk}(Z, C) \leq \text{rk}(Z, A, [C])$ whenever A is closed in Z .

We start by constructing a specific map $A \rightarrow \mathcal{N}(C)$ from the class $[C]$. Let $C = \{C_1, \dots, C_n\}$ and let $\mathcal{N}(C)$ be a subcomplex in Δ^{n-1} (as in the first part of the proof). Since A is normal, there exists an open cover $\mathcal{U} = \{U_1, \dots, U_n\}$ of A such that U_i contains C_i for each i and the nerve of \mathcal{U} coincides with that of C (see eg [25, Theorem 1.3; 16, pages 31–33]). The Urysohn lemma for normal spaces implies that, for each i , there exists a continuous function $f_i: A \rightarrow [0, 1]$ with $f_i(C_i) = 1$ and $f_i(A \setminus U_i) = 0$. Then $\Phi = \{\varphi_1, \dots, \varphi_n\}$, where $\varphi_i := f_i / \sum_j f_j$, is a partition of unity subordinate to \mathcal{U} such that $\varphi_i^{-1}[1/n, 1]$ contains C_i for all i . Let

$$h_{\mathcal{U}, \Phi}(x) := \sum_{i=1}^n \varphi_i(x) v_i$$

be the corresponding map $A \rightarrow \mathcal{N}(C)$ representing the class $[C] = [\mathcal{U}]$ (see the construction preceding Definition 7).

Now, let $p := \text{rk}(Z, A, [C])$. Then, by the definition of the EP rank, there exists a continuous map $F: Z \rightarrow \mathcal{N}(C) \cup \text{sk}_p(\Delta^{n-1})$ such that the restriction $F|_A$ is homotopic to $h_{\mathcal{U}, \Phi}$ in $\mathcal{N}(C) \cup \text{sk}_p(\Delta^{n-1})$. The generalizations of Borsuk’s homotopy extension theorem obtained in [26; 34] imply that, since A is closed in Z , there exists a continuous map $G: Z \rightarrow \mathcal{N}(C) \cup \text{sk}_p(\Delta^{n-1})$ with $G|_A = h_{\mathcal{U}, \Phi}$. Then the collection of subsets

$$\mathcal{G} := \{G_1^{-1}[1/n, 1], \dots, G_n^{-1}[1/n, 1]\},$$

where G_1, \dots, G_n are the coordinate functions of G , is a closed cover of Z such that $G_i^{-1}[1/n, 1]$ contains C_i for all i and the nerve $\mathcal{N}(\mathcal{G})$ is contained in $\mathcal{N}(C) \cup \text{sk}_p(\Delta^{n-1})$, so the dimension of $\mathcal{N}(\mathcal{G}) \setminus \mathcal{N}(C)$ is at most p . By the definition of the KKM rank, this means that $\text{rk}(Z, [C]) \leq p = \text{rk}(Z, A, [C])$. □

Now we state and prove Lemmas 52 and 54, and then deduce Theorem 36 from Lemmas 33, 52 and 54.

Lemma 52 *Let $(Z, \mathcal{C} = \{C_1, \dots, C_n\})$ be a KKM system of rank $r > 0$, let $f: Z \rightarrow Z'$ be a continuous map to a topological space Z' , and let $\mathcal{C}' = \{C'_1, \dots, C'_n\}$ be a family of subsets in Z' such that $f(C_i) \subset C'_i$ for all i . Then either $\{1, \dots, n\}$ contains a subset J of cardinality $r + 1$ such that $\bigcap_{j \in J} C_j = \emptyset$ and $\bigcap_{j \in J} C'_j \neq \emptyset$ or (Z', \mathcal{C}') is a KKM system of rank at least r .*

Remark 53 In Lemma 52, two key special cases are $C'_i = f(C_i)$ and $f = \text{id}$.

Proof If neither $\text{rk}(Z', \mathcal{C}') \geq r$ nor $\{1, \dots, n\}$ contains J with $|J| = r + 1$ such that $\bigcap_{j \in J} C_j = \emptyset$ and $\bigcap_{j \in J} C'_j \neq \emptyset$, then

- (i) there exists a closed cover $\mathcal{E}' = \{E'_1, \dots, E'_n\}$ of Z' with $C'_i \subset E'_i$ for all i such that the dimension of $\mathcal{N}(\mathcal{E}') \setminus \mathcal{N}(C')$ is less than r (by definition), and
- (ii) the dimension of $\mathcal{N}(C') \setminus \mathcal{N}(C)$ is less than r .

Consequently, the dimension of $\mathcal{N}(\mathcal{E}') \setminus \mathcal{N}(C)$ is less than r . The collection $\mathcal{E} = \{E_1, \dots, E_n\}$ with $E_i := f^{-1}(E'_i)$ is a closed cover of Z such that $C_i \subset E_i$ for all i . The nerve $\mathcal{N}(\mathcal{E}')$ contains $\mathcal{N}(\mathcal{E})$. Therefore, the dimension of $\mathcal{N}(\mathcal{E}) \setminus \mathcal{N}(C)$ is less than r . This contradicts the assumption that $r = \text{rk}(Z, C)$. □

Lemma 54 *Let $(Z, \mathcal{C} = \{C_1, \dots, C_n\})$ be a KKM system of rank $r > 0$ with metrizable Z and all C_i compact, and let d be a metric on Z . Then there exists a closed metric ball*

$$B_R(x) := \{z \in Z \mid d(z, x) \leq R\}, \quad x \in Z, R > 0,$$

whose interior intersects no element of \mathcal{C} and whose boundary sphere touches at least $r + 1$ elements of \mathcal{C} .

Proof If z is a point and N is a subset in Z , we write

$$d(z, N) := \inf_{p \in N} d(z, p).$$

Let A denote the union $\bigcup_{i=1}^n C_i$. For each $i \in \{1, \dots, n\}$, we set

$$E_i := \{z \in Z \mid d(z, C_i) = d(z, A)\}.$$

Observe that E_i contains C_i and is closed, so $\{E_1, \dots, E_n\}$ is a closed cover of Z . Since (Z, \mathcal{C}) is a KKM system of rank $r > 0$, it follows by the definition of KKM rank that the set $\{1, \dots, n\}$ contains a subset J of cardinality $r + 1$ such that $\bigcap_{j \in J} C_j = \emptyset$ and $\bigcap_{j \in J} E_j \neq \emptyset$. Let x be a point in $\bigcap_{j \in J} E_j \neq \emptyset$. Then the ball $B_{d(x,A)}(x)$ of radius $d(x, A)$ centered at x meets the requirements of the lemma (since each of C_i is compact). □

Proof of Theorem 36 Since $(Z, A, [\mathcal{C}])$ is an Eilenberg–Pontryagin triple, it follows by Lemma 33 that (Z, \mathcal{C}) is a KKM system of rank $\text{rk}(Z, \mathcal{C}) = \text{rk}(Z, A, [\mathcal{C}])$.

Let $\mathcal{C} = \{C_1, \dots, C_n\}$. Set $\mathcal{F} := \{F(C_1), \dots, F(C_n)\}$. Then Lemma 52 implies that we have two possibilities:

- (1) The dimension of $\mathcal{N}(\mathcal{F}) \setminus \mathcal{N}(\mathcal{C})$ is at least $\text{rk}(Z, \mathcal{C})$, so $\{1, \dots, n\}$ contains a subset J of cardinality $\text{rk}(Z, \mathcal{C}) + 1$ such that $\bigcap_{j \in J} C_j = \emptyset$ and $\bigcap_{j \in J} F(C_j) \neq \emptyset$.
- (2) The pair (M, \mathcal{F}) is a KKM system of rank at least $\text{rk}(Z, \mathcal{C})$.

In case (1), for any point $x \in \bigcap_{j \in J} F(C_j)$, the set $F^{-1}(x)$ is a set of spherical $F|_A$ -neighbors that intersects all elements of $\{C_j\}_{j \in J}$, which proves the theorem.

In case (2), the required statement follows by Lemma 54 applied to (M, \mathcal{F}) . □

Proof of Corollary 41 We use a spherical version of Theorem 4. Let T be a regular triangulation of the unit sphere S^n and let $\tilde{\Delta}_1, \dots, \tilde{\Delta}_{n+2}$, be the n -simplices of T ; all of $\tilde{\Delta}_i$ are regular spherical simplices with Euclidean distances between vertices

$$(1) \quad d_{n,\text{Eu}} = \sqrt{\frac{2(n+2)}{n+1}}$$

and angular edge length

$$(2) \quad d_{n,A} = 2 \arcsin\left(\frac{1}{2}d_{n,\text{Eu}}\right) = 2 \arcsin \sqrt{\frac{n+2}{2(n+1)}} = \arccos\left(\frac{-1}{n+1}\right).$$

We recall that the *circumradius* of a compact set Q in a metric space is defined as the radius of a least metric ball containing Q . If Q is a compact subset of S^n we denote by $\text{circ}_A Q$ and $\text{diam}_A Q$, respectively, the circumradius and diameter of Q with respect to the angular metric, and $\text{diam}_{\text{Eu}} Q$ will stand for the Euclidean diameter of Q in $\mathbb{R}^{n+1} \supset S^n$. Under this notation, Dekster’s extension [9] of the Jung theorem says that, for any compact subset Q of S^n ,

$$2 \arcsin\left(\sqrt{\frac{n+1}{2n}} \sin(\text{circ}_A Q)\right) \leq \text{diam}_A Q.$$

This immediately implies that, in the case where $\text{circ}_A Q \leq \frac{\pi}{2}$,

$$(3) \quad \sqrt{\frac{2(n+1)}{n}} \sin(\text{circ}_A Q) \leq \text{diam}_{\text{Eu}} Q.$$

Another auxiliary fact we need is that

$$(4) \quad \text{diam}_A \tilde{\Delta}_i = \frac{1}{2}(\pi - d_{n,A}).$$

Indeed, observe that $\tilde{\Delta}_i$ is the intersection of a finite number of closed hemispheres and hence its boundary is composed of fragments of great hyperspheres, which are geodesic in S^n . Therefore, if a and b are two points in $\tilde{\Delta}_i$ such that neither a nor b is a vertex of $\tilde{\Delta}_i$, then $\text{diam}_A \{a, b\} < \text{diam}_A \tilde{\Delta}_i$ because $\tilde{\Delta}_i$ contains two geodesic arcs⁴ α and β such that α contains a in its relative interior and β contains b in its relative interior. Since $\tilde{\Delta}_i$ is contained in the interior of a hemisphere, so a and b are not antipodal, it follows that there exist $a' \in \alpha \subset \tilde{\Delta}_i$ and $b' \in \beta \subset \tilde{\Delta}_i$ with $\text{diam}_A \{a, b\} < \text{diam}_A \{a', b'\}$ (imagine the interposition of α, β and the metric ball $D \subset S^n$ of diameter $\text{diam}_A \{a, b\}$ containing a and b). Thus, if a and b are two points in $\tilde{\Delta}_i$ such that $\text{diam}_A \{a, b\} = \text{diam}_A \tilde{\Delta}_i$, then one of a and b is a vertex of $\tilde{\Delta}_i$ and we easily obtain (4) by considering the regular triangulation of S^n dual (antipodal) to T .

Now, we pass to the proof of Corollary 41. If we have a continuous map $f: S^n \rightarrow M$, then Theorem 4 implies that a finite set \mathcal{P} of spherical f -neighbors intersects all of $\tilde{\Delta}_i$. We need to prove that

$$(5) \quad \text{diam}_{\text{Eu}} \mathcal{P} \geq \sqrt{\frac{n+2}{n}}.$$

Let $B \subset S^n$ be a metric ball with angular radius $\text{circ}_A \mathcal{P}$ containing \mathcal{P} , let $C \in S^n$ be the center of B , let $A \in S^n$ be the antipode of C , let $\tilde{\Delta}_k$ be a simplex of T containing A and

⁴By geodesic arcs in S^n we mean arcs of great circles.

let $B_2 \subset \mathbb{S}^n$ be the metric ball centered at A of angular radius $\text{diam}_A \tilde{\Delta}_k = \frac{1}{2}(\pi - d_{n,A})$ (see (4)). Then B_2 contains $\tilde{\Delta}_k$. Since \mathcal{P} intersects $\tilde{\Delta}_k$ while $\mathcal{P} \subset B$ and $\tilde{\Delta}_k \subset B_2$, it follows that B intersects B_2 . Therefore,

$$(6) \quad \text{circ}_A \mathcal{P} = \text{circ}_A B \geq \pi - \text{circ}_A B_2 = \frac{1}{2}d_{n,A}.$$

The situation splits into two cases:

- (i) $\text{circ}_A \mathcal{P} > \frac{\pi}{2}$ (ie no hemisphere contains \mathcal{P}).
- (ii) $\text{circ}_A \mathcal{P} \leq \frac{\pi}{2}$.

In case (i), we observe that, since no hemisphere contains \mathcal{P} , it follows that no Euclidean ball in \mathbb{R}^n of radius less than 1 contains \mathcal{P} . Then the Jung theorem⁵ says that $\text{diam}_{\text{Eu}} \mathcal{P} \geq d_{n,\text{Eu}}$, which implies the required (5).

In case (ii), (3) is applicable and yields

$$\begin{aligned} \text{diam}_{\text{Eu}} \mathcal{P} &\geq \sqrt{\frac{2(n+1)}{n}} \cdot \sin(\text{circ}_A \mathcal{P}) && \text{(by (3))} \\ &\geq \sqrt{\frac{2(n+1)}{n}} \cdot \sin\left(\frac{1}{2}d_{n,A}\right) && \text{(by (6) and (ii))} \\ &= \sqrt{\frac{2(n+1)}{n}} \cdot \frac{1}{2}d_{n,\text{Eu}} && \text{(by (2))} \\ &= \sqrt{\frac{2(n+1)}{n}} \cdot \sqrt{\frac{n+2}{2(n+1)}} = \sqrt{\frac{n+2}{n}} && \text{(by (1)).} \quad \square \end{aligned}$$

Remark 55 It would be interesting to find a way to upgrade the above proof of Corollary 41 by considering the family of all regular triangulations of the unit sphere \mathbb{S}^n .

Proof of Corollary 51 (1) Corollary 44 implies that, if M is a contractible metric space and $f : \partial P \rightarrow M$ is a continuous map, then each cubic hemisphere in ∂P contains a facet that is a member of a pair of disjoint facets that are spherical f -neighbors. The statement follows.

(2) Proposition 56 below implies that, if the lighthouse independence number of a flag 3-polytope P is k , then ∂P contains k cubic hemispheres with pairwise disjoint interiors. This implies the required assertion by assertion (1) of the corollary.

(3)–(4) These follow from assertion (2) and Proposition 57 below.

⁵For a discussion and materials concerning the Jung theorem and containment in hemispheres, see [39; 15; 8, pages 112, 113, 132–136, 38; 19; 4; 1, Proposition 2.4].

(5)–(6) These follow from (2) because a direct check shows that the lighthouse independence number of the regular dodecahedron is 4 and the lighthouse independence number of the regular icosahedron is 3.

(7) Corollary 44 shows that the weak Radon rank of the cube is at least 1 and an example where $\partial[0, 1]^3$ is mapped to an oblate spheroid in \mathbb{R}^3 shows that it is at most 1. \square

We say that a vertex v of a polytope is *cubical* if the union of the facets containing v is a cubic hemisphere.

Proposition 56 *All vertices of a flag 3–polytope are cubical.*

Proof Let v be a vertex of a flag 3–polytope P . Observe that no facet of P is a triangle (because any triangular facet together with the three adjacent ones form a collection of four pairwise intersecting facets with no common point). Therefore, each facet of P containing v has a vertex that is not adjacent to v . Let v_1, v_2 and v_3 be three such vertices lying on three distinct facets containing v . Let D denote the union of the facets of P that do not contain v . Then D is a topological disk with the points v_1, v_2 and v_3 on its boundary. Since P is flag, we see that

- no facet contained in D intersects three of the facets not contained in D ,
- no facet of P splits D (in the sense that $D \setminus F$ is connected for each facet F).

This implies that

- each of the vertices v_1, v_2 and v_3 is incident to an edge of P whose second endpoint is contained in the interior of D (in particular, the interior of D contains at least one vertex of P); and
- the subgraph G_D in the 1–skeleton P_1 of P induced by the vertices of P contained in the interior of D is connected.

Thus, each of v_1, v_2 and v_3 is adjacent to a vertex of the connected subgraph G_D in P_1 . This easily implies that P_1 contains a Y –homeomorphic subgraph Y' that is contained in D and intersects the boundary ∂D exactly in the set $\{v_1, v_2, v_3\}$.

Furthermore, since v_1, v_2 and v_3 belong to three distinct facets containing v , it follows that there exists a triple of edges in P_1 incident to v whose endpoints split ∂D into three arcs each of which contains exactly one of v_1, v_2 and v_3 . Clearly, the union of these edges with ∂D and Y' is a graph homeomorphic to the cube 1–skeleton. This shows that v is cubical. \square

Proposition 57 (1) *Let P be a flag simple 3–polytope with ψ facets and let g be the largest number of edges in a facet of P . Then the lighthouse independence number of P is at least*

$$\left\lfloor \frac{2\psi - 7}{3g - 8} \right\rfloor.$$

(2) *If P is a fullerene with ψ facets, then the lighthouse independence number of P is at least*

$$\left\lfloor \frac{1}{3}(\psi - 3) \right\rfloor.$$

Proof In the proof, if v is a vertex of P , we denote by $L(v)$ the union of facets of P that contain v .

We construct a lighthouse independent set by the following algorithm. First we choose a vertex v_1 of P such that the number of vertices in $L(v_1)$ is the least possible and set $W_1 = L(v_1)$. The number of vertices in $L(v_1)$ is at most $3g - 5$.

At each next step, being given $W_i \subset P$ such that a vertex of P is not in W_i , we take a vertex v_{i+1} of P in $P \setminus W_i$ such that the number of vertices in $L(v_{i+1}) \setminus W_i$ is the least possible and set $W_{i+1} = W_i \cup L(v_{i+1})$. Observe that, if a vertex v of P is not in W_i and adjacent to a vertex in W_i , then $L(v)$ shares at least three vertices with W_i . This implies that the number of vertices in $L(v_{i+1}) \setminus W_i$ is at most $3g - 8$.

Therefore, if P has N vertices, this algorithm produces a lighthouse independent set v_1, v_2, \dots with at least

$$1 + \left\lfloor \frac{N - (3g - 5)}{3g - 8} \right\rfloor = \left\lfloor \frac{N - 3}{3g - 8} \right\rfloor$$

elements. Since P is simple, Euler’s formula yields $N = 2\psi - 4$. This proves the required estimate.

The case of fullerenes follows if we observe that, when v_1 is a vertex of a pentagon, the number of vertices in $L(v_1)$ is at most 12. □

4 Concluding remarks

Now we discuss several concepts and open questions.

(1) **The Hopf theorem** The trefoil curve in Figure 5 shows that there exists a continuous map $f : S^1 \rightarrow \mathbb{R}^2$ with no pair of spherical f –neighbors having distance less than $\sqrt{3}$ between them. This means that the direct analog of the aforementioned Hopf theorem for spherical f –neighbors does not hold for small distances. It would be




Figure 5

interesting to find more properties of the set of distances between spherical f -neighbors for a continuous map f of given metric spaces. For example:

Question *Is it true that, for any continuous map $f : \mathbb{S}^n \rightarrow \mathbb{R}^{n+k}$, the set*

$$\Gamma_f := \{\delta \in \mathbb{R} \mid \delta = d(p, q) \text{ for a pair } \{p, q\} \text{ of spherical } f\text{-neighbors}\}$$

contains a nondegenerate interval? Is there a nonzero lower bound for the diameter of Γ_f ?

(2) **Topological Tverberg theorems** Projecting a Euclidean n -sphere $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ into a hyperplane in \mathbb{R}^{n+1} shows that there exists a continuous map $f : \mathbb{S}^n \rightarrow \mathbb{R}^n$ with no set of spherical f -neighbors of cardinality exceeding 2. Consequently, each convex n -polytope P has a map $f : \partial P \rightarrow \mathbb{R}^{n-1}$ with no set of spherical f -neighbors intersecting three disjoint faces of P . This means that no direct analog of the topological Tverberg theorems with three or more disjoint faces holds for spherical f -neighbors. This correlates with the property (see Remark 16) that no principal cover has disjoint elements. Nevertheless, we have some analogs of the topological Radon theorem, which is the topological Tverberg theorem for two disjoint faces; see Corollaries 44, 46 and 51.

Problem *Find extensions of topological Tverberg theorems for spherical f -neighbors with additional restrictions.*

(See also Van Kampen–Flores- and Conway–Gordon–Sachs-type results [33].)

(3) **Weak Radon rank** It would be interesting to:

Problem *Describe the set of polyhedra that are not weakly Radon. Find the weak Radon rank for fullerenes.*

(4) **Minimaxes, I** Let (X, ρ) and (M, d) be metric spaces and let $f : X \rightarrow M$ be a continuous map. Let P_f be the set of all pairs of spherical f -neighbors in X . We set

$$D_f := \sup_{\{x,y\} \in P_f} \rho(x, y), \quad \mu(X, M) := \inf_{f \in C(X, M)} D_f,$$

where $C(X, M)$ stands for continuous maps. Suppose $X = \mathbb{S}^n$ and $M = \mathbb{R}^m$. If $m \leq n$, then $\mu(\mathbb{S}^n, \mathbb{R}^m) = 2$ by the Borsuk–Ulam theorem. For $n < m$, it is shown in [21] that

$$\mu(\mathbb{S}^n, \mathbb{R}^m) = \sqrt{\frac{2(n+2)}{n+1}}.$$

Problem Find $\mu(X, M)$ and its lower bounds in general and some special cases. In particular, find D_f and μ for the case where $M = \mathbb{R}^n$ and X is an n -dimensional Riemannian manifold.

(5) **Minimaxes, II** Let us fix $[C]$ in $[X, \mathbb{S}^{n-2}]$ (see Definition 7), for instance $[C] \neq 0$ in $\pi_3(\mathbb{S}^2)$.

Question What is the min–max distance between the points of a set intersecting each element of a cover of this class?

(6) **Widths, distortion, filling radius, etc** Similarly to $\mu(X, M)$, we consider infima of D_f over families of homotopic maps, over all continuous maps of a given space to certain classes of spaces (eg contractible spaces), etc. This generates a series of new metric “ μ -invariants” of maps and metric spaces. These μ -invariants are similar to such invariants as distortion, filling radius and various widths (see [37; 12; 13; 10; 31; 18; 3]).

Problem Find and describe relations between μ -invariants and classical ones.

(7) **Topological and visual f -neighbors** Let $f: X \rightarrow Y$ be a map of topological spaces. We say that two points a and b in X are *topological f -neighbors* if $f(a)$ and $f(b)$ belong to the boundary of the same connected component of the complement $Y \setminus f(X)$. If Y is a geodesic metric space, we say that a and b in X are *visual f -neighbors* if $f(a)$ and $f(b)$ are connected by a geodesic, in Y , whose interior does not meet $f(X)$.

Problem Translate the above constructions and questions to these new types of f -neighbors.

(8) **Helly-type sufficient conditions for principal covers** Remark 18 implies some Helly-type sufficient conditions for principal covers. For example, if $\mathcal{C} = \{C_1, \dots, C_n\}$ is a closed cover of a normal space Y such that $\mathcal{N}(\mathcal{C}) = \partial\Delta^{n-1}$ and, for each $J \subset \{1, \dots, n\}$ with $|J| \leq n - 2$, any continuous map $\mathbb{S}^{n-2-|J|} \rightarrow \bigcap_{j \in J} C_j$ is null-homotopic, then there exists a map $f: \partial\Delta^{n-1} \rightarrow Y$ such that the image of each

facet is contained in an element of \mathcal{C} , so \mathcal{C} is principal. (See the proofs of Theorems 5 and 6 in [7].)

Question Which of the other versions of topological Helly theorem (see eg [7; 24]) give sufficient conditions for principal and non-nullhomotopic covers?

References

- [1] **C Adams, F Morgan, J M Sullivan**, *When soap bubbles collide*, Amer. Math. Monthly 114 (2007) 329–337 MR Zbl
- [2] **H Adams, J Bush, F Frick**, *Metric thickenings, Borsuk–Ulam theorems, and orbitopes*, Mathematika 66 (2020) 79–102 MR Zbl
- [3] **A Akopyan, R Karasev, A Volovikov**, *Borsuk–Ulam type theorems for metric spaces*, preprint (2012) arXiv 1209.1249
- [4] **R Alexander**, *The width and diameter of a simplex*, Geometriae Dedicata 6 (1977) 87–94 MR Zbl
- [5] **I Bárány, P V M Blagojević, G M Ziegler**, *Tverberg’s theorem at 50: extensions and counterexamples*, Not. Amer. Math. Soc. 63 (2016) 732–739 MR Zbl
- [6] **I Bárány, P Soberón**, *Tverberg’s theorem is 50 years old: a survey*, Bull. Amer. Math. Soc. 55 (2018) 459–492 MR Zbl
- [7] **S A Bogatyĭ**, *The topological Helly theorem*, Fundam. Prikl. Mat. 8 (2002) 365–405 MR Zbl In Russian
- [8] **L Danzer, B Grünbaum, V Klee**, *Helly’s theorem and its relatives*, from “Convexity”, Proc. Sympos. Pure Math. 7, Amer. Math. Soc., Providence, RI (1963) 101–180 MR Zbl
- [9] **B V Dekster**, *The Jung theorem for spherical and hyperbolic spaces*, Acta Math. Hungar. 67 (1995) 315–331 MR Zbl
- [10] **E Denne, J M Sullivan**, *The distortion of a knotted curve*, Proc. Amer. Math. Soc. 137 (2009) 1139–1148 MR Zbl
- [11] **F Frick**, *Counterexamples to the topological Tverberg conjecture*, from “Geometric and algebraic combinatorics” (G Kalai, I Novik, F Santos, V Welker, editors), Oberwolfach Rep. 12 (2015) 318–322 MR Zbl
- [12] **M Gromov**, *Homotopical effects of dilatation*, J. Differential Geometry 13 (1978) 303–310 MR Zbl
- [13] **M Gromov**, *Filling Riemannian manifolds*, J. Differential Geom. 18 (1983) 1–147 MR Zbl

- [14] **B Grünbaum**, *On the facial structure of convex polytopes*, Bull. Amer. Math. Soc. 71 (1965) 559–560 MR Zbl
- [15] **W Gustin**, Review of [39], Math. Reviews (1952) <https://mathscinet.ams.org/mathscinet/article?mr=51539>
- [16] **N V Ivanov**, *Leray theorems in bounded cohomology theory*, preprint (2020) arXiv 2012.08038
- [17] **R N Karasëv**, *Topological methods in combinatorial geometry*, Uspekhi Mat. Nauk 63 (2008) 39–90 MR Zbl In Russian; translated in Russian Math. Surveys 63 (2008) 1031–1078
- [18] **R N Karasev**, *A topological central point theorem*, Topology Appl. 159 (2012) 864–868 MR Zbl
- [19] **V Klee**, Review of [38], Math. Reviews (1965) <https://mathscinet.ams.org/mathscinet/article?mr=183000>
- [20] **A V Malyutin**, *On the question of genericity of hyperbolic knots*, Int. Math. Res. Not. 2020 (2020) 7792–7828 MR Zbl
- [21] **A V Malyutin, O R Musin**, *Borsuk–Ulam type theorems for Delaunay maps*, in preparation
- [22] **J Matoušek**, *Using the Borsuk–Ulam theorem*, Springer (2003) MR Zbl
- [23] **B Matschke**, *A survey on the square peg problem*, Not. Amer. Math. Soc. 61 (2014) 346–352 MR Zbl
- [24] **L Montejano**, *A new topological Helly theorem and some transversal results*, Discrete Comput. Geom. 52 (2014) 390–398 MR Zbl
- [25] **K Morita**, *On the dimension of normal spaces, II*, J. Math. Soc. Japan 2 (1950) 16–33 MR Zbl
- [26] **K Morita**, *On generalizations of Borsuk’s homotopy extension theorem*, Fund. Math. 88 (1975) 1–6 MR Zbl
- [27] **O R Musin**, *Homotopy invariants of covers and KKM-type lemmas*, Algebr. Geom. Topol. 16 (2016) 1799–1812 MR Zbl
- [28] **O R Musin**, *KKM type theorems with boundary conditions*, J. Fixed Point Theory Appl. 19 (2017) 2037–2049 MR Zbl
- [29] **O R Musin, J Wu**, *Cobordism classes of maps and covers for spheres*, Topology Appl. 237 (2018) 21–25 MR Zbl
- [30] **N Noble**, *Poorly separated infinite normal products*, preprint (2020) arXiv 2002.02483
- [31] **J Pardon**, *On the distortion of knots on embedded surfaces*, Ann. of Math. 174 (2011) 637–646 MR Zbl
- [32] **E Schechter**, *Handbook of analysis and its foundations*, Academic, San Diego, CA (1997) MR Zbl

- [33] **A B Skopenkov**, *Topological Tverberg conjecture*, Uspekhi Mat. Nauk 73 (2018) 141–174 MR Zbl In Russian; translated in Russian Math. Surveys 73 (2018) 323–353
- [34] **M Starbird**, *The Borsuk homotopy extension theorem without the binormality condition*, Fund. Math. 87 (1975) 207–211 MR Zbl
- [35] **H Steinlein**, *Borsuk’s antipodal theorem and its generalizations and applications: a survey*, from “Topological methods in nonlinear analysis” (A Granas, editor), Sémin. Math. Sup. 95, Presses Univ. Montréal, Montreal, QC (1985) 166–235 MR Zbl
- [36] **H Steinlein**, *Spheres and symmetry: Borsuk’s antipodal theorem*, Topol. Methods Nonlinear Anal. 1 (1993) 15–33 MR Zbl
- [37] **V M Tikhomirov**, *Some questions of the approximation theory*, Izdat. Moskov. Univ., Moscow (1976) MR In Russian
- [38] **T A Timan**, *Proof of a geometric theorem of Jung, and its analogue in the theory of stochastic processes*, Uspehi Mat. Nauk 20 (1965) 213–218 MR Zbl In Russian
- [39] **S Verblunsky**, *On the circumradius of a bounded set*, J. London Math. Soc. 27 (1952) 505–507 MR Zbl

*St Petersburg Department of Steklov Institute of Mathematics
St Petersburg, Russia*

*School of Mathematical and Statistical Sciences, University of Texas – Rio Grande Valley
Brownsville, TX, United States*

malyutin@pdmi.ras.ru, oleg.musin@utrgv.edu

Received: 13 April 2021 Revised: 8 January 2022

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Kathryn Hess
kathryn.hess@epfl.ch
École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Norihiko Minami	Nagoya Institute of Technology nori@nitech.ac.jp
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Thomas Nikolaus	University of Münster nikolaus@uni-muenster.de
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr		


See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2023 is US \$650/year for the electronic version, and \$940/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 23 Issue 7 (pages 2925–3415) 2023

- Differential geometric invariants for time-reversal symmetric Bloch bundles, II: The low-dimensional “quaternionic” case 2925
GIUSEPPE DE NITTIS and KIYONORI GOMI
- Detecting isomorphisms in the homotopy category 2975
KEVIN ARLIN and J DANIEL CHRISTENSEN
- Mod 2 power operations revisited 2993
DYLAN WILSON
- The Devinatz–Hopkins theorem via algebraic geometry 3015
ROK GREGORIC
- Neighboring mapping points theorem 3043
ANDREI V MALYUTIN and OLEG R MUSIN
- Stable cohomology of the universal degree d hypersurface in \mathbb{P}^n 3071
ISHAN BANERJEE
- On the wheeled PROP of stable cohomology of $\text{Aut}(F_n)$ with bivariant coefficients 3089
NARIYA KAWAZUMI and CHRISTINE VESPA
- Anchored foams and annular homology 3129
ROSTISLAV AKHMECHET and MIKHAIL KHOVANOV
- On a problem of Hopf for circle bundles over aspherical manifolds with hyperbolic fundamental groups 3205
CHRISTOFOROS NEOFYTIDIS
- The mod 2 cohomology of the infinite families of Coxeter groups of type B and D as almost-Hopf rings 3221
LORENZO GUERRA
- Operads in unstable global homotopy theory 3293
MIGUEL BARRERO
- On some p -differential graded link homologies, II 3357
YOU QI and JOSHUA SUSSAN
- Leighton’s theorem and regular cube complexes 3395
DANIEL J WOODHOUSE