

AG  
T

*Algebraic & Geometric  
Topology*

Volume 24 (2024)

**Computing the Morava  $K$ -theory of real Grassmannians  
using chromatic fixed point theory**

NICHOLAS J KUHN

CHRISTOPHER J R LLOYD



# Computing the Morava $K$ –theory of real Grassmannians using chromatic fixed point theory

NICHOLAS J KUHN  
CHRISTOPHER J R LLOYD

We study  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$ , the 2–local Morava  $K$ –theories of the real Grassmannians, about which very little has been previously computed. We conjecture that the Atiyah–Hirzebruch spectral sequences computing these all collapse after the first possible nonzero differential  $d_{2n+1-1}$ , and give much evidence that this is the case.

We use a novel method to show that higher differentials can’t occur: we get a lower bound on the size of  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$  by constructing a  $C_4$ –action on our Grassmannians and then applying the chromatic fixed point theory of the authors’ previous paper. In essence, we bound the size of  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$  by computing  $K(n-1)^*(\mathrm{Gr}_d(\mathbb{R}^m)^{C_4})$ .

Meanwhile, the size of  $E_{2n+1}$  is given by  $Q_n$ –homology, where  $Q_n$  is Milnor’s  $n^{\mathrm{th}}$  primitive mod 2 cohomology operation. Whenever we are able to calculate this  $Q_n$ –homology, we have found that the size of  $E_{2n+1}$  agrees with our lower bound for the size of  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$ . We have two general families where we prove this:  $m \leq 2^{n+1}$  and all  $d$ , and  $d = 2$  and all  $m$  and  $n$ . Computer calculations have allowed us to check many other examples with larger values of  $d$ .

55M35, 55N20; 55P91, 57S17

## 1 Introduction

Let  $\mathrm{Gr}_d(\mathbb{R}^m)$  be the real Grassmannian of  $k$ –planes in  $\mathbb{R}^m$ , a much studied compact manifold of dimension  $d(m-d)$  admitting the structure of a CW complex with  $\binom{m}{d}$  “Schubert cells”.

Much is known about the ordinary cohomology of these spaces:

- (1)  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$  is generated by Stiefel–Whitney classes satisfying standard relations. It has total dimension  $\binom{m}{d}$ .
- (2)  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Q})$  is generated by Pontryagin classes, along with, in some cases, an odd-dimensional class. For fixed  $d$ , and  $\epsilon = 0$  or 1, the total dimension of  $H^*(\mathrm{Gr}_d(\mathbb{R}^{2-\epsilon+2l}); \mathbb{Q})$  is polynomial of degree  $\lfloor d/2 \rfloor$  as a function of  $l \geq 0$ .
- (3) If  $m$  is even, then  $\mathrm{Gr}_d(\mathbb{R}^m)$  is oriented. Furthermore, the inclusion  $\mathrm{Gr}_d(\mathbb{R}^{m-1}) \hookrightarrow \mathrm{Gr}_d(\mathbb{R}^m)$  induces an epimorphism in rational cohomology.

- (4) Nontrivial torsion in  $H^*(Gr_d(\mathbb{R}^m); \mathbb{Z})$  has order 2. The mod 2 Bockstein spectral sequence (BSS) collapses after the first differential. Equivalently, the mod 2 Adams spectral sequence (ASS) converging to  $H^*(Gr_d(\mathbb{R}^m); \mathbb{Z})$  collapses at  $E_2$ .

Much less is known about other cohomology theories applied to these Grassmannians. In this paper, we study  $K(n)^*(Gr_d(\mathbb{R}^m))$  for  $n \geq 1$ . Here  $K(n)^*(X)$  denotes the 2-local  $n^{\text{th}}$  Morava  $K$ -theory of a space  $X$ , a graded vector space over the graded field  $K(n)_* = \mathbb{Z}/2[v_n^{\pm}]$  with  $|v_n| = 2^{n+1} - 2$ . We let  $k(n)$  denote the connective cover of  $K(n)$ :  $k(n)_* = \mathbb{Z}/2[v_n]$ .

Viewing  $H\mathbb{Q}$  as  $K(0)$  and  $H\mathbb{Z}$  as  $k(0)$ , our discovery is that analogues of statements (2)–(4) above appear to hold for all  $n$ , with the Atiyah–Hirzebruch spectral sequence (AHSS) replacing the Bockstein spectral sequence in statement (4). Furthermore, the analogue of statement (1) holds through a much bigger range than one would expect from dimension considerations.

In the next two subsections, we describe our main results.

### 1.1 Results proved using chromatic fixed point theory

Given a finite complex  $X$  and  $n \geq 0$ , we let  $k_n(X) = \dim_{K(n)_*} K(n)^*(X)$ .

**Theorem 1.1** *If  $m \leq 2^{n+1}$ , then  $k_n(Gr_d(\mathbb{R}^m)) = \binom{m}{d}$ . Thus, in this range, the AHSS converging to  $K(n)^*(Gr_d(\mathbb{R}^m))$  collapses at  $E_2$ .*

We note that this collapsing range is surprisingly large, as dimension considerations just imply collapsing if  $d(m - d) < 2^{n+1}$ .

For larger  $m$ , we have the following lower bound.

**Theorem 1.2** *Let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . Then*

$$k_n(Gr_d(\mathbb{R}^m)) \geq \sum_{i=0}^{\lfloor d/2 \rfloor} \binom{2^{n+1} - \epsilon}{d - 2i} \binom{l}{i}.$$

**Conjecture 1.3** *Equality always holds in this last theorem.*

The biggest novelty of this paper is our method for proving Theorems 1.1 and 1.2: we make use of chromatic fixed point theory to prove these nonequivariant results.

The blue shift theorem of Barthel, Hausmann, Naumann, Nikolaus, Noel and Stapleton [2] says that if  $C$  is a finite cyclic  $p$ -group and  $X$  is a finite  $C$ -CW complex, then

$$\tilde{K}(n)^*(X) = 0 \implies \tilde{K}(n - 1)^*(X^C) = 0;$$

see also Balderrama and the first author [1]. In [8], we upgraded this. Specialized to cyclic groups, [8, Theorem 2.17] says the following.

**Theorem 1.4** *If  $C$  is a finite cyclic  $p$ -group, and  $X$  is a finite  $C$ -CW complex, then*

$$k_n(X) \geq k_{n-1}(X^C).$$

Note that, in these statements,  $K(n)_*$  means Morava  $K$ -theory at the prime  $p$ .

As  $\binom{m}{d}$  is an evident upper bound for  $k_n(\text{Gr}_d(\mathbb{R}^m))$ , to prove [Theorem 1.1](#), it suffices to show that  $k_n(\text{Gr}_d(\mathbb{R}^m)) \geq \binom{m}{d}$  in the stated range. Using [Theorem 1.4](#), we will show this by induction on  $n$  using a  $C_2$ -action on  $\text{Gr}_d(\mathbb{R}^m)$  induced by an  $m$ -dimensional real representation of  $C = C_2$ .

We will similarly prove [Theorem 1.2](#) for  $n \geq 1$  by using a  $C_4$ -action on  $\text{Gr}_d(\mathbb{R}^m)$  induced by an  $m$ -dimensional real representation of  $C = C_4$ .

In both cases, it will be quite easy to compute  $k_{n-1}(\text{Gr}_d(\mathbb{R}^m)^C)$ .

Details of this will be in [Section 2](#).

## 1.2 Results about the $Q_n$ -homology of the Grassmannians

[Conjecture 1.3](#) follows from a conjectural calculation that only involves  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ , viewed as a module over the Steenrod algebra.

Let  $Q_n$  for  $n = 0, 1, 2, \dots$  be the Milnor primitives: the elements in the mod 2 Steenrod algebra recursively defined by  $Q_0 = \text{Sq}^1$ , and  $Q_n = [Q_{n-1}, \text{Sq}^{2^n}]$ . These satisfy  $Q_n^2 = 0$ , and we let  $k_{Q_n}(X)$  denote the total dimension of the  $Q_n$ -homology of  $X$ ,

$$H^*(X; Q_n) = \frac{Z^*(X; Q_n)}{B^*(X; Q_n)},$$

where

$$Z^*(X; Q_n) = \ker\{Q_n: H^*(X; \mathbb{Z}/2) \rightarrow H^{*+2^{n+1}-1}(X; \mathbb{Z}/2)\},$$

$$B^*(X; Q_n) = \text{im}\{Q_n: H^{*-2^{n+1}+1}(X; \mathbb{Z}/2) \rightarrow H^*(X; \mathbb{Z}/2)\}.$$

As will be reviewed in [Section 3.1](#), the first differential in the AHSS converging to  $K(n)^*(X)$  is  $d_{2^{n+1}-1}$ , with formula

$$d_{2^{n+1}-1}(x) = Q_n(x)v_n$$

for all  $x \in E_2^{*,0}(X) = H^*(X; \mathbb{Z}/2)$ . This makes it not hard to check the next lemma.

**Lemma 1.5** *If  $X$  is a finite complex,  $k_{Q_n}(X) \geq k_n(X)$  is always true, and the following are equivalent:*

- (a)  $k_{Q_n}(X) = k_n(X)$ ;
- (b) the AHSS, when  $n \geq 1$ , or the BSS, when  $n = 0$ , computing  $K(n)^*(X)$  collapses at  $E_{2^{n+1}}$ ;
- (c) the ASS computing  $k(n)^*(X)$  collapses at  $E_2$ .

We apply this to our situation. First, [Theorem 1.1](#) has the following nontrivial algebraic consequence.

**Corollary 1.6** *If  $m \leq 2^{n+1}$ , then  $Q_n$  acts trivially on  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .*

For an algebraic proof of this result using the methods of [Section 3.5](#), see the second author’s thesis [[10](#), page 75].

For  $m > 2^{n+1}$ , we believe the following is true.

**Conjecture 1.7** *Let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . Then*

$$k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) = \sum_{i=0}^{\lfloor d/2 \rfloor} \binom{2^{n+1} - \epsilon}{d - 2i} \binom{l}{i}.$$

Comparison with [Theorem 1.2](#) shows that when [Conjecture 1.7](#) is true, one can conclude

- $k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) = k_n(\text{Gr}_d(\mathbb{R}^m))$ , and [Conjecture 1.3](#) is true;
- the AHSS computing  $K(n)^*(\text{Gr}_d(\mathbb{R}^m))$  collapses at  $E_{2^{n+1}}$ ;
- the ASS computing  $k(n)^*(\text{Gr}_d(\mathbb{R}^m))$  collapses at  $E_2$ ;
- $k_n(\text{Gr}_d(\mathbb{R}^{2^{n+1} - \epsilon + 2l}))$  is polynomial of degree  $\lfloor d/2 \rfloor$  as a function of  $l$ .

Known rational calculations imply that the conjecture is true when  $n = 0$ . It is also easy to show that the conjecture is true when  $d = 1$ , and one calculates

$$k_n(\text{Gr}_1(\mathbb{R}^m)) = \begin{cases} m & \text{if } 1 \leq m \leq 2^{n+1}, \\ 2^{n+1} - \epsilon & \text{if } m = 2^{n+1} - \epsilon + 2l. \end{cases}$$

With much more work we prove the following.

**Theorem 1.8** *[Conjecture 1.7](#) is true when  $d = 2$ . Thus the Atiyah–Hirzebruch spectral sequence computing  $K(n)^*(\text{Gr}_2(\mathbb{R}^m))$  collapses at  $E_{2^{n+1}}$ , the Adams spectral sequence computing  $k(n)^*(\text{Gr}_2(\mathbb{R}^m))$  collapses at  $E_2$ , and we have the calculation*

$$k_n(\text{Gr}_2(\mathbb{R}^m)) = \begin{cases} \binom{m}{2} & \text{if } 2 \leq m \leq 2^{n+1}, \\ \binom{2^{n+1} - \epsilon}{2} + l & \text{if } m = 2^{n+1} - \epsilon + 2l. \end{cases}$$

We are firm believers in our conjectures. For more evidence, the second author has made extensive computer calculations verifying [Conjecture 1.7](#) in hundreds more cases with larger values of  $d$ ; see the tables in the [appendix](#).

For  $d \geq 2$ , computing the size of  $H^*(\text{Gr}_d(\mathbb{R}^m); Q_n)$  seems tricky. We have organized our efforts by studying how these numbers change as  $m$  is increased as follows.

Let  $C_d(\mathbb{R}^m)$  denote the cofiber of the inclusion  $\text{Gr}_d(\mathbb{R}^{m-1}) \rightarrow \text{Gr}_d(\mathbb{R}^m)$ , so there is a cofiber sequence  $\text{Gr}_d(\mathbb{R}^{m-1}) \xrightarrow{i} \text{Gr}_d(\mathbb{R}^m) \xrightarrow{p} C_d(\mathbb{R}^m)$ . In [Section 3.3](#),  $C_d(\mathbb{R}^m)$  is identified as the Thom space of the canonical normal bundle over  $\text{Gr}_{d-1}(\mathbb{R}^{m-1})$ , and in [Section 3.4](#) we study the  $Q_n$ -module

$\tilde{H}^*(C_d(\mathbb{R}^m); \mathbb{Z}/2)$ , viewed as  $H^*(\mathrm{Gr}_{d-1}(\mathbb{R}^{m-1}); \mathbb{Z}/2)$  equipped with an explicit twisted  $Q_n$ -action. One has an induced short exact sequence of modules over the Steenrod algebra

$$0 \rightarrow \tilde{H}^*(C_d(\mathbb{R}^m); \mathbb{Z}/2) \xrightarrow{p^*} H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2) \xrightarrow{i^*} H^*(\mathrm{Gr}_d(\mathbb{R}^{m-1}); \mathbb{Z}/2) \rightarrow 0,$$

inducing a long exact sequence on  $Q_n$ -homology.

When  $m$  is even, we see much orderly behavior.

**Theorem 1.9** *Let  $m$  be even.*

(a)  $H^{d(m-d)}(\mathrm{Gr}_d(\mathbb{R}^m); Q_n) \simeq \mathbb{Z}/2$ , ie the nonzero top-dimensional cohomology class is not in the image of  $Q_n$  for all  $n$ .

(b) The chain complex  $(\tilde{H}^*(C_d(\mathbb{R}^m); Q_n))$  is dual to the chain complex

$$(H^{d(m-d)-*}(\mathrm{Gr}_{d-1}(\mathbb{R}^{m-1}); Q_n)).$$

(c) If [Conjecture 1.7](#) is true for  $(n, d, m-1)$  and  $(n, d-1, m-1)$ , then it is true for  $(n, d, m)$ . Furthermore,  $\mathrm{Gr}_d(\mathbb{R}^m)$  will then be  $k(n)$ -oriented, and the cofiber sequence above will induce short exact sequences

$$\begin{aligned} 0 \rightarrow \tilde{H}^*(C_d(\mathbb{R}^m); Q_n) &\xrightarrow{p^*} H^*(\mathrm{Gr}_d(\mathbb{R}^m); Q_n) \xrightarrow{i^*} H^*(\mathrm{Gr}_d(\mathbb{R}^{m-1}); Q_n) \rightarrow 0, \\ 0 \rightarrow \tilde{K}(n)^*(C_d(\mathbb{R}^m)) &\xrightarrow{p^*} K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m)) \xrightarrow{i^*} K(n)^*(\mathrm{Gr}_d(\mathbb{R}^{m-1})) \rightarrow 0. \end{aligned}$$

We prove [Theorem 1.9](#) in [Section 4](#). We make use of the additive basis  $\{s_\lambda\}$  dual to the classical Schubert cells. Here  $\lambda$  runs through partitions having at most  $d$  parts, each no bigger than  $m-d$ . In [\[9\]](#), Cristian Lennart gave a combinatorial formula for  $Q_n(s_\lambda)$ , and we use this to prove (a). Duality statement (b) follows quite formally from (a), and (c) follows easily from (b).

When  $m$  is odd, the analogues of statements (a) and (b) are false, and, for  $d \geq 3$ , the full behavior of the connecting map in the  $Q_n$ -homology long exact sequence,

$$\delta: H^*(\mathrm{Gr}_d(\mathbb{R}^{m-1}); Q_n) \rightarrow \tilde{H}^{*+2^{n+1}-1}(C_d(\mathbb{R}^m); Q_n),$$

is as yet unclear to the authors. In [Section 6](#), we will prove analogues of [Theorems 1.1](#) and [1.2](#) for  $C_d(\mathbb{R}^m)$ , and then speculate on behavior of  $\delta$  that would be compatible with all of our computations.

However, when  $d = 2$ , we have the following result.

**Theorem 1.10** *Let  $m > 2^{n+1}$  be odd. Then  $k_{Q_n}(C_2(\mathbb{R}^m)) = 2^{n+1} - 2$  and the map*

$$\tilde{H}^*(C_2(\mathbb{R}^m); Q_n) \xrightarrow{p^*} H^*(\mathrm{Gr}_2(\mathbb{R}^m); Q_n)$$

*is zero, so there is a short exact sequence*

$$0 \rightarrow H^*(\mathrm{Gr}_2(\mathbb{R}^m); Q_n) \xrightarrow{i^*} H^*(\mathrm{Gr}_2(\mathbb{R}^{m-1}); Q_n) \xrightarrow{\delta} \tilde{H}^{*+2^{n+1}-1}(C_2(\mathbb{R}^m); Q_n) \rightarrow 0.$$

From this, [Theorem 1.8](#) quickly follows and one can deduce that, in this case, there is a short exact sequence

$$0 \rightarrow K(n)^*(\mathrm{Gr}_2(\mathbb{R}^m)) \xrightarrow{i^*} K(n)^*(\mathrm{Gr}_2(\mathbb{R}^{m-1})) \xrightarrow{\delta} \tilde{K}(n)^{*+1}(C_2(\mathbb{R}^m)) \rightarrow 0.$$

We prove [Theorem 1.10](#) in [Section 5](#). The tools we use are very different from those used in proving [Theorem 1.9](#): we work with the classical presentation of  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$  as a ring of Stiefel–Whitney classes.

### 1.3 Comparison with other work

When comparing our work to what has come before, the first thing to say is that the outcome of our calculations — though not the methods — are in line with the classical calculations first made by C Ehresmann in 1937 [\[3\]](#). He determined the additive structure of both  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$  and  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Q})$ . He also showed that all the torsion in  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z})$  was of order 2; in modern terms this is equivalent to showing that the Bockstein spectral sequence computing  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z})$  collapses after the first nonzero differential given by  $Q_0 = \mathrm{Sq}^1 = \beta$ .

Calculating the Morava  $K$ -theories of  $\mathrm{Gr}_d(\mathbb{R}^\infty) = \mathrm{BO}(d)$  was done first by Kono and Yagita [\[7\]](#), and then, with a simpler proof, by Kitchloo and Wilson [\[6\]](#). Again, the AHSS computing  $K(n)^*(\mathrm{BO}(d))$  collapses after the first nonzero differential, but the collapsing is for an elementary reason:  $H^*(\mathrm{BO}(d); Q_n)$  is concentrated in even degrees. Indeed, one quickly learns that the complexification map  $\mathrm{BO}(d) \rightarrow \mathrm{BU}(d)$  induces an epimorphism  $K(n)^*(\mathrm{BU}(d)) \rightarrow K(n)^*(\mathrm{BO}(d))$ , so  $K(n)^*(\mathrm{BO}(d))$  is generated by Chern classes  $c_1, \dots, c_d$ .

An equivalent statement is that  $H^*(\mathrm{BO}(d); Q_n)$  is generated by the classes  $w_1^2, \dots, w_d^2$ . These will still be permanent classes in the AHSS converging to  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$ , but now we have odd-dimensional classes as well, with the number of these seemingly growing as  $d$  and  $m$  grow.

Finally, we point out that we do not attempt to describe  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$  as a  $K(n)^*$ -algebra. Our results do tell us something about this, however. In the situation of [Theorem 1.1](#), the known algebra  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2) \otimes K(n)^*$  will be an associated graded. Similarly, whenever our conjecture is valid,  $H^*(\mathrm{Gr}_d(\mathbb{R}^m); Q_n) \otimes K(n)^*$  would be an associated graded of the  $K(n)^*$ -algebra  $K(n)^*(\mathrm{Gr}_d(\mathbb{R}^m))$ . What is still needed, and might be necessary to prove our conjectural collapsing in general, are sensible constructions of classes in odd degrees.

### Acknowledgements

Kuhn is a PI of RTG NSF grant DMS-1839968, which partially supported the research of Lloyd.

## 2 The proofs of Theorems 1.1 and 1.2

In this section we prove [Theorems 1.1](#) and [1.2](#) by using our chromatic fixed point theorem [Theorem 1.4](#).

### 2.1 A fixed point formula

Let  $G$  be a finite group, and let  $V$  be an  $m$ -dimensional real representation of  $G$ . Then  $\text{Gr}_d(V)$ , the space of  $d$ -planes in  $V$ , is a model for  $\text{Gr}_d(\mathbb{R}^m)$  with an evident  $G$ -action. Here we describe  $\text{Gr}_d(V)^G$ , its space of  $G$ -fixed points.

To state this, we need some notation. Let  $V_1, \dots, V_k$  be the irreducible real representations of  $G$ , let  $r_i = \dim_{\mathbb{R}} V_i$ , and let  $\mathbb{D}_i = \text{End}_{\mathbb{R}[G]}(V_i, V_i)$ . Each of the endomorphism algebras  $\mathbb{D}_i$  will be a finite-dimensional real division algebra, and thus isomorphic to  $\mathbb{R}, \mathbb{C}$ , or  $\mathbb{H}$ , and  $\dim_{\mathbb{R}} \mathbb{D}_i$  will divide  $r_i$ .

**Proposition 2.1** *If  $V = V_1^{m_1} \oplus \dots \oplus V_k^{m_k}$ , then there is a homeomorphism*

$$\text{Gr}_d(V)^G = \bigsqcup_{j_1 r_1 + \dots + j_k r_k = d} \text{Gr}_{j_1}(\mathbb{D}_1^{m_1}) \times \dots \times \text{Gr}_{j_k}(\mathbb{D}_k^{m_k}).$$

**Proof** The fixed point space  $\text{Gr}_d(V)^G$  will be the space of sub- $G$ -modules  $W < V$  of real dimension  $d$ . Such a  $G$ -module  $W$  will decompose canonically as  $W = W_1 \oplus \dots \oplus W_k$ , with  $W_i < V_i^{m_i}$ . If  $d_i = \dim_{\mathbb{R}} W_i$ , then  $d_1 + \dots + d_k = d$ . Thus we have a decomposition

$$\text{Gr}_d(V)^G = \bigsqcup_{d_1 + \dots + d_k = d} \text{Gr}_{d_1}(V_1^{m_1})^G \times \text{Gr}_{d_2}(V_2^{m_2})^G \times \dots \times \text{Gr}_{d_k}(V_k^{m_k})^G.$$

A submodule  $W_i$  of  $V_i^{m_i}$  must be isomorphic to  $V_i^j$  for some  $j$ ; thus  $\text{Gr}_{d_i}(V_i^{m_i})^G$  will be empty unless  $d_i = j_i r_i$  for some  $j_i$ .

Finally, using that  $\text{Hom}_{\mathbb{R}[G]}(V_i^{j_i}, V_i^{m_i}) = \text{Hom}_{\mathbb{D}}(\mathbb{D}^{j_i}, \mathbb{D}^{m_i})$ , one deduces that the submodules of  $V_i^{m_i}$  isomorphic to  $V_i^{j_i}$  correspond to the  $\mathbb{D}$ -subspaces of  $\mathbb{D}^{m_i}$  of dimension  $j_i$  over  $\mathbb{D}$ . Thus there is a homeomorphism

$$\text{Gr}_{j_i r_i}(V_i^{m_i})^G = \text{Gr}_{j_i}(\mathbb{D}_i^{m_i}). \quad \square$$

**Corollary 2.2** *If  $V = V_1^{m_1} \oplus \dots \oplus V_k^{m_k}$ , then, for any  $n$ ,*

$$k_n(\text{Gr}_d(V)^G) = \sum_{j_1 r_1 + \dots + j_k r_k = d} k_n(\text{Gr}_{j_1}(\mathbb{D}_1^{m_1})) \cdots k_n(\text{Gr}_{j_k}(\mathbb{D}_k^{m_k})).$$

**Proof** A consequence of the Künneth theorem for  $K(n)_*$  is that  $k_n(X \times Y) = k_n(X)k_n(Y)$ . Thus the corollary follows from the proposition. □

**Remark 2.3** If  $\mathbb{D} = \mathbb{C}$  or  $\mathbb{H}$ , then  $\text{Gr}_d(\mathbb{D}^m)$  has a CW structure with  $\binom{m}{d}$  cells that are all even-dimensional, and thus  $k_n(\text{Gr}_d(\mathbb{D}^m)) = \binom{m}{d}$  for all  $n$ .



## 2.2 Proof of Theorem 1.1

Theorem 1.1 says that if  $m \leq 2^{n+1}$  then  $k_n(\text{Gr}_d(\mathbb{R}^m)) = \binom{m}{d}$ . Using Theorem 1.4 and Proposition 2.1, we prove this by induction on  $n$ .

The  $n = 0$  case of the theorem is easy to check, as

$$\text{Gr}_d(\mathbb{R}^0) = \begin{cases} * & \text{if } d = 0, \\ \emptyset & \text{otherwise,} \end{cases} \quad \text{and} \quad \text{Gr}_d(\mathbb{R}^1) = \begin{cases} * & \text{if } d = 0, 1, \\ \emptyset & \text{otherwise.} \end{cases}$$

For the inductive step, assume that if  $p \leq 2^n$  then  $k_{n-1}(\text{Gr}_d(\mathbb{R}^p)) = \binom{p}{d}$ .

Let  $m \leq 2^{n+1}$ . As it is clear that  $k_n(\text{Gr}_d(\mathbb{R}^m)) \leq \binom{m}{d}$ , our goal is to show that  $k_n(\text{Gr}_d(\mathbb{R}^m)) \geq \binom{m}{d}$ .

Let  $C_2$  be the cyclic group of order 2. To get our needed lower bound, our strategy will be to make  $\mathbb{R}^m$  into a  $C_2$ -module, and then apply Theorem 1.4.

The group  $C_2$  has two irreducible 1-dimensional real representations; call them  $L_1$  and  $L_2$ . Since  $m \leq 2^{n+1}$ , we can write  $m$  as  $m = p + q$  with both  $p \leq 2^n$  and  $q \leq 2^n$ . Now let  $V = L_1^p \oplus L_2^q$ , an  $m$ -dimensional real representation of  $C_2$ .

Applying Proposition 2.1, we see that

$$\text{Gr}_d(V)^{C_2} = \bigsqcup_{i+j=d} \text{Gr}_i(\mathbb{R}^p) \times \text{Gr}_j(\mathbb{R}^q).$$

Applying Theorem 1.4 to this, we learn that

$$\begin{aligned} k_n(\text{Gr}_d(\mathbb{R}^m)) &\geq \sum_{i+j=d} k_{n-1}(\text{Gr}_i(\mathbb{R}^p))k_{n-1}(\text{Gr}_j(\mathbb{R}^q)) \\ &= \sum_{i+j=d} \binom{p}{i} \binom{q}{j} \quad (\text{by inductive hypothesis}) \\ &= \binom{m}{d}. \end{aligned}$$

**Remark 2.4** The same inductive proof can be used to prove the classical result that

$$\dim_{\mathbb{Z}/2} H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2) = \binom{m}{d}$$

for all  $m$  and  $d$ , with our chromatic fixed point theorem Theorem 1.4 replaced by the classical theorem of Ed Floyd [4, Theorem 4.4]: if the cyclic group  $C_p$  acts on a finite CW complex  $X$ , then  $\dim_{\mathbb{Z}/p} H^*(X; \mathbb{Z}/p) \geq \dim_{\mathbb{Z}/p} H^*(X^{C_p}; \mathbb{Z}/p)$ . It would be interesting to know if this argument was known to Floyd, or others, like Bob Stong, who regularly worked with these sorts of group actions.

### 2.3 Proof of Theorem 1.2

The strategy of the proof of Theorem 1.2 is the same as the proof in the last subsection: we get a lower bound on  $k_n(\text{Gr}_d(\mathbb{R}^m))$  by letting a cyclic 2-group act on  $\mathbb{R}^m$  and applying Theorem 1.4.

In this case, the representation theory of  $C_2$  is not rich enough to give us a big enough lower bound, but a well chosen real representation of the group  $C_4$  of order 4 works better. Curiously, in our calculation of  $k_{n-1}$  of the resulting fixed point space, we are able to use our already proven Theorem 1.1, so the proof is not by induction, but more direct.

The group  $C_4$  has three irreducible real representations:  $L_1$  and  $L_2$  of dimension 1, and  $R$  of real dimension 2. Note that  $\text{End}_{\mathbb{R}[C_4]}(R) \simeq \mathbb{C}$ .

Now let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . We define an  $m$ -dimensional real representation  $V$  of  $C_4$  by  $V = L_1^{2^n} \oplus L_2^{2^n - \epsilon} \oplus R^l$ .

Applying Proposition 2.1, we see that

$$\text{Gr}_d(V)^{C_4} = \bigsqcup_{j+k+2i=d} \text{Gr}_j(\mathbb{R}^{2^n}) \times \text{Gr}_k(\mathbb{R}^{2^n - \epsilon}) \times \text{Gr}_i(\mathbb{C}^l).$$

Applying Theorem 1.4 to this, we learn that

$$\begin{aligned} k_n(\text{Gr}_d(\mathbb{R}^m)) &\geq \sum_{j+k+2i=d} k_{n-1}(\text{Gr}_j(\mathbb{R}^{2^n}))k_{n-1}(\text{Gr}_k(\mathbb{R}^{2^n - \epsilon}))k_{n-1}(\text{Gr}_i(\mathbb{C}^l)) \\ &= \sum_{j+k+2i=d} \binom{2^n}{j} \binom{2^n - \epsilon}{k} \binom{l}{i} \quad (\text{using Theorem 1.1}) \\ &= \sum_i \left[ \sum_{j+k=d-2i} \binom{2^n}{j} \binom{2^n - \epsilon}{k} \right] \binom{l}{i} \\ &= \sum_i \binom{2^{n+1} - \epsilon}{d-2i} \binom{l}{i}. \end{aligned}$$

## 3 The $Q_n$ homology of $\text{Gr}_d(\mathbb{R}^m)$ : background material

### 3.1 The AHSS and the ASS for Morava $K$ -theory

Let  $n \geq 1$ . We recall the structure of the AHSS converging to  $K(n)^*(X)$  (as always, in this paper, with  $p = 2$ ). It is a spectral sequence of graded  $K(n)^* = \mathbb{Z}/2[v_n^\pm]$  algebras with

$$E_2^{*,*}(X) = H^*(X; K(n)^*) = H^*(X; \mathbb{Z}/2)[v_n^\pm].$$

Here  $v_n$  has cohomological degree  $2 - 2^{n+1}$ .

Sparseness of the rows implies that the differential  $d_r$  will be zero unless  $r = s(2^{n+1} - 2) + 1$  for some  $s$ . The first possible nonzero differential,  $d_{2^{n+1}-1}$ , satisfies the following formula [15]: given  $x \in E_2^{*,0}(X) = H^*(X; \mathbb{Z}/2)$ ,

$$d_{2^{n+1}-1}(x) = Q_n(x)v_n.$$

It follows that  $E_{2^{n+1}}(X) \simeq H^*(X; Q_n)[v_n^\pm]$ , and so the dimension of  $E_{2^{n+1}}(X)$  as a  $K(n)^*$ -vector space will equal  $k_{Q_n}(X)$ , the dimension of the  $Q_n$ -homology of  $X$ . One immediately deduces part of Lemma 1.5:  $k_{Q_n}(X) = k_n(X)$  if and only if the AHSS converging to  $K(n)^*(X)$  collapses at  $E_{2^{n+1}}(X)$ .

To continue with the proof of Lemma 1.5, let  $cE_r^{*,*}(X)$  denote the terms of the AHSS computing  $k(n)^*(X)$ , a 4<sup>th</sup> quadrant spectral sequence. Note that  $cE_2^{*,*} = H^*(X; \mathbb{Z}/2)[v_n]$  embeds in  $E_2^{*,*}(X) = H^*(X; \mathbb{Z}/2)[v_n^\pm]$ , and equals it for  $\star \leq 0$ , and that the latter spectral sequence is obtained from the former by inverting  $v_n$ .

It follows that  $cE_{2^{n+1}}^{*,*}(X) = E_{2^{n+1}}^{*,*}(X)$  for  $\star < 0$ , with the map on the 0-line between the spectral sequences corresponding to the epimorphism  $Z^*(X; Q_n) \twoheadrightarrow H^*(X; Q_n)$ . From this, one sees that any higher differential in the  $k(n)^*(X)$  AHSS would be detected in the  $K(n)^*(X)$  AHSS. Since this second spectral sequence is the localization of the first, we can conclude that the  $K(n)^*(X)$  AHSS collapses at  $E_{2^{n+1}}(X)$  if and only if the  $k(n)^*(X)$  AHSS collapses at  $cE_{2^{n+1}}(X)$ .

Next we note that the AHSS spectral sequence  $cE_r^{*,*}(X)$  identifies with the ASS computing  $k(n)^*(X)$  with suitable reindexing, with  $cE_{2^{n+1}}^{*,*}(X)$  corresponding to the Adams  $E_2$  term. Firstly, a result of CRF Maunder [11] implies that the AHSS converging to  $[X, k(n)]_*$  can be constructed by taking the Postnikov filtration of the spectrum  $k(n)$ . But the Postnikov tower for  $k(n)$  is also an Adams tower: as described in the survey paper [14, Section 5], there is a cofibration sequence

$$\Sigma^{2^{n+1}-2}k(n) \xrightarrow{v_n} k(n) \xrightarrow{\pi} H\mathbb{Z}/2 \xrightarrow{\bar{Q}_n} \Sigma^{2^{n+1}-1}k(n)$$

such that  $\Sigma^{2^{n+1}-1}\pi \circ \bar{Q}_n = Q_n$  and  $\pi$  induces the epimorphism  $A \rightarrow A/AQ_n$  on mod 2 cohomology.

Finally, we note that, when  $n = 0$ , one still has the cofibration sequence as above, now with  $v_0 = 2$ , so that the ASS for  $k(0) = H\mathbb{Z}$  is similarly related to the Bockstein spectral sequence.

### 3.2 The description of $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ via Stiefel–Whitney classes

We recall classical results that are either explicitly in [12] or can easily be deduced from the material there.

Let  $w_1, \dots, w_d$  denote the Stiefel–Whitney classes of the canonical  $d$ -dimensional bundle  $\gamma_d$  over  $\text{Gr}_d(\mathbb{R}^\infty)$ . One has

$$H^*(\text{Gr}_d(\mathbb{R}^\infty); \mathbb{Z}/2) = \mathbb{Z}/2[w_1, \dots, w_d].$$

Dual classes  $\bar{w}_1, \bar{w}_2, \dots$  are defined by the equation

$$(1 + w_1 + \dots + w_d)(1 + \bar{w}_1 + \bar{w}_2 + \dots) = 1,$$

and this allows one to write the classes  $\bar{w}_k$  as polynomials in  $w_1, \dots, w_d$ .

The inclusion  $\text{Gr}_d(\mathbb{R}^m) \hookrightarrow \text{Gr}_d(\mathbb{R}^\infty)$  then induces a surjective ring homomorphism

$$H^*(\text{Gr}_d(\mathbb{R}^\infty); \mathbb{Z}/2) \rightarrow H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$$

with kernel  $J(d, m - d) = (\bar{w}_k \mid k > m - d)$ . Now  $\bar{w}_k$  can be interpreted as  $w_k(\gamma_d^\perp)$ , where  $\gamma_d^\perp$  is the  $(m - d)$ -dimensional bundle complementary to  $\gamma_d$ .

We record some useful consequences. To state these, it is useful to let

$$i : \text{Gr}_d(\mathbb{R}^{m-1}) \hookrightarrow \text{Gr}_d(\mathbb{R}^m)$$

be the inclusion induced by the inclusion  $\mathbb{R}^{m-1} \hookrightarrow \mathbb{R}^m$ , and to let

$$j : \text{Gr}_{d-1}(\mathbb{R}^{m-1}) \hookrightarrow \text{Gr}_d(\mathbb{R}^m)$$

be the inclusion sending  $V \subset \mathbb{R}^{m-1}$  to  $V \oplus \mathbb{R} \subset \mathbb{R}^m$ .

**Lemma 3.1** (a) *The ideal  $J(d, m - d)$  is generated by the  $d$  classes  $\bar{w}_{m-d+1}, \bar{w}_{m-d+2}, \dots, \bar{w}_m$ .*

(b) *In  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ ,  $w_d \bar{w}_{m-d} = 0$ .*

(c)  *$\ker\{i^*\} = (\bar{w}_{m-d}) \subset H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .*

(d)  *$\ker\{j^*\} = (w_d) \subset H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .*

**Proof** Statement (a) follows from the recursive relations among the  $\bar{w}_k$ . Statement (b) follows from the equation

$$(1 + w_1 + \dots + w_d)(1 + \bar{w}_1 + \dots + \bar{w}_{m-d}) = 1,$$

which holds in  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ . Statement (c) follows from the fact that

$$J(d, m - 1 - d) = J(d, m) + (\bar{w}_{m-d}),$$

and (d) follows from (c), noting that  $j$  can be written as the composite

$$\text{Gr}_{d-1}(\mathbb{R}^{m-1}) \simeq \text{Gr}_{m-d}(\mathbb{R}^{m-1}) \xrightarrow{i} \text{Gr}_{m-d}(\mathbb{R}^m) \simeq \text{Gr}_d(\mathbb{R}^m),$$

where the indicated homeomorphisms are given by taking complementary subspaces (and, in cohomology, these maps swap each  $w_i$  with a  $\bar{w}_j$ ). □

We end this subsection with a couple more facts about  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .

An additive basis for  $H^q(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$  is given by the monomials

$$\left\{ w_1^{r_1} w_2^{r_2} \cdots w_d^{r_d} \mid \sum_{i=1}^d r_i \leq m - d \right\},$$

so the top-dimensional class is  $w_d^{m-d}$  in degree  $d(m - d)$ ; see [5].

The Wu formulae [12, page 94] are closed formulae for  $\text{Sq}^i w_j$ , and, in theory, formulae for  $Q_n(w_j)$  follow.

### 3.3 A description of the cofiber $C_d(\mathbb{R}^m)$ and its cohomology

Recall that  $C_d(\mathbb{R}^m)$  is defined as the cofiber of the inclusion  $\text{Gr}_d(\mathbb{R}^{m-1}) \xrightarrow{i} \text{Gr}_d(\mathbb{R}^m)$ . This cofiber can be identified as a Thom space as follows.

**Proposition 3.2** *Let  $S(\gamma_{d-1}^\perp)$  and  $D(\gamma_{d-1}^\perp)$  be the sphere and disk bundles associated to*

$$\gamma_{d-1}^\perp \rightarrow \text{Gr}_{d-1}(\mathbb{R}^{m-1}).$$

*There is a pushout*

$$\begin{array}{ccc} S(\gamma_{d-1}^\perp) & \xrightarrow{f} & \text{Gr}_d(\mathbb{R}^{m-1}) \\ \downarrow & & \downarrow \\ D(\gamma_{d-1}^\perp) & \xrightarrow{f} & \text{Gr}_d(\mathbb{R}^m), \end{array}$$

*inducing a homeomorphism  $f : \text{Th}(\gamma_{d-1}^\perp) \xrightarrow{\sim} C_d(\mathbb{R}^m)$ , such that the composite*

$$\text{Gr}_{d-1}(\mathbb{R}^{m-1}) \xrightarrow[\sim]{0\text{-section}} D(\gamma_{d-1}^\perp) \xrightarrow{f} \text{Gr}_d(\mathbb{R}^m)$$

*is the map  $j$  of Lemma 3.1.*

**Proof** Recall that

$$\begin{aligned} D(\gamma_{d-1}^\perp) &= \{(V, v) \mid V \in \text{Gr}_{d-1}(\mathbb{R}^{m-1}), v \in V^\perp, |v| \leq 1\}, \\ S(\gamma_{d-1}^\perp) &= \{(V, v) \mid V \in \text{Gr}_{d-1}(\mathbb{R}^{m-1}), v \in V^\perp, |v| = 1\}. \end{aligned}$$

We define  $f : D(\gamma_{d-1}^\perp) \rightarrow \text{Gr}_d(\mathbb{R}^m)$  by the formula

$$f(V, v) = V + \langle v + \sqrt{1 - |v|^2} e_m \rangle,$$

where  $e_m$  is the  $m^{\text{th}}$  standard basis vector in  $\mathbb{R}^m$ . We claim this  $f$  has the needed properties.

First, note that  $f(V, 0) = V + \langle e_m \rangle = V \oplus \mathbb{R} = j(V)$ .

Second,  $(V, v) \in S(\gamma_{d-1}^\perp)$  if and only if  $f(V, v) = V + \langle v \rangle$ , and so is an element of  $\text{Gr}_d(\mathbb{R}^{m-1})$ . Furthermore,  $f : S(\gamma_{d-1}^\perp) \rightarrow \text{Gr}_d(\mathbb{R}^{m-1})$  is surjective: given any  $W \in \text{Gr}_d(\mathbb{R}^{m-1})$ , if we choose any  $(d-1)$ -dimensional subspace  $V$  of  $W$ , and a unit length vector  $v \in W$  in the 1-dimensional orthogonal complement, then  $f(V, v) = W$ .

Finally, we need to check that  $f$  is bijective on  $\mathring{D}(\gamma_{d-1}^\perp) = D(\gamma_{d-1}^\perp) - S(\gamma_{d-1}^\perp)$ . To check this, let  $W \in \text{Gr}_d(\mathbb{R}^m)$  be a  $d$ -dimensional subspace of  $\mathbb{R}^m$  not contained in  $\mathbb{R}^{m-1}$ , so that

$$V = W \cap \mathbb{R}^{m-1} \in \text{Gr}_{d-1}(\mathbb{R}^{m-1}).$$

Let  $V^\perp$  be the complement of  $V$  in  $\mathbb{R}^m$  so that  $W \cap V^\perp$  is one-dimensional, and let  $v$  be the unique unit vector  $v \in W \cap V^\perp$  such that  $v$  has positive  $m^{\text{th}}$  coordinate. Let  $\pi: \mathbb{R}^m \rightarrow \mathbb{R}^{m-1}$  be the standard projection. We claim that  $f(V, \pi(v)) = W$  and  $(V, \pi(v))$  is the unique point in  $\mathring{D}(\gamma_{d-1}^\perp)$  with this property: since  $|v| = 1$  the  $m^{\text{th}}$  component of  $v$  is  $\sqrt{1 - |\pi(v)|^2}$ ; thus  $v = \pi(v) + \sqrt{1 - |\pi(v)|^2}e_m$  and so  $f(V, \pi(v)) = V + \langle v \rangle = W$ . □

Let  $u_{\gamma_{d-1}^\perp} \in \tilde{H}^{m-d}(\text{Th}(\gamma_{d-1}^\perp))$  be the Thom class of  $\gamma_{d-1}^\perp \rightarrow \text{Gr}_{d-1}(\mathbb{R}^{m-1})$ . Then  $\tilde{H}^{m-d}(\text{Th}(\gamma_{d-1}^\perp))$  is a free rank 1  $H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$ -module on  $u_{\gamma_{d-1}^\perp}$ . Meanwhile,  $H^*(\text{Gr}_d(\mathbb{R}^m))$  is a  $H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$ -module via  $j^*$ , and the ideal  $(\bar{w}_{m-d}) = \tilde{H}^*(C_d(\mathbb{R}^m))$  is a submodule. The proposition thus implies the following.

**Corollary 3.3** *The map  $f^*: \tilde{H}^*(C_d(\mathbb{R}^m)) \xrightarrow{\sim} \tilde{H}^*(\text{Th}(\gamma_{d-1}^\perp))$  is an isomorphism of free rank 1  $H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$ -modules, and  $f^*(\bar{w}_{m-d}) = u_{\gamma_{d-1}^\perp}$ .*

### 3.4 The characteristic class associated to $Q_n$ and a twisted $Q_n$ -module

Let  $\alpha_n \in H^{2^{n+1}-1}(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$  be defined as the element satisfying

$$Q_n(\bar{w}_{m-d}) = \alpha_n \bar{w}_{m-d} \in \tilde{H}^*(C_d(\mathbb{R}^m)).$$

Then define

$$\hat{Q}_n: H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1})) \rightarrow H^{*+2^{n+1}-1}(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$$

by the formula

$$\hat{Q}_n(x) = Q_n(x) + x\alpha_n.$$

**Proposition 3.4**  $\hat{Q}_n^2 = 0$ , and the chain complex  $(H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1})), \hat{Q}_n)$  is isomorphic to the chain complex  $(\tilde{H}^{*+m-d}(C_d(\mathbb{R}^m)), Q_n)$ .

**Proof** Let  $\Theta: H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1})) \rightarrow \tilde{H}^{*+m-d}(C_d(\mathbb{R}^m))$  be the isomorphism established in the last subsection,  $\Theta(x) = x\bar{w}_{m-d}$ . The proposition follows once we check that  $\Theta(\hat{Q}_n(x)) = Q_n(\Theta(x))$ ;

$$\begin{aligned} \Theta(\hat{Q}_n(x)) &= \hat{Q}_n(x)\bar{w}_{m-d} \\ &= (Q_n(x) + x\alpha_n)\bar{w}_{m-d} \\ &= Q_n(x)\bar{w}_{m-d} + x(\alpha_n\bar{w}_{m-d}) \\ &= Q_n(x)\bar{w}_{m-d} + xQ_n(\bar{w}_{m-d}) \\ &= Q_n(x\bar{w}_{m-d}) \\ &= Q_n(\Theta(x)). \end{aligned}$$

□

It is useful to put the class  $\alpha_n$  in context. Given any element  $a$  in the Steenrod algebra  $\mathcal{A}$ , one gets a characteristic class  $w_a(\xi) \in H^{|a|}(B; \mathbb{Z}/2)$  associated to any real vector bundle  $\xi \rightarrow B$ ;  $w_a(\xi)$  is defined as the element satisfying  $a(u_\xi) = w_a(\xi)u_\xi \in \tilde{H}^{\dim \xi + |a|}(\text{Th}(\xi); \mathbb{Z}/2)$ , where  $u_\xi$  is the Thom class of  $\xi$ . So, for example,  $w_{\text{Sq}^n}(\xi) = w_n(\xi)$ , and, relevant for us, our class  $\alpha_n$  equals  $w_{Q_n}(\xi)$  when  $\xi = \gamma_{d-1}^\perp \rightarrow \text{Gr}_{d-1}(\mathbb{R}^{m-1})$ .

We have the following characterization of  $w_{Q_n}$ .

**Proposition 3.5**  *$w_{Q_n}$  is the unique characteristic class satisfying the following two properties:*

- (a)  $w_{Q_n}(\xi \oplus \nu) = w_{Q_n}(\xi) + w_{Q_n}(\nu)$ ;
- (b) if  $\gamma \rightarrow B$  is one-dimensional, then  $w_{Q_n}(\gamma) = w_1(\gamma)^{2^{n+1}-1}$ .

**Proof** Property (a) follows from the fact that  $Q_n$  is primitive in  $\mathcal{A}$  (or, equivalently, that  $Q_n$  acts a derivation). To see property (b), one first calculates that  $Q_n(t) = t^{2^{n+1}} \in \mathbb{Z}/2[t] = H^*(\mathbb{R}P^\infty; \mathbb{Z}/2)$ , recalling that  $Q_0 = \text{Sq}^1$ , and  $Q_n = \text{Sq}^{2^n} Q_{n-1} + Q_{n-1} \text{Sq}^{2^n}$ . Then property (b) follows, since if  $\gamma$  is the universal line bundle over  $\mathbb{R}P^\infty$ , then  $u_\gamma = t$ . Uniqueness follows from the splitting principle.  $\square$

**Remark 3.6** Thus  $w_{Q_n}(\xi)$  agrees with the “ $s$ -class”  $s_{2^{n+1}-1}(\xi)$ , analogous to the class of the same name for complex vector bundles as defined in [12, Section 16]. (These  $s_I$  are *not* the same as the  $s_\lambda$  of the next subsection; these are two conflicting and standard usages.)

Since  $\gamma_{d-1}^\perp \oplus \gamma_{d-1}$  is trivial, property (b) has the following consequence.

**Corollary 3.7**  $\alpha_n = w_{Q_n}(\gamma_{d-1}) \in H^{2^{n+1}-1}(\text{Gr}_{d-1}(\mathbb{R}^{m-1}); \mathbb{Z}/2)$ .

### 3.5 The description of $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ via Schubert cells, and Lenart’s formula

For the purposes of proving [Theorem 1.9](#), we use an alternative description of  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .

We recall the cell structure of  $\text{Gr}_d(\mathbb{R}^{d+c})$  as described in [12, Section 6]. A *Schubert symbol*  $\lambda = (\lambda_1, \dots, \lambda_d)$  of  $\text{Gr}_d(\mathbb{R}^m)$  is a sequence of integers

$$m - d \geq \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d > 0.$$

The *weight* of  $\lambda$  is defined to be  $\sum_i \lambda_i$  and is denoted  $|\lambda|$ . Such a  $\lambda$  is a partition contained inside of a  $d \times (m - d)$  grid when depicted as Young diagrams — diagrams with  $\lambda_i$  boxes in the  $i^{\text{th}}$  row.


To each such partition is associated a Schubert cell  $e(\lambda)$  of dimension  $|\lambda|$  in  $\text{Gr}_d(\mathbb{R}^m)$  defined by

$$e(\lambda) = \{V \in \text{Gr}_d(\mathbb{R}^m) \mid \dim(V \cap \mathbb{R}^{i+\lambda_{d+1-i}}) \geq i \text{ for } 1 \leq i \leq d\}.$$


This cell decomposition of the Grassmannian leads to the dual Schubert cell basis for  $H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$  with basis elements  $s_\lambda \in H^{|\lambda|}(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .

With this notation, one has that  $w_i = s_{(1^i)}$  and  $\bar{w}_j = s_{(j)}$ . Although we don't use this here, it is worth noting that the cohomology ring structure in this basis is described by the Littlewood–Richardson rule of symmetric function theory.


To state Lenart's formula for calculating  $Q_n$  on a Schubert basis element [9], we need some combinatorial definitions. Given a Young diagram  $\lambda$  that includes into another Young diagram  $\mu$ , one can form the complement  $\mu/\lambda$ . For example,




The *content* of a box  $b$  of  $\mu$  in row  $i$  and column  $j$  is defined to be  $c(b) = j - i$ . For a box  $b$  in the skew shape  $\mu/\lambda$ , we define its content to be the content of  $b$  embedded in  $\mu$ . Here we fill in the contents of the diagrams from above:



A skew-shape is said to be *connected* when each pair of boxes in the diagram is connected by a sequence of boxes that each share an edge. A shape  $\lambda$  is called a *border strip*, if it is connected and does not contain a  $2 \times 2$  block of boxes. A shape satisfying just the second condition is called a *broken border strip*, and in particular, a border strip is an example of a broken border strip with just one connected component. If  $\lambda$  is a broken border strip, then we denote by  $cc(\lambda)$  the number of connected components of  $\lambda$ . If  $\lambda$  is not a broken border strip, then we define  $cc(\lambda) = \infty$ . For example, in the next diagram,  $\lambda_1$  is a border strip,  $\lambda_2$  is a broken border strip that is not a border strip, and  $\lambda_3$  is an example of a shape that is neither:



A *sharp corner* of a broken border strip is a box with no north, no west and no northwest neighbors. A *dull corner* is a box with both north and west neighbors, but no northwest neighbor. Let  $C(\mu/\lambda)$  denote the set of sharp and dull corners of  $\mu/\lambda$ . For example, in the following diagram the sharp corners have been labeled  $S$  and the dull corners have been labeled  $D$ :



We are now ready to state Lenart's formula from [9]:

$$(3-1) \quad Q_n(s_\lambda) = \sum_{\substack{\mu \supset \lambda: |\mu| - |\lambda| = 2^{n+1} - 1 \\ cc(\mu/\lambda) \leq 2}} d_{\lambda\mu} s_\mu,$$



where  $\mu/\lambda$  must be a broken border strip and

$$(3-2) \quad d_{\lambda\mu} = \begin{cases} \sum_{b \in C(\mu/\lambda)} c(b) & \text{if } \mu/\lambda \text{ is connected,} \\ 1 & \text{if } \mu/\lambda \text{ is disconnected.} \end{cases}$$

**Example 3.8** As an example we compute  $Q_1$  on  $w_1 = s_{\square}$  in the Schubert basis in  $\text{Gr}_2(\mathbb{R}^6)$ . There are three basis elements in degree four,

$$\mu_1 = \square\square\square\square, \quad \mu_2 = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \quad \mu_3 = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}.$$

To compute  $Q_n(s_{\square})$  using (3-1) we must consider each complement. Let  $\lambda = \square$ . For  $\mu_1$ ,

$$\mu_1/\lambda = \begin{array}{|c|c|c|c|} \hline 0 & 1 & 2 & 3 \\ \hline \end{array} / \begin{array}{|c|} \hline \square \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}$$

The complement is a border strip and there is just one sharp corner (the left most corner) and no dull corners. The content of the sharp corner is 1 modulo two; hence  $d_{\lambda\mu_1} = 1$ , and so  $s_{\mu_1}$  is in the expansion of  $Q_1(s_{\lambda})$ . Next we consider

$$\mu_2/\lambda = \begin{array}{|c|c|c|} \hline 0 & 1 & 2 \\ \hline -1 & & \\ \hline \end{array} / \begin{array}{|c|} \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline & 1 & 2 \\ \hline -1 & & \\ \hline \end{array}$$

This is a disconnected broken border strip; hence  $d_{\lambda\mu_2} = 1$ , and so  $s_{\mu_2}$  is in the expansion. Finally,

$$\mu_3/\lambda = \begin{array}{|c|c|} \hline 0 & 1 \\ \hline -1 & -2 \\ \hline \end{array} / \begin{array}{|c|} \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline & 1 \\ \hline -1 & -2 \\ \hline \end{array}$$

There are two sharp corners, one of content  $-1$  and the other of content 1. There is also one dull corner of content  $-2$ . This means  $d_{\lambda\mu_3} = (-1) + 1 + 2 \equiv 0$ , and so  $s_{\mu_3}$  is not in the expansion. Hence,

$$Q_1(s_{\square}) = s_{\square\square\square\square} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}.$$

### 4 Results about $H^*(\text{Gr}_d(\mathbb{R}^m); Q_n)$ when $m$ is even

**Proof of Theorem 1.9(a)** We are going to show that  $Q_n(s_{\lambda}) = 0$  for each Schubert basis element  $s_{\lambda}$  in degree  $d(m-d) - 2^{n+1} + 1$ . Since  $s_{(d(m-d))}$  is the only class in degree  $d(m-d)$ ,

$$Q_n(s_{\lambda}) = d_{\lambda(d(m-d))} s_{(d(m-d))},$$

where  $d_{\lambda(d(m-d))}$  is given by (3-2). We must only consider  $\lambda$  such that  $(d^{(m-d)})/\lambda$  is a broken border strip. As  $(d^{(m-d)})$  is a  $d \times (m-d)$  grid the complement  $(d^{(m-d)})/\lambda$  is always connected and so if  $(d^{(m-d)})/\lambda$  is a broken border strip it must be, in particular, a border strip. If  $(d^{(m-d)})/\lambda$  is a border strip, then it must be one of three types:

- (1)  $(d^{(m-d)})/\lambda$  is the last row of  $(d^{(m-d)})$ ,
- (2)  $(d^{(m-d)})/\lambda$  is the last column of  $(d^{(m-d)})$ ,
- (3)  $(d^{(m-d)})/\lambda$  is the union of the last row and last column of  $(d^{(m-d)})$ .

We will show that  $d_{\lambda(d^{(m-d)})} = 0$  in each of these cases. As  $m$  was assumed to be even, the content of the right most bottom box of  $(d^{(m-d)})$  is also even.

- (1) For the first case, there is just one sharp corner, namely the left most box, and there are no dull corners. Since the strip is of odd length, namely,  $2^{n+1} - 1$ , the leftmost box and the rightmost box have the same content modulo two. Hence, the content of this sharp corner is zero modulo two, and so  $d_{\lambda(d^{(m-d)})} = 0$ .
- (2) For the second case, the argument is exactly the same, but with the sharp corner on the top.
- (3) For the third case, the content of the sharp corner on the bottom left and the content of the sharp corner on the top right agree modulo two, because the border strip is of odd length. There is one dull corner in the bottom right and it is zero modulo two. Thus, the two sharp corners cancel and the dull corner contributes nothing.

Thus, in all cases  $Q_n(s_\lambda) = 0$  for  $s_\lambda$  in degree  $d(m-d) - 2^{n+1} + 1$ . This completes the proof that the top class is not in the image of  $Q_n$  for even  $m$ . □

**Proof of Theorem 1.9(b)** We wish to prove that, when  $m$  is even, the chain complexes  $(\tilde{H}^*(C_d(\mathbb{R}^m)); Q_n)$  and  $(H^{d(m-d)-*}(\text{Gr}_{d-1}(\mathbb{R}^{m-1})); Q_n)$  are dual.

By Proposition 3.4,  $(\tilde{H}^{*+m-d}(C_d(\mathbb{R}^m)); Q_n)$  is isomorphic to the chain complex

$$(H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1})); \hat{Q}_n),$$

where we recall that  $\hat{Q}_n(y) = Q_n(y) + y\alpha_n$ , and that  $\alpha_n \bar{w}_{m-d} = Q_n(\bar{w}_{m-d}) \in H^*(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ .

So we need to check that the chain complexes

$$(H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1})); \hat{Q}_n) \quad \text{and} \quad (H^{(d-1)(m-d)-*}(\text{Gr}_{d-1}(\mathbb{R}^{m-1})); Q_n)$$

are dual. This means we need to show that, if  $x, y \in H^*(\text{Gr}_{d-1}(\mathbb{R}^{m-1}); \mathbb{Z}/2)$  satisfy

$$|x| + |y| + |Q_n| = (d-1)(m-1),$$

then

$$Q_n(x)y = x\hat{Q}_n(y).$$

By Theorem 1.9(a), we know that

$$Q_n(xy\bar{w}_{m-d}) = 0 \in H^{d(m-d)}(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2).$$

Thus, in  $H^{d(m-d)}(\text{Gr}_d(\mathbb{R}^m); \mathbb{Z}/2)$ ,

$$\begin{aligned} 0 &= Q_n(xy\bar{w}_{m-d}) \\ &= Q_n(x)y\bar{w}_{m-d} + xQ_n(y)\bar{w}_{m-d} + xyQ_n(\bar{w}_{m-d}) \\ &= Q_n(x)y\bar{w}_{m-d} + xQ_n(y)\bar{w}_{m-d} + xy\alpha_n\bar{w}_{m-d} \\ &= (Q_n(x)y + xQ_n(y) + xy\alpha_n)\bar{w}_{m-d} \\ &= (Q_n(x)y + x\hat{Q}_n(y))\bar{w}_{m-d}, \end{aligned}$$

and we conclude that  $0 = Q_n(x)y + x\hat{Q}_n(y) \in H^{(d-1)(m-d)}(\text{Gr}_{d-1}(\mathbb{R}^{m-1}); \mathbb{Z}/2)$ . □

**Proof of Theorem 1.9(c)** Recall that  $k_{Q_n}(X)$  denotes the rank of the  $Q_n$ -homology  $H^*(X; Q_n)$ . Similarly, let  $\bar{k}_{Q_n}(X)$  denote the rank of  $\tilde{H}^*(X; Q_n)$ .

Let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . Let

$$k_n^G(d, m) = \sum_{i=0}^{\lfloor d/2 \rfloor} \binom{2^{n+1} - \epsilon}{d - 2i} \binom{l}{i}.$$

We start with the first part of Theorem 1.9(c). This asserts that, when  $m$  is even, if we assume that

$$k_{Q_n}(\text{Gr}_d(\mathbb{R}^{m-1})) = k_n^G(d, m - 1) \quad \text{and} \quad k_{Q_n}(\text{Gr}_{d-1}(\mathbb{R}^{m-1})) = k_n^G(d - 1, m - 1),$$

then we can conclude that  $k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) = k_n^G(d, m)$ .

Theorem 1.2 tells us that  $k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) \geq k_n^G(d, m)$ .

Since we have a short exact sequence

$$0 \rightarrow \tilde{H}^*(C_d(\mathbb{R}^m)) \rightarrow H^*(\text{Gr}_d(\mathbb{R}^m)) \rightarrow H^*(\text{Gr}_d(\mathbb{R}^{m-1})) \rightarrow 0,$$

we see that

$$k_{Q_n}(\text{Gr}_d(\mathbb{R}^{m-1})) + \bar{k}_{Q_n}(C_d(\mathbb{R}^m)) \geq k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)),$$

with equality if and only if the associated long exact  $Q_n$ -homology sequence is still short exact.

Since  $m$  is even, Theorem 1.9(b) applies, and tells us that  $\bar{k}_{Q_n}(C_d(\mathbb{R}^m)) = k_{Q_n}(\text{Gr}_{d-1}(\mathbb{R}^{m-1}))$ .

Putting this all together, under our assumptions,

$$k_n^G(d, m - 1) + k_n^G(d - 1, m - 1) \geq k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) \geq k_n^G(d, m).$$

That these would be, in fact, equalities, follows from the next lemma.

**Lemma 4.1** *If  $m = 2^{n+1} + 2l$  with  $l \geq 0$ , then*

$$k_n^G(d, m - 1) + k_n^G(d - 1, m - 1) = k_n^G(d, m).$$

**Proof** We compute

$$\begin{aligned} k_n^G(d, m-1) + k_n^G(d-1, m-1) &= \sum_i \left[ \binom{2^{n+1}-1}{d-2i} + \binom{2^{n+1}-1}{d-2i-1} \right] \binom{l}{i} \\ &= \sum_i \binom{2^{n+1}}{d-2i} \binom{l}{i} \\ &= k_n^G(d, m). \end{aligned} \quad \square$$

When all of this happens, we then see that the  $Q_n$ -homology long exact sequence really is still short exact, and also that the  $K(n)$ -AHSS must collapse for these three spaces. Thus there is also a short exact sequence

$$0 \rightarrow \tilde{K}(n)^*(C_d(\mathbb{R}^m)) \xrightarrow{p^*} K(n)^*(Gr_d(\mathbb{R}^m)) \xrightarrow{i^*} K(n)^*(Gr_d(\mathbb{R}^{m-1})) \rightarrow 0.$$

Finally, the top cohomology class in  $H^{d(m-d)}(Gr_d(\mathbb{R}^m); \mathbb{Z}/2)$  will be a permanent cycle in the AHSS computing  $K(n)^*(Gr_d(\mathbb{R}^m))$  and thus also in the AHSS computing  $k(n)^*(Gr_d(\mathbb{R}^m))$ , and this is equivalent to saying that  $Gr_d(\mathbb{R}^m)$  is  $k(n)$ -oriented.  $\square$

## 5 Results about $H^*(Gr_d(\mathbb{R}^m); Q_n)$ when $d = 2$

In this section we present our results about the  $Q_n$ -homology of  $Gr_2(\mathbb{R}^m)$ , with the focus on understanding the case when  $m$  has the form  $2^{n+1} - 1 + 2l$ .

To begin with, we know that

- $H^*(Gr_2(\mathbb{R}^m); \mathbb{Z}/2) = \mathbb{Z}/2[w_1, w_2]/(\bar{w}_{m-1}, \bar{w}_m)$ ;
- in  $H^*(Gr_2(\mathbb{R}^m); \mathbb{Z}/2)$ , the ideal  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$  has an additive basis  $\{w_1^i \bar{w}_{m-2} \mid 0 \leq i \leq m-2\}$ .

Now we collect results that hold in  $H^*(Gr_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ .

**Lemma 5.1** In  $H^*(Gr_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ ,

- $\bar{w}_0 = 1, \bar{w}_1 = w_1$ , and, recursively,  $\bar{w}_k = w_1 \bar{w}_{k-1} + w_2 \bar{w}_{k-2}$ ;
- $w_2^j \bar{w}_k = \sum_i \binom{j}{i} w_1^{j-i} \bar{w}_{k+j+i}$ ;
- $\bar{w}_k = \sum_j \binom{k-j}{j} w_1^{k-2j} w_2^j$ ;
- $\bar{w}_{2^b-1} = w_1^{2^b-1}$  for all  $b \geq 0$ ;
- $\bar{w}_{2^b-2} = \sum_{c=0}^{b-1} w_1^{2^b-2^{c+1}} w_2^{2^c-1}$  for all  $b \geq 1$ .

**Proof** The homogeneous components of the equation  $0 = (1 + w_1 + w_2)(1 + \bar{w}_1 + \bar{w}_2 + \dots)$  give statement (a).

Statement (b) is proved by induction on  $j$ . The case when  $j = 0$  is trivial, and statement (a) rewrites as  $w_2 \bar{w}_k = w_1 \bar{w}_{k+1} + \bar{w}_{k+2}$ , which is the case when  $j = 1$ . One then computes

$$\begin{aligned} w_2^j \bar{w}_k &= w_2(w_2^{j-1} \bar{w}_k) \\ &= \sum_i \binom{j-1}{i} w_1^{j-1-i} w_2 \bar{w}_{k+j-1+i} \\ &= \sum_i \binom{j-1}{i} [w_1^{j-i} \bar{w}_{k+j+i} + w_1^{j-1-i} \bar{w}_{k+j+i+1}] \\ &= \sum_i \left[ \binom{j-1}{i} + \binom{j-1}{i-1} \right] w_1^{j-i} \bar{w}_{k+j+i} \\ &= \sum_i \binom{j}{i} w_1^{j-i} \bar{w}_{k+j+i}. \end{aligned}$$

For (c), note that  $\bar{w}_k$  is the homogeneous component of degree  $k$  in

$$\bar{w} = (1 + w_1 + w_2)^{-1} = \sum_{t=0}^{\infty} (w_1 + w_2)^t.$$

Statement (d) follows from (c):

$$\bar{w}_{2^b-1} = \sum_j \binom{2^b-1-j}{j} w_1^{k-2j} w_2^j = w_1^{2^b-1},$$

using that  $\binom{2^b-1-j}{j} \equiv 1 \pmod 2$  only if  $j = 0$ .

Similarly, statement (e) follows from (c):

$$\bar{w}_{2^b-2} = \sum_j \binom{2^b-2-j}{j} w_1^{k-2j} w_2^j = \sum_{c=0}^{b-1} w_1^{2^b-2^{c+1}} w_2^{2^c-1},$$

using that  $\binom{2^b-2-j}{j} \equiv 1 \pmod 2$  if and only if  $j = 2^c - 1$  with  $0 \leq j \leq b - 1$ . □

Now we determine the action of  $Q_n$  on various classes.

**Lemma 5.2** In  $H^*(\text{Gr}_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ ,  $Q_n(w_1) = w_1^{2^{n+1}} = w_1 \bar{w}_{2^{n+1}-1}$ .

**Proof** The first equality here was already noted in the proof of Proposition 3.5, and the second follows from Lemma 5.1(d). □

**Lemma 5.3** In  $H^*(\text{Gr}_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ ,

$$Q_n(w_2) = \sum_{c=0}^n w_1^{2^{n+1}-2^{c+1}+1} w_2^{2^c} = w_1 w_2 \bar{w}_{2^{n+1}-2}.$$

**Proof** The second equality here follows from Lemma 5.1(e), so we just need to check the first. We do this by induction on  $n$ , where the  $n = 0$  case is the easily checked:  $Q_0(w_2) = \text{Sq}^1(w_2) = w_1 w_2$ .

Before proceeding with the inductive step, we make two observations.

The first is that for  $n \geq 1$ ,  $Q_n(w_2) = \text{Sq}^{2^n} Q_{n-1}(w_2)$  because the other term,  $Q_{n-1} \text{Sq}^{2^n}(w_2)$ , will be zero. This is clear if  $n \geq 2$  as then  $\text{Sq}^{2^n}(w_2) = 0$ , and when  $n = 1$ , we observe that  $Q_0 \text{Sq}^2(w_2) = \text{Sq}^1(w_2^2) = 0$ .

The second observation is that  $\text{Sq}(w_2) = w_2(1 + w_1 + w_2)$ , so  $\text{Sq}(w_2^{2^c}) = w_2^{2^c}(1 + w_1^{2^c} + w_2^{2^c})$ , and thus

$$\text{Sq}^j(w_2^{2^c}) = \begin{cases} w_2^{2^c} & \text{if } j = 0, \\ w_1^{2^c} w_2^{2^c} & \text{if } j = 2^c, \\ w_2^{2^{c+1}} & \text{if } j = 2^{c+1}, \\ 0 & \text{otherwise.} \end{cases}$$

Now we check the inductive step of our proof.

$$\begin{aligned} Q_n(w_2) &= \text{Sq}^{2^n} Q_{n-1}(w_2) && \text{(by our first observation)} \\ &= \sum_{c=0}^{n-1} \text{Sq}^{2^n}(w_1^{2^n-2^{c+1}+1} w_2^{2^c}) && \text{(by inductive hypothesis)} \\ &= \sum_{c=0}^{n-1} \sum_j \text{Sq}^{2^n-j}(w_1^{2^n-2^{c+1}+1}) \text{Sq}^j(w_2^{2^c}). \end{aligned}$$

By our second observation, the only possible nonzero terms in this double sum are when  $j = 0, 2^c, 2^{c+1}$ . The terms with  $j = 0$  are all zero, as  $\text{Sq}^{2^n}(w_1^{2^n-2^{c+1}+1}) = 0$  by the unstable condition. Similarly, the only nonzero term with  $j = 2^c$  is the term  $w_1^{2^{n+1}-1} w_2$ , when  $c = 0$ . Finally, one gets  $w_1^{2^{n+1}-2^{c+2}+1} w_2^{2^{c+1}}$  when  $j = 2^{c+1}$  for all  $0 \leq c \leq n - 1$ . One is left with

$$Q_n(w_2) = \sum_{c=0}^n w_1^{2^{n+1}-2^{c+1}+1} w_2^{2^c},$$

completing our induction. □

**Remark 5.4** The referee has pointed out that the first equality in the last lemma appears in [13, page 508].<sup>1</sup>

We now turn our attention to the behavior of  $Q_n$  on  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$ .

**Lemma 5.5** In  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$ ,  $Q_n(\bar{w}_{m-2}) = w_1^{2^{n+1}-1} \bar{w}_{m-2}$ .

**Proof** By Corollary 3.7,  $Q_n(\bar{w}_{m-2}) = w_{Q_n(\gamma_1)} \bar{w}_{m-2}$ , where  $\gamma_1 \rightarrow \text{Gr}_1(\mathbb{R}^{m-1})$  is the canonical line bundle, and Proposition 3.5 tells us that  $w_{Q_n(\gamma_1)} = w_1^{2^{n+1}-1}$ . □

<sup>1</sup>There is a slight misprint, and a proof is just hinted at.

**Remark 5.6** This lemma also admits a proof using the Schubert cell perspective; see [10, Lemma 4.9.13].

As  $Q_n$  is a derivation, the lemma, together with the calculation  $Q_n(w_1) = w_1^{2^{n+1}}$ , allows one to easily compute the  $Q_n$ -homology of  $C_2(\mathbb{R}^m)$ . What results is the following.

**Proposition 5.7** (a) In  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$ ,

$$Q_n(w_1^i \bar{w}_{m-2}) = \begin{cases} w_1^{2^{n+1}-1+i} \bar{w}_{m-2} & \text{if } i \text{ is even,} \\ 0 & \text{if } i \text{ is odd.} \end{cases}$$

(b) If  $m \leq 2^{n+1}$ , then  $Q_n$  acts as zero on  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$ . Thus  $\tilde{k}_{Q_n}(C_2(\mathbb{R}^m)) = m - 1$ .

(c) If  $m > 2^{n+1}$  and is even, then the classes

$$\{w_1^{2j-1} \bar{w}_{m-2} \mid 1 \leq j \leq 2^n - 1\} \quad \text{and} \quad \{w_1^{m-2j} \bar{w}_{m-2} \mid 1 \leq j \leq 2^n\}$$

represent the  $Q_n$ -homology classes. Thus  $\tilde{k}_{Q_n}(C_2(\mathbb{R}^m)) = 2^{n+1} - 1$ .

(d) If  $m > 2^{n+1}$  and is odd, then the classes

$$\{w_1^{2j-1} \bar{w}_{m-2} \mid 1 \leq j \leq 2^n - 1\} \quad \text{and} \quad \{w_1^{m-1-2j} \bar{w}_{m-2} \mid 1 \leq j \leq 2^n - 1\}$$

represent the  $Q_n$ -homology classes. Thus  $\tilde{k}_{Q_n}(C_2(\mathbb{R}^m)) = 2^{n+1} - 2$ .

**Proof of Theorem 1.10** Let  $m = 2^{n+1} + 1 + 2l$ . We need to prove that the map

$$\tilde{H}^*(C_2(\mathbb{R}^m); Q_n) \xrightarrow{P^*} H^*(\text{Gr}_2(\mathbb{R}^m); Q_n)$$

is zero; ie we need to show that representatives of the  $Q_n$ -homology classes in  $\tilde{H}^*(C_2(\mathbb{R}^m); \mathbb{Z}/2)$  are in the image of  $Q_n$  when regarded in  $H^*(\text{Gr}_2(\mathbb{R}^m); \mathbb{Z}/2)$ .

By Proposition 5.7(d), these representatives are in two families,

$$w_1^{1+2j} \bar{w}_{2^{n+1}-1+2l} \quad \text{and} \quad w_1^{2l+2+2j} \bar{w}_{2^{n+1}-1+2l},$$

both with  $0 \leq j \leq 2^n - 2$ .

If we can find  $a, b \in H^*(\text{Gr}_2(\mathbb{R}^m); \mathbb{Z}/2)$  such that

$$Q_n(a) = w_1 \bar{w}_{2^{n+1}-1+2l} \quad \text{and} \quad Q_n(b) = w_1^{2l+2} \bar{w}_{2^{n+1}-1+2l},$$

we will be done, as then

$$Q_n(w_1^{2j} a) = w_1^{1+2j} \bar{w}_{2^{n+1}-1+2l} \quad \text{and} \quad Q_n(w_1^{2j} b) = w_1^{2l+2+2j} \bar{w}_{2^{n+1}-1+2l}.$$

Thus the next two propositions finish the proof. □

**Proposition 5.8** In  $H^*(\text{Gr}_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ ,

$$Q_n(w_1 \bar{w}_{2l}) = w_1 \bar{w}_{2^{n+1}-1+2l}.$$

**Proposition 5.9** In  $H^*(\mathrm{Gr}_2(\mathbb{R}^{2^{n+1}+1+2l}); \mathbb{Z}/2)$ ,

$$Q_n(w_2^{2l+1}) = w_1^{2l+2} \bar{w}_{2^{n+1}-1+2l}.$$

Before proving these, we first run through how [Theorem 1.10](#) leads to the proof of [Theorem 1.8](#).

**Proof of Theorem 1.8** Our goal is to show that if  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ , then  $k_{Q_n}(\mathrm{Gr}_2(\mathbb{R}^m)) = \binom{2^{n+1}-\epsilon}{2} + l$ , the lower bound coming from [Theorem 1.2](#).

We prove this by induction on  $m$ , with the two cases when  $l = 0$  already covered by [Theorem 1.1](#). The case when  $m$  is even is covered by [Theorem 1.9\(c\)](#), as we know our calculations are right for  $(n, 1, m - 1)$ , and by induction we can assume the theorem for  $(n, 2, m - 1)$ .

Suppose  $m$  is odd, so  $\epsilon = 1$  and  $m - 1 = 2^{n+1} + 2(l - 1)$ . By induction, we can assume that  $k_{Q_n}(\mathrm{Gr}_2(\mathbb{R}^{m-1})) = \binom{2^{n+1}}{2} + (l - 1)$ . Then

$$\begin{aligned} k_{Q_n}(\mathrm{Gr}_2(\mathbb{R}^m)) &= k_{Q_n}(\mathrm{Gr}_2(\mathbb{R}^{m-1})) - \bar{k}_{Q_n}(C_2(\mathbb{R}^m)) \quad (\text{by Theorem 1.10}) \\ &= \binom{2^{n+1}}{2} + (l - 1) - (2^{n+1} - 2) \quad (\text{by Proposition 5.7(d)}) \\ &= \binom{2^{n+1}-1}{2} + l. \end{aligned}$$

□

It remains to prove [Propositions 5.8](#) and [5.9](#).

**Proof of Proposition 5.8** We prove by induction on  $l$  that

$$Q_n(w_1 \bar{w}_{2l}) = w_1 \bar{w}_{2^{n+1}-1+2l}$$

holds in  $H^*(\mathrm{Gr}_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ .

We start the induction by checking both the  $l = 0$  and  $l = 1$  cases.

When  $l = 0$ , this reads  $Q_n(w_1) = w_1 \bar{w}_{2^{n+1}}$ , which was proved in [Lemma 5.2](#).

We check the  $l = 1$  case using both [Lemmas 5.2](#) and [5.3](#),

$$\begin{aligned} Q_n(w_1 \bar{w}_2) &= Q_n(w_1(w_2 + w_1^2)) \\ &= Q_n(w_1 w_2 + w_1^3) \\ &= Q_n(w_1)w_2 + w_1 Q_n(w_2) + w_1^2 Q_n(w_1) \\ &= w_1 w_2 \bar{w}_{2^{n+1}-1} + w_1^2 w_2 \bar{w}_{2^{n+1}-2} + w_1^3 \bar{w}_{2^{n+1}-1} \\ &= w_1 [w_2 \bar{w}_{2^{n+1}-1} + w_1 (w_2 \bar{w}_{2^{n+1}-2} + w_1 \bar{w}_{2^{n+1}-1})] \\ &= w_1 [w_2 \bar{w}_{2^{n+1}-1} + w_1 \bar{w}_{2^{n+1}}] \\ &= w_1 \bar{w}_{2^{n+1}+1}. \end{aligned}$$



For the inductive case, we use the identity  $\bar{w}_k = w_2^2 \bar{w}_{k-4} + w_1^2 \bar{w}_{k-2}$  which holds for all  $k \geq 4$ . Then

$$\begin{aligned} Q_n(w_1 \bar{w}_{2l}) &= Q_n(w_1 w_2^2 \bar{w}_{2(l-2)} + w_1^3 \bar{w}_{2(l-1)}) \\ &= w_2^2 Q_n(w_1 \bar{w}_{2(l-2)}) + w_1^2 Q_n(w_1 \bar{w}_{2(l-1)}) \\ &= w_1 w_2^2 \bar{w}_{2^{n+1}+2(l-2)-1} + w_1^3 \bar{w}_{2^{n+1}+2(l-1)-1} \\ &= w_1 [w_2^2 \bar{w}_{2^{n+1}+2(l-2)-1} + w_1^2 \bar{w}_{2^{n+1}+2(l-1)-1}] \\ &= w_1 \bar{w}_{2^{n+1}+2l-1}. \end{aligned}$$

□

**Proof of Proposition 5.9** We wish to prove that

$$Q_n(w_2^{2l+1}) = w_1^{2l+2} \bar{w}_{2^{n+1}-1+2l}$$

holds in  $H^*(\text{Gr}_2(\mathbb{R}^{2^{n+1}+1+2l}); \mathbb{Z}/2)$ .

We begin with a calculation in  $H^*(\text{Gr}_2(\mathbb{R}^\infty); \mathbb{Z}/2)$ ,

$$\begin{aligned} Q_n(w_2^{2l+1}) &= w_2^{2l} Q_n(w_2) \\ &= w_1 w_2^{2l+1} \bar{w}_{2^{n+1}-2} && \text{(using Lemma 5.3)} \\ &= \sum_i \binom{2l+1}{i} w_1^{2l+2-i} \bar{w}_{2^{n+1}+2l-1+i} && \text{(using Lemma 5.1(b)).} \end{aligned}$$

When we project this sum onto

$$H^*(\text{Gr}_2(\mathbb{R}^{2^{n+1}+1+2l}); \mathbb{Z}/2) = \mathbb{Z}/2[w_1, w_2]/(\bar{w}_k \mid k \geq 2^{n+1} + 2l),$$

only the term with  $i = 0$  is not zero. In other words

$$Q_n(w_2^{2l+1}) = w_1^{2l+2} \bar{w}_{2^{n+1}-1+2l}$$

holds in  $H^*(\text{Gr}_2(\mathbb{R}^{2^{n+1}+1+2l}); \mathbb{Z}/2)$ .

□

## 6 Towards the conjectures

As organized in this paper, we are trying to calculate  $H^*(\text{Gr}_d(\mathbb{R}^m); Q_n)$  by induction on  $m$  (and  $d$ ) with two steps:

- calculate  $\tilde{H}^*(C_d(\mathbb{R}^m); Q_n)$ , recalling that  $C_d(\mathbb{R}^m)$  is the Thom space of a bundle over  $\text{Gr}_{d-1}(\mathbb{R}^{m-1})$ ;
- calculate  $\delta: H^*(\text{Gr}_d(\mathbb{R}^{m-1}); Q_n) \rightarrow \tilde{H}^{*+2^{n+1}-1}(C_d(\mathbb{R}^m); Q_n)$ .

When  $m$  is even, **Theorem 1.9** says we can carry through with this plan. In this section we speculate about how things might go when  $m$  is odd.

Firstly, we have the analogues of Theorems 1.1 and 1.2 for

$$\bar{k}_n(C_d(\mathbb{R}^m)) = \dim_{K(n)_*} \tilde{K}(n)^*(C_d(\mathbb{R}^m)).$$

**Theorem 6.1** If  $m \leq 2^{n+1}$ , then  $\bar{k}_n(C_d(\mathbb{R}^m)) = \binom{m-1}{d-1}$ .

**Proof** [Theorem 1.1](#) implies that, if  $m \leq 2^{n+1}$ , the inclusion  $\text{Gr}_d(\mathbb{R}^{m-1}) \rightarrow \text{Gr}_d(\mathbb{R}^m)$  induces an inclusion  $K(n)_*(\text{Gr}_d(\mathbb{R}^{m-1})) \rightarrow K(n)_*(\text{Gr}_d(\mathbb{R}^m))$ , as this is true in mod  $p$  homology. Thus

$$\bar{k}_n(C_d(\mathbb{R}^m)) = k_n(\text{Gr}_d(\mathbb{R}^m)) - k_n(\text{Gr}_d(\mathbb{R}^{m-1})) = \binom{m}{d} - \binom{m-1}{d} = \binom{m-1}{d-1}. \quad \square$$

**Theorem 6.2** Let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . Then

$$k_n(C_d(\mathbb{R}^m)) \geq \sum_{i=0}^{\lfloor d/2 \rfloor} \binom{2^{n+1} - 1 - \epsilon}{d-1-2i} \binom{l}{i}.$$

**Proof** The proof is similar to the proof of [Theorem 1.2](#), with a little tweak.

If  $V$  is a real representation of  $C_4$ , and  $W$  is a subrepresentation, let  $C_d(V, W)$  denote the cofiber of the inclusion  $\text{Gr}_d(W) \hookrightarrow \text{Gr}_d(V)$ ; this is a based  $C_4$  space.

If  $\dim V = m$  and  $\dim W = m - 1$ , then  $C_d(\mathbb{R}^m) = C_d(V, W)$  and thus  $\bar{k}_n(C_d(\mathbb{R}^m)) \geq \bar{k}_n(C_d(V, W)^{C_4})$ , by our chromatic fixed point theorem, [Theorem 1.4](#). Furthermore,  $C_d(V, W)^{C_4}$  will be the cofiber of the inclusion  $\text{Gr}_d(W)^{C_4} \hookrightarrow \text{Gr}_d(V)^{C_4}$ .

Now we choose  $V$  and  $W$ . Recall that  $L_1$  and  $L_2$  were the one-dimensional real representations of  $C_4$  and  $R$  was the two-dimensional irreducible. We let  $V = L_1^{2^n} \oplus L_2^{2^n - \epsilon} \oplus R^l$  and  $W = L_1^{2^n - 1} \oplus L_2^{2^n - \epsilon} \oplus R^l$ .

[Proposition 2.1](#) tells us that

$$\text{Gr}_d(V)^{C_4} = \bigsqcup_{j+k+2i=d} \text{Gr}_j(\mathbb{R}^{2^n}) \times \text{Gr}_k(\mathbb{R}^{2^n - \epsilon}) \times \text{Gr}_i(\mathbb{C}^l)$$

and

$$\text{Gr}_d(W)^{C_4} = \bigsqcup_{j+k+2i=d} \text{Gr}_j(\mathbb{R}^{2^n - 1}) \times \text{Gr}_k(\mathbb{R}^{2^n - \epsilon}) \times \text{Gr}_i(\mathbb{C}^l),$$

so

$$C_d(V, W)^{C_4} = \bigvee_{j+k+2i=d} C_j(\mathbb{R}^{2^n}) \wedge \text{Gr}_k(\mathbb{R}^{2^n - \epsilon})_+ \wedge \text{Gr}_i(\mathbb{C}^l)_+.$$

Thus,

$$\begin{aligned} \bar{k}_n(C_d(\mathbb{R}^m)) &\geq \sum_{j+k+2i=d} \bar{k}_{n-1}(C_j(\mathbb{R}^{2^n})) k_{n-1}(\text{Gr}_k(\mathbb{R}^{2^n - \epsilon})) k_{n-1}(\text{Gr}_i(\mathbb{C}^l)) \\ &= \sum_{j+k+2i=d} \binom{2^n - 1}{j-1} \binom{2^n - \epsilon}{k} \binom{l}{i} \quad (\text{using Theorems 6.1 and 1.1}) \\ &= \sum_i \left[ \sum_{j+k=d-2i} \binom{2^n - 1}{j-1} \binom{2^n - \epsilon}{k} \right] \binom{l}{i} \\ &= \sum_i \binom{2^{n+1} - 1 - \epsilon}{d-1-2i} \binom{l}{i}. \end{aligned} \quad \square$$

**Conjecture 6.3** Equality holds in [Theorem 6.2](#).

As before, this would be implied by a conjectural calculation of the  $Q_n$  homology of  $C_d(\mathbb{R}_n)$ .

**Conjecture 6.4** Let  $m = 2^{n+1} - \epsilon + 2l$  with  $\epsilon = 0$  or  $1$ , and  $l \geq 0$ . Then

$$\bar{k}_{Q_n}(C_d(\mathbb{R}^m)) = \sum_i \binom{2^{n+1}-1-\epsilon}{d-1-2i} \binom{l}{i}.$$

Our various conjectures imply a conjecture about the behavior of the boundary map

$$\delta: H^*(\text{Gr}_d(\mathbb{R}^{m-1}); Q_n) \rightarrow \tilde{H}^{*+2^{n+1}-1}(C_d(\mathbb{R}^m); Q_n)$$

when  $m = 2^{n+1} - \epsilon + 2l$ . Let  $k_n^\delta(d, m)$  denote the dimension of the image of this map.

[Conjecture 1.7](#) says that  $k_{Q_n}(\text{Gr}_d(\mathbb{R}^m)) = k_n^G(d, m)$ , where

$$k_n^G(d, m) = \sum_i \binom{2^{n+1}-\epsilon}{d-2i} \binom{l}{i}.$$

[Conjecture 6.4](#) similarly says that  $\bar{k}_{Q_n}(C_d(\mathbb{R}^m)) = \bar{k}_n^C(d, m)$ , where

$$\bar{k}_n^C(d, m) = \sum_i \binom{2^{n+1}-1-\epsilon}{d-1-2i} \binom{l}{i}.$$

If these conjectures are true, then the exactness of the  $Q_n$ -homology long exact sequence would imply that

$$k_n^G(d, m) + 2k_n^\delta(d, m) = k_n^G(d, m-1) + \bar{k}_n^C(d, m),$$

so that

$$k_n^\delta(d, m) = \frac{1}{2}[k_n^G(d, m-1) + \bar{k}_n^C(d, m) - k_n^G(d, m)].$$

As expected, the right hand side here is zero if  $m$  is even, ie  $\epsilon = 0$ .

When  $m$  is odd, so  $\epsilon = 1$ , the right hand side is not zero, but can be rearranged as in the following lemma.

**Lemma 6.5** If  $m = 2^{n+1} - 1 + 2l$  and  $l > 0$ , then

$$\frac{1}{2}[k_n^G(d, m-1) + \bar{k}_n^C(d, m) - k_n^G(d, m)] = \sum_i \binom{2^{n+1}-2}{d-1-2i} \binom{l-1}{i}.$$

**Proof** We expand  $k_n^G(d, m-1)$ :

$$\begin{aligned} k_n^G(d, m-1) &= \sum_i \binom{2^{n+1}}{d-2i} \binom{l-1}{i} \\ &= \sum_i \left[ \binom{2^{n+1}-2}{d-2i} + 2\binom{2^{n+1}-2}{d-1-2i} + \binom{2^{n+1}-2}{d-2-2i} \right] \binom{l-1}{i}. \end{aligned}$$

We rewrite  $k_n^G(d, m) - \bar{k}_n^C(d, m)$ :

$$\begin{aligned} k_n^G(d, m) - \bar{k}_n^C(d, m) &= \sum_i \left[ \binom{2^{n+1}-1}{d-2i} - \binom{2^{n+1}-2}{d-1-2i} \right] \binom{l}{i} \\ &= \sum_i \binom{2^{n+1}-2}{d-2i} \binom{l}{i} \\ &= \sum_i \binom{2^{n+1}-2}{d-2i} \left[ \binom{l-1}{i} + \binom{l-1}{i-1} \right] \\ &= \sum_i \left[ \binom{2^{n+1}-2}{d-2i} + \binom{2^{n+1}-2}{d-2-2i} \right] \binom{l-1}{i}. \end{aligned}$$

Subtracting our second expression from the first, and dividing by two, proves the lemma. □

Thus we can add the following to our conjectures.

**Conjecture 6.6** *If  $m = 2^{n+1} - 1 + 2l$  and  $l > 0$ , then*

$$k_n^\delta(d, m) = \sum_i \binom{2^{n+1}-2}{d-1-2i} \binom{l-1}{i}.$$

**Example 6.7** Suppose that  $n = 0$ , so  $m = 2l + 1$ . [Conjecture 6.4](#) predicts that

$$\dim_{\mathbb{Q}} H^*(C_d(\mathbb{R}^{2l+1}); \mathbb{Q}) = \begin{cases} 0 & \text{if } d \text{ is even,} \\ \binom{l}{c} & \text{if } d = 2c + 1. \end{cases}$$

Similarly, [Conjecture 6.6](#) predicts that

$$k_0^\delta(d, 2l + 1) = \begin{cases} 0 & \text{if } d \text{ is even,} \\ \binom{l-1}{c} & \text{if } d = 2c + 1. \end{cases}$$

Noting that  $k_0^\delta(d, 2l + 1)$  can be viewed as the dimension of the cokernel of the map

$$i^*: H^*(Gr_d(\mathbb{R}^{2l+1}); \mathbb{Q}) \rightarrow H^*(Gr_d(\mathbb{R}^{2l}); \mathbb{Q}),$$

one can check that our conjectures do correspond to the known behavior of  $i^*$  — it takes Pontryagin classes to Pontryagin classes — together with the computations

$$\dim_{\mathbb{Q}} H^*(Gr_d(\mathbb{R}^m); \mathbb{Q}) = \begin{cases} \binom{l}{c} & \text{if } m = 2l + 1 \text{ and } d = 2c \text{ or } 2c + 1, \\ 2\binom{l-1}{c} & \text{if } m = 2l \text{ and } d = 2c + 1. \end{cases}$$

## Appendix Tables

We present some tables of calculations made by the second author that support [Conjecture 1.7](#). Computational algorithms used are documented in [\[10, Appendix B\]](#). For larger Grassmannians the authors used the University of Virginia Rivanna high-performance computing system. The white cells are the conjectured values which have not been checked due to computational limitations. The tables are necessarily symmetric in  $c$  and  $d$ .

| $d \backslash c$ | 1 | 2  | 3   | 4   | 5    | 6    | 7     | 8     | 9     | 10     | 11     |
|------------------|---|----|-----|-----|------|------|-------|-------|-------|--------|--------|
| 1                | 2 | 3  | 4   | 3   | 4    | 3    | 4     | 3     | 4     | 3      | 4      |
| 2                | 3 | 6  | 4   | 7   | 5    | 8    | 6     | 9     | 7     | 10     | 8      |
| 3                | 4 | 4  | 8   | 7   | 12   | 10   | 16    | 13    | 20    | 16     | 24     |
| 4                | 3 | 7  | 7   | 14  | 12   | 22   | 18    | 31    | 25    | 41     | 33     |
| 5                | 4 | 5  | 12  | 12  | 24   | 22   | 40    | 35    | 60    | 51     | 84     |
| 6                | 3 | 8  | 10  | 22  | 22   | 44   | 40    | 75    | 65    | 116    | 98     |
| 7                | 4 | 6  | 16  | 18  | 40   | 40   | 80    | 75    | 140   | 126    | 224    |
| 8                | 3 | 9  | 13  | 31  | 35   | 75   | 75    | 150   | 140   | 266    | 238    |
| 9                | 4 | 7  | 20  | 25  | 60   | 65   | 140   | 140   | 280   | 266    | 504    |
| 10               | 3 | 10 | 16  | 41  | 51   | 116  | 126   | 266   | 266   | 532    | 504    |
| 11               | 4 | 8  | 24  | 33  | 84   | 98   | 224   | 238   | 504   | 504    | 1008   |
| 12               | 3 | 11 | 19  | 52  | 70   | 168  | 196   | 434   | 462   | 966    | 966    |
| 13               | 4 | 9  | 28  | 42  | 112  | 140  | 336   | 378   | 840   | 882    | 1848   |
| 14               | 3 | 12 | 22  | 64  | 92   | 232  | 288   | 666   | 750   | 1632   | 1716   |
| 15               | 4 | 10 | 32  | 52  | 144  | 192  | 480   | 570   | 1320  | 1452   | 3168   |
| 16               | 3 | 13 | 25  | 77  | 117  | 309  | 405   | 975   | 1155  | 2607   | 2871   |
| 17               | 4 | 11 | 36  | 63  | 180  | 255  | 660   | 825   | 1980  | 2277   | 5148   |
| 18               | 3 | 14 | 28  | 91  | 145  | 400  | 550   | 1375  | 1705  | 3982   | 4576   |
| 19               | 4 | 12 | 40  | 75  | 220  | 330  | 880   | 1155  | 2860  | 3432   | 8008   |
| 20               | 3 | 15 | 31  | 106 | 176  | 506  | 726   | 1881  | 2431  | 5863   | 7007   |
| 21               | 4 | 13 | 44  | 88  | 264  | 418  | 1144  | 1573  | 4004  | 5005   | 12012  |
| 22               | 3 | 16 | 34  | 122 | 210  | 628  | 936   | 2509  | 3367  | 8372   | 10374  |
| 23               | 4 | 14 | 48  | 102 | 312  | 520  | 1456  | 2093  | 5460  | 7098   | 17472  |
| 24               | 3 | 17 | 37  | 139 | 247  | 767  | 1183  | 3276  | 4550  | 11648  | 14924  |
| 25               | 4 | 15 | 52  | 117 | 364  | 637  | 1820  | 2730  | 7280  | 9828   | 24752  |
| 26               | 3 | 18 | 40  | 157 | 287  | 924  | 1470  | 4200  | 6020  | 15848  | 20944  |
| 27               | 4 | 16 | 56  | 133 | 420  | 770  | 2240  | 3500  | 9520  | 13328  | 34272  |
| 28               | 3 | 19 | 43  | 176 | 330  | 1100 | 1800  | 5300  | 7820  | 21148  | 28764  |
| 29               | 4 | 17 | 60  | 150 | 480  | 920  | 2720  | 4420  | 12240 | 17748  | 46512  |
| 30               | 3 | 20 | 46  | 196 | 376  | 1296 | 2176  | 6596  | 9996  | 27744  | 38760  |
| 31               | 4 | 18 | 64  | 168 | 544  | 1088 | 3264  | 5508  | 15504 | 23256  | 62016  |
| 32               | 3 | 21 | 49  | 217 | 425  | 1513 | 2601  | 8109  | 12597 | 35853  | 51357  |
| 33               | 4 | 19 | 68  | 187 | 612  | 1275 | 3876  | 6783  | 19380 | 30039  | 81396  |
| 34               | 3 | 22 | 52  | 239 | 477  | 1752 | 3078  | 9861  | 15675 | 45714  | 67032  |
| 35               | 4 | 20 | 72  | 207 | 684  | 1482 | 4560  | 8265  | 23940 | 38304  | 105336 |
| 36               | 3 | 23 | 55  | 262 | 532  | 2014 | 3610  | 11875 | 19285 | 57589  | 86317  |
| 37               | 4 | 21 | 76  | 228 | 760  | 1710 | 5320  | 9975  | 29260 | 48279  | 134596 |
| 38               | 3 | 24 | 58  | 286 | 590  | 2300 | 4200  | 14175 | 23485 | 71764  | 109802 |
| 39               | 4 | 22 | 80  | 250 | 840  | 1960 | 6160  | 11935 | 35420 | 60214  | 170016 |
| 40               | 3 | 25 | 61  | 311 | 651  | 2611 | 4851  | 16786 | 28336 | 88550  | 138138 |
| 41               | 4 | 23 | 84  | 273 | 924  | 2233 | 7084  | 14168 | 42504 | 74382  | 212520 |
| 42               | 3 | 26 | 64  | 337 | 715  | 2948 | 5566  | 19734 | 33902 | 108284 | 172040 |
| 43               | 4 | 24 | 88  | 297 | 1012 | 2530 | 8096  | 16698 | 50600 | 91080  | 263120 |
| 44               | 3 | 27 | 67  | 364 | 782  | 3312 | 6348  | 23046 | 40250 | 131330 | 212290 |
| 45               | 4 | 25 | 92  | 322 | 1104 | 2852 | 9200  | 19550 | 59800 | 110630 | 322920 |
| 46               | 3 | 28 | 70  | 392 | 852  | 3704 | 7200  | 26750 | 47450 | 158080 | 259740 |
| 47               | 4 | 26 | 96  | 348 | 1200 | 3200 | 10400 | 22750 | 70200 | 133380 | 393120 |
| 48               | 3 | 29 | 73  | 421 | 925  | 4125 | 8125  | 30875 | 55575 | 188955 | 315315 |
| 49               | 4 | 27 | 100 | 375 | 1300 | 3575 | 11700 | 26325 | 81900 | 159705 | 475020 |
| 50               | 3 | 30 | 76  | 451 | 1001 | 4576 | 9126  | 35451 | 64701 | 224406 | 380016 |

   $cd \leq 2^{n+1} - 1$ 
  projective spaces
   Theorem 1.8  
  conjecture verified
   conjecture

Table 1:  $k_1(\text{Gr}_d(\mathbb{R}^{d+c}))$ .

| $d \backslash c$ | 1 | 2  | 3   | 4   | 5    | 6     | 7     | 8     | 9      | 10     | 11      |
|------------------|---|----|-----|-----|------|-------|-------|-------|--------|--------|---------|
| 1                | 2 | 3  | 4   | 5   | 6    | 7     | 8     | 7     | 8      | 7      | 8       |
| 2                | 3 | 6  | 10  | 15  | 21   | 28    | 22    | 29    | 23     | 30     | 24      |
| 3                | 4 | 10 | 20  | 35  | 56   | 42    | 64    | 49    | 72     | 56     | 80      |
| 4                | 5 | 15 | 35  | 70  | 56   | 98    | 78    | 127   | 101    | 157    | 125     |
| 5                | 6 | 21 | 56  | 56  | 112  | 98    | 176   | 147   | 248    | 203    | 328     |
| 6                | 7 | 28 | 42  | 98  | 98   | 196   | 176   | 323   | 277    | 480    | 402     |
| 7                | 8 | 22 | 64  | 78  | 176  | 176   | 352   | 323   | 600    | 526    | 928     |
| 8                | 7 | 29 | 49  | 127 | 147  | 323   | 323   | 646   | 600    | 1126   | 1002    |
| 9                | 8 | 23 | 72  | 101 | 248  | 277   | 600   | 600   | 1200   | 1126   | 2128    |
| 10               | 7 | 30 | 56  | 157 | 203  | 480   | 526   | 1126  | 1126   | 2252   | 2128    |
| 11               | 8 | 24 | 80  | 125 | 328  | 402   | 928   | 1002  | 2128   | 2128   | 4256    |
| 12               | 7 | 31 | 63  | 188 | 266  | 668   | 792   | 1794  | 1918   | 4046   | 4046    |
| 13               | 8 | 25 | 88  | 150 | 416  | 552   | 1344  | 1554  | 3472   | 3682   | 7728    |
| 14               | 7 | 32 | 70  | 220 | 336  | 888   | 1128  | 2682  | 3046   | 6728   | 7092    |
| 15               | 8 | 26 | 96  | 176 | 512  | 728   | 1856  | 2282  | 5328   | 5964   | 13056   |
| 16               | 7 | 33 | 77  | 253 | 413  | 1141  | 1541  | 3823  | 4587   | 10551  | 11679   |
| 17               | 8 | 27 | 104 | 203 | 616  | 931   | 2472  | 3213  | 7800   | 9177   | 20856   |
| 18               | 7 | 34 | 84  | 287 | 497  | 1428  | 2038  | 5251  | 6625   | 15802  | 18304   |
| 19               | 8 | 28 | 112 | 231 | 728  | 1162  | 3200  | 4375  | 11000  | 13552  | 31856   |
| 20               | 7 | 35 | 91  | 322 | 588  | 1750  | 2626  | 7001  | 9251   | 22803  | 27555   |
| 21               | 8 | 29 | 120 | 260 | 848  | 1422  | 4048  | 5797  | 15048  | 19349  | 46904   |
| 22               | 7 | 36 | 98  | 358 | 686  | 2108  | 3312  | 9109  | 12563  | 31912  | 40118   |
| 23               | 8 | 30 | 128 | 290 | 976  | 1712  | 5024  | 7509  | 20072  | 26858  | 66976   |
| 24               | 7 | 37 | 105 | 395 | 791  | 2503  | 4103  | 11612 | 16666  | 43524  | 56784   |
| 25               | 8 | 31 | 136 | 321 | 1112 | 2033  | 6136  | 9542  | 26208  | 36400  | 93184   |
| 26               | 7 | 38 | 112 | 433 | 903  | 2936  | 5006  | 14548 | 21672  | 58072  | 78456   |
| 27               | 8 | 32 | 144 | 353 | 1256 | 2386  | 7392  | 11928 | 33600  | 48328  | 126784  |
| 28               | 7 | 39 | 119 | 472 | 1022 | 3408  | 6028  | 17956 | 27700  | 76028  | 106156  |
| 29               | 8 | 33 | 152 | 386 | 1408 | 2772  | 8800  | 14700 | 42400  | 63028  | 169184  |
| 30               | 7 | 40 | 126 | 512 | 1148 | 3920  | 7176  | 21876 | 34876  | 97904  | 141032  |
| 31               | 8 | 34 | 160 | 420 | 1568 | 3192  | 10368 | 17892 | 52768  | 80920  | 221952  |
| 32               | 7 | 41 | 133 | 553 | 1281 | 4473  | 8457  | 26349 | 43333  | 124253 | 184365  |
| 33               | 8 | 35 | 168 | 455 | 1736 | 3647  | 12104 | 21539 | 64872  | 102459 | 286824  |
| 34               | 7 | 42 | 140 | 595 | 1421 | 5068  | 9878  | 31417 | 53211  | 155670 | 237576  |
| 35               | 8 | 36 | 176 | 491 | 1912 | 4138  | 14016 | 25677 | 78888  | 128136 | 365712  |
| 36               | 7 | 43 | 147 | 638 | 1568 | 5706  | 11446 | 37123 | 64657  | 192793 | 302233  |
| 37               | 8 | 37 | 184 | 528 | 2096 | 4666  | 16112 | 30343 | 95000  | 158479 | 460712  |
| 38               | 7 | 44 | 154 | 682 | 1722 | 6388  | 13168 | 43511 | 77825  | 236304 | 380058  |
| 39               | 8 | 38 | 192 | 566 | 2288 | 5232  | 18400 | 35575 | 113400 | 194054 | 574112  |
| 40               | 7 | 45 | 161 | 727 | 1883 | 7115  | 15051 | 50626 | 92876  | 286930 | 472934  |
| 41               | 8 | 39 | 200 | 605 | 2488 | 5837  | 20888 | 41412 | 134288 | 235466 | 708400  |
| 42               | 7 | 46 | 168 | 773 | 2051 | 7888  | 17102 | 58514 | 109978 | 345444 | 582912  |
| 43               | 8 | 40 | 208 | 645 | 2696 | 6482  | 23584 | 47894 | 157872 | 283360 | 866272  |
| 44               | 7 | 47 | 175 | 820 | 2226 | 8708  | 19328 | 67222 | 129306 | 412666 | 712218  |
| 45               | 8 | 41 | 216 | 686 | 2912 | 7168  | 26496 | 55062 | 184368 | 338422 | 1050640 |
| 46               | 7 | 48 | 182 | 868 | 2408 | 9576  | 21736 | 76798 | 151042 | 489464 | 863260  |
| 47               | 8 | 42 | 224 | 728 | 3136 | 7896  | 29632 | 62958 | 214000 | 401380 | 1264640 |
| 48               | 7 | 49 | 189 | 917 | 2597 | 10493 | 24333 | 87291 | 175375 | 576755 | 1038635 |
| 49               | 8 | 43 | 232 | 771 | 3368 | 8667  | 33000 | 71625 | 247000 | 473005 | 1511640 |
| 50               | 7 | 50 | 196 | 967 | 2793 | 11460 | 27126 | 98751 | 202501 | 675506 | 1241136 |

   $cd \leq 2^{n+1} - 1$ 
  projective spaces
   Theorem 1.8  
  conjecture verified
   conjecture
   Theorem 1.1

Table 2:  $k_2(\text{Gr}_d(\mathbb{R}^{d+c}))$ .

| $d \backslash c$ | 1  | 2   | 3   | 4    | 5     | 6     | 7      | 8      | 9       | 10      | 11       |
|------------------|----|-----|-----|------|-------|-------|--------|--------|---------|---------|----------|
| 1                | 2  | 3   | 4   | 5    | 6     | 7     | 8      | 9      | 10      | 11      | 12       |
| 2                | 3  | 6   | 10  | 15   | 21    | 28    | 36     | 45     | 55      | 66      | 78       |
| 3                | 4  | 10  | 20  | 35   | 56    | 84    | 120    | 165    | 220     | 286     | 364      |
| 4                | 5  | 15  | 35  | 70   | 126   | 210   | 330    | 495    | 715     | 1001    | 1365     |
| 5                | 6  | 21  | 56  | 126  | 252   | 462   | 792    | 1287   | 2002    | 3003    | 4368     |
| 6                | 7  | 28  | 84  | 210  | 462   | 924   | 1716   | 3003   | 5005    | 8008    | 6370     |
| 7                | 8  | 36  | 120 | 330  | 792   | 1716  | 3432   | 6435   | 11440   | 9438    | 15808    |
| 8                | 9  | 45  | 165 | 495  | 1287  | 3003  | 6435   | 12870  | 11440   | 20878   | 17810    |
| 9                | 10 | 55  | 220 | 715  | 2002  | 5005  | 11440  | 11440  | 22880   | 20878   | 38688    |
| 10               | 11 | 66  | 286 | 1001 | 3003  | 8008  | 9438   | 20878  | 20878   | 41756   | 38688    |
| 11               | 12 | 78  | 364 | 1365 | 4368  | 6370  | 15808  | 17810  | 38688   | 38688   | 77376    |
| 12               | 13 | 91  | 455 | 1820 | 3458  | 9828  | 12896  | 30706  | 33774   | 72462   | 72462    |
| 13               | 14 | 105 | 560 | 1470 | 4928  | 7840  | 20736  | 25650  | 59424   | 64338   | 136800   |
| 14               | 15 | 120 | 470 | 1940 | 3928  | 11768 | 16824  | 42474  | 50598   | 114936  | 123060   |
| 15               | 16 | 106 | 576 | 1576 | 5504  | 9416  | 26240  | 35066  | 85664   | 99404   | 222464   |
| 16               | 15 | 121 | 485 | 2061 | 4413  | 13829 | 21237  | 56303  | 71835   | 171239  | 194895   |
| 17               | 16 | 107 | 592 | 1683 | 6096  | 11099 | 32336  | 46165  | 118000  | 145569  | 340464   |
| 18               | 15 | 122 | 500 | 2183 | 4913  | 16012 | 26150  | 72315  | 97985   | 243554  | 292880   |
| 19               | 16 | 108 | 608 | 1791 | 6704  | 12890 | 39040  | 59055  | 157040  | 204624  | 497504   |
| 20               | 15 | 123 | 515 | 2306 | 5428  | 18318 | 31578  | 90633  | 129563  | 334187  | 422443   |
| 21               | 16 | 109 | 624 | 1900 | 7328  | 14790 | 46368  | 73845  | 203408  | 278469  | 700912   |
| 22               | 15 | 124 | 530 | 2430 | 5958  | 20748 | 37536  | 111381 | 167099  | 445568  | 589542   |
| 23               | 16 | 110 | 640 | 2010 | 7968  | 16800 | 54336  | 90645  | 257744  | 369114  | 958656   |
| 24               | 15 | 125 | 545 | 2555 | 6503  | 23303 | 44039  | 134684 | 211138  | 580252  | 800680   |
| 25               | 16 | 111 | 656 | 2121 | 8624  | 18921 | 62960  | 109566 | 320704  | 478680  | 1279360  |
| 26               | 15 | 126 | 560 | 2681 | 7063  | 25984 | 51102  | 160668 | 262240  | 740920  | 1062920  |
| 27               | 16 | 112 | 672 | 2233 | 9296  | 21154 | 72256  | 130720 | 392960  | 609400  | 1672320  |
| 28               | 15 | 127 | 575 | 2808 | 7638  | 28792 | 58740  | 189460 | 320980  | 930380  | 1383900  |
| 29               | 16 | 113 | 688 | 2346 | 9984  | 23500 | 82240  | 154220 | 475200  | 763620  | 2147520  |
| 30               | 15 | 128 | 590 | 2936 | 8228  | 31728 | 66968  | 221188 | 387948  | 1151568 | 1771848  |
| 31               | 16 | 114 | 704 | 2460 | 10688 | 25960 | 92928  | 180180 | 568128  | 943800  | 2715648  |
| 32               | 15 | 129 | 605 | 3065 | 8833  | 34793 | 75801  | 255981 | 463749  | 1407549 | 2235597  |
| 33               | 16 | 115 | 720 | 2575 | 11408 | 28535 | 104336 | 208715 | 672464  | 1152515 | 3388112  |
| 34               | 15 | 130 | 620 | 3195 | 9453  | 37988 | 85254  | 293969 | 549003  | 1701518 | 2784600  |
| 35               | 16 | 116 | 736 | 2691 | 12144 | 31226 | 116480 | 239941 | 788944  | 1392456 | 4177056  |
| 36               | 15 | 131 | 635 | 3326 | 10088 | 41314 | 95342  | 335283 | 644345  | 2036801 | 3428945  |
| 37               | 16 | 117 | 752 | 2808 | 12896 | 34034 | 129376 | 273975 | 918320  | 1666431 | 5095376  |
| 38               | 15 | 132 | 650 | 3458 | 10738 | 44772 | 106080 | 380055 | 750425  | 2416856 | 4179370  |
| 39               | 16 | 118 | 768 | 2926 | 13664 | 36960 | 143040 | 310935 | 1061360 | 1977366 | 6156736  |
| 40               | 15 | 133 | 665 | 3591 | 11403 | 48363 | 117483 | 428418 | 867908  | 2845274 | 5047278  |
| 41               | 16 | 119 | 784 | 3045 | 14448 | 40005 | 157488 | 350940 | 1218848 | 2328306 | 7375584  |
| 42               | 15 | 134 | 680 | 3725 | 12083 | 52088 | 129566 | 480506 | 997474  | 3325780 | 6044752  |
| 43               | 16 | 120 | 800 | 3165 | 15248 | 43170 | 172736 | 394110 | 1391584 | 2722416 | 8767168  |
| 44               | 15 | 135 | 695 | 3860 | 12778 | 55948 | 142344 | 536454 | 1139818 | 3862234 | 7184570  |
| 45               | 16 | 121 | 816 | 3286 | 16064 | 46456 | 188800 | 440566 | 1580384 | 3162982 | 10347552 |
| 46               | 15 | 136 | 710 | 3996 | 13488 | 59944 | 155832 | 596398 | 1295650 | 4458632 | 8480220  |
| 47               | 16 | 122 | 832 | 3408 | 16896 | 49864 | 205696 | 490430 | 1786080 | 3653412 | 12133632 |
| 48               | 15 | 137 | 725 | 4133 | 14213 | 64077 | 170045 | 660475 | 1465695 | 5119107 | 9945915  |
| 49               | 16 | 123 | 848 | 3531 | 17744 | 53395 | 223440 | 543825 | 2009520 | 4197237 | 14143152 |
| 50               | 15 | 138 | 740 | 4271 | 14953 | 68348 | 184998 | 728823 | 1650693 | 5847930 | 11596608 |

   $cd \leq 2^{n+1} - 1$ 
  projective spaces
   Theorem 1.8  
  conjecture verified
   conjecture
   Theorem 1.1

Table 3:  $k_3(\text{Gr}_d(\mathbb{R}^{d+c}))$ .

| $d \backslash c$ | 1  | 2   | 3    | 4     | 5      | 6       | 7       | 8        | 9        | 10        |
|------------------|----|-----|------|-------|--------|---------|---------|----------|----------|-----------|
| 1                | 2  | 3   | 4    | 5     | 6      | 7       | 8       | 9        | 10       | 11        |
| 2                | 3  | 6   | 10   | 15    | 21     | 28      | 36      | 45       | 55       | 66        |
| 3                | 4  | 10  | 20   | 35    | 56     | 84      | 120     | 165      | 220      | 286       |
| 4                | 5  | 15  | 35   | 70    | 126    | 210     | 330     | 495      | 715      | 1001      |
| 5                | 6  | 21  | 56   | 126   | 252    | 462     | 792     | 1287     | 2002     | 3003      |
| 6                | 7  | 28  | 84   | 210   | 462    | 924     | 1716    | 3003     | 5005     | 8008      |
| 7                | 8  | 36  | 120  | 330   | 792    | 1716    | 3432    | 6435     | 11440    | 19448     |
| 8                | 9  | 45  | 165  | 495   | 1287   | 3003    | 6435    | 12870    | 24310    | 43758     |
| 9                | 10 | 55  | 220  | 715   | 2002   | 5005    | 11440   | 24310    | 48620    | 92378     |
| 10               | 11 | 66  | 286  | 1001  | 3003   | 8008    | 19448   | 43758    | 92378    | 184756    |
| 11               | 12 | 78  | 364  | 1365  | 4368   | 12376   | 31824   | 75582    | 167960   | 352716    |
| 12               | 13 | 91  | 455  | 1820  | 6188   | 18564   | 50388   | 125970   | 293930   | 646646    |
| 13               | 14 | 105 | 560  | 2380  | 8568   | 27132   | 77520   | 203490   | 497420   | 1144066   |
| 14               | 15 | 120 | 680  | 3060  | 11628  | 38760   | 116280  | 319770   | 817190   | 1961256   |
| 15               | 16 | 136 | 816  | 3876  | 15504  | 54264   | 170544  | 490314   | 1307504  | 3268760   |
| 16               | 17 | 153 | 969  | 4845  | 20349  | 74613   | 245157  | 735471   | 2042975  | 5311735   |
| 17               | 18 | 171 | 1140 | 5985  | 26334  | 100947  | 346104  | 1081575  | 3124550  | 8436285   |
| 18               | 19 | 190 | 1330 | 7315  | 33649  | 134596  | 480700  | 1562275  | 4686825  | 13123110  |
| 19               | 20 | 210 | 1540 | 8855  | 42504  | 177100  | 657800  | 2220075  | 6906900  | 20030010  |
| 20               | 21 | 231 | 1771 | 10626 | 53130  | 230230  | 888030  | 3108105  | 10015005 | 30045015  |
| 21               | 22 | 253 | 2024 | 12650 | 65780  | 296010  | 1184040 | 4292145  | 14307150 | 44352165  |
| 22               | 23 | 276 | 2300 | 14950 | 80730  | 376740  | 1560780 | 5852925  | 20160075 | 64512240  |
| 23               | 24 | 300 | 2600 | 17550 | 98280  | 475020  | 2035800 | 7888725  | 28048800 | 52240890  |
| 24               | 25 | 325 | 2925 | 20475 | 118755 | 593775  | 2629575 | 10518300 | 22789650 | 75030540  |
| 25               | 26 | 351 | 3276 | 23751 | 142506 | 736281  | 3365856 | 8625006  | 31414656 | 60865896  |
| 26               | 27 | 378 | 3654 | 27405 | 169911 | 906192  | 2799486 | 11424492 | 25589136 | 86455032  |
| 27               | 28 | 406 | 4060 | 31465 | 201376 | 767746  | 3567232 | 9392752  | 34981888 | 70258648  |
| 28               | 29 | 435 | 4495 | 35960 | 174406 | 942152  | 2973892 | 12366644 | 28563028 | 98821676  |
| 29               | 30 | 465 | 4960 | 31930 | 206336 | 799676  | 3773568 | 10192428 | 38755456 | 80451076  |
| 30               | 31 | 496 | 4526 | 36456 | 178932 | 978608  | 3152824 | 13345252 | 31715852 | 112166928 |
| 31               | 32 | 466 | 4992 | 32396 | 211328 | 832072  | 3984896 | 11024500 | 42740352 | 91475576  |
| 32               | 31 | 497 | 4557 | 36953 | 183489 | 1015561 | 3336313 | 14360813 | 35052165 | 126527741 |
| 33               | 32 | 467 | 5024 | 32863 | 216352 | 864935  | 4201248 | 11889435 | 46941600 | 103365011 |
| 34               | 31 | 498 | 4588 | 37451 | 188077 | 1053012 | 3524390 | 15413825 | 38576555 | 141941566 |
| 35               | 32 | 468 | 5056 | 33331 | 221408 | 898266  | 4422656 | 12787701 | 51364256 | 116152712 |
| 36               | 31 | 499 | 4619 | 37950 | 192696 | 1090962 | 3717086 | 16504787 | 42293641 | 158446353 |
| 37               | 32 | 469 | 5088 | 33800 | 226496 | 932066  | 4649152 | 13719767 | 56013408 | 129872479 |
| 38               | 31 | 500 | 4650 | 38450 | 197346 | 1129412 | 3914432 | 17634199 | 46208073 | 176080552 |
| 39               | 32 | 470 | 5120 | 34270 | 231616 | 966336  | 4880768 | 14686103 | 60894176 | 144558582 |
| 40               | 31 | 501 | 4681 | 38951 | 202027 | 1168363 | 4116459 | 18802562 | 50324532 | 194883114 |
| 41               | 32 | 471 | 5152 | 34741 | 236768 | 1001077 | 5117536 | 15687180 | 66011712 | 160245762 |
| 42               | 31 | 502 | 4712 | 39453 | 206739 | 1207816 | 4323198 | 20010378 | 54647730 | 214893492 |
| 43               | 32 | 472 | 5184 | 35213 | 241952 | 1036290 | 5359488 | 16723470 | 71371200 | 176969232 |
| 44               | 31 | 503 | 4743 | 39956 | 211482 | 1247772 | 4534680 | 21258150 | 59182410 | 236151642 |
| 45               | 32 | 473 | 5216 | 35686 | 247168 | 1071976 | 5606656 | 17795446 | 76977856 | 194764678 |
| 46               | 31 | 504 | 4774 | 40460 | 216256 | 1288232 | 4750936 | 22546382 | 63933346 | 258698024 |
| 47               | 32 | 474 | 5248 | 36160 | 252416 | 1108136 | 5859072 | 18903582 | 82836928 | 213668260 |
| 48               | 31 | 505 | 4805 | 40965 | 221061 | 1329197 | 4971997 | 23875579 | 68905343 | 282573603 |
| 49               | 32 | 475 | 5280 | 36635 | 257696 | 1144771 | 6116768 | 20048353 | 88953696 | 233716613 |
| 50               | 31 | 506 | 4836 | 41471 | 225897 | 1370668 | 5197894 | 25246247 | 74103237 | 307819850 |

   $cd \leq 2^{n+1} - 1$ 
  projective spaces
   Theorem 1.8  
  conjecture verified
   conjecture
   Theorem 1.1

Table 4:  $k_4(\text{Gr}_d(\mathbb{R}^{d+c}))$ .



## References

- [1] **W Balderrama, N J Kuhn**, *An elementary proof of the chromatic Smith fixed point theorem*, Homology Homotopy Appl. 26 (2024) 131–140 [MR](#) [Zbl](#)
- [2] **T Barthel, M Hausmann, N Naumann, T Nikolaus, J Noel, N Stapleton**, *The Balmer spectrum of the equivariant homotopy category of a finite abelian group*, Invent. Math. 216 (2019) 215–240 [MR](#) [Zbl](#)
- [3] **C Ehresmann**, *Sur la topologie de certaines variétés algébriques réelles*, J. Math. Pures Appl. 16 (1937) 69–100 [Zbl](#)
- [4] **E E Floyd**, *On periodic maps and the Euler characteristics of associated spaces*, Trans. Amer. Math. Soc. 72 (1952) 138–147 [MR](#) [Zbl](#)
- [5] **J Jaworowski**, *An additive basis for the cohomology of real Grassmannians*, from “Algebraic topology Poznań 1989” (S Jackowski, B Oliver, K Pawałowski, editors), Lecture Notes in Math. 1474, Springer (1991) 231–234 [MR](#) [Zbl](#)
- [6] **N Kitchloo, W S Wilson**, *The Morava  $K$ -theory of  $BO(q)$  and  $MO(q)$* , Algebr. Geom. Topol. 15 (2015) 3049–3058 [MR](#) [Zbl](#)
- [7] **A Kono, N Yagita**, *Brown–Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups*, Trans. Amer. Math. Soc. 339 (1993) 781–798 [MR](#) [Zbl](#)
- [8] **N J Kuhn, C J R Lloyd**, *Chromatic fixed point theory and the Balmer spectrum for extraspecial 2-groups*, preprint (2020) [arXiv 2008.00330](#) To appear in Amer. J. Math.
- [9] **C Lenart**, *The combinatorics of Steenrod operations on the cohomology of Grassmannians*, Adv. Math. 136 (1998) 251–283 [MR](#) [Zbl](#)
- [10] **C J R Lloyd**, *Applications of chromatic fixed point theory*, PhD thesis, University of Virginia (2021) Available at [https://libraetd.lib.virginia.edu/public\\_view/h702q715z](https://libraetd.lib.virginia.edu/public_view/h702q715z)
- [11] **C R F Maunder**, *The spectral sequence of an extraordinary cohomology theory*, Proc. Cambridge Philos. Soc. 59 (1963) 567–574 [MR](#) [Zbl](#)
- [12] **J W Milnor, J D Stasheff**, *Characteristic classes*, Annals of Mathematics Studies No. 76, Princeton Univ. Press (1974) [MR](#) [Zbl](#)
- [13] **B Schuster**, *Morava  $K$ -theory of groups of order 32*, Algebr. Geom. Topol. 11 (2011) 503–521 [MR](#) [Zbl](#)
- [14] **U Würigler**, *Morava  $K$ -theories: a survey*, from “Algebraic topology” (S Jackowski, B Oliver, K Pawałowski, editors), Lecture Notes in Math. 1474, Springer (1991) 111–138 [MR](#) [Zbl](#)
- [15] **N Yagita**, *On the Steenrod algebra of Morava  $K$ -theory*, J. London Math. Soc. 22 (1980) 423–438 [MR](#) [Zbl](#)

Department of Mathematics, University of Virginia

Charlottesville, VA, United States

Arlington, VA, United States

[nj4x@virginia.edu](mailto:njk4x@virginia.edu), [cjl8zf@virginia.edu](mailto:cjl8zf@virginia.edu)

Received: 16 November 2021      Revised: 14 July 2022

# ALGEBRAIC & GEOMETRIC TOPOLOGY

[msp.org/agt](https://msp.org/agt)

## EDITORS

### PRINCIPAL ACADEMIC EDITORS

John Etnyre  
[etnyre@math.gatech.edu](mailto:etnyre@math.gatech.edu)  
Georgia Institute of Technology

Kathryn Hess  
[kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)  
École Polytechnique Fédérale de Lausanne

### BOARD OF EDITORS

|                        |   |                   |   |
|------------------------|---|-------------------|---|
| Julie Bergner          | University of Virginia<br><a href="mailto:jeb2md@eservices.virginia.edu">jeb2md@eservices.virginia.edu</a>          | Robert Lipshitz   | University of Oregon<br><a href="mailto:lipshitz@uoregon.edu">lipshitz@uoregon.edu</a>                  |
| Steven Boyer           | Université du Québec à Montréal<br><a href="mailto:cohf@math.rochester.edu">cohf@math.rochester.edu</a>             | Norihiko Minami   | Nagoya Institute of Technology<br><a href="mailto:nori@nitech.ac.jp">nori@nitech.ac.jp</a>              |
| Tara E Brendle         | University of Glasgow<br><a href="mailto:tara.brendle@glasgow.ac.uk">tara.brendle@glasgow.ac.uk</a>                 | Andrés Navas      | Universidad de Santiago de Chile<br><a href="mailto:andres.navas@usach.cl">andres.navas@usach.cl</a>    |
| Indira Chatterji       | CNRS & Univ. Côte d'Azur (Nice)<br><a href="mailto:indira.chatterji@math.cnrs.fr">indira.chatterji@math.cnrs.fr</a> | Thomas Nikolaus   | University of Münster<br><a href="mailto:nikolaus@uni-muenster.de">nikolaus@uni-muenster.de</a>         |
| Alexander Dranishnikov | University of Florida<br><a href="mailto:dranish@math.ufl.edu">dranish@math.ufl.edu</a>                             | Robert Oliver     | Université Paris 13<br><a href="mailto:bobol@math.univ-paris13.fr">bobol@math.univ-paris13.fr</a>       |
| Tobias Ekholm          | Uppsala University, Sweden<br><a href="mailto:tobias.ekholm@math.uu.se">tobias.ekholm@math.uu.se</a>                | Jessica S Purcell | Monash University<br><a href="mailto:jessica.purcell@monash.edu">jessica.purcell@monash.edu</a>         |
| Mario Eudave-Muñoz     | Univ. Nacional Autónoma de México<br><a href="mailto:mario@matem.unam.mx">mario@matem.unam.mx</a>                   | Birgit Richter    | Universität Hamburg<br><a href="mailto:birgit.richter@uni-hamburg.de">birgit.richter@uni-hamburg.de</a> |
| David Futer            | Temple University<br><a href="mailto:dfuter@temple.edu">dfuter@temple.edu</a>                                       | Jérôme Scherer    | École Polytech. Féd. de Lausanne<br><a href="mailto:jerome.scherer@epfl.ch">jerome.scherer@epfl.ch</a>  |
| John Greenlees         | University of Warwick<br><a href="mailto:john.greenlees@warwick.ac.uk">john.greenlees@warwick.ac.uk</a>             | Vesna Stojanoska  | Univ. of Illinois at Urbana-Champaign<br><a href="mailto:vesna@illinois.edu">vesna@illinois.edu</a>     |
| Ian Hambleton          | McMaster University<br><a href="mailto:ian@math.mcmaster.ca">ian@math.mcmaster.ca</a>                               | Zoltán Szabó      | Princeton University<br><a href="mailto:szabo@math.princeton.edu">szabo@math.princeton.edu</a>          |
| Matthew Hedden         | Michigan State University<br><a href="mailto:mhedden@math.msu.edu">mhedden@math.msu.edu</a>                         | Maggy Tomova      | University of Iowa<br><a href="mailto:maggy-tomova@uiowa.edu">maggy-tomova@uiowa.edu</a>                |
| Hans-Werner Henn       | Université Louis Pasteur<br><a href="mailto:henn@math.u-strasbg.fr">henn@math.u-strasbg.fr</a>                      | Nathalie Wahl     | University of Copenhagen<br><a href="mailto:wahl@math.ku.dk">wahl@math.ku.dk</a>                        |
| Daniel Isaksen         | Wayne State University<br><a href="mailto:isaksen@math.wayne.edu">isaksen@math.wayne.edu</a>                        | Chris Wendl       | Humboldt-Universität zu Berlin<br><a href="mailto:wendl@math.hu-berlin.de">wendl@math.hu-berlin.de</a>  |
| Thomas Koberda         | University of Virginia<br><a href="mailto:thomas.koberda@virginia.edu">thomas.koberda@virginia.edu</a>              | Daniel T Wise     | McGill University, Canada<br><a href="mailto:daniel.wise@mcgill.ca">daniel.wise@mcgill.ca</a>           |
| Christine Lescop       | Université Joseph Fourier<br><a href="mailto:lescop@ujf-grenoble.fr">lescop@ujf-grenoble.fr</a>                     |                   |   |

---

See inside back cover or [msp.org/agt](https://msp.org/agt) for submission instructions.

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

---

AGT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<https://msp.org/>

© 2024 Mathematical Sciences Publishers

# ALGEBRAIC & GEOMETRIC TOPOLOGY

Volume 24 Issue 2 (pages 595–1223) 2024

---

|   |      |
|---|------|
| Comparing combinatorial models of moduli space and their compactifications                | 595  |
| DANIELA EGAS SANTANDER and ALEXANDER KUPERS   |      |
| Towards a higher-dimensional construction of stable/unstable Lagrangian laminations       | 655  |
| SANGJIN LEE   |      |
| A strong Haken theorem  | 717  |
| MARTIN SCHARLEMANN  |      |
| Right-angled Artin subgroups of right-angled Coxeter and Artin groups                     | 755  |
| PALLAVI DANI and IVAN LEVCOVITZ   |      |
| Filling braided links with trisected surfaces   | 803  |
| JEFFREY MEIER   |      |
| Equivariantly slicing strongly negative amphichiral knots                                 | 897  |
| KEEGAN BOYLE and AHMAD ISSA   |      |
| Computing the Morava $K$ -theory of real Grassmannians using chromatic fixed point theory | 919  |
| NICHOLAS J KUHN and CHRISTOPHER J R LLOYD   |      |
| Slope gap distributions of Veech surfaces   | 951  |
| LUIS KUMANDURI, ANTHONY SANCHEZ and JANE WANG   |      |
| Embedding calculus for surfaces   | 981  |
| MANUEL KRANNICH and ALEXANDER KUPERS  |      |
| Vietoris–Rips persistent homology, injective metric spaces, and the filling radius        | 1019 |
| SUNHYUK LIM, FACUNDO MÉMOLI and OSMAN BERAT OKUTAN  |      |
| Slopes and concordance of links   | 1101 |
| ALEX DEGTYAREV, VINCENT FLORENS and ANA G LECUONA   |      |
| Cohomological and geometric invariants of simple complexes of groups                      | 1121 |
| NANSEN PETROSYAN and TOMASZ PRYTUŁA   |      |
| On the decategorification of some higher actions in Heegaard Floer homology               | 1157 |
| ANDREW MANION   |      |
| A simplicial version of the 2–dimensional Fulton–MacPherson operad                        | 1183 |
| NATHANIEL BOTTMAN   |      |
| Intrinsically knotted graphs with linklessly embeddable simple minors                     | 1203 |
| THOMAS W MATTMAN, RAMIN NAIMI, ANDREI PAVELESCU and ELENA PAVELESCU                       |      |