

Algebra & Number Theory

Volume 3

2009

No. 1

Frobenius splittings of toric varieties

Sam Payne

 mathematical sciences publishers

Frobenius splittings of toric varieties

Sam Payne

We discuss a characteristic free version of Frobenius splittings for toric varieties and give a polyhedral criterion for a toric variety to be diagonally split. We apply this criterion to show that section rings of nef line bundles on diagonally split toric varieties are normally presented and Koszul, and that Schubert varieties are not diagonally split in general.

1. Introduction

Fix an integer q greater than one. Let $T = \text{Spec } \mathbb{Z}[M]$ be the torus with character lattice M , and let N be the dual lattice. Let Σ be a complete fan in $N_{\mathbb{R}}$, with $X = X(\Sigma)$ the associated toric variety over \mathbb{Z} . Multiplication by q preserves the fan and maps the lattice N into itself, and therefore gives an endomorphism

$$F : X \rightarrow X.$$

Each T -orbit in X is a torus that is preserved by F , which acts by taking a point t to t^q . For example, if X is projective space, then F is given in homogeneous coordinates by

$$[x_0 : \cdots : x_n] \mapsto [x_0^q : \cdots : x_n^q].$$

If q is prime and k is the field with q elements, then the restriction of F to the variety X_k is the absolute Frobenius morphism. Pulling back functions by F gives a natural inclusion of \mathbb{O}_X -algebras $F^* : \mathbb{O}_X \hookrightarrow F_*\mathbb{O}_X$.

Definition 1.1. A *splitting of X* is an \mathbb{O}_X -module map $\pi : F_*\mathbb{O}_X \rightarrow \mathbb{O}_X$ such that the composition $\pi \circ F^*$ is the identity on \mathbb{O}_X .

Standard results from the theory of Frobenius splittings generalize in a straightforward way to these splittings of toric varieties. See [Section 2](#) for details.

If Y is a subvariety of X cut out by an ideal sheaf I_Y and $\pi(F_*I_Y)$ is contained in I_Y then we say that π is *compatible* with Y . If Y is a toric variety embedded equivariantly in X , the closure of a subtorus of an orbit in X , then a splitting compatible with Y induces a splitting of Y . We say that X is *diagonally split* if

MSC2000: primary 14M25; secondary 13A35, 14M15, 16S37.

Keywords: Frobenius splitting, toric variety, diagonal splitting, Koszul.

Supported by the Clay Mathematics Institute.

there is a splitting of $X \times X$ that is compatible with the diagonal, for some q . Such splittings are of particular interest; by classic arguments of Mehta, Ramanan and Ramanathan, if X is diagonally split then every ample line bundle on X is very ample and defines a projectively normal embedding.

Our main result is a polyhedral criterion for a toric variety to be diagonally split. Let v_ρ denote the primitive generator of a ray, or one-dimensional cone, in Σ . Let $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$, and let the *diagonal splitting polytope* \mathbb{F}_X be defined by

$$\mathbb{F}_X = \{u \in M_{\mathbb{R}} \mid -1 \leq \langle u, v_\rho \rangle \leq 1 \text{ for all } \rho \in \Sigma\}.$$

We write $\frac{1}{q}M$ for the subgroup of $M_{\mathbb{R}}$ consisting of *fractional lattice points* u such that qu is in M .

Theorem 1.2. *The toric variety X is diagonally split if and only if the interior of \mathbb{F}_X contains representatives of every equivalence class in $\frac{1}{q}M/M$.*

While the existence of a compatible splitting of the diagonal in $X \times X$ implies that section rings of ample line bundles on X are generated in degree one, compatible splittings of large semidiagonals in products of multiple copies of X give further information on these section rings. For example, if the union of $\Delta \times X$ and $X \times \Delta$ is compatibly split in $X \times X \times X$, where Δ is the diagonal in $X \times X$, then it follows from standard arguments that the section ring of each ample line bundle on X is *normally presented*, that is, generated in degree one with relations generated in degree two. For fixed n greater than one, let Δ_i be the large semidiagonal

$$\Delta_i = X^{i-1} \times \Delta \times X^{n-i-1},$$

for $1 \leq i < n$.

Theorem 1.3. *Let X be a diagonally split toric variety. Then $\Delta_1 \cup \dots \cup \Delta_{n-1}$ is compatibly split in X^n .*

In particular, if X is diagonally split then the union of $\Delta \times X$ and $X \times \Delta$ is compatibly split in $X \times X \times X$, so the section ring of any ample line bundle on X is normally presented. Analogous results hold for any finite collection of nef line bundles on X , as we now discuss.

For line bundles L_1, \dots, L_r on X , let $R(L_1, \dots, L_r)$ be the section ring

$$R(L_1, \dots, L_r) = \bigoplus_{(\alpha_1, \dots, \alpha_r) \in \mathbb{N}^r} H^0(X, L_1^{\alpha_1} \otimes \dots \otimes L_r^{\alpha_r}).$$

We consider $R(L_1, \dots, L_r)$ as a graded ring, where the degree of

$$H^0(X, L_1^{\alpha_1} \otimes \dots \otimes L_r^{\alpha_r})$$

is $\alpha_1 + \dots + \alpha_r$. In particular, the degree zero part of $R(L_1, \dots, L_r)$ is \mathbb{Z} . Recall that a graded ring R is *Koszul* if the ideal generated by elements of positive degree

has a linear resolution as an R -module. See [Polishchuk and Positselski 2005] for background on Koszul rings and further details.

Theorem 1.4. *Let X be a complete, diagonally split toric variety, and let*

$$L_1, \dots, L_r$$

be nef line bundles on X . Then the section ring $R(L_1, \dots, L_r)$ is normally presented and Koszul.

In particular, if X is diagonally split then the section ring of any ample line bundle on X is normally presented and Koszul. Well-known open problems ask whether every ample line bundle on a smooth projective toric variety gives a projectively normal embedding [Oda 1997] and, if so, whether its section ring is normally presented [Sturmfels 1996, Conjecture 13.19]. When the section ring is normally presented, it is natural to ask whether it is also Koszul. Addressing these questions and their analogues for singular toric varieties in as many cases as possible is one of the main motivations behind this work.

Remark 1.5. The section ring $R(L_1, \dots, L_r)$ associated to a finite collection of line bundles is canonically identified with the section ring of the line bundle $\mathcal{O}(1)$ on the projectivized vector bundle $\mathbb{P}(L_1 \oplus \dots \oplus L_r)$, which is also a toric variety. If L_1, \dots, L_r are nef and correspond to polytopes P_1, \dots, P_r , then the Cayley sum is the polytope associated to $\mathcal{O}(1)$. Cayley sums have also appeared prominently in recent work related to boundedness questions in toric mirror symmetry [Batyrev and Nill 2008; 2007; Haase et al. 2008].


Remark 1.6. Frobenius morphisms and their lifts to characteristic zero have been used powerfully in several other contexts related to the geometry of toric varieties, including by Buch, Lauritzen, Mehta and Thomsen [1997] to prove Bott vanishing and degeneration of the Hodge to de Rham spectral sequence, by Totaro [199?] to give a splitting of the weight filtration on Borel–Moore homology, by Smith [2000] to prove global F -regularity, by Brylinski and Zhang [2003] to prove degeneration of a spectral sequence computing equivariant cohomology with rational coefficients, and by Fujino [2007] to prove vanishing theorems for vector bundles and reflexive sheaves. Frobenius splittings have also played a role in unsuccessful attempts to show that section rings of ample line bundles on smooth toric varieties are normally presented [Bøgvad 1995]. We hope that this work will help revive the insight of Bøgvad and others into the potential usefulness of Frobenius splittings as a tool for understanding ample line bundles on toric varieties.

We conclude the introduction with an example illustrating Theorem 1.2 for Hirzebruch surfaces. As mentioned earlier, the proofs that section rings of ample line bundles on Schubert varieties are normally presented and Koszul via Frobenius

splittings involved compatible splittings of semidiagonals in $(G/B)^n$. It has been an open question for over twenty years whether Schubert varieties themselves are diagonally split (see [Ramanathan 1987, Remark 3.6] and [Brion and Kumar 2005, p. 81]). The following example gives a negative answer: the Hirzebruch surface F_3 is a Schubert variety in the G_2 -flag variety, and F_3 is not diagonally split.

Remark 1.7. To see that F_3 occurs as a Schubert variety in the G_2 -flag variety, first note that for any G , G/B is a \mathbb{P}^1 -bundle over G/P , where P is a minimal parabolic subgroup. If $w = s_1s_2$ is an element of length two in the Weyl group of G , and P is the minimal parabolic corresponding to s_1 , then X_w is a \mathbb{P}^1 -bundle over its image, which is a rational curve in G/P . In particular, X_w is a Hirzebruch surface. Then X_{s_1} is a rational curve in X_w with self-intersection $\langle \alpha_2, \alpha_1^\vee \rangle$, where α_i is the simple root corresponding to s_i , and $\alpha_i^\vee = 2\alpha_i / \langle \alpha_i, \alpha_i \rangle$. See [Kempf 1976, Section 2] for details. For G_2 , we can choose coordinates identifying the root lattice with the sublattice of \mathbb{Z}^3 consisting of those (a_1, a_2, a_3) such that $a_1 + a_2 + a_3 = 0$, with simple roots $\alpha_1 = (1, -1, 0)$ and $\alpha_2 = (-1, 2, -1)$. Then X_{s_1} is a curve of self-intersection -3 in X_w , and hence X_w is isomorphic to F_3 . See also [Anderson 2007] for a detailed study of the G_2 -flag variety and its Schubert varieties.

Example 1.8. Let a be a nonnegative integer, and let Σ be the complete fan in \mathbb{R}^2 whose rays are spanned by $(1, 0)$, $(0, 1)$, $(0, -1)$, and $(-1, a)$. Then $X(\Sigma)$ is isomorphic to the Hirzebruch surface F_a , the projectivization of the vector bundle $\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(a)$ [Fulton 1993, pp. 7–8]. Let $q \geq 2$ be an integer. By Theorem 1.2, X is diagonally split if and only if the fractional lattice points in the interior of \mathbb{F}_X represent every equivalence class in $\frac{1}{q}\mathbb{Z}^2/\mathbb{Z}^2$. The polytopes \mathbb{F}_X for different values of a are shown below.




If a is equal to 0 or 1, the interior of \mathbb{F}_X contains the half open unit square $[0, 1) \times [0, 1)$, which contains representatives of every equivalence class in $\frac{1}{q}\mathbb{Z}^2/\mathbb{Z}^2$. Therefore, F_0 and F_1 are diagonally split for all q .

If $a = 2$, then \mathbb{F}_X is the parallelogram with vertices $(\pm 1, 0)$, $\pm(1, 1)$. For $0 \leq m < q$, the interior of this intersection contains the fractional lattice points

$(m/q, n/q)$ for $(m - q)/2 < n < (m + q)/2$. If q is odd, then these represent every equivalence class in $\frac{1}{q}\mathbb{Z}^2/\mathbb{Z}^2$. In particular, F_2 is diagonally split for q odd.

If a is greater than two, then \mathbb{F}_X is the parallelogram with vertices $(\pm 1, 0)$, $\pm(1, 2/a)$. The only points in the interior of this intersection whose first coordinate is integral, are of the form $(0, y)$ for $-1/a < y < 1/a$. In particular, the equivalence class of $(0, \lfloor q/2 \rfloor / q)$ in $\frac{1}{q}\mathbb{Z}^2/\mathbb{Z}^2$ is not represented by any point in the interior of this intersection. Therefore, F_a is not diagonally split for a greater than two.



2. Preliminaries

Frobenius splittings were introduced and developed by Mehta, Ramanathan, and their collaborators in the 1980s. The original paper of Mehta and Ramanathan is exceedingly well written and remains an excellent first introduction to the subject [1985]. Frobenius splittings were rapidly applied to give elegant unified proofs that all ample line bundles on generalized Schubert varieties of all types are very ample and give projectively normal embeddings whose images are cut out by quadrics [Ramanan and Ramanathan 1985; Ramanathan 1987]. Inamdar and Mehta [1994], and independently Bezrukavnikov [1995], later showed that the homogeneous coordinate rings of these embeddings are Koszul. In characteristic zero, these results are deduced from the positive characteristic case using general semicontinuity theorems. See the recent book of Brion and Kumar [2005] for a unified exposition of these results, along with further details, references, and applications. On toric varieties, the Frobenius endomorphisms lift to endomorphisms over \mathbb{Z} , and it seems easiest and most natural to work independently of the characteristic using these lifted endomorphisms. One feature of this approach is that we can prove results about section rings of toric varieties over \mathbb{Z} , or an arbitrary field, by producing a splitting of the diagonal in $X \times X$ for a single q .

We begin by considering the structure of $F_*\mathbb{O}_X$ as an \mathbb{O}_X -module. As a sheaf of groups, $F_*\mathbb{O}_X$ evaluated on the invariant affine open set U_σ associated to a cone $\sigma \in \Sigma$ is the coordinate ring $\mathbb{Z}[U_\sigma]$, which is usually identified with the semigroup ring $\mathbb{Z}[\sigma^\vee \cap M]$. However, the module structure on $F_*\mathbb{Z}[U_\sigma]$ is different from the

action of $\mathbb{Z}[U_\sigma]$ on itself. For this reason, we identify $F_*\mathbb{Z}[U_\sigma]$ with the semigroup ring of fractional lattice points

$$F_*\mathbb{Z}[U_\sigma] = \mathbb{Z}[\sigma^\vee \cap \frac{1}{q}M],$$

taking a monomial $x^u \in \mathbb{Z}[U_\sigma]$ to $x^{u/q}$. The action of $\mathbb{Z}[U_\sigma]$ on $F_*\mathbb{Z}[U_\sigma]$ is then induced by the natural action of M on $\frac{1}{q}M$, so

$$x^u \cdot x^{u'} = x^{u+u'},$$

for $u \in M$ and $u' \in \frac{1}{q}M$. If Y is a toric variety embedded equivariantly in X , then a splitting π is compatible with Y if and only if the induced map

$$\mathbb{Z}[U_\sigma] \xrightarrow{\sim} \mathbb{Z}[\sigma^\vee \cap \frac{1}{q}M] \xrightarrow{\pi} \mathbb{Z}[U_\sigma]$$

maps $I_Y(U_\sigma)$ into $I_Y(U_\sigma)$ for every $\sigma \in \Sigma$.

We now summarize some basic properties of compatible splittings and their applications to section rings of ample line bundles. Let X be a complete toric variety, and let L be a line bundle on X . A splitting π of X makes \mathbb{O}_X a direct summand of $F_*\mathbb{O}_X$ and hence L a direct summand of $L \otimes F_*\mathbb{O}_X$. By the projection formula, $L \otimes F_*\mathbb{O}_X$ is isomorphic to $F_*(F^*L)$, and we claim that F^*L is isomorphic to L^q . To see this, note that there is a T -invariant Cartier divisor D such that L is isomorphic to $\mathbb{O}(D)$ [Fulton 1993, Section 3.4]. The restriction of D to an invariant affine open U_σ is the divisor of a rational function x^u for some $u \in M$, and hence the restriction of F^*D to U_σ is the divisor of x^{qu} . It follows that F^*L is isomorphic to L^q , as claimed. Now, since cohomology commutes with direct sums, π induces a split injection

$$H^i(X, L) \hookrightarrow H^i(X, L^q),$$

for every i . Iterating this argument gives split injections of $H^i(X, L)$ in $H^i(X, L^{q^r})$ for all positive integers r . In particular, if $H^i(X, L^{q^r})$ vanishes for some r , as is the case when L is ample, then $H^i(X, L)$ vanishes as well.

The proofs of the following five propositions are essentially identical to the standard proofs of the analogous results for Frobenius splittings, and are omitted. See [Brion and Kumar 2005]; Proposition 1.2.1, Theorem 1.2.8, and Exercises 1.5.E.1, 1.5.E.2, and 1.5.E.3, respectively, for the case where the line bundles in question are ample. The extensions to nef bundles can be deduced following the arguments in [Inamdar 1994], using the fact that any nef line bundle on X is the pullback of an ample line bundle on some toric variety X' under a proper birational toric morphism $f : X \rightarrow X'$.

Proposition 2.1. *Let Y and Y' be toric varieties equivariantly embedded in X . If $Y \cup Y'$ is split compatibly in X then Y, Y' , and $Y \cap Y'$ are split compatibly in X .*

Proposition 2.2. *Let Y be a compatibly split subvariety of X . If L is a nef line bundle on X then the restriction map*

$$H^0(X, L) \rightarrow H^0(Y, L)$$

is surjective and $H^1(X, I_Y \otimes L) = 0$.

Proposition 2.3. *Let L_1, \dots, L_r be nef line bundles on X . If the diagonal is compatibly split in $X \times X$ then the section ring $R(L_1, \dots, L_r)$ is normally generated.*

Proposition 2.4. *Let L_1, \dots, L_r be nef line bundles on X . If the union of $\Delta \times X$ and $X \times \Delta$ is compatibly split in $X \times X \times X$ then the section ring $R(L_1, \dots, L_r)$ is normally presented.*

Proposition 2.5. *Let L_1, \dots, L_r be nef line bundles on X . If $\Delta_1 \cup \dots \cup \Delta_{n-1}$ is compatibly split in X^n for every n then the section ring $R(L_1, \dots, L_r)$ is normally presented and Koszul.*

3. Canonical splittings

Every toric variety has a splitting, and among all splittings of X there is a unique one that extends to every toric compactification $X' \supset X$ and lifts to every proper birational toric modification $X'' \rightarrow X'$ of such a compactification. If q is prime and k is the field with q elements, then the restriction of this splitting to X_k is the unique Frobenius splitting that is canonical in the sense of Mathieu [Brion and Kumar 2005, Chapter 4]. We now describe this *canonical splitting*, starting with its restriction to the dense torus T .

Let π_0 be the map of $\mathbb{Z}[T]$ -modules from $F_*\mathbb{Z}[T]$ to $\mathbb{Z}[T]$ given by

$$\pi_0(x^u) = \begin{cases} x^u & \text{if } u \in M, \\ 0 & \text{otherwise.} \end{cases}$$

The pullback map $F^* : \mathbb{Z}[T] \rightarrow F_*\mathbb{Z}[T]$ is induced by the inclusion of M in $\frac{1}{q}M$; if q is prime and k is the field with q elements, then the induced map $k[T] \rightarrow F_*k[T]$ may be identified with the inclusion of $k[T]$ in $k[T]^{1/q}$.

In particular, $\pi_0 \circ F_*$ is the identity, and hence gives a splitting of T .

Proposition 3.1. *For any toric variety X , π_0 extends to a splitting of X .*

Proof. The composition $\pi_0 \circ F^*$ is the identity, and for each affine open U_σ , π_0 maps $\mathbb{Z}[\sigma^\vee \cap \frac{1}{q}M]$ into $\mathbb{Z}[\sigma^\vee \cap M]$. \square

Properties of this canonical splitting π_0 are closely related to Smith's proof [2000, Proposition 6.3] that toric varieties are globally F -regular.

Proposition 3.2. *The canonical splitting π_0 is compatible with every T -invariant subvariety.*

Proof. First we claim that π_0 is compatible with the union of the T -invariant divisors. To see this, note that the ideal sheaf I of the union of the invariant divisors is given by

$$I(U_\sigma) = \mathbb{Z}[\text{int}(\sigma^\vee) \cap M],$$

where $\text{int}(\sigma^\vee)$ is the interior of σ^\vee . If u is a fractional lattice point in the interior of σ^\vee then $\pi_0(x^u)$ is either zero or x^u , and so is contained in $I(U_\sigma)$, which proves the claim. The proposition then follows from [Proposition 2.1](#), since every T -invariant subvariety is an intersection of invariant divisors. \square

However, if X is positive dimensional then the canonical splitting π_0 of $X \times X$ is not compatible with the diagonal Δ . To see this, observe that if $u \in \frac{1}{q}M$ is not in M , then $1 - x^u \otimes x^{-u}$ is in F_*I_Δ , but

$$\pi_0(1 - x^u \otimes x^{-u}) = 1,$$

which is not in I_Δ . To apply the standard techniques relating splittings to section rings of ample line bundles discussed in [Section 2](#), we must look for other splittings of $X \times X$, and X^r for r greater than two, that are compatible with the diagonal and the union of the large semidiagonals, respectively.

4. Splittings of diagonals

We now describe the space of all splittings of a toric variety and use this description to characterize diagonally split toric varieties. First, it is helpful to consider the structure of $F_*\mathbb{O}_X$ as an \mathbb{O}_X -module in more detail.

Recall that an equivariant structure, or T -linearization, on a coherent sheaf \mathcal{F} on X is an isomorphism of sheaves on $T \times X$,

$$\varphi : \mu^*\mathcal{F} \rightarrow p^*\mathcal{F},$$

where $\mu : T \times X \rightarrow X$ is the torus action and p is the second projection, that satisfies the usual cocycle condition [[Brion and Kumar 2005](#), Section 2.1]. For example, the natural equivariant structure on \mathbb{O}_X is given by $1 \otimes x^u \mapsto x^{-u} \otimes x^u$. In general, the push forward of an equivariant sheaf under an equivariant morphism does not carry a natural equivariant structure. However, the equivariant endomorphism F has the property that $F_*\mathbb{O}_X$ is *equivariantizable* [[Bøgvad 1998](#); [Thomsen 2000](#)]; it is possible to choose an equivariant structure as follows. First, choose representatives u_1, \dots, u_s of the cosets in $\frac{1}{q}M/M$. Let φ be the map from $\mu^*F_*\mathbb{O}_X$ to $p^*F_*\mathbb{O}_X$ that takes $1 \otimes x^u$ to $x^{u_i - u} \otimes x^u$, for u in the coset $u_i + M$. It is straightforward to check that φ is an isomorphism and gives an equivariant structure on $F_*\mathbb{O}_X$, as required.

A splitting of X restricts to a splitting of T , and two splittings of X agree if and only if they agree on T , so we describe the space of all splittings of X in terms of

splittings of T that extend to X , as follows. For fractional lattice points $a \in \frac{1}{q}M$, let

$$\pi_a : F_*\mathbb{Z}[T] \rightarrow \mathbb{Z}[T]$$

be the map given by

$$\pi_a(x^u) = \begin{cases} x^{a+u} & \text{if } a+u \text{ is in } M, \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 4.1. *The set of maps π_a , for a in $\frac{1}{q}M$, is a \mathbb{Z} -basis for*

$$\text{Hom}(F_*\mathbb{Z}[T], \mathbb{Z}[T]).$$

Proof. The maps π_a are independent, and the free generators x^{u_1}, \dots, x^{u_s} for $F_*\mathbb{Z}[T]$ can be sent to an arbitrary s -tuple of elements of $\mathbb{Z}[T]$ by a suitable linear combination of the maps π_a . \square

If we choose an equivariant structure for $\text{Hom}(F_*\mathbb{Z}[T], \mathbb{Z}[T])$, as above, then the maps π_a form a T -eigenbasis. Therefore, a rational section

$$\pi = c_1\pi_{a_1} + \dots + c_r\pi_{a_r}$$

of the sheaf $\mathcal{H}om(F_*\mathcal{O}_X, \mathcal{O}_X)$, with each c_i nonzero, extends to X if and only if each π_{a_i} is regular on X . For a ray, or one-dimensional cone ρ in Σ , we write v_ρ for the primitive generator of ρ .

Proposition 4.2. *Let U_σ be an affine toric variety. Then π_a is regular on U_σ if and only if $\langle a, v_\rho \rangle$ is greater than minus one for each ray ρ in σ .*

Proof. The map π_a is regular on U_σ if and only if it takes

$$\mathbb{Z}[\sigma^\vee \cap \frac{1}{q}M]$$

into $\mathbb{Z}[\sigma^\vee \cap M]$. Suppose $\langle a, v_\rho \rangle$ is greater than minus one for each ray ρ in σ and u is in $\sigma^\vee \cap \frac{1}{q}M$. Either $\pi_a(x^u)$ is zero or $a+u$ is in M and $\langle u, v_\rho \rangle$ is a nonnegative integer for all rays ρ in σ , and hence $a+u$ is in σ^\vee . Therefore π_a extends to U_σ . Conversely, if $\langle a, v_\rho \rangle$ is less than or equal to minus one for some ρ , then it is straightforward to produce points $u \in \sigma^\vee$ such that $a+u$ is in M , but not in σ^\vee . In this case, π_a is not regular on U_σ . \square

We follow the usual toric convention fixing $K = -\sum D_\rho$, the sum of the prime T -invariant divisors each with multiplicity minus one, as a convenient representative of the canonical class. The polytope associated to a divisor $D = \sum d_\rho D_\rho$ is

$$P_D = \{u \in M_{\mathbb{R}} \mid \langle u, v_\rho \rangle \geq -d_\rho \text{ for all } \rho\}.$$

In particular, the polytope P_{-K} associated to the anticanonical divisor is

$$P_{-K} = \{u \in M_{\mathbb{R}} \mid \langle u, v_\rho \rangle \geq -1 \text{ for all } \rho\}.$$

The interior of the polytope P_{-K} controls the space of \mathbb{C}_X -module maps from $F_*\mathbb{C}_X$ to \mathbb{C}_X as follows.

Proposition 4.3. *The set of maps π_a for fractional lattice points a in the interior of P_{-K} is a basis for $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$.*

Proof. If a is not in the interior of P_{-K} then $\langle a, v_\rho \rangle$ is less than or equal to minus one for some ray $\rho \in \Sigma$ and then π_a is not regular on U_ρ . Conversely, if a is in the interior of P_{-K} , then π_a extends to every invariant affine open subvariety of X , by [Proposition 4.2](#), and therefore is regular on X . \square

Remark 4.4. When X is smooth, [Proposition 4.3](#) corresponds to the natural identification between $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$ and $H^0(X, K_X^{1-q})$ given by duality for finite flat morphisms [[Brion and Kumar 2005](#), Section 1.3].

Proposition 4.5. *A map $\sum c_a \pi_a$ in $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$ is a splitting if and only if $c_0 = 1$.*

Proof. Zero is the only lattice point in the interior of $P_{-K} \cap M$, so the image of x^u under $\pi = \sum c_a \pi_a$ is equal to $c_0 x^u$ for $u \in M$. In particular, since F^* maps x^u to x^u in $F_*\mathbb{C}_X$, $\pi \circ F^*$ is the identity if and only if c_0 is equal to one. \square

The set of splittings of X is an affine hyperplane in $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$ by [Proposition 4.5](#). For any subvariety $Y \subset X$, the condition that $\pi(F_*I_Y)$ is contained in I_Y cuts out a linear subspace of $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$. So the set of splittings of X that are compatible with Y is an affine subspace of $\text{Hom}(F_*\mathbb{C}_X, \mathbb{C}_X)$, which may be empty. We now prove [Theorem 1.2](#), which gives a necessary and sufficient condition for the space of splittings of $X \times X$ that are compatible with the diagonal to be nonempty.

Proof of Theorem 1.2. Suppose

$$\pi = \sum c_{a,a'} \pi_{a,a'}$$

is a splitting of $X \times X$ that is compatible with the diagonal. Then the restriction of π to the dense torus is a splitting compatible with the diagonal in $T \times T$. For any $u \in \frac{1}{q}M$, we have

$$1 - x^u \otimes x^{-u}$$

in F_*I_Δ , where I_Δ is the ideal of the diagonal in $T \times T$. Since $\pi(1)$ is equal to one, the restriction of $\pi(x^u \otimes x^{-u})$ to the diagonal must also be equal to one. Now the restriction of $\pi(x^u \otimes x^{-u})$ to the diagonal is a Laurent polynomial in $\mathbb{Z}[T]$ whose constant term is

$$\sum_{a \in [u]} c_{-a,a},$$

where $[u]$ is the coset of u in $\frac{1}{q}M/M$. Since the polytope associated to $-K_{X \times X}$ is $P_{-K_X} \times P_{-K_X}$, there must be a representative a of $[u]$ such that both a and $-a$ are

contained in the interior of P_{-K_X} , which means that a is contained in the interior of the diagonal splitting polytope

$$\mathbb{F}_X = P_{-K_X} \cap -P_{-K_X}.$$

For the converse, suppose that every nonzero equivalence class $[u_i]$ in $\frac{1}{q}M/M$ has a representative a_i in the interior of \mathbb{F}_X . Then

$$\pi_\Delta = \pi_0 + \sum_{i=1}^s \pi_{a_i, -a_i}$$

is a splitting, and we claim that π_Δ is compatible with the diagonal. To see this, note that the ideal of the diagonal in $U_\sigma \times U_\sigma$ is generated by the Laurent polynomials $1 - x^u \otimes x^{-u}$ for u in $\sigma^\vee \cap M$. Then F_*I_Δ is generated as a $\mathbb{Z}[U_\sigma \times U_\sigma]$ -module by the

$$x^b - x^b \cdot (x^u \otimes x^{-u}),$$

as $b = (b_1, b_2)$ ranges over $\frac{1}{q}(M \times M)$ and u ranges over $\frac{1}{q}M$. Now the restriction of $\pi_\Delta(x^b)$ to the diagonal is $x^{b_1+b_2}$ if $b_1 + b_2$ is in M and zero otherwise. In particular, the restriction of $\pi_\Delta(x^b - x^b \cdot (x^u \otimes x^{-u}))$ to the diagonal vanishes, as required. \square

Proof of Theorem 1.3. Suppose the diagonal is compatibly split in $X \times X$. Then every nonzero equivalence class $[u_j]$ in $\frac{1}{q}M/M$ is represented by a fractional lattice point a_j in the interior of \mathbb{F}_X , by Theorem 1.2.

A splitting of X^n is compatible with the union $\Delta_1 \cup \dots \cup \Delta_{n-1}$ if it is compatible with each Δ_i . For $u \in M$, let $u^{(i)}$ denote the lattice point in M^n whose only nonzero coordinate is the i -th one, which is equal to u . The ideal of Δ_i is generated by the functions

$$1 - x^{u^{(i)}} \cdot x^{-u^{(i+1)}}.$$

We claim that the splitting

$$\pi = \pi_0 + \sum_{i=1}^{n-1} \sum_j \pi_{a_j^{(i)} - a_j^{(i+1)}}$$

is compatible with Δ_i for $1 \leq i < n$, and hence with the union $\Delta_1 \cup \dots \cup \Delta_{n-1}$. The proof of the claim is then similar to the proof of Theorem 1.2 above, and the theorem follows. \square

Proof of Theorem 1.4. Suppose the diagonal is compatibly split in $X \times X$. Then $\Delta_1 \cup \dots \cup \Delta_{n-1}$ is compatibly split in X^n for every n , by Theorem 1.3. Therefore, for any nef line bundles L_1, \dots, L_r on X the section ring $R(L_1, \dots, L_r)$ is normal and Koszul, by Proposition 2.5. \square

Remark 4.6. Pairs of opposite lattice points u and $-u$ in the polytopes associated to anticanonical divisors have also appeared in relation to the classification of smooth toric Fano varieties. In particular, Ewald [1988] conjectured twenty years ago that if X is a smooth toric Fano variety then \mathbb{F}_X contains a basis for the character lattice M . Ewald’s conjecture has been verified for smooth toric Fano varieties of dimension less than or equal to seven by Øbro [2007, Section 4.1]. However, it remains unknown in higher dimensions whether there exists a single nonzero lattice point u in \mathbb{F}_X [Kreuzer and Nill 2007, Section 4.6].

Acknowledgments

This work grew out of an expository talk at a mini-workshop on normality of lattice polytopes at Oberwolfach. I thank the organizers, C. Haase, T. Hibi, and D. Maclagan, as well as the other participants, for the chance to be part of such a stimulating event, and the MFO for their hospitality. While preparing this paper, I benefited from helpful discussions with D. Anderson, D. Eisenbud, N. Fakhruddin, B. Howard, and A. Paffenholz, and from the referee’s thoughtful comments. Finally, I am grateful to N. Lauritzen and J. Thomsen for pointing out that the methods in a first version of this paper can be used to give the example of the nondiagonally split Schubert variety presented here.

References

- [Anderson 2007] D. Anderson, “Chern class formulas for G_2 Schubert loci”, preprint, 2007. [arXiv 0712.2641](#)
- [Batyrev and Nill 2007] V. Batyrev and B. Nill, “Multiples of lattice polytopes without interior lattice points”, *Mosc. Math. J.* **7**:2 (2007), 195–207, 349. [MR 2008g:52018](#) [Zbl 1134.52020](#)
- [Batyrev and Nill 2008] V. Batyrev and B. Nill, “Combinatorial aspects of mirror symmetry”, pp. 35–66 in *Integer points in polyhedra—geometry, number theory, representation theory, algebra, optimization, statistics* (Snowbird, UT, 2006), edited by M. Beck et al., Contemp. Math. **452**, Amer. Math. Soc., Providence, RI, 2008. [MR 2405763](#) [arXiv math/0703456](#)
- [Bezrukavnikov 1995] R. Bezrukavnikov, “Koszul property and Frobenius splitting of Schubert varieties”, preprint, 1995. [arXiv alg-geom/9502021](#)
- [Bøgvad 1995] R. Bøgvad, “On the homogeneous ideal of a projective nonsingular toric variety”, preprint, 1995. [arXiv alg-geom/9501012](#)
- [Bøgvad 1998] R. Bøgvad, “Splitting of the direct image of sheaves under the Frobenius”, *Proc. Amer. Math. Soc.* **126**:12 (1998), 3447–3454. [MR 99h:14051](#) [Zbl 0902.14038](#)
- [Brion and Kumar 2005] M. Brion and S. Kumar, *Frobenius splitting methods in geometry and representation theory*, Progress in Mathematics **231**, Birkhäuser, Boston, 2005. [MR 2005k:14104](#) [Zbl 1072.14066](#)
- [Brylinski and Zhang 2003] J.-L. Brylinski and B. Zhang, “Equivariant Todd classes for toric varieties”, preprint, 2003. [arXiv math/0311318](#)
- [Buch et al. 1997] A. Buch, J. F. Thomsen, N. Lauritzen, and V. Mehta, “The Frobenius morphism on a toric variety”, *Tohoku Math. J. (2)* **49**:3 (1997), 355–366. [MR 98j:14020](#) [Zbl 0899.14026](#)

- [Ewald 1988] G. Ewald, “On the classification of toric Fano varieties”, *Discrete Comput. Geom.* **3**:1 (1988), 49–54. MR 88m:14037 Zbl 0686.14042
- [Fujino 2007] O. Fujino, “Multiplication maps and vanishing theorems for toric varieties”, *Math. Z.* **257**:3 (2007), 631–641. MR 2008g:14092 Zbl 1129.14029
- [Fulton 1993] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies **131**, Princeton University Press, Princeton, NJ, 1993. MR 94g:14028 Zbl 0813.14039
- [Haase et al. 2008] C. Haase, B. Nill, and S. Payne, “Cayley decompositions of lattice polytopes and upper bounds for h^* -polynomials”, [arXiv 0804.3667](https://arxiv.org/abs/0804.3667)
- [Inamdar 1994] S. P. Inamdar, “A note on Frobenius splitting of Schubert varieties and linear syzygies”, *Amer. J. Math.* **116**:6 (1994), 1587–1590. MR 96a:14055 Zbl 0819.14020
- [Inamdar and Mehta 1994] S. P. Inamdar and V. B. Mehta, “Frobenius splitting of Schubert varieties and linear syzygies”, *Amer. J. Math.* **116**:6 (1994), 1569–1586. MR 96a:14054 Zbl 0817.14032
- [Kempf 1976] G. R. Kempf, “Linear systems on homogeneous spaces”, *Ann. of Math. (2)* **103**:3 (1976), 557–591. MR 53 #13229 Zbl 0327.14016
- [Kreuzer and Nill 2007] M. Kreuzer and B. Nill, “Classification of toric Fano 5-folds”, preprint, 2007. [arXiv math/0702890](https://arxiv.org/abs/math/0702890)
- [Mehta and Ramanathan 1985] V. B. Mehta and A. Ramanathan, “Frobenius splitting and cohomology vanishing for Schubert varieties”, *Ann. of Math. (2)* **122**:1 (1985), 27–40. MR 86k:14038 Zbl 0601.14043
- [Øbro 2007] M. Øbro, *Classification of smooth Fano polytopes*, PhD Thesis, University of Århus, Århus, Denmark, 2007.
- [Oda 1997] T. Oda, “Problems on Minkowski sums of convex lattice polytopes”, preprint, 1997. [arXiv 0812.1418](https://arxiv.org/abs/0812.1418)
- [Polishchuk and Positselski 2005] A. Polishchuk and L. Positselski, *Quadratic algebras*, University Lecture Series **37**, American Mathematical Society, Providence, RI, 2005. MR 2006f:16043 Zbl 1145.16009
- [Ramanan and Ramanathan 1985] S. Ramanan and A. Ramanathan, “Projective normality of flag varieties and Schubert varieties”, *Invent. Math.* **79**:2 (1985), 217–224. MR 86j:14051 Zbl 0553.14023
- [Ramanathan 1987] A. Ramanathan, “Equations defining Schubert varieties and Frobenius splitting of diagonals”, *Inst. Hautes Études Sci. Publ. Math.* **65** (1987), 61–90. MR 88k:14032 Zbl 0634.14035
- [Smith 2000] K. E. Smith, “Globally F -regular varieties: applications to vanishing theorems for quotients of Fano varieties”, *Michigan Math. J.* **48** (2000), 553–572. MR 2001k:13007 Zbl 0994.14012
- [Sturmfels 1996] B. Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series **8**, American Mathematical Society, Providence, RI, 1996. MR 97b:13034 Zbl 0856.13020
- [Thomsen 2000] J. F. Thomsen, “Frobenius direct images of line bundles on toric varieties”, *J. Algebra* **226**:2 (2000), 865–874. MR 2001f:14095 Zbl 0957.14036
- [Totaro 199?] B. Totaro, “Chow groups, Chow cohomology, and linear varieties”, preprint, 199?, Available at <http://www.dpmms.cam.ac.uk/~bt219/linear.dvi>. To appear in *J. Algebraic Geom.*

Communicated by Vasudevan Srinivas

Received 2008-05-08

Revised 2008-10-15

Accepted 2008-11-22

spayne@stanford.edu

Department of Mathematics, Building 380, Sloan Hall,
Stanford University, Stanford, CA 94305, United States