

Algebra & Number Theory

Volume 3

2009

No. 4

Syzygies of the secant variety of a curve

Jessica Sidman and Peter Vermeire

 mathematical sciences publishers

Syzygies of the secant variety of a curve

Jessica Sidman and Peter Vermeire

We show the secant variety of a linearly normal smooth curve of degree at least $2g + 3$ is arithmetically Cohen–Macaulay, and we use this information to study the graded Betti numbers of the secant variety.

1. Introduction

We work throughout over an algebraically closed field of characteristic 0. A well-known result dating back to Castelnuovo states that if $C \subset \mathbb{P}^n$ is a linearly normal curve of genus g with $\deg C \geq 2g + 1$, then C is projectively normal and hence is arithmetically Cohen–Macaulay (ACM). Our main result is this:

Theorem 1. *If $C \subset \mathbb{P}^n$ is a smooth linearly normal curve of genus g and degree $d \geq 2g + 3$, then its secant variety Σ is ACM.*

Using the Auslander–Buchsbaum theorem [Eisenbud 1995, §19], this tells us that a minimal free resolution of the coordinate ring of Σ , S_Σ , has length equal to $\text{codim } \Sigma$, and the remainder of this paper is devoted to studying the syzygies among the defining equations of Σ .

To describe our results on syzygies more precisely, we set up some notation. Let $S = k[x_0, \dots, x_n]$. Any finitely generated S -module M has a minimal free resolution

$$0 \rightarrow \bigoplus S(-j)^{\beta_{r,j}} \rightarrow \dots \rightarrow \bigoplus S(-j)^{\beta_{1,j}} \rightarrow \bigoplus S(-j)^{\beta_{0,j}} \rightarrow M \rightarrow 0,$$

where the *graded Betti numbers* $\beta_{i,j}$ are uniquely determined by minimality. It is convenient to display the $\beta_{i,j}$ in a graded Betti diagram in which the (i, j) entry is $\beta_{i,i+j}$.

	0	1	2	3
0	$\beta_{0,0}$	$\beta_{1,1}$	\cdots	
1	$\beta_{0,1}$	$\beta_{1,2}$	\cdots	
2	$\beta_{0,2}$	$\beta_{1,3}$	\cdots	

MSC2000: primary 13D02; secondary 14N05, 14H99, 14F05.

Keywords: syzygies, secant varieties, projective curves, graded Betti numbers.

Sidman is partially supported by NSF grant DMS 0600471 and the Clare Boothe Luce Program.

As in [Eisenbud 2005] we say that the Betti numbers $\beta_{i,i+k}$ in the i th row of the Betti diagram form the degree $k + 1$ linear strand if $M = S/I$ for some homogeneous ideal I . In this case, $\beta_{1,k+1}$ is the number of minimal generators of I in degree $k + 1$. (For an arbitrary module, M , it might make more sense to call this the degree k linear strand.)

It is useful to recast several notions from the geometric literature in terms of the graded Betti diagram. Suppose that a variety $X \subset \mathbb{P}^n$ is projectively normal with an ideal generated by quadrics. Then for $p \geq 1$ it satisfies Green’s condition N_p [Green 1984] if for all $i \leq p$, $\beta_{ij} = 0$ unless $j = i + 1$. Eisenbud et al. [2005] extended this notion to a variety X with ideal generated in degree k , so that X satisfies $N_{k,p}$ if for all $i \leq p$, $\beta_{ij} = 0$ unless $j = i + k - 1$. Thus, the only nonzero entries in columns one through p of a Betti diagram of a variety satisfying $N_{k,p}$ are in row $k - 1$. The Castelnuovo–Mumford regularity [Mumford 1966], or simply *regularity*, of a module can also be defined in terms of graded Betti numbers. A module is m -regular if $\beta_{i,m+i} = 0$ for all $i > 0$, which is equivalent to stating that its Betti diagram is zero in all rows greater than m .

If $C \subset \mathbb{P}^n$ is a linearly normal curve of genus g and degree $d \geq 2g + 3$, we obtain several results as consequences of the Cohen–Macaulay condition. In Corollary 3.9 we show that if $\text{reg } I_\Sigma < 5$, then C is rational and $\text{reg } I_\Sigma = 3$. We give explicit formulas for several graded Betti numbers in Corollary 4.1 and Proposition 4.4, showing that

- $\beta_{1,3} = \binom{n+1}{3} - (d - 2)n - 3g + 1$.
- $\beta_{2,4} = \beta_{1,4} + \beta_{1,3}(n + 1) - \binom{n+4}{n} + P_\Sigma(4)$.
- $\beta_{n-3,n+1} = \binom{g+1}{2}$.

Note that via Theorem 1 there are exactly $n - 3$ syzygy modules in the resolution of S_Σ , and if $g \geq 1$, then Corollary 3.9 implies that the final syzygy module is generated by elements of degree $\leq n + 1$. Thus, $\beta_{n-3,n+1}$ is the bottom right corner of the graded Betti diagram, and it depends only on the genus of the curve.

We compute the Hilbert polynomial of S_Σ by relating it to the Hilbert polynomial of a curve of degree D and genus G gotten by intersecting Σ with a plane of codimension 2.

Theorem 1.1. *The Hilbert polynomial of S_Σ agrees with its Hilbert function for all positive integers and is given by*

$$D \binom{m+2}{3} + (1 - G) \binom{m+1}{2} + \alpha_1 m + \alpha_0,$$

where $\alpha_1 = \binom{n+2}{2} - (n+1) - 3D - 2(1 - G)$ and $\alpha_0 = -\binom{n+2}{2} + 2(n+1) + 2D + 1 - G$.

We also obtain a nonvanishing result on the graded Betti numbers of higher secant varieties.

Theorem 1.2. *Let C be a smooth curve of genus g embedded into \mathbb{P}^n via a line bundle L of degree $d \geq 2g + 2k + p + 1$ and Σ_k be its variety of secant k -planes. Suppose that $L = L_1 \otimes L_2$ where $|L_1| = s \leq |L_2| = t$. If $s + 1 \geq k + 2$, then the length of the degree $k + 2$ linear strand of S_{Σ_k} is at least $s + t - 2k - 1$. In particular, if L is a general line bundle of degree $d \geq 2g + 2k + p + 1$, then $\beta_{s+t-2k-1, s+t-k}(\Sigma_k) \neq 0$.*

We briefly sketch part of the picture of what is known about syzygies of high degree curves to put our results in context. The homogeneous coordinate ring of a curve of degree at least $2g + 1$ is 1-regular if $g = 0$ and has regularity two otherwise. From [Green 1984; Green and Lazarsfeld 1985; 1988] we know that if $d \geq 2g + 1 + p$, then the curve satisfies N_p . By a result of Schreyer [Eisenbud 2005, Theorem 8.17], we know that $\beta_{p+\lfloor g/2 \rfloor, p+\lfloor g/2 \rfloor+1}$ is nonzero. Furthermore, as a consequence of duality, the “last” graded Betti number is $\beta_{n-1, n+1} = g$. (See [Eisenbud 2005, Chapter 8] for a nice discussion.)

Based on what we have seen we extend and refine the conjectures in [Vermeire 2008b] as follows:

Conjecture 1.3. Suppose that $C \subset \mathbb{P}^n$ is a smooth linearly normal curve of genus g and degree $d \geq 2g + 2k + 1 + p$, where $p, k \geq 0$. Then

- (1) Σ_k is ACM and has regularity $(2k + 2)$ unless $g = 0$, in which case the regularity is $k + 1$.
- (2) $\beta_{n-2k-1, n+1} = \binom{g+k}{k+1}$.
- (3) Σ_k satisfies $N_{k+2, p}$.

As described above, the full conjecture is known to hold for $k = 0$. Further, by [Graf von Bothmer and Hulek 2004] and [Fisher 2006] it holds for $g \leq 1$. In this work, we show that parts (1) and (2) hold for $k = 1$. After the completion of this work, progress on part (3) was made for $k = 1$ [Vermeire 2008a]. We illustrate the behavior that we have seen with the example below.

Example 1.4. At the suggestion of D. Eisenbud we used ideas of F. Schreyer to compute the ideal of a genus 2 curve embedded in \mathbb{P}^7 . Let \bar{C} be a plane curve of degree 5 with 4 nodes. If we blow up the four nodes in \mathbb{P}^2 and consider the linear system $|5H - 2\Sigma E_i|$, where H is the proper transform of a hyperplane and the E_i are the exceptional divisors of the blow-up, the restriction of this system to the proper transform C of \bar{C} has degree $9 = 2g + 5$, and embeds $C \subset \mathbb{P}^7$ as a smooth curve of genus 2. Using Macaulay 2 [Grayson and Stillman] we can write down a basis of $|5H - 2\Sigma E_i|$ over the rationals. We display the Betti diagram of the

coordinate ring, where “—” denotes a zero entry.

	0	1	2	3	4	5	6
0	1	—	—	—	—	—	—
1	—	19	58	75	44	5	—
2	—	—	—	—	—	6	2

Note that the quadratic strand of the resolution has length 5 but that the curve satisfies N_4 but not N_5 .

Using code developed for [Sidman and Sullivant 2006], we computed the ideal of Σ . From the Betti diagram we see that the cubic strand of the resolution has length 2 and that $\beta_{4,8} = 3$ as predicted by Conjecture 1.3.

	0	1	2	3	4
0	1	—	—	—	—
1	—	—	—	—	—
2	—	12	16	—	—
3	—	—	—	4	—
4	—	—	—	4	3

Comparing the diagram to the statement of Corollary 4.7, we see that the three unknowns at the tail of the resolution are all zero here as in Example 4.8. \square

We give a brief outline of the structure of the paper. The ACM condition is treated in §3. To understand the ACM condition, we work geometrically to show that cohomology groups vanish. The key observation is that there is a desingularization $\tilde{\Sigma} \rightarrow \Sigma$ such that $\tilde{\Sigma}$ is a \mathbb{P}^1 -bundle over the symmetric square of C , which we denote by S^2C , and hence the cohomology of the structure sheaf of $\tilde{\Sigma}$ is the same as that of S^2C , which is easier to understand. As Σ has nonrational singularities, the higher direct image sheaves of the ideal of $\tilde{\Sigma}$ do not vanish, but there is another divisor whose ideal sheaf has the same direct image and whose higher direct images do vanish. (See Lemma 2.4). Making the exact relationships between these objects precise is the bulk of our work. The technical preliminaries are summarized in Section 2. We examine the graded Betti diagram of S_Σ in Section 4.

To improve readability we have written out some arguments which are surely well-known to experts, but are perhaps not easily available in the standard references.

2. Setup and notation

Suppose that $X \subset \mathbb{P}^n$ is a variety. We let \mathcal{O}_X and \mathcal{F}_X denote the structure sheaf and ideal sheaf of X . The homogeneous coordinate ring of \mathbb{P}^n is $S = k[x_0, \dots, x_n]$. We

let $I_X = \bigoplus H^0(\mathbb{P}^n, \mathcal{I}_X(d))$ and $S_X = S/I_X$. We let H denote a general hyperplane in \mathbb{P}^n and its pullback under a morphism. We write $\mathcal{O}(k)$ for $\mathcal{O}(kH)$ when no confusion will arise. We may write $H^i(\mathcal{F})$ for $H^i(X, \mathcal{F})$ and $h^i(\mathcal{F})$ for $h^i(X, \mathcal{F})$ if the meaning is clear.

Let C be a smooth curve of genus g . Throughout, L is a very ample line bundle on C embedding it as a linearly normal curve in $\mathbb{P}^n = \mathbb{P}(H^0(C, L))$ with degree $d = \deg L$.

A line bundle L on a smooth curve C is said to *separate k points* if

$$h^0(C, L(-Z)) = h^0(C, L) - k \quad \text{for all } Z \in S^k C,$$

where $S^k C$ is the k th symmetric product of C . We let Σ_k denote the variety of $(k + 1)$ -secant k -planes to C and write Σ for the variety Σ_1 .

We recall the first stages of a construction of Aaron Bertram which provides the geometric framework for our results.

Theorem 2.1 [Bertram 1992, Theorem 1]. *Suppose that L separates 4 points. Let $g : B_1 \rightarrow B_0 = \mathbb{P}^n$ be the blowup of B_0 along C with $\tilde{\Sigma}$ the proper transform of Σ . Let $h : B_2 \rightarrow B_1$ be the blowup of B_1 along $\tilde{\Sigma}$ and E_i be the proper transform in B_i of each exceptional divisor. We further let $f = g \circ h$.*

Then $\tilde{\Sigma} \subset B_1$ is smooth and irreducible, and transverse to E_1 , so in particular B_2 is smooth. Moreover, by Terracini recursiveness, if $x \in \Sigma \setminus C$, then $f^{-1}(x) \cong \mathbb{P}(H^0(C, L(-2V)))$, where V is the unique divisor of degree 2 whose span contains x . If $x \in C$, then $f^{-1}(x)$ is isomorphic to the blowup of $\mathbb{P}(H^0(C, L(-2x)))$ along the image of C embedded by $L(-2x)$. \square

Remark 2.2. Bertram’s construction continues, blowing up the strict transform of each Σ_k successively, so that a fiber over a point of C of the composition is \mathbb{P}^{n-2} in which we have blown up copies of Σ_i for $i = 0, \dots, k - 1$ and the degree of $\Sigma_0 = C$ is two less than the degree of the original embedding. We will abuse notation in the hopes of highlighting the recursive nature of the construction and denote the restriction of E_i to a fiber F of the composition using the notation of our setup relative to the blowing up that has occurred within F . For example, if $x \in C$ and $F = f^{-1}(x)$, we will write $\mathcal{O}_{B_2}(E_2)|_F = \mathcal{O}_F(E_1)$, keeping in mind that “ $E_1 \subset F$ ” is the exceptional divisor of \mathbb{P}^{n-2} blown up at C where the degree has already dropped by two.

A key point in what follows is that $\tilde{\Sigma}$ is a resolution of singularities of Σ , and is a \mathbb{P}^1 -bundle over $S^2 C$ in a natural way. We summarize this relationship:

Lemma 2.3. *The variety $\tilde{\Sigma} \subset B_1$ is a resolution of singularities $g : \tilde{\Sigma} \rightarrow \Sigma$ with the following properties:*

- (1) $g_* \mathcal{O}_{\tilde{\Sigma}} = \mathcal{O}_{\Sigma}$.

- (2) $Z := E_1 \cap \widetilde{\Sigma} \cong C \times C$.
- (3) The restriction $g : C \times C \rightarrow C$ is projection onto one factor.
- (4) The restriction of the linear system $|2H - E_1|$ to $\widetilde{\Sigma}$ yields a morphism $\pi : \widetilde{\Sigma} \rightarrow S^2C$ realizing $\widetilde{\Sigma}$ as a \mathbb{P}^1 -bundle over S^2C . The restriction of this morphism to $Z \cong C \times C$ is the canonical double cover $d : C \times C \rightarrow S^2C$.
- (5) If we define δ by $d^*\mathbb{O}_{S^2C}(\frac{\delta}{2}) = \mathbb{O}_{C \times C}(\Delta)$, then $d_*\mathbb{O}_{C \times C} = \mathbb{O}_{S^2C} \oplus \mathbb{O}_{S^2C}(-\frac{\delta}{2})$.
- (6) If F is a fiber of the \mathbb{P}^1 -bundle $\pi : \widetilde{\Sigma} \rightarrow S^2C$, then

$$\mathbb{O}_F(aH - bE_1) = \mathbb{O}_{\mathbb{P}^1}(a - 2b).$$

Proof. The first is [Vermeire 2002, 3.2], the second and third are [Vermeire 2001, 3.7], the fourth is [Vermeire 2001, 3.8]. Part (5) follows from [Barth et al. 2004, V.22]. For (6), note that each fiber F is the proper transform of a secant line, hence the intersection with a hyperplane is 1, while the intersection with the exceptional divisor is 2 (since each secant or tangent line intersects C in a scheme of length two). □

Lemma 2.4. *With hypotheses and notation as above:*

- (1) $\Sigma \subset B_0$ is normal and is smooth away from C .
- (2) $f_*\mathbb{O}_{B_2} = \mathbb{O}_{B_0}$ and $R^j f_*\mathbb{O}_{B_2} = 0$ for $j \geq 1$.
- (3) $R^i f_*\mathbb{O}_{B_2}(-E_2) = \begin{cases} \mathcal{F}_\Sigma & i = 0, \\ H^1(C, \mathbb{O}_C) \otimes \mathbb{O}_C & i = 2, \\ 0 & i \neq 0, 2. \end{cases}$
- (4) $R^i g_*\mathbb{O}_{B_1}(-mE_1) = R^i h_*\mathbb{O}_{B_2}(-mE_2) = 0$ for $i > 0$ and $m \geq 0$.
- (5) $R^i g_*\mathcal{F}_{\widetilde{\Sigma}} = R^i f_*\mathbb{O}_{B_2}(-E_2)$.
- (6) $R^i f_*\mathbb{O}_{B_2}(-E_1 - E_2) = \mathcal{F}_{\Sigma/\mathbb{P}^n}$ for $i = 0$ and is zero otherwise.

Proof. The first two can be found in [Vermeire 2002, 3.2], while the third is [Vermeire 2008b, Proposition 9] and the fourth is [Lazarsfeld 2004, Lemma 4.3.16]. Part (5) follows immediately from (4) and a degenerate case of Grothendieck’s composition of functors spectral sequence [Grothendieck 1957].

For the sixth item, we compute sheaves $R^i f_*\mathbb{O}_{E_1}(-E_2)$ and use them to show the claim via

$$0 \rightarrow \mathbb{O}_{B_2}(-E_1 - E_2) \rightarrow \mathbb{O}_{B_2}(-E_2) \rightarrow \mathbb{O}_{E_1}(-E_2) \rightarrow 0.$$

Since $E_1 \rightarrow C$ is flat, the locally free sheaf $\mathbb{O}_{E_1}(-E_2)$ is also flat over C . Thus, we can compute higher direct images via cohomology along the fibers of f restricted to E_1 by [Hartshorne 1977, Corollary III.12.9]. By the Terracini recursive-ness portion of Theorem 2.1, if $x \in C$, a fiber $F = f^{-1}(x)$ is the blowup of C in

$\mathbb{P}H^0(C, L(-2x))$ and E_2 intersects F in the exceptional divisor E_1 of this blowup. As $H^i(F, \mathcal{O}_F(-E_1)) = H^i(\mathbb{P}(H^0(C, L(-2x))), \mathcal{F}_C)$, $H^i(F, \mathcal{O}_F(-E_1))$ vanishes for $i = 0, 1$, and $h^2(\mathbb{P}(H^0(C, L(-2x))), \mathcal{F}_C) = h^1(C, \mathcal{O}_C) = g$. We conclude that $R^i f_* \mathcal{O}_{E_1}(-E_2) = 0$ for $i = 0, 1$ and that for $i = 2$ it is locally free of rank g . Note that by part (5), $R^2 f_* \mathcal{O}_{B_2}(-E_2)$ is also locally free of rank g . Therefore, if the map between them is a surjection, it is an isomorphism.

To get the surjectivity above we show $R^3 f_* \mathcal{O}_{B_2}(-E_1 - E_2) = 0$ by looking at

$$0 \rightarrow \mathcal{O}_{B_2}(-E_1 - E_2) \rightarrow \mathcal{O}_{B_2}(-E_1) \rightarrow \mathcal{O}_{E_2}(-E_1) \rightarrow 0. \quad (1)$$

Applying h_* , the projection formula and the observation that $E_2 \rightarrow \tilde{\Sigma}$ is a projective bundle, we see that

$$0 \rightarrow \mathcal{F}_{\tilde{\Sigma}}(-E_1) \rightarrow \mathcal{O}_{B_1}(-E_1) \rightarrow \mathcal{O}_{\tilde{\Sigma}}(-E_1) \rightarrow 0$$

is exact and all higher direct images vanish. If we apply g_* we get

$$\rightarrow R^2 g_* \mathcal{O}_{\tilde{\Sigma}}(-E_1) \rightarrow R^3 g_* \mathcal{F}_{\tilde{\Sigma}}(-E_1) \rightarrow R^3 g_* \mathcal{O}_{B_1}(-E_1) \rightarrow,$$

where the left-hand term vanishes because $\tilde{\Sigma} \rightarrow \Sigma$ has fibers of dimension at most one, and the right-hand term vanishes by (4). \square

We will use [Lemma 2.5](#) to show that $H^1(\Sigma, \mathcal{O}_{\Sigma}(2)) = 0$ in [Theorem 3.3](#).

Lemma 2.5. *Let L be a very ample line bundle on a variety X with $H^i(X, L) = 0$ for $i > 0$, E a locally free sheaf on X . Let $\varphi : X \rightarrow \mathbb{P}^n = \mathbb{P}(H^0(X, L))$ be the induced morphism.*

$$(1) \quad H^i(X \times X, (L \boxtimes E) \otimes \mathcal{F}_{\Delta}) = H^i(X, \varphi^* \Omega_{\mathbb{P}^n}^1 \otimes L \otimes E).$$

$$(2) \quad H^i(X \times X, (L \boxtimes E) \otimes \mathcal{F}_{\Delta}^2) = H^i(X, N_{X/\mathbb{P}^n}^* \otimes L \otimes E).$$

Proof. Applying $(\pi_2)_*$ to the exact sequence

$$0 \rightarrow (L \boxtimes E) \otimes \mathcal{F}_{\Delta} \rightarrow L \boxtimes E \rightarrow (L \boxtimes E) \otimes \mathcal{O}_{\Delta} \rightarrow 0$$

yields a twist of the Euler sequence on X :

$$0 \rightarrow \varphi^* \Omega_{\mathbb{P}^n}^1 \otimes L \otimes E \rightarrow H^0(X, L) \otimes E \rightarrow L \otimes E \rightarrow 0$$

Note that the hypothesis $H^i(X, L) = 0$ and the fact that L is globally generated imply that all higher direct images vanish, and part (1) follows immediately.

As $\mathcal{O}_{\Delta} \otimes \mathcal{F}_{\Delta} = N_{\Delta}^* = \Omega_X^1$; applying $(\pi_2)_*$ to the exact sequence

$$0 \rightarrow (L \boxtimes E) \otimes \mathcal{F}_{\Delta}^2 \rightarrow (L \boxtimes E) \otimes \mathcal{F}_{\Delta} \rightarrow (L \boxtimes E) \otimes N_{\Delta}^* \rightarrow 0$$

yields a twist of the conormal sequence on X :

$$0 \rightarrow N_{X/\mathbb{P}^n}^* \otimes L \otimes E \rightarrow \varphi^* \Omega_{\mathbb{P}^n}^1 \otimes L \otimes E \rightarrow \Omega_X^1 \otimes L \otimes E \rightarrow 0$$

Note that the hypothesis $H^i(X, L) = 0$ and the fact that L is very ample imply that all higher direct images vanish, and part (2) follows similarly. \square

3. Σ is ACM

The main goal of this section is the proof of [Theorem 1](#). As a consequence of our work we get [Corollary 3.4](#) showing that Σ is projectively normal. We will work throughout with the following hypothesis.

Hypothesis 3.1. *Let $C \subset \mathbb{P}^n$ be a smooth linearly normal curve of genus g and degree $d \geq 2g + 3$.*

Using the Serre–Grothendieck correspondence between local and global cohomology, the depth of the maximal ideal on the homogeneous coordinate ring of $\Sigma \subset \mathbb{P}^n$ can be measured by vanishings of global cohomology groups. We see that Σ is ACM if and only if $H^i(\mathbb{P}^n, \mathcal{F}_\Sigma(k)) = 0$ for all k and for $0 < i \leq \dim \Sigma$ (for example, [[Eisenbud 1995](#), Example 18.16]). In light of [[Vermeire 2008b](#)] where it is shown that \mathcal{F}_Σ is 5-regular, in order to show that Σ is ACM we are left to show that $H^i(\Sigma, \mathcal{O}_\Sigma(k)) = 0$ for $i = 1, 2$ and all $k \leq 3 - i$. In what follows we handle the required cohomological vanishing cases individually.

3A. Vanishings for $k < 0$. The vanishings needed for $k < 0$ follow easily from Kawamata–Viehweg vanishing together with part (3) of [Lemma 2.4](#). We write the 5-term sequence associated to the Leray spectral sequence (applying [Theorem 2.1](#)) to the map $g : \tilde{\Sigma} \rightarrow \Sigma$ as it will be crucial in what follows (note that the first and fourth terms follow by part (1) of [Lemma 2.3](#)).

$$0 \rightarrow H^1(\Sigma, \mathcal{O}_\Sigma(k)) \rightarrow H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(k)) \rightarrow H^0(\Sigma, R^1 g_* \mathcal{O}_{\tilde{\Sigma}}(k)) \\ \rightarrow H^2(\Sigma, \mathcal{O}_\Sigma(k)) \rightarrow H^2(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(k)) \quad (2)$$

Theorem 3.2. *If C satisfies [Hypothesis 3.1](#), then $H^i(\Sigma, \mathcal{O}_\Sigma(k)) = 0$ for $k < 0$ and $i = 1, 2$.*

Proof. We know that $g^* \mathcal{O}_\Sigma(1) = \mathcal{O}_{\tilde{\Sigma}}(1)$ is big and nef on $\tilde{\Sigma}$; hence $H^i(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(k))$ is 0 for $k < 0$ and $i < 3$ by Kawamata–Viehweg vanishing. Using the sequence (2), we have the claimed vanishing for $i = 1$ immediately. As $R^1 g_* \mathcal{O}_{\tilde{\Sigma}} \cong H^1(C, \mathcal{O}_C) \otimes \mathcal{O}_C$ by [Lemma 2.4](#) (3–5), we have $H^0(\Sigma, R^1 g_* \mathcal{O}_{\tilde{\Sigma}}(k)) = H^1(C, \mathcal{O}_C) \otimes H^0(C, \mathcal{O}_C(k)) = 0$, and the vanishing for $i = 2$ also follows. \square

3B. Vanishings of $H^1(\Sigma, \mathcal{O}_\Sigma(k))$ for $k > 0$. All of the remaining vanishings exploit the structure of $\tilde{\Sigma}$ as a \mathbb{P}^1 -bundle over $S^2 C$. Given work of the second author in [[Vermeire 2008b](#)], the projective normality of Σ follows by exploiting Terracini recursion as a corollary of the next result.

Theorem 3.3. *If C satisfies [Hypothesis 3.1](#), then $H^1(\Sigma, \mathcal{O}_\Sigma(2)) = 0$.*

Proof. We show that $H^2(\mathbb{P}^n, \mathcal{F}_{\tilde{\Sigma}}(2)) = 0$.

Since $\mathcal{O}(2H - E)$ is trivial along the fibers of $\pi : \tilde{\Sigma} \rightarrow S^2C$, $\mathcal{O}_{\tilde{\Sigma}}(2H - E) = \pi^*M$ for some line bundle M on S^2C [Hartshorne 1977, Exercise III.12.4]. We know from [Vermeire 2002, 3.6] that

$$\mathcal{O}_{\tilde{\Sigma}}(2H - E) \otimes \mathcal{O}_Z \cong \pi^*M \otimes \mathcal{O}_Z \cong L \boxtimes L \otimes \mathcal{O}_Z(-2\Delta).$$

Further restricting π to the double cover $d : C \times C \rightarrow S^2C$, by the projection formula and part (5) of Lemma 2.3 we have

$$H^i(Z, L \boxtimes L \otimes \mathcal{O}_Z(-2\Delta)) = H^i(S^2C, M) \oplus H^i\left(S^2C, M \otimes \mathcal{O}_{S^2C}\left(-\frac{\delta}{2}\right)\right).$$

Again by the projection formula, we know that $H^i(\tilde{\Sigma}, \mathcal{O}(2H - E)) = H^i(S^2C, M)$. By Lemma 2.5, we have $H^i(Z, L \boxtimes L \otimes \mathcal{O}_Z(-2\Delta)) \cong H^i(C, N_C^*(2))$. Thus we immediately have $H^2(Z, L \boxtimes L \otimes \mathcal{O}_Z(-2\Delta)) = 0$, but this in turn implies $H^2(S^2C, M) = H^2(\tilde{\Sigma}, \mathcal{O}(2H - E)) = 0$.

Let \mathcal{L}_L be the line bundle on S^2C such that $d^*\mathcal{L}_L = L \boxtimes L$ (see [Kouvidakis 2002, §2.1], for example). Now, as $L \boxtimes L \otimes \mathcal{O}_Z(-\Delta) = d^*(\mathcal{L}_L \otimes \mathcal{O}_{S^2C}(-\frac{\delta}{2}))$, we know that

$$\begin{aligned} d_*((L \boxtimes L) \otimes \mathcal{O}_Z(-\Delta)) &= \left[\mathcal{L}_L \otimes \mathcal{O}_{S^2C}\left(-\frac{\delta}{2}\right)\right] \oplus \left[\mathcal{L}_L \otimes \mathcal{O}_{S^2C}\left(-2\frac{\delta}{2}\right)\right] \\ &= \left[\mathcal{L}_L \otimes \mathcal{O}_{S^2C}\left(-\frac{\delta}{2}\right)\right] \oplus M. \end{aligned}$$

Again by Lemma 2.5 we know that

$$H^1(C \times C, L \boxtimes L \otimes \mathcal{O}_Z(-\Delta)) = H^1(C, \Omega_{\mathbb{P}^n}^1(2) \otimes \mathcal{O}_C) = 0,$$

where the vanishing comes from quadratic normality of the embedding of C . Thus $H^1(S^2C, M) = H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(2H - E)) = 0$.

We see immediately that $H^2(B_1, \mathcal{F}_{\tilde{\Sigma}}(2H)) = H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(2H))$, and from the sequence

$$0 \rightarrow \mathcal{O}_{\tilde{\Sigma}}(2H - E) \rightarrow \mathcal{O}_{\tilde{\Sigma}}(2H) \rightarrow \mathcal{O}_{\tilde{\Sigma}}(2H) \otimes \mathcal{O}_E \rightarrow 0$$

and the (just proved) fact that $H^i(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(2H - E)) = 0$ for $i = 1, 2$ implies further that $H^2(B_1, \mathcal{F}_{\tilde{\Sigma}}(2)) = H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(2) \otimes \mathcal{O}_E)$. A straightforward computation gives

$$\begin{aligned} h^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(2H) \otimes \mathcal{O}_E) &= h^1(C \times C, L^2 \boxtimes \mathcal{O}_C) \\ &= h^0(C, L^2) \cdot h^1(C, \mathcal{O}_C) \\ &= h^0(C, H^1(C, \mathcal{O}_C) \otimes L^2) \\ &= h^0(\mathbb{P}^n, R^2g_*\mathcal{F}_{\tilde{\Sigma}}(2)). \end{aligned}$$

Therefore, $h^2(B_1, \mathcal{F}_{\tilde{\Sigma}}(2)) = h^0(\mathbb{P}^n, R^2g_*\mathcal{F}_{\tilde{\Sigma}}(2))$.

Interpreting what we have just shown in terms of the Leray–Serre spectral sequence associated to $g_*\mathcal{F}_{\Sigma}(2)$, we have $h^2(B_1, \mathcal{F}_{\Sigma}(2)) = \dim E_2^{0,2}$. We also know that $R^1g_*\mathcal{F}_{\Sigma}(2) = 0$ by the projection formula and [Lemma 2.4](#) (3) and (5). Thus, at the E_2 level, where we have

$$0 \rightarrow E_2^{0,1} \xrightarrow{d_2} E_2^{2,0} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow E_2^{0,2} \xrightarrow{d_2} E_2^{2,1} \rightarrow 0,$$

we see that $E_2^{2,0} = E_{\infty}^{2,0}$ and $E_2^{0,2} = E_{\infty}^{0,2}$ because $H^i(\mathbb{P}^n, R^1g_*\mathcal{F}_{\Sigma}(2)) = 0$. Recall [[Weibel 1994](#), 5.2.6] that $H^2 := H^2(B_1, \mathcal{F}_{\Sigma}(2))$ has a finite filtration

$$0 = F^3H^2 \subseteq F^2H^2 \subseteq F^1H^2 \subseteq F^0H^2 = H^2,$$

where $F^2H^2 \cong E_{\infty}^{2,0}$ and $H^2/F^1H^2 \cong E_{\infty}^{0,2}$.

Now, because $\dim H^2 = \dim E_2^{0,2} = \dim E_{\infty}^{0,2}$, we have $F^1H^2 = 0$, but this implies that $F^2H^2 = E_{\infty}^{2,0} = 0$, and hence that $E_2^{2,0} = 0$. □

In [[Vermeire 2008b](#)] it was shown that for the general embedding of degree at least $2g+3$, Σ is projectively normal; the only vanishing that could not be shown to always hold was $H^1(\mathbb{P}^n, \mathcal{F}_{\Sigma}(2)) = 0$. [Theorem 3.3](#) allows us remove the hypothesis that the embedding must be general. The idea in [[Vermeire 2008b](#)] was to obtain a vanishing statement for direct image sheaves, and then to use those vanishings along with [[Mumford 1966](#), p. 52, Corollary 1 $\frac{1}{2}$] to show that the cohomology groups along the fibers vanish. Of course, to make this work, we must find a flat morphism and a locally free sheaf so that the restriction of the sheaf to the fiber is precisely the vanishing statement we want. This is done using [Theorem 2.1](#). However, note that in the proof we need to increase the degree of the embedding to at least $2g+5$, so that curves of degree $2g+3$ occur in the fibers.

Corollary 3.4. *Let $C \subset \mathbb{P}^n$ be a smooth curve embedded by a line bundle L of degree at least $2g+3$. Then Σ is projectively normal.*

Proof. We know by combining [[Vermeire 2008b](#), Proposition 12] with [[Wahl 1997](#), 1.16] that $H^1(\mathbb{P}^n, \mathcal{F}_{\Sigma}(k)) = 0$ for $k = 1, 3$, and by [[Vermeire 2008b](#), Corollary 11] that $H^1(\mathbb{P}^n, \mathcal{F}_{\Sigma}(k)) = 0$ for $k \geq 4$. Clearly, $H^1(\Sigma, \mathcal{O}_{\Sigma}(2)) = H^2(\mathbb{P}^n, \mathcal{F}_{\Sigma}(2))$. As these vanish by [Theorem 3.3](#), we note that by [Lemma 2.4](#) we have

$$H^2(B_2, \mathcal{O}(2H - E_1 - E_2)) = 0.$$

We further have $H^i(B_2, \mathcal{O}(2H - E_1 - E_2)) = 0$ for $i \geq 3$ by 5-regularity of \mathcal{F}_{Σ} .

By [Lemma 2.4](#), along the fibers of $E_1 \rightarrow C$ we are computing $H^i(\mathbb{P}^{n-2}, \mathcal{F}_C(1))$; thus $R^i f_*\mathcal{O}_{E_1}(2H - E_1 - E_2) = 0$ for $i \geq 0$. Hence $H^i(B_2, \mathcal{O}_{E_1}(2H - E_1 - E_2))$ vanishes, showing that $H^i(B_2, \mathcal{O}(2H - 2E_1 - E_2)) = 0$ for $i \geq 2$.

Fixing a point $p \in C$, and applying an extension of [Theorem 2.1](#) to $L(2p)$ (which now separates 6 points as L is nonspecial), we may blow up three times to get a resolution of Σ_2 . In the notation of [[Vermeire 2008b](#), Theorem 15], the

previous paragraph gives $R^i f_* \mathbb{O}_{E_1}(kH - 2E_1 - 2E_2 - E_3) = 0$ for $i \geq 2$, since the restriction of $\mathbb{O}_{E_1}(kH - 2E_1 - 2E_2 - E_3)$ to a fiber of $E_1 \rightarrow C$ is $\mathbb{O}(2H - 2E_1 - E_2)$ using the convention of [Remark 2.2](#). It was shown in [[Vermeire 2008b](#), Theorem 15] that $R^1 f_* \mathbb{O}_{E_1}(kH - 2E_1 - 2E_2 - E_3) = 0$, and so we know that H^1 along the fibers vanishes by [[Mumford 1966](#), page 52, Corollary 1 $\frac{1}{2}$]. Thus we have $H^1(B_2, \mathbb{O}(2H - 2E_1 - E_2)) = 0$ and so, as above,

$$H^1(B_2, \mathbb{O}(2H - E_1 - E_2)) = H^1(\mathbb{P}^n, \mathcal{F}_\Sigma(2)) = 0. \quad \square$$

Theorem 3.5. *If C satisfies [Hypothesis 3.1](#), then $H^i(\Sigma, \mathbb{O}_\Sigma(1)) = 0$ for $i = 1, 2$.*

Proof. For $i = 1, 2$, we have $h^i(\Sigma, \mathbb{O}_\Sigma(1)) = h^{i+1}(\mathbb{P}^n, \mathcal{F}_\Sigma(1))$ and $H^{i+1}(\mathbb{P}^n, \mathcal{F}_\Sigma(1))$ is isomorphic to $H^{i+1}(B_2, \mathbb{O}_{B_2}(H - E_1 - E_2))$ by the last part of [Lemma 2.4](#). Using [Equation \(1\)](#) twisted by H , the projection formula gives $R^i h_* (\mathbb{O}_{E_2}(H - E_1)) = R^i h_* (\mathbb{O}_{E_2}) \otimes \mathbb{O}_{\tilde{\Sigma}}(H - E_1)$. By part (6) of [Lemma 2.3](#) the restriction of $\mathbb{O}(H - E_1)$ to the fibers of $\tilde{\Sigma} \rightarrow S^2C$ is isomorphic to $\mathbb{O}_{\mathbb{P}^1}(-1)$, hence $h^i(\tilde{\Sigma}, \mathbb{O}(H - E_1)) = 0$ for all i , which implies that $h^i(E_2, \mathbb{O}_{E_2}(H - E_1)) = 0$. We therefore have $h^{i+1}(B_2, \mathbb{O}_{B_2}(H - E_1 - E_2)) = h^{i+1}(B_2, \mathbb{O}_{B_2}(H - E_1))$.

We see that $R^i f_* (\mathbb{O}_{B_2}(H - E_1)) = 0$ for $i \geq 1$ and $f_* (\mathbb{O}_{B_2}(H - E_1)) = \mathcal{F}_C(1)$ by [[Bertram et al. 1991](#), 1.2,1.4]. Thus $h^{i+1}(B_2, \mathbb{O}_{B_2}(H - E_1)) = h^{i+1}(\mathbb{P}^n, \mathcal{F}_C(1)) = 0$. □

Remark 3.6. In the case of a canonical curve, we have

$$h^0(\Sigma, R^1 g_* \mathbb{O}_{\tilde{\Sigma}}(H)) = h^1(C, \mathbb{O}_C) \cdot h^0(C, \mathbb{O}_C(1)) = g^2$$

while

$$h^1(\tilde{\Sigma}, \mathbb{O}_{\tilde{\Sigma}}(H)) = h^1(\mathbb{O}_C) \cdot h^0(\mathbb{O}_C(1)) + h^0(\mathbb{O}_C) \cdot h^1(\mathbb{O}_C(1)) = g^2 + 1.$$

Therefore using the 5-term sequence [\(2\)](#) again we see that $h^1(\Sigma, \mathbb{O}_\Sigma(1)) \geq 1$ (in fact, equality can be shown to hold). Thus the secant variety to a canonical curve of Clifford index at least 3 (for example, the generic curve of genus ≥ 7) is *never* ACM.

Note the secant variety of a canonical curve $C \subset \mathbb{P}^4$ is a hypersurface of degree 16, hence is ACM, but such curves have Clifford index ≤ 2 .

3C. Vanishings for $k = 0$. We now consider the vanishing of $H^i(\Sigma, \mathbb{O}_\Sigma)$ where $i = 1, 2$.

Proposition 3.7. *If C satisfies [Hypothesis 3.1](#), then $H^1(\Sigma, \mathbb{O}_\Sigma) = 0$.*

Proof. Associated to the morphism $g : B_1 \rightarrow \mathbb{P}^n$ we have

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & \downarrow & \downarrow & & \downarrow & \\
 0 & \longrightarrow & H^1(g_*\mathbb{O}_{\tilde{\Sigma}}) & \longrightarrow & H^1(\mathbb{O}_{\tilde{\Sigma}}) & \xrightarrow{\gamma} & H^0(R^1g_*\mathbb{O}_{\tilde{\Sigma}}) \longrightarrow \\
 & & \downarrow & & \downarrow \alpha & & \downarrow \\
 0 & \longrightarrow & H^1(g_*\mathbb{O}_Z) & \longrightarrow & H^1(\mathbb{O}_Z) & \xrightarrow{\beta} & H^0(R^1g_*\mathbb{O}_Z) \longrightarrow 0
 \end{array}$$

where the horizontal maps come from 5-term exact sequences.

As $Z \cong C \times C$, we see that the inclusion and projection in the bottom row come from the Künneth formula. The map $\alpha : H^1(\mathbb{O}_{\tilde{\Sigma}}) \rightarrow H^1(\mathbb{O}_Z)$ is an inclusion because it is the diagonal mapping $\alpha : H^1(\mathbb{O}_{S^2C}) \rightarrow H^1(C, \mathbb{O}_C) \oplus H^1(C, \mathbb{O}_C)$ induced by the pull-back of $d : Z \rightarrow S^2C$ to $\tilde{\Sigma}$. We conclude that the composition $\beta \circ \alpha$ is an isomorphism. Moreover, as $H^0(R^1g_*\mathbb{O}_{\tilde{\Sigma}}) \rightarrow H^0(R^1g_*\mathbb{O}_Z)$ is an isomorphism, we see that γ is an isomorphism by commutativity of the diagram. Hence, $H^1(g_*\mathbb{O}_{\tilde{\Sigma}}) = H^1(\Sigma, \mathbb{O}_{\Sigma}) = 0$. \square

Proposition 3.8. *If C satisfies Hypothesis 3.1, then $H^2(\Sigma, \mathbb{O}_{\Sigma}) = 0$.*

Proof. We note that $h^i(\Sigma, \mathbb{O}_{\Sigma}) = h^{i+1}(\mathbb{P}^n, \mathcal{F}_{\Sigma})$ for $i = 1, 2$. Moreover, by part (6) of Lemma 2.4 we have $h^j(\mathbb{P}^n, \mathcal{F}_{\Sigma}) = h^j(B_2, \mathbb{O}_{B_2}(-E_1 - E_2))$. Therefore, the result follows if we can show that $h^2(B_2, \mathbb{O}_{B_2}(-E_1 - E_2)) = h^3(B_2, \mathbb{O}_{B_2}(-E_1 - E_2))$, since we know by Proposition 3.7 that $h^2(B_2, \mathbb{O}_{B_2}(-E_1 - E_2)) = 0$.

To this end, consider the long exact sequence associated to Equation (1) on page 451. The result will follow if $h^2(B_2, \mathbb{O}_{B_2}(-E_1)) = h^2(E_2, \mathbb{O}_{E_2}(-E_1))$ is equal to g and $h^1(E_2, \mathbb{O}_{E_2}(-E_1)) = h^3(B_2, \mathbb{O}_{B_2}(-E_1)) = 0$.

From the sequence $0 \rightarrow \mathbb{O}_{B_2}(-E_1) \rightarrow \mathbb{O}_{B_2} \rightarrow \mathbb{O}_{E_1} \rightarrow 0$ we see immediately that $h^i(B_2, \mathbb{O}_{B_2}(-E_1)) = g$ if $i = 2$ and is zero otherwise as $R^j f_*\mathbb{O}_{B_2} = 0$ for $j > 0$ from Lemma 2.4 (2) and $h^j(\mathbb{O}_{E_1}) = h^j(\mathbb{O}_C)$ for all j .

We compute the cohomology of $\mathbb{O}_{E_2}(-E_1)$ using Equation (1). Using the projection formula and part (4) of Lemma 2.4, we see that $R^i h_*\mathbb{O}_{E_2}(-E_1) = 0$ for $i > 0$. Thus, $H^i(\mathbb{O}_{E_2}(-E_1)) \cong H^i(\mathbb{O}_{\tilde{\Sigma}}(-E_1))$.

To compute $H^i(\mathbb{O}_{\tilde{\Sigma}}(-E_1))$, observe that

$$0 \rightarrow \pi_*\mathbb{O}_{\tilde{\Sigma}}(-E_1) \rightarrow \pi_*\mathbb{O}_{\tilde{\Sigma}} \rightarrow \pi_*\mathbb{O}_Z \rightarrow R^1\pi_*\mathbb{O}_{\tilde{\Sigma}}(-E_1) \rightarrow 0$$

with all remaining higher direct images vanishing by parts (2) and (4) of Lemma 2.3 and $\pi_*\mathbb{O}_{\tilde{\Sigma}}(-E_1) = 0$ by part (6).

As $\text{Hom}_{\mathbb{O}_{S^2C}}(\mathbb{O}_{S^2C}, \mathbb{O}_{S^2C}(-\frac{\delta}{2}))$ is trivial, this gives rise to the natural inclusion

$$\pi_*\mathbb{O}_{\tilde{\Sigma}} \cong \mathbb{O}_{S^2C} \hookrightarrow \mathbb{O}_{S^2C} \oplus \mathbb{O}_{S^2C} \left(-\frac{\delta}{2}\right) \cong \pi_*\mathbb{O}_Z,$$

and we see that $H^i(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}) \hookrightarrow H^i(Z, \mathcal{O}_Z)$. In fact, using the long exact sequence on $\tilde{\Sigma}$, these inclusions imply that $H^i(Z, \mathcal{O}_Z) \cong H^i(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}) \oplus H^{i+1}(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(-E_1))$.

As $h^1(S^2C, \mathcal{O}_{S^2C}) = g$ and $h^2(S^2C, \mathcal{O}_{S^2C}) = \binom{g}{2}$ by [Macdonald 1962], using the sequence $0 \rightarrow \mathcal{O}_{\tilde{\Sigma}}(-E_1) \rightarrow \mathcal{O}_{\tilde{\Sigma}} \rightarrow \mathcal{O}_Z \rightarrow 0$ together with the Künneth formula and that $H^i(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}) \cong H^i(S^2C, \mathcal{O}_{S^2C})$, implies that $h^2(E_2, \mathcal{O}_{E_2}(-E_1)) = g$, and that $h^3(E_2, \mathcal{O}_{E_2}(-E_1)) = \binom{g+1}{2}$. Further, we see immediately that $H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(-E_1))$ is 0, since $H^0(Z, \mathcal{O}_Z) \cong H^0(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}) \oplus H^1(\tilde{\Sigma}, \mathcal{O}_{\tilde{\Sigma}}(-E_1))$. \square

Proof of Theorem 1. As explained at the beginning of the section, in order to show that Σ is ACM we are left to show that $H^i(\Sigma, \mathcal{O}_{\Sigma}(k)) = 0$ for $i = 1, 2$ and all $k \leq 3 - i$.

The vanishings for $k < 0$ were shown in Theorem 3.2. The vanishing for $i = 1$ and $k = 0$ is Proposition 3.7, while $i = 2$ and $k = 0$ is Proposition 3.8. Both vanishings for $k = 1$ are found in Theorem 3.5. Finally, the vanishing for $i = 1$ and $k = 2$ is found in Theorem 3.3. \square

As an immediate consequence of the proof of Proposition 3.8 we get a sharpening of the regularity result of the second author in [Vermeire 2008b].

Corollary 3.9. *If C satisfies Hypothesis 3.1, then \mathcal{F}_{Σ} has regularity 3 if C is rational and regularity 5 otherwise.*

Proof. Running the long exact sequence associated to Equation (1) in the proof of Proposition 3.8 shows that $h^4(\mathbb{P}^n, \mathcal{F}_{\Sigma}) = \binom{g+1}{2}$. \square

4. Betti diagrams

In this section we attempt to paint a picture of the shape of the Betti diagram of S_{Σ} that parallels the discussion of the Betti diagram of a high degree curve in Chapter 8 of [Eisenbud 2005]. In Section 4A we use the fact that Σ is ACM to use duality and algebraic techniques to compute the extremal nontrivial Betti numbers, $\beta_{1,3}$ (Proposition 4.4) and $\beta_{n-3,n+1}$ (Corollary 4.1) as well as the Hilbert polynomial. Independent of the Cohen–Macaulay property, we prove a nonvanishing result about the length of the degree $(k+2)$ linear strand of S_{Σ_k} using determinantal methods and Koszul homology (Proposition 4.10 and Theorem 1.2) in Section 4B.

4A. Computing Betti numbers. We begin with a simple consequence of duality. As Σ is ACM, dualizing a resolution of S_{Σ} and shifting by $-n-1$ gives a resolution of the canonical module, which is defined to be $\omega_{\Sigma} = \text{Ext}^{n-3}(S_{\Sigma}, S(-n-1)) = \bigoplus_{d \in \mathbb{Z}} H^0(\Sigma, \omega_{\Sigma}^{\circ} \otimes L^d)$ where $\omega_{\Sigma}^{\circ} = \mathcal{E}xt_{\mathbb{P}^n}^{n-3}(\mathcal{O}_{\Sigma}, \mathcal{O}_{\mathbb{P}^n}(-n-1))$ is the dualizing sheaf of Σ . Therefore, the last few Betti numbers of S_{Σ} are the first few of ω_{Σ} . As an immediate consequence of Corollary 3.9 we see that the number of minimal generators of ω_{Σ} in degree 0 is $\binom{g+1}{2}$ and hence depends only on g , independent of the embedding (as long as the degree is at least $2g+3$).

Corollary 4.1. *If C satisfies Hypothesis 3.1, then $\beta_{n-3,n+1} = \binom{g+1}{2}$.*

Proof. If $g = 0$, we know that $\beta_{n-3,n+1} = 0$. If $g > 0$, then Corollary 3.9 shows that $\text{reg } S_\Sigma = 4$. Hence, the a -invariant of S_Σ is 0, so $h^0(\omega_\Sigma^\circ) = \beta_{0,0}(\omega_\Sigma) = \beta_{n-3,n+1}(S_\Sigma)$. By Serre duality,

$$h^0(\Sigma, \omega_\Sigma^\circ) = h^3(\Sigma, \mathbb{C}_\Sigma) = h^4(\mathbb{P}^n, \mathcal{F}_\Sigma) = \binom{g+1}{2}. \quad \square$$

Knowing $\beta_{n-3,n+1}$ allows us to compute the Hilbert polynomial of S_Σ and to gather information about other Betti numbers inductively. To begin this process, fix general linear forms $H_1, H_2, H_3, H_4 \in S$. Let X be the intersection of Σ with the hyperplanes determined by H_1 and H_2 and $M = S_\Sigma / \langle H_1, H_2, H_3, H_4 \rangle$. Using Corollary 4.1 we may compute the genus of X from which formulae for the Hilbert polynomial of S_Σ and $\beta_{1,3}$ follow. First we gather together basic facts about X .

Lemma 4.2. *If C satisfies Hypothesis 3.1, the variety X is a smooth curve of degree $D = \binom{d-1}{2} - g$ embedded in \mathbb{P}^{n-2} via the complete linear series associated to a line bundle A and $S_X = S_\Sigma / \langle H_1, H_2 \rangle$.*

Proof. All the statements follow immediately from the fact that Σ is ACM. The only thing that may not be immediate to the reader is that $\text{deg}(\Sigma) = \binom{d-1}{2} - g$, though this is certainly well-known to experts.

To see this, take a generic $L = \mathbb{P}^{n-3} \subset \mathbb{P}^n$ and consider the induced projection $\pi : \mathbb{P}^n \dashrightarrow \mathbb{P}^2$. Every point of intersection of L with Σ corresponds to a node of $\pi(C)$. It is well-known that the number of nodes is $\binom{d-1}{2} - g$. \square

We will denote the genus of X by G . To compute G we compare the Hilbert function of S_X to that of successive quotients by H_1 and H_2 .

Proposition 4.3. *If C satisfies Hypothesis 3.1, the genus of X is*

$$G = \frac{1}{2}(d-2)(d+2g-3).$$

Proof. Since S_X is 4-regular, $h^0(X, A^m) = mD - G + 1$ for $m \geq 3$. We also know that the ideal of Σ is empty in degree less than three, since a quadric hypersurface vanishing on Σ must vanish twice on C , but this is not possible since C is nondegenerate. Therefore, we can fill in the table of Hilbert functions below where each entry in the first two columns of the table is the sum of the entries directly above and to the right.

	$S_\Sigma / \langle H_1, H_2 \rangle$	$S_\Sigma / \langle H_1, H_2, H_3 \rangle$	M
2	$\binom{n}{2}$	$\binom{n-1}{2}$	$\binom{n-2}{2}$
3	$3D - G + 1$	$3D - G + 1 - \binom{n}{2}$	$3D - G + 1 - \binom{n}{2} - \binom{n-1}{2}$
4	$4D - G + 1$	D	$G - 2D - 1 + \binom{n}{2}$

But computing graded Betti numbers via Koszul homology as in Proposition 2.7 in [Eisenbud 2005] shows that $\dim M_4 = \beta_{n-3,n+1} = \binom{g+1}{2}$. Substituting $n = d - g$ and simplifying $G = 2D + 1 - \binom{d-g}{2} + \binom{g+1}{2}$ gives the desired result. \square

The computation of the Hilbert polynomial $P_\Sigma(m)$ follows easily.

Proof of Theorem 1.1. Using [Eisenbud 2005, Theorem 4.2], the Hilbert polynomial and Hilbert function of S_Σ agree for $m \geq \text{reg } S_\Sigma + \text{proj-dim } S_\Sigma - n \geq 4 - 3 = 1$. Write

$$P_\Sigma(m) = \sum_{i=0}^3 \alpha_i \binom{m+i-1}{i}.$$

As X is gotten by cutting down by a regular sequence of two hyperplanes, $P_X(m) = P_\Sigma(m) - P_\Sigma(m-1) - P_\Sigma(m-2) = \alpha_3 m + \alpha_2$. Since X is a curve of degree D and genus, G , we see that $\alpha_3 = D$ and $\alpha_2 = 1 - G$. Since the ideal of Σ is empty in degrees 1 and 2, we see that $P_\Sigma(1) = n + 1$ and $P_\Sigma(2) = \binom{n+2}{2}$ and the result follows. \square

We compute $\beta_{1,3}$ and get a relationship on Betti numbers at the beginning of the resolution.

Proposition 4.4. *If C satisfies Hypothesis 3.1, we have*

$$\beta_{1,3} = \binom{n+1}{3} - (d-2)n - 3g + 1 \quad \text{and} \quad \beta_{2,4} = \beta_{1,4} + \beta_{1,3}(n+1) - \binom{n+4}{n} + P_\Sigma(4).$$

Proof. As observed above, the Hilbert polynomial and function of S_X agree in degree 3 and higher. Since $\beta_{1,3} = \binom{n+1}{3} - (S_X)_3$ we get $\beta_{1,3} = \binom{n+1}{3} - 3D + G - 1$, and this simplifies to the given formula.

By [Eisenbud 2005, Corollary 1.10] we get a formula for the Hilbert function of S_Σ in terms of graded Betti numbers:

$$(S_\Sigma)_m = \sum_{i \geq 0, j \in \mathbb{Z}} (-1)^i \beta_{i,j} \binom{n+m-j}{n}.$$

When $m = 4$, we must have $j \leq 4$ for $\beta_{i,j}$ to contribute to the sum. As we know that the ideal of X does not contain any forms of degree < 3 , the result follows. \square

Remark 4.5. In the formula for $\beta_{2,4}$ we have an explicit formula for each term except $\beta_{1,4}$, which is the number of quartic minimal generators of I_Σ . For large d , we know $\beta_{1,4} = 0$, as the ideal of Σ is generated by cubics [Vermeire 2008a].

Using duality, we get a similar result for the tail of the resolution.

Theorem 4.6. *If C satisfies Hypothesis 3.1, the tail of the graded Betti diagram of S_Σ has the form*

	$n - 5$	$n - 4$	$n - 3$
0	—	—	—
1	—	—	—
2	*	*	A
3	*	$A + B + \binom{g+1}{2} \binom{n}{2} - \binom{g}{2} (n-3)(n-1) - G$	C
4	B	$C + \binom{g}{2} (n-3)$	$\binom{g+1}{2}$

Proof. Let $A = \beta_{n-3,n-1}$, $B = \beta_{n-5,n-1}$ and $C = \beta_{n-3,n}$. We know that the canonical module ω_X is $\bigoplus_{n \in \mathbb{Z}} H^0(K_X \otimes A^n)$, where K_X is the canonical line bundle of X . By duality, $\beta_{i,j}(\omega_X) = \beta_{n-3-i,n-1-j}(S_\Sigma)$.

By [Eisenbud 2005, Corollary 1.10] we get a formula for the Hilbert function of ω_X in terms of graded Betti numbers:

$$h^0(K_X \otimes A^m) = \sum_{i \geq 0, j \in \mathbb{Z}} (-1)^i \beta_{i,j}(\omega_X) \binom{n-2+m-j}{n-2}.$$

By Serre duality and Riemann–Roch $h^0(K_X \otimes A^{-1}) = h^1(A) = g(d-2)$. Thus, $g(d-2) = (n-1) \binom{g+1}{2} + C - \beta_{n-4,n}$, which gives the desired statement. The second statement follows from the equation

$$G = \binom{g+1}{2} \binom{n}{2} - \binom{g}{2} (n-3)(n-1) + B - \beta_{n-4,n-1} + A. \quad \square$$

In particular, if $g = 2$, we have the following immediate corollary.

Corollary 4.7. *If C satisfies Hypothesis 3.1 and $g = 2$, the tail has the form*

	$n - 5$	$n - 4$	$n - 3$
0	—	—	—
1	—	—	—
2	*	*	A
3	*	$A + B + d - 5$	C
4	B	$C + d - 5$	$\binom{g+1}{2}$

Based on Example 1.4 and the following example, we expect $A = B = C = 0$.

Example 4.8. Suppose C is a genus 2 curve of degree 12 in \mathbb{P}^{10} . We use Example (c) of [Eisenbud et al. 1988] to compute the ideal of the curve determinantly in Macaulay 2 over the field of rational numbers. We then used the code created to implement ideas in [Sidman and Sullivant 2006] to compute the least degree pieces of the ideals of the secant varieties. Computing the degree, dimension, and

projective dimension of the resulting ideals showed that we had actually computed the secant ideals.

	0	1	2	3	4	5	6	7	8	9
total:	1	43	222	558	840	798	468	147	17	2
0:	1	-	-	-	-	-	-	-	-	-
1:	-	43	222	558	840	798	468	147	8	-
2:	-	-	-	-	-	-	-	-	9	2

The Betti diagrams for S_{Σ_1} and S_{Σ_2} are

	0	1	2	3	4	5	6	7			0	1	2	3	4	5
total:	1	70	283	483	413	155	14	3	total:	1	41	94	61	11	4	
0:	1	-	-	-	-	-	-	-	0:	1	-	-	-	-	-	-
1:	-	-	-	-	-	-	-	-	1:	-	-	-	-	-	-	-
2:	-	70	283	483	413	155	-	-	2:	-	-	-	-	-	-	-
3:	-	-	-	-	-	-	7	-	3:	-	41	94	61	-	-	-
4:	-	-	-	-	-	-	7	3	4:	-	-	-	-	-	-	-
									5:	-	-	-	-	6	-	
									6:	-	-	-	-	5	4	

4B. The length of the first nonzero strand. We now turn to the consideration of a lower bound on the length of the minimal degree linear strand of the ideal of Σ_k , essentially following Chapter 8B.2 of [Eisenbud 2005], building on Green and Lazarsfeld’s proof of nonvanishing for curves. In this section we will assume the following:

Hypothesis 4.9. *C is a smooth curve of genus g and degree d embedded into \mathbb{P}^n via a line bundle L that factors as $L = L_1 \otimes L_2$, where $|L_1| = s$ and $|L_2| = t$, with $1 \leq s \leq t$.*

First note that part of the proof of [Eisenbud 2005, Theorem 8.12] which is given in the case $k = 0$ goes through for arbitrary k and allows us to see easily that the degree $k + 2$ linear strand of the Betti diagram of Σ_k has length at least p .

Proposition 4.10. *Under the conditions of Hypothesis 4.9, if $d \geq 2g + 2k + 1 + p$, then $\beta_{p,k+1+p} \neq 0$.*

Proof. Factor L so that $\deg L_1 \geq g+k+1$ and $\deg L_2 = g+k+p$. By Riemann–Roch $h^0(C, L_1) \geq k+2$ and $h^0(C, L_2) \geq k+p+1$. Thus multiplication of sections gives rise to a 1-generic matrix of linear forms with at least $(k+2)$ rows and $(k+1+p)$ columns. Delete rows and columns to get a $(k+2) \times (k+1+p)$ matrix which is still 1-generic as an equation making a generalized entry of the smaller matrix zero also makes a generalized entry of the larger matrix zero. The maximal minors of the smaller matrix are resolved by an Eagon–Northcott complex of length p . The resolution of this ideal is a subcomplex of the ideal of Σ_k . The result follows. \square

We can get a better lower bound by exhibiting an explicit nontrivial cycle in the Koszul homology of S_{Σ_k} to show that $\beta_{s+t-2k-1,s+t-k}$ does not vanish.

In [Eisenbud 2005, Theorem 8.15], the following result is stated for $k = 1$:

Theorem 4.11 [Eisenbud 2005, Theorem 8.15]. *If $I \subset S$ is a homogenous ideal which contains no forms of degree less than or equal to k , then $\beta_{i,i+k} \neq 0$ if and only if there exists $\gamma \in \wedge^i S^{n+1}(-i)$ of degree $i + k$ whose image under the differential of the Koszul complex is nonzero and lies in $I \otimes \wedge^{i-1} S^{n+1}(-i + 1)$.*

Proof. The proof goes through as in [Eisenbud 2005], replacing one by k everywhere. □

We show that [Eisenbud 2005, Theorem 8.13] can be extended to the case of minors of arbitrary size.

Theorem 4.12. *Suppose that A is an $(s + 1) \times (t + 1)$ matrix of linear forms with $s + 1 \geq k + 2$. If the $s + t + 1$ elements in the union of the entries of the zeroth row and column are linearly independent and some $(k + 2)$ minor involving the zeroth row or column does not vanish, then $\beta_{s+t-2k-1,s+t-k}(S/I_{k+2})$ does not vanish.*

Proof. By Theorem 4.11 it suffices to construct an explicit cycle

$$\gamma \in \wedge^{s+t-2k-1} S^{n+1}(-s - t + 2k + 1)$$

of degree $s + t - k$ whose image under the differential is a nonzero element of $I_{k+1} \otimes \wedge^{s+t-2k-2} S^{n+1}(-s - t + 2k + 2)$. To do this we set some notation.

By our hypotheses, the matrix A has the form

$$A = \begin{pmatrix} a_{0,0} & a_{0,1} & \cdots & a_{0,t} \\ a_{1,0} & a_{1,1} & \cdots & a_{1,t} \\ \cdot & & & \\ a_{s,0} & a_{s,1} & \cdots & a_{s,t} \end{pmatrix} = \begin{pmatrix} x_0 & x_1 & \cdots & x_t \\ x_{1+t} & a_{1,1} & \cdots & x_{1,t} \\ \cdot & & & \\ x_{s+t} & a_{s,1} & \cdots & a_{s,t} \end{pmatrix}.$$

Since the x_i are linearly independent they may be chosen as part of a basis for S_1 , and we may choose a basis $\{e_i\}$ for S^{n+1} so that $\partial(e_i) = x_i$ for $i = 0, \dots, s + t$.

Let $\sigma \subset \{1, \dots, s\}$ and $\tau \subset \{0, \dots, t\}$ be sets of size $k + 1$ and σ_t denote the set gotten by adding t to each element of σ . Let $e_{\sigma_t, \tau}$ be the wedge product of $\{e_0, \dots, e_{s+t}\} \setminus (\sigma_t \cup \tau)$ in the standard order. Note that $e_{\sigma_t, \tau} \in \wedge^{s+t-2k-2} S^{n+1}$.

We define an element γ which will serve as our nonzero cycle. Informally, it is the signed sum of all of the $(k + 1)$ -minors of A which do not involve the top row, each indexed by an element $e_{\sigma_t, \tau}$ in a natural way. More precisely,

$$\gamma = \sum_{\sigma, \tau} (-1)^{(\sigma+\tau)+t(k+1)} \det(\sigma | \tau) e_{\sigma_t, \tau},$$

where we define $\sigma + \tau$ to be the sum of the union of the elements in σ and τ and $\det(\sigma | \tau)$ is the minor of A gotten by using the rows in σ and the columns in τ .

To complete the proof we need to show that the coefficients of $\partial(\gamma)$ are all of the $(k + 2)$ -minors of A involving the zeroth row or column. The only basis elements which can have nonzero coefficients are $e_{\sigma',\tau}$, where $\sigma' \subset \{1, \dots, s\}$ and $|\sigma'| = k + 2$ and $e_{\sigma_i,\tau'}$ where $\tau' \subset \{0, \dots, t\}$ also has size $k + 2$.

To understand the coefficient of $e_{\sigma_i,\tau'}$, note that there are $k + 2$ basis elements $e_{\sigma_i,\tau}$ whose images under the differential could contain $e_{\sigma_i,\tau'}$ with nonzero coefficient. Since $\partial(e_i) = x_i$ for $i = 0, \dots, t$, we see that the coefficient of $e_{\sigma_i,\tau'}$ is $\pm \det(\sigma_i \cup \{0\} \mid \tau')$ where the differential expands the determinant along the zeroth row.

Similarly, the coefficient of $e_{\sigma',\tau}$ is $\pm \det(\sigma' \mid \tau \cup \{0\})$, the differential expands the determinant along the zeroth column. (If $0 \in \tau$, we repeat the zeroth column twice and get coefficient zero.) □

We are now ready to prove [Theorem 1.2](#), which is analogous to [[Eisenbud 2005](#), Theorem 8.12].

Proof of Theorem 1.2. We will construct a matrix A corresponding to the factorization of $L = L_1 \otimes L_2$ by choosing bases carefully as in the proof of [[Eisenbud 2005](#), Theorem 8.12]. Let B_i be the base locus of L_i . Fix a basis β_0, \dots, β_t of $H^0(L_2)$ so that the divisor of β_i is $B_2 + D_i$ where D_i and B_2 have disjoint support. Let D be the divisor consisting of the union of the points in the divisors determined by β_0, \dots, β_t . Since $L_1(-B_1)$ is base-point free, a general element is disjoint from D and from B_1 . Therefore we can pick a basis $\alpha_0, \dots, \alpha_s$ so that the divisor of each α_i is $B_1 + E_i$ where E_i is disjoint from D and from B_1 .

We will show that the $s + t + 1$ elements in the union of any row and any column of the corresponding matrix A are linearly independent. Without loss of generality, consider the top row and leftmost column. We know that the elements of the column $\alpha_0\beta_0, \alpha_1\beta_0, \dots, \alpha_s\beta_0$ are linearly independent, as are the elements $\alpha_0\beta_0, \alpha_0\beta_1, \dots, \alpha_0, \beta_t$. Suppose γ is an element in the intersection of the two vector spaces with these bases. This implies that the divisor of γ contains the divisor of α_0 and of β_0 . This implies that it must contain D_0 and E_0 as well as the base loci B_1 and B_2 . Since $\gamma \in H^0(L)$ and $\alpha_0\beta_0 \in H^0(L)$, then one is a scalar multiple of the other. Therefore, we conclude that the union of the elements in the top row and first column form a set of $s + t + 1$ linearly independent elements.

As the matrix A is 1-generic, we know that the ideal generated by its maximal minors has the expected codimension and hence some $(k + 2)$ -minor does not vanish. Permuting rows and columns we can assume it is in the upper lefthand corner. Since $I_{k+2} \subseteq I_{\Sigma_k}$, the result follows from [Theorems 4.11 and 4.12](#).

If $\deg L \geq 2g + 2k + p + 1$, then L can be factored as the product of line bundles L_1 with degree at least $g + k + \lfloor (1 + p)/2 \rfloor$ and L_2 with degree greater than or equal to $\deg L_1$. If L_1 and L_2 are generic, then each has at least $k + 2$ sections. □

Acknowledgements

We thank David Eisenbud for his suggestions, and we thank two anonymous referees for suggestions which improved this paper. We would not have discovered the statements of our main theorems without help from Macaulay 2 [Grayson and Stillman].

In computing secant equations we used code developed with the help of Mike Stillman in conjunction with the first author's work with Seth Sullivant on [Sidman and Sullivant 2006].

References

- [Barth et al. 2004] W. P. Barth, K. Hulek, C. A. M. Peters, and A. Van de Ven, *Compact complex surfaces*, 2nd ed., *Ergebnisse der Math. (3)* **4**, Springer, Berlin, 2004. MR 2004m:14070 Zbl 1036.14016
- [Bertram 1992] A. Bertram, “Moduli of rank-2 vector bundles, theta divisors, and the geometry of curves in projective space”, *J. Differential Geom.* **35**:2 (1992), 429–469. MR 93g:14037 Zbl 0787.14014
- [Bertram et al. 1991] A. Bertram, L. Ein, and R. Lazarsfeld, “Vanishing theorems, a theorem of Severi, and the equations defining projective varieties”, *J. Amer. Math. Soc.* **4**:3 (1991), 587–602. MR 92g:14014 Zbl 0762.14012
- [Graf von Bothmer and Hulek 2004] H.-C. Graf von Bothmer and K. Hulek, “Geometric syzygies of elliptic normal curves and their secant varieties”, *Manuscripta Math.* **113**:1 (2004), 35–68. MR 2006b:14009 Zbl 1053.14032
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra: with a view toward algebraic geometry*, Graduate Texts in Math. **150**, Springer, New York, 1995. MR 97a:13001 Zbl 0819.13001
- [Eisenbud 2005] D. Eisenbud, *The geometry of syzygies: a second course in commutative algebra and algebraic geometry*, Graduate Texts in Math. **229**, Springer, New York, 2005. MR 2005h:13021 Zbl 1066.14001
- [Eisenbud et al. 1988] D. Eisenbud, J. Koh, and M. Stillman, “Determinantal equations for curves of high degree”, *Amer. J. Math.* **110**:3 (1988), 513–539. MR 89g:14023 Zbl 0681.14027
- [Eisenbud et al. 2005] D. Eisenbud, M. Green, K. Hulek, and S. Popescu, “Restricting linear syzygies: algebra and geometry”, *Compos. Math.* **141**:6 (2005), 1460–1478. MR 2006m:14072 Zbl 1086.14044
- [Fisher 2006] T. Fisher, “The higher secant varieties of an elliptic normal curve”, preprint, 2006, Available at <http://www.dpmms.cam.ac.uk/~taf1000/papers/hsecenc.html>.
- [Grayson and Stillman] D. R. Grayson and M. E. Stillman, *Macaulay 2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2>.
- [Green 1984] M. L. Green, “Koszul cohomology and the geometry of projective varieties”, *J. Differential Geom.* **19**:1 (1984), 125–171. MR 85e:14022 Zbl 0559.14008
- [Green and Lazarsfeld 1985] M. Green and R. Lazarsfeld, “On the projective normality of complete linear series on an algebraic curve”, *Invent. Math.* **83**:1 (1985), 73–90. MR 87g:14022 Zbl 0594.14010
- [Green and Lazarsfeld 1988] M. Green and R. Lazarsfeld, “Some results on the syzygies of finite sets and algebraic curves”, *Compositio Math.* **67**:3 (1988), 301–314. MR 90d:14034 Zbl 0671.14010

- [Grothendieck 1957] A. Grothendieck, “Sur quelques points d’algèbre homologique”, *Tôhoku Math. J. (2)* **9** (1957), 119–221. [MR 21 #1328](#) [Zbl 0118.26104](#)
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, New York, 1977. [MR 57 #3116](#) [Zbl 0531.14001](#)
- [Kouvidakis 2002] A. Kouvidakis, “On some results of Morita and their application to questions of ampleness”, *Math. Z.* **241**:1 (2002), 17–33. [MR 2003i:14006](#) [Zbl 1031.14010](#)
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, *Ergebnisse der Math. (3)* **48**, Springer, Berlin, 2004. [MR 2005k:14001a](#) [Zbl 1093.14501](#)
- [Macdonald 1962] I. G. Macdonald, “Symmetric products of an algebraic curve”, *Topology* **1** (1962), 319–343. [MR 27 #1445](#) [Zbl 0121.38003](#)
- [Mumford 1966] D. Mumford, *Lectures on curves on an algebraic surface*, *Annals of Mathematics Studies* **59**, Princeton University Press, 1966. [MR 35 #187](#) [Zbl 0187.42701](#)
- [Sidman and Sullivant 2006] J. Sidman and S. Sullivant, “Prolongations and computational algebra”, preprint, 2006. to appear in *Canadian Math. J.* [arXiv arxiv:math.AC/0611696](#)
- [Vermeire 2001] P. Vermeire, “Some results on secant varieties leading to a geometric flip construction”, *Compositio Math.* **125**:3 (2001), 263–282. [MR 2002c:14027](#) [Zbl 1056.14016](#)
- [Vermeire 2002] P. Vermeire, “On the regularity of powers of ideal sheaves”, *Compositio Math.* **131**:2 (2002), 161–172. [MR 2003f:14016](#) [Zbl 1083.14503](#)
- [Vermeire 2008a] P. Vermeire, “Generation and syzygies of the first secant variety”, preprint, 2008. [arXiv 0809.4463](#)
- [Vermeire 2008b] P. Vermeire, “Regularity and normality of the secant variety to a projective curve”, *J. Algebra* **319**:3 (2008), 1264–1270. [MR 2009b:14099](#) [Zbl 1132.14027](#)
- [Wahl 1997] J. Wahl, “On cohomology of the square of an ideal sheaf”, *J. Algebraic Geom.* **6**:3 (1997), 481–511. [MR 99b:14046](#) [Zbl 0892.14022](#)
- [Weibel 1994] C. A. Weibel, *An introduction to homological algebra*, *Cambridge Studies in Advanced Mathematics* **38**, Cambridge University Press, 1994. [MR 95f:18001](#) [Zbl 0797.18001](#)

Communicated by David Eisenbud

Received 2008-09-01

Revised 2009-04-01

Accepted 2009-04-29

jsidman@mtholyoke.edu

Mount Holyoke College, Department of Mathematics and Statistics, 415A Clapp Lab, South Hadley, MA 01075, United States

verme1pj@cmich.edu

Central Michigan University, Department of Mathematics, 214 Pearce, Mount Pleasant, MI 48859, United States