

Algebra & Number Theory

Volume 3

2009

No. 5

A general homological Kleiman–Bertini theorem

Susan J. Sierra



mathematical sciences publishers

A general homological Kleiman–Bertini theorem

Susan J. Sierra

Let G be a smooth algebraic group acting on a variety X . Let \mathcal{F} and \mathcal{E} be coherent sheaves on X . We show that if all the higher $\mathcal{T}or$ sheaves of \mathcal{F} against G -orbits vanish, then for generic $g \in G$, the sheaf $\mathcal{T}or_j^X(g\mathcal{F}, \mathcal{E})$ vanishes for all $j \geq 1$. This generalizes a result of Miller and Speyer for transitive group actions and a result of Speiser, itself generalizing the classical Kleiman–Bertini theorem, on generic transversality, under a general group action, of smooth subvarieties over an algebraically closed field of characteristic 0.

1. Introduction

All schemes that we consider in this paper are of finite type over a fixed field k ; we make no assumptions on the characteristic of k .

Our starting point is this:

Theorem 1.1 [Miller and Speyer 2008]. *Let X be a variety with a transitive left action of a smooth algebraic group G . Let \mathcal{F} and \mathcal{E} be coherent sheaves on X , and for all k -points $g \in G$, let $g\mathcal{F}$ denote the push-forward of \mathcal{F} along multiplication by g . Then there is a dense Zariski open subset U of G such that, for all k -rational points $g \in U$ and for all $j \geq 1$, the sheaf $\mathcal{T}or_j^X(g\mathcal{F}, \mathcal{E})$ is zero.*

As Miller and Speyer remark, their result is a homological generalization of the Kleiman–Bertini theorem: in characteristic 0, if $\mathcal{F} = \mathcal{O}_W$ and $\mathcal{E} = \mathcal{O}_Y$ are structure sheaves of smooth subvarieties of X and G acts transitively on X , then gW and Y meet transversally for generic g , implying that $\mathcal{O}_{gW} = g\mathcal{O}_W$ and \mathcal{O}_Y have no higher $\mathcal{T}or$. Motivated by this, if \mathcal{F} and \mathcal{E} are quasicoherent sheaves on X with $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E}) = 0$ for $j \geq 1$, we will say that \mathcal{F} and \mathcal{E} are *homologically transverse*; if $\mathcal{E} = \mathcal{O}_Y$ for some closed subscheme Y of X , we will simply say that \mathcal{F} and Y are homologically transverse.

MSC2000: primary 14L30; secondary 16S38.

Keywords: generic transversality, homological transversality, Kleiman’s theorem, group action.

Partially supported by NSF grants DMS-0502170 and DMS-0802935. This paper is part of the author’s Ph.D. thesis at the University of Michigan under the direction of J. T. Stafford.

Homological transversality has a geometric meaning if $\mathcal{F} = \mathcal{O}_W$ and $\mathcal{E} = \mathcal{O}_Y$ are structure sheaves of closed subschemes of X . If P is a component of $Y \cap W$, then Serre's formula for the multiplicity of the intersection of Y and W at P [Hartshorne 1977, p. 427] is

$$i(Y, W; P) = \sum_{j \geq 0} (-1)^j \text{len}_P(\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})),$$

where the length is taken over the local ring at P . Thus if Y and W are homologically transverse, their intersection multiplicity at P is simply the length of their scheme-theoretic intersection over the local ring at P .

It is natural to ask what conditions on the action of G are necessary to conclude that homological transversality is generic in the sense of Theorem 1.1. In particular, the restriction to transitive actions is unfortunately strong, as it excludes important situations such as the torus action on \mathbb{P}^n . On the other hand, suppose that \mathcal{F} is the structure sheaf of the closure of a nondense orbit. Then for all k -points $g \in G$, we have $\mathcal{T}or_1^X(g\mathcal{F}, \mathcal{F}) = \mathcal{T}or_1^X(\mathcal{F}, \mathcal{F}) \neq 0$, and so the conclusion of Theorem 1.1 fails (as long as $G(k)$ is dense in G). Thus for nontransitive group actions some additional hypothesis is necessary.

The main result of this paper is that there is a simple condition for homological transversality to be generic:

Theorem 1.2. *Let X be a scheme with a left action of a smooth algebraic group G , and let \mathcal{F} be a coherent sheaf on X . Let \bar{k} be an algebraic closure of k . Consider the following conditions:*

- (1) *For all closed points $x \in X \times \text{Spec } \bar{k}$, the pull-back of \mathcal{F} to $X \times \text{Spec } \bar{k}$ is homologically transverse to the closure of the $G(\bar{k})$ -orbit of x .*
- (2) *For all coherent sheaves \mathcal{E} on X , there is a Zariski open and dense subset U of G such that for all k -rational points $g \in U$, the sheaf $g\mathcal{F}$ is homologically transverse to \mathcal{E} .*

Then (1) implies (2). If k is algebraically closed, then (1) and (2) are equivalent.

If g is not k -rational, the sheaf $g\mathcal{F}$ can still be defined; in Section 2 we give this definition and a generalization of (2) that is equivalent to (1) in any setting (see Theorem 2.1).

If G acts transitively on X in the sense of [Miller and Speyer 2008], then the action is geometrically transitive, and so (1) is trivially satisfied. Thus Theorem 1.1 follows from Theorem 1.2. Since transversality of smooth subvarieties in characteristic 0 implies homological transversality, Theorem 1.2 also generalizes the following result of Robert Speiser:

Theorem 1.3 [Speiser 1988, Theorem 1.3]. *Suppose that k is algebraically closed of characteristic 0. Let X be a smooth variety, and let G be a (necessarily smooth) algebraic group acting on X . Let W be a smooth closed subvariety of X . If W is transverse to every G -orbit in X , then for any smooth closed subvariety $Y \subseteq X$, there is a dense open subset U of G such that if $g \in U$, then gW and Y are transverse.*

Speiser’s result implies that the generic intersection $gW \cap Y$, for $g \in U$, is also smooth. We also give a more general homological version of this. For simplicity, we state it here for algebraically closed fields, although in the body of the paper (see Theorem 4.2) we remove this assumption.

Theorem 1.4. *Assume that $k = \bar{k}$. Let X be a scheme with a left action of a smooth algebraic group G , and let W be a Cohen–Macaulay (alternatively, Gorenstein) closed subscheme of X such that W is homologically transverse to the G -orbit closure of every closed point $x \in X$. Then for any Cohen–Macaulay (Gorenstein) closed subscheme Y of X , there is a dense open subset $U \subseteq G$ so that gW is homologically transverse to Y and $gW \cap Y$ is Cohen–Macaulay (Gorenstein) for all closed points $g \in U$.*

Theorem 1.2 was proved in the course of an investigation of certain rings, determined by geometric data, that arise in the study of noncommutative algebraic geometry. Given a variety X , an automorphism σ of X and an invertible sheaf \mathcal{L} on X , Artin and Van den Bergh [1990] construct a *twisted homogeneous coordinate ring* $B = B(X, \mathcal{L}, \sigma)$. The graded ring B is defined via

$$B_n = H^0(X, \mathcal{L} \otimes_X \sigma^* \mathcal{L} \otimes_X \cdots \otimes_X (\sigma^{n-1})^* \mathcal{L})$$

with multiplication of sections given by the action of σ . A closed subscheme W of X determines a graded right ideal I of B , generated by sections vanishing on W . In [Sierra 2008], we study the *idealizer* of I ; that is, the maximal subring R of B such that I is a two-sided ideal of R . It turns out that quite subtle properties of W and its motion under σ control many of the properties of R ; in particular, for R to be left Noetherian one needs that for any closed subscheme Y , all but finitely many $\sigma^n W$ are homologically transverse to Y . (For details, we refer the reader to [Sierra 2008].) Thus we were naturally led to ask how often homological transversality can be considered “generic” behavior, and what conditions on W ensure this.

We make some remarks on notation. If x is any point of a scheme X , we denote the skyscraper sheaf at x by k_x . For schemes X and Y , we will write $X \times Y$ for the product $X \times_k Y$. Finally, if X is a scheme with a (left) action of an algebraic group G , we will always denote the multiplication map by

$$\mu : G \times X \rightarrow X.$$

2. Generalizations

We begin this section by defining homological transversality more generally. If W and Y are schemes over a scheme X , with (quasi)coherent sheaves \mathcal{F} on W and \mathcal{E} on Y respectively, then for all $j \geq 0$ there is a (quasi)coherent sheaf $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})$ on $W \times_X Y$. This sheaf is defined locally. Suppose that $X = \text{Spec } R$, $W = \text{Spec } S$, and $Y = \text{Spec } T$ are affine. Let $(_)^\sim$ denote the functor that takes an R -module (or S - or T -module, respectively) to the associated quasicoherent sheaf on X (or W or Y). If F is an S -module and E is a T -module, we define $\mathcal{T}or_j^X(\tilde{F}, \tilde{E})$ to be $(\text{Tor}_j^R(F, E))^\sim$. That these glue properly to give sheaves on $W \times_X Y$ for general W , Y , and X is the result in [Grothendieck 1963, 6.5.3]. As before, we will say that \mathcal{F} and \mathcal{E} are *homologically transverse* if the sheaf $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})$ is zero for all $j \geq 1$.

We caution the reader that the maps from W and Y to X are implicit in the definition of $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})$; at times we will write $\mathcal{T}or_j^{W \rightarrow X \leftarrow Y}(\mathcal{F}, \mathcal{E})$ to make this more obvious. We also remark that if $Y = X$, then $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})$ is a sheaf on $W \times_X X = W$. As localization commutes with Tor , for any $w \in W$ lying over $x \in X$ we have in this case that $\mathcal{T}or_j^X(\mathcal{F}, \mathcal{E})_w = \text{Tor}_j^{O_{X,x}}(\mathcal{F}_w, \mathcal{E}_x)$.

Now suppose that $f : W \rightarrow X$ is a morphism of schemes and G is an algebraic group acting on X . Let \mathcal{F} be a (quasi)coherent sheaf on W and let g be any point of G . We will denote the pull-back of \mathcal{F} to $\{g\} \times W$ by $g\mathcal{F}$. There is a map

$$\{g\} \times W \longrightarrow G \times W \xrightarrow{1 \times f} G \times X \xrightarrow{\mu} X.$$

If Y is a scheme over X and \mathcal{E} is a (quasi)coherent sheaf on Y , the (quasi)coherent sheaf $\mathcal{T}or_j^{\{g\} \times W \rightarrow X \leftarrow Y}(g\mathcal{F}, \mathcal{E})$ on $W \times_X Y \times k(g)$ will be written $\mathcal{T}or_j^X(g\mathcal{F}, \mathcal{E})$. Note that if $W = X$ and g is k -rational, then $g\mathcal{F}$ is simply the push-forward of \mathcal{F} along multiplication by g .

In this context, we prove the following relative version of Theorem 1.2:

Theorem 2.1. *Let X be a scheme with a left action of a smooth algebraic group G , let $f : W \rightarrow X$ be a morphism of schemes, and let \mathcal{F} be a coherent sheaf on W . We define maps*

$$\begin{array}{ccc} G \times W & \xrightarrow{\rho} & X \\ p \downarrow & & \\ & & W, \end{array}$$

where ρ is the map $\rho(g, w) = gf(w)$ induced by the action of G and p is projection onto the second factor.

The following conditions are equivalent:

- (1) For all closed points $x \in X \times \text{Spec } \bar{k}$, the pull-back of \mathcal{F} to $W \times \text{Spec } \bar{k}$ is homologically transverse to the closure of the $G(\bar{k})$ -orbit of x .

- (2) For all schemes $r : Y \rightarrow X$ and all coherent sheaves \mathcal{E} on Y , there is a Zariski open and dense subset U of G such that for all closed points $g \in U$, the sheaf $g^*\mathcal{F}$ on $\{g\} \times W$ is homologically transverse to \mathcal{E} .
- (3) The sheaf $p^*\mathcal{F}$ on $G \times W$ is ρ -flat over X .

A related relative version of Theorem 1.3 is given in [Speiser 1988].

Our general approach to Theorem 2.1 mirrors that of [Speiser 1988], although the proof techniques are quite different. We first generalize Theorem 1.1 to apply to any flat map $f : W \rightarrow X$; this is a homological version of [Kleiman 1974, Lemma 1] and may be of independent interest.

Theorem 2.2. *Let X, Y , and W be schemes, let A be a generically reduced scheme, and suppose that there are morphisms*

$$\begin{array}{ccc}
 & & Y \\
 & & \downarrow r \\
 W & \xrightarrow{f} & X \\
 q \downarrow & & \\
 & & A.
 \end{array}$$

Let \mathcal{F} be a coherent sheaf on W that is f -flat over X , and let \mathcal{E} be a coherent sheaf on Y . For all $a \in A$, let W_a denote the fiber of W over a , and let $\mathcal{F}_a = \mathcal{F} \otimes_W \mathbb{O}_{W_a}$ be the fiber of \mathcal{F} over a .

Then there is a dense open $U \subseteq A$ such that if $a \in U$, then \mathcal{F}_a is homologically transverse to \mathcal{E} .

We note that we have not assumed that X, Y, W , or A are smooth.

3. Proofs

In this section we prove Theorems 1.2, 2.1, and 2.2.

Lemma 3.1. *Let*

$$X_1 \xrightarrow{\alpha} X_2 \xrightarrow{\gamma} X_3$$

be morphisms of schemes, and assume that γ is flat. Let \mathcal{G} be a quasicohherent sheaf on X_1 that is flat over X_3 . Let \mathcal{H} be any quasicohherent sheaf on X_3 . Then for all $j \geq 1$, we have $\mathcal{T}or_j^{X_2}(\mathcal{G}, \gamma^*\mathcal{H}) = 0$.

Proof. We may reduce to the local case. Thus let $x \in X_1$ and let $y = \alpha(x)$ and $z = \gamma(y)$. Let $S = \mathbb{O}_{X_2, y}$ and let $R = \mathbb{O}_{X_3, z}$. Then $(\gamma^*\mathcal{H})_y \cong S \otimes_R \mathcal{H}_z$. Since S is flat over R , we have

$$\mathcal{T}or_j^R(\mathcal{G}_x, \mathcal{H}_z) \cong \mathcal{T}or_j^S(\mathcal{G}_x, S \otimes_R \mathcal{H}_z) = \mathcal{T}or_j^{X_2}(\mathcal{G}, \gamma^*\mathcal{H})_x$$

by flat base change. The left-hand side is 0 for $j \geq 1$ since \mathcal{G} is flat over X_3 . Thus for $j \geq 1$ we have $\mathcal{T}or_j^{X_2}(\mathcal{G}, \gamma^*\mathcal{L}) = 0$. \square

To prove Theorem 2.2, we show that a suitable modification of the spectral sequences used by Miller and Speyer will work in our situation. Our key computation is the following lemma; compare to [Miller and Speyer 2008, Proposition 2].

Lemma 3.2. *Given the notation of Theorem 2.2, there is an open dense $U \subseteq A$ such that for all $a \in U$ and for all $j \geq 0$ we have*

$$\mathcal{T}or_j^W(\mathcal{F} \otimes_X \mathcal{E}, q^*k_a) \cong \mathcal{T}or_j^X(\mathcal{F}_a, \mathcal{E})$$

as sheaves on $W \times_X Y$.

Note that $\mathcal{F} \otimes_X \mathcal{E}$ is a sheaf on $W \times_X Y$ and thus $\mathcal{T}or_j^W(\mathcal{F} \otimes_X \mathcal{E}, q^*k_a)$ is a sheaf on $W \times_X Y \times_W W = W \times_X Y$ as required.

Proof. Since A is generically reduced, we may apply generic flatness to the morphism $q : W \rightarrow A$. Thus there is an open dense subset U of A such that both W and \mathcal{F} are flat over U . Let $a \in U$. Away from $q^{-1}(U)$, both sides of the equality we seek to establish are zero, and so the result is trivial. Since $\mathcal{F}|_{q^{-1}(U)}$ is still flat over X , without loss of generality we may replace W by $q^{-1}(U)$; that is, we may assume that both W and \mathcal{F} are flat over A .

The question is local, so assume that $X = \text{Spec } R, Y = \text{Spec } T$, and $W = \text{Spec } S$ are affine. Let $E = \Gamma(Y, \mathcal{E})$ and let $F = \Gamma(W, \mathcal{F})$. Let $Q = \Gamma(W, q^*k_a)$; then $\Gamma(W, \mathcal{F}_a) = F \otimes_S Q$. We seek to show that

$$\text{Tor}_j^S(F \otimes_R E, Q) \cong \text{Tor}_j^R(F \otimes_S Q, E)$$

as $S \otimes_R T$ -modules.

We will work on $W \times X$. For clarity, we lay out the various morphisms and corresponding ring maps in our situation. We have morphisms of schemes

$$\begin{array}{ccc} W \times X & & Y \\ p \downarrow & \searrow \phi & \downarrow r \\ W & \xrightarrow{f} & X, \end{array}$$

where p is projection onto the first factor and the morphism ϕ splitting p is given by the graph of f . Letting $B = S \otimes_k R$, we have corresponding maps of rings

$$\begin{array}{ccc} B & & T \\ p^\# \uparrow & \searrow \phi^\# & \uparrow r^\# \\ S & \xleftarrow{f^\#} & R, \end{array}$$

where $p^\#(s) = s \otimes 1$ and $\phi^\#(s \otimes r) = s \cdot f^\#(r)$. We make the trivial observation that

$$B \otimes_R E = (S \otimes_k R) \otimes_R E \cong S \otimes_k E.$$

Let $K_\bullet \rightarrow F$ be a projective resolution of F , considered as a B -module via the map $\phi^\# : B \rightarrow S$. As E is an R -module via the map $r^\# : R \rightarrow T$, there is a B -action on $S \otimes_k E$; let $L_\bullet \rightarrow S \otimes_k E$ be a projective resolution over B .

Let $P_{\bullet,\bullet}$ be the double complex $K_\bullet \otimes_B L_\bullet$. We claim the total complex of $P_{\bullet,\bullet}$ resolves $F \otimes_B (S \otimes_k E)$. To see this, note that the rows of $P_{\bullet,\bullet}$, which are of the form $K_\bullet \otimes_B L_j$, are acyclic, except in degree 0, where the homology is $F \otimes_B L_j$. The degree 0 horizontal homology forms a vertical complex whose homology computes $\text{Tor}_j^B(F, S \otimes_k E)$. But $S \otimes_k E \cong B \otimes_R E$, and B is a flat R -module. Therefore $\text{Tor}_j^B(F, S \otimes_k E) \cong \text{Tor}_j^B(F, B \otimes_R E) \cong \text{Tor}_j^R(F, E)$ by the formula for flat base change for Tor. Since F is flat over R , this is zero for all $j \geq 1$. Thus, via the spectral sequence

$$H_j^v(H_i^h P_{\bullet,\bullet}) \Rightarrow H_{i+j} \text{Tot } P_{\bullet,\bullet}$$

we see that the total complex of $P_{\bullet,\bullet}$ is acyclic, except in degree 0, where the homology is $F \otimes_B S \otimes_k E \cong F \otimes_R E$.

Consider the double complex $P_{\bullet,\bullet} \otimes_S Q$. Since $\text{Tot } P_{\bullet,\bullet}$ is a B -projective and therefore S -projective resolution of $F \otimes_R E$, the homology of the total complex of this double complex computes $\text{Tor}_j^S(F \otimes_R E, Q)$.

Now consider the row $K_\bullet \otimes_B L_j \otimes_S Q$. As L_j is B -projective and therefore B -flat, the i th homology of this row is isomorphic to $\text{Tor}_i^S(F, Q) \otimes_B L_j$. Since W and \mathcal{F} are flat over A , by Lemma 3.1 we have $\text{Tor}_i^S(F, Q) = 0$ for all $i \geq 1$. Thus this row is acyclic except in degree 0, where the homology is $F \otimes_B L_j \otimes_S Q$. The vertical differentials on the degree 0 homology give a complex whose j th homology is isomorphic to $\text{Tor}_j^B(F \otimes_S Q, S \otimes_k E)$. As before, this is simply $\text{Tor}_j^R(F \otimes_S Q, E)$.

Thus (via a spectral sequence) we see that the homology of the total complex of $P_{\bullet,\bullet} \otimes_S Q$ computes $\text{Tor}_j^R(F \otimes_S Q, E)$. But we have already seen that the homology of this total complex is isomorphic to $\text{Tor}_j^S(F \otimes_R E, Q)$. Thus the two are isomorphic. □

Proof of Theorem 2.2. By generic flatness, there is no loss of generality in restricting to W flat over A . Since \mathcal{F} and \mathcal{E} are coherent sheaves on W and Y respectively, $\mathcal{F} \otimes_X \mathcal{E}$ is a coherent sheaf on $W \times_X Y$. Applying generic flatness to the composition $W \times_X Y \rightarrow W \rightarrow A$, we obtain a dense open $V \subseteq A$ such that $\mathcal{F} \otimes_X \mathcal{E}$ is flat over V . Therefore, by Lemma 3.1, if $a \in V$ and $j \geq 1$, we have $\text{Tor}_j^W(\mathcal{F} \otimes_X \mathcal{E}, q^*k_a) = 0$.

We apply Lemma 3.2 to choose a dense open $U \subseteq A$ such that for all $j \geq 1$, if $a \in U$, then $\text{Tor}_j^W(\mathcal{F} \otimes_X \mathcal{E}, q^*k_a) \cong \text{Tor}_j^X(\mathcal{F}_a, \mathcal{E})$. Thus if a is in the dense open

set $U \cap V$, then for all $j \geq 1$ we have

$$\mathcal{T}or_j^X(\mathcal{F}_a, \mathcal{E}) \cong \mathcal{T}or_j^W(\mathcal{F} \otimes_X \mathcal{E}, q^*k_a) = 0,$$

as required. □

We now turn to the proof of Theorem 2.1; for the remainder of this paper, we will adopt the hypotheses and notation given there.

Lemma 3.3. *Let $R, R', S,$ and T be commutative rings, and let*

$$\begin{array}{ccc} R' & \longrightarrow & T \\ \uparrow & & \uparrow \\ R & \longrightarrow & S \end{array}$$

be a commutative diagram of ring homomorphisms such that R'_R and T_S are flat. Let N be an R -module. Then for all $j \geq 0$, we have that

$$\mathcal{T}or_j^{R'}(N \otimes_R R', T) \cong \mathcal{T}or_j^R(N, S) \otimes_S T.$$

Proof. Let $P_\bullet \rightarrow N$ be a projective resolution of N . Consider the complex

$$P_\bullet \otimes_R R' \otimes_{R'} T \cong P_\bullet \otimes_R T \cong P_\bullet \otimes_R S \otimes_S T. \tag{3.4}$$

Since R'_R is flat, $P_\bullet \otimes_R R'$ is a projective resolution of $N \otimes_R R'$. Thus the j th homology of (3.4) computes $\mathcal{T}or_j^{R'}(N \otimes_R R', T)$. Since T_S is flat, this homology is isomorphic to $H_j(P_\bullet \otimes_R S) \otimes_S T$. Thus $\mathcal{T}or_j^{R'}(N \otimes_R R', T) \cong \mathcal{T}or_j^R(N, S) \otimes_S T$. □

Lemma 3.5. *Let x be a closed point of X . Consider the multiplication map*

$$\mu_x : G \times \{x\} \rightarrow X.$$

Then for all $j \geq 0$ we have

$$\mathcal{T}or_j^X(\mathcal{F}, \mathcal{O}_{G \times \{x\}}) \cong \mathcal{T}or_j^{G \times X}(p^*\mathcal{F}, \mu^*k_x). \tag{3.6}$$

If k is algebraically closed, then we also have

$$\mathcal{T}or_j^{G \times X}(p^*\mathcal{F}, \mu^*k_x) \cong \mathcal{T}or_j^X(\mathcal{F}, \mathcal{O}_{\overline{Gx}}) \otimes_X \mathcal{O}_{G \times \{x\}}. \tag{3.7}$$

All isomorphisms are of sheaves on $G \times W$.

Proof. Note that μ_x maps $G \times \{x\}$ onto a locally closed subscheme of X , which we will denote Gx . Since all computations may be done locally, without loss of generality we may assume that Gx is in fact a closed subscheme of X .

Let $\nu : G \rightarrow G$ be the inverse map, and let $\psi = \nu \times \mu : G \times X \rightarrow G \times X$. Consider the commutative diagram

$$\begin{array}{ccccc}
 G \times W & \xrightarrow{1 \times f} & G \times X & \xleftarrow{\psi} & G \times \{x\} \\
 p \downarrow & & p \downarrow & \swarrow \mu_x & \downarrow \pi \\
 W & \xrightarrow{f} & X & \xleftarrow{\cong} & Gx,
 \end{array} \tag{3.8}$$

where π is the induced map and p is projection onto the second factor. Since $\psi^2 = \text{Id}_{G \times X}$ and $\mu = p \circ \psi$, we obtain $\mu^*k_x \cong \psi^*p^*k_x \cong \psi_*\mathcal{O}_{G \times \{x\}}$, considered as sheaves on $G \times X$. Then the isomorphism (3.6) is a direct consequence of the flatness of p and Lemma 3.3. If k is algebraically closed, then π is also flat, and so the isomorphism (3.7) also follows from Lemma 3.3. \square

Proof of Theorem 2.1. (3) \Rightarrow (2). Assume (3). Let \mathcal{E} be a coherent sheaf on Y . Consider the maps

$$\begin{array}{ccc}
 & & Y \\
 & & \downarrow r \\
 G \times W & \xrightarrow{\rho} & X \\
 q \downarrow & & \\
 G, & &
 \end{array}$$

where q is projection on the first factor.

Since G is smooth, it is generically reduced. Thus we may apply Theorem 2.2 to the ρ -flat sheaf $p^*\mathcal{F}$ to obtain a dense open $U \subseteq G$ such that if $g \in U$ is a closed point, then ρ makes $(p^*\mathcal{F})_g$ homologically transverse to \mathcal{E} . But $\rho|_{\{g\} \times W}$ is the map used to define $\mathcal{T}or_j^X(g\mathcal{F}, \mathcal{E})$; that is, $(p^*\mathcal{F})_g \cong g\mathcal{F}$ as sheaves over X . Thus (2) holds.

(2) \Rightarrow (3). The morphism ρ factors as

$$G \times W \xrightarrow{1 \times f} G \times X \xrightarrow{\mu} X.$$

Since the multiplication map μ is the composition of an automorphism of $G \times X$ and a projection, it is flat.

Therefore for any quasicohherent \mathcal{N} on X and \mathcal{M} on $G \times W$ and for any closed point $z \in G \times W$, we have

$$\mathcal{T}or_j^{G \times X}(\mathcal{M}, \mu^*\mathcal{N})_z \cong \mathcal{T}or_j^{\mathcal{O}_{X, \rho(z)}}(\mathcal{M}_z, \mathcal{N}_{\rho(z)}), \tag{3.9}$$

as in the proof of Lemma 3.1.

If $p^*\mathcal{F}$ fails to be flat over X , then flatness fails against the structure sheaf of some closed point $x \in X$, by the local criterion for flatness [Eisenbud 1995,

Theorem 6.8]. Thus to check that $p^*\mathcal{F}$ is flat over X , it is equivalent to test flatness against structure sheaves of closed points of X . By (3.9), we see that $p^*\mathcal{F}$ is ρ -flat over X if and only if

$$\mathcal{T}or_j^{G \times X}(p^*\mathcal{F}, \mu^*k_x) = 0 \quad \text{for all closed points } x \in X \text{ and for all } j \geq 1. \quad (3.10)$$

Applying Lemma 3.5, we see that the flatness of $p^*\mathcal{F}$ is equivalent to the vanishing

$$\mathcal{T}or_j^X(\mathcal{F}, \mathbb{O}_{G \times \{x\}}) = 0 \quad \text{for all closed points } x \in X \text{ and for all } j \geq 1. \quad (3.11)$$

Assume (2). We will show that (3.11) holds for all $x \in X$. Fix a closed point $x \in X$ and consider the morphism $\mu_x : G \times \{x\} \rightarrow X$. By assumption, there is a closed point $g \in G$ such that $g^*\mathcal{F}$ is homologically transverse to $\mathbb{O}_{G \times \{x\}}$. Let $k' = k(g)$ and let g' be the canonical k' -point of $G \times \text{Spec } k'$ lying over g . Let $G' = G \times \text{Spec } k'$ and let $X' = X \times \text{Spec } k'$. Let \mathcal{F}' be the pull-back of \mathcal{F} to $W' = W \times \text{Spec } k'$. Consider the commutative diagram

$$\begin{array}{ccccc} G \times \{x\} \times \text{Spec } k' & \xrightarrow{\mu_x \times 1} & X' & \xleftarrow{\rho} & \{g'\} \times_{k'} W' \\ \downarrow & & \downarrow & & \cong \downarrow \\ G \times \{x\} & \xrightarrow{\mu_x} & X & \xleftarrow{\rho} & \{g\} \times W. \end{array}$$

Since the vertical maps are faithfully flat and the left-hand square is a fiber square, by Lemma 3.3 $g'^*\mathcal{F}'$ is homologically transverse to

$$G \times \{x\} \times \text{Spec } k' \cong G' \times \{x\}.$$

By $G(k')$ -equivariance, \mathcal{F}' is homologically transverse to $(g')^{-1}G' \times \{x\} = G' \times \{x\}$. Since

$$G' \times \{x\} \longrightarrow X' \xleftarrow{f} W'$$

is base-extended from

$$G \times \{x\} \longrightarrow X \xleftarrow{f} W,$$

we obtain that \mathcal{F} is homologically transverse to $G \times \{x\}$. Thus (3.11) holds.

(1) \Rightarrow (3). The ρ -flatness of \mathcal{F} is not affected by base extension, so without loss of generality we may assume that k is algebraically closed. Then (3) follows directly from Lemma 3.5 and the criterion (3.10) for flatness.

(3) \Rightarrow (1). As before, we may assume that k is algebraically closed. Let x be a closed point of X . We have seen that (3) and (2) are equivalent; by applying (2) to $\mathcal{E} = \overline{\mathbb{O}_{Gx}}$ there is a closed point $g \in G$ such that $g^*\mathcal{F}$ and \overline{Gx} are homologically transverse. By $G(k)$ -equivariance, \mathcal{F} and $g^{-1}\overline{Gx} = \overline{Gx}$ are homologically transverse. □

Proof of Theorem 1.2. If \mathcal{F} is homologically transverse to orbit closures upon extension to \bar{k} , then, using Theorem 2.1(2), for any \mathcal{E} there is a dense open $U \subseteq G$ such that, in particular, for any k -rational $g \in U$ the sheaves $g\mathcal{F}$ and \mathcal{E} are homologically transverse.

The equivalence of (1) and (2) in the case that k is algebraically closed follows directly from Theorem 2.1. \square

Theorem 1.2 is a statement about k -rational points in $U \subseteq G$. However, the proof shows that for any extension k' of k and any k' -rational $g \in U \times \text{Spec } k'$, $g\mathcal{F}$ will be homologically transverse to \mathcal{E} on $X \times \text{Spec } k'$. Further, in many situations U will automatically contain a k -rational point of G . This holds, in particular, if k is infinite, G is connected and affine, and either k is perfect or G is reductive, by [Borel 1991, Corollary 18.3].

4. Singularities of generic intersections

We now specialize to consider generic intersections of two subschemes of X . That is, let X be a scheme with a left action of a smooth algebraic group G . Let Y and W be closed subschemes of X . By Theorem 1.3, if k is algebraically closed of characteristic 0, W is transverse to G -orbit closures, and X , Y , and W are smooth, then for generic $g \in G$ the subschemes gW and Y meet transversally, and so by definition $gW \cap Y$ is smooth. Here we remark that a homological version of this result holds more generally: if W is homologically transverse to G -orbit closures and Y and W are both Cohen–Macaulay or both Gorenstein, their generic intersection has the same property. We use the following result from commutative algebra:

Theorem 4.1. *Let $A \rightarrow B$ be a local homomorphism of Noetherian local rings, and let \mathfrak{m} be the maximal ideal of A and $F = B/\mathfrak{m}B$. Assume that B is flat over A . Then B is Cohen–Macaulay (alternatively, Gorenstein) if and only if B and F are both Cohen–Macaulay (Gorenstein).*

Proof. See [Matsumura 1989, Corollary 23.3, Theorem 23.4]. \square

Theorem 4.2. *Let X be a scheme with a left action of a smooth algebraic group G . Suppose that $f : W \rightarrow X$ and $r : Y \rightarrow X$ are morphisms of schemes and that $W \times \text{Spec } \bar{k}$ is homologically transverse to the $G(\bar{k})$ -orbit of x for all closed points $x \in X \times \text{Spec } \bar{k}$. Further suppose that Y and W are Cohen–Macaulay (alternatively, Gorenstein). Then there is a dense open subset $U \subseteq G$ so that for all closed points $g \in U$, the scheme $\{g\} \times W$ is homologically transverse to Y and the fiber product $(\{g\} \times W) \times_X Y$ is Cohen–Macaulay (Gorenstein).*

Proof. Let $\rho : G \times W \rightarrow X$ be the map $\rho(g, w) = gf(w)$ induced by f and the action of G . Let $q : G \times W \rightarrow G$ be projection to the first factor. Thus there is a

commutative diagram

$$\begin{array}{ccc}
 G \times W \times_X Y & \xrightarrow{\rho \times 1} & Y \\
 1 \times r \downarrow & & \downarrow r \\
 G \times W & \xrightarrow{\rho} & X \\
 q \downarrow & & \\
 G & &
 \end{array}$$

By Theorem 2.1 applied to $\mathcal{F} = \mathbb{O}_W$, ρ is flat. Now, $G \times W$ is Cohen–Macaulay or Gorenstein, and so by Theorem 4.1, the fibers of ρ have the same property. Since Y is Cohen–Macaulay (Gorenstein) and $\rho \times 1$ is flat, applying Theorem 4.1 again, we see that $G \times W \times_X Y$ is also Cohen–Macaulay (Gorenstein). Now, by generic flatness and Theorem 2.1, there is a dense open $U \subset G$ such that $q \circ (1 \times r)$ is flat over U and $\{g\} \times W$ is homologically transverse to Y for all $g \in U$. For $g \in U$, the fiber $(\{g\} \times W) \times_X Y$ of $q \circ (1 \times r)$ is Cohen–Macaulay (Gorenstein), by Theorem 4.1 again. □

We note that, although we did not assume that X is Cohen–Macaulay or Gorenstein, it follows from the flatness of ρ and from Theorem 4.1.

We also remark that if Y and W are homologically transverse local complete intersections in a smooth X , it is not hard to show directly that $Y \cap W$ is also a local complete intersection. We do not know if it is true in general that the homologically transverse intersection of two Cohen–Macaulay subschemes is Cohen–Macaulay, although it follows, for example, from [Fulton and Pragacz 1998, Lemma, p. 108] if X is smooth.

Theorem 1.4 follows directly from Theorem 4.2.

Thus we may refine Theorem 1.1 to obtain a result on transitive group actions that echoes the Kleiman–Bertini theorem even more closely.

Corollary 4.3. *Let X be a scheme with a geometrically transitive left action of a smooth algebraic group G . Let Y and W be Cohen–Macaulay (alternatively, Gorenstein) closed subschemes of X . Then there is a dense Zariski open subset U of G such that gW is homologically transverse to Y and $gW \cap Y$ is Cohen–Macaulay (Gorenstein) for all k -rational points $g \in U$.* □

Acknowledgments

I am grateful to Ezra Miller for his extraordinarily careful reading of an earlier version of this paper and for several corrections and discussions, to David Speyer for many informative conversations, and to Mel Hochster, Kyle Hofmann, Gopal Prasad, and Karen Smith for their suggestions and assistance with references. I particularly thank Brian Conrad for finding an error in an earlier version of this

paper and for several helpful discussions. I also thank Susan Colley and Gary Kennedy for calling my attention to [Speiser 1988].

References

- [Artin and Van den Bergh 1990] M. Artin and M. Van den Bergh, “Twisted homogeneous coordinate rings”, *J. Algebra* **133**:2 (1990), 249–271. MR 91k:14003 Zbl 0717.14001
- [Borel 1991] A. Borel, *Linear algebraic groups*, 2nd ed., Graduate Texts in Mathematics **126**, Springer, New York, 1991. MR 92d:20001 Zbl 0726.20030
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer, New York, 1995. MR 97a:13001 Zbl 0819.13001
- [Fulton and Pragacz 1998] W. Fulton and P. Pragacz, *Schubert varieties and degeneracy loci*, Lecture Notes in Mathematics **1689**, Springer, Berlin, 1998. MR 99m:14092 Zbl 0913.14016
- [Grothendieck 1963] A. Grothendieck, “Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. II”, *Inst. Hautes Études Sci. Publ. Math.* **17** (1963), 91. MR 29 #1210 Zbl 0122.16102
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, New York, 1977. MR 57 #3116 Zbl 0531.14001
- [Kleiman 1974] S. L. Kleiman, “The transversality of a general translate”, *Compositio Math.* **28** (1974), 287–297. MR 50 #13063 Zbl 0288.14014
- [Matsumura 1989] H. Matsumura, *Commutative ring theory*, 2nd ed., Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, 1989. MR 90i:13001 Zbl 0603.13001
- [Miller and Speyer 2008] E. Miller and D. E. Speyer, “A Kleiman–Bertini theorem for sheaf tensor products”, *J. Algebraic Geom.* **17**:2 (2008), 335–340. MR 2008k:14044 Zbl 1138.14011 arXiv math.AG/0601202
- [Sierra 2008] S. J. Sierra, “Geometric idealizers”, preprint, 2008. To appear in *Trans. Amer. Math. Soc.* arXiv 0809.3971
- [Speiser 1988] R. Speiser, “Transversality theorems for families of maps”, pp. 235–252 in *Algebraic geometry (Sundance, UT, 1986)*, edited by A. Holme and R. Speiser, Lecture Notes in Math. **1311**, Springer, Berlin, 1988. MR 89k:14005 Zbl 0683.14003

Communicated by David Eisenbud

Received 2009-03-09 Accepted 2009-07-21

sjsierra@math.washington.edu *Department of Mathematics, University of Washington,
Seattle, WA 98195, United States*
<http://www.math.washington.edu/~sjsierra/>

