

# *Algebra & Number Theory*

Volume 7

2013

No. 2

**Base change behavior of the relative canonical sheaf  
related to higher dimensional moduli**

Zsolt Patakfalvi



# Base change behavior of the relative canonical sheaf related to higher dimensional moduli

Zsolt Patakfalvi

We show that the compatibility of the relative canonical sheaf with base change fails generally in families of normal varieties. Furthermore, it always fails if the general fiber of a family of pure dimension  $n$  is Cohen–Macaulay and the special fiber contains a strictly  $S_{n-1}$  point. In particular, in moduli spaces with functorial relative canonical sheaves Cohen–Macaulay schemes can not degenerate to  $S_{n-1}$  schemes. Another, less immediate consequence is that the canonical sheaf of an  $S_{n-1}$ ,  $G_2$  scheme of pure dimension  $n$  is not  $S_3$ .

## 1. Introduction

The canonical sheaf plays a crucial role in the classification of varieties of characteristic zero. Global sections of its powers define the canonical map, which is birational onto its image for varieties of general type with mild singularities. The image is called the canonical model, and it is a unique representative of the birational equivalence class of the original variety. In particular, the canonical model can be used to construct a moduli space that classifies varieties of general type up to birational equivalence. This moduli space  $\overline{\mathcal{M}}_h$  of stable schemes is the higher dimensional generalization of the intensely investigated space  $\overline{\mathcal{M}}_g$  of stable curves. In order to build  $\overline{\mathcal{M}}_h$ , it is important to understand when the canonical sheaf behaves functorially in families, that is, when it is compatible with base change.

More precisely, to obtain a compact moduli space, in  $\overline{\mathcal{M}}_h$ , not only canonical models are allowed, but also their generalizations, the semi-log canonical models [Kollár 2010, Definition 15]. By definition these are projective schemes with semi-log canonical singularities [Hacon and Kovács 2010, Definition 3.13.5] and ample canonical bundles. The first naive definition of the moduli functor of stable schemes

---

*MSC2010:* primary 14J10; secondary 14D06, 14F10, 14E30.

*Keywords:* canonical sheaf, relative canonical sheaf, dualizing complex, relative dualizing complex, base change, depth, moduli of stable varieties.

with Hilbert function  $h$  is then as follows. Here  $h : \mathbb{Z} \rightarrow \mathbb{Z}$  is an arbitrary function.

$$\overline{\mathfrak{M}}_h(B) = \left\{ f : X \rightarrow B \left| \begin{array}{l} f \text{ is flat, proper, } X_{\bar{b}} \text{ is a semi-} \\ \text{log canonical model } (\forall b \in B), \\ h(m) = \chi(\omega_{X_b}^{[m]}) \text{ } (\forall m \in \mathbb{Z}, b \in B) \end{array} \right. \right\} / \cong \text{ over } B \quad (1.0.a)$$

As usual, the naive definition works only in the naive cases but not in general. More precisely, (1.0.a) is insufficient to prove the existence of a projective coarse moduli space or a proper Deligne–Mumford stack structure on  $\overline{\mathfrak{M}}_h$ ; see [Kollár 2008; 2010]. In general, (1.0.a) has to be complemented with

$$\omega_{X/B}^{[m]}|_{X_b} \cong \omega_{X_b}^{[m]} \quad \text{for every integer } m \text{ and } b \in B. \quad (1.0.b)$$

Usually (1.0.b) is referred to as *Kollár’s condition* (for instance, in [Hassett and Kovács 2004, page 238]). Note also that (1.0.b) is not necessary for reduced  $B$ , but it does add important extra restrictions when  $B$  is nonreduced.

Currently, it is not understood in every aspect why and how deeply this condition is needed. For example it is not known if in characteristic zero it is equivalent or not to the other possible choice, called *Viehweg’s condition* (see [Viehweg 1995, Assumption 8.30; Hassett and Kovács 2004, page 238]):

$$\text{There is an integer } m \text{ such that } \omega_{X/B}^{[m]} \text{ is a line bundle.} \quad (1.0.c)$$

The starting point of this article is the  $m = 1$  case of (1.0.b), that is, the compatibility of the relative canonical sheaf with base change. We will try to understand how restrictive this condition is on flat families. The results will also yield statements about how Serre’s  $S_n$  condition behaves in families and for the canonical sheaves of single schemes.

Recently it has been proven in [Kollár and Kovács 2010, Theorem 7.9.3] that the relative canonical sheaf of flat families of projective schemes (over  $\mathbb{C}$ ) with Du Bois fibers is compatible with base change. According to [Kollár and Kovács 2010, Theorem 1.4] this pertains to families with semi-log canonical fibers as well. Furthermore, compatibility holds whenever the fibers are Cohen–Macaulay [Conrad 2000, Theorem 3.6.1].

It is important to note at this point that the  $m = 1$  case of (1.0.b) behaves differently than the rest. For  $m > 1$  there are examples of families of normal surfaces for which (1.0.b) does not hold; see [Hacon and Kovács 2010, Section 14.A]. However, since normal surfaces are Cohen–Macaulay, condition (1.0.b) with  $m = 1$  holds for every flat family of normal surfaces. Hence, any incompatibility can be observed only in higher dimensions. Partly due to this fact, there has been a common misbelief, sometimes even stated in articles, that the relative canonical sheaf is compatible with base change for flat families of normal varieties. The question if this compatibility

holds indeed was asked about the same time independently by János Kollár and the author.

**Question 1.1** (Kollár). Is  $\omega_{X/B}|_{X_b} \cong \omega_{X_b}$  for every flat family  $X \rightarrow B$  of normal varieties?

Here we construct examples showing that the answer is no. That is, there are flat families of normal varieties over smooth curves such that the relative canonical sheaves are not compatible with base change. The examples also show that the known results are optimal in many senses. That is, the fibers of the given families can be chosen to be  $S_j$  for any  $n > j \geq 2$  and their relative canonical sheaves to be  $\mathbb{Q}$ -line bundles. The precise statement is as follows.

**Theorem 1.2.** *For each  $n \geq 3$  and  $n > j \geq 2$  there is a flat family  $\mathcal{H} \rightarrow B$  of  $S_j$  (but not  $S_{j+1}$ ) normal varieties of dimension  $n$  over some open set  $B \subseteq \mathbb{P}^1$ , with  $\omega_{\mathcal{H}/B}$  a  $\mathbb{Q}$ -line bundle, such that*

$$\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \not\cong \omega_{\mathcal{H}_0}, \tag{1.2.a}$$

(Here  $\mathcal{H}_0$  is the central fiber of  $\mathcal{H}$ .)

Moreover, the general fiber of  $\mathcal{H}$  can be chosen to be smooth and the central fiber to have only one singular point.

When  $j = n - 1$  and the general fiber is Cohen–Macaulay, somewhat surprisingly, the incompatibility of (1.2.a) always holds. Furthermore, one can allow  $S_{n-1}$  points also in the general fibers provided the relative  $S_{n-1}$  locus has a component in the central fiber. The precise statement is as follows. (See Section 2 for the assumptions of the article, for instance, scheme is always separated and of finite type over  $k = \bar{k}$ , etc.)

**Theorem 1.3.** *If  $f : \mathcal{H} \rightarrow B$  is a flat family of schemes of pure dimension  $n$  over a smooth curve, such that a component of the locus*

$$\overline{\{x \in \mathcal{H} \mid x \text{ is closed, } \text{depth } \mathbb{O}_{\mathcal{H}(x),x} = n - 1\}} \tag{1.3.a}$$

*is contained in the special fiber  $\mathcal{H}_0$ , then the restriction homomorphism  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is not an isomorphism.*

In particular, the contrapositive of Theorem 1.3 when the general fiber is Cohen–Macaulay yields the following corollary.

**Corollary 1.4.** *If  $f : \mathcal{H} \rightarrow B$  is a flat family of schemes of pure dimension  $n$  such that  $\omega_{\mathcal{H}/B}$  is compatible with base change and the general fiber of  $f$  is Cohen–Macaulay, then the central fiber of  $f$  cannot have a closed point  $x$  such that  $\text{depth } \mathbb{O}_{\mathcal{H}(x),x} = n - 1$ .*

Corollary 1.4 has many geometric consequences with respect to building moduli spaces with functorial relative canonical sheaves. For example, cone singularities over abelian surfaces cannot be smoothed over irreducible bases. It also generalizes some aspects of theorems by Kollár and Kovács [2010, Theorem 7.12] and Hassett [2001, Theorem 1.1] stating that if all fibers are Du Bois schemes or log canonical surfaces and the general fiber is  $S_k$  or Cohen–Macaulay, respectively, then so is the central fiber.

Interestingly, the nonexistence of a depth  $n - 1$  point is the strongest implication of the compatibility of the relative canonical sheaf with base change.

**Proposition 1.5.** *Corollary 1.4 is sharp in the sense that  $n - 1$  cannot be replaced by  $i$  for any  $i < n - 1$ .*

Summarizing, Corollary 1.4 and Proposition 1.5 state that in moduli spaces satisfying Kollár’s condition,  $S_{n-1}$  schemes do not appear in the irreducible components containing Cohen–Macaulay schemes. However,  $S_j$  schemes can possibly show up for some  $j < n - 1$ .

If a scheme  $X$  is Cohen–Macaulay, which by definition means that  $\mathbb{O}_X$  is Cohen–Macaulay, then  $\omega_X$  is Cohen–Macaulay as well [Kollár and Mori 1998, Corollary 5.70]. One would expect that if  $\mathbb{O}_X$  is only  $S_{n-1}$ , then typically  $\omega_X$  is also  $S_{n-1}$  or at least it can be  $S_{n-1}$ . Surprisingly the truth is quite the opposite. The following application of Theorem 1.3 states that in certain cases an  $S_{n-1}$  scheme cannot have even an  $S_3$  canonical sheaf.

**Theorem 1.6.** *If  $X$  is an  $S_3, G_2$  scheme of pure dimension  $n$ , which has a closed point with depth  $n - 1$ , then  $\omega_X$  is not  $S_3$ .*

The most immediate consequences of Theorem 1.6 deal with compatibility of restriction to subvarieties. For example, one can show that on a cone  $X$  over a Calabi–Yau threefold  $Y$  with  $h^2(\mathbb{O}_Y) \neq 0$ , for an effective, normal Cartier divisor  $D$ ,

$$\omega_X(D)|_D \cong \omega_D \iff D \text{ does not pass through the vertex.}$$

Or more generally, for an  $S_{n-1}$ , normal variety  $X$  and an effective, normal Cartier divisor  $D$ ,

$$\omega_X(D)|_D \cong \omega_D \iff D \text{ does not pass through any closed point with depth } n - 1.$$

Theorem 1.6 can also be related to log canonical centers. If  $(X, D)$  is a log canonical pair,  $D \sim_{\mathbb{Q}} -K_X$  and  $\omega_X$  is not  $S_3$  at  $x \in X$ , then  $x$  is a log canonical center of the pair  $(X, D)$  [Kollár 2011, Theorem 3]. Hence by Theorem 1.6, if  $X$  is  $S_{n-1}$  and  $(X, D)$  log canonical such that  $D \sim_{\mathbb{Q}} -K_X$ , then  $(X, D)$  has a log-canonical center at all closed points with depth  $n - 1$ . This statement is of course obvious if we know that the depth  $n - 1$  closed points are already log-canonical centers of  $X$ . However, that is not always the case. For example, let  $X$  be the

cone, with high enough polarization, over the product  $Y$  of a K3 surface with the projective line and let  $D$  be the cone over an anti canonical divisor of  $Y$ . Then,  $(X, D)$  is log-canonical,  $X$  is  $S_{n-1}$  and the cone point is the only closed point with depth  $n - 1$ ; see Lemma 4.3. Still, the vertex is not a log-canonical center of  $X$ , because  $K_X$  is not  $\mathbb{Q}$ -Cartier.

Theorem 1.6 raises the following question as well.

**Question 1.7.** Is it true that if  $X$  is a pure  $n$ -dimensional scheme such that  $\mathbb{O}_X$  is  $S_l$ , but not  $S_{l+1}$ , and  $\omega_X$  is  $S_j$ , but not  $S_{j+1}$ , for some  $j, l < n$ , then  $j + l \leq n + 1$ ?

**Remark 1.8.** By the methods of Section 4, the answer to Question 1.7 is positive if  $X$  is a cone over a smooth projective variety.

There are a couple of intuitive reasons for the failure of compatibility in (1.2.a). First, compatibility holds for the relative dualizing complex if the base is smooth by Proposition 3.3.(1). Hence  $\omega_{\mathcal{H}/B}$  is a nonfunctorial component, the  $-n$ -th cohomology sheaf, of the functorial object  $\omega_{\mathcal{H}/B}^\bullet$ . For example, by the proof of Theorem 1.3, if the general fiber is Cohen–Macaulay and the central fiber is  $S_{n-1}$ , the restriction homomorphism fits into an exact sequence as follows, with a nonzero term on the right.

$$0 \rightarrow \omega_{\mathcal{H}/B}|_{X_b} \rightarrow \omega_{\mathcal{H}_b} \rightarrow \mathcal{T}or^1(h^{-(n-1)}(\omega_{\mathcal{H}/B}^\bullet), \mathbb{O}_{\mathcal{H}_0}) \rightarrow 0 \tag{1.8.a}$$

This shows in a precise way how the functoriality might be destroyed by passing to the lowest cohomology sheaf of  $\omega_{\mathcal{H}/B}^\bullet$ .

Another explanation for the incompatibility (1.2.a) is that  $\mathcal{H}_0$  is too singular. Using stable reduction one may find a replacement for  $\mathcal{H}_0$  with the mildest possible singularities. The reduction steps consist of blow-ups, finite surjective normalized base changes and contractions on the total space of the family. The output is a family, the relative canonical sheaf of which is compatible with base change by [Kollár and Kovács 2010]. At the end of the article, we also present the stable reduction of our construction using a straightforward ad hoc method. The algorithmic, and lengthy, method can be found in the preprint version of the article.

In Section 3, we start with a short background overview on the base-change properties of relative dualizing complexes and relative canonical sheaves. The proofs of the main theorems can be found in Section 6 and Section 7. Some of these results are based on the existence of projective cones with appropriately chosen singularities. In Section 4 we give a cohomological characterization of when certain sheaves on a cone are  $S_d$ . Then in Section 5 we use this characterization to give the desired examples of projective cones. In Section 8 we compute the stable limit of our construction.

## 2. Notation and assumptions

Unless otherwise stated, *scheme* means a separated scheme of finite type over a fixed field  $k$  of characteristic zero and every morphism is separated. A *variety* is an integral scheme. A *projective* or *quasiprojective* scheme means a projective or quasiprojective scheme over  $k$ . A *curve* is a quasiprojective, integral scheme of dimension one. If  $Y$  is a subscheme of  $X$ , then  $\mathcal{I}_{Y,X}$  is the ideal sheaf of  $Y$  in  $X$ . If  $\mathcal{I}_{Y,X}$  is a line bundle (that is, a locally free sheaf of rank one), then we define  $\mathbb{O}_X(-Y) := \mathcal{I}_{Y,X}$  and  $\mathbb{O}_X(Y) := \mathbb{O}_X(-Y)^{-1}$ . Notice that  $\mathcal{I}_{Y,X}$  being a line bundle is equivalent to  $Y$  being defined around every point  $P$  by a single nonzero divisor element of  $\mathbb{O}_{X,P}$ .

A hypersurface of a quasiprojective scheme  $X \subseteq \mathbb{P}^N$  is a subscheme  $H \subseteq X$  defined by a section of  $\mathbb{O}_X(d)$  for some  $d > 0$ . If  $H$  and  $H'$  are hypersurfaces of a quasiprojective scheme  $X \subseteq \mathbb{P}^N$ , defined by  $f_0$  and  $f_\infty \in H^0(\mathbb{P}^N, \mathbb{O}_{\mathbb{P}^N}(d))$ , respectively, then the *pencil* generated by  $H$  and  $H'$  is the subscheme  $\mathcal{H} \subseteq X \times \mathbb{P}^1$  defined by the section  $f_0 t_0 + f_\infty t_1$  of  $H^0(X \times \mathbb{P}^1, \mathbb{O}(d, 1))$ . Here  $t_0$  and  $t_1$  are the usual parameters of  $\mathbb{P}^1$ , and  $f_0$  and  $f_\infty$  are viewed as elements of  $H^0(X, \mathbb{O}_X(d))$  via the natural homomorphism  $H^0(\mathbb{P}^N, \mathbb{O}_{\mathbb{P}^N}(d)) \rightarrow H^0(X, \mathbb{O}_X(d))$ .

For a complex  $\mathcal{C}^\bullet$  of sheaves,  $h^i(\mathcal{C}^\bullet)$  is the  $i$ -th cohomology sheaf of  $\mathcal{C}$ . For a morphism  $f : X \rightarrow Y$ ,  $\omega_{X/Y}^\bullet := f^! \mathbb{O}_Y$ , where  $f^!$  is the functor obtained in [Hartshorne 1966, Corollary VII.3.4.a]. If  $f$  has equidimensional fibers of dimension  $n$ , then  $\omega_{X/Y} := h^{-n}(\omega_{X/Y}^\bullet)$ . Every complex and morphism of complexes is considered in the derived category  $D(qc/\cdot)$  of quasicohherent sheaves up to the equivalences defined there. If  $Z$  is a closed subscheme of  $X$ , where  $\iota : Z \rightarrow X$  is the embedding morphism, then the map  $\mathcal{R}\iota_* \cong \iota_*$  identifies  $D(qc/Z)$  with a full subcategory of  $D(qc/X)$ . We use this identification at multiple places, equating  $\mathcal{C}^\bullet$  and  $\mathcal{R}\iota_* \mathcal{C}^\bullet$  for every  $\mathcal{C}^\bullet \in D(qc/Z)$ . If  $Z$  is a closed subscheme of a scheme  $X$ , then  $(\cdot)|_Z^L$  denotes the derived restriction functor, which is naturally isomorphic to  $\cdot \otimes^L \mathbb{O}_Z$  via the identification mentioned above. A *line bundle* is a locally free sheaf of rank one.

If  $X \rightarrow B$  is a morphism of schemes, then  $X_b$  is the scheme theoretic fiber of  $X$  over  $B$ . If a sheaf  $\mathcal{F}$  on  $X$  is given, then  $\mathcal{F}_b := \mathcal{F}|_{X_b}$ . The dimension  $\dim_X P$  of a point  $P \in X$  is the dimension of its closure in  $X$ . The acronym *slc* stands for semi-log canonical [Hacon and Kovács 2010, Definition 3.13.5].

The *depth* of a coherent sheaf  $\mathcal{F}$  at a point  $x \in X$  is by definition the depth of  $\mathcal{F}_x$  with respect to the maximal ideal  $m_{X,x}$  at  $x$  and is denoted by  $\text{depth } \mathcal{F}_x$ . The depth of a scheme  $X$  at  $x$  is  $\text{depth } \mathbb{O}_{X,x}$ . A coherent sheaf  $\mathcal{F}$  is  $S_d$  on  $X$  if for every  $x \in X$ ,

$$\text{depth } \mathcal{F}_x \geq \min\{d, \dim \mathbb{O}_{X,x}\}. \quad (2.0.b)$$

Note that there is an ambiguity in the literature about the definition of  $S_d$  sheaves. Many sources replace  $\mathbb{O}_{X,x}$  in (2.0.b) by  $\mathcal{F}_x$ , thus gaining a weaker notion. Since every sheaf of this article has full support, or equivalently every sheaf is considered over its support, the two definitions are equivalent for all cases considered here. Hence, we decided to include the stronger notion, but the reader should feel free to think about the other one as well. For a morphism  $f : X \rightarrow B$ ,  $\mathcal{F}$  is *relative  $S_d$*  if  $\mathcal{F}|_{X_b}$  is  $S_d$  for all  $b \in B$ . The word (relative) *Cohen–Macaulay* is a synonym for (relative)  $S_{\dim X}$ .

A scheme  $X$  is  $G_r$  for some  $r \geq 0$  if it is Gorenstein in codimension  $r$ . A point  $P \in X$  is an *associated point* of a coherent sheaf  $\mathcal{F}$  if  $m_{X,P}$  is the annihilator of some element of  $\mathcal{F}_P$ . An *associated component* of a coherent sheaf is the closure of an associated point. One can show that if  $Q \in X$ ,  $\mathcal{F}_Q \neq 0$  and  $\mathcal{P}$  is the set of prime ideals of  $\mathbb{O}_{X,Q}$  corresponding to generalizations of  $Q$  that are associated points of  $\mathcal{F}$ , then

$$\bigcup_{P \in \mathcal{P}} P = \{x \in \mathbb{O}_{X,Q} \mid \text{there exists } 0 \neq m \in \mathcal{F}_Q \text{ with } xm = 0\}$$

Consequently, if  $s$  is a section of a line bundle, then it does not vanish on any associated component of  $X$  (that is, of  $\mathbb{O}_X$ ) if and only if  $s_P$  is not a zero divisor for every  $P \in X$ . That is, if  $H$  is the subscheme of  $X$  cut out by  $s$ , then  $\mathcal{I}_{H,X}$  is a line bundle if and only if  $s$  does not vanish on any associated component of  $X$ .

For an  $S_2, G_1$  scheme and an arbitrary coherent sheaf  $\mathcal{F}$ , the  $n$ -th *reflexive power* is

$$\mathcal{F}^{[n]} := \begin{cases} (\mathcal{F}^{\otimes n})^{**} & \text{if } n \geq 0, \\ (\mathcal{F}^{\otimes (-n)})^* & \text{if } n < 0. \end{cases}$$

That is, it is the reflexive hull of the  $n$ -th tensor power. A coherent sheaf  $\mathcal{F}$  is a  $\mathbb{Q}$ -line bundle if  $\mathcal{F}^{[n]}$  is a line bundle for some  $n > 0$ . Note that if  $f : X \rightarrow B$  is a family with  $\omega_{X_b}$  a  $\mathbb{Q}$ -line bundle for all  $b \in B$ , then  $\omega_{X/B}$  is not necessarily a  $\mathbb{Q}$ -line bundle [Hacon and Kovács 2010, Section 14.A]. However, if the  $X_b$  are  $S_2, G_1$  schemes and  $\omega_{X/B}$  a  $\mathbb{Q}$ -line bundle then  $\omega_{X_b}$  is a  $\mathbb{Q}$ -line bundle for all  $b \in B$ ; see [Hassett and Kovács 2004, Lemma 2.6].

### 3. Background on base change for dualizing complexes

This section contains a general overview on the base change properties of relative dualizing complexes and relative canonical sheaves. For experts, some of the statements might be well known, but they are included here for completeness and easier reference. Readers more interested in geometric arguments and willing to accept the statements of this section without proof should feel free to skip to the next section.



Recall that the relative dualizing complex  $\omega_{X/B}^\bullet$  of a quasiprojective family  $f : X \rightarrow B$  is defined as  $f^! \mathbb{O}_B$ . Here  $f^!$  is the functor constructed in [Hartshorne 1966, Corollary VII.3.4.a]. The following technical point should be noted here.

**Remark 3.1.** There is also another definition of  $f^!$  in [Neeman 1996] as the right adjoint of  $Rf_*$ . The two definitions coincide for proper morphisms by [Hartshorne 1966, Theorem VII.3.3; Neeman 1996, Section 6], but not in general. For example, if  $X$  is smooth affine variety over  $B = \text{Spec } k$  and  $f$  is the structure map, then Hartshorne's definition of  $f^! \mathbb{O}_{\text{Spec } k}$  lives in cohomological degree  $-\dim X$  while Neeman's is in cohomological degree zero. See [Lipman and Hashimoto 2009, Part I, Exercise 4.2.3.d] for more details on the differences (Neeman's  $f^!$  is denoted  $f^\times$  there). We use Hartshorne's definition in this article.

The dualizing complex of a single scheme  $Y$  is  $\omega_Y^\bullet := \omega_{Y/\text{Spec } k}^\bullet$ . The following fact is needed in the proof of Proposition 3.3(11). It follows from the invariance of the length of maximal regular sequences [Bruns and Herzog 1993, Theorem 1.2.5].

**Fact 3.2.** *Let  $P$  be a point of a subscheme  $H$  of a scheme  $X$  such that  $(\mathcal{F}_{H,X})_P$  is a line bundle,  $d$  is an integer, and  $\mathcal{F}$  is a coherent  $S_1$  sheaf with full support (that is,  $\text{supp } \mathcal{F} = X$ ) on  $X$ .*

- (1)  $\text{depth } \mathcal{F}_P \geq d \iff \text{depth}(\mathcal{F}|_H)_P \geq d - 1$  (here  $\mathcal{F}|_H$  is regarded as a sheaf on  $H$ , not on  $X$ ),
- (2)  $\text{depth } \mathcal{F}_P \geq \min\{d, \dim \mathbb{O}_{X,P}\} \iff \text{depth}(\mathcal{F}|_H)_P \geq \min\{d - 1, \dim \mathbb{O}_{H,P}\}$ .

**Proposition 3.3.** *Suppose we have a flat family  $f : \mathcal{H} \rightarrow B$  of schemes of pure dimension  $n$  over a smooth base, a point  $0 \in B$  and a single quasiprojective scheme  $X$  of pure dimension  $n$ .*

- (1) *There is an isomorphism*

$$\omega_{\mathcal{H}/B}^\bullet|_{\mathcal{H}_0}^L \cong \omega_{\mathcal{H}_0}^\bullet. \quad (3.3.a)$$

- (2) *Fixing any isomorphism in (3.3.a) yields natural homomorphism*

$$\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}. \quad (3.3.b)$$

- (3) *If  $B$  is of pure dimension  $d$  with  $\mathbb{O}_B \cong \omega_B$ , then  $\omega_{\mathcal{H}/B}^\bullet \cong \omega_{\mathcal{H}}^\bullet[-d]$ .*
- (4) *If  $V \subseteq X$  is any open set, then  $\omega_V^\bullet \cong \omega_X^\bullet|_V$ .*
- (5) *If  $U \subseteq \mathcal{H}$  is any open set, then  $\omega_{U/B}^\bullet \cong \omega_{\mathcal{H}/B}^\bullet|_U$ .*
- (6) *If  $P \in X$  is a point, then  $\text{depth}_P \mathbb{O}_X = d$  if and only if  $h^i(\omega_X^\bullet)_P$  is zero for  $i > -d - \dim_X P$  and nonzero for  $i = -d - \dim_X P$ .*
- (7) *If  $P \in \mathcal{H}$  is a point, then  $\text{depth}_P \mathbb{O}_{\mathcal{H}_{f(P)}} = d$  if and only if  $h^i(\omega_{\mathcal{H}/B}^\bullet)_P$  is zero for  $i > -d - \dim_{\mathcal{H}_{f(P)}} P$  and nonzero for  $i = -d - \dim_{\mathcal{H}_{f(P)}} P$ .*

- (8)  $\omega_X$  is  $S_2$ .
- (9)  $\omega_{\mathcal{H}/B}$  is  $S_2$ .
- (10) If the fibers of  $f$  are Cohen–Macaulay then  $\omega_{\mathcal{H}/B} \cong \omega_{\mathcal{H}/B}^\bullet$  and consequently (3.3.b) is an isomorphism.
- (11) If  $\mathcal{H}_0$  is  $S_2$  and  $G_1$ , then (3.3.b) is isomorphism if and only if

$$\text{depth } \omega_{\mathcal{H}/B,P} \geq \min\{3, \dim \mathbb{O}_{\mathcal{H},P}\} \quad \text{for every } P \in \mathcal{H}_0. \quad (3.3.c)$$

Furthermore if (3.3.c) is not satisfied then not only is (3.3.b) not an isomorphism, but  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \not\cong \omega_{\mathcal{H}_0}$ .

*Proof.* First, we prove point (1). It will be an ad hoc proof, since we have not found the exact statement in the literature. The statements we found are either only for flat base change morphisms [Hartshorne 1966, Corollary VII.3.4.a] or for proper  $f$  [Lipman and Hashimoto 2009, Part I, Corollary 4.4.3]. Note that, however, it might seem that point (1) follows from base change for proper  $f$ , to the best knowledge of the author, it is not clear whether one can compactify a flat morphism to a flat morphism.

First, by [Hartshorne 1966, Corollary VII.3.4.a],  $\omega_{\mathcal{H}/B}^\bullet$  is compatible with flat base change. So, since  $\text{Spec } \widehat{\mathbb{O}}_{B,0}$  is flat over  $B$ , we may assume that  $B$  is the spectrum of a complete local ring of a smooth scheme and  $0$  is the unique closed point. In particular, then  $B \cong \text{Spec } k[[x_1, \dots, x_m]]$ . Hence, by induction on  $m$ , it is enough to prove that

$$\omega_{\mathcal{H}/B}|_Y \cong \omega_{Y/C}^\bullet, \quad (3.3.d)$$

where  $C := \text{Spec } k[[x_1, \dots, x_{m-1}]]$  and  $Y := \mathcal{H} \times_B C$ . To prove (3.3.d), first consider the usual exact triangle

$$\mathbb{O}_{\mathcal{H}} \xrightarrow{\mu} \mathbb{O}_{\mathcal{H}} \rightarrow \mathbb{O}_Y \xrightarrow{+1}, \quad (3.3.e)$$

where  $\mu$  is multiplication by  $x_m$ . Tensoring (3.3.e) by  $\omega_{\mathcal{H}/B}^\bullet$  yields

$$\omega_{\mathcal{H}/B}^\bullet \xrightarrow{\mu \otimes \text{id}_{\omega_{\mathcal{H}/B}^\bullet}} \omega_{\mathcal{H}/B}^\bullet \rightarrow \omega_{\mathcal{H}/B}^\bullet|_Y \xrightarrow{+1}. \quad (3.3.f)$$

On the other hand, applying  $\mathcal{R}\mathcal{H}om(\cdot, \omega_{\mathcal{H}/B}^\bullet)$  and a rotation to (3.3.e) yields

$$\omega_{\mathcal{H}/B}^\bullet \xrightarrow{\mu \otimes \text{id}_{\omega_{\mathcal{H}/B}^\bullet}} \omega_{\mathcal{H}/B}^\bullet \rightarrow \mathcal{R}\mathcal{H}om(\mathbb{O}_Y, \omega_{\mathcal{H}/B}^\bullet)[1] \xrightarrow{+1}. \quad (3.3.g)$$

So, (3.3.f) and (3.3.g) together imply that

$$\mathcal{R}\mathcal{H}om(\mathbb{O}_Y, \omega_{\mathcal{H}/B}^\bullet)[1] \cong \omega_{\mathcal{H}/B}^\bullet|_Y. \quad (3.3.h)$$

Denote by  $\iota$  and  $g$  the maps  $Y \rightarrow X$  and  $Y \rightarrow C$ , respectively. The following stream of isomorphisms finishes then the proof of point (1).

$$\begin{aligned} \omega_{\mathcal{H}/B}^\bullet|_Y &\cong \underbrace{\mathcal{R}\mathcal{H}om_X(\mathcal{R}\iota_*\mathbb{O}_Y, \omega_{\mathcal{H}/B}^\bullet)[1]}_{\text{by (3.3.h)}} \cong \underbrace{\mathcal{R}\mathcal{H}om_Y(\mathbb{O}_Y, \iota^!\omega_{\mathcal{H}/B}^\bullet)[1]}_{\text{by Grothendieck duality}} \\ &\cong \iota^!\omega_{\mathcal{H}/B}^\bullet[1]! \cong \underbrace{\iota^!f^!\mathbb{O}_B[1]}_{\text{definition of } \omega_{\mathcal{H}/B}^\bullet} \cong \underbrace{\iota^!f^!\omega_B^\bullet[-(m-1)]}_{\omega_B^\bullet[-m] \cong \mathbb{O}_B} \\ &\cong \underbrace{\omega_Y^\bullet[-(m-1)]}_{\omega_Y^\bullet \cong (f \circ \iota)!\omega_B^\bullet} \cong \underbrace{g^!\omega_C^\bullet[-(m-1)]}_{\omega_Y^\bullet \cong g^!\omega_C^\bullet} \cong \underbrace{g^!\mathbb{O}_C}_{\omega_C^\bullet[-(m-1)] \cong \mathbb{O}_C} \cong \omega_{Y/C}^\bullet. \end{aligned}$$

To prove point (2), notice that since  $\omega_{\mathcal{H}/B} := h^{-n}(\omega_{\mathcal{H}/B}^\bullet)$  is the lowest cohomology sheaf of  $\omega_{\mathcal{H}/B}^\bullet$ , there is a homomorphism

$$\omega_{\mathcal{H}/B}[n] \rightarrow \omega_{\mathcal{H}/B}^\bullet. \tag{3.3.i}$$

Applying  $(\cdot)|_{\mathcal{H}_0}^L$  to (3.3.i) and then composing with the isomorphism given by (3.3.a) yields a homomorphism

$$\omega_{\mathcal{H}/B}[n]|_{\mathcal{H}_0}^L \rightarrow \omega_{\mathcal{H}_0}^\bullet. \tag{3.3.j}$$

Finally taking  $-n$ -th cohomology sheaves of (3.3.j) yields the restriction homomorphism of (3.3.b).

Point (3) is shown by the following line of isomorphisms:

$$\omega_{\mathcal{H}/B}^\bullet = f^!\mathbb{O}_B \cong f^!\omega_B \cong f^!\omega_B^\bullet[-d] \cong \omega_{\mathcal{H}}^\bullet[-d].$$

To prove point (4), consider the following commutative diagram.

$$\begin{array}{ccc} V & \xrightarrow{j} & X \\ & \searrow v & \downarrow \mu \\ & & \text{Spec } k \end{array}$$

Since  $j$  is smooth of relative dimension 0, using the notation of [Hartshorne 1966], we have  $j^! \cong j^\# \cong j^*$ , and then

$$\omega_V^\bullet = v^!\mathbb{O}_{\text{Spec } k} \cong j^!\mu^!\mathbb{O}_{\text{Spec } k} \cong j^!\omega_X^\bullet \cong j^*\omega_X^\bullet = \omega_X^\bullet|_V.$$

Point (5) follows from points (4) and (3).

Point (6) is proved in [Kovács 2011, Proposition 3.2] (by taking  $\mathcal{F} := \mathbb{O}_X$ ). To prove point (7), let  $b := f(P)$  and consider the following Cartesian square.

$$\begin{array}{ccc} \mathcal{H} & \longleftarrow & \mathcal{H}' \\ \downarrow f & \lambda' & \downarrow f' \\ B & \longleftarrow_{\lambda} & \text{Spec } \mathbb{O}_{B,b} \end{array}$$

By flat base change,

$$(\lambda')^* \omega_{\mathcal{H}/B}^\bullet \cong \omega_{\mathcal{H}'/\text{Spec } \mathbb{O}_{B,b}}^\bullet. \tag{3.3.k}$$

That is,

$$h^i(\omega_{\mathcal{H}/B}^\bullet)_P \cong \underbrace{h^i(\omega_{\mathcal{H}'/\text{Spec } \mathbb{O}_{B,b}}^\bullet)_P}_{\text{by (3.3.k)}} \cong \underbrace{h^i(\omega_{\mathcal{H}'}^\bullet[-\dim \mathbb{O}_{B,b}])_P}_{\text{by point (3)}} \cong h^{i-\dim \mathbb{O}_{B,b}}(\omega_{\mathcal{H}'}^\bullet)_P.$$

Hence,

$$\begin{array}{c} h^i(\omega_{\mathcal{H}/B}^\bullet)_P \text{ is } \begin{cases} 0 & \text{if } i > -d - \dim_{\mathcal{H}_b} P, \\ \neq 0 & \text{if } i = -d - \dim_{\mathcal{H}_b} P. \end{cases} \\ \updownarrow \\ h^i(\omega_{\mathcal{H}'}^\bullet)_P \text{ is } \begin{cases} 0 & \text{if } i > -d - \dim_{\mathcal{H}_b} P - \dim \mathbb{O}_{B,b}, \\ \neq 0 & \text{if } i = -d - \dim_{\mathcal{H}_b} P - \dim \mathbb{O}_{B,b}. \end{cases} \\ \updownarrow \\ \text{depth}_P \mathbb{O}_{\mathcal{H}'} = d + \dim \mathbb{O}_{B,b} \\ \updownarrow \text{ (by Fact 3.2)} \\ (\text{depth}_P \mathbb{O}_{(\mathcal{H}')_{f(P)}} =) \text{depth}_P \mathbb{O}_{\mathcal{H}_{f(P)}} = d. \end{array}$$

To prove point (8), by point (4) we may assume that  $X$  is affine. Using point (4) again we may also assume that it is projective. Then [Kollár and Mori 1998, Corollary 5.69] concludes the proof of point (8). Point (9) is a consequence of point (8) and point (3). Point (10) is shown in [Conrad 2000, Theorem 3.5.1].

To prove point (11), notice that by point (8),  $\omega_{\mathcal{H}_0}$  is  $S_2$ . Also since  $\mathcal{H}_0$  is  $G_1$ , using point (10), the homomorphism  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is isomorphism in codimension one. Then by [Hartshorne 1994, Theorem 1.9 and Theorem 1.12], using that  $\mathcal{H}_0$  is  $S_2$  and  $G_1$ ,  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is an isomorphism if and only if  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$  is  $S_2$ . Finally, by Fact 3.2(2), this is equivalent to (3.3.c).

Notice that if (3.3.c) is not satisfied, then  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$  is not  $S_2$  over  $\mathcal{H}_0$ . Hence in this case not only can (3.3.b) not be isomorphism, but any isomorphism between  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$  and  $\omega_{\mathcal{H}_0}$  is impossible.  $\square$

**Remark 3.4.** A priori, saying that (3.3.b) is an isomorphism is a stronger statement than that  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$  is isomorphic to  $\omega_{\mathcal{H}_0}$ . However, if  $B$  is smooth,  $\mathcal{H}_0$  is projective,  $S_2$  and  $G_1$ , they are equivalent by the following argument. In this case  $\omega_{\mathcal{H}_0}$  is  $S_2$  and is a line bundle over the Gorenstein locus  $U$ . Assume that  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \cong \omega_{\mathcal{H}_0}$  via an arbitrary isomorphism  $\alpha$ . Then  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$  is also  $S_2$  and a line bundle over  $U$ . Since both are  $S_2$ , homomorphisms  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  are determined in codimension one, e.g., over  $U$ . Furthermore, any two isomorphisms over  $U$  between any two line bundles differ by multiplication with an element of  $H^0(U, \mathcal{O}_{\mathcal{H}_0})$ , where  $H^0(U, \mathcal{O}_{\mathcal{H}_0}) \cong k^*$ , by  $\mathcal{H}_0$  being  $S_2$  and projective. Since the restriction of the natural morphism  $\beta : \omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  over  $U$  is an isomorphism,  $\alpha$  differs from  $\beta$  over  $U$  by a multiplication with an element of  $k^*$ . However, then the same is true over entire  $X$ , by the codimension one determination. Hence  $\beta$  is also an isomorphism.

Finally, we conclude with a statement about restriction behavior of relative dualizing complexes and relative canonical sheaves to hypersurfaces. For that we also need a lemma about flatness of hypersurfaces.

**Lemma 3.5.** *If  $f : \mathcal{X} \rightarrow B$  is a flat morphism onto a smooth curve and  $\mathcal{H} \subseteq \mathcal{X}$  is a subscheme for which  $\mathcal{I}_{\mathcal{H},\mathcal{X}}$  is a line bundle, then the following are equivalent:*

- (1)  $\mathcal{I}_{\mathcal{H}_b,\mathcal{X}_b}$  is a line bundle for every  $b \in B$ .
- (2)  $\mathcal{H}$  is flat over  $B$ .

*In particular, if  $f : \mathcal{X} \rightarrow B$  is flat with fibers of pure dimension  $n$  and  $\mathcal{H} \subseteq \mathcal{X}$  is also flat with  $\mathcal{I}_{\mathcal{H},\mathcal{X}}$  a line bundle, then fibers of  $\mathcal{H}$  are of pure dimension  $n - 1$ .*

*Proof.* We prove only the equivalence statement, since the addendum follows from the ideals  $\mathcal{I}_{\mathcal{H}_b,\mathcal{X}_b}$  being line bundles.

The statement is local on  $\mathcal{H}$ . So, fix  $P \in \mathcal{H}$  and let  $Q := f(P)$ . By [Hartshorne 1977, Proposition 9.1A.a],  $\mathcal{H}$  and  $\mathcal{X}$  are flat over  $B$  at  $P$  if and only if the respective homomorphisms  $\mathcal{O}_{\mathcal{H},P} \rightarrow \mathcal{O}_{\mathcal{H},P}$  and  $\mathcal{O}_{\mathcal{H},P} \rightarrow \mathcal{O}_{\mathcal{X},P}$  induced by multiplication with some power of the local parameter  $t$  of  $\mathcal{O}_{B,Q}$  are injective. Furthermore, by induction this is equivalent to the injectivity of multiplication with the first power  $t$ .

The assumptions of the lemma state that  $(\mathcal{I}_{\mathcal{H},\mathcal{X}})_P \subseteq \mathcal{O}_{\mathcal{X},P}$  is generated by a nonzero divisor element  $s$ . Hence there is a commutative diagram with exact rows

and columns as follows.

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \uparrow & & \uparrow & \\
 0 & \longrightarrow & \ker(\cdot s) & \longrightarrow & \mathbb{O}_{\mathcal{X}_Q, P} & \xrightarrow{\cdot s} & \mathbb{O}_{\mathcal{X}_Q, P} \\
 & & & \uparrow & & \uparrow & \\
 & & & \mathbb{O}_{\mathcal{X}, P} & \xrightarrow{\cdot s} & \mathbb{O}_{\mathcal{X}, P} & \xrightarrow{\cdot s} & \mathbb{O}_{\mathcal{H}, P} & \longrightarrow & 0 \\
 & & & \uparrow & & \uparrow & & \uparrow & & \\
 0 & \longrightarrow & \mathbb{O}_{\mathcal{X}, P} & \xrightarrow{\cdot s} & \mathbb{O}_{\mathcal{X}, P} & \xrightarrow{\cdot s} & \mathbb{O}_{\mathcal{H}, P} & \longrightarrow & 0 \\
 & & & \uparrow & & \uparrow & & \uparrow & & \\
 & & & 0 & & 0 & & \ker(\cdot t) & & \\
 & & & & & & & \uparrow & & \\
 & & & & & & & 0 & & 
 \end{array}$$

By the snake lemma applied vertically,  $\ker(\cdot t) = \ker(\cdot s)$ . In particular,  $\ker(\cdot t) = 0$  if and only if  $\ker(\cdot s) = 0$ . The former is equivalent to flatness of  $\mathcal{H} \rightarrow B$  at  $P$  while the latter is equivalent to  $\mathcal{I}_{\mathcal{H}, \mathcal{X}_Q}$  being a line bundle at  $P$ .  $\square$

**Proposition 3.6.** *If  $\mathcal{X} \rightarrow B$  is a flat family of pure  $n$ -dimensional schemes, and  $\mathcal{H} \subseteq \mathcal{X}$  a flat subscheme such that  $\mathcal{I}_{\mathcal{H}, \mathcal{X}}$  is a line bundle, then*

(1) *there is an isomorphism*

$$\omega_{\mathcal{X}/B}^\bullet(\mathcal{H})|_{\mathcal{H}}^L[-1] \cong \omega_{\mathcal{H}/B}^\bullet, \tag{3.6.a}$$

(2) *there is a homomorphism*

$$\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}} \rightarrow \omega_{\mathcal{H}/B}, \tag{3.6.b}$$

*which is isomorphism over the relative Cohen–Macaulay locus of  $\mathcal{H} \rightarrow B$ .*

*Proof.* Notice first that by Lemma 3.5,  $\mathcal{H}$  has equidimensional fibers and hence  $\omega_{\mathcal{H}/B}$  is defined indeed. To prove point (1), consider the exact sequence

$$0 \rightarrow \mathbb{O}_{\mathcal{X}} \rightarrow \mathbb{O}_{\mathcal{X}}(\mathcal{H}) \rightarrow \mathbb{O}_{\mathcal{H}}(\mathcal{H}) \rightarrow 0. \tag{3.6.c}$$

Applying  $(\cdot) \otimes^L \omega_{\mathcal{X}/B}^\bullet$  to (3.6.c) and then translating yields the exact triangle

$$\omega_{\mathcal{X}/B}^\bullet(\mathcal{H})|_{\mathcal{H}}^L[-1] \rightarrow \omega_{\mathcal{X}/B}^\bullet \rightarrow \omega_{\mathcal{X}/B}^\bullet(\mathcal{H}) \xrightarrow{+1} . \tag{3.6.d}$$

On the other hand if  $\iota : \mathcal{H} \rightarrow \mathcal{X}$  is the embedding morphism, then

$$\omega_{\mathcal{X}/B}^\bullet \cong \iota^! \omega_{\mathcal{X}/B}^\bullet = R\mathcal{H}om_{\mathcal{X}}(\mathbb{O}_{\mathcal{H}}, \iota^! \omega_{\mathcal{X}/B}^\bullet) \cong \underbrace{R\mathcal{H}om_{\mathcal{X}}(\mathbb{O}_{\mathcal{H}}, \omega_{\mathcal{X}/B}^\bullet)}_{\text{by Grothendieck duality}}.$$

Now, applying  $R\mathcal{H}om_{\mathcal{X}}(\cdot, \omega_{\mathcal{X}/B}^\bullet)$  to the twist of (3.6.c) by  $\mathbb{O}_{\mathcal{X}}(-\mathcal{H})$  yields the exact triangle

$$\omega_{\mathcal{X}/B}^\bullet \cong R\mathcal{H}om_{\mathcal{X}}(\mathbb{O}_{\mathcal{H}}, \omega_{\mathcal{X}/B}^\bullet) \rightarrow \omega_{\mathcal{X}/B}^\bullet \rightarrow \omega_{\mathcal{X}/B}^\bullet(\mathcal{H}) \xrightarrow{+1}. \tag{3.6.e}$$

Putting together (3.6.d) and (3.6.e) finishes the proof of point (1).

To prove (2), take the natural map  $\omega_{\mathcal{X}/B}[n] \rightarrow \omega_{\mathcal{X}/B}^\bullet$ , twist it with  $\mathbb{O}_{\mathcal{X}}(\mathcal{H})$  and then restrict to  $\mathcal{H}$ . This yields the commutative diagram

$$\begin{array}{ccc} & \xrightarrow{\quad \quad \quad} & \\ \omega_{\mathcal{X}/B}[n-1](\mathcal{H})|_{\mathcal{H}}^L & \longrightarrow & \omega_{\mathcal{X}/B}^\bullet(\mathcal{H})[-1]|_{\mathcal{H}}^L \xrightarrow{\cong} \omega_{\mathcal{X}/B}^\bullet \\ & & \underbrace{\qquad \qquad \qquad}_{\text{by point (1)}} \end{array} \tag{3.6.f}$$

Applying then  $h^{-(n-1)}(\cdot)$  to the long composition arrow of (3.6.f) yields the homomorphism (3.6.b).

Let  $P$  be a point of  $\mathcal{H}$  that is relatively Cohen–Macaulay over  $B$ , and let  $b$  be the image of  $P$  in  $B$ . By the openness of the relative Cohen–Macaulay locus, there is a neighborhood  $U$  of  $P$  where  $X \rightarrow B$  is relatively Cohen–Macaulay. In particular, then  $\omega_{\mathcal{X}/B}[n-1] \rightarrow \omega_{\mathcal{X}/B}^\bullet[-1]$  is an isomorphism over  $U$  by Proposition 3.3(10) and hence so is the first arrow of (3.6.f). This proves that (3.6.b) is an isomorphism in a neighborhood of  $P$ , which finishes the proof of point (2) as well.  $\square$

**Remark 3.7.** The homomorphisms constructed in Propositions 3.3 and 3.6, for example the isomorphisms (3.3.a) and (3.6.a), are not canonical in any sense.

### 4. Serre’s condition on projective cones

In this section we consider sheaves on projective cones that are isomorphic to pullbacks from the base outside the vertex. Lemma 4.3 gives a cohomological description of when such sheaves are  $S_d$ . Before that we also need a short lemma, Lemma 4.2, about how the property  $S_d$  pulls back in flat relatively Cohen–Macaulay families.

We cite the following fact separately here, because it is used at many places throughout the article, including the aforementioned Lemma 4.2.

**Fact 4.1** [Grothendieck 1965, Proposition 6.3.1]. *Let  $X$  and  $Y$  be two noetherian schemes,  $f : X \rightarrow Y$  a flat morphism,  $P \in X$  arbitrary and  $\mathcal{F}$  a coherent  $Y$  module. In this situation,*

$$\text{depth}_{\mathbb{O}_{X,P}}(f^*\mathcal{F})_P = \text{depth}_{\mathbb{O}_{Y,f(P)}} \mathcal{F}_{f(P)} + \text{depth}_{\mathbb{O}_{X_{f(P)},P}} \mathbb{O}_{X_{f(P)},P}.$$

**Lemma 4.2.** *If  $\mathcal{G}$  is a full-dimensional coherent  $S_d$  sheaf on the scheme  $X$ , and  $f : \mathcal{X} \rightarrow X$  is a flat, relatively Cohen–Macaulay family, then  $\mathcal{F} := f^*\mathcal{G}$  is  $S_d$  as well.*

*Proof.* For every  $x \in X$ ,

$$\begin{aligned} \text{depth } \mathcal{F}_x &= \underbrace{\text{depth } \mathcal{G}_{f(x)} + \text{depth } \mathbb{O}_{\mathcal{X}_{f(x)},x}}_{\text{Fact 4.1}} = \underbrace{\text{depth } \mathcal{G}_{f(x)} + \dim \mathbb{O}_{\mathcal{X}_{f(x)},x}}_{\mathcal{X}_{f(x)} \text{ is Cohen–Macaulay}} \\ &\geq \underbrace{\min\{d, \dim \mathbb{O}_{X,f(x)}\} + \dim \mathbb{O}_{\mathcal{X}_{f(x)},x}}_{\mathcal{G} \text{ is } S_d} \\ &\geq \min\{d, \dim \mathbb{O}_{X,f(x)} + \dim \mathbb{O}_{\mathcal{X}_{f(x)},x}\} \\ &= \underbrace{\min\{d, \dim \mathbb{O}_{\mathcal{X},x}\}}_{\substack{\dim \mathbb{O}_{X,f(x)} + \dim \mathbb{O}_{\mathcal{X}_{f(x)},x} = \dim \mathbb{O}_{\mathcal{X},x} \\ \text{by [Matsumura 1989, Theorem 15.1.ii]}}} \quad \square \end{aligned}$$

**Lemma 4.3.** *Assume that we are in the following situation:*

- $Y$  is a projective scheme,
- $X$  is the projectivized cone over  $Y$ ,
- $P$  is the vertex of  $X$  and  $V := X \setminus P$ ,
- $d$  is an integer such that  $2 \leq d \leq \dim X$ , and
- $\mathcal{F}$  is a coherent sheaf on  $X$ , such that  $\mathcal{F}|_V = \pi^*\mathcal{G}$  for some  $S_d$  coherent sheaf  $\mathcal{G}$  on  $Y$ , where  $\pi : V \rightarrow Y$  is the natural projection.

*Then the following conditions are equivalent:*

- (1)  $\text{depth } \mathcal{F}_P \geq d$ .
- (2)  $\text{depth } \mathcal{F}_P \geq \min\{d, \dim \mathbb{O}_{X,P}\}$ .
- (3)  $\mathcal{F}$  is  $S_d$ .
- (4)  $\mathcal{F}$  is  $S_2$  and  $H^i(Y, \mathcal{G}(n)) = 0$  for all  $0 < i < d - 1$  and  $n \in \mathbb{Z}$ .

*Proof.* Since  $\mathcal{G}$  is  $S_d$ ,  $\mathcal{F}$  is  $S_d$  everywhere except at the vertex  $P$  by Lemma 4.2. Hence, using the assumption  $d \leq \dim X$ ,

$$\begin{array}{c} \mathcal{F} \text{ is } S_d, \\ \updownarrow \\ \text{depth } \mathcal{F}_P \geq \min\{d, \dim \mathbb{O}_{X,P}\} \\ \updownarrow \\ \text{depth } \mathcal{F}_P \geq d \\ \updownarrow \\ H_P^i(Z, \mathcal{F}) = 0 \text{ for all } i < d \text{ and for the affine cone } Z, \end{array}$$



where the latter equivalence follows from [Hartshorne 1977, Exercises III.3.4.b and III.2.5]. So, we are left to show that the condition  $H^i_p(Z, \mathcal{F}) = 0$  for all  $i < d$  is equivalent to point (4). Define  $U := Z \setminus P$ . Then there is a long exact sequence

$$\dots \rightarrow H^i_p(Z, \mathcal{F}) \rightarrow H^i(Z, \mathcal{F}) \rightarrow H^i(U, \mathcal{F}) \rightarrow \dots$$

Since  $Z$  is affine  $H^i(Z, \mathcal{F}) = 0$  for all  $i > 0$ . Hence

$$H^i(U, \mathcal{F}) \cong H^{i+1}_p(Z, \mathcal{F}) \text{ for all } i > 0. \tag{4.3.a}$$

So, since  $H^0_p(Z, \mathcal{F}) = H^1_p(Z, \mathcal{F}) = 0$  is assumed in point (4), it is enough to show that for all  $0 < i < d - 1$ ,

$$H^i(U, \mathcal{F}) \cong \bigoplus_{n \in \mathbb{Z}} H^i(Y, \mathcal{G}(n)). \tag{4.3.b}$$

In fact we will prove this for all  $i$ . First, notice that  $U \cong \text{Spec}_Y(\bigoplus_{n \in \mathbb{Z}} \mathbb{O}_Y(n))$  and the natural projection  $\text{Spec}_Y(\bigoplus_{n \in \mathbb{Z}} \mathbb{O}_Y(n)) \rightarrow Y$  can be identified with  $\pi|_U$  via this isomorphism. Hence  $(\pi|_U)_* \mathbb{O}_U \cong \bigoplus_{n \in \mathbb{Z}} \mathbb{O}_Y(n)$  and  $R^i(\pi|_U)_* \mathbb{O}_U = 0$  for  $i > 0$ . So:

$$\begin{aligned} H^i(U, \mathcal{F}) &\cong H^i(Y, (\pi|_U)_* \mathcal{F}|_U) \cong H^i(Y, (\pi|_U)_* (\pi|_U)^* \mathcal{G}) \cong \\ &\cong H^i(Y, \bigoplus_{n \in \mathbb{Z}} \mathcal{G}(n)) \cong \bigoplus_{n \in \mathbb{Z}} H^i(Y, \mathcal{G}(n)) \end{aligned}$$

as claimed in (4.3.b). □

### 5. Construction of varieties with prescribed singularities

In this section, normal  $S_j$  (but not  $S_{j+1}$ ) varieties of dimension  $n \geq 3$  with  $S_l$  (but not  $S_{l+1}$ ),  $\mathbb{Q}$ -line bundle canonical sheaves are constructed for certain values of  $j$  and  $l$ . They are going to be used in Section 6 and in Section 7 to build families with prescribed base change behavior for the relative canonical sheaves. First we need some lemmas.

**Lemma 5.1.** *If  $H$  is a general, high enough degree hypersurface in a projective variety  $X$ , then  $H^i(H, \mathbb{O}_H) \cong H^i(X, \mathbb{O}_X)$  for every  $i$  such that  $0 < i < \dim H$ .*

*Proof.* We start with the usual exact sequence

$$0 \rightarrow \mathbb{O}_X(-H) \rightarrow \mathbb{O}_X \rightarrow \mathbb{O}_H \rightarrow 0. \tag{5.1.a}$$

Since  $\deg H \gg 0$ ,

$$H^i(X, \mathbb{O}_X(-H)) = 0 \text{ whenever } i < \dim X. \tag{5.1.b}$$

Taking the cohomology long exact sequence of (5.1.a) and using (5.1.b) finishes the proof. □

Iterated use of Lemma 5.1 yields the following:

**Lemma 5.2.** *If  $H$  is a general, high enough degree complete intersection (that is, it is the intersection of hypersurfaces, all of which are high enough degree) in a smooth projective variety  $X$ , then  $H^i(H, \mathbb{O}_H) \cong H^i(X, \mathbb{O}_X)$  for every  $i$  such that  $0 < i < \dim H$ .*

Finally, iterated use of the adjunction formula yields the following:

**Lemma 5.3.** *If  $H$  is a complete intersection in a smooth projective variety  $X$ , then  $\omega_H \cong \omega_X(m)|_H$  for some  $m > 0$  (here  $\mathbb{O}_X(1)$  is the very ample line bundle given by the projective embedding of  $X$ ).*

**Proposition 5.4.** *For each  $n \geq 2$  and  $2 \leq d, l \leq n$  such that  $l \leq d$  and  $d + l \leq n + 2$  there is an  $(n + 1)$ -dimensional projective variety  $X_{n+1}$  for which*

- $X_{n+1}$  is the projective cone over a smooth projective variety  $Y_n$  with vertex  $P$ ,
- $X_{n+1}$  is  $S_d$  and  $\text{depth } \mathbb{O}_{X_{n+1}, P} = d$ ,
- $\omega_{X_{n+1}}$  is  $S_l$  and  $\text{depth } \omega_{X_{n+1}, P} = l$ , and
- $\omega_{X_{n+1}}$  is a  $\mathbb{Q}$ -line bundle.

*Proof.* Take first two Calabi–Yau hypersurfaces  $Z$  and  $W$  of dimensions  $d - 1$  and  $n + 1 - l$ , respectively. Let  $Y := Y_n$  be a general high enough degree complete intersection of codimension  $d - l$  in  $Z \times W$ . Notice that  $d - l \geq 0$  by assumption. Finally, let  $X_{n+1}$  be the projective cone over  $Y$  polarized by  $\mathbb{O}_Y(1) := \mathbb{O}_{Z \times W}(p)|_Y$  for some  $p \gg 0$  (after fixing  $Y$ ). Here  $\mathbb{O}_{Z \times W}(1)$  is the very ample line bundle on  $Z \times W$  coming from its projective embedding.

The Künneth isomorphism yields

$$H^q(Z \times W, \mathbb{O}_{Z \times W}) \cong \bigoplus_{r=0}^q H^r(Z, \mathbb{O}_Z) \otimes H^{q-r}(W, \mathbb{O}_W).$$

Since  $Z$  and  $W$  are Calabi–Yau hypersurfaces of dimension  $d - 1$  and  $n + 1 - l$ , respectively, the following holds for their cohomology table:

$$\begin{aligned} H^q(Z, \mathbb{O}_Z) \neq 0 &\iff q = 0 \text{ or } d - 1, \\ H^s(W, \mathbb{O}_W) \neq 0 &\iff s = 0 \text{ or } n + 1 - l. \end{aligned}$$

Hence

$$H^q(Z \times E, \mathbb{O}_{Z \times E}) \neq 0 \iff q = 0, d - 1, n + 1 - l \text{ or } n - l + d.$$

Using Lemma 5.2 yields, for  $0 < q < n$ ,

$$H^q(Y, \mathbb{O}_Y) \neq 0 \iff q = d - 1 \text{ or } n + 1 - l. \tag{5.4.a}$$

Since  $p \gg 0$ , also,

$$H^q(Y, \mathbb{O}_Y(r)) = 0 \text{ for every } r \text{ and } 0 < q < n.$$

Then by Lemma 4.3 using that  $d - 1 \leq n + 1 - l$  by assumption,  $X_{n+1}$  is  $S_d$  and  $\text{depth } \mathcal{O}_{X_{n+1}, P} = d$  ( $X_{n+1}$  is  $S_2$  at the vertex, because  $p \gg 0$  and hence  $Y$  is projectively normal).

Serre duality implies that

$$H^q(Y, \omega_Y) \cong (H^{n-q}(Y, \mathcal{O}_Y))^*.$$

So, by (5.4.a), for  $0 < q < n$ ,

$$H^q(Y, \omega_Y) \neq 0 \iff q = l - 1 \text{ or } n + 1 - d.$$

Since  $X_{n+1}$  is an affine bundle over  $Y$ ,  $\omega_{X_{n+1}}$  is isomorphic to the pullback of  $\omega_Y$  outside of the vertex. Then by Lemma 4.3 using that  $l - 1 \leq n + 1 - d$ ,  $\omega_{X_{n+1}}$  is  $S_l$  and  $\text{depth } \omega_{X_{n+1}, P} = l$  ( $\omega_{X_{n+1}}$  is always  $S_2$  by Proposition 3.3(8)).

We have left to show that the  $\omega_{X_{n+1}}$  are  $\mathbb{Q}$ -Cartier. By Lemma 5.3,

$$\omega_Y^{\otimes p} \cong (\omega_{Z \times E}(m)|_Y)^{\otimes p} \cong (\mathcal{O}_{Z \times E}(m)|_Y)^{\otimes p} \cong \mathcal{O}_Y(m).$$

That is,  $\omega_Y^{\otimes p}$  is an integer multiple of the polarization of  $Y$  used at the construction of  $X_{n+1}$ . Hence, [Hacon and Kovács 2010, Exercise 3.5] concludes the proof.  $\square$

### 6. Construction of families without the base change property

In this section we present the proof of Theorem 1.2. The following lemma contains the key argument. It is also used in the proofs of Proposition 1.5 and Theorem 1.6.

**Lemma 6.1.** *Let  $f : \mathcal{H} \rightarrow B = \mathbb{P}^1$  be a flat pencil of hypersurfaces of a quasi-projective, equidimensional scheme  $X$ , such that  $\mathcal{I}_{\mathcal{H}, X \times B}$  is a line bundle and  $\mathcal{H}$  and the closed fibers of  $f$  are  $S_2$  and  $G_1$ .*

- (1) *If  $\omega_X$  is  $S_3$ , the restriction map  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is an isomorphism.*
- (2) *If  $\text{depth } \omega_{X, P} \not\geq \min\{3, \dim \mathcal{O}_{X, P}\}$  for some  $P \in X$ , such that  $P \in \mathcal{H}_0$ , but  $P \notin \mathcal{H}_\infty$ , then  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \not\cong \omega_{\mathcal{H}_0}$ .*

*Proof.* Notice that by flatness of  $\mathcal{H}$  and by Lemma 3.5, it does make sense to talk about  $\omega_{\mathcal{H}/B}$ . Define  $\mathcal{X} := X \times B$ . Then  $\mathcal{H}$  is a hypersurface of  $\mathcal{X}$ . By Proposition 3.6.(2) there is a homomorphism  $\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}} \rightarrow \omega_{\mathcal{H}/B}$ , which is an isomorphism in codimension one, over the Gorenstein locus of  $\mathcal{H}$ . Fix this homomorphism for the course of the proof.

Now, we show point (1). If  $\omega_X$  is  $S_3$ , then so is  $\omega_{\mathcal{X}/B} \cong p_1^* \omega_X$  by Lemma 4.2. Hence, by Fact 3.2.(2),  $\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}}$  is  $S_2$ . Then, since  $\omega_{\mathcal{H}/B}$  is  $S_2$  by Proposition 3.3(9),  $\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}} \rightarrow \omega_{\mathcal{H}/B}$  is an isomorphism everywhere by [Hartshorne 1994,

Theorems 1.9 and 1.12]. However, for every  $P \in \mathcal{X}_0$ ,

$$\begin{aligned} \text{depth } \omega_{\mathcal{X}/B, P} &= \underbrace{\text{depth } \omega_{X, p_1(P)} + 1}_{\text{Fact 4.1, applied to } \omega_{\mathcal{X}/B} \cong p_1^* \omega_X} \\ &\geq \underbrace{\min\{3, \dim \mathbb{O}_{X, p_1(P)}\} + 1}_{\omega_X \text{ is } S_3} = \min\{4, \dim \mathbb{O}_{\mathcal{X}, P}\}. \end{aligned} \quad (6.1.a)$$

But then, for every  $P \in \mathcal{H}_0$ ,

$$\text{depth } \omega_{\mathcal{H}/B, P} = \underbrace{\text{depth}(\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}})_P}_{\omega_{\mathcal{X}/B} \cong \omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}}} \geq \underbrace{\min\{3, \dim \mathbb{O}_{\mathcal{H}, P}\}}_{\text{Fact 3.2.(2) and (6.1.a)}}$$

which implies point (1) by Proposition 3.3(11).

To prove point (2), denote by  $U$  the open set  $p_1^{-1}(X \setminus (\mathcal{H}_0 \cap \mathcal{H}_\infty)) \subseteq \mathcal{X}$ . This is the set of points, the first coordinates of which are not contained in every element of the pencil  $\mathcal{H} \rightarrow B$ . By Proposition 3.3(4) and 3.3(5), we may replace  $\mathcal{X}$  by  $U$ , or with other words,  $X$  by  $X \setminus (\mathcal{H}_0 \cap \mathcal{H}_\infty)$ . In particular, then  $\mathcal{H}_0 \cap \mathcal{H}_\infty = \emptyset$  and  $P$  is an arbitrary point of  $\mathcal{H}_0$ , such that

$$\text{depth } \omega_{X, P} \not\geq \min\{3, \dim \mathbb{O}_{X, P}\}. \quad (6.1.b)$$

Then all fibers of the projection  $p_1|_{\mathcal{H}} : \mathcal{H} \rightarrow X$  have dimension zero. So, for every  $Q \in \mathcal{H}$ ,

$$\begin{aligned} \text{depth } \omega_{\mathcal{X}/B, Q} &= \underbrace{\text{depth } \omega_{X, p_1(Q)} + 1}_{\text{Fact 4.1, applied to } \omega_{\mathcal{X}/B} \cong p_1^* \omega_X} \\ &\geq \underbrace{\min\{2, \dim \mathbb{O}_{X, p_1(Q)}\} + 1}_{\text{Proposition 3.3.(8)}} = \min\{3, \dim \mathbb{O}_{\mathcal{X}, Q}\}. \end{aligned}$$

Then, repeating the argument of the previous paragraph  $\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}} \cong \omega_{\mathcal{X}/B}$ . Also, at the fixed  $P \in \mathcal{H}_0$ , the following computation estimates the depth more precisely.

$$\begin{aligned} \text{depth } \omega_{\mathcal{X}/B, P} &= \underbrace{\text{depth } \omega_{X, P} + 1}_{\text{Fact 4.1, applied to } \omega_{\mathcal{X}/B} \cong p_1^* \omega_X} \\ &\not\geq \underbrace{\min\{3, \dim \mathbb{O}_{X, P}\} + 1}_{(6.1.b)} = \min\{4, \dim \mathbb{O}_{\mathcal{X}, P}\} \end{aligned} \quad (6.1.c)$$

However, then

$$\text{depth } \omega_{\mathcal{H}/B, P} = \underbrace{\text{depth}(\omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}})_P}_{\omega_{\mathcal{X}/B} \cong \omega_{\mathcal{X}/B}(\mathcal{H})|_{\mathcal{H}}} \not\geq \underbrace{\min\{3, \dim \mathbb{O}_{\mathcal{H}, P}\}}_{\text{by Fact 3.2.(2)}}$$

which concludes the proof by Proposition 3.3(11). □

**Remark 6.2.** The condition of  $\mathcal{I}_{\mathcal{H}, \mathcal{X}}$  being a line bundle in Lemma 6.1 might look superfluous at first sight, since  $\mathcal{H}$  is a hypersurface in  $\mathcal{X}$ . However, according to Section 2, the latter only means that  $\mathcal{H}$  is the zero locus of some special section of a line bundle. That is,  $\mathcal{H}$  or  $\mathcal{H}_b$  for some  $b \in B$  could contain an entire irreducible component of  $\mathcal{X}$  or  $\mathcal{X}_b$ , respectively. Then Proposition 3.6 would not apply. Such situations should definitely be avoided.

The following is the main construction to which Lemma 6.1 is applied in this section.

**Construction 6.3.** Consider a projective cone  $X$  over a variety  $Y$ . Let  $P$  be the vertex of  $X$ . Take two hypersurfaces in  $X$ . The first one  $H$  is a projective cone over a degree  $d$  generic hypersurface  $D$  of  $Y$ . The second one  $\tilde{H}$  is a general degree  $d$  hypersurface of  $X$ . Denote by  $\mathcal{H} \rightarrow B$  the pencil generated by  $H$  and  $\tilde{H}$  (for which  $H = \mathcal{H}_0$  and  $\tilde{H} = \mathcal{H}_\infty$ ). Throughout the paper we allow ourselves to replace this family by its restriction to a small enough open neighborhood of  $0 \in B$ . Furthermore, when we compute stable reduction in Section 8, we will assume that  $d \gg 0$ .

**Lemma 6.4.** *In the situation of Construction 6.3, if  $X$  is  $S_3$  and  $Y$  is  $R_1$ , then*

- (1)  $\mathcal{H}$  and the closed fibers of  $f$  are normal varieties,
- (2)  $\mathcal{I}_{\mathcal{H}, X \times B}$  is a line bundle,
- (3)  $f$  is flat.

*Proof.* We use the notation  $\mathcal{X} := X \times B$ . Since  $Y$  is a variety (that is, integral), so are  $D$ ,  $X$ ,  $\mathcal{X}$ ,  $\mathcal{H}_0$  and  $\mathcal{H}_\infty$ . By the definition of a pencil,  $\mathcal{H}$  is defined by a single nonzero equation locally on  $\mathcal{X}$ . So, since  $\mathcal{X}$  is integral, point (2) follows. Similarly, for every  $b \in B$ ,  $\mathcal{H}_b$  is defined locally by a single nonzero equation locally. Hence by integrality of  $X$ ,  $\mathcal{I}_{\mathcal{H}_b, \mathcal{X}_b}$  is also a line bundle for every  $b \in B$ . Thus, Lemma 3.5 yields point (3).

To prove point (1), note that  $\mathcal{X}$  is  $S_3$  by Lemma 4.2 and by the assumption of the lemma. So, by Fact 3.2,  $\mathcal{H}$  and the closed fibers of  $\mathcal{H}$  are  $S_2$ . (Remember, in Construction 6.3 we allowed ourselves to shrink  $B$  around  $0 \in B$ ). Since  $D$  is general and  $Y$  is  $R_1$ ,  $D$  is  $R_1$  as well by Bertini’s theorem; see [Harris 1992, Theorem 17.16]. Therefore, so is  $H$ . Then, by possibly shrinking  $B$ , each closed fiber of  $\mathcal{H}$  is  $R_1$ . Thus all closed fibers of  $\mathcal{H}$ , and  $\mathcal{H}$  itself, are normal. □

**Theorem 6.5.** *In the situation of Construction 6.3, if  $\dim X \geq 3$ ,  $X$  is  $S_3$ ,  $Y$  is  $R_1$ , and  $\text{depth } \omega_{X, P} = 2$ , then*

$$\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \not\cong \omega_{\mathcal{H}_0}. \tag{6.5.a}$$

*In addition:*

- (1) If  $\omega_X$  is a  $\mathbb{Q}$ -line bundle, then  $\omega_{\mathcal{H}/B}$  is a  $\mathbb{Q}$ -line bundle. In particular then  $\omega_{\mathcal{H}_b}$  is a  $\mathbb{Q}$ -line bundle for all  $b \in B$ .
- (2) If  $X$  is  $S_d$  and  $\text{depth } \mathcal{O}_{X,P} = d$ , then  $\mathcal{H}_b$  is  $S_{d-1}$  for all  $b \in B$ , and

$$\text{depth } \mathcal{O}_{\mathcal{H}_0,P} = d - 1.$$

*Proof.* By Lemma 6.4, we may apply Lemma 6.1(2) to obtain the main statement of the theorem.

To prove addendum (1), note that the normality of  $\mathcal{H}$  and  $\mathcal{H}_b$  for every  $b \in B$ , [Hartshorne 1994, Theorem 1.12] and Proposition 3.6 imply that

$$\omega_{\mathcal{H}_b}^{[n]} \cong (\omega_{\mathcal{H}_b}(\mathcal{H}_b)|_{\mathcal{H}_b})^{[n]} \text{ for any } b \in B, \text{ and } \omega_{\mathcal{H}/B}^{[n]} \cong (\omega_{\mathcal{H}/B}(\mathcal{H})|_{\mathcal{H}})^{[n]} \quad (6.5.b)$$

for all  $n \in \mathbb{Z}$ . Hence if  $\omega_X$  is a  $\mathbb{Q}$ -line bundle, then (6.5.b) implies that so is  $\omega_{\mathcal{H}/B}$  and  $\omega_{\mathcal{H}_b}$  for all  $b \in B$ . To prove (2) we use Fact 3.2 once again.  $\square$

Theorem 1.2 now follows as a corollary:

*Proof of Theorem 1.2.* It follows by combining Proposition 5.4 (setting  $d = j + 1$  and  $l = 2$ ), Construction 6.3 and Theorem 6.5.  $\square$

### 7. Degenerations and Serre’s condition

We turn to proving the statements relating Serre’s condition  $S_d$  to degenerations of flat families. The first half of the section is devoted to Theorem 1.3.

**Remark 7.1.** By the restriction homomorphism  $\omega_{\mathcal{H}/B} \rightarrow \omega_{\mathcal{H}_0}$  we mean any homomorphism obtained as in Proposition 3.3(2).

Theorem 1.3 might look technical, but it applies for example to the special case, when the general fiber is Cohen–Macaulay and the central fiber contains at least one closed point with depth  $n - 1$ .

*Proof of Corollary 1.4 and Proposition 1.5.* Fix a  $2 \leq i < n - 1$ . Consider the projective cone  $X$  given by Proposition 5.4, setting  $d = i + 1$  and  $l = 3$ . Use then Construction 6.3 for  $X$ . By Lemma 6.4, this yields a flat family  $f : \mathcal{H} \rightarrow B$  of normal varieties for which Lemma 6.1(1) applies. That is, the restriction homomorphisms  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_b} \rightarrow \omega_{\mathcal{H}_b}$  are isomorphisms for every  $b \in B$ . Finally, since  $X$  is Cohen–Macaulay outside of  $P$  and  $\text{depth } \mathcal{O}_{X,P} = i + 1$ , by Fact 3.2,  $\mathcal{H}_b$  is Cohen–Macaulay outside of  $P$ , where  $\text{depth } \mathcal{O}_{\mathcal{H}_0,P} = i$ .  $\square$

We also need the following lemma in the proof of Theorem 1.3.

**Lemma 7.2.** *If  $f : \mathcal{H} \rightarrow B$  is a flat morphism of schemes onto a smooth curve,  $\mathcal{F}$  is a coherent  $\mathcal{O}_{\mathcal{H}}$ -module on  $\mathcal{H}$ , and  $P \in \mathcal{H}_0$ , then*

- (1)  $\text{Tor}_{\mathcal{H}}^1(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0})_P \neq 0$  if and only if  $\mathcal{F}$  has an associated component  $W$  such that  $P \in W \subseteq \mathcal{H}_0$ , and

(2)  $\text{Tor}_{\mathcal{H}}^i(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0}) = 0$  for  $i > 1$ .

*Proof.* By restricting  $B$ , we may assume that  $\mathcal{I}_{0,B} \cong \mathcal{O}_B$ . Denote by  $s$  a generator of  $\mathcal{I}_{0,B}$  and consider the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathcal{H}} \xrightarrow{s} \mathcal{O}_{\mathcal{H}} \rightarrow \mathcal{O}_{\mathcal{H}_0} \rightarrow 0. \tag{7.2.a}$$

Then the long exact sequence of  $\text{Tor}_{\mathcal{H}}^\bullet(\mathcal{F}, \cdot)$  applied to (7.2.a) yields

$$\text{Tor}_{\mathcal{H}}^1(\mathcal{F}, \mathcal{O}_{\mathcal{H}}) = 0 \rightarrow \text{Tor}_{\mathcal{H}}^1(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0}) \rightarrow \mathcal{F} \xrightarrow{s} \mathcal{F}.$$

Hence  $\text{Tor}_{\mathcal{H}}^1(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0})_P \neq 0$  if and only if  $s$  annihilates something in  $\mathcal{F}_P$ , if and only if  $\mathcal{F}$  has an associated component  $W$  such that  $P \in W \subseteq \mathcal{H}_0$ .

Another part of the long exact sequence of  $\text{Tor}_{\mathcal{H}}^\bullet(\mathcal{F}, \cdot)$  applied to (7.2.a) yields the following for  $i > 1$ :

$$\text{Tor}_{\mathcal{H}}^i(\mathcal{F}, \mathcal{O}_{\mathcal{H}}) = 0 \rightarrow \text{Tor}_{\mathcal{H}}^i(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0}) \rightarrow \text{Tor}_{\mathcal{H}}^{i-1}(\mathcal{F}, \mathcal{O}_{\mathcal{H}}) = 0.$$

Hence,  $\text{Tor}_{\mathcal{H}}^i(\mathcal{F}, \mathcal{O}_{\mathcal{H}_0}) = 0$  indeed if  $i > 1$ . □

*Proof of Theorem 1.3.* Fix a closed point  $x \in \mathcal{H}_0$  with  $\text{depth } \mathcal{O}_{\mathcal{H}_0,x} = n - 1$ , contained in a component  $W \subseteq \mathcal{H}_0$  of the locus (1.3.a). The locus (1.3.a) is  $\text{supp}(h^{-(n-1)}(\omega_{\mathcal{H}/B}^\bullet))$  by Proposition 3.3(7); hence  $W$  is also an associated component of  $h^{-(n-1)}(\omega_{X/B}^\bullet)$ . Consider an open neighborhood of  $x$ , where every closed point has depth at least  $n - 1$ . Replacing  $\mathcal{H}$  by this neighborhood, all assumptions of the theorem stay valid, and moreover we may assume that every closed point of  $\mathcal{H}$  has depth at least  $n - 1$ . In particular, then

$$h^i(\omega_{\mathcal{H}/B}^\bullet) \neq 0 \iff i = -n \text{ or } -(n - 1). \tag{7.2.b}$$

Define  $\mathcal{E} := h^{-(n-1)}(\omega_{\mathcal{H}/B}^\bullet)$ . By (7.2.b), there is an exact triangle

$$\omega_{\mathcal{H}/B}[n] \rightarrow \omega_{\mathcal{H}/B}^\bullet \rightarrow \mathcal{E}[n - 1] \xrightarrow{+1} . \tag{7.2.c}$$

Applying  $\cdot \otimes_L \mathcal{O}_{\mathcal{H}_0}$  to (7.2.c) and then considering the long exact sequence of cohomology sheaves yields

$$\begin{aligned} h^{-n-1}(\mathcal{E}[n - 1] \otimes_L \mathcal{O}_{\mathcal{H}_0}) &\rightarrow h^{-n}(\omega_{\mathcal{H}/B}[n] \otimes_L \mathcal{O}_{\mathcal{H}_0}) \rightarrow h^{-n}(\omega_{\mathcal{H}/B}^\bullet \otimes_L \mathcal{O}_{\mathcal{H}_0}) \\ &\rightarrow h^{-n}(\mathcal{E}[n - 1] \otimes_L \mathcal{O}_{\mathcal{H}_0}) \rightarrow h^{-n+1}(\omega_{\mathcal{H}/B}[n] \otimes_L \mathcal{O}_{\mathcal{H}_0}), \end{aligned} \tag{7.2.d}$$

where

- $h^{-n-1}(\mathcal{E}[n - 1] \otimes_L \mathcal{O}_{\mathcal{H}_0}) \cong \text{Tor}_{\mathcal{H}}^2(\mathcal{E}, \mathcal{O}_{\mathcal{H}_0}) = 0$  by Lemma 7.2,
- $h^{-n}(\omega_{\mathcal{H}/B}[n] \otimes_L \mathcal{O}_{\mathcal{H}_0}) \cong \omega_{\mathcal{H}/B}|_{\mathcal{H}_0}$ ,
- $h^{-n}(\omega_{\mathcal{H}/B}^\bullet \otimes_L \mathcal{O}_{\mathcal{H}_0}) \cong h^{-n}(\omega_{\mathcal{H}_0}^\bullet) \cong \omega_{\mathcal{H}_0}$  by Proposition 3.3(1),
- $h^{-n}(\mathcal{E}[n - 1] \otimes_L \mathcal{O}_{\mathcal{H}_0}) \cong \text{Tor}_{\mathcal{H}}^1(\mathcal{E}, \mathcal{O}_{\mathcal{H}_0})$  and

- $h^{-n+1}(\omega_{\mathcal{H}/B}[n] \otimes_L \mathbb{O}_{\mathcal{H}_0}) \cong h^1(\omega_{\mathcal{H}/B} \otimes_L \mathbb{O}_{\mathcal{H}_0}) = 0$  since  $\cdot \otimes_L \mathbb{O}_{\mathcal{H}_0}$  is a left derived functor, so  $\omega_{\mathcal{H}/B} \otimes_L \mathbb{O}_{\mathcal{H}_0}$  is supported in negative cohomological degrees.

Therefore, (7.2.d) is isomorphic to the exact sequence

$$0 \rightarrow \omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0} \rightarrow \mathcal{T}or_{\mathcal{H}}^1(\mathcal{E}, \mathbb{O}_{\mathcal{H}_0}) \rightarrow 0.$$

Since  $\mathcal{E}$  has an associated component through  $x$  contained in  $\mathcal{H}_0$ , we know that  $\mathcal{T}or_{\mathcal{H}}^1(\mathcal{E}, \mathbb{O}_{\mathcal{H}_0})_x \neq 0$  by Lemma 7.2, which concludes our proof.  $\square$

Having finished the proof of Theorem 1.3, the rest of the section is devoted its consequence, Theorem 1.6. See Section 1 for its motivation.

*Proof of Theorem 1.6.* Since the statement of the theorem is local, we may assume  $X$  is affine and hence quasiprojective. Restricting to a sufficiently small neighborhood of a point with depth  $n - 1$ , all assumptions of the theorem stay valid and we may assume that all closed points of  $X$  have depth at least  $n - 1$ . We use the notation  $\mathcal{X} := X \times B$ . Let  $X = \bigcup_{i=1}^r X_i$  be the decomposition into irreducible components.

Consider a pencil  $f : \mathcal{H} \rightarrow B = \mathbb{P}^1$  of hypersurfaces of  $X$  such that

- (1)  $\mathcal{H}_0$  contains the entire non-Gorenstein locus,
- (2)  $\emptyset \neq \mathcal{H}_0 \cap X_j \neq X_j$  for every  $1 \leq j \leq r$ ,
- (3)  $\mathcal{H}_\infty$  is a general hypersurface.

In particular then,

$$(\mathcal{H}_0 \setminus \mathcal{H}_\infty) \cap X_j \neq \emptyset \quad \text{for every } 1 \leq j \leq r. \tag{7.2.e}$$

By definition of the pencil, if  $P \in \mathcal{H}_0 \setminus \mathcal{H}_\infty$ , then  $P \notin \mathcal{H}_b$  for any  $b \neq 0$ . Hence assumption (2) and (7.2.e) imply that for all  $b \in B$ , there is a point of  $X_j$  not contained in  $\mathcal{H}_b$ . Note now, that since  $X$  is  $S_1$ , by Lemma 4.2, so is  $\mathcal{X}$ . In particular, then all associated points of  $X$  and  $\mathcal{X}$  are general points of components. So, since none of the  $\mathcal{H}_b$  contains any of the  $X_j$ ,  $\mathcal{I}_{\mathcal{X}, \mathcal{X}}$  and  $\mathcal{I}_{\mathcal{H}_b, \mathcal{H}_b}$  for every  $b \in B$  have nonzero divisor local generators and hence are line bundles. Then  $\mathcal{H}$  is flat over  $B$  by Lemma 3.5.

Define the loci

$$Z := \overline{\{x \in X \mid x \text{ is closed, depth } \mathbb{O}_{X,x} = n - 1\}},$$

$$W := \overline{\{x \in X \mid x \text{ is closed, depth } \mathbb{O}_{\mathcal{H}_f(x),x} = n - 2\}}.$$

By construction and by Fact 3.2,  $W_0 = Z$  and  $W = (p^{-1}Z)_{\text{red}}$ , where  $p : \mathcal{H} \rightarrow X$  is the natural projection. Let  $Z'$  be an irreducible component of  $Z$  of the highest dimension. By the choice of  $\mathcal{H}_0$  and  $\mathcal{H}_\infty$ , we have  $Z' \subseteq \mathcal{H}_0$ , and  $Z' \not\subseteq \mathcal{H}_\infty$ . Furthermore,  $\mathcal{H}_\infty$  does not contain any of the irreducible components of  $Z$ . Hence, the general fiber of the map  $W \rightarrow B$  will have dimension at most  $\dim Z' - 1$ . So,



$W$  has dimension  $\dim Z'$ . Hence,  $Z' \subseteq W_0$  is an irreducible component of  $W$ . In particular, by Theorem 1.3, the restriction morphism  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is not an isomorphism.

On the other hand assume that  $\omega_X$  is  $S_3$ . Since  $X$  is  $G_2$ ,  $\mathcal{H}$  and  $\mathcal{H}_b$  are  $G_1$  for every  $b \in B$ . In fact,  $\mathcal{H}$ , and  $\mathcal{H}_b$  for a general  $b \in B$ , are  $G_2$  also, but for  $\mathcal{H}_0$  only  $G_1$  can be guaranteed. Also,  $X$  is  $S_3$  by assumption and  $\mathcal{X}$  is  $S_3$  because of Lemma 4.2. Then  $\mathcal{H}$  and  $\mathcal{H}_b$  are  $S_2$  for every  $b \in B$  by Fact 3.2. That is, we may apply Lemma 6.1(1), which states that the restriction homomorphism  $\omega_{\mathcal{H}/B}|_{\mathcal{H}_0} \rightarrow \omega_{\mathcal{H}_0}$  is an isomorphism. This is a contradiction; hence  $\omega_X$  cannot be  $S_3$ .  $\square$

### 8. Stable reduction

In Construction 6.3, although the general fiber of  $\mathcal{H} \rightarrow B$  has mild, that is, log canonical, singularities,  $\mathcal{H}_0$  is very singular. The failure of base change for  $\omega_{\mathcal{H}/B}$  implies that by [Kollár and Kovács 2010, Theorem 7.9]  $\mathcal{H}_0$  is not Du Bois. By [ibid., Theorem 1.4], it is also not log canonical. In this section, we compute the stable limit of  $\mathcal{H} \rightarrow B$ . It is the limit at 0 of some stable family  $\mathcal{H}' \rightarrow \tilde{B}$ . This family has two important properties. First,  $\mathcal{H} \times_B \tilde{B}|_{\tilde{B} \setminus \{0\}} \cong \mathcal{H}'|_{\tilde{B} \setminus \{0\}}$  for a finite cover  $\phi : (\tilde{B}, 0) \rightarrow (B, 0)$  totally ramified at 0. Second,  $(\mathcal{H}')_0$  is log canonical, and hence by [ibid., Theorems 1.4 and 7.9],  $\omega_{\mathcal{H}'/\tilde{B}}$  commutes with base change. So,  $(\mathcal{H}')_0$  is the “right” limit of  $\mathcal{H}$ , and the incompatibility of Theorem 1.2 can be thought of as a consequence of using the wrong limit in Construction 6.3.

**Proposition 8.1.** *Assuming that  $Y$  is smooth, the stable limit of Construction 6.3 is the  $d$ -fold cyclic cover of  $Y$  ramified exactly over  $D$ , with eigen-line bundles  $\mathcal{O}_Y(-i)$  for  $i = 0, \dots, d - 1$ .*

*Proof.* First, shrink  $B$  if necessary so that  $\infty \notin B$  and that every fiber apart from  $\mathcal{H}_0$  is log canonical. This is possible because the general fiber of  $\mathcal{H}$  is smooth by Bertini’s theorem. Also, since we assumed that  $d \gg 0$ , the family  $\mathcal{H} \rightarrow B$  has canonically polarized fibers and hence is stable over  $B^* := B \setminus \{0\}$ . Define  $\mathcal{X} := X \times B$ .

The closed embedding  $Y \subseteq \mathbb{P}^{N-1}$  induces a natural closed embedding  $X \subseteq \mathbb{P}^N$ . This yields very ample line bundles  $\mathcal{O}_{\mathbb{P}^N}(1)$  and  $\mathcal{O}_X(1)$ . Then,  $\mathcal{H}$  is the zero locus of a section  $f_0 + tf_\infty$  of  $\mathcal{O}_{\mathcal{X}}(d) := p_1^* \mathcal{O}_X(d)$  for some  $f_0, f_\infty \in H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(d))$ , as explained in Section 2.

Choose a basis  $z_0, \dots, z_N$  of  $H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(1))$ , such that  $z_0, \dots, z_{N-1}$  form a basis of  $H^0(\mathbb{P}^{N-1}, \mathcal{O}_{\mathbb{P}^{N-1}}(1))$ . Then  $f_0$  and  $f_\infty$  correspond to degree  $d$  homogeneous polynomials in variables  $z_0, \dots, z_{N-1}$  and  $z_0, \dots, z_N$ , respectively. Also, the fact that  $P \notin \mathcal{H}_\infty$  implies that the coefficient of  $z_N^d$  in  $f_\infty$  is nonzero, say  $c$ .

Let  $\phi : (\tilde{B}, 0) \rightarrow (B, 0)$  be the degree  $d$  cyclic cover branched only at 0, where it is totally ramified, and let  $s$  be a local parameter of  $\tilde{B}$  at 0, such that  $s^d = t$ .

Consider the subscheme  $\mathcal{H}' \subseteq \mathcal{X} \times_B \tilde{B} =: \mathcal{X}_\phi$  defined by

$$f_0(z_0, \dots, z_{N-1}) + s^d f_\infty(z_0, \dots, z_{N-1}, \frac{1}{s}z_N) \in H^0(\mathcal{X}_\phi, \mathcal{O}_{\mathcal{X}_\phi}(d)),$$

where  $\mathcal{O}_{\mathcal{X}_\phi}(d)$  is the pullback of  $\mathcal{O}_{\mathcal{X}}(d)$  to  $\mathcal{X}_\phi$ .

By the uniqueness of stable limit,  $\mathcal{H}'$  is a stable reduction of  $\mathcal{H}$  (that is, a stable family isomorphic generically to the pullback of  $\mathcal{H}$ ), if

- (1)  $(\mathcal{H}')_0$  is a canonically polarized manifold, and
- (2)  $\mathcal{H}'|_{\tilde{B}^*} \cong \mathcal{H}_\phi|_{\tilde{B}^*}$ , where  $\mathcal{H}_\phi := \mathcal{H} \times_B \tilde{B}$  and  $\tilde{B}^* := \tilde{B} \setminus \{0\}$ .

To prove point (1), notice that  $(\mathcal{H}')_0$  is defined by the zero locus of  $s$  on  $\mathcal{H}'$  or equivalently by the zero locus of the following section of  $\mathcal{O}_X(d)$  on  $X$ :

$$f_0(z_0, \dots, z_{N-1}) + cz_N^d.$$

Hence it is the cyclic cover of  $Y$  of degree  $d$  branched along  $D$  with eigensheaves  $\mathcal{O}_Y(-i)$  for  $0 \leq i \leq d - 1$ . So first, it is smooth by [Kollár and Mori 1998, Lemma 2.51]. Second, since  $(\mathcal{H}')_0$  is contained in the smooth part of  $X$ , we have  $\omega_{(\mathcal{H}')_0} \cong \omega_X(d)|_{(\mathcal{H}')_0}$  by Proposition 3.6 and it is a line bundle. So, since  $d \gg 0$ ,  $(\mathcal{H}')_0$  is a canonically polarized manifold indeed.

To prove point (2), notice that the equation of  $\mathcal{H}_\phi$  in  $\mathcal{X}_\phi$  is

$$f_0(z_0, \dots, z_{N-1}) + s^d f_\infty(z_0, \dots, z_{N-1}, z_N) \in H^0(\mathcal{X}_\phi, \mathcal{O}_{\mathcal{X}_\phi}(d)).$$

Hence,  $\mathcal{H}_\phi|_{\tilde{B}^*} \cong \mathcal{H}'|_{\tilde{B}^*}$  via the isomorphism induced by the following automorphism of  $\mathbb{P}^N \times \tilde{B}^*$ :

$$(z_0, \dots, z_{N+1}, z_N) \mapsto (z_0, \dots, z_{N-1}, sz_N).$$

We proved both points (1) and (2). Consequently,  $\mathcal{H}'$  is a stable reduction of  $\mathcal{H}$  indeed. Through the course of the proof of point (1), we also proved that  $(\mathcal{H}')_0$  is indeed the cyclic cover described in the statement of the proposition. □

### Acknowledgements

The discussion contains ideas that originated from János Kollár and my advisor Sándor Kovács. I would like to thank both of them for their help. I would also like to thank Joseph Lipman for useful comments.

### References

[Bruns and Herzog 1993] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1993. MR 95h:13020 Zbl 0788.13005

[Conrad 2000] B. Conrad, *Grothendieck duality and base change*, Lecture Notes in Mathematics **1750**, Springer, Berlin, 2000. MR 2002d:14025 Zbl 0992.14001

- [Grothendieck 1965] A. Grothendieck, “Éléments de géométrie algébrique, IV: Étude locale des schémas et des morphismes de schémas, II”, *Inst. Hautes Études Sci. Publ. Math.* **24** (1965), 1–231. MR 33 #7330 Zbl 0135.39701
- [Hacon and Kovács 2010] C. D. Hacon and S. J. Kovács, *Classification of higher dimensional algebraic varieties*, Oberwolfach Seminars **41**, Birkhäuser, Basel, 2010. MR 2011f:14025 Zbl 1204.14001
- [Harris 1992] J. Harris, *Algebraic geometry: A first course*, Graduate Texts in Mathematics **133**, Springer, New York, 1992. MR 93j:14001 Zbl 0779.14001
- [Hartshorne 1966] R. Hartshorne, *Residues and duality*, Lecture Notes in Mathematics **20**, Springer, Berlin, 1966. MR 36 #5145 Zbl 0212.26101
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, New York, 1977. MR 57 #3116 Zbl 0367.14001
- [Hartshorne 1994] R. Hartshorne, “Generalized divisors on Gorenstein schemes”, *K-Theory* **8**:3 (1994), 287–339. MR 95k:14008 Zbl 0826.14005
- [Hassett 2001] B. Hassett, “Stable limits of log surfaces and Cohen–Macaulay singularities”, *J. Algebra* **242**:1 (2001), 225–235. MR 2002f:14051 Zbl 1064.14037
- [Hassett and Kovács 2004] B. Hassett and S. J. Kovács, “Reflexive pull-backs and base extension”, *J. Algebraic Geom.* **13**:2 (2004), 233–247. MR 2005b:14028 Zbl 1081.14017
- [Kollár 2008] J. Kollár, “Hulls and husks”, preprint, 2008. arXiv 0805.0576
- [Kollár 2010] J. Kollár, “Moduli of varieties of general type”, preprint, 2010. arXiv 1008.0621
- [Kollár 2011] J. Kollár, “A local version of the Kawamata–Viehweg vanishing theorem”, *Pure Appl. Math. Q.* **7**:4 (2011), 1477–1494. MR 2918170 Zbl 06107785
- [Kollár and Kovács 2010] J. Kollár and S. J. Kovács, “Log canonical singularities are Du Bois”, *J. Amer. Math. Soc.* **23**:3 (2010), 791–813. MR 2011m:14061 Zbl 1202.14003
- [Kollár and Mori 1998] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics **134**, Cambridge University Press, 1998. MR 2000b:14018 Zbl 0926.14003
- [Kovács 2011] S. Kovács, “Irrational centers”, *Pure Appl. Math. Q.* **7**:4 (2011), 1495–1515. MR 2918171 Zbl 06107786
- [Lipman and Hashimoto 2009] J. Lipman and M. Hashimoto, *Foundations of Grothendieck duality for diagrams of schemes*, Lecture Notes in Mathematics **1960**, Springer, Berlin, 2009. MR 2010b:18001 Zbl 1163.14001
- [Matsumura 1989] H. Matsumura, *Commutative ring theory*, 2nd ed., Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, 1989. MR 90i:13001 Zbl 0666.13002
- [Neeman 1996] A. Neeman, “The Grothendieck duality theorem via Bousfield’s techniques and Brown representability”, *J. Amer. Math. Soc.* **9**:1 (1996), 205–236. MR 96c:18006 Zbl 0864.14008
- [Viehweg 1995] E. Viehweg, *Quasi-projective moduli for polarized manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* **30**, Springer, Berlin, 1995. MR 97j:14001 Zbl 0844.14004

Communicated by Ravi Vakil

Received 2011-06-09

Revised 2012-01-26

Accepted 2012-03-09

pzs@princeton.edu

*Department of Mathematics, Princeton University, Fine Hall, Washington Road, Princeton, NJ 08540, United States*  
[www.math.princeton.edu/~pzs](http://www.math.princeton.edu/~pzs)

# Algebra & Number Theory

msp.org/ant

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Karl Rubin	University of California, Irvine, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Ehud Hrushovski	Hebrew University, Israel	Bernd Sturmfels	University of California, Berkeley, USA
Craig Huneke	University of Virginia, USA	Richard Taylor	Harvard University, USA
Mikhail Kapranov	Yale University, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Efim Zelmanov	University of California, San Diego, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.


The subscription price for 2013 is US \$200/year for the electronic version, and \$350/year (+\$40, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 7    No. 2    2013

---

The system of representations of the Weil–Deligne group associated to an abelian variety	243
RUTGER NOOT	
Fourier–Jacobi coefficients of Eisenstein series on unitary groups	283
BEI ZHANG	
The phase limit set of a variety	339
MOUNIR NISSE and FRANK SOTTILE	
Base change behavior of the relative canonical sheaf related to higher dimensional moduli	353
ZSOLT PATAKFALVI	
Two ways to degenerate the Jacobian are the same	379
JESSE LEO KASS	
Arithmetic motivic Poincaré series of toric varieties	405
HELENA COBO PABLOS and PEDRO DANIEL GONZÁLEZ PÉREZ	
Maximal ideals and representations of twisted forms of algebras	431
MICHAEL LAU and ARTURO PIANZOLA	
Higher Chow groups of varieties with group action	449
AMALENDU KRISHNA	



1937-0652(2013)7:2;1-B