

Algebra & Number Theory

Volume 8

2014

No. 2

Wild models of curves

Dino Lorenzini



Wild models of curves

Dino Lorenzini

Let K be a complete discrete valuation field with ring of integers \mathcal{O}_K and algebraically closed residue field k of characteristic $p > 0$. Let X/K be a smooth proper geometrically connected curve of genus $g > 0$ with $X(K) \neq \emptyset$ if $g = 1$. Assume that X/K does not have good reduction and that it obtains good reduction over a Galois extension L/K of degree p . Let $\mathcal{Y}/\mathcal{O}_L$ be the smooth model of X_L/L . Let $H := \text{Gal}(L/K)$.

In this article, we provide information on the regular model of X/K obtained by desingularizing the wild quotient singularities of the quotient \mathcal{Y}/H . The most precise information on the resolution of these quotient singularities is obtained when the special fiber \mathcal{Y}_k/k is ordinary. As a corollary, we are able to produce for each odd prime p an infinite class of wild quotient singularities having pairwise distinct resolution graphs. The information on the regular model of X/K also allows us to gather insight into the p -part of the component group of the Néron model of the Jacobian of X .

1. Introduction

Let K be a complete discrete valuation field with valuation v , ring of integers \mathcal{O}_K and residue field k of characteristic $p > 0$, assumed to be *algebraically closed*. Let X/K be a smooth proper geometrically connected curve of genus $g > 0$ with $X(K) \neq \emptyset$ if $g = 1$.

Assume that X/K does not have good reduction and that it obtains good reduction over a Galois extension L/K . Let $\mathcal{Y}/\mathcal{O}_L$ be the smooth model of X_L/L . Let $H := \text{Gal}(L/K)$, and let $\mathcal{Z}/\mathcal{O}_K$ denote the quotient \mathcal{Y}/H . A regular model for X/K can be obtained by resolving the singularities of the scheme \mathcal{Z} . Our goal is to obtain information on this regular model when p divides $[L : K]$. Since the presence of wild ramification renders the subject quite challenging, we will restrict our attention in this article to the case where $[L : K] = p$.

Lorenzini was supported by NSF Grant 0902161.

MSC2010: primary 14G20; secondary 14G17, 14K15, 14J17.


Keywords: model of a curve, ordinary curve, cyclic quotient singularity, wild ramification, arithmetical tree, resolution graph, component group, Néron model.

Beyond our interest in models of curves per se, our motivation for understanding these regular models is twofold. First, since \mathcal{X} is obtained by desingularizing certain quotient singularities, we hope to gain more insight in the general theory of resolutions of wild quotient singularities by producing interesting classes of examples where the singularities can be resolved explicitly. Second, since from a regular model of the curve one can compute much of the Néron model of its Jacobian, we hope to bring new insight into the structure of the rather mysterious p -part of the component group of the Néron model of a general abelian variety from an increased understanding of the special case of Jacobians of curves.

Let us introduce some notation needed to state our theorems. Let σ denote a generator of $H := \text{Gal}(L/K)$. Denote also by σ the automorphism of \mathcal{Y}_k induced by the action of H on \mathcal{Y} . The scheme \mathcal{Z} is singular exactly at the images Q_1, \dots, Q_d of the ramification points P_1, \dots, P_d of the map $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ (5.2). Consider the regular model $\mathcal{X} \rightarrow \mathcal{Z}$ obtained from \mathcal{Z} by a minimal desingularization. Let $\mathcal{X}' \rightarrow \mathcal{X}$ denote the regular model of X/K minimal with the property that \mathcal{X}'_k has smooth components and normal crossings. Let f denote the composition $\mathcal{X}' \rightarrow \mathcal{Z}$. Let C_0/k denote the strict transform in \mathcal{X}' of the irreducible closed subscheme $\mathcal{Z}_k^{\text{red}}$ of \mathcal{Z} . Let D_1, \dots, D_d denote the irreducible components of \mathcal{X}'_k that meet C_0 . Let r_i denote the multiplicity of D_i , $i = 1, \dots, d$, in \mathcal{X}'_k .

Recall that to any connected curve $\bigcup_{\ell=1}^n C_\ell$ on a regular model \mathcal{X} we associate a graph G as follows: the vertices are the irreducible components C_ℓ , and in G , the vertices C_i and C_j ($i \neq j$) are linked by exactly $(C_i \cdot C_j)_\mathcal{X}$ edges, where $(C_i \cdot C_j)_\mathcal{X}$ denotes the intersection number of C_i and C_j on the regular scheme \mathcal{X} . Recall that the *degree* of a vertex v of a graph is the number of edges attached to v . A *node* on a graph is a vertex of degree at least 3. A vertex of degree 1 is a *terminal vertex*. A *chain* is a subgraph of G with vertices C_0, C_1, \dots, C_n , $n \geq 1$, such that C_i is linked to C_{i+1} by exactly one edge in G when $i = 0, \dots, n - 1$ and the degree of C_i is 2 when $i = 1, \dots, n - 1$. If the chain contains a terminal vertex (which can only be C_0 or C_n), the chain is called a *terminal chain*.

Let G denote the graph associated with \mathcal{X}'_k . We assume $d \geq 1$. For each $i = 1, \dots, d$, let G_{Q_i} denote the graph associated with the curve $f^{-1}(Q_i)$. In particular, D_i corresponds to a vertex of G_{Q_i} . We have the following configuration on the graph G (where a positive integer next to a vertex denotes the multiplicity of the corresponding irreducible component in \mathcal{X}'):




Theorem 5.3. *Let X/K be a curve with potentially good reduction after a wildly ramified extension L/K of degree p , as above. Keep the above notation. Then, for all $i = 1, \dots, d$, the graph G_{Q_i} contains a node of G and p divides r_i .*

In contrast, when H is of prime order $q \neq p$, then it is known that $q > r_i$ and that the graph G_{Q_i} does not contain a node of G . In particular, when L/K is tame and $d \geq 3$, the graph G has only a single node, the component C_0 (see, e.g., [Lorenzini 1990a, Theorem 2.1]).

We propose in 6.1 a combinatorial measure $\gamma_{Q_i} g_{Q_i}$ of the complexity of the graph G_{Q_i} , which we conjecturally relate in 6.2 to the higher ramification data of the morphism $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$. This conjectural relationship expresses the fact that the graph G_{Q_i} is “complicated” only if the higher ramification above Q_i is “large”. We prove this conjecture in the ordinary case (Theorem 6.4).

Recall that a smooth proper curve Y/k of genus g is called *ordinary* if its Jacobian J/k is an ordinary abelian variety (that is, $J(k)$ has exactly p^g points of order dividing p). When \mathcal{Y}_k is ordinary, the morphism $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ has the smallest possible ramification data at each Q_i (2.2), and in this case, we can use Theorem 5.3 to describe the graph G_{Q_i} explicitly, as in the following theorem, whose statement is slightly strengthened in the version given in Section 6. In the graph below, a bullet \bullet represents an irreducible component of the desingularization of Q_i . A negative number next to a vertex is the self-intersection of the component. A positive number next to a vertex is the multiplicity of the corresponding component in \mathcal{X}'_k .

Theorem (see Theorem 6.8). *Let X/K be a curve with potentially good reduction after a Galois extension L/K of degree p , as above. Assume that \mathcal{Y}_k ordinary. Then, for all $i = 1, \dots, d$, we have $r_i = p$ and G_{Q_i} is a graph with a single node C_i of degree 3:*




The intersection matrix $N(p, \alpha_i, r_1(i))$ of the resolution of Q_i is uniquely determined as in 4.7 by the two integers α_i and $r_1(i)$ with $1 \leq r_1(i) < p$. The integer α_i denotes the number of vertices of self-intersection -2 (including the node C_i) on the chain in G_{Q_i} connecting the node C_0 to the single node C_i of G_{Q_i} , and the integer α_i is divisible by p .

To further determine the regular model, one would need to determine explicitly the integers α_i and $r_1(i)$. We address this issue in [Lorenzini 2013b]. In all cases where we have been able to compute α_i and $r_1(i)$, we found them to be related to the valuation of the different of L/K . More precisely, let $(s_{L/K} + 1)(p - 1)$

denote the valuation of the different of L/K . In [Lorenzini 2013b, Theorem 1.1], we present some instances where $\alpha_i = ps_{L/K}$ and $r_1(i) \equiv -s_{L/K}^{-1}$ modulo p . We also show in [Lorenzini 2013b, Theorem 4.1] that the singularities Q_i are rational.

Remark 1.1. The same type of intersection matrix, $N(p, \alpha_i, r_1(i))$, also occurs in the resolution of the singularities of the model \mathcal{Z} when X/K has genus $p - 1$ and $\text{Jac}(X)/K$ has purely toric reduction after an extension of degree p [Lorenzini 2010, Theorem 2.2].

Remark 1.2. The special fiber of the model $\mathcal{X}/\mathcal{O}_K$ of X/K in Theorem 6.8 has thus a graph with a central vertex to which d branches are attached, of the form described below, where we picture the case $d = 4$.



Fix any $d > 1$. We establish in Theorem 6.8 and Example 6.13 the existence of some field K of residue characteristic $p > 0$ and of some smooth proper curve X/K with a regular model whose special fiber has a graph of the above type. This is clearly a weak existence result, but our understanding of models in the presence of wild ramification is so limited that even this weak existence result does not follow from the general existence results of Viehweg [1977] and Winters [1974].

An immediate but surprising corollary to Theorem 6.8 is as follows.

Corollary (see Corollary 6.10). *Let X/K be a curve of genus $g > 1$ with potentially good reduction after a Galois extension L/K of degree p , as above. Assume that \mathcal{Y}_k is ordinary. Then $X(K) \neq \emptyset$.*

The information on the regular model of X/K obtained in Theorem 6.8, while incomplete to fully describe the special fiber of the model, suffices to compute several invariants of arithmetical interest. For instance, the set of components of multiplicity 1 on the special fiber of the model is determined, and this information is one of the ingredients needed to apply the Chabauty–Coleman method to bound the number of \mathbb{Q} -rational points on a curve X/\mathbb{Q} using the reduction at a small prime p , as in [Lorenzini and Tucker 2002, Theorem 1.1]. Let A/K denote the Jacobian of X/K with Néron model $\mathcal{A}/\mathcal{O}_K$ and component group $\Phi_{A/K}$. The information obtained in Theorem 6.8 suffices to compute $\Phi_{A/K}$ and a new canonical subgroup $\Phi_{A/K}^0$ of $\Phi_{A/K}$ that we now define.

1.3. Let A/K be an abelian variety with Néron model $\mathcal{A}/\mathcal{O}_K$. Let L/K be any finite extension, and let $\mathcal{A}'/\mathcal{O}_L$ denote the Néron model of A_L/L . Denote by

$$\eta : \mathcal{A} \times_{\mathcal{O}_K} \mathcal{O}_L \rightarrow \mathcal{A}'$$

the canonical map induced by the functoriality property of Néron models. The special fiber \mathcal{A}_k is an extension of a finite group $\Phi_{A/K}$, called the *group of components*, by the *connected component of zero* \mathcal{A}_k^0 of \mathcal{A}_k :

$$0 \rightarrow \mathcal{A}_k^0 \rightarrow \mathcal{A}_k \rightarrow \Phi_{A/K} \rightarrow 0.$$

Assume that A_L/L has semistable reduction, and consider the natural map $\Phi_{A/K} \rightarrow \mathcal{A}'_k/\eta(\mathcal{A}_k^0)$. We let

$$\Phi_{A/K}^0 := \text{Ker}(\Phi_{A/K} \rightarrow \mathcal{A}'_k/\eta(\mathcal{A}_k^0)).$$

The subgroup $\Phi_{A/K}^0$ does not depend on the choice of such an extension L/K and is functorial in A . Our interest in this subgroup stems from the following conjectures.

When A/K has potentially good reduction and, more generally, when the toric rank of \mathcal{A}_k^0 is trivial, we conjecture that the order of the group $\Phi_{A/K}$ is bounded by a constant depending only on the dimension g of A/K [Lorenzini 1990b, p. 146]. This statement is true when A/K is a Jacobian [Lorenzini 1990b, Theorem 2.4] and for the prime-to- p part of $\Phi_{A/K}$ [Lorenzini 1993, Theorem 2.15]. Since $[L : K]^2$ kills the group $\Phi_{A/K}$ when the toric rank of \mathcal{A} is trivial [Liu and Lorenzini 2001, Proposition 1.8], we find that, to prove the conjecture that $\Phi_{A/K}$ is bounded by a constant depending only on g , it suffices to prove that the minimal number of generators of $\Phi_{A/K}$ can be bounded by a constant depending on g only. We guess, under the above hypotheses, that $\Phi_{A/K}$ can be generated by $2g$ elements.

Assume now that A/K has potentially good reduction. The p -torsion in \mathcal{A}'_k can always be generated by at most g elements. Thus, the above conjecture is proved if the p -part of the kernel $\Phi_{A/K}^0$ can be generated by a number of elements bounded by a constant depending on g only (possibly $2g$). In the ordinary case, where the p -torsion in \mathcal{A}'_k is minimally generated by g elements, one may wonder if $\Phi_{A/K}^0$ can also be generated by g elements. Our next corollary gives some evidence that this latter question may have a positive answer for all abelian varieties with potentially good ordinary reduction.

Let A/K be the Jacobian of a curve X/K with $X(K) \neq \emptyset$. We denote by $\langle \cdot, \cdot \rangle : \Phi_{A/K} \times \Phi_{A/K} \rightarrow \mathbb{Q}/\mathbb{Z}$ Grothendieck's pairing, which is nondegenerate [Bosch and Lorenzini 2002, Theorem 4.6]. We denote by $(\Phi_{A/K}^0)^\perp$ the orthogonal of $\Phi_{A/K}^0$ under Grothendieck's pairing.

Corollary (see Corollary 6.12). *Let A/K be the Jacobian of a curve X/K of genus $g > 1$ having potentially good ordinary reduction after a Galois extension L/K of degree p , as above. Then $\Phi_{A/K}$ is a $\mathbb{Z}/p\mathbb{Z}$ -vector space of dimension $2d - 2$, and $\Phi_{A/K}^0$ is a subspace of dimension $d - 1$. Moreover, $\Phi_{A/K}^0 = (\Phi_{A/K}^0)^\perp$.*

It is natural in view of [Corollary 6.12](#) to wonder whether the same result holds for all principally polarized abelian varieties A/K having potentially good ordinary reduction after a Galois extension L/K of degree p . We may also wonder, for any principally polarized abelian variety A/K with potentially good reduction, whether the order of $\Phi_{A/K}^0 \cap (\Phi_{A/K}^0)^\perp$ can be bounded by a constant depending only on the p -rank of \mathcal{A}'_k . We hope to return to these questions in the future.

1.4. Our explicit computation of a regular model of a curve having potentially good ordinary reduction also has an application to quotient singularities. Our current understanding of wild $\mathbb{Z}/p\mathbb{Z}$ -quotient singularities of surfaces is quite limited, and few explicit examples are known (see, e.g., [\[Artin 1975\]](#), [\[Katsura 1978\]](#) for $p = 2$ and [\[Peskin 1983\]](#) for $p = 3$). In contrast to the case of a tame cyclic quotient singularity, where the number of possible resolution graphs is finite once the order of the group is fixed, we show below that, for any fixed odd prime p , there are infinitely many graphs that can occur as the resolution graphs of a wild $\mathbb{Z}/p\mathbb{Z}$ -quotient singularity in both mixed characteristic and in the equicharacteristic case. The analogous result when $p = 2$ is discussed in [\[Lorenzini 2013a, Theorem 4.1\]](#).

Corollary 6.14. *Fix any odd prime p . For each integer $m > 0$, there exist a 2-dimensional regular local ring B of equicharacteristic p endowed with an action of $H := \mathbb{Z}/p\mathbb{Z}$ and a 2-dimensional regular local ring B' of mixed characteristic $(0, p)$ endowed with an action of $\mathbb{Z}/p\mathbb{Z}$ such that $\text{Spec } B^H$ and $\text{Spec}(B')^H$ are singular exactly at their closed point and the graphs associated with a minimal resolution of $\text{Spec } B^H$ and $\text{Spec}(B')^H$ have one node and more than m vertices.*

This article is organized as follows. The proof of [Theorem 5.3](#), in [Section 5](#), is of a global nature and includes in particular a study of the natural map $\Phi_{A/K} \rightarrow \mathcal{A}'_k/\eta(\mathcal{A}_k^0)$. The proof uses two auxiliary results of independent interest. The first result, [Proposition 2.5](#), is discussed in [Section 2](#) and is a relation between torsion points in a quotient of two Jacobians. This proposition is one place in our arguments where the tame and wild cases can be seen to differ in an explicit way. The second result, [Proposition 3.5](#), is the main result of [Section 3](#) and is a general relation between elements in the component group Φ_M of an arithmetical tree.

[Section 4](#) presents further results of a combinatorial nature on arithmetical trees that are needed in the proof of [Theorem 6.8](#). [Section 6](#) contains the proof of [Theorem 6.8](#) and of its applications.

2. Cyclic morphisms and torsion

Let k be an algebraically closed field of characteristic p . Let $f : D \rightarrow C$ be a ramified Galois morphism of smooth connected projective curves over k . Our main result in this section is [Proposition 2.5](#), which will be applied to the case of the quotient morphism $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ in the course of the proof of [Theorem 5.3](#).

2.1. Assume that the Galois group H of f is cyclic of degree q^s with q prime. Let P_1, \dots, P_d in $D(k)$ be the ramification points. Assume that, at each P_i , the morphism is totally ramified, and let $Q_i := f(P_i)$, $i = 1, \dots, d$, be the branch points.

When $q \neq p$, the Riemann–Hurwitz formula is

$$2g(D) - 2 = q^s(2g(C) - 2) + d(q^s - 1). \quad (2.1.1)$$

Moreover, $d \geq 2$. When $g(C) = 0$, this follows immediately from the formula; the general case requires a separate proof.

Assume now that $q = p$. For $P \in D(k)$, let $H_0(P) \supseteq H_1(P) \supseteq \dots$ denote the sequence of higher ramification groups. If P is a ramification point, then $|H_0(P)| = |H_1(P)| = p^s$. Set

$$\delta(P) := \sum_i (|H_i(P)| - 1).$$

Then the *Riemann–Hurwitz formula* is

$$2g(D) - 2 = p^s(2g(C) - 2) + \sum_{P \in D(k)} \delta(P), \quad (2.1.2)$$

and it may happen that $d = 1$.

2.2. Let $\gamma(D)$ denote the p -rank of D (i.e., the p -rank of $\text{Jac}(D)$). The *Deuring–Shafarevich formula* relates the p -ranks of C and D :

$$\gamma(D) - 1 = p^s(\gamma(C) - 1) + d(p^s - 1). \quad (2.2.1)$$

The curve D is *ordinary* when $\gamma(D) = g(D)$. When D is ordinary, we find, comparing the formulas (2.1.2) and (2.2.1), that $|H_2(P)| = 1$ for all P and that C is also ordinary. Moreover, when $g(D) > 0$, Equation (2.2.1) shows that $p \leq g(D) + 1$.

When a ramification point P of a Galois morphism $f : D \rightarrow C$ is such that $H_2(P) = (0)$, we will say that the morphism is *weakly ramified* at P .

2.3. We record here the following well-known fact (see [Hasse 1934, p. 42], or [Singh 1974, Lemma 1.3], when $K = k(x)$). Let K be a field with $\text{char}(K) = p$. Let (A, \mathcal{M}) be a discrete valuation ring with field of fractions K , valuation v_K and uniformizer π_K . Assume that the residue field k of A is algebraically closed. Let L/K be a cyclic ramified Galois extension of degree p with Galois group H . Let (B, \mathcal{N}) denote the integral closure of A in L . Let $H = H_0 \supseteq H_1 \supseteq \dots$ denote the sequence of ramification groups. Then $\sum_{i=0}^{\infty} (|H_i| - 1) = (m + 1)(p - 1)$ for some integer m prime to p .

2.4. Examples of curves with an automorphism of degree p in characteristic p can be given in Artin–Schreier form. Consider the curve $y^p - y = \prod_{i=1}^d (x - a_i)^{-n_i}$, where $a_1, \dots, a_s \in k$ are distinct and the n_i are positive integers coprime to p . The automorphism $y \mapsto y + 1$ has order p . The genus g of the smooth complete curve defined by the above equation is given by the Riemann–Hurwitz formula $2g - 2 = -2p + (p - 1)(\sum_{i=1}^d (n_i + 1))$ (see [Subrao 1975, p. 8]).

The following simple proposition exhibits a key difference between the tame and wild cases:

Proposition 2.5. *Let q be a prime. Let $f : D \rightarrow C$ be a ramified cyclic morphism of degree q^s between smooth connected projective curves over k . Let P_1, \dots, P_d , $d \geq 2$, denote the ramification points, assumed to be totally ramified. For $i \neq j$, the image ω_{ij} of $P_i - P_j$ in $\text{Jac}(D)/f^*(\text{Jac}(C))$ is of finite order q^s . Let T denote the finite subgroup $\text{Jac}(D)/f^*(\text{Jac}(C))$ generated by $\{\omega_{id} \mid i = 1, \dots, d - 1\}$.*

- (a) *If $q = p$, then T is isomorphic to $(\mathbb{Z}/p^s\mathbb{Z})^{d-1}$ and is generated by the set $\{\omega_{id} \mid i = 1, \dots, d - 1\}$.*
- (b) *If $q \neq p$, then T is isomorphic to $(\mathbb{Z}/q^s\mathbb{Z})^{d-2}$ and is generated by the set $\{\omega_{id} \mid i = 1, \dots, d - 2\}$.*

Proof. Let S denote the subgroup of $\text{Div}^0(D)$ with support on the set $\{P_1, \dots, P_d\}$. It is clear that $\{P_i - P_d \mid i = 1, \dots, d - 1\}$ is a \mathbb{Z} -basis for S . Let $S \rightarrow T$ denote the natural surjective map. This map factors through $S/q^s S$ since $q^s(\sum_i b_i P_i) = f^*(\sum_i b_i Q_i)$ with $\sum_i b_i Q_i \in \text{Div}^0(C)$.

Let σ be a generator of $\text{Aut}(D/C)$. Suppose that $\sigma(\text{div}_D(g)) = \text{div}_D(g)$ for some $g \in k(D)^*$. Then $g^\sigma = cg$ for some $c \in k^*$. Since σ has finite order q^s , we find that $c^{q^s} = 1$.

Consider first the case where $q = p$. Then $c = 1$. Thus, $g^\sigma = g$ and $g \in k(C)^*$. Suppose that the divisor $(\sum_i b_i P_i)$ has trivial image in T . Then it is possible to write $(\sum_i b_i P_i) = f^*(\sum_j R_j) + \text{div}_D(h)$ for some $R_j \in C(k)$ and $h \in k(D)^*$. Then we have $\sigma(\text{div}_D(h)) = \text{div}_D(h)$, and we conclude that $h \in k(C)^*$. Therefore, we have an equality of divisors of the form $(\sum_i b_i P_i) = f^*(E)$ for some $E \in \text{Div}^0(C)$. It follows that $E = \sum_i c_i Q_i$ for some c_i . Hence, the map $S/p^s S \rightarrow T$ is an isomorphism, proving Part (a).

Suppose now that $q \neq p$. Fix a primitive q^s -th root ξ of 1. Then $k(D)/k(C)$ is a Kummer extension, generated by the root α of $y^{q^s} - a \in k(C)[y]$ such that $\alpha^\sigma = \xi\alpha$. It is easy to check that, for each $i = 0, \dots, q^s - 1$,

$$\{\beta \in k(D) \mid \beta^\sigma = \xi^i \beta\} = k(C)\alpha^i.$$

The equality $\alpha^\sigma = \xi\alpha$ implies that $\text{div}_D(\alpha)$ can be written as

$$\left(\sum_{i=1}^d a_i P_i\right) + \sum_j c_j \left(\sum_{i=0}^{q^s-1} \sigma^i(S_j)\right)$$

for some integers a_i and c_j , and some $S_j \in D(k) \setminus \{P_1, \dots, P_d\}$. It follows that q^s divides $\sum_{i=1}^d a_i$ since $\deg(\operatorname{div}_D(\alpha)) = 0$. This means that the divisor

$$\sum_j c_j \left(\sum_{i=0}^{q^s-1} \sigma^i(S_j) \right) + \left(\sum_{i=1}^d a_i \right) P_d$$

defines an element in $f^*(\operatorname{Jac}(C))$. Hence, the image v of $(\sum_i a_i P_i) - (\sum_i a_i) P_d$ in T is trivial. We thus have a map

$$\varphi : S / \left\langle q^s S, \left(\sum_i a_i P_i \right) - \left(\sum_i a_i \right) P_d \right\rangle \rightarrow T.$$

Let us note that $(\sum_i a_i P_i) - (\sum_i a_i) P_d \notin q^s S$ because, otherwise, the morphism f given by the Kummer equation $y^{q^s} - a$ would not be totally ramified at P_1, \dots, P_d .

Suppose that the divisor $(\sum_i b_i P_i)$ has trivial image in T . Then it is possible to write $(\sum_i b_i P_i) = f^*(\sum_j R_j) + \operatorname{div}_D(h)$ for some $R_j \in C(k)$ and $h \in k(D)^*$. Thus, we have $\sigma(\operatorname{div}_D(h)) = \operatorname{div}_D(h)$, and we conclude that $h^\sigma = \xi^j h$ for some $j \in \{0, \dots, q^s - 1\}$. Therefore, there exists $b \in k(C)^*$ such that $h = b\alpha^j$. Hence, we have an equality of divisors of the form $(\sum_i b_i P_i) = f^*(E) + j[(\sum_i a_i P_i) - (\sum_i a_i) P_d]$ for some $E \in \operatorname{Div}^0(C)$. It follows that $E = \sum_i c_i Q_i$ for some c_i . Hence, the map φ is an isomorphism, proving Part (b). \square

Corollary 2.6. *Assume that $p \neq 2$. Let D/k be a smooth projective connected hyperelliptic curve of genus g . Denote by τ the hyperelliptic involution. Let σ be an automorphism of order p . Then either σ has a single fixed point, fixed by τ , or it has exactly two fixed points, permuted by τ .*

Proof. The hyperelliptic involution commutes with σ , and hence, it permutes the fixed points $\{P_1, \dots, P_d\}$. If $d \geq 2$ and two fixed points P_1 and P_2 of σ are fixed by τ , then the divisor class $P_1 - P_2$ is fixed by τ . [Proposition 2.5](#) shows that the class of $P_1 - P_2$ is not trivial and, since $p > 2$, this divisor class is not equal to the class of $-(P_1 - P_2)$. This is a contradiction since τ acts as the $[-1]$ -map on $\operatorname{Jac}(D)$. Thus, τ fixes at most one point P_i .

If $d \geq 3$, then we may assume that either $\tau(P_1) = P_2$ and P_3 is fixed or that $\tau(P_1) = P_2$ and $\tau(P_3) = P_4$. In the first case, we find that $\tau(P_1 - P_3) = (P_2 - P_3) = -(P_1 - P_2) + (P_1 - P_3)$. Using the fact that τ acts as the $[-1]$ -map on $\operatorname{Jac}(D)$, we find the relation $-(P_1 - P_3) = -(P_1 - P_2) + (P_1 - P_3)$ in $\operatorname{Jac}(D)$. Looking at this relation in T contradicts [Proposition 2.5](#). The other case is similar and is left to the reader. \square

Example 2.7. Assume that $p \neq 2$. Consider a smooth hyperelliptic curve C/k given by an affine equation $y^2 = f(x)$, and let D be its Galois cover given by the equation $z^p - z = x$. The automorphism $\sigma : D \rightarrow D$ with $\sigma(z) = z + 1$ has one fixed point P with $\delta(P) = 3(p - 1)$ when $\deg(f)$ is odd, and it has two fixed points P_1 and P_2 with $\delta(P_1) = \delta(P_2) = 2(p - 1)$ when $\deg(f)$ is even.

3. Arithmetical trees

Our main result in this section is [Proposition 3.5](#), which will be needed in the proof of [Theorem 5.3](#). This proposition pertains to arithmetical graphs, and we now recall how one associates such an object to any regular model of a curve.

Let X/K be any smooth, proper, geometrically connected curve of genus g . Let $\mathcal{X}/\mathcal{O}_K$ be a regular model of X/K . Let $\mathcal{X}_k := \sum_{i=1}^v r_i C_i$ denote the special fiber of \mathcal{X} , where C_i is an irreducible component and r_i is its multiplicity. Let $M := ((C_i \cdot C_j))_{1 \leq i, j \leq v}$ be the associated *intersection matrix*. Denote by G the associated graph. Let ${}^tR := (r_1, \dots, r_v)$ so that $MR = 0$. We call the triple (G, M, R) an *arithmetical graph* (in [\[Lorenzini 1989\]](#), the additional condition that $\gcd(r_1, \dots, r_v) = 1$ is assumed, and it is $(G, -M, R)$ that is called an arithmetical graph). For the purpose of simplifying the statements of some definitions, we sometimes think of G as a metric space with the natural topology where each edge of G with its two endpoints is homeomorphic to the closed unit interval $[0, 1]$.

Let (G, M, R) be any arithmetical graph on v vertices. Let $M : \mathbb{Z}^v \rightarrow \mathbb{Z}^v$ and ${}^tR : \mathbb{Z}^v \rightarrow \mathbb{Z}$ be the linear maps associated to the matrices M and R . The *group of components* of (G, M, R) is defined as

$$\Phi_M := \text{Ker}({}^tR) / \text{Im}(M) = (\mathbb{Z}^v / \text{Im}(M))_{\text{tors}}.$$

Motivated by the case of degenerations of curves, we shall denote by $(C, r(C))$ a vertex of G , where $r(C)$ is the coefficient of R corresponding to C . The integer $r(C)$, also denoted simply by r , is called the multiplicity of C . The matrix M is written as $M := ((C_i \cdot C_j))_{1 \leq i, j \leq v}$, and we write $|C_i \cdot C_i| := |(C_i \cdot C_i)|$.

3.1. Denote by $\langle \cdot, \cdot \rangle : \Phi_M \times \Phi_M \rightarrow \mathbb{Q}/\mathbb{Z}$ the perfect pairing $\langle \cdot, \cdot \rangle_M$ attached in [\[Bosch and Lorenzini 2002, Lemma 1.1\]](#) to the symmetric matrix M . Explicit values of this pairing are computed as follows. Let (C, r) and (C', r') be two distinct vertices of G . Define

$$E(C, C') := \left(0, \dots, 0, \frac{r'}{\gcd(r, r')}, 0, \dots, 0, \frac{-r}{\gcd(r, r')}, 0, \dots, 0 \right) \in \mathbb{Z}^v,$$

where the first nonzero coefficient of $E(C, C')$ is in the column corresponding to the vertex C and, similarly, the second nonzero coefficient is in the column corresponding to the vertex C' . We say that the pair (C, C') is *uniquely connected* if there exists a path \mathcal{P} in G between C and C' such that, for each edge e on \mathcal{P} , the graph $G \setminus \{e\}$ is disconnected. Note that, when a pair (C, C') is uniquely connected, then the path \mathcal{P} is the unique shortest path between C and C' . A graph is a tree if and only if every pair of vertices of G is uniquely connected.

Let (C, r) and (C', r') be a uniquely connected pair with associated path \mathcal{P} . While walking on $\mathcal{P} \setminus \{C, C'\}$ from C to C' , label each encountered vertex consecutively

by $(C_1, r_1), (C_2, r_2), \dots, (C_n, r_n)$. Let G_i denote the connected component of C_i in $G \setminus \{\text{edges of } \mathcal{P}\}$. The graph G_i is reduced to a single vertex if and only if C_i is not a node of G . For convenience, we write $(C, r) = (C_0, r_0)$ and $(C', r') = (C_{n+1}, r_{n+1})$ and define G_0 and G_{n+1} accordingly.

3.2. The following facts are proved in [Bosch and Lorenzini 2002, Proposition 5.1]. Let (G, M, R) be any arithmetical graph. Let C and C' be two vertices such that (C, C') is a *uniquely connected* pair of G . Let γ denote the image of $E(C, C')$ in Φ_M . For (D, s) and (D', s') any two distinct vertices on G , let δ denote the image of $E(D, D')$ in Φ_M . Writing \mathcal{P} for the oriented shortest path from C to C' as above, let C_α denote the vertex of \mathcal{P} closest to D in G , and let C_β denote the vertex of \mathcal{P} closest to D' . In other words, $D \in G_\alpha$ and $D' \in G_\beta$. Assume that $\alpha \leq \beta$. (Note that we may have $\alpha = \beta$, and we may have $D = C_\alpha$ or $D' = C_\beta$.) Then if $\alpha < \beta$,

$$\langle \gamma, \delta \rangle = -\text{lcm}(r, r') \text{lcm}(s, s') \left(\frac{1}{r_\alpha r_{\alpha+1}} + \frac{1}{r_{\alpha+1} r_{\alpha+2}} + \dots + \frac{1}{r_{\beta-1} r_\beta} \right) \pmod{\mathbb{Z}}, \quad (3.2.1)$$

and if $C_\alpha = C_\beta$, then $\langle \gamma, \delta \rangle = 0$. Moreover,

$$\langle \gamma, \gamma \rangle = -\text{lcm}(r, r')^2 \left(\frac{1}{r r_1} + \frac{1}{r_1 r_2} + \dots + \frac{1}{r_n r'} \right) \pmod{\mathbb{Z}}. \quad (3.2.2)$$

Note that the negative signs in the expressions (3.2.1) and (3.2.2) are missing in [Bosch and Lorenzini 2002, Proposition 5.1]. Thus, all expressions for $\langle \gamma, \delta \rangle$ computed in Section 5 of [Bosch and Lorenzini 2002] using Proposition 5.1 are correct only after having been multiplied by -1 . Similar sign mistakes occurred in [Lorenzini 2000]. The proof of [Bosch and Lorenzini 2002, Proposition 5.1] is correct except that its last line produces the opposite of the stated values for $\langle \gamma, \delta \rangle$ since we assume $\alpha \leq \beta$.

3.3. Let (C, r) be a vertex of G of degree $d \geq 2$. Let $(D_i, r_i), i = 1, \dots, d$, denote the neighbors of C , that is, the vertices of G linked to C . Let τ_i denote the image of $E(D_i, D_d)$ in Φ_M for $i \in \{1, \dots, d-1\}$. We will use repeatedly the following expressions computed using (3.2.1) and (3.2.2):

$$\langle \tau_i, \tau_i \rangle = -\text{lcm}(r_i, r_d)^2 \frac{r_i + r_d}{r_i r_d r} \pmod{\mathbb{Z}}$$

and, when $i \neq j$,

$$\langle \tau_i, \tau_j \rangle = -\text{lcm}(r_i, r_d) \text{lcm}(r_j, r_d) \frac{1}{r_d r} \pmod{\mathbb{Z}}.$$

These formulas allow us to easily show that τ_i may not always be trivial. For example, let p be a prime dividing r . When $p \nmid r_i r_d (r_i + r_d)$, we find that $\langle \tau_i, \tau_i \rangle \neq 0$

and, thus, $\tau_i \neq 0$. Similarly, when for three distinct indices i, j and d we have $p \nmid r_i r_j r_d$, we find that $\langle \tau_i, \tau_j \rangle \neq 0$, showing that both τ_i and τ_j are not trivial.

We claim that r kills τ_i . Indeed, we find, using [Lorenzini 2000, Lemma 2.2], that the images in Φ_M of $E(D_i, C)$ and $E(C, D_d)$ have order dividing $\gcd(r_i, r)$ and $\gcd(r, r_d)$, respectively. Consider the following easy relation between vectors in \mathbb{Z}^v [Lorenzini 2000, Remark 3.5]: given any three vertices (A, a) , (B, b) and (C, c) ,

$$bE(A, C) = \frac{c}{\gcd(a, c)} \gcd(a, b)E(A, B) + \frac{a}{\gcd(a, c)} \gcd(b, c)E(B, C). \quad (3.3.1)$$

Using this relation, we find that $r\tau_i = 0$.

Lemma 3.4. *Let (G, M, R) be an arithmetical graph. Consider any two distinct vertices (A, a) and (A', a') , and let $\alpha_{AA'}$ denote the image of $E(A, A')$ in Φ_M . Then the set $\{\alpha_{AA'} \mid A \neq A'\}$ is a set of generators for Φ_M .*

Proof. Let us note first that the statement is proved for (G, M, R) as soon as it is proved for $(G, M, R/\gcd(r_1, \dots, r_v))$. We will thus assume now that $\gcd(r_1, \dots, r_v) = 1$. Fix a vertex A , and consider the subgroup $(\Phi_M)_A$ of Φ_M generated by $\{\alpha_{AA'} \mid \text{all } A' \neq A\}$. We claim that $a\Phi_M \subseteq (\Phi_M)_A$. Indeed, an element $\phi \in \Phi_M$ is represented by the class of a vector $(f_D \mid D \in G)$ such that $\sum f_D r(D) = 0$. It follows that $a\phi = -\sum \gcd(a, r(D)) f_D \alpha_{AD}$. Since $\gcd(r_1, \dots, r_v) = 1$, ϕ can be expressed in terms of elements of the form $\alpha_{AA'}$. \square

The following is a key relation between the τ_i 's:

Proposition 3.5. *Let (G, M, R) be an arithmetical tree. Let (C, r) be a vertex of degree $d \geq 2$. Keep the notation introduced in 3.3. Then $\sum_{i=1}^{d-1} \gcd(r_i, r_d) \tau_i = 0$.*

Proof. Consider any two distinct vertices (A, a) and (A', a') , and let α denote the image of $E(A, A')$ in Φ_M . The previous lemma shows that the group Φ_M is generated by such elements α .

Let $\tau := \sum_{i=1}^{d-1} \gcd(r_i, r_d) \tau_i$. We claim that $\langle \tau, \alpha \rangle = 0$ for all such elements α . This claim, proved below, implies immediately that $\tau = 0$. Indeed, recall that, $\langle \cdot, \cdot \rangle$ being perfect, the element τ is trivial if and only if $\langle \tau, \phi \rangle = 0$ for all $\phi \in \Phi_M$.

Let us now prove our claim. Assume first that the path \mathcal{Q} between A and A' contains the vertices D_i and D_d with $i \neq d$. We use (3.2.1) to compute modulo \mathbb{Z} that

$$\begin{aligned} \langle \tau, \alpha \rangle &= \pm \operatorname{lcm}(a, a') \\ &\times \left(\gcd(r_i, r_d) \operatorname{lcm}(r_i, r_d) \left(\frac{1}{r_i r} + \frac{1}{r r_d} \right) + \sum_{j \neq i, d} \gcd(r_j, r_d) \operatorname{lcm}(r_j, r_d) \left(\frac{1}{r r_d} \right) \right), \end{aligned}$$

which simplifies to

$$\langle \tau, \alpha \rangle = \pm \operatorname{lcm}(a, a') \left(\sum_{j=1}^d r_j \right) \frac{1}{r}.$$

Since $\sum_{j=1}^d r_j = |C \cdot C| r$, we find that $\langle \tau, \alpha \rangle = 0$. When \mathcal{Q} contains D_i and D_j with $i, j \neq d$ and $i \neq j$, we find that modulo \mathbb{Z}

$$\begin{aligned} \langle \tau, \alpha \rangle &= \pm \operatorname{lcm}(a, a') \left(\operatorname{gcd}(r_i, r_d) \operatorname{lcm}(r_i, r_d) \frac{1}{r_i r} - \operatorname{gcd}(r_j, r_d) \operatorname{lcm}(r_j, r_d) \frac{1}{r_j r} \right) \\ &= \pm \operatorname{lcm}(a, a') \left(\frac{r_d}{r} - \frac{r_d}{r} \right) = 0. \end{aligned}$$

It is clear that if the path \mathcal{Q} contains no vertices D_i , or if it contains exactly one vertex D_i and does not contain the vertex C , then $\langle \tau, \alpha \rangle = 0$. It remains to consider the case where the path \mathcal{Q} contains exactly one vertex D_i and the vertex C . Then C is an endpoint of \mathcal{Q} , and thus, r divides $\operatorname{lcm}(a, a')$. When $i \neq d$, we find that

$$\langle \tau, \alpha \rangle = \pm \operatorname{lcm}(a, a') \operatorname{lcm}(r_i, r_d) \operatorname{gcd}(r_i, r_d) \frac{1}{r_i r}$$

is 0 modulo \mathbb{Z} , and when $i = d$, we find that

$$\langle \tau, \alpha \rangle = \pm \operatorname{lcm}(a, a') \left(\sum_{i=1}^{d-1} \operatorname{lcm}(r_i, r_d) \operatorname{gcd}(r_i, r_d) \frac{1}{r_i r} \right)$$

is also 0 modulo \mathbb{Z} . □

4. Some combinatorics

Let (G, M, R) be an arithmetical graph. We introduce below a measure $\gamma_D g_D$ of how “complicated” certain subgraphs G_D of G are, and we describe G_D in [Proposition 4.3](#) when $\gamma_D g_D$ is as small as possible. This result is needed in the proof of [Theorem 6.8](#). A geometric motivation for the introduction of the quantity $\gamma_D g_D$ is found in the genus formula [\(6.1.1\)](#).

4.1. Let (G, M, R) be an arithmetical graph. Fix a vertex $(C_0, r(C_0))$ of G . Assume that C_0 is linked to a vertex $(D, r(D))$ by a single edge e and that, when the edge e is removed from G , then D and C_0 are not in the same connected component of the resulting graph. Let G_D denote the connected component of $G \setminus \{e\}$ that contains D . Consider the minor N_D of M corresponding to the vertices in G_D . Let

$$\gamma_D := \operatorname{gcd}(r(A) \mid A \text{ a vertex of } G_D).$$

Then γ_D divides $r(C_0)$. Indeed, γ_D divides the multiplicity of D and of all vertices linked to D except possibly that of C_0 . But the relation $MR = 0$ implies then that γ_D divides the multiplicity of C_0 . Let R_D denote the vector R restricted to the vertices of G_D . By definition, we find that R_D / γ_D is an integer vector.

Let $\beta(G)$ denote the first Betti number of the graph G . Letting $d_G(A)$ denote the degree of a vertex A in the graph G , we have

$$2\beta(G) - 2 = \sum_{\text{vertices } A \text{ of } G} (d_G(A) - 2).$$

Associated with any arithmetical graph (G, M, R) is the following integer invariant $g_0(G) \geq \beta(G)$ [Lorenzini 1989, Theorem 4.10], defined by the formula

$$2g_0(G) = 2\beta(G) + \sum_{\text{vertices } A \text{ of } G} (r(A) - 1)(d_G(A) - 2). \tag{4.1.1}$$

Let C_0 and D be as above. We now associate to the pair (N_D, R_D) an integer g_D , defined so that the formula below holds:

$$\gamma_D g_D = r(C_0) + r(D) + \sum_{\text{vertices } A \text{ of } G_D} r(A)(d_{G_D}(A) - 2).$$

Since γ_D divides $r(C_0)$, the invariant g_D is indeed an integer. We can rewrite this formula as

$$\begin{aligned} \gamma_D g_D = 2\beta(G_D) + (r(C_0) - 1) + (r(D) - 1) \\ + \sum_{\text{vertices } A \text{ of } G_D} (r(A) - 1)(d_{G_D}(A) - 2), \end{aligned} \tag{4.1.2}$$

and we find that

$$\begin{aligned} g_D = 2\beta(G_D) + \left(\frac{r(C_0)}{\gamma_D} - 1\right) + \left(\frac{r(D)}{\gamma_D} - 1\right) \\ + \sum_{\text{vertices } A \text{ of } G_D} \left(\frac{r(A)}{\gamma_D} - 1\right)(d_{G_D}(A) - 2). \end{aligned} \tag{4.1.3}$$

4.2. We will make use below of the following facts. Suppose that, on G , the vertices D_0, D_1, \dots, D_n are consecutive vertices on a terminal chain and D_n is the terminal vertex on this chain (in other words, D_i is linked by one edge to D_{i+1} for $i = 0, \dots, n - 1$, $d_G(D_i) = 2$ for $i = 1, \dots, n - 1$ and $d_G(D_n) = 1$). Then $\gcd(r(D_0), r(D_1)) = r(D_n)$, and if $|D_i \cdot D_i| > 1$ for all $i = 1, \dots, n$, then


$$r(D_0) > r(D_1) > \dots > r(D_n).$$

Indeed, the equality $|D_n \cdot D_n| r(D_n) = r(D_{n-1})$ obtained from the relation $MR = 0$ shows that $r(D_n)$ divides $r(D_{n-1})$ and $r(D_n) < r(D_{n-1})$ if $|D_n \cdot D_n| > 1$. Suppose that, for some i , we have $r(D_i) > r(D_{i+1})$. Then $r(D_{i-1}) > r(D_i)$ because $|D_i \cdot D_i| r(D_i) = r(D_{i-1}) + r(D_{i+1})$ and $|D_i \cdot D_i| \geq 2$. The equality $|D_i \cdot D_i| r(D_i) = r(D_{i-1}) + r(D_{i+1})$ implies that $\gcd(r(D_{i-1}), r(D_i)) = \gcd(r(D_i), r(D_{i+1}))$.

Proposition 4.3. *Let (G, M, R) be an arithmetical tree containing a vertex C_0 of prime multiplicity p . Assume that a vertex D linked to C_0 by an edge e has multiplicity divisible by p . Let G_D denote the connected component of $G \setminus \{e\}$ that contains D . Assume in addition that G_D does not contain any vertex A of degree 1 or 2 in G with $|A \cdot A| = 1$. Then*

$$\gamma_D g_D \geq 2(p - 1).$$

If $\gamma_D g_D = 2(p - 1)$, then $\gamma_D = 1$ and G_D is a graph of the shape depicted below, containing one node C of G only, of multiplicity p and degree 3 in G . The two terminal vertices of G that belong to G_D have multiplicity 1.



Let α denote the number of vertices of G_D on the chain linking C_0 to the node C of G_D (including the node C). Let C_1 and C'_1 denote the vertices linked to C on the two terminal chains. Then $1 \leq r(C_1) < p$, and the minor of M corresponding to the vertices of G_D is completely determined by p , α and $r(C_1)$.

The proof of Proposition 4.3 is given in 4.6. We start with a preliminary lemma.

4.4. Let (G, M, R) be an arithmetical tree. For each node $(C, r(C))$ of degree $d(C) \geq 3$ in G , we define an invariant $\mu(C)$ as follows. Let $\rho(C)$ denote the number of terminal chains attached to C , and let $D_1(C), \dots, D_{\rho(C)}(C)$ be the vertices of G linked to C that belong each to one terminal chain attached to C . Let $r_i(C)$ denote the multiplicity of $D_i(C)$. The multiplicity of the terminal vertex on the chain containing $D_i(C)$ is $\gcd(r(C), r_i(C))$. If no vertex A on the terminal chain has $|A \cdot A| = 1$, then $r_i(C) < r(C)$ (see 4.2). When a chain attached to C is not terminal, we will call it a *connecting chain*. As in [Lorenzini 1989, Theorem 4.7], we let, when $\rho(C) > 0$,

$$\mu(C) := (d(C) - 2)(r(C) - 1) - \sum_{j=1}^{\rho(C)} (\gcd(r(C), r_j(C)) - 1).$$

When $\rho(C) = 0$, we let $\mu(C) := (d(C) - 2)(r(C) - 1)$. It is clear that, if $r(C) = 1$, then $\mu(C) = 0$.

Lemma 4.5. *Assume that the terminal chains attached to C do not contain a vertex A with $|A \cdot A| = 1$. Then $\mu(C) \geq 0$, and $\mu(C) = 0$ if and only if $r(C) = 1$ and $\rho(C) = 0$.*

Proof. It is clear that, if a node C has $\rho(C) = 0$, then $\mu(C) \geq 0$, and $\mu(C) = 0$ only when $r(C) = 1$. Assume now that $\rho(C) > 0$. Our hypothesis implies that $r(C) > \gcd(r(C), r_i(C))$ for each vertex $D_i(C)$, $i = 1, \dots, \rho(C)$. In particular, $r(C) > 1$, and we need to prove that $\mu(C) > 0$. Let

$$s := \gcd(r(C), r_1(C), \dots, r_d(C)).$$

Assume first that $\rho(C) = d(C)$ so that G has a single node. It is proven in [Lorenzini 1989, Proposition 4.1] that, if $\rho(C) = d(C)$ and $s = 1$, then $\mu(C) \geq 0$. When $s > 1$, define

$$\mu_s(C) := (d(C) - 2) \left(\frac{r(C)}{s} - 1 \right) - \sum_{j=1}^{\rho(C)} \left(\frac{\gcd(r(C), r_j(C))}{s} - 1 \right).$$

The integer $\mu_s(C)$ is nothing but the μ -invariant of the node on the arithmetical graph obtained from G by dividing all its multiplicities by s . Thus, $\mu_s(C)$ is even [Lorenzini 1989, Definition 3.6] and $\mu_s(C) \geq 0$. Since

$$\mu(C) = -2(s - 1) + s\mu_s(C),$$

we find that $\mu(C) > 0$ if $\mu_s(C) > 0$. We claim that, under our hypotheses, $\mu(C) > 0$ when $s = 1$. Indeed, our hypotheses imply that $r(C) > \gcd(r(C), r_i(C))$ for each vertex $D_i(C)$, $i = 1, \dots, \rho(C)$. Dropping the reference to C , we can write

$$\begin{aligned} \mu(C) &:= (d - 2)(r - 1) - \sum_{j=1}^d (\gcd(r, r_j) - 1) \\ &\geq (d - 2)(r - 1) - d(r/2 - 1) = (d - 4)r/2 + 2. \end{aligned}$$

Thus, $\mu(C) > 0$ if $d \geq 4$. Assume now that $d = 3$. Then $cr = r_1 + r_2 + r_3$ for some c . Let $h_i = \gcd(r, r_i)$, and assume that $h_1 \geq h_2 \geq h_3$. Then $(h_1, h_2, h_3) = (r/2, r/2, r/2), (r/2, r/2, r/3), (r/2, r/3, r/3), (r/2, r/3, r/4)$ cannot occur due to the divisibility $r \mid (r_1 + r_2 + r_3)$. Since the cases $(h_1, h_2, h_3) = (r/3, r/3, r/3), (r/2, r/4, r/4), (r/2, r/3, r/6)$ have $\mu(C) > 0$, we need only consider $(h_1, h_2, h_3) = (r/2, r/3, r/5)$. In this case, $r_1 = r/2, r_2 = r/3$ or $2r/3$ and $r_3 = ar/5$ with $a = 1, \dots, 4$. The reader will check that $cr = r_1 + r_2 + r_3$ is impossible to achieve with these values, and our claim is proved.

Let us assume now that $0 < \rho(C) < d(C)$. Then

$$\begin{aligned} \mu(C) &:= (d - 2)(r - 1) - \sum_{j=1}^{\rho} (\gcd(r, r_j) - 1) \\ &\geq (d - 2)(r - 1) - (d - 1)(r/2 - 1) = (d - 3)r/2 + 1 > 0. \quad \square \end{aligned}$$

4.6. Proof of Proposition 4.3. We claim that G_D contains a node of G . (This node is also a node of G_D unless it is D itself and $d_G(D) = 3$.) Indeed, the hypotheses that $r(C_0) \leq r(D)$ and $|D \cdot D| > 1$ imply that $d_G(D) > 1$ because the relation $MR = 0$

provides otherwise for the equality $|D \cdot D|r(D) = r(C_0)$, which is a contradiction. Suppose then that D is connected in G_D to D_1 . If $d_G(D) = 2$, then we find from the relation $|D \cdot D|r(D) = r(C_0) + r(D_1)$ that $r(D) \leq r(D_1)$. Repeating this discussion with D and D_1 instead of C_0 and D , we find that the graph G_D has a chain of increasing multiplicities $r(D) \leq r(D_1) \leq \dots$, which eventually leads to a node of G_D (and of G).

In G , C_0 and D are adjacent vertices. Consider the connected component \mathcal{G} of $G \setminus \{D\}$ that contains C_0 . Two cases can occur: either (a) \mathcal{G} contains a node of G , or (b) \mathcal{G} does not contain a node of G , in which case we will call \mathcal{G} a terminal chain of G . In the latter case, the terminal vertex on this chain has multiplicity $\gcd(r(C_0), r(D))$ (see 4.2), which equals $r(C_0)$ by hypothesis. The definition of $\gamma_D g_D$ in (4.1.2), along with the fact that we assume that G is a tree, allow us to write

$$\gamma_D g_D = (r(C_0) - 1) + \sum_{\text{vertices } A \text{ of } G_D} (r(A) - 1)(d_G(A) - 2).$$

In case (a), C_0 is not on a terminal chain of G so that, by definition of $\mu(C)$ in 4.4, we can write

$$\gamma_D g_D = (r(C_0) - 1) + \sum_{\text{nodes } C \text{ of } G \text{ in } G_D} \mu(C) \tag{4.6.1}$$

(where $\mu(C)$ is computed viewing C as a node of G and not of G_D). In case (b) where C_0 is on a terminal chain of G whose terminal vertex has multiplicity $r(C_0)$, we have

$$\gamma_D g_D = 2(r(C_0) - 1) + \sum_{\text{nodes } C \text{ of } G \text{ in } G_D} \mu(C).$$

We prove below case (a). The arguments to prove (b) are similar and are left to the reader. Case (b) will not be used in the remainder of this article.

Assume that we are in case (a). We can apply Lemma 4.5, and we obtain that each term $\mu(C)$ in the above sum is nonnegative. In view of (4.6.1), since $r(C_0) = p$ by hypothesis, we need to show that $\sum_{\text{nodes } C} \mu(C) \geq p - 1$, and we need to describe the graphs for which $\sum_{\text{nodes } C} \mu(C) = (p - 1)$.

Denote by C the node of G closest to C_0 in G_D . (This node could be D .) The multiplicity of C is divisible by p since p divides the consecutive multiplicities $r(C_0)$ and $r(D)$ (similar argument as in 4.2). Let np denote the multiplicity of C .

Suppose that C (of degree d in G) has only one connecting chain. If $n = 1$, then all terminal multiplicities at C equal 1 and $\mu(C) = (d - 2)(p - 1)$. The case $d = 3$ leads to the case described in the statement of Proposition 4.3 with $\mu(C) = (p - 1)$, $\gamma_D g_D = 2(p - 1)$ and $\gamma_D = 1$. When $d > 3$, we have $\mu(C) > p - 1$, as desired.

When $n > 1$, the inequality

$$\begin{aligned}\mu(C) &\geq (d-2)(np-1) - (d-1)(np/2-1) \\ &= (d-2)np/2 - np/2 + 1\end{aligned}$$

shows that we have $\mu(C) > p-1$ unless $d=3$. When $n > 1$ and $d=3$, every vertex on the chain linking C to C_0 has multiplicity divisible by p . Thus, either (i) both terminal multiplicities of C are coprime to p (call them n_1 and n_2), or (ii) both are divisible by p (call them m_1p and m_2p).

In case (i), $\mu(C) = np - n_1 - n_2 + 1$ with n_1 and n_2 dividing n . It follows that $\mu(C) \geq n(p-2) + 1$. Clearly, $\mu(C) > p-1$ unless $p=2$. Assume that $p=2$. If $(n_1, n_2) \neq (n, n)$, we find that $\mu(C) = n(p-1) + 1 > (p-1)$. The case $(n_1, n_2) = (n, n)$ cannot happen because, in that case, n divides the multiplicity of all the components linked to C , which implies then that $n=2$. But a node of multiplicity 4 cannot have exactly three vertices of multiplicity 2 attached to it.

In case (ii), $\mu(C) = (n - m_1 - m_2)p + 1$ with m_1 and m_2 dividing n . The equality $(n - m_1 - m_2) = 0$ is not possible. Indeed, it is only possible if $m_1 = m_2 = n/2$. But since $\gcd(m_1, m_2) = 1$, this case can happen only if $n=2$. But then $|C \cdot C|$ would equal $3/2$, a contradiction. It follows that $\mu(C) = (n - m_1 - m_2)p + 1 > p-1$.

Suppose now that C , of multiplicity np , has at least two connecting chains. If $n > 1$, then

$$\mu(C) \geq (d-2)(np-1) - (d-2)(np/2-1) = (d-2)np/2 > p-1,$$

as desired. If $n=1$, then $\mu(C) = (d-2)(p-1)$. Thus, $\mu(C) > p-1$ if $d > 3$. Suppose now that $d=3$. Since G_D is a tree with a node C of degree 3, G_D must have at least three terminal vertices. Thus, there must exist at least one additional node C' on the graph G_D that has a terminal chain. It follows that $\mu(C') \geq 1$ (Lemma 4.5) and, therefore, $\mu(C) + \mu(C') > p-1$, as desired.

4.7. To conclude the proof of Proposition 4.3, we now specify the intersection matrix in the case where $\gamma_D g_D = 2(p-1)$. Let (G, M, R) be as in Proposition 4.3, and assume that the vertex D is such that $\gamma_D g_D = 2(p-1)$. Let N_D denote the matrix M restricted to the vertices of G_D . Let α denote the number of vertices of G_D on the chain linking C_0 to the node C of G_D (including the node C). Each of these vertices except C is of degree 2. The multiplicity of C is p . Since we assume that no vertex of degree 2 has self-intersection -1 , we find that the multiplicity of each of these vertices must be p . It follows that each of these vertices except possibly C must have self-intersection -2 .

Let C_1 and C'_1 denote the vertices linked to C on the two terminal chains. Since they have degree 1 or 2 and cannot have self-intersection -1 , we find that $1 \leq r(C_1) < r(C) = p$ and $r(C'_1) < r(C)$. Moreover, from $MR = 0$, we find

that $p + r(C_1) + r(C'_1) = p|C \cdot C|$. It follows that $|C \cdot C| = 2$, and $r(C'_1) = p - r(C_1)$. We claim that N_D depends only on p , α and $r(C_1)$, and we write it as $N_D = N(p, \alpha, r(C_1))$. Indeed, the pair $(p, r(C_1))$ completely determines all multiplicities and all self-intersections on the terminal chain containing C_1 : use $(r, s) = (p, r(C_1))$ in 4.8 below to determine the self-intersections and multiplicities of the terminal chain. Similarly, the pair $(p, r(C'_1))$ completely determines all multiplicities and all self-intersections on the terminal chain containing C'_1 . This concludes the proof of Proposition 4.3. The matrix N_D is an intersection matrix also introduced in [Lorenzini 2013a, Example 3.18]. \square

4.8. Recall the following standard construction. Given an ordered pair of positive integers $r > s$ with $\gcd(r, s) = 1$, we construct an associated intersection matrix $N = N(r, s)$ with vector $R = R(r, s)$ and $NR = -re_1$ as follows (where e_1 denotes the first standard basis vector of \mathbb{Z}^n). Using the division algorithm, we can find positive integers b_1, \dots, b_m and $s_1 = s > s_2 > \dots > s_m = 1$ such that $r = b_1s - s_2$, $s_1 = b_2s_2 - s_3$ and so on until we get $s_{m-1} = b_ms_m$. These equations are best written in matrix form:

$$\begin{pmatrix} -b_1 & 1 & \dots & 0 \\ 1 & -b_2 & \ddots & \\ & \ddots & \ddots & 1 \\ 0 & \dots & 1 & -b_m \end{pmatrix} \begin{pmatrix} s_1 \\ \vdots \\ \vdots \\ s_m \end{pmatrix} = \begin{pmatrix} -r \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$


We let $N(r, s)$ denote the above square matrix and $R(r, s)$ be the column matrix on the left of the “equals” sign. It is well-known that $\det(N(r, s)) = \pm r$ (see [Lorenzini 2000, Lemma 2.6]). We recall also for use in Corollary 6.12 that

$$\frac{1}{rs} + \frac{1}{ss_2} + \dots + \frac{1}{s_{m-1}s_m} = \frac{c}{r}, \quad (4.8.1)$$

where $0 < c < r$ is such that $r \mid cs - 1$ (see [Lorenzini 2000, Lemmas 2.8 and 2.6]).


Remark 4.9. In Proposition 4.3, the hypothesis that $\gamma_D g_D = 2(p - 1)$ allowed us to completely describe the graph G_D . For a fixed $\gamma_D g_D > 2(p - 1)$, the situation is much more complicated and several possible types of graphs G_D may occur. It would follow from our guess in 6.2 that, for applications to models of curves, it suffices to classify the cases where $\gamma_D g_D$ is a multiple of $p - 1$. We give below several possible types of graphs G_D with $\gamma_D g_D = 3(p - 1)$ when p is odd.

- (a) G_D is a graph with one node of G only, of multiplicity p and degree 4 in G . The three terminal vertices of G that belong to G_D have multiplicity 1.




To completely determine the intersection matrix N_D and the vector R_D , one needs to also provide the multiplicities r_1, r_2 and r_3 , of the first vertices on each of the three terminal chains, with the conditions $1 \leq r_1, r_2, r_3 < p$ and $r_1 + r_2 + r_3$ divisible by p . Such data can only be provided when p is odd. The self-intersection of the node is then $-(p + r_1 + r_2 + r_3)/p = -2$ or -3 .

- (b) G_D is a graph with one node of G only, of multiplicity $2p$ and degree 3 in G . The two terminal vertices of G that belong to G_D have multiplicity 1 and 2, respectively.



- (c) G_D is a graph with 2 nodes C and C' of G . Let C be the node closest to C_0 in G_D . It has multiplicity p and degree 3 in G , and it has a single terminal chain with terminal multiplicity 1. The node C' is connected to C by a connecting chain that contains a vertex of multiplicity coprime to p .



We conclude this section with some general remarks concerning the invariant g_D introduced in (4.1.2).

Remark 4.10. Let (G, M, R) be an arithmetical graph. As at the beginning of this section, fix a vertex $(C_0, r(C_0))$ of G . Assume that C_0 is linked to a vertex $(D, r(D))$ by a single edge e and that, when the edge e is removed from G , then D and C_0 are not in the same connected component of the resulting graph. Let G_D denote the connected component of $G \setminus \{e\}$ that contains D . Consider the minor $N = N_D$ of M corresponding to the vertices in G_D . Let n denote the number of vertices of G_D .

(a) *The integer g_D depends only on the matrix N_D and the vertex D on the graph G_D .* To prove this statement, we show that the vector R_D/γ_D is completely determined by N_D and the vertex D . Indeed, let us number the vertices of G_D such that D is the first vertex numbered. Then R_D/γ_D is a vector with positive coefficients such that $N_D(R_D/\gamma_D) = {}^t(-r(C_0)/\gamma_D, 0, \dots, 0)$ (where the superscript t indicates the transpose vector). The existence of such a relation insures that N_D is negative-definite (see [Lorenzini 2013a, §3.3]), and the vector R_D/γ_D is a rational multiple of the first column of the unique matrix N^* such that $NN^* = N^*N = \det(N) \text{Id}_n$ [Lorenzini 2013a, Definition 3.4]. The integer $r(C_0)/\gamma_D$ is the order in $\mathbb{Z}^n / \text{Im}(N)$ of the class of the first basis vector e_1 [Lorenzini 2013a, Lemma 3.5].

(b) *The integer g_D is nonnegative.* More precisely,

$$g_D - 2\beta(G_D) \geq \left(\frac{r(C_0)}{\gamma_D}\right) + \gcd\left(\frac{r(D)}{\gamma_D}, \frac{r(C_0)}{\gamma_D}\right) - 2 \geq 0. \tag{4.10.1}$$

To prove the first inequality, complete the pair $(N, R_D/\gamma_D)$ into an arithmetical graph (G', M', R') by adding a chain attached to D , as in [Lorenzini 2013a, §3.15]. Clearly, $\beta(G') = \beta(G_D)$. The graphs G' and G_D differ in only two vertices of degree not equal to 2: the terminal vertex on the new terminal chain on G' has terminal multiplicity $\gcd(r(D)/\gamma_D, r(C_0)/\gamma_D)$, and $d_{G'}(D) = d_{G_D}(D) + 1$. Using (4.1.1) and (4.1.3), it is easy to show that

$$\begin{aligned} &2g_0(G', M', R') - 2\beta(G') \\ &= g_D - 2\beta(G_D) - \left(\frac{r(C_0)}{\gamma_D} - 1\right) - \left(\gcd\left(\frac{r(D)}{\gamma_D}, \frac{r(C_0)}{\gamma_D}\right) - 1\right). \end{aligned} \tag{4.10.2}$$

The integer $g_0(G') - \beta(G')$ is always nonnegative [Lorenzini 1989, Theorem 4.10], and the statement follows.

(c) In analogy with the arithmetic genus of curves on surfaces, we define, given $Z \in \mathbb{Z}^n$, a (possibly negative) integer $p_a(Z)$ as follows. If $Z = C_i$ is a vertex of G_D , we let $p_a(Z) = 0$. We let $p_a(rC_i)$ be defined by the formula $2p_a(rC_i) - 2 = r^2C_i^2 + r(|C_i^2| - 2)$ (where we have abbreviated $C_i \cdot C_i$ by C_i^2). Since $r^2 - r$ is always even, $p_a(rC_i)$ is an integer. In general, when $Z = \sum_{i=1}^n r_i C_i$, we let

$$Z^2 := \sum_{1 \leq i, j \leq n} r_i r_j (C_i \cdot C_j)$$

and set

$$2p_a(Z) - 2 := Z^2 + \sum_{i=1}^n r_i (|C_i^2| - 2).$$

We leave it to the reader to check that

$$g_D = 2p_a(R_D/\gamma_D) - 2 + \frac{r(D)}{\gamma_D} \left(\frac{r(C_0)}{\gamma_D} + 1\right).$$

(d) The integer g_D is even when either $r(C_0)$ is odd or $r(D)$ is even. This can be seen from the formula for g_D in (c) or from (4.10.2).

(e) Assume that G_D is a tree. Then the order $|\det(N)|$ of the group $\Phi_N := \mathbb{Z}^n / N(\mathbb{Z}^n)$ can be computed completely in terms of the vector R_D/γ_D and of the graph G_D (see [Lorenzini 2013a, Theorem 3.14]), and we find that

$$|\det(N)| = \frac{r(D)}{\gamma_D} \frac{r(C_0)}{\gamma_D} \prod_{\text{vertices } A \text{ of } G_D} \left(\frac{r(A)}{\gamma_D} \right)^{d_{G_D}(A)-2},$$

where $d_{G_D}(A)$ is the degree of the vertex A in the graph G_D . Recall now the formula (4.1.3):

$$g_D = \left(\frac{r(D)}{\gamma_D} - 1 \right) + \left(\frac{r(C_0)}{\gamma_D} - 1 \right) + \sum_{\text{vertices } A \text{ of } G_D} \left(\frac{r(A)}{\gamma_D} - 1 \right) (d_{G_D}(A) - 2).$$

This last expression is surprisingly similar to the expression for $|\det(N)|$. This motivates the following result.

Let $x > 0$ be any integer, and define the function $\ell(x) := \sum_{q \text{ prime}} \text{ord}_q(x)(q - 1)$. Then

$$\ell(|\det(N)|) \leq g_D. \tag{4.10.3}$$

This result is not used in the remainder of this paper, and we will provide here only a sketch of proof.

Sketch of proof. We complete the pair $(N, R_D/\gamma_D)$ into an arithmetical graph (G', M', R') by adding a chain attached to D , as in [Lorenzini 2013a, §3.15]. The order of the component group $\Phi(M')$ is given in [Lorenzini 1989, Corollary 2.5], and the relation between $\det(N)$ and $|\Phi(M')|$ is discussed in the proof of Theorem 3.14 in [Lorenzini 2013a]. We can then bound $|\Phi(M')|$ in terms of $g_0(G', M', R')$ using [Lorenzini 1989, Corollary 4.8], which states that $\ell(|\Phi(M')|) \leq 2g_0(G', M', R')$. The inequality $\ell(|\det(N)|) \leq g_D$ follows then from (4.10.2). \square

5. The quotient construction

Let K be a complete discrete valuation field with valuation v , ring of integers \mathcal{O}_K , uniformizer π_K and residue field k of characteristic $p > 0$, assumed to be algebraically closed. Let X/K be a smooth proper geometrically connected curve of genus $g > 0$. When $g = 1$, assume in addition that $X(K) \neq \emptyset$. Assume that X/K does not have semistable reduction over \mathcal{O}_K and that it achieves good reduction after a cyclic extension L/K of prime degree q .

Let H denote the Galois group of L/K . Let $\mathcal{Y}/\mathcal{O}_L$ be the smooth model of X_L/L . Let σ denote a generator of H . By minimality of the model \mathcal{Y} , σ defines an automorphism of \mathcal{Y} also denoted by σ (but note that $\sigma : \mathcal{Y} \rightarrow \mathcal{Y}$ is not a morphism

of \mathcal{O}_L -schemes). We also denote by σ the automorphism of \mathcal{Y}_k induced by the action of σ on \mathcal{Y} . Let $\mathcal{Z}/\mathcal{O}_K$ denote the quotient \mathcal{Y}/H , and let $\alpha : \mathcal{Y} \rightarrow \mathcal{Z}$ denote the quotient map. The scheme \mathcal{Z} is normal. The map α induces a natural map $\mathcal{Y}_k \rightarrow \mathcal{Z}_k^{\text{red}}$ that factors as follows:

$$\mathcal{Y}_k \xrightarrow{\rho} \mathcal{Y}_k/\langle\sigma\rangle \rightarrow \mathcal{Z}_k^{\text{red}}.$$

5.1. We claim that the first map is Galois of order $|H|$ and that the second map is the normalization map of $\mathcal{Z}_k^{\text{red}}$. Indeed, let $\text{Spec}(B)$ denote a dense open set of \mathcal{Y} invariant under the action of H . Then $\text{Spec}(B^H)$ is a dense open set of \mathcal{Z} . Let $A := B^H$. Let $P_B = (\pi_L)$ denote the prime ideal of B corresponding to \mathcal{Y}_k , and let $P_A := P_B \cap A$. We have the natural maps

$$B^H/P_A \hookrightarrow (B/P_B)^H \hookrightarrow B/P_B.$$

The extension of discrete valuation rings $(B^H)_{P_A} \rightarrow B_{P_B}$ induces an extension of residue fields $(B^H)_{P_A}/P_A(B^H)_{P_A} \rightarrow B_{P_B}/P_B B_{P_B}$. We claim that this extension has degree $|H|$. Indeed, our assumption is that the curve X/K does not have good reduction over \mathcal{O}_K . If the residue extension is trivial, the normalization of the curve $\mathcal{Z}_k^{\text{red}}$ is isomorphic to \mathcal{Y}_k and, thus, is of genus g . In addition, we find that $P_A B_{P_B} = (P_B B_{P_B})^{|H|}$ so that $\pi_K A_{P_A} = (P_A A_{P_A})$. The special fiber of \mathcal{Z} is then reduced, and the curve X/K has good reduction over \mathcal{O}_K , a contradiction. It follows then that $P_A B_{P_B} = P_B B_{P_B}$ so that $\pi_K A_{P_A} = (P_A A_{P_A})^{|H|}$. Hence, the multiplicity in \mathcal{Z} of the irreducible component $\mathcal{Z}_k^{\text{red}}$ equals $|H|$.

It is easy to check that, for any $x \in (B/P_B)^H$, $|H|x$ and $x^{|H|}$ belong to A/P_A . Thus, when $|H| \neq p$, A/P_A and $(B/P_B)^H$ have the same field of fractions. When $|H| = p$, it could happen that A/P_A and $(B/P_B)^H$ do not have the same field of fractions, and then the extension of fields of fractions is purely inseparable of degree p with $(B/P_B)^H = B/P_B$. It follows that the special fiber of \mathcal{Z} also has genus g . When $g > 1$, this is not possible since the multiplicity of \mathcal{Z}_k is p . When $g = 1$, it could happen that \mathcal{Z} is the minimal model of X/K with a multiple special fiber. This case cannot happen in our situation because we assumed $X(K) \neq \emptyset$: a K -rational point always reduces to a smooth point in the special fiber. Thus, the automorphism $\sigma : \mathcal{Y}_k \rightarrow \mathcal{Y}_k$ is not trivial. We find that A/P_A and $(B/P_B)^H$ have the same fields of fractions so that the Dedekind domain $(B/P_B)^H$ is the integral closure of A/P_A .

5.2. Let P_1, \dots, P_d be the ramification points of the map $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$. Let Q_1, \dots, Q_d be their images in \mathcal{Z} . The normal scheme \mathcal{Z} is singular exactly at Q_1, \dots, Q_d . Indeed, the morphism $\mathcal{Y} \rightarrow \mathcal{Z}$ is unramified outside these points. If the point Q_i were regular, the morphism would be flat above Q_i [Altman and Kleiman 1970, Corollary V.3.6] and the branch locus would then be pure of codimension 1 [Altman and Kleiman 1970, Theorem VI.6.8], a contradiction.

Consider the regular model $\mathcal{X} \rightarrow \mathcal{Z}$ obtained from \mathcal{Z} by a minimal desingularization. After finitely many blow-ups $\mathcal{X}' \rightarrow \mathcal{X}$, we can assume that the model \mathcal{X}' is such that \mathcal{X}'_k has smooth components and normal crossings and is minimal with this property. Let f denote the composition $\mathcal{X}' \rightarrow \mathcal{Z}$. Let C_0/k denote the strict transform in \mathcal{X}' of the irreducible closed subscheme $\mathcal{Z}_k^{\text{red}}$ of \mathcal{Z} . The curve C_0 has multiplicity $|H|$ in \mathcal{X}' . Let D_1, \dots, D_d denote the irreducible components of \mathcal{X}'_k that meet C_0 . Let r_i denote the multiplicity of D_i , $i = 1, \dots, d$. We assume $d \geq 1$. Our main theorem in this section is this:

Theorem 5.3. *Let X/K be a smooth proper geometrically connected curve of genus $g > 0$ with $X(K) \neq \emptyset$ if $g = 1$. Assume that X/K does not have semistable reduction over \mathcal{O}_K and that it achieves good reduction after a cyclic extension L/K with Galois group H of prime degree p . Keep the above notation, and let Q_i be a singular point of the quotient $\mathcal{Z} := \mathcal{Y}/H$. Let G_{Q_i} denote the graph associated with the curve $f^{-1}(Q_i)$. Let G denote the graph associated with the special fiber \mathcal{X}'_k . Then, for all $i = 1, \dots, d$, the graph G_{Q_i} contains a node of G and p divides r_i .*

Proof. When $d = 1$, the theorem is immediate: the component C_0 of multiplicity p is a terminal vertex of the graph of \mathcal{X}' , and thus, $p|C_0 \cdot C_0| = r_1$. Assume that G_{Q_1} does not contain a node of G . Then since $d = 1$, G does not contain a node. Since the resolution is minimal with normal crossings, none of the components of \mathcal{X}'_k can have self-intersection -1 except possibly for C_0 . It is clear that the graph G is not reduced to a single vertex since the model \mathcal{Z} is singular. Thus, the graph G has a second terminal vertex C' in addition to C_0 . But then, walking on G from C' towards C_0 , we find that the multiplicities can only be strictly increasing. This is a contradiction since all multiplicities on G are divisible by p (because two consecutive ones are), and G must contain a node. We assume from now on that $d > 1$.

Let $A := \text{Jac}(X/K)$. Let $\mathcal{A}_K/\mathcal{O}_K$ denote the Néron model of A/K . Let $\mathcal{A}_L/\mathcal{O}_L$ denote the Néron model of A_L/L , and denote by $\eta : \mathcal{A}_K \times_{\mathcal{O}_K} \mathcal{O}_L \rightarrow \mathcal{A}_L$ the canonical map induced by the functoriality property of Néron models. The special fiber $(\mathcal{A}_K)_k$ is an extension of a finite group $\Phi_{A/K}$, called the group of components, by the connected component of zero $(\mathcal{A}_K)_k^0$ of $(\mathcal{A}_K)_k$:

$$0 \rightarrow (\mathcal{A}_K)_k^0 \rightarrow (\mathcal{A}_K)_k \rightarrow \Phi_{A/K} \rightarrow 0.$$

Assume by contradiction that p is coprime to one of the r_i 's. Without loss of generality, we may assume that $p \nmid r_d$. For each $i = 1, \dots, d$, choose a point $x_i \in D_i$ such that x_i is a regular point of $(\mathcal{X}'_k)^{\text{red}}$. Since K is complete, we can find a closed point R_i of X of degree r_i over K and such that the closure of R_i in \mathcal{X}' meets the special fiber \mathcal{X}'_k exactly in x_i (see, e.g., [Gabber et al. 2013, Proposition 8.4(3)]). For each $i = 1, \dots, d - 1$, consider the following divisor of degree 0 on X :

$$S_i := \frac{r_d}{\gcd(r_i, r_d)} R_i - \frac{r_i}{\gcd(r_i, r_d)} R_d.$$

We also denote by S_i its image in $\text{Jac}(X)/K$. We recall below Raynaud's description of the Néron model of a Jacobian in order to be able to describe explicitly the image of S_i under both the reduction map $\text{Jac}(X)(K) \rightarrow \Phi_{A/K}$ and the reduction map $\text{Jac}(X)(L) \rightarrow (\mathcal{A}_L)_k(k)$. We will be able to contradict the hypothesis that $p \nmid r_d$ by considering the reductions of $\sum_{i=1}^{d-1} \gcd(r_i, r_d) S_i$.

Raynaud [1970] exhibited an explicit separated quotient Q_K/\mathcal{O}_K of the open subfunctor of $\text{Pic}_{\mathcal{X}'/\mathcal{O}_K}$ consisting of line bundles of total degree 0, and he showed that, when the residue field k is algebraically closed, Q_K/\mathcal{O}_K is isomorphic to the Néron model of A/K [Bosch et al. 1990, Theorem 9.5.4(a)]. The canonical map $Q_K(K) \rightarrow \Phi_{Q_K}$ is described as follows [Bosch et al. 1990, Lemma 9.5.9, Theorem 9.6.1]. Represent an element of $Q_K(K)$ by a line bundle \mathcal{L} on X of degree 0. Let $\bar{\mathcal{L}}$ denote an extension of \mathcal{L} to \mathcal{X}' . Number the irreducible components of \mathcal{X}'_k as C_1, \dots, C_v . Consider the map $\bigoplus_i \mathbb{Z} C_i \rightarrow \text{Hom}(\bigoplus_i \mathbb{Z} C_i, \mathbb{Z})$ that sends C_i to the map δ_{C_i} with $\delta_{C_i}(C_j) := (C_i \cdot C_j)$. The group Φ_M is isomorphic to the torsion subgroup of the cokernel of this map. The group of components Φ_{Q_K} is isomorphic to Φ_M , and under this isomorphism, the image of \mathcal{L} under $Q_K(K) \rightarrow \Phi_{Q_K}$ is the map $\delta_{\mathcal{L}}$ with $\delta_{\mathcal{L}}(C_i) := (C_i \cdot \bar{\mathcal{L}})$. It follows immediately from these facts that the image in Φ_{Q_K} of $S_i \in \text{Jac}(X)(K)$ can be identified with the image τ_i of the vector $E(D_i, D_d)$ in Φ_M (notation as in 3.1 and 3.3).

Consider now the reduction map $Q_L(L) \rightarrow (Q_L)_k(k)$. The closure of any point in the preimage under $X_L \rightarrow X$ of the closed point R_i meets the special fiber of the smooth model \mathcal{Y} of X_L only at the point P_i . The line bundle \mathcal{L} corresponding to the divisor S_i pulls back to a line bundle \mathcal{L}_L on X_L . We find that the reduction of $\mathcal{L}_L \in \text{Jac}(X_L)(L)$ is the point of $\text{Jac}(\mathcal{Y}_k)(k)$ corresponding to the divisor $\text{lcm}(r_i, r_d)(P_i - P_d)$.

We may now find a contradiction to the assertion that $p \nmid r_d$ when the quotient of \mathcal{Y}_k by the action of H has genus 0. As we indicated above, the element $\sum_{i=1}^{d-1} \gcd(r_i, r_d) S_i$ in $\text{Jac}(X)(K)$ reduces to the element $\sum_{i=1}^{d-1} \gcd(r_i, r_d) \tau_i$ in Φ_M . Proposition 3.5 shows that the latter element is zero in Φ_M . Thus, $\sum_{i=1}^{d-1} \gcd(r_i, r_d) S_i$ reduces in the connected component $(Q_K)_k^0$. Our additional hypothesis implies that this connected component is unipotent. This follows from [Bosch et al. 1990, Theorem 9.5.4] if the greatest common divisor of the multiplicities of the components of \mathcal{X}'_k is 1 and from [Liu et al. 2004, Proposition 7.1] in general. It follows that the image of $(Q_K)_k^0$ under the canonical map $\eta: \mathcal{A}_K \times_{\mathcal{O}_K} \mathcal{O}_L \rightarrow \mathcal{A}_L$ is trivial.

Consider now the element $\sum_{i=1}^{d-1} \gcd(r_i, r_d) S_i$ in $\text{Jac}(X_L)(L)$. Our discussion above shows that it reduces to the element $r_d(\sum_{i=1}^{d-1} r_i(P_i - P_d))$ in $\text{Jac}(\mathcal{Y}_k)(k)$. We have thus proved that $r_d(\sum_{i=1}^{d-1} r_i(P_i - P_d)) = 0$ in $\text{Jac}(\mathcal{Y}_k)(k)$. Our hypothesis

on the quotient of \mathcal{Y}_k by H implies that each $P_i - P_d$ has order p (Proposition 2.5). Since $r_d(\sum_{i=1}^{d-1} r_i(P_i - P_d)) = 0$ and we assume that p does not divide r_d , we can conclude that $\sum_{i=1}^{d-1} r_i(P_i - P_d) = 0$. Then Proposition 2.5 implies that p divides r_i for all $i = 1, \dots, d - 1$. Since $|C_0 \cdot C_0|_p = r_1 + \dots + r_d$, it follows that p divides r_d , which contradicts our assumption.

When the quotient of \mathcal{Y}_k by the action of H has positive genus, the image of $(\mathcal{Q}_K)_k^0$ under the canonical map $\eta : \mathcal{A}_K \times_{\mathcal{O}_K} \mathcal{O}_L \rightarrow \mathcal{A}_L$ is not trivial, and the following additional considerations must be discussed. Let $\text{Norm}(\mathcal{X}')$ denote the normalization of \mathcal{X}' in the field of fractions of \mathcal{Y} . Since \mathcal{Y} is integral over \mathcal{Z} , we have a natural map $\text{Norm}(\mathcal{X}') \rightarrow \mathcal{Y}$. All components of \mathcal{X}' are rational except possibly the component C_0 [Lorenzini 2013a, Lemma 2.10].

By construction, we have a natural map $\text{Norm}(\mathcal{X}') \rightarrow \mathcal{X}' \times_{\mathcal{O}_K} \mathcal{O}_L$. Let $\mathcal{N} \rightarrow \text{Norm}(\mathcal{X}')$ denote a resolution of the singularities of $\text{Norm}(\mathcal{X}')$. Consider the commutative diagram of \mathcal{O}_L -morphisms

$$\begin{array}{ccccc} \mathcal{N} & \longrightarrow & \text{Norm}(\mathcal{X}') & \longrightarrow & \mathcal{Y} \\ & & \downarrow & & \downarrow \\ & & \mathcal{X}' \times_{\mathcal{O}_K} \mathcal{O}_L & \longrightarrow & \mathcal{Z} \times_{\mathcal{O}_K} \mathcal{O}_L \end{array}$$

The maps $\mathcal{N} \rightarrow \text{Norm}(\mathcal{X}') \rightarrow \mathcal{X}' \times_{\mathcal{O}_K} \mathcal{O}_L$ induce maps of the associated Picard functors

$$\text{Pic}_{\mathcal{X}'/\mathcal{O}_K} \times_{\mathcal{O}_K} \mathcal{O}_L \cong \text{Pic}_{\mathcal{X}' \times_{\mathcal{O}_K} \mathcal{O}_L/\mathcal{O}_L} \rightarrow \text{Pic}_{\text{Norm}(\mathcal{X}')/\mathcal{O}_L} \rightarrow \text{Pic}_{\mathcal{N}/\mathcal{O}_L},$$

whose composition induces the canonical map of Néron models

$$\eta : \mathcal{Q}_K \times_{\mathcal{O}_K} \mathcal{O}_L \rightarrow \mathcal{Q}_L.$$

Considering the special fibers over k , we obtain a commutative diagram

$$\begin{array}{ccc} \text{Pic}_{\mathcal{N}_k/k}^0 & \longrightarrow & (\mathcal{Q}_L)_k^0 \\ \uparrow & & \uparrow \\ \text{Pic}_{\mathcal{X}'_k/k}^0 & \longrightarrow & (\mathcal{Q}_K)_k^0 \end{array}$$

Since we do not have additional information on the special fiber \mathcal{X}'_k , we cannot conclude that the bottom horizontal map is an isomorphism. It is however faithfully flat [Raynaud 1970, Corollaire 4.1.2]. Since the special fiber of \mathcal{Y} is reduced, we find that the top horizontal map is an isomorphism [Bosch et al. 1990, Theorem 9.5.4].

Let D denote the irreducible component of \mathcal{N}_k lying above \mathcal{Y}_k . The composition $D \hookrightarrow \mathcal{N}_k \rightarrow \mathcal{Y}_k$ is an isomorphism. The image of D in $(\mathcal{X}')_k^{\text{red}}$ is the curve C_0 , and we will identify the map $D \rightarrow C_0$ with the quotient map $\rho : \mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle \sigma \rangle$. Consider the following diagram, whose top right horizontal morphism is an isomorphism:

$$\begin{array}{ccccc}
 \mathrm{Pic}_D^0(k) & \longleftarrow & \mathrm{Pic}_{\mathcal{N}_k}^0(k) & \xrightarrow{\sim} & (Q_L)_k^0(k) \\
 \rho^* \uparrow & & \uparrow & & \uparrow \\
 \mathrm{Pic}_{C_0}^0(k) & \longleftarrow & \mathrm{Pic}_{\mathcal{X}_k}^0(k) & \longrightarrow & (Q_K)_k^0(k)
 \end{array}$$

We may now conclude the proof of [Theorem 5.3](#) using the same method as in the case where the reduction of $\mathrm{Jac}(X)/K$ is purely unipotent. Consider again the element $\sum_{i=1}^{d-1} \mathrm{gcd}(r_i, r_d) S_i$ in $\mathrm{Jac}(X)(K)$, which reduces to the element $\sum_{i=1}^{d-1} \mathrm{gcd}(r_i, r_d) \tau_i$ in Φ_M . [Proposition 3.5](#) shows that the latter element is zero in Φ_M . Thus, $\sum_{i=1}^{d-1} \mathrm{gcd}(r_i, r_d) S_i$ reduces in the connected component $(Q_K)_k^0$. Consider now the element $\sum_{i=1}^{d-1} \mathrm{gcd}(r_i, r_d) S_i$ in $\mathrm{Jac}(X_L)(L)$. Our discussion above shows that it reduces to the element $r_d(\sum_{i=1}^{d-1} r_i (P_i - P_d))$ in $\mathrm{Jac}(\mathcal{Y}_k)(k)$.

Since $\mathrm{Pic}_{\mathcal{X}'_k/k}^0 \rightarrow (Q_K)_k^0$ is a faithfully flat morphism and each of the above squares commutes, the element $\sum_{i=1}^{d-1} \mathrm{gcd}(r_i, r_d) S_i$, which reduces to $r_d(\sum_{i=1}^{d-1} r_i (P_i - P_d))$ in $\mathrm{Pic}_{\mathcal{Y}_k/k}^0(k)$, in fact reduces to an element in $\rho^*(\mathrm{Jac}(\mathcal{Y}_k/\langle\sigma\rangle))$. Thus, the image of $r_d(\sum_{i=1}^{d-1} r_i (P_i - P_d))$ in $\mathrm{Jac}(\mathcal{Y}_k)/\rho^*(\mathrm{Jac}(\mathcal{Y}_k/\langle\sigma\rangle))$ is trivial. Each $P_i - P_d$ defines an element of order p in $\mathrm{Jac}(\mathcal{Y}_k)/\rho^*(\mathrm{Jac}(\mathcal{Y}_k/\langle\sigma\rangle))$ ([Proposition 2.5](#)). Since $r_d(\sum_{i=1}^{d-1} r_i (P_i - P_d)) = 0$, we conclude that $\sum_{i=1}^{d-1} r_i (P_i - P_d) = 0$. Then [Proposition 2.5](#) implies that p divides r_i for all $i = 1, \dots, d-1$, and since $|C_0 \cdot C_0| p = r_1 + \dots + r_d$, we find that p divides r_d , which contradicts our assumption.

Now that we know that p divides r_i , we see that the multiplicities on the chain of G that leaves C_0 starting with D_i can only be increasing or constant because this chain of vertices of degree 2 contains no vertex of self-intersection -1 . If D_i is not a node of G , we continue along this chain and find either a terminal vertex or a node of G . We cannot find a terminal vertex because the multiplicity of a terminal vertex can only be at most the multiplicity of its unique neighbor with equality only if the self-intersection of the terminal vertex is -1 . Thus, G_{Q_i} contains a node of G . \square

Remark 5.4. Let N_i denote the intersection matrix of the exceptional divisor, with smooth components and normal crossings, of a resolution of the $\mathbb{Z}/p\mathbb{Z}$ -quotient singularity Q_i . We recall here some properties of N_i :

- (a) It is negative definite (attributed to Du Val in [\[Lipman 1969, Lemma 14.1\]](#)).
- (b) The graph $G(N_i)$ associated with N_i is a tree, and all components of the exceptional divisor are rational [\[Lorenzini 2013a, Theorem 2.8\]](#).
- (c) Let n_i denote the number of irreducible components in the exceptional divisor. The Smith group $\Phi_{N_i} := \mathbb{Z}^{n_i} / \mathrm{Im}(N_i)$ is killed by p [\[Lorenzini 2013a, Theorem 2.6\]](#).
- (d) The fundamental cycle Z of N_i is such that $|Z^2| \leq p$ [\[Lorenzini 2013a, Theorem 2.3, Remark 2.4\]](#).

6. The weakly ramified case

We present in this section some applications of [Theorem 5.3](#). Let us recall our notation. Let K be a complete discrete valuation field with valuation v , ring of integers \mathcal{O}_K , uniformizer π_K and residue field k of characteristic $p > 0$, assumed to be algebraically closed. Let X/K be a smooth proper geometrically connected curve of genus $g > 0$. When $g = 1$, we assume in addition that $X(K) \neq \emptyset$.

Assume that X/K does not have semistable reduction over \mathcal{O}_K and that it achieves good reduction after a cyclic extension L/K of prime degree p . Let $H = \langle \sigma \rangle$ denote the Galois group of L/K . Let $\mathcal{Y}/\mathcal{O}_L$ be the smooth model of X_L/L . Let $\mathcal{Z}/\mathcal{O}_K$ denote the quotient \mathcal{Y}/H with singular points Q_1, \dots, Q_d and $d \geq 1$. Recall the regular model $f : \mathcal{X}' \rightarrow \mathcal{Z}$ introduced in [5.2](#).

6.1. The resolution of a singularity Q of \mathcal{Z} is a local process and depends only on the local ring $\mathcal{O}_{\mathcal{Z}, Q}$. It seems therefore natural to try to relate the “complexity” of the resolution graph to some local invariants of $\mathcal{O}_{\mathcal{Z}, Q}$. In this respect, we propose the following.

Consider the Galois morphism $\rho : \mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle \sigma \rangle$. Associated with any point $Q \in \mathcal{Y}_k/\langle \sigma \rangle$ is the following measure of the ramification of ρ over Q :

$$v(Q) := \delta(P) = \sum_{j=0}^{\infty} (|H_j(P)| - 1),$$

where P is the preimage of Q in \mathcal{Y}_k and $H_j(P)$ denotes the j -th higher ramification group at P . (For more general morphisms, we would define $v(Q) := \sum_{P \in \rho^{-1}(Q)} \delta(P)$.) Recall from [2.2](#) that the morphism is *weakly ramified* at P if $\delta(P) = 2(p-1)$. Our guess is that $v(Q)$ should also be an important measure of how complicated the exceptional divisor of the resolution of Q is. To formulate this guess more precisely, we compare the expressions of the genus g in the Riemann–Hurwitz formula and in the adjunction formula. The Riemann–Hurwitz formula for the morphism ρ can be rephrased as

$$2g = 2g(\mathcal{Y}_k) = 2|H|g(C_0) - 2(|H| - 1) + \sum_{i=1}^d v(Q_i).$$

Consider now the model \mathcal{X}' . By hypothesis, it is minimal with the property that the special fiber has smooth components and normal crossings. Thus, none of the vertices A in the graph $G := G(\mathcal{X}')$ with degree 1 or 2 can have self-intersection -1 (we use here also the fact that only the curve C_0 can have positive genus [[Lorenzini 2013a](#), Lemma 2.10]). Moreover, since the curve X/K has potentially good reduction, the graph $G(\mathcal{X}')$ is a tree [[Lorenzini 2013a](#), Lemma 2.10].

The adjunction formula

$$2g - 2 = \mathcal{X}'_k \cdot \mathcal{X}'_k + \mathcal{X}'_k \cdot \Omega,$$

with Ω a relative canonical divisor of $\mathcal{X}'/\mathcal{O}_K$, can be rewritten as

$$\begin{aligned} 2g &= 2|H|g(C_0) + \sum_{\text{vertex } A \text{ of } G} (r(A) - 1)(d_G(A) - 2) \\ &= 2|H|g(C_0) - 2(|H| - 1) + \sum_{i=1}^d \left(|H| - 1 + \sum_{\text{vertex } A \text{ of } G_{Q_i}} (r(A) - 1)(d_G(A) - 2) \right) \\ &= 2|H|g(C_0) - 2(|H| - 1) + \sum_{i=1}^d \gamma_{D_i} g_{D_i}, \end{aligned} \tag{6.1.1}$$

where D_1, \dots, D_d are the vertices attached to C_0 in the tree $G(\mathcal{X}')$ and the integers γ_{D_i} and g_{D_i} are defined as in 4.1 and (4.1.2). Since the graph G_{D_i} is nothing but the graph G_{Q_i} of the desingularization of Q_i , we define our measure of the desingularization of Q_i to be $\gamma_{Q_i} g_{Q_i} := \gamma_{D_i} g_{D_i}$ for each $i = 1, \dots, d$. The integer $g_{Q_i} := g_{D_i}$ depends only on the intersection matrix of the desingularization and the marked vertex D_i on its graph. Since $r(C_0) = p$ and is divisible by γ_{Q_i} , we find that $\gamma_{Q_i} = 1$ or p .

6.2. Our guess regarding the resolution $\mathcal{X}' \rightarrow \mathcal{Z}$ of the singularities of \mathcal{Z} is that

$$\gamma_{Q_i} g_{Q_i} = v(Q_i) \text{ holds for all } i = 1, \dots, d.$$

This equality would have interesting implications. For instance, since $H = \mathbb{Z}/p\mathbb{Z}$, we always have $v(Q)$ divisible by $p - 1$ so that $p - 1$ divides $\gamma_{Q_i} g_{Q_i}$ when $\gamma_{Q_i} g_{Q_i} = v(Q_i)$. Since $\gamma_{Q_i} = 1$ or p , we find that

$$p - 1 \text{ divides } g_{Q_i} \text{ when } \gamma_{Q_i} g_{Q_i} = v(Q_i).$$

Examples where $g_{Q_i} = 2(p - 1)$ and $3(p - 1)$ are given in 4.7 and Remark 4.9. It immediately follows from the Riemann–Hurwitz formula and the adjunction formula that:

Lemma 6.3. *With the above notation and hypotheses,*

$$\sum_{i=1}^d v(Q_i) = \sum_{i=1}^d \gamma_{Q_i} g_{Q_i}. \tag{6.3.1}$$

We now prove the equality $\gamma_{Q_i} g_{Q_i} = v(Q_i) = 2(p - 1)$ for all $i = 1, \dots, d$ in the weakly ramified case, using Theorem 5.3.

Theorem 6.4. *Let X/K be a curve with potentially good reduction after a ramified extension L/K of prime degree p . Keep the above notation. Then for all $i = 1, \dots, d$:*

- (a) *We have $\gamma_{Q_i} g_{Q_i} \geq 2(p-1)$ and $v(Q_i) \geq 2(p-1)$.*
- (b) *If the ramification points of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ are all weakly ramified (in particular, if \mathcal{Y}_k is ordinary), then $\gamma_{Q_i} g_{Q_i} = v(Q_i) = 2(p-1)$.*

Proof. (a) The fact that $v(Q_i) \geq 2(p-1)$ follows immediately from the properties of a wildly ramified extension: the higher ramification groups H_0 and H_1 must be nontrivial. To prove that $\gamma_{Q_i} g_{Q_i} \geq 2(p-1)$, we note first that [Theorem 5.3](#) shows that $p \mid r_i$. The inequality follows then from [Proposition 4.3](#).

(b) When the ramification points of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ are all weakly ramified, we have $v(Q_i) = 2(p-1)$ ([2.2](#)). It follows from ([6.3.1](#)) and from the fact that $\gamma_{Q_i} g_{Q_i} \geq 2(p-1)$ proven in (a) that $\gamma_{Q_i} g_{Q_i} = 2(p-1)$. \square

Remark 6.5. Without the use of [Theorem 5.3](#), we could only argue that $\gamma_{Q_i} g_{Q_i} \geq p-1$. Indeed, if $r(C_0)$ does not divide $r(D_i)$, then $\gamma_{D_i} = 1$. Then we can use the fact that $g_{Q_i} \geq r(C_0) - 1$ established in [Remark 4.10](#).

Using the notation γ_{Q_i} introduced in this section, we may now state a corollary to [Theorem 5.3](#).


Corollary 6.6. *Let X/K be a curve with potentially good reduction after a wildly ramified Galois extension L/K of degree p as in [Theorem 5.3](#). Let N_i denote the intersection matrix associated with the resolution of Q_i . Assume that $\gamma_{Q_i} = 1$. Then p^2 divides $\det(N_i)$.*

Proof. The graph associated with the matrix N_i is G_{Q_i} with a marked vertex D_i on it. Let R_{D_i} denote the vector of multiplicities of the components of the resolution of Q_i . Then the determinant of N_i can be computed in terms of the coefficients of R_{D_i}/γ_{D_i} (see [[Lorenzini 2013a](#), Theorem 3.14]). In particular, it is known that $(r(C_0)/\gamma_{D_i}) \gcd(r(C_0)/\gamma_{D_i}, r(D_i)/\gamma_{D_i})$ divides $\det(N_i)$. Under our hypotheses, $r(C_0) = p$, p divides $r(D_i)$ ([Theorem 5.3](#)) and $\gamma_{D_i} = 1$. \square

Remark 6.7. Let X/K be a curve with potentially good reduction after a wildly ramified extension L/K of degree p as in [Theorem 5.3](#). Let N_i denote the intersection matrix associated with the resolution of Q_i . Then p kills the Smith group Φ_{N_i} [[Lorenzini 2013a](#), Theorem 2.6], and thus, $|\det(N_i)|$ is a power of p . It follows from ([4.10.3](#)) that $\text{ord}_p(|\det(N_i)|)(p-1) \leq g_{D_i}$.

In the examples of graphs and matrices N_i given in [Remark 4.9](#) with $g_{D_i} = 3(p-1)$, we find that both $|\det(N_i)| = p^2$ and $|\det(N_i)| = p^3$ can occur: the former in (b) and (c)(ii), and the latter in (a) and (c)(i).

Theorem 6.8. *Let X/K be a curve with potentially good reduction after a wildly ramified Galois extension L/K of degree p . Assume that all ramification points of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ are weakly ramified (this is the case if \mathcal{Y}_k is ordinary). Keep the above notation. Then, for all $i = 1, \dots, d$, we have $r_i = p$, and G_{Q_i} is a graph¹ with a single node C_i of degree 3:*



The intersection matrix $N(p, \alpha_i, r_1(i))$ of the resolution of Q_i is uniquely determined as in 4.7 by the two integers α_i and $r_1(i)$ with $1 \leq r_1(i) < p$. The integer α_i is the number of vertices of self-intersection -2 (including the node C_i) on the chain in G_{Q_i} connecting the node C_0 to the single node C_i of G_{Q_i} , and this integer α_i is divisible by p .

Proof. Theorem 6.4(b) shows that $\gamma_{Q_i} g_{Q_i} = 2(p-1)$ for all $i = 1, \dots, d$. Proposition 4.3 classifies the graphs with $\gamma_{Q_i} g_{Q_i} = 2(p-1)$, and the statement on the shape of the graph follows.

The Smith group of the intersection matrix $N(p, \alpha_i, r_1(i))$ is computed in [Lorenzini 2013a, §3.19, Lemma 3.21] and is found to be of order p^2 and killed by p if and only if p divides α_i . Theorem 2.6(c) of [Lorenzini 2013a] shows that this Smith group must be killed by p . The divisibility $p \mid \alpha_i$ follows. \square

Remark 6.9. It is natural to wonder whether the statements of Theorems 6.4(b) and 6.8 hold for the resolution of Q_i when P_i is a weakly ramified ramification point of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ without also assuming as we do in Theorems 6.4(b) and 6.8 that all ramification points are weakly ramified.

Corollary 6.10. *Let X/K be a curve with potentially good reduction after a wildly ramified Galois extension L/K of degree p as in Theorem 6.8. Suppose that $g > 1$ and that all ramification points of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle\sigma\rangle$ are weakly ramified. Then:*

- (a) $X(K) \neq \emptyset$.
- (b) Let A/K denote the Jacobian of X/K . Let $\mathcal{A}/\mathcal{O}_K$ be its Néron model. Then the unipotent part U/k of the connected component of the identity in \mathcal{A}_k/k is a product of additive groups $\mathbb{G}_{a,k}$.
- (c) The group of components $\Phi_{A,K}$ of the Néron model is isomorphic to $(\mathbb{Z}/p\mathbb{Z})^{2d-2}$.

¹A bullet \bullet represents an irreducible component of the desingularization of Q_i . A positive number next to a vertex is the multiplicity of the corresponding component while a negative number next to a vertex is the self-intersection of the component.

Proof. Part (a) is immediate since it follows from [Theorem 6.8](#) that a regular model of X/K contains a component of multiplicity 1. It follows from [[Penniston 2000](#), [Theorem 2.4](#)] that p kills U since the maximal multiplicity in the regular model $\mathcal{X}'/\mathcal{O}_K$ is equal to p . That U is now split follows from [[Serre 1959](#), [Proposition VII.11.11](#)]. This proves (b).

The order of $\Phi_{A,K}$ can be computed using the intersection matrix of the regular model \mathcal{X}' . Since the associated graph is a tree, we find using [[Lorenzini 1989](#), [Corollary 2.5](#)] that $|\Phi_{A,K}| = p^{2d-2}$. Part (c) follows since $\Phi_{A,K}$ is killed by $[L : K]$ because A/K has potentially good reduction [[Edixhoven et al. 1996](#)]. \square

Note that in general the special fiber \mathcal{A}_k/k need not be killed by p even when its subgroup U and quotient $\Phi_{A,K}$ are both killed by p (see [[Liu and Lorenzini 2001](#)] for a general discussion of such phenomena).

6.11. Let A/K be the Jacobian of a smooth proper and geometrically connected curve X/K having a K -rational point. For use in our next corollary, we recall below the main result of [[Bosch and Lorenzini 2002](#), [Theorem 4.6](#)]. Identify A/K with its dual A'/K via the map $-\varphi_{[\Theta]} : A \rightarrow A'$ as in [[Bosch and Lorenzini 2002](#)] just before [Theorem 4.6](#). Let $\mathcal{X}'/\mathcal{O}_K$ denote a regular model of X/K . Let M be the intersection matrix of \mathcal{X}'_k . Identify, as recalled in [[Bosch and Lorenzini 2002](#), [Theorem 2.3](#)], the component group $\Phi_{A/K}$ with the group of components Φ_M of M (Φ_M is the torsion subgroup of $\mathbb{Z}^v/\text{Im}(M)$). Then Grothendieck's pairing

$$\langle \cdot, \cdot \rangle_K : \Phi_{A/K} \times \Phi_{A/K} \rightarrow \mathbb{Q}/\mathbb{Z}$$

coincides with the pairing $\langle \cdot, \cdot \rangle_M : \Phi_{A/K} \times \Phi_{A/K} \rightarrow \mathbb{Q}/\mathbb{Z}$ considered in [3.1](#). In particular, this pairing is nondegenerate. Recall also the definition of the functorial subgroup $\Phi_{A/K}^0$ of $\Phi_{A/K}$ in [1.3](#). We denote by $(\Phi_{A/K}^0)^\perp$ the orthogonal of $\Phi_{A/K}^0$ under Grothendieck's pairing.

Corollary 6.12. *Let A/K be the Jacobian of a curve X/K of genus $g > 1$ having potentially good reduction after a Galois extension L/K of degree p as in [Theorem 6.8](#). Assume that all ramification points of $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/\langle \sigma \rangle$ are weakly ramified. Then $\Phi_{A/K}$ is a $\mathbb{Z}/p\mathbb{Z}$ -vector space of dimension $2d - 2$, and $\Phi_{A/K}^0$ is a subspace of dimension $d - 1$. Moreover, $\Phi_{A/K}^0 = (\Phi_{A/K}^0)^\perp$.*

Proof. It follows from [Corollary 6.10](#) that $X(K) \neq \emptyset$. We can thus use the results of [[Bosch and Lorenzini 2002](#)] recalled above. We produce below explicit generators for the groups $\Phi_{A/K}$ and $\Phi_{A/K}^0$. For each singular point Q_i on the model $\mathcal{Z}/\mathcal{O}_K$, denote by A_i and B_i the terminal components of multiplicity 1 in the exceptional divisor of the resolution of Q_i in \mathcal{X}' . Let α_i denote the image in $\Phi_{A/K}$ of the vector $E(A_i, B_i)$, $i = 1, \dots, d - 1$ (notation as in [3.1](#)). Let β_i denote the image in $\Phi_{A/K}$ of the vector $E(A_i, A_d)$, $i = 1, \dots, d - 1$. We have seen in [Corollary 6.10](#) that $\Phi_{A/K}$ is a $\mathbb{Z}/p\mathbb{Z}$ -vector space of dimension $2(d - 1)$.

We claim that

$$\{\alpha_1, \dots, \alpha_{d-1}, \beta_1, \dots, \beta_{d-1}\}$$

is a basis for $\Phi_{A/K}$ and that $\{\alpha_1, \dots, \alpha_{d-1}\}$ is a basis for $\Phi_{A/K}^0$. To prove our claim, consider the matrix $V := (\langle \alpha_i, \beta_j \rangle)_{1 \leq i, j \leq d-1}$ with coefficients in \mathbb{Q}/\mathbb{Z} . We can use the computation (4.8.1) to show that V is the diagonal matrix

$$\text{diag}(c_1/p \pmod{\mathbb{Z}}, \dots, c_{d-1}/p \pmod{\mathbb{Z}}),$$

where, for each $i = 1, \dots, d-1$, $0 < c_i < p$ and p divides $c_i r_1(i) - 1$. In particular, $c_i/p \neq 0$ in \mathbb{Q}/\mathbb{Z} . It follows that the set $\{\alpha_1, \dots, \alpha_{d-1}, \beta_1, \dots, \beta_{d-1}\}$ is linearly independent in $(\mathbb{Z}/p\mathbb{Z})^{2d-2}$. Hence, it is a basis.

It follows from the explicit computations in [Lorenzini 2000, Proposition 3.7(a)], that $\langle \alpha_i, \alpha_j \rangle = 0$ for all $1 \leq i, j \leq d-1$. Since the pairing $\langle \cdot, \cdot \rangle$ is perfect on $(\mathbb{Z}/p\mathbb{Z})^{2d-2}$, we find that $\{\alpha_1, \dots, \alpha_{d-1}\}$ generates a maximal isotropic subspace.

It remains to show that $\alpha_1, \dots, \alpha_{d-1}$ belong to $\Phi_{A/K}^0$ and that neither $\beta_1, \dots, \beta_{d-1}$ nor any nontrivial linear combination of $\beta_1, \dots, \beta_{d-1}$ belong to $\Phi_{A/K}^0$. For this, since K is complete, we can pick for each $i = 1, \dots, d-1$ two K -rational points a_i and b_i of X whose closure in \mathcal{X}' intersects \mathcal{X}'_k in a smooth point of A_i and B_i , respectively (see, e.g., [Bosch et al. 1990, Corollary 9.1.9]). Then $a_i - b_i$ and $a_i - a_d$ are divisors of degree 0 on X , which we identify with K -rational points in the Jacobian A/K of X/K . These rational points reduce in the component group $\Phi_{A/K}$ of the Néron model of A/K to the points α_i and β_i , respectively. Since $A(K) \subset A(L)$, we can reduce $a_i - b_i$ in the special fiber of the Néron model $\mathcal{A}'/\mathcal{O}_L$. This special fiber is isomorphic to the Jacobian of the special fiber \mathcal{Y}_k of the smooth model $\mathcal{Y}/\mathcal{O}_L$ of X_L/L . It is clear that, by construction, the reduction of $a_i - b_i$ is trivial so that $\alpha_i \in \Phi_{A/K}^0$ for $i = 1, \dots, d-1$. On the other hand, the reduction of $a_i - a_d$ is the divisor $P_i - P_d$, which is a nontrivial p -torsion point when viewed in the quotient $\mathcal{A}'_k/\eta(\mathcal{A}_k)$. This shows that $\beta_i \notin \Phi_{A/K}^0$ for $i = 1, \dots, d-1$. Moreover, any nontrivial linear combination of the images of the divisors $P_i - P_d$ is not zero in $\mathcal{A}'_k/\eta(\mathcal{A}_k)$ (Proposition 2.5), so no nontrivial linear combination of $\beta_1, \dots, \beta_{d-1}$ belongs to $\Phi_{A/K}^0$. \square

Example 6.13. Examples of curves having good reduction after an extension of degree p can be obtained as twists as follows. Choose a smooth proper curve C/k having an automorphism σ_k of order p . Over an appropriate ring \mathcal{O}_K with residue field k , there exists a smooth scheme $\mathcal{Y}^0/\mathcal{O}_K$ with an \mathcal{O}_K -automorphism σ such that C is k -isomorphic to \mathcal{Y}_k^0 and σ restricted to \mathcal{Y}_k^0 induces the given automorphism σ_k . It is shown in [Sekiguchi et al. 1989, §IV, Theorem 2.2] that one can take \mathcal{O}_K to be the Witt ring $W(k)(\zeta_p)$ with ζ_p a primitive p -th root of unity. If one wants a lift in equicharacteristic p , one can trivially take $\mathcal{O}_K = k[[t]]$.

Choose any cyclic (ramified) extension L/K of degree p . The twist of \mathcal{Y}_K^0/K by L/K and σ is a curve X/K that achieves good reduction over L . Starting with an ordinary curve C/k produces a curve X/K having potentially good ordinary reduction over L .

Corollary 6.14. *Fix any odd prime p . For each integer $m > 0$, there exist a regular local ring B of equicharacteristic p endowed with an action of $H := \mathbb{Z}/p\mathbb{Z}$ and a regular local ring B' of mixed characteristic $(0, p)$ endowed with an action of $\mathbb{Z}/p\mathbb{Z}$ such that $\text{Spec } B^H$ and $\text{Spec}(B')^H$ are singular exactly at their closed point, and the graphs associated with a minimal resolution of $\text{Spec } B^H$ and $\text{Spec}(B')^H$ have one node and more than m vertices.*

Proof. As we noted in [Example 6.13](#), there exist a field K of either mixed characteristic $(0, p)$ or of equicharacteristic p and a curve X/K without good reduction over K and with good ordinary reduction over a Galois extension L/K of degree p . Let $H := \text{Gal}(L/K)$. Let $\mathcal{Y}/\mathcal{O}_L$ denote the smooth model of X_L/L . Let $\mathcal{Z}/\mathcal{O}_K$ denote the quotient \mathcal{Y}/H . Let P denote a ramification point of the morphism $\mathcal{Y}_k \rightarrow \mathcal{Y}_k/H$, and let $B := \mathcal{O}_{\mathcal{Y},P}$. [Theorem 6.8](#) shows that the resolution of singularity of $\text{Spec } B^H$ has an intersection matrix of type $N(p, \alpha, r_1)$ for some $\alpha \geq 1$ and $0 < r_1 < p$.

Immediately after the statement of [Theorem 6.8](#) given in the introduction, we briefly alluded to the fact that the integer α is likely to be related to the valuation of the different of L/K . Thus, in principle, by choosing K and L/K appropriately, the above method will produce examples with α as large, as desired. Since at this time we do not know how to prove in general that α is related to the valuation of the different of L/K (except when $p = 2$ and $g = 1$; see [\[Lorenzini 2013a, Theorem 4.1\]](#)), we proceed below with a different argument to prove the existence of resolutions with α as large, as desired.

Consider a quadratic extension K'/K . Since p is odd by hypothesis, the extension K'/K is tame, and one knows how to compute a regular model of $X_{K'}/K'$ from the model $\mathcal{X}/\mathcal{O}_K$ of X/K obtained in [Theorem 6.8](#): simply normalize the base change $\mathcal{X} \times_{\mathcal{O}_K} \mathcal{O}_{K'}$ and resolve its singularities. A singularity on the normalization can only be the preimage of a closed point of \mathcal{X}_k that belongs to two irreducible components of \mathcal{X}'_k and such that both components have odd multiplicity. This singular point is resolved by a single smooth rational curve.

Let $L' := LK'$ with $[L' : K'] = p$. The curve $X_{K'}/K'$ achieves good ordinary reduction over L' . The model $\mathcal{Y}'/\mathcal{O}_{L'} := \mathcal{Y} \times_{\mathcal{O}_L} \mathcal{O}_{L'}$ is smooth, and we let P' denote the preimage of P under the natural map $\mathcal{Y}' \rightarrow \mathcal{Y}$. Let $B' := \mathcal{O}_{\mathcal{Y}',P'}$. We leave it to the reader to check, using [\[Halle 2010, Proposition 4.3\]](#) and the desingularization of the normalization of $\mathcal{X} \times_{\mathcal{O}_K} \mathcal{O}_{K'}$, that the resolution of the singularity of $\text{Spec}(B')^H$ has an intersection matrix of type $N(p, 2\alpha, r'_1)$, where $r'_1 := r_1/2$ if r_1 is even and $r'_1 := (r_1 + p)/2$ if r_1 is odd.

Since we can make an infinite chain of quadratic extensions $K \subset K' \subset K'' \subset \dots$ and since the graph associated with $N(p, \beta, r_1)$ has at least β irreducible components, the corollary is proved. \square

Remark 6.15. Consider an intersection matrix N , and assume that, for some prime p , it satisfies all the conditions listed in [Remark 5.4](#), conditions that would have to be satisfied if this intersection matrix was associated with the resolution of a $\mathbb{Z}/p\mathbb{Z}$ -singularity: its graph $G(N)$ is a tree, $|\det(N)|$ is a power of p , the Smith group Φ_N is killed by p and the fundamental cycle Z has $|Z^2| \leq p$. If $\det(N) = 1$ and $G(N)$ is a tree, then the above conditions are satisfied for every prime at least equal to $|Z^2|$. In particular, when $\det(N) = 1$, the matrix N could potentially be associated with the resolution of a $\mathbb{Z}/p\mathbb{Z}$ -singularity for infinitely many primes p .

An interesting consequence of our guess in [6.2](#) that $\gamma_{Q_i} g_{Q_i} = v(Q_i)$ holds for all $i = 1, \dots, d$ is that a matrix N as above can be associated with the resolution of a $(\mathbb{Z}/p\mathbb{Z})$ -quotient singularity $\mathcal{X}' \rightarrow \mathcal{Z}$ occurring in models of curves as at the beginning of this section *only for finitely many primes* p . Indeed, the choice of a vertex D on N lets us define the integer g_D associated with N and D . If N is the intersection matrix of the resolution of a singularity Q_i of \mathcal{Z} with the marked vertex D linked to C_0 , we noted in [6.2](#) that $p - 1$ must then divide g_D when the equality $\gamma_{Q_i} g_{Q_i} = v(Q_i)$ holds. Since there are only finitely many vertices D , the set of integers g_D is finite, and hence, any prime p larger than the maximum of the integers g_D cannot have the property that $p - 1$ divides some g_D .

Remark 6.16. Let X/K be a curve with potentially good reduction over an extension L/K of degree p as at the beginning of this section. Let Q_i be a singular point of the quotient \mathcal{Z} , and consider the graph G_{Q_i} associated with the resolution of Q_i in $\mathcal{X}' \rightarrow \mathcal{Z}$. One may wonder whether a node of G in G_{Q_i} could have its multiplicity in \mathcal{X}'_k divisible by p^2 . Similar considerations are found in [[Lorenzini 2010](#), Question 1.4].

Acknowledgements

It is my pleasure to thank Qing Liu, Werner Lütkebohmert and Michel Raynaud for helpful comments. I also thank the referee for a careful reading of the article.

References

- [Altman and Kleiman 1970] A. Altman and S. Kleiman, *Introduction to Grothendieck duality theory*, Lecture Notes in Mathematics **146**, Springer, Berlin, 1970. [MR 43 #224](#) [Zbl 0215.37201](#)
- [Artin 1975] M. Artin, “Wildly ramified $\mathbb{Z}/2$ actions in dimension two”, *Proc. Amer. Math. Soc.* **52** (1975), 60–64. [MR 51 #10336](#) [Zbl 0315.14015](#)
- [Bosch and Lorenzini 2002] S. Bosch and D. Lorenzini, “Grothendieck’s pairing on component groups of Jacobians”, *Invent. Math.* **148**:2 (2002), 353–396. [MR 2003b:14053](#) [Zbl 1061.14042](#)

- [Bosch et al. 1990] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **21**, Springer, Berlin, 1990. MR 91i:14034 Zbl 0705.14001
- [Edixhoven et al. 1996] B. Edixhoven, Q. Liu, and D. Lorenzini, “The p -part of the group of components of a Néron model”, *J. Algebraic Geom.* **5**:4 (1996), 801–813. MR 98m:14051 Zbl 0898.14007
- [Gabber et al. 2013] O. Gabber, Q. Liu, and D. Lorenzini, “The index of an algebraic variety”, *Invent. Math.* **192**:3 (2013), 567–626. MR 3049930 Zbl 1268.13009
- [Halle 2010] L. H. Halle, “Stable reduction of curves and tame ramification”, *Math. Z.* **265**:3 (2010), 529–550. MR 2011e:14030 Zbl 1195.14011
- [Hasse 1934] H. Hasse, “Theorie der relativ-zyklischen algebraischen Funktionenkörper, insbesondere bei endlichem Konstantenkörper”, *J. Reine Angew. Math.* **172** (1934), 37–54. Zbl 0010.00501
- [Katsura 1978] T. Katsura, “On Kummer surfaces in characteristic 2”, pp. 525–542 in *Proceedings of the International Symposium on Algebraic Geometry* (Kyoto, 1977), edited by M. Nagata, Kinokuniya Book Store, Tokyo, 1978. MR 82d:14022 Zbl 0411.14010
- [Lipman 1969] J. Lipman, “Rational singularities, with applications to algebraic surfaces and unique factorization”, *Inst. Hautes Études Sci. Publ. Math.* **36** (1969), 195–279. MR 43 #1986 Zbl 0181.48903
- [Liu and Lorenzini 2001] Q. Liu and D. Lorenzini, “Special fibers of Néron models and wild ramification”, *J. Reine Angew. Math.* **532** (2001), 179–222. MR 2001m:11094 Zbl 0970.14025
- [Liu et al. 2004] Q. Liu, D. Lorenzini, and M. Raynaud, “Néron models, Lie algebras, and reduction of curves of genus one”, *Invent. Math.* **157**:3 (2004), 455–518. MR 2005m:14039 Zbl 1060.14037
- [Lorenzini 1989] D. Lorenzini, “Arithmetical graphs”, *Math. Ann.* **285**:3 (1989), 481–501. MR 91b:14026 Zbl 0662.14008
- [Lorenzini 1990a] D. Lorenzini, “Dual graphs of degenerating curves”, *Math. Ann.* **287**:1 (1990), 135–150. MR 91j:14018 Zbl 0668.14019
- [Lorenzini 1990b] D. Lorenzini, “Groups of components of Néron models of Jacobians”, *Compositio Math.* **73**:2 (1990), 145–160. MR 92d:14019 Zbl 0737.14008
- [Lorenzini 1993] D. Lorenzini, “On the group of components of a Néron model”, *J. Reine Angew. Math.* **445** (1993), 109–160. MR 94k:11065 Zbl 0781.14029
- [Lorenzini 2000] D. Lorenzini, “Reduction of points in the group of components of the Néron model of a Jacobian”, *J. Reine Angew. Math.* **527** (2000), 117–150. MR 2001i:11073 Zbl 0983.14022
- [Lorenzini 2010] D. Lorenzini, “Models of curves and wild ramification”, *Pure Appl. Math. Q.* **6**:1 (2010), 41–82. MR 2012a:11068 Zbl 1200.14052
- [Lorenzini 2013a] D. Lorenzini, “Wild quotient singularities of surfaces”, *Math. Z.* **275**:1–2 (2013), 211–232. MR 3101805 Zbl 06224338
- [Lorenzini 2013b] D. Lorenzini, “Wild quotients of products of curves”, preprint, 2013, available at <http://www.math.uga.edu/~lorenz/Paper3.pdf>.
- [Lorenzini and Tucker 2002] D. Lorenzini and T. J. Tucker, “Thue equations and the method of Chabauty–Coleman”, *Invent. Math.* **148**:1 (2002), 47–77. MR 2003d:11088 Zbl 1048.11023
- [Penniston 2000] D. Penniston, “Unipotent groups and curves of genus two”, *Math. Ann.* **317**:1 (2000), 57–78. MR 2001b:14050 Zbl 1005.14010
- [Peskin 1983] B. R. Peskin, “Quotient-singularities and wild p -cyclic actions”, *J. Algebra* **81**:1 (1983), 72–99. MR 84k:13018 Zbl 0517.14017
- [Raynaud 1970] M. Raynaud, “Spécialisation du foncteur de Picard”, *Inst. Hautes Études Sci. Publ. Math.* **38** (1970), 27–76. MR 44 #227 Zbl 0207.51602

- [Sekiguchi et al. 1989] T. Sekiguchi, F. Oort, and N. Suwa, “On the deformation of Artin–Schreier to Kummer”, *Ann. Sci. École Norm. Sup. (4)* **22**:3 (1989), 345–375. MR 91g:14041 Zbl 0714.14024
- [Serre 1959] J.-P. Serre, *Groupes algébriques et corps de classes*, Publications de l’institut de mathématique de l’université de Nancago, VII. Hermann, Paris, 1959. MR 21 #1973 Zbl 0097.35604
- [Singh 1974] B. Singh, “On the group of automorphisms of function field of genus at least two”, *J. Pure Appl. Algebra* **4** (1974), 205–229. MR 50 #13047 Zbl 0284.12007
- [Subrao 1975] D. Subrao, “The p -rank of Artin–Schreier curves”, *Manuscripta Math.* **16**:2 (1975), 169–193. MR 51 #12868 Zbl 0321.14017
- [Viehweg 1977] E. Viehweg, “Invarianten der degenerierten Fasern in lokalen Familien von Kurven”, *J. Reine Angew. Math.* **293/294** (1977), 284–308. MR 58 #16655 Zbl 0349.14017
- [Winters 1974] G. B. Winters, “On the existence of certain families of curves”, *Amer. J. Math.* **96** (1974), 215–228. MR 50 #9874 Zbl 0334.14004

Communicated by Brian Conrad

Received 2013-01-03

Revised 2013-06-06

Accepted 2013-07-16

lorenzin@uga.edu

*Department of Mathematics, University of Georgia,
Athens, GA 30602, United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Karl Rubin	University of California, Irvine, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Roger Heath-Brown	Oxford University, UK	Bernd Sturmfels	University of California, Berkeley, USA
Ehud Hrushovski	Hebrew University, Israel	Richard Taylor	Harvard University, USA
Craig Huneke	University of Virginia, USA	Ravi Vakil	Stanford University, USA
Mikhail Kapranov	Yale University, USA	Michel van den Bergh	Hasselt University, Belgium
Yujiro Kawamata	University of Tokyo, Japan	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Yuri Manin	Northwestern University, USA	Efim Zelmanov	University of California, San Diego, USA
Barry Mazur	Harvard University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2014 is US \$225/year for the electronic version, and \$400/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 8 No. 2 2014

Large self-injective rings and the generating hypothesis	257
LEIGH SHEPPERSON and NEIL STRICKLAND	
On lower ramification subgroups and canonical subgroups	303
SHIN HATTORI	
Wild models of curves	331
DINO LORENZINI	
Geometry of Wachspress surfaces	369
COREY IRVING and HAL SCHENCK	
Groups with exactly one irreducible character of degree divisible by p	397
DANIEL GOLDSTEIN, ROBERT M. GURALNICK, MARK L. LEWIS, ALEXANDER MORETÓ, GABRIEL NAVARRO and PHAM HUU TIEP	
The homotopy category of injectives	429
AMNON NEEMAN	
Essential dimension of spinor and Clifford groups	457
VLADIMIR CHERNOUSOV and ALEXANDER MERKURJEV	
On Deligne's category $\text{Rep}^{ab}(S_d)$	473
JONATHAN COMES and VICTOR OSTRIK	
Algebraicity of the zeta function associated to a matrix over a free group algebra	497
CHRISTIAN KASSEL and CHRISTOPHE REUTENAUER	