

Algebra & Number Theory

Volume 10

2016

No. 4

Nef cones of Hilbert schemes of points on surfaces

Barbara Bolognese, Jack Huizenga, Yinbang Lin, Eric Riedl,
Benjamin Schmidt, Matthew Woolf and Xiaolei Zhao



Nef cones of Hilbert schemes of points on surfaces

Barbara Bolognese, Jack Huizenga, Yinbang Lin, Eric Riedl,
Benjamin Schmidt, Matthew Woolf and Xiaolei Zhao

Let X be a smooth projective surface of irregularity 0. The Hilbert scheme $X^{[n]}$ of n points on X parametrizes zero-dimensional subschemes of X of length n . We discuss general methods for studying the cone of ample divisors on $X^{[n]}$. We then use these techniques to compute the cone of ample divisors on $X^{[n]}$ for several surfaces where the cone was previously unknown. Our examples include families of surfaces of general type and del Pezzo surfaces of degree 1. The methods rely on Bridgeland stability and the positivity lemma of Bayer and Macrì.

A correction was posted on 16 June 2018 in an [online supplement](#).

1. Introduction	907
2. Preliminaries	910
3. Gieseker walls and the nef cone	915
4. Picard rank 1 examples	922
5. Del Pezzo surfaces of degree 1	925
Acknowledgements	928
References	928

1. Introduction

If X is a projective variety, the cone $\text{Amp}(X) \subset N^1(X)$ of ample divisors controls the various projective embeddings of X . It is one of the most important invariants of X , and carries detailed information about the geometry of X . Its closure is the *nef cone* $\text{Nef}(X)$, which is dual to the Mori cone of curves (see for example [Lazarsfeld 2004]). In this paper, we will study the nef cone of the Hilbert scheme of points $X^{[n]}$, where X is a smooth projective surface over \mathbb{C} .

Nef divisors on Hilbert schemes of points on surfaces $X^{[n]}$ are sometimes easy to construct by classical methods. If L is an $(n - 1)$ -very ample line bundle

J. Huizenga was partially supported by an NSF Mathematical Sciences Postdoctoral Research Fellowship. B. Schmidt was partially supported by NSF grant DMS-1523496 (PI Emanuele Macrì) and a Presidential Fellowship of the Ohio State University.

MSC2010: primary 14C05; secondary 14E30, 14J29, 14J60.

Keywords: Hilbert schemes, surfaces, nef cone, ample cone, birational geometry, Bridgeland stability.

on X , then for any $Z \in X^{[n]}$ we have an inclusion $H^0(L \otimes I_Z) \rightarrow H^0(L)$ which defines a morphism from $X^{[n]}$ to the Grassmannian $G(h^0(L) - n, h^0(L))$. The pullback of an ample divisor on the Grassmannian is nef on $X^{[n]}$. It is frequently possible to construct extremal nef divisors by this method. For example, this method completely computes the nef cone of $X^{[n]}$ when X is a del Pezzo surface of degree ≥ 2 or a Hirzebruch surface (see [Arcara et al. 2013; Bertram and Coskun 2013]). Unfortunately, this approach to computing the nef cone is insufficient in general. At the very least, to study nef cones of more interesting surfaces it would be necessary to study an analog of k -very ampleness for higher rank vector bundles, which is considerably more challenging than line bundles.

More recently, many nef cones have been computed by making use of Bridgeland stability conditions and the positivity lemma of Bayer and Macrì (see [Bridgeland 2007; 2008; Arcara and Bertram 2013; Bayer and Macrì 2014b] for background on these topics, which will be reviewed in Section 2). Let $\mathbf{v} = \text{ch}(I_Z) \in K_0(X)$, where $Z \in X^{[n]}$. In the stability manifold $\text{Stab}(X)$ for X there is an open Gieseker chamber \mathcal{C} such that if $\sigma \in \mathcal{C}$ then $M_\sigma(\mathbf{v}) \cong X^{[n]}$, where $M_\sigma(\mathbf{v})$ is the moduli space of σ -semistable objects with invariants \mathbf{v} . The positivity lemma associates to any $\sigma \in \bar{\mathcal{C}}$ a nef divisor on $X^{[n]}$. Stability conditions in the boundary $\partial\mathcal{C}$ frequently give rise to extremal nef divisors. The positivity lemma also classifies the curves orthogonal to a nef divisor constructed in this way, and so gives a tool for checking extremality.

The stability manifold is rather large in general, so computation of the full Gieseker chamber can be unwieldy. We deal with this problem by focusing on a small *slice* of the stability manifold parametrized by a half-plane. Up to scale, the corresponding divisors in $N^1(X^{[n]})$ form an affine ray. The nef cone $\text{Nef}(X^{[n]})$ is spanned by a codimension-1 subcone identified with $\text{Nef}(X)$ and other more interesting classes which are positive on curves contracted by the Hilbert–Chow morphism. Since $\text{Nef}(X^{[n]})$ is convex, we can study $\text{Nef}(X^{[n]})$ by looking at positivity properties of divisors along rays in $N^1(X^{[n]})$ starting from a class in $\text{Amp}(X) \subset \text{Nef}(X^{[n]})$. The positivity lemma gives us an effective criterion for testing when divisors along the ray are nef.

The slices of the stability manifold that we consider are given by a pair of divisors (H, D) on X with H ample and $-D$ effective. The following is a weak version of one of our main theorems.

Theorem 1.1. *Let X be a smooth projective surface. If $n \gg 0$, then there is an extremal nef divisor on $X^{[n]}$ coming from the (H, D) -slice. It can be explicitly computed if both the intersection pairing on $\text{Pic}(X)$ and the set of effective classes in $\text{Pic}(X)$ are known. An orthogonal curve class is given by n points moving in a g_n^1 on a curve of a particular class.*

See [Section 3](#), especially [Corollary 3.7](#) and [Theorem 3.11](#), for more explicit statements. Stronger statements can also be shown under strong assumptions on $\text{Pic}(X)$; for example, we study the Picard rank 1 case in detail in [Section 4](#). Recall that if X is a surface of irregularity $q := H^1(\mathcal{O}_X) = 0$, then $N^1(X^{[n]})$ is spanned by the divisor B of nonreduced schemes and divisors $L^{[n]}$ induced by divisors $L \in \text{Pic}(X)$; see [Section 2A](#) for details.

Theorem 1.2. *Let X be a smooth projective surface with $\text{Pic } X \cong \mathbb{Z}H$, where H is an ample divisor. Let $a > 0$ be the smallest integer such that aH is effective. If*

$$n \geq \max\{a^2 H^2, p_a(aH) + 1\},$$

then $\text{Nef}(X^{[n]})$ is spanned by the divisor $H^{[n]}$ and the divisor

$$\frac{1}{2}K_X^{[n]} + \left(\frac{a}{2} + \frac{n}{aH^2}\right)H^{[n]} - \frac{1}{2}B. \quad (*)$$

An orthogonal curve class is given by letting n points move in a g_n^1 on a curve in X of class aH .

Note that in the Picard rank 1 case the divisor class $(*)$ is frequently of the form $\lambda H^{[n]} - \frac{1}{2}B$ for a noninteger number $\lambda \in \mathbb{Q}$. Any divisor constructed from an $(n-1)$ -very ample line bundle will be of the form $\lambda H^{[n]} - \frac{1}{2}B$ with $\lambda \in \mathbb{Z}$, so in general the edge of the nef cone cannot be obtained from line bundles in this way.

The required lower bound on n in [Theorem 1.2](#) can be improved in specific examples where special linear series on hyperplane sections are better understood.

Theorem 1.3. *Let X be one of the following surfaces:*

- (1) *a very general hypersurface in \mathbb{P}^3 of degree $d \geq 4$, or*
- (2) *a very general degree- d cyclic branched cover of \mathbb{P}^2 of general type.*

In either case, $\text{Pic}(X) \cong \mathbb{Z}H$ with H effective. Suppose $n \geq d-1$ in the first case, and $n \geq d$ in the second case. Then $\text{Nef}(X^{[n]})$ is spanned by $H^{[n]}$ and the divisor class $()$ with $a = 1$.*

Finally, in [Section 5](#) we compute the nef cone of $X^{[n]}$ where X is a smooth del Pezzo surface of degree 1 and $n \geq 2$ is arbitrary. This computation was an open problem posed by Bertram and Coskun [2013]; they noted that the method of k -very ample line bundles would not be sufficient to prove the expected answer. Since X has Picard rank 9, this computation makes full use of the general methods developed in [Section 3](#). If $C \subset X$ is a reduced, irreducible curve which admits a g_n^1 , we write $C_{[n]}$ for the curve in the Hilbert scheme $X^{[n]}$ given by letting n points move in a g_n^1 on C .

Theorem 1.4. *Let X be a smooth del Pezzo surface of degree 1. The Mori cone of curves $\text{NE}(X^{[n]})$ is spanned by the 240 classes $E_{[n]}$ given by (-1) -curves $E \subset X$,*

the class of a curve contracted by the Hilbert–Chow morphism, and the class $F_{[n]}$, where $F \in |-K_X|$ is an anticanonical curve. The nef cone is determined by duality.

Many previous authors have used Bridgeland stability conditions to study nef cones and wall-crossing for Hilbert schemes $X^{[n]}$ and moduli spaces of sheaves $M_H(\mathbf{v})$ for various classes of surfaces. For instance, the program was studied for \mathbb{P}^2 in [Arcara et al. 2013; Coskun and Huizenga 2016; Bertram et al. 2014; Li and Zhao 2013], Hirzebruch and del Pezzo surfaces in [Bertram and Coskun 2013], abelian surfaces in [Yanagida and Yoshioka 2014; Maciocia and Meachan 2013], K3 surfaces in [Bayer and Macrì 2014b; 2014a; Hassett and Tschinkel 2010], and Enriques surfaces in [Nuer 2014]. Our results unify several of these approaches. Additionally, nef cones were classically studied in the context of k -very ample line bundles in papers such as [Ellingsrud et al. 2001; Beltrametti and Sommese 1991; Beltrametti et al. 1989; Catanese and Göttsche 1990].

2. Preliminaries

Throughout the paper, we let X be a smooth projective surface over \mathbb{C} .

2A. Divisors and curves on $X^{[n]}$. For simplicity we assume that X has irregularity $q = h^1(\mathcal{O}_X) = 0$ in this subsection. By work of Fogarty [1968], the Hilbert scheme $X^{[n]}$ is a smooth projective variety of dimension $2n$ which resolves the singularities in the symmetric product $X^{(n)}$ via the Hilbert–Chow morphism $X^{[n]} \rightarrow X^{(n)}$. A line bundle L on X induces the S_n -equivariant line bundle $L^{\boxtimes n}$ on X^n which descends to a line bundle $L^{(n)}$ on the symmetric product $X^{(n)}$. The pullback of $L^{(n)}$ by the Hilbert–Chow morphism $X^{[n]} \rightarrow X^{(n)}$ defines a line bundle on $X^{[n]}$ which we will denote by $L^{[n]}$. Intuitively, if $L \cong \mathcal{O}_X(D)$ for a reduced effective divisor $D \subset X$, then $L^{[n]}$ can be represented by the divisor $D^{[n]}$ of schemes $Z \subset X$ which meet D .

Fogarty shows that

$$\text{Pic}(X^{[n]}) \cong \text{Pic}(X) \oplus \mathbb{Z}(B/2),$$

where $\text{Pic}(X) \subset \text{Pic}(X^{[n]})$ is embedded by $L \mapsto L^{[n]}$ and B is the locus of non-reduced schemes, i.e., the exceptional divisor of the Hilbert–Chow morphism [Fogarty 1973]. Tensoring by the real numbers, the Néron–Severi space $N^1(X^{[n]})$ is therefore spanned by $N^1(X)$ and B .

There are also curve classes in $X^{[n]}$ induced by curves in X . Two different constructions are immediate. Let $C \subset X$ be a reduced and irreducible curve.

- (1) There is a curve $\tilde{C}_{[n]}$ in $X^{[n]}$ given by fixing $n - 1$ general points of X and letting an n -th point move along C .
- (2) If C admits a g_n^1 , i.e., a degree- n map to \mathbb{P}^1 , then the fibers of $C \rightarrow \mathbb{P}^1$ give a rational curve $\mathbb{P}^1 \rightarrow X^{[n]}$. We write $C_{[n]}$ for this class.

These constructions preserve intersection numbers, in the sense that if $D \subset X$ is a divisor and $C \subset X$ is a curve then

$$D^{[n]} \cdot \tilde{C}_{[n]} = D^{[n]} \cdot C_{[n]} = D \cdot C.$$

Part of the nef cone $\text{Nef}(X^{[n]})$ is easily described in terms of the nef cone of X . If D is an ample divisor, then $D^{(n)}$ is ample so $D^{[n]}$ is nef. In the limit, we find that if D is nef then $D^{[n]}$ is nef. Conversely, if D is not nef then there is an irreducible curve C with $D \cdot C < 0$, so $D^{[n]} \cdot \tilde{C}_{[n]} < 0$ and $D^{[n]}$ is not nef. Under the Fogarty isomorphism,

$$\text{Nef}(X^{[n]}) \cap N^1(X) = \text{Nef}(X).$$

The hyperplane $N^1(X) \subset N^1(X^{[n]})$ is orthogonal to any curve contracted by the Hilbert–Chow morphism, so all the divisors in $\text{Nef}(X) \subset \text{Nef}(X^{[n]})$ are extremal. Since B is the exceptional locus of the Hilbert–Chow morphism, we see that any nef class must have nonpositive coefficient of B . After scaling, then, we see that computation of the cone $\text{Nef}(X^{[n]})$ reduces to describing the nef classes of the form $L^{[n]} - \frac{1}{2}B$ lying outside $\text{Nef}(X) \subset \text{Nef}(X^{[n]})$.

2B. Bridgeland stability conditions. We now recall some basic definitions and properties of Bridgeland stability conditions. We fix a polarization $H \in \text{Pic}(X)_{\mathbb{R}}$. For any divisor $D \in \text{Pic}(X)_{\mathbb{R}}$, the twisted Chern character $\text{ch}^D = e^{-D} \text{ch}$ can be expanded as

$$\text{ch}_0^D = \text{ch}_0, \quad \text{ch}_1^D = \text{ch}_1 - D \text{ch}_0, \quad \text{ch}_2^D = \text{ch}_2 - D \cdot \text{ch}_1 + \frac{1}{2} D^2 \text{ch}_0.$$

Recall that a Bridgeland stability condition is a pair $\sigma = (Z, \mathcal{A})$ for which $Z : K_0(X) \rightarrow \mathbb{C}$ is an additive homomorphism and $\mathcal{A} \subset D^b(X)$ is the heart of a bounded t-structure. In particular, \mathcal{A} is an abelian category. Moreover, Z maps any nontrivial object in \mathcal{A} to the upper half-plane or the negative real line. The σ -slope function is defined by

$$\nu_{\sigma} = -\frac{\Re Z}{\Im Z},$$

and σ -(semi)stability of objects of \mathcal{A} is defined in terms of this slope function. More technical requirements are the existence of Harder–Narasimhan filtrations and the support property. We recommend [Bridgeland 2007] for a more precise definition. The support property is well explained in Appendix A of [Bayer et al. 2014].

In the case of surfaces, Bridgeland [2008] and Arcara and Bertram [2013] showed how to construct Bridgeland stability conditions in a slice corresponding to a choice of an ample divisor $H \in \text{Pic}(X)_{\mathbb{R}}$ and arbitrary twisting divisor $D \in \text{Pic}(X)_{\mathbb{R}}$. The

classical Mumford slope function for twisted Chern characters is defined by

$$\mu_{H,D} = \frac{H \cdot \text{ch}_1^D}{H^2 \text{ch}_0^D},$$

where torsion sheaves are interpreted as having positive infinite slope. Given a real number $\beta \in \mathbb{R}$, there are two categories defined as

$$\begin{aligned} \mathcal{T}_\beta &= \{E \in \text{Coh}(X) : \text{any quotient } E \twoheadrightarrow G \text{ satisfies } \mu_{H,D}(G) > \beta\}, \\ \mathcal{F}_\beta &= \{E \in \text{Coh}(X) : \text{any subsheaf } F \hookrightarrow E \text{ satisfies } \mu_{H,D}(F) \leq \beta\}. \end{aligned}$$

A new heart of a bounded t-structure is defined as the extension closure $\mathcal{A}_\beta := \langle \mathcal{F}_\beta[1], \mathcal{T}_\beta \rangle$. We fix an additional positive real number α and define the homomorphism as

$$Z_{\beta,\alpha} = -\text{ch}_2^{D+\beta H} + \frac{1}{2}\alpha^2 H^2 \text{ch}_0^{D+\beta H} + iH \cdot \text{ch}_1^{D+\beta H}.$$

The pair $\sigma_{\beta,\alpha} := (Z_{\beta,\alpha}, \mathcal{A}_\beta)$ is then a Bridgeland stability condition. The (H, D) -slice of stability conditions is the family of stability conditions

$$\{\sigma_{\beta,\alpha} : \beta, \alpha \in \mathbb{R}, \alpha > 0\}$$

parametrized by the (β, α) upper half-plane.

Definition 2.1. Fix a set of invariants $\mathbf{v} \in K_0(X)$.

- (1) Let $\mathbf{w} \in K_0(X)$ be a vector such that \mathbf{v} and \mathbf{w} do not have the same $\sigma_{\beta,\alpha}$ -slope everywhere in the (H, D) -slice. The *numerical wall* for \mathbf{v} given by \mathbf{w} is the set of points (β, α) where \mathbf{v} and \mathbf{w} have the same $\sigma_{\beta,\alpha}$ -slope.
- (2) A numerical wall for \mathbf{v} given by a vector \mathbf{w} as above is a *wall* (or *actual wall*) if there is a point (β, α) on the wall and an exact sequence $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ in \mathcal{A}_β , where $\text{ch } F = \mathbf{w}$, $\text{ch } E = \mathbf{v}$, and F, E , and G are $\sigma_{\beta,\alpha}$ -semistable objects (of the same $\sigma_{\beta,\alpha}$ -slope).

We write $K_{\text{num}}(X)$ for the numerical Grothendieck group of classes in $K_0(X)$ modulo numerical equivalence. Note that numerical walls for $\mathbf{v} \in K_0(X)$ only depend on the numerical class of \mathbf{v} , while actual walls a priori depend on $c_1(\mathbf{v}) \in \text{Pic}(X)$. The structure of walls in a slice is heavily restricted by Bertram’s nested wall theorem. This was first observed for Picard rank 1 with $D = 0$, but the proof immediately generalizes by replacing ch by ch^D everywhere.

Theorem 2.2 [Maciocia 2014]. *Let $\mathbf{v} \in K_0(X)$.*

- (1) *Numerical walls for \mathbf{v} can either be semicircles with center on the β -axis or the unique vertical line given by $\beta = \mu_{H,D}(\mathbf{v})$. Moreover, the apex of each semicircle lies on the hyperbola $\Re Z_{\beta,\alpha}(\mathbf{v}) = 0$.*

- (2) Numerical walls for \mathbf{v} are disjoint, and the semicircular walls on either side of the vertical wall are nested.
- (3) If W_1 and W_2 are two semicircular numerical walls left of the vertical wall with centers $(s_{W_1}, 0)$ and $(s_{W_2}, 0)$, then W_2 is nested inside W_1 if and only if $s_{W_1} < s_{W_2}$.
- (4) Suppose $0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$ is an exact sequence destabilizing an object E with $\text{ch}(E) = \mathbf{v}$ at a point (β, α) on a numerical wall W , in the sense that all three objects have the same $\sigma_{\beta, \alpha}$ -slope and this is an exact sequence in \mathcal{A}_β . Then it is an exact sequence of objects in $\mathcal{A}_{\beta'}$ with the same $\sigma_{\beta', \alpha'}$ -slope for all $(\beta', \alpha') \in W$. That is, E is destabilized along the entire wall.

2C. Slope and discriminant. The explicit geometry of walls is frequently best understood in terms of slopes and discriminants; the formulas presented here previously appeared in [Coskun and Huizenga 2014] in the context of \mathbb{P}^2 . When the rank is nonzero, we define

$$\Delta_{H,D} = \frac{1}{2}\mu_{H,D}^2 - \frac{\text{ch}_2^D}{H^2 \text{ch}_0^D}.$$

The Bogomolov inequality gives $\Delta_{H,D}(E) \geq 0$ whenever E is an (H, D) -twisted Gieseker semistable sheaf. Observe that $\Delta_{H,D+\beta H} = \Delta_{H,D}$ for every $\beta \in \mathbb{R}$. A straightforward calculation shows that for vectors of nonzero rank the slope function for the stability condition $\sigma_{\beta, \alpha}$ in the (H, D) -slice is given by

$$v_{\sigma_{\beta, \alpha}} = \frac{(\mu_{H,D} - \beta)^2 - \alpha^2 - 2\Delta_{H,D}}{(\mu_{H,D} - \beta)}. \quad (1)$$

Suppose \mathbf{v}, \mathbf{w} are two classes with positive rank, and let their slopes and discriminants be $\mu_{H,D}, \Delta_{H,D}$ and $\mu'_{H,D}, \Delta'_{H,D}$, respectively. The numerical wall W in the (H, D) -slice where \mathbf{v} and \mathbf{w} have the same slope is computed as follows.

- If $\mu_{H,D} = \mu'_{H,D}$ and $\Delta_{H,D} = \Delta'_{H,D}$, then \mathbf{v} and \mathbf{w} have the same slope everywhere in the slice, so there is no numerical wall.
- If $\mu_{H,D} = \mu'_{H,D}$ and $\Delta_{H,D} \neq \Delta'_{H,D}$, then W is the vertical wall $\beta = \mu_{H,D}$.
- If $\mu_{H,D} \neq \mu'_{H,D}$, then (1) implies W is the semicircle with center $(s_W, 0)$ and radius ρ_W , where

$$s_W = \frac{1}{2}(\mu_{H,D} + \mu'_{H,D}) - \frac{\Delta_{H,D} - \Delta'_{H,D}}{\mu_{H,D} - \mu'_{H,D}}, \quad (2)$$

$$\rho_W^2 = (s_W - \mu_{H,D})^2 - 2\Delta_{H,D}, \quad (3)$$

provided that the expression defining ρ_W^2 is positive; if it is negative then the wall is empty.

Notice that if $\Delta_{H,D}(\mathbf{v}) \geq 0$ then numerical walls for \mathbf{v} left of the vertical wall accumulate at the point

$$(\mu_{H,D}(\mathbf{v}) - \sqrt{2\Delta_{H,D}(\mathbf{v})}, 0) \tag{4}$$

as their radii go to 0.

2D. Nef divisors and the positivity lemma. In this section, we describe the positivity lemma of Bayer and Macrì. Let $\sigma = (Z, \mathcal{A})$ be a stability condition on X , $\mathbf{v} \in K_{\text{num}}(X)$ and S a proper algebraic space of finite type over \mathbb{C} . Let $\mathcal{E} \in D^b(X \times S)$ be a flat family of σ -semistable objects of class \mathbf{v} , i.e., for every \mathbb{C} -point $p \in S$, the derived restriction $\mathcal{E}|_{\pi_S^{-1}(\{p\})}$ is σ -semistable of class \mathbf{v} . Then Bayer and Macrì define a numerical divisor class $D_{\sigma,\mathcal{E}} \in N^1(S)$ on the space S by assigning its intersection with any projective integral curve $C \subset S$:

$$D_{\sigma,\mathcal{E}} \cdot C = \mathfrak{S} \left(- \frac{Z((pX)_* \mathcal{E}|_{C \times X})}{Z(\mathbf{v})} \right).$$

The positivity lemma shows that this divisor inherits positivity properties from the homomorphism Z , and classifies the curve classes orthogonal to the divisor. Recall that two σ -semistable objects are *S-equivalent* with respect to σ if their sets of Jordan–Hölder factors are the same.

Theorem 2.3 (positivity lemma [Bayer and Macrì 2014b, Lemma 3.3]). *The divisor $D_{\sigma,\mathcal{E}} \in N^1(S)$ is nef. Moreover, if $C \subset S$ is a projective integral curve then $D_{\sigma,\mathcal{E}} \cdot C = 0$ if and only if two general objects parametrized by C are S-equivalent with respect to σ .*

Our primary use of the positivity lemma is to attempt to construct extremal nef divisors on Hilbert schemes of points. Thus it is important to recover Hilbert schemes of points as Bridgeland moduli spaces. Recall that a torsion-free coherent sheaf E is (H, D) -twisted Gieseker semistable if for every $F \subset E$ we have

$$\frac{\chi(F \otimes \mathcal{O}_X(mH - D))}{\text{rk}(F)} \leq \frac{\chi(E \otimes \mathcal{O}_X(mH - D))}{\text{rk}(E)}$$

for all $m \gg 0$, where the Euler characteristic is computed formally via Riemann–Roch; see [Matsuki and Wentworth 1997]. For any class $\mathbf{v} \in K_0(X)$, there are projective moduli spaces $M_{H,D}(\mathbf{v})$ of S -equivalence classes of (H, D) -twisted Gieseker semistable sheaves with class \mathbf{v} . If $\mathbf{v} = (1, 0, -n)$ is the Chern character of an ideal sheaf of n points then $M_{H,D}(\mathbf{v}) = X^{[n]}$. Note that if the irregularity of X is nonzero, then it is crucial to fix the determinant.

Fix an (H, D) -slice in the stability manifold, and fix a vector $\mathbf{v} \in K_0(X)$ with positive rank. If β lies to the left of the vertical wall $\beta = \mu_{H,D}(\mathbf{v})$ for \mathbf{v} , then for $\alpha \gg 0$ the moduli space coincides with a twisted Gieseker moduli space.

Proposition 2.4 (the large volume limit [Bridgeland 2008; Maciocia 2014]). *Fix divisors (H, D) giving a slice in $\text{Stab}(X)$. Let $\mathbf{v} \in K_0(X)$ be a vector with positive rank, and let $\beta \in \mathbb{R}$ be such that $\mu_{H,D}(\mathbf{v}) > \beta$. If $E \in \mathcal{A}_\beta$ has $\text{ch}(E) = \mathbf{v}$ then E is $\sigma_{\beta,\alpha}$ -semistable for all $\alpha \gg 0$ if and only if E is an $(H, D - \frac{1}{2}K_X)$ -twisted Gieseker semistable sheaf.*

Moreover, in the quadrant of the (H, D) -slice left of the vertical wall there is a largest semicircular wall for \mathbf{v} , called the Gieseker wall. For all (β, α) between this wall and the vertical wall, the moduli space $M_{\sigma_{\beta,\alpha}}(\mathbf{v})$ coincides with the moduli space $M_{H,D-K_X/2}(\mathbf{v})$ of $(H, D - \frac{1}{2}K_X)$ -twisted Gieseker semistable sheaves.

We use these results as follows. Let $\mathbf{v} = (1, 0, -n) \in K_0(X)$ be the vector for the Hilbert scheme $X^{[n]}$, and let σ_+ be a stability condition in the (H, D) -slice lying above the Gieseker wall, so that $M_{\sigma_+}(\mathbf{v}) \cong X^{[n]}$. Let $\mathcal{E}/(X \times X^{[n]})$ be the universal ideal sheaf, and let σ_0 be a stability condition on the Gieseker wall. By the definition of the Gieseker wall, \mathcal{E} is a family of σ_0 -semistable objects, so there is an induced nef divisor $D_{\sigma_0,\mathcal{E}}$ on $X^{[n]}$. Furthermore, curves orthogonal to $D_{\sigma_0,\mathcal{E}}$ are understood in terms of destabilizing sequences along the wall, so it is possible to test for extremality.

3. Gieseker walls and the nef cone

Fix an ample divisor $H \in \text{Pic}(X)$ with $H^2 = d$ and an *antieffective* divisor D . In this section we study the nef divisor arising from the Gieseker wall (i.e., the largest wall where some ideal sheaf is destabilized) in the slice of the stability manifold given by the pair (H, D) . We first compute the Gieseker wall, and then investigate when the corresponding nef divisor is in fact extremal.

3A. Bounding higher rank walls. The main difficulty in computing extremal rays of the nef cone is to show that a destabilizing subobject along the Gieseker wall is a line bundle, and not some higher rank sheaf. We first prove a lemma which generalizes Proposition 8.3 of [Coskun and Huizenga 2016] from $X = \mathbb{P}^2$ to an arbitrary surface. We prove the result in slightly more generality than we will need here as we expect it to be useful in future work.

Lemma 3.1. *Let σ_0 be a stability condition in the (H, D) -slice, and suppose*

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0$$

is an exact sequence of σ_0 -semistable objects of the same σ_0 -slope, where E is an (H, D) -twisted Gieseker semistable torsion-free sheaf. If the map $F \rightarrow E$ of sheaves is not injective, then the radius ρ_W of the wall W defined by this sequence satisfies

$$\rho_W^2 \leq \frac{(\min\{\text{rk}(F) - 1, \text{rk}(E)\})^2}{2 \text{rk}(F)} \Delta_{H,D}(E).$$

Proof. The proof is similar to the proof in [Coskun and Huizenga 2016] given in the case of \mathbb{P}^2 ; we present it for completeness. The object F is a torsion-free sheaf by the standard cohomology sequence and the fact that the heart of the t-structure in the slice we are working in consists of objects which only have nonzero cohomology sheaves in degrees 0 and -1 . The exact sequence along W gives an exact sequence

$$0 \rightarrow K \rightarrow F \rightarrow E \rightarrow C \rightarrow 0$$

of sheaves of ranks k, f, e, c , respectively. By assumption, $k, f, e > 0$. Let $(s_W, 0)$ be the center of W . As F is in the category \mathcal{T}_β whenever (β, α) is on W , we find that $\mu_{H,D}(F) \geq s_W + \rho_W$, so

$$\begin{aligned} df(s_W + \rho_W) &\leq df\mu_{H,D}(F) = \text{ch}_1^D(F) \cdot H = (\text{ch}_1^D(K) + \text{ch}_1^D(E) - \text{ch}_1^D(C)) \cdot H \\ &= dk\mu_{H,D}(K) + de\mu_{H,D}(E) - \text{ch}_1^D(C) \cdot H. \end{aligned}$$

Similarly, $K \in \mathcal{F}_\beta$ along W , so $\mu_{H,D}(K) \leq s_W - \rho_W$ and

$$df(s_W + \rho_W) \leq dk(s_W - \rho_W) + de\mu_{H,D}(E) - \text{ch}_1^D(C) \cdot H,$$

which gives

$$d(k + f)\rho_W \leq d(k - f)s_W + de\mu_{H,D}(E) - \text{ch}_1^D(C) \cdot H. \tag{5}$$

We now wish to eliminate the term $\text{ch}_1^D(C) \cdot H$ in inequality (5). If C is either 0 or torsion, then $\text{ch}_1^D(C) \cdot H \geq 0$ and $-e = k - f$, and we deduce

$$(k + f)\rho_W \leq (k - f)(s_W - \mu_{H,D}(E)). \tag{6}$$

Suppose instead that C is not torsion. Since C is a quotient of the semistable sheaf E , we have $\mu_{H,D}(C) \geq \mu_{H,D}(E)$, so $\text{ch}_1^D(C) \cdot H = dc\mu_{H,D}(C) \geq dc\mu_{H,D}(E)$. As $k - f = c - e$, we find that inequality (6) also holds in this case.

Both sides of inequality (6) are positive, so squaring both sides gives

$$(k + f)^2 \rho_W^2 \leq (k - f)^2 (s_W - \mu_{H,D}(E))^2.$$

Formula (3) for ρ_W^2 shows this is equivalent to

$$(k + f)^2 \rho_W^2 \leq (k - f)^2 (\rho_W^2 + 2\Delta_{H,D}(E)),$$

from which we obtain

$$\rho_W^2 \leq \frac{(k - f)^2}{2kf} \Delta_{H,D}(E).$$

Since $k = f - e + c$, we see that $k \geq \max\{1, f - e\}$. By taking derivatives in k , we see that $(k - f)^2 / (2kf)$ is decreasing for $k + f > 0$, and so the maximum possible value of the right-hand side must occur when $k = \max\{1, f - e\}$. The denominator

will be at least $2f$ in this case, and the numerator is $\min\{(f-1)^2, e^2\}$. The result follows. \square

For our present work we will only need the next consequence of [Lemma 3.1](#), which follows immediately from computing $\Delta_{H,D}(I_Z)$.

Corollary 3.2. *With the hypotheses of [Lemma 3.1](#), if E is an ideal sheaf $I_Z \in X^{[n]}$ and F has rank at least 2, then the radius of the corresponding wall satisfies*

$$\rho_W^2 \leq \frac{2nd + (H \cdot D)^2 - dD^2}{8d^2} := \varrho_{H,D,n}.$$

The number $\varrho_{H,D,n}$ therefore bounds the squares of the radii of higher rank walls for $X^{[n]}$.

3B. Rank-1 walls and critical divisors. In the cases where we compute the Gieseker wall, the ideal sheaf that is destabilized along the wall will be destabilized by a rank-1 subobject. We first compute the numerical walls given by rank-1 subobjects.

Lemma 3.3. *Consider a rank-1 torsion-free sheaf $F = I_{Z'}(-L)$, where Z' is a zero-dimensional scheme of length w and L is an effective divisor. In the (H, D) -slice, the numerical wall W for $X^{[n]}$ where F has the same slope as an ideal I_Z of n points has center $(s_W, 0)$ given by*

$$s_W = -\frac{2(n-w) + L^2 + 2(D \cdot L)}{2(H \cdot L)}.$$

Proof. This is an immediate consequence of (2) for the center of a wall. \square

Recalling that walls for $X^{[n]}$ left of the vertical wall get larger as their centers decrease, we deduce the following consequence.

Lemma 3.4. *If the Gieseker wall in the (H, D) -slice is given by a rank-1 subobject, then it is a line bundle $\mathcal{O}_X(-L)$ for some effective divisor L .*

Proof. Suppose some $I_Z \in X^{[n]}$ is destabilized along the Gieseker wall W by a sheaf of the form $I_{Z'}(-L)$, where Z' is a nonempty zero-dimensional scheme and L is effective. By [Lemma 3.3](#), the numerical wall W' given by $\mathcal{O}_X(-L)$ is strictly larger than W . Since $\mathcal{O}_X(-L)$ has the same $\mu_{H,D}$ -slope as $I_{Z'}(-L)$, which is in the categories along W , we find that $\mathcal{O}_X(-L)$ is in at least some of the categories along W' . But then W' is an actual wall, since any ideal sheaf I_Z where Z lies on a curve $C \in |L|$ is destabilized along it. This contradicts the fact that W is the Gieseker wall. \square

Less trivially, there is a further minimality condition automatically satisfied by a line bundle $\mathcal{O}_X(-L)$ which gives the Gieseker wall. We define the set of *critical effective divisors* with respect to H and D by

$$\text{CrDiv}(H, D) = \{-D\} \cup \{L \in \text{Pic}(X) \text{ effective} : H \cdot L < H \cdot (-D)\}.$$

By [Hartshorne 1977, Exercise V.1.11], the set $\text{CrDiv}(H, D)/\sim$ of critical divisors modulo numerical equivalence is finite. Therefore, the set of numerical walls for $X^{[n]}$ given by line bundles $\mathcal{O}_X(-L)$ with $L \in \text{CrDiv}(H, D)$ is also finite. Note that the inequality $H \cdot L < H \cdot (-D)$ is equivalent to the inequality $\mu_{H,D}(\mathcal{O}_X(-L)) > 0$. The next proposition demonstrates the importance of critical divisors.

Proposition 3.5. *Assume $2n > D^2$, and suppose the subobject giving the Gieseker wall for $X^{[n]}$ in the (H, D) -slice is a line bundle. Then the Gieseker wall is computed by $\mathcal{O}_X(-L)$, where $L \in \text{CrDiv}(H, D)$ is chosen so that the numerical wall given by $\mathcal{O}_X(-L)$ is as large as possible.*

Proof. First, consider the numerical wall W given by $\mathcal{O}_X(D)$. By Lemma 3.3, the center $(s_W, 0)$ has

$$s_W = \frac{2n - D^2}{2(H \cdot D)} < 0, \tag{7}$$

since $2n > D^2$ and D is antieffective. Since $\mu_{H,D}(\mathcal{O}_X(D)) = \Delta_{H,D}(\mathcal{O}_X(D)) = 0$, formula (3) for the radius of W gives $\rho_W^2 = s_W^2$. In particular, W is nonempty, and $\mathcal{O}_X(D)$ lies in at least some of the categories along W . Since D is antieffective, there are exact sequences of the form

$$0 \rightarrow \mathcal{O}_X(D) \rightarrow I_Z \rightarrow I_{Z \subset C} \rightarrow 0,$$

where $C \in |-D|$ and $Z \subset C$ is a collection of n points. If no actual wall is larger than W , it follows that W is an actual wall and it is the Gieseker wall.

Suppose the Gieseker wall is larger than W and computed by a line bundle $\mathcal{O}_X(-L)$ with L effective. Since W passes through the origin in the (β, α) -plane, $\mathcal{O}_X(-L)$ must lie in the category \mathcal{T}_0 . Therefore $\mu_{H,D}(\mathcal{O}_X(-L)) > 0$, and $L \in \text{CrDiv}(H, D)$.

Conversely, suppose $L \in \text{CrDiv}(H, D)$ is chosen to maximize the wall W' given by $\mathcal{O}_X(-L)$. Then no actual wall is larger than W' . Since $s_W < 0$ and $\mu_{H,D}(\mathcal{O}_X(-L)) \geq 0$, we find that $\mathcal{O}_X(-L)$ is in at least some of the categories along W , and hence in at least some of the categories along W' . We conclude that W' is an actual wall, and therefore that it is the Gieseker wall. \square

Combining Corollary 3.2 and Proposition 3.5 gives our primary tool to compute the Gieseker wall.

Theorem 3.6. *Assume $2n > D^2$, and let $L \in \text{CrDiv}(H, D)$ be a critical divisor such that the wall for $X^{[n]}$ given by $\mathcal{O}_X(-L)$ is as large as possible. If this wall has radius ρ satisfying $\rho^2 \geq \varrho_{H,D,n}$, then it is the Gieseker wall.*

Conversely, if the Gieseker wall has radius satisfying $\rho^2 \geq \varrho_{H,D,n}$ then it is obtained in this way.

While the theorem is our sharpest result, it is useful to lose some generality to get a more explicit version. Since $-D \in \text{CrDiv}(H, D)$, if the wall given by $\mathcal{O}_X(D)$ satisfies $\rho^2 \geq \varrho_{H,D,n}$ then the Gieseker wall is computed by [Theorem 3.6](#). This allows us to compute the Gieseker wall so long as n is large enough, depending only on the intersection numbers of H and D .

Corollary 3.7. *Let*

$$\eta_{H,D} := \frac{(H \cdot D)^2 + dD^2}{2d}.$$

If $n \geq \eta_{H,D}$ then the Gieseker wall is the largest wall given by a critical divisor.

Furthermore, if $n > \eta_{H,D}$ then every I_Z destabilized along the Gieseker wall fits into an exact sequence

$$0 \rightarrow \mathcal{O}_X(-C) \rightarrow I_Z \rightarrow I_{Z \subset C} \rightarrow 0$$

for some curve $C \in |L|$, where L is a critical divisor computing the Gieseker wall. If the critical divisor computing the Gieseker wall is unique, then $\mathcal{O}_X(-C)$ and $I_{Z \subset C}$ are the Jordan–Hölder factors of any I_Z destabilized along the Gieseker wall.

Proof. Observe that the inequality $n \geq \eta_{H,D}$ automatically implies the inequality $2n > D^2$ needed to apply [Theorem 3.6](#).

Let W be the wall for $X^{[n]}$ in the (H, D) -slice corresponding to $\mathcal{O}_X(D)$. The center $(s_W, 0)$ of W was computed in [\(7\)](#), and $\rho_W^2 = s_W^2$. We find that $\rho_W^2 \geq \varrho_{H,D,n}$ holds when $n \geq \eta_{H,D}$, with strict inequality when $n > \eta_{H,D}$.

When $n > \eta_{H,D}$ there can be no higher-rank destabilizing subobject of an I_Z destabilized along the Gieseker wall, so there is an exact sequence as claimed. Furthermore, if there is only one critical divisor computing the wall, then there is a unique destabilizing subobject along the wall, so the Jordan–Hölder filtration has length two. □

3C. Classes of divisors. In this subsection we give an elementary computation of the class of the divisor corresponding to a wall in a given slice of the stability manifold. Similar results have been obtained by Liu [\[2015\]](#), but the result is critical to our discussion so we include the proof. See [\[Bayer and Macrì 2014b, §4\]](#) for more details on the definitions and results we use here.

Throughout this subsection, let $\mathbf{v} \in K_0(X)$ be a vector such that the moduli space $M_{H,D}(\mathbf{v})$ of (H, D) -Gieseker semistable sheaves admits a (quasi-)universal family \mathcal{E} which is unique up to equivalence (Hilbert schemes $X^{[n]}$ are examples of such spaces). We also let $\sigma = (Z, \mathcal{A})$ be a stability condition in the closure of the Gieseker chamber for \mathbf{v} in the (H, D) -slice. Then there is a well-defined corresponding divisor $D_\sigma \in N^1(M_{H,D-K_X/2}(\mathbf{v}))$ which is independent of the choice of \mathcal{E} .

Let $(\mathbf{v}, \mathbf{w}) = \chi(\mathbf{v}\mathbf{w})$ be the Euler pairing on $K_{\text{num}}(X)_{\mathbb{R}}$, and write $\mathbf{v}^{\perp} \subset K_{\text{num}}(X)_{\mathbb{R}}$ for the orthogonal complement with respect to this pairing. The correspondence between stability conditions and divisor classes is understood in terms of the Donaldson homomorphism

$$\lambda : \mathbf{v}^{\perp} \rightarrow N^1(M_{H,D-K_X/2}(\mathbf{v})).$$

Since the Euler pairing is nondegenerate, there is a unique vector $\mathbf{w}_{\sigma} \in \mathbf{v}^{\perp}$ such that

$$\mathfrak{S}\left(-\frac{Z(\mathbf{w}')}{Z(\mathbf{v})}\right) = (\mathbf{w}', \mathbf{w}_{\sigma}),$$

for all $\mathbf{w}' \in K_{\text{num}}(X)_{\mathbb{R}}$. Bayer and Macrì show that $D_{\sigma} = \lambda(\mathbf{w}_{\sigma})$. In what follows, we write vectors in $K_{\text{num}}(X)_{\mathbb{R}}$ as $(\text{ch}_0, \text{ch}_1, \text{ch}_2)$.

Proposition 3.8. *With the above assumptions, suppose σ lies on a numerical wall W in the (H, D) -slice with center $(s_W, 0)$. Then \mathbf{w}_{σ} is a multiple of*

$$\left(-1, -\frac{1}{2}K_X + s_W H + D, m\right) \in \mathbf{v}^{\perp},$$

where m is determined by the requirement $\mathbf{w}_{\sigma} \in \mathbf{v}^{\perp}$.

In particular, if X has irregularity 0 and $\mathbf{v} = (1, 0, -n)$ is the vector for $X^{[n]}$, then the divisor D_{σ} is a multiple of

$$\frac{1}{2}K_X^{[n]} - s_W H^{[n]} - D^{[n]} - \frac{1}{2}B.$$

Remark 3.9. Suppose X has irregularity 0. Up to scale, the divisors induced by stability conditions in the (H, D) -slice give a ray in $N^1(X^{[n]})$ emanating from the class $H^{[n]} \in \text{Nef}(X) \subset \text{Nef}(X^{[n]})$. The particular ray is determined by the choice of the twisting divisor D .

Proof of Proposition 3.8. Since σ is in the (H, D) -slice, write $\sigma = \sigma_{\beta,\alpha}$ and $(Z, \mathcal{A}) = (Z_{\beta,\alpha}, \mathcal{A}_{\beta})$ for short. Put $z = -1/Z(\mathbf{v}) = u + iv$. We evaluate the identity

$$\mathfrak{S}(zZ(\mathbf{w}')) = (\mathbf{w}', \mathbf{w}_{\sigma})$$

defining \mathbf{w}_{σ} on various classes \mathbf{w}' to compute \mathbf{w}_{σ} .

Write the Chern character $\mathbf{w}_{\sigma} = (r, C, d)$. Then

$$-v = \mathfrak{S}(zZ(0, 0, 1)) = ((0, 0, 1), \mathbf{w}_{\sigma}) = r,$$

so $r = -v$. Next, for any curve class C' ,

$$\begin{aligned} (u + \beta v)(C' \cdot H) + v(C' \cdot D) &= \mathfrak{S}(zZ(0, C', 0)) \\ &= ((0, C', 0), \mathbf{w}_{\sigma}) = \chi(0, -vC', C' \cdot C). \end{aligned}$$

By Riemann–Roch and adjunction,

$$\begin{aligned} \chi(0, -vC', C' \cdot C) &= -v \left(\left(-\frac{1}{v}(C' \cdot C) + \frac{1}{2}(C')^2 \right) - \frac{1}{2}(C')^2 - \frac{1}{2}(C' \cdot K_X) \right) \\ &= C' \cdot C + \frac{v}{2}(K_X \cdot C'), \end{aligned}$$

so

$$C' \cdot C = (u + \beta v)(C' \cdot H) + v(C' \cdot D) - \frac{v}{2}(C' \cdot K_X)$$

for every class C' . Thus for any class C' with $C' \cdot H = 0$, we have $C' \cdot C = v(C' \cdot D) - \frac{1}{2}v(C' \cdot K_X)$; it follows that there is some number a with

$$C = -\frac{v}{2}K_X + aH + vD.$$

Considering $C = H$ shows that $a = u + \beta v$. Therefore,

$$w_\sigma = \left(-v, -\frac{v}{2}K_X + (u + \beta v)H, m \right),$$

where m is chosen such that $w_\sigma \in v^\perp$.

Finally, a straightforward calculation shows that

$$\frac{u}{v} + \beta = v_\sigma(v) + \beta = s_W$$

holds for all (β, α) along W . The follow-up statement for Hilbert schemes follows by computing the Donaldson homomorphism. \square

3D. Dual curves. Suppose D_{σ_0} is the nef divisor corresponding to the Gieseker wall for $X^{[n]}$ in the (H, D) -slice. Showing that D_{σ_0} is an extremal nef divisor amounts to showing that there is some curve $\gamma \subset X^{[n]}$ with $D_{\sigma_0} \cdot \gamma = 0$. By the positivity lemma, this happens when γ parametrizes objects of $X^{[n]}$ which are generically S -equivalent with respect to σ_0 .

In every case where we computed the Gieseker wall, the wall can be given by a destabilizing subobject which is a line bundle $\mathcal{O}_X(-C)$ with C an effective curve. If Z is a length- n subscheme of C , then there is a destabilizing sequence

$$0 \rightarrow \mathcal{O}_X(-C) \rightarrow I_Z \rightarrow I_{Z \subset C} \rightarrow 0.$$

If $\text{ext}^1(I_{Z \subset C}, \mathcal{O}_X(-C)) \geq 2$, then curves of objects of $X^{[n]}$ which are generically S -equivalent with respect to σ_0 are obtained by varying the extension class. We obtain the following general result.

Lemma 3.10. *Suppose the Gieseker wall for $X^{[n]}$ in the (H, D) -slice is computed by the subobject $\mathcal{O}_X(-C)$, where C is an effective curve class of arithmetic genus $p_a(C)$. If $n \geq p_a(C) + 1$, then the corresponding nef divisor D_{σ_0} is extremal.*

Proof. Bilinearity of the Euler characteristic $\chi(\cdot, \cdot)$ and Serre duality show that

$$\chi(I_{Z \subset C}, \mathcal{O}_X(-C)) = p_a(C) - 1 - n.$$

Therefore, once $n \geq p_a(C) + 1$ we will have $\chi(I_{Z \subset C}, \mathcal{O}_X(-C)) \leq -2$, and curves orthogonal to D_{σ_0} can be constructed by varying the extension class. \square

Combining [Lemma 3.10](#) with our previous results on the computation of the Gieseker wall gives us the following asymptotic result.

Theorem 3.11. *Fix a slice (H, D) for $\text{Stab}(X)$. There is some $L \in \text{CrDiv}(H, D)$ such that for all $n \gg 0$ the Gieseker wall is computed by $\mathcal{O}_X(-L)$. Furthermore, the corresponding nef divisor is extremal.*

Proof. Recall that the set $\text{CrDiv}(H, D)/\sim$ of critical divisors modulo numerical equivalence is finite; say $\{L_1, \dots, L_m\}$ is a set of representatives. For $1 \leq i \leq m$, let $(s_i(n), 0)$ be the center of the wall $\mathcal{O}_X(-L_i)$ for $X^{[n]}$. Then $s_i(n)$ is a linear function of n by [Lemma 3.3](#), so there is some i with $s_i(n) \leq s_j(n)$ for all $1 \leq j \leq m$ and $n \gg 0$. Then by [Corollary 3.7](#) the Gieseker wall is given by $\mathcal{O}_X(-L_i)$. Again increasing n if necessary, the divisor D_{σ_0} corresponding to the Gieseker wall is extremal by [Lemma 3.10](#). \square

Remark 3.12. The requirement $n \geq p_a(C) + 1$ in [Lemma 3.10](#) is not typically sharp. For example, if $|C|$ contains a smooth curve we may as well assume C is smooth. Then $I_{Z \subset C}$ is a line bundle on C , and

$$\text{Ext}^1(I_{Z \subset C}, \mathcal{O}_X(-C)) \cong H^0(\mathcal{O}_C(Z)).$$

Thus a g_n^1 on C gives a curve which is orthogonal to D_{σ_0} . The following fact from Brill–Noether theory therefore provides curves on $X^{[n]}$ for smaller values on n .

Lemma 3.13 [[Arbarello et al. 1985](#)]. *If C is smooth of genus g , then it has a g_n^1 for any $n \geq \lceil (g + 2)/2 \rceil$.*

For specific surfaces, some curves in $|C|$ may have highly special linear series giving better constructions of curves on $X^{[n]}$.

4. Picard rank 1 examples

For the rest of the paper, we will apply the methods of [Section 3](#) to compute $\text{Nef}(X^{[n]})$ for several interesting surfaces X . These applications form the heart of the paper.

4A. Picard rank 1 in general. Suppose $\text{Pic}(X) \cong \mathbb{Z}H$ for some ample divisor H . If we choose $D = -aH$, where $a > 0$ is the smallest positive integer such that aH is effective, then $\text{CrDiv}(H, D) = \{-D\}$.

Lemma 4.1. *Suppose $\text{Pic}(X) = \mathbb{Z}H$ and aH is the minimal effective class. If $n \geq (aH)^2 = a^2d$, then the Gieseker wall for $X^{[n]}$ is the wall given by $\mathcal{O}_X(-aH)$.*

Proof. Apply [Corollary 3.7](#) with $D = -aH$. \square

Note that when $n > a^2d$, additional information about the Jordan–Hölder filtration can be obtained as in [Corollary 3.7](#). We use formula (7) to see that the wall W given by $\mathcal{O}_X(-aH)$ has center $(s_W, 0)$ with

$$s_W = \frac{a}{2} - \frac{n}{ad}.$$

Combining [Lemmas 4.1](#) and [3.10](#) with [Proposition 3.8](#), we have proved the following general result.

Theorem 4.2. *Suppose $\text{Pic } X \cong \mathbb{Z}H$ and aH is the minimal effective class. If $n \geq a^2d$ then the divisor*

$$\frac{1}{2}K_X^{[n]} + \left(\frac{a}{2} + \frac{n}{ad}\right)H^{[n]} - \frac{1}{2}B \quad (8)$$

is nef. Additionally, if $n \geq p_a(aH) + 1$ then this divisor is extremal, so $\text{Nef}(X^{[n]})$ is spanned by this divisor and $H^{[n]}$. An orthogonal curve is given by letting n points move in a g_n^1 on a curve of class aH .

Remark 4.3. If $\text{Pic}(X) = \mathbb{Z}H$ and H is already effective, then a different argument computes the Gieseker wall so long as $2n > d$, improving the bound in [Lemma 4.1](#). However, fine information about the Jordan–Hölder filtration of a destabilized ideal sheaf is not obtained. In fact, if $n \leq d$ then the destabilizing behavior can be complicated. For instance, a scheme Z contained in the complete intersection of two curves of class H will admit an interesting map from $\mathcal{O}_X(-H)^{\oplus 2}$.

Proposition 4.4. *Suppose $\text{Pic } X = \mathbb{Z}H$ and H is effective. If $2n > d$, then the Gieseker wall for $X^{[n]}$ in the $(H, -H)$ -slice is the wall given by $\mathcal{O}_X(-H)$. Thus the divisor (8) with $a = 1$ is nef.*

Proof. Let W be the numerical wall given by $\mathcal{O}_X(-H)$. If no actual wall is larger than W then, by the proof of [Proposition 3.5](#), W is an actual wall, and hence the Gieseker wall. If there is a destabilizing sequence

$$0 \rightarrow F \rightarrow I_Z \rightarrow G \rightarrow 0$$

giving a wall W' larger than W , then $F, G \in \mathcal{A}_0$ since W passes through the origin in the (β, α) -plane. Fix $\alpha > 0$ such that $(0, \alpha)$ lies on W' . We have

$$H \cdot \text{ch}_1^{-H}(F) = \mathfrak{S}Z_{0,\alpha}(F) \geq 0 \quad \text{and} \quad H \cdot \text{ch}_1^{-H}(G) = \mathfrak{S}Z_{0,\alpha}(G) \geq 0.$$

Since d is the smallest intersection number of H with an integral divisor and

$$d = \mathfrak{S}Z_{0,\alpha}(I_Z) = \mathfrak{S}Z_{0,\alpha}(F) + \mathfrak{S}Z_{0,\alpha}(G),$$

we conclude that either $\mathfrak{S}Z_{0,\alpha}(F) = 0$ or $\mathfrak{S}Z_{0,\alpha}(G) = 0$. Thus either F or G has infinite $\sigma_{0,\alpha}$ -slope, contradicting the fact that $(0, \alpha)$ is on W' . \square

We now further relax the lower bound on n needed to guarantee the existence of orthogonal curve classes in special cases.

4B. Surfaces in \mathbb{P}^3 . By the Noether–Lefschetz theorem, a very general surface $X \subset \mathbb{P}^3$ of degree $d \geq 4$ is smooth of Picard rank 1 and irregularity 0. Let H be the hyperplane class and put $D = -H$. We have $K_X = (d - 4)H$, so [Proposition 4.4](#) shows that if $2n > d$ then the divisor

$$\left(\frac{d}{2} - \frac{3}{2} + \frac{n}{d}\right)H^{[n]} - \frac{1}{2}B$$

is nef. If C is any smooth hyperplane section then the projection from a point on C gives a map of degree $d - 1$ to \mathbb{P}^1 , so C carries a g_n^1 for any $n \geq d - 1$. We have proved the following result.

Proposition 4.5. *Let X be a smooth degree- d hypersurface in \mathbb{P}^3 with Picard rank 1. The divisor*

$$\left(\frac{d}{2} - \frac{3}{2} + \frac{n}{d}\right)H^{[n]} - \frac{1}{2}B$$

on $X^{[n]}$ is nef if $2n > d$. If $n \geq d - 1$, then it is extremal, and together with $H^{[n]}$ it spans $\text{Nef}(X^{[n]})$.

Remark 4.6. The behavior of $\text{Nef}(X^{[n]})$ for smaller n in [Proposition 4.5](#) is more mysterious. Even the cases $d = 5$ and $n = 2, 3$ are interesting.

Remark 4.7. The case $d = 4$ of [Proposition 4.5](#) recovers a special case of [\[Bayer and Macrì 2014b, Proposition 10.3\]](#) for K3 surfaces. The case $d = 1$ recovers the computation of the nef cone of $\mathbb{P}^{2[n]}$ [\[Arcara et al. 2013\]](#).

4C. Branched covers of \mathbb{P}^2 . Next we consider cyclic branched covers of \mathbb{P}^2 . Let X be a very general cyclic degree- d cover of \mathbb{P}^2 , branched along a degree- e curve. Note that this means that d necessarily divides e . We can view these covers as hypersurfaces in a weighted projective space, which gives us a Noether–Lefschetz type theorem: $\text{Pic } X = \mathbb{Z}H$, generated by the pullback H of the hyperplane class on \mathbb{P}^2 , provided that X has positive geometric genus. The canonical bundle of X is

$$K_X = -3H + e\left(\frac{d-1}{d}\right)H = \left(\frac{e(d-1)}{d} - 3\right)H.$$

Then X will have positive geometric genus if $e \geq 3d/(d - 1)$.

Setting $D = -H$, we see that if $2n > d$ then the divisor class

$$\left(\frac{e(d-1)}{2d} - 1 + \frac{n}{d}\right)H^{[n]} - \frac{1}{2}B$$

is nef by [Proposition 4.4](#). The preimage of a line is a curve of class H , and it carries a g_d^1 given by the map to \mathbb{P}^2 . Therefore, the above divisor is extremal once $n \geq d$.

Proposition 4.8. *Let X be a very general degree- d cyclic cover of \mathbb{P}^2 ramified along a degree- e curve, where d divides e and $e \geq 3d/(d-1)$. The divisor*

$$\left(\frac{e(d-1)}{2d} - 1 + \frac{n}{d}\right)H^{[n]} - \frac{1}{2}B$$

on $X^{[n]}$ is nef if $2n > d$. For $n \geq d$, this class is extremal, and together with $H^{[n]}$ it spans $\text{Nef}(X^{[n]})$.

5. Del Pezzo surfaces of degree 1

Bertram and Coskun [\[2013\]](#) studied the birational geometry of $X^{[n]}$ when X is a minimal rational surface or a del Pezzo surface. In particular, they completely computed the nef cones of all these Hilbert schemes except in the case of a del Pezzo surface of degree 1. The constructions they gave were classical: they produced nef divisors from k -very ample line bundles, and dual curves by letting collections of points move in linear pencils on special curves.

In this section, we will compute the nef cone of $X^{[n]}$, where X is a smooth del Pezzo surface of degree 1. Then $X \cong \text{Bl}_{p_1, \dots, p_8} \mathbb{P}^2$ for distinct points p_1, \dots, p_8 with the property that $-K_X$ is ample (see [\[Manin 1974, Theorem 24.4\]](#) or [\[Beauville 1996, Exercise V.21.1\]](#)). This application exhibits the full strength of the methods of [Section 3](#).

5A. Notation and statement of results. Let H be the class of a line and let the 8 exceptional divisors over the p_i be E_1, \dots, E_8 , so $\text{Pic}(X) \cong \mathbb{Z}H \oplus \mathbb{Z}E_1 \oplus \dots \oplus \mathbb{Z}E_8$ and $K_X = -3H + \sum_i E_i$. Recall that a (-1) -curve on X is a smooth rational curve of self-intersection -1 . It is simplest to describe the dual cone of effective curves. We recommend reviewing [Section 2A](#) for notation.

Theorem 5.1. *The cone of curves $\text{NE}(X^{[n]})$ is spanned by all the classes $E_{[n]}$ given by (-1) -curves $E \subset X$, the class of a curve contracted by the Hilbert–Chow morphism, and the class $F_{[n]}$, where $F \in |-K_X|$ is an anticanonical curve.*

The 240 (-1) -curves E on X are well-known. The possible classes are

$$(0; 1), \quad (1; 1^2), \quad (2; 1^5), \quad (3; 2, 1^6), \quad (4; 2^3, 1^5), \quad (5; 2^6, 1^2), \quad (6; 3, 2^7),$$

where, e.g., $(4; 2^3, 1^5)$ denotes any class equivalent to

$$4H - 2E_1 - 2E_2 - 2E_3 - E_4 - E_5 - E_6 - E_7 - E_8$$

under the natural action of S_8 on $\text{Pic}(X)$. The cone of curves $\text{NE}(X)$ is spanned by the classes of the (-1) -curves. The Weyl group action on $\text{Pic}(X)$ acts transitively

on (-1) -curve classes. It also acts transitively on systems of 8 pairwise disjoint (-1) -curves; dually, it acts transitively on the extremal rays of the nef cone $\text{Nef}(X)$. We refer the reader to [Manin 1974, §26] for details.

Consider the divisor class $(n - 1)(-K_X)^{[n]} - \frac{1}{2}B$. If E is any (-1) -curve on X , then $-K_X \cdot E = 1$, so

$$E_{[n]} \cdot \left((n - 1)(-K_X)^{[n]} - \frac{1}{2}B \right) = (n - 1)(-K_X \cdot E) - (n - 1) = 0.$$

Let $\Lambda \subset N^1(X^{[n]})$ be the cone spanned by divisors which are nonnegative on all classes $E_{[n]}$ and curves contracted by the Hilbert–Chow morphism. It follows that $\Lambda \supset \text{Nef}(X^{[n]})$ is spanned by $\text{Nef}(X) \subset \text{Nef}(X^{[n]})$ and the single additional class $(n - 1)(-K_X)^{[n]} - \frac{1}{2}B$.

However, $\text{Nef}(X^{[n]}) \subset \Lambda$ is a *proper* subcone. Indeed, if $F \in |-K_X|$ is an anti-canonical curve then by Riemann–Hurwitz $F_{[n]} \cdot B = 2n$, so $F_{[n]} \cdot \left((n - 1)(-K_X)^{[n]} - \frac{1}{2}B \right) = -1$. Let $\Lambda' \subset \Lambda$ be the subcone of $F_{[n]}$ -nonnegative divisors. Taking duals, we see that Theorem 5.1 is equivalent to the next result.

Theorem 5.2. *The nef cone of $X^{[n]}$ is Λ' .*

To prove Theorem 5.2, we must show that all the extremal rays of Λ' are actually nef. Suppose $N \in \text{Nef}(X)$ spans an extremal ray of $\text{Nef}(X)$. Then the cone spanned by $N^{[n]}$ and $(n - 1)(-K_X)^{[n]} - \frac{1}{2}B$ contains a single ray of $F_{[n]}$ -orthogonal divisors, and this ray is an extremal ray of Λ' . Conversely, due to our description of the cone Λ , the extremal rays of Λ' which are not in $\text{Nef}(X)$ are all obtained in this way.

5B. Choosing a slice. More concretely, making use of the Weyl group action we may as well assume our extremal nef class $N \in \text{Nef}(X)$ is $H - E_1$. The corresponding $F_{[n]}$ -orthogonal ray described in the previous paragraph is spanned by

$$(n - 1)(-K_X)^{[n]} + \frac{1}{2}(H^{[n]} - E_1^{[n]}) - \frac{1}{2}B; \tag{9}$$

our job is to show that this class is nef. We will prove this by exhibiting this divisor as the nef divisor on $X^{[n]}$ corresponding to the Gieseker wall for a suitable choice of slice of $\text{Stab}(X)$.

To apply the methods of Section 3, it is convenient to choose our polarization to be

$$P = \left(n - \frac{3}{2} \right)(-K_X) + \frac{1}{2}(H - E_1)$$

(which depends on n !) and our antieffective class to be $D = K_X$. Observe that P is ample since it is the sum of an ample and a nef class. If we show that the Gieseker wall W in the (P, K_X) -slice has center $(s_W, 0) = (-1, 0)$, then Proposition 3.8 implies the divisor class (9) is nef.

5C. Critical divisors. Our plan is to apply [Corollary 3.7](#) to compute the Gieseker wall in the (P, K_X) -slice. We must first identify the set $\text{CrDiv}(P, K_X)$ of critical divisors.

Lemma 5.3. *If $n > 2$, then the set $\text{CrDiv}(P, K_X)$ consists of $-K_X$ and the classes L of (-1) -curves on X with $L \cdot (H - E_1) \leq 1$.*

When $n = 2$, the above classes are still critical. Additionally, the class $H - E_1$ is critical, as is any sum of two (-1) -curves L_1, L_2 with $L_i \cdot (H - E_1) = 0$.

Proof. Write $2P = A + N$, where $A = (2n - 3)(-K_X)$ is ample and $N = H - E_1$ is nef. Then $A \cdot (-K_X) = 2n - 3$ and $N \cdot (-K_X) = 2$, so an effective curve class $L \neq -K_X$ is in $\text{CrDiv}(P, K_X)$ if and only if $L \cdot (2P) < 2n - 1$.

First suppose $n > 2$, and let $L \in \text{CrDiv}(P, K_X)$. If $L \cdot (-K_X) \geq 2$, then $L \cdot (2P) \geq 4n - 6 > 2n - 1$, so L is not critical. Therefore, $L \cdot (-K_X) = 1$. Thus any curve of class L is reduced and irreducible. By the Hodge index theorem,

$$L^2 = L^2 \cdot (-K_X)^2 \leq (L \cdot (-K_X))^2 = 1,$$

with equality if and only if $L = -K_X$. If the inequality is strict, then by adjunction we must have $L^2 = -1$ and L is a (-1) -curve. Since $L \cdot (2P) < 2n - 1$, we further have $L \cdot N \leq 1$.

Suppose instead that $n = 2$ and $L \in \text{CrDiv}(P, K_X)$. The cases $L \cdot (2P) \leq 1$ and $L \cdot (2P) \geq 3$ follow as in the previous case. The only other possibility is that $L \cdot (-K_X) = 2$ and $L \cdot N = 0$. Since $L \cdot N = 0$, the curve L is a sum of curves in fibers of the projection $X \rightarrow \mathbb{P}^1$ given by $|N|$. This easily implies the result. \square

The next application of [Corollary 3.7](#) completes the proof of [Theorems 5.1](#) and [5.2](#).

Proposition 5.4. *The Gieseker wall for $X^{[n]}$ in the (P, K_X) -slice has center $(-1, 0)$, and is given by the subobject $\mathcal{O}_X(K_X)$. It coincides with the wall given by $\mathcal{O}_X(-L)$, where L is any (-1) -curve with $L \cdot (H - E_1) = 0$.*

Proof. By [\(7\)](#), the center of the wall for $\mathcal{O}_X(K_X)$ is $(s_W, 0)$ with

$$s_W = \frac{2n - K_X^2}{(2P) \cdot K_X} = -1.$$

A straightforward computation shows $\eta_{P, K_X} < n$ for all $n \geq 2$. Therefore, by [Corollary 3.7](#), the Gieseker wall is computed by a critical divisor.

We only need to verify that no other critical divisor gives a larger wall. Let $L \in \text{CrDiv}(P, K_X)$. By [Lemma 3.3](#), the center of the wall given by $\mathcal{O}_X(-L)$ lies at the point $(s_L, 0)$, where

$$s_L = -\frac{2n + L^2 + 2(K_X \cdot L)}{(2P) \cdot L}.$$

If L is a (-1) -curve, then

$$s_L = -\frac{2n-3}{(2P) \cdot L} = -\frac{2n-3}{2n-3 + L \cdot (H - E_1)} \geq -1,$$

with equality if and only if $L \cdot (H - E_1) = 0$. This proves the result if $n > 2$.

To complete the proof when $n = 2$, we only need to consider the additional critical classes mentioned in [Lemma 5.3](#). For every such $L \in \text{CrDiv}(P, K_X)$ we have $L \cdot K_X = -2$ and $L^2 \leq 0$. Thus $s_L \geq 0$ for every such divisor. \square

Acknowledgements

This work was initiated at the 2015 Algebraic Geometry Bootcamp preceding the Algebraic Geometry Summer Research Institute organized by the AMS and the University of Utah. We would like to thank the organizers of both programs for providing the wonderful environment where this collaboration could happen. Additionally, we would like to thank Arend Bayer, Izzet Coskun, and Emanuele Macrì for many valuable discussions on Bridgeland stability.

References

- [Arbarello et al. 1985] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of algebraic curves*, vol. I, Grundlehren der Math. Wissenschaften **267**, Springer, New York, 1985. [MR 770932](#) [Zbl 0559.14017](#)
- [Arcara and Bertram 2013] D. Arcara and A. Bertram, “Bridgeland-stable moduli spaces for K -trivial surfaces”, *J. Eur. Math. Soc. (JEMS)* **15**:1 (2013), 1–38. [MR 2998828](#) [Zbl 1259.14014](#)
- [Arcara et al. 2013] D. Arcara, A. Bertram, I. Coskun, and J. Huizenga, “The minimal model program for the Hilbert scheme of points on \mathbb{P}^2 and Bridgeland stability”, *Adv. Math.* **235** (2013), 580–626. [MR 3010070](#) [Zbl 1267.14023](#)
- [Bayer and Macrì 2014a] A. Bayer and E. Macrì, “MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations”, *Invent. Math.* **198**:3 (2014), 505–590. [MR 3279532](#) [Zbl 1308.14011](#)
- [Bayer and Macrì 2014b] A. Bayer and E. Macrì, “Projectivity and birational geometry of Bridgeland moduli spaces”, *J. Amer. Math. Soc.* **27**:3 (2014), 707–752. [MR 3194493](#) [Zbl 1314.14020](#)
- [Bayer et al. 2014] A. Bayer, E. Macrì, and P. Stellari, “The Space of Stability Conditions on Abelian Threefolds, and on some Calabi–Yau Threefolds”, preprint, 2014. [arXiv 1410.1585](#)
- [Beauville 1996] A. Beauville, *Complex algebraic surfaces*, 2nd ed., London Mathematical Society Student Texts **34**, Cambridge University Press, 1996. [MR 1406314](#) [Zbl 0849.14014](#)
- [Beltrametti and Sommese 1991] M. Beltrametti and A. J. Sommese, “Zero cycles and k -th order embeddings of smooth projective surfaces”, pp. 33–48 in *Problems in the theory of surfaces and their classification* (Cortona, 1988), edited by F. Catanese et al., Sympos. Math. **XXXII**, Academic Press, London, 1991. [MR 1273371](#) [Zbl 0827.14029](#)
- [Beltrametti et al. 1989] M. Beltrametti, P. Francia, and A. J. Sommese, “On Reider’s method and higher order embeddings”, *Duke Math. J.* **58**:2 (1989), 425–439. [MR 1016428](#) [Zbl 0702.14010](#)

- [Bertram and Coskun 2013] A. Bertram and I. Coskun, “The birational geometry of the Hilbert scheme of points on surfaces”, pp. 15–55 in *Birational geometry, rational curves, and arithmetic*, edited by F. Bogomolov et al., Springer, New York, 2013. MR 3114922 Zbl 1273.14032
- [Bertram et al. 2014] A. Bertram, C. Martinez, and J. Wang, “The birational geometry of moduli spaces of sheaves on the projective plane”, *Geom. Dedicata* **173** (2014), 37–64. MR 3275289 Zbl 1314.14087
- [Bridgeland 2007] T. Bridgeland, “Stability conditions on triangulated categories”, *Ann. of Math. (2)* **166**:2 (2007), 317–345. MR 2373143 Zbl 1137.18008
- [Bridgeland 2008] T. Bridgeland, “Stability conditions on $K3$ surfaces”, *Duke Math. J.* **141**:2 (2008), 241–291. MR 2376815 Zbl 1138.14022
- [Catanese and Göttsche 1990] F. Catanese and L. Göttsche, “ d -very-ample line bundles and embeddings of Hilbert schemes of 0-cycles”, *Manuscripta Math.* **68**:3 (1990), 337–341. MR 1065935 Zbl 0729.14006
- [Coskun and Huizenga 2014] I. Coskun and J. Huizenga, “Interpolation, Bridgeland stability and monomial schemes in the plane”, *J. Math. Pures Appl. (9)* **102**:5 (2014), 930–971. MR 3271294 Zbl 06361756
- [Coskun and Huizenga 2016] I. Coskun and J. Huizenga, “The ample cone of moduli spaces of sheaves on the plane”, *Algebr. Geom.* **3**:1 (2016), 106–136. MR 3455422
- [Ellingsrud et al. 2001] G. Ellingsrud, L. Göttsche, and M. Lehn, “On the cobordism class of the Hilbert scheme of a surface”, *J. Algebraic Geom.* **10**:1 (2001), 81–100. MR 1795551 Zbl 0976.14002
- [Fogarty 1968] J. Fogarty, “Algebraic families on an algebraic surface”, *Amer. J. Math* **90** (1968), 511–521. MR 0237496 Zbl 0176.18401
- [Fogarty 1973] J. Fogarty, “Algebraic families on an algebraic surface, II: The Picard scheme of the punctual Hilbert scheme”, *Amer. J. Math.* **95** (1973), 660–687. MR 0335512 Zbl 0299.14020
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, New York, 1977. MR 0463157 Zbl 0367.14001
- [Hassett and Tschinkel 2010] B. Hassett and Y. Tschinkel, “Intersection numbers of extremal rays on holomorphic symplectic varieties”, *Asian J. Math.* **14**:3 (2010), 303–322. MR 2755719 Zbl 1216.14012
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, *Ergebnisse der Mathematik (3)* **48**, Springer, Berlin, 2004. MR 2095471 Zbl 1093.14501
- [Li and Zhao 2013] C. Li and X. Zhao, “The MMP for deformations of Hilbert schemes of points on the projective plane”, preprint, 2013. arXiv 1312.1748
- [Liu 2015] W. Liu, “Bayer–Macrì decomposition on Bridgeland moduli spaces over surfaces”, preprint, 2015. arXiv 1501.06397
- [Maciocia 2014] A. Maciocia, “Computing the walls associated to Bridgeland stability conditions on projective surfaces”, *Asian J. Math.* **18**:2 (2014), 263–279. MR 3217637 Zbl 1307.14022
- [Maciocia and Meachan 2013] A. Maciocia and C. Meachan, “Rank 1 Bridgeland stable moduli spaces on a principally polarized abelian surface”, *Int. Math. Res. Not.* **2013**:9 (2013), 2054–2077. MR 3053413 Zbl 1312.14036
- [Manin 1974] Y. I. Manin, *Cubic forms: algebra, geometry, arithmetic*, North-Holland, Amsterdam, 1974. MR 0460349 Zbl 0277.14014
- [Matsuki and Wentworth 1997] K. Matsuki and R. Wentworth, “Mumford–Thaddeus principle on the moduli space of vector bundles on an algebraic surface”, *Internat. J. Math.* **8**:1 (1997), 97–148. MR 1433203 Zbl 0879.14002

[Nuer 2014] H. Nuer, “Projectivity and Birational Geometry of Bridgeland Moduli spaces on an Enriques Surface”, preprint, 2014. [arXiv 1406.0908](#)

[Yanagida and Yoshioka 2014] S. Yanagida and K. Yoshioka, “Bridgeland’s stabilities on abelian surfaces”, *Math. Z.* **276**:1-2 (2014), 571–610. [MR 3150219](#) [Zbl 1292.14012](#)

Communicated by János Kollár

Received 2015-11-01

Revised 2016-03-12

Accepted 2016-05-07

bolognese.b@husky.neu.edu *Department of Mathematics, Northeastern University, Boston, MA 02115, United States*

huizenga@psu.edu *Department of Mathematics, The Pennsylvania State University, University Park, PA 16802, United States*

lin.yinb@husky.neu.edu *Department of Mathematics, Northeastern University, Boston, MA 02115, United States*

ebriedl@uic.edu *Department of Mathematics, Statistics, and CS, University of Illinois, Chicago, IL 60607, United States*

schmidt.707@osu.edu *Department of Mathematics, The Ohio State University, Columbus, OH 43210, United States*

mwoolf@ed.ac.uk *Department of Mathematics, Statistics, and CS, University of Illinois, Chicago, IL 60607, United States*

x.zhao@neu.edu *Department of Mathematics, Northeastern University, Boston, MA 02115, United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
Craig Huneke	University of Virginia, USA	Marie-France Vignéras	Université Paris VII, France
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Kei-Ichi Watanabe	Nihon University, Japan
János Kollár	Princeton University, USA	Efim Zelmanov	University of California, San Diego, USA
Yuri Manin	Northwestern University, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2016 is US \$290/year for the electronic version, and \$485/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 10 No. 4 2016

Moduli of morphisms of logarithmic schemes	695
JONATHAN WISE	
Residual intersections and the annihilator of Koszul homologies	737
SEYED HAMID HASSANZADEH and JOSE NAÉLITON	
The Prym map of degree-7 cyclic coverings	771
HERBERT LANGE and ANGELA ORTEGA	
Local bounds for L^p norms of Maass forms in the level aspect	803
SIMON MARSHALL	
Hasse principle for Kummer varieties	813
YONATAN HARPAZ and ALEXEI N. SKOROBOGATOV	
Analytic continuation on Shimura varieties with μ-ordinary locus	843
STÉPHANE BIJAKOWSKI	
A note on secondary K-theory	887
GONÇALO TABUADA	
Nef cones of Hilbert schemes of points on surfaces	907
BARBARA BOLOGNESE, JACK HUIZENGA, YINBANG LIN, ERIC RIEDL, BENJAMIN SCHMIDT, MATTHEW WOOLF and XIAOLEI ZHAO	
Interpolation for restricted tangent bundles of general curves	931
ERIC LARSON	