

# *Algebra & Number Theory*

Volume 11

2017

No. 3

**A duality in Buchsbaum rings  
and triangulated manifolds**

Satoshi Murai, Isabella Novik and Ken-ichi Yoshida





# A duality in Buchsbaum rings and triangulated manifolds

Satoshi Murai, Isabella Novik and Ken-ichi Yoshida

Let  $\Delta$  be a triangulated homology ball whose boundary complex is  $\partial\Delta$ . A result of Hochster asserts that the canonical module of the Stanley–Reisner ring  $\mathbb{F}[\Delta]$  of  $\Delta$  is isomorphic to the Stanley–Reisner module  $\mathbb{F}[\Delta, \partial\Delta]$  of the pair  $(\Delta, \partial\Delta)$ . This result implies that an Artinian reduction of  $\mathbb{F}[\Delta, \partial\Delta]$  is (up to a shift in grading) isomorphic to the Matlis dual of the corresponding Artinian reduction of  $\mathbb{F}[\Delta]$ . We establish a generalization of this duality to all triangulations of connected orientable homology manifolds with boundary. We also provide an explicit algebraic interpretation of the  $h''$ -numbers of Buchsbaum complexes and use it to prove the monotonicity of  $h''$ -numbers for pairs of Buchsbaum complexes as well as the unimodality of  $h''$ -vectors of barycentric subdivisions of Buchsbaum polyhedral complexes. We close with applications to the algebraic manifold  $g$ -conjecture.

## 1. Introduction

In this paper, we study an algebraic duality of Stanley–Reisner rings of triangulated homology manifolds with nonempty boundary. Our starting point is the following (unpublished) result of Hochster — see [Stanley 1996, Chapter II, §7]. (We defer most definitions until later sections.) Let  $\Delta$  be a triangulated  $(d - 1)$ -dimensional homology ball whose boundary complex is  $\partial\Delta$ , let  $\mathbb{F}[\Delta]$  be the Stanley–Reisner ring of  $\Delta$ , and let  $\mathbb{F}[\Delta, \partial\Delta]$  be the Stanley–Reisner module of  $(\Delta, \partial\Delta)$ . (Throughout the paper  $\mathbb{F}$  denotes an infinite field.) Hochster’s result asserts that the canonical module  $\omega_{\mathbb{F}[\Delta]}$  of  $\mathbb{F}[\Delta]$  is isomorphic to  $\mathbb{F}[\Delta, \partial\Delta]$ . In the last decade or so, this result had a lot of impact on the study of face numbers of simplicial complexes, especially in connection with the  $g$ -conjecture for spheres; see, for instance, the proof of Theorem 3.1 in the recent survey paper [Swartz 2014].

---

Murai’s research is partially supported by Grant-in-Aid for Scientific Research (C) 25400043.

Novik’s research is partially supported by NSF grant DMS-1361423.

Yoshida’s research is partially supported by Grant-in-Aid for Scientific Research (C) 25400050.

*MSC2010*: primary 13F55; secondary 13H10, 05E40, 52B05, 57Q15.

*Keywords*: triangulated manifolds, Buchsbaum rings,  $h$ -vectors, Stanley–Reisner rings.

One numerical consequence of Hochster's result is the following symmetry of  $h$ -numbers of homology balls:  $h_i(\Delta, \partial\Delta) = h_{d-i}(\Delta)$ . The  $h$ -numbers are certain linear combinations of the face numbers; they are usually arranged in a vector called the  $h$ -vector. In fact, Hochster's result implies a stronger statement: it implies that there is an isomorphism

$$\mathbb{F}[\Delta, \partial\Delta]/\Theta\mathbb{F}[\Delta, \partial\Delta] \cong (\mathbb{F}[\Delta]/\Theta\mathbb{F}[\Delta])^\vee(-d), \quad (1)$$

where  $\Theta$  is a linear system of parameters for  $\mathbb{F}[\Delta]$  and  $N^\vee$  is the (graded) Matlis dual of  $N$  (see, e.g., [Murai and Yanagawa 2014, Lemma 3.6]). As the  $\mathbb{F}$ -dimensions of the  $i$ -th graded components of modules in (1) are equal to  $h_i(\Delta, \partial\Delta)$  and  $h_{d-i}(\Delta)$ , respectively, the above-mentioned symmetry,  $h_i(\Delta, \partial\Delta) = h_{d-i}(\Delta)$ , follows.

Hochster's result was generalized to homology manifolds with boundary by Gräbe [1984]. To state Gräbe's result, we recall the definition of homology manifolds. We denote by  $\mathbb{S}^d$  and  $\mathbb{B}^d$  the  $d$ -dimensional sphere and ball, respectively. A pure  $d$ -dimensional simplicial complex  $\Delta$  is an  $\mathbb{F}$ -homology  $d$ -manifold without boundary if the link of each nonempty face  $\tau$  of  $\Delta$  has the homology of  $\mathbb{S}^{d-|\tau|}$  (over  $\mathbb{F}$ ). An  $\mathbb{F}$ -homology  $d$ -manifold with boundary is a pure  $d$ -dimensional simplicial complex  $\Delta$  such that

- (i) the link of each nonempty face  $\tau$  of  $\Delta$  has the homology of either  $\mathbb{S}^{d-|\tau|}$  or  $\mathbb{B}^{d-|\tau|}$ , and
- (ii) the set of all *boundary faces*, that is,

$$\partial\Delta := \{\tau \in \Delta : \text{the link of } \tau \text{ has the same homology as } \mathbb{B}^{d-|\tau|}\} \cup \{\emptyset\}$$

is a  $(d-1)$ -dimensional  $\mathbb{F}$ -homology manifold without boundary.

A connected  $\mathbb{F}$ -homology  $d$ -manifold with boundary is said to be *orientable* if the top homology  $\tilde{H}_d(\Delta, \partial\Delta)$  is isomorphic to  $\mathbb{F}$ .

Gräbe proved [1984] that if  $\Delta$  is an orientable homology manifold with boundary, then  $\mathbb{F}[\Delta, \partial\Delta]$  is the canonical module of  $\mathbb{F}[\Delta]$ . Gräbe also established [1987] a symmetry of  $h$ -numbers for such a  $\Delta$ . While Gräbe's original statement of symmetry is somewhat complicated, it was recently observed by the first two authors [Murai and Novik 2016] that it takes the following simple form when expressed in the language of  $h''$ -numbers.

**Theorem 1.1.** *Let  $\Delta$  be a connected orientable  $\mathbb{F}$ -homology  $(d-1)$ -manifold with nonempty boundary  $\partial\Delta$ . Then  $h''_i(\Delta, \partial\Delta) = h''_{d-i}(\Delta)$  for all  $i = 0, 1, \dots, d$ .*

The  $h''$ -numbers are certain modifications of  $h$ -numbers (see Section 3 for their definition). Similarly to the  $h$ -numbers, the  $h''$ -numbers are usually arranged in a vector, called the  $h''$ -vector. For homology manifolds, this vector appears to be a "correct" analog of the  $h$ -vector. Indeed, many properties of  $h$ -vectors of homology

balls and spheres are now known to hold for the  $h''$ -vectors of homology manifolds (with and without boundary); see recent survey articles [Klee and Novik 2016; Swartz 2014]. In light of Gräbe’s result [1984] and Theorem 1.1, it is natural to ask if Theorem 1.1 can be explained by Matlis duality. The first goal of this paper is to provide such an explanation.

To this end, the key object is the submodule  $\Sigma(\Theta; M)$  defined by Goto [1983]. Several definitions are in order. Let  $S = \mathbb{F}[x_1, \dots, x_n]$  be a graded polynomial ring over a field  $\mathbb{F}$  with  $\deg x_i = 1$  for  $i = 1, 2, \dots, n$ . Let  $M$  be a finitely generated graded  $S$ -module of Krull dimension  $d$  and let  $\Theta = \theta_1, \dots, \theta_d$  be a homogeneous system of parameters for  $M$ . The module  $\Sigma(\Theta; M)$  is defined as follows:

$$\Sigma(\Theta; M) := \Theta M + \sum_{i=1}^d ((\theta_1, \dots, \hat{\theta}_i, \dots, \theta_d)M :_M \theta_i) \subseteq M.$$

This module was introduced by Goto [1983] and has been used in the study of Buchsbaum local rings. Note that if  $M$  is a Cohen–Macaulay module, then  $\Sigma(\Theta; M) = \Theta M$ . We first show that this submodule is closely related to the  $h''$ -vectors. Specifically, we establish the following explicit algebraic interpretation of  $h''$ -numbers.

**Theorem 1.2.** *Let  $(\Delta, \Gamma)$  be a Buchsbaum relative simplicial complex of dimension  $d - 1$  and let  $\Theta$  be a linear system of parameters for  $\mathbb{F}[\Delta, \Gamma]$ . Then*

$$\dim_{\mathbb{F}}(\mathbb{F}[\Delta, \Gamma]/\Sigma(\Theta; \mathbb{F}[\Delta, \Gamma]))_j = h''_j(\Delta, \Gamma) \quad \text{for all } j = 0, 1, \dots, d.$$

Theorems 1.1 and 1.2 suggest that when  $\Delta$  is a homology manifold with boundary there might be a duality between the quotients of  $\mathbb{F}[\Delta]$  and  $\mathbb{F}[\Delta, \partial\Delta]$  by  $\Sigma(\Theta; \mathbb{F}[\Delta])$  and  $\Sigma(\Theta; \mathbb{F}[\Delta, \partial\Delta])$ , respectively. We prove that this is indeed the case. In fact, we prove a more general algebraic result on canonical modules of Buchsbaum graded algebras. Let  $\mathfrak{m} = (x_1, \dots, x_n)$  be the graded maximal ideal of  $S$ . If  $M$  is a finitely generated graded  $S$ -module of Krull dimension  $d$ , then the *canonical module* of  $M$  is the module

$$\omega_M := (H_{\mathfrak{m}}^d(M))^{\vee},$$

where  $H_{\mathfrak{m}}^i(M)$  denotes the  $i$ -th local cohomology module of  $M$ . We prove that the following isomorphism holds for *all* Buchsbaum graded algebras.

**Theorem 1.3.** *Let  $R = S/I$  be a Buchsbaum graded  $\mathbb{F}$ -algebra of Krull dimension  $d \geq 2$ , let  $\Theta = \theta_1, \dots, \theta_d \in S$  be a homogeneous system of parameters for  $R$ , and let  $\delta = \sum_{i=1}^d \deg \theta_i$ . If  $\text{depth } R \geq 2$ , then*

$$\omega_R/\Sigma(\Theta; \omega_R) \cong (R/\Sigma(\Theta; R))^{\vee}(-\delta).$$

As we mentioned above, if  $\Delta$  is a connected orientable homology manifold, then (by Gräbe’s result) the module  $\mathbb{F}[\Delta, \partial\Delta]$  is the canonical module of  $\mathbb{F}[\Delta]$ ; furthermore it is not hard to see that  $\mathbb{F}[\Delta]$  satisfies the assumptions of Theorem 1.3. (Indeed, the connectivity of  $\Delta$  implies that  $\text{depth } \mathbb{F}[\Delta] \geq 2$ , and the fact that  $\mathbb{F}[\Delta]$  is Buchsbaum follows from Schenzel’s theorem — see Theorem 3.1 below.) Hence we obtain the following corollary that generalizes (1).

**Corollary 1.4.** *Let  $\Delta$  be a connected orientable  $\mathbb{F}$ -homology  $(d - 1)$ -manifold with nonempty boundary  $\partial\Delta$ , and let  $\Theta$  be a linear system of parameters for  $\mathbb{F}[\Delta]$ . Then*

$$\mathbb{F}[\Delta, \partial\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta, \partial\Delta]) \cong (\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]))^\vee(-d).$$

We also consider combinatorial and algebraic applications of Theorems 1.2 and 1.3. Specifically, we prove the monotonicity of  $h''$ -vectors for pairs of Buchsbaum simplicial complexes, establish the unimodality of  $h''$ -vectors of barycentric subdivisions of Buchsbaum polyhedral complexes, provide a combinatorial formula for the  $a$ -invariant of Buchsbaum Stanley–Reisner rings, and extend [Swartz 2014, Theorem 3.1] as well as [Böhm and Papadakis 2015, Corollary 4.5], which is related to the sphere  $g$ -conjecture, to the generality of the manifold  $g$ -conjecture. More precisely, Swartz’s result asserts that most bistellar flips when applied to a homology sphere preserve the weak Lefschetz property, while Böhm and Papadakis’ result asserts that stellar subdivisions at large-dimensional faces of homology spheres preserve the weak Lefschetz property; we extend both of these results to bistellar flips and stellar subdivisions performed on connected orientable homology manifolds.

The structure of the paper is as follows. In Section 2 we prove Theorem 1.3 (although we defer part of the proof to the Appendix). In Section 3, we study Stanley–Reisner rings and modules of Buchsbaum simplicial complexes. There, after reviewing basics of simplicial complexes and Stanley–Reisner rings and modules, we verify Theorem 1.2 and derive several combinatorial consequences. Section 4 is devoted to applications of our results to the manifold  $g$ -conjecture. Finally, in the Appendix, we prove a graded version of [Goto 1983, Proposition 3.6] — a result on which our proof of Theorem 1.3 is based.

## 2. Duality in Buchsbaum rings

In this section, we prove Theorem 1.3. We start by recalling some definitions and results pertaining to Buchsbaum rings and modules.

Let  $S = \mathbb{F}[x_1, \dots, x_n]$  be a graded polynomial ring with  $\deg x_i = 1$ ,  $i = 1, 2, \dots, n$ , and let  $\mathfrak{m} = (x_1, \dots, x_n)$  be the graded maximal ideal of  $S$ . Given a graded  $S$ -module  $N$ , we denote by  $N(a)$  the module  $N$  with grading shifted by  $a \in \mathbb{Z}$ , that is,  $N(a)_j = N_{a+j}$ . If  $M$  is a finitely generated graded  $S$ -module of Krull dimension  $d$ , then a *homogeneous system of parameters* (or h.s.o.p.) for  $M$  is a sequence

$\Theta = \theta_1, \dots, \theta_d \in \mathfrak{m}$  of homogeneous elements such that  $\dim_{\mathbb{F}} M/\Theta M < \infty$ . A sequence  $\theta_1, \dots, \theta_r \in \mathfrak{m}$  of homogeneous elements is said to be a *weak  $M$ -sequence* if

$$(\theta_1, \dots, \theta_{i-1})M :_M \theta_i = (\theta_1, \dots, \theta_{i-1})M :_M \mathfrak{m}$$

for all  $i = 1, 2, \dots, r$ . We say that  $M$  is *Buchsbaum* if every h.s.o.p. for  $M$  is a weak  $M$ -sequence.

Let  $M$  be a finitely generated graded  $S$ -module and let  $\Theta$  be its h.s.o.p. The function  $P_{\Theta, M} : \mathbb{Z} \rightarrow \mathbb{Z}$  defined by

$$P_{\Theta, M}(n) := \dim_{\mathbb{F}}(M/(\Theta)^{n+1}M)$$

is called the *Hilbert–Samuel function of  $M$  with respect to the ideal  $(\Theta)$* . It is known that there is a polynomial in  $n$  of degree  $d$ , denoted by  $p_{\Theta, M}(n)$ , such that  $P_{\Theta, M}(n) = p_{\Theta, M}(n)$  for  $n \gg 0$  (see [Bruns and Herzog 1993, Proposition 4.6.2]). The leading coefficient of the polynomial  $p_{\Theta, M}(n)$  multiplied by  $d!$  is called the *multiplicity of  $M$  with respect to the ideal  $(\Theta)$* , and is denoted by  $e_{\Theta}(M)$ . We will make use of the following known characterization of the Buchsbaum property; see [Stückrad and Vogel 1986, Theorem I.1.12 and Proposition I.2.6].

**Lemma 2.1.** *A finitely generated graded  $S$ -module  $M$  of Krull dimension  $d$  is Buchsbaum if and only if, for every h.s.o.p.  $\Theta$  of  $M$ ,*

$$\dim_{\mathbb{F}}(M/\Theta M) - e_{\Theta}(M) = \sum_{i=0}^{d-1} \binom{d-1}{i} \dim_{\mathbb{F}} H_{\mathfrak{m}}^i(M).$$

We also recall some known results on canonical modules of Buchsbaum modules. For a finitely generated graded  $S$ -module  $M$ , the *depth* of  $M$  is defined by  $\text{depth}(M) := \min\{i : H_{\mathfrak{m}}^i(M) \neq 0\}$ .

**Lemma 2.2.** *Let  $M$  be a finitely generated graded  $S$ -module of Krull dimension  $d$ . If  $M$  is Buchsbaum, then the following properties hold:*

- (i)  $\omega_M$  is Buchsbaum.
- (ii)  $H_{\mathfrak{m}}^i(\omega_M) \cong (H_{\mathfrak{m}}^{d-i+1}(M))^{\vee}$  (as graded modules) for all  $i = 2, 3, \dots, d - 1$ .
- (iii) If  $\text{depth}(M) \geq 2$ , then  $(H_{\mathfrak{m}}^d(\omega_M))^{\vee} \cong M$  (as graded modules).
- (iv) If  $\Theta$  is an h.s.o.p. for  $M$ , then  $\Theta$  is also an h.s.o.p. for  $\omega_M$  and  $e_{\Theta}(\omega_M) = e_{\Theta}(M)$ .

See [Stückrad and Vogel 1986, Theorem II.4.9] for (i) and (ii), [Schenzel 1982, Korollar 3.13] or [Aoyama and Goto 1986, (1.16)] for (iii), and [Suzuki 1981, Lemma 2.2] for (iv).

The following theorem is a graded version of [Goto 1983, Proposition 3.6]. The original result by Goto is a statement about Buchsbaum *local rings*. It may be possible to prove Theorem 2.3 in the same way as in [Goto 1983] by replacing rings

with modules and by carefully keeping track of grading. However, since we could not find any literature allowing us to easily check this statement, we will provide its proof in the Appendix.

**Theorem 2.3.** *Let  $M$  be a finitely generated graded  $S$ -module of Krull dimension  $d > 0$ . Assume further that  $M$  is Buchsbaum and that  $\Theta = \theta_1, \dots, \theta_d$  is an h.s.o.p. for  $M$  with  $\deg \theta_i = \delta_i$ . Let  $\delta_C := \sum_{i \in C} \delta_i$  for  $C \subseteq [d] = \{1, 2, \dots, d\}$ . Then*

- (i)  $\Sigma(\Theta; M)/\Theta M \cong \bigoplus_{C \subsetneq [d]} H_m^{|C|}(M)(-\delta_C)$ , and
- (ii) *there is an injection  $M/\Sigma(\Theta; M) \rightarrow H_m^d(M)(-\delta_{[d]})$ .*

We are now in a position to prove Theorem 1.3.

*Proof of Theorem 1.3.* We first prove that  $R/\Sigma(\Theta; R)$  and  $\omega_R/\Sigma(\Theta; \omega_R)$  have the same  $\mathbb{F}$ -dimension. Indeed,

$$\begin{aligned} \dim_{\mathbb{F}}(R/\Sigma(\Theta; R)) &= \dim_{\mathbb{F}}(R/\Theta R) - \dim_{\mathbb{F}}(\Sigma(\Theta; R)/\Theta R) \\ &= e_{\Theta}(R) + \sum_{i=0}^{d-1} \binom{d-1}{i} \dim_{\mathbb{F}} H_m^i(R) - \sum_{i=0}^{d-1} \binom{d}{i} \dim_{\mathbb{F}} H_m^i(R) \\ &= e_{\Theta}(R) - \sum_{i=1}^{d-1} \binom{d-1}{i-1} \dim_{\mathbb{F}} H_m^i(R), \end{aligned}$$

where we use Lemma 2.1 and Theorem 2.3(i) for the second equality. Similarly, since  $\omega_R$  is Buchsbaum, the same computation yields

$$\dim_{\mathbb{F}}(\omega_R/\Sigma(\Theta; \omega_R)) = e_{\Theta}(\omega_R) - \sum_{i=1}^{d-1} \binom{d-1}{i-1} \dim_{\mathbb{F}} H_m^i(\omega_R).$$

Now, since  $\text{depth}(R) \geq 2$  by the assumptions of the theorem and since  $\text{depth}(\omega_R) \geq 2$  always holds (see [Aoyama 1980, Lemma 1]), parts (ii) and (iv) of Lemma 2.2 guarantee that

$$\dim_{\mathbb{F}}(R/\Sigma(\Theta; R)) = \dim_{\mathbb{F}}(\omega_R/\Sigma(\Theta; \omega_R)).$$

Thus, to complete the proof of the statement, it suffices to show that there is a surjection from  $R/\Sigma(\Theta; R)(+\delta)$  to  $(\omega_R/\Sigma(\Theta; \omega_R))^\vee$ . By Theorem 2.3(ii), there is an injection

$$\omega_R/\Sigma(\Theta; \omega_R) \rightarrow H_m^d(\omega_R)(-\delta).$$

Dualizing and using Lemma 2.2(iii), we obtain a surjection

$$R(+\delta) \cong (H_m^d(\omega_R)(-\delta))^\vee \rightarrow (\omega_R/\Sigma(\Theta; \omega_R))^\vee. \tag{2}$$

Since  $\omega_R$  is an  $R$ -module, it follows from the definition of  $\Sigma(\Theta; \omega_R)$  that

$$\Sigma(\Theta; R) \cdot \omega_R \subseteq \Sigma(\Theta; \omega_R).$$



Hence

$$\Sigma(\Theta; R) \cdot (\omega_R / \Sigma(\Theta; \omega_R)) = 0,$$

which in turn implies

$$\Sigma(\Theta; R) \cdot (\omega_R / \Sigma(\Theta; \omega_R))^\vee = 0. \tag{3}$$

Now (2) and (3) put together guarantee the existence of a surjection

$$(R / \Sigma(\Theta; R))(+\delta) \rightarrow (\omega_R / \Sigma(\Theta; \omega_R))^\vee,$$

as desired. □

**Remark 2.4.** The above proof also works in the local setting: it shows that if  $R$  is a Buchsbaum Noetherian local ring with depth  $R \geq 2$  (and if the canonical module of  $R$  exists), then  $\omega_R / \Sigma(\Theta; \omega_R)$  is isomorphic to the Matlis dual of  $R / \Sigma(\Theta; R)$ .

### 3. Duality in Stanley–Reisner rings of manifolds

In this section, we study Buchsbaum Stanley–Reisner rings and modules. Some objects in this section such as homology groups, Betti numbers, and  $h'$ - and  $h''$ -numbers depend on the characteristic of  $\mathbb{F}$ ; however, we fix a field  $\mathbb{F}$  throughout this section and omit  $\mathbb{F}$  from our notation.

We start by reviewing basics of simplicial complexes and Stanley–Reisner rings. A *simplicial complex*  $\Delta$  on  $[n]$  is a collection of subsets of  $[n]$  that is closed under inclusion. A *relative simplicial complex*  $\Psi$  on  $[n]$  is a collection of subsets of  $[n]$  with the property that there are simplicial complexes  $\Delta \supseteq \Gamma$  such that  $\Psi = \Delta \setminus \Gamma$ . We identify such a pair of simplicial complexes  $(\Delta, \Gamma)$  with the relative simplicial complex  $\Delta \setminus \Gamma$ . Also, a simplicial complex  $\Delta$  will be identified with  $(\Delta, \emptyset)$ . A *face* of  $(\Delta, \Gamma)$  is an element of  $\Delta \setminus \Gamma$ . The *dimension* of a face  $\tau$  is its cardinality minus one, and the dimension of  $(\Delta, \Gamma)$  is the maximal dimension of its faces. A relative simplicial complex is said to be *pure* if all its maximal faces have the same dimension.

We denote by  $\tilde{H}_i(\Delta, \Gamma)$  the  $i$ -th reduced homology group of the pair  $(\Delta, \Gamma)$  computed with coefficients in  $\mathbb{F}$ : when  $\Gamma \neq \emptyset$ ,  $\tilde{H}_*(\Delta, \Gamma)$  is the usual relative homology of a pair, and when  $\Gamma = \emptyset$ ,  $\tilde{H}_*(\Delta, \Gamma) = \tilde{H}_*(\Delta)$  is the reduced homology of  $\Delta$ . The Betti numbers of  $(\Delta, \Gamma)$  are defined by  $\tilde{\beta}_i(\Delta, \Gamma) := \dim_{\mathbb{F}} \tilde{H}_i(\Delta, \Gamma)$ . If  $\Delta$  is a simplicial complex on  $[n]$  and  $\tau \in \Delta$  is a face of  $\Delta$ , then the *link* of  $\tau$  in  $\Delta$  is

$$\text{lk}_\Delta(\tau) := \{\sigma \in \Delta : \tau \cup \sigma \in \Delta, \tau \cap \sigma = \emptyset\}.$$

For convenience, we also define  $\text{lk}_\Delta(\tau) = \emptyset$  if  $\tau \notin \Delta$ .

Let  $\Delta$  be a simplicial complex on  $[n]$ . The Stanley–Reisner ideal of  $\Delta$  (in  $S$ ) is the ideal

$$I_\Delta = (x^\tau : \tau \subseteq [n], \tau \notin \Delta) \subseteq S,$$

where  $x^\tau = \prod_{i \in \tau} x_i$ . If  $(\Delta, \Gamma)$  is a relative simplicial complex, then the *Stanley–Reisner module* of  $(\Delta, \Gamma)$  is the  $S$ -module

$$\mathbb{F}[\Delta, \Gamma] = I_\Gamma/I_\Delta.$$

When  $\Gamma = \emptyset$ , the ring  $\mathbb{F}[\Delta] = \mathbb{F}[\Delta, \emptyset] = S/I_\Delta$  is called the *Stanley–Reisner ring* of  $\Delta$ .

A relative simplicial complex  $(\Delta, \Gamma)$  is said to be *Buchsbaum* if  $\mathbb{F}[\Delta, \Gamma]$  is a Buchsbaum module. The following characterization of the Buchsbaum property was given in [Schenzel 1981, Theorem 3.2]; a proof for relative simplicial complexes appears in [Adiprasito and Sanyal 2016, Theorem 1.11].

**Theorem 3.1.** *A pure relative simplicial complex  $(\Delta, \Gamma)$  of dimension  $d$  is Buchsbaum if and only if*

$$\tilde{H}_i(\text{lk}_\Delta(\tau), \text{lk}_\Gamma(\tau)) = 0$$

for every nonempty face  $\tau \in \Delta \setminus \Gamma$  and all  $i \neq d - |\tau|$ .

In particular, if  $\Delta$  is a homology manifold with boundary, then  $\Delta$  and  $(\Delta, \partial\Delta)$  are Buchsbaum (relative) simplicial complexes. (However, most Buchsbaum complexes are not homology manifolds.)

Next, we discuss face numbers of Buchsbaum simplicial complexes. For a relative simplicial complex  $(\Delta, \Gamma)$  of dimension  $d - 1$ , let  $f_i(\Delta, \Gamma)$  be the number of  $i$ -dimensional faces of  $(\Delta, \Gamma)$  and let

$$h_j(\Delta, \Gamma) = \sum_{i=0}^j (-1)^{j-i} \binom{d-i}{d-j} f_{i-1}(\Delta, \Gamma) \quad \text{for } j = 0, 1, \dots, d.$$

For convenience, we also define  $h_j(\Delta, \Gamma) = 0$  for  $j > \dim(\Delta, \Gamma) + 1$ . The  $h$ -numbers play a central role in the study of face numbers of Cohen–Macaulay simplicial complexes. On the other hand, for Buchsbaum simplicial complexes, the following modifications of  $h$ -numbers, called  $h'$ -numbers and  $h''$ -numbers, behave better than the usual  $h$ -numbers.

Recall that a *linear system of parameters* (or l.s.o.p.) is an h.s.o.p.  $\Theta = \theta_1, \dots, \theta_d$  consisting of linear forms. Note that when  $\mathbb{F}$  is infinite, any finitely generated graded  $S$ -module has an l.s.o.p. The following result, established in [Schenzel 1981, §4] (a proof for Stanley–Reisner modules appears in [Adiprasito and Sanyal 2016, Theorem 2.5]), is known as Schenzel’s formula.

**Theorem 3.2.** *Let  $(\Delta, \Gamma)$  be a Buchsbaum relative simplicial complex of dimension  $d - 1$  and let  $\Theta$  be an l.s.o.p. for  $\mathbb{F}[\Delta, \Gamma]$ . Then, for  $j = 0, 1, \dots, d$ ,*

$$\dim_{\mathbb{F}}(\mathbb{F}[\Delta, \Gamma]/\Theta\mathbb{F}[\Delta, \Gamma])_j = h_j(\Delta, \Gamma) - \binom{d}{j} \sum_{i=1}^{j-1} (-1)^{j-i} \tilde{\beta}_{i-1}(\Delta, \Gamma).$$

In view of Schenzel’s formula, we define the  $h'$ -numbers of a  $(d - 1)$ -dimensional relative simplicial complex  $(\Delta, \Gamma)$  by

$$h'_j(\Delta, \Gamma) = h_j(\Delta, \Gamma) - \binom{d}{j} \sum_{i=1}^{j-1} (-1)^{j-i} \tilde{\beta}_{i-1}(\Delta, \Gamma).$$

Furthermore, we define the  $h''$ -numbers of  $(\Delta, \Gamma)$  by

$$h''_j(\Delta, \Gamma) = \begin{cases} h'_j(\Delta, \Gamma) - \binom{d}{j} \tilde{\beta}_{j-1}(\Delta, \Gamma) & \text{if } 0 \leq j < d, \\ h'_d(\Delta, \Gamma) & \text{if } j = d. \end{cases}$$

We are now ready to prove Theorem 1.2.

*Proof of Theorem 1.2.* Let  $M = \mathbb{F}[\Delta, \Gamma]$ . Observe that

$$\dim_{\mathbb{F}}(M/\Sigma(\Theta; M))_j = \dim_{\mathbb{F}}(M/\Theta M)_j - \dim_{\mathbb{F}}(\Sigma(\Theta; M)/\Theta M)_j.$$

Since  $H_m^i(M) = (H_m^i(M))_0 \cong \tilde{H}_{i-1}(\Delta, \Gamma)$  for  $i < d$  (see [Adiprasito and Sanyal 2016, Theorem 1.8]), Theorem 2.3(i) implies

$$\dim_{\mathbb{F}}(\Sigma(\Theta; M)/\Theta M)_j = \begin{cases} \binom{d}{j} \tilde{\beta}_{j-1}(\Delta, \Gamma) & \text{if } 0 \leq j \leq d - 1, \\ 0 & \text{if } j \geq d. \end{cases}$$

The statement then follows from Theorem 3.2, asserting that  $\dim_{\mathbb{F}}(M/\Theta M)_j = h'_j(\Delta, \Gamma)$  for all  $j$ . □

It was proved in [Novik and Swartz 2009b, Theorem 3.4] that the  $j$ -th graded component of the socle of  $\mathbb{F}[\Delta]/\Theta\mathbb{F}[\Delta]$  has dimension at least  $\binom{d}{j} \tilde{\beta}_{j-1}(\Delta)$ . This implies that the  $h''$ -numbers of a Buchsbaum simplicial complex form the Hilbert function of some quotient of its Stanley–Reisner ring. A new contribution and the significance of Theorem 1.2 is that it provides an explicit algebraic interpretation of  $h''$ -numbers via a submodule  $\Sigma(\Theta; -)$ .

In the rest of this section we discuss a few algebraic and combinatorial applications of our results. As was proved by Kalai and, independently, Stanley [1993; 1996, Chapter III, §9], if  $\Delta \supseteq \Gamma$  are Cohen–Macaulay simplicial complexes of the same dimension, then  $h_i(\Delta) \geq h_i(\Gamma)$  for all  $i$ . The interpretation of the  $h''$ -numbers given in Theorem 1.2 allows us to prove the following generalization of this fact.

**Theorem 3.3.** *Let  $\Delta \supseteq \Gamma$  be Buchsbaum simplicial complexes of the same dimension. Then  $h''_i(\Delta) \geq h''_i(\Gamma)$  for all  $i$ .*

*Proof.* We may assume that  $\Delta$  and  $\Gamma$  are simplicial complexes on  $[n]$  and that  $\mathbb{F}$  is infinite. Let  $d = \dim \Delta + 1$ . Then there is a common linear system of parameters

$\Theta = \theta_1, \dots, \theta_d$  for  $\mathbb{F}[\Delta]$  and  $\mathbb{F}[\Gamma]$ . By the definition of  $\Sigma(\Theta; -)$ ,

$$\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]) = S / \left( (\Theta) + \sum_{k=1}^d ((\theta_1, \dots, \hat{\theta}_k, \dots, \theta_d) + I_\Delta) :_S \theta_k \right),$$

and an analogous formula holds for  $\mathbb{F}[\Gamma]/\Sigma(\Theta; \mathbb{F}[\Gamma])$ . Since  $I_\Gamma \supseteq I_\Delta$ , the above formula implies that  $\mathbb{F}[\Gamma]/\Sigma(\Theta; \mathbb{F}[\Gamma])$  is a quotient ring of  $\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta])$ . The desired statement then follows from Theorem 1.2.  $\square$

**Corollary 3.4.** *Let  $\Delta$  be a Buchsbaum simplicial complex and let  $\tau \in \Delta$  be any nonempty face. Then  $h''_i(\Delta) \geq h_i(\text{lk}_\Delta(\tau))$  for all  $i$ .*

*Proof.* Consider the star of  $\tau$ ,  $\text{st}_\Delta(\tau) = \{\sigma \in \Delta : \sigma \cup \tau \in \Delta\}$ . Then  $\text{st}_\Delta(\tau)$  is Cohen–Macaulay (by Theorem 3.1 and Reisner’s criterion) and has the same dimension as  $\Delta$ , and so by Theorem 3.3 we have  $h''_i(\Delta) \geq h''_i(\text{st}_\Delta(\tau))$  for all  $i$ . The result follows since  $\text{lk}_\Delta(\tau)$  and  $\text{st}_\Delta(\tau)$  have the same  $h$ -numbers (this is because  $\text{st}_\Delta(\tau)$  is just a cone over  $\text{lk}_\Delta(\tau)$ ; cf. [Stanley 1996, Corollary III.9.2]) and since for Cohen–Macaulay simplicial complexes the  $h''$ -numbers coincide with the  $h$ -numbers.  $\square$

Let  $R = S/I$  be a graded  $\mathbb{F}$ -algebra. The  $a$ -invariant of  $R$  is the number

$$a(R) = -\min\{k : (\omega_R)_k \neq 0\}.$$

This number is an important invariant in commutative algebra. When  $R$  is Cohen–Macaulay, it is well-known that  $a(R) = \max\{i : h_i(R) \neq 0\} - \dim R$ , where  $h_i(R)$  is the  $i$ -th  $h$ -number of  $R$ . (See [Bruns and Herzog 1993, §4.1] for the definition of  $h$ -numbers for modules.) The following result provides a generalization of this fact.

**Theorem 3.5.** *Let  $R = S/I$  be a Buchsbaum graded  $\mathbb{F}$ -algebra of Krull dimension  $d$  with  $\text{depth}(R) \geq 2$ , and let  $\Theta = \theta_1, \dots, \theta_d$  be an l.s.o.p. for  $R$ . Then*

$$a(R) = \max\{k : (R/\Sigma(\Theta; R))_k \neq 0\} - d.$$

*In particular, for any connected Buchsbaum simplicial complex  $\Delta$  of dimension  $d - 1$ , we have  $a(\mathbb{F}[\Delta]) = \max\{k : h''_k(\Delta) \neq 0\} - d$ .*

*Proof.* Let  $m = \max\{k : (R/\Sigma(\Theta; R))_k \neq 0\}$ . Then by Theorem 1.3,

$$\min\{k : (\omega_R/\Sigma(\Theta; \omega_R))_k \neq 0\} = d - m.$$

As  $\min\{k : (\omega_R)_k \neq 0\} = \min\{k : (\omega_R/\Theta\omega_R)_k \neq 0\}$ , the theorem would follow if we prove that  $\min\{k : (\omega_R/\Theta\omega_R)_k \neq 0\} = d - m$ . To this end, it is enough to show that

$$(\Sigma(\Theta; \omega_R)/\Theta\omega_R)_k = 0 \quad \text{for } k \leq d - m - 1. \tag{4}$$

Since  $\text{m}H_m^i(R) = 0$  for  $i < d$  (see [Stückrad and Vogel 1986, Proposition I.2.1]), we conclude from Theorem 2.3(i) that  $\text{m}(\Sigma(\Theta; R)/\Theta R) = 0$ . Furthermore, since

$(R/\Theta R)_k = (\Sigma(\Theta; R)/\Theta R)_k$  for  $k \geq m + 1$ , it follows that, for  $k \geq m + 2$ ,

$$(\Sigma(\Theta; R)/\Theta R)_k = (R/\Theta R)_k = (\mathfrak{m}(R/\Theta R))_k = (\mathfrak{m}(\Sigma(\Theta; R)/\Theta R))_k = 0.$$

The isomorphism

$$\Sigma(\Theta; R)/\Theta R \cong \bigoplus_{C \subsetneq [d]} H_m^{|C|}(R)(-|C|)$$

established in Theorem 2.3(i) then implies that

$$H_m^i(R)_j = 0 \quad \text{for all } i \leq d - 1, j \geq m + 2 - i.$$

Therefore,

$$(H_m^{d-i+1}(R)^\vee(-i))_k = 0 \quad \text{for all } 2 \leq i \leq d - 1, -(k - i) \geq m + 2 - (d - i + 1),$$

and so

$$(H_m^{d-i+1}(R)^\vee(-i))_k = 0 \quad \text{for all } 2 \leq i \leq d - 1, k \leq d - m - 1. \tag{5}$$

Finally, since  $\text{depth}(\omega_R) \geq 2$ , we infer from Theorem 2.3(i) and Lemma 2.2(ii) that

$$\Sigma(\Theta; \omega_R)/\Theta \omega_R \cong \bigoplus_{C \subsetneq [d], |C| \geq 2} H_m^{|C|}(\omega_R)(-|C|) \cong \bigoplus_{C \subsetneq [d], |C| \geq 2} H_m^{d-|C|+1}(R)^\vee(-|C|).$$

The above isomorphisms and (5) then yield the desired property (4). □

Observe that, by Hochster’s formula on local cohomology [Bruns and Herzog 1993, Theorem 5.3.8], if  $\Delta$  is a  $(d - 1)$ -dimensional Buchsbaum simplicial complex, then  $-a(\mathbb{F}[\Delta])$  equals the minimum cardinality of a face whose link has a nonvanishing top homology. Thus the “in particular” part of Theorem 3.5 can be equivalently restated as follows: if, for every face  $\tau \in \Delta$  of dimension  $< d - k$ , the link of  $\tau$  has vanishing top homology, then  $h_k''(\Delta) = 0$ . For the case of homology manifolds with boundary, faces with vanishing top homology are precisely the boundary faces; in this case, the above statement reduces to [Murai and Nevo 2014, Theorem 3.1].

A sequence  $h_0, h_1, \dots, h_m$  of numbers is said to be *unimodal* if there is an index  $p$  such that  $h_0 \leq h_1 \leq \dots \leq h_p \geq \dots \geq h_m$ . It was proved in [Murai 2010] that the  $h''$ -numbers of the barycentric subdivision of any connected Buchsbaum simplicial complex form a unimodal sequence. Here we use the Matlis duality established in Theorem 1.3 to generalize this result to Buchsbaum polyhedral complexes. (We refer our readers to [Murai 2010] for the definition of barycentric subdivisions.)

**Theorem 3.6.** *Let  $\Delta$  be the barycentric subdivision of a connected polyhedral complex  $\Gamma$  of dimension  $d - 1$ . Suppose that the characteristic of  $\mathbb{F}$  is zero and*

$\Delta$  is Buchsbaum. Then, for a generic choice of linear forms  $\Theta = \theta_1, \dots, \theta_d$ , and  $\theta_{d+1} \in \mathbb{F}[\Delta]$ , the multiplication

$$\times_{\theta_{d+1}} : (\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]))_i \rightarrow (\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]))_{i+1}$$

is injective for  $i \leq \frac{d}{2} - 1$  and is surjective for  $i \geq \frac{d}{2}$ . In particular, the sequence  $h''_0(\Delta), h''_1(\Delta), \dots, h''_d(\Delta)$  is unimodal.

*Proof.* By genericity of linear forms,  $\Theta = \theta_1, \dots, \theta_d$  is a common l.s.o.p. for  $\mathbb{F}[\Delta]$  and  $\omega_{\mathbb{F}[\Delta]}$ . Let  $P$  be the face poset of  $\Gamma$ . Then the Stanley–Reisner ring  $\mathbb{F}[\Delta]$  is a squarefree  $P$ -module (a notion introduced in [Murai and Yanagawa 2014, Definition 2.1]); furthermore, by [Murai and Yanagawa 2014, Theorem 3.1]  $\omega_{\mathbb{F}[\Delta]}$  is also a squarefree  $P$ -module. It then follows from [Murai and Yanagawa 2014, Theorem 6.2(ii)] that the multiplication maps

$$\times_{\theta_{d+1}} : (\mathbb{F}[\Delta]/\Theta\mathbb{F}[\Delta])_i \rightarrow (\mathbb{F}[\Delta]/\Theta\mathbb{F}[\Delta])_{i+1} \tag{6}$$

and

$$\times_{\theta_{d+1}} : (\omega_{\mathbb{F}[\Delta]}/(\Theta\omega_{\mathbb{F}[\Delta]}))_i \rightarrow (\omega_{\mathbb{F}[\Delta]}/(\Theta\omega_{\mathbb{F}[\Delta]}))_{i+1}$$

are surjective for  $i \geq \frac{d}{2}$ . (While Cohen–Macaulayness was assumed in [Murai and Yanagawa 2014, Theorem 6.2], one can see from their proof of the theorem that this assumption was used only in the proof of part (i) and is unnecessary to derive surjectivity.) Since  $\Theta\mathbb{F}[\Delta]$  is contained in  $\Sigma(\Theta; \mathbb{F}[\Delta])$ , the map in (6) remains surjective if we replace  $\Theta\mathbb{F}[\Delta]$  with  $\Sigma(\Theta; \mathbb{F}[\Delta])$ , and a similar statement holds for  $\omega_{\mathbb{F}[\Delta]}$ . These surjectivities and the Matlis duality  $\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]) \cong (\omega_{\mathbb{F}[\Delta]}/\Sigma(\Theta; \omega_{\mathbb{F}[\Delta]}))^\vee(d)$  of Theorem 1.3 yield the desired statement.  $\square$

In fact, in view of results from [Juhnke-Kubitzke and Murai 2015], it is tempting to conjecture that if  $\Delta$  is a barycentric subdivision of a Buchsbaum regular CW-complex of dimension  $d - 1$ , then even the sequence

$$h''_0(\Delta)/\binom{d}{0}, h''_1(\Delta)/\binom{d}{1}, h''_2(\Delta)/\binom{d}{2}, \dots, h''_d(\Delta)/\binom{d}{d}$$

is unimodal.

We close this section with a couple of remarks.

**Remark 3.7.** Our results on  $h''$ -vectors can be generalized to the following setting. Consider a finitely generated graded  $S$ -module  $M$  of Krull dimension  $d$  such that

$$H_m^i(M) = (H_m^i(M))_0 \quad \text{for all } 0 \leq i < d. \tag{7}$$

Define the  $h'$ - and  $h''$ -numbers of  $M$  in the same way as for the Stanley–Reisner modules but with  $\dim_{\mathbb{F}}(H_m^j(M))$  used as a replacement for  $\tilde{\beta}_{j-1}(\Delta, \Gamma)$ . Then suitably modified statements of Theorems 1.2 and 3.2 continue to hold for such an  $M$ . Furthermore, if  $M$  satisfies (7), then  $M$  must be Buchsbaum (see [Stückrad and Vogel

1986, Proposition I.3.10]), in which case  $\omega_M$  also satisfies (7) by Lemma 2.2(ii). In particular, if  $\Delta$  is an arbitrary connected Buchsbaum simplicial complex of dimension  $d - 1$ , then

$$h''_i(\omega_{\mathbb{F}[\Delta]}) = h''_{d-i}(\mathbb{F}[\Delta]) \quad \text{for all } i = 0, 1, \dots, d.$$

**Remark 3.8.** A statement analogous to Corollary 1.4 also holds for homology manifolds without boundary. Indeed, if  $\Delta$  is an orientable homology manifold without boundary, then by Gräbe’s result [1984],  $\mathbb{F}[\Delta]$  is isomorphic to its own canonical module, and hence, by Theorem 1.3,  $\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta])$  is an Artinian Gorenstein algebra. This fact was essentially proved in [Novik and Swartz 2009a, Theorem 1.4].

Let us also point out that if  $\Delta$  is a connected orientable homology manifold with or without boundary, then for  $M = \mathbb{F}[\Delta]$  the statement of part (ii) of Lemma 2.2 is a simple consequence of the Poincaré–Alexander–Lefschetz duality along with Gräbe’s result [1984] that  $\omega_M \cong \mathbb{F}[\Delta, \partial\Delta]$ .

#### 4. Applications to the manifold $g$ -conjecture

In this section we discuss connected orientable homology manifolds without boundary. One of the most important open problems in algebraic combinatorics is the algebraic  $g$ -conjecture; it asserts that every homology sphere has the weak Lefschetz property (the WLP, for short). Kalai proposed [Novik 1998, Conjecture 7.5] a far-reaching generalization of this conjecture to homology manifolds. Using the  $\Sigma(\Theta, -)$  module allows us to restate Kalai’s conjecture as follows.

**Conjecture 4.1.** *Let  $\Delta$  be a connected orientable  $\mathbb{F}$ -homology  $(d - 1)$ -manifold without boundary. Then, for a generic choice of linear forms  $\Theta = \theta_1, \dots, \theta_d$ , the ring  $\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta])$  has the WLP; that is, for a generic linear form  $\omega$ , the multiplication*

$$\times \omega : (\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]))_{\lfloor d/2 \rfloor} \rightarrow (\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta]))_{\lfloor d/2 \rfloor + 1}$$

*is surjective.*

It is worth mentioning that while Kalai’s original statement of the conjecture did not involve  $\Sigma(\Theta, -)$ , the two statements are equivalent (see [Novik and Swartz 2009a] and Remark 3.8 above). Somewhat informally, we say that  $\Delta$  (or  $\mathbb{F}[\Delta]$ ) has the WLP if  $\Delta$  satisfies the conclusions of the above conjecture. Enumerative consequences of this conjecture are discussed in [Novik and Swartz 2009a, §1].

Given a finite set  $A$ , we denote by  $\bar{A}$  the simplex on  $A$ , i.e., the simplicial complex whose set of faces consists of all subsets of  $A$ . When  $A = \{a\}$  consists of a single vertex, we write  $\bar{a}$  to denote the vertex  $a$ , viewed as a 0-dimensional simplex. If

$\Delta$  and  $\Gamma$  are two simplicial complexes on disjoint vertex sets, then the *join* of  $\Delta$  and  $\Gamma$ ,  $\Delta * \Gamma$ , is the simplicial complex defined by

$$\Delta * \Gamma := \{\sigma \cup \tau : \sigma \in \Delta, \tau \in \Gamma\}.$$

Finally, if  $\Delta$  is a simplicial complex and  $W$  is a set, then  $\Delta_W = \{\sigma \in \Delta : \sigma \subseteq W\}$  is the subcomplex of  $\Delta$  induced by  $W$ .

Let  $\Delta$  be a  $(d - 1)$ -dimensional homology manifold, and let  $A$  and  $B$  be disjoint subsets such that  $|A| + |B| = d + 1$ . If  $\Delta_{A \cup B} = \bar{A} * \partial \bar{B}$ , then the operation of removing  $\bar{A} * \partial \bar{B}$  from  $\Delta$  and replacing it with  $\partial \bar{A} * \bar{B}$  is called a  $(|B| - 1)$ -*bistellar flip*. (For instance, a 0-flip is simply a stellar subdivision at a facet.) The resulting complex is a homology manifold homeomorphic to the original complex.

The following surprising property was proved by Pachner [1987; 1991]: if  $\Delta_1$  and  $\Delta_2$  are two PL homeomorphic combinatorial manifolds without boundary, then they can be connected by a sequence of bistellar flips. Since the boundary complex of a simplex has the WLP, Pachner's result suggests the following inductive approach to the algebraic  $g$ -conjecture for PL spheres: prove that bistellar flips applied to PL spheres preserve the WLP. Swartz showed [2014, §3] that most bistellar flips applied to homology *spheres* preserve the WLP. Here we extend Swartz's results to the generality of orientable homology manifolds without boundary:

**Theorem 4.2.** *Let  $\Delta$  be a  $(d - 1)$ -dimensional, connected, orientable homology manifold without boundary. Suppose  $\Delta'$  is obtained from  $\Delta$  via a  $(p - 1)$ -bistellar flip with  $p \neq \frac{1}{2}(d + 1)$  if  $d$  is odd and with  $p \notin \{\frac{d}{2}, \frac{1}{2}(d + 2)\}$  if  $d$  is even, and let  $\Theta$  be a common l.s.o.p. for  $\mathbb{F}[\Delta]$  and  $\mathbb{F}[\Delta']$ . Then  $\mathbb{F}[\Delta'] / \Sigma(\Theta; \mathbb{F}[\Delta'])$  has the WLP if and only if  $\mathbb{F}[\Delta] / \Sigma(\Theta; \mathbb{F}[\Delta])$  has the WLP.*

If  $\Delta$  is a simplicial complex and  $\sigma$  is a face of  $\Delta$ , then the *stellar subdivision* of  $\Delta$  at  $\sigma$  consists of (i) removing  $\sigma$  and all faces containing it from  $\Delta$ , (ii) introducing a new vertex  $a$ , and (iii) adding new faces in  $\bar{a} * \partial \bar{\sigma} * \text{lk}_\Delta(\sigma)$  to  $\Delta$ :

$$\text{sd}_\sigma(\Delta) := (\Delta \setminus \text{st}_\Delta(\sigma)) \cup (\bar{a} * \partial \bar{\sigma} * \text{lk}_\Delta(\sigma)).$$

A classical result due to Alexander [1930] asserts that two simplicial complexes are PL homeomorphic if and only if they are stellar equivalent; that is, one of them can be obtained from another by a sequence of stellar subdivisions and their inverses. Thus, a different approach to the algebraic  $g$ -conjecture (at least for PL manifolds) is to show that the WLP is preserved by stellar subdivisions and their inverses. Böhm and Papadakis proved [2015, Corollary 4.5] that this is the case if one applies stellar subdivisions at faces of sufficiently large dimension (or their inverses) to homology spheres. We extend their result to the generality of orientable homology manifolds without boundary:



**Theorem 4.3.** *Let  $\Delta$  be a  $(d - 1)$ -dimensional, connected, orientable  $\mathbb{F}$ -homology manifold without boundary and let  $\sigma$  be a face of  $\Delta$  with  $\dim \sigma > \frac{d}{2}$ . Then  $\mathbb{F}[\Delta]$  has the WLP if and only if  $\mathbb{F}[\text{sd}_\sigma(\Delta)]$  has the WLP.*

The structure of the proofs of both theorems is similar to the proof of [Swartz 2014, Theorem 3.1]. The new key ingredient is given by the following lemma. Recall that  $\Delta$  is an  $\mathbb{F}$ -homology  $(d - 1)$ -sphere if  $\Delta$  is an  $\mathbb{F}$ -homology manifold whose homology over  $\mathbb{F}$  coincides with that of  $\mathbb{S}^{d-1}$ . Similarly,  $\Delta$  is an  $\mathbb{F}$ -homology ball if (i)  $\Delta$  is an  $\mathbb{F}$ -homology manifold with boundary, (ii) the homology of  $\Delta$  (over  $\mathbb{F}$ ) vanishes, and (iii) the boundary of  $\Delta$  is an  $\mathbb{F}$ -homology sphere.

**Lemma 4.4.** *Let  $\Delta$  be a  $(d - 1)$ -dimensional  $\mathbb{F}$ -homology manifold without boundary,  $\Gamma$  a full-dimensional subcomplex of  $\Delta$ , and  $\Theta$  an l.s.o.p. for  $\mathbb{F}[\Delta]$ . Assume further that  $\Gamma$  is an  $\mathbb{F}$ -homology ball, and let  $D$  be the simplicial complex obtained from  $\Delta$  by removing the interior faces of  $\Gamma$ . Then  $D$  is an  $\mathbb{F}$ -homology manifold with boundary and the natural surjection  $\mathbb{F}[\Delta] \rightarrow \mathbb{F}[\Gamma]$  induces the short exact sequence*

$$0 \rightarrow \mathbb{F}[D, \partial D] / \Sigma(\Theta; \mathbb{F}[D, \partial D]) \rightarrow \mathbb{F}[\Delta] / \Sigma(\Theta; \mathbb{F}[\Delta]) \rightarrow \mathbb{F}[\Gamma] / (\Theta \mathbb{F}[\Gamma]) \rightarrow 0.$$

*Proof.* The fact that  $D$  is a homology manifold with boundary follows by a standard Mayer–Vietoris argument (note that  $\partial D = \partial \Gamma$  is a homology  $(d - 2)$ -sphere). Furthermore, by excision and since  $\Gamma$  has vanishing homology,

$$\tilde{\beta}_i(D, \partial D) = \tilde{\beta}_i(\Delta, \Gamma) = \tilde{\beta}_i(\Delta) \quad \text{for all } i. \tag{8}$$

Now, using that the homology ball  $\Gamma$  is a full-dimensional subcomplex of  $\Delta$ , we conclude as in the proof of Theorem 3.3 that there is a natural surjection

$$\phi : \mathbb{F}[\Delta] / \Sigma(\Theta; \mathbb{F}[\Delta]) \rightarrow \mathbb{F}[\Gamma] / (\Theta \mathbb{F}[\Gamma]) = \mathbb{F}[\Gamma] / \Sigma(\Theta; \mathbb{F}[\Gamma]). \tag{9}$$

Thus, to finish the proof of the lemma, it suffices to show that the kernel of  $\phi$  is  $\mathbb{F}[D, \partial D] / \Sigma(\Theta; \mathbb{F}[D, \partial D])$ . This, in turn, would follow if we verify that

- (i)  $\text{Ker}(\phi)$  is a quotient of  $\mathbb{F}[D, \partial D] / \Sigma(\Theta; \mathbb{F}[D, \partial D])$ , and
- (ii)  $\dim_{\mathbb{F}}(\text{Ker}(\phi))_j = \dim_{\mathbb{F}}(\mathbb{F}[D, \partial D] / \Sigma(\Theta; \mathbb{F}[D, \partial D]))_j$  for all  $j$ .

For brevity, let  $R = \mathbb{F}[\Delta]$  and let  $I = \mathbb{F}[D, \partial D]$ . Since  $(D, \partial D) = (\Delta, \Gamma)$  as relative simplicial complexes, it follows that  $I$  is an ideal of  $R$ , and  $R/I$  is  $\mathbb{F}[\Gamma]$ . Thus our surjection  $\phi$  is the projection  $\phi : R / \Sigma(\Theta; R) \rightarrow R / (I + \Theta R)$ . Hence

$$\text{Ker}(\phi) = (I + \Theta R) / \Sigma(\Theta; R) = (I + \Sigma(\Theta; R)) / \Sigma(\Theta; R) = I / (I \cap \Sigma(\Theta; R)),$$

which together with an observation that  $I \cap \Sigma(\Theta; R)$  contains  $\Sigma(\Theta; I)$  yields assertion (i). To see that  $I \cap \Sigma(\Theta; R) \supseteq \Sigma(\Theta; I)$ , note that, since  $I$  is a subset of  $R$ ,  $\Theta I$  is contained in  $\Theta R$ , and  $(\theta_1, \dots, \hat{\theta}_i, \dots, \theta_d) I :_I \theta_i$  is contained in  $(\theta_1, \dots, \hat{\theta}_i, \dots, \theta_d) R :_R \theta_i$  for all  $i \in [d]$ ; therefore  $\Sigma(\Theta; R) \supseteq \Sigma(\Theta; I)$ .

As for assertion (ii), the dimension of the kernel of  $\phi$  can be computed as follows: by (9) and Theorem 1.2,

$$\dim_{\mathbb{F}}(\text{Ker}(\phi))_j = h_j''(\Delta) - h_j(\Gamma). \tag{10}$$

Now, since each face of  $\Delta$  is either a face of  $(D, \partial D)$  or a face of  $\Gamma$  (but not of both),  $f_i(\Delta) = f_i(D, \partial D) + f_i(\Gamma)$  for all  $i \geq -1$ , and so

$$h_j(D, \partial D) = h_j(\Delta) - h_j(\Gamma) \quad \text{for all } j \geq 0. \tag{11}$$

Equations (8), (10) and (11) yield

$$\dim_{\mathbb{F}}(\text{Ker}(\phi))_j = h_j''(D, \partial D) = \dim_{\mathbb{F}}(\mathbb{F}[D, \partial D]/\Sigma(\Theta; \mathbb{F}[D, \partial D]))_j \quad \text{for all } j \geq 0,$$

where the last equality is another application of Theorem 1.2. The lemma follows.  $\square$

Another ingredient needed for both proofs is the following immediate consequence of the snake lemma.

**Lemma 4.5.** *Let  $0 \rightarrow L \rightarrow N \rightarrow M \rightarrow 0$  be an exact sequence of graded  $S$ -modules, let  $\omega \in S$  be a linear form, and let  $k$  be a fixed integer. Assume also that the map  $\times \omega : M_k \rightarrow M_{k+1}$  is bijective. Then the map  $\times \omega : N_k \rightarrow N_{k+1}$  is surjective if and only if the map  $\times \omega : L_k \rightarrow L_{k+1}$  is surjective.*

We are now ready to prove both of the theorems.

*Proof of Theorem 4.2.* We are given that  $\Delta' = (\Delta \setminus (\bar{A} * \partial \bar{B})) \cup (\partial \bar{A} * \bar{B})$ , where  $|B| = p$ . Let  $D$  be the simplicial complex obtained from  $\Delta$  by removing the interior faces of the ball  $\Gamma_1 = \bar{A} * \partial \bar{B}$ ; equivalently,  $D$  is obtained from  $\Delta'$  by removing the interior faces of  $\Gamma_2 = \partial \bar{A} * \bar{B}$ . Since  $\Gamma_1$  and  $\Gamma_2$  are full-dimensional subcomplexes of  $\Delta$  and  $\Delta'$ , and  $\Theta$  is an l.s.o.p. for both  $\mathbb{F}[\Delta]$  and  $\mathbb{F}[\Delta']$ , it is also an l.s.o.p. for both  $\mathbb{F}[\Gamma_1]$  and  $\mathbb{F}[\Gamma_2]$ . Hence

$$\dim_{\mathbb{F}}(\mathbb{F}[\Gamma_1]/\Theta\mathbb{F}[\Gamma_1])_j = h_j(\bar{A} * \partial \bar{B}) = h_j(\partial \bar{B}) = \begin{cases} 1 & \text{if } j \leq p - 1, \\ 0 & \text{if } j > p - 1, \end{cases}$$

which implies that  $\mathbb{F}[\Gamma_1]/\Theta\mathbb{F}[\Gamma_1] \cong \mathbb{F}[x]/(x^p)$ . Similarly, we have  $\mathbb{F}[\Gamma_2]/\Theta\mathbb{F}[\Gamma_2] \cong \mathbb{F}[x]/(x^{d-p+1})$ . These isomorphisms together with the assumption  $p \notin \{\frac{d}{2}, \frac{d+1}{2}, \frac{d+2}{2}\}$  yield that, for a generic choice of a linear form  $w$ , the multiplication map

$$\times w : (\mathbb{F}[\Gamma_i]/\Theta\mathbb{F}[\Gamma_i])_{\lfloor d/2 \rfloor} \rightarrow (\mathbb{F}[\Gamma_i]/\Theta\mathbb{F}[\Gamma_i])_{\lfloor d/2 \rfloor + 1}$$

is bijective for  $i \in \{1, 2\}$ .

Applying Lemma 4.4 to  $\Delta$  and  $\Gamma_1$ , we can conclude from Lemma 4.5 that  $\mathbb{F}[\Delta]/\Sigma(\Theta; \mathbb{F}[\Delta])$  has the WLP if and only if, for a generic linear form  $w$ , the multiplication

$$\times w : (\mathbb{F}[D, \partial D]/\Sigma(\Theta; \mathbb{F}[D, \partial D]))_{\lfloor d/2 \rfloor} \rightarrow (\mathbb{F}[D, \partial D]/\Sigma(\Theta; \mathbb{F}[D, \partial D]))_{\lfloor d/2 \rfloor + 1}$$

is surjective. On the other hand, by Lemma 4.4 applied to  $\Delta'$  and  $\Gamma_2$ , the latter condition is equivalent to the WLP of  $\mathbb{F}[\Delta']/\Sigma(\Theta; \mathbb{F}[\Delta'])$ .  $\square$

*Proof of Theorem 4.3.* Let  $D$  be the homology manifold obtained from  $\Delta$  by removing the interior faces of the homology ball  $\Gamma_1 = \text{st}_\Delta(\sigma)$ ; equivalently,  $D$  is obtained from  $\text{sd}_\sigma(\Delta)$  by removing the interior faces of the homology ball  $\Gamma_2 = \bar{a} * \partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$ . As the proof of Theorem 4.2 shows, to verify Theorem 4.3, it suffices to check that, for a generic choice of linear forms  $\Theta = \theta_1, \dots, \theta_d$  and another generic linear form  $w$ , the multiplication

$$\times w : (\mathbb{F}[\Gamma_i]/\Theta\mathbb{F}[\Gamma_i])_{\lfloor d/2 \rfloor} \rightarrow (\mathbb{F}[\Gamma_i]/\Theta\mathbb{F}[\Gamma_i])_{\lfloor d/2 \rfloor + 1} \tag{12}$$

is bijective for each  $i \in \{1, 2\}$ .

In the case of  $i = 1$ , the desired bijection is an immediate consequence of the fact that

$$\dim_{\mathbb{F}}(\mathbb{F}[\Gamma_1]/\Theta\mathbb{F}[\Gamma_1])_j = h_j(\text{st}_\Delta(\sigma)) = h_j(\text{lk}_\Delta(\sigma)) = 0 \quad \text{for } j \geq d - |\sigma| + 1$$

and the assumption  $|\sigma| > \frac{d}{2} + 1$ . We now prove that the map in (12) is also bijective for  $i = 2$ . To this end, observe that the homology  $(d - 2)$ -sphere  $\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$  is the boundary of the homology  $(d - 1)$ -ball  $\bar{\sigma} * \text{lk}_\Delta(\sigma)$ , and hence  $\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$  is  $(d - |\sigma|)$ -stacked. (A homology  $(d - 2)$ -sphere is called  $(d - k)$ -stacked if it is the boundary of a homology ball that has no interior faces of size  $\leq k - 1$ .) It then follows from [Swartz 2014, Corollary 6.3] that  $\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$  has the WLP. This, in turn, implies that the map in (12) is surjective, as  $\mathbb{F}[\Gamma_2]$  is a polynomial ring over  $\mathbb{F}[\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)]$ . Also, since  $|\sigma| > \frac{d}{2} + 1$ , we infer from [McMullen and Walkup 1971, Theorem 2] and the  $(d - |\sigma|)$ -stackedness of  $\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$  that  $h_{\lfloor d/2 \rfloor}(\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)) = h_{\lfloor d/2 \rfloor + 1}(\partial\bar{\sigma} * \text{lk}_\Delta(\sigma))$ . As  $\Gamma_2$  and  $\partial\bar{\sigma} * \text{lk}_\Delta(\sigma)$  have the same  $h$ -vector, we conclude that  $h_{\lfloor d/2 \rfloor}(\Gamma_2) = h_{\lfloor d/2 \rfloor + 1}(\Gamma_2)$ , and therefore that the map in (12) is bijective.  $\square$

Let  $\Delta$  be a triangulation of a closed surface and assume that one of the vertices of  $\Delta$  is connected to all other vertices of  $\Delta$ . Then by [Novik and Swartz 2009a, Theorem 1.6],  $\Delta$  has the WLP. However, the problem of whether *every* triangulation of a closed surface other than the sphere possesses the WLP is at present wide open.

**Remark 4.6.** Note that if  $\Delta$  is an arbitrary *odd-dimensional* Buchsbaum complex (e.g., a homology manifold), then there exists a simplicial complex  $\Gamma$  that is PL homeomorphic to  $\Delta$  and has the WLP in characteristic 0. Simply take  $\Gamma$  to be the barycentric subdivision of  $\Delta$ : the WLP of  $\Gamma$  is guaranteed by Theorem 3.6.

### Appendix: Proof of Theorem 2.3

The goal of this appendix is to verify Theorem 2.3. Our proof is based on the proof of [Novik and Swartz 2009b, Theorem 2.2], and so some details are omitted.

Let  $M$  be a finitely generated Buchsbaum graded  $S$ -module of Krull dimension  $d$ , and let  $\Theta = \theta_1, \dots, \theta_d$  be an h.s.o.p. for  $M$  with  $\deg \theta_i = \delta_i$ . We write  $\delta_C = \sum_{i \in C} \delta_i$  for  $C \subseteq [d]$ . We need the following result (see [Stückrad and Vogel 1986, Lemma II.4.14']).

**Lemma A.1.** *There is an isomorphism*

$$H_m^i(M/((\theta_1, \dots, \theta_j)M)) \cong \bigoplus_{C \subseteq [j]} H_m^{|C|+i}(M)(-\delta_C) \quad \text{for all } i, j \text{ with } i + j < d.$$

We use the following notation: if  $C \subseteq [d]$ , then  $M\langle C \rangle$  is defined by

$$M\langle C \rangle = M/((\theta_i : i \notin C)M).$$

By [Stückrad and Vogel 1986, Proposition I.2.1], if  $C \subsetneq [d]$  and  $s \in [d] \setminus C$ , then  $H_m^0(M\langle C \cup \{s\} \rangle) = 0 :_{M\langle C \cup \{s\} \rangle} \theta_s$ . This leads to the short exact sequence

$$0 \longrightarrow M\langle C \cup \{s\} \rangle / H_m^0(M\langle C \cup \{s\} \rangle)(-\delta_s) \xrightarrow{\times \theta_s} M\langle C \cup \{s\} \rangle \xrightarrow{\pi_s} M\langle C \rangle \longrightarrow 0, \quad (13)$$

where  $\pi_s$  is a natural projection. This short exact sequence gives rise to exact sequences

$$0 \longrightarrow H_m^k(M\langle C \cup \{s\} \rangle) \xrightarrow{\pi_s^*} H_m^k(M\langle C \rangle) \xrightarrow{\varphi_s^*} H_m^{k+1}(M\langle C \cup \{s\} \rangle)(-\delta_s) \longrightarrow 0 \quad (14)$$

for  $k < |C|$ , and

$$0 \longrightarrow H_m^{|C|}(M\langle C \cup \{s\} \rangle) \xrightarrow{\pi_s^*} H_m^{|C|}(M\langle C \rangle) \xrightarrow{\varphi_s^*} H_m^{|C|+1}(M\langle C \cup \{s\} \rangle)(-\delta_s), \quad (15)$$

where we denote by  $\varphi_s^*$  the connecting homomorphism.

Similarly, if  $C \subsetneq [d]$ ,  $|C| \leq d - 2$ , and  $s, t \in [d] \setminus C$ , we obtain the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & M\langle C \cup \{s\} \rangle / H_m^0(M\langle C \cup \{s\} \rangle)(-\delta_s) & \xrightarrow{\times \theta_s} & M\langle C \cup \{s\} \rangle & \xrightarrow{\pi_s} & M\langle C \rangle \longrightarrow 0 \\ & & \uparrow \pi_t & & \uparrow \pi_t & & \uparrow \pi_t \\ 0 & \longrightarrow & M\langle C \cup \{s, t\} \rangle / H_m^0(M\langle C \cup \{s, t\} \rangle)(-\delta_s) & \xrightarrow{\times \theta_s} & M\langle C \cup \{s, t\} \rangle & \xrightarrow{\pi_s} & M\langle C \cup \{t\} \rangle \longrightarrow 0 \end{array}$$

This diagram, in turn, induces the commutative diagram

$$\begin{array}{ccc} H_m^i(M\langle C \rangle) & \xrightarrow{\varphi_s^*} & H_m^{i+1}(M\langle C \cup \{s\} \rangle)(-\delta_s) \\ \uparrow \pi_t^* & & \uparrow \pi_t^* \\ H_m^i(M\langle C \cup \{t\} \rangle) & \xrightarrow{\varphi_s^*} & H_m^{i+1}(M\langle C \cup \{s, t\} \rangle)(-\delta_s) \end{array} \quad (16)$$

Now, for  $k = 2, 3, \dots, d$ , define the maps  $\phi_k$  and  $\psi_k$  as compositions:

$$\begin{aligned} \phi_k &: H_m^0(M\langle\emptyset\rangle) \xrightarrow{\varphi_1^*} H_m^1(M\langle[1]\rangle)(-\delta_{[1]}) \xrightarrow{\varphi_2^*} \dots \xrightarrow{\varphi_{k-1}^*} H_m^{k-1}(M\langle[k-1]\rangle)(-\delta_{[k-1]}), \\ \psi_k &: H_m^0(M\langle\{k}\rangle) \xrightarrow{\varphi_1^*} H_m^1(M\langle[1]\cup\{k}\rangle)(-\delta_{[1]}) \xrightarrow{\varphi_2^*} \dots \xrightarrow{\varphi_{k-1}^*} H_m^{k-1}(M\langle[k]\rangle)(-\delta_{[k-1]}). \end{aligned}$$

Then the commutativity of (16) implies the commutativity of

$$\begin{array}{ccc} H_m^0(M\langle\emptyset\rangle) & \xrightarrow{\phi_k} & H_m^{k-1}(M\langle[k-1]\rangle)(-\delta_{[k-1]}) \\ \uparrow \pi_k^* & & \uparrow \pi_k^* \\ H_m^0(M\langle\{k}\rangle) & \xrightarrow{\psi_k} & H_m^{k-1}(M\langle[k]\rangle)(-\delta_{[k-1]}) \end{array} \tag{17}$$

To prove part (ii) of Theorem 2.3, we consider the diagram

$$\begin{array}{ccccccc} & & & & H_m^0(M\langle\{1}\rangle) & \xrightarrow{\pi_1^*} & H_m^0(M\langle\emptyset\rangle) \\ & & & & & & \downarrow \varphi_1^* \\ H_m^0(M\langle\{2}\rangle) & \xrightarrow{\psi_2} & H_m^1(M\langle[2]\rangle)(-\delta_{[1]}) & \xrightarrow{\pi_2^*} & H_m^1(M\langle[1]\rangle)(-\delta_{[1]}) & & \\ & & & & & & \downarrow \varphi_2^* \\ & & & & & & \vdots \\ & & & & & & \downarrow \varphi_{d-1}^* \\ H_m^0(M\langle\{d}\rangle) & \xrightarrow{\psi_d} & H_m^{d-1}(M\langle[d]\rangle)(-\delta_{[d-1]}) & \xrightarrow{\pi_d^*} & H_m^{d-1}(M\langle[d-1]\rangle)(-\delta_{[d-1]}) & & \\ & & & & & & \downarrow \varphi_d^* \\ & & & & & & H_m^d(M\langle[d]\rangle)(-\delta_{[d]}) \end{array}$$

The surjectivity of  $\varphi_s^*$  in (14) implies that  $\psi_k$  is surjective. Hence, in each horizontal line of the diagram,

$$\text{Im}(\pi_k^* \circ \psi_k) = \text{Im}(\pi_k^*). \tag{18}$$

Also, since the sequence in (15) is exact, it follows that in the diagram we have

$$\text{Ker}(\varphi_k^*) = \text{Im}(\pi_k^*). \tag{19}$$

Define  $\phi_{d+1} = \varphi_d^* \circ \dots \circ \varphi_1^*$  to be the composition of the vertical maps in the diagram. By (15), each  $\pi_k^*$  (in the diagram) is injective, and we conclude that

$$\text{Ker}(\phi_{d+1}) \cong \bigoplus_{k=1}^d \text{Im}(\pi_k^*) \cong \bigoplus_{k=1}^d H_m^{k-1}(M\langle[k]\rangle)(-\delta_{[k-1]}) \tag{20}$$

as  $\mathbb{F}$ -vector spaces. In addition, using (18) and the commutativity of (17), we obtain that  $\text{Ker}(\phi_{d+1})$  is the sum of the images of

$$\pi_k^* : H_m^0(M\langle\{k\}\rangle) \rightarrow H_m^0(M\langle\emptyset\rangle) = M/\Theta M.$$

Finally, since

$$H_m^0(M\langle\{k\}\rangle) = ((\theta_1, \dots, \hat{\theta}_k, \dots, \theta_d)M :_M \theta_k) / (\theta_1, \dots, \hat{\theta}_k, \dots, \theta_d)M$$

and since  $\pi_k^*$  is a natural projection, it follows that

$$\text{Ker}(\phi_{d+1}) = \Sigma(\Theta; M)/\Theta M.$$

This proves Theorem 2.3(ii) since  $\phi_{d+1}$  is a map from  $M/\Theta M$  to  $H_m^d(M)(-\delta_{[d]})$ .

It remains to verify Theorem 2.3(i). Recall that  $\text{Ker}(\phi_{d+1})$  is the sum of images of  $H_m^0(M\langle\{k\}\rangle)$  and that, by [Stückrad and Vogel 1986, Proposition I.2.1],  $\mathfrak{m} \cdot H_m^0(M\langle\{k\}\rangle) = 0$ . Hence the modules in (20) are in fact isomorphic as  $S$ -modules (since they are direct sums of copies of  $\mathbb{F}$ ). Thus we infer from (20) and Lemma A.1 that

$$\begin{aligned} \Sigma(\Theta; M)/\Theta M = \text{Ker}(\phi_{d+1}) &\cong \bigoplus_{k=1}^d H_m^{k-1}(M\langle[k]\rangle)(-\delta_{[k-1]}) \\ &\cong \bigoplus_{k=1}^d \left[ \bigoplus_{C \subseteq [d] \setminus [k]} H_m^{|C|+k-1}(M)(-\delta_{C \cup [k-1]}) \right] \\ &\cong \bigoplus_{C \subsetneq [d]} H_m^{|C|}(M)(-\delta_C), \end{aligned}$$

as desired. □

**Remark.** As the proof in this appendix is based on the proof of [Novik and Swartz 2009b, Theorem 2.2], it is worth pointing out that their paper contains a minor mistake. Indeed, the short “exact” sequence that appears three lines after their statement of Theorem 2.4 is not necessarily exact. However, this mistake can be easily corrected by replacing this sequence with the short exact sequence of (13).

### Acknowledgments

We thank Ed Swartz for raising the question of whether there is an algebraic version of Theorem 1.1 and for bringing the result of Böhm and Papadakis to our attention. We are also grateful to Shiro Goto for helpful conversations.

## References

- [Adiprasito and Sanyal 2016] K. A. Adiprasito and R. Sanyal, “Relative Stanley–Reisner theory and upper bound theorems for Minkowski sums”, *Publ. Math. Inst. Hautes Études Sci.* **124** (2016), 99–163. MR Zbl
- [Alexander 1930] J. W. Alexander, “The combinatorial theory of complexes”, *Ann. of Math. (2)* **31**:2 (1930), 292–320. MR Zbl
- [Aoyama 1980] Y. Aoyama, “On the depth and the projective dimension of the canonical module”, *Japan. J. Math. (N.S.)* **6**:1 (1980), 61–66. MR Zbl
- [Aoyama and Goto 1986] Y. Aoyama and S. Goto, “Some special cases of a conjecture of Sharp”, *J. Math. Kyoto Univ.* **26**:4 (1986), 613–634. MR Zbl
- [Böhm and Papadakis 2015] J. Böhm and S. A. Papadakis, “Weak Lefschetz property and stellar subdivisions of Gorenstein complexes”, preprint, 2015. arXiv
- [Bruns and Herzog 1993] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1993. MR Zbl
- [Goto 1983] S. Goto, “On the associated graded rings of parameter ideals in Buchsbaum rings”, *J. Algebra* **85**:2 (1983), 490–534. MR Zbl
- [Gräbe 1984] H.-G. Gräbe, “Über den Stanley–Reisner–Ring von Quasimannigfaltigkeiten”, *Math. Nachr.* **117** (1984), 161–174. MR Zbl
- [Gräbe 1987] H.-G. Gräbe, “Generalized Dehn–Sommerville equations and an upper bound theorem”, *Beiträge Algebra Geom.* **25** (1987), 47–60. MR Zbl
- [Juhnke-Kubitzke and Murai 2015] M. Juhnke-Kubitzke and S. Murai, “Balanced generalized lower bound inequality for simplicial polytopes”, preprint, 2015. arXiv
- [Klee and Novik 2016] S. Klee and I. Novik, “Face enumeration on simplicial complexes”, pp. 653–686 in *Recent Trends in Combinatorics*, edited by A. Beveridge et al., The IMA Volumes in Mathematics and its Applications **159**, Springer, 2016. Zbl
- [McMullen and Walkup 1971] P. McMullen and D. W. Walkup, “A generalized lower-bound conjecture for simplicial polytopes”, *Mathematika* **18** (1971), 264–273. MR Zbl
- [Murai 2010] S. Murai, “On face vectors of barycentric subdivisions of manifolds”, *SIAM J. Discrete Math.* **24**:3 (2010), 1019–1037. MR Zbl
- [Murai and Nevo 2014] S. Murai and E. Nevo, “On  $r$ -stacked triangulated manifolds”, *J. Algebraic Combin.* **39**:2 (2014), 373–388. MR Zbl
- [Murai and Novik 2016] S. Murai and I. Novik, “Face numbers of manifolds with boundary”, preprint, 2016. To appear in *Int. Math. Res. Not. IMRN*. arXiv
- [Murai and Yanagawa 2014] S. Murai and K. Yanagawa, “Squarefree  $P$ -modules and the  $\mathbf{cd}$ -index”, *Adv. Math.* **265** (2014), 241–279. MR Zbl
- [Novik 1998] I. Novik, “Upper bound theorems for homology manifolds”, *Israel J. Math.* **108** (1998), 45–82. MR Zbl
- [Novik and Swartz 2009a] I. Novik and E. Swartz, “Gorenstein rings through face rings of manifolds”, *Compos. Math.* **145**:4 (2009), 993–1000. MR Zbl
- [Novik and Swartz 2009b] I. Novik and E. Swartz, “Socles of Buchsbaum modules, complexes and posets”, *Adv. Math.* **222**:6 (2009), 2059–2084. MR Zbl
- [Pachner 1987] U. Pachner, “Konstruktionsmethoden und das kombinatorische Homöomorphieproblem für Triangulationen kompakter semilinearer Mannigfaltigkeiten”, *Abh. Math. Sem. Univ. Hamburg* **57** (1987), 69–86. MR Zbl

- [Pachner 1991] U. Pachner, “P.L. homeomorphic manifolds are equivalent by elementary shellings”, *European J. Combin.* **12**:2 (1991), 129–145. MR Zbl
- [Schenzel 1981] P. Schenzel, “On the number of faces of simplicial complexes and the purity of Frobenius”, *Math. Z.* **178**:1 (1981), 125–142. MR Zbl
- [Schenzel 1982] P. Schenzel, *Dualisierende Komplexe in der lokalen Algebra und Buchsbaum–Ringe*, Lecture Notes in Mathematics **907**, Springer, 1982. MR Zbl
- [Stanley 1993] R. P. Stanley, “A monotonicity property of  $h$ -vectors and  $h^*$ -vectors”, *European J. Combin.* **14**:3 (1993), 251–258. MR Zbl
- [Stanley 1996] R. P. Stanley, *Combinatorics and commutative algebra*, 2nd ed., Progress in Mathematics **41**, Birkhäuser, Boston, 1996. MR Zbl
- [Stückrad and Vogel 1986] J. Stückrad and W. Vogel, *Buchsbaum rings and applications: an interaction between algebra, geometry and topology*, Springer, 1986. MR Zbl
- [Suzuki 1981] N. Suzuki, “The Koszul complex of Buchsbaum modules”, pp. 15–25 in *Commutative Algebra and Algebraic Geometry* (Kyoto University, 1981), vol. 446, RIMS Kokyuroku, 1981. Zbl
- [Swartz 2014] E. Swartz, “Thirty-five years and counting”, preprint, 2014. arXiv

Communicated by Victor Reiner

Received 2016-03-08      Accepted 2016-10-08

s-murai@ist.osaka-u.ac.jp

*Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Suita 565-0871, Japan*

novik@math.washington.edu

*Department of Mathematics, University of Washington, Box 354350, Seattle, WA 98195-4350, United States*

yoshida@math.chs.nihon-u.ac.jp

*Department of Mathematics, College of Humanities and Sciences, Nihon University, Setagaya-ku, Tokyo 156-8550, Japan*



# Algebra & Number Theory

msp.org/ant

## EDITORS

### MANAGING EDITOR

Bjorn Poonen  
Massachusetts Institute of Technology  
Cambridge, USA

### EDITORIAL BOARD CHAIR

David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
John H. Coates	University of Cambridge, UK	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2017 is US \$325/year for the electronic version, and \$520/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.


---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 11    No. 3    2017

---

The umbral moonshine module for the unique unimodular Niemeier root system JOHN F. R. DUNCAN and JEFFREY A. HARVEY	505
Geometry on totally separably closed schemes STEFAN SCHRÖER	537
The sixth moment of automorphic $L$ -functions VORRAPAN CHANDEE and XIANNAN LI	583
A duality in Buchsbaum rings and triangulated manifolds SATOSHI MURAI, ISABELLA NOVIK and KEN-ICHI YOSHIDA	635
Analytic functions on tubes of nonarchimedean analytic spaces FLORENT MARTIN	657
Automatic sequences and curves over finite fields ANDREW BRIDY	685
On an analogue of the Ichino–Ikeda conjecture for Whittaker coefficients on the metaplectic group EREZ LAPID and ZHENGYU MAO	713



1937-0652(2017)11:3;1-I