

Algebra & Number Theory

Volume 12

2018

No. 1

**Governing singularities
of symmetric orbit closures**

Alexander Woo, Benjamin J. Wyser and Alexander Yong



Governing singularities of symmetric orbit closures

Alexander Woo, Benjamin J. Wyser and Alexander Yong

We develop *interval pattern avoidance* and *Mars–Springer ideals* to study singularities of symmetric orbit closures in a flag variety. This paper focuses on the case of the Levi subgroup $\mathrm{GL}_p \times \mathrm{GL}_q$ acting on the classical flag variety. We prove that all reasonable singularity properties can be classified in terms of interval patterns of *clans*.

1. Introduction

1A. Overview. Let G/B be a generalized flag variety where G is a complex, reductive algebraic group and B is a choice of Borel subgroup. A subgroup K of G is *symmetric* if $K = G^\theta$ is the fixed point subgroup for an involutive automorphism θ of G . Such a subgroup is *spherical*, which means that its action on G/B by left translations has finitely many orbits.

The study of orbits of a symmetric subgroup on the flag variety was initiated in [Lusztig and Vogan 1983; Vogan 1983], where the singularities of closures of the orbits were related to characters of particular infinite-dimensional representations of a certain real form $G_{\mathbb{R}}$ of G . Since then, there has been a stream of results on the combinatorics and geometry of these orbit closures. Notably, R. W. Richardson and T. A. Springer [Richardson and Springer 1990] gave a description of the partial order given by inclusions of orbit closures, and M. Brion [2001] studied general properties of their singularities, showing that, in many cases, including the one addressed in this paper, all these orbit closures are normal and Cohen–Macaulay with rational singularities. One might also hope that the study of singularities on closures of symmetric subgroup orbits would lead to better understanding of the general relationship between the combinatorics associated to spherical varieties and singularities of orbit closures on them; N. Perrin has written a survey of this topic [Perrin 2014].

We initiate a combinatorial approach, backed by explicit commutative algebra computations, to the study of the singularities of these orbit closures. This paper considers the case of the symmetric subgroup $K = \mathrm{GL}_p \times \mathrm{GL}_q$ of block diagonal matrices in GL_n , where $n = p + q$. Here θ is defined by

$$\theta(M) = I_{p,q} M I_{p,q}, \quad (1)$$

MSC2010: primary 14M17; secondary 05E40, 14M15, 14M27.

Keywords: flag variety, Schubert variety, symmetric subgroup, interval pattern avoidance.

where $I_{p,q}$ is the diagonal matrix with diagonal consisting of 1 p times followed by -1 q times. In this case, $G_{\mathbb{R}} = U(p, q)$ is the *indefinite unitary group* of signature (p, q) .

For this symmetric subgroup K , the finitely many K -orbits \mathcal{O}_γ and their closures Y_γ can be parametrized by (p, q) -clans γ [Matsuki and Ōshima 1990, Theorem 4.1] (see also [Yamamoto 1997, Theorem 2.2.8]). These clans are partial matchings of vertices $\{1, 2, \dots, n\}$, where each unmatched vertex is assigned a sign of $+$ or $-$; the difference in the number of $+$ and $-$ signs must be $p - q$. We represent clans by length n strings in $\mathbb{N} \cup \{+, -\}$, with pairs of equal numbers indicating a matching. Let $\text{Clans}_{p,q}$ denote the set of all such clans. For example, three clans from $\text{Clans}_{7,3}$ are represented by the strings:

$$1+++2+1-2+, \quad +1-++1+2+2, \quad \text{and} \quad ++-++++-+-.$$

Note that strings that differ by a permutation of the natural numbers, such as 1212 and 2121, represent the same clan.

One inspiration for this work is W. M. McGovern's characterization [2009] of singular K -orbit closures in terms of *pattern avoidance* of clans. Suppose $\gamma \in \text{Clans}_{p,q}$ and $\theta \in \text{Clans}_{r,s}$. Then $\theta = \theta_1 \cdots \theta_{r+s}$ is said to (*pattern*) *contain* $\gamma = \gamma_1 \cdots \gamma_{p+q}$ if there are indices $i_1 < i_2 < \cdots < i_{p+q}$ such that:

- (1) if $\gamma_j = \pm$ then $\theta_{i_j} = \gamma_j$; and
- (2) if $\gamma_k = \gamma_\ell$ then $\theta_{i_k} = \theta_{i_\ell}$.

For example, the clan $\gamma = 1++--1$ contains the pattern $\theta = 1++-1$, taking (i_1, \dots, i_5) to be either $(1, 2, 3, 4, 6)$ or $(1, 2, 3, 5, 6)$.

Say that θ (*pattern*) *avoids* γ if θ does not contain γ . The main theorem of [McGovern 2009] asserts that Y_γ is smooth if and only if γ avoids the patterns

$$1+-1, \quad 1-+1, \quad 1212, \quad 1+221, \quad 1-221, \quad 122+1, \quad 122-1, \quad 122331.$$

On the other hand, in [Woo and Wyser 2015, Section 3.3] it is noted that Y_{1++-1} is non-Gorenstein, while Y_{1++--1} is Gorenstein, even though $1++--1$ pattern contains $1++-1$. Therefore, a more general notion will sometimes be required to characterize which K -orbits satisfy a particular singularity property.

Suppose that \mathcal{P} is any *singularity mildness property*, by which we mean a local property of varieties that holds on open subsets and is stable under smooth morphisms. Many singularity properties, such as being Gorenstein, being a local complete intersection (lci), being factorial, having Cohen–Macaulay rank $\leq k$, or having Hilbert–Samuel multiplicity $\leq k$, satisfy these conditions. For such a \mathcal{P} , consider two related problems:

- (I) Which K -orbit closures Y_γ are globally \mathcal{P} ?
- (II) What is the non- \mathcal{P} -locus of Y_γ ?

This paper gives a universal combinatorial language, *interval pattern avoidance* of clans, to answer these questions for any singularity mildness property, at the cost of potentially requiring an infinite number of patterns. This language is also useful for collecting and analyzing data and partial results. We present

explicit equations for computing whether a property holds at a specific orbit \mathcal{O}_α on an orbit closure Y_γ . Note that all points of $\mathcal{O}_\alpha \subseteq Y_\gamma$ are locally isomorphic to one another, since the K -action can be used to move any point of \mathcal{O}_α isomorphically to any other point. Since the non- \mathcal{P} -locus is closed, it is a union of K -orbit closures. Consequently, for any given clan γ , (II) can be answered by finding a finite set of clans $\{\alpha\}$, namely those indexing the irreducible components of the non- \mathcal{P} -locus. (I) asks if this set is nonempty.

This situation parallels that for Schubert varieties. In that setting, the first and third authors introduced interval pattern avoidance for permutations, showing that it provides a common perspective to study all reasonable singularity measures [Woo and Yong 2006; 2008; Woo 2010]. This paper gives the first analogue of those results for K -orbit closures.

Some properties \mathcal{P} hold globally on every Y_γ . For those cases, the above questions are unnecessary. For example, this is true when $\mathcal{P} = \text{“normal”}$ and $\mathcal{P} = \text{“Cohen–Macaulay”}$ in the case $(G, K) = (\mathrm{GL}_n, \mathrm{GL}_p \times \mathrm{GL}_q)$ of this paper. (This is not the case for all symmetric pairs (G, K) .)

As is explained in [McGovern 2009], the above answer to (I) for $\mathcal{P} = \text{“smooth”}$ also is the answer for the property $\mathcal{P} = \text{“rationally smooth”}$. (Recall rational smoothness means that the local intersection cohomology of Y_γ (at a point of \mathcal{O}_α) is trivial.) However, the answer to question (II), which asks for a combinatorial description of the (rationally) singular locus, is unsolved except in some special cases [Woo and Wyser 2015]. Actually, it is unknown whether the singular locus and rationally singular locus coincide for all orbit closures (but see Conjecture 7.5).

For most finer singularity mildness properties, answers to both (I) and (II) are unknown. Perhaps the most famous such property comes from the *Kazhdan–Lusztig–Vogan* (KLV) polynomials $P_{\gamma,\alpha}(q)$. These polynomials are the link to representation theory that originally motivated the study of K -orbit closures. For $(G, K) = (\mathrm{GL}_n, \mathrm{GL}_p \times \mathrm{GL}_q)$, the KLV polynomial $P_{\gamma,\alpha}(q) \in \mathbb{Z}_{\geq 0}[q]$ is the Poincaré polynomial for the local intersection cohomology of Y_γ at any point of the orbit \mathcal{O}_α [Lusztig and Vogan 1983]. Rational smoothness of Y_γ along \mathcal{O}_α is hence equivalent to the equality $P_{\gamma,\alpha}(q) = 1$. More generally, for any fixed $k > 1$, the property $\mathcal{P} = \text{“}P_{\gamma,\alpha}(1) \leq k\text{”}$ behaves as a singularity mildness property on K -orbit closures by recent work of McGovern [2015], but his proof uses representation theory, and the algebraic geometry is not well-understood.

1B. Main ideas. Each K -orbit closure Y_γ is a union of K -orbits \mathcal{O}_α ; let \leq denote the *Bruhat (closure) order* on K -orbit closures on GL_n/B . (This means $\alpha \leq \gamma$ if and only if $Y_\alpha \subseteq Y_\gamma$.) Also, let $[\alpha, \gamma]$ and $[\beta, \theta]$ be intervals in Bruhat order on $\mathrm{Clans}_{p,q}$ and $\mathrm{Clans}_{r,s}$, respectively.

Define $[\beta, \theta]$ to *interval pattern contain* $[\alpha, \gamma]$ (or equivalently $[\alpha, \gamma]$ to *interval pattern embed* into $[\beta, \theta]$), and write $[\alpha, \gamma] \hookrightarrow [\beta, \theta]$ if:

- (a) there are indices

$$I : i_1 < i_2 < \cdots < i_{p+q}$$

which commonly witness the containment of γ into θ and α into β ;

- (b) θ and β agree outside of these indices; and
- (c) $\ell(\theta) - \ell(\beta) = \ell(\gamma) - \ell(\alpha)$, where if $\gamma = \gamma_1\gamma_2 \dots \gamma_{p+q}$ then

$$\ell(\gamma) := \sum_{\gamma_i = \gamma_j \in \mathbb{N}} j - i - \#\{\gamma_s = \gamma_t \mid s < i < t < j\}.$$

Notice β is determined by α, γ, θ , and the set of indices I . In particular, β is the unique clan $\Phi(\alpha)$ that agrees with α on I and agrees with θ on $\{1, \dots, n\} \setminus I$. Thus, we define a clan θ to *interval pattern contain* $[\alpha, \gamma]$ if $[\alpha, \gamma]$ is contained in $[\Phi(\alpha), \theta]$. Similarly, we can speak of θ avoiding a list of intervals.

Example 1.1. Let $[\alpha, \gamma] = [+--+, 1212]$ and $[\beta, \theta] = [1+---+1, 123231]$. Then one can check $[\beta, \theta]$ interval contains $[\alpha, \gamma]$, using the middle four positions.

Example 1.2. Let $[\alpha, \gamma] = [+-, 11]$ and $[\beta, \theta] = [\underline{+}+\underline{-}, \underline{1}+\underline{1}]$. If $[\beta, \theta]$ contains $[\alpha, \gamma]$ it must be using the underlined positions. However, $\ell(\gamma) - \ell(\alpha) = 1$ while $\ell(\theta) - \ell(\beta) = 2$. Thus $[\beta, \theta]$ does not contain $[\alpha, \gamma]$.

Define

$$\mathcal{C} := \{[\alpha, \gamma] \mid \alpha \leq \gamma \text{ in some } \text{Clans}_{p,q}\} \subseteq \text{Clans} \times \text{Clans},$$

where

$$\text{Clans} = \bigcup_{p,q} \text{Clans}_{p,q}.$$

Declare $\leq_{\mathcal{C}}$ to be the poset relation on \mathcal{C} generated by

- $[\alpha, \gamma] \leq_{\mathcal{C}} [\beta, \theta]$ if $[\beta, \theta]$ interval pattern contains $[\alpha, \gamma]$, and
- $[\alpha, \gamma] \leq_{\mathcal{C}} [\alpha', \gamma]$ if $\alpha' \leq \alpha$.

For any poset (\mathcal{S}, \leq) , an *upper order ideal* is a subset \mathcal{I} of \mathcal{S} having the property that whenever $x \in \mathcal{I}$, we also have $y \in \mathcal{I}$ for all $y \geq x$. The following theorem provides a basic language to express answers to (I) and (II).

Theorem 1.3. *Let \mathcal{P} be a singularity mildness property (see Definition 2.1).*

- (I) *The set of intervals $[\alpha, \gamma] \in \mathcal{C}$ such that \mathcal{P} fails on each point of $\mathcal{O}_{\alpha} \subseteq Y_{\gamma}$ is an upper order ideal in $(\mathcal{C}, \leq_{\mathcal{C}})$.*
- (II) *The set of clans γ such that \mathcal{P} holds at all points of Y_{γ} are those that avoid all the intervals $[\alpha_i, \gamma_i]$ constituting some (possibly infinite) set $\mathcal{A}_{\mathcal{P}} \subseteq \mathcal{C}$.*

For simplicity, we work over \mathbb{C} , but our results are valid over any field of characteristic not 2, with the caveat that even for the same \mathcal{P} , the set $\mathcal{A}_{\mathcal{P}}$ may be field dependent.

Although stated in combinatorial language, as we will see, Theorem 1.3 follows from a geometric result, Theorem 4.6, which establishes a local isomorphism between certain “slices” of the orbit closures. Rather than working directly with the K -orbit closures on G/B , it is easier to establish this isomorphism using particular slices of B -orbit closures on G/K given by J. G. M. Mars and Springer [1998].

The space $G/K = \mathrm{GL}_n / (\mathrm{GL}_p \times \mathrm{GL}_q)$ is the configuration space of all splittings of \mathbb{C}^n as a direct sum $V_1 \oplus V_2$ of subspaces, with $\dim(V_1) = p$ and $\dim(V_2) = q$. Indeed, $G = \mathrm{GL}_n(\mathbb{C})$ acts transitively on the set of all such splittings, and $K = \mathrm{GL}_p \times \mathrm{GL}_q$ is the stabilizer of the “standard” splitting of \mathbb{C}^n as

$$\langle e_1, \dots, e_p \rangle \oplus \langle e_{p+1}, \dots, e_n \rangle,$$

where e_1, \dots, e_n is the standard basis of \mathbb{C}^n . A point $gK/K \in G/K$, with $g \in \mathrm{GL}_n(\mathbb{C})$ an invertible $n \times n$ matrix, is identified with the splitting whose p -dimensional component is the span of the first p columns of g and whose q -dimensional component is the span of the last q columns of g .

B -orbits on G/K are in bijection with K -orbits on G/B (both orbit sets being in bijection with the $B \times K$ -orbits on G). In addition, this bijection preserves all mildness properties. This is because the orbit closures correspond via the two locally trivial fibrations $G \rightarrow G/K$ and $G \rightarrow G/B$, each of which has smooth fiber. The B -orbit on G/K corresponding to the clan γ will be denoted by Q_γ , and its closure will be denoted by W_γ . For any singularity mildness property \mathcal{P} , this discussion implies:

Observation 1.4. Y_γ is \mathcal{P} along \mathcal{O}_α if and only if W_γ is \mathcal{P} along Q_α .

In [Wyser and Yong 2014], the second and third authors considered certain open affine subsets of K -orbit closures on G/B which they called *patches*. That work also introduced *patch ideals* of equations which set-theoretically cut out the patches. Experimental computation using patch ideals led to several of the conjectures which appear in [Wyser and Yong 2014; Woo and Wyser 2015].

In contrast, the Mars–Springer slices are not open affine pieces of the orbit closures. However, as we will see, they are essentially as good, in that they carry all of the local information that we are interested in. They are analogues of *Kazhdan–Lusztig varieties*, which play a similar role in the study of Schubert varieties.

In this paper we introduce *Mars–Springer ideals* (see Section 4) with explicit equations that set-theoretically cut out the Mars–Springer slices. These equations are conjectured to also be scheme-theoretically correct; see Conjecture 7.1. These are the K -orbit versions of *Kazhdan–Lusztig ideals*, which define the aforementioned Kazhdan–Lusztig varieties. The latter ideals have been of use in both computational and theoretical analysis of Schubert varieties (see, for example, [Woo and Yong 2012; Ulfarsson and Woo 2013]). In the same vein, we mention a practical advantage of the Mars–Springer ideal over the patch ideal. Gröbner basis calculations with the former are several times faster than those of the latter, as fewer variables are involved.

1C. Organization. In Section 2, we present some preliminaries. In particular, we more precisely define “mildness properties,” giving examples and establishing some basic facts that we will need. We next recall the attractive slices of Mars and Springer [1998]. Finally, we describe the quasiprojective variety structure of G/K for the case $(G, K) = (\mathrm{GL}_n, \mathrm{GL}_p \times \mathrm{GL}_q)$ of this paper. In Section 3, we give explicit affine coordinates for the Mars–Springer slice (Theorem 4.3). Section 4 defines the *Mars–Springer variety* and its ideal. It culminates with Theorem 4.6, which asserts that certain Mars–Springer varieties are isomorphic to others. This theorem is the key to our proof of Theorem 1.3. In order to prove Theorem 4.6,

we develop combinatorics of interval embeddings using earlier work of the second author [Wyser 2016] on the Bruhat order on clans; this is Section 5. We then give our proofs of Theorem 4.6 and Theorem 1.3 in Section 6. We conclude with problems and conjectures in Section 7.

2. Preliminaries

2A. Singularity mildness properties. We define the class of local properties that we are interested in, with specific examples.

Definition 2.1. Suppose that \mathcal{P} is a local property of algebraic varieties, meaning that \mathcal{P} is verified at a point w of an algebraic variety W based solely on the local ring $\mathcal{O}_{W,w}$. We say that \mathcal{P} is a *singularity mildness property* (or simply a *mildness property*) if \mathcal{P} is

- (1) open, meaning that the \mathcal{P} -locus of any variety is open; and
- (2) stable under smooth morphisms, i.e., if X and Y are any varieties, and $f : X \rightarrow Y$ is any smooth morphism between them, then for any $x \in X$, X is \mathcal{P} at x if and only if Y is \mathcal{P} at $f(x)$.

For any smooth variety S and for any variety X , the projection map $S \times X \rightarrow X$ is a smooth morphism. Thus:

Observation 2.2. *If \mathcal{P} is a mildness property, S is a smooth variety, and X is any variety, then for any $x \in X$ and for any $s \in S$, X is \mathcal{P} at x if and only if $S \times X$ is \mathcal{P} at (s, x) .*

The next lemma gives a (nonexhaustive) list of examples of mildness properties \mathcal{P} .

Lemma 2.3. *Examples of mildness properties \mathcal{P} include:*

- (i) *reducedness*;
- (ii) *normality*;
- (iii) *smoothness*;
- (iv) *lci-ness*;
- (v) *Gorensteinness*.

Proof. That these properties are open (on varieties) is well-known. That they are stable under smooth morphisms follows from results of [Matsumura 1986, §23]. Indeed, let X and Y be varieties, with $f : X \rightarrow Y$ a smooth morphism between them. Let $x \in X$ be given, and let $y = f(x)$. Now, f is flat, and also the fiber F_y over y is smooth over the residue field $\kappa(y)$, hence regular. Thus $\mathcal{O}_{F_y,x}$ is regular, hence also lci and Gorenstein. Now (iii) and (v) follow from Theorems 23.7 and 23.4 of [Matsumura 1986, §23], respectively, while (iv) is mentioned in the remark following Theorem 23.6 of [loc. cit.]; as indicated there, further details can be found in [Avramov 1975].

For (i) and (ii), we appeal to [Matsumura 1986, Corollary to Theorem 23.9]. Note that this result requires that the fiber ring corresponding to any prime ideal of $\mathcal{O}_{Y,y}$ be reduced (resp. normal), rather than just the fiber ring of the maximal ideal. In our case, all such fiber rings are once again regular

(hence both reduced and normal). Indeed, if p is any prime of $\mathcal{O}_{Y,y}$, then p corresponds to an irreducible closed subvariety Y' of Y . The inverse image scheme $f^{-1}(Y')$ is smooth over Y' , and the fiber ring $\mathcal{O}_{X,x} \otimes_{\mathcal{O}_{Y,y}} \kappa(p)$ is the generic fiber of $f^{-1}(Y') \rightarrow Y'$. \square

2B. Mars–Springer slices. Let W be any irreducible variety with an action of a Borel group B , let $w \in W$ be given, and let $Q = B \cdot w$.

Definition 2.4 [Mars and Springer 1998]. An *attractive slice* to Q at w in W is a locally closed subset S of W containing w such that:

- (1) the restriction of the action map $B \times S \rightarrow W$ is a smooth morphism;
- (2) $\dim(S) + \dim(Q) = \dim(W)$; and
- (3) there exists a map $\lambda : \mathbb{G}_m \rightarrow B$ such that
 - (a) S is stable under $\text{Im}(\lambda)$;
 - (b) w is a fixed point of $\text{Im}(\lambda)$; and
 - (c) the action map $\mathbb{G}_m \times S \rightarrow S$ extends to a morphism $\mathbb{A}^1 \times S \rightarrow S$ which sends $\{0\} \times S$ to w .

The following result justifies our use of slices in studying singularities, since taking slices preserves the local properties that we are interested in.

Lemma 2.5. (Using the notation of Definition 2.4.) *Given a mildness property \mathcal{P} , the variety W is \mathcal{P} at w if and only if S is \mathcal{P} at w .*

Proof. Indeed, the action map $B \times S \rightarrow W$ is smooth, so W is \mathcal{P} at w if and only if $B \times S$ is \mathcal{P} at $(1, w)$, which is the case if and only if S is \mathcal{P} at w , by Observation 2.2. \square

Let G be a complex algebraic group with an involutive automorphism θ , with $K = G^\theta$ the corresponding symmetric subgroup. Suppose that $T \subseteq B$ are a θ -stable maximal torus and Borel subgroup of G , respectively. Let N be the normalizer of T in G , and let $\mathcal{W} = N/T$ be the Weyl group.

Given a B -orbit Q on G/K , we now define the *Mars–Springer slice* $S_{\bar{x}}$ to Q at a specially chosen point $\bar{x} \in Q$. Letting U^- be the unipotent radical of the Borel subgroup opposite to B , this slice is of the form

$$S_{\bar{x}} := (U^- \cap \psi(U^-)) \cdot \bar{x}, \tag{2}$$

where ψ is a certain involution on G .

To be precise, consider the set

$$\mathcal{V} := \{x \in G \mid x(\theta x)^{-1} \in N\}.$$

In (2), we pick any

$$\bar{x} := xK/K \in Q \quad \text{with } x \in \mathcal{V};$$

such a choice exists by [Richardson and Springer 1990, Theorem 1.3]. Let η be the image in \mathcal{W} of $x\theta(x)^{-1} \in N$. The element $x\theta(x)^{-1} \in N$ may depend upon the choice of \bar{x} , but η , its class modulo T ,

does not. Then in (2) we take

$$\psi := c_\eta \circ \theta,$$

where c_η denotes conjugation by η . With these choices, it is shown in [Mars and Springer 1998, Section 6.4] that $S_{\bar{x}}$, as defined in (2), is indeed an attractive slice to Q at \bar{x} in G/K , in the sense of Definition 2.4.

The following lemma is more or less formal; we include a proof for completeness:

Lemma 2.6. *With notation as above, if W is any B -orbit closure on G/K containing Q , then:*

- (i) *the scheme-theoretic intersection $W \cap S_{\bar{x}}$ is reduced, and*
- (ii) *$W \cap S_{\bar{x}}$ is an attractive slice to Q at \bar{x} in W .*

Proof. It is easy to see that the diagram

$$\begin{array}{ccc} B \times (W \cap S_{\bar{x}}) & \longrightarrow & W \\ \downarrow & & \downarrow \\ B \times S_{\bar{x}} & \longrightarrow & G/K \end{array}$$

is Cartesian. Therefore, since the action map along the bottom is smooth, the (restricted) action map along the top is also smooth, since smooth morphisms are stable under base extension. Since W is reduced and reducedness is a mildness property by Lemma 2.3, $B \times (W \cap S_{\bar{x}})$ is reduced, and thus $W \cap S_{\bar{x}}$ is reduced, by Observation 2.2. This proves (i).

For (ii), the preceding observation regarding the smoothness of the map along the top of the above diagram verifies Definition 2.4(1) for $S = W \cap S_{\bar{x}}$. This map furthermore has the same relative dimension as the one along the bottom, which implies that

$$\dim(G/K) - \dim(W) = \dim(S_{\bar{x}}) - \dim(W \cap S_{\bar{x}}).$$

Since $S_{\bar{x}}$ is an attractive slice to Q in G/K , we know that

$$\dim(S_{\bar{x}}) + \dim(Q) = \dim(G/K),$$

and combining these two equations gives

$$\dim(Q) + \dim(W \cap S_{\bar{x}}) = \dim(W),$$

as required by Definition 2.4(2).

That $W \cap S_{\bar{x}}$ satisfies Definition 2.4(3) is obvious: Let $\lambda : \mathbb{G}_m \rightarrow B$ be a one-parameter subgroup having properties (a) and (b) relative to the slice $S_{\bar{x}}$. Then clearly $W \cap S_{\bar{x}}$ is stable under $\text{Im}(\lambda)$, \bar{x} is still a fixed point of $\text{Im}(\lambda)$, and the action map $\mathbb{G}_m \times (W \cap S_{\bar{x}}) \rightarrow W \cap S_{\bar{x}}$ still extends to a map $\mathbb{A}^1 \times (W \cap S_{\bar{x}}) \rightarrow W \cap S_{\bar{x}}$ sending $\{0\} \times (W \cap S_{\bar{x}})$ to \bar{x} (simply restrict the original extension to the smaller subset). \square

Corollary 2.7. *For $(G, K) = (\mathrm{GL}_n, \mathrm{GL}_p \times \mathrm{GL}_q)$, let clans α and γ be given, with $\alpha \leq \gamma$ in Bruhat order. Let X_α denote the Mars–Springer slice to Q_α at a particular point $x_\alpha \in Q_\alpha$ in G/K . Then the slice $W_\gamma \cap X_\alpha$ is irreducible.*

Proof. It is observed in [Brion 2001] that all B -orbit closures on G/K for this particular case are “multiplicity-free,” which implies by [Brion 2001, Theorem 5] that W_γ has rational singularities. In particular, it is normal. Since normality is a mildness property by Lemma 2.3, $W_\gamma \cap X_\alpha$ is also normal. Now, the “attractive” condition of slices (part (3) of Definition 2.4) ensures that $W_\gamma \cap X_\alpha$ is connected because x_α must lie in every connected component of it. Being both normal and connected, $W_\gamma \cap X_\alpha$ must be irreducible. \square

Remark. The result of Corollary 2.7 in fact holds for any symmetric pair (G, K) such that all of the B -orbit closures on G/K are multiplicity-free. Other such cases include $(\mathrm{GL}_{2n}, \mathrm{Sp}_{2n})$ and $(\mathrm{SO}_{2n}, \mathrm{GL}_n)$, as noted in [Brion 2001]. However, not all symmetric pairs (G, K) have this property. For those that do not, B -orbit closures on G/K can be nonnormal, and the Mars–Springer slices can in fact be reducible. One example is (GL_n, O_n) . \square

2C. $\mathrm{GL}_{p+q}/(\mathrm{GL}_p \times \mathrm{GL}_q)$ as a quasiprojective variety. We now discuss the structure of $G/K = \mathrm{GL}_{p+q}/(\mathrm{GL}_p \times \mathrm{GL}_q)$ as a quasiprojective variety and its affine patches. The statements of this section could be extracted from standard definitions and results, but to the best of our knowledge, this has never been made explicit in the literature, so we take the opportunity to do so here.

As stated in the Introduction, G/K is the configuration space whose points correspond to splittings of \mathbb{C}^n as a direct sum $V_1 \oplus V_2$ of subspaces, with $\dim(V_1) = p$ and $\dim(V_2) = q$. Hence it is natural to identify G/K with a subset of the product of Grassmannians $\mathrm{Gr}(p, \mathbb{C}^n) \times \mathrm{Gr}(q, \mathbb{C}^n)$. We can realize $\mathrm{Gr}(p, \mathbb{C}^n) \times \mathrm{Gr}(q, \mathbb{C}^n)$ as a subvariety of $\mathbb{P}^{\binom{n}{p}-1} \times \mathbb{P}^{\binom{n}{q}-1}$ using the Plücker embedding into each factor. The Segre embedding then realizes this product as a subvariety of $\mathbb{P}^{\binom{n}{p}\binom{n}{q}-1}$, so $\mathrm{Gr}(p, \mathbb{C}^n) \times \mathrm{Gr}(q, \mathbb{C}^n)$ is a projective variety.

We let p_S and q_T denote the Plücker coordinates on $\mathrm{Gr}(p, \mathbb{C}^n)$ and $\mathrm{Gr}(q, \mathbb{C}^n)$ respectively, where S and T respectively range over all subsets of $\{1, \dots, n\}$ of size p (respectively q).

Not every pair V_1 and V_2 of subspaces of \mathbb{C}^n gives a splitting. In order for V_1 and V_2 to form a direct sum, a basis for V_1 must be linearly independent of a basis of V_2 . Let M be a matrix whose first p columns are a basis for V_1 and whose last q columns are a basis for V_2 . If we take the Laplace expansion of $\det M$ using the first p columns and identify determinants of submatrices with Plücker coordinates, then we get

$$\det M = \sum_{S \subset \{1, \dots, n\}, \#S=p} (-1)^S p_S q_{\bar{S}}, \quad (3)$$

where $\bar{S} := \{1, \dots, n\} \setminus S$ and $(-1)^S := (-1)^{(\sum_{s \in S} s) - \binom{p+1}{2}}$. Hence we can identify G/K with the open subset of $\mathrm{Gr}(p, \mathbb{C}^n) \times \mathrm{Gr}(q, \mathbb{C}^n)$ where (3) is nonzero. This gives G/K the structure of a quasiprojective variety.

The product $\mathbb{P}^{\binom{n}{p}-1} \times \mathbb{P}^{\binom{n}{q}-1}$ of projective spaces has a covering by affine patches $\{U_{S,T} := U_S \times U_T\}$, where U_S is the patch where p_S is nonzero and U_T is the patch where q_T is nonzero. Thus $\text{Gr}(p, \mathbb{C}^n) \times \text{Gr}(q, \mathbb{C}^n)$ is covered by these affine patches. As explained in [Fulton 1997, Section 9.1], any subspace $V_1 \in \text{Gr}(p, \mathbb{C}^n)$ having Plücker coordinate $p_S(V_1) \neq 0$ has a basis $\{e_s + \sum_{r \notin S} a_{s,r}(V_1)e_r\}_{s \in S}$ where e_i denotes the i -th standard basis vector, and the functions $a_{s,r}$ are local coordinates on the patch $U_S \cap \text{Gr}(p, \mathbb{C}^n)$. Over this patch, there is an algebraic section ϕ_S of the projection map $M_{n \times p}^0 \rightarrow \text{Gr}(p, \mathbb{C}^n)$, where $M_{n \times p}^0$ denotes the $n \times p$ matrices of full rank p . The section ϕ sends the subspace V_1 to the matrix whose columns are precisely the aforementioned basis.

Likewise, any subspace $V_2 \in \text{Gr}(q, \mathbb{C}^n)$ having Plücker coordinate $q_T(V_2) \neq 0$ has a basis

$$\left\{ e_t + \sum_{r \notin T} b_{t,r}(V_2)e_r \right\}_{t \in T},$$

and the $b_{t,r}$ are local coordinates on the patch $U_T \cap \text{Gr}(q, \mathbb{C}^n)$. Over this patch, there is a similar section ϕ_T which sends V_2 to the full rank $n \times q$ matrix whose columns are this basis.

Taking the products of the sections ϕ_S and ϕ_T , over $U_{S,T} \cap (\text{Gr}(p, \mathbb{C}^n) \times \text{Gr}(q, \mathbb{C}^n))$ we have an algebraic map $\phi_{S,T}$ from G/K into $M_{n \times p}^0 \times M_{n \times q}^0$, which naturally embeds into $M_{n \times n}$ by simple concatenation of matrices. Over G/K , this map takes values in $G = \text{GL}_n$ and gives an algebraic section $\phi_{S,T}$ of the projection map $G \rightarrow G/K$.

3. Affine coordinates for the Mars–Springer slice

We provide explicit coordinates for the Mars–Springer slice for the case that

$$(G, K) = (\text{GL}_n, \text{GL}_p \times \text{GL}_q).$$

3A. The affine space S_α . We define an affine space of matrices associated to each clan α . First, let $w_\alpha \in S_n$ (in one line notation) be defined as follows. From left to right, 1 through p are assigned to the +’s and left ends of matchings. Assign $\{p + 1, \dots, n\}$ as we read the clan from left to right. When we encounter a $-$, we assign the smallest unused number, and when we encounter the *left* end of a matching, we immediately assign the smallest unused number from $\{p + 1, \dots, n\}$ to the corresponding *right* end of the matching.

Example 3.1. If $\alpha = 122133 \in \text{Clans}_{3,3}$ then $w_\alpha = 125436$.

We construct, in stages, the generic matrix $M_\alpha(z)$ of S_α . First, for $i = 1, \dots, n$:

- (O.1) If $\alpha_i = \pm$, then row i has a 1 in position $w_\alpha(i)$.
- (O.2) If $(i < j)$ is a matching of α , then row i has a 1 in positions $w_\alpha(i)$ and $w_\alpha(j)$.
- (O.3) If $(j < i)$ is a matching of α , then row i has a -1 in position $w_\alpha(j)$ and a 1 in position $w_\alpha(i)$.

A *pivot* is the northmost 1 in each column. (Note that w_α is chosen precisely so that the pivots in the first p columns, and in the last q columns, occur northwest to southeast.) In the first p columns only, set to 0 all entries

- (Z.1) in the same row as a pivot;
- (Z.2) above a pivot (and in that pivot’s column);
- (Z.3) between the pivot 1 and the corresponding -1 of a column (for all columns to which this applies);
or
- (Z.4) right of a -1 .

In the rightmost q columns, set to 0 all entries

- (Z.5) in the same row as a pivot;
- (Z.6) above a pivot (and in that pivot’s column);
- (Z.7) between the pivot 1 and the corresponding 1 below it in the same column (for all columns to which this applies); or
- (Z.8) right of a 1 which is the second 1 in its column.

The remaining entries z_{ij} of $M_\alpha(z)$ are arbitrary, with an exception for each pair of matchings of α which are “in the pattern 1212”. For each such pair of matchings $(i < k)$, $(j < \ell)$ with $i < j < k < \ell$:

- (Z.9) position $(\ell, w_\alpha(i))$ has entry $z_{\ell, w_\alpha(i)}$ (as usual) whereas its “partner” position $(\ell, w_\alpha(k))$ has entry $-z_{\ell, w_\alpha(i)}$.

Example 3.2. Let $\alpha = 1+12-2 \in \text{Clans}_{3,3}$. Then $w_\alpha = 124365$. The 1’s and -1 ’s are first placed into a 6×6 matrix, followed by placement of 0’s as follows:

$$\begin{pmatrix} 1 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ -1 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & -1 & \cdot & 1 & \cdot \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ \cdot & \cdot & 0 & 0 & 0 & 1 \\ \cdot & \cdot & -1 & \cdot & 1 & 0 \end{pmatrix}.$$

Since α has no 1212-patterns, the remaining unspecialized entries of the matrix are arbitrary. Thus S_{1+12-2} is an affine space of dimension 5 (the codimension of Q_{1+12-2} in GL_6/K), with its generic entry being the matrix

$$M_{1+12-2}(z) = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ z_{5,1} & z_{5,2} & 0 & 0 & 0 & 1 \\ z_{6,1} & z_{6,2} & -1 & z_{6,4} & 1 & 0 \end{pmatrix}.$$

Example 3.3. Now, consider $\alpha = 1+21-2$. Here $w_\alpha = 123465$. The placement of 1's and -1 's, and then 0's, is achieved by:

$$\begin{pmatrix} 1 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & 1 & \cdot \\ -1 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & -1 & \cdot & 1 & \cdot \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ \cdot & \cdot & 0 & 0 & 0 & 1 \\ \cdot & \cdot & -1 & \cdot & 1 & 0 \end{pmatrix}.$$

The 1212-pattern in positions $1 < 3 < 4 < 6$ dictates that the unspecialized positions $(6, w_\alpha(1)) = (6, 1)$ and $(6, w_\alpha(4)) = (6, 4)$ are negatives of one another. Thus S_{1+21-2} is an affine space of dimension 4 (the codimension of Q_{1+21-2} in GL_6/K), and

$$M_{1+21-2}(\mathbf{z}) = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ z_{5,1} & z_{5,2} & 0 & 0 & 0 & 1 \\ z_{6,1} & z_{6,2} & -1 & -z_{6,1} & 1 & 0 \end{pmatrix}.$$

Example 3.4. Each 1212-pattern gives rise to a separate identification of coordinates. For instance, when $\alpha = 123123$, the reader can verify that

$$M_{123123}(\mathbf{z}) = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \\ -1 & 0 & 0 & 1 & 0 & 0 \\ z_{5,1} & -1 & 0 & -z_{5,1} & 1 & 0 \\ z_{6,1} & z_{6,2} & -1 & -z_{6,1} & -z_{6,2} & 1 \end{pmatrix}.$$

The following result will be used in Section 4.

Lemma 3.5. $M_\alpha(\mathbf{z})$ invertible over $\mathbb{Q}[\mathbf{z}]$.

Proof. By construction, the matrix is invertible over the field of rational functions $\mathbb{Q}(\mathbf{z})$. The lemma would follow if we could prove that the determinant is a constant (which will be necessarily nonzero). We argue by induction on $n = p' + q'$ that $\det M_{\alpha'}(\mathbf{z}) \in \mathbb{Q}$ for $\alpha' \in \text{Clans}_{p',q'}$. The base case $n = 1, 2$ is easy to check.

Suppose $\gamma \in \text{Clans}_{p,q}$ where $n = p + q > 2$. If $\gamma_1 = +$ consider the submatrix of $M_\alpha(\mathbf{z})$ obtained by striking the first row and column. One checks that this submatrix M' is equal to $M_{\alpha'}(\mathbf{z})$, where α' is α with the first $+$ removed. By induction, $\det M' \in \mathbb{Q}$. Then by cofactor expansion along the first row, since the unique nonzero entry is a 1 in position $(1, 1)$, we have that $\det M = \det M' \in \mathbb{Q}$, as desired.

A similar argument may be used if $\alpha_1 = -$.

Finally, suppose α_1 is the left end of a matching. Let α_r be the right end of said matching. Observe that the submatrix M' obtained by deleting the first row and first column of $M = M_\alpha$ is the matrix $M_{\alpha'}$ where α' is obtained by deleting α_1 and replacing α_r by $-$. Thus by induction $\det M' \in \mathbb{Q}$.

Similarly, consider the submatrix M'' obtained by deleting the first row and $w(r)$ -th column. We observe that by

- cyclically reordering rows $1, 2, \dots, w(r) - 1$ of M'' ; and then
- multiplying the first row by -1 ,

we obtain the matrix $M_{\alpha''}$ where α'' is obtained from α by deleting α_r and replacing α_1 by $+$. Hence, by cofactor expansion along the first row of M , we obtain that $\det M = \det M' \pm \det M'' \in \mathbb{Q}$, as desired. \square

3B. Identification of S_α with the Mars–Springer slice. Let B be the Borel subgroup of upper triangular matrices in GL_n , and let T be the maximal torus of diagonal matrices of GL_n . Recall from Section 2 that in order to define the slice, we must choose a point $x_\alpha = xK/K \in Q_\alpha$ such that $x\theta(x)^{-1} \in N$, where N is the normalizer of T , and where θ is the involution (1). Here, we choose x to be the origin of S_α , meaning the matrix where all z -variables are set to 0. The fact that $x_\alpha = xK/K$ is actually a point of Q_α follows from [Yamamoto 1997, Section 2.2-2], while the fact that $x\theta(x)^{-1} \in N$ is easy to check. Denote by X_α the Mars–Springer slice to Q_α at x_α in G/K , so

$$X_\alpha := (U^- \cap \psi(U^-)) \cdot x_\alpha = ((U^- \cap \psi(U^-))x)/K.$$

Theorem 3.6. *We have an isomorphism $X_\alpha \cong S_\alpha$.*

Proof. Recall from Section 2B that the map $\psi : G \rightarrow G$ is defined by $c_\eta \circ \theta$, where $\eta = x\theta(x)^{-1} \pmod{T}$.

Given a clan γ , let $\mathcal{I}(\alpha)$ denote the *underlying involution* for α ; this is the permutation in S_n such that $\mathcal{I}(\alpha)(j) = j$ if $\alpha_j = \pm$, while, if $(i < j)$ is a matching of α , $\mathcal{I}(\alpha)(i) = j$ and $\mathcal{I}(\alpha)(j) = i$. The map \mathcal{I} is the concrete realization for $(G, K) = (\mathrm{GL}_n, \mathrm{GL}_p \times \mathrm{GL}_q)$ of the map defined by Richardson and Springer [1990] associating a twisted involution (which is an involution in this case) to every orbit.

Claim 3.7. *As an element of $\mathcal{W} = S_n$, we have $\eta = \mathcal{I}(\alpha)$.*

Proof. The matrix computation $x\theta(x)^{-1} = xI_{p,q}x^{-1}I_{p,q}$ is straightforward, and results in a monomial matrix representing the involution described above. \square

Claim 3.8. *The group $U^- \cap \psi(U^-)$ consists of the unipotent matrices having 1’s on the diagonal, 0’s above the diagonal, 0’s below the diagonal in all positions (i, j) with $i > j$ such that $\eta(i) < \eta(j)$, and arbitrary entries in all other positions below the diagonal.*

Proof. U^- is the set of lower triangular unipotent $n \times n$ matrices. U^- is θ -stable, so θ maps it isomorphically to itself, meaning that $\psi(U^-) = c_\eta(U^-)$. By Claim 3.7, $\eta = \eta^{-1}$. Thus conjugation by η sends the affine coordinate at position (i, j) ($i > j$) to position $(\eta(i), \eta(j))$. The claim follows. \square

Denote by S'_α the set of $n \times n$ matrices that satisfy (O.1), (O.2), (O.3), (Z.2), (Z.4), (Z.6), (Z.8), (Z.9), and the following condition:

(Z.10) If $(i < j)$ is any matching of α , and $i < k < j$, then:

- if k is the left end of a matching $(k < \ell)$ with $j < \ell$, then the entries at positions $(k, w_\alpha(i))$ and $(k, w_\alpha(j))$ are equal;
- otherwise, the entries at positions $(k, w_\alpha(i))$ and $(k, w_\alpha(j))$ are both zero.

Claim 3.9. $S'_\alpha = (U^- \cap \psi(U^-))x.$

Proof. By a direct but tedious matrix computation using Claim 3.8, one checks that $(U^- \cap \psi(U^-))x \subseteq S'_\alpha$. The space $(U^- \cap \psi(U^-))x$ is evidently an affine space of dimension $\binom{n}{2} - \ell(\eta)$, where $\ell(\cdot)$ denotes Coxeter length. The space S'_α is also visibly an affine space, and a straightforward combinatorial argument shows that its dimension is also $\binom{n}{2} - \ell(\eta)$. Thus the aforementioned containment is in fact an equality. \square

Claim 3.10. $S_\alpha \subseteq S'_\alpha.$

Proof. This follows from the fact that (Z.3) and (Z.7) together imply (Z.10). \square

The right action of K on G is by column operations which separately act on the first p columns and the last q columns of any matrix in G . Given a matrix $M' \in S'_\alpha$, we now describe an algorithm for column reducing M' via this K -action to another matrix M of a particular form.

Algorithm 3.11. Begin with the first p columns of M' . For each row $i = 1, 2, \dots, n$, determine whether there is a pivot in row i . If not, move to the next row. If so, then this pivot is at $(i, w_\alpha(i))$. Now start with column $j = 1$, and move right toward column $j = w_\alpha(i) - 1$. For each such j , if entry (i, j) is zero, move to the next column. Otherwise, entry (i, j) is $\lambda_{i,j} \neq 0$. Replace column C_j with $C_j - \lambda_{i,j} \cdot C_i$. Repeat the same procedure using the last q columns.

Let M be the output of Algorithm 3.11.

Claim 3.12. *The first p columns and the last q columns of M are in reduced column echelon form, meaning that all entries in the same row as a pivot (except for the pivot) are zero. M is the unique matrix in the K -orbit of M' with this property.*

Proof. Consider first the leftmost p columns. Algorithm 3.11 brings to zero any entry in the same row as a pivot at $(i, w_\alpha(i))$ and to the left of $(i, w_\alpha(i))$. Since M' satisfies (Z.2), any entry to the right of a pivot is already zero: there is a pivot in every column, and by our choice of w_α , any pivot right of column $w_\alpha(i)$ occurs in a row $j > i$.

For the rightmost q columns, one argues identically, using the fact that M' satisfies (Z.6).

Uniqueness of M follows from the uniqueness of reduced column echelon form (for each of the two submatrices of M). \square

Claim 3.13. *The matrix M is an element of S_α .*

Proof. By Claim 3.12, M satisfies (Z.1) and (Z.5). We show that each step performed by Algorithm 3.11 preserves properties (O.1), (O.2), (O.3), (Z.2), (Z.4), (Z.6), and (Z.8), all of which are satisfied by M' . Hence M also possesses these properties. We also show that once Algorithm 3.11 is complete, the

resulting matrix also satisfies (Z.3), (Z.7), and (Z.9). This will show that M possesses all the properties (O.1)–(O.3) and (Z.1)–(Z.9).

(O.1): Suppose that $c_i = +$. There is a pivot 1 at $(i, w_\alpha(i))$. Any step of Algorithm 3.11 affecting column $C_{w_\alpha(i)}$ is performed to make zero an entry $\lambda \neq 0$ at $(j, w_\alpha(i))$ for $j > i$ (using the pivot in row j). Such a step replaces column $C_{w_\alpha(i)}$ by $C_{w_\alpha(i)} - \lambda \cdot C_{w_\alpha(j)}$. Since $j > i$, by (Z.2), the entry at $(i, w_\alpha(j))$ is zero, so the 1 at $(i, w_\alpha(i))$ remains a 1 after this column operation.

When $c_i = -$, one argues identically using the last q columns and (Z.6). Thus (O.1) is preserved by all steps of our algorithm, as desired.

(O.2): The argument is identical to that for (O.1).

(O.3): Suppose $(i < j)$ is a matching. First consider the -1 at $(j, w_\alpha(i))$. Since entries to the right of this -1 are zero by (Z.4), the -1 is clearly unchanged by steps of the algorithm. Similarly, the 1 at $(j, w_\alpha(j))$ is unchanged, by (Z.8).

(Z.2): Given a zero entry above a pivot 1, any entry to its right is also zero by (Z.2) since such an entry is above a different pivot 1. Thus any step of the algorithm leaves the first of these two zero entries unchanged, as needed.

(Z.3): Suppose $(i < j)$ is a matching. There are a 1 and a -1 in positions $(i, w_\alpha(i))$ and $(j, w_\alpha(i))$, respectively. Let $i < k < j$. We must check that M is zero at $(k, w_\alpha(i))$. There are two cases based on (Z.10):

Case 1: (The $(k, w_\alpha(i))$ entry of M' is arbitrary.) This case occurs if and only if α_k is the left end of a matching $(k < \ell)$ such that $i < k < j < \ell$. Thus there is a pivot 1 in row k at position $(k, w_\alpha(k))$. Since M satisfies (Z.1), entry $(k, w_\alpha(i))$ of M must be zero.

Case 2: (The $(k, w_\alpha(i))$ entry of M' is required to be zero.) Consider a step of Algorithm 3.11 altering column $C_{w_\alpha(i)}$. It does so by replacing $C_{w_\alpha(i)}$ with $C_{w_\alpha(i)} - \lambda \cdot C_{w_\alpha(\ell)}$ for ℓ such that $\ell > i$ and $w_\alpha(\ell) > w_\alpha(i)$. This step is done if and only if there is a pivot 1 at $(\ell, w_\alpha(\ell))$ and a nonzero entry λ in position $(\ell, w_\alpha(i))$. We examine the effect of this step on entry $(k, w_\alpha(i))$.

If $\ell > k$, then entry $(k, w_\alpha(\ell))$ is zero by (Z.2), so entry $(k, w_\alpha(i))$ remains zero after the step in question. So suppose that $i < \ell < k$. If entry $(\ell, w_\alpha(i))$ is $\lambda \neq 0$, then by (Z.10) this must be because α_ℓ is the left end of a matching $(\ell < r)$ such that $i < \ell < j < r$. If the entry at $(k, w_\alpha(\ell))$ is zero, then the entry at $(k, w_\alpha(i))$ remains zero after our step, and we are done. So suppose it is not. Then by (Z.10) again, α_k is the left end of a matching $(k < s)$ with $\ell < k < r < s$. But then $i < k < j < s$, contradicting our assumption that (Z.10) required entry $(k, w_\alpha(i))$ of M' to be zero. Thus this situation cannot occur.

(Z.4): Suppose that M' has a -1 at position (i, j) with $1 \leq j \leq p$. Since M' satisfies (Z.4), any entry in a position (i, k) with $j < k \leq p$ is zero at the outset. Moreover, again by (Z.4), any entry in a position (i, ℓ) with $k < \ell \leq p$ is also zero at the outset. Thus all steps of Algorithm 3.11 leave the 0 in position (i, k) unchanged.

(Z.6): We argue as with (Z.2) above, but using the rightmost q columns.

(Z.7): We argue as with (Z.3).

(Z.8): We argue as with (Z.4).

(Z.9): Suppose α has a 1212 pattern consisting of matchings $(i < j)$ and $(k < \ell)$ with $i < k < j < \ell$. Then by (Z.10), in positions $(k, w_\alpha(i))$ and $(k, w_\alpha(j))$ of M' , we have arbitrary but equal entries; assume that both entries are equal to λ_1 . Moreover, by (Z.9), in positions $(\ell, w_\alpha(i))$ and $(\ell, w_\alpha(j))$, we have arbitrary entries that are negatives of one another. Suppose that the entry at $(\ell, w_\alpha(i))$ is λ_2 , and that the entry at $(\ell, w_\alpha(j))$ is $-\lambda_2$. Consider two steps performed by Algorithm 3.11:

(i) Replace column $C_{w_\alpha(i)}$ by $C_{w_\alpha(i)} - \lambda_1 \cdot C_{w_\alpha(k)}$ to make the entry at $(k, w_\alpha(i))$ zero.

(ii) Replace column $C_{w_\alpha(j)}$ by $C_{w_\alpha(j)} - \lambda_1 \cdot C_{w_\alpha(\ell)}$ to make the entry at $(k, w_\alpha(j))$ zero.

Step (i) replaces the λ_2 at $(\ell, w_\alpha(i))$ by $\lambda_2 + \lambda_1$ (since there is a -1 in position $(\ell, w_\alpha(k))$). Meanwhile, step (ii) replaces the $-\lambda_2$ at $(\ell, w_\alpha(j))$ by $-\lambda_2 - \lambda_1$ (since there is a 1 in position $(\ell, w_\alpha(\ell))$). Thus after (i) and (ii), (Z.9) holds for the 1212 pattern $i < k < j < \ell$.

To conclude (Z.9) holds, it remains to show:

Subclaim 3.14. *Steps (i) and (ii) are the only steps of Algorithm 3.11 that can change either of the entries $(\ell, w_\alpha(i))$ or $(\ell, w_\alpha(j))$.*

Proof. Suppose there is a pivot 1 (among the first p columns) at $(r, w_\alpha(r))$, and suppose that there is a nonzero entry at $(r, w_\alpha(i))$. There are two possibilities. First, suppose $w_\alpha(r) > w_\alpha(k)$. In this case, $(\ell, w_\alpha(r))$ must be zero by (Z.4). Thus a column operation involving columns $w_\alpha(i)$ and $w_\alpha(r)$ leaves $(\ell, w_\alpha(i))$ unchanged, as desired. Otherwise, $w_\alpha(i) < w_\alpha(r) < w_\alpha(k)$. Then $i < r < j$, so by (Z.10) the entry at $(r, w_\alpha(i))$ can be nonzero only if there is a 1212 pattern formed by matchings $(i < j)$ and $(r < s)$ with $i < r < j < s$. However, this situation is covered by the argument preceding this subclaim. A similar analysis of the last q columns (using either (Z.8) or (Z.10)) shows that there are no steps of our algorithm that alter position $(\ell, w_\alpha(j))$. \square

This completes the proof of Claim 3.13. \square

By Claim 3.13, we can now define a map

$$\phi : X_\alpha \rightarrow S_\alpha$$

which sends the point $M' \cdot K$ of X_α to the matrix M , retaining the notation above. Its inverse, as a set map, is the restriction of the natural projection $\pi : G \rightarrow G/K$ to S_α . (Note that this restriction does take values in $X_\alpha = S'_\alpha/K$ by Claim 3.10.) Abusing notation, we denote this restriction simply by π . Then letting M' and M be as above, it is indeed clear that

$$\pi(\phi(M' \cdot K)) = \pi(\phi(M \cdot K)) = \pi(M) = M \cdot K = M' \cdot K$$

and that

$$\phi(\pi(M)) = \phi(M \cdot K) = M,$$

so that π and ϕ are inverse as set maps.

It remains to check that ϕ and π are actually morphisms of algebraic varieties. Clearly π is a morphism, since it is a restriction of the natural map $G \rightarrow G/K$. The map ϕ , on the other hand, is the restriction of an algebraic section of π to X_α . Indeed, it takes a splitting $V_1 \oplus V_2$ and produces a matrix M whose first p columns are a basis for V_1 and whose last q columns are a basis for V_2 . These two bases are of the form described in Section 2C when we take S to be the positions of the $+$'s and left endpoints of α and T to be the position of the $-$'s and left endpoints of α . Thus $X_\alpha \subseteq (U_S \times U_T) \cap G/K$, and indeed the map ϕ is the restriction to X_α of the section $\phi_{S,T} : (U_S \times U_T) \cap G/K \rightarrow G$ described in Section 2C. \square

4. The Mars–Springer variety and ideal

For $\gamma \in \text{Clans}_{p,q}$, define the following collections of nonnegative integers [Wyser 2016]:

- (1) $\gamma(i; +) :=$ number of matchings and $+$ signs among the first i positions of γ (for $i = 1, \dots, n$);
- (2) $\gamma(i; -) :=$ number of matchings and $-$ signs among the first i positions of γ (for $i = 1, \dots, n$); and
- (3) $\gamma(i; j) :=$ number of matchings whose left endpoint is weakly left of i and whose right endpoint is strictly right of j (for $1 \leq i < j \leq n$).

Given any invertible $n \times n$ matrix M and any pair of indices $i < j$, we define an auxiliary $n \times (i + j)$ matrix $M^{[i;j]}$ as follows. The first i columns of $M^{[i;j]}$ are formed by taking the first i columns of M^{-1} and zeroing out all entries in the last q rows. The last j columns of $M^{[i;j]}$ are the first j columns of M^{-1} , unaltered.

Example 4.1. Consider $\alpha = 1+--+1 \in \text{Clans}_{3,3}$. Then

$$M_{1+--+1}(\mathbf{z}) = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{3,2} & 0 & 0 & 1 & 0 \\ 0 & z_{4,2} & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & z_{5,5} & z_{5,6} \\ -1 & 0 & 0 & 1 & 0 & 0 \end{pmatrix},$$

as the reader can verify. The inverse of this matrix is

$$M_{1+--+1}(\mathbf{z})^{-1} = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 & 0 & -\frac{1}{2} \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & z_{3,2}z_{5,5} + z_{4,2}z_{5,6} & -z_{5,5} & -z_{5,6} & \frac{1}{2} & 0 \\ \frac{1}{2} & 0 & 0 & 0 & 0 & \frac{1}{2} \\ 0 & -z_{3,2} & 1 & 0 & 0 & 0 \\ 0 & -z_{4,2} & 0 & 1 & 0 & 0 \end{pmatrix}.$$

To obtain the auxiliary matrix $M_{1+--+1}(\mathbf{z})^{[2;4]}$, one takes the first 2 columns of $M_{1+--+1}(\mathbf{z})^{-1}$, zeroes out their last $q = 3$ rows, and then concatenates to this the first 4 columns of $M_{1+--+1}(\mathbf{z})^{-1}$, unaltered.

The result is

$$M_{1+---+1}(\mathbf{z})^{[2;4]} = \begin{pmatrix} \frac{1}{2} & 0 & \frac{1}{2} & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & z_{3,2}z_{5,5} + z_{4,2}z_{5,6} & 0 & z_{3,2}z_{5,5} + z_{4,2}z_{5,6} & -z_{5,5} & -z_{5,6} \\ 0 & 0 & \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -z_{3,2} & 1 & 0 \\ 0 & 0 & 0 & -z_{4,2} & 0 & 1 \end{pmatrix}.$$

Now define the *Mars–Springer variety* $\mathcal{N}_{\gamma,\alpha}$ to be the reduced subscheme of S_α whose points are the matrices M satisfying all of the following rank conditions:

- (R.1) the rank of the southwest $(n - i) \times p$ submatrix of M is at most $p - \gamma(i; +)$ (for $i = 1, \dots, n$);
- (R.2) the rank of the southeast $(n - i) \times q$ submatrix of M is at most $q - \gamma(i; -)$ (for $i = 1, \dots, n$); and
- (R.3) the rank of $M^{[i;j]}$ is at most $j + \gamma(i; j)$ (for all $1 \leq i < j \leq n$).

Note that $\mathcal{N}_{\gamma,\alpha}$ is clearly defined *set-theoretically* by a certain collection of determinants. Indeed, define the *Mars–Springer ideal* $\mathcal{I}_{\gamma,\alpha}$ to be the ideal of $\mathbb{C}[\mathbf{z}]$ generated by:

- (1) all minors of the southwest $(n - i) \times p$ submatrix of $M_\alpha(\mathbf{z})$ of size $p - \gamma(i; +) + 1$ (for $i = 1, \dots, n$);
- (2) all minors of the southeast $(n - i) \times q$ submatrix of $M_\alpha(\mathbf{z})$ of size $q - \gamma(i; -) + 1$ (for $i = 1, \dots, n$); and
- (3) all minors of $M_\alpha(\mathbf{z})^{[i;j]}$ of size $j + \gamma(i; j) + 1$ (for all $1 \leq i < j \leq n$).

(By Lemma 3.5, the minors of (3) above are in fact polynomials.)

Example 4.2. Consider the ideal $\mathcal{I}_{123231,1+---+1}$. One can check that, with the exception of the condition that the rank of $M^{[2;4]}$ be at most 5, the conditions defining $\mathcal{N}_{123231,1+---+1}$ are all actually nonconditions. Thus the ideal $\mathcal{I}_{123231,1+---+1}$ is generated by the determinant of the 6×6 matrix $M_{1+---+1}(\mathbf{z})^{[2;4]}$ computed above. One checks that this determinant equals (up to a scalar multiple) $z_{3,2}z_{5,5} + z_{4,2}z_{5,6}$. Thus

$$\mathcal{I}_{123231,1+---+1} = \langle z_{3,2}z_{5,5} + z_{4,2}z_{5,6} \rangle,$$

which defines a singular quadric hypersurface.

Theorem 4.3. $\mathcal{N}_{\gamma,\alpha} \cong W_\gamma \cap X_\alpha$, the *Mars–Springer slice* to Q_α at x_α in W_γ .

Proof. Recall that points of G/K are splittings $S = V_1 \oplus V_2$ of \mathbb{C}^{p+q} where $\dim(V_1) = p$ and $\dim(V_2) = q$. The B -orbit closure $W_\gamma \subseteq G/K$ has a description due to the second author, which we now state. Define $\pi_S : \mathbb{C}^n \rightarrow V_1$ to be the projection onto the first summand of S with kernel V_2 , the second summand of S .

Theorem 4.4 [Wyser 2016]. *Let E_\bullet be the standard coordinate flag on \mathbb{C}^n , with E_i the span of the first i standard basis vectors. Then $W_\gamma \subseteq G/K$ is precisely the set of splittings $S = V_1 \oplus V_2$ satisfying all of the following incidence conditions with E_\bullet :*

- (C.1) $\dim(V_1 \cap E_i) \geq \gamma(i; +)$ for $i = 1, \dots, n$;

(C.2) $\dim(V_2 \cap E_i) \geq \gamma(i; -)$ for $i = 1, \dots, n$;

(C.3) $\dim(\pi_S(E_i) + E_j) \leq j + \gamma(i; j)$ for $1 \leq i < j \leq n$.

Let $\phi : X_\alpha \rightarrow S_\alpha$ be the isomorphism from Theorem 3.6. It suffices to show that the image of

$$\phi|_{W_\gamma \cap X_\alpha} : (W_\gamma \cap X_\alpha) \rightarrow S_\alpha$$

is precisely $\mathcal{N}_{\gamma, \alpha}$.

The map ϕ is a bijection, taking a splitting $\mathcal{S} = V_1 \oplus V_2 \in X_\alpha$ to a matrix $M \in S_\alpha$, the span of whose first p columns is V_1 , and the span of whose last q columns is V_2 . Therefore it suffices to show that under this bijection, \mathcal{S} satisfies (C.1)–(C.3) if and only if $M = \phi(\mathcal{S})$ satisfies (R.1)–(R.3).

The equivalence of (R.1) to (C.1) and of (R.2) to (C.2) is immediate.

To see the equivalence of (C.3) to (R.3) we argue as follows. Condition (C.3) states that

$$\dim(\pi_S(E_i) + E_j) \leq j + \gamma(i; j).$$

If $M \in S_\alpha$ represents a splitting \mathcal{S} , then the matrix for π_S can be written as MPM^{-1} , where P is the diagonal matrix with p 1's on the diagonal followed by q 0's. Hence $\pi_S(E_i) + E_j$ is the span of the first i columns of MPM^{-1} and the first j columns of the identity matrix I . Therefore,

$$\mathcal{S} \text{ satisfies (C.3)} \iff \text{rank}(M^{\overline{i;j}}) \leq j + \gamma(i; j), \quad (4)$$

where $M^{\overline{i;j}}$ is the $n \times (i+j)$ matrix whose first i columns are the first i columns of MPM^{-1} and whose last j columns are the first j columns of I .

Since M^{-1} is invertible,

$$\text{rank}(M^{[i;j]}) = \text{rank}(M^{-1}M^{\overline{i;j}}) = \text{rank}(M^{\overline{i;j}}). \quad (5)$$

Combining (4) and (5) we conclude

$$\begin{aligned} \mathcal{S} \text{ satisfies (C.3)} &\iff \text{rank}(M^{[i;j]}) \leq j + \gamma(i; j) \quad \text{for all } i, j \text{ with } 1 \leq i < j \leq n \\ &\iff M \text{ satisfies (R.3),} \end{aligned}$$

as desired. □

Corollary 4.5. *For any clans $\alpha \leq \gamma$, the Mars–Springer variety $\mathcal{N}_{\gamma, \alpha}$ is irreducible.*

Proof. This is immediate by Theorem 4.3 combined with Corollary 2.7. □

The key to our proof of Theorem 1.3 is:

Theorem 4.6 (isomorphism theorem). *If $[\beta, \theta]$ interval pattern contains $[\alpha, \gamma]$, then*

$$\mathcal{N}_{\theta, \beta} \cong \mathcal{N}_{\gamma, \alpha}.$$

This will be proved in Section 6.

5. Combinatorics of interval embeddings

Denote Bruhat order on clans by $<$. Let $\gamma \triangleleft \tau$ denote a covering relation, so $\gamma \triangleleft \tau$ if $\gamma < \tau$ and there is no clan β such that $\gamma < \beta < \tau$. Equivalently, $\gamma \triangleleft \tau$ if $\gamma < \tau$ and $\ell(\tau) - \ell(\gamma) = 1$.

We need the following result of the second author describing the covering relations \triangleleft :

Theorem 5.1 [Wyser 2016]. *Let $\gamma, \tau \in \text{Clans}_{p,q}$ such that $\gamma < \tau$. Then there exists $\gamma' \in \text{Clans}_{p,q}$ such that $\gamma < \gamma' \leq \tau$, where γ' is obtained from γ by one of the following operations on patterns in γ :*

- (T.1) $+- \mapsto 11$,
- (T.2) $-+ \mapsto 11$,
- (T.3) $11+ \mapsto 1+1$,
- (T.4) $11- \mapsto 1-1$,
- (T.5) $+11 \mapsto 1+1$,
- (T.6) $-11 \mapsto 1-1$,
- (T.7) $1122 \mapsto 1212$,
- (T.8) $1122 \mapsto 1+-1$,
- (T.9) $1122 \mapsto 1-+1$,
- (T.10) $1212 \mapsto 1221$.

Any ordered pair $\gamma \mapsto \gamma'$ of clans obtained by one of the operations (T.1)–(T.10) is called a *transposition*. Thus:

Corollary 5.2 [Wyser 2016]. *$\gamma \triangleleft \gamma'$ if and only if $\gamma \mapsto \gamma'$ is a transposition such that $\ell(\gamma') = \ell(\gamma) + 1$.*

For any clan γ , denote by M_γ the set of matchings of γ . Call a matching $(a < b) \in M_\gamma$ with $a < i < b < j$ incoming to $(i < j) \in M_\gamma$. For $(i < j) \in M_\gamma$, let

$$I(i, j, \gamma) := \#\{(a < b) \in M_\gamma \mid (a < b) \text{ is incoming to } (i < j)\},$$

and let

$$C(i, j, \gamma) := j - i - I(i, j, \gamma).$$

Then by [Yamamoto 1997],

$$\ell(\gamma) := \sum_{(a < b) \in M_\gamma} C(a, b, \gamma).$$

We will need the following combinatorial fact:

Proposition 5.3. *Suppose $\gamma \mapsto \gamma'$ is a transposition. Then $\ell(\gamma) < \ell(\gamma')$.*

Proof. For brevity, we prove the hardest case (T.7). Arguments for the remaining cases of (T.1)–(T.10) are similar in nature.

Suppose $\gamma \mapsto \gamma'$ via (T.7). Say the 1122 pattern occurs at positions $i < j < k < \ell$. In order to compare $\ell(\gamma)$ to $\ell(\gamma')$, we consider other matchings ($a < b$) and compare $I(a, b, \gamma')$ with $I(a, b, \gamma)$, $I(i, k, \gamma')$ with $I(i, j, \gamma)$, and $I(j, \ell, \gamma')$ with $I(k, \ell, \gamma)$.

There are 15 different possible orientations of a matching ($a < b$) relative to ($i < j$) and ($k < \ell$) to consider.

Case 1: (The configuration $a < i < j < b < k < \ell$.) Here

$$I(a, b, \gamma') = I(a, b, \gamma), \quad I(j, \ell, \gamma') = I(k, \ell, \gamma) + 1,$$

and

$$I(i, k, \gamma') = I(i, j, \gamma) + 1.$$

Indeed, ($a < b$) was incoming to neither ($i < j$) nor ($k < \ell$) in γ , but it is incoming to both ($i < k$) and ($j < \ell$) in γ' . So each such matching ($a < b$) accounts for a decrease in length of 2 as we transpose from γ to γ' .

Case 2: (The configuration $i < j < a < k < \ell < b$.) Now,

$$I(a, b, \gamma') = I(a, b, \gamma) + 2.$$

Indeed, in γ neither ($i < j$) nor ($k < \ell$) are incoming to ($a < b$), but in γ' both ($i < k$) and ($j < \ell$) are incoming to it. On the other hand, the move in this configuration does not introduce any new incoming matchings to ($i < k$) or ($j < \ell$). So each matching ($a < b$) in this configuration also accounts for a decrease in length of 2 as we transpose from γ to γ' .

Case 3: (The configuration $i < a < j < b < k < \ell$.) When we transpose from γ to γ' , ($a < b$) loses an incoming matching: ($i < j$) is incoming to ($a < b$) in γ , but ($i < k$) is not incoming to ($a < b$) in γ' . However, ($j < \ell$) gains an incoming matching compared to ($k < \ell$), as ($a < b$) is not incoming to ($k < \ell$) in γ , but it is incoming to ($j < \ell$) in γ' . Thus while the transposition causes a change in the number of matchings incoming to ($a < b$) and in the number of matchings to which ($a < b$) is incoming, the net effect is zero.

By arguing similarly to Case 3 for the remaining 12 configurations, one sees that Case 1 and Case 2 are the only two which result in a net change in length coming from changes in the number of incoming matchings.

The only change in the number of incoming matchings which we have not accounted for is that after the move, ($j < \ell$) has also added the incoming matching ($i < k$) in γ' , whereas ($k < \ell$) did not have the incoming matching ($i < j$) in γ . This causes an additional decrease by 1 in length as we perform this transposition.

Finally, we must account for the change in total lengths of matchings caused by the transposition. This is $(k - i) + (\ell - j) - ((j - i) + (\ell - k)) = 2(k - j)$.

Putting this all together, we conclude that

$$\begin{aligned} \ell(\gamma') - \ell(\gamma) = 2(k - j) - 1 - 2(\#\{(a < b) \in M_\gamma \mid a < i < j < b < k < \ell\} \\ + \#\{(a < b) \in M_\gamma \mid i < j < a < k < \ell < b\}). \end{aligned} \quad (6)$$

Now, the sum in parentheses is clearly at most $k - j - 1$, which implies that the right-hand side is at least 1. Thus $\ell(\gamma') > \ell(\gamma)$, as desired. \square

Say $[\alpha, \gamma]$ *merely embeds* into $[\beta, \theta]$ if the pair of intervals satisfies the definition of an interval pattern embedding, *except* possibly for the length requirement

$$\ell(\gamma) - \ell(\alpha) = \ell(\theta) - \ell(\beta).$$

Given any α , we define $\Phi(\alpha)$ to be the unique clan such that $[\alpha, \gamma]$ merely embeds in $[\Phi(\alpha), \theta]$, so $\Phi(\alpha)$ agrees with α on the set of embedding indices I and agrees with θ on $\{1, \dots, n\} \setminus I$.

Theorem 5.4. *Assume $[\alpha, \gamma]$ merely embeds into $[\Phi(\alpha), \theta]$.*

(I) $\ell(\gamma) - \ell(\alpha) \leq \ell(\theta) - \ell(\Phi(\alpha))$.

(II) *Further suppose $[\alpha, \gamma]$ interval embeds into $[\Phi(\alpha), \theta]$. If $\alpha \triangleleft \alpha' \leq \gamma$ then*

$$\Phi(\alpha) \triangleleft \Phi(\alpha') \leq \theta.$$

Proof. (I): Pick a chain of covering transpositions

$$\alpha = \alpha^{(0)} \mapsto \alpha^{(1)} \mapsto \dots \mapsto \alpha^{(m)} = \gamma.$$

This induces a chain from $\Phi(\alpha)$ to θ by using the same transpositions (relative to the embedding). By Proposition 5.3, each transposition in the latter chain increases the length by at least one, from which the statement follows.

(II): There exists a covering transposition from α to α' , followed by a chain of covering transpositions from α' to γ . Thus $\Phi(\alpha)$ is at least related to $\Phi(\alpha')$ by the same transposition (relative to the embedding), and the chain from α' to γ induces a chain $\Phi(\alpha') \rightarrow \dots \rightarrow \theta$. Hence we have

$$\Phi(\alpha) < \Phi(\alpha') \leq \theta.$$

It remains to show

$$\Phi(\alpha') \triangleleft \Phi(\alpha).$$

By Proposition 5.3, we have

$$\ell(\Phi(\alpha')) - \ell(\Phi(\alpha)) \geq 1.$$

We are done unless

$$\ell(\Phi(\alpha')) - \ell(\Phi(\alpha)) \geq 2,$$

so assume this. Let

$$k := \ell(\gamma) - \ell(\alpha) = \ell(\theta) - \ell(\Phi(\alpha)).$$

Since $[\alpha', \gamma]$ merely embeds into $[\Phi(\alpha'), \theta]$, we obtain a contradiction:

$$\begin{aligned} k - 1 &= \ell(\gamma) - \ell(\alpha') \leq \ell(\theta) - \ell(\Phi(\alpha')) \\ &= \ell(\theta) - \ell(\Phi(\alpha)) - (\ell(\Phi(\alpha')) - \ell(\Phi(\alpha))) \\ &\leq k - 2, \end{aligned}$$

where for the first inequality we have used (I). □

Corollary 5.5. *If $[\alpha, \gamma] \hookrightarrow [\Phi(\alpha), \theta]$ and $\alpha \leq \alpha' \leq \gamma$, then $[\alpha', \gamma] \hookrightarrow [\Phi(\alpha'), \theta]$.*

Proof. By Theorem 5.4(II),

$$\Phi(\alpha) \leq \Phi(\alpha') \leq \theta.$$

Hence $[\alpha', \gamma]$ and $[\Phi(\alpha'), \theta]$ have the same length difference, as desired. □

Corollary 5.6. *If θ avoids $[\alpha', \gamma]$ then θ avoids $[\alpha, \gamma]$ for all $\alpha \leq \alpha'$.*

Proof. This is the contrapositive of Corollary 5.5, combined with induction on $\ell(\alpha') - \ell(\alpha)$. □

For the next lemma, assume that $[\alpha, \gamma]$ interval embeds into $[\beta, \theta]$, let I be the set of indices for the embedding, and let $J = \{1, \dots, n\} \setminus I$ be the set of indices not involved in the embedding.

Lemma 5.7. (I) *If β_j (and hence θ_j) is a sign for some $j \in J$, then*

$$\beta(j; +) = \theta(j; +) \quad \text{and} \quad \beta(j; -) = \theta(j; -).$$

(II) *If β_a and β_b are a matched pair for some $a, b \in J$, then*

$$\beta(a; b) = \theta(a; b).$$

Example 5.8. Let

$$[\alpha, \gamma] = [1212, 1221] \subseteq \text{Clans}_{2,2} \quad \text{and} \quad [\beta, \theta] = [1+212, 1+221] \subseteq \text{Clans}_{3,2}.$$

Both are intervals of length 1, and hence $[\alpha, \gamma] \hookrightarrow [\beta, \theta]$. Here

$$n = 5, \quad I = \{1, 3, 4, 5\}, \quad \text{and} \quad J = \{2\}.$$

This is an instance of Lemma 5.7(I) since $\beta_2 = \theta_2 = +$, $\beta(2; +) = \theta(2; +) = 1$, and $\beta(2; -) = \theta(2; -) = 0$.

Example 5.9. Recall Example 1.1, where

$$[\alpha, \gamma] = [+--+ , 1212] \quad \text{and} \quad [\beta, \theta] = [1+--+1, 123231].$$

Here

$$n = 6, \quad I = \{2, 3, 4, 5\}, \quad \text{and} \quad J = \{1, 6\}.$$

This is an instance of Lemma 5.7(II) since $a = 1$ and $b = 6$ are matched and $\beta(1; 6) = \theta(1; 6) = 0$.

Proof of Lemma 5.7. The proof is by induction on

$$k = \ell(\gamma) - \ell(\alpha) = \ell(\theta) - \ell(\beta).$$

The base case, $k = 0$, when $\theta = \beta$, is trivial, so we may assume $k \geq 1$. Suppose that

$$\alpha \triangleleft \alpha' \leq \gamma.$$

By Corollary 5.5,

$$[\alpha', \gamma] \hookrightarrow [\beta', \theta], \quad \text{where } \beta' = \Phi(\alpha').$$

For the statement (I), by induction,

$$\beta'(j; \pm) = \theta(j; \pm).$$

Next, clearly

$$[\alpha, \alpha'] \hookrightarrow [\beta, \beta'].$$

If we know that

$$\beta(j; \pm) = \beta'(j; \pm),$$

then we are done. Hence we are reduced to the case $k = 1$. The statement (II) similarly reduces to this situation.

Therefore we prove both claims when $k = 1$. To do this, one must analyze each possible covering transposition $\beta \triangleleft \beta'$. As with the proof of Proposition 5.3, for brevity we give the details only for the hardest case (T.7). The other cases are similar.

Suppose the 1122 pattern of β occurs at

$$i < j < k < \ell.$$

From the proof of Proposition 5.3, we recall from (6) that $\ell(\beta') - \ell(\beta) = 1$ if and only if each position between j and k is either the right endpoint of a matching ($a < b$) with

$$a < i < j < b < k < \ell$$

or the left endpoint of a matching ($a < b$) with

$$i < j < a < k < \ell < b.$$

For (I), let $m \in J$ be such that $\beta_m = \beta'_m$ are the same sign. If $m < j$ or $m > k$, then

$$\beta(m; \pm) = \beta'(m; \pm).$$

Since $\beta \triangleleft \beta'$, by the observation of the previous paragraph, we cannot have $j < m < k$.

For (II), for $(a < b) \in M_\beta$, one checks that $\beta(a; b) = \beta'(a; b)$ unless $(a < b)$ is in one of the three configurations

$$i < a < j < b < k < \ell, \quad i < j < a < b < k < \ell, \quad \text{or} \quad i < j < a < k < b < \ell.$$

Since $\beta < \beta'$, none of these configurations are possible, again by the above observation. \square

6. Proof of Theorems 4.6 and 1.3

Assume that $[\alpha, \gamma]$ interval embeds into $[\beta, \theta]$ and that J is the set of indices *not* involved in this embedding. Let $w := w_\beta$ be the permutation associated to the clan β as described in Section 3A.

Proposition 6.1. *For any point $M \in \mathcal{N}_{\theta, \beta}$ and $j \in J$, all the entries in the j -th row and the $w(j)$ -th column of M are 0 except for the required ± 1 specified by (O.1), (O.2) or (O.3).*

Assuming Proposition 6.1, whose proof is deferred to Section 6C, we prove Theorems 4.6 and 1.3 in Sections 6A and 6B.

6A. Proof of Theorem 4.6: Define a map

$$\Psi : \mathcal{N}_{\theta, \beta} \rightarrow S_\alpha$$

by deleting the rows j and columns $w(j)$ for $j \in J$. Proposition 6.1 says that Ψ is injective as a map of sets. Hence it is injective as a map of varieties, since it is the restriction of a linear map.

If we can show that

$$\text{Im}(\Psi) \subseteq \mathcal{N}_{\gamma, \alpha}, \tag{7}$$

we are done. Indeed, this containment is then an equality since $\mathcal{N}_{\gamma, \alpha}$ is irreducible (by Corollary 4.5) and since

$$\dim(\mathcal{N}_{\theta, \beta}) = \ell(\theta) - \ell(\beta) = \ell(\gamma) - \ell(\alpha) = \dim(\mathcal{N}_{\gamma, \alpha}),$$

where the middle equality is by the interval embedding hypothesis.

To prove (7), we need to show $\Psi(m)$ satisfies (R.1), (R.2) and (R.3) whenever $m \in \mathcal{N}_{\theta, \beta}$. By (strong) induction on $\#J \geq 0$, we reduce to the case where θ is obtained from γ (or, equivalently, β is obtained from α) by adding a single $+$, a single $-$, or a single matching. (The base case $\#J = 0$ is trivial.)

Case 1: (θ is obtained from γ by adding a single $+$.) Let us suppose that γ and α are (p, q) -clans of length n , so that θ and β are $(p+1, q)$ -clans of length $n+1$. Let us further suppose that the $+$ is added between the $(\ell-1)$ -th and ℓ -th characters of α and γ . Thus

$$J = \{\ell\}, \quad \text{and} \quad \beta_\ell = \theta_\ell = +.$$

This $+$ in β corresponds to the ℓ -th row R and $w(\ell)$ -th column C of a matrix $m \in \mathcal{N}_{\theta, \beta}$; C is among the first p columns of m . (It is R and C which are deleted by the map Ψ .) The column C has a single 1, which is the only nonzero entry in C and in R .

(R.1): For $i = 1, \dots, n$, let R_i^+ denote the rank of the southwest $(n-i) \times p$ submatrix of $\Psi(m)$. For $j = 1, \dots, n+1$, let \mathcal{R}_j^+ denote the rank of the southwest $(n+1-j) \times (p+1)$ submatrix of m . We want to show that

$$R_i^+ \leq p - \gamma(i; +) \quad \text{for } i = 1, \dots, n,$$

knowing that

$$\mathcal{R}_j^+ \leq p + 1 - \theta(j; +) \quad \text{for } j = 1, \dots, n + 1.$$

There are two cases: Either $i < \ell$, or $i \geq \ell$.

If $i < \ell$, then we know that

$$\mathcal{R}_i^+ \leq p + 1 - \theta(i; +).$$

Furthermore, we know that $\mathcal{R}_i^+ = R_i^+ + 1$ since the row R is included among the last $n + 1 - i$ rows of m and hence contributes 1 to the rank of the southwest $(n + 1 - i) \times (p + 1)$ submatrix of m relative to the rank of the southwest $(n - i) \times p$ submatrix of $\Psi(m)$. Finally, we know that

$$\theta(i; +) = \gamma(i; +)$$

since θ and γ are the same up to position i . Putting these facts together gives us that

$$R_i^+ = \mathcal{R}_i^+ - 1 \leq p + 1 - \theta(i; +) - 1 = p - \gamma(i; +),$$

the desired conclusion.

On the other hand, if $i \geq \ell$, then we have that

$$\mathcal{R}_{i+1}^+ \leq p + 1 - \theta(i + 1; +).$$

We also have that

$$\mathcal{R}_{i+1}^+ = R_i^+$$

since the row R is now not among the last $n + 1 - (i + 1)$ rows of m and hence does not contribute 1 to the rank of the southwest $(n + 1 - (i + 1)) \times (p + 1)$ submatrix of m relative to the rank of the southwest $(n - i) \times p$ submatrix of $\Psi(m)$. Finally, we also clearly have that

$$\theta(i + 1; +) = \gamma(i; +) + 1.$$

Putting these facts together, we see that

$$R_i^+ = \mathcal{R}_{i+1}^+ \leq p + 1 - \theta(i + 1; +) = p + 1 - (\gamma(i; +) + 1) = p - \gamma(i; +),$$

as desired.

(R.2): Now, we let R_i^- denote the rank of the southeast $(n - i) \times q$ submatrix of $\Psi(m)$ for $i = 1, \dots, n$, and we let \mathcal{R}_j^- denote the rank of the southeast $(n + 1 - j) \times q$ submatrix of m . We want to show that

$$R_i^- \leq q - \gamma(i; -) \quad \text{for } i = 1, \dots, n,$$

knowing that

$$\mathcal{R}_j^- \leq q - \theta(j; -) \quad \text{for } j = 1, \dots, n + 1.$$

Again, we consider the cases $i < \ell$ and $i \geq \ell$.

In this case, the map Ψ simply deletes the row R of 0's from the submatrix of m formed by the last q columns, which has no effect on the rank whether this row is included among the last $n + 1 - i$ rows or not. Thus for $i < \ell$, we know that

$$\mathcal{R}_i^- \leq q - \theta(i; -), \quad \mathcal{R}_i^- = R_i^-, \quad \text{and} \quad \gamma(i; -) = \theta(i; -),$$

so

$$R_i^- \leq q - \gamma(i; -),$$

as desired.

When $i \geq \ell$, we have

$$\mathcal{R}_{i+1}^- \leq q - \theta(i + 1; -), \quad \mathcal{R}_{i+1}^- = R_i^-, \quad \text{and} \quad \gamma(i; -) = \theta(i + 1; -).$$

Thus

$$R_i^- \leq q - \gamma(i; -),$$

again as desired.

(R.3): We now must consider the ranks of the (i, j) -auxiliary matrices for all possible $i < j$. Given $i < j$, we denote by $\text{aux}(\Psi(m), i, j)$ the $n \times (i + j)$ auxiliary matrix formed from $\Psi(m)$, and we denote by $R_{i,j}$ the rank of this matrix. Similarly, we denote by $\text{aux}(m, i, j)$ the $(n + 1) \times (i + j)$ auxiliary matrix formed from m and by $\mathcal{R}_{i,j}$ the rank of this matrix. We want to see that

$$R_{i,j} \leq j + \gamma(i; j) \quad \text{for all } 1 \leq i < j \leq n,$$

given that

$$\mathcal{R}_{i,j} \leq j + \theta(i; j) \quad \text{for all } 1 \leq i < j \leq n + 1.$$

Since m is obtained from $\Psi(m)$ by adding the row ℓ and column $w(\ell)$, containing a 1 in position $(\ell, w(\ell))$ and zeros elsewhere, m^{-1} is obtained from $\Psi(m)^{-1}$ by adding the row $w(\ell)$ and the column ℓ , containing a 1 in position $(w(\ell), \ell)$ and zeros elsewhere. Since the auxiliary matrices are built from m^{-1} and $\Psi(m)^{-1}$, our analysis relies primarily on this observation.

Now, there are three cases to consider:

$$i < j < \ell, \quad i < \ell \leq j, \quad \text{and} \quad \ell \leq i < j.$$

When $i < j < \ell$, we compare $\text{aux}(\Psi(m), i, j)$ to $\text{aux}(m, i, j)$. Note that since the lone 1 added to $\Psi(m)^{-1}$ to form m^{-1} is located in column $\ell > j$, it does not appear in the latter auxiliary matrix; therefore, $\text{aux}(m, i, j)$ is obtained from $\text{aux}(\Psi(m), i, j)$ by simply adding a single row of zeros. Hence

$$R_{i,j} = \mathcal{R}_{i,j},$$

and since

$$\mathcal{R}_{i,j} \leq j + \theta(i; j), \quad \text{and} \quad \gamma(i; j) = \theta(i; j),$$

it follows that

$$R_{i,j} \leq j + \gamma(i; j),$$

as desired.

Now consider the case where $i < \ell \leq j$. In this case, we compare $\text{aux}(\Psi(m), i, j)$ to $\text{aux}(m, i, j + 1)$. Here, the latter matrix is obtained from the former by adding a single row and column, with a lone 1 at the intersection of this row and column, and 0's elsewhere. (Note that the 1 does in fact appear since it is in row $w(\ell) \leq p$, by the definition of w .) Thus

$$\mathcal{R}_{i,j+1} = R_{i,j} + 1, \quad \text{and clearly} \quad \theta(i; j + 1) = \gamma(i; j).$$

So since

$$\mathcal{R}_{i,j+1} \leq j + 1 + \theta(i; j + 1),$$

we have

$$R_{i,j} = \mathcal{R}_{i,j+1} - 1 \leq j + 1 + \theta(i; j + 1) - 1 = j + \gamma(i; j),$$

as desired.

Finally, consider the case $\ell \leq i < j$. Here we compare $\text{aux}(\Psi(m), i, j)$ to $\text{aux}(m, i + 1, j + 1)$. The latter matrix is obtained from the former this time by adding one row and two columns, with the row containing two 1's, one at its intersection with each of the two columns. The row and the two columns have zeros in all other entries. Note that this adds 1 (*not* 2) to the rank, so

$$\mathcal{R}_{i+1,j+1} = R_{i+1,j+1} + 1.$$

Note further that

$$\gamma(i; j) = \theta(i + 1; j + 1),$$

so since

$$\mathcal{R}_{i+1,j+1} \leq j + 1 + \theta(i + 1; j + 1),$$

we have

$$R_{i+1,j+1} = \mathcal{R}_{i+1,j+1} - 1 \leq j + 1 + \theta(i + 1; j + 1) - 1 = j + \gamma(i; j),$$

as required.

Case 2: (θ is obtained from γ by adding a single $-$.) Here the arguments are similar to those of Case 1; we simply interchange signs and the roles of the last q columns and the first p columns in the previous argument. We omit the details.

Case 3: (θ is obtained from γ by adding a single matching.) Suppose that the $(p+1, q+1)$ -clan θ is obtained from γ by adding to the (p, q) -clan γ a single matching, the left endpoint of which is added between positions $\ell - 1$ and ℓ , and the right endpoint of which is added between positions $\ell' - 1$ and ℓ' . Thus $J = \{\ell, \ell'\}$.

(R.1): As above, let R_i^+ denote the rank of the southwest $(n - i) \times p$ submatrix of $\Psi(m)$, and let \mathcal{R}_j^+ denote the rank of the southwest $(n + 2 - j) \times (p + 1)$ submatrix of m . Now the submatrix of m formed by its first $p + 1$ columns differs from the submatrix of $\Psi(m)$ formed by its first p columns in that the former has two extra rows and one extra column. The northernmost extra row (row ℓ) contains a 1 in

column $w(\ell)$ and 0's elsewhere, and the southernmost extra row (row ℓ') contains a -1 in column $w(\ell)$ and 0's elsewhere. Besides the aforementioned 1 and -1 , column $w(\ell)$ has all of its other entries zero.

We want to see that

$$R_i^+ \leq p - \gamma(i; +) \quad \text{for all } i$$

given that

$$\mathcal{R}_j^+ \leq p + 1 - \theta(j; +) \quad \text{for all } j.$$

We consider three cases:

$$i < \ell, \quad \ell \leq i < \ell', \quad \text{and} \quad \ell' \leq i.$$

If $i < \ell'$, then we know that

$$\mathcal{R}_i^+ \leq p + 1 - \theta(i; +) \quad \text{and that} \quad \theta(i; +) = \gamma(i; +).$$

Also,

$$\mathcal{R}_i^+ = R_i^+ + 1$$

since the southwest $(n + 2 - i) \times (p + 1)$ submatrix of m contains both the aforementioned extra rows relative to the southwest $(n - i) \times p$ submatrix of $\Psi(m)$, which causes the rank of the former to be 1 (*not* 2) higher than the rank of the latter (since the 1 and -1 occur in the same column). Thus

$$R_i^+ = \mathcal{R}_i^+ - 1 \leq p + 1 - \theta(i; +) - 1 = p - \gamma(i; +),$$

as required.

Now, if $\ell \leq i < \ell'$, then we know that

$$\mathcal{R}_{i+1}^+ \leq p + 1 - \theta(i + 1; +) \quad \text{and that} \quad \theta(i + 1; +) = \gamma(i; +).$$

Note here that

$$\mathcal{R}_{i+1}^+ = R_i^+ + 1,$$

because the southwest $(n + 2 - (i + 1)) \times (p + 1)$ submatrix of m still contains one of the two extra rows (namely row ℓ') relative to the southwest $(n - i) \times p$ submatrix of $\Psi(m)$; this causes the rank of the former to be 1 larger than the rank of the latter. Then

$$R_i^+ = \mathcal{R}_{i+1}^+ - 1 \leq p + 1 - \theta(i + 1; +) - 1 = p - \gamma(i; +).$$

Finally, if $\ell' \leq i$, then we know that

$$\mathcal{R}_{i+2} \leq p + 1 - \theta(i + 2; +) \quad \text{and that} \quad \theta(i + 2; +) = \gamma(i; +) + 1.$$

Here,

$$\mathcal{R}_{i+2}^+ = R_i^+$$

because the southwest $(n + 2 - (i + 2)) \times (p + 1)$ submatrix of m now does not contain either of the additional rows, so it differs from the southwest $(n - i) \times p$ submatrix of $\Psi(m)$ only in that it has an extra column of zeros. Thus we have

$$R_i^+ = \mathcal{R}_{i+2}^+ \leq p + 1 - \theta(i + 2; +) = p + 1 - (\gamma(i; +) + 1) = p - \gamma(i; +),$$

as required.

(R.2): The argument is virtually identical to that for (R.1).

(R.3): We again must consider the ranks of the (i, j) -auxiliary matrices for all possible $i < j$. As above, given $i < j$, we denote by $\text{aux}(\Psi(m), i, j)$ the $n \times (i + j)$ auxiliary matrix formed from $\Psi(m)$, and by $R_{i,j}$ the rank of this matrix. We denote by $\text{aux}(m, i, j)$ the $(n + 2) \times (i + j)$ auxiliary matrix formed from m , and by $\mathcal{R}_{i,j}$ the rank of this matrix. We want to see that

$$R_{i,j} \leq j + \gamma(i; j) \quad \text{for all } 1 \leq i < j \leq n$$

given that

$$\mathcal{R}_{i,j} \leq j + \theta(i; j) \quad \text{for all } 1 \leq i < j \leq n + 2.$$

Recall that m is obtained from $\Psi(m)$ by adding rows ℓ and ℓ' and columns $w(\ell)$ and $w(\ell')$, with 1 in positions $(\ell, w(\ell))$, $(\ell, w(\ell'))$, and $(\ell', w(\ell'))$, a -1 in position $(\ell', w(\ell))$, and 0's elsewhere. Thus m^{-1} is obtained from $\Psi(m)^{-1}$ by adding rows $w(\ell)$ and $w(\ell')$ and columns ℓ and ℓ' , with $\frac{1}{2}$ in positions $(w(\ell), \ell)$, $(w(\ell'), \ell)$, and $(w(\ell'), \ell')$, a $-\frac{1}{2}$ in position $(w(\ell), \ell')$, and 0's elsewhere. The analysis below relies primarily on this observation.

There are several cases to consider, depending upon the values of $i < j$ relative to ℓ and ℓ' . Given $i < j$, we define $i_\theta < j_\theta$ respectively to be the positions in θ of the i -th and j -th characters of γ relative to the embedding. So for a given i , there are three possibilities:

- (1) $i_\theta < \ell$, in which case $i_\theta = i$;
- (2) $\ell < i_\theta < \ell'$, in which case $i_\theta = i + 1$; or
- (3) $\ell' < i_\theta$, in which case $i_\theta = i + 2$.

Now, we consider the following cases:

$(i_\theta < j_\theta < \ell)$: In this case,

$$i_\theta = i, \quad j_\theta = j, \quad \text{and} \quad \gamma(i; j) = \theta(i_\theta; j_\theta).$$

Furthermore, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ only by the addition of two rows of zeros, so

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i,j}.$$

Thus

$$R_{i,j} = \mathcal{R}_{i_\theta, j_\theta} \leq j_\theta + \theta(i_\theta, j_\theta) = j + \gamma(i; j),$$

as required.

$(i_\theta < \ell < j_\theta < \ell')$: Then

$$i_\theta = i, \quad j_\theta = j + 1, \quad \text{and} \quad \gamma(i; j) = \theta(i_\theta, j_\theta).$$

Here, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ by the addition of two rows and one column. The column contains two $\frac{1}{2}$'s, each of which is contained in one of the two added rows. All other entries of this column and these two rows are zero. Thus

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i, j} + 1.$$

Then

$$R_{i, j} = \mathcal{R}_{i_\theta, j_\theta} - 1 \leq j_\theta + \theta(i_\theta; j_\theta) - 1 = j + 1 + \gamma(i; j) - 1 = j + \gamma(i; j).$$

$(i_\theta < \ell < \ell' < j_\theta)$: Then

$$i_\theta = i, \quad j_\theta = j + 2, \quad \text{and} \quad \gamma(i; j) = \theta(i_\theta, j_\theta).$$

Here, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ by the addition of two rows and two columns. The first column contains two $\frac{1}{2}$'s, each of which is contained in one of the two added rows. The second column contains a $-\frac{1}{2}$ and a $\frac{1}{2}$, again with each of these entries occurring in the two added rows. All other entries of the added rows and columns are zero. Thus

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i, j} + 2.$$

Then

$$R_{i, j} = \mathcal{R}_{i_\theta, j_\theta} - 2 \leq j_\theta + \theta(i_\theta; j_\theta) - 2 = j + 2 + \gamma(i; j) - 2 = j + \gamma(i; j).$$

$(\ell < i_\theta < j_\theta < \ell')$: Then

$$i_\theta = i + 1, \quad j_\theta = j + 1, \quad \text{and} \quad \gamma(i; j) + 1 = \theta(i_\theta, j_\theta).$$

Here, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ by the addition of two rows and two columns. The first column contains a single $\frac{1}{2}$ in the northmost added row. The second column contains two $\frac{1}{2}$'s, with each of these entries occurring in the two added rows. All other entries of the added rows and columns are zero. Thus

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i, j} + 2.$$

Then

$$R_{i, j} = \mathcal{R}_{i_\theta, j_\theta} - 2 \leq j_\theta + \theta(i_\theta; j_\theta) - 2 = j + 1 + \gamma(i; j) + 1 - 2 = j + \gamma(i; j).$$

$(\ell < i_\theta < \ell' < j_\theta)$: Then

$$i_\theta = i + 1, \quad j_\theta = j + 2, \quad \text{and} \quad \gamma(i; j) = \theta(i_\theta, j_\theta).$$

Here, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ by the addition of two rows and three columns; one checks that again we have

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i, j} + 2.$$

Thus

$$R_{i,j} = \mathcal{R}_{i_\theta, j_\theta} - 2 \leq j_\theta + \theta(i_\theta; j_\theta) - 2 = j + 2 + \gamma(i; j) - 2 = j + \gamma(i; j).$$

($\ell < \ell' < i_\theta < j_\theta$): Then

$$i_\theta = i + 2, \quad j_\theta = j + 2, \quad \text{and} \quad \gamma(i; j) = \theta(i_\theta, j_\theta).$$

Here, $\text{aux}(m, i_\theta, j_\theta)$ differs from $\text{aux}(\Psi(m), i, j)$ by the addition of two rows and four columns; again, it is the case that

$$\mathcal{R}_{i_\theta, j_\theta} = R_{i,j} + 2.$$

Then

$$R_{i,j} = \mathcal{R}_{i_\theta, j_\theta} - 2 \leq j_\theta + \theta(i_\theta; j_\theta) - 2 = j + 2 + \gamma(i; j) - 2 = j + \gamma(i; j).$$

This completes the proof of Theorem 4.6, having assumed Proposition 6.1. \square

6B. Proof of Theorem 1.3: (I): Let $[\alpha, \gamma] \in \mathcal{C}$ be such that Y_γ is non- \mathcal{P} along \mathcal{O}_α . Now suppose

$$[\alpha, \gamma] \preceq_{\mathcal{C}} [\beta, \theta]. \quad (8)$$

Clearly, we may assume this is a covering relation, of which there are two kinds.

The first possibility is that $[\beta, \theta]$ interval pattern contains $[\alpha, \gamma]$. We know from Observation 1.4 that W_γ is non- \mathcal{P} along \mathcal{Q}_α , so in particular, W_γ is non- \mathcal{P} at x_α . Thus $\mathcal{N}_{\gamma, \alpha}$ is non- \mathcal{P} at the origin, since (non-) \mathcal{P} is stable under taking slices by Lemma 2.5. Then $\mathcal{N}_{\theta, \beta}$ is non- \mathcal{P} at the origin as well, by Theorem 4.6. (Note that the isomorphism Ψ of Theorem 4.6 carries the origin on $\mathcal{N}_{\theta, \beta}$ to the origin on $\mathcal{N}_{\gamma, \alpha}$.) Thus W_θ is non- \mathcal{P} at x_β , again by slice-stability of (non-) \mathcal{P} , and then in fact W_θ is non- \mathcal{P} along \mathcal{Q}_β , by homogeneity. Using Observation 1.4 once more, we conclude that Y_θ is non- \mathcal{P} along \mathcal{O}_β .

The other possibility is that $\theta = \gamma$ and $\beta \preceq \alpha$. Then $\mathcal{O}_\beta \subseteq \overline{\mathcal{O}_\alpha} \subseteq Y_\gamma$, by the definition of the closure order. Since Y_γ is non- \mathcal{P} along \mathcal{O}_α and non- \mathcal{P} is a closed property, Y_γ is non- \mathcal{P} on $\overline{\mathcal{O}_\alpha}$, and hence on \mathcal{O}_β , as desired.

(II): This is the contrapositive of (I). \square

6C. Proof of Proposition 6.1. Recall that $w := w_\beta$ is the permutation associated to the clan β , as defined at the beginning of Section 3A. Denote by $\mathcal{I} = \mathcal{I}(\beta)$ the underlying involution of β , as defined prior to Claim 3.7. This means \mathcal{I} is the permutation that fixes j if β_j is a sign and interchanges i and j if $(i < j)$ is a matching of β .

We split the proof into three main cases (A, B and C) depending on the value of θ_j .

6C1. Case A: ($\theta_j = +$). We prove Case A via two claims.

Claim 6.2. *The entries of column $w(j)$ (except the 1 in row j required by (O.1)) vanish.*

Proof. By Lemma 5.7(I), $\theta(j; +) = \beta(j; +)$. Define

$$r := p - \theta(j; +) = p - \beta(j; +). \quad (9)$$

Since $\beta(n; +) = p$, it follows from the definitions that

$$r = \#\{k \mid k > j, \text{ and } \beta_k = + \text{ or } \beta_k \text{ is the right end of a matching}\}. \quad (10)$$

Let $k_1 < \dots < k_r$ be the r indices of the set given in (10). Also define

$$c_i = \begin{cases} w(k_i) & \text{if } \beta_{k_i} = +, \\ w(\mathcal{I}(k_i)) & \text{if } \beta_k \text{ is the right end of a matching,} \end{cases}$$

for $i = 1, \dots, r$.

Subclaim 6.3. (I) $c_i \leq p$ for $1 \leq i \leq r$.

(II) Column c_i of M has a fixed ± 1 in row k_i . Besides any fixed ± 1 in this row, any of the leftmost p columns of M strictly right of column c_i has zero in row k_i .

Proof. First suppose $\beta_{k_i} = +$. Then by definition of w ,

$$c_i = w(k_i) \leq p,$$

proving (I) in this case. Case (II) holds by (O.1) and (Z.1) combined.

Otherwise, β_{k_i} is the right end of a matching. For (I), here $\mathcal{I}(k_i)$ is the left end of that matching, and

$$c_i = w(\mathcal{I}(k_i)) \leq p,$$

again by the definition of w . Now, by (O.2), columns $w(\mathcal{I}(k_i))$ and $w(k_i)$ of M are assigned -1 and 1 respectively in row k_i . Then (II) holds by (Z.4). This completes the proof of Subclaim 6.3. \square

Now, let \vec{v}_i denote the vector consisting of the last $n - j$ entries in column c_i . By Subclaim 6.3(II), it follows that $\{\vec{v}_1, \dots, \vec{v}_r\}$ is a linearly independent set. Since $M \in \mathcal{N}_{\theta, \beta}$, by (R.1) (with $i = j$), the southwest $(n - j) \times p$ submatrix M° of M has rank at most r (see (9)). Thus

$$\vec{v}_1, \dots, \vec{v}_r \text{ is a basis of } \text{colspace}(M^\circ). \quad (11)$$

Since we assume $\theta_j = +$, by (Z.2), all entries of column $w(j)$ strictly north of row j are zero. Thus it remains to show the vector \vec{v} consisting of the last $n - j$ entries in column $w(j) \leq p$ is the zero vector. By (11) we have

$$\vec{v} \in \text{Span}(\vec{v}_1, \dots, \vec{v}_r). \quad (12)$$

Now, for $i = 1, \dots, r$, let

$$k'_i = \begin{cases} \mathcal{I}(k_i) & \text{if } \mathcal{I}(k_i) > j, \\ k_i & \text{otherwise.} \end{cases}$$

Notice that k'_i is either the row of a pivot, or it is a row containing a -1 which is located southwest of entry $(j, w(j))$. Thus either by (Z.1) in the former case, or by (Z.4) in the latter, \vec{v} has a 0 in row k'_i for all i . In view of (12) and Subclaim 6.3(II), we see that $\vec{v} = \vec{0}$, as desired. This completes the proof of Claim 6.2. \square

Claim 6.4. *The entries of row j (except the 1 in column $w(j)$ required by (O.1)) vanish.*

Proof. By (Z.1), the claim holds for the leftmost p columns of M . Thus, it remains to check the conclusion for the rightmost q columns $p + 1, \dots, n$.

Let M° be the southeast $n - (j - 1) \times q$ submatrix of M . By (R.2) for $i = j - 1$, the rank of M° is at most

$$q - \theta(j - 1; -) = q - \theta(j; -) = q - \beta(j; -) =: r,$$

where the second equality is by Lemma 5.7. As in the proof of Claim 6.2, there are r positions $k_1, \dots, k_r > j$ with β_{k_i} being either a $-$ or the right end of a matching. Each index corresponds to a 1 in row $k_i > j$ in one of the rightmost q columns of M . More precisely, if β_{k_i} is a $-$, this 1 is the pivot of row k_i , and if β_{k_i} is the right end of a matching, the 1 is the second 1 in its column. By (Z.5) and (Z.8) respectively, each of these 1's have all 0's to their right (and in the same row). None of these 1's appear in the same column, by (O.1) and (O.3). So, these r column vectors $\vec{v}_1, \dots, \vec{v}_r$ of M° are linearly independent, and therefore

$$\vec{v}_1, \dots, \vec{v}_r \text{ is a basis of } \text{colspace}(M^\circ). \quad (13)$$

In row j of M , the entry in each \vec{v}_i is zero, by (Z.6) or (Z.7), since each such position is above either a pivot 1 or is between two 1's. So if any entry in row j among the last q were nonzero, its column in M° would be linearly independent of $\{\vec{v}_1, \dots, \vec{v}_r\}$, contradicting (13). The claim therefore holds. \square

6C2. *Case B:* ($\theta_j = -$). The argument is nearly identical to that of Case A; we omit the details.

6C3. *Case C:* ($j, j' \in J$ (with $j < j'$) is a matched pair). We start with an observation we will use repeatedly:

Claim 6.5. *If $(k < k')$ is a matching of β , then*

$$1 \leq a < w(k) \implies w^{-1}(a) < k,$$

and

$$p + 1 \leq a < w(k') \implies \mathcal{I}(w^{-1}(a)) < k.$$

Proof. Since k is the left end of a matching, $w(k) \leq p$. If $1 \leq a < w(k) \leq p$, then by definition, a is the label assigned by w to either a $+$ or the left end of a matching appearing to the left of k , so $w^{-1}(a) < k$.

Similarly, since k' is the right end of a matching, we have $p + 1 \leq w(k') \leq n$. If $p + 1 \leq a < w(k')$, then a is assigned by w to either a $-$ or the right end f' of a matching ($f < f'$). By definition of w , in the former case, the “ $-$ ” must appear left of k , so that $w^{-1}(a) < k$. Since $\mathcal{I}(w^{-1}(a)) = w^{-1}(a)$ in this case, we have $\mathcal{I}(w^{-1}(a)) < k$, as claimed. In the latter case, where f' occurs at position $w^{-1}(a)$ and f at position $\mathcal{I}(w^{-1}(a))$, again by the definition of w , we must have that f is left of k , so $\mathcal{I}(w^{-1}(a)) < k$. \square

Denote by \vec{v}_i the i -th column vector of M .

Since M is invertible, $\{\vec{v}_1, \dots, \vec{v}_n\}$ is a basis of \mathbb{C}^n . Thus, for any $1 \leq k \leq n$, we have a linear dependence relation

$$\vec{e}_k + \sum_{a=p+1}^n \lambda_{k,a} \vec{v}_a = \sum_{a=1}^p \lambda_{k,a} \vec{v}_a \quad (14)$$

for some scalars $\lambda_{k,1}, \dots, \lambda_{k,n} \in \mathbb{C}$.

Claim 6.6. *Given any k such that $\beta_k = +, -, \text{ or the left end of a matching, we have$*

$$\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = 0 \quad \text{for any } b < k.$$

If β_k is the right end of a matching, then

$$\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = 0 \quad \text{for any } b < \mathcal{I}(k).$$

Proof. Fix k . We prove both assertions by a common induction on $b < \min\{k, \mathcal{I}(k)\}$.

In the base case, when $b = 1$, β_1 cannot be the right end of a matching. So we check the remaining three possibilities in turn.

If $\beta_1 = +$, then the first coordinate of \vec{v}_a is 1 for $a = 1$ and is 0 otherwise. This follows from the definition of w together with (O.1), (Z.1), and (Z.6). Since $k > b = 1$, the first row of (14) thus says that $0 = \lambda_{k,1}$. However,

$$1 = w(1) = w(\mathcal{I}(1))$$

in this case, so we are done. The argument when $\beta_1 = -$ is similar.

Now, suppose that β_1 is the left end of a matching. Then the right end of this matching is at position $\mathcal{I}(1)$. From w 's definition, along with (O.2), (Z.1), and (Z.5), the first row of (14) asserts

$$0 + \lambda_{k,p+1} = \lambda_{k,1}.$$

Meanwhile, by (O.3), (Z.4), and (Z.8), row $\mathcal{I}(1)$ reads

$$0 + \lambda_{k,p+1} = -\lambda_{k,1}.$$

Putting these together, we have

$$\lambda_{k,w(1)} = \lambda_{k,1} = \lambda_{k,p+1} = \lambda_{k,w(\mathcal{I}(1))} = 0,$$

as desired.

For the inductive step, now suppose that $b > 1$, and that the claims hold for indices less than b .

Case 1: ($\beta_b = +$.) By (Z.1) in row b , all the entries among the first p columns are 0 except for a 1 in column $w(b)$. Therefore, the b -th row of (14) is

$$0 + \sum_{a=p+1}^n \lambda_{k,a} z_{b,a} = \lambda_{k,w(b)}. \quad (15)$$

If

$$\mathcal{I}(w^{-1}(a)) > b,$$

then $z_{b,a} = 0$ by (Z.6), since this entry is above the pivot in column a . If on the other hand

$$\mathcal{I}(w^{-1}(a)) < b,$$

then $\lambda_{k,a} = 0$ by the inductive hypothesis. Hence $\lambda_{k,w(b)} = 0$. Since $w(\mathcal{I}(b)) = w(b)$ in this case, we have $\lambda_{k,w(\mathcal{I}(b))} = 0$ as well.

Case 2: ($\beta_b = -$.) This is similar to Case 1.

Case 3: (β_b is left end of a matching.) Let $b' = \mathcal{I}(b)$ be the right end of the matching. Let

$$c = w(b) \quad \text{and} \quad c' = w(b').$$

In row b , (14) states that

$$\lambda_{k,c'} = \lambda_{k,c}, \tag{16}$$

since all other entries in row b are 0, by (Z.1) and (Z.5).

Now, note that if β_k is a $+$, a $-$, or the left end of a matching, then we clearly have that $k \neq b'$. If β_k is the right end of a matching, then since $b < \mathcal{I}(k)$, we again have that $k \neq b'$. Thus in row b' , (14) states that

$$0 + \sum_{a=p+1}^{c'-1} \lambda_{k,a} z_{b',a} + \lambda_{k,c'} = \sum_{a=1}^{c-1} \lambda_{k,a} z_{b',a} - \lambda_{k,c}, \tag{17}$$

by (Z.4) and (Z.8).

Now,

$$\mathcal{I}(w^{-1}(a)) < w^{-1}(c) = b$$

for a in the first sum, whereas

$$w^{-1}(a) < w^{-1}(c) = b$$

for a in the second sum. Hence by induction, $\lambda_{k,a} = 0$ for $1 \leq a \leq c-1$ and $p+1 \leq a \leq c'-1$. So (17) reduces to

$$\lambda_{k,c'} = -\lambda_{k,c}.$$

Combining this with (16) gives

$$\lambda_{k,c} = \lambda_{k,c'} = 0,$$

as desired.

Case 4: (β_b is the right end of a matching.) The left end of the matching is at position $b' = \mathcal{I}(b)$. By Case 3,

$$\lambda_{k,w(b')} = \lambda_{k,w(\mathcal{I}(b'))} = 0,$$

which is the same as

$$\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = 0,$$

as required. □

Corollary 6.7. *If $(k < k')$ is a matching of β , then*

$$\lambda_{k,a} = \lambda_{k',a} = 0 \quad \text{for } a = 1, \dots, w(k) - 1 \text{ and for } a = p + 1, \dots, w(k') - 1.$$

Proof. By Claim 6.5, for $1 \leq a < w(k)$, we have $w^{-1}(a) < k$, whereas for $p+1 \leq a < w(k')$, we have $\mathcal{I}(w^{-1}(a)) < k$. In either event, $\lambda_{k,a} = \lambda_{k',a} = 0$ by Claim 6.6. \square

Claim 6.8. *If $\beta_k = +$, then $\lambda_{k,w(k)} = 1$. If $\beta_k = -$, then $\lambda_{k,w(k)} = -1$.*

Proof. First suppose $\beta_k = +$. Consider row k of (14). By (Z.1) and (O.1), this equation is

$$1 + \sum_{a=p+1}^n \lambda_{k,a} z_{k,a} = \lambda_{k,w(k)}.$$

If $w^{-1}(a) < k$, then $\lambda_{k,a} = 0$ by Claim 6.6. On the other hand, if $w^{-1}(a) > k$, then $z_{k,a} = 0$ by (Z.6) or (Z.7). Hence the equation reduces to $1 = \lambda_{k,w(k)}$, as desired.

Now, if $\beta_k = -$, then row k of (14) is

$$1 + \lambda_{k,w(k)} = \sum_{a=1}^p \lambda_{k,a} z_{k,a}.$$

Again, if $w^{-1}(a) < k$, then $\lambda_{k,a} = 0$ by Claim 6.6. If $w^{-1}(a) > k$, then $z_{k,a} = 0$ by (Z.2). Hence the equation reduces to $1 + \lambda_{k,w(k)} = 0$, whence $\lambda_{k,w(k)} = -1$ as claimed. \square

Claim 6.9. *Suppose $(k < k')$ is a matching of β . Then*

$$\lambda_{k,w(k)} = \frac{1}{2} \quad \text{and} \quad \lambda_{k,w(k')} = -\frac{1}{2},$$

and

$$\lambda_{k',w(k)} = \lambda_{k',w(k')} = -\frac{1}{2}.$$

Proof. First consider row k of (14). By (O.2), (Z.1), and (Z.5), it reads

$$1 + \lambda_{k,w(k')} = \lambda_{k,w(k)}. \tag{18}$$

Now consider row k' of (14). By (O.3), (Z.4), and (Z.8), it reads

$$\sum_{a=p+1}^{w(k')-1} \lambda_{k,a} z_{k',a} + \lambda_{k,w(k')} = \sum_{a=1}^{w(k)-1} \lambda_{k,a} z_{k',a} - \lambda_{k,w(k)}. \tag{19}$$

Applying Corollary 6.7 to (19) reduces it to

$$\lambda_{k,w(k')} = -\lambda_{k,w(k)}. \tag{20}$$

Solving (18) and (20) simultaneously gives $\lambda_{k,w(k)} = \frac{1}{2}$ and $\lambda_{k,w(k')} = -\frac{1}{2}$, as claimed.

Similarly, now consider (14), but with k replaced by k' ; looking at row k gives

$$\lambda_{k',w(k')} = \lambda_{k',w(k)}. \tag{21}$$

If we examine row k' , we obtain

$$1 + \lambda_{k',w(k')} = -\lambda_{k',w(k)}. \tag{22}$$

Solving (22) and (21) simultaneously gives the remainder of the claim. \square

Claim 6.10. *If $(k < k')$ is a matching of β , then*

$$\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = \lambda_{k',w(b)} = \lambda_{k',w(\mathcal{I}(b))} = 0$$

for $b = k + 1, \dots, k' - 1$.

Proof. We only give the proof of the claim that

$$\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = 0.$$

The proof that

$$\lambda_{k',w(b)} = \lambda_{k',w(\mathcal{I}(b))} = 0$$

is identical.

We induct on b . For b in the appropriate range, the inductive hypothesis is that the claim holds for all a satisfying $k < a < b$.

Case 1: ($\beta_b = +$.) In this case, row b of (14) reads

$$0 + \sum_{a=p+1}^n \lambda_{k,a} z_{b,a} = \lambda_{k,w(b)}. \quad (23)$$

Now, if $\mathcal{I}(w^{-1}(a)) > b$, then $z_{b,a} = 0$ by (Z.6), since this entry is above the pivot in column a . Otherwise, either

$$\mathcal{I}(w^{-1}(a)) < k, \quad \mathcal{I}(w^{-1}(a)) = k, \quad \text{or} \quad k < \mathcal{I}(w^{-1}(a)) < b.$$

In the first case, we have that $\lambda_{k,a} = 0$ by Claim 6.6. In the last case, we have that $\lambda_{k,a} = 0$ by the inductive hypothesis. In the middle case, we have that $z_{b,a} = 0$ by (Z.7), since this entry is between the two 1's in positions (k, a) and (k', a) . Thus the left-hand side of (23) is actually zero, implying that $\lambda_{k,w(b)} = \lambda_{k,w(\mathcal{I}(b))} = 0$, as claimed.

Case 2: ($\beta_b = -$.) This is very similar to Case 1; we omit the details.

Case 3: (β_b is the left end of a matching.) Let the right end of this matching be $\beta_{b'}$. Let $c = w(b)$, and let $c' = w(b')$.

By (Z.1) and (Z.5), in row b , (14) states that

$$\lambda_{k,c} = \lambda_{k,c'}. \quad (24)$$

By (Z.4), (Z.8) and (O.3), in row b' , we have

$$0 + \sum_{a=p+1}^{c'-1} \lambda_{k,a} z_{b',a} + \lambda_{k,c'} = \sum_{a=1}^{c-1} \lambda_{k,a} z_{b',a} - \lambda_{k,c} \quad (25)$$

Subcase 3.1: ($b' < k'$.) In fact each term of the sum on the left-hand side of (25) vanishes for one of three reasons:

(L.3.1.1) $\mathcal{I}(w^{-1}(a)) < k$: $\lambda_{k,a} = 0$ by Claim 6.6.

(L.3.1.2) $\mathcal{I}(w^{-1}(a)) = k$: $z_{b',a} = 0$ by (Z.7), since in column $a = w(k')$, the entry in row b' is located between two 1's in rows k and k' , since we have assumed that $b' < k'$.

(L.3.1.3) $\mathcal{I}(w^{-1}(a)) > k$: $\lambda_{k,a} = 0$ by the inductive hypothesis, since $p + 1 \leq a < w(b')$ implies that $\mathcal{I}(w^{-1}(a)) < b$ by Claim 6.5.

Similarly, each term of the sum on the right hand side of (25) vanishes for one of these three reasons:

(R.3.1.1) $w^{-1}(a) < k$: $\lambda_{k,a} = 0$ by Claim 6.6.

(R.3.1.2) $w^{-1}(a) = k$: $z_{b',a} = 0$ by (Z.3), since in column $a = w(k)$, the entry in row b' is located between a 1 in row k and a -1 in row k' , again because we have assumed that $b' < k'$.

(R.3.1.3) $w^{-1}(a) > k$: $\lambda_{k,a} = 0$ by the inductive hypothesis, since $1 \leq a < w(b)$ implies that $w^{-1}(a) < b$ by Claim 6.5.

Thus by (L.3.1.1)–(L.3.1.3) and (R.3.1.1)–(R.3.1.3) combined, (25) reduces to $\lambda_{k,c'} = -\lambda_{k,c}$. The simultaneous solution of this with (24) is $\lambda_{k,c} = \lambda_{k,c'} = 0$, as desired.

Subcase 3.2: ($b' > k'$.) The two matchings ($k < k'$) and ($b < b'$) form a 1212 pattern. We analyze each term of the sum on the left-hand side of (25):

(L.3.2.1) $\mathcal{I}(w^{-1}(a)) < k$: $\lambda_{k,a} = 0$ by Claim 6.6.

(L.3.2.2) $\mathcal{I}(w^{-1}(a)) = k$: We cannot conclude that $z_{b',a} = 0$. We do know by Claim 6.9 that $\lambda_{k,a} = \lambda_{k,w(k')} = -\frac{1}{2}$, so all we can say is that this particular term is equal to $-\frac{1}{2}z_{b',w(k')}$.

(L.3.2.3) $\mathcal{I}(w^{-1}(a)) > k$: $\lambda_{k,a} = 0$ by the inductive hypothesis, as in (L.3.1.3).

For the right hand side of (25), we see:

(R.3.2.1) $w^{-1}(a) < k$: $\lambda_{k,a} = 0$ by Claim 6.6.

(R.3.2.2) $w^{-1}(a) = k$: We do not know that $z_{b',a} = 0$, but we do at least know by Claim 6.9 that $\lambda_{k,a} = \lambda_{k,w(k)} = \frac{1}{2}$. Thus this term is equal to $\frac{1}{2}z_{b',w(k)}$.

(R.3.2.3) $w^{-1}(a) > k$: $\lambda_{k,a} = 0$ by the inductive hypothesis, as in (R.3.1.3).

In view of the 1212 pattern occurring in positions $k < b < k' < b'$, by (Z.9),

$$z_{b',w(k')} = -z_{b',w(k)}.$$

Thus the terms from (L.3.2.2) and (R.3.2.2) may be canceled in (25). Therefore (25) reduces to $\lambda_{k,c'} = -\lambda_{k,c}$. The simultaneous solution of this with (24) is

$$\lambda_{k,c} = \lambda_{k,c'} = 0,$$

as required.

Case 4: (β_b is the right end of a matching.) If the left end of this matching occurs prior to position k , we are done by Claim 6.6. Otherwise it occurs after position k , and we may apply Case 3. \square

Let

$$\pi_M : \mathbb{C}^n \rightarrow V_1$$

be the projection onto

$$V_1 := \langle \vec{v}_1, \dots, \vec{v}_p \rangle$$

with kernel

$$V_2 := \langle \vec{v}_{p+1}, \dots, \vec{v}_n \rangle.$$

Applying π_M to both sides of (14) gives

$$\pi_M(\vec{e}_k) = \sum_{a=1}^p \lambda_{k,a} \vec{v}_a. \quad (26)$$

Claim 6.11. *Suppose $(k < k')$ is a matching of β . Then*

- (I) $\pi_M(\vec{e}_k)$ is equal to $-1/2$ in row k' .
- (II) If ℓ is the left end of a matching with $\ell > k$, then $\pi_M(\vec{e}_\ell)$ has entry 0 in row k' .
- (III) $\pi_M(\vec{e}_k)$ is zero in row ℓ for any $\ell < k'$ except $\ell = k$.

Proof. (I) We have

$$\pi_M(\vec{e}_k) = \sum_{a=1}^p \lambda_{k,a} \vec{v}_a = \sum_{a=w(k)}^p \lambda_{k,a} \vec{v}_a = \frac{1}{2} \vec{v}_{w(k)} + \sum_{a=w(k)+1}^p \lambda_{k,a} \vec{v}_a. \quad (27)$$

The first equality is (26). The second equality is by Corollary 6.7. The third equality applies Claim 6.9. Now, by (O.3), $z_{k',w(k)} = -1$, and by (Z.4), $z_{k',a} = 0$ for $w(k) < a \leq p$. Therefore row k' of (27) is clearly $-\frac{1}{2}$, as claimed.

(II) By (Z.4) and (26), entry k' of $\pi_M(\vec{e}_\ell)$ is $\sum_{a=1}^{w(k)} \lambda_{\ell,a} z_{k',a}$. But if $a \leq w(k)$, we have $w^{-1}(a) \leq k < \ell$ by Claim 6.5. Thus by Claim 6.6, all $\lambda_{\ell,a}$ in this sum are zero.

(III) $\pi_M(\vec{e}_k)$ in row ℓ is $\sum_{a=1}^p \lambda_{k,a} z_{\ell,a}$. However, $\lambda_{k,a} = 0$ if $w^{-1}(a) < k'$ and $w^{-1}(a) \neq k$, either by Claim 6.6 (if $w^{-1}(a) < k$) or Claim 6.10 (if $k < w^{-1}(a) < k'$).

Now suppose $w^{-1}(a) = k$, so that $a = w(k)$. Then there is a 1 in position (k, a) and a -1 in position (k', a) . If $\ell < k$, then $z_{\ell,a} = 0$ by (Z.2), while if $k < \ell < k'$, we have $z_{\ell,a} = 0$ by (Z.3).

Finally, if $w^{-1}(a) > k'$, then since $1 \leq a \leq p$ (and hence β has either a $+$ or a left endpoint at position $w^{-1}(a)$), there is a pivot at position $(w^{-1}(a), a)$. Then since $\ell < k'$, we have $z_{\ell,a} = 0$ by (Z.2). So in fact every term of the sum is zero. \square

Set

$$r := \theta(j; j').$$

By Lemma 5.7,

$$r = \beta(j; j') = \#\{(k < k') \text{ a matching of } \beta \mid k < j < j' < k'\}.$$

Let the r matchings of the above set be

$$(k_1 < k'_1), (k_2 < k'_2), \dots, (k_r < k'_r), \quad \text{with } k_1 < k_2 < \dots < k_r < j.$$

Claim 6.12. (I) *The set*

$$\{\pi_M(\vec{e}_{k_1}), \pi_M(\vec{e}_{k_2}), \dots, \pi_M(\vec{e}_{k_r})\} \quad (\text{modulo } E_{j'})$$

forms a basis of $\pi_M(E_j) + E_{j'}/E_{j'}$.

(II) $\pi_M(\vec{e}_j) = \vec{0}$ as an element of $\pi_M(E_j) + E_{j'}/E_{j'}$. In other words, for any $t > j'$, $\pi_M(\vec{e}_j)$ is 0 in row t .

Proof. (I) Suppose in the quotient $\pi_M(E_j) + E_{j'}/E_{j'}$ that we have a dependence relation of the form

$$\sum_{i=1}^r \lambda_i (\pi_M(\vec{e}_{k_i}) + E_{j'}) = \sum_{i=1}^r \lambda_i \pi_M(\vec{e}_{k_i}) + E_{j'} = 0 \quad (\text{modulo } E_{j'}).$$

Then $\sum_{i=1}^r \lambda_i \pi_M(\vec{e}_{k_i}) \in E_{j'}$, meaning that for any index $\ell > j'$, the vector $\sum_{i=1}^r \lambda_i \pi_M(\vec{e}_{k_i})$ has entry 0 in position ℓ . In particular, this vector has entry 0 in positions k'_1, \dots, k'_r . Then applying parts (I) and (II) of Claim 6.11, a triangularity argument gives that all the scalars λ_i are equal to zero. Hence the vectors $\pi_M(\vec{e}_{k_1}), \dots, \pi_M(\vec{e}_{k_r})$ descend to a linearly independent set in the quotient $\pi_M(E_j) + E_{j'}/E_{j'}$.

Then by (C.3) and the discussion following the statement of Theorem 4.4, we know that

$$\dim(\pi_M(E_j) + E_{j'}) \leq j' + \theta(j; j') = j' + \beta(j; j') = j' + r.$$

Then $\dim(\pi_M(E_j) + E_{j'}/E_{j'}) \leq r$, so in fact the linearly independent set $\{\pi_M(\vec{e}_{k_1}), \dots, \pi_M(\vec{e}_{k_r})\}$ is a basis for $\pi_M(E_j) + E_{j'}/E_{j'}$, as claimed.

(II) From Part (I), we have that $\pi_M(\vec{e}_j)$, viewed as an element of $\pi_M(E_j) + E_{j'}/E_{j'}$, is a linear combination of $\{\pi_M(\vec{e}_{k_1}), \pi_M(\vec{e}_{k_2}), \dots, \pi_M(\vec{e}_{k_r})\}$, or equivalently,

$$\pi_M(\vec{e}_j) + E_{j'} = \sum_{i=1}^r \lambda_i \pi_M(\vec{e}_{k_i}) + E_{j'}$$

for some scalars $\lambda_1, \dots, \lambda_r$. We want to show that $\lambda_i = 0$ for each i . Since we know that $k_i < j < k'_i$ for $1 \leq i \leq r$, we have by Claim 6.11(II) that $\pi_M(\vec{e}_j)$ has entry 0 in row $k'_i > j'$ for $1 \leq i \leq r$. We also know by Claim 6.11(I) that $\pi_M(\vec{e}_{k_i})$ is nonzero in position k'_i for $1 \leq i \leq r$. Combining these two facts gives that each $\lambda_i = 0$, and hence we have $\pi_M(\vec{e}_j) = 0$ in $\pi_M(E_j) + E_{j'}/E_{j'}$, as desired. Since we thus have $\pi_M(\vec{e}_j) \in E_{j'}$ the remaining assertion also follows. \square

We also record the following related claim for later use:

Claim 6.13. (I) *The set*

$$\{\pi_M(\vec{e}_{k_1}), \pi_M(\vec{e}_{k_2}), \dots, \pi_M(\vec{e}_{k_r})\} \quad (\text{modulo } E_{j'-1})$$

forms a basis of $(\pi_M(E_{j-1}) + E_{j'-1})/E_{j'-1}$.

(II) $\pi_M(\vec{v}) = 0$ in row j' for any vector \vec{v} .

Proof. (I) Since $(j < j')$ is a matching, we have

$$\theta(j-1; j'-1) = \theta(j; j') = r,$$

since a matching $(k < k')$ satisfies

$$k \leq j < j' < k' \iff k \leq j-1 < j'-1 < k'.$$

Hence the claim follows by exactly the same argument as for Claim 6.12(I).

(II) By (I), it suffices to prove this for \vec{e}_{k_i} for $i = 1, \dots, r$. Since $j' < k'_i$ by definition, this follows from Claim 6.11(III). \square

Now, by (O.2),

$$z_{j,w(j)} = 1 \quad \text{and} \quad z_{j,w(j')} = 1.$$

By (O.3),

$$z_{j',w(j)} = -1 \quad \text{and} \quad z_{j',w(j')} = 1.$$

By (Z.2) and (Z.3),

$$z_{b,w(j)} = 0 \quad \text{if } b < j \text{ or } j < b < j'.$$

Finally, by (Z.6) and (Z.7),

$$z_{b,w(j')} = 0 \quad \text{if } b < j \text{ or } j < b < j'.$$

Hence the first assertion of our next claim is what remains to obtain the desired conclusion of Proposition 6.1 about columns $w(j)$ and $w(j')$. The second assertion is a technical strengthening for the induction argument we give.

Claim 6.14 (Case C: column zeroness). $z_{b,w(j)} = z_{b,w(j')} = 0$ and $\lambda_{j,w(b)} = \lambda_{j,w(\mathcal{I}(b))} = 0$ for all $b > j'$.

Proof. Our argument is by induction on $b > j'$. The inductive hypothesis is that the claims hold for all q satisfying $j' < q < b$. Our inductive step is to prove that this implies that the claims hold for b . Our argument depends upon the value of β_b .

($\beta_b = +$): The expression for $\pi_M(\vec{e}_j)$ in row b (see (26)) is

$$\sum_{a=1}^p \lambda_{j,a} z_{b,a}. \tag{28}$$

However, by (O.1) and (Z.1) we know

$$z_{b,w(b)} = 1 \quad \text{and} \quad z_{b,t} = 0 \quad \text{for } 1 \leq t \leq p \quad \text{and} \quad t \neq w(b).$$

Thus, in particular we obtain the claim's assertion that $z_{b,w(j)} = 0$. Moreover, (28) reduces to $\lambda_{j,w(b)}$. By Claim 6.12(II), all entries of $\pi_M(\vec{e}_j)$ beyond position j' are zero. Since we assume $b > j'$, we therefore conclude that $\lambda_{j,w(b)} = 0$. Since $\beta_b = +$, by definition we have

$$w(\mathcal{I}(b)) = w(b),$$

so

$$\lambda_{j,w(\mathcal{I}(b))} = \lambda_{j,w(b)} = 0,$$

proving the claim's second assertion in this case.

It remains to show that $z_{b,w(j')} = 0$. For this, we use (14), taking $k = j$, which in row b gives

$$0 + \sum_{a=p+1}^n \lambda_{j,a} z_{b,a} = \sum_{a=1}^p \lambda_{j,a} z_{b,a} = \lambda_{j,w(b)} = 0, \quad (29)$$

where the second and third equalities were shown in the previous paragraph.

We wish to isolate the $\lambda_{j,w(j')} z_{b,w(j')}$ term on the left-hand side of (29) by showing all other terms there vanish.

Subclaim 6.15. *For all $a \neq w(j')$ on the left-hand side of (29), either $\lambda_{j,a} = 0$ or $z_{b,a} = 0$.*

Proof of Subclaim 6.15. Given $p + 1 \leq a \leq n$, we prove the subclaim by case analysis on the position of the pivot in column a .

Since $\beta_b = +$ and $p + 1 \leq a \leq n$, the pivot of M in column a cannot be in row b . This leaves two other cases.

First, if the pivot of M in column a is in a row strictly north of row b , then this pivot occurs in row $\mathcal{I}(w^{-1}(a)) < b$. Note that we cannot have $\mathcal{I}(w^{-1}(a)) = j'$, since this would imply that $a = w(j)$, contradicting the fact that $p + 1 \leq a \leq n$ while $1 \leq w(j) \leq p$. Furthermore, we cannot have $\mathcal{I}(w^{-1}(a)) = j$, since this implies that $a = w(j')$, and we are excluding this case from consideration. Thus we need only consider the cases where $\mathcal{I}(w^{-1}(a)) < j$, where $j < \mathcal{I}(w^{-1}(a)) < j'$, and where $j' < \mathcal{I}(w^{-1}(a)) < b$. In the first case, we have that $\lambda_{j,a} = \lambda_{j,w(\mathcal{I}(w^{-1}(a)))} = 0$ by Claim 6.6. In the second case, we have $\lambda_{j,a} = 0$ by Claim 6.10. And in the third case, we have that $\lambda_{j,a} = 0$ by the inductive hypothesis.

The second possibility is that the pivot of M in column a is in a row strictly south of row b . But in this case (Z.6) implies that $z_{b,a} = 0$, so we are done.

This completes the proof of Subclaim 6.15. □

By Claim 6.9, we have $\lambda_{j,w(j')} = -\frac{1}{2}$. Hence, by Subclaim 6.15, (29) reduces to

$$\lambda_{j,w(j')} z_{b,w(j')} = -\frac{1}{2} z_{b,w(j')} = 0.$$

Therefore, $z_{b,w(j')} = 0$, as needed.

(β has a $-$ at b): The reasoning here is very similar to the previous case. We omit the details.

(β has the left end of a matching at b): By (O.2) in row b of M , there are 1's in positions $w(b)$ and $w(\mathcal{I}(b))$. In addition, by (Z.1) and (Z.5), M has 0's elsewhere in row b . Thus we obtain the claim's assertion that

$$z_{b,w(j)} = z_{b,w(j')} = 0.$$

Using this, taking $k = j$ in (14) and examining row b , we see that

$$\lambda_{j,w(b)} = \lambda_{j,w(\mathcal{I}(b))}. \quad (30)$$

Claim 6.14's hypothesis that $b > j'$, combined with Claim 6.12(II), indicates that $\lambda_{j,w(b)} = 0$. Hence, by (30) we obtain the required $\lambda_{j,w(b)} = \lambda_{j,w(\mathcal{I}(b))} = 0$.

(β has the right end of a matching at b and its left end is at $\mathcal{I}(b) < j$): Since this subcase assumes $\mathcal{I}(b) < j$, by Claim 6.6, we have that

$$\lambda_{j,w(\mathcal{I}(b))} = \lambda_{j,w(b)} = 0,$$

as desired.

By definition of w ,

$$w(\mathcal{I}(b)) < w(j) \leq p \quad \text{and} \quad p + 1 \leq w(b) < w(j'). \quad (31)$$

Now, $z_{b,w(\mathcal{I}(b))} = -1$ by (O.3), and column $w(j)$ is right of column $w(\mathcal{I}(b))$ by the first inequality of (31), so by (Z.4), we know that $z_{b,w(j)} = 0$. Similarly, $z_{b,w(b)} = 1$ by (O.3), and using the last inequality of (31) combined with (Z.8), we see that $z_{b,w(j')} = 0$ as well.

(β has the right end of a matching at b and its left end $\mathcal{I}(b)$ satisfies $j < \mathcal{I}(b) < j'$): By Claim 6.10 we know $\lambda_{j,w(b)} = \lambda_{j,w(\mathcal{I}(b))} = 0$. Here, the matchings $(\mathcal{I}(b) < b)$ and $(j < j')$ are in a 1212 pattern, so by (Z.9),

$$z_{b,w(j)} = -z_{b,w(j')}.$$

Thus we only need to see that $z_{b,w(j)} = 0$. For this, consider row b of (26), where we take $k = j$. Since $b > j'$, using Claim 6.12(II), (O.3), and (Z.4), we see that

$$\left(\sum_{a=1}^{w(\mathcal{I}(b))-1} \lambda_{j,a} z_{b,a} \right) - \lambda_{j,w(\mathcal{I}(b))} = 0. \quad (32)$$

It was argued already, in the proof of Claim 6.10 (see (R.3.2.1)–(R.3.2.3)), that the left-hand side of (32) reduces to $\frac{1}{2}z_{b,w(j)} - \lambda_{j,w(\mathcal{I}(b))}$, and we have noted above that $\lambda_{j,w(\mathcal{I}(b))} = 0$. So we have $\frac{1}{2}z_{b,w(j)} = 0$, whence $z_{b,w(j)} = 0$, as desired.

(β has the right end of a matching at b and its left end $\mathcal{I}(b)$ satisfies $j' < \mathcal{I}(b)$): Consider row $\mathcal{I}(b)$ of (14), where we take $k = j$. Since we assume $j' < \mathcal{I}(b)$, by (O.2), (Z.1) and (Z.5), (14) reads

$$\lambda_{j,w(b)} = \lambda_{j,w(\mathcal{I}(b))}. \quad (33)$$

By Claim 6.12(II) and the assumption $j' < \mathcal{I}(b)$, we see that $\lambda_{j,w(\mathcal{I}(b))} = 0$, and so by (33),

$$\lambda_{j,w(b)} = 0. \quad (34)$$

Let us show $z_{b,w(j)} = 0$. For this, we first analyze row b of (26), taking $k = j$. By (O.3), (Z.8) and Claim 6.12(II), this row is

$$\sum_{a=1}^{w(\mathcal{I}(b))-1} \lambda_{j,a} z_{b,a} - \lambda_{j,w(\mathcal{I}(b))} = 0.$$

Since by (33) and (34) we know $\lambda_{j,w(\mathcal{I}(b))} = 0$, this equation reduces to

$$\sum_{a=1}^{w(\mathcal{I}(b))-1} \lambda_{j,a} z_{b,a} = 0. \quad (35)$$

By Claim 6.5 (where $k = \mathcal{I}(b)$), all a indexing the summation in (35) satisfy

$$w^{-1}(a) < \mathcal{I}(b) < b.$$

Therefore, all $\lambda_{j,a}$ except for $\lambda_{j,w(j)}$ vanish, either by Claim 6.6, Claim 6.10, or the inductive hypothesis. In view of Claim 6.9,

$$\lambda_{j,w(j)} z_{b,w(j)} = \frac{1}{2} z_{b,w(j)}$$

does appear in the left-hand side of (35). Thus (35) actually states

$$\frac{1}{2} z_{b,w(j)} = 0,$$

whence $z_{b,w(j)} = 0$.

We now show $z_{b,w(j')} = 0$. For this, consider row b of (14) (taking $k = j$), which says that

$$0 + \sum_{a=p+1}^{w(b)-1} \lambda_{j,a} z_{b,a} + \lambda_{j,w(b)} = \sum_{a=1}^{w(\mathcal{I}(b))-1} \lambda_{j,a} z_{b,a} - \lambda_{j,w(\mathcal{I}(b))},$$

or equivalently, by (33) and (34),

$$\sum_{a=p+1}^{w(b)-1} \lambda_{j,a} z_{b,a} = \sum_{a=1}^{w(\mathcal{I}(b))-1} \lambda_{j,a} z_{b,a}.$$

By (35), the right hand side of this equation is zero, so we are reduced to

$$\sum_{a=p+1}^{w(b)-1} \lambda_{j,a} z_{b,a} = 0. \quad (36)$$

By Claim 6.5, all a indexing the sum satisfy $\mathcal{I}(w^{-1}(a)) < b$. Thus, all $\lambda_{j,a}$ vanish with the exception of $\lambda_{j,w(j')}$, either by Claim 6.6, Claim 6.10, or the inductive hypothesis. Thus (36) reduces to

$$\lambda_{j,w(j')} z_{b,w(j')} = -\frac{1}{2} z_{b,w(j')} = 0,$$

by Claim 6.9. Hence $z_{b,w(j')} = 0$.

This concludes the proof of Claim 6.14. \square

Now, we turn to rows j and j' . By (Z.1) and (Z.5), every entry in row j other than $z_{j,w(j)} = z_{j,w(j')} = 1$ is 0. Furthermore, by (Z.4) and (Z.8),

$$z_{j',a} = 0 \quad \text{except when } 1 \leq a \leq w(j) \quad \text{or} \quad p+1 \leq a \leq w(j').$$

Thus what remains to complete the proof of Proposition 6.1 is the following:

Claim 6.16 (Case C: row zeroness). $z_{j',a} = 0$ for all a satisfying either $1 \leq a < w(j)$ or $p+1 \leq a < w(j')$.

Proof. By Claim 6.5, it suffices to prove

$$z_{j',w(b)} = 0 \quad \text{and} \quad z_{j',w(\mathcal{I}(b))} = 0 \quad \text{for all } b < j.$$

We prove this by *reverse* induction on b . Our inductive hypothesis is that

$$z_{j',w(c)} = z_{j',w(\mathcal{I}(c))} = 0 \quad \text{for all } c \text{ with } b < c < j.$$

Now, in view of (O.3), (Z.1), and (Z.5), row j of Equation (14) (taking $k = b$) gives

$$\lambda_{b,w(j)} = \lambda_{b,w(j')}. \tag{37}$$

Now consider row j' of (14), again taking $k = b$. By (O.3), (Z.4) and (Z.8),

$$\sum_{a=1}^{w(j)-1} \lambda_{b,a} z_{j',a} - \lambda_{b,w(j)} = \sum_{a=p+1}^{w(j')-1} \lambda_{b,a} z_{j',a} + \lambda_{b,w(j')}. \tag{38}$$

Hence, by Claim 6.13(II), we have

$$\sum_{a=1}^{w(j)-1} \lambda_{b,a} z_{j',a} - \lambda_{b,w(j)} = 0, \tag{39}$$

so by (38),

$$\sum_{a=p+1}^{w(j')-1} \lambda_{b,a} z_{j',a} + \lambda_{b,w(j')} = 0 \tag{40}$$

as well.

We now consider multiple cases depending on the value of β_b .

($\beta_b = +$): By Claim 6.5, for a indexing the sum in (40), we have $\mathcal{I}(w^{-1}(a)) < j$. Now, if $\mathcal{I}(w^{-1}(a)) < b$, then $\lambda_{b,a} = 0$ by Claim 6.6. If $b < \mathcal{I}(w^{-1}(a)) < j$, then $z_{j',a} = 0$ by the inductive hypothesis. Note that $\mathcal{I}(w^{-1}(a)) = b$ is not possible, since this would imply that $a = w(b)$, contradicting the fact that $p+1 \leq a < w(j')$ while $1 \leq w(b) \leq p$. Thus (40) reduces to $\lambda_{b,w(j')} = 0$. Then by (37), $\lambda_{b,w(j)} = 0$ also. Then Equation (39) gives

$$\sum_{a=1}^{w(j)-1} \lambda_{b,a} z_{j',a} = 0. \tag{41}$$

Similarly to the previous paragraph,

$$\lambda_{b,a} = 0 \quad \text{if } w^{-1}(a) < b, \quad \text{and} \quad z_{j',a} = 0 \quad \text{if } w^{-1}(a) > b.$$

Hence (41) reduces to $\lambda_{b,w(b)}z_{j',w(b)} = 0$. Since $\lambda_{b,w(b)} \neq 0$ by Claim 6.8, we have $z_{j',w(b)} = 0$. Since $\beta_b = +$, $\mathcal{I}(b) = b$, so $z_{j',w(\mathcal{I}(b))} = 0$, as desired.

($\beta_b = -$): This is similar to the previous case, except the roles of Equations (39) and (40) are switched.

(β_b is a right endpoint): First, consider (39). Since $1 \leq a < w(j)$, by Claim 6.5 we have that $w^{-1}(a) < j$. So consider the cases $w^{-1}(a) < \mathcal{I}(b)$, $w^{-1}(a) = \mathcal{I}(b)$, $\mathcal{I}(b) < w^{-1}(a) < b$, and $b < w^{-1}(a) < j$. (Note that $w^{-1}(a) = b$ is not possible, since $1 \leq a < w(j) \leq p$, while $p + 1 \leq w(b) \leq n$.) In the first case, we have that $\lambda_{b,a} = 0$ by Claim 6.6. In the second case, we simply have that $a = w(\mathcal{I}(b))$. In the third case, we have that $\lambda_{b,a} = 0$ by Claim 6.10. Finally, in the fourth case, we have that $z_{j',a} = 0$ by the inductive hypothesis. Thus by Claim 6.9, (39) reduces to

$$-\frac{1}{2}z_{j',w(\mathcal{I}(b))} - \lambda_{b,w(j)} = 0. \tag{42}$$

Now, consider (40). Since $p + 1 \leq a < w(j')$, Claim 6.5 says that $\mathcal{I}(w^{-1}(a)) < j$. So consider the cases $\mathcal{I}(w^{-1}(a)) < \mathcal{I}(b)$, $\mathcal{I}(w^{-1}(a)) = \mathcal{I}(b)$, $\mathcal{I}(b) < \mathcal{I}(w^{-1}(a)) < b$, and $b < \mathcal{I}(w^{-1}(a)) < j$. (As above, $\mathcal{I}(w^{-1}(a)) = b$ cannot occur, since this would imply that $a = w(\mathcal{I}(b))$, but $p + 1 \leq a < w(j')$ while $1 \leq w(\mathcal{I}(b)) < w(j) \leq p$.) In the first case, $\lambda_{b,a} = 0$ by Claim 6.6. In the second case, we simply have that $a = w(b)$. In the third case, $\lambda_{b,a} = 0$ by Claim 6.10. Lastly, in the fourth case, we have that $z_{j',a} = 0$ by induction. Thus using Claim 6.9 again, (40) reduces to

$$-\frac{1}{2}z_{j',w(b)} + \lambda_{b,w(j')} = 0. \tag{43}$$

We can make arguments similar to the preceding ones when examining rows j and j' of (14), but taking $k = \mathcal{I}(b)$ instead. Doing so, analogously to (37), we obtain

$$\lambda_{\mathcal{I}(b),w(j)} = \lambda_{\mathcal{I}(b),w(j')}. \tag{44}$$

Analogously to (42) and (43), we obtain

$$\frac{1}{2}z_{j',w(\mathcal{I}(b))} - \lambda_{\mathcal{I}(b),w(j)} = 0 \tag{45}$$

and

$$-\frac{1}{2}z_{j',w(b)} + \lambda_{\mathcal{I}(b),w(j')} = 0. \tag{46}$$

The only simultaneous solution to (37), (44), (42), (43), (45), and (46) is

$$z_{j',w(\mathcal{I}(b))} = z_{j',w(b)} = \lambda_{\mathcal{I}(b),w(j)} = \lambda_{\mathcal{I}(b),w(j')} = \lambda_{b,w(j)} = \lambda_{b,w(j')} = 0.$$

(β_b is a left endpoint, and its right end satisfies $j' < \mathcal{I}(b)$): We have $z_{j',w(b)} = z_{j',w(\mathcal{I}(b))} = 0$ by (Z.3) and (Z.7) applied to columns $w(b)$ and $w(\mathcal{I}(b))$ respectively.

(β_b is a left endpoint, and its right end satisfies $j < \mathcal{I}(b) < j'$): Since the matchings $(b < \mathcal{I}(b))$ and $(j < j')$ are in a 1212 pattern, by (Z.9),

$$z_{j',w(b)} = -z_{j',w(\mathcal{I}(b))}. \quad (47)$$

By precisely the arguments that preceded (42) and (43), we see that (39) reduces to

$$\lambda_{b,w(b)} z_{j',w(b)} - \lambda_{b,w(j)} = 0,$$

while (40) reduces to

$$\lambda_{b,w(\mathcal{I}(b))} z_{j',w(\mathcal{I}(b))} + \lambda_{b,w(j')} = 0.$$

Using Claim 6.9, (37), and (47), we have

$$\frac{1}{2} z_{j',w(b)} - \lambda_{b,w(j)} = 0 \quad \text{and} \quad \frac{1}{2} z_{j',w(b)} + \lambda_{b,w(j)} = 0.$$

This implies $z_{j',w(b)} = \lambda_{b,w(j)} = 0$. Then we also have $z_{j',w(\mathcal{I}(b))} = 0$ by (47), as required.

(β_b is a left endpoint, and its right end satisfies $\mathcal{I}(b) < j$): In this case, the necessary claims are already proved by induction, since $b < \mathcal{I}(b) < j$.

This completes the proof of Claim 6.16, and hence the proof of Proposition 6.1. \square

7. Conjectures and problems

Conjecture 7.1. $\mathcal{I}_{\gamma,\alpha}$ is a radical ideal.

Conjecture 7.1 has been verified using Macaulay 2 through $p + q = 6$. This conjecture would follow from a solution to:

Problem 7.2. Find a Gröbner basis for $\mathcal{I}_{\gamma,\alpha}$ with square-free lead terms.

The analogous problem for *Kazhdan–Lusztig ideals* was solved in [Woo and Yong 2012]; it was shown that the defining generators of the ideal form a Gröbner basis. In contrast, we have:

Example 7.3. Let $\alpha = -1221+$ and $\gamma = 12-+12$. The defining generators of $\mathcal{I}_{\gamma,\alpha}$ are

$$z_{4,4}z_{6,6} + z_{5,4}z_{6,5} - z_{6,4}, \quad -\frac{1}{8}z_{4,4}z_{6,6} - \frac{1}{8}z_{5,4}z_{6,5} + \frac{1}{4}z_{6,4}.$$

Thus $z_{6,4} \in \mathcal{I}_{\gamma,\alpha}$, and hence $z_{4,4}z_{6,4} + z_{5,4}z_{6,5} \in \mathcal{I}_{\gamma,\alpha}$. Under any term order, the initial ideal must contain $z_{6,4}$ and either $z_{4,4}z_{6,4}$ or $z_{5,4}z_{6,5}$. However, at most one of these monomials can be realized as a lead term of the defining generators (for a fixed choice of term order). Hence the defining generators cannot be a Gröbner basis under any term order.

The *maximal singular locus* $\text{Maxsing}(Y_\gamma)$ is the set of K -orbits \mathcal{O}_α such that Y_γ is singular along \mathcal{O}_α and \mathcal{O}_α is maximal in Bruhat order with respect to this property.

Whereas the theorem of McGovern [2009] recalled in the introduction combinatorially characterizes which Y_γ are singular, the following question is open:

Problem 7.4. Give a combinatorial description of $\text{Maxsing}(Y_\gamma)$.

In [Woo and Wyser 2015], a solution to Problem 7.4 is given for clans γ which are “1212-avoiding”. This gives a complete solution for the case $K = \mathrm{GL}_{n-1} \times \mathrm{GL}_1$ (since all $(n-1, 1)$ -clans are 1212-avoiding). For the case of $K = \mathrm{GL}_{n-2} \times \mathrm{GL}_2$, we have a conjectural solution, but it is lengthy to state, so we omit it here.

The orbit closure Y_γ is *rationally smooth* along \mathcal{O}_α if the Kazhdan–Lusztig–Vogan polynomial $P_{\gamma,\alpha}(q)$ equals 1, and Y_γ is (globally) rationally smooth if $P_{\gamma,\alpha}(q) = 1$ for every $\alpha \leq \gamma$. It is known [McGovern 2009] that Y_γ is rationally smooth if and only if Y_γ is smooth. Using data from the ATLAS project, we have verified the following conjecture for $(p, q) = (2, 2), (3, 2)$:

Conjecture 7.5. *For $\alpha \leq \gamma$, Y_γ is rationally smooth on \mathcal{O}_α if and only if Y_γ is smooth on \mathcal{O}_α .*

Let (R, \mathfrak{m}) be the local ring of a point p in a projective variety X . The *associated graded ring* is

$$\mathrm{gr}_{\mathfrak{m}} R = \bigoplus_{i \geq 0} \mathfrak{m}^i / \mathfrak{m}^{i+1}.$$

Conjecture 7.6. *Let (R, \mathfrak{m}) be the local ring associated to any point $p \in \mathcal{O}_\alpha \subseteq Y_\gamma$. Then $\mathrm{gr}_{\mathfrak{m}} R$ is Cohen–Macaulay and reduced. Moreover, it is Gorenstein whenever Y_γ is Gorenstein along \mathcal{O}_α .*

This has been checked for $(p, q) = (2, 2), (3, 2), (3, 3), (4, 3)$. This conjecture does not follow from Conjecture 7.1, since reducedness, Cohen–Macaulayness, and Gorensteinness may be lost on degenerating to the associated graded ring.

The *projectivized tangent cone* is $\mathrm{Proj}(\mathrm{gr}_{\mathfrak{m}} R)$. The (*Hilbert–Samuel*) *multiplicity* of $p \in X$ is

$$\mathrm{mult}_p(X) = \mathrm{deg}(\mathrm{Proj}(\mathrm{gr}_{\mathfrak{m}} R)).$$

This statistic provides singularity information. Specifically, $\mathrm{mult}_p(X) = 1$ if and only if p is a smooth point of X . Let $\mathrm{mult}_{\gamma,\alpha}$ be $\mathrm{mult}_p(Y_\gamma)$ for any point $p \in \mathcal{O}_\alpha$.

Define

$$\mathrm{maxmult}(Y_\gamma) = \max_{\alpha \leq \gamma} \mathrm{mult}_{\gamma,\alpha}.$$

Since multiplicity is a semicontinuous numerical invariant, $\mathrm{maxmult}(Y_\gamma) = \mathrm{mult}_{\gamma,\alpha}$ for some matchless $\alpha \leq \gamma$ (so \mathcal{O}_α is a closed orbit). A search for a combinatorial rule for $\mathrm{mult}_{\gamma,\alpha}$ might be partially guided by a solution to the following problem:

Problem 7.7. *Is the maximum value of $\mathrm{maxmult}(Y_\gamma)$ for $\gamma \in \mathrm{Clans}_{p,q}$ achieved at $\gamma_{\max} = 1 + p^{-1} - q^{-1} 1 \in \mathrm{Clans}_{p,q}$?*

We have checked that the answer to Problem 7.7 is affirmative $p + q \leq 6$. Note the maximizer in Problem 7.7 need not be unique. For instance, when $p = q = 2$, the maximum is achieved at $\gamma = 1212$, $\gamma = 1+-1$, and $\gamma = 1-+1$, with $\mathrm{maxmult}(Y_\gamma) = 2$.

Example 7.8 (Analogue of Lusztig’s conjecture is false). Note that $P_{1+-1,++--}(q) = q + 1$, whereas $P_{1212,++--}(q) = 1$. However, $[++--, 1+-1] \cong [++--, 1212]$ as posets (see Figure 1). Thus, the KLV polynomial is not an invariant of the poset in Bruhat order. This contrasts with the situation for Schubert




Figure 1. Bruhat order on $(\mathrm{GL}(4, \mathbb{C}), \mathrm{GL}(2, \mathbb{C}) \times \mathrm{GL}(2, \mathbb{C}))$.

varieties, where a conjecture attributed to Lusztig asserts that the Kazhdan–Lusztig polynomial is a poset invariant; see for example [Brenti 2002/04]. Similarly, $\mathrm{mult}_{1+-1,++--} = 2$ whereas $\mathrm{mult}_{1212,++--} = 1$.

The *Cohen–Macaulay type* of a local Cohen–Macaulay ring R is $\dim_{\mathbb{k}} \mathrm{Ext}_R^{\dim R}(\mathbb{k}, R)$. A ring R is *Gorenstein* if its Cohen–Macaulay type is 1.

Define the *maximal non-Gorenstein locus* $\mathrm{MaxnonGor}(Y_\gamma) = \{\mathcal{O}_\alpha\}$ to be the set of K -orbits $\mathcal{O}_\alpha \subseteq Y_\gamma$ such that Y_γ is non-Gorenstein at any point of \mathcal{O}_α and \mathcal{O}_α is maximal in Bruhat order with respect to this property.

Conjecture 7.9. $\mathrm{MaxnonGor}(Y_\gamma) \subseteq \mathrm{Maxsing}(Y_\gamma)$.

If Y_γ is non-Gorenstein along $\mathcal{O}_\alpha \in \mathrm{Maxsing}(Y_\gamma)$ then $\mathcal{O}_\alpha \in \mathrm{MaxnonGor}(Y_\gamma)$. However, if Y_γ is Gorenstein along some $\mathcal{O}_\alpha \in \mathrm{Maxsing}(Y_\gamma)$, there a priori may be $\mathcal{O}_\beta \in \mathrm{MaxnonGor}(Y_\gamma)$ with $\beta < \alpha$. Conjecture 7.9 asserts that this does not occur. This conjecture is an analogue of [Woo and Yong 2008, Conjecture 6.7].

We verified Conjecture 7.9 for all (p, q) with $p + q \leq 7$ and for many cases of $(p, q) = (4, 4)$. In Table 1, we give the singularity data $(\gamma, \ell(\gamma), \mathrm{Maxsing}(Y_\gamma), \mathrm{MaxnonGor}(Y_\gamma))$ for $p = 3, q = 2$ for all the clans γ where Y_γ is singular.

Problem 7.10. *When is Y_γ (globally) Gorenstein?*

An answer to this question has been given by the first two authors in [Woo and Wyser 2015] in the event that γ is 1212-avoiding. As mentioned in the introduction, it is shown in [loc. cit.] that Gorensteinness cannot be characterized by ordinary pattern avoidance in general. However, for small q , it apparently can.

γ	$\ell(\gamma)$	$\text{Maxsing}(Y_\gamma)$	$\text{MaxnonGor}(Y_\gamma)$
122+1	5	{1-1++}	{}
1+221	5	{++1-1}	{}
12+12	5	{-++++-, +1-1+}	{}
1+212	4	{++--+, 11++-}	{++--+}
1-++1	4	{---+++}	{---+++}
121+2	4	{+--+++, -++11}	{+--+++}
1+-+1	4	{11-+++, +-11}	{}
1++-1	4	{+++--}	{+++--}
1+-1+	3	{++--+}	{}
+1+-1	3	{+++--}	{}
+1-+1	3	{+--++}	{}
1212+	3	{+--+++, -++-+}	{}
+1212	3	{+-++-, +-+-+}	{}
1-+1+	3	{---+++}	{}

Table 1. The singularity data for $p = 3, q = 2$ for all the clans γ where Y_γ is singular.

When $q = 1$, all orbit closures are smooth (hence Gorenstein). Any $(p, 1)$ -clan ($p \geq 1$) has at most one matching, and if it has a matching, it can have no $-$ signs. Thus it necessarily avoids all of McGovern’s singular patterns recalled in the introduction.

Next consider the case $q = 2$. In this case, the Gorensteinness criterion of [Woo and Wyser 2015] is easily seen to amount to the following ordinary pattern avoidance criterion.

Fact [Woo and Wyser 2015, Proposition 3.4.2]. *If γ is a 1212-avoiding $(p, 2)$ -clan with $p \geq 2$, then Y_γ is Gorenstein if and only if γ avoids $1++-1, 1-++1, 1++221$, and $122++1$.*

Experimentally, it seems that when $q = 2$, Gorensteinness can be characterized by ordinary pattern avoidance even if we allow 1212-including clans.

Conjecture 7.11. *If γ is any $(p, 2)$ -clan with $p \geq 2$, then Y_γ is Gorenstein if and only if γ avoids $1++-1, 1-++1, 1++221, 122++1, 1+212$, and $121+2$.*

This conjecture holds (by computation) for $n \leq 7$. The example recalled in the introduction says that there is no ordinary pattern avoidance characterization when $q > 2$.

Recall that a local ring R is said to be a *local complete intersection* (lci) if it is the quotient of a regular local ring by an ideal generated by a regular sequence. A variety is lci if each of its local rings are lci. Every smooth variety is lci, while every lci variety is Gorenstein (and hence Cohen–Macaulay).

At present, we do not have a characterization of which Y_γ are lci in general. An ordinary pattern avoidance criterion is given in [Woo and Wyser 2015] for 1212-avoiding clans. For general p and q , there are many non-lci patterns that must be avoided. However, when $q = 2$, the list of non-lci patterns that can actually occur is much smaller.

Fact [Woo and Wyser 2015, Proposition 3.3.3]. *If γ is a 1212-avoiding $(p, 2)$ -clan with $p \geq 2$, then Y_γ is lci if and only if γ avoids $1++-1$, $1-++1$, $1++221$, and $122++1$. Thus Y_γ is lci if and only if it is Gorenstein.*

It is conjectured in [Woo and Wyser 2015] that ordinary pattern avoidance can be used to characterize lci-ness in general, whether γ is 1212-avoiding or not. The complete list of non-lci patterns is not known, even conjecturally. However, for $q = 2$, we have:

Conjecture 7.12. *If γ is any $(p, 2)$ -clan with $p \geq 2$, then Y_γ is lci if and only if γ avoids $1++-1$, $1-++1$, $1++221$, $122++1$, $1+212$, and $121+2$. Equivalently, we conjecture that Y_γ is lci if and only if it is Gorenstein.*

This conjecture has been checked for $n \leq 7$. The equivalence of lci-ness and Gorensteinness breaks down when $q > 2$; for example, Y_{122331} is Gorenstein but not lci.

Acknowledgements

We thank Frank Sottile for a conversation years ago about [Woo and Yong 2008] that ultimately led to this project. We thank Michel Brion for many helpful comments and in particular for his suggestion to consider the slices of [Mars and Springer 1998], as well as his assistance with Lemmas 2.3 and 2.6. Woo was supported by NSA grant H98230-13-1-0242 and Simons Collaboration Grant 359792. Wyser was supported by NSF International Research Fellowship 1159045 and hosted by Institut Fourier in Grenoble. Yong was supported by NSF grants DMS 1201595 and DMS 1500691. This work was partially completed while Yong and Woo were visitors at the Fields Institute program on “Combinatorial algebraic geometry”; they thank the organizers and the institute for their hospitality.

References

- [Avramov 1975] L. L. Avramov, “Flat morphisms of complete intersections”, *Dokl. Akad. Nauk SSSR* **225**:1 (1975), 11–14. In Russian; translated in *Soviet Math. Dokl.* **16**:6 (1976), 1413–1417. MR Zbl
- [Brenti 2002/04] F. Brenti, “Kazhdan–Lusztig polynomials: history problems, and combinatorial invariance”, *Sém. Lothar. Combin.* **49** (2002/04), Art. B49b, 30. MR Zbl
- [Brion 2001] M. Brion, “On orbit closures of spherical subgroups in flag varieties”, *Comment. Math. Helv.* **76**:2 (2001), 263–299. MR Zbl
- [Fulton 1997] W. Fulton, *Young tableaux: with applications to representation theory and geometry*, London Mathematical Society Student Texts **35**, Cambridge University Press, 1997. MR Zbl
- [Lusztig and Vogan 1983] G. Lusztig and D. A. Vogan, Jr., “Singularities of closures of K -orbits on flag manifolds”, *Invent. Math.* **71**:2 (1983), 365–379. MR Zbl
- [Mars and Springer 1998] J. G. M. Mars and T. A. Springer, “Hecke algebra representations related to spherical varieties”, *Represent. Theory* **2** (1998), 33–69. MR Zbl
- [Matsuki and Ōshima 1990] T. Matsuki and T. Ōshima, “Embeddings of discrete series into principal series”, pp. 147–175 in *The orbit method in representation theory* (Copenhagen, 1988), edited by M. Duflo et al., Progr. Math. **82**, Birkhäuser, Boston, 1990. MR Zbl
- [Matsumura 1986] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, 1986. MR Zbl

- [McGovern 2009] W. M. McGovern, “Closures of K -orbits in the flag variety for $U(p, q)$ ”, *J. Algebra* **322**:8 (2009), 2709–2712. MR Zbl
- [McGovern 2015] W. M. McGovern, “Upper semicontinuity of KLV polynomials for certain blocks of Harish-Chandra modules”, pp. 437–441 in *Representations of reductive groups*, edited by M. Nevins and P. E. Trapa, Progr. Math. **312**, Springer, 2015. MR Zbl
- [Perrin 2014] N. Perrin, “On the geometry of spherical varieties”, *Transform. Groups* **19**:1 (2014), 171–223. MR Zbl
- [Richardson and Springer 1990] R. W. Richardson and T. A. Springer, “The Bruhat order on symmetric varieties”, *Geom. Dedicata* **35**:1-3 (1990), 389–436. MR Zbl
- [Ulfarsson and Woo 2013] H. Ulfarsson and A. Woo, “Which Schubert varieties are local complete intersections?”, *Proc. Lond. Math. Soc.* (3) **107**:5 (2013), 1004–1052. MR Zbl
- [Vogan 1983] D. A. Vogan, “Irreducible characters of semisimple Lie groups, III: Proof of Kazhdan–Lusztig conjecture in the integral case”, *Invent. Math.* **71**:2 (1983), 381–417. MR Zbl
- [Woo 2010] A. Woo, “Interval pattern avoidance for arbitrary root systems”, *Canad. Math. Bull.* **53**:4 (2010), 757–762. MR Zbl
- [Woo and Wyser 2015] A. Woo and B. J. Wyser, “Combinatorial results on $(1, 2, 1, 2)$ -avoiding $GL(p, \mathbb{C}) \times GL(q, \mathbb{C})$ -orbit closures on $GL(p+q, \mathbb{C})/B$ ”, *Int. Math. Res. Not.* **2015**:24 (2015), 13148–13193. MR Zbl
- [Woo and Yong 2006] A. Woo and A. Yong, “When is a Schubert variety Gorenstein?”, *Adv. Math.* **207**:1 (2006), 205–220. MR Zbl
- [Woo and Yong 2008] A. Woo and A. Yong, “Governing singularities of Schubert varieties”, *J. Algebra* **320**:2 (2008), 495–520. MR Zbl
- [Woo and Yong 2012] A. Woo and A. Yong, “A Gröbner basis for Kazhdan–Lusztig ideals”, *Amer. J. Math.* **134**:4 (2012), 1089–1137. MR Zbl
- [Wyser 2016] B. J. Wyser, “The Bruhat order on clans”, *J. Algebraic Combin.* **44**:3 (2016), 495–517. MR Zbl
- [Wyser and Yong 2014] B. J. Wyser and A. Yong, “Polynomials for $GL_p \times GL_q$ orbit closures in the flag variety”, *Selecta Math. (N.S.)* **20**:4 (2014), 1083–1110. MR Zbl
- [Yamamoto 1997] A. Yamamoto, “Orbits in the flag variety and images of the moment map for classical groups, I”, *Represent. Theory* **1** (1997), 329–404. MR Zbl

Communicated by Victor Reiner

Received 2017-06-10 Accepted 2017-08-17

awoo@uidaho.edu

Department of Mathematics, University of Idaho, Moscow, ID, United States

bwyser@okstate.edu

Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL, United States

Current address:

Department of Mathematics, Oklahoma State University, Stillwater, OK, United States

ayong@uiuc.edu

Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL, United States

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Martin Olsson	University of California, Berkeley, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Raman Parimala	Emory University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	University of California, Santa Cruz, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Michael Rapoport	Universität Bonn, Germany
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Hubert Flenner	Ruhr-Universität, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Christopher Skinner	Princeton University, USA
Joseph Gubeladze	San Francisco State University, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Roger Heath-Brown	Oxford University, UK	J. Toby Stafford	University of Michigan, USA
Craig Huneke	University of Virginia, USA	Pham Huu Tiep	University of Arizona, USA
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Ravi Vakil	Stanford University, USA
János Kollár	Princeton University, USA	Michel van den Bergh	Hasselt University, Belgium
Yuri Manin	Northwestern University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Shou-Wu Zhang	Princeton University, USA
Shigefumi Mori	RIMS, Kyoto University, Japan		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2018 is US \$340/year for the electronic version, and \$535/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 12 No. 1 2018

Local positivity of linear series on surfaces ALEX KÜRONYA and VICTOR LOZOVANU	1
The mean value of symmetric square L -functions OLGA BALKANOVA and DMITRY FROLENKOV	35
F -signature and Hilbert–Kunz multiplicity: a combined approach and comparison THOMAS POLSTRA and KEVIN TUCKER	61
Regular pairs of quadratic forms on odd-dimensional spaces in characteristic 2 IGOR DOLGACHEV and ALEXANDER DUNCAN	99
Graded Steinberg algebras and their representations PERE ARA, ROOZBEH HAZRAT, HUANHUAN LI and AIDAN SIMS	131
Governing singularities of symmetric orbit closures ALEXANDER WOO, BENJAMIN J. WYSER and ALEXANDER YONG	173



1937-0652(2018)12:1;1-G