

Algebra & Number Theory

Volume 17
2023
No. 10

Separation of periods of quartic surfaces

Pierre Lairez and Emre Can Sertöz



Separation of periods of quartic surfaces

Pierre Lairez and Emre Can Sertöz

We give a computable lower bound for the distance between two distinct periods of a given quartic surface defined over the algebraic numbers. The main ingredient is the determination of height bounds on components of the Noether–Lefschetz loci. This makes it possible to study the Diophantine properties of periods of quartic surfaces and to certify a part of the numerical computation of their Picard groups.

1. Introduction

Periods are a countable set of complex numbers containing all the algebraic numbers, as well as many of the transcendental constants of nature. In light of the ubiquity of periods in mathematics and the sciences, Kontsevich and Zagier [2001] asked for the development of an algorithm to check for the equality of two given periods. We solve this problem for periods coming from quartic surfaces by giving a computable separation bound, that is, a lower bound on the minimum distance between distinct periods.

Let $f \in \mathbb{C}[w, x, y, z]_4$ be a homogeneous quartic polynomial defining a smooth quartic X_f in $\mathbb{P}^3(\mathbb{C})$. The periods of X_f are the integrals of a nowhere vanishing holomorphic 2-form on X_f over integral 2-cycles in X_f . The periods can also be given in the form of integrals of a rational function

$$\frac{1}{2\pi i} \oint_{\gamma} \frac{dx \, dy \, dz}{f(1, x, y, z)}, \quad (1)$$

where γ is a 3-cycle in $\mathbb{C}^3 \setminus X_f$. The integral (1) depends only on the homology class of γ . These periods form a group under addition. The geometry of quartic surfaces dictates that there are only 21 independent 3-cycles in $\mathbb{C}^3 \setminus X_f$. These give 21 periods $\alpha_1, \dots, \alpha_{21} \in \mathbb{C}$ such that the integral over any other 3-cycle is an integer linear combination of these periods.

It is possible to compute the periods to high precision [Sertöz 2019], typically to thousands of decimal digits, and to deduce from them interesting algebraic invariants such as the Picard group of X_f [Lairez and Sertöz 2019]. This point of view has been fruitful for computing algebraic invariants for algebraic curves from their periods [van Wamelen 1999; Costa et al. 2019; Bruin et al. 2019; Booker et al. 2016].

For quartic surfaces, the computation of the Picard group reduces to computing the lattice in \mathbb{Z}^{21} of integer relations $x_1\alpha_1 + \dots + x_{21}\alpha_{21} = 0$, where $x_i \in \mathbb{Z}$. A basis for this lattice can be guessed from approximate α_i 's using lattice reduction algorithms. But is it possible to *prove* that all guessed relations are true relations? Previous work related to this question [Simpson 2008] required explicit construction of

MSC2020: primary 14Q10, 14J28, 32G20; secondary 11Y16, 14Q20, 11J99.

Keywords: K3 surfaces, periods, Diophantine approximation, Hodge loci, effective mathematics.

algebraic curves on X_f , which becomes challenging very quickly. Instead, we give a method of proving relations by checking them at a predetermined finite precision. At the moment, this is equally challenging, but we conjecture that the numerical approach can be made asymptotically faster, see Section 4.4 for details.

The Lefschetz theorem on $(1, 1)$ -classes (Section 2.2) associates a divisor on X_f to any integer relation between the periods of X_f . In turn, the presence of a divisor imposes algebraic conditions on the coefficients of f . Such algebraic conditions define the *Noether–Lefschetz loci* on the space of quartic polynomials (Section 3). In addition to the degree computations of Maulik and Pandharipande [2013], we give height bounds on the polynomial equations defining the Noether–Lefschetz loci (Theorem 14). These lead to our main result (Theorem 17): Assume f has algebraic coefficients, then for $x_i \in \mathbb{Z}$,

$$x_1\alpha_1 + \cdots + x_{21}\alpha_{21} = 0 \quad \text{or} \quad |x_1\alpha_1 + \cdots + x_{21}\alpha_{21}| > 2^{-c^{\max_i |x_i|^9}} \quad (2)$$

for some constant $c > 0$ depending only on f and the choice of the 21 independent 3-cycles (see Theorem 17 for a coordinate-free formulation). The constant c is computable in rather simple terms and without prior knowledge of the Picard group of X_f . We use the term “computable” in the sense of “computable with a Turing machine”, not “primitive recursive”, as our suggested algorithm to compute c depends, through Lemma 1, on the numerical computation of a nonzero constant (depending on f), whose magnitude is not known *a priori*, only the fact that it is nonzero.

The expression (2) is essentially a lower bound for the linear independence measure [Shidlovskii 1989, Chapter 11] for the periods of X_f . Our construction of this bound bears a loose resemblance to the ideas involved in the statement of the analytic subgroup theorem [Wüstholz 1989], and in particular, to the Masser–Wüstholz period theorem [1993]. We briefly comment on this analogy in Section 5.4.

As a consequence of the separation bound (2), we apply a construction in the manner of Liouville [1851] and prove, for instance, that the number

$$\sum_{n \geq 0} (2 \uparrow\uparrow 3n)^{-1} \quad (3)$$

is not a quotient of two periods of a single quartic surface that is defined over $\overline{\mathbb{Q}}$, where $2 \uparrow\uparrow 3n$ denotes an exponentiation tower with $3n$ twos (Theorem 19, with $\theta_{i+1} = 2^{2^{\theta_i}}$).

The methods we employed to attain the period separation bound (2) can, in principle, be generalized to separate the periods of some other algebraic varieties, e.g., of cubic fourfolds. We discuss these and other generalizations in Section 5.

2. Periods and deformations

2.1. Construction of the period map. For any nonzero homogeneous polynomial f in $\mathbb{C}[w, x, y, z]$, let X_f denote the surface in \mathbb{P}^3 defined as the zero locus of f . Let $R \doteq \mathbb{C}[w, x, y, z]$, and let $R_4 \subset R$ be the subspace of degree 4 homogeneous polynomials. Let $U_4 \subset R_4$ denote the dense open subset of all homogeneous polynomials f of degree 4 such that X_f is smooth. For our purposes, it will be useful

to consider not only the periods of a single quartic surface X_f , but also the *period map*, to study the dependence of periods on f .

The topology of X_f does not depend on f as long as X_f is smooth: given two polynomials $f, g \in U_4$, we can connect them by a continuous path in U_4 , and the surface X_f deforms continuously along this path, giving a homeomorphism $X_f \simeq X_g$, which is uniquely defined up to isotopy. In particular, if we fix a base point $b \in U_4$, then for every $f \in \tilde{U}_4$, where \tilde{U}_4 is a universal covering of U_4 , we have a uniquely determined isomorphism of cohomology groups $H^2(X_b, \mathbb{Z}) \simeq H^2(X_f, \mathbb{Z})$. Let $H_{\mathbb{Z}}$ denote the second cohomology group of X_b , which is isomorphic to \mathbb{Z}^{22} , e.g., [Huybrechts 2016, §1.3.3].

The hyperplane class and its multiples are redundant for the problem we are interested in, as their periods are 0. In practice, therefore, we work with a rank 21 quotient lattice. The map (7) below identifies this quotient with the cohomology of the complement of the quartic.

An element of \tilde{U}_4 determines a polynomial $f \in U_4$ together with an identification of $H^2(X_f, \mathbb{Z})$ with $H_{\mathbb{Z}}$. We often work locally around a given polynomial f and, in that case, we do not actively distinguish between U_4 and its universal covering.

The group $H_{\mathbb{Z}}$ is endowed with an even unimodular pairing

$$(x, y) \in H_{\mathbb{Z}} \times H_{\mathbb{Z}} \rightarrow x \cdot y \in \mathbb{Z}, \tag{4}$$

given by the intersection form on cohomology. Through this pairing, the second homology and cohomology groups are canonically identified with one another. For K3 surfaces, such as smooth quartic surfaces in \mathbb{P}^3 , the structure of the lattice $H_{\mathbb{Z}}$ with its intersection form is explicitly known [Huybrechts 2016, Proposition 1.3.5]. The fundamental class of a generic hyperplane section of X_f gives an element of $H_{\mathbb{Z}}$ denoted by h .

Further, the complex cohomology group $H^2(X_f, \mathbb{C})$, which is just $H_{\mathbb{C}} \doteq H_{\mathbb{Z}} \otimes \mathbb{C}$, is isomorphic to the corresponding de Rham cohomology $H_{\text{dR}}^2(X_f, \mathbb{C})$ group as follows: Elements of $H_{\text{dR}}^2(X_f, \mathbb{C})$ are represented by differential 2-forms. To a form Ω , one associates the element $\Theta(\Omega)$ of $H^2(X_f, \mathbb{C})$ given by the map

$$\Theta(\Omega): [\gamma] \in H_2(X_f, \mathbb{C}) \mapsto \int_{\gamma} \Omega \in \mathbb{C}. \tag{5}$$

The group $H_{\text{dR}}^2(X_f, \mathbb{C})$ has a distinguished element Ω_f , a nowhere vanishing holomorphic 2-form, described below. Every other holomorphic 2-form on X_f is a scalar multiple of Ω_f [Huybrechts 2016, Example 1.1.3]. Mapping Ω_f to $H_{\mathbb{C}}$ gives rise to the *period map*

$$\mathcal{P}: f \in \tilde{U}_4 \mapsto \omega_f \doteq \Theta(\Omega_f) \in H_{\mathbb{C}}. \tag{6}$$

The coordinates of the *period vector* ω_f , in some fixed basis of $H_{\mathbb{Z}}$, generates the group of periods of X_f .

There is a standard Thom–Gysin type map in homology

$$T: H_2(X_f, \mathbb{Z}) \rightarrow H_3(\mathbb{P}^3 \setminus X_f, \mathbb{Z}), \tag{7}$$

see [Voisin 2003, p. 159] for a modern description. Roughly speaking, T takes the class of a 2-cycle in X_f and returns the class of a narrow S^1 -bundle around the cycle lying entirely in $\mathbb{P}^3 \setminus X_f$. See [Griffiths 1969a, §3] for this classical interpretation. The map T is a surjective morphism, and its kernel is generated by the class of a hyperplane section of X_f .

We choose Ω_f so that the following identity holds:

$$\int_{\gamma} \Omega_f = \frac{1}{2\pi i} \int_{T(\gamma)} \frac{dx dy dz}{f(1, x, y, z)}. \tag{8}$$

Therefore, in view of (5), the coefficients of ω_f in a basis of $H_{\mathbb{Z}}$ coincides with periods as defined in (1).

The image \mathcal{D} of the period map \mathcal{P} is called the *period domain*. It admits a simple description

$$\mathcal{D} \doteq \mathcal{P}(\tilde{U}_4) = \{w \in H_{\mathbb{C}} \setminus \{0\} \mid w \cdot h = 0, w \cdot w = 0, w \cdot \bar{w} > 0\}, \tag{9}$$

where “ \cdot ” denotes the intersection form on $H_{\mathbb{Z}}$, extended to $H_{\mathbb{C}}$ by \mathbb{C} -linearity, and h denotes the fundamental class of a hyperplane section, as introduced above [Huybrechts 2016, Chapter 6]. Moreover, by the local Torelli theorem for K3 surfaces [Huybrechts 2016, Proposition 6.2.8], the map \mathcal{P} is a submersion: its derivative at any point of \tilde{U}_4 is surjective.

2.2. The Lefschetz (1,1)-theorem. Lefschetz proved that the linear integer relations between the periods of a quartic surface X_f are in correspondence with homology classes coming from algebraic curves in X_f . We now explain this statement in more detail. Let $C \subset X_f$ be an algebraic curve. Its fundamental class is the element $[C]$ of $H_{\mathbb{Z}}$ obtained as the Poincaré dual of the homology class of C . Here, we identify $H_{\mathbb{Z}}$ with $H^2(X_f, \mathbb{Z})$ by fixing a preimage of f in \tilde{U}_4 . The Picard group $\text{Pic}(X_f)$ of X_f is the sublattice of $H_{\mathbb{Z}}$ spanned by the fundamental classes of algebraic curves.

It follows from the definition that for any class $[\Omega] \in H_{\text{dR}}^2(X_f)$ of a differential 2-form on X_f ,

$$[C] \cdot \Theta(\Omega) = \int_C \Omega. \tag{10}$$

Moreover, if Ω is a holomorphic 2-form, then $\int_C \Omega = 0$, because the restriction of Ω to the complex 1-dimensional subvariety C vanishes. In particular $[C] \cdot \omega_f = 0$. It turns out that this condition characterizes the elements of $\text{Pic}(X_f)$.

More precisely, let $H^{1,1}(X_f) \subset H_{\mathbb{C}}$ denote the space orthogonal to ω_f and $\bar{\omega}_f$, the conjugate of ω_f , with respect to the intersection form. This space is a direct summand in the Hodge decomposition of $H^2(X_f, \mathbb{C})$.

The Lefschetz (1,1)-theorem [Griffiths and Harris 1978, p. 163] asserts that the lattice of integer relations coincide with the Picard group

$$\text{Pic}(X_f) = H_{\mathbb{Z}} \cap H^{1,1}(X_f). \tag{11}$$

Noting that for any $\gamma \in H_{\mathbb{Z}}$, we have $\bar{\gamma} = \gamma$, where $\bar{\gamma}$ denotes the complex conjugate, it follows that $\overline{\omega_f \cdot \bar{\gamma}} = \bar{\omega}_f \cdot \gamma$, so that (11) becomes

$$\text{Pic}(X_f) = \{\gamma \in H_{\mathbb{Z}} \mid \gamma \cdot \omega_f = 0\}. \tag{12}$$

2.3. A deformation argument. Let $\gamma_1, \dots, \gamma_{22}$ be a basis of $H_{\mathbb{Z}}$. The space $H_{\mathbb{R}}$ (respectively, $H_{\mathbb{C}}$) is endowed with the coefficientwise Euclidean (respectively, Hermitian) norm

$$\left\| \sum_{i=1}^{22} x_i \gamma_i \right\|^2 \doteq \sum_{i=1}^{22} |x_i|^2. \tag{13}$$

For $\gamma \in H_{\mathbb{Z}}$, if $|\gamma \cdot \omega_f|$ is small enough, then γ is close to being an integer relation between the periods of X_f . We want to argue that, in this case, γ is a genuine integer relation between the periods of X_g for some polynomial $g \in U_4$ close to f .

Recall $f, g \in \tilde{U}_4$ means f and g are smooth quartics with second cohomology identified with $H_{\mathbb{Z}}$. The space \tilde{U}_4 inherits a metric from U_4 , so that $\tilde{U}_4 \rightarrow U_4$ is locally isometric. The metric on $U_4 \subset R_4 \simeq \mathbb{C}^{35}$ is induced by an inner product. The choice of an inner product will change the distances, but this is absorbed into the constants in the statements below.

Let $f \in \tilde{U}_4$ be fixed. For any $g \in R_4$ and $t \in \mathbb{C}$ small enough, the polynomials $f + tg \in R_4$ lift canonically to \tilde{U}_4 . For any $\gamma \in H_{\mathbb{C}}$, we consider the map

$$\phi_{\gamma, g}(t) \doteq \gamma \cdot \mathcal{P}(f + tg), \tag{14}$$

which is well defined and analytic in a neighborhood of 0 in \mathbb{C} .

Lemma 1. *There is a constant $C > 0$, depending only on f , such that for any $\gamma \in H_{\mathbb{C}}$ satisfying $\gamma \cdot h = 0$ and $|\gamma \cdot \bar{\omega}_f| \|\omega_f\| \leq \frac{1}{2} \|\gamma\| (\omega_f \cdot \bar{\omega}_f)$, there is a monomial $m \in R_4$ for which $|\phi'_{\gamma, m}(0)| \geq C \|\gamma\|$.*

Proof. Observe that for any monomial $m \in R_4$, we have $\phi'_{\gamma, m}(0) = \gamma \cdot d_f \mathcal{P}(m)$, where $d_f \mathcal{P}$ is the derivative at f of \mathcal{P} . Let Q be the positive semidefinite Hermitian form defined on $H_{\mathbb{C}}$ by

$$Q(\gamma) \doteq \sum_m |\gamma \cdot d_f \mathcal{P}(m)|^2, \tag{15}$$

where the sum is taken over the monomials in m . Since $\max_m |\phi'_{\gamma, m}(0)|^2 \geq (1/\dim R_4) Q(\gamma)$, it is enough to prove that $Q(\gamma) \geq C \|\gamma\|^2$ for some constant $C > 0$, when $\gamma \cdot h = 0$ and $|\gamma \cdot \bar{\omega}_f| \|\omega_f\| \leq \frac{1}{2} \|\gamma\| (\omega_f \cdot \bar{\omega}_f)$.

The form Q vanishes exactly on the orthogonal complement (for the intersection product) of the tangent space $T_{\omega_f} \mathcal{D}$ of \mathcal{D} at ω_f . By (9),

$$T_{\omega_f} \mathcal{D} = \{w \in H_{\mathbb{C}} \mid w \cdot h = w \cdot \omega_f = 0\}. \tag{16}$$

So the kernel of Q is $K \doteq \mathbb{C}h + \mathbb{C}\omega_f$. Moreover, let E be the orthogonal complement of $\mathbb{C}h + \mathbb{C}\bar{\omega}_f$ (still for the intersection product). Since $h \cdot \omega_f = h \cdot \bar{\omega}_f = 0$, $h \cdot h = 4$ and $\omega_f \cdot \bar{\omega}_f > 0$, we check that $E \cap K = 0$. In particular, the form Q is positive definite on E , so there is a constant $C > 0$ such that $Q(\eta) \geq C \|\eta\|^2$ for

any $\eta \in E$. This constant is easily computable as the smallest eigenvalue of the matrix of the restriction of Q on that space, in a unitary basis, for the Hermitian norm $\| - \|$.

Now, let γ such that $\gamma \cdot h = 0$ and

$$|\gamma \cdot \bar{\omega}_f| \|\omega_f\| \leq \frac{1}{2} \|\gamma\| (\omega_f \cdot \bar{\omega}_f). \tag{17}$$

Let $a \doteq (\gamma \cdot \bar{\omega}_f) / (\omega_f \cdot \bar{\omega}_f)$ and $\eta \doteq \gamma - a\omega_f$, so that $\eta \cdot \bar{\omega}_f = 0$ and $\eta \cdot h = 0$, that is, $\eta \in E$. Since ω_f is in the kernel of Q , we have $Q(\eta) = Q(\gamma)$, and thus $Q(\gamma) \geq C\|\eta\|$. Lastly, we compute that

$$\|\eta\| \geq \|\gamma\| - |a| \|\omega_f\| = \|\gamma\| - \left| \frac{\gamma \cdot \bar{\omega}_f}{\omega_f \cdot \bar{\omega}_f} \right| \|\omega_f\| \geq \frac{1}{2} \|\gamma\|, \tag{18}$$

using (17). So $Q(\gamma) \geq \frac{1}{2} C \|\gamma\|$. □

The next statement is proved using the following result of [Smale 1986]. Let ϕ be an analytic function on a maximal open disc around 0 in \mathbb{C} with $\phi'(0) \neq 0$. We define

$$\gamma_{\text{Smale}}(\phi) \doteq \sup_{k \geq 2} \left| \frac{1}{k!} \frac{\phi^{(k)}(0)}{\phi'(0)} \right|^{1/(k-1)} \quad \text{and} \quad \beta_{\text{Smale}}(\phi) \doteq \left| \frac{\phi(0)}{\phi'(0)} \right|. \tag{19}$$

If $\beta_{\text{Smale}}(\phi) \gamma_{\text{Smale}}(\phi) \leq \frac{1}{34}$, then there is a $t \in \mathbb{C}$ such that $|t| \leq 2\beta_{\text{Smale}}(\phi)$ and $\phi(t) = 0$ [Smale 1986], see also [Blum et al. 1998, Chapter 8, Theorem 2].

Proposition 2. *For any $f \in \tilde{U}_4$, there exists C_f and $\varepsilon_f > 0$ such that for all $\varepsilon < \varepsilon_f$ the following holds: For any $\gamma \in H_{\mathbb{R}}$, if $\gamma \cdot h = 0$ and $|\gamma \cdot \omega_f| \leq \varepsilon \|\gamma\|$, then there is a monomial $m \in R_4$ and $t \in \mathbb{C}$ such that $|t| \leq C_f \varepsilon$ and $\gamma \cdot \omega_{f+tm} = 0$.*

Proof. Let $\gamma \in H_{\mathbb{R}}$ such that $\gamma \cdot h = 0$ and

$$|\gamma \cdot \omega_f| \leq \left(\frac{\omega_f \cdot \bar{\omega}_f}{2\|\omega_f\|} \right) \|\gamma\|. \tag{20}$$

Since γ has real coefficients, we have $|\gamma \cdot \omega_f| = |\gamma \cdot \bar{\omega}_f|$ and we may apply Lemma 1 to obtain a monomial m and a constant C such that

$$|\phi'_{\gamma,m}(0)| \geq C \|\gamma\|. \tag{21}$$

It follows, in particular, that

$$\beta_{\text{Smale}}(\phi_{\gamma,m}) \leq \frac{|\gamma \cdot \omega_f|}{C \|\gamma\|}. \tag{22}$$

Moreover, for any $k \geq 2$, and using $C \leq 1$,

$$\begin{aligned} \left| \frac{1}{k!} \frac{\phi_{\gamma,m}^{(k)}(0)}{\phi'_{\gamma,m}(0)} \right|^{1/(k-1)} &\leq C^{-1} \left| \frac{\phi_{\gamma,m}^{(k)}(0)}{k! \|\gamma\|} \right|^{1/(k-1)} \\ &= C^{-1} \left| \frac{\gamma}{\|\gamma\|} \cdot \frac{1}{k!} d_f^k \mathcal{P}(m, \dots, m) \right|^{1/(k-1)} \tag{23} \\ &\leq C^{-1} \left\| \frac{1}{k!} d_f^k \mathcal{P} \right\|^{1/(k-1)}, \tag{24} \end{aligned}$$

where $d_f^k \mathcal{P} : R_4^k \rightarrow H_{\mathbb{C}}$ is the k -th higher derivative of \mathcal{P} at f and where $\|\cdot\|$ is the operator norm defined as

$$\left\| \frac{1}{k!} d_f^k \mathcal{P} \right\| \doteq \sup_{\gamma \in H_{\mathbb{C}}} \sup_{h_1, \dots, h_k} \frac{|\gamma \cdot (1/k!) d_f^k \mathcal{P}(h_1, \dots, h_k)|}{\|\gamma\| \|h_1\| \cdots \|h_k\|}, \tag{25}$$

with supremum taken over $h_1, \dots, h_n \in \mathbb{C}[w, x, y, z]_4$. It follows that

$$\gamma_{\text{Smale}}(\phi_{\gamma, m}) \leq C^{-1} \sup_{k \geq 2} \left\| \frac{1}{k!} d_f^k \mathcal{P} \right\|^{1/(k-1)}. \tag{26}$$

Let Γ denote the supremum on the right-hand side of (26). By Smale’s theorem, together with (22) and (26), if $|\gamma \cdot \omega_f| \leq \frac{1}{34} C^2 \Gamma^{-1} \|\gamma\|$, then there is a $t \in \mathbb{C}$ such that $|t| \leq 2C^{-1} |\gamma \cdot \omega_f| \|\gamma\|^{-1}$ and $\gamma \cdot \mathcal{P}(f + tm) = 0$. The claim follows with $C_f \doteq 2C^{-1}$ and

$$\varepsilon_f \doteq \min \left(\frac{1}{34} C^2 \Gamma^{-1}, \frac{\omega_f \cdot \bar{\omega}_f}{2\|\omega_f\|} \right). \tag{27}$$

This concludes the proof. □

The constants C_f and ε_f are actually computable with simple algorithms. The constant from Lemma 1 is not hard to get with elementary linear algebra. It only remains to compute an upper bound for Γ . We address this issue in Section 2.4.

Corollary 3. *For any $f \in \tilde{U}_4$, any $\varepsilon < \varepsilon_f$ and any $\gamma \in H_{\mathbb{Z}}$, if $|\gamma \cdot \omega_f| \leq \frac{1}{4}\varepsilon$, then there exists a monomial $m \in R_4$ and $t \in \mathbb{C}$ such that $|t| \leq C_f \varepsilon$ and $\gamma \in \text{Pic}(X_{f+tm})$.*

Proof. We may assume that $\gamma \cdot \omega_f \neq 0$ (otherwise, choose any m and $t = 0$). Let $\gamma' = \gamma - \frac{1}{4}(\gamma \cdot h)h$. Since $h \cdot h = 4$, we have $\gamma' \cdot h = 0$. Moreover, we have $\gamma' \cdot \omega_f = \gamma \cdot \omega_f \neq 0$. In particular, $\gamma' \neq 0$, and since $\gamma' \in \frac{1}{4}H_{\mathbb{Z}}$, we have $\|\gamma'\| \geq \frac{1}{4}$. Then

$$|\gamma' \cdot \omega_f| \leq 4\|\gamma'\| |\gamma \cdot \omega_f| \leq \varepsilon \|\gamma'\|, \tag{28}$$

and Proposition 2 applies. □

2.4. Effective bounds for the higher derivatives of the period map. In the proof of Proposition 2, only the quantity Γ is not clearly computable. We show in this section how to compute an upper bound for Γ using the Griffiths–Dwork reduction. We follow here [Griffiths 1969a; Griffiths 1969b].

Firstly, as a variant of (8) avoiding dehomogenization, we write

$$\mathcal{P}(f) = \left(\frac{1}{2\pi i} \int_{T(\gamma_i)} \frac{\text{Vol}}{f} \right)_{1 \leq i \leq 22}, \tag{29}$$

where Vol is the projective volume form

$$\text{Vol} \doteq w \, dx \, dy \, dz - x \, dw \, dy \, dz + y \, dw \, dx \, dz - z \, dw \, dx \, dy. \tag{30}$$

For any $k > 0$ and $a \in R_{4k-4}$, we denote

$$\int \frac{a \text{Vol}}{f^k} \doteq \left(\frac{1}{2\pi i} \int_{T(\gamma_i)} \frac{a \text{Vol}}{f^k} \right)_{1 \leq i \leq 22} \in H_{\mathbb{C}}. \tag{31}$$

For any $h \in R_4$ close enough to 0, we have the power series expansion

$$\int \frac{\text{Vol}}{f+h} = \sum_{k \geq 1} (-1)^{k-1} \int \frac{h^{k-1} \text{Vol}}{f^k}. \tag{32}$$

Proposition 4. *For any $k \geq 3$, there is a linear map $G_k : R_{4k-4} \rightarrow R_8$ such that*

$$\int \frac{a}{f^k} \text{Vol} = \int \frac{G_k(a)}{f^3} \text{Vol}.$$

Moreover, there is a computable constant C , which depends only on f , such that for any $k \geq 3$, we have $\|G_k\| \leq C^{k-3}$, where R is endowed with the 1-norm (57).

Before we begin the proof of proposition, let us show that this is enough to bound Γ . Let

$$A : a \in R_8 \mapsto \int (a/f^3) \text{Vol} \in H_{\mathbb{C}},$$

then, using (32), we obtain

$$\int \frac{\text{Vol}}{f+h} = \sum_{k \geq 1} (-1)^{k-1} A(G_k(h^{k-1})), \tag{33}$$

and it follows that

$$\frac{1}{k!} d_f^k \mathcal{P}(h_1, \dots, h_k) = (-1)^k A(G_{k+1}(h_1 \cdots h_k)). \tag{34}$$

In particular,

$$\left\| \frac{1}{k!} d_f^k \mathcal{P}(h_1, \dots, h_k) \right\| \leq \|A\| \|G_{k+1}\| \|h_1 \cdots h_n\|_1 \tag{35}$$

$$\leq \|A\| \|G_{k+1}\| \|h_1\|_1 \cdots \|h_n\|_1, \tag{36}$$

and therefore, $\|(1/k!)d_f^k \mathcal{P}\| \leq \|A\| C^{k+1}$, from which we get

$$\Gamma \leq C \max(\|A\| C^2, 1). \tag{37}$$

Let us remark on how to bound the operator norm of A in practice. The period integrals can be approximated to arbitrary precision and with rigorous error bounds as in [Sertöz 2019]. This construction gives a small neighborhood of A in the matrix space. In practice, we represent this neighborhood as a matrix A' of complex balls and compute the operator norm of A' as usual but using complex ball arithmetic. This will return a real open interval containing $\|A\| \neq 0$. If the precision is high enough, 0 will not be contained in the closure of this interval, and we can take the lower bound of the interval.

2.4.1. Proof of Proposition 4. Let $R = \mathbb{C}[w, x, y, z]$. We define two families of maps for this proof. First, for $d \geq 12$, a multivariate division map $Q_d : R_d \rightarrow R_{d-3}^4$, such that for any $a \in R_d$,

$$a = \sum_{i=0}^3 Q_d(a)_i \partial_i f. \tag{38}$$

Note that such a map exists as soon as $d \geq 12$ by a theorem due to Macaulay, see [Lazard 1977, Corollaire, p. 169]. The choice of Q_d is not unique. We fix Q_{12} arbitrarily and define $Q_d(a)$, for $d > 12$ and $a \in R_{12}$,

as follows: Write $a = \sum_{i=0}^3 x_i a_i$, in such a way that the terms of the sum have disjoint monomial support, and define

$$Q_d(a) = \sum_{i=0}^3 x_i Q_{d-1}(a_i). \tag{39}$$

It is easy to check that this definition satisfies (38).

Second, for $k \geq 3$, we define $G_k: R_{4k-4} \rightarrow R_8$ as follows: Begin with $G_3 = \text{id}$, and then define G_k for $k \geq 4$ inductively. For $a \in R_{4k-4}$, we write $(b_0, \dots, b_3) = Q_{4k-4}(a)$ and define

$$G_k(a) \doteq G_{k-1} \left(\frac{1}{k-1} (\partial_0 b_0 + \dots + \partial_3 b_3) \right). \tag{40}$$

This map is the Griffiths–Dwork reduction, and it satisfies

$$\int_{\gamma} \frac{a\Omega}{f^k} = \int_{\gamma} \frac{G_k(a)\Omega}{f^3}. \tag{41}$$

Lemma 5. For any $d \geq 12$, we have $\| \| Q_d \| \| \leq \| \| Q_{12} \| \|$, where R is endowed with the 1-norm and R^4 with the norm $\|(f_0, \dots, f_3)\|_1 \doteq \|f_0\|_1 + \dots + \|f_3\|_1$.

Proof. For any $a \in R_d$,

$$\| \| Q_d(a) \| \|_1 = \sum_{i=0}^3 \| \| Q_d(a)_i \| \|_1 \leq \sum_{i=0}^3 \sum_{j=0}^3 \| \| x_j Q_{d-1}(a_j)_i \| \|_1 \tag{42}$$

$$= \sum_{i=0}^3 \sum_{j=0}^3 \| \| Q_{d-1}(a_j)_i \| \|_1 = \sum_j \| \| Q_{d-1}(a_j) \| \|_1 \tag{43}$$

$$\leq \| \| Q_{d-1} \| \| \sum_j \| \| a_j \| \|_1 = \| \| Q_{d-1} \| \| \| \| a \| \|_1, \tag{44}$$

using, for the last equality, that the terms a_j have disjoint monomial support. □

Lemma 6. For any $k \geq 3$, we have $\| \| G_k \| \| \leq (4 \| \| Q_{12} \| \|)^{k-3}$, where R is endowed with the 1-norm.

Proof. We proceed by induction on k (the base case $k = 3$ is trivial since $G_3 = \text{id}$). Let $a \in R_{4k-4}$ and $(b_0, \dots, b_3) = Q_{4k-4}(a)$. By (40), we have

$$\| \| G_k(a) \| \|_1 \leq \frac{\| \| G_{k-1} \| \|}{k-1} (\| \| \partial_0 b_0 \| \|_1 + \dots + \| \| \partial_3 b_3 \| \|_1). \tag{45}$$

By the induction hypothesis, $\| \| G_{k-1} \| \| \leq (4 \| \| Q_{12} \| \|)^{k-4}$, and moreover, since each b_i has degree $4k - 7$, we have $\| \| \partial_i b_i \| \|_1 \leq (4k - 7) \| \| b_i \| \|_1$. It follows that

$$\| \| G_k(a) \| \|_1 \leq (4 \| \| Q_{12} \| \|)^{k-4} \frac{4k-7}{k-1} (\| \| b_0 \| \|_1 + \dots + \| \| b_3 \| \|_1). \tag{46}$$

Next, we note that $\| \| b_0 \| \|_1 + \dots + \| \| b_3 \| \|_1 = \| \| Q_{4k-4}(a) \| \|_1$ and, by Lemma 5, we have $\| \| Q_{4k-4}(a) \| \| \leq \| \| Q_{12} \| \|$. Therefore,

$$\| \| G_k(a) \| \|_1 \leq (4 \| \| Q_{12} \| \|)^{k-3} \| \| a \| \|_1, \tag{47}$$

and the claim follows. □

3. The Noether–Lefschetz locus

3.1. Basic properties. We define the Noether–Lefschetz locus for quartic surfaces and review a few classical properties, especially algebraicity, with a view towards Theorem 14 about the degree and the height of the equations defining the components of the Noether–Lefschetz locus.

3.1.1. Definition. The Noether–Lefschetz locus of quartics \mathcal{NL} is the set of all $f \in U_4$ such that the rank of $\text{Pic}(X_f)$ is at least 2. Equivalently, in view of (12), \mathcal{NL} is the set of quartic polynomials f whose primitive periods (1) are \mathbb{Z} -linearly dependent.

The set \mathcal{NL} is locally the union of smooth analytic hypersurfaces in U_4 . To see this, let $\widetilde{\mathcal{NL}}$ be the lift of \mathcal{NL} in the universal covering \widetilde{U}_4 of U_4 . Recall that $\mathcal{P}: \widetilde{U}_4 \rightarrow \mathcal{D}$ is the period map. The Lefschetz (1,1)-theorem implies

$$\widetilde{\mathcal{NL}} = \bigcup_{\gamma \in H_{\mathbb{Z}} \setminus \mathbb{Z}h} \mathcal{P}^{-1}\{w \in \mathcal{D} \mid w \cdot \gamma = 0\}. \tag{48}$$

That is, $\widetilde{\mathcal{NL}}$ is the pullback of smooth hyperplane sections of \mathcal{D} . Since \mathcal{P} is a submersion, $\widetilde{\mathcal{NL}}$ is the union of smooth analytic hypersurfaces. It follows that \mathcal{NL} is locally the union of smooth analytic hypersurfaces.

We break \mathcal{NL} into algebraic pieces as follows: For any integers d and g , let $\mathcal{NL}_{d,g}$ be the set

$$\mathcal{NL}_{d,g} = \{f \in U_4 \mid \exists \gamma \in \text{Pic}(X_f) \setminus \mathbb{Z}h : \gamma \cdot h = d \text{ and } \gamma \cdot \gamma = 2g - 2\}. \tag{49}$$

By replacing γ by $\gamma + h$ or $-\gamma$, we observe that

$$\mathcal{NL}_{d,g} = \mathcal{NL}_{d+4,g+d+2} = \mathcal{NL}_{-d,g}. \tag{50}$$

In particular, $\mathcal{NL}_{d,g}$ is equal to some $\mathcal{NL}_{d',g'}$ with $d' > 0$ and $g' \geq 0$, so that

$$\mathcal{NL} = \bigcup_{d>0} \bigcup_{g \geq 0} \mathcal{NL}_{d,g}. \tag{51}$$

For $\gamma \in H_{\mathbb{Z}}$, let $\Delta(\gamma) = (h \cdot \gamma)^2 - 4\gamma \cdot \gamma$. It is the negative of the discriminant of the lattice generated by h and γ in $H_{\mathbb{Z}}$, with respect to the intersection product (and it is zero if $\gamma \in \mathbb{Z}h$). It follows from the Hodge index theorem, see [Hartshorne 1977, Theorem V.1.9] that for any $f \in U_4$ and any $\gamma \in \text{Pic}(X_f)$, where $\Delta(\gamma) \geq 0$, with equality if and only if $\gamma \in \mathbb{Z}h$. If $\gamma \cdot h = d$ and $\gamma \cdot \gamma = 2g - 2$, then $\Delta(\gamma) = d^2 - 8g + 8$. We obtain, therefore, that for any $d > 0$ and $g \geq 0$,

$$\mathcal{NL}_{d,g} = \begin{cases} \{f \in U_4 \mid \exists \gamma \in \text{Pic}(X_f) : \gamma \cdot h = d, \\ \gamma \cdot \gamma = 2g - 2\}, & \text{if } d^2 > 8g - 8, \\ \emptyset, & \text{otherwise.} \end{cases} \tag{52}$$

It is, in fact, more natural to introduce, for $\Delta > 0$, the locus

$$\mathcal{NL}_{\Delta} \doteq \{f \in U_4 \mid \exists \gamma \in \text{Pic}(X_f) : \Delta(\gamma) = \Delta\} \tag{53}$$

$$= \bigcup_{\substack{d>0, \\ d^2 \equiv \Delta \pmod{8}}} \mathcal{NL}_{d,(d^2-\Delta)/8+1}. \tag{54}$$

Due to (50), \mathcal{NL}_Δ reduces to a single $\mathcal{NL}_{d,g}$. Namely,

$$\mathcal{NL}_\Delta = \begin{cases} \mathcal{NL}_{4t, 2t^2+(8-\Delta)/8}, & \text{if } \Delta \equiv 0 \pmod{8}, \\ \mathcal{NL}_{4t+1, 2t^2+t+(9-\Delta)/8}, & \text{if } \Delta \equiv 1 \pmod{8}, \\ \mathcal{NL}_{4t+2, 2t^2+2t+(12-\Delta)/8}, & \text{if } \Delta \equiv 4 \pmod{8}, \\ \emptyset, & \text{otherwise,} \end{cases} \tag{55}$$

where $t = \lceil \frac{1}{4}\sqrt{\Delta} \rceil$. Conversely, each $\mathcal{NL}_{d,g} = \mathcal{NL}_{d^2-8g+8}$.

3.1.2. Algebraicity. For any $d > 0$ and $g \geq 0$, the set $\mathcal{NL}_{d,g}$ is either empty or an algebraic hypersurface in U_4 . This is a classical result, e.g., [Voisin 2003, Theorem 3.32], which we recall here to obtain an explicit algebraic description of $\mathcal{NL}_{d,g}$.

Lemma 7. *For any $f \in U_4$, $d > 0$ and $g \geq 0$, we have: $f \in \mathcal{NL}_{d,g}$ if and only if X_f contains an effective divisor with Hilbert polynomial $t \mapsto dt + 1 - g$.*

Proof. Assume that X_f contains an effective divisor C with Hilbert polynomial $t \mapsto td + 1 - g$. Since X_f is smooth, C is a locally principal divisor and gives an element γ of $\text{Pic } X_f$. The integer d is the degree of C , so it is the number of points in the intersection with a generic hyperplane, that is, $d = \gamma \cdot h$. Moreover, g is the arithmetic genus of C , which is determined by $2g - 2 = \gamma \cdot \gamma$ [Hartshorne 1977, Exercises III.5.3(b) and V.1.3(a)]. So, $f \in \mathcal{NL}_{d,g}$.

Conversely, let $f \in \mathcal{NL}_{d,g}$. By definition, there is a divisor C on X_f such that its class γ in $\text{Pic } X_f$ satisfies $\gamma \cdot h = d$ and $\gamma \cdot \gamma = 2g - 2$. From the Riemann–Roch theorem for surfaces [Hartshorne 1977, Theorem V.1.6], we get

$$\dim H^0(X, \mathcal{O}_X(C)) + \dim H^0(X, \mathcal{O}_X(-C)) \geq \frac{1}{2}\gamma \cdot \gamma + 2 = g + 1 > 0,$$

so that either C or $-C$ must be linearly equivalent to an effective divisor. Since $\gamma \cdot h > 0$, it follows that $-C$ cannot be effective, and therefore, C must be. As above, the Hilbert polynomial of C is given by $t \mapsto dt + 1 - g$. □

In light of Lemma 7, the algebraicity of $\mathcal{NL}_{d,g}$ is proved by using the Hilbert scheme $\mathcal{H}_{d,g}$. The Hilbert scheme $\mathcal{H}_{d,g}$ of degree d and genus g curves in \mathbb{P}^3 is a projective scheme that parametrizes all the subschemes of \mathbb{P}^3 whose Hilbert polynomial is $t \mapsto dt + 1 - g$.

The Hilbert scheme $\mathcal{H}_{d,g}$ may contain components that are not desirable for our purposes. For example, $\mathcal{H}_{3,0}$, which contains twisted cubics in \mathbb{P}^3 , contains two irreducible components [Piene and Schlessinger 1985]: a 12-dimensional component that is the closure of the space of all smooth cubic rational curves in \mathbb{P}^3 and a 15-dimensional component parametrizing the union of a plane cubic curve with a point in \mathbb{P}^3 . We would be only interested in the first, not in the second component. So we introduce $\mathcal{H}'_{d,g}$, the union of components of $\mathcal{H}_{d,g}$ obtained by removing the components that do not correspond to locally complete-intersection pure-dimensional subschemes of \mathbb{P}^3 .

When $d^2 > 8g - 8$, Lemma 7 can be rephrased as

$$\mathcal{NL}_{d,g} = \text{proj}_1 \{(f, C) \in U_4 \times \mathcal{H}'_{d,g} \mid C \subset X_f\}, \tag{56}$$

where proj_1 denotes the projection $U_4 \times \mathcal{H}'_{d,g} \rightarrow U_4$. Since $\mathcal{H}'_{d,g}$ is a projective variety, and the condition $C \subset X_f$ is algebraic, this shows that $\mathcal{NL}_{d,g}$ is a closed subvariety of U_4 (for more details about this construction, see [Voisin 2003, §3.3]).

We note, furthermore, that $\mathcal{NL}_{d,g}$ is clearly invariant under the action of the Galois group of algebraic numbers. Therefore, it can be defined over the rational numbers.

As a consequence, for any nonnegative integers d and g , there is a squarefree primitive homogeneous polynomial $\text{NL}_{d,g} \in \mathbb{Z}[u_1, \dots, u_{35}]$ in the 35 coefficients of the general quartic polynomial that is unique up to sign and whose zero locus is $\mathcal{NL}_{d,g}$ in U_4 . Similarly, we define NL_Δ up to sign as the unique squarefree primitive polynomial vanishing exactly on \mathcal{NL}_Δ .

3.2. Height of multiprojective varieties. The mainstay of our results is a bound on the degree and size of the coefficients of the polynomials $\text{NL}_{d,g}$. The determination of these bounds is based on (56) and involves the theory of heights of multiprojective varieties as developed by D’Andrea et al. [2013], and, before them, [Bost et al. 1991; Philippon 1995; Krick et al. 2001; Rémond 2001a; 2001b], among others. We recall here the results that we need, following [D’Andrea et al. 2013].

3.2.1. Heights of polynomials. Let $f = \sum_\alpha c_\alpha \mathbf{x}^\alpha \in \mathbb{C}[x_1, \dots, x_n]$. We recall the following different measures of height of f :

$$\|f\|_1 \doteq \sum_\alpha |c_\alpha|, \tag{57}$$

$$\|f\|_{\text{sup}} \doteq \sup_{|x_1|=\dots=|x_n|=1} |f(\mathbf{x})|, \tag{58}$$

$$m(f) \doteq \int_{[0,1]^n} \log |f(e^{2\pi i t_1}, \dots, e^{2\pi i t_n})| dt_1 \dots dt_n. \tag{59}$$

Lemma 8 [D’Andrea et al. 2013, Lemma 2.30]. *For any homogeneous polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$,*

$$\exp(m(f)) \leq \|f\|_{\text{sup}} \leq \|f\|_1 \leq \exp(m(f))(n + 1)^{\deg f}.$$

3.2.2. Extended Chow ring. The extended Chow ring [D’Andrea et al. 2013, Definition 2.50] is a tool to track a measure of height of multiprojective varieties when performing intersections and projections. We present here a very brief summary. Bold letters refer to multiindices, and all varieties are considered over \mathbb{Q} . Let $\mathbf{n} \in \mathbb{N}^r$, and let $\mathbb{P}^{\mathbf{n}}$ be the multiprojective space $\mathbb{P}^{\mathbf{n}} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_m}$.

An *algebraic cycle* is a finite \mathbb{Z} -linear combination $\sum_V n_V V$ of irreducible subvarieties of $\mathbb{P}^{\mathbf{n}}$. The *irreducible components* of an algebraic cycle, as above, are the irreducible varieties V such that $n_V \neq 0$. An algebraic cycle is *equidimensional* if all its irreducible components have the same dimension. An algebraic cycle is *effective* if $n_V \geq 0$ for all V . The *support* of X , denoted by $\text{supp } X$, is the union of the irreducible components of X .

Let $A^*(\mathbb{P}^{\mathbf{n}}; \mathbb{Z})$ be the extended Chow ring, namely

$$A^*(\mathbb{P}^{\mathbf{n}}; \mathbb{Z}) \doteq \mathbb{R}[\eta, \theta_1, \dots, \theta_m] / (\eta^2, \theta_1^{n_1+1}, \dots, \theta_m^{n_m+1}), \tag{60}$$

where θ_i is the class of the pullback of a hyperplane from \mathbb{P}^{n_i} and η is used to keep track of heights of varieties. For two elements a and b of this ring, we write $a \leq b$ when the coefficients of $b - a$ in the monomial basis are nonnegative.

To an algebraic cycle X of \mathbb{P}^n , we associate an element $[X]_{\mathbb{Z}}$ of $A^*(\mathbb{P}^n; \mathbb{Z})$ [D’Andrea et al. 2013, Definition 2.50]. If X is effective, then $[X]_{\mathbb{Z}} \geq 0$. The coefficients of the terms in $[X]_{\mathbb{Z}}$ for monomials not involving η record the usual multidegrees of X . The terms involving η record mixed canonical heights of X . The definition of these heights is based on the heights of various Chow forms associated to X [D’Andrea et al. 2013, §2.3]. For the computations in this paper, we only need the following results:

Let $f \in \mathbb{Z}[\mathbf{x}_\Delta, \dots, \mathbf{x}_r]$ be a nonzero multihomogeneous polynomial with respect to the group of variables $\mathbf{x}_1, \dots, \mathbf{x}_n$. We assume that f is *primitive*, that is, the gcd of the coefficients of f is 1. The element associated in $A^*(\mathbb{P}^n; \mathbb{Z})$ to the hypersurface $V(f) \subset \mathbb{P}^n$ is [D’Andrea et al. 2013, Proposition 2.53 (2)]

$$[V(f)]_{\mathbb{Z}} = m(f)\eta + \deg_{\mathbf{x}_1}(f)\theta_1 + \dots + \deg_{\mathbf{x}_r}(f)\theta_r. \tag{61}$$

To such a polynomial f , we also associate [D’Andrea et al. 2013, Equation (2.57)]

$$[f]_{\text{sup}} \doteq \log(\|f\|_{\text{sup}})\eta + \deg_{\mathbf{x}_1}(f)\theta_1 + \dots + \deg_{\mathbf{x}_r}(f)\theta_r. \tag{62}$$

3.2.3. Arithmetic Bézout theorem. Let X be an effective cycle and H a hypersurface in \mathbb{P}^n . They *intersect properly* if no irreducible component of X is in H . When X and H intersect properly, one defines an intersection product $X \cdot H$, that is an effective cycle supported on $X \cap H$. If X is equidimensional of dimension r , then $X \cdot H$ is equidimensional of dimension $r - 1$.

The following statement is an arithmetic Bézout bound that not only bounds the degree, as with the classical Bézout bound, but also the height of an intersection:

Theorem 9 [D’Andrea et al. 2013, Theorem 2.58]. *Let X be an effective equidimensional cycle on \mathbb{P}^n and $f \in \mathbb{Z}[\mathbf{x}_1, \dots, \mathbf{x}_m]$. If X and $V(f)$ intersect properly, then $[X \cdot V(f)]_{\mathbb{Z}} \leq [X]_{\mathbb{Z}} \cdot [f]_{\text{sup}}$.*

This theorem can be applied (as in [D’Andrea et al. 2013, Corollary 2.61]) to bound the height of the irreducible components of a variety in terms of its defining equations.

Proposition 10. *Let $Z \subset \mathbb{P}^n$ be an equidimensional variety, and let X be $V(f_1, \dots, f_s) \cap Z$, where f_i is a multihomogeneous polynomial of multidegree at most \mathbf{d} and sup-norm at most L . Let X_r be the union of all the irreducible components of X of codimension r in Z . Then*

$$[X_r]_{\mathbb{Z}} \leq [Z]_{\mathbb{Z}} \left(\log(sL)\eta + \sum_{i=1}^m d_i \theta_i \right)^r.$$

Proof. Let (y_{ij}) be a new group of variables, with $1 \leq i \leq r$ and $1 \leq j \leq s$. Let $g_i \doteq \sum_{j=1}^s y_{ij} f_j$ and $X' \doteq V(g_1, \dots, g_r)$ in $\mathbb{P}^k \times Z$, with $k = rs - 1$. We first claim that $\mathbb{P}^k \times X_r$ is a union of components of X' . Indeed, let ξ_0 be the generic point of \mathbb{P}^k and ξ_1 be the generic point of a component Y of X_r , so that $\xi = (\xi_0, \xi_1)$ is the generic point of the component $\mathbb{P}^k \times Y$ of $\mathbb{P}^k \times X_r$. Since X has codimension r at ξ_1 , the generic linear combinations g_1, \dots, g_r form a regular sequence at ξ (in other words, they

form a regular sequence at ξ_1 for generic values of the v_{ij}). Therefore, X' has codimension r at ξ . Since $\mathbb{P}^k \times Y \subseteq X'$, it follows that $\mathbb{P}^k \times Y$ is a component of X' .

Let X'_r be the union of the components of codimension r of X' . The argument above shows that $[\mathbb{P}^k \times X_r]_{\mathbb{Z}} \leq [X'_r]_{\mathbb{Z}}$. Besides, by repeated application of [D'Andrea et al. 2013, Corollary 2.61],

$$[X'_r]_{\mathbb{Z}} \leq [\mathbb{P}^k \times Z]_{\mathbb{Z}} \prod_{i=1}^r [g_i]_{\text{sup}}, \tag{63}$$

where θ_0 is the variable attached to \mathbb{P}^k in the extended Chow ring of $\mathbb{P}^k \times \mathbb{P}^n$. We compute, using (61), that

$$[g_i]_{\text{sup}} \leq \log(sL)\eta + \theta_0 + \sum_{i=1}^s d_i \theta_i. \tag{64}$$

Finally, we note that $[\mathbb{P}^k \times X_r]_{\mathbb{Z}} = [X_r]_{\mathbb{Z}}$ and $[\mathbb{P}^k \times Z]_{\mathbb{Z}} = [Z]_{\mathbb{Z}}$ by [D'Andrea et al. 2013, Propositions 2.51.3 and 2.66]. □

Proposition 11. *Let X be an equidimensional closed subvariety of $\mathbb{P}^k \times \mathbb{P}^n$, and let $Y \subset \mathbb{P}^n$ be the projection of X . If Y is equidimensional, then*

$$\theta_0^k [Y]_{\mathbb{Z}} \leq \theta_0^{\dim X - \dim Y} [X]_{\mathbb{Z}} \in A^*(\mathbb{P}^k \times \mathbb{P}^n; \mathbb{Z}),$$

where θ_0 is the variable attached to \mathbb{P}^k in the extended Chow ring of $\mathbb{P}^k \times \mathbb{P}^n$.

Proof. We will argue by induction on $r \doteq \dim X - \dim Y$. When $r = 0$, this is [D'Andrea et al. 2013, Proposition 2.64].

Suppose now that $r > 0$ and X is irreducible. Let $\mathbb{Q}[y, \mathbf{x}_1, \dots, \mathbf{x}_m]$ denote the multihomogeneous coordinate ring of $\mathbb{P}^k \times \mathbb{P}^n$. There is an i , with $0 \leq i \leq k$, such that $H \doteq V(y_i) \subset \mathbb{P}^k \times \mathbb{P}^n$ intersects X properly (otherwise, X would be included in all $V(y_i)$ and would be empty). Since the fibers of $X \rightarrow Y$ are positive dimensional, H intersects each fiber. In particular, the set-theoretical projections of X and $X \cap H$ coincide. As X is irreducible, so is Y . In particular, there is an irreducible component $X' \subset X \cap H$ that maps to Y . By the induction hypothesis applied to X' , we have $\theta_0^k [Y]_{\mathbb{Z}} \leq \theta_0^{\dim X' - \dim Y} [X']_{\mathbb{Z}}$. Moreover, $[X']_{\mathbb{Z}} \leq [X]_{\mathbb{Z}} [y_i]_{\text{sup}}$, and, in view of (62), $[y_i]_{\text{sup}} = \theta_0$. The claim follows.

If X is reducible, then we apply the inequality above to each of the irreducible components of Y together with an irreducible component of X mapping onto that component. □

3.3. Explicit equations for the Noether–Lefschetz loci. Following Gotzmann [1978], Bayer [1982] and the exposition of Lella [2012], we describe the equations defining the Hilbert schemes of curves in \mathbb{P}^3 . An explicit description of the Noether–Lefschetz loci $\mathcal{NL}_{d,g}$ follows.

3.3.1. Hilbert schemes of curves. For $d > 0$ and $g \geq 0$, let $\mathcal{H}_{d,g}$ be the Hilbert scheme of curves of degree d and genus g in \mathbb{P}^3 . It parametrizes subschemes of \mathbb{P}^3 with Hilbert polynomial $p(m) \doteq dm + 1 - g$. Smooth curves in \mathbb{P}^3 of degree d and genus g , in particular, have Hilbert polynomial $p(m)$. Let $R = \mathbb{C}[w, x, y, z]$ be the homogeneous coordinate ring of \mathbb{P}^3 . For $m \geq 0$, let R_m denote the m -th homogeneous part of R , and let $q(m) = \dim R_m - p(m)$.

The Hilbert scheme $\mathcal{H}_{d,g}$ can be realized in a Grassmannian variety as follows: A subscheme X of \mathbb{P}^3 is uniquely defined by a saturated homogeneous ideal I of R . If the Hilbert polynomial of X is p , then I is the saturation of the ideal generated by the degree r slice $I_r \doteq I \cap R_r$ [Gotzmann 1978] and [Bayer 1982, §II.10], where

$$r = \binom{d}{2} + 1 - g \tag{65}$$

is the Gotzmann number of p [Bayer 1982, §II.1.17]. For practical reasons, we need $r \geq 4$, so we define instead

$$r = \max\left(\binom{d}{2} + 1 - g, 4\right). \tag{66}$$

So X is entirely determined by I_r , which is a $q(r)$ -dimensional subspace of R_r .

Let \mathbb{G} be the Grassmannian variety of $q(r)$ -dimensional subspaces of R_r . As a set, one can construct $\mathcal{H}_{d,g}$ as the subset of all $\Xi \in \mathbb{G}$ such that the ideal generated by Ξ in R defines a subscheme of \mathbb{P}^3 with Hilbert polynomial p . In fact, $\mathcal{H}_{d,g}$ is a subvariety that is defined by the following condition [Bayer 1982, §VI.1]:

$$\mathcal{H}_{d,g} = \{ \Xi \in \mathbb{G} \mid \dim(R_1 \Xi) \leq q(r+1) \}, \tag{67}$$

where R_1 is the space of linear forms in w, x, y, z , so that $R_1 \Xi$ is a subspace of R_{r+1} .

Several authors gave explicit equations for $\mathcal{H}_{d,g}$ in the Plücker coordinates [Bayer 1982; Grothendieck 1966; Gotzmann 1978; Brachat et al. 2016]. We will prefer here a more direct path that avoids the Plücker embedding.

3.4. Equations for the relative Hilbert scheme. Define the relative Hilbert scheme of curves inside quartic surfaces

$$\mathcal{H}_{d,g}(4) \doteq \{ (f, C) \in \mathbb{P}(R_4) \times \mathcal{H}_{d,g} \mid C \subset V(f) \}, \tag{68}$$

for each $d > 0$ and $g \geq 0$.

We define the following auxiliary spaces to better describe (68): First, define the ambient space

$$\mathcal{A} \doteq \mathbb{P}(R_4) \times \mathbb{P}(\text{End}(\mathbb{C}^{q(r)-N_{r-4}}, R_r)) \times \mathbb{P}(\text{End}(R_{r+1}, \mathbb{C}^{p(r+1)})). \tag{69}$$

Second, let $\mathcal{B} = \{ (f, \phi, \psi) \in \mathcal{A} \}$ be the set of all triples satisfying the conditions

- (i) $R_{r-3}f \subseteq \ker \psi$,
- (ii) $R_1 \text{im}(\phi) \subseteq \ker \psi$,
- (iii) $\text{im} \phi \cap R_{r-4}f = 0$,
- (iv) ϕ and ψ are full rank.

Finally, we denote by $\bar{\mathcal{B}}$ the Zariski closure of \mathcal{B} .

Lemma 12. *The map $\mathcal{B} \rightarrow \mathcal{H}_{d,g}(4)$ defined by $(f, \phi, \psi) \mapsto (f, R_{r-4}f + \text{im} \phi)$ is well defined and surjective.*

Proof. Let $(f, \phi, \psi) \in \mathcal{B}$, and let $\Xi = R_{r-4}f + \text{im } \phi$. Constraint (iv) implies that $\text{im } \phi$ has dimension $q(r) - N_{r-4}$. Together with Constraint (iii), we have $\dim \Xi = q(r)$. Moreover, Constraint (iv) implies that $\ker \psi$ has dimension $q(r + 1)$. In particular, since $R_1 \Xi = R_{r-3}f + R_1 \text{im } \phi$, Constraints (i) and (ii) imply that $R_1 \Xi$ has dimension at most $q(r + 1)$. So, $\Xi \in \mathcal{H}_{d,g}(4)$. Since $R_{r-4}f \subseteq \Xi$, the polynomial f is in the saturation of the ideal generated by Ξ . Hence, $(f, \Xi) \in \mathcal{H}_{d,g}(4)$.

Conversely, let $(f, \Xi) \in \mathcal{H}_{d,g}(4)$, then $R_{r-4}f \subset \Xi$ and there is a full rank map $\phi: \mathbb{C}^{q(r)-N_{r-4}} \rightarrow R_r$ such that $\text{im } \phi$ complements $R_{r-4}f$ in Ξ . Furthermore, $\dim R_1 \Xi \leq q(r + 1)$, because $\Xi \in \mathcal{H}_{d,g}$, so there is a full rank map $\psi: R_{r+1} \rightarrow \mathbb{C}^{p(r+1)}$ such that $R_1 \Xi \subseteq \ker \psi$. So, (f, Ξ) is the image of $(f, \phi, \psi) \in \mathcal{B}$. \square

Lemma 13. *For any $a \geq 0$, let $\bar{\mathcal{B}}_a$ be the union of the codimension a components of $\bar{\mathcal{B}}$. Then*

$$[\bar{\mathcal{B}}_a]_{\mathbb{Z}} \leq (15 \log(d + 2)\eta + \theta_1 + \theta_2 + \theta_3)^a$$

Proof. Let \mathcal{B}' be the closed set defined by Constraints (i) and (ii). Constraints (iii) and (iv) are open, so any component of $\bar{\mathcal{B}}$ is a component of \mathcal{B}' . In particular, $[\bar{\mathcal{B}}_a]_{\mathbb{Z}} \leq [\mathcal{B}'_a]_{\mathbb{Z}}$.

Constraint (i) is expressed with $p(r + 1)N_{r-3}$ polynomial equations of multidegree $(1, 0, 1)$ (with respect to f, ϕ and ψ , respectively). Namely, $\psi(mf) = 0$ for every monomial m in R_{r-3} . Each $p(r + 1)$ components of the equation $\psi(mf) = 0$ involves a sum of 35 terms (since f , as a quartic polynomial, contains only 35 terms) with coefficients 1. So the 1-norm of these constraints is at most 35 (which is also at most N_r , since $r \geq 4$).

Constraint (ii) is expressed with $4p(r + 1)(q(r) - N_{r-4})$ polynomial equations of multidegree $(0, 1, 1)$. Namely, $\psi(v\phi(e)) = 0$ for any basis vector e and any variable $v \in \{w, x, y, z\}$. Each $p(r + 1)$ component of the equation $\psi(v\phi(e)) = 0$ involves a sum of N_r terms with coefficients 1. So the 1-norm of these constraints is at most N_r .

The claim is then a consequence of Proposition 10, with

$$s = p(r + 1)N_{r-3} + 4p(r + 1)(q(r) - N_{r-4}) \quad \text{and} \quad L = N_r.$$

We check routinely, with Mathematica, that $sL \leq (d + 2)^{15}$. \square

Theorem 14. *There is an absolute constant $A > 0$ such that for any $d > 0$ and $g \geq 0$, we have*

$$\deg(\text{NL}_{d,g}) \leq A^{d^9} \quad \text{and} \quad \|\text{NL}_{d,g}\|_1 \leq 2^{A^{d^9}}.$$

Proof. We assume $\mathcal{NL}_{d,g}$ is nonempty, since these inequalities are trivially satisfied if $\mathcal{NL}_{d,g} = \emptyset$ with $\text{NL}_{d,g} = 1$. Let $P_2 \doteq \mathbb{P}(\text{End}(\mathbb{C}^{q(r)-N_{r-4}}, R_r))$ and $P_3 \doteq \mathbb{P}(\text{End}(R_{r+1}, \mathbb{C}^{p(r+1)}))$ denote the second and third factors of \mathcal{A} , respectively. Let $\alpha \doteq (q(r) - N_{r-4})N_r - 1$ and $\beta \doteq p(r + 1)N_{r+1} - 1$ denote the dimensions of P_2 and P_3 , respectively. Let \mathcal{E} be the projection of $\bar{\mathcal{B}}$ on $\mathbb{P}(R_4) \times P_2$. The fibers of the map $\bar{\mathcal{B}} \rightarrow \mathcal{E}$ are projective subspaces of P_3 since Constraints (i) and (ii) are linear in ψ . The dimension of these fibers are $\beta' \doteq p(r + 1)^2 - 1$. So, by Proposition 11,

$$\theta_3^\beta[\mathcal{E}]_{\mathbb{Z}} \leq \theta_3^{\beta'}[\bar{\mathcal{B}}]_{\mathbb{Z}}. \tag{70}$$

Next, the map $\mathcal{B} \rightarrow \mathcal{H}_{d,g}(4)$ factors through \mathcal{E} , and the fibers of the corresponding map $\mathcal{E} \rightarrow \mathcal{H}_{d,g}(4)$ have dimension $\alpha' \doteq (q(r) - N_{r-4})q(r) - 1$. Finally, let e be the dimension of the fibers of the map $\mathcal{H}_{d,g}(4) \rightarrow \mathcal{NL}_{d,g}$. (If this dimension is not generically constant, we work one component at a time.) Once again, by Proposition 11, we obtain

$$\theta_2^\alpha [\mathcal{NL}_{d,g}]_{\mathbb{Z}} \leq \theta_2^{\alpha'+e} [\mathcal{E}]_{\mathbb{Z}}. \tag{71}$$

Since $[\mathcal{NL}_{d,g}]_{\mathbb{Z}} = m(\text{NL}_{d,g})\eta + \text{deg}(\text{NL}_{d,g})\theta_1$, taking $L = 15 \log(d + 2)$, we get

$$\text{deg NL}_{d,g} \leq \text{coeff. of } \theta_1 \theta_2^{\alpha-\alpha'-e} \theta_3^{\beta-\beta'} \text{ in } (L\eta + \theta_1 + \theta_2 + \theta_3)^{\alpha+\beta-\alpha'-\beta'-e+1} \tag{72}$$

$$\leq 3^{\alpha+\beta-\alpha'-\beta'-e+1}. \tag{73}$$

The exponent is a polynomial in d and g . Unless $d^2 \geq 8g - 8$, we have that $\mathcal{NL}_{d,g}$ is empty. So, we may bound the exponent with a polynomial only in d , which turns out to be of degree 9. Therefore, $\text{deg NL}_{d,g} \leq A^{d^9}$ for some constant $A > 0$.

Similarly,

$$m(\text{NL}_{d,g}) \leq \text{coeff. of } \eta \theta_2^{\alpha-\alpha'-e} \theta_3^{\beta-\beta'} \text{ in } (L\eta + \theta_1 + \theta_2 + \theta_3)^{\alpha+\beta-\alpha'-\beta'-e+1} \tag{74}$$

$$\leq (\alpha + \beta - \alpha' - \beta' - e + 1)L3^{\alpha+\beta-\alpha'-\beta'-e} \tag{75}$$

$$\leq 2^{O(d^9)}. \tag{76}$$

By [D'Andrea et al. 2013, Lemma 2.30.3],

$$\|\text{NL}_{d,g}\|_1 \leq \exp(m(\text{NL}_{d,g}))36^{\text{deg NL}_{d,g}}, \tag{77}$$

and this implies the claim, for some other constant $A > 0$. □

For the following, we write $a \uparrow b$ for a^b . This is a right-associative operation.

Corollary 15. *There is an absolute constant $A > 0$ such that for any $\Delta > 0$,*

$$\text{deg}(\text{NL}_{\Delta}) \leq A \uparrow \Delta \uparrow \frac{9}{2} \quad \text{and} \quad \|\text{NL}_{\Delta}\|_1 \leq 2 \uparrow A \uparrow \Delta \uparrow \frac{9}{2}.$$

In fact, one can obtain the following explicit bounds:

$$\text{deg}(\text{NL}_{\Delta}) \leq 3^{(\Delta+20)^{9/2}} \quad \text{and} \quad \log_2 \|\text{NL}_{\Delta}\|_1 \leq (\Delta + 60)^5 3^{(\Delta+20)^{9/2}}.$$

Proof. The first statement follows directly from (55) and Theorem 14 using a different A . The second statement is found by carrying out the arguments in the proof of Theorem 14 with the help of a computer algebra system. □

3.5. How good are these bounds? We can compare our degree bounds for NL_{Δ} to the exact degrees computed by Maulik and Pandharipande [2013], from which it actually follows that

$$\text{deg NL}_{\Delta} = O(\Delta^{19/2}). \tag{78}$$

This sharper bound does not directly imply a sharper bound on the height of NL_Δ , but it suggests the following conjecture. This would improve subsequently Theorems 17 and 19. In particular, (2) would be exponential in the size of the coefficients, as opposed to being doubly exponential.

Conjecture 16. *As Δ goes to ∞ , we have*

$$\log \|NL_\Delta\|_1 \leq \Delta^{19/2+o(1)}.$$

Now we turn to the details of (78). Following Maulik and Pandharipande [2013] (but replacing q by q^8), consider the power series

$$A \doteq \sum_{n \in \mathbb{Z}} q^{n^2}, \quad B \doteq \sum_{n \in \mathbb{Z}} (-1)^n q^{n^2}, \quad \Psi = 108 \sum_{n > 0} q^{8n^2}, \tag{79}$$

and Θ defined by

$$\begin{aligned} 2^{22}\Theta \doteq & 3A^{21} - 81A^{19}B^2 - 627A^{18}B^3 - 14436A^{17}B^4 - 20007A^{16}B^5 \\ & - 169092A^{15}B^6 - 120636A^{14}B^7 - 621558A^{13}B^8 \\ & - 292796A^{12}B^9 - 1038366A^{11}B^{10} - 346122A^{10}B^{11} \\ & - 878388A^9B^{12} - 207186A^8B^{13} - 361908A^7B^{14} - 56364A^6B^{15} \\ & - 60021A^5B^{16} - 4812A^4B^{17} - 1881A^3B^{18} - 27A^2B^{19} + B^{21}. \end{aligned} \tag{80}$$

From [Maulik and Pandharipande 2013, Corollary 2], we have, for any $\Delta > 0$,

$$\deg NL_\Delta \leq \text{coeff. of } q^\Delta \text{ in } \Theta - \Psi. \tag{81}$$

In fact, this is an equality when the components of \mathcal{NL}_Δ are given appropriate multiplicities. Let $\Theta[k]$ denote the coefficient of q^k in Θ . By (81), we only need to bound $\Theta[\Delta]$ in order to bound $\deg NL_\Delta$. To do so, replace every negative sign in the definition of Θ by a positive sign, including those in B , to obtain the *coefficientwise* inequality

$$\Theta \leq 6 \left(\sum_{n \in \mathbb{Z}} q^{n^2} \right)^{21}. \tag{82}$$

The coefficient of q^k in $(\sum_{n \in \mathbb{Z}} q^{n^2})^{21}$ is

$$r_{21}(k) \doteq \#\left\{ (a_1, \dots, a_{21}) \in \mathbb{Z}^{21} \mid \sum_i a_i^2 = k \right\}. \tag{83}$$

The asymptotic bound $r_d(k) = O(k^{d/2-1})$, for $d > 4$, is well known, e.g., [Krätzel 2000, Satz 5.8].

4. Separation bound

We now state and prove the main results. Recall that $a \uparrow b = a^b$ is right associative, and for $\gamma \in H_\mathbb{Z}$, we defined the discriminant $\Delta(\gamma)$ as $(\gamma \cdot h)^2 - 4\gamma \cdot \gamma$.

Theorem 17. *For any $f \in U_4$ with algebraic coefficients, there is a computable constant $c > 1$ such that for any $\gamma \in H^2(X_f, \mathbb{Z})$, if $\gamma \cdot \omega_f \neq 0$, then*

$$|\gamma \cdot \omega_f| > (2 \uparrow c \uparrow \Delta(\gamma) \uparrow \frac{9}{2})^{-1}.$$

To make the connection with (1), recall the map T introduced in (7). We choose a basis $\gamma_1, \dots, \gamma_{21}$ of $H_3(\mathbb{P}^3 \setminus X_f, \mathbb{Z}) \simeq H_{\mathbb{Z}}/\mathbb{Z}h$, write $T(\gamma) = \sum_i x_i \gamma_i$ and observe that $\Delta(\gamma)$ is a quadratic function of the coordinates x_i , so that $\Delta(\gamma)^{1/2} \leq C \max_i |x_i|$ for some constant C depending on the choice of basis.

4.1. Multiplicity of Noether–Lefschetz loci. The multiplicity at a point $p \in \mathbb{C}^s$ of some nonzero polynomial $F \in \mathbb{C}[x_1, \dots, x_s]$ is the unique integer k such that all partial derivatives of F of order $< k$ vanish at p and some partial derivative of order k does not. It is denoted by $\text{mult}_p F$.

The multiplicity of NL_{Δ} at some $f \in U_4$ is related to the elements of $\text{Pic}(X_f)$ with discriminant Δ . For $\Delta > 0$, let E_{Δ} be a set of representatives for elements of discriminant Δ modulo the equivalence relation \sim on $H_{\mathbb{Z}}$ defined by

$$\gamma \sim \gamma', \quad \text{if } \exists a \in \mathbb{Q}^*, b \in \mathbb{Q} : \gamma' = a\gamma + bh. \tag{84}$$

Lemma 18. *For any $f \in U_4$ and any $\Delta > 0$,*

$$\text{mult}_f \text{NL}_{\Delta} = \#(\text{Pic } X_f \cap E_{\Delta}).$$

Proof. Let $\widetilde{\mathcal{NL}}_{\Delta}$ be the lift of \mathcal{NL}_{Δ} in \widetilde{U}_4 . Arguing as in Section 3.1.1, $\widetilde{\mathcal{NL}}_{\Delta}$ is the union of smooth analytic hypersurfaces

$$\widetilde{\mathcal{NL}}_{\Delta} = \bigcup_{\eta \in E_{\Delta}} \mathcal{P}^{-1}\{w \in \mathcal{D} \mid w \cdot \eta = 0\}. \tag{85}$$

Then the same holds locally for \mathcal{NL}_{Δ} .

For any $f \in U_4$, it follows from the smoothness of branches of \mathcal{NL}_{Δ} that $\text{mult}_f \text{NL}_{\Delta}$ is exactly the number of branches meeting at f . The branches meeting at f are described by the elements of $\text{Pic } X_f$ with discriminant Δ . Two elements γ and γ' describe the same branch (that is, the same hyperplane section of \mathcal{D}) if and only if $\gamma' \sim \gamma$. So $\text{mult}_f \text{NL}_{\Delta}$ is exactly the number of equivalence classes in $\{\gamma \in \text{Pic } X_f \mid \Delta(\gamma) = \Delta\}$ for this relation. \square

4.2. Proof of Theorem 17. We first apply Corollary 3. Let $\varepsilon = 4|\gamma \cdot \omega_f|$. The corollary gives constants $C_f > 0$ and $\varepsilon_f > 0$ (depending only on f) such that if $\varepsilon < \varepsilon_f$, then there exists a monomial $m \in R_4$ and $t \in \mathbb{C}$ such that

$$|t| \leq C_f \varepsilon \tag{86}$$

and

$$\gamma \in \text{Pic } X_{f+tm}. \tag{87}$$

Assume $\varepsilon < \varepsilon_f$. As u varies, the number $\#(\text{Pic}(X_{f+um}) \cap E_{\Delta})$ has a strict local maximum at $u = t$, where t and m are as above. By Lemma 18, so does $\text{mult}_{f+um} \text{NL}_{\Delta(\gamma)}$. In particular, there is some higher-order

partial derivative of NL_Δ which vanishes at $f + tm$ but not at $f + um$, for u close to but not equal to t . Let $\alpha \in \mathbb{N}^{35}$ be the multiindex for which

$$P \doteq \frac{1}{\alpha_1! \cdots \alpha_{35}!} \frac{\partial^{|\alpha|} NL_\Delta}{\partial u^\alpha} \in \mathbb{Z}[u_1, \dots, u_{35}] \tag{88}$$

is this derivative. For a monomial $u^\beta \doteq u_1^{\beta_1} \cdots u_{35}^{\beta_{35}}$, we have

$$\frac{1}{\alpha_1! \cdots \alpha_{35}!} \frac{\partial^{|\alpha|} u^\beta}{\partial u^\alpha} = \prod_{i=1}^{35} \binom{\beta_i}{\alpha_i} u^{\beta-\alpha}. \tag{89}$$

Since $\binom{\beta_i}{\alpha_i} \leq 2^{\beta_i}$, it follows that

$$\left\| \frac{1}{\alpha_1! \cdots \alpha_{35}!} \frac{\partial^{|\alpha|} NL_\Delta}{\partial u^\alpha} \right\|_1 \leq 2^{\deg NL_\Delta} \|NL_\Delta\|_1. \tag{90}$$

Let $Q \in \overline{\mathbb{Q}}[s]$ be the polynomial $Q(s) \doteq P(f + sm)$. By construction, $Q \neq 0$ and $Q(t) = 0$. Clearly $\deg Q \leq \deg NL_\Delta$, and we check that

$$\|Q\|_1 \leq \|P\|_1 (\|f\|_1 + 1)^{\deg P}. \tag{91}$$

Then

$$\|Q\|_1 \leq 2^{\deg NL_\Delta} \|NL_\Delta\|_1 (\|f\|_1 + 1)^{\deg NL_\Delta}. \tag{92}$$

From Corollary 15, we find a constant c depending only on f such that

$$\deg Q \leq c \uparrow \Delta \uparrow \frac{9}{2} \quad \text{and} \quad \|Q\|_1 \leq 2 \uparrow c \uparrow \Delta \uparrow \frac{9}{2}. \tag{93}$$

We write $Q = \sum_{i=0}^{\deg Q} q_i s^i$. Let k be the smallest integer such that $q_k \neq 0$. Since $Q(t) = 0$, it follows that

$$|q_k t^k| \leq \sum_{i=k+1}^{\deg Q} |q_i t^i|. \tag{94}$$

If $\varepsilon < C_f^{-1}$, we have $|t| < 1$, by (86), and it follows that

$$|t| \geq \frac{|q_k|}{\|Q\|_1}. \tag{95}$$

Let $D \geq 1$ be the degree of the number field generated by the coefficients of f . Let $H > 0$ be an upper bound for the absolute logarithmic Weil height for the coefficient vector of f [Waldschmidt 2000, p. 77]. Then q_k is an algebraic number defined by a polynomial expression $\tilde{q}_k(f)$ in the coefficients of f , with \tilde{q}_k having integer coefficients. Liouville’s inequality [Waldschmidt 2000, Proposition 3.14] gives

$$|q_k| \geq \|\tilde{q}_k\|_1^{-D+1} e^{-DH \deg \tilde{q}_k}. \tag{96}$$

It is easy to see that $\deg \tilde{q}_k \leq \deg NL_\Delta$ and $\|\tilde{q}_k\|_1 \leq 2^{\deg NL_\Delta} \|NL_\Delta\|_1$, the latter can be bounded by $\|Q\|_1$.

By (86), this leads to

$$\varepsilon \geq (2 \uparrow c \uparrow \Delta \uparrow \frac{9}{2})^{-D(1+H)}, \tag{97}$$

for some other constant c depending only on f . Recall that (97) holds with the assumptions that $\varepsilon \leq \varepsilon_f$ and $\varepsilon < C_f^{-1}$. However, we can choose c large enough so that the right-hand side of (97) is smaller than ε_f and C_f^{-1} . Then (97) holds unconditionally. Absorb the outer exponent of (97) into c to conclude the proof of Theorem 17. \square

4.3. Numbers à la Liouville. Let $(\theta_i)_{i \geq 0}$ be a sequence of positive integers such that θ_i is a strict divisor of θ_{i+1} for all $i \geq 0$ (in particular, $\theta_i \geq 2^i$.) Consider the number

$$L_\theta \doteq \sum_{i=0}^{\infty} \theta_i^{-1}.$$

As a corollary to the separation bound obtained in Theorem 17, the following result states that L_θ is not a ratio of periods of quartic surfaces when θ grows fast enough:

Theorem 19. *If $\theta_{i+1} \geq 2 \uparrow 2 \uparrow \theta_i \uparrow 10$, for all i large enough, then L_θ is not equal to $(\gamma_1 \cdot \omega_f)/(\gamma_2 \cdot \omega_f)$ for any $\gamma_1, \gamma_2 \in H_{\mathbb{Z}}$ and any $f \in U_4$ with algebraic coefficients.*

Proof. Let $l_k = \sum_{i=0}^k \theta_i^{-1}$. Since θ_i divides θ_{i+1} , we can write $l_k = u_k/\theta_k$ for some integer u_k . And since the divisibility is strict, $\theta_i \geq 2^i$ and $u_k \leq 2\theta_k$. Moreover,

$$0 < L_\theta - l_k \leq 2\theta_{k+1}^{-1}, \tag{98}$$

using $\theta_{k+i+1} \geq 2^i \theta_{k+1}$, for any $i \geq 0$. Assume now that $L_\theta = (\gamma_1 \cdot \omega_f)/(\gamma_2 \cdot \omega_f)$ for some $\gamma_1, \gamma_2 \in H_{\mathbb{Z}}$ and some $f \in U_4$ with rational coefficients. Then, with

$$\gamma_k \doteq \theta_k \gamma_1 - u_k \gamma_2, \tag{99}$$

we check that $\Delta(\gamma_k) = O(\theta_k^2)$ and that

$$0 < |\theta_k| |\gamma_2 \cdot \omega_f| (L_\theta - l_k) = |\gamma_k \cdot \omega_f| \leq C \frac{\theta_k}{\theta_{k+1}}, \tag{100}$$

for some constant C . By Theorem 17, we obtain

$$(2 \uparrow c \uparrow \theta_k \uparrow 9)^{-1} \leq C \frac{\theta_k}{\theta_{k+1}}, \tag{101}$$

for some constant $c > 0$ which depends only on f . This contradicts the assumption on the growth of θ . \square

4.4. Computational complexity. Given a polynomial $f \in \overline{\mathbb{Q}}[w, x, y, z] \cap U_4$ and a cohomology class $\gamma \in H^2(X_f, \mathbb{Z})$, we can decide if $\gamma \in \text{Pic}(X_f)$ (that is, $\gamma \cdot \omega_f = 0$) as follows:

- (a) Compute the constant c in Theorem 17.
- (b) Let $\varepsilon = (2 \uparrow c \uparrow \Delta(\gamma) \uparrow \frac{9}{2})^{-1}$ and compute an approximation $s \in \mathbb{C}$ of the period $\gamma \cdot \omega_f$ such that $|s - \gamma \cdot \omega_f| < \frac{1}{2}\varepsilon$.

Then γ is in $\text{Pic}(X_f)$ if and only if $|s| < \frac{1}{2}\varepsilon$.

Computing the Picard group itself is an interesting application of this procedure. Algorithms for computing the Picard group of X_f , or even just the rank of it, break the problem into two: a part gives larger and larger lattices inside $\text{Pic}(X_f)$ while the other part gets finer and finer upper bounds on the rank of $\text{Pic}(X_f)$ [Charles 2014; Hassett et al. 2013; Poonen et al. 2015]. The computation stops when the two parts meet. Approximations from the inside are based on finding sufficiently many elements of $\text{Pic}(X_f)$. So while deciding the membership of γ in $\text{Pic}(X_f)$ can be solved by computing $\text{Pic}(X_f)$ first, it makes sense not to assume prior knowledge of the Picard group and to study the complexity of deciding membership as $\Delta(\gamma) \rightarrow \infty$, with f fixed.

Step (a) does not depend on γ , so only the complexity of Step (b) matters, that is, the numerical approximation of $\gamma \cdot \omega_f$. This approximation amounts to numerically solving a Picard–Fuchs differential equation [Sertöz 2019] and the complexity is $(\log(1/\varepsilon))^{1+o(1)}$ [Beeler et al. 1972; van der Hoeven 2001; Mezzarobba 2010; 2016]. With the value of ε in Step (b), we have a complexity bound of $\exp(\Delta(\gamma)^{O(1)})$ for deciding membership.

For the sake of comparison, we may speculate about an approach that would decide the membership of γ in $\text{Pic}(X_f)$ by trying to construct an explicit algebraic divisor on X_f whose cohomology class is equal to γ . It would certainly need to decide the existence of a point satisfying some algebraic conditions in some Hilbert scheme $\mathcal{H}_{d,g}$, with $d = O(\Delta^{\frac{1}{2}})$ and $g = O(\Delta)$ (see Section 3.1.1). Embedding $\mathcal{H}_{d,g}$ (or some fibration over it, as we did in Section 3.4) in some affine chart of a projective space of dimension $d^{O(1)}$ will lead to a complexity of $\exp(\Delta(\gamma)^{O(1)})$ for deciding membership in this way.

However, if Conjecture 16 holds true, then the complexity of the numerical approach for deciding membership would reduce to $\Delta(\gamma)^{O(1)}$.

5. Concluding remarks

5.1. Going beyond quartic surfaces. There are two directions in which the main result, Theorem 17, can, in principle, be generalized beyond quartic surfaces.

In the first direction, our effective methods naturally extend to complete intersections in complete simplicial toric varieties, provided the complete intersection has a K3 type middle cohomology satisfying the integral Hodge conjecture. By this last condition, we mean that a single period should govern if a homology cycle is algebraic. For instance, cubic fourfolds satisfy all of these conditions [Voisin 2013]. Of course, polarized K3 surfaces of degrees 2, 6 and 8 also work, in addition to the degree 4 case covered here.

To generalize the result to this context, one needs to compute two ingredients. The height and degree bounds for the image of a Hilbert schemes, and the “spread” of the period map (as in Section 2.4). Our use of effective Nullstellensatz to compute heights clearly extends. To compute the spread, we used the Griffiths–Dwork reduction, which continues to work for complete intersections in compact simplicial toric varieties [Batyrev and Cox 1994; Dimca 1995; Mavlyutov 1999].

The second direction one could generalize the result is to stick with surfaces in \mathbb{P}^3 but to increase the degree. In this case, we do not know how to control the vanishing of individual period integrals. However, the Lefschetz (1, 1)-theorem can be used to relate algebraic cycles to the simultaneous vanishing of a vector of periods coming from all holomorphic forms. For instance, on quintic surfaces one can separate 4-dimensional (holomorphic) period vectors from one another. The deduction of the separation bounds would be possible from a parallel discussion to the one provided here. This application would make it possible to prove our heuristic Picard group computations of surfaces [Lairez and Sertöz 2019].

It would also be highly desirable to be able to numerically verify arbitrary, nonlinear, relations between periods of quartics. However, in order to generalize our approach to this setup, one would need the integral Hodge conjecture on products of quartic surfaces.

5.2. Closed formulae for the bounds. It is possible to determine a closed formula, involving the height of f , that bounds the constant c in Theorem 17. We removed the deduction of such a formula due to the excessive technical complexity it presents. In addition, the pursuit of a human readable bound gets us further and further from the optimal bounds. We envisioned using the constant c on computer calculations where an algorithmic deduction of c is possible and preferable. We designed our proofs so that such an algorithm is explicit in the proofs. An implementation of this algorithm would be beneficial after the bounds for the heights of the Noether–Lefschetz loci are brought down significantly.

5.3. Optimal bounds. We conjectured by analogy (Conjecture 16) that our bounds for the height of the Noether–Lefschetz locus can be lowered by one level of exponentiation. One can be more optimistic based on the following observation: For many example quartics X_f , we determined the equations for the Hilbert scheme of lines over each pencil X_{f+tm} for monomials m . Then, going through the algorithm in the proofs, we computed sharper separation bounds on these example quartics. On these examples, the separation bound was around 10^{-60} . In other words, it was sufficient to deduce whether a homology cycle was the class of a line using only 60 digits of precision. This suggests that for homology cycles of small discriminant, optimal separation bounds may be small enough to be used in practice. It would be interesting to see if generalizing the work of Maulik and Pandharipande from degrees to heights by using the modularity of arithmetic Chow rings [Kudla 2003] would give close to optimal bounds.

5.4. Analogies with related work. Our construction bears a remote resemblance to the analytic subgroup theorem of Wüstholz [1989] and the period theorem of Masser and Wüstholz [1993]. The analytic subgroup theorem and its applications work with the exponential map $\exp_A : T_0A \rightarrow A$ of a (principally polarized) abelian variety A over $\bar{\mathbb{Q}} \subset \mathbb{C}$. The periods of A form a lattice $\Lambda \doteq \ker \exp A$. Let $P = \exp_A^{-1} A(\bar{\mathbb{Q}})$ be the periods of all algebraic points on A .

The analytic subgroup theorem implies that $\bar{\mathbb{Q}}$ -linear relations between any set of elements $S \subset P$ are determined by abelian subvarieties of B : there is an abelian subvariety such that T_0B coincides with the span of S . Observe that the linear relations live on the domain of the transcendental map \exp_A and are converted to an algebraic subvariety on the codomain. When $S = \{\gamma\} \subset \Lambda$, the Masser–Wüstholz period theorem bounds the degree of smallest B whose tangent space contains γ using the height of A and the norm of γ .

In our work, we consider the space U_4 of smooth homogeneous quartic polynomials of degree 4 and its universal cover $\tilde{U}_4 \rightarrow U_4$. We then take the (transcendental) period map $\mathcal{P}: \tilde{U}_4 \rightarrow H_{\mathbb{C}}$. Note that the \mathbb{Z} -relations between periods are realized as linear subspaces of the period domain, whereas the preimage of these linear spaces are the Noether–Lefschetz loci. These Noether–Lefschetz loci map to algebraic hypersurfaces on the space U_4 .

Superficially, the main difference between the two approaches is the direction of the naturally appearing transcendental maps that linearize relations between periods. However, the nature of the two transcendental maps appearing in both constructions also differs substantially.

Acknowledgements

We thank Bjorn Poonen for suggesting the use of heights of Noether–Lefschetz loci and Gavril Farkas for suggesting the paper by Maulik and Pandharipande. We also thank Alin Bostan and Matthias Schütt for numerous helpful comments. We thank the referee for a careful reading of the paper.

Sertöz was supported by the Max Planck Institute for Mathematics in the Sciences, Leibniz University Hannover, and the Max Planck Institute for Mathematics. Lairez was supported by the project *De Rerum Natura* ANR-19-CE40-0018 of the French National Research Agency (ANR).

References

- [Batyrev and Cox 1994] V. V. Batyrev and D. A. Cox, “On the Hodge structure of projective hypersurfaces in toric varieties”, *Duke Math. J.* **75**:2 (1994), 293–338. MR Zbl
- [Bayer 1982] D. A. Bayer, *The division algorithm and the Hilbert scheme*, Ph.D. thesis, Harvard University, 1982, available at <https://www.proquest.com/docview/303209159>.
- [Beeler et al. 1972] M. Beeler, R. W. Gosper, and R. Schroepfel, “HAKMEM”, AI memo 239, MIT AI Lab., 1972, available at <https://tinyurl.com/hakmem239>.
- [Blum et al. 1998] L. Blum, F. Cucker, M. Shub, and S. Smale, *Complexity and real computation*, Springer, 1998. MR Zbl
- [Booker et al. 2016] A. R. Booker, J. Sijlsing, A. V. Sutherland, J. Voight, and D. Yasaki, “A database of genus-2 curves over the rational numbers”, *LMS J. Comput. Math.* **19**:suppl. A (2016), 235–254. MR Zbl
- [Bost et al. 1991] J.-B. Bost, H. Gillet, and C. Soulé, “Un analogue arithmétique du théorème de Bézout”, *C. R. Acad. Sci. Paris Sér. I Math.* **312**:11 (1991), 845–848. MR Zbl
- [Brachat et al. 2016] J. Brachat, P. Lella, B. Mourrain, and M. Roggero, “Extensors and the Hilbert scheme”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **16**:1 (2016), 65–96. MR Zbl
- [Bruin et al. 2019] N. Bruin, J. Sijlsing, and A. Zotine, “Numerical computation of endomorphism rings of Jacobians”, pp. 155–171 in *Proceedings of the Thirteenth Algorithmic Number Theory Symposium* (Madison, WI, 2018), edited by R. Scheidler and J. Sorenson, Open Book Ser. **2**, Math. Sci. Publ., Berkeley, CA, 2019. MR
- [Charles 2014] F. Charles, “On the Picard number of K3 surfaces over number fields”, *Algebra Number Theory* **8**:1 (2014), 1–17. MR Zbl
- [Costa et al. 2019] E. Costa, N. Mascot, J. Sijlsing, and J. Voight, “Rigorous computation of the endomorphism ring of a Jacobian”, *Math. Comp.* **88**:317 (2019), 1303–1339. MR Zbl
- [D’Andrea et al. 2013] C. D’Andrea, T. Krick, and M. Sombra, “Heights of varieties in multiprojective spaces and arithmetic Nullstellensätze”, *Ann. Sci. École Norm. Sup. (4)* **46**:4 (2013), 549–627. MR Zbl
- [Dimca 1995] A. Dimca, “Residues and cohomology of complete intersections”, *Duke Math. J.* **78**:1 (1995), 89–100. MR Zbl

- [Gotzmann 1978] G. Gotzmann, “Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes”, *Math. Z.* **158**:1 (1978), 61–70. MR Zbl
- [Griffiths 1969a] P. A. Griffiths, “On the periods of certain rational integrals, I”, *Ann. of Math. (2)* **90**:3 (1969), 460–495. MR Zbl
- [Griffiths 1969b] P. A. Griffiths, “On the periods of certain rational integrals, II”, *Ann. of Math. (2)* **90**:3 (1969), 496–541. MR Zbl
- [Griffiths and Harris 1978] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Wiley, New York, 1978. MR Zbl
- [Grothendieck 1966] A. Grothendieck, “Techniques de construction et théorèmes d’existence en géométrie algébrique, IV: Les schémas de Hilbert”, exposé 221 in *Séminaire Bourbaki: 1960/1961*, W. A. Benjamin, New York, 1966. Reprinted as pp. 249–276 in *Séminaire Bourbaki* **6**, Soc. Math. France, Paris, 1995. MR
- [Hartshorne 1977] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics **52**, Springer, 1977. MR Zbl
- [Hassett et al. 2013] B. Hassett, A. Kresch, and Y. Tschinkel, “Effective computation of Picard groups and Brauer–Manin obstructions of degree two $K3$ surfaces over number fields”, *Rend. Circ. Mat. Palermo (2)* **62**:1 (2013), 137–151. MR Zbl
- [van der Hoeven 2001] J. van der Hoeven, “Fast evaluation of holonomic functions near and in regular singularities”, *J. Symbolic Comput.* **31**:6 (2001), 717–743. MR Zbl
- [Huybrechts 2016] D. Huybrechts, *Lectures on $K3$ surfaces*, Cambridge Stud. Adv. Math. **158**, Cambridge Univ. Press, 2016. MR Zbl
- [Kontsevich and Zagier 2001] M. Kontsevich and D. Zagier, “Periods”, pp. 771–808 in *Mathematics unlimited: 2001 and beyond*, edited by B. Engquist and W. Schmid, Springer, 2001. MR Zbl
- [Krätzel 2000] E. Krätzel, *Analytische Funktionen in der Zahlentheorie*, Teubner-Texte Math. **139**, Teubner, Stuttgart, 2000. MR Zbl
- [Krick et al. 2001] T. Krick, L. M. Pardo, and M. Sombra, “Sharp estimates for the arithmetic Nullstellensatz”, *Duke Math. J.* **109**:3 (2001), 521–598. MR Zbl
- [Kudla 2003] S. S. Kudla, “Modular forms and arithmetic geometry”, pp. 135–179 in *Current developments in mathematics (Cambridge, 2002)*, edited by D. Jerison et al., Int. Press, Somerville, MA, 2003. MR Zbl
- [Lairez and Sertöz 2019] P. Lairez and E. C. Sertöz, “A numerical transcendental method in algebraic geometry: computation of Picard groups and related invariants”, *SIAM J. Appl. Algebra Geom.* **3**:4 (2019), 559–584. MR Zbl
- [Lazard 1977] D. Lazard, “Algèbre linéaire sur $K[X_1, \dots, X_n]$, et élimination”, *Bull. Soc. Math. France* **105**:2 (1977), 165–190. MR Zbl
- [Lella 2012] P. Lella, *Computable Hilbert schemes*, Ph.D. thesis, Università degli Studi di Torino, 2012. arXiv 1202.4384
- [Liouville 1851] J. Liouville, “Sur des classes très-étendues de quantités dont la valeur n’est ni algébrique, ni même réductible à des irrationnelles algébriques”, *J. Math. Pures Appl.* **16** (1851), 133–142.
- [Masser and Wüstholz 1993] D. Masser and G. Wüstholz, “Periods and minimal abelian subvarieties”, *Ann. of Math. (2)* **137**:2 (1993), 407–458. MR Zbl
- [Maulik and Pandharipande 2013] D. Maulik and R. Pandharipande, “Gromov–Witten theory and Noether–Lefschetz theory”, pp. 469–507 in *A celebration of algebraic geometry* (Cambridge, 2011), edited by B. Hassett et al., Clay Math. Proc. **18**, Amer. Math. Soc., Providence, RI, 2013. MR Zbl
- [Mavlyutov 1999] A. R. Mavlyutov, “Cohomology of complete intersections in toric varieties”, *Pacific J. Math.* **191**:1 (1999), 133–144. MR Zbl
- [Mezzarobba 2010] M. Mezzarobba, “NumGfun: a package for numerical and analytic computation with D-finite functions”, pp. 139–146 in *ISSAC 2010: Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation* (Munich, 2010), edited by S. M. Watt, ACM, New York, 2010. MR Zbl
- [Mezzarobba 2016] M. Mezzarobba, “Rigorous multiple-precision evaluation of D-finite functions in SageMath”, lecture notes, 2016. arXiv 1607.01967
- [Philippon 1995] P. Philippon, “Sur des hauteurs alternatives, III”, *J. Math. Pures Appl. (9)* **74**:4 (1995), 345–365. MR Zbl
- [Piene and Schlessinger 1985] R. Piene and M. Schlessinger, “On the Hilbert scheme compactification of the space of twisted cubics”, *Amer. J. Math.* **107**:4 (1985), 761–774. MR Zbl

- [Poonen et al. 2015] B. Poonen, D. Testa, and R. van Luijk, “Computing Néron–Severi groups and cycle class groups”, *Compos. Math.* **151**:4 (2015), 713–734. MR Zbl
- [Rémond 2001a] G. Rémond, “Élimination multihomogène”, pp. 53–81 in *Introduction to algebraic independence theory*, edited by Y. V. Nesterenko and P. Philippon, Lecture Notes in Math. **1752**, Springer, 2001. MR
- [Rémond 2001b] G. Rémond, “Géométrie diophantienne multiprojective”, pp. 95–131 in *Introduction to algebraic independence theory*, edited by Y. V. Nesterenko and P. Philippon, Lecture Notes in Math. **1752**, Springer, 2001. MR
- [Sertöz 2019] E. C. Sertöz, “Computing periods of hypersurfaces”, *Math. Comp.* **88**:320 (2019), 2987–3022. MR Zbl
- [Shidlovskii 1989] A. B. Shidlovskii, *Transcendental numbers*, de Gruyter Stud. Math. **12**, de Gruyter, Berlin, 1989. MR Zbl
- [Simpson 2008] C. Simpson, “Algebraic cycles from a computational point of view”, *Theoret. Comput. Sci.* **392**:1-3 (2008), 128–140. MR Zbl
- [Smale 1986] S. Smale, “Newton’s method estimates from data at one point”, pp. 185–196 in *The merging of disciplines: new directions in pure, applied, and computational mathematics* (Laramie, WY, 1985), edited by R. E. Ewing et al., Springer, 1986. MR Zbl
- [Voisin 2003] C. Voisin, *Hodge theory and complex algebraic geometry, II*, Cambridge Stud. Adv. Math. **77**, Cambridge Univ. Press, 2003. MR Zbl
- [Voisin 2013] C. Voisin, “Abel–Jacobi map, integral Hodge classes and decomposition of the diagonal”, *J. Algebraic Geom.* **22**:1 (2013), 141–174. MR Zbl
- [Waldschmidt 2000] M. Waldschmidt, *Diophantine approximation on linear algebraic groups*, Grundlehren der Math. Wissenschaften **326**, Springer, 2000. MR Zbl
- [van Wamelen 1999] P. van Wamelen, “Examples of genus two CM curves defined over the rationals”, *Math. Comp.* **68**:225 (1999), 307–320. MR Zbl
- [Wüstholz 1989] G. Wüstholz, “Algebraische Punkte auf analytischen Untergruppen algebraischer Gruppen”, *Ann. of Math. (2)* **129**:3 (1989), 501–517. MR Zbl

Communicated by Antoine Chambert-Loir

Received 2021-06-18 Revised 2022-05-19 Accepted 2022-09-21

pierre.lairez@inria.fr

INRIA Saclay, Palaiseau, France

emre@serto.com

*Faculty of Mathematics and Physics, Leibniz Universität Hannover,
Hannover, Germany*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR
Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR
David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2023 is US \$485/year for the electronic version, and \$705/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 17 No. 10 2023

Special cycles on the basic locus of unitary Shimura varieties at ramified primes YOUSHENG SHI	1681
Hybrid subconvexity bounds for twists of $GL(3) \times GL(2)$ L -functions BINGRONG HUANG and ZHAO XU	1715
Separation of periods of quartic surfaces PIERRE LAIREZ and EMRE CAN SERTÖZ	1753
Global dimension of real-exponent polynomial rings NATHAN GEIST and EZRA MILLER	1779
Differences between perfect powers: prime power gaps MICHAEL A. BENNETT and SAMIR SIKSEK	1789
On fake linear cycles inside Fermat varieties JORGE DUQUE FRANCO and ROBERTO VILLAFLOR LOYOLA	1847