

# *Algebra & Number Theory*

Volume 17  
2023  
No. 11

**Bézoutians and the  $\mathbb{A}^1$ -degree**

Thomas Brazelton, Stephen McKean and Sabrina Pauli





# Bézoutians and the $\mathbb{A}^1$ -degree

Thomas Brazelton, Stephen McKean and Sabrina Pauli

We prove that both the local and global  $\mathbb{A}^1$ -degree of an endomorphism of affine space can be computed in terms of the multivariate Bézoutian. In particular, we show that the Bézoutian bilinear form, the Scheja–Storch form, and the  $\mathbb{A}^1$ -degree for complete intersections are isomorphic. Our global theorem generalizes Cazanave’s theorem in the univariate case, and our local theorem generalizes Kass–Wickelgren’s theorem on EKL forms and the local degree. This result provides an algebraic formula for local and global degrees in motivic homotopy theory.

## 1. Introduction

Morel’s  $\mathbb{A}^1$ -Brouwer degree [25] assigns a bilinear form-valued invariant to a given endomorphism of affine space. However, Morel’s construction is not explicit. In order to make computations and applications, we would like algebraic formulas for the  $\mathbb{A}^1$ -degree. Such formulas were constructed by Cazanave for the global  $\mathbb{A}^1$ -degree in dimension 1 [9], Kass and Wickelgren for the local  $\mathbb{A}^1$ -degree at rational points and étale points [16], and Brazelton, Burklund, McKean, Montoro and Opie for the local  $\mathbb{A}^1$ -degree at separable points [7]. In this paper, we give a general algebraic formula for the  $\mathbb{A}^1$ -degree in both the global and local cases. In the global case, we remove Cazanave’s dimension restriction, while in the local case, we remove previous restrictions on the residue field of the point at which the local  $\mathbb{A}^1$ -degree is taken.

Let  $k$  be a field, and let  $f = (f_1, \dots, f_n) : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  be an endomorphism of affine space with isolated zeros, so that  $Q := k[x_1, \dots, x_n]/(f_1, \dots, f_n)$  is a complete intersection. We now recall the definition of the Bézoutian of  $f$ , as well as a special bilinear form determined by the Bézoutian. Introduce new variables  $X := (X_1, \dots, X_n)$  and  $Y := (Y_1, \dots, Y_n)$ . For each  $1 \leq i, j \leq n$ , define the quantity

$$\Delta_{ij} := \frac{f_i(Y_1, \dots, Y_{j-1}, X_j, \dots, X_n) - f_i(Y_1, \dots, Y_j, X_{j+1}, \dots, X_n)}{X_j - Y_j}.$$

**Definition 1.1.** The *Bézoutian* of  $f$  is the image  $\text{Béz}(f_1, \dots, f_n)$  of the determinant  $\det(\Delta_{ij})$  in  $k[X, Y]/(f(X), f(Y))$ . Given a basis  $\{a_1, \dots, a_m\}$  of  $Q$  as a  $k$ -vector space, there exist scalars  $B_{i,j}$  for which

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i,j=1}^m B_{i,j} a_i(X) a_j(Y).$$

MSC2020: primary 14F42; secondary 55M25.

Keywords: Bézoutian, Brouwer degree,  $\mathbb{A}^1$ -degree, motivic homotopy.

We define the *Bézoutian form* of  $f$  to be the class  $\beta_f$  in the Grothendieck–Witt ring  $\mathrm{GW}(k)$  determined by the bilinear form  $Q \times Q \rightarrow k$  with Gram matrix  $(B_{i,j})$ .

For any isolated zero of  $f$  corresponding to a maximal ideal  $\mathfrak{m}$ , there is an analogous bilinear form  $\beta_{f,\mathfrak{m}}$  on the local algebra  $Q_{\mathfrak{m}}$ . We refer to  $\beta_{f,\mathfrak{m}}$  as the *local Bézoutian form* of  $f$  at  $\mathfrak{m}$ . We will demonstrate that both  $\beta_f$  and  $\beta_{f,\mathfrak{m}}$  yield well-defined classes in  $\mathrm{GW}(k)$ . Our main theorem is that the Bézoutian form of  $f$  agrees with the  $\mathbb{A}^1$ -degree in both the local and global contexts.

**Theorem 1.2.** *Let  $\mathrm{char} k \neq 2$ . Let  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  have an isolated zero at a closed point  $\mathfrak{m}$ . Then  $\beta_{f,\mathfrak{m}}$  is isomorphic to the local  $\mathbb{A}^1$ -degree of  $f$  at  $\mathfrak{m}$ . If we further assume that all the zeros of  $f$  are isolated, then  $\beta_f$  is isomorphic to the global  $\mathbb{A}^1$ -degree of  $f$ .*

Because the Bézoutian form can be explicitly computed using commutative algebraic tools, Theorem 1.2 provides a tractable formula for  $\mathbb{A}^1$ -degrees and Euler classes in motivic homotopy theory. Using the Bézoutian formula for the  $\mathbb{A}^1$ -degree, we are able to deduce several computational rules for the degree. We also provide a Sage implementation for calculating local and global  $\mathbb{A}^1$ -degrees via the Bézoutian at [8].

**Remark 1.3.** The key contribution of this article is computability. Building on the work of Kass and Wickelgren [16], Bachmann and Wickelgren [2] show that the  $\mathbb{A}^1$ -degree agrees with the Scheja–Storch form as elements of  $\mathrm{KO}^0(k)$ . In Theorem 5.1, we show how this immediately implies that the  $\mathbb{A}^1$ -degree and Scheja–Storch form determine the same element of  $\mathrm{GW}(k)$ . Scheja and Storch [30] showed that their form is a Bézoutian bilinear form (in the sense of Definition 3.8; see also Lemma 4.4 and Remark 4.8), which was further explored by Becker, Cardinal, Roy and Szafraniec [4]. Putting these results together shows that the isomorphism class of the Bézoutian bilinear form is the  $\mathbb{A}^1$ -degree.

In dimension 1, Cazanave [9] gives a simple formula for computing the  $\mathbb{A}^1$ -degree as a Bézoutian bilinear form in the global setting. However, it is not immediately clear how to adapt this to higher dimensions or the local setting. Becker, Cardinal, Roy and Szafraniec show how to compute Bézoutian bilinear forms in terms of “dualizing forms,” but this method is computationally analogous to using the Eisenbud–Khimshiashvili–Levine form to compute the  $\mathbb{A}^1$ -degree [16]. In the proof of Theorem 1.2 (found in Section 5), we show that our two notions of Bézoutian bilinear forms (Definitions 1.1 and 3.8) agree up to isomorphism. Since Definition 1.1 is the desired generalization of Cazanave’s formula, this enables us to calculate  $\mathbb{A}^1$ -degrees in full generality.

**1A. Outline.** Before proving Theorem 1.2, we recall some classical results on Bézoutians (following [4]) in Section 3, as well as the work of Scheja and Storch on residue pairings [30] in Section 4. We then discuss a local decomposition procedure for the Scheja–Storch form and show that the global Scheja–Storch form is isomorphic to the Bézoutian form in Section 4A. In Section 5, we complete the proof of Theorem 1.2 by applying the work of Kass and Wickelgren [16] and Bachmann and Wickelgren [2] on the local  $\mathbb{A}^1$ -degree and the Scheja–Storch form. Using Theorem 1.2, we give an algorithm for computing the local and global  $\mathbb{A}^1$ -degree at the end of Section 5A, available at [8]. In Section 6, we establish some basic properties for

computing degrees. In Section 7, we provide a step-by-step illustration of our ideas by working through some explicit examples. Finally, we implement our code to compute some examples of  $\mathbb{A}^1$ -Euler characteristics of Grassmannians in Section 8. We check our computations by proving a general formula for the  $\mathbb{A}^1$ -Euler characteristic of a Grassmannian in Theorem 8.4. The  $\mathbb{A}^1$ -Euler characteristic of Grassmannians is essentially a folklore result that follows from the work of Hoyois, Levine, and Bachmann and Wickelgren.

**1B. Background.** Let  $\mathrm{GW}(k)$  denote the Grothendieck–Witt group of isomorphism classes of symmetric, nondegenerate bilinear forms over a field  $k$ . Morel’s  $\mathbb{A}^1$ -Brouwer degree [25, Corollary 1.24]

$$\mathrm{deg} : [\mathbb{P}_k^n / \mathbb{P}_k^{n-1}, \mathbb{P}_k^n / \mathbb{P}_k^{n-1}]_{\mathbb{A}^1} \rightarrow \mathrm{GW}(k),$$

which is a group isomorphism (in fact, a ring isomorphism [24, Lemma 6.3.8]) for  $n \geq 2$ , demonstrates that bilinear forms play a critical role in motivic homotopy theory. However, Morel’s  $\mathbb{A}^1$ -degree is nonconstructive. Kass and Wickelgren addressed this problem by expressing the  $\mathbb{A}^1$ -degree as a sum of local degrees [17, Lemma 19] and providing an explicit formula (building on the work of Eisenbud and Levine [11] and Khimshiashvili [13]) for the local  $\mathbb{A}^1$ -degree [16] at rational points and étale points. This explicit formula can also be used to compute the local  $\mathbb{A}^1$ -degree at points with separable residue field by [7]. Together, these results allow one to compute the global  $\mathbb{A}^1$ -degree of a morphism  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  with only isolated zeros by computing the local  $\mathbb{A}^1$ -degrees of  $f$  over its zero locus, so long as the residue field of each point in the zero locus is separable over the base field. In the local case, Theorem 1.2 gives a commutative algebraic formula for the local  $\mathbb{A}^1$ -degree at any closed point.

Cazanave showed that the Bézoutian gives a formula for the global  $\mathbb{A}^1$ -degree of any endomorphism of  $\mathbb{P}_k^1$  [9]. An advantage to Cazanave’s formula is that one does not need to determine the zero locus or other local information about  $f$ . We extend Cazanave’s formula for morphisms  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  with isolated zeros. The work of Scheja and Storch on global complete intersections [30] is central to both [16] and our result. We also rely on the work of Becker, Cardinal, Roy and Szafraniec [4], who describe a procedure for recovering the global version of the Scheja–Storch form.

Theorem 1.2 has applications wherever Morel’s  $\mathbb{A}^1$ -degree is used. One particularly successful application of the  $\mathbb{A}^1$ -degree has been the  $\mathbb{A}^1$ -enumerative geometry program. The goal of this program is to enrich enumerative problems over arbitrary fields by producing  $\mathrm{GW}(k)$ -valued enumerative equations and interpreting them geometrically over various fields. Notable results in this direction include Srinivasan and Wickelgren’s count of lines meeting four lines in three-space [31], Larson and Vogt’s count of bitangents to a smooth plane quartic [19], and Bethea, Kass, and Wickelgren’s enriched Riemann–Hurwitz formula [5]. See [22; 26] for other related works. For a more detailed account of recent developments in  $\mathbb{A}^1$ -enumerative geometry; see [6; 28].

## 2. Notation and conventions

In this section, we fix some standard terminology and notation. Let  $k$  denote an arbitrary field. We will always use  $f = (f_1, \dots, f_n) : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  to denote an endomorphism of affine space, assumed to have

isolated zeros when we work with it in the global context. We denote by  $Q$  the global algebra associated to this endomorphism

$$Q := \frac{k[x_1, \dots, x_n]}{(f_1, \dots, f_n)}.$$

The maximal ideals of  $Q$  correspond to the maximal ideals of  $k[x_1, \dots, x_n]$  on which  $f$  vanishes. For any maximal ideal  $\mathfrak{m}$  of  $k[x_1, \dots, x_n]$  on which  $f$  vanishes, we denote by  $Q_{\mathfrak{m}}$  the local algebra

$$Q_{\mathfrak{m}} := \frac{k[x_1, \dots, x_n]_{\mathfrak{m}}}{(f_1, \dots, f_n)}.$$

If  $\lambda : V \rightarrow k$  is a  $k$ -linear form on any  $k$ -algebra, we will denote by  $\Phi_{\lambda}$  the associated bilinear form given by

$$\begin{aligned} \Phi_{\lambda} : V \times V &\rightarrow k \\ (a, b) &\mapsto \lambda(ab). \end{aligned}$$

**Definition 2.1.** We say that  $\lambda$  is a *dualizing linear form* if  $\Phi_{\lambda}$  is nondegenerate as a symmetric bilinear form [4, 2.1]. If  $\lambda$  is dualizing, then we say that two vector space bases  $\{a_i\}$  and  $\{b_i\}$  of  $V$  are *dual with respect to  $\lambda$*  if

$$\lambda(a_i b_j) = \delta_{ij},$$

where  $\delta_{ij} = 1$  for  $i = j$  and  $\delta_{ij} = 0$  for  $i \neq j$ . We show in Remark 3.6 that if  $\{a_i\}$  and  $\{b_i\}$  are dual with respect to  $\lambda$ , then  $\lambda$  is a dualizing linear form.

More notation will be introduced as we provide an overview of Bézoutians and the Scheja–Storch bilinear form. We will borrow and clarify notation from both [30] and [4].

### 3. Bézoutians

We first provide an overview of the construction of the *Bézoutian*, following [4]. Given one of our  $n$  polynomials  $f_i$ , we introduce two sets of auxiliary indeterminants and study how  $f_i$  changes when we incrementally exchange one set of indeterminants for the other. Explicitly, consider variables  $X := (X_1, \dots, X_n)$  and  $Y := (Y_1, \dots, Y_n)$ . For any  $1 \leq i, j \leq n$ , we denote by  $\Delta_{ij}$  the quantity

$$\Delta_{ij} := \frac{f_i(Y_1, \dots, Y_{j-1}, X_j, \dots, X_n) - f_i(Y_1, \dots, Y_j, X_{j+1}, \dots, X_n)}{X_j - Y_j}.$$

Note that  $\Delta_{ij}$  is a multivariate polynomial. Indeed,

$$f_i(Y_1, \dots, Y_{j-1}, X_j, \dots, X_n) \quad \text{and} \quad f_i(Y_1, \dots, Y_j, X_{j+1}, \dots, X_n)$$

differ only in the terms in which  $X_j$  or  $Y_j$  appear, so we can expand the difference

$$f_i(Y_1, \dots, Y_{j-1}, X_j, \dots, X_n) - f_i(Y_1, \dots, Y_j, X_{j+1}, \dots, X_n) = \sum_{\ell \geq 1} g_{\ell} \cdot (X_j - Y_j)^{\ell},$$

where  $g_{\ell} \in k[Y_1, \dots, Y_{j-1}, X_{j+1}, \dots, X_n]$ . In this notation,  $\Delta_{ij} = \sum_{\ell \geq 1} g_{\ell} \cdot (X_j - Y_j)^{\ell-1}$ .

We view  $\Delta_{ij}$  as living in the tensor product ring  $Q \otimes_k Q$ , under the isomorphism

$$\varepsilon : \frac{k[X, Y]}{(f(X), f(Y))} \xrightarrow{\cong} Q \otimes_k Q,$$

given by sending  $X_i$  to  $x_i \otimes 1$ , and  $Y_i$  to  $1 \otimes x_i$ .

**Definition 3.1.** We define the *Bézoutian* of the polynomials  $f_1, \dots, f_n$  to be the image  $\text{Béz}(f_1, \dots, f_n)$  of the determinant  $\det(\Delta_{ij})$  in  $Q \otimes_k Q$ .

**Example 3.2.** Let  $(f_1, f_2, f_3) = (x_1^2, x_2^2, x_3^2)$ . Then we have that

$$\begin{aligned} \text{Béz}(f_1, f_2, f_3) &= \varepsilon \left( \det \begin{pmatrix} X_1 + Y_1 & 0 & 0 \\ 0 & X_2 + Y_2 & 0 \\ 0 & 0 & X_3 + Y_3 \end{pmatrix} \right) \\ &= \varepsilon((X_1 + Y_1)(X_2 + Y_2)(X_3 + Y_3)) \\ &= x_1x_2x_3 \otimes 1 + x_1x_2 \otimes x_3 + x_1x_3 \otimes x_2 + x_2x_3 \otimes x_1 \\ &\quad + x_1 \otimes x_2x_3 + x_2 \otimes x_1x_3 + x_3 \otimes x_1x_2 + 1 \otimes x_1x_2x_3. \end{aligned}$$

There is a natural multiplication map  $\delta : Q \otimes_k Q \rightarrow Q$ , defined by  $\delta(a \otimes b) = ab$ , that sends the Bézoutian of  $f$  to the image of the Jacobian of  $f$  in  $Q$ .

**Proposition 3.3.** Let  $\text{Jac}(f_1, \dots, f_n)$  be the image of the Jacobian determinant  $\det\left(\frac{\partial f_i}{\partial x_j}\right)$  in  $Q$ . Then

$$\delta(\text{Béz}(f_1, \dots, f_n)) = \text{Jac}(f_1, \dots, f_n) \in Q.$$

*Proof.* Note that  $(\delta \circ \varepsilon)(a(X, Y)) = a(x, x)$  and  $\delta \circ \varepsilon$  is an algebra homomorphism. In particular,  $\delta \circ \varepsilon$  preserves the multiplication and addition occurring in the determinant which defines  $\text{Béz}(f_1, \dots, f_n)$ . Therefore it suffices for us to verify that

$$(\delta \circ \varepsilon)(\Delta_{ij}) = \frac{\partial f_i}{\partial x_j}.$$

Recall that

$$\Delta_{ij} = \frac{f_i(Y_1, \dots, Y_{j-1}, X_j, \dots, X_n) - f_i(Y_1, \dots, Y_j, X_{j+1}, \dots, X_n)}{X_j - Y_j}.$$

Taking the  $x_j$ -Taylor expansion of  $f(x_1, \dots, x_n)$  about  $Y_j$  gives us

$$f_i(x_1, \dots, x_n) = f_i(x_1, \dots, Y_j, \dots, x_n) + \sum_{\ell \geq 1} \frac{\partial^\ell f_i}{\partial x_j^\ell} \cdot (x_j - Y_j)^\ell.$$

We now subtract  $f_i(x_1, \dots, Y_j, \dots, x_n)$  from both sides, evaluate  $x_j \mapsto X_j$ , and divide by  $X_j - Y_j$  to deduce

$$\frac{f_i(x_1, \dots, X_j, \dots, x_n) - f_i(x_1, \dots, Y_j, \dots, x_n)}{(X_j - Y_j)} = \frac{\partial f_i}{\partial x_j} + \sum_{\ell \geq 2} \frac{\partial^\ell f_i}{\partial x_j^\ell} \cdot (X_j - Y_j)^{\ell-1}.$$

Finally, evaluating  $X_j \mapsto x_j$  and  $Y_j \mapsto x_j$  gives us  $(\delta \circ \varepsilon)(\Delta_{ij}) = \frac{\partial f_i}{\partial x_j}$ , as desired. □

**Lemma 3.4.** *Let  $a_1, \dots, a_m$  be any vector space basis for  $Q$ , and write the Bézoutian as*

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i=1}^m a_i \otimes b_i$$

*for some  $b_1, \dots, b_n \in Q$ . Then  $\{b_i\}_{i=1}^m$  is a basis for  $Q$ .*

*Proof.* This is [4, 2.10(iii)]. □

This allows us to associate to the Bézoutian a pair of vector space bases for  $Q$ . Given any such pair of bases, we will construct a unique linear form for which the bases are dual. Before doing so, we establish some equivalent conditions for the duality of a linear form given a pair of bases.

**Proposition 3.5.** *Let  $\{a_i\}$  and  $\{b_i\}$  be a pair of bases for  $B$ . Consider the induced  $k$ -linear isomorphism*

$$\begin{aligned} \Theta : \text{Hom}_k(Q, k) &\rightarrow Q \\ \varphi &\mapsto \sum_i \varphi(a_i) b_i. \end{aligned}$$

*Given a linear form  $\lambda : Q \rightarrow k$ , the following are equivalent:*

- (1) *We have that  $\Theta(\lambda) = \sum_i \lambda(a_i) b_i = 1$ .*
- (2) *For any  $a \in Q$ , we have  $a = \sum_i \lambda(aa_i) b_i$ .*
- (3) *We have that  $\{a_i\}$  and  $\{b_i\}$  are dual with respect to  $\lambda$ .*

*Proof.* Note that (2) implies (1) by setting  $a = 1$ . Next, we remark that  $\Theta$  is a  $Q$ -module isomorphism by [30, 3.3 Satz], where the  $Q$ -module structure on  $\text{Hom}_k(Q, k)$  is given by  $a \cdot \varphi = \varphi(a \cdot -)$ . This allows us to conclude that  $a \cdot \Theta(\lambda) = \Theta(a \cdot \lambda)$  for any linear form  $\lambda$ . In particular, we have

$$a \sum_i \lambda(a_i) b_i = \sum_i \lambda(aa_i) b_i.$$

It follows from this identity that (1) implies (2). Now suppose that (2) holds. By setting  $a = b_j$  for some  $j$ , we have

$$\sum_i \lambda(a_i b_j) b_i = b_j.$$

Since  $\{b_i\}$  is a basis, it follows that  $\lambda(a_i b_j) = \delta_{ij}$ . Thus the bases  $\{a_i\}$  and  $\{b_i\}$  are dual with respect to  $\lambda$ . Finally, suppose that (3) holds, so that  $\lambda(a_i b_j) = \delta_{ij}$ . For any  $a \in Q$ , write  $a$  as  $a := \sum_j c_j b_j$  for some scalars  $c_j$ . Then

$$\sum_i \lambda(aa_i) b_i = \sum_i \lambda\left(a_i \sum_j c_j b_j\right) b_i = \sum_i \left(\sum_j c_j \lambda(a_i b_j)\right) b_i = \sum_i c_i b_i = a.$$

Thus (3) implies (2). □



**Remark 3.6.** If  $\{a_i\}$  and  $\{b_i\}$  are dual with respect to  $\lambda$ , then  $\lambda$  is a dualizing form. Indeed, suppose there exists  $x \in Q$  such that  $\Phi_\lambda(x, y) = 0$  for all  $y \in Q$ . Write  $x = \sum_i x_i a_i$  with  $x_i \in k$ . Then

$$0 = \lambda(xb_j) = \lambda\left(\sum_i x_i a_i b_j\right) = \sum_i x_i \lambda(a_i b_j) = x_j$$

for all  $j$ , so  $x = 0$ .

**Corollary 3.7.** *Let  $\{a_i\}$  and  $\{b_i\}$  be two  $k$ -vector space bases for  $Q$ . Then there exists a unique dualizing linear form  $\lambda : Q \rightarrow k$  such that  $\{a_i\}$  and  $\{b_i\}$  are dual with respect to  $\lambda$ .*

*Proof.* As  $\Theta$  is a  $k$ -algebra isomorphism, it admits a unique preimage of 1. Thus, given any pair of bases  $\{a_i\}$  and  $\{b_i\}$  of  $Q$ , there is a unique dualizing linear form with respect to which  $\{a_i\}$  and  $\{b_i\}$  are dual.  $\square$

**Definition 3.8.** We call  $\Phi_\lambda$  a *Bézoutian bilinear form* if  $\lambda : Q \rightarrow k$  is a dualizing linear form such that

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i=1}^m a_i \otimes b_i,$$

where  $\{a_i\}$  and  $\{b_i\}$  are dual bases with respect to  $\lambda$ .

A priori this is different than the Bézoutian form detailed in Definition 1.1, although we will prove that they define the same class in  $\text{GW}(k)$  in Section 5A.

**Proposition 3.9.** *Given a function  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  with isolated zeros, its Bézoutian bilinear form is a well-defined class in  $\text{GW}(k)$ .*

*Proof.* Let  $\Phi_\lambda$  be a Bézoutian bilinear form for  $f$ . Recall that  $\Phi_\lambda : Q \times Q \rightarrow k$  is defined by  $\Phi_\lambda(a, b) = \lambda(ab)$ . Since  $\lambda$  is a dualizing linear form,  $\Phi_\lambda$  is nondegenerate and as  $Q$  is commutative,  $\Phi_\lambda$  is symmetric. Lemma 3.4 implies that given a basis  $a_1, \dots, a_m$  for  $Q$ , we can write

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i=1}^m a_i \otimes b_i,$$

and obtain a second basis  $b_1, \dots, b_m$  for  $Q$ . By Corollary 3.7, there is a dualizing linear form for the two bases  $\{a_i\}_{i=1}^m$  and  $\{b_i\}_{i=1}^m$ . It remains to show that if

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i=1}^m a_i \otimes b_i = \sum_{i=1}^m a'_i \otimes b'_i,$$

for some bases  $\{a_i\}, \{b_i\}$  dual with respect to  $\lambda$  and  $\{a'_i\}, \{b'_i\}$  dual with respect to  $\lambda'$ , then  $\Phi_\lambda$  and  $\Phi_{\lambda'}$  are isomorphic. We will in fact show that  $\lambda = \lambda'$ , so that  $\Phi_\lambda = \Phi_{\lambda'}$ . Write  $a_i = \sum_s \alpha_{is} a'_s$  and  $b_i = \sum_s \beta_{is} b'_s$ . Then

$$\sum_{i=1}^m a'_i \otimes b'_i = \sum_{i=1}^m a_i \otimes b_i = \sum_i \left( \sum_s \alpha_{is} a'_s \right) \otimes \left( \sum_t \beta_{it} b'_t \right) = \sum_{s,t} \left( \sum_i \alpha_{is} \beta_{it} \right) a'_s \otimes b'_t.$$

Since  $\{a'_s \otimes b'_t\}$  is a basis for  $Q \otimes_k Q$ , we conclude that  $\sum_i \alpha_{is} \beta_{it} = \delta_{st}$ . In particular,  $(\alpha_{ij})^{-1} = (\beta_{ij})^T$ , so  $(\beta_{ij})(\alpha_{ij})^T$  is the identity matrix. Thus  $\sum_j \alpha_{sj} \beta_{tj} = \delta_{st}$ .

Now given  $g = \sum_i c_i a_i = \sum_i c'_i a'_i \in Q$  and  $1 = \sum_i d_i b_i = \sum_i d'_i b'_i$ , we have that

$$\lambda(g) = \lambda\left(\sum_i (c_i a_i) \cdot \sum_j d_j b_j\right) = \sum_{i,j} c_i d_j \lambda(a_i b_j) = \sum_i c_i d_i.$$

Similarly, we have  $\lambda'(g) = \sum_i c'_i d'_i$ . By our change of bases, we have  $c'_j = \sum_i c_j \alpha_{ij}$  and  $d'_j = \sum_i d_i \beta_{ij}$ . Thus

$$\lambda'(g) = \sum_j c'_j d'_j = \sum_j \left(\sum_s c_s \alpha_{sj}\right) \left(\sum_t d_t \beta_{tj}\right) = \sum_{s,t} c_s d_t \left(\sum_j \alpha_{sj} \beta_{tj}\right) = \sum_s c_s d_s = \lambda(g).$$

Therefore  $\lambda = \lambda'$ , as desired. □

**Example 3.10.** Continuing Example 3.2, let  $f = (x_1^2, x_2^2, x_3^2)$ , so that

$$\begin{aligned} \varepsilon^{-1}(\text{Béz}(f_1, f_2, f_3)) &= (X_1 + Y_1)(X_2 + Y_2)(X_3 + Y_3) \\ &= X_1 X_2 X_3 + X_1 X_2 Y_3 + X_1 Y_2 X_3 + X_1 Y_2 Y_3 + Y_1 X_2 X_3 + Y_1 X_2 Y_3 + Y_1 Y_2 X_3 + Y_1 Y_2 Y_3. \end{aligned}$$

We give two bases for  $k[Z_1, Z_2, Z_3]/(Z_1^2, Z_2^2, Z_3^2)$  in the following table, where we replace  $Z$  by either  $X$  or  $Y$ . We pair off these bases in a convenient way.

$i$	$a_i$	$b_i$
1	1	$Y_1 Y_2 Y_3$
2	$X_1$	$Y_2 Y_3$
3	$X_2$	$Y_1 Y_3$
4	$X_3$	$Y_1 Y_2$
5	$X_1 X_2$	$Y_3$
6	$X_1 X_3$	$Y_2$
7	$X_2 X_3$	$Y_1$
8	$X_1 X_2 X_3$	1

The Bézoutian we computed is in the desired form  $\sum_{i=1}^8 a_i \otimes b_i$ , so we now need to compute the dualizing linear form  $\lambda$  for  $\{a_i\}$  and  $\{b_i\}$ . Since  $1 = 1 \cdot b_8 + \sum_{i=1}^7 0 \cdot b_i$ , we define  $\lambda$  by  $\lambda(a_i) = 0$  for  $1 \leq i \leq 7$  and  $\lambda(a_8) = \lambda(X_1 X_2 X_3) = 1$ . Now let  $g \in k[X_1, X_2, X_3]/(X_1^2, X_2^2, X_3^2)$  be arbitrary. We can write  $g$  as

$$g = c_1 + c_2 X_1 + c_3 X_2 + c_4 X_3 + c_5 X_1 X_2 + c_6 X_1 X_3 + c_7 X_2 X_3 + c_8 X_1 X_2 X_3.$$

Then  $\lambda$  is the dualizing linear form sending

$$\begin{aligned} \lambda : \frac{k[X_1, X_2, X_3]}{(X_1^2, X_2^2, X_3^2)} &\rightarrow k \\ g &\mapsto c_8. \end{aligned}$$

Finally we can compute the Gram matrix of  $\Phi_\lambda$  in the basis  $\{a_i\}$ . Note that  $a_i a_j$  is a scalar multiple of  $X_1 X_2 X_3$  if and only if  $i + j - 1 = 8$ . Thus the Gram matrix is

$$\Phi_\lambda = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \cong \bigoplus_{i=1}^4 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

#### 4. The Scheja–Storch bilinear form

Associated to any polynomial with an isolated zero, Eisenbud and Levine [11] and Khimshiashvili [13] used the Scheja–Storch construction [30] to produce a bilinear form on the local algebra  $Q_m$ . Kass and Wickelgren proved that this Eisenbud–Khimshiashvili–Levine bilinear form computes the local  $\mathbb{A}^1$ -degree [16]. The machinery of Scheja and Storch works in great generality; in particular, one may produce a Scheja–Storch bilinear form on the global algebra  $Q$  as well as the local algebras  $Q_m$ . We will provide a brief account of the Scheja–Storch construction before comparing it with the Bézoutian.

In [30],  $k\langle X \rangle := k\langle X_1, \dots, X_n \rangle$  denotes either a polynomial ring  $k[X_1, \dots, X_n]$  or a power series ring  $k[[X_1, \dots, X_n]]$ . We will also use this notation, although we will focus on the situation where  $k\langle X \rangle$  is a polynomial ring. Let  $\rho : k\langle X \rangle \rightarrow Q$  denote the map obtained by quotienting out by the ideal  $(f_1, \dots, f_n)$ , let  $\mu_1 : k\langle X \rangle \otimes_k k\langle X \rangle \rightarrow k\langle X \rangle$  denote the multiplication map, and let  $\mu : Q \otimes_k Q \rightarrow Q$  denote the multiplication map on the global algebra, fitting into a commutative diagram:

$$\begin{array}{ccc} k\langle X \rangle \otimes_k k\langle X \rangle & \xrightarrow{\mu_1} & k\langle X \rangle \\ \rho \otimes \rho \downarrow & & \downarrow \rho \\ Q \otimes_k Q & \xrightarrow{\mu} & Q \end{array}$$

We remark that  $f_j \otimes 1 - 1 \otimes f_j$  lies in  $\ker(\mu_1)$ , and that  $\ker(\mu_1)$  is generated by elements of the form  $X_i \otimes 1 - 1 \otimes X_i$ . Thus for any  $j$ , there are elements  $a_{ij} \in k\langle X \rangle \otimes_k k\langle X \rangle$  such that

$$f_j \otimes 1 - 1 \otimes f_j = \sum_{i=1}^n a_{ij} (X_i \otimes 1 - 1 \otimes X_i). \tag{4-1}$$

We denote by  $\Delta$  the following distinguished element in the tensor algebra  $Q \otimes_k Q$

$$\Delta := (\rho \otimes \rho)(\det(a_{ij})),$$

which corresponds to the Bézoutian which we will later demonstrate. It is true that  $\Delta$  is independent of the choice of  $a_{ij}$ , as shown by Scheja and Storch [30, 3.1 Satz]. We now define an important isomorphism

$\chi$  of  $k$ -algebras used in the Scheja–Storch construction. However, we will phrase this more categorically than in [30], as it will benefit us later.

**Proposition 4.1.** *Consider two endofunctors  $F, G : \text{Alg}_k^{f.g.} \rightarrow \text{Alg}_k^{f.g.}$  on the category of finitely generated  $k$ -algebras, where  $F(A) = A \otimes_k A$  and  $G(A) = \text{Hom}_k(\text{Hom}_k(A, k), A)$ . Then there is a natural isomorphism  $\chi : F \rightarrow G$  whose component at a  $k$ -algebra  $A$  is*

$$\begin{aligned} \chi_A : A \otimes_k A &\rightarrow \text{Hom}_k(\text{Hom}_k(A, k), A) \\ b \otimes c &\mapsto [\varphi \mapsto \varphi(b)c]. \end{aligned}$$

*Proof.* This canonical isomorphism is given in [30, page 181], so it will suffice for us to verify naturality. Let  $g : A \rightarrow B$  be any morphism of  $k$ -algebras. Consider the induced maps  $g \otimes g : A \otimes_k A \rightarrow B \otimes_k B$  and

$$\begin{aligned} g_* : \text{Hom}_k(\text{Hom}_k(A, k), A) &\rightarrow \text{Hom}_k(\text{Hom}_k(B, k), B) \\ \psi &\mapsto [\epsilon \mapsto g \circ \psi(\epsilon \circ g)]. \end{aligned}$$

It remains to show that the following diagram commutes:

$$\begin{array}{ccc} A \otimes_k A & \xrightarrow{\chi_A} & \text{Hom}_k(\text{Hom}_k(A, k), A) \\ g \otimes g \downarrow & & \downarrow g_* \\ B \otimes_k B & \xrightarrow{\chi_B} & \text{Hom}_k(\text{Hom}_k(B, k), B) \end{array}$$

To see this, we compute  $g_* \circ \chi_A = [b \otimes c \mapsto [\epsilon \mapsto g((\epsilon \circ g)(b) \cdot c)]]$ . Note that  $\epsilon \circ g : B \rightarrow k$ , so  $(\epsilon \circ g)(b) \in k$ . Since  $g$  is  $k$ -linear, we have  $g((\epsilon \circ g)(b) \cdot c) = \epsilon(g(b)) \cdot g(c)$ . Next, we compute  $\chi_B \circ (g \otimes g) = [b \otimes c \mapsto [\epsilon \mapsto \epsilon(g(b)) \cdot g(c)]]$ . Thus  $g_* \circ \chi_A = \chi_B \circ (g \otimes g)$ , so the diagram commutes.  $\square$

We now let  $\Theta := \chi_Q(\Delta)$  denote the image of  $\Delta$  under the component of this natural isomorphism at the global algebra  $Q$ . We have that  $\Theta$  is a  $k$ -linear map  $\Theta : \text{Hom}_k(Q, k) \rightarrow Q$ . Letting  $\eta$  denote  $\Theta^{-1}(1)$ , we obtain a well-defined linear form  $\eta : Q \rightarrow k$  by [30, 3.3 Satz].

**Definition 4.2.** We refer to  $\Phi_\eta : Q \times Q \rightarrow k$  as the *global Scheja–Storch bilinear form*.

The Bézoutian gives us an explicit formula for  $\Delta$ . As a result, the global Scheja–Storch form agrees with the Bézoutian form.

**Proposition 4.3.** *In  $Q \otimes_k Q$ , we have  $\Delta = \text{Béz}(f_1, \dots, f_n)$ .*

*Proof.* We first compute

$$\begin{aligned} \sum_{i=1}^n \Delta_{ji}(X_i - Y_i) &= \sum_{i=1}^n \frac{f_j(Y_1, \dots, Y_{i-1}, X_i, \dots, X_n) - f_j(Y_1, \dots, Y_i, X_{i+1}, \dots, X_n)}{(X_i - Y_i)} \cdot (X_i - Y_i) \\ &= \sum_{i=1}^n f_j(Y_1, \dots, Y_{i-1}, X_i, \dots, X_n) - f_j(Y_1, \dots, Y_i, X_{i+1}, \dots, X_n) \\ &= f_j(X_1, \dots, X_n) - f_j(Y_1, \dots, Y_n). \end{aligned}$$

Let  $\varphi : k\langle X \rangle \otimes_k k\langle X \rangle \xrightarrow{\sim} k\langle X, Y \rangle$  be the ring isomorphism given by  $\varphi(b \otimes c) = b(X)c(Y)$ . Note that  $\varphi(x_i \otimes 1) = X_i$  and  $\varphi(1 \otimes x_i) = Y_i$ , so the inverse of  $\varphi$  is characterized by  $\varphi^{-1}(X_i) = x_i \otimes 1$  and  $\varphi^{-1}(Y_i) = 1 \otimes x_i$ . It follows that

$$f_j \otimes 1 - 1 \otimes f_j = \varphi^{-1}(f_j(X) - f_j(Y)) = \sum_{i=1}^n \varphi^{-1}(\Delta_{ji}(X_i - Y_i)) = \sum_{i=1}^n \varphi^{-1}(\Delta_{ji})(x_i \otimes 1 - 1 \otimes x_i).$$

We may thus set  $a_{ij} = \varphi^{-1}(\Delta_{ji})$ , and [30, 3.1 Satz] implies that  $\Delta = (\rho \otimes \rho)(\det(a_{ij}))$ . On the other hand,  $(\rho \otimes \rho)(\varphi^{-1}(\det(\Delta_{ji}))) = \text{Béz}(f_1, \dots, f_n)$  by Definition 3.1.  $\square$

**Lemma 4.4.** *The Bézoutian bilinear form and the global Scheja–Storch bilinear form are identical.*

*Proof.* We showed in Proposition 4.3 that  $\Delta$  is the Bézoutian in  $Q \otimes_k Q$ . We now show that the associated forms are identical. Pick bases  $\{a_i\}$  and  $\{b_i\}$  of  $Q$  such that

$$\Delta = \text{Béz}(f_1, \dots, f_n) = \sum_{i=1}^m a_i \otimes b_i.$$

Since the natural isomorphism  $\chi$  has  $k$ -linear components,  $\Delta$  is mapped to

$$\Theta := \chi_Q(\Delta) = \left[ \varphi \mapsto \sum_{i=1}^m \varphi(a_i)b_i \right].$$

Thus  $\eta := \Theta^{-1}(1)$  is the linear form  $\eta : Q \rightarrow k$  satisfying  $\sum_{i=1}^m \eta(a_i)b_i = 1$ . By Proposition 3.5, this implies that  $\eta$  is the form for which  $\{a_i\}$  and  $\{b_i\}$  are dual bases. As in Definition 3.8, this tells us that  $\eta$  is the linear form producing the Bézoutian bilinear form.  $\square$

**4A. Local decomposition.** While our discussion of the Scheja–Storch form in the previous section was global, it is perfectly valid to localize at a maximal ideal and repeat the story again [30, pages 180–181]. The fact that  $Q$  is an Artinian ring then gives a convenient way to relate the global version of  $\eta$  to the local version of  $\eta$ . This local decomposition has been utilized previously, for example in [16].

Let  $\mathfrak{m}$  be a maximal ideal in  $k[x_1, \dots, x_n]$  at which the morphism  $f = (f_1, \dots, f_n)$  has an isolated root. Letting  $\rho_{\mathfrak{m}}$  denote the quotient map  $k\langle X \rangle_{\mathfrak{m}} \rightarrow Q_{\mathfrak{m}}$ , we have a commutative diagram:

$$\begin{array}{ccc} k\langle X \rangle_{\mathfrak{m}} \otimes_k k\langle X \rangle_{\mathfrak{m}} & \xrightarrow{\mu'_1} & k\langle X \rangle_{\mathfrak{m}} \\ \rho_{\mathfrak{m}} \otimes \rho_{\mathfrak{m}} \downarrow & & \downarrow \rho_{\mathfrak{m}} \\ Q_{\mathfrak{m}} \otimes_k Q_{\mathfrak{m}} & \xrightarrow{\mu'} & Q_{\mathfrak{m}} \end{array}$$

In  $k\langle X \rangle_{\mathfrak{m}} \otimes_k k\langle X \rangle_{\mathfrak{m}}$ , we can again write

$$f_j \otimes 1 - 1 \otimes f_j = \sum_{i=1}^n \tilde{a}_{ij}(X_i \otimes 1 - 1 \otimes X_i)$$

to obtain the local Bézoutian  $\Delta_m := (\rho_m \otimes \rho_m)(\det(\tilde{a}_{ij})) \in Q_m \otimes_k Q_m$ . Let  $\lambda_m : Q \rightarrow Q_m$  be the localization map. From [30, page 181] we have  $(\lambda_m \otimes \lambda_m)(\Delta) = \Delta_m$ . Via the natural isomorphism  $\chi$  in Proposition 4.1, we have a commutative diagram of the form:

$$\begin{array}{ccc} Q \otimes_k Q & \xrightarrow{\chi_Q} & \text{Hom}_k(\text{Hom}_k(Q, k), Q) \\ \lambda_m \otimes \lambda_m \downarrow & & \downarrow \lambda_{m*} \\ Q_m \otimes_k Q_m & \xrightarrow{\chi_{Q_m}} & \text{Hom}_k(\text{Hom}_k(Q_m, k), Q_m) \end{array}$$

Tracing  $\Delta$  through this diagram, we see that

$$\begin{array}{ccc} \Delta & \longmapsto & \Theta \\ \downarrow & & \downarrow \\ \Delta_m & \longmapsto & \Theta_m \end{array}$$

where  $\Theta_m = \chi_{Q_m}(\Delta_m)$ . Unwinding  $\Theta_m = \lambda_{m*}(\Theta)$ , we find that  $\Theta_m$  is the map

$$\begin{aligned} \Theta_m : \text{Hom}_k(Q_m, k) &\rightarrow Q_m \\ \psi &\mapsto \lambda_m \circ \Theta(\psi \circ \lambda_m). \end{aligned}$$

Recall that as  $Q$  is a zero-dimensional Noetherian commutative  $k$ -algebra, the localization maps induce a  $k$ -algebra isomorphism:<sup>1</sup>

$$(\lambda_m)_m : Q \xrightarrow{\sim} \prod_m Q_m.$$

This is reflected by an internal decomposition of  $Q$  in terms of orthogonal idempotents [4, 2.13], which we now describe; see also [32, Lemma 00JA]. By the Chinese remainder theorem, we may pick a collection of pairwise orthogonal idempotents  $\{e_m\}_m$  such that  $\sum_m e_m = 1$ . The internal decomposition of  $Q$  is then

$$Q = \bigoplus_m Q \cdot e_m,$$

and the localization maps restrict to isomorphisms  $\lambda_m|_{Q \cdot e_m} : Q \cdot e_m \xrightarrow{\sim} Q_m$  with  $\lambda_m(e_m) = 1$ . Moreover,  $\lambda_m(Q \cdot e_n) = 0$  for any  $n \neq m$ .

**Proposition 4.5.** *Suppose  $\ell : Q \rightarrow k$  is a linear form which factors through the localization  $\lambda_m : Q \rightarrow Q_m$  for some maximal ideal  $\mathfrak{m}$ . Then  $\Theta(\ell)$  lies in  $Q \cdot e_m$ .*

*Proof.* Recall that  $\lambda_m|_{Q \cdot e_n} = 0$  for  $n \neq m$ . Since  $e_m \cdot e_n = 0$  for  $n \neq m$  and  $e_m$  is idempotent, the localization  $\lambda_m : Q \rightarrow Q_m$  can be written as the following composition:

$$\lambda_m : Q \xrightarrow{-e_m} Q \xrightarrow{\lambda_m} Q_m.$$

Since  $\ell$  factors through the localization, it can be written as a composite

$$\ell : Q \xrightarrow{-e_m} Q \xrightarrow{\lambda_m} Q_m \xrightarrow{\ell_m} k.$$

<sup>1</sup> $Q$  is Artinian by [32, Lemma 00KH], so the claimed isomorphism exists by [32, Lemma 00JA].

Thus  $\Theta(\ell) = \Theta(\ell_m \circ \lambda_m \circ (e_m \cdot -))$ . Scheja–Storch proved that  $\Theta$  respects the  $Q$ -module structure on  $\text{Hom}_k(Q, k)$  given by  $a \cdot \sigma = \sigma(a \cdot -)$  [30, 3.3 Satz]. That is,  $\Theta(\sigma(a \cdot -)) = \Theta(a \cdot \sigma) = a\Theta(\sigma)$  for any  $a \in Q$  and  $\sigma \in \text{Hom}_k(Q, k)$ . Thus

$$\Theta(\ell) = e_m \cdot \Theta(\ell_m \circ \lambda_m),$$

so  $\Theta(\ell) \in Q \cdot e_m$ . □

Returning to the Scheja–Storch form, we have the following commutative diagram relating  $\Theta_m$  and  $\Theta$ :

$$\begin{array}{ccc} \text{Hom}_k(Q, k) & \xrightarrow{\Theta} & Q \\ \uparrow -\circ\lambda_m & & \downarrow \lambda_m \\ \text{Hom}_k(Q_m, k) & \xrightarrow{\Theta_m} & Q_m \end{array}$$

This coherence between  $\Theta$  and  $\Theta_m$  allows us to relate the local linear forms  $\eta_m := \Theta_m^{-1}(1)$  to the global linear form  $\eta := \Theta^{-1}(1)$  in the following way.

**Proposition 4.6.** *For each maximal ideal  $\mathfrak{m}$  of  $Q$ , let  $\eta_m := \Theta_m^{-1}(1) : Q_m \rightarrow k$ , and let  $\eta := \Theta^{-1}(1) : Q \rightarrow k$ . Then  $\eta = \sum_{\mathfrak{m}} \eta_m \circ \lambda_m$ .*

*Proof.* It suffices to show that  $\Theta(\sum_{\mathfrak{m}} \eta_m \circ \lambda_m) = 1$ . Since  $\eta_m = \Theta_m^{-1}(1)$  by definition, we have  $1 = \Theta_m(\eta_m) := \lambda_m(\Theta(\eta_m \circ \lambda_m))$ . By Proposition 4.5, we have  $\Theta(\eta_m \circ \lambda_m) \in Q \cdot e_m$ . Since  $\lambda_m(\Theta(\eta_m \circ \lambda_m)) = 1$  and  $\lambda_m|_{Q \cdot e_m}$  is an isomorphism sending  $e_m$  to 1, it follows that  $\Theta(\eta_m \circ \lambda_m) = e_m$ . Finally, since  $\Theta$  is  $k$ -linear, we have

$$\Theta\left(\sum_{\mathfrak{m}} \eta_m \circ \lambda_m\right) = \sum_{\mathfrak{m}} \Theta(\eta_m \circ \lambda_m) = \sum_{\mathfrak{m}} e_m = 1. \quad \square$$

Using this local decomposition procedure for the linear forms  $\eta_m$  and  $\eta$ , we obtain a local decomposition for Scheja–Storch bilinear forms.

**Lemma 4.7. (Local decomposition of Scheja–Storch forms)** *Let  $\eta$  and  $\eta_m$  be as in Proposition 4.6. Then  $\Phi_\eta = \bigoplus_{\mathfrak{m}} \Phi_{\eta_m}$ . In particular, the global Scheja–Storch form is a sum over local Scheja–Storch forms*

$$\text{SS}(f) = \sum_{\mathfrak{m}} \text{SS}_{\mathfrak{m}}(f).$$

*Proof.* For each maximal ideal  $\mathfrak{m}$ , let  $\{w_{\mathfrak{m},i}\}_i$  be a  $k$ -vector space basis for  $Q_m$ . Let  $\{v_{\mathfrak{m},i}\}_{\mathfrak{m},i}$  (ranging over all  $i$  and all maximal ideals) be a basis of  $Q$  such that  $\lambda_{\mathfrak{m}}(v_{\mathfrak{m},i}) = w_{\mathfrak{m},i}$  for each  $i$  and  $\mathfrak{m}$ , and  $\lambda_{\mathfrak{m}}(v_{\mathfrak{n},i}) = 0$  for  $\mathfrak{m} \neq \mathfrak{n}$ . We now compare the Gram matrix for  $\eta : Q \rightarrow k$  and the Gram matrices for  $\eta_m : Q_m \rightarrow k$  in these bases. Via the internal decomposition consisting of pairwise orthogonal idempotents, we have  $v_{\mathfrak{m},i} \cdot v_{\mathfrak{n},j} = 0$  if  $\mathfrak{m} \neq \mathfrak{n}$ . Thus

$$\eta(v_{\mathfrak{m},i} \cdot v_{\mathfrak{n},j}) = 0,$$

so the Gram matrix for  $\Phi_\eta$  will be a block sum indexed over the maximal ideals. If  $m = n$ , then Proposition 4.6 implies

$$\eta(v_{m,i} \cdot v_{m,j}) = \sum_n \eta_n(\lambda_n(v_{m,i} \cdot v_{m,j})) = \eta_m(\lambda_m(v_{m,i} \cdot v_{m,j})) = \eta_m(w_{m,i} \cdot w_{m,j}).$$

Thus the Gram matrices of  $\Phi_\eta$  and  $\bigoplus_m \Phi_{\eta_m}$  are equal, so  $\Phi_\eta = \bigoplus_m \Phi_{\eta_m}$ . □

**Remark 4.8.** The local Scheja–Storch bilinear form is given by  $\Phi_{\eta_m} : Q_m \times Q_m \rightarrow k$ . Given a basis  $\{a_1, \dots, a_m\}$  of  $Q_m$ , we may write  $\Delta_m = \sum a_i \otimes b_i$  and define the local Bézoutian bilinear form as a suitable dualizing form. Replacing  $Q, \Delta, \Theta,$  and  $\eta$  with  $Q_m, \Delta_m, \Theta_m,$  and  $\eta_m$ , the results of Sections 3 and 4 also hold for local Bézoutians and the local Scheja–Storch form. In particular, the local analog of Lemma 4.4 implies that the local Scheja–Storch form is equal to the local Bézoutian form.

### 5. Proof of Theorem 1.2

We now relate the Scheja–Storch form to the  $\mathbb{A}^1$ -degree. The following theorem was first proven in the case where  $p$  is a rational zero by Kass and Wickelgren [16], and then in the case where  $p$  has finite separable residue field over the ground field in [7, Corollary 1.4]. Recent work of Bachmann and Wickelgren [2] gives a general result about the relation between local  $\mathbb{A}^1$ -degrees and Scheja–Storch forms.

**Theorem 5.1.** *Let  $\text{char } k \neq 2$ . Let  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  be an endomorphism of affine space with an isolated zero at a closed point  $p$ . Then we have that the local  $\mathbb{A}^1$ -degree of  $f$  at  $p$  and the Scheja–Storch form of  $f$  at  $p$  coincide as elements of  $\text{GW}(k)$ :*

$$\text{deg}_p^{\mathbb{A}^1}(f) = \text{SS}_p(f).$$

*Proof.* We may rewrite  $f$  as a section of the trivial rank  $n$  bundle over affine space  $\mathcal{O}_{\mathbb{A}_k^n}^n \rightarrow \mathbb{A}_k^n$ . Under the hypothesis that  $p$  is isolated, we may find a neighborhood  $X \subseteq \mathbb{A}_k^n$  of  $p$  where the section  $f$  is nondegenerate (meaning it is cut out by a regular sequence). By [2, Corollary 8.2], the local index of  $f$  at  $p$  with the trivial orientation, corresponding to the representable Hermitian  $K$ -theory spectrum  $\text{KO}$ , agrees with the local Scheja–Storch form as elements of  $\text{KO}^0(k)$ :

$$\text{ind}_p(f, \rho_{\text{triv}}, \text{KO}) = \text{SS}_p(f). \tag{5-1}$$

Let  $\mathbb{S}$  denote the sphere spectrum in the stable motivic homotopy category  $\mathcal{SH}(k)$ . It is a well-known fact that Hermitian  $K$ -theory receives a map from the sphere spectrum, inducing an isomorphism  $\pi_0(\mathbb{S}) \xrightarrow{\sim} \pi_0(\text{KO})$  if  $\text{char } k \neq 2$  (see for example [14, 6.9] for more detail); this is the only place where we use the assumption that  $\text{char } k \neq 2$ . Combining this with the fact that  $\pi_0(\mathbb{S}) = \text{GW}(k)$  under Morel’s degree isomorphism, we observe that (5-1) is really an equality in  $\text{GW}(k)$ . By [2, Theorem 7.6, Example 7.7], the local index associated to the representable theory agrees with the local  $\mathbb{A}^1$ -degree:

$$\text{ind}_p(f, \rho_{\text{triv}}, \text{KO}) = \text{deg}_p^{\mathbb{A}^1}(f).$$

Combining these equalities gives the desired equality in  $\text{GW}(k)$ . □



**Remark 5.2.** Bachmann and Wickelgren in fact show that  $\deg_Z^{\mathbb{A}^1}(f) = \text{SS}_Z(f)$  for any isolated zero locus  $Z$  of  $f$  [2, Corollary 8.2]. This gives an alternate viewpoint on the local decomposition described in Lemma 4.7

**Corollary 5.3.** *Let  $\text{char } k \neq 2$ . The local Bézoutian bilinear form is the local  $\mathbb{A}^1$ -degree.*

*Proof.* As discussed in Remark 4.8, we can modify Lemma 4.4 to the local case by replacing  $Q, \Delta, \Theta,$  and  $\eta$  with  $Q_m, \Delta_m, \Theta_m,$  and  $\eta_m$ . The local Bézoutian form is thus equal to the local Scheja–Storch form, which is equal to the local  $\mathbb{A}^1$ -degree by Theorem 5.1.  $\square$

In contrast to previous techniques for computing the local  $\mathbb{A}^1$ -degree at rational or separable points, Corollary 5.3 gives an algebraic formula for the local  $\mathbb{A}^1$ -degree at any closed point.

As a result of the local decomposition of Scheja–Storch forms, the Bézoutian form agrees with the  $\mathbb{A}^1$ -degree globally as well.

**Corollary 5.4.** *Let  $\text{char } k \neq 2$ . The Bézoutian bilinear form is the global  $\mathbb{A}^1$ -degree.*

*Proof.* Let  $\Phi_\eta$  denote the Bézoutian bilinear form, which is equal to the global Scheja–Storch bilinear form by Lemma 4.4. By Lemma 4.7, the global Scheja–Storch form decomposes as a block sum of local Scheja–Storch forms. By Theorem 5.1, the local Scheja–Storch bilinear form agrees with the local  $\mathbb{A}^1$ -degree. Finally, we have that the sum of local  $\mathbb{A}^1$ -degrees is the global  $\mathbb{A}^1$ -degree. Putting this all together, we have

$$\Phi_\eta = \text{SS}(f) = \sum_m \text{SS}_m(f) = \sum_m \deg_m^{\mathbb{A}^1}(f) = \deg^{\mathbb{A}^1}(f). \quad \square$$

**Remark 5.5.** It is not known if GW is represented by KO over fields of characteristic 2, which is the source of our assumption that  $\text{char } k \neq 2$ . If this problem is resolved, one can remove any characteristic restrictions from our results. Alternately, Lemma 4.7 implies Corollaries 5.3 and 5.4 if all roots of  $f$  satisfy  $\deg_p^{\mathbb{A}^1}(f) = \text{SS}_p(f)$ . By [16], [7], and [17, Proposition 34], Corollaries 5.3 and 5.4 are true in any characteristic if all roots of  $f$  are rational, étale, or separable.

**5A. Computing the Bézoutian bilinear form.** We now prove Theorem 1.2 by describing a method for computing the class in  $\text{GW}(k)$  of the Bézoutian bilinear form in terms of the Bézoutian.

*Proof of Theorem 1.2.* Let  $R$  denote either a global algebra  $Q$  or a local algebra  $Q_m$ . Let  $\{\alpha_i\}$  be any basis for  $R$ , and express

$$\text{Béz}(f_1, \dots, f_n) = \sum_{i,j} B_{i,j} \alpha_i \otimes \alpha_j. \quad (5-2)$$

Rewriting this, we have

$$\text{Béz}(f_1, \dots, f_n) = \sum_i \alpha_i \otimes \left( \sum_j B_{i,j} \alpha_j \right).$$

Let  $\beta_i := \sum_j B_{i,j} \alpha_j$ , so that  $\{\alpha_i\}$  and  $\{\beta_i\}$  are dual bases. Then for any linear form  $\lambda : R \rightarrow k$  for which  $\{\alpha_i\}$  and  $\{\beta_i\}$  are dual, we will have that  $\Phi_\lambda$  agrees with the global or local  $\mathbb{A}^1$ -degree (depending on our

choice of  $R$ ) by Corollaries 5.3 and 5.4. Let  $\lambda$  be such a form. The product of  $\alpha_i$  and  $\beta_j$  is given by

$$\alpha_i \beta_j = \alpha_i \cdot \sum_s B_{j,s} \alpha_s.$$

Applying  $\lambda$  to each side, we get an indicator function

$$\delta_{ij} = \lambda(\alpha_i \beta_j) = \lambda\left(\alpha_i \sum_s B_{j,s} \alpha_s\right) = \sum_s B_{j,s} \lambda(\alpha_i \alpha_s).$$

Varying over all  $i, j, s$ , this equation above tells us that the identity matrix is equal to the product of the matrix  $(B_{j,s})$  and the matrix  $(\lambda(\alpha_i \alpha_s)) = (\lambda(\alpha_s \alpha_i))$ . Explicitly, we have that

$$\begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix} = \begin{pmatrix} B_{1,1} & B_{1,2} & \cdots & B_{1,m} \\ B_{2,1} & B_{2,2} & \cdots & B_{2,m} \\ \vdots & \vdots & \ddots & \vdots \\ B_{m,1} & B_{m,2} & \cdots & B_{m,m} \end{pmatrix} \begin{pmatrix} \lambda(\alpha_1^2) & \lambda(\alpha_1 \alpha_2) & \cdots & \lambda(\alpha_1 \alpha_m) \\ \lambda(\alpha_2 \alpha_1) & \lambda(\alpha_2^2) & \cdots & \lambda(\alpha_2 \alpha_m) \\ \vdots & \vdots & \ddots & \vdots \\ \lambda(\alpha_m \alpha_1) & \lambda(\alpha_m \alpha_2) & \cdots & \lambda(\alpha_m^2) \end{pmatrix}.$$

Thus the Gram matrix for  $\Phi_\lambda$  in the basis  $\{\alpha_i\}$  is  $(B_{i,j})^{-1}$ . We conclude by proving that  $(B_{i,j})$  and  $(B_{i,j})^{-1}$  represent the same element of  $\text{GW}(k)$ . Since any symmetric bilinear form can be diagonalized, there is an invertible  $m \times m$  matrix  $S$  such that  $S^T \cdot (B_{i,j}) \cdot S$  is diagonal. Since  $(S^T \cdot (B_{i,j}) \cdot S) \cdot (S^{-1} \cdot (B_{i,j})^{-1} \cdot (S^{-1})^T)$  is equal to the identity matrix, it follows that  $S^{-1} \cdot (\lambda(\alpha_i \alpha_j)) \cdot (S^{-1})^T$  is diagonal with entries inverse to the diagonal entries of  $S^T \cdot (B_{i,j}) \cdot S$ . Applying the equality  $\langle a \rangle = \langle 1/a \rangle$  along the diagonals, it follows that  $(B_{i,j})^{-1}$  and  $(B_{i,j})$  define the same element in  $\text{GW}(k)$ . Theorem 1.2 now follows from Corollaries 5.3 and 5.4.  $\square$

The following tables describe algorithms for computing the global and local  $\mathbb{A}^1$ -degrees in terms of the Bézoutian bilinear form. A Sage implementation of these algorithms is available at [8].

### Computing the global $\mathbb{A}^1$ -degree via the Bézoutian

- (1) Compute the  $\Delta_{ij}$  and the image of their determinant  $\text{Béz}(f) = \det(\Delta_{ij})$  in  $k[X, Y]/(f(X), f(Y))$ .
- (2) Pick a  $k$ -vector space basis  $a_1, \dots, a_m$  of  $Q = k[X_1, \dots, X_n]/(f_1, \dots, f_n)$ . Find  $B_{i,j} \in k$  such that

$$\text{Béz}(f) = \sum_{i=1}^m B_{i,j} a_i(X) a_j(Y).$$

- (3) The matrix  $B = (B_{i,j})$  represents  $\text{deg}^{\mathbb{A}^1}(f)$ . Diagonalize  $B$  to write its class in  $\text{GW}(k)$ .

### Computing the local $\mathbb{A}^1$ -degree via the Bézoutian

- (1) Compute the  $\Delta_{ij}$  and the image of their determinant  $\text{Béz}(f) = \det(\Delta_{ij})$  in  $k[X, Y]/(f(X), f(Y))$ .
- (2) Pick a  $k$ -vector space basis  $a_1, \dots, a_m$  of  $Q_m = k[X_1, \dots, X_n]_m/(f_1, \dots, f_n)$ . Find  $B_{i,j} \in k$  such that

$$\text{Béz}(f) = \sum_{i=1}^m B_{i,j} a_i(X) a_j(Y).$$

- (3) The matrix  $B = (B_{i,j})$  represents  $\text{deg}_m^{\mathbb{A}^1}(f)$ . Diagonalize  $B$  to write its class in  $\text{GW}(k)$ .

### 6. Calculation rules

Using the Bézoutian characterization of the  $\mathbb{A}^1$ -degree, we are able to establish various calculation rules for local and global  $\mathbb{A}^1$ -degrees. See [18; 29] for related results in the local case.

Our ultimate goal in this section is the product rule for the  $\mathbb{A}^1$ -degree (see Proposition 6.5), which was already known by the work of Morel. See the paragraph preceding Proposition 6.5 for a more detailed discussion.

**Proposition 6.1.** *Suppose that  $f = (f_1, \dots, f_n)$  and  $g = (g_1, \dots, g_n)$  are endomorphisms of affine space that generate the same ideal*

$$I = (f_1, \dots, f_n) = (g_1, \dots, g_n) \triangleleft k[x_1, \dots, x_n].$$

*If  $\text{Béz}(f) = \text{Béz}(g)$  in  $k[X, Y]$ , then  $\text{deg}^{\mathbb{A}^1}(f) = \text{deg}^{\mathbb{A}^1}(g)$ , and  $\text{deg}_p^{\mathbb{A}^1}(f) = \text{deg}_p^{\mathbb{A}^1}(g)$  for all  $p$ .*

*Proof.* We may choose the same basis for  $Q = k[x_1, \dots, x_n]/I$  (or  $Q_p$  in the local case) in our computation for the degrees of  $f$  and  $g$ . The Bézoutians  $\text{Béz}(f) = \text{Béz}(g)$  will have the same coefficients in this basis, so their Gram matrices will coincide. □

The following result is the global analogue of [29, Lemma 14].

**Lemma 6.2.** *Let  $f = (f_1, \dots, f_n) : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  be an endomorphism of  $\mathbb{A}_k^n$  with only isolated zeros. Let  $A \in k^{n \times n}$  be an invertible matrix. Then*

$$\langle \det A \rangle \cdot \text{deg}^{\mathbb{A}^1}(f) = \text{deg}^{\mathbb{A}^1}(A \circ f)$$

*as elements of  $\text{GW}(k)$ .*

*Proof.* Write  $A = (a_{ij})$  and

$$\Delta_{ij}^g = \frac{g_i(X_1, \dots, X_j, Y_{j+1}, \dots, Y_n) - g_i(X_1, \dots, X_{j-1}, Y_j, \dots, Y_n)}{X_j - Y_j},$$

where  $g$  is either  $f$  or  $A \circ f$ . Then  $\Delta_{ij}^{A \circ f} = \sum_{l=1}^n a_{il} \Delta_{lj}^f$ , and thus  $(\Delta_{ij}^{A \circ f}) = A \cdot (\Delta_{ij}^f)$  as matrices over  $k[X, Y]$ . The ideals generated by  $A \circ (f_1, \dots, f_n)$  and  $(f_1, \dots, f_n)$  are equal, and the images in  $Q \otimes_k Q$  of  $\det(\Delta_{ij}^{A \circ f})$  and  $\det A \cdot \det(\Delta_{ij}^f)$  are equal. Thus the Gram matrix of the Bézoutian bilinear form for  $A \circ f$  is  $\det A$  times the Gram matrix of the Bézoutian bilinear form for  $f$ . Proposition 6.1 then proves the claim. □

**Example 6.3.** We may apply Lemma 6.2 in the case where  $A$  is a permutation matrix associated to some permutation  $\sigma \in \Sigma_n$ . Letting  $f_\sigma := (f_{\sigma(1)}, \dots, f_{\sigma(n)})$ , we observe that

$$\text{deg}_p^{\mathbb{A}^1}(f_\sigma) = \langle \text{sgn}(\sigma) \rangle \cdot \text{deg}_p^{\mathbb{A}^1}(f)$$

at any isolated zero  $p$  of  $f$ , and an analogous statement is true for global degrees as well.

Next, we prove a lemma inspired by [18, Lemma 12].

**Lemma 6.4.** *Let  $f, g : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  be two endomorphisms of  $\mathbb{A}_k^n$ . Assume that  $f$  and  $g$  are quasifinite. Let  $L \in M_n(k)$  be an invertible  $n \times n$  matrix, which defines a morphism  $L : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  given by  $(x_1, \dots, x_n) \mapsto (x_1, \dots, x_n) \cdot L^T$ . Let  $I_n$  denote the  $n \times n$  identity matrix, and assume that  $\det(I_n + t(L - I_n)) \in k[t]$  is in fact an element of  $k^\times$ . Then  $\deg^{\mathbb{A}^1}(f \circ g) = \deg^{\mathbb{A}^1}(f \circ L \circ g)$ .*

*Proof.* Quasifinite morphisms have isolated zero loci by [32, Definition 01TD (3)]. The composition of quasifinite morphisms is again quasifinite [32, Lemma 01TL], so  $f \circ g$  has isolated zero locus.

Next, we show that  $L$  is also quasifinite. We will actually prove a stronger statement. Let  $A_t \in M_n(k[t])$  be an invertible  $n \times n$  matrix, which implies that  $\det A_t \in k[t]^\times = k^\times$ . This matrix determines a family of morphisms  $A_t : \mathbb{A}_k^n \times \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^n$  by  $(x_1, \dots, x_n, t) \mapsto (x_1, \dots, x_n) \cdot A_t^T$ . Given  $t_0 \in \mathbb{A}_k^1$ , the morphism  $A_{t_0}$  has Jacobian determinant  $\det\left(\frac{\partial(A_{t_0})_i}{\partial x_j}\right) = \det A_{t_0}$ , which is a unit. In particular,  $A_{t_0}$  is unramified for each  $t_0 \in \mathbb{A}_k^1$ . Thus  $A_{t_0}$  is locally quasifinite [32, Lemma 02VF]. Since  $\mathbb{A}_k^n$  is Noetherian,  $A_{t_0} : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  is quasicompact. Quasicompact and locally quasifinite morphisms are quasifinite [32, Lemma 01TJ], so we conclude that  $A_{t_0}$  is quasifinite for each  $t_0 \in \mathbb{A}_k^1$ .

Just as in [18, Lemma 12], we now define  $L_t = I_n + t \cdot (L - I_n)$ . Our assumption on  $\det(I_n + t(L - I_n))$  implies that  $L_t$  is invertible. Thus  $L_t$  is quasifinite, so  $f \circ L_t \circ g$  is quasifinite and hence only has isolated zeros for all  $t$ . Set

$$\tilde{Q} = \frac{k[t][x_1, \dots, x_n]}{(f \circ L_t \circ g)}.$$

Then [30, page 182] gives us a Scheja–Storch form  $\tilde{\eta} : \tilde{Q} \rightarrow k[t]$  such that the bilinear form  $\Phi_{\tilde{\eta}} : \tilde{Q} \times \tilde{Q} \rightarrow k[t]$  is symmetric and nondegenerate. By Harder’s theorem [16, Lemma 30], the stable isomorphism class of  $\Phi_{\tilde{\eta}} \otimes_k k(t_0) \in \text{GW}(k)$  is independent of  $t_0 \in \mathbb{A}_k^1(k)$ . In particular, the Scheja–Storch bilinear forms of  $f \circ L_0 \circ g = f \circ g$  and  $f \circ L_1 \circ g = f \circ L \circ g$  are isomorphic.  $\square$

The following product rule is a consequence of Morel’s proof that the  $\mathbb{A}^1$ -degree is a ring isomorphism [24, Lemma 6.3.8]. We give a more hands-on proof of this product rule. See [18, Theorem 13] and [29, Theorem 26] for an analogous proof of the product rule for local degrees at rational points.

**Proposition 6.5** (product rule). *Let  $f, g : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  be two quasifinite endomorphisms of  $\mathbb{A}_k^n$ . Then  $\deg^{\mathbb{A}^1}(f \circ g) = \deg^{\mathbb{A}^1}(f) \cdot \deg^{\mathbb{A}^1}(g)$ .*

*Proof.* We follow the proofs of [18, Theorem 13] and [29, Theorem 26]. The general idea is to mimic the Eckmann–Hilton argument [10]. Let  $x := (x_1, \dots, x_n)$  and  $y := (y_1, \dots, y_n)$ . Define  $\tilde{f}, \tilde{g} : \mathbb{A}^n \times \mathbb{A}^n \rightarrow \mathbb{A}^n \times \mathbb{A}^n$  by  $\tilde{f}(x, y) = (f(x), y)$  and  $\tilde{g}(x, y) = (g(x), y)$ , and note that  $\tilde{f}$  and  $\tilde{g}$  are both quasifinite because  $f$  and  $g$  are quasifinite. Since  $(f \circ g, y)$  and  $\tilde{f} \circ \tilde{g}$  define the same ideal in  $k[x, y]$  and have the same Bézoutian, we have  $\deg^{\mathbb{A}^1}(f \circ g) = \deg^{\mathbb{A}^1}(\tilde{f} \circ \tilde{g})$  by Proposition 6.1.

Let  $g \times f : \mathbb{A}_k^n \times \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n \times \mathbb{A}_k^n$  be given by  $(g \times f)(x, y) = (g(x), f(y))$ . Using Lemma 6.4 repeatedly, we will show that  $\deg^{\mathbb{A}^1}(\tilde{f} \circ \tilde{g}) = \deg^{\mathbb{A}^1}(g \times f)$ . Let  $I_n$  be the  $n \times n$  identity matrix, and let

$$L_1 = \begin{pmatrix} I_n & 0 \\ -I_n & I_n \end{pmatrix}, \quad L_2 = \begin{pmatrix} I_n & I_n \\ 0 & I_n \end{pmatrix}, \quad A = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}.$$

By construction,  $\det(I_{2n} + t(L_1 - I_{2n})) = \det(I_{2n} + t(L_2 - I_{2n})) = 1$ , so Lemma 6.4 implies that

$$\deg^{\mathbb{A}^1}(\tilde{f} \circ \tilde{g}) = \deg^{\mathbb{A}^1}(\tilde{f} \circ L_1 \circ \tilde{g}) = \deg^{\mathbb{A}^1}(\tilde{f} \circ L_2 \circ (L_1 \circ \tilde{g})) = \deg^{\mathbb{A}^1}(\tilde{f} \circ L_1 \circ (L_2 \circ L_1 \circ \tilde{g})).$$

One can check that  $A \circ \tilde{f} \circ L_1 \circ L_2 \circ L_1 \circ \tilde{g} = g \times f$ . By Lemma 6.2, we have

$$\langle \det A \rangle \cdot \deg^{\mathbb{A}^1}(\tilde{f} \circ \tilde{g}) = \langle \det A \rangle \cdot \deg^{\mathbb{A}^1}(\tilde{f} \circ L_1 \circ L_2 \circ L_1 \circ \tilde{g}) = \deg^{\mathbb{A}^1}(g \times f).$$

Since  $\det A = 1$ , it just remains to show that  $\deg^{\mathbb{A}^1}(g \times f) = \deg^{\mathbb{A}^1}(g) \cdot \deg^{\mathbb{A}^1}(f)$ . Let  $a_1, \dots, a_m$  be a basis for  $k[x_1, \dots, x_n]/(g_1, \dots, g_n)$  and  $a'_1, \dots, a'_{m'}$  be a basis for  $k[y_1, \dots, y_n]/(f_1, \dots, f_n)$ . Write  $\text{Béz}(g) = \sum_{i,j=1}^m B_{ij} a_i \otimes a_j$  and  $\text{Béz}(f) = \sum_{i',j'=1}^{m'} B'_{i'j'} a'_{i'} \otimes a'_{j'}$ . By Theorem 1.2,  $(B_{ij})$  and  $(B'_{i'j'})$  are the Gram matrices for  $\deg^{\mathbb{A}^1}(g)$  and  $\deg^{\mathbb{A}^1}(f)$ , respectively. Next, we have  $\text{Béz}(g \times f) = \text{Béz}(g) \cdot \text{Béz}(f)$ , since

$$(\Delta_{ij}^{g \times f}) = \begin{pmatrix} (\Delta_{ij}^g) & 0 \\ 0 & (\Delta_{ij}^f) \end{pmatrix}.$$

Note that  $\{a_i(x) a'_{i'}(y)\}_{i,i'=1}^{m,m'}$  is a basis of  $k[x_1, \dots, x_n, y_1, \dots, y_n]/(g_1(x), \dots, g_n(x), f_1(y), \dots, f_n(y))$ . In this basis, we have

$$\text{Béz}(g) \cdot \text{Béz}(f) = \sum_{i,j=1}^m \sum_{i',j'=1}^{m'} B_{ij} B'_{i'j'} a_i a'_{i'} \otimes a_j a'_{j'},$$

so the Gram matrix of  $\deg^{\mathbb{A}^1}(g \times f)$  is the tensor product  $(B_{ij}) \otimes (B'_{i'j'})$ . We thus we have an equality  $\deg^{\mathbb{A}^1}(g \times f) = \deg^{\mathbb{A}^1}(g) \cdot \deg^{\mathbb{A}^1}(f)$  in  $\text{GW}(k)$ .  $\square$

## 7. Examples

We now give a few remarks and examples about computing the Bézoutian.

**Remark 7.1.** It is not always the case that the determinant  $\det(\Delta_{ij}) \in k[X, Y]$  is symmetric. For example, consider the morphism  $f : \mathbb{A}_k^2 \rightarrow \mathbb{A}_k^2$  sending  $(x_1, x_2) \mapsto (x_1 x_2, x_1 + x_2)$ . Then the Bézoutian is given by

$$\text{Béz}(f) = \det \begin{pmatrix} X_2 & Y_1 \\ 1 & 1 \end{pmatrix} = X_2 - Y_1.$$

However, the Bézoutian is symmetric once we pass to the quotient  $k[X, Y]/(f(X), f(Y))$  [4, 2.12]. Continuing the present example, let  $\{1, x_2\}$  be a basis for the algebra  $Q = k[x_1, x_2]/(x_1 x_2, x_1 + x_2)$ . Then we have that

$$\text{Béz}(f) = X_2 - Y_1 = X_2 + Y_2,$$

which is symmetric. Moreover, the Bézoutian bilinear form is represented by  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ , so  $\deg^{\mathbb{A}^1}(f) = \mathbb{H}$ .

**Example 7.2.** Let  $k = \mathbb{F}_p(t)$ , where  $p$  is an odd prime, and consider the endomorphism of the affine plane given by

$$\begin{aligned} f : \text{Spec } \mathbb{F}_p(t)[x_1, x_2] &\rightarrow \text{Spec } \mathbb{F}_p(t)[x_1, x_2] \\ (x_1, x_2) &\mapsto (x_1^p - t, x_1 x_2). \end{aligned}$$

As the residue field of the zero of  $f$  is not separable over  $k$ , existing strategies for computing the local  $\mathbb{A}^1$ -degree are insufficient. Our results allow us to compute this  $\mathbb{A}^1$ -degree. The Bézoutian is given by

$$\begin{aligned} \text{Béz}(f) &= \det \begin{pmatrix} \frac{X_1^p - Y_1^p}{X_1 - Y_1} & 0 \\ X_2 & Y_1 \end{pmatrix} \\ &= X_1^{p-1}Y_1 + X_1^{p-2}Y_1^2 + \dots + X_1Y_1^{p-1} + Y_1^p \\ &= X_1^{p-1}Y_1 + X_1^{p-2}Y_1^2 + \dots + X_1Y_1^{p-1} + t. \end{aligned}$$

In the basis  $\{1, x_1, \dots, x_1^{p-1}\}$  of  $Q$ , the Bézoutian bilinear form consists of a  $t$  in the upper left corner and a 1 in each entry just below the antidiagonal. Thus

$$\deg^{\mathbb{A}^1}(f) = \deg_{(t^{1/p}, 0)}^{\mathbb{A}^1}(f) = \langle t \rangle + \frac{1}{2}(p-1)\mathbb{H}.$$

**Example 7.3.** Let  $f_1 = (x_1 - 1)x_1x_2$  and  $f_2 = (ax_1^2 - bx_2^2)$  for some  $a, b \in k^\times$  with  $\frac{a}{b}$  not a square in  $k$ . Then  $f = (f_1, f_2)$  has isolated zeros at  $\mathfrak{m} := (x_1 - 0, x_2 - 0)$  and  $\mathfrak{n} := (x_1 - 1, x_2^2 - a/b)$ . We will use Bézoutians to compute the local degrees  $\deg_{\mathfrak{m}}^{\mathbb{A}^1}(f)$  and  $\deg_{\mathfrak{n}}^{\mathbb{A}^1}(f)$ , as well as the global degree  $\deg^{\mathbb{A}^1}(f)$ . Let

$$Q = \frac{k[x_1, x_2]}{((x_1 - 1)x_1x_2, ax_1^2 - bx_2^2)}.$$

We first compute the global Bézoutian as

$$\begin{aligned} \text{Béz}(f) &= \det \begin{pmatrix} (X_1 + Y_1 - 1)X_2 & a(X_1 + Y_1) \\ Y_1^2 - Y_1 & -b(X_2 + Y_2) \end{pmatrix} \\ &= -a(X_1Y_1^2 - X_1Y_1 + Y_1^3 - Y_1^2) - b(X_1X_2^2 + X_2^2Y_1 - X_2^2 + X_1X_2Y_2 + X_2Y_1Y_2 - X_2Y_2). \end{aligned}$$

In the basis  $\{1, x_1, x_2, x_1^2, x_1x_2, x_1^3\}$  of  $Q$ , the Bézoutian is given by

$$\text{Béz}(f) = -a(X_1Y_1^2 - X_1Y_1 + Y_1^3 - Y_1^2 + X_1^3 + X_1^2Y_1 - X_1^2) - b(X_1X_2Y_2 + X_2Y_1Y_2 - X_2Y_2).$$

We now write the Bézoutian matrix given by the coefficients of  $\text{Béz}(f)$ :

	1	$X_1$	$X_2$	$X_1^2$	$X_1X_2$	$X_1^3$
1	0	0	0	$a$	0	$-a$
$Y_1$	0	$a$	0	$-a$	0	0
$Y_2$	0	0	$b$	0	$-b$	0
$Y_1^2$	$a$	$-a$	0	0	0	0
$Y_1Y_2$	0	0	$-b$	0	0	0
$Y_1^3$	$-a$	0	0	0	0	0

One may check (e.g., with a computer) that this is equal to  $3\mathbb{H}$  in  $\text{GW}(k)$ .

In  $Q_{\mathfrak{m}}$ , we have that  $x_1^2x_2 = x_1x_2 = 0$  and  $x_1^3 = \frac{b}{a}x_1x_2^2 = 0$ . In the basis  $\{1, x_1, x_2, x_1^2\}$  of  $Q_{\mathfrak{m}}$ , the global Bézoutian reduces to

$$\text{Béz}(f) = -a(X_1Y_1^2 - X_1Y_1 - Y_1^2 + X_1^2Y_1 - X_1^2) + bX_2Y_2.$$

We thus get the Bézoutian matrix at  $m$ :

$$\begin{array}{c|cccc}
 & 1 & X_1 & X_2 & X_1^2 \\
 \hline
 1 & 0 & 0 & 0 & a \\
 Y_1 & 0 & a & 0 & -a \\
 Y_2 & 0 & 0 & b & 0 \\
 Y_1^2 & a & -a & 0 & 0
 \end{array}$$

This is  $\mathbb{H} + \langle a, b \rangle$  in  $\text{GW}(k)$ .

In  $Q_n$ , we have  $x_1 = 1$ . In the basis  $\{1, x_2\}$  for  $Q_n$ , the Bézoutian reduces to

$$\text{Béz}(f) = -a - bX_2Y_2.$$

We can then write the Bézoutian matrix at  $n$ :

$$\begin{array}{c|cc}
 & 1 & X_2 \\
 \hline
 1 & -a & 0 \\
 Y_2 & 0 & -b
 \end{array}$$

This is  $\langle -a, -b \rangle$  in  $\text{GW}(k)$ . Note that  $\langle -a, -b \rangle$  need not be equal to  $\mathbb{H}$ . However, this does not contradict [29, Theorem 2], since  $n$  is a nonrational point.

Putting these computations together, we see that

$$\text{deg}_m^{\mathbb{A}^1}(f) + \text{deg}_n^{\mathbb{A}^1}(f) = \mathbb{H} + \langle a, b \rangle + \langle -a, -b \rangle = 3\mathbb{H} = \text{deg}^{\mathbb{A}^1}(f).$$

### 8. Application: The $\mathbb{A}^1$ -Euler characteristic of Grassmannians

As an application of Theorem 1.2, we compute the  $\mathbb{A}^1$ -Euler characteristic of various low-dimensional Grassmannians in Example 8.2 and Figure 1. These computations suggest a recursive formula for the  $\mathbb{A}^1$ -Euler characteristic of an arbitrary Grassmannian, which we prove in Theorem 8.4. This formula is analogous to the recursive formulas for the Euler characteristics of complex and real Grassmannians. Theorem 8.4 is probably well-known, and the proof is essentially a combination of results of Hoyois, Levine, and Bachmann–Wickelgren.

**8A. The  $\mathbb{A}^1$ -Euler characteristic.** Let  $X$  be a smooth, proper  $k$ -variety of dimension  $n$  with structure map  $\pi : X \rightarrow \text{Spec } k$ . Let  $p : T_X \rightarrow X$  denote the tangent bundle of  $X$ . The  $\mathbb{A}^1$ -Euler characteristic  $\chi^{\mathbb{A}^1}(X) \in \text{GW}(k)$  is a refinement of the classical Euler characteristic. In particular, if  $k = \mathbb{R}$ , then  $\text{rank } \chi^{\mathbb{A}^1}(X) = \chi(X(\mathbb{C}))$  and  $\text{sgn } \chi^{\mathbb{A}^1}(X) = \chi(X(\mathbb{R}))$ . There exist several equivalent definitions of the  $\mathbb{A}^1$ -Euler characteristic [20; 21; 1]. For example, we may define  $\chi^{\mathbb{A}^1}(X)$  to be the  $\pi$ -pushforward of the  $\mathbb{A}^1$ -Euler class

$$e(T_X) := z^*z_*1_X \in \widetilde{\text{CH}}^n(X, \omega_{X/k}),$$

of the tangent bundle [20], where  $z : X \rightarrow T_X$  is the zero section and  $\widetilde{\text{CH}}^d(X, \omega_{X/k})$  is the Chow–Witt group defined by Barge and Morel [3; 12]. That is,

$$\chi^{\mathbb{A}^1}(X) := \pi_*(e(T_X)) \in \widetilde{\text{CH}}^0(\text{Spec } k) = \text{GW}(k).$$

Analogous to the classical case [23], the  $\mathbb{A}^1$ -Euler characteristic can be computed as the sum of local  $\mathbb{A}^1$ -degrees at the zeros of a general section of the tangent bundle using the work of Kass and Wickelgren [2; 17; 20]. We now describe this process. Let  $\sigma$  be a section of  $T_X$  which only has isolated zeros. For a zero  $x$  of  $\sigma$ , choose Nisnevich coordinates  $\psi : U \rightarrow \mathbb{A}_k^n$  around  $x$ .<sup>2</sup> Since  $\psi$  is étale, it induces an isomorphism of tangent spaces and thus yields local coordinates around  $x$ . Shrinking  $U$  if necessary, we can trivialize  $T_X|_U \cong U \times \mathbb{A}_k^n$ . The chosen Nisnevich coordinates  $(\psi, U)$  and trivialization  $\tau : T_X|_U \cong U \times \mathbb{A}_k^n$  each define distinguished elements  $d_\psi, d_\tau \in \det T_X|_U$ . In turn, this yields a distinguished section  $d$  of  $\mathcal{H}om(\det T_X|_U, \det T_X|_U)$ , which is defined by  $d_\psi \mapsto d_\tau$ . We say that a trivialization  $\tau$  is *compatible* with the chosen coordinates  $(\psi, U)$  if the image of the distinguished section  $d$  under the canonical isomorphism  $\rho : \mathcal{H}om(\det T_X|_U, \det T_X|_U) \cong \mathcal{O}_U$  is a square [17, Definition 21].

Given a compatible trivialization  $\tau : T_X|_U \cong U \times \mathbb{A}_k^n$ , the section  $\sigma$  trivializes to  $\sigma : U \rightarrow \mathbb{A}_k^n$ . We can then define the *local index*  $\text{ind}_x \sigma$  at  $x$  to be the  $\mathbb{A}^1$ -degree of the composite

$$\frac{\mathbb{P}_k^n}{\mathbb{P}_k^{n-1}} \rightarrow \frac{\mathbb{P}_k^n}{\mathbb{P}_k^n \setminus \{\psi(x)\}} \cong \frac{\mathbb{A}_k^n}{\mathbb{A}_k^n \setminus \{\psi(x)\}} \cong \frac{U}{U \setminus \{x\}} \xrightarrow{\sigma} \frac{\mathbb{A}_k^n}{\mathbb{A}_k^n \setminus \{0\}} \cong \frac{\mathbb{P}_k^n}{\mathbb{P}_k^{n-1}}.$$

Here, the first map is the collapse map, the second map is excision, the third map is induced by the Nisnevich coordinates  $(\psi, U)$ , and the fifth map is purity; see e.g., [2, Definition 7.1]. By [17, Theorem 3], the  $\mathbb{A}^1$ -Euler characteristic is then the sum of local indices

$$\chi^{\mathbb{A}^1}(X) = \sum_{x \in \sigma^{-1}(0)} \text{ind}_x \sigma \in \text{GW}(k).$$

By Theorem 1.2, we may thus compute the  $\mathbb{A}^1$ -Euler characteristic by computing the global Bézoutian bilinear form of an appropriate map  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$ .

**Remark 8.1.** If all the zeros of  $f : \mathbb{A}_k^n \rightarrow \mathbb{A}_k^n$  are simple, then each local ring  $Q_{\mathfrak{m}_i}$  in the decomposition of  $Q = k[x_1, \dots, x_n]/(f_1, \dots, f_n) = Q_{\mathfrak{m}_1} \times \dots \times Q_{\mathfrak{m}_s}$  is equal to the residue field of the corresponding zero. If each residue field  $Q_{\mathfrak{m}_i}$  is a separable extension of  $k$ , then the  $\mathbb{A}^1$ -degree of  $f$  is equal to sum of the scaled trace forms  $\text{Tr}_{Q_{\mathfrak{m}_i}/k}(\langle J(f)|_{\mathfrak{m}_i} \rangle)$  (see e.g., [7, Definition 1.2]), where  $J(f)|_{\mathfrak{m}_i}$  is the determinant of the Jacobian of  $f$  evaluated at the point  $\mathfrak{m}_i$ . In [27] the last named author uses the scaled trace form for several  $\mathbb{A}^1$ -Euler number computations. However, Theorem 1.2 yields a formula for  $\text{deg}^{\mathbb{A}^1}(f)$  for any  $f$  with only isolated zeros and without any restriction on the residue field of each zero. Moreover, we can even compute  $\text{deg}^{\mathbb{A}^1}(f)$  without solving for the zero locus of  $f$ .

**8B. The  $\mathbb{A}^1$ -Euler characteristic of Grassmannians.** Let  $G := \text{Gr}_k(r, n)$  be the Grassmannian of  $r$ -planes in  $k^n$ . In order to compute  $\chi^{\mathbb{A}^1}(G)$ , we first need to describe Nisnevich coordinates and compatible trivializations for  $G$  and  $T_G$ . We then need to choose a convenient section of  $T_G$  and describe the resulting endomorphism  $\mathbb{A}_k^{r(n-r)}$ . The tangent bundle  $T_G \rightarrow G$  is isomorphic to  $p : \mathcal{H}om(\mathcal{S}, \mathcal{Q}) \rightarrow G$ , where  $\mathcal{S} \rightarrow G$  and  $\mathcal{Q} \rightarrow G$  are the universal sub- and quotient bundles.

<sup>2</sup>Nisnevich coordinates consist of an open neighborhood  $U$  of  $x$  and an étale map  $\psi : U \rightarrow \mathbb{A}_k^n$  that induces an isomorphism of residue fields  $k(x) \cong k(\psi(x))$  [17, Definition 18].



We now describe Nisnevich coordinates on  $G$  and a compatible trivialization of  $T_G$ , following [31]. Let  $d = r(n - r)$  be the dimension of  $G$ , and let  $\{e_1, \dots, e_n\}$  be the standard basis of  $k^n$ . Let  $\mathbb{A}_k^d = \text{Spec } k[\{x_{i,j}\}_{i,j=1}^{r,n-r}] \cong U \subset G$  be the open affine subset consisting of the  $r$ -planes

$$H(\{x_{i,j}\}_{i,j=1}^{r,n-r}) := \text{span} \left\{ e_{n-r+i} + \sum_{j=1}^{n-r} x_{i,j} e_j \right\}_{i=1}^r.$$

The map  $\psi : U \rightarrow \mathbb{A}_k^d$  given by  $\psi(H(\{x_{i,j}\}_{i,j=1}^{r,n-r})) = (\{x_{i,j}\}_{i,j=1}^{n-r,r})$  yields Nisnevich coordinates  $(\psi, U)$  centered at  $\psi(\text{span}\{e_{n-r+1}, \dots, e_n\}) = (0, \dots, 0)$ . For the trivialization of  $T_G|_U$ , let

$$\tilde{e}_i = \begin{cases} e_i & i \leq n-r, \\ e_i + \sum_{j=1}^{n-r} x_{i-(n-r),j} e_j & i \geq n-r+1. \end{cases}$$

Then  $\{\tilde{e}_1, \dots, \tilde{e}_n\}$  is a basis for  $k^n$ , and we denote the dual basis by  $\{\tilde{\phi}_1, \dots, \tilde{\phi}_n\}$ . Over  $U$ , the bundles  $\mathcal{S}^*$  and  $\mathcal{Q}$  are trivialized by  $\{\tilde{\phi}_{n-r+1}, \dots, \tilde{\phi}_n\}$  and  $\{\tilde{e}_1, \dots, \tilde{e}_{n-r}\}$ , respectively. Since

$$T_G \cong \mathcal{H}om(\mathcal{S}, \mathcal{Q}) \cong \mathcal{S}^* \otimes \mathcal{Q},$$

we get a trivialization of  $T_G|_U$  given by  $\{\tilde{\phi}_{n-r+i} \otimes \tilde{e}_j\}_{i,j=1}^{r,n-r}$ . By construction, our Nisnevich coordinates  $(\psi, U)$  induce this local trivialization of  $T_G$ . It follows that the distinguished element of  $\text{Hom}(\det T_G|_U, \det T_G|_U)$  sending the distinguished element of  $\det T_G|_U$  (determined by the Nisnevich coordinates) to the distinguished element of  $T_G|_U$  (determined by our local trivialization) is just the identity, which is a square.

Next, we describe sections of  $T_G \rightarrow G$  and the resulting endomorphisms  $\mathbb{A}_k^d \rightarrow \mathbb{A}_k^d$ . Let  $\{\phi_1, \dots, \phi_n\}$  be the dual basis of the standard basis  $\{e_1, \dots, e_n\}$  of  $k^n$ . A homogeneous degree 1 polynomial  $\alpha \in k[\phi_1, \dots, \phi_n]$  gives rise to a section  $s$  of  $\mathcal{S}^*$ , defined by evaluating  $\alpha$ . In particular, given a vector  $t = \sum_{i=1}^n t_i \tilde{e}_i$  in  $H(\{x_{i,j}\}_{i,j=1}^{r,n-r})$ , we use the dual change of basis

$$\phi_j = \begin{cases} \tilde{\phi}_j + \sum_{i=1}^r x_{i,j} \tilde{\phi}_{n-r+i} & j \leq n-r, \\ \tilde{\phi}_j & j \geq n-r+1 \end{cases}$$

to set

$$s(t) = \alpha \left( t_1 + \sum_{i=1}^r x_{i,1} t_{n-r+i}, \dots, t_{n-r} + \sum_{i=1}^r x_{i,n-r} t_{n-r+i}, t_{n-r+1}, \dots, t_n \right).$$

Note that  $t_1 = \dots = t_{n-r} = 0$  if and only if  $t \in H(\{x_{i,j}\}_{i,j=1}^{r,n-r})$ , so  $s(t) \in k[t_{n-r+1}, \dots, t_n]$ . Taking  $n$  sections  $s_1, \dots, s_n$  of  $\mathcal{S}^*$ , we get a section of  $T_G \cong \mathcal{H}om(\mathcal{S}, \mathcal{Q})$  given by

$$\mathcal{S} \xrightarrow{(s_1, \dots, s_n)} \mathbb{A}_k^n \rightarrow \mathcal{Q},$$

where the second map is quotienting by  $\{\tilde{e}_{n-r+1}, \dots, \tilde{e}_n\}$ . We obtain our map  $\mathbb{A}_k^d \rightarrow \mathbb{A}_k^d$  by applying the trivializations  $\{\tilde{\phi}_{n-r+i} \otimes \tilde{e}_j\}_{i,j=1}^{r,n-r}$  of  $T_G$ . Explicitly, take  $n$  sections  $s_1, \dots, s_n$  of  $\mathcal{S}^*$ . Since  $e_i =$

$\tilde{e}_i - \sum_{j=1}^{n-r} x_{i-(n-r),j} e_j$  for  $i > n - r$ , we have

$$s_j e_j \equiv s_j e_j - \sum_{i=1}^r x_{i,j} s_{n-r+i} e_j \pmod{(\tilde{e}_{n-r+1}, \dots, \tilde{e}_n)},$$

for all  $j \leq n - r$ . Recall that  $e_j = \tilde{e}_j$  for  $j \leq n - r$ . The coordinate of  $\mathbb{A}_k^d \rightarrow \mathbb{A}_k^d$  corresponding to  $\tilde{\phi}_{n-r+i} \otimes \tilde{e}_j$  is thus the coefficient of  $t_{n-r+i}$  in  $s_j(t) - \sum_{\ell=1}^r x_{\ell,j} s_{n-r+\ell}(t)$ .

For a general section  $\sigma$  of  $p : T_G \rightarrow G$ , the finitely many zeros of  $\sigma$  will all lie in  $U$ . In this case, the  $\mathbb{A}^1$ -Euler characteristic of  $G$  is equal to the global  $\mathbb{A}^1$ -degree of the resulting map  $\mathbb{A}_k^d \rightarrow \mathbb{A}_k^d$ , which can be computed using the Bézoutian.

**Example 8.2** ( $\text{Gr}_k(2, 4)$ ). Let

$$\begin{aligned} \alpha_1 &= \phi_2 = \tilde{\phi}_2 + x_{1,2} \tilde{\phi}_3 + x_{2,2} \tilde{\phi}_4, \\ \alpha_2 &= \phi_3 = \tilde{\phi}_3, \\ \alpha_3 &= \phi_4 = \tilde{\phi}_4, \\ \alpha_4 &= \phi_1 = \tilde{\phi}_1 + x_{1,1} \tilde{\phi}_3 + x_{2,1} \tilde{\phi}_4. \end{aligned}$$

Evaluating at  $t = (0, 0, t_3, t_4)$  in the basis  $\{\tilde{e}_i\}$ , we have

$$\begin{aligned} s_1 &= x_{1,2} t_3 + x_{2,2} t_4, \\ s_2 &= t_3, \\ s_3 &= t_4, \\ s_4 &= x_{1,1} t_3 + x_{2,1} t_4. \end{aligned}$$

It remains to read off the coefficients of  $t_3$  and  $t_4$  of

$$\begin{aligned} s_1 - x_{1,1} s_3 - x_{2,1} s_4 &= (x_{1,2} - x_{1,1} x_{2,1}) t_3 + (x_{2,2} - x_{1,1} - x_{2,1}^2) t_4, \\ s_2 - x_{1,2} s_3 - x_{2,2} s_4 &= (1 - x_{1,1} x_{2,2}) t_3 + (-x_{1,2} - x_{2,1} x_{2,2}) t_4. \end{aligned}$$

We thus have our endomorphism  $\sigma : \mathbb{A}_k^4 \rightarrow \mathbb{A}_k^4$  defined by

$$\sigma = (x_{1,2} - x_{1,1} x_{2,1}, x_{2,2} - x_{1,1} - x_{2,1}^2, 1 - x_{1,1} x_{2,2}, -x_{1,2} - x_{2,1} x_{2,2}).$$

Using the Sage implementation of the Bézoutian formula for the  $\mathbb{A}^1$ -degree [8], we can calculate  $\chi^{\mathbb{A}^1}(\text{Gr}_k(2, 4)) = \deg^{\mathbb{A}^1}(\sigma) = 2\mathbb{H} + \langle 1, 1 \rangle$ .

Using a computer, we performed computations analogous to Example 8.2 for  $r \leq 5$  and  $n \leq 7$ . These  $\mathbb{A}^1$ -Euler characteristics of Grassmannians are recorded in Figure 1.

Recall that the Euler characteristics of real and complex Grassmannians are given by binomial coefficients. In particular, these Euler characteristics satisfy certain recurrence relations related to Pascal's rule. The computations in Figure 1 indicate that an analogous recurrence relation is true for the  $\mathbb{A}^1$ -Euler characteristic of Grassmannians over an arbitrary field. In fact, this recurrence relation is a direct consequence of a result of Levine [20].

$n$	$r = 1$	2	3	4	5
2	$\mathbb{H}$	$\langle 1 \rangle$			
3	$\mathbb{H} + \langle 1 \rangle$	$\mathbb{H} + \langle 1 \rangle$	$\langle 1 \rangle$		
4	$2\mathbb{H}$	$2\mathbb{H} + \langle 1, 1 \rangle$	$2\mathbb{H}$	$\langle 1 \rangle$	
5	$2\mathbb{H} + \langle 1 \rangle$	$4\mathbb{H} + \langle 1, 1 \rangle$	$4\mathbb{H} + \langle 1, 1 \rangle$	$2\mathbb{H} + \langle 1 \rangle$	$\langle 1 \rangle$
6	$3\mathbb{H}$	$6\mathbb{H} + \langle 1, 1, 1 \rangle$	$10\mathbb{H}$	$6\mathbb{H} + \langle 1, 1, 1 \rangle$	$3\mathbb{H}$
7	$3\mathbb{H} + \langle 1 \rangle$	$9\mathbb{H} + \langle 1, 1, 1 \rangle$	$16\mathbb{H} + \langle 1, 1, 1 \rangle$	$16\mathbb{H} + \langle 1, 1, 1 \rangle$	$9\mathbb{H} + \langle 1, 1, 1 \rangle$

**Figure 1.** More examples of  $\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n))$ .

**Proposition 8.3.** *Let  $1 \leq r < n$  be integers. Then*

$$\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n)) = \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r - 1, n - 1)) + \langle -1 \rangle^r \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n - 1)).$$

*Proof.* Fix a line  $L$  in  $k^n$ . Let  $Z$  be the closed subvariety consisting of all  $r$ -planes containing  $L$  (which is isomorphic to  $\mathrm{Gr}_k(r - 1, n - 1)$ ), and let  $U$  be its open complement (which is isomorphic to an affine rank  $r$  bundle over  $\mathrm{Gr}_k(r, n - 1)$ ). We then get a decomposition  $\mathrm{Gr}_k(r, n) = Z \cup U$ . Since  $\mathrm{Gr}_k(l, m) \cong \mathrm{Gr}_k(m - l, m)$ , we have  $\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(l, m)) = \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(m - l, m))$ . We can thus apply [20, Proposition 1.4(3)] to obtain

$$\begin{aligned} \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n)) &= \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(n - r, n)) \\ &= \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(n - r, n - 1)) + \langle -1 \rangle^r \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(n - r - 1, n - 1)) \\ &= \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r - 1, n - 1)) + \langle -1 \rangle^r \chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n - 1)). \end{aligned} \quad \square$$

We can now apply a theorem of Bachmann and Wickelgren [2] to completely characterize  $\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n))$ .

**Theorem 8.4.** *Let  $k$  be field of characteristic not equal to 2. Let  $n_{\mathbb{C}} := \binom{n}{r}$ , and let  $n_{\mathbb{R}} := \binom{\lfloor n/2 \rfloor}{\lfloor r/2 \rfloor}$ . Then*


$$\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n)) = \frac{n_{\mathbb{C}} + n_{\mathbb{R}}}{2} \langle 1 \rangle + \frac{n_{\mathbb{C}} - n_{\mathbb{R}}}{2} \langle -1 \rangle.$$

*Proof.* By [2, Theorem 5.8], we can restrict this computation to two different possibilities. We will prove by induction that  $\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(r, n)) \bmod \mathbb{H}$  has no  $\langle 2 \rangle$  summand. The desired result will then follow from [2, Theorem 5.8] by noting that  $n_{\mathbb{C}}$  and  $n_{\mathbb{R}}$  are the Euler characteristics of  $\mathrm{Gr}_{\mathbb{C}}(r, n)$  and  $\mathrm{Gr}_{\mathbb{R}}(r, n)$ , respectively.

Since  $\mathbb{A}_k^n$  is  $\mathbb{A}^1$ -homotopic to  $\mathrm{Spec} k$ , we have  $\chi^{\mathbb{A}^1}(\mathbb{A}_k^n) = \chi^{\mathbb{A}^1}(\mathrm{Spec} k) = \langle 1 \rangle$ . Using this observation and the decomposition  $\mathbb{P}_k^n = \bigcup_{i=0}^n \mathbb{A}_k^i$  (and a result analogous to [20, Proposition 1.4(3)]), Hoyois computed the  $\mathbb{A}^1$ -Euler characteristic of projective space [15, Example 1.7]:

$$\chi^{\mathbb{A}^1}(\mathbb{P}_k^n) = \begin{cases} \frac{n}{2} \mathbb{H} + \langle 1 \rangle & n \text{ is even,} \\ \frac{n+1}{2} \mathbb{H} & n \text{ is odd.} \end{cases}$$

Note that  $\mathrm{Gr}_k(0, n) \cong \mathrm{Gr}_k(n, n) \cong \mathrm{Spec} k$  and  $\mathrm{Gr}_k(1, n) \cong \mathrm{Gr}_k(n - 1, n) \cong \mathbb{P}_k^{n-1}$ . In particular,  $\chi^{\mathbb{A}^1}(\mathrm{Gr}_k(i, n)) \bmod \mathbb{H}$  is either trivial or  $\langle 1 \rangle$  for  $i = 0, 1, n - 1$ , or  $n$ . This forms the base case of our



**Figure 2.** Addition rules for modified Pascal's triangle.

induction, with the inductive step given by Proposition 8.3—namely, if  $\chi^{\mathbb{A}^1}(\text{Gr}_k(r-1, n-1)) \bmod \mathbb{H}$  and  $\chi^{\mathbb{A}^1}(\text{Gr}_k(r, n-1)) \bmod \mathbb{H}$  only have  $\langle 1 \rangle$  and  $\langle -1 \rangle$  summands, then

$$(\chi^{\mathbb{A}^1}(\text{Gr}_k(r-1, n-1)) + \langle -1 \rangle^r \chi^{\mathbb{A}^1}(\text{Gr}_k(r, n-1))) \bmod \mathbb{H}$$

only has  $\langle 1 \rangle$  and  $\langle -1 \rangle$  summands. □


**8C. Modified Pascal's triangle for  $\chi^{\mathbb{A}^1}(\text{Gr}_k(r, n))$ .** Pascal's triangle gives a mnemonic device for binomial coefficients and hence for the Euler characteristics of complex and real Grassmannians. The recurrence relation of Proposition 8.3 indicates that a modification of Pascal's triangle can also be used to calculate the  $\mathbb{A}^1$ -Euler characteristics of Grassmannians. Explicitly, each entry in the modified Pascal's triangle is an element of  $\text{GW}(k)$ . The two diagonal edges of this triangle correspond to  $\chi^{\mathbb{A}^1}(\text{Gr}_k(0, n)) = \langle 1 \rangle$  and  $\chi^{\mathbb{A}^1}(\text{Gr}_k(n, n)) = \langle 1 \rangle$ . Elements of each row of the modified Pascal's triangle are obtained from the previous row by the addition rule illustrated in Figure 2.

We rewrite the data recorded in Figure 1 in a modified Pascal's triangle in Figure 3. The rows correspond to the dimension  $n$  of the ambient affine space  $k^n$ , while the southwest-to-northeast diagonals correspond to the dimension  $r$  of the planes  $k^r$  in the ambient space.

### Acknowledgements

We thank Tom Bachmann, Gard Helle, Kyle Ormsby, Paul Arne Østvær, and Kirsten Wickelgren for helpful comments. We thank Kirsten Wickelgren for suggesting that we make available a code implementation for computing local and global  $\mathbb{A}^1$ -degrees. Finally, we thank the anonymous referee for their helpful comments, suggestions, and corrections.

Brazelton is supported by an NSF Graduate Research Fellowship (DGE-1845298). McKean received support from Kirsten Wickelgren's NSF CAREER grant (DMS-1552730). Pauli acknowledges support from a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 832833).



**Figure 3.** Modified Pascal’s triangle for  $\chi^{\mathbb{A}^1}(\text{Gr}_k(r, n))$  (see Section 8C).

**References**

[1] N. Arcila-Maya, C. Bethea, M. Opie, K. Wickelgren, and I. Zakharevich, “Compactly supported  $\mathbb{A}^1$ -Euler characteristic and the Hochschild complex”, *Topology Appl.* **316** (2022), Paper No. 108108, 24. MR

[2] T. Bachmann and K. Wickelgren, “Euler classes: six-functors formalism, dualities, integrality and linear subspaces of complete intersections”, *J. Inst. Math. Jussieu* **22:2** (2023), 681–746. MR Zbl

[3] J. Barge and F. Morel, “Groupe de Chow des cycles orientés et classe d’Euler des fibrés vectoriels”, *C. R. Acad. Sci. Paris Sér. I Math.* **330:4** (2000), 287–290. MR Zbl

[4] E. Becker, J. P. Cardinal, M.-F. Roy, and Z. Szafraniec, “Multivariate Bezoutians, Kronecker symbol and Eisenbud–Levine formula”, pp. 79–104 in *Algorithms in algebraic geometry and applications* (Santander, 1994), edited by L. González-Vega and T. Recio, Progr. Math. **143**, Birkhäuser, Basel, 1996. MR Zbl

[5] C. Bethea, J. L. Kass, and K. Wickelgren, “Examples of wild ramification in an enriched Riemann–Hurwitz formula”, pp. 69–82 in *Motivic homotopy theory and refined enumerative geometry*, edited by F. Binda et al., Contemp. Math. **745**, Amer. Math. Soc., Providence, RI, [2020] ©2020. MR Zbl

[6] T. Brazelton, “An introduction to  $\mathbb{A}^1$ -enumerative geometry”, pp. 11–47 in *Homotopy theory and arithmetic geometry—motivic and Diophantine aspects*, edited by F. Neumann and A. Pál, Lecture Notes in Math. **2292**, Springer, [2021] ©2021. MR Zbl

[7] T. Brazelton, R. Burklund, S. McKean, M. Montoro, and M. Opie, “The trace of the local  $\mathbb{A}^1$ -degree”, *Homology Homotopy Appl.* **23:1** (2021), 243–255. MR

- [8] T. Brazelton, S. McKean, and S. Pauli,  $\mathbb{A}^1$ -degree.sage, available at <https://github.com/shmckean/A1-degree/>.
- [9] C. Cazanave, “Algebraic homotopy classes of rational functions”, *Ann. Sci. Éc. Norm. Supér. (4)* **45**:4 (2012), 511–534. MR Zbl
- [10] B. Eckmann and P. J. Hilton, “Group-like structures in general categories, I: Multiplications and comultiplications”, *Math. Ann.* **145** (1961/62), 227–255. MR Zbl
- [11] D. Eisenbud and H. I. Levine, “An algebraic formula for the degree of a  $C^\infty$  map germ”, *Ann. of Math. (2)* **106**:1 (1977), 19–44. MR Zbl
- [12] J. Fasel, “Groupes de Chow–Witt”, *Mém. Soc. Math. Fr. (N.S.)* 113 (2008), viii+197. MR Zbl
- [13] G. N. Himšiašvili, “The local degree of a smooth mapping”, *Sakharth. SSR Mecn. Akad. Moambe* **85**:2 (1977), 309–312. MR
- [14] J. Hornbostel, “ $\mathbb{A}^1$ -representability of Hermitian  $K$ -theory and Witt groups”, *Topology* **44**:3 (2005), 661–687. MR Zbl
- [15] M. Hoyois, “A quadratic refinement of the Grothendieck–Lefschetz–Verdier trace formula”, *Algebr. Geom. Topol.* **14**:6 (2014), 3603–3658. MR Zbl
- [16] J. L. Kass and K. Wickelgren, “The class of Eisenbud–Khimshiashvili–Levine is the local  $\mathbb{A}^1$ -Brouwer degree”, *Duke Math. J.* **168**:3 (2019), 429–469. MR Zbl
- [17] J. L. Kass and K. Wickelgren, “An arithmetic count of the lines on a smooth cubic surface”, *Compos. Math.* **157**:4 (2021), 677–709. MR Zbl
- [18] J. Knight, A. A. Swaminathan, and D. Tseng, “On the EKL-degree of a Weyl cover”, *J. Algebra* **565** (2021), 64–81. MR Zbl
- [19] H. Larson and I. Vogt, “An enriched count of the bitangents to a smooth plane quartic curve”, *Res. Math. Sci.* **8**:2 (2021), Paper No. 26, 21. MR Zbl
- [20] M. Levine, “Aspects of enumerative geometry with quadratic forms”, *Doc. Math.* **25** (2020), 2179–2239. MR Zbl
- [21] M. Levine and A. Raksit, “Motivic Gauss–Bonnet formulas”, *Algebra Number Theory* **14**:7 (2020), 1801–1851. MR
- [22] S. McKean, “An arithmetic enrichment of Bézout’s Theorem”, *Math. Ann.* **379**:1-2 (2021), 633–660. MR Zbl
- [23] J. W. Milnor, *Topology from the differentiable viewpoint*, University Press of Virginia, Charlottesville, VA, 1965. MR Zbl
- [24] F. Morel, “An introduction to  $\mathbb{A}^1$ -homotopy theory”, pp. 357–441 in *Contemporary developments in algebraic K-theory* (Trieste, 2002), edited by M. Karoubi et al., ICTP Lecture Notes **15**, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004. MR Zbl
- [25] F. Morel, “ $\mathbb{A}^1$ -algebraic topology”, pp. 1035–1059 in *International Congress of Mathematicians*, vol. 2, edited by M. Sanz-Solé et al., Eur. Math. Soc., Zürich, 2006. MR Zbl
- [26] S. Pauli, “Quadratic types and the dynamic Euler number of lines on a quintic threefold”, *Adv. Math.* **405** (2022), Paper No. 108508, 37. MR Zbl
- [27] S. Pauli, “Computing  $\mathbb{A}^1$ -Euler numbers with Macaulay2”, *Res. Math. Sci.* **10**:3 (2023), Paper No. 26, 14. MR Zbl
- [28] S. Pauli and K. Wickelgren, “Applications to  $\mathbb{A}^1$ -enumerative geometry of the  $\mathbb{A}^1$ -degree”, *Res. Math. Sci.* **8**:2 (2021), Paper No. 24, 29. MR Zbl
- [29] G. Quick, T. Strand, and G. M. Wilson, “Representability of the local motivic Brouwer degree”, *Math. Scand.* **128**:1 (2022), 54–77. MR Zbl
- [30] G. Scheja and U. Storch, “Über Spurfunktionen bei vollständigen Durchschnitten”, *J. Reine Angew. Math.* **278(279)** (1975), 174–190. MR Zbl
- [31] P. Srinivasan and K. Wickelgren, “An arithmetic count of the lines meeting four lines in  $\mathbb{P}^3$ ”, *Trans. Amer. Math. Soc.* **374**:5 (2021), 3427–3451. MR Zbl
- [32] “The Stacks project”, electronic reference, available at <http://stacks.math.columbia.edu>. Zbl

Communicated by Vasudevan Srinivas

Received 2022-04-11

Revised 2022-09-29

Accepted 2023-01-20

tbraz@math.upenn.edu

*Department of Mathematics, University of Pennsylvania, Philadelphia, PA, United States*

smckean@math.harvard.edu

*Department of Mathematics, Harvard University, Cambridge, MA, United States*

sabrina.pauli@uni-due.de

*Fakultät für Mathematik, Universität Duisburg-Essen, Essen, Germany*

# Algebra & Number Theory

msp.org/ant

## EDITORS

MANAGING EDITOR  
Antoine Chambert-Loir  
Université Paris-Diderot  
France

EDITORIAL BOARD CHAIR  
David Eisenbud  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/ant](http://msp.org/ant) for submission instructions.

---

The subscription price for 2023 is US \$485/year for the electronic version, and \$705/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.


---

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

---

ANT peer review and production are managed by EditFLOW<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

# Algebra & Number Theory

Volume 17    No. 11    2023

---

On self-correspondences on curves JOËL BELLAÏCHE	1867
Fitting ideals of class groups for CM abelian extensions MAHIRO ATSUTA and TAKENORI KATAOKA	1901
The behavior of essential dimension under specialization, II ZINOVY REICHSTEIN and FEDERICO SCAVIA	1925
Differential operators, retracts, and toric face rings CHRISTINE BERKESCH, C-Y. JEAN CHAN, PATRICIA KLEIN, LAURA FELICIA MATUSEVICH, JANET PAGE and JANET VASSILEV	1959
Bézoutians and the $\mathbb{A}^1$ -degree THOMAS BRAZELTON, STEPHEN MCKEAN and SABRINA PAULI	1985
Axiomatizing the existential theory of $\mathbb{F}_q((t))$ SYLVY ANSCOMBE, PHILIP DITTMANN and ARNO FEHM	2013
The diagonal coinvariant ring of a complex reflection group STEPHEN GRIFFETH	2033