

Algebra & Number Theory

Volume 17

2023

No. 6

**Correction to the article
Height bounds and the Siegel property**

Martin Orr and Christian Schnell



Correction to the article Height bounds and the Siegel property

Martin Orr and Christian Schnell

This is a correction to the paper “Height bounds and the Siegel property” (*Algebra Number Theory* **12:2** (2018), 455–478). We correct an error in the proof of Theorem 4.1. Theorem 4.1 as stated in the original paper is correct, but the correction affects additional information about the theorem which is important for applications.

There is an error in the proof of [Orr 2018, Theorem 4.1]. The statement of Theorem 4.1 is correct, but [loc. cit., Lemma 4.4] is incorrect under the conditions on K_G stated above it.

Subsequent applications [Bakker et al. 2020, Theorem 1.1(2); Daw and Orr 2021, Lemma 2.3] have required greater control of the maximal compact subgroup K_G than is given by the statement of [Orr 2018, Theorem 4.1]. As a result of the error in the proof, the choice of K_G is more constrained than it appears in [loc. cit.]. We therefore state a version of [loc. cit., Theorem 4.1], extended to correctly describe the constraints on K_G .

Theorem 1. *Let G and H be reductive \mathbb{Q} -algebraic groups, with $H \subset G$. Let \mathfrak{S}_H be a Siegel set in $H(\mathbb{R})$ with respect to the Siegel triple (P_H, S_H, K_H) . Let $K_G \subset G(\mathbb{R})$ be a maximal compact subgroup such that*

- (i) $K_H \subset K_G$; and
- (ii) the Cartan involution of G associated with K_G stabilises S_H .

Then there exist subgroups $P_G, S_G \subset G$ forming a Siegel triple (P_G, S_G, K_G) , a Siegel set $\mathfrak{S}_G \subset G(\mathbb{R})$ with respect to this Siegel triple, and a finite set $C \subset G(\mathbb{Q})$ such that

$$\mathfrak{S}_H \subset C \cdot \mathfrak{S}_G.$$

Furthermore, $R_u(P_H) \subset R_u(P_G)$ and $S_H = S_G \cap H$.

Remark 2. In the setting of Theorem 1, let Θ be the Cartan involution of G associated with K_G . We now compare (ii) with:

- (ii') Θ stabilises H .

MSC2020: primary 11F06; secondary 11G18.

Keywords: reduction theory, Siegel sets.

If (i) and (ii') are satisfied, then the restriction $\Theta|_{\mathbf{H}}$ is the Cartan involution of \mathbf{H} associated with $K_{\mathbf{H}}$. Hence, by the definition of Siegel triple, (ii) is satisfied. However, if (i) and (ii) are satisfied, then (ii') does not necessarily hold. This may be seen in the example $\mathbf{G} = \mathrm{SL}_2$, $\mathbf{H} = \left\{ \begin{pmatrix} a & b \\ db & a \end{pmatrix} : a^2 - db^2 = 1 \right\}$ where d is a nonsquare positive rational number, $K_{\mathbf{G}} = \mathrm{SO}_2(\mathbb{R})$, $\mathbf{S}_{\mathbf{H}} = \{1\}$, $K_{\mathbf{H}} = \{1\}$.

In this note, we explain how to correct the proof of [Orr 2018, Theorem 4.1] and prove Theorem 1. We also give examples showing that condition (ii) of Theorem 1 cannot be deleted from the statement of the theorem: first an example in which \mathbf{H} is a torus, then a more sophisticated example in which \mathbf{H} is semisimple. At the end of the note, we correct some unrelated minor errors in [loc. cit.].

A. Correction to proof of [Orr 2018, Theorem 4.1]. On [Orr 2018, page 470], item (2) (the choice of $K_{\mathbf{G}}$) should be replaced by:

- (2) $K_{\mathbf{G}}$, a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$ containing $K_{\mathbf{H}}$, such that the Cartan involution of \mathbf{G} associated with $K_{\mathbf{G}}$ stabilises $\mathbf{S}_{\mathbf{H}}$.

Paragraph 1 of the proof of [loc. cit., Lemma 4.4] is incorrect: neither the original constraint on $K_{\mathbf{G}}$, nor the corrected constraint, are sufficient to guarantee that Θ restricts to an involution of \mathbf{H} (see Remark 2). With the corrected constraint, that paragraph can be ignored and paragraph 2 of the proof of [loc. cit., Lemma 4.4] is valid. Hence the lemma is true under the corrected constraint on $K_{\mathbf{G}}$.

The remainder of the proof of [loc. cit., Theorem 4.1] is valid without any changes related to the choice of $K_{\mathbf{G}}$ (but see unrelated minor corrections in Section E of this note). No further conditions are imposed on $K_{\mathbf{G}}$, so this proves Theorem 1.

In order to establish [loc. cit., Theorem 4.1], it is necessary to verify the existence of $K_{\mathbf{G}}$ satisfying (2) above. To show this, choose a faithful representation $\rho : \mathbf{G}_{\mathbb{R}} \rightarrow \mathrm{GL}(V)$ for some real vector space V . By [Mostow 1955, Theorem 7.3], there exists a positive definite symmetric form ψ on V with respect to which the groups $K_{\mathbf{H}} \subset \mathbf{H}(\mathbb{R}) \subset \mathbf{G}(\mathbb{R}) \subset \mathrm{GL}(V)$ are simultaneously self-adjoint. In other words, if Θ denotes the Cartan involution of $\mathrm{GL}(V)$ associated with ψ , then Θ restricts to Cartan involutions of \mathbf{G} , \mathbf{H} and $K_{\mathbf{H}}$.

Letting $K_{\mathbf{G}}$ denote the stabiliser of ψ in $\mathbf{G}(\mathbb{R})$, we obtain $K_{\mathbf{H}} \subset K_{\mathbf{G}}$. By Remark 2, Θ stabilises $\mathbf{S}_{\mathbf{H}}$.

B. Counterexample in which condition (ii) of Theorem 1 is not satisfied: a torus. Let $\mathbf{G} = \mathrm{SL}_2$ and let $(\mathbf{P}_0, \mathbf{S}_0, K_{\mathbf{G}})$ be the standard Siegel triple for \mathbf{G} , that is, \mathbf{P}_0 is the subgroup of upper triangular matrices in \mathbf{G} , \mathbf{S}_0 is the subgroup of diagonal matrices in \mathbf{G} and $K_{\mathbf{G}} = \mathrm{SO}_2(\mathbb{R})$.

Let

$$\mathbf{H} = \left\{ \begin{pmatrix} x & x^{-1} - x \\ 0 & x^{-1} \end{pmatrix} \right\} \subset \mathbf{G}.$$

This is a \mathbb{Q} -split torus so it possesses a unique Siegel triple, namely $\mathbf{P}_{\mathbf{H}} = \mathbf{S}_{\mathbf{H}} = \mathbf{H}$, $K_{\mathbf{H}} = \{\pm 1\}$, and a unique Siegel set, $\mathfrak{S}_{\mathbf{H}} = \mathbf{H}(\mathbb{R})$.

Clearly $K_{\mathbf{H}} = \{\pm 1\} \subset K_{\mathbf{G}}$. Thus $K_{\mathbf{G}}$ satisfies condition (i) of Theorem 1. However by [Orr 2018, Lemma 2.1], \mathbf{S}_0 is the only \mathbb{Q} -split torus in \mathbf{P}_0 stabilised by the Cartan involution of \mathbf{G} associated with $K_{\mathbf{G}}$.

Hence this Cartan involution does not stabilise S_H . In other words, K_G does not satisfy condition (ii) of [Theorem 1](#).

Now we shall show that this \mathfrak{S}_H and K_G do not satisfy the conclusion of [Theorem 1](#). Suppose for contradiction that there exist subgroups $P_G, S_G \subset G$ forming a Siegel triple (P_G, S_G, K_G) , a Siegel set $\mathfrak{S}_G \subset G(\mathbb{R})$ with respect to this Siegel triple, and a finite set $C \subset G(\mathbb{Q})$ such that $\mathfrak{S}_H \subset C \cdot \mathfrak{S}_G$.

By [[Borel and Tits 1965](#), Théorème 4.13], there exists $g \in G(\mathbb{Q})$ such that $P_0 = gP_Gg^{-1}$. Writing $g = pk$ where $p \in P_0(\mathbb{R})$ and $k \in K_G$, (P_0, kS_Gk^{-1}, K_G) is a Siegel triple and $g\mathfrak{S}_G$ is a Siegel set with respect to (P_0, kS_Gk^{-1}, K_G) . Hence we can replace P_G by P_0 , S_G by kS_Gk^{-1} , \mathfrak{S}_G by $g\mathfrak{S}_G$ and C by Cg^{-1} . We can thus assume that $P_G = P_0$. By the uniqueness of the torus in a Siegel triple, this implies that $S_G = S_0$ and \mathfrak{S}_G is a standard Siegel set in $G(\mathbb{R})$.

The image of $\mathfrak{S}_H = S_H(\mathbb{R})$ in $G(\mathbb{R})/K_0$, identified with the upper half-plane, is the ray

$$R = \{(1 - y) + yi : y \in \mathbb{R}_{>0}\}.$$

Write \mathcal{F}_G for the image of \mathfrak{S}_G in the upper half-plane.

Since $R \subset C\mathcal{F}_G$ and C is finite, there exists $\gamma \in C \subset G(\mathbb{Q})$ such that $R \cap \gamma\mathcal{F}_G$ contains points z where both $\text{Im } z, |\text{Re } z| \rightarrow \infty$. But this is impossible because:

- (i) If $\gamma \notin P_0(\mathbb{Q})$, then $\gamma\mathcal{F}_G$ lies below a horizontal line.
- (ii) If $\gamma \in P_0(\mathbb{Q})$, then $\gamma\mathcal{F}_G$ lies within a vertical strip of finite width.

C. Counterexample in which condition (ii) of [Theorem 1](#) is not satisfied: a semisimple subgroup. Let $G = \text{SL}_3$ and let (P_0, S_0, K_G) be the standard Siegel triple for G . Let

$$H_0 = \text{SO}_3(J) \quad \text{where } J = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Let Q_J denote the quadratic form on \mathbb{R}^3 represented by J . This form is negative definite on the 1-dimensional subspace $L = \mathbb{R}(1, 0, -1)^t \subset \mathbb{R}^3$ and positive definite on the 2-dimensional subspace $M = \mathbb{R}(0, 1, 0)^t + \mathbb{R}(1, 0, 1)^t$. Let

$$K_H = \{h \in H_0(\mathbb{R}) : h(L) = L \text{ and } h(M) = M\}.$$

This is a maximal compact subgroup of $H_0(\mathbb{R})$ and is isomorphic to $O_2(\mathbb{R})$ via restriction to its action on M .

Let $c \in \mathbb{Q} \setminus \{0, \pm 1\}$. Let $\eta \in \text{GL}_3(\mathbb{Q})$ be the linear map which acts as multiplication by c on L and as the identity on M . Explicitly,

$$\eta = \begin{pmatrix} \frac{1}{2}(1+c) & 0 & \frac{1}{2}(1-c) \\ 0 & 1 & 0 \\ \frac{1}{2}(1-c) & 0 & \frac{1}{2}(1+c) \end{pmatrix}.$$

Let

$$H = \eta H_0 \eta^{-1} = \text{SO}_3(\eta J \eta^t).$$

By construction, η centralises K_H . It follows that $\eta K_H \eta^{-1} = K_H = K_G \cap \mathbf{H}(\mathbb{R})$ and K_H is a maximal compact subgroup of $\mathbf{H}(\mathbb{R})$.

Let Q_0 denote the standard quadratic form on \mathbb{R}^3 . The spaces L and M are orthogonal with respect to Q_0 and $Q_{0|M} = Q_{J|M}$. Hence $K_H \subset \mathrm{SO}_3(Q_0) = K_G$. Thus condition (i) of [Theorem 1](#) is satisfied.

Let $\mathbf{P}_H = \eta(\mathbf{P}_0 \cap \mathbf{H}_0)\eta^{-1}$ and $\mathbf{S}_H = \eta(\mathbf{S}_0 \cap \mathbf{H}_0)\eta^{-1}$. As in [\[Borel 1969, 11.16\]](#), $\mathbf{P}_0 \cap \mathbf{H}_0$ is a minimal \mathbb{Q} -parabolic subgroup of \mathbf{H}_0 so $(\mathbf{P}_H, \mathbf{S}_H, K_H)$ is a Siegel triple in \mathbf{H} . Let $\mathfrak{S}_H = \Omega_H A_{H,t} K_H$ be a Siegel set in $\mathbf{H}(\mathbb{R})$ with respect to this Siegel triple.

We shall show that \mathfrak{S}_H and K_G do not satisfy the conclusion of [Theorem 1](#). Suppose for contradiction that there exist subgroups $\mathbf{P}_G, \mathbf{S}_G \subset \mathbf{G}$ forming a Siegel triple $(\mathbf{P}_G, \mathbf{S}_G, K_G)$, a Siegel set $\mathfrak{S}_G \subset \mathbf{G}(\mathbb{R})$ with respect to this Siegel triple, and a finite set $C \subset \mathbf{G}(\mathbb{Q})$ such that $\mathfrak{S}_H \subset C \cdot \mathfrak{S}_G$. By the same argument as in [Section B](#), we may assume that $\mathbf{P}_G = \mathbf{P}_0$ and $\mathbf{S}_G = \mathbf{S}_0$.

Let $\sigma_s = \mathrm{diag}(s, 1, s^{-1})$ for $s \in \mathbb{R}_{>0}$. Now

$$\{\eta\sigma_s\eta^{-1} : s \geq t\} = A_{H,t} \subset \mathfrak{S}_H \subset C\mathfrak{S}_G.$$

Since C is finite, there exists some $\gamma \in C$ such that $\gamma\mathfrak{S}_G$ contains elements of the form $\eta\sigma_s\eta^{-1}$ for arbitrarily large s . Consequently $\eta^{-1}\gamma\mathfrak{S}_G\eta$ contains σ_s for arbitrarily large s . Furthermore the standard Siegel set \mathfrak{S}_G contains $\{\sigma_s : s \geq t'\}$ for some $t' \in \mathbb{R}_{>0}$.

Let χ_1, χ_2 denote the simple roots of \mathbf{G} with respect to \mathbf{S}_0 , using the ordering induced by \mathbf{P}_0 . Then $\chi_1(\sigma_s) = \chi_2(\sigma_s) = s$ so the previous paragraph shows that $\mathfrak{S}_G \cap \eta^{-1}\gamma\mathfrak{S}_G\eta$ contains elements σ_s with arbitrarily large values for χ_1 and χ_2 . Applying [Lemma 3](#) below (with $\Omega_G = K_G \cup K_G\eta^{-1}$), we deduce that $\eta^{-1}\gamma$ is contained in the standard parabolic subgroup ${}_{\mathbb{Q}}\mathbf{P}_{0,\emptyset} = \mathbf{P}_0$.

Let $U_0 = R_u(\mathbf{P}_0)$. Write the Iwasawa decomposition of η^{-1} as

$$\eta^{-1} = \mu\alpha\kappa \quad \text{where } \mu \in U_0(\mathbb{R}), \alpha \in \mathbf{S}_0(\mathbb{R}), \kappa \in K_G.$$

For arbitrarily large real numbers s , we have

$$\sigma_s \mu \sigma_s^{-1} \cdot \sigma_s \alpha \cdot \kappa = \sigma_s \eta^{-1} \in \mathfrak{S}_G \eta^{-1} \cap \eta^{-1} \gamma \mathfrak{S}_G \subset \eta^{-1} \gamma \mathfrak{S}_G.$$

By the definition of Siegel sets and since $\eta^{-1}\gamma \in \mathbf{P}_0(\mathbb{R})$, the $U_0(\mathbb{R})$ -component in the Iwasawa decomposition of every element of $\eta^{-1}\gamma\mathfrak{S}_G$ is bounded. Thus $\sigma_s \mu \sigma_s^{-1}$ lies in a bounded set for arbitrarily large real numbers s . By direct calculation, this implies that $\mu = 1$. (This is the opposite situation to [\[Borel 1969, Lemme 12.2\]](#), adapted to our conventions about Siegel sets.) Hence $\eta^{-1} = \alpha\kappa \in \mathbf{S}_0(\mathbb{R})K_G$.

It follows that $\eta^t \eta = (\alpha^{-1})^t (\kappa^{-1})^t \kappa^{-1} \alpha^{-1} = \alpha^{-2}$ is diagonal. But $\eta^t \eta$ is not diagonal, as can be seen either by direct calculation or by noting that η is symmetrical so $\eta^t \eta = \eta^2$ has L as a 1-dimensional eigenspace yet L is not a coordinate axis.

D. Siegel sets with noncompact intersection. In this section, we prove a generalisation of [\[Borel 1969, Proposition 12.6\]](#), replacing a Siegel set $\mathfrak{S} = \Omega_P A_t K$ by a set of the form $\Omega_P A_t \Omega_G$ where Ω_G may be any compact subset of $\mathbf{G}(\mathbb{R})$. This generalisation was used in [Section C](#).

Let G be a reductive \mathbb{Q} -algebraic group. Let P be a minimal parabolic \mathbb{Q} -subgroup of G and let U be the unipotent radical of P . Let S be a maximal \mathbb{Q} -split torus in S and let M be the maximal \mathbb{Q} -anisotropic subgroup of $Z_G(S)$. Let t be a positive real number and let A_t be the subset of $S(\mathbb{R})$ defined in [Orr 2018, Section 2B]. Let \mathfrak{g} and \mathfrak{u} denote the Lie algebras of G and U respectively (over \mathbb{R}).

Let Δ be the set of simple roots of G with respect to S , using the ordering induced by P . For $\theta \in \Delta$, let Ψ_θ denote the set of roots ϕ such that the expression of ϕ as a linear combination of elements of Δ has a positive coefficient for at least one element of θ .

For each character $\chi \in X^*(S)$, there is a unique continuous group homomorphism $P(\mathbb{R}) \rightarrow \mathbb{R}_{>0}$, which we denote f_χ , with the properties $f_\chi(s) = |\chi(s)|$ for all $s \in S(\mathbb{R})$ and $f_\chi = 1$ on $U(\mathbb{R})M(\mathbb{R})$. (This is because $S(\mathbb{R}) \cap U(\mathbb{R})M(\mathbb{R})$ is finite, so $|\chi(s)| = 1$ for all $s \in S(\mathbb{R}) \cap U(\mathbb{R})M(\mathbb{R})$, and S normalises UM .) Choose a maximal compact subgroup $K \subset G(\mathbb{R})$. Then $f_\chi(P(\mathbb{R}) \cap K)$ is a compact subgroup of $\mathbb{R}_{>0}$, so is trivial. Therefore we can extend f_χ to a continuous function $G(\mathbb{R}) = P(\mathbb{R})K \rightarrow \mathbb{R}_{>0}$ by setting $f_\chi(pk) = f_\chi(p)$ for all $p \in P(\mathbb{R})$ and $k \in K$. These functions f_χ are not necessarily “of type (P, χ) ” as defined in [Borel 1969, 14.1] because $\chi \in X^*(S)$ might not extend to a character of P , but the argument in [loc. cit., 14.2(c)] still applies to the functions f_χ .

Lemma 3. *Let Ω_P and Ω_G be compact subsets of $P(\mathbb{R})$ and $G(\mathbb{R})$ respectively. Let $\gamma \in G(\mathbb{R})$. If $\Omega_P A_t \Omega_G \cap \gamma \Omega_P A_t \Omega_G$ is noncompact, then γ is contained in a proper parabolic \mathbb{Q} -algebraic subgroup of G containing P . More precisely, let*

$$\theta = \{\chi \in \Delta : f_\chi \text{ is bounded above on } \Omega_P A_t \Omega_G \cap \gamma \Omega_P A_t \Omega_G\}.$$

Then γ lies in the standard parabolic subgroup ${}_{\mathbb{Q}}P_\theta$ in the notation of [Borel and Tits 1965, 5.12].

Proof. Let

$$\Omega = \left(\bigcup_{a \in A_t} a^{-1} \Omega_P a \right) \Omega_G \subset G(\mathbb{R}).$$

By [Borel 1969, Lemme 12.2], Ω is compact. From the definitions, $\Omega_P A_t \Omega_G \subset A_t \Omega$. Hence, for all $\chi \in \Delta \setminus \theta$, f_χ is unbounded on $A_t \Omega \cap \gamma A_t \Omega$.

Let ${}_{\mathbb{Q}}U_\theta$ denote the unipotent radical of ${}_{\mathbb{Q}}P_\theta$ and let ${}_{\mathbb{Q}}\mathfrak{u}_\theta = \text{Lie}({}_{\mathbb{Q}}U_\theta)$. Let

$$Y = \{v \in \mathfrak{g} : (\text{Ad } \xi_n^{-1})v \rightarrow 0 \text{ for some sequence } (\xi_n) \text{ in } A_t \Omega \cap \gamma A_t \Omega\}.$$

Let $\langle Y \rangle$ denote the subspace of \mathfrak{g} generated by Y . We shall show that

$${}_{\mathbb{Q}}\mathfrak{u}_\theta \subset \langle Y \rangle \subset (\text{Ad } \gamma)\mathfrak{u}. \tag{1}$$

To prove the first inclusion of (1), note that ${}_{\mathbb{Q}}\mathfrak{u}_\theta$ is the direct sum of the root spaces \mathfrak{u}_ϕ for $\phi \in \Psi_{\Delta \setminus \theta}$, so it suffices to prove that $\mathfrak{u}_\phi \subset Y$ for each $\phi \in \Psi_{\Delta \setminus \theta}$.

Let $\phi \in \Psi_{\Delta \setminus \theta}$ and write ϕ as a linear combination of simple roots: $\phi = \sum_{\psi \in \Delta} m_\psi \psi$. By the definition of $\Psi_{\Delta \setminus \theta}$, there exists some $\chi \in \Delta \setminus \theta$ such that $m_\chi > 0$.

By the definition of θ , f_χ is unbounded on $\Omega_P A_t \Omega_G \cap \gamma \Omega_P A_t \Omega_G \subset A_t \Omega \cap \gamma A_t \Omega$. Choose a sequence (ξ_n) in $A_t \Omega \cap \gamma A_t \Omega$ such that $f_\chi(\xi_n) \rightarrow +\infty$. Write $\xi_n = \alpha_n \kappa_n$ where $\alpha_n \in A_t$ and $\kappa_n \in \Omega$.

The argument of [Borel 1969, 14.2(c)] shows that $f_\chi(\xi_n)/f_\chi(\alpha_n)$ is bounded both above and below independently of n . Hence

$$|\chi(\alpha_n)| = f_\chi(\alpha_n) \rightarrow +\infty.$$

Since ϕ is a positive root, $m_\psi \geq 0$ for all $\psi \in \Delta$. Since $\alpha_n \in A_t$ and $m_\chi > 0$, it follows that $\phi(\alpha_n) \rightarrow +\infty$.

Hence for every $v \in \mathfrak{u}_\phi$, we have $(\text{Ad } \alpha_n^{-1})v \rightarrow 0$. Since Ω is compact, after replacing (ξ_n) by a subsequence, we may assume that κ_n converges, say to $\kappa \in \Omega$. Then $(\text{Ad } \xi_n^{-1})v \rightarrow (\text{Ad } \kappa)^{-1}0 = 0$. Thus $\mathfrak{u}_\phi \subset Y$.

To prove the second inclusion of (1), consider an element $v \in Y$. Let (ξ_n) be a sequence in $A_t \Omega \cap \gamma A_t \Omega$ such that $(\text{Ad } \xi_n^{-1})v \rightarrow 0$. Write $\xi_n = \gamma \beta_n \lambda_n$ with $\beta_n \in A_t$, $\lambda_n \in \Omega$. Since Ω is compact, after replacing (ξ_n) by a subsequence, we may assume that λ_n converges, say to $\lambda \in \Omega$. Then

$$(\text{Ad } \beta_n^{-1})(\text{Ad } \gamma^{-1})v = (\text{Ad } \lambda_n)(\text{Ad } \xi_n^{-1})v \rightarrow (\text{Ad } \lambda)0 = 0.$$

Hence, when we decompose $(\text{Ad } \gamma^{-1})v$ using the root space decomposition of \mathfrak{g} , nonzero components can occur only for those roots ϕ satisfying $|\phi(\beta_n)| \rightarrow +\infty$. Since $\beta_n \in A_t$, such roots ϕ must be positive roots. Thus $(\text{Ad } \gamma^{-1})v \in \bigoplus_{\phi \in \Phi^+} \mathfrak{u}_\phi = \mathfrak{u}$.

We have proved both parts of (1). Passing from Lie algebras to groups, we obtain

$$\mathbb{Q}U_\theta \subset \gamma U \gamma^{-1} \subset \gamma P \gamma^{-1} \subset \gamma \mathbb{Q}P_\theta \gamma^{-1}.$$

By [Borel and Tits 1965, Corollaire 4.5], it follows that $\mathbb{Q}P_\theta = \gamma \mathbb{Q}P_\theta \gamma^{-1}$. Since a parabolic subgroup of G is its own normaliser, we conclude that $\gamma \in \mathbb{Q}P_\theta(\mathbb{R})$. □

E. Additional minor corrections to [Orr 2018]. The following are additional corrections to [Orr 2018]:

- (page 461, Section 2D) (F2) should begin “For every $g \in G(\mathbb{Q})$.”
- (page 474, proof of Proposition 4.7) On the fifth line from the end, should say “ $\chi|_{S_H} \in \Phi_\alpha \cup \{0\}$.” instead of “ $\chi|_{S_H} \in \Phi_\alpha$ ”.
- (page 474, proof of Lemma 4.10) The first paragraph should say “Let T_G be a maximal \mathbb{R} -split torus in G which contains S_G and is stabilised by the Cartan involution of G associated with K_G .” This is necessary to apply [Borel and Tits 1965, Section 14].

Acknowledgements

Orr is grateful to Dave Witte Morris for informing him of the errors in [Orr 2018, page 474] which are corrected in Section E.

We are grateful to the referee for very careful reading of this note and for suggesting alternative arguments for Sections C and D, including a more general statement for Lemma 3 than in the original version.

References

- [Bakker et al. 2020] B. Bakker, B. Klingler, and J. Tsimerman, “Tame topology of arithmetic quotients and algebraicity of Hodge loci”, *J. Amer. Math. Soc.* **33**:4 (2020), 917–939. [MR](#) [Zbl](#)
- [Borel 1969] A. Borel, *Introduction aux groupes arithmétiques*, Publ. Inst. Math. Univ. Strasbourg **1341**, Hermann, Paris, 1969. [MR](#) [Zbl](#)
- [Borel and Tits 1965] A. Borel and J. Tits, “Groupes réductifs”, *Inst. Hautes Études Sci. Publ. Math.* **27** (1965), 55–150. [MR](#) [Zbl](#)
- [Daw and Orr 2021] C. Daw and M. Orr, “Unlikely intersections with $E \times \text{CM}$ curves in \mathcal{A}_2 ”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **22**:4 (2021), 1705–1745. [MR](#) [Zbl](#)
- [Mostow 1955] G. D. Mostow, “Self-adjoint groups”, *Ann. of Math. (2)* **62** (1955), 44–55. [MR](#) [Zbl](#)
- [Orr 2018] M. Orr, “Height bounds and the Siegel property”, *Algebra Number Theory* **12**:2 (2018), 455–478. [MR](#) [Zbl](#)

Communicated by Jonathan Pila

Received 2021-10-14 Revised 2022-06-22 Accepted 2022-07-06

martin.orr@manchester.ac.uk

Department of Mathematics, University of Manchester, Manchester, United Kingdom

christian.schnell@stonybrook.edu

Department of Mathematics, Stony Brook University, Stony Brook, NY, United States

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2023 is US \$485/year for the electronic version, and \$705/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 17 No. 6 2023

On Héthelyi–Külshammer’s conjecture for principal blocks	1127
NGUYEN NGOC HUNG and A. A. SCHAEFFER FRY	
Shintani–Barnes cocycles and values of the zeta functions of algebraic number fields	1153
HOHTO BEKKI	
On the commuting probability of p-elements in a finite group	1209
TIMOTHY C. BURNES, ROBERT GURALNICK, ALEXANDER MORETÓ and GABRIEL NAVARRO	
Correction to the article Height bounds and the Siegel property	1231
MARTIN ORR and CHRISTIAN SCHNELL	