

Algebra & Number Theory

Volume 18

2024

No. 4

**Fundamental exact sequence
for the pro-étale fundamental group**

Marcin Lara



Fundamental exact sequence for the pro-étale fundamental group

Marcin Lara

The pro-étale fundamental group of a scheme, introduced by Bhatt and Scholze, generalizes formerly known fundamental groups — the usual étale fundamental group $\pi_1^{\text{ét}}$ defined in SGA1 and the more general π_1^{SGA3} . It controls local systems in the pro-étale topology and leads to an interesting class of “geometric coverings” of schemes, generalizing finite étale coverings.

We prove exactness of the fundamental sequence for the pro-étale fundamental group of a geometrically connected scheme X of finite type over a field k , i.e., that the sequence

$$1 \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k \rightarrow 1$$

is exact as abstract groups and the map $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ is a topological embedding.

On the way, we prove a general van Kampen theorem and the Künneth formula for the pro-étale fundamental group.

1. Introduction	631
2. Infinite Galois categories, Noohi groups and $\pi_1^{\text{proét}}$	637
3. Seifert–van Kampen theorem for $\pi_1^{\text{proét}}$ and its applications	649
4. Fundamental exact sequence	664
Acknowledgements	682
References	683

1. Introduction

Bhatt and Scholze [2015] introduced the pro-étale topology for schemes. The main motivation was that the definitions of ℓ -adic sheaves and cohomologies in the usual étale topology are rather indirect. In contrast, the naive definition of, e.g., a constant \mathbb{Q}_ℓ -sheaf in the pro-étale topology as $X_{\text{proét}} \ni U \mapsto \text{Maps}_{\text{cts}}(U, \mathbb{Q}_\ell)$ is a sheaf and if X is a variety over an algebraically closed field, then $H^i(X_{\text{ét}}, \mathbb{Q}_\ell) = H^i(X_{\text{proét}}, \mathbb{Q}_\ell)$, where the right-hand side is defined “naively” by applying the derived functor $\text{R}\Gamma(X_{\text{proét}}, -)$ to the described constant sheaf.

Along with the new topology, Bhatt and Scholze [2015] introduced a new fundamental group — *the pro-étale fundamental group*. It is defined for a connected locally topologically noetherian scheme X with

MSC2010: primary 14F35; secondary 14D10, 14F20, 20E06.

Keywords: pro-étale topology, pro-étale fundamental group, étale fundamental group, homotopy exact sequence, fundamental exact sequence, Noohi groups.

a geometric point \bar{x} and denoted $\pi_1^{\text{proét}}(X, \bar{x})$. The name “pro-étale” is justified by the fact that there is an equivalence $\pi_1^{\text{proét}}(X, \bar{x}) - \text{Sets} \simeq \text{Loc}_{X_{\text{proét}}}$ between the categories of (possibly infinite) discrete sets with continuous action by $\pi_1^{\text{proét}}(X, \bar{x})$ and locally constant sheaves of (discrete) sets in $X_{\text{proét}}$. This is analogous to the classical fact that $\pi_1^{\text{ét}}(X, \bar{x}) - \text{FSets}$ is equivalent to the category of lcc sheaves on $X_{\text{ét}}$, where $G - \text{FSets}$ denotes *finite* sets with a continuous G action. This is the first striking difference between these fundamental groups: $\pi_1^{\text{proét}}$ allows working with sheaves of infinite sets. In fact, Bhatt and Scholze [2015] study abstract “infinite Galois categories”, which are pairs (\mathcal{C}, F) satisfying certain axioms that (together with an additional tameness condition) turn out to be equivalent to a pair $(G - \text{Sets}, F_G : G - \text{Sets} \rightarrow \text{Sets})$ for a Hausdorff topological group G and the forgetful functor F_G . In fact, one takes $G = \text{Aut}(F)$ with a suitable topology. This generalizes the usual Galois categories, introduced by Grothendieck to define $\pi_1^{\text{ét}}(X, \bar{x})$. In Grothendieck’s approach, one takes the category FÉt_X of finite étale coverings together with the fiber functor $F_{\bar{x}}$ and obtains that $\pi_1^{\text{ét}}(X, \bar{x}) - \text{FSets} \simeq \text{FÉt}_X$. Discrete sets with a continuous $\pi_1^{\text{proét}}(X, \bar{x})$ -action correspond to a larger class of coverings, namely “geometric coverings”, which are defined to be schemes Y over X such that $Y \rightarrow X$:

- (1) Is étale (not necessarily quasicompact!).
- (2) Satisfies the valuative criterion of properness.

We denote the category of geometric coverings by Cov_X (seen as a full subcategory of Sch_X). It is clear that $\text{FÉt} \subset \text{Cov}_X$. As Y is not assumed to be of finite type over X , the valuative criterion does not imply that $Y \rightarrow X$ is proper (otherwise we would get finite étale morphisms again) and so in general we get more. A basic example of a nonfinite covering in Cov_X can be obtained by viewing an infinite chain of (suitably glued) \mathbb{P}_k^1 s as a covering of the nodal curve $X = \mathbb{P}^1 / \{0, 1\}$ obtained by gluing 0 and 1 on \mathbb{P}_k^1 (to formalize the gluing one can use [Schwede 2005]). Then, if $k = \bar{k}$, $\pi_1^{\text{proét}}(X, \bar{x}) = \mathbb{Z}$ and $\pi_1^{\text{ét}}(X, \bar{x}) = \widehat{\mathbb{Z}}$. In this example, the prodiscrete group π_1^{SGA3} defined in Chapter X.6 of [SGA 3_{II} 1970] would give the same answer. This is essentially because our infinite covering is a torsor under a discrete group in $X_{\text{ét}}$. However, for more general schemes (e.g., an elliptic curve with two points glued), the category Cov_X contains more. So far, all the new examples were coming from nonnormal schemes. This is not a coincidence, as for a normal scheme X , any $Y \in \text{Cov}_X$ is a (possibly infinite) disjoint union of finite étale coverings. In this case, $\pi_1^{\text{proét}}(X, \bar{x}) = \pi_1^{\text{SGA3}}(X, \bar{x}) = \pi_1^{\text{ét}}(X, \bar{x})$. In general $\pi_1^{\text{ét}}$ can be recovered as the profinite completion of $\pi_1^{\text{proét}}$ and π_1^{SGA3} is the prodiscrete completion of $\pi_1^{\text{proét}}$.

The groups $\pi_1^{\text{proét}}$ belong in general to a class of *Noohi groups*. These can be characterized as Hausdorff topological groups G that are Raïkov complete and such that the open subgroups form a basis of neighborhoods at 1_G . However, *open normal subgroups do not necessarily form a basis* of open neighborhoods of 1_G in a Noohi group. In the case of $\pi_1^{\text{proét}}$, this means that there might exist a connected $Y \in \text{Cov}_X$ that do not have a Galois closure. Examples of Noohi groups include: profinite groups, (pro)discrete groups, but also \mathbb{Q}_ℓ and $\text{GL}_n(\mathbb{Q}_\ell)$. A slightly different example would be $\text{Aut}(S)$, where S is a discrete set and Aut has the compact-open topology.

The fact that groups like $GL_n(\mathbb{Q}_\ell)$ are Noohi (but not profinite or prodiscrete) makes $\pi_1^{\text{proét}}$ better suited to work with \mathbb{Q}_ℓ (or $\overline{\mathbb{Q}}_\ell$) local systems. Indeed, denoting by $\text{Loc}_{X_{\text{proét}}}(\mathbb{Q}_\ell)$ the category of \mathbb{Q}_ℓ -local systems on $X_{\text{proét}}$, i.e., locally constant sheaves of finite-dimensional \mathbb{Q}_ℓ -vector spaces (again, the “naive” definition works in $X_{\text{proét}}$), one has an equivalence $\text{Rep}_{\text{cts}, \mathbb{Q}_\ell}(\pi_1^{\text{proét}}(X, \bar{x})) \simeq \text{Loc}_{X_{\text{proét}}}(\mathbb{Q}_\ell)$. This fails for $\pi_1^{\text{ét}}$, as any \mathbb{Q}_ℓ -representation of a profinite group must stabilize a \mathbb{Z}_ℓ -lattice, while \mathbb{Q}_ℓ -local systems (in the above sense) stabilize lattices only étale locally. The group $\pi_1^{\text{SGA}3}$ is not enough either; as shown by [Bhatt and Scholze 2015, Example 7.4.9] (due to Deligne), if X is the scheme obtained by gluing two points on a smooth projective curve of suitably large genus, there are \mathbb{Q}_ℓ -local systems on X that do not come from a representation of $\pi_1^{\text{SGA}3}(X)$.

We will often drop \bar{x} from the notation for brevity. This usually does not matter much, as a different choice of the base point leads to an isomorphic group.

Classical results. In [SGA 1 1971], Grothendieck proved some foundational results regarding the étale fundamental group. Among them:

(1) The fundamental exact sequence, i.e., the comparison between the “arithmetic” and “geometric” fundamental groups:

Theorem 1.1 [SGA 1 1971, Exposé IX, Théorème 6.1]. *Let k be a field with algebraic closure \bar{k} . Let X be a quasicompact and quasiseparated scheme over k . If the base change $X_{\bar{k}}$ is connected, then there is a short exact sequence*

$$1 \rightarrow \pi_1^{\text{ét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{ét}}(X) \rightarrow \text{Gal}_k \rightarrow 1$$

of profinite topological groups.

(2) The homotopy exact sequence:

Theorem 1.2 [SGA 1 1971, Exposé X, Corollaire 1.4]. *Let $f : X \rightarrow S$ be a flat proper morphism of finite presentation whose geometric fibers are connected and reduced. Assume S is connected and let \bar{s} be a geometric point of S . Then there is an exact sequence*

$$\pi_1^{\text{ét}}(X_{\bar{s}}) \rightarrow \pi_1^{\text{ét}}(X) \rightarrow \pi_1^{\text{ét}}(S) \rightarrow 1$$

of fundamental groups.

(3) “Künneth formula”:

Proposition 1.3 [SGA 1 1971, Exposé X, Corollary 1.7]. *Let X, Y be two connected schemes locally of finite type over an algebraically closed field k and assume that Y is proper. Let \bar{x}, \bar{y} be geometric points of X and Y respectively with values in the same algebraically closed field extension K of k . Then the map induced by the projections is an isomorphism*

$$\pi_1^{\text{ét}}(X \times_k Y, (\bar{x}, \bar{y})) \xrightarrow{\sim} \pi_1^{\text{ét}}(X, \bar{x}) \times \pi_1^{\text{ét}}(Y, \bar{y}).$$

- (4) Invariance of $\pi_1^{\acute{e}t}$ under extensions of algebraically closed fields for proper schemes [SGA 1 1971, Exposé X, Corollaire 1.8].
- (5) General van Kampen theorem (proved in a special case in [SGA 1 1971, IX Section 5] and generalized in [Stix 2006]).

The aim of this and the subsequent article [Lara 2022] is to generalize statements (1) and (2), correspondingly, to the case of $\pi_1^{\text{proét}}$. In the present article, we also establish the generalizations of all the other points besides (2). The main difficulties in trying to directly generalize the proofs of Grothendieck are as follows:

- Geometric coverings of schemes (i.e., elements of Cov_X defined above) are often not quasicompact, unlike elements of FÉt_X . For example, for X a variety over a field k and connected $Y \in \text{Cov}_{X_{\bar{k}}}$, there may be no finite extension l/k such that Y would be defined over l . Similarly, some useful constructions (like Stein factorization) no longer work (at least without significant modifications).
- For a connected geometric covering $Y \in \text{Cov}_X$, there is in general no Galois geometric covering dominating it. Equivalently, there might exist an open subgroup $U < \pi_1^{\text{proét}}(X)$ that does not contain an open normal subgroup. This prevents some proofs that would work for π_1^{SGA3} to carry over to $\pi_1^{\text{proét}}$.
- The topology of $\pi_1^{\text{proét}}$ is more complicated than the one of $\pi_1^{\acute{e}t}$, e.g., it is not necessarily compact, which complicates the discussion of exactness of sequences.

Our results. Our main theorem is the generalization of the fundamental exact sequence. More precisely, we prove the following:

Theorem (Theorem 4.14). *Let X be a geometrically connected scheme of finite type over a field k . Then the sequence*

$$1 \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k \rightarrow 1$$

is exact as abstract groups.

Moreover, the map $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ is a topological embedding and the map $\pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k$ is a quotient map of topological groups.

The most difficult part is showing that $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ is injective or, more precisely, a topological embedding. This is Theorem 4.13.

As in the case of usual Galois categories, statements about exactness of sequences of Noohi groups translate to statements on the corresponding categories of G – Sets. If the groups involved are the pro-étale fundamental groups, this translates to statements about geometric coverings. We give a detailed dictionary in Proposition 2.37. As Noohi groups are not necessarily compact, the statements on coverings are equivalent to some weaker notions of exactness (e.g., preserving connectedness of coverings is equivalent to the map of groups having dense image). In fact, we first prove a “near-exact” version of Theorem 4.14 and obtain the above one as a corollary using an extra argument.

For $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ to be a topological embedding boils down to the following statement: Every geometric covering Y of $X_{\bar{k}}$ can be dominated by a covering Y' that embeds into a base-change to \bar{k} of a geometric covering Y'' of X (i.e., defined over k):

$$\begin{array}{ccc} Y' \subset Y''_{\bar{k}} & \longrightarrow & Y'' \\ \downarrow & & \\ Y & & \end{array}$$

For finite coverings, the analogous statement is easy to prove; by finiteness, the given covering is defined over a finite field extension l/k and one concludes quickly. This is also the case for infinite coverings detected by $\pi_1^{\text{SGA}3}$, see [Proposition 4.8](#). But for general geometric coverings, the situation is much less obvious; as we show by counterexamples ([Examples 4.5](#) and [4.6](#)), it is **not true in general that a connected geometric covering of $X_{\bar{k}}$ is isomorphic to a base-change of a covering of X_l for some finite extension l/k** . This property is crucially used in the proof of [\[SGA 1 1971, Exposé IX, Theorem 6.1\]](#), and thus *trying to carry the classical proof of SGA over to $\pi_1^{\text{proét}}$ fails*. This last statement is, however, stronger than what we need to prove, and so does not contradict our theorem.

A useful technical tool across the article is the *van Kampen theorem* for $\pi_1^{\text{proét}}$. Its abstract form is proven by adapting the proof in [\[Stix 2006\]](#) to the case of Noohi groups and infinite Galois categories. For a morphism of schemes $X' \rightarrow X$ of effective descent for Cov (satisfying some extra conditions), it allows one to write the pro-étale fundamental group of X in terms of the pro-étale fundamental groups of the connected components of X' and certain relations. By the results of [\[Rydh 2010\]](#), one can take $X' = X^\nu \rightarrow X$ to be the normalization morphism of a Nagata scheme X . As $\pi_1^{\text{proét}}$ and $\pi_1^{\text{ét}}$ coincide for normal schemes, this allows us to present $\pi_1^{\text{proét}}(X)$ in terms of $\pi_1^{\text{ét}}(X_w^\nu)$, where $X^\nu = \bigsqcup_w X_w^\nu$, and the (discrete) topological fundamental group of a suitable graph. In this case, the van Kampen theorem takes on concrete form and generalizes [\[Lavanda 2018, Theorem 1.17\]](#).

Theorem (van Kampen theorem, [Corollary 3.19](#), [Remark 3.21](#), [Proposition 3.12](#); compare [\[Stix 2006\]](#)). *Let X be a Nagata scheme and $X^\nu = \bigsqcup_w X_w^\nu$ its normalization written as a union of connected components. Then, after a choice of geometric points, étale paths between them and a maximal tree T within a suitable “intersection” graph Γ , there is an isomorphism*

$$\pi_1^{\text{proét}}(X, \bar{x}) \simeq ((*_w^{\text{top}} \pi_1^{\text{ét}}(X_w^\nu, \bar{x}_w) *_w^{\text{top}} \pi_1^{\text{top}}(\Gamma, T)) / \langle R_1, R_2 \rangle)^{\text{Noohi}}$$

where R_1, R_2 are two sets of relations described in [Corollary 3.19](#) and $(-)^{\text{Noohi}}$ is the **Noohi completion** defined in [Section 2](#).

In the proof of the main theorem, the van Kampen theorem allows us to construct $\pi_1^{\text{proét}}(X_{\bar{k}})$ - and $\pi_1^{\text{proét}}(X)$ -sets in more concrete terms of graphs of groups involving the $\pi_1^{\text{ét}}$. We “explicitly” construct a Galois invariant open subgroup of a given open subgroup $U < \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ in terms of “regular loops” (with respect to U), see [Definition 4.20](#).

In fact, the existence of elements that are too far from being a product of regular loops is tacitly behind the counterexamples Examples 4.5 and 4.6, while the fact that, despite this, there is still an abundance of (products of) regular loops (i.e., their closure is open) is behind our main proof. We also sketch a quicker but less constructive approach in Remark 4.27.

Another interesting result proven with the help of the van Kampen theorem is the Künneth formula.

Proposition (Künneth formula for $\pi_1^{\text{proét}}$, Proposition 3.29). *Let X, Y be two connected schemes locally of finite type over an algebraically closed field k and assume that Y is proper. Let \bar{x}, \bar{y} be geometric points of X and Y respectively with values in the same algebraically closed field extension K of k . Then the map induced by the projections is an isomorphism*

$$\pi_1^{\text{proét}}(X \times_k Y, (\bar{x}, \bar{y})) \xrightarrow{\sim} \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y}).$$

Along the way, we prove the invariance of $\pi_1^{\text{proét}}$ under extensions of algebraically closed fields for proper schemes (see Proposition 3.31) and give a short direct proof of the fact that $\pi_1^{\text{SGA}^3}(X_{\bar{k}}, \bar{x}) \hookrightarrow \pi_1^{\text{SGA}^3}(X, \bar{x})$, see Corollary 4.10.

In a separate article [Lara 2022], we discuss the homotopy exact sequence for $\pi_1^{\text{proét}}$. It is proven by constructing an infinite (i.e., nonquasicompact) analogue of the Stein factorization. Although the construction does not use the main results of this article, the auxiliary results on Noohi groups and $\pi_1^{\text{proét}}$ have proven to be very handy.

We hope that our techniques, with some extra tweaks and work, will allow to draw similar conclusions about other Noohi fundamental groups arising from the infinite Galois formalism. One such example could be the de Jong fundamental group π_1^{DJ} , defined in the rigid-analytic setting in [de Jong 1995]. In a later joint work [Achinger et al. 2022], we have proven the existence of a specialization morphism between $\pi_1^{\text{proét}}$ and π_1^{DJ} , relating $\pi_1^{\text{proét}}$ to this more established fundamental group.

1A. Conventions and notations.

- For us, compact = quasicompact + Hausdorff.
- $H <^\circ G$ will mean that H is an open subgroup of G .
- For subgroups $H < G$, H^{nc} will denote the normal closure of H in G , i.e., the smallest normal subgroup of G containing H . We will use $\langle\langle - \rangle\rangle$ to denote the normal closure of the subgroup generated by some subset of G , i.e., $\langle\langle - \rangle\rangle = \langle - \rangle^{nc}$.
- For a field k , we will use \bar{k} to denote its (fixed) algebraic closure and k^{sep} or k^s to denote its separable closure (in \bar{k}).
- The topological groups are assumed to be Hausdorff unless specified otherwise or appearing in a context where it is not automatically satisfied (e.g., as a quotient by a subgroup that is not necessarily closed). We will usually comment whenever a non-Hausdorff group appears.
- We assume (almost) every base scheme to be locally topologically noetherian. This does not cause problems when considering geometric coverings, as a geometric covering of a locally topologically

noetherian scheme is locally topologically noetherian again — this is [Bhatt and Scholze 2015, Lemma 6.6.10].

- A “ G -set” for a topological group G will mean a discrete set with a continuous action of G unless specified otherwise. We will denote the category of G -sets by $G - \text{Sets}$. We will denote the category of sets by Sets .
- We will often omit the base points from the statements and the discussion; by Corollary 3.18, this usually does not change much. In some proofs (e.g., involving the van Kampen theorem), we keep track of the base points.

2. Infinite Galois categories, Noohi groups and $\pi_1^{\text{proét}}$

2A. Overview of the results in [Bhatt and Scholze 2015]. Throughout the entire article we use the language and results of [Bhatt and Scholze 2015], especially of Chapter 7, as this is where the pro-étale fundamental group was defined. Some familiarity with the results of [loc. cit., Section 7] is a prerequisite to read this article. We are going to give a quick overview of some of these results below, but we recommend keeping a copy of [loc. cit.] at hand.

Definition 2.1 [Bhatt and Scholze 2015, Definition 7.1.1]. Fix a topological group G . Let $G - \text{Sets}$ be the category of discrete sets with a continuous G -action, and let $F_G : G - \text{Sets} \rightarrow \text{Sets}$ be the forgetful functor. We say that G is a *Noohi group* if the natural map induces an isomorphism $G \rightarrow \text{Aut}(F_G)$ of topological groups. Here, $S \in \text{Sets}$ are considered with the discrete topology, $\text{Aut}(S)$ with the compact-open topology and $\text{Aut}(F_G)$ is topologized using $\text{Aut}(F_G(S))$ for $S \in G - \text{Sets}$. More precisely, the stabilizers $\text{Stab}_{F(S),s}^{\text{Aut}(F_G)}$ for connected $S \in G - \text{Sets}$, $s \in F(S)$, form a basis of neighborhoods of $1 \in \text{Aut}(F_G)$.

In particular, it follows from the definition that open subgroups form a basis of neighborhoods of 1 in a Noohi group. Now, by [Bhatt and Scholze 2015, Proposition 7.1.5], it follows that a topological group is Noohi if and only if it satisfies the following conditions:

- Its open subgroups form a basis of open neighborhoods of $1 \in G$.
- It is Raïkov complete.

A topological group G is Raïkov complete if it is complete for its two-sided uniformity (see [Dikranjan 2013] or [Arhangel’skii and Tkachenko 2008, Chapter 3.6] for an introduction to the Raïkov completion). Using the above proposition it is easy to give examples of Noohi groups.

Example 2.2. The following classes of topological groups are Noohi: discrete groups, profinite groups, $\text{Aut}(S)$ with the compact-open topology for S a discrete set (see [Bhatt and Scholze 2015, Lemma 7.1.4]), groups containing an open subgroup which is Noohi; see [loc. cit., Lemma 7.1.8].

The following groups are Noohi: $\mathbb{Q}_\ell, \overline{\mathbb{Q}_\ell}$ for the colimit topology induced by expressing $\overline{\mathbb{Q}_\ell}$ as a union of finite extensions (in contrast with the situation for the ℓ -adic topology), $\text{GL}_n(E)$ for any algebraic extension E/\mathbb{Q}_ℓ and the colimit topology on $\text{GL}_n(E)$; see [loc. cit., Example 7.1.7].

The notion of a Noohi group is tightly connected to a notion of an infinite Galois category, which we are about to introduce. Here, an object $X \in \mathcal{C}$ is called connected if it is not empty (i.e., initial), and for every subobject $Y \rightarrow X$ (i.e., $Y \xrightarrow{\sim} Y \times_X Y$), either Y is empty or $Y = X$.

Definition 2.3 [Bhatt and Scholze 2015, Definition 7.2.1]. An *infinite Galois category* \mathcal{C} is a pair $(\mathcal{C}, F : \mathcal{C} \rightarrow \text{Sets})$ satisfying:

- (1) \mathcal{C} is a category admitting colimits and finite limits.
- (2) Each $X \in \mathcal{C}$ is a disjoint union of connected (in the sense explained above) objects.
- (3) \mathcal{C} is generated under colimits by a set of connected objects.
- (4) F is faithful, conservative, and commutes with colimits and finite limits.

The *fundamental group* of (\mathcal{C}, F) is the topological group $\pi_1(\mathcal{C}, F) := \text{Aut}(F)$, topologized by the compact-open topology on $\text{Aut}(S)$ for any $S \in \text{Sets}$.

An infinite Galois category (\mathcal{C}, F) is *tame* if for any connected $X \in \mathcal{C}$, $\pi_1(\mathcal{C}, F)$ acts transitively on $F(X)$.

Example 2.4. If G is a topological group, then $(G - \text{Sets}, F_G)$ is a tame infinite Galois category.

Theorem 2.5 [Bhatt and Scholze 2015, Theorem 7.2.5]. Fix an infinite Galois category (\mathcal{C}, F) and a Noohi group G . Then:

- (1) $\pi_1(\mathcal{C}, F)$ is a Noohi group.
- (2) There is a natural identification of $\text{Hom}_{\text{cont}}(G, \pi_1(\mathcal{C}, F))$ with the groupoid of functors $\mathcal{C} \rightarrow G - \text{Sets}$ that commute with the fiber functors.
- (3) If (\mathcal{C}, F) is tame, then F induces an equivalence $\mathcal{C} \simeq \pi_1(\mathcal{C}, F) - \text{Sets}$.

The “tameness” assumption cannot be dropped as there exist infinite Galois categories that are not of the form $(G - \text{Sets}, F_G)$; see [Bhatt and Scholze 2015, Example 7.2.3]. This was overlooked in [Noohi 2008], where a similar formalism was considered.

Remark 2.6. The above formalism was also studied in [Lepage 2010, Chapter 4] under the names of “quasiprodiscrete” groups and “pointed classifying categories”.

In Section 2B below we will study “Noohi completion” and the dictionary between Noohi groups and $G - \text{Sets}$ (see Section 2C). For now, let us return to gathering the results from [Bhatt and Scholze 2015].

Pro-étale topology and the definition of $\pi_1^{\text{proét}}(X)$.

Definition 2.7. Let X be a locally topologically noetherian scheme. Let $Y \rightarrow X$ be a morphism of schemes such that:

- (1) It is étale (not necessarily quasicompact!).
- (2) It satisfies the valuative criterion of properness.

We will call Y a *geometric covering* of X . We will denote the category of geometric coverings by Cov_X .

As Y is not assumed to be of finite type over X , the valuative criterion does not imply that $Y \rightarrow X$ is proper (otherwise we would simply get a finite étale morphism).

Example 2.8. For an algebraically closed field \bar{k} , the category $\text{Cov}_{\text{Spec}(\bar{k})}$ consists of (possibly infinite) disjoint unions of $\text{Spec}(\bar{k})$ and we have $\text{Cov}_{\text{Spec}(\bar{k})} \simeq \text{Sets}$.

More generally, one has:

Lemma 2.9 [Bhatt and Scholze 2015, Lemma 7.3.8]. *If X is a henselian local scheme, then any $Y \in \text{Cov}_X$ is a disjoint union of finite étale X -schemes.*

Let us choose a geometric point $\bar{x} : \text{Spec}(\bar{k}) \rightarrow X$ on X . By Example 2.8, this gives a fiber functor $F_{\bar{x}} : \text{Cov}_X \rightarrow \text{Sets}$. By [Bhatt and Scholze 2015, Lemma 7.4.1], the pair $(\text{Cov}_X, F_{\bar{x}})$ is a tame infinite Galois category. Then one defines:

Definition 2.10. The *pro-étale fundamental group* is defined as

$$\pi_1^{\text{proét}}(X, \bar{x}) = \pi_1(\text{Cov}_X, F_{\bar{x}}).$$

In other words, $\pi_1^{\text{proét}}(X, \bar{x}) = \text{Aut}(F_{\bar{x}})$ and this group is topologized using the compact-open topology on $\text{Aut}(S)$ for any $S \in \text{Sets}$.

One can compare the groups $\pi_1^{\text{proét}}(X, \bar{x})$, $\pi_1^{\text{ét}}(X, \bar{x})$ and $\pi_1^{\text{SGA}^3}(X, \bar{x})$, where the last group is the group introduced in Chapter X.6 of [SGA 3 II 1970].

Lemma 2.11. *For a scheme X , the following relations between the fundamental groups hold:*

- (1) *The group $\pi_1^{\text{ét}}(X, \bar{x})$ is the profinite completion of $\pi_1^{\text{proét}}(X)$.*
- (2) *The group $\pi_1^{\text{SGA}^3}(X, \bar{x})$ is the prodiscrete completion of $\pi_1^{\text{proét}}(X, \bar{x})$.*

Proof. This follows from [Bhatt and Scholze 2015, Lemma 7.4.3 and 7.4.6]. □

As shown in [loc. cit., Example 7.4.9], $\pi_1^{\text{proét}}(X, \bar{x})$ is indeed more general than $\pi_1^{\text{SGA}^3}(X, \bar{x})$. This can be also seen by combining Example 4.5 with Proposition 4.8 below.

The following lemma is extremely important to keep in mind and will be used many times throughout the paper. Recall that, for example, a normal scheme is geometrically unibranch.

Lemma 2.12 [Bhatt and Scholze 2015, Lemma 7.4.10]. *If X is geometrically unibranch, then*

$$\pi_1^{\text{proét}}(X, \bar{x}) \simeq \pi_1^{\text{ét}}(X, \bar{x}).$$

There is another way of looking at the pro-étale fundamental group, which justifies the name “pro-étale”.

Definition 2.13. (1) A map $f : Y \rightarrow X$ of schemes is called *weakly étale* if f is flat and the diagonal $\Delta_f : Y \rightarrow Y \times_X Y$ is flat.

- (2) The pro-étale site $X_{\text{proét}}$ is the site of weakly étale X -schemes, with covers given by fpqc covers.

This definition of the pro-étale site is justified by a foundational theorem — part (c) of the following fact.

Fact 2.14. Let $f : A \rightarrow B$ be a map of rings:

- (a) f is étale if and only if f is weakly étale and finitely presented.
- (b) If f is ind-étale, i.e., B is a filtered colimit of étale A -algebras, then f is weakly étale.
- (c) [Bhatt and Scholze 2015, Theorem 2.3.4] If f is weakly étale, then there exists a faithfully flat ind-étale $g : B \rightarrow C$ such that $g \circ f$ is ind-étale.

Definition 2.15 [Bhatt and Scholze 2015, Definition 7.3.1]. We say that $F \in \text{Shv}(X_{\text{proét}})$ is *locally constant* if there exists a cover $\{Y_i \rightarrow X\}$ in $X_{\text{proét}}$ with $F|_{Y_i}$ constant. We write Loc_X for the corresponding full subcategory of $\text{Shv}(X_{\text{proét}})$.

We are ready to state the following important result.

Theorem 2.16 [Bhatt and Scholze 2015, Lemma 7.3.9]. *Let X to be locally topologically noetherian scheme. One has $\text{Loc}_X = \text{Cov}_X$ as subcategories of $\text{Shv}(X_{\text{proét}})$.*

Topological invariance of the pro-étale fundamental group. We note that universal homeomorphisms of schemes induce equivalences on the corresponding categories of geometric coverings.

Proposition 2.17 [Bhatt and Scholze 2015, Lemma 5.4.2]. *Let $h : X' \rightarrow X$ be a universal homeomorphism of topologically noetherian schemes (i.e., induces a homeomorphism of topological spaces after any base-change). Then the pullback*

$$h^* : \text{Cov}_X \rightarrow \text{Cov}_{X'}, \quad Y \mapsto Y' = Y \times_X X'$$

is an equivalence of categories.

Proof. As $\text{Cov}_X \simeq \text{Loc}_X$, the theorem follows by the same proof as in [loc. cit., Lemma 5.4.2].

Alternatively, one can argue more directly (i.e., avoiding the equivalence with Loc_X) as follows. By [Stacks 2020, Theorem 04DZ], $V \mapsto V' = V \times_X X'$ induces an equivalence of categories of schemes étale over X and schemes étale over X' . By [Rydh 2010, Proposition 5.4.], this induces an equivalence between schemes étale and separated over respectively X and X' . The only thing left to be shown is that if for an étale separated scheme $Y \rightarrow X$, the map $Y \times_X X' \rightarrow X'$ satisfies the existence part of the valuative criterion of properness, then so does $Y \rightarrow X$. But this property can be characterized in purely topological terms (see [Stacks 2020, Lemma 01KE]) and so the result follows from the fact that h is a universal homeomorphism. \square

2B. Noohi completion. Let HausdGps denote the category of Hausdorff topological groups (recall that we assume all topological groups to be Hausdorff, unless stated otherwise) and NoohiGps to be the full subcategory of Noohi groups. Let G be a topological group. Denote $\mathcal{C}_G = G - \text{Sets}$ and let $F_G : \mathcal{C}_G \rightarrow \text{Sets}$ be the forgetful functor. Observe that (\mathcal{C}_G, F_G) is a tame infinite Galois category. Thus, the group $\text{Aut}(F_G)$ is a Noohi group. It is easy to see that a morphism $G \rightarrow H$ defines an induced morphism of groups $\text{Aut}(F_G) \rightarrow \text{Aut}(F_H)$ and check that it is continuous. Let $\psi_N : \text{HausdGps} \rightarrow \text{NoohiGps}$ be the functor defined by $G \mapsto \text{Aut}(F_G)$. Denote also the inclusion $i_N : \text{NoohiGps} \rightarrow \text{HausdGps}$.

Definition 2.18. We call $\psi_N(G)$ the Noohi completion of G and will denote it G^{Noohi} .

Example 2.19. In [Bhatt and Scholze 2015, Example 7.2.6], it was explained that the category of Noohi groups admits coproducts. Let G_1, G_2 be two Noohi groups and let $G_1 *^N G_2$ denote their coproduct as Noohi groups. Let $G_1 *^{\text{top}} G_2$ be their topological coproduct. It exists and it is a Hausdorff group [Graev 1948]. Then $G_1 *^N G_2 = (G_1 *^{\text{top}} G_2)^{\text{Noohi}}$.

Let $\alpha_G : G \rightarrow \text{Aut}(F_G) = G^{\text{Noohi}}$ denote the obvious morphism.

Proposition 2.20. For a topological group G , the functor F_G induces an equivalence of categories

$$\tilde{F}_G : G\text{-Sets} \xrightarrow{\sim} G^{\text{Noohi}}\text{-Sets}.$$

Moreover, $\alpha_G^* \circ \tilde{F}_G \simeq \text{id}$, and thus α^* is an equivalence of categories, too.

Proof. The first part follows directly from [Bhatt and Scholze 2015, Theorem 7.2.5]. The natural isomorphism $\alpha_G^* \circ \tilde{F}_G \simeq \text{id}$ is clear from the definitions. It follows that α_G^* is an equivalence. \square

The following lemma is in contrast with [Noohi 2008, Remark 2.13], but agrees with [Lepage 2010, Proposition 4.1.1].

Lemma 2.21. For any topological group G , the image of $\alpha_G : G \rightarrow G^{\text{Noohi}}$ is dense.

Proof. Let $U \subset G^{\text{Noohi}}$ be open. As G^{Noohi} is Noohi, there exists $q \in G^{\text{Noohi}}$ and an open subgroup $V <^\circ G^{\text{Noohi}}$ such that $qV \subset U$. The quotient G^{Noohi}/V gives a G^{Noohi} -set. It is connected in the category $G^{\text{Noohi}}\text{-Sets}$ and, by Proposition 2.20, $\alpha_G^*(G^{\text{Noohi}}/V)$ is connected. Thus, the action of G on G^{Noohi}/V is transitive and so there exists $g \in G$ such that $\alpha_G(g) \cdot [V] = [qV]$, i.e., $\alpha_G(g) \in qV$. Thus, the image of α_G is dense. \square

Observation. Let $f : H \rightarrow G$ be a map of topological groups. Directly from the definitions, one sees that the following diagram commutes:

$$\begin{array}{ccc} H & \xrightarrow{f} & G \\ \alpha_H \downarrow & & \alpha_G \downarrow \\ H^{\text{Noohi}} & \xrightarrow{f^{\text{Noohi}}} & G^{\text{Noohi}} \end{array}$$

Lemma 2.22 (universal property of Noohi completion). Let $f : H \rightarrow G$ be a continuous morphism from a topological group to a Noohi group. Then there exists a unique map $f' : H^{\text{Noohi}} \rightarrow G$ such that $f' \circ \alpha_H = f$.

Proof. By the definition of a Noohi group, α_G is an isomorphism. Defining $f' := \alpha_G^{-1} \circ f^{\text{Noohi}}$ gives the existence. The uniqueness follows from α_H having dense image. Alternatively, one can combine Proposition 2.20 with [Bhatt and Scholze 2015, Theorem 7.2.5(2)]. \square

Corollary 2.23. The functor ψ_N is a left adjoint of i_N .

Remark 2.24. There are few places, where we write G^{Noohi} for a non-Hausdorff group G . This is mostly to avoid a large overline sign over a subgroup described by generators. In these cases, we mean

$$G^{\text{Noohi}} := (G^{\text{Hausd}})^{\text{Noohi}}$$

where G^{Hausd} is the maximal Hausdorff quotient. As $(-)^{\text{Hausd}}$ is a left adjoint as well, this usually does not cause problems. This also provides a left adjoint to the forgetful functor $\text{NoohiGps} \rightarrow \text{TopGps}$ to all topological groups.

We now move towards a more explicit description of the Noohi completion.

Lemma 2.25. *Let (G, τ) be a topological group. Denote by \mathcal{B} the collection of sets of the form*

$$x_1 \Gamma_1 y_1 \cap x_2 \Gamma_2 y_2 \cap \dots \cap x_m \Gamma_m y_m$$

where $m \in \mathbb{N}$, $x_i, y_i \in G$ and $\Gamma_i < G$ are open subgroups of G . Then \mathcal{B} is a basis of a group topology τ' on G that is weaker than τ and open subgroups of (G, τ) form a basis of open neighborhoods of 1_G in (G, τ') .

Moreover, the natural map $i' : (G, \tau) \rightarrow (G, \tau')$ induces an equivalence of categories $(G, \tau') - \text{Sets} \rightarrow (G, \tau) - \text{Sets}$. If $\{1_G\} \subset (G, \tau)$ is thickly closed, i.e., $\bigcap_{U <_G} U = \{1_G\}$ (see [Definition 2.30](#)), then (G, τ') is Hausdorff and $(G, \tau)^{\text{Noohi}} \xrightarrow{\simeq} (G, \tau')^{\text{Noohi}}$ is an isomorphism.

Proof. The first statement follows from [[Bourbaki 1966](#), Proposition III.1.1] by taking the filter of subsets of G containing an open subgroup. It is also proven in [[Lavanda 2018](#), Lemma 1.13] (the proposition is stated there in a particular case, but the proof works for any topological group). The second statement follows from the fact that for a discrete set S , any continuous morphism $(G, \tau) \rightarrow \text{Aut}(S)$ factorizes through $i' : (G, \tau) \rightarrow (G, \tau')$. □

Fact 2.26 [[Bhatt and Scholze 2015](#), Proposition 7.1.5]. Let G be a topological group such that its open subgroups form a basis of open neighborhoods of 1_G . Then $G^{\text{Noohi}} \simeq \widehat{G}$, where \widehat{G} denotes the Raïkov completion of G .

Proposition 2.27. *Let (G, τ) be a topological group. Assume that $\{1_G\} \subset (G, \tau)$ is thickly closed (see [Definition 2.30](#)). Then there is a natural isomorphism of groups*

$$G^{\text{Noohi}} \simeq \widehat{(G, \tau)},$$

where τ' denotes the topology described in the previous lemma and $\widehat{\cdot}$ denotes the Raïkov completion.

Proof. We combine [Fact 2.26](#) with the last lemma and get $(G, \tau)^{\text{Noohi}} \simeq (G, \tau')^{\text{Noohi}} \simeq \widehat{(G, \tau)}$. □

Observation 2.28. Let G be a topological group and H a normal subgroup. Then the full subcategory of $G - \text{Sets}$ of objects on which H acts trivially is equal to the full subcategory of $G - \text{Sets}$ on which its closure \bar{H} acts trivially and it is equivalent to the category of $G/\bar{H} - \text{Sets}$. So, it is an infinite Galois category with the fundamental group equal to $(G/\bar{H})^{\text{Noohi}}$.

Lemma 2.29. *Let X be a connected, locally path-connected, semilocally simply connected topological space and $x \in X$ a point. Let F_x be the functor taking a covering space $Y \rightarrow X$ to the fiber Y_x over the point $x \in X$. Then $(\text{TopCov}(X), F_x)$ is a tame infinite Galois category and $\pi_1(\text{TopCov}(X), F_x) = \pi_1^{\text{top}}(X, x)$, where we consider $\pi_1^{\text{top}}(X, x)$ with the discrete topology. Here, $\text{TopCov}(X)$ denotes the category of covering spaces of X .*

Proof. We first claim that there is an isomorphism: $(\text{TopCov}(X), F_x) \simeq (\pi_1^{\text{top}}(X, x) - \text{Sets}, F_{\pi_1^{\text{top}}(X, x)})$. This is in fact a classical result in algebraic topology, which can be recovered from [Fulton 1995, Chapter 13] or [Hatcher 2002, Chapter 1] and is stated explicitly in [Çakar 2014, Corollary 4.1]. This finishes the proof, as discrete groups are Noohi. □

2C. Dictionary between Noohi groups and $G - \text{Sets}$.

Definition 2.30. Let $H \subset G$ be a subgroup of a topological group G . Then we define a “thick closure” $\overline{\overline{H}}$ of H in G to be the intersection of all open subgroups of G containing H , i.e., $\overline{\overline{H}} := \bigcap_{H \subset U <^\circ G} U$. If a subgroup satisfies $H = \overline{\overline{H}}$ we will call it thickly closed in G .

In a topological group open subgroups are also closed, so a thickly closed subgroup is also an intersection of closed subgroups, so it is closed in G . Observe also that an arbitrary intersection of thickly closed subgroups is thickly closed. This justifies, for example, the existence of the smallest normal thickly closed subgroup containing a given group. In fact, we can formulate a more precise observation.

Observation 2.31. Let $H < G$ be a subgroup of a topological group G . Then the smallest normal thickly closed subgroup of G containing H is equal to $\overline{\overline{H^{nc}}}$, where H^{nc} is the normal closure of H in G .

Observation 2.32. Let G be a topological group such that the open subgroups form a local base at 1_G . Let $W \subset G$ be a subset. Then the topological closure of W can be written as $\overline{W} = \bigcap_{V <^\circ G} WV$.

The following lemma can be found on page 79 of [Lepage 2010].

Lemma 2.33. *Let G be a topological group such that the open subgroups form a basis of neighborhoods of 1_G . Let $H \triangleleft G$ be a normal subgroup. Then*

$$\overline{H} = \overline{\overline{H}}$$

i.e., the usual topological closure and the thick closure coincide.

Proof. We compute that $\overline{H} = \bigcap_{V <^\circ G} HV \stackrel{(*)}{\supset} \bigcap_{H < U <^\circ G} U = \overline{\overline{H}} \supset \overline{H}$. The inclusion $(*)$ follows from the fact that HV is an (open) subgroup of G as H is normal. □

Let us make an easy observation, that will be useful to keep in mind while reading the proof of the technical proposition below.

Observation 2.34. Let $U < G$ be an open subgroup of a topological group and let $g_0 \in G$. Then the mapping $G/g_0Ug_0^{-1} \rightarrow G/U$ given by $[gg_0Ug_0^{-1}] \mapsto [g_0U]$ is an isomorphism of G -sets.

Given open subgroups $U, V < G$ and some surjective map of G -sets $\phi : G/V \rightarrow G/U$ we can assume that it is the standard quotient map (i.e., $V \subset U$) up to replacing U by a conjugate open subgroup (more precisely by $g_0 U g_0^{-1}$, where g_0 is such that $\phi([V]) = [g_0 U]$).

Remark 2.35. A map $Y' \rightarrow Y$ in an infinite Galois category (\mathcal{C}, F) is an epimorphism/monomorphism if and only if the map $F(Y') \rightarrow F(Y)$ is surjective/injective. Similarly Y is an initial object if and only if $F(Y) = \emptyset$ and so on. The proofs of those facts are the same as the proofs in [Stacks 2020, Tag 0BN0]. This justifies using words “injective” or “surjective” when speaking about maps in (\mathcal{C}, F) .

Recall the following fact.

Observation. Let $f : G' \rightarrow G$ be a surjective map of topological groups. Then the induced morphism $G'/\ker(f) \rightarrow G$ is an isomorphism if and only if f is open. In such case, we say that f is a quotient map. In the language of [Bourbaki 1966, III.2.8] we would call f *strict* and surjective.

Definition 2.36. We will say that an object of a tame infinite Galois category is *completely decomposed* if it is a (possibly infinite) disjoint union of final objects.

Proposition 2.37. Let $G'' \xrightarrow{h'} G' \xrightarrow{h} G$ be maps between Noohi groups and $\mathcal{C}_{G''} \xleftarrow{H'} \mathcal{C}_{G'} \xleftarrow{H} \mathcal{C}_G$ the corresponding maps of the infinite Galois categories. Then the following hold:

- (1) The map $h' : G'' \rightarrow G'$ is a topological embedding if and only if for every connected object X in $\mathcal{C}_{G''}$, there exist connected objects $X' \in \mathcal{C}_{G''}$ and $Y \in \mathcal{C}_{G'}$ and maps $X' \twoheadrightarrow X$ and $X' \hookrightarrow H'(Y)$.
- (2) The following are equivalent:
 - (a) The morphism $h : G' \rightarrow G$ has dense image.
 - (b) The functor H maps connected objects to connected objects.
 - (c) The functor H is fully faithful.
- (3) The thick closure of $\text{Im}(h') \subset G'$ is normal if and only if for every connected object Y of $\mathcal{C}_{G'}$ such that $H'(Y)$ contains a final object of $\mathcal{C}_{G''}$, $H'(Y)$ is completely decomposed.
- (4) $h'(G'') \subset \text{Ker}(h)$ if and only if the composition $H' \circ H$ maps any object to a completely decomposed object.
- (5) Assume that $h'(G'') \subset \text{ker}(h)$ and that $h : G' \rightarrow G$ has dense image. Then the following conditions are equivalent:
 - (a) The induced map $(G'/\text{ker}(h))^{\text{Noohi}} \rightarrow G$ is an isomorphism and the smallest normal thickly closed subgroup containing $\text{Im}(h')$ is equal to $\text{ker}(h)$.
 - (b) For any connected $Y \in \mathcal{C}_{G'}$ such that $H'(Y)$ is completely decomposed, Y is in the essential image of H .
 - (c) The induced map $(G'/\text{ker}(h))^{\text{Noohi}} \rightarrow G$ is an isomorphism and for any connected $Y \in \mathcal{C}_{G'}$ such that $H'(Y)$ is completely decomposed, there exists $Z \in \mathcal{C}_G$ and an epimorphism $H(Z) \twoheadrightarrow Y$.

Proof. (1) The proof is virtually the same as for usual Galois categories, but there every injective map is automatically a topological embedding (as profinite groups are compact). Assume that $G'' \rightarrow G'$

is a topological embedding. Let $X \in \mathcal{C}_{G''}$ be connected and write $X \simeq G''/U$ for an open subgroup $U < G''$. Then there exists an open subset $\tilde{V} \subset G'$ such that $\tilde{V} \cap G'' = U$ (as $G'' \rightarrow G'$ is a topological embedding) and an open subgroup $V < G'$ such that $V \subset \tilde{V}$ (as G' is Noohi). Denote $W = V \cap G''$. Then $X' := G''/W \rightarrow X$ and $X' \hookrightarrow H'(G'/V)$, so we conclude by setting $Y := G'/V$. For the other implication: we want to prove that $G'' \rightarrow G'$ is a topological embedding under the assumption from the statement. It is enough to check that the set of preimages $h'^{-1}(\mathcal{B})$ of some basis \mathcal{B} of opens of $e_{G'}$ forms a basis of opens of $e_{G''}$. Indeed, assume that this is the case. Firstly, observe that it implies that h' is injective, as both G'' and G' are Hausdorff (and in particular T_0). If U is an open subset of G'' , then we can write $U = \bigcup g''_\alpha U_\alpha$ for some $g''_\alpha \in G''$ and $U_\alpha \in h'^{-1}(\mathcal{B})$. We can write $U_\alpha = h'^{-1}(V_\alpha)$ for some $V_\alpha \in \mathcal{B}$. Then $V = \bigcup h'(g''_\alpha)V_\alpha$ satisfies $h'^{-1}(V) = U$ because $h'^{-1}(h'(g''_\alpha)V_\alpha) = g''_\alpha U_\alpha$ (by injectivity of h'). So this will prove that the topology on G'' is induced from G' via h' . Let $\mathcal{B} = \{U < G' \mid U \text{ is open}\}$. This is a basis of opens of $e_{G'}$ (as G' is Noohi). We want to check that $h'^{-1}(\mathcal{B})$ is a basis of opens of $e_{G''}$. As open subgroups of G'' form a basis of opens of $e_{G''}$ it is enough to show that for any open subgroup $U < G''$ there exists an open subgroup $V < G'$ such that $h'^{-1}(V) \subset U$. From the assumption we know that there exist open subgroups $\tilde{U} < G''$ and $V < G'$ such that $G''/\tilde{U} \rightarrow G''/U$ and $G''/\tilde{U} \hookrightarrow G'/V$. The surjectivity of the first map means that we can assume (up to replacing \tilde{U} by a conjugate) $\tilde{U} \subset U$. The injectivity of the second means that we can assume (up to replacing V by a conjugate) that $h'^{-1}(V) \subset \tilde{U}$. Indeed, the injectivity implies that if $h'(g'')V = V$, then $g''\tilde{U} = \tilde{U}$ which translates immediately to $h'^{-1}(V) \subset \tilde{U}$. So we have also $h'^{-1}(V) \subset U$, which is what we wanted to prove.

(2) The equivalence between (a) and (b) follows from the observation that a map between Noohi groups $G' \rightarrow G$ has a dense image if and only if for any open subgroup U of G , the induced map on sets $G' \rightarrow G/U$ is surjective. Here, we only use that open subgroups form a basis of open neighborhoods of $1_G \in G$.

Now, the functor H is automatically faithful and conservative (because $F_{G'} \circ H = F_G$ is faithful and conservative). Assume that (b) holds. Let $S, T \in G - \text{Sets}$ and let $g \in \text{Hom}_{G' - \text{Sets}}(H(S), H(T))$. We have to show that g comes from $g_0 \in \text{Hom}_{G - \text{Sets}}(S, T)$. We can and do assume S, T connected for that. Let $\Gamma_g \subset H(S) \times H(T)$ be the graph of g . It is a connected subobject. As $H(S) \times H(T) = H(S \times T)$, the assumption (b) implies that each connected component of $H(S) \times H(T)$ is the pullback of a connected component Γ_0 of $S \times T$. Thus, Γ_g is the pullback of some $\Gamma_0 \subset S \times T$. By conservativity of H , the projection $p_{\Gamma_0} : \Gamma_0 \rightarrow S$ is an isomorphism, as this is true for $p_{\Gamma_g} : \Gamma_g \rightarrow H(S)$. The composition $q_{\Gamma_0} \circ p_{\Gamma_0}^{-1} : S \rightarrow T$ maps via H to g .

Conversely, assume (c) holds. Let $S \in G - \text{Sets}$ be connected. We want to show that $H(S)$ is connected. Suppose on the contrary that $H(S) = A \sqcup B$ with $A, B \in G' - \text{Sets}$. Let $T = \bullet \sqcup \bullet \in G - \text{Sets}$ be a two-element set with a trivial G -action. Then $\text{Hom}_{G - \text{Sets}}(S, T)$ has precisely two elements, while $\text{Hom}_{G' - \text{Sets}}(H(S), H(T)) = \text{Hom}_{G' - \text{Sets}}(A \sqcup B, \bullet \sqcup \bullet)$ has at least four.

(3) Assume first that the thick closure of $\text{im}(h')$ is normal. Let $Y = G'/U$ be an element of $\mathcal{C}_{G'}$ whose pull-back to $G'' - \text{Sets}$ contains the final object. This means that G'' fixes one of the classes, let's say $[g'U]$.

This is equivalent to $g'^{-1}h'(G'')g'$ fixing $[U]$, i.e., $g'^{-1}h'(G'')g' \subset U$. But this implies immediately that $(g'^{-1}h'(G'')g') \subset U$. Let $\tilde{g} \in G'$ be any element. We have $(g'^{-1}h'(G'')g') = g'^{-1}\overline{h'(G'')}g' = \overline{h'(G'')} = \tilde{g}^{-1}\overline{h'(G'')}\tilde{g}$ from the assumption that $\overline{h'(G'')}$ is normal. So $\tilde{g}^{-1}h'(G'')\tilde{g} \subset \tilde{g}^{-1}\overline{h'(G'')}\tilde{g} \subset U$ and we conclude that $h'(G'')$ fixes an arbitrary class $[\tilde{g}U]$. This shows that G'/U pulls back to a completely decomposed object.

The other way round: assume that for every connected object Y of $\mathcal{C}_{G'}$ such that $H'(Y)$ contains a final object, $H'(Y)$ is completely decomposed. Let U be an open subgroup of G' containing $h'(G'')$. Then G'' fixes $[U] \in G'/U$ and so, by assumption, fixes every $[g'U] \in G'/U$. This implies that for any $g' \in G'$ $g'^{-1}h'(G'')g' \subset U$ which easily implies that also $h'(G'')^{nc} \subset U$. As this is true for any U containing $h'(G'')$ we get that $\overline{h'(G'')} = \overline{(h'(G'')^{nc})}$ and the last group is the smallest normal thickly closed subgroup of G' containing $h'(G'')$ (**Observation 2.31**).

(4) The same as for usual Galois categories, we use that $\bigcap_{U < \circ G} U = 1_G$.

(5) (b) \Rightarrow (c): Assume (b). We only need to show, that $(G'/\ker(h))^{Noohi} \rightarrow G$ is an isomorphism. This is equivalent to showing that H induces an equivalence $G'/\ker(h) - \text{Sets} \simeq G - \text{Sets}$. As $G'/\ker(h) - \text{Sets} \simeq \{S \in G' - \text{Sets} \mid \ker(h) \text{ acts trivially on } S\} \subset \{S \in G - \text{Sets} \mid G'' \text{ acts trivially on } S\}$, the assumption of (b) implies that the functor $G - \text{Sets} \rightarrow G'/\ker(h) - \text{Sets}$ is essentially surjective. By the global assumption that $G' \rightarrow G$ has dense image, it is fully faithful (see (2)).

(c) \Rightarrow (b): Assume (c). Let $Y \in \mathcal{C}_{G'}$ be connected and such that $H'(Y)$ is completely decomposed. We have $Z \in \mathcal{C}_G$ and an epimorphism $H(Z) \twoheadrightarrow Y$. As $\ker(h)$ acts trivially on $H(Z)$, we conclude that it also acts trivially on Y . Thus, by abuse of notation, $Y \in G'/\ker(h) - \text{Sets}$. But $G'/\ker(h) - \text{Sets} \simeq (G'/\ker(h))^{Noohi} - \text{Sets} \simeq G - \text{Sets}$ from the assumption. Thus, we see that Y is in the essential image of H .

(b) \Rightarrow (a): Assume (b). We give two proofs of this fact.

First proof. We have proven above that (b) $\Rightarrow (G'/\ker(h))^{Noohi} \simeq G$. Let N be the smallest normal thickly closed subgroup of G' containing $h'(G'')$. Observe that $N \subset \ker h$ (as $\ker(h)$ is thickly closed). Let U be an open subgroup containing N . We want to show that U contains $\ker h$. This will finish the proof as both N and $\ker h$ are thickly closed. Write $Y = G'/U$. Observe that G'/U pulls back to a completely decomposed G'' -set if and only if for any $g' \in G'$ there is $g'h'(G'')g'^{-1} \subset U$. Indeed, $h'(G'')$ fixes $[g'U] \in G'/U$ if and only if $g'h'(G'')g'^{-1}$ fixes $[U]$. So $N \subset U$ implies that Y pulls back to a completely decomposed G'' -set and, by assumption, Y is isomorphic to a pull-back of some G -set and so $\ker(h)$ acts trivially on Y . This implies that $\ker h \subset U$, which finishes the proof. \square

Alternative proof. We already know that (b) $\Rightarrow (G'/\ker(h))^{Noohi} \simeq G$. Let $N \subset \ker(h)$ be as in the first proof above. Consider the map $G/N \twoheadrightarrow G/\ker(h)$. The assumption (b) and full faithfulness of H (by the global assumption and using (2)) imply that $(G'/N)^{Noohi} \rightarrow G$ is an isomorphism. Thus, $(G'/N)^{Noohi} \simeq (G'/\ker(h))^{Noohi}$. Using **Proposition 2.27**, we check that the canonical maps $G'/N \rightarrow$

$(G'/N)^{\text{Noohi}}$ and $G'/\ker(h) \rightarrow (G'/\ker(h))^{\text{Noohi}}$ are injective. Thus, $G'/N \rightarrow G'/\ker(h)$ is injective and so $N = \ker(h)$. \square

(a) \Rightarrow (b): Assume (a). Let $Y = G'/U$ be a connected G' -set that pulls back via h' to a completely decomposed object. As we have seen while proving “(b) \Rightarrow (a)”, this implies that for any $g' \in G'$ $g'h'(G'')g'^{-1} \subset U$, so $H^{nc} \subset U$ and so also $\overline{(H^{nc})} \subset U$. But, by [Observation 2.31](#), there is $N = \overline{(H^{nc})}$. By assumption, we have $N = \ker h$ and so we conclude that $\ker h \subset U$. But then, by assumption $(G'/\ker(h))^{\text{Noohi}} \simeq G$, Y is in the essential image of H . \square

To distinguish between exactness in the usual sense (i.e., on the level of abstract groups) and notions of exactness appearing in [Proposition 2.37](#), we introduce a new notion. It will be mainly used in the context of Noohi groups.

Definition 2.38. Let $G'' \xrightarrow{h'} G' \xrightarrow{h} G \rightarrow 1$ be a sequence of topological groups such that $\text{im}(h') \subset \ker(h)$. Then we will say that the sequence is:

- (1) *Nearly exact on the right* if h has dense image,
- (2) *Nearly exact in the middle* if $\overline{\text{im}(h')} = \ker(h)$, i.e., the thick closure of the image of h' in G' is equal to the kernel of h .
- (3) *Nearly exact* if it is both nearly exact on the right and nearly exact in the middle.

We end this subsection with a lemma on topological groups and their Noohi completions that will be used later in the proof of the main theorem.

Lemma 2.39. Let G be a topological group and \tilde{G} be a subgroup of G^{Noohi} such that the canonical map $G \rightarrow G^{\text{Noohi}}$ factorizes through \tilde{G} :

$$G \rightarrow \tilde{G} \subset G^{\text{Noohi}}.$$

Let $V_0 < \tilde{G}$ be a subgroup. Let $S = (\tilde{G}/V_0, \text{discr})$ be the discrete set that comes naturally with an **abstract** action by \tilde{G} .

If the induced abstract G -action on S is continuous, then V_0 is open in \tilde{G} . Moreover, in such case, denoting $V = \text{Stab}_{G^{\text{Noohi}}}([V_0] \in \tilde{G}/V_0)$, there is

$$V = V_0^{\text{Noohi}} = \overline{V_0}^{G^{\text{Noohi}}} \quad \text{and} \quad V_0 = V \cap \tilde{G}.$$

Proof. By the universal property, the G -action on S extends to G^{Noohi} and this action is transitive. Then V_0 is the preimage of the stabilizer $V = \text{Stab}_{G^{\text{Noohi}}}([V_0] \in \tilde{G}/V_0)$, which is open.

The group V is open in a Noohi group, thus Noohi; see [\[Bhatt and Scholze 2015, Lemma 7.1.8.\]](#) By the universal property, there is a factorization $V_0^{\text{Noohi}} \rightarrow V$. But as V_0 is a subgroup of a Noohi group, its open subgroups form a basis of 1_{V_0} . Thus, the Noohi completion of V_0 is just the Raïkov completion. But as the canonical map from a group to its Raïkov completion is a topological embedding, [\[Arhangel'skii and Tkachenko 2008, Corollary 3.6.18\]](#) implies that $V_0^{\text{Noohi}} \rightarrow V$ is a topological embedding. By a characterization of Raïkov completeness (see [\[Dikranjan 2013, Proposition 6.2.7\]](#)), it follows that V_0^{Noohi}

is closed in V . But as \tilde{G} contains the image of G , it is dense in G^{Noohi} , and from the definition of V it follows that V_0 has to be dense in V . Putting this together, we get that $V_0^{\text{Noohi}} = V = \overline{V_0}^{G^{\text{Noohi}}}$. \square

2D. A remark on valuative criteria. We will sometimes shorten “the valuative criterion of properness” to “VCoP”. It is useful to keep in mind the precise statements of different parts of the valuative criterion, see [Stacks 2020, Lemmas 01KE, 01KC and Section 01KY]. Let us prove a lemma (which is implicit in [Bhatt and Scholze 2015]), that VCoP can be checked fpqc-locally.

Lemma 2.40. *Let $g : X \rightarrow S$ be a map of schemes. The properties*

- (a) *g is étale,*
- (b) *g is separated,*
- (c) *g satisfies the existence part of VCoP,*

can be checked fpqc-locally on S . Moreover, property (c) can be also checked after a surjective proper base-change.

Proof. The cases of étale and separated morphisms are proven in [Stacks 2020, Section 02YJ]. For the last part, satisfying the existence part of VCoP is equivalent to specializations lifting along any base-change of g [Stacks 2020, Lemma 01KE]. It is easy to see that this property can be checked Zariski locally. Thus, if $S' \rightarrow S$ is an fpqc cover such that the base-change $g' : X' \rightarrow S'$ satisfies specialization lifting for any base-change, we can assume that S, S' are affine with $S' \rightarrow S$ faithfully flat. Let $T \rightarrow S$ be any morphism. Consider the diagram:

$$\begin{array}{ccccc}
 X' & \longleftarrow & S' \times_S X \times_S T & \longrightarrow & T \times_S X \\
 \downarrow & & \downarrow & & \downarrow \\
 S' & \longleftarrow & S' \times_S T & \longrightarrow & T
 \end{array}$$

Let $\xi' \in T \times_S X$, let ξ be its image in T and let $t \in T$ be such that $\xi \rightsquigarrow t$. We need to find $t' \in T \times_S X$ over t such that $\xi' \rightsquigarrow t'$. Let $Z = \overline{\{\xi'\}} \subset T \times_S X$ be the closure of $\{\xi'\}$. We need to show that the set-theoretic image $W \subset T$ of Z in T contains t . It is enough to show, that W is stable under specialization or, equivalently, that $T \setminus W$ is stable under generalization. But, from flatness [Stacks 2020, Lemma 03HV], generalizations lift along $S' \times_S T \rightarrow T$. Thus, it is enough to show that the preimage of $T \setminus W$ in $S' \times_S T$ is stable under generalizations or, equivalently (using the surjectivity of $S' \times_S T \rightarrow T$), that the preimage of W in $S' \times_S T$ is closed under specializations. But an easy diagram chasing (using the fact that the right square of the diagram above is cartesian) shows that the preimage of W in $S' \times_S T$ is the image of a closed subset of $S' \times_S X \times_S T$. We conclude, because specializations lift along $S' \times_S X \times_S T \rightarrow S' \times_S T$ by assumption.

The last part of the statement is proven in an analogous way. \square

Lemma 2.41. *Let $f : Y \rightarrow X$ be a geometric covering of a locally topologically noetherian scheme. Then f is separated.*

Proof. By [Bhatt and Scholze 2015, Remark 7.3.3], f is quasiseparated. A quasiseparated morphism satisfying VCoP is separated; see [Stacks 2020, Tag 01KY]. \square

3. Seifert–van Kampen theorem for $\pi_1^{\text{proét}}$ and its applications

3A. Abstract Seifert–van Kampen theorem for infinite Galois categories. We aim at recovering a general version of van Kampen theorem, proven in [Stix 2006], in the case of the pro-étale fundamental group. Most of the definitions and proofs are virtually the same as in [loc. cit.], after replacing “Galois category” with “(tame) infinite Galois category” and “profinite” with “Noohi”, but still some additional technical difficulties appear here and there. We make the necessary changes in the definitions and deal with those difficulties below.

Denote by $\Delta_{\leq 2}$ a category whose objects are $[0] = \{0\}$, $[1] = \{0, 1\}$, $[2] = \{0, 1, 2\}$ and has strictly increasing maps as morphisms. There are face maps $\partial_i : [n - 1] \rightarrow [n]$ for $n = 1, 2$ and $0 \leq i \leq n$ which omit the value i and vertices $v_i : [0] \rightarrow [2]$ with image i .

The category of 2-complexes in a category \mathcal{C} is the category of contravariant functors $T_\bullet : \Delta_{\leq 2} \rightarrow \mathcal{C}$. We denote $T_n = T_\bullet([n])$ and call it the n -simplices of T_\bullet . $T(\partial_i)$ is called the i -th boundary map.

By a 2-complex E we mean a 2-complex in the category of sets. We often think of E as a category: its objects are the elements of E_n for $n = 0, 1, 2$ and its morphisms are obtained by defining $\partial : s \rightarrow t$ where $s \in E_n$ and $t = E(\partial)(s)$. Let $\Delta_n = \{ \sum_{i=0}^n \lambda_i e_i \in \mathbb{R}_{\geq 0}^{n+1} \mid \sum_i \lambda_i = 1 \}$ denote the topological n -simplex. Then we define $|E| = \bigsqcup E_n \times \Delta_n / \sim$, where \sim identifies $(s, d(x))$ with $(E(\partial)(s), x)$ for all $\partial : [m] \rightarrow [n]$ and its corresponding linear map $d : \Delta_m \rightarrow \Delta_n$ sending e_i to $e_{\partial(i)}$, and $s \in E_n$ and $x \in \Delta_m$. We call E connected if $|E|$ is a connected topological space.

Definition 3.1. *Noohi group data (\mathcal{G}, α) on a 2-complex E consists of the following:*

- (1) A mapping (not necessarily a functor!) \mathcal{G} from the category E to the category of Noohi groups: to a complex $s \in E_n$ is attributed a Noohi group $\mathcal{G}(s)$ and to a map $\partial : s \rightarrow t$ is attached a continuous morphism $\mathcal{G}(\partial) : \mathcal{G}(s) \rightarrow \mathcal{G}(t)$.
- (2) For every triple $v \in E_0$, $e \in E_1$, $f \in E_2$ and boundary maps ∂', ∂ such that $\partial'(f) = e$, $\partial(e) = v$, an element $\alpha_{vef} \in \mathcal{G}(v)$ (its existence is a part of the definition) such that the following diagram commutes:

$$\begin{array}{ccc}
 \mathcal{G}(f) & \xrightarrow{\mathcal{G}(\partial')} & \mathcal{G}(e) \\
 \mathcal{G}(\partial\partial') \downarrow & & \downarrow \mathcal{G}(\partial) \\
 \mathcal{G}(v) & \xrightarrow{\alpha_{vef}(\cdot)\alpha_{vef}^{-1}} & \mathcal{G}(v)
 \end{array}$$

Definition 3.2. Let (\mathcal{G}, α) be Noohi group data on the 2-complex E . A (\mathcal{G}, α) -system M on E consists of the following:

- (1) For every simplex $s \in E$ a $\mathcal{G}(s)$ -set M_s .
- (2) For every boundary map $\partial : s \rightarrow t$ a map of $\mathcal{G}(s)$ -sets $m_\partial : M_s \rightarrow \mathcal{G}(\partial)^*(M_t)$, such that.
- (3) For every triple $v \in E_0, e \in E_1, f \in E_2$ and boundary maps ∂', ∂ such that $\partial'(f) = e, \partial(e) = v$ the following diagram commutes:

$$\begin{array}{ccc}
 M_f & \xrightarrow{m_{\partial'}} & M_e \\
 m_{\partial\partial'} \downarrow & & \downarrow m_\partial \\
 M_v & \xrightarrow{\alpha_{vef}} & M_v
 \end{array}$$

Definition 3.3. A (\mathcal{G}, α) -system is called *locally constant* if all the maps m_∂ are bijections.

Observe that $\alpha \cdot : m \mapsto \alpha m$ is a $\mathcal{G}(v)$ -equivariant map $M_v \rightarrow (\alpha(\alpha^{-1})^*)M_v$. Observe that there is an obvious notion of a morphism of (\mathcal{G}, α) -systems: a collection of $\mathcal{G}(s)$ -equivariant maps that commute with the m . Let us denote by $\text{lcs}(E, (\mathcal{G}, \alpha))$ the category of locally constant (\mathcal{G}, α) -systems.

Let $M \in \text{lcs}(\mathcal{G}, \alpha)$ for Noohi group data (\mathcal{G}, α) on some 2-complex E . We define oriented graphs $E_{\leq 1}$ and $M_{\leq 1}$ (which will be an oriented graph over $E_{\leq 1}$) as in [Stix 2006], but our graphs $M_{\leq 1}$ are possibly infinite. For $E_{\leq 1}$ the vertices are E_0 and edges E_1 such that ∂_0 (resp. ∂_1) map an edge to its target (resp. origin). For $M_{\leq 1}$ the vertices are $\bigsqcup_{v \in E_0} M_v$ and edges are $\bigsqcup_{e \in E_1} M_e$ serves as the set of edges. The target/origin maps are induced by the m_∂ and the map $M_{\leq 1} \rightarrow E_{\leq 1}$ is the obvious one.

There is an obvious topological realization functor for graphs $|\cdot|$. By applying this functor to the above construction we get a *topological covering* (because M is locally constant) $|M_{\leq 1}| \rightarrow |E_{\leq 1}|$. This gives a functor

$$|\cdot|_{\leq 1} : \text{lcs}(E, (\mathcal{G}, \alpha)) \rightarrow \text{TopCov}(|E_{\leq 1}|).$$

Choosing a maximal subtree T of $|E_{\leq 1}|$ gives a fiber functor $F_T : \text{TopCov}(|E_{\leq 1}|) \rightarrow \text{Sets}$ by $(p : Y \rightarrow |E_{\leq 1}|) \mapsto \pi_0(p^{-1}(|T|))$. The pair $(\text{TopCov}(|E_{\leq 1}|), F_T)$ is an infinite Galois category and the resulting fundamental group $\pi_1(\text{Cov}(|E_{\leq 1}|), F_T)$ is isomorphic to $\pi_1^{\text{top}}(|E_{\leq 1}|)$ (see Lemma 2.29) which is in turn isomorphic to $\text{Fr}(E_1) / \langle\langle \{\vec{e} \mid e \in T\}^{\text{Fr}(E_1)} \rangle\rangle = \text{Fr}(\vec{e} \mid e \in E_1 \setminus T)$, where $\text{Fr}(\cdot)$ denotes a free group on the given set of generators and $\langle\langle \{\vec{e} \mid e \in T\}^{\text{Fr}(E_1)} \rangle\rangle$ denotes the normal closure in $\text{Fr}(E_1)$ of the subgroup generated by $\{\vec{e} \in T\}$. Here, \vec{e} acts on $F_T(M)$ via

$$\pi_0(p^{-1}(|T|)) \cong \pi_0(p^{-1}(\partial_0(e))) \cong \pi_0(p^{-1}(|e|)) \cong \pi_0(p^{-1}(\partial_1(e))) \cong \pi_0(p^{-1}(|T|)).$$

As in [Stix 2006], for every $s \in E_0$ and $M \in \text{lcs}(E, (\mathcal{G}, \alpha))$ we have that $F_T(M)$ can be seen canonically as a $\mathcal{G}(s)$ -module by $M_s = \pi_0(p^{-1}(s)) \cong \pi_0(p^{-1}(T))$. Denote $\pi_1(E_{\leq 1}, T) = \text{Fr}(E_1) / \langle\langle \{\vec{e} \mid e \in T\}^{\text{Fr}(E_1)} \rangle\rangle$. Putting the above together we get a functor

$$Q : \text{lcs}(E, (\mathcal{G}, \alpha)) \rightarrow (*_{v \in E_0}^N \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T)) - \text{sets}.$$

Remark 3.4. In the setting of usual (“finite”) Galois categories, it is usually enough to say that a particular morphism between two Galois categories is exact, because of the following fact [Stacks 2020, Tag 0BMV]:

Let G be a topological group. Let $F : \text{Finite-}G\text{-Sets} \rightarrow \text{Sets}$ be an exact functor with $F(X)$ finite for all X . Then F is isomorphic to the forgetful functor.

As we do not know if an analogous fact is true for infinite Galois categories, given two infinite Galois categories (\mathcal{C}, F) , (\mathcal{C}', F') and a morphism $\phi : \mathcal{C} \rightarrow \mathcal{C}'$, we are usually more interested in checking whether $F \simeq F' \circ \phi$. If ϕ satisfies this condition, it also commutes with finite limits and arbitrary colimits. Indeed, we have a map $\text{colim } \phi(X_i) \rightarrow \phi(\text{colim } X_i)$ that becomes an isomorphism after applying F' (as F' and $F = F' \circ \phi$ commute with colimits) and we conclude by conservativity of F' . Similarly for finite limits.

Proposition 3.5. *Let $(E, (\mathcal{G}, \alpha))$ be a connected 2-complex with Noohi group data. Define a functor $F : \text{lcs}(E, (\mathcal{G}, \alpha)) \rightarrow \text{Sets}$ in the following way: pick any simplex s and define F by $M \mapsto M_s$. Then $(\text{lcs}(E, (\mathcal{G}, \alpha)), F)$ is a tame infinite Galois category.*

Moreover, the obtained functor

$$Q : \text{lcs}(E, (\mathcal{G}, \alpha)) \rightarrow (*_{v \in E_0}^N \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T)) - \text{sets}$$

satisfies $F \simeq F_{\text{forget}} \circ Q$ and maps connected objects to connected objects.

Proof. We first check conditions (1), (2) and (4) of [Bhatt and Scholze 2015, Definition 7.2.1]. Then we show that Q maps connected objects to connected objects and we use the proof of this last fact to show the condition (3).

Colimits and finite limits: they exist simplex-wise and taking limits and colimits is functorial so we get a system as candidate for a colimit/finite limit. This will be a locally constant system, as the colimit/finite limit of bijections between some G -sets is a bijection.

Each M is a disjoint union of connected objects: let us call $N \in \text{lcs}(\mathcal{G}, \alpha)$ a *subsystem* of M if there exists a morphism $N \rightarrow M$ such that for any simplex s the map $N_s \rightarrow M_s$ is injective (we then identify, for any simplex s , N_s with a subset of M_s). We can intersect such subsystems in an obvious way and observe that it gives another subsystem. So for any element $a \in M_v$ there exists the smallest subsystem N of M such that $a \in N_v$. We see readily that for any vertices v, v' and $a \in M_v, a' \in M_{v'}$ the smallest subsystems N and N' containing one of them are either equal or disjoint (in the sense that, for each simplex s , N_s and N'_s are disjoint as subsets of M_s). It is easy to see that in this way we have obtained a decomposition of M into a disjoint union of connected objects.

F is faithful, conservative and commutes with colimits and finite limits. Observe that $\phi_s : \text{lcs}(E, (\mathcal{G}, \alpha)) \ni M \mapsto M_s \in \mathcal{G}(s)\text{-Sets}$ is faithful, conservative and commutes with colimits and finite limits and $F = F_s \circ \phi_s$, where F_s is the usual forgetful functor on $\mathcal{G}(s)\text{-Sets}$.

It is obvious that $F \simeq F_{\text{forget}} \circ Q$. We are now going to show that Q preserves connected objects. Take a connected object $M \in \text{lcs}(E, (\mathcal{G}, \alpha))$ and suppose that N is a nonempty subset of $F_T(M)$ stable under the action of $\pi_1(E_{\leq 1}, T)$ and $\mathcal{G}(v)$ for $v \in E_0$. Stability under the action of $\pi_1(E_{\leq 1}, T)$ shows that N can be extended to a subgraph $N_{\leq 1} \subset M_{\leq 1}$: for an edge e of $M_{\leq 1}$ we declare it to be an edge of $N_{\leq 1}$ if one of its ends touches a connected component of $p^{-1}(|T|)$ corresponding to an element of N . This is

well defined, as in this case both ends touch such a component — this is because the action of $m_{\partial_1} m_{\partial_0}^{-1}$ equals the action of $\vec{e} \in \pi_1(E_{\leq 1}, T)$.

Now we want to show that it extends to 2-simplices. This is a local question and we can restrict to simplices in the boundary of a given face $f \in E_2$. Define N_f as a preimage of N_s via any ∂ such that $\partial(f) = s$. We see that if the choice is independent of s , then we have extended N to a locally constant system. To see the independence it is enough to prove that if (vef) is a barycentric subdivision (i.e., we have ∂ and ∂' such that $\partial'(f) = e$ and $\partial(e) = v$), then $m_{\partial\partial'}^{-1}(N_v) = m_{\partial'}^{-1}(N_e)$. But from the $\mathcal{G}(v)$ -invariance we have $N_v = \alpha_{vef}^{-1}(N_v)$ and so

$$m_{\partial\partial'}^{-1}(N_v) = m_{\partial\partial'}^{-1}(\alpha_{vef}^{-1}(N_v)) = m_{\partial'}^{-1} m_{\partial}^{-1}(N_v) = m_{\partial'}(N_e)$$

and thus N can be seen as an element of $\text{lcs}(E, (\mathcal{G}, \alpha))$ which is a subobject of M , which contradicts connectedness of M .

To see that $\text{lcs}(E, (\mathcal{G}, \alpha))$ is generated under colimits by a set of connected objects, observe that in the above proof of the fact that Q preserves connected objects, we have in fact shown the following statement.

Fact 3.6. Let $M \in \text{lcs}(\mathcal{G}, \alpha)$ and let Z be a connected component of $Q(M)$. Then there exists a subsystem $W \subset M$ such that $Q(W) = Z$.

We want to show that there exists a set of connected objects in $\text{lcs}(\mathcal{G}, \alpha)$ such that any connected object of $\text{lcs}(\mathcal{G}, \alpha)$ is isomorphic to an element in that set. As an analogous fact is true in $(*_v \in E_0 \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T))$ – sets, it is easy to see that it is enough to check that, for any X, Y , if $QX \simeq QY$, then $X \simeq Y$. Let $X, Y \in \text{lcs}(\mathcal{G}, \alpha)$ be connected. Assume that $QX \simeq QY$. Looking at the graph of this isomorphism, we find a connected subobject $Z \subset QX \times QY$ that maps isomorphically on QX and QY via the respective projections. By the above fact, we know that there exists $W \subset X \times Y$ such that $QW = Z$. Because $F \simeq F_{\text{forget}} \circ Q$ and F is conservative, we see that the projections $W \rightarrow X$ and $W \rightarrow Y$ must be isomorphisms. This shows $X \simeq Y$ as desired.

The only claim left is that $\text{lcs}(E(\mathcal{G}, \alpha))$ is tame, but this follows from tameness of $(*_v \in E_0 \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T))$ – Sets, the equality $F \simeq F_{\text{forget}} \circ Q$ and the fact that Q maps connected objects to connected objects. □

Let us denote by $\pi_1(E, \mathcal{G}, s)$ the fundamental group of the infinite Galois category $(\text{lcs}(E, \mathcal{G}), F_s)$. The proposition above tells us that there is a continuous map of Noohi groups with dense image $*_{v \in E_0} \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T) \rightarrow \pi_1(E, \mathcal{G}, s)$. We now proceed to describe the kernel.

Recall that $\pi_1(E_{\leq 1}, T) = \text{Fr}(E_1) / \langle\langle \{\vec{e} \mid e \in T\}^{\text{Fr}(E_1)} \rangle\rangle$.

Theorem 3.7 (abstract Seifert–van Kampen theorem for infinite Galois categories). *E be a connected 2-complex with group data (\mathcal{G}, α) . With notations as above, the functor Q induces an isomorphism of Noohi groups*

$$(*_{v \in E_0} \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T) / \overline{H})^{\text{Noohi}} \rightarrow \pi_1(E, \mathcal{G}, s)$$

where \bar{H} is the closure of the group

$$H = \left\langle \left\langle \begin{array}{l} \mathcal{G}(\partial_1)(g)\vec{e} = \vec{e}\mathcal{G}(\partial_0)(g) \\ (\partial_2 f)\alpha_{102}^{(f)}(\alpha_{120}^{(f)})^{-1}(\partial_0 f)\alpha_{210}^{(f)}(\alpha_{201}^{(f)})^{-1}((\partial_1 f))^{-1}\alpha_{021}^{(f)}(\alpha_{012}^{(f)})^{-1} \end{array} \middle| \begin{array}{l} e \in E_1, g \in \mathcal{G}(e) \\ f \in E_2 \end{array} \right\rangle \right\rangle$$

where $\langle\langle - \rangle\rangle$ denotes the normal closure of the subgroup generated by the indicated elements and the α come from the definition of a (\mathcal{G}, α) -system for each given f .

Proof. The same proof as the proof of [Stix 2006, Theorem 3.2(2)] shows that Q induces an equivalence of categories between the infinite Galois categories $(\text{lcs}(E, \mathcal{G}), F_S)$ and the full subcategory of objects of $*_{v \in E_0}^N \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T)$ – Sets on which H acts trivially. We conclude by Observation 2.28. \square

Remark 3.8. It is important to note that we can replace free Noohi products by free topological products in the statement above, as we take the Noohi completion of the quotient anyway. More precisely, the canonical map

$$(*_{v \in E_0}^{\text{top}} \mathcal{G}(v) *^{\text{top}} \pi_1(E_{\leq 1}, T) / \bar{H}_1)^{\text{Noohi}} \rightarrow (*_{v \in E_0}^N \mathcal{G}(v) *^N \pi_1(E_{\leq 1}, T) / \bar{H})^{\text{Noohi}}$$

is an isomorphism, where H_1 is the normal closure in $*_{v \in E_0}^{\text{top}} \mathcal{G}(v) *^{\text{top}} \pi_1(E_{\leq 1}, T)$ of a group having the same generators as H . This is because the categories of G – Sets are the same for those two Noohi groups.

Fact 3.9. The topological free product $*_i^{\text{top}} G_i$ of topological groups has as an underlying space the free product of abstract groups $*_i G_i$. This follows from the original construction of Graev [1948].

3B. Application to the pro-étale fundamental group.

Descent data. Let T_\bullet be a 2-complex in a category \mathcal{C} and let $\mathcal{F} \rightarrow \mathcal{C}$ be a category fibered over \mathcal{C} , with $\mathcal{F}(S)$ as a category of sections above the object S .

Definition 3.10. The category $\text{DD}(T_\bullet, \mathcal{F})$ of descent data for \mathcal{F}/\mathcal{C} relative T_\bullet has as objects pairs (X', ϕ) where $X' \in \mathcal{F}(T_0)$ and ϕ is an isomorphism $\partial_0^* X' \xrightarrow{\sim} \partial_1^* X'$ in $\mathcal{F}(T_1)$ such that the cocycle condition holds, i.e., the following commutes in $\mathcal{F}(T_2)$:

$$\begin{array}{ccc} v_2^* X' & \xrightarrow{\partial_0^* \phi} & v_1^* X' \\ & \searrow \partial_1^* \phi & \swarrow \partial_2^* \phi \\ & v_0^* X' & \end{array}$$

Morphisms $F : (X', \phi) \rightarrow (Y', \psi)$ in $\text{DD}(T_\bullet, \mathcal{F})$ are morphisms $F : X' \rightarrow Y'$ in $\mathcal{F}(T_0)$ such that its two pullbacks $\partial_0^* f$ and $\partial_1^* f$ are compatible with ϕ, ψ , i.e., $\partial_1^* f \circ \phi = \psi \circ \partial_0^* f$.

Let $h : S' \rightarrow S$ be a map of schemes. There is an associated 2-complex of schemes

$$S_\bullet(h) : S' \rightrightarrows S' \times_S S' \rightleftarrows S' \times_S S' \times_S S'.$$

The value of ∂_i is the projection under omission of the i -th component. We abbreviate $\mathrm{DD}(S_\bullet(h), \mathcal{F})$ by $\mathrm{DD}(h, \mathcal{F})$. Observe that h^* gives a functor $h^* : \mathcal{F}(S) \rightarrow \mathrm{DD}(h, \mathcal{F})$.

Definition 3.11. In the above context $h : S' \rightarrow S$ is called an *effective descent* morphism for \mathcal{F} if h^* is an equivalence of categories.

Proposition 3.12 [Lavanda 2018, Proposition 1.16]. *Let $g : S' \rightarrow S$ be a proper, surjective morphism of finite presentation, then g is a morphism of effective descent for geometric coverings.*

Proof. This was proven by Lavanda and relies on the results of [Rydh 2010]. More precisely, this follows from Proposition 5.4 and Theorem 5.19 of [loc. cit.], then checking that the obtained algebraic space is a scheme (using étaleness and separatedness, see [Stacks 2020, Tag 0417]) and that it still satisfies the valuative criterion (see Lemma 2.40). \square

Discretization of descent data. We would like to apply the procedure described in [Stix 2006, Section 4.3] but to the pro-étale fundamental group. However, in the classical setting of Galois categories, given a category \mathcal{C} and functors $F, F' : \mathcal{C} \rightarrow \mathrm{Sets}$ such that (\mathcal{C}, F) and (\mathcal{C}, F') are Galois categories (i.e., F, F' are fiber functors), there exists an isomorphism (not unique) between F and F' . Choosing such an isomorphism is called “choosing a path” between F and F' . However, it is not clear whether an analogous statement is true for tame infinite Galois categories as the proof does not carry over to this case (see the proof of [Stacks 2020, Lemma 0BN5] or in [SGA 1 1971]—these proofs are essentially the same and rely on the pro-representability result of Grothendieck [1960, Proposition A.3.3.1]).

Question 3.13. Let \mathcal{C} be a category and $F, F' : \mathcal{C} \rightarrow \mathrm{Sets}$ be two functors such that (\mathcal{C}, F) and (\mathcal{C}, F') are tame infinite Galois categories. Is it true that F and F' are isomorphic?

As we do not know the answer to this question, we have to make an additional assumption when trying to discretize the descent data. Fortunately, it will always be satisfied in the geometric setting, which is our main case of interest.

Definition 3.14. Let $(\mathcal{C}, F), (\mathcal{C}', F')$ be two infinite Galois categories and let $\phi : \mathcal{C} \rightarrow \mathcal{C}'$ be a functor. We say that ϕ is *compatible* if there exists an isomorphism of functors $F \simeq F' \circ \phi$.

Let $\mathcal{F} \rightarrow \mathcal{C}$ be fibered in tame infinite Galois categories. More precisely, we have a notion of connected objects in \mathcal{C} and any $T \in \mathcal{C}$ is a coproduct of connected components. Over connected objects \mathcal{F} takes values in tame infinite Galois categories (i.e., over a connected $Y \in \mathcal{C}$ there exists a functor $F_Y : \mathcal{F}(Y) \rightarrow \mathrm{Sets}$ such that $(\mathcal{F}(Y), F_Y)$ is a tame infinite Galois category but we do not fix the functor).

Definition 3.15. Let T_\bullet be a 2-complex in \mathcal{C} . Let $E = \pi_0(T_\bullet)$ be its 2-complex of connected components: the 2-complex in Sets built by degree-wise application of the connected component functor. We will say that T_\bullet is a *compatible* 2-complex if one can fix fiber functors F_s of $\mathcal{F}(s)$ for each simplex $s \in E$ such that $(\mathcal{F}(s), F_s)$ is tame and for any boundary map $\partial : s \rightarrow s'$ there exists an isomorphism of fiber functors $F_s \circ T(\partial)^* \xrightarrow{\sim} F_{s'}$.

The 2-complexes that will appear in the (geometric) applications below will always be compatible. From now on, we will assume all 2-complexes to be compatible, even if not stated explicitly. Let T_\bullet be a compatible 2-complex in \mathcal{C} . Fix fiber functors F_s and isomorphisms between them as in the definition of a compatible 2-complex. For any ∂ , denote the fixed isomorphism by $\vec{\partial}$. For a 2-simplex (vef) of the barycentric subdivision with $\partial' : f \rightarrow e$ and $\partial : e \rightarrow v$ we define

$$\alpha_{vef} = \vec{\partial}' \vec{\partial} (\vec{\partial} \partial')^{-1}$$

or, more precisely,

$$\alpha_{vef} = T(\partial)(\vec{\partial}') \vec{\partial} (\vec{\partial} \partial')^{-1} \in \text{Aut}(F_v) = \pi_1(\mathcal{F}(s), F_s).$$

We define Noohi group data (\mathcal{G}, α) on E in the following way: $\mathcal{G}(s) = \pi_1(\mathcal{F}(s), F_s)$ for any simplex $s \in E$ and to $\partial : s \rightarrow s'$ is associated $\mathcal{G}(\partial) : \pi_1(\mathcal{F}(s), F_s) \xrightarrow{T(\partial)^*} \pi_1(\mathcal{F}(s'), F_{s'}) \xrightarrow{\vec{\partial}(\cdot) \vec{\partial}^{-1}} \pi_1(\mathcal{F}(s'), F_{s'})$. We define elements α as described above and we easily check that this gives Noohi group data.

Proposition 3.16. *The choice of functors F_s and the choice of $\vec{\partial}$ as above fix a functor*

$$\text{discr} : \text{DD}(T_\bullet, \mathcal{F}) \rightarrow \text{lcs}(E, (\mathcal{G}, \alpha))$$

which is an equivalence of categories.

Proof. Given a descent datum (X', ϕ) relative T_\bullet we have to attach a locally constant (\mathcal{G}, α) -system on E in a functorial way. For $v \in E_0, e \in E_1$ and $f \in E_2$, the definition of suitable $\mathcal{G}(v)$ (or $\mathcal{G}(e)$ or $\mathcal{G}(f)$) sets and maps m_∂ between them can be given by the same formulas as in [Stix 2006, Proposition 4.4] and also the same computations as in [loc. cit., Proposition 4.4] show that we obtain an element of $\text{lcs}(E, (\mathcal{G}, \alpha))$. Again, the reasoning of [loc. cit., Proposition 4.4] gives a functor in the opposite direction: given $M \in \text{lcs}(E, (\mathcal{G}, \alpha))$ we define $X' \in \mathcal{F}(T_0) = \prod_{v \in E_0} \mathcal{F}(v)$ as $X'|_v$ corresponding to M_v for all $v \in E_0$. Maps from edges to vertices define a map $\phi : T(\partial_0)^* X' \rightarrow T(\partial_1)^* X'$ and to check the cocycle condition one reverses the argument of the proof that discr gives a locally constant system. \square

To apply the last proposition we need to know that the compatibility condition holds in the setting we are interested in.

Lemma 3.17 [Bhatt and Scholze 2015, Lemma 7.4.1]. *Let $f : X' \rightarrow X$ be a morphism of two connected locally topologically noetherian schemes and let \bar{x}', \bar{x} be geometric points on X', X , correspondingly. Then the functor $f^* : \text{Cov}_X \rightarrow \text{Cov}_{X'}$ is a compatible functor between infinite Galois categories $(\text{Cov}_X, F_{\bar{x}})$ and $(\text{Cov}_{X'}, F_{\bar{x}'})$, i.e., the functors $F_{\bar{x}}$ and $F_{\bar{x}'} \circ f^*$ are isomorphic.*

Proof. Looking at the image of \bar{x}' (as a geometric point) on X , we reduce to the case when both \bar{x}' and \bar{x} lie on the same scheme X . In that case we proceed as in the second part of the proof of [loc. cit., Lemma 7.4.1]. \square

Corollary 3.18 [Bhatt and Scholze 2015, Lemma 7.4.1]. *Let X be a connected topologically noetherian scheme. Let \bar{x}_1, \bar{x}_2 be two geometric points on X . Then there is an isomorphism $\pi_1^{\text{proét}}(X, \bar{x}_1) \simeq \pi_1^{\text{proét}}(X, \bar{x}_2)$. It is unique (only) up to an inner automorphism.*

The above results combine to recover the analogue of [Stix 2006, Corollary 5.3] in the pro-étale setting.

Corollary 3.19. *Let $h : S' \rightarrow S$ be an effective descent morphism for geometric coverings. Assume that S is connected and $S, S', S' \times_S S', S' \times_S S' \times_S S'$ are locally topologically noetherian. Let $S' = \bigsqcup_v S'_v$ be the decomposition into connected components. Let \bar{s} be a geometric point of S , let $\bar{s}(t)$ be a geometric point of the simplex $t \in \pi_0(S_*(h))$, and let T be a maximal tree in the graph $\Gamma = \pi_0(S_*(h))_{\leq 1}$. For every boundary map $\partial : t \rightarrow t'$ let $\gamma_{t',t} : \bar{s}(t') \rightarrow S_*(h)(\partial)\bar{s}(t)$ be a fixed path (i.e., an isomorphism of fiber functors as in Lemma 3.17). Then canonically with respect to all these choices*

$$\pi_1^{\text{proét}}(S, \bar{s}) \cong \left(\left(\prod_{v \in E_0}^N \pi_1^{\text{proét}}(S'_v, \bar{s}(v)) \right) *^N \pi_1(\Gamma, T) \right) / \bar{H}^{\text{Noohi}}$$

where H is the normal subgroup generated by the cocycle and edge relations

$$\pi_1^{\text{proét}}(\partial_1)(g)\bar{e} = \bar{e}\pi_1^{\text{proét}}(\partial_0)(g), \tag{1}$$

$$\overrightarrow{(\partial_2 f)}\alpha_{102}^{(f)}(\alpha_{120}^{(f)})^{-1}\overrightarrow{(\partial_0 f)}\alpha_{210}^{(f)}(\alpha_{201}^{(f)})^{-1}(\overrightarrow{(\partial_1 f)})^{-1}\alpha_{021}^{(f)}(\alpha_{012}^{(f)})^{-1} = 1, \tag{2}$$

for all parameter values $e \in S_1(h)$, $g \in \pi_1^{\text{proét}}(e, \bar{s}(e))$, and $f \in S_2(h)$. The map $\pi_1^{\text{proét}}(\partial_i)$ uses the fixed path $\gamma_{\partial_i(e),e}$ and $\alpha_{ijk}^{(f)}$ is defined using $v \in S_0(h)$ and $e \in S_1(h)$ determined by $v_i(f) = v, \{\partial_0(e), \partial_1(e)\} = \{v_i(f), v_j(f)\}$ as

$$\alpha_{ijk}^{(f)} = \gamma_{v,e}\gamma_{e,f}\gamma_{v,f}^{-1} \in \pi_1^{\text{proét}}(v, \bar{s}(v)).$$

Remark 3.20. Similarly as in Remark 3.8, we could replace $*^N$ by $*^{\text{top}}$ in the above, as we take the Noohi completion of the whole quotient anyway.

Remark 3.21. We will often use Corollary 3.19 for h — the normalization map (or similar situations), where the connected components S'_v are normal. In this case $\pi_1^{\text{proét}}(S'_v, \bar{s}(v)) = \pi_1^{\text{ét}}(S'_v, \bar{s}_v)$. This implies that $\pi_1^{\text{proét}}(\partial_i)$ factorizes through the profinite completion of $\pi_1^{\text{proét}}(e, \bar{s}(e))$, which can be identified with $\pi_1^{\text{ét}}(e, \bar{s}(e))$. Moreover, the map $\pi_1^{\text{proét}}(e, \bar{s}(e)) \rightarrow \pi_1^{\text{ét}}(e, \bar{s}(e))$ has dense image and, in the end, we take the closure \bar{H} of H . The upshot of this discussion is that in the definition of generators of H we might consider $g \in \pi_1^{\text{ét}}(e, \bar{s}(e))$ instead of $g \in \pi_1^{\text{proét}}(e, \bar{s}(e))$ and $\pi_1^{\text{ét}}(\partial_i)$ instead of $\pi_1^{\text{proét}}(\partial_i)$, $i \in \{0, 1\}$, i.e.,

$$\pi_1^{\text{proét}}(S, \bar{s}) \cong \left(\left(\prod_{v \in E_0}^{\text{top}} \pi_1^{\text{ét}}(S'_v, \bar{s}(v)) \right) *^{\text{top}} \pi_1(\Gamma, T) \right) / \bar{H}^{\text{Noohi}}$$

where H is the normal subgroup generated by

$$\pi_1^{\text{ét}}(\partial_1)(g)\bar{e}\pi_1^{\text{ét}}(\partial_0)(g)^{-1}\bar{e}^{-1} \quad \text{for all } e \in S_1(h), g \in \pi_1^{\text{ét}}(e, \bar{s}(e)) \tag{R_1}$$

and

$$\overrightarrow{(\partial_2 f)}\alpha_{102}^{(f)}(\alpha_{120}^{(f)})^{-1}\overrightarrow{(\partial_0 f)}\alpha_{210}^{(f)}(\alpha_{201}^{(f)})^{-1}(\overrightarrow{(\partial_1 f)})^{-1}\alpha_{021}^{(f)}(\alpha_{012}^{(f)})^{-1} \quad \text{for all } f \in S_2(h). \tag{R_2}$$

Let us move on to some applications.

Ordered descent data. Let \mathcal{F} be a category fibered over \mathcal{C} (with a fixed cleavage, for convenience). Assume that \mathcal{C} is some subcategory of the category of locally topologically noetherian schemes with the property that finite fiber products in \mathcal{C} are the same as the finite fiber products as schemes. Let $h = \bigsqcup_{i \in I} h_i : S' = \bigsqcup_i S'_{i \in I} \rightarrow S$ be a morphism of schemes and let $<$ be a total order on the set of indices I . Let $S_{\bullet}^{<}(h) \subset S_{\bullet}(h)$ be the open and closed sub-2-complex of schemes in \mathcal{C} of ordered partial products

$$S_0^{<}(h) = S', \quad S_1^{<}(h) = \bigsqcup_{i < j} S'_i \times_S S'_j, \quad S_2^{<}(h) = \bigsqcup_{i < j < k} S'_i \times_S S'_j \times_S S'_k.$$

Proposition 3.22. *Let $h = \bigsqcup_{i \in I} h_i : S' = \bigsqcup_i S'_{i \in I} \rightarrow S$ be a morphism of schemes such that, for every $i, j \in I$, the maps induced by the diagonal morphisms $\Delta_i^* : \mathcal{F}(S'_i \times_S S'_i) \rightarrow \mathcal{F}(S'_i)$ and $(\Delta_i \times \text{id}_{S'_j})^* : \mathcal{F}(S'_i \times_S S'_j \times_S S'_j) \rightarrow \mathcal{F}(S'_i \times_S S'_j)$ are fully faithful. Then the natural open and closed immersion $S_{\bullet}^{<}(h) \hookrightarrow S_{\bullet}(h)$ induces an equivalence of categories*

$$\text{DD}(h, \mathcal{F}) \xrightarrow{\cong} \text{DD}(S_{\bullet}^{<}(h), \mathcal{F}).$$

Proof. For the problem at hand, we can and do replace \mathcal{F} by an equivalent category that admits a splitting cleavage (i.e., the associated pseudofunctor is a functor). Let $Y \in \mathcal{F}(S'_i)$ and consider $\partial_0^* Y, \partial_1^* Y \in \mathcal{F}(S'_i \times_S S'_i)$ obtained via maps induced by the projections $\mathcal{F}(S'_i) \rightarrow \mathcal{F}(S'_i \times_S S'_i)$. We first claim that there is *exactly one* isomorphism $\partial_0|_{S'_i \times_S S'_i}^* Y \rightarrow \partial_1|_{S'_i \times_S S'_i}^* Y$ as in the definition of descent data. Observe that $\Delta_i^* \partial_0^* Y = Y, \Delta_i^* \partial_1^* Y = Y$ and from the assumption any isomorphism $\phi : \partial_0^* Y|_{S'_i} \rightarrow \partial_1^* Y|_{S'_i}$ corresponds to precisely one isomorphism $\psi \in \text{Hom}_{S'_i}(Y|_{S'_i}, Y|_{S'_i})$. Pulling back the cocycle condition via the diagonal $\Delta_{2,i}^* : \mathcal{F}(S'_i \times_S S'_i \times_S S'_i) \rightarrow \mathcal{F}(S'_i)$ we get $\psi = \text{id}_{Y|_{S'_i}}$, so there is at most one map ϕ as above. Moreover, our assumptions imply that $\Delta_{2,i}^*$ is fully faithful as well, which shows that $\phi : \partial_0^* Y|_{S'_i} \rightarrow \partial_1^* Y|_{S'_i}$ corresponding to $\text{id}_{Y|_{S'_i}}$ will satisfy the condition. A similar reasoning shows that if we have ϕ_{ij} specified for $i < j$, then ϕ_{ji} is uniquely determined and if the ϕ_{ij} satisfy the cocycle condition on S_{ijk} for $i < j < k$, then the ϕ_{ij} together with the ϕ_{ji} obtained will satisfy the cocycle condition on any $S_{\alpha\beta\gamma}, \alpha, \beta, \gamma \in \{i, j, k\}$. \square

Observation 3.23. If the map of schemes $S'_i \rightarrow S$ is injective, i.e., if the diagonal map $S'_i \rightarrow S'_i \times_S S'_i$ is an isomorphism, then the assumptions of the proposition are satisfied.

Two examples.

Example 3.24. Let k be a field and C be \mathbb{P}_k^1 with two k -rational closed points p_0 and p_1 glued (see [Schwede 2005] for results on gluing schemes). Denote by p the node (i.e., the image of the p_i in C). We want to compute $\pi_1^{\text{proét}}(C)$. By the definition of C , we have a map $h : \tilde{C} = \mathbb{P}^1 \rightarrow C$ (which is also the normalization). It is finite, so it is an effective descent map for geometric coverings. Thus, we can use the van Kampen theorem. This goes as follows:

- Check that $\tilde{C} \times_C \tilde{C} \simeq \tilde{C} \sqcup p_{01} \sqcup p_{10}$ as schemes over C , where $p_{\alpha\beta}$ are equal to $\text{Spec}(k)$ and map to the node of C via the structural map. This can be done by checking that $\text{Hom}_C(Y, \tilde{C} \sqcup p_{01} \sqcup p_{10}) \simeq \text{Hom}_C(Y, \tilde{C}) \times \text{Hom}_C(Y, \tilde{C})$.

- Similarly, check that $\tilde{C} \times_C \tilde{C} \times_C \tilde{C} \simeq \tilde{C} \sqcup p_{001} \sqcup p_{010} \sqcup p_{011} \sqcup p_{100} \sqcup p_{101} \sqcup p_{110}$, where the projection $\tilde{C} \times_C \tilde{C} \times_C \tilde{C} \rightarrow \tilde{C} \times_C \tilde{C}$ omitting the first factor maps p_{abc} to p_{bc} and so on.
- We fix a geometric point $\bar{b} = \text{Spec}(\bar{k})$ over the base scheme $\text{Spec}(k)$ and fix geometric points \bar{p}_0 and \bar{p}_1 over p_0 and p_1 that map to \bar{b} . Then we fix geometric points on \tilde{C} , $p_{01}, p_{10} \subset \tilde{C} \sqcup p_{01} \sqcup p_{10} \simeq \tilde{C} \times_C \tilde{C}$ in a compatible way and similarly for connected components of $\tilde{C} \times_C \tilde{C} \times_C \tilde{C}$ (i.e., let us say that $\bar{p}_{\alpha\beta\gamma} \mapsto \bar{p}_\alpha$ via v_0 and $\bar{p}_{\alpha\beta} \mapsto \bar{p}_\alpha$). We fix a path γ from \bar{p}_0 to \bar{p}_1 that becomes trivial on $\text{Spec}(k)$ via the structural map (this can be done by viewing \bar{p}_0 and \bar{p}_1 as geometric points on $\tilde{C}_{\bar{k}}$, choosing the path on $\tilde{C}_{\bar{k}}$ first and defining γ to be its image). Let \bar{p} be the fixed geometric point on C given by the image of \bar{p}_0 (or, equivalently, \bar{p}_1).
- We want to use [Corollary 3.19](#) to compute $\pi_1^{\text{proét}}(C, \bar{p})$. We choose \bar{p}_0 as the base point $\bar{s}(\tilde{C})$ for $\tilde{C} \in \pi_0(S_0(h))$, $\tilde{C} \in \pi_0(S_1(h))$ and $\tilde{C} \in \pi_0(S_2(h))$. Then for any $t, t' \in \pi_0(S_\bullet(h))$ and the boundary map $\partial : t \rightarrow t'$, we use either the identity or γ to define $\gamma_{t',t} : \bar{s}(t') \rightarrow S_\bullet(h)(\partial)\bar{s}(t)$ as all the points \bar{p}_{abc} map ultimately either to \bar{p}_0 or \bar{p}_1 .

With this setup, the $\alpha_{ijk}^{(f)}$ (defined as in [Corollary 3.19](#)) are trivial for any f and so the relation (2) in this corollary reads $\overrightarrow{(\partial_2 f)} \overrightarrow{(\partial_0 f)} \overrightarrow{((\partial_1 f))}^{-1} = 1$. Applying this to different faces $f \in \pi_0(\tilde{C} \times_C \tilde{C} \times_C \tilde{C})$ gives that the image of $\pi_1(\Gamma, T) \simeq \mathbb{Z}^{*3}$ in $\pi_1^{\text{proét}}(C, \bar{p})$ is generated by a single edge (in our case only one maximal tree can be chosen – containing a single vertex). The choice of paths made guarantees $\pi_1^{\text{proét}}(\partial_0)(g) = \pi_1^{\text{proét}}(\partial_1)(g)$ in $\pi_1^{\text{proét}}(\tilde{C}, \bar{p}_0)$ for any $g \in \pi_1^{\text{proét}}(p_{ab}, \bar{p}_{ab}) = \text{Gal}(k)$. So relation (1) in [Corollary 3.19](#) implies that the image of $\pi_1^{\text{proét}}(\tilde{C}, \bar{p}_0) \simeq \text{Gal}(k)$ in $\pi_1^{\text{proét}}(C, \bar{p}_0)$ commutes with the elements of the image of $\pi_1(\Gamma, T)$. Putting this together we get

$$\begin{aligned} &\pi_1^{\text{proét}}(C, \bar{p}) \\ &\simeq ((\pi_1^{\text{proét}}(\tilde{C}, \bar{p}_0) *^{\text{top}} \pi_1(\Gamma, T)) / \langle \langle \pi_1^{\text{proét}}(\partial_1)(g)\bar{e} = \bar{e}\pi_1^{\text{proét}}(\partial_2)(g), \overrightarrow{(\partial_2 f)} \overrightarrow{(\partial_0 f)} \overrightarrow{((\partial_1 f))}^{-1} = 1 \rangle \rangle)^{\text{Noohi}} \\ &\simeq (\text{Gal}_k \times \mathbb{Z})^{\text{Noohi}} \\ &= \text{Gal}_k \times \mathbb{Z}. \end{aligned}$$

Example 3.25. Let X_1, \dots, X_m be geometrically connected normal curves over a field k and let Y_{m+1}, \dots, Y_n be nodal curves over k as in [Example 3.24](#). Let $x_i : \text{Spec}(k) \rightarrow X_i$ be rational points and let y_j denote the node of Y_j . Let $X := \cup_\bullet X_i \cup_\bullet Y_j$ be a scheme over k obtained via gluing of the X_i and Y_j along the rational points x_i and y_j (in the sense of [\[Schwede 2005\]](#)). The notation \cup_\bullet denotes gluing along the obvious points. The point of gluing gives a rational point $x : \text{Spec}(k) \rightarrow X$. We choose a geometric point $\bar{b} = \text{Spec}(\bar{k})$ over the base $\text{Spec}(k)$ and choose a geometric point \bar{x} over x such that it maps to \bar{b} . The maps $X_i \rightarrow X$ and $Y_j \rightarrow X$ are closed immersions (this is basically [\[Schwede 2005, Lemma 3.8\]](#)). We also get geometric points \bar{x}_i and \bar{y}_j over x_i and y_j that map to \bar{b} as well. Denote $\bar{X}_i = (X_i)_{\bar{k}}$. Let $\text{Gal}_{k,i} = \pi_1^{\text{ét}}(x_i, \bar{x}_i)$. It is a copy of Gal_k in the sense that the induced map $\pi_1^{\text{ét}}(x_i, \bar{x}_i) \rightarrow \pi_1^{\text{ét}}(\text{Spec}(k), \bar{b})$ is an isomorphism. Let us denote by $\iota_i : \text{Gal}_k \rightarrow \text{Gal}_{k,i}$ the inverse of this isomorphism. The group $\pi_1^{\text{ét}}(x_i, \bar{x}_i)$ acts on $\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i)$ and allows to write $\pi_1^{\text{ét}}(X_i, \bar{x}_i) \simeq \pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \rtimes \text{Gal}_{k,i}$.

After some computations (as in the previous example), using [Corollary 3.19](#) and [Example 3.24](#), one gets

$$\pi_1^{\text{proét}}(X, \bar{x}) \simeq \left(\ast_{1 \leq i \leq m}^N (\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \rtimes \text{Gal}_{k,i}) \ast_{m+1 \leq j \leq n}^N (\mathbb{Z} \times \text{Gal}_{k,j}) / \langle \langle \iota_i(\sigma) = \iota_{i'}(\sigma) \mid \sigma \in \text{Gal}_k, i, i' = 1, \dots, n \rangle \rangle \right)^{\text{Noohi}}.$$

Let us describe the category of group-sets:

$$\begin{aligned} & \left(\ast_{1 \leq i \leq m}^N (\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \rtimes \text{Gal}_{k,i}) \ast_{m+1 \leq j \leq n}^N (\mathbb{Z} \times \text{Gal}_{k,j}) / \langle \langle \iota_i(\sigma) = \iota_{i'}(\sigma) \mid \sigma \in \text{Gal}_k, i, i' = 1, \dots, n \rangle \rangle \right)^{\text{Noohi}} \text{-Sets} \\ & \simeq \{ S \in \left(\ast_{1 \leq i \leq m}^N (\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \rtimes \text{Gal}_{k,i}) \ast_{m+1 \leq j \leq n}^N (\mathbb{Z} \times \text{Gal}_{k,j}) \right)^{\text{Noohi}} \text{-Sets} \mid \forall i, i', \sigma \forall s \in S \iota_i(\sigma) \cdot s = \iota_{i'}(\sigma) \cdot s \} \\ & \simeq \{ S \in \left(\ast_{1 \leq i \leq m}^N \pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \ast^N \mathbb{Z}^{\ast n-m} \ast^N \text{Gal}_k \right)^{\text{Noohi}} \text{-Sets} \mid (\spadesuit) \} \end{aligned}$$

where the condition (\spadesuit) reads

$$\forall \sigma \in \text{Gal}_k, s \in S, 1 \leq i \leq m \forall \gamma \in \pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i), w \in \mathbb{Z}^{\ast n-m} (\sigma \cdot (\gamma \cdot s) =^\sigma \gamma \cdot (\sigma \cdot s) \text{ and } \sigma \cdot (w \cdot s) = w \cdot (\sigma \cdot s)).$$

We have used [Observation 2.28](#) and [Lemma 3.26](#) below.

Lemma 3.26. *Let K and Q be topological groups and assume we have a continuous action $K \times Q \rightarrow K$ respecting multiplication in K . Then $K \rtimes Q$ with the product topology (on $K \times Q$) is a topological group and there is an isomorphism*

$$K \ast^{\text{top}} Q / \langle \langle qkq^{-1} = {}^q k \rangle \rangle \rightarrow K \rtimes Q.$$

Proof. That $K \rtimes Q$ becomes a topological group is easy from the continuity assumption of the action. The isomorphism is obtained as follows: from the universal property we have a continuous homomorphism $K \ast^{\text{top}} Q \rightarrow K \rtimes Q$ and the kernel of this map is the smallest normal subgroup containing the elements $qkq^{-1}({}^q k)^{-1}$ (this follows from the fact that the underlying abstract group of $K \ast^{\text{top}} Q$ is the abstract free product of the underlying abstract groups, similarly for $K \rtimes Q$ and that we know the kernel in this case). So we have a continuous map that is an isomorphism of abstract groups. We have to check that the inverse map $K \rtimes Q \ni kq \mapsto kq \in K \ast^{\text{top}} Q / \langle \langle qkq^{-1} = {}^q k \rangle \rangle$ is continuous. It is enough to check that the map $K \times Q \ni (k, q) \mapsto kq \in K \ast^{\text{top}} Q$ (of topological spaces) is continuous, but this follows from the fact that the maps $K \rightarrow K \ast^{\text{top}} Q$ and $Q \rightarrow K \ast^{\text{top}} Q$ are continuous and that the multiplication map $(K \ast^{\text{top}} Q) \times (K \ast^{\text{top}} Q) \rightarrow K \ast^{\text{top}} Q$ is continuous. \square

Let us also state a technical lemma concerning the “functoriality” of the van Kampen theorem. It is important that the diagram formed by the schemes $X_1, X_2, \tilde{X}, \tilde{X}_1$ in the statement is cartesian.

Lemma 3.27. *Let $f : X_1 \rightarrow X_2$ be a morphism of connected schemes and $h : \tilde{X} \rightarrow X_2$ be a morphism of schemes. Denote by $h_1 : \tilde{X}_1 \rightarrow X_1$ the base-change of h via f . Assume that h and h_1 are effective descent morphisms for geometric coverings and that local topological noetherianity assumptions are satisfied for the schemes involved as in the statement of [Corollary 3.19](#). Assume that for any connected component $W \in \pi_0(S_*(h))$, the base-change W_1 of W via f is connected. Choose the geometric points on $W_1 \in \pi_0(S_*(h_1))$ and paths between the obtained fiber functors as in [Corollary 3.19](#) and choose the*

geometric points and paths on $W \in \pi_0(S_\bullet(h))$ as the images of those chosen for \tilde{X}_1 . Identify the graphs $\Gamma = \pi_0(S_\bullet(h))_{\leq 1}$ and $\Gamma_1 = \pi_0(S_\bullet(h_1))_{\leq 1}$ (it is possible thanks to the assumption made) and choose a maximal tree T in Γ . Using the above choices, use [Corollary 3.19](#). to write the fundamental groups

$$\pi_1^{\text{proét}}(X_1) \simeq \left(\left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W_1) \right) \ast^{\text{top}} \pi_1(\Gamma_1, T) / \langle R' \rangle \right)^{\text{Noohi}}$$

and

$$\pi_1^{\text{proét}}(X_2) \simeq \left(\left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W) \right) \ast^{\text{top}} \pi_1(\Gamma, T) / \langle R \rangle \right)^{\text{Noohi}}.$$

Then the map of fundamental groups $\pi_1^{\text{proét}}(f) : \pi_1^{\text{proét}}(X_1) \rightarrow \pi_1^{\text{proét}}(X_2)$ is the Noohi completion of the map

$$\left(\left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W_1) \right) \ast^{\text{top}} \pi_1(\Gamma_1, T) \right) / \langle R' \rangle \rightarrow \left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W) \ast^{\text{top}} \pi_1(\Gamma, T) \right) / \langle R \rangle,$$

which is induced by the maps $\pi_1^{\text{proét}}(W_1) \rightarrow \pi_1^{\text{proét}}(W)$ and the identity on $\pi_1(\Gamma_1, T)$ (which makes sense after identification of Γ_1 with Γ).

Proof. It is clear that on (the image of) $\pi_1^{\text{proét}}(W_1)$ (in $\pi_1^{\text{proét}}(X_1)$) the map is the one induced from $f_W : W_1 \rightarrow W$. The part about $\pi_1(\Gamma_1, T)$ follows from the fact that $\pi_1(\Gamma_1, T) < \pi_1^{\text{proét}}(X_1)$ acts in the same way as $\pi_1(\Gamma, T) < \pi_1^{\text{proét}}(X_2)$ on any geometric covering of X_2 . This follows from the choice of points and paths on $W \in \pi_0(S_\bullet(h))$ as the images of the points and paths on the corresponding connected components $W_1 \in \pi_0(S_\bullet(h_1))$. The maps as in the statement give a morphism $\phi : \left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W_1) \right) \ast^{\text{top}} \pi_1(\Gamma_1, T) \rightarrow \left(\ast_{W \in \pi_0(\tilde{X})}^{\text{top}} \pi_1^{\text{proét}}(W) \right) \ast^{\text{top}} \pi_1(\Gamma, T)$ and it is easy to check that $\phi(R') \subset R$, which finishes the proof. \square

3C. Künneth formula. In this subsection we use the van Kampen formula to prove the Künneth formula for $\pi_1^{\text{proét}}$.

Let X, Y be two connected schemes locally of finite type over an algebraically closed field k and assume that Y is proper. Let \bar{x}, \bar{y} be geometric points of X and Y respectively with values in the same algebraically closed field extension K of k . With these assumptions, the classical statement says that the “Künneth formula” for $\pi_1^{\text{ét}}$ holds, i.e., the following fact:

Fact 3.28 [[SGA 1 1971](#), Exposé X, Corollary 1.7]. With the above assumptions, the map induced by the projections is an isomorphism

$$\pi_1^{\text{ét}}(X \times_k Y, (\bar{x}, \bar{y})) \xrightarrow{\sim} \pi_1^{\text{ét}}(X, \bar{x}) \times \pi_1^{\text{ét}}(Y, \bar{y}).$$

We want to establish analogous statement for $\pi_1^{\text{proét}}$.

Proposition 3.29. *Let X, Y be two connected schemes locally of finite type over an algebraically closed field k and assume that Y is proper. Let \bar{x}, \bar{y} be geometric points of X and Y respectively with values in the same algebraically closed field extension K of k . Then the map induced by the projections is an isomorphism*

$$\pi_1^{\text{proét}}(X \times_k Y, (\bar{x}, \bar{y})) \xrightarrow{\sim} \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y}).$$

Choosing a path between (\bar{x}, \bar{y}) and some fixed k -point of $X \times_k Y$ (seen as a geometric point) and looking at the images of this path via projections onto X and Y reduces us (by [Corollary 3.18](#) and compatibility of the chosen paths), to the situation where we can assume that \bar{x} and \bar{y} are k -points. We are going to assume this in the proof. Before we start, let us state and prove the surjectivity of the above map as a lemma. Properness is not needed for this.

Lemma 3.30. *Let X, Y be two connected schemes over an algebraically closed field k with k -points on them: \bar{x} on X and \bar{y} on Y . Then the map induced by the projections*

$$\pi_1^{\text{proét}}(X \times_k Y, (\bar{x}, \bar{y})) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y})$$

is surjective.

Proof. Consider the map $(\text{id}_X, \bar{y}) : X = X \times_k \bar{y} \rightarrow X \times_k Y$. It is easy to check that the map induced on fundamental groups $\pi_1^{\text{proét}}(X, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X \times_k Y, (\bar{x}, \bar{y})) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y})$ is given by $(\text{id}_{\pi_1^{\text{proét}}(X, \bar{x})}, 1_{\pi_1^{\text{proét}}(Y, \bar{y})}) : \pi_1^{\text{proét}}(X, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y})$. Analogous fact holds if we consider $(\bar{x}, \text{id}_Y) : Y \rightarrow X \times_k Y$. As a result, the image $\text{im}(\pi_1^{\text{proét}}(X \times_k Y, (\bar{x}, \bar{y})) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y}))$ contains the set $(\pi_1^{\text{proét}}(X, \bar{x}) \times \{1_{\pi_1^{\text{proét}}(Y, \bar{y})}\}) \cup (\{1_{\pi_1^{\text{proét}}(X, \bar{x})}\} \times \pi_1^{\text{proét}}(Y, \bar{y}))$. This finishes the proof, as this set generates $\pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y})$. \square

Proof of Proposition 3.29. As X, Y are locally of finite type over a field, the normalization maps are finite and we can apply [Proposition 3.12](#). Let $\tilde{X} \rightarrow X$ be the normalization of X and let $\tilde{X} = \bigsqcup_v \tilde{X}_v$ be its decomposition into connected components and let us fix a closed point $x_v \in \tilde{X}_v$ for each v . Similarly, let $\bigsqcup_u \tilde{Y}_u = \tilde{Y} \rightarrow Y$ be the decomposition into connected components of the normalization of Y with closed points $y_u \in \tilde{Y}_u$.

We first deal with a particular case.

Claim. *The statement of Proposition 3.29 holds under the additional assumption that*

- either, for any v , the projections induce isomorphisms

$$\pi_1^{\text{proét}}(\tilde{X}_v \times_k Y, (x_v, \bar{y})) \xrightarrow{\sim} \pi_1^{\text{proét}}(\tilde{X}_v, x_v) \times \pi_1^{\text{proét}}(Y, \bar{y}),$$

- or, for any u , the projections induce isomorphisms

$$\pi_1^{\text{proét}}(X \times_k \tilde{Y}_u, (\bar{x}, y_u)) \xrightarrow{\sim} \pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(\tilde{Y}_u, y_u).$$

Proof of the claim. Apply [Corollary 3.19](#) to $h : \tilde{X} \rightarrow X$. We choose \bar{x} and x_v 's as geometric points $\bar{s}(t)$ of the corresponding simplexes $t \in \pi_0(S_\bullet(h))_0$ and choose $\bar{s}(t)$ to be arbitrary closed points (of suitable double and triple fiber products) for $t \in \pi_0(S_\bullet(h))_2$. We fix a maximal tree T in $\Gamma = \pi_0(S_\bullet(h))_{\leq 1}$ and fix paths $\gamma_{t',t} : \bar{s}(t') \rightarrow S_\bullet(h)(\partial)\bar{s}(t)$. Thus, we get $\pi_1^{\text{proét}}(X, \bar{x}) \cong ((*_v^N \pi_1^{\text{proét}}(\tilde{X}_v, x_v) *^N \pi_1(\Gamma, T))/\bar{H})^{\text{Noohi}}$ where H is defined as in [Corollary 3.19](#).

Observe now that $\tilde{X}_v \times_k Y$ are connected (as k is algebraically closed) and that $h \times \text{id}_Y : \tilde{X} \times Y \rightarrow X \times Y$ is an effective descent morphism for geometric coverings. So we might use [Corollary 3.19](#) in this setting. As

$(\tilde{X}_v \times Y) \times_{X \times Y} (\tilde{X}_w \times Y) = (\tilde{X}_v \times_X X_w) \times_k Y$, and similarly for triple products, we can identify in a natural way $i^{-1} : \pi_0(S_*(h \times \text{id}_Y)) \xrightarrow{\sim} \pi_0(S_*(h))$. In particular we can identify the graph $\Gamma_Y = \pi_0(S_*(h \times \text{id}_Y))_{\leq 1}$ with Γ and we choose the maximal tree T_Y of Γ_Y as the image of T via this identification. For $t \in \pi_0(S_*(h))$ choose $(\bar{s}(t), \bar{y})$ as the closed base points for $i(t) \in \pi_0(S_*(h \times \text{id}_Y))$. Denote by α_{ijk} elements of various $\pi_1^{\text{proét}}(\tilde{X}_v)$ defined as in [Corollary 3.19](#) and by \vec{e} elements of $\pi_1(\Gamma, T)$. By the choices and identifications above we can identify $\pi_1(\Gamma_Y, T_Y)$ with $\pi_1(\Gamma, T)$. Using van Kampen and the assumption, we write

$$\begin{aligned} \pi_1^{\text{proét}}(X \times Y, (\bar{x}, \bar{y})) &\cong ((*_v^N \pi_1^{\text{proét}}(\tilde{X}_v \times Y, (x_v, \bar{y})) *^N \pi_1(\Gamma_Y, T_Y)) / \overline{H_Y})^{\text{Noohi}} \\ &\cong ((*_v^N (\pi_1^{\text{proét}}(\tilde{X}_v, x_v) \times \pi_1^{\text{proét}}(Y, \bar{y})) *^N \pi_1(\Gamma, T)) / \overline{H_Y})^{\text{Noohi}}. \end{aligned}$$

Here $\pi_1^{\text{proét}}(Y, \bar{y})_v$ denotes a ‘‘copy’’ of $\pi_1^{\text{proét}}(Y, \bar{y})$ for each v . By [Lemma 3.30](#), for $T \in \pi_0(S_*(h))$ the natural map $\pi_1^{\text{proét}}(T \times Y, (\bar{s}(T), \bar{y})) \rightarrow \pi_1^{\text{proét}}(T, \bar{s}(T)) \times \pi_1^{\text{proét}}(Y, \bar{y})$ is surjective. It follows that the relations defining H_Y (as in [Corollary 3.19](#)) can be written as

$$\pi_1^{\text{proét}}(\partial_1)(g)h_{y,1}\vec{e} = \vec{e}\pi_1^{\text{proét}}(\partial_0)(g)h_{y,0},$$

for $e \in e(\Gamma)$, $g \in \pi_1^{\text{proét}}(e, \bar{s}(e))$, $e \in S_1(h)$, $h_y \in \pi_1^{\text{proét}}(Y, \bar{y})$, and

$$\overrightarrow{(\partial_2 f)} \alpha_{102} \alpha_{120}^{-1} \overrightarrow{(\partial_0 f)} \alpha_{210} \alpha_{201}^{-1} (\overrightarrow{(\partial_1 f)})^{-1} \alpha_{021} \alpha_{012}^{-1} = 1,$$

for $f \in S_2(h)$, where the α in the second relation are elements of suitable the $\pi_1^{\text{proét}}(\tilde{X}_v)$ and are the same as in the corresponding generators of H . The $h_{y,i}$ denotes a copy of element $h_y \in \pi_1^{\text{proét}}(Y, \bar{y})$ in a suitable $\pi_1^{\text{proét}}(Y, \bar{y})_v$. Varying e and h_y while choosing $g = 1 \in \pi_1^{\text{proét}}(e, \bar{s}(e))$ for every e , gives that $h_{y,1}\vec{e} = \vec{e}h_{y,0}$. For $e \in T$ we have $\vec{e} = 1$ and so the first relation reads $h_{y,1} = h_{y,0}$, i.e., it identifies $\pi_1^{\text{proét}}(Y, \bar{y})_v$ with $\pi_1^{\text{proét}}(Y, \bar{y})_w$ for v, w — the ends of the edge e . As T is a maximal tree in Γ , it contains all the vertices, so the first relation identifies $\pi_1^{\text{proét}}(Y, \bar{y})_v = \pi_1^{\text{proét}}(Y, \bar{y})_w$ for any two vertices v, w and we will denote this subgroup (of the quotient) by $\pi_1^{\text{proét}}(Y, \bar{y})$. This way $h_{y,1}\vec{e} = \vec{e}h_{y,0}$ reads simply $h_y\vec{e} = \vec{e}h_y$, so elements of $\pi_1^{\text{proét}}(Y, \bar{y})$ commute with elements of $\pi_1(\Gamma, T)$. Moreover, elements of $\pi_1^{\text{proét}}(Y, \bar{y})$ commute with elements of each $\pi_1^{\text{proét}}(\tilde{X}_v, x_v)$, as this was true for $\pi_1^{\text{proét}}(Y, \bar{y})_v$. On the other hand, choosing $h_y = 1$ in the first relation and looking at the second relation, we see that H_Y contains all the relations of H . Using notations from the above discussion, we can sum it up by writing

$$H_Y = \langle\langle \text{relations generating } H, h_{y,0} = h_{y,1}, h_y\vec{e} = \vec{e}h_y, h_yg = gh_y (g \in \pi_1^{\text{proét}}(\tilde{X}_v, x_v)) \rangle\rangle.$$

Putting this together, we get equivalences of categories

$$\begin{aligned} &((*_v^N \pi_1^{\text{proét}}(\tilde{X}_v, x_v) \times \pi_1^{\text{proét}}(Y, \bar{y})_v) *^N \pi_1(\Gamma, T)) / \overline{H_Y} - \text{Sets} \\ &\cong \{S \in (*_v^N (\pi_1^{\text{proét}}(\tilde{X}_v, x_v) \times \pi_1^{\text{proét}}(Y, \bar{y})) *^N \pi_1(\Gamma, T)) - \text{Sets} \mid H_Y \text{ acts trivially on } S\} \\ &\cong \{S \in ((*_v^N \pi_1^{\text{proét}}(\tilde{X}_v, x_v) *^N \pi_1(\Gamma, T)) \times \pi_1^{\text{proét}}(Y, \bar{y})) - \text{Sets} \mid H \text{ acts trivially on } S\} \\ &\cong (((*_v^N \pi_1^{\text{proét}}(\tilde{X}_v, x_v) *^N \pi_1(\Gamma, T)) / \overline{H}) \times \pi_1^{\text{proét}}(Y, \bar{y})) - \text{Sets} \\ &\cong (\pi_1^{\text{proét}}(X, \bar{x}) \times \pi_1^{\text{proét}}(Y, \bar{y})) - \text{Sets}, \end{aligned}$$

where equality \spadesuit follows from the fact that for topological groups G_1, G_2 there is an equivalence $(G_1 \times G_2) - \text{Sets} \cong \{S \in G_1 *^N G_2 - \text{Sets} \mid \forall_{g_1 \in G_1, g_2 \in G_2} \forall_{s \in S} g_1 g_2 s = g_2 g_1 s\}$ (see [Lemma 3.26](#)).

This finishes the proof of the Claim in the “either” case. After noting that each \tilde{Y}_u is still proper, the “or” case follows in a completely symmetrical manner. We have proven a particular case of the proposition. Let us now go ahead and prove the full statement. \square

General case. The general case follows from the claim proven above in the following way: Let $\bigsqcup_v \tilde{X}_v = \tilde{X} \rightarrow X$ and $\bigsqcup_u \tilde{Y}_u = \tilde{Y} \rightarrow Y$ be decompositions into connected components of the normalizations of X and Y . Fix v and note that $\pi_1^{\text{proét}}(\tilde{X}_v \times_k Y) = \pi_1^{\text{proét}}(\tilde{X}_v) \times \pi_1^{\text{proét}}(Y)$ by applying the claim to Y and \tilde{X}_v . This is possible, as the \tilde{Y}_u, \tilde{X}_v and the products $\tilde{Y}_u \times_k \tilde{X}_v$ (for all u) are normal varieties and so their pro-étale fundamental groups are equal to the usual étale fundamental groups (by [Lemma 2.12](#)) for which the equality $\pi_1^{\text{ét}}(\tilde{Y}_u \times_k \tilde{X}_v) = \pi_1^{\text{ét}}(\tilde{Y}_u) \times \pi_1^{\text{ét}}(\tilde{X}_v)$ is known (see [Fact 3.28](#)). Thus, for any v , we have that $\pi_1^{\text{proét}}(\tilde{X}_v \times_k Y) = \pi_1^{\text{proét}}(\tilde{X}_v) \times \pi_1^{\text{proét}}(Y)$. We can now apply the claim to X and Y and finish the proof in the general case. \square

3D. Invariance of $\pi_1^{\text{proét}}$ of a proper scheme under a base-change $K \supset k$ of algebraically closed fields.

Proposition 3.31. *Let X be a proper scheme over an algebraically closed field k . Let $K \supset k$ be another algebraically closed field. Then the pullback induces an equivalence of categories*

$$F : \text{Cov}_X \rightarrow \text{Cov}_{X_K} .$$

In particular, if X is connected, $X_K \rightarrow X$ induces an isomorphism

$$\pi_1^{\text{proét}}(X_K) \xrightarrow{\sim} \pi_1^{\text{proét}}(X) .$$

Proof. Let $X^\nu \rightarrow X$ be the normalization. It is finite, and thus a morphism of effective descent for geometric coverings. Let us show that the functor F is essentially surjective. Let $Y' \in \text{Cov}_{X_K}$. As k is algebraically closed and X^ν is normal, we conclude that X^ν is geometrically normal, and thus the base change $(X^\nu)_K$ is normal as well; see [[Stacks 2020](#), Tag 038O]. Pulling Y' back to $(X^\nu)_K$ we get a disjoint union of schemes finite étale over $(X^\nu)_K$ with a descent datum. It is a classical result [[SGA 1 1971](#), Exposé X, Corollary 1.8] that the pullback induces an equivalence $\text{Fét}_{X^\nu} \rightarrow \text{Fét}_{X^\nu_K}$ of finite étale coverings and similarly for the double and triple products $X_2^\nu = X^\nu \times_X X^\nu, X_3^\nu = X^\nu \times_X X^\nu \times_X X^\nu$. These equivalences obviously extend to categories whose objects are (possibly infinite) disjoint unions of finite étale schemes (over X^ν, X_2^ν, X_3^ν respectively) with étale morphisms as arrows. These categories can be seen as subcategories of Cov_{X^ν} and so on. These subcategories are moreover stable under pullbacks between $\text{Cov}_{X^\nu_i}$. Putting this together we see, that $Y'' = Y' \times_{X_K} (X^\nu)_K$ with its descent datum is isomorphic to a pullback of a descent datum from X^ν . Thus, we conclude that there exists $Y \in \text{Cov}_X$ such that $Y' \simeq Y_K$. Full faithfulness of F is shown in the same way. If X is connected, it can be also proven more directly, as F being fully faithful is equivalent to preserving connectedness of geometric coverings, but any connected $Y \in \text{Cov}_X$ is geometrically connected, and thus Y_K remains connected by [Proposition 2.37\(2\)](#). Note that in the above argument we do not claim that the double and triple intersections X_2^ν, X_3^ν are

normal, as this is in general false. Instead, we are only using that all the considered geometric coverings of those schemes came as pullbacks from X^ν , and thus were already split-to-finite. \square

4. Fundamental exact sequence

4A. Statement of the results and examples. The main result of this chapter is the following theorem.

Theorem (see [Theorem 4.14](#) below). *Let k be a field and fix an algebraic closure \bar{k} . Let X be a geometrically connected scheme of finite type over k . Then the sequence of abstract groups*

$$1 \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k \rightarrow 1$$

is exact.

Moreover, the map $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ is a topological embedding and the map $\pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k$ is a quotient map of topological groups.

One shows the near exactness first and obtains the above version as a corollary with an extra argument. The most difficult part of the sequence is exactness on the left. We will prove it as a separate theorem and its proof occupies an entire subsection.

Theorem (see [Theorem 4.13](#) below). *Let k be a field and fix an algebraic closure \bar{k} of k . Let X be a scheme of finite type over k such that the base change $X_{\bar{k}}$ is connected. Then the induced map*

$$\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$$

is a topological embedding.

By [Proposition 2.37](#), it translates to the following statement in terms of coverings: every geometric covering of $X_{\bar{k}}$ can be dominated by a covering that embeds into a base-change to \bar{k} of a geometric covering of X (i.e., defined over k). In practice, we prove that every connected geometric covering of $X_{\bar{k}}$ can be dominated by a (base-change of a) covering of X_l for l/k finite.

For finite coverings, the analogous statement is very easy to prove simply by finiteness condition. But for general geometric coverings this is nontrivial and maybe even slightly surprising as we show by counterexamples ([Examples 4.5](#) and [4.6](#)) that it is not always true that a connected geometric covering of $X_{\bar{k}}$ is isomorphic to a base-change of a covering of X_l for some finite extension l/k . This last statement is, however, stronger than what we need to prove, and thus does not contradict our theorem. Observe, that the stronger statement is true for finite coverings and, even more generally, whenever $\pi_1^{\text{proét}}(X_{\bar{k}})$ is prodiscrete, as proven in [Proposition 4.8](#).

Let us proceed to proving the easier part of the sequence first.

Observation 4.1. By [Proposition 2.17](#), the category of geometric coverings is invariant under universal homeomorphisms. In particular, for a connected X over a field and k'/k purely inseparable, there is $\pi_1^{\text{proét}}(X_{k'}) = \pi_1^{\text{proét}}(X)$. Similarly, we can replace X by X_{red} and so assume X to be reduced when convenient. In this case, base change to separable closure X_{k^s} is reduced as well. We will often use this observation without an explicit reference.

We start with the following lemmas.

Lemma 4.2. *Let k be a field. Let $k \subset k'$ be a (possibly infinite) Galois extension. Let X be a connected scheme over k . Let $\bar{T}_0 \subset \pi_0(X_{k'})$ be a nonempty closed subset preserved by the $\text{Gal}(k'/k)$ -action. Then $\bar{T}_0 = \pi_0(X_{k'})$.*

Proof. Let \bar{T} be the preimage of \bar{T}_0 in $X_{k'}$ (with the reduced induced structure). By [Stacks 2020, Lemma 038B], \bar{T} is the preimage of a closed subset $T \subset X$ via the projection morphism $p : X_{k'} \rightarrow X$. On the other hand, by [loc. cit., Lemma 04PZ], the image $p(\bar{T})$ equals the entire X . Thus, $T = X$ and $\bar{T} = X_{k'}$, and so $\bar{T}_0 = \pi_0(X_{k'})$. □

Lemma 4.3. *Let X be a connected scheme over a field k with an l' -rational point with l'/k a finite field extension. Then $\pi_0(X_{k^{\text{sep}}})$ is finite, the Gal_k action on $\pi_0(X_{k^{\text{sep}}})$ is continuous and there exists a finite separable extension l/k such that the induced map $\pi_0(X_{k^{\text{sep}}}) \rightarrow \pi_0(X_l)$ is a bijection. Moreover, there exists the smallest field (contained in k^{sep}) with this property and it is Galois over k .*

Proof. Let us first show the continuity of the Gal_k -action. The morphism $\text{Spec}(l') \rightarrow X$ gives a Gal_k -equivariant morphism $\text{Spec}(l' \otimes_k k^{\text{sep}}) \rightarrow X_{k^{\text{sep}}}$ and a Gal_k -equivariant map $\pi_0(\text{Spec}(l' \otimes_k k^{\text{sep}})) \rightarrow \pi_0(X_{k^{\text{sep}}})$. Denote by $M \subset \pi_0(X_{k^{\text{sep}}})$ the image of the last map. It is finite and Gal_k -invariant, and by Lemma 4.2, $M = \pi_0(X_{k'})$. We have tacitly used that M is closed, as $\pi_0(X_{k'})$ is Hausdorff (as the connected components are closed). As Gal_k acts continuously on $\pi_0(\text{Spec}(l' \otimes_k k^{\text{sep}}))$ (for example by [Stacks 2020, Lemma 038E]), we conclude that it acts continuously on $\pi_0(X_{k^{\text{sep}}})$ as well. From Lemma 4.2 again and from [loc. cit., Tag 038D], we easily see that the fields $l \subset k^{\text{sep}}$ such that $\pi_0(X_{k^{\text{sep}}}) \rightarrow \pi_0(X_l)$ is a bijection are precisely those that Gal_l acts trivially on $\pi_0(X_{k^{\text{sep}}})$. To get the minimal field with this property we choose l such that $\text{Gal}_l = \ker(\text{Gal}_k \rightarrow \text{Aut}(\pi_0(X_{k^{\text{sep}}}))$). □

Theorem 4.4. *Let k be a field and fix an algebraic closure \bar{k} . Let X be a geometrically connected scheme of finite type over k . Let $\bar{x} : \text{Spec}(\bar{k}) \rightarrow X_{\bar{k}}$ be a geometric point on $X_{\bar{k}}$. Then the induced sequence*

$$\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \xrightarrow{\iota} \pi_1^{\text{proét}}(X, \bar{x}) \xrightarrow{p} \text{Gal}_k \rightarrow 1$$

of topological groups is nearly exact in the middle (i.e., the thick closure of $\text{im}(\iota)$ equals $\ker(p)$) and $\pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k$ is a topological quotient map.

Proof. (1) The map p is surjective and open: let $U < \pi_1^{\text{proét}}(X)$ be an open subgroup. There is a geometric covering Y of X with a \bar{k} -point \bar{y} such that the morphism $\pi_1^{\text{proét}}(Y, \bar{y}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$ is equal to $U \subset \pi_1^{\text{proét}}(X, \bar{x})$. As Y is locally of finite type over k , the image of \bar{y} in Y has a finite extension l of k as the residue field. Thus, we get $\text{Gal}_l \rightarrow \pi_1^{\text{proét}}(Y) \rightarrow \text{Gal}_k$ and we see that the image $\pi_1^{\text{proét}}(Y) \rightarrow \text{Gal}_k$ contains an open subgroup, so is open. We have shown that p is an open morphism. In particular the image of $\pi_1^{\text{proét}}(X)$ in Gal_k is open and so also closed. On the other hand, this image is dense as we have

the following diagram:

$$\begin{array}{ccc}
 \pi_1^{\text{proét}}(X) & \longrightarrow & \pi_1^{\text{proét}}(\text{Spec}(k)) \\
 \downarrow & & \parallel \\
 \widehat{\pi_1^{\text{proét}}(X)}^{\text{prof}} = \pi_1^{\text{ét}}(X) & \twoheadrightarrow & \pi_1^{\text{ét}}(\text{Spec}(k))
 \end{array}$$

where $\widehat{\cdot}^{\text{prof}}$ means the profinite completion. In the diagram, the left vertical map has dense image and the lower horizontal is surjective. This shows that $\pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k$ is surjective.

(2) The composition $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \rightarrow \text{Gal}_k$ is trivial — this is clear thanks to [Proposition 2.37](#) and the fact that the map $X_{\bar{k}} \rightarrow \text{Spec}(k)$ factorizes through $\text{Spec}(\bar{k})$.

(3) The thick closure of $\text{im}(\iota)$ is normal: as remarked above, $\pi_1^{\text{proét}}(X_{\bar{k}}) = \pi_1^{\text{proét}}(X_{k^s})$, where k^s denotes the separable closure. Thus, we are allowed to replace \bar{k} with k^s in the proof of this point. Moreover, by the same remark, we can and do assume X to be reduced. Let $Y \rightarrow X$ be a connected geometric covering such that there exists a section $s : X_{k^s} \rightarrow Y \times_X X_{k^s} = Y_{k^s}$ over X_{k^s} . Observe that any such section is a clopen immersion: this follows immediately from the equivalence of categories of $\pi_1^{\text{proét}}(X_{k^s}) - \text{Sets}$ and geometric coverings. Define $\bar{T} := \bigcup_{\sigma \in \text{Gal}(k)} \sigma s(X_{k^s}) \subset Y_{k^s}$. Observe that two images of sections in the sum either coincide or are disjoint as X_{k^s} is connected and they are clopen. Now, \bar{T} is obviously open, but we claim that it is also a closed subset. This follows from [Lemma 4.3](#) (which implies that $\pi_0(Y_{k^s})$ is finite), but one can also argue directly by using that Y_{k^s} is locally noetherian and $\sigma s(X_{k^s})$ are clopen. Now by [\[Stacks 2020, Tag 038B\]](#), \bar{T} descends to a closed subset $T \subset Y$. It is also open as T is the image of \bar{T} via projection $Y_{k^s} \rightarrow Y$ which is surjective and open map. Indeed, surjectivity is clear and openness is easy as well and is a particular case of a general fact, that any map from a scheme to a field is universally open [\[loc. cit., Tag 0383\]](#) By connectedness of Y we see that $T = Y$. So $Y_{k^s} = \bar{T}$. But this last one is a disjoint union of copies of X_{k^s} , which is what we wanted to show by [Proposition 2.37](#).

(4) The smallest normal thickly closed subgroup of $\pi_1^{\text{proét}}(X)$ containing $\text{im}(\iota)$ is equal to $\ker(p)$: as we already know that this image is contained in the kernel and that the map $\pi_1^{\text{proét}}(X) \rightarrow \text{Gal}_k$ is a quotient map of topological groups, we can apply [Proposition 2.37](#). Let Y be a connected geometric covering of X such that $Y_{\bar{k}} = Y \times_X X_{\bar{k}}$ splits completely. Denote $Y_{\bar{k}} = \bigsqcup_{\alpha} X_{\bar{k}, \alpha}$, where by $X_{\bar{k}, \alpha}$ we label different copies of $X_{\bar{k}}$. By [Lemma 4.3](#), $\pi_0(Y_{\bar{k}})$ is finite, and thus the indexing set $\{\alpha\}$ and the covering $Y \rightarrow X$ are finite. But in this case, the statement follows from the classical exact sequence of étale fundamental groups due to Grothendieck. \square


As promised above, we give examples of geometric coverings of $X_{\bar{k}}$ that cannot be defined over any finite field extension l/k .

Example 4.5. Let $X_i = \mathbb{G}_{m, \mathbb{Q}}$, $i = 1, 2$. Define X to be the gluing $X = \cup_{\bullet} X_i$ of these schemes at the rational points $1_i : \text{Spec}(\mathbb{Q}) \rightarrow X_i$ corresponding to 1. Fix an algebraic closure $\bar{\mathbb{Q}}$ of \mathbb{Q} and so a geometric point \bar{b} over the base $\text{Spec}(\mathbb{Q})$. This gives geometric points \bar{x}_i on $\bar{X}_i = X_{i, \bar{\mathbb{Q}}}$ and X_i lying over 1_i , which we choose as base points for the fundamental groups involved. Similarly, we get a geometric point \bar{x}

over the point of gluing x that maps to \bar{b} . Then [Example 3.25](#) gives us a description of the fundamental group $\pi_1^{\text{proét}}(X, \bar{x}) \simeq \left(\ast_{i=1,2}^{\text{top}} (\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \rtimes \text{Gal}_{\mathbb{Q},i}) / \langle \langle \iota_1(\sigma) = \iota_2(\sigma) \mid \sigma \in \text{Gal}_{\mathbb{Q}} \rangle \rangle \right)^{\text{Noohi}}$ and of its category of sets:

$$\pi_1^{\text{proét}}(X, \bar{x}) - \text{Sets} \simeq \left\{ S \in \left(\ast^{\text{top}} \pi_1^{\text{ét}}(\bar{X}_1) \ast^{\text{top}} \pi_1^{\text{ét}}(\bar{X}_2) \ast^{\text{top}} \text{Gal}_{\mathbb{Q}} \right) - \text{Sets} \mid \forall \sigma \in \text{Gal}_{\mathbb{Q}} \forall_i \forall_{\gamma \in \pi_1^{\text{ét}}(\bar{X}_i)} \forall_{s \in S} \sigma \cdot (\gamma \cdot s) = \sigma \gamma \cdot (\sigma \cdot s) \right\}.$$

For the base change \bar{X} to $\bar{\mathbb{Q}}$, we have $\pi_1^{\text{proét}}(\bar{X}, \bar{x}) \simeq \pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1) \ast^N \pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$. Recall that the groups $\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i)$ are isomorphic to $\widehat{\mathbb{Z}}(1) = \varprojlim \mu_n$ as $\text{Gal}_{\mathbb{Q}}$ -modules. Fix these isomorphisms. Let $S = \mathbb{N}_{>0}$. Let us define a $\pi_1^{\text{proét}}(\bar{X}, \bar{x})$ -action on S , which means giving actions by $\pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1)$ and $\pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$ (no compatibilities of the actions required). Let ℓ be a fixed odd prime number (e.g., $\ell = 3$). We will give two different actions of $\mathbb{Z}_{\ell}(1)$ on S which will define actions of $\widehat{\mathbb{Z}}(1)$ by projections on $\mathbb{Z}_{\ell}(1)$. We start by dividing S into consecutive intervals labeled a_1, a_3, a_5, \dots of cardinality $\ell^1, \ell^3, \ell^5, \dots$ respectively. These will be the orbits under the action of $\pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1)$. Similarly, we divide S into consecutive intervals b_2, b_4, b_6, \dots of cardinality ℓ^2, ℓ^4, \dots .



We still have to define the action on each a_m and b_m . We choose arbitrary identifications $b_m \simeq \mu_{\ell^m}$ as $\mathbb{Z}_{\ell}(1)$ -modules. Now, fix a compatible system of ℓ^n -th primitive roots of unity $\zeta = (\zeta_{\ell^n}) \in \mathbb{Z}(1)$. For a_m 's, we choose the identifications with μ_{ℓ^m} arbitrarily with one caveat: we demand that for any even number m , the intersection $b_m \cap a_{m+1}$ contains the elements $1, \zeta_{\ell^{m+1}} \in \mu_{\ell^{m+1}}$ via the chosen identification $a_{m+1} \simeq \mu_{\ell^{m+1}}$. As $|b_m \cap a_{m+1}| > 0$ and $|b_m \cap a_{m+1}| \equiv 0 \pmod{\ell}$, the intersection $b_m \cap a_{m+1}$ contains at least two elements and we see that choosing such a labeling is always possible.

Assume that S corresponds to a covering that can be defined over a finite Galois extension K/\mathbb{Q} . Fix $s_0 \in a_1 \cap b_2$. By increasing K , we might and do assume that Gal_K fixes s_0 . Let p be a prime number $\neq \ell$ that splits completely in K and \mathfrak{p} be a prime of \mathcal{O}_K lying above p . Let $\phi_{\mathfrak{p}} \in \text{Gal}_K$ be a Frobenius element (which depends on the choice of the decomposition group and the coset of the inertia subgroup). It acts on $\mathbb{Z}_{\ell}(1)$ via $t \mapsto t^p$ and this action is independent of the choice of $\phi_{\mathfrak{p}}$. Choose $N > 0$ such that $p^N \equiv 1 \pmod{\ell^2}$ and let m be the biggest number such that $p^N \equiv 1 \pmod{\ell^m}$. If m is odd, we look at $p^{\ell N}$ instead. In this case $m + 1$ is the biggest number such that $p^{\ell N} \equiv 1 \pmod{\ell^{m+1}}$ and so, by changing N if necessary, we can assume that m is even, > 1 . The whole point of the construction is the following: if $s \in a_i \cap b_j$ with $i, j < m$ is fixed by $\phi_{\mathfrak{p}}^N$, then so are $g \cdot s$ and $h \cdot s$ (for $h \in \pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1)$ and $g \in \pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$). Then moving such s with the g and h to $b_m \cap a_{m+1}$ leads to a contradiction. Indeed, let $s_1 \in b_m \cap a_{m+1} \subset S$ correspond to $1 \in \mu_{\ell^{m+1}} \simeq a_{m+1}$ (it is possible by the choices made in the construction of S). Write $s_1 = g_m h_{m-1} \cdots h_3 g_2 h_1 \cdot s_0$ with $h_i \in \pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1)$ and $g_j \in \pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$ (this form is not unique, of course). This is possible thanks to the fact that the sets a_i, b_j form consecutive intervals separately such that b_j intersects nontrivially a_{j-1} and a_{j+1} . By the construction of S again, there is an element $s_2 \in b_m \cap a_{m+1}$

corresponding to $\zeta_{\ell^{m+1}} \in \mu_{\ell^{m+1}}$ via $a_{m+1} \simeq \mu_{\ell^{m+1}}$. We can now write s_2 in two ways:

$$s_2 = \zeta \cdot s_1 = g \cdot s_1,$$

where $g \in \pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$ and ζ is the chosen element in $\pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1) \simeq \widehat{\mathbb{Z}}(1)$. By the choices made, the action of ϕ_p^N fixes the elements s_1 and $g \cdot s_1$, while it moves $\zeta \cdot s_1$. Indeed, $\phi_p^N \cdot (\zeta \cdot s_1) = (\phi_p^N \zeta \phi_p^{-N}) \cdot (\phi_p^N \cdot s_1) = \zeta^{p^N} \cdot (\phi_p^N \cdot s_1) = \zeta^{p^N} \cdot s_1 = \zeta_{\ell^{m+1}}^{p^N} \neq \zeta_{\ell^{m+1}} = \zeta \cdot s_1 \in \mu_{\ell^{m+1}} \simeq a_{m+1}$ — a contradiction.

Example 4.6. Let $X_i = \mathbb{G}_{m, \mathbb{Q}}, i = 1, 2, 3$ and let X_4, X_5 be the nodal curves obtained from gluing 1 and -1 on $\mathbb{P}_{\mathbb{Q}}^1$ (see Example 3.24). Define X to be the gluing $X = \cup_{\bullet} X_i$ of all these schemes at the rational points corresponding to 1 (or the image of 1 in the case of the nodal curves). We fix an algebraic closure $\bar{\mathbb{Q}}$ of \mathbb{Q} and so fix a geometric point \bar{b} over the base $\text{Spec}(\mathbb{Q})$. We get geometric points \bar{x}_i on $\bar{X}_i = X_i \times_{\mathbb{Q}} \bar{\mathbb{Q}}$ lying over 1. We have and fix the following isomorphisms of $\text{Gal}_{\mathbb{Q}}$ -modules. For $1 \leq i \leq 3$, $\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \simeq \widehat{\mathbb{Z}}(1)$ and for $4 \leq j \leq 5$, we have $\pi_1^{\text{proét}}(\bar{X}_j, \bar{x}_j) \simeq \langle t^{\mathbb{Z}} \rangle$ (i.e., \mathbb{Z} written multiplicatively). Let $t_i \in \pi_1^{\text{proét}}(\bar{X}_i, \bar{x}_i)$ be the elements corresponding via these isomorphisms to a fixed inverse system of primitive roots $\zeta \in \widehat{\mathbb{Z}}(1)$ (for $i = 1, 2, 3$) and to $t \in \langle t^{\mathbb{Z}} \rangle$ (for $i = 4, 5$). Example 3.25 gives a description of the fundamental group

$$\pi_1^{\text{proét}}(X, \bar{x}) \simeq \left(\ast_{i=1,2,3}^N (\widehat{\mathbb{Z}}(1)_i \rtimes \text{Gal}_{\mathbb{Q}, i}) \ast_{j=4,5}^N (\langle t^{\mathbb{Z}} \rangle \times \text{Gal}_{\mathbb{Q}, j}) / \overline{\left\langle \iota_i(\sigma) = \iota_{i'}(\sigma) \mid i, i' = 1, \dots, 5 \right\rangle} \right)^{\text{Noohi}}$$

and of its category of sets:

$$\pi_1^{\text{proét}}(X, \bar{x}) - \text{Sets} \simeq \{ S \in (\ast_{1 \leq i \leq 3}^{\text{top}} \widehat{\mathbb{Z}}(1)_i) \ast^{\text{top}} \langle t^{\mathbb{Z}} \rangle^{\ast 2} \ast^{\text{top}} \text{Gal}_{\mathbb{Q}}) - \text{Sets} \mid \forall \sigma \in \text{Gal}_{\mathbb{Q}} \forall 1 \leq i \leq 3 \forall \gamma \in \mathbb{Z}(1)_i \forall s \in S \sigma \cdot (\gamma \cdot s) = \sigma \gamma \cdot (\sigma \cdot s) \text{ and } \sigma \cdot (w \cdot s) = w \cdot (\sigma \cdot s) \}.$$

Let $G = \left\{ \begin{pmatrix} \ast & \ast \\ \ast & \ast \end{pmatrix} \right\} \subset \text{GL}_2(\mathbb{Q}_{\ell})$ be the subgroup of upper triangular matrices. Fix $u_1 \in \mathbb{Z}_{\ell}^{\times}$ such that $u_1^p \neq u_1$. Let $H = \ast_i^{\text{top}} \pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i)$ and define a continuous homomorphism $\psi : H \rightarrow G$ by:

$$\psi(t_1) = \begin{pmatrix} u_1 & \\ & 1 \end{pmatrix}, \quad \psi(t_2) = \begin{pmatrix} 1 & \\ & u_1 \end{pmatrix}, \quad \psi(t_3) = \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}, \quad \psi(t_4) = \begin{pmatrix} \ell & \\ & 1 \end{pmatrix}, \quad \psi(t_5) = \begin{pmatrix} 1 & \\ & \ell \end{pmatrix}.$$

It is easy to see that ψ is surjective.

Let $U \subset G$ be the subgroup of matrices with elements in \mathbb{Z}_{ℓ} , i.e., $U = \left\{ \begin{pmatrix} \ast & \ast \\ & \ast \end{pmatrix} \right\} \subset \text{GL}_2(\mathbb{Z}_{\ell})$. It is an open subgroup of G . Thus, using ψ and the fact that $H^{\text{Noohi}} = \pi_1^{\text{proét}}(\bar{X}, \bar{x})$, we get that $S := G/U$ defines a $\pi_1^{\text{proét}}(\bar{X}, \bar{x})$ -set. It is connected (i.e., transitive) and so corresponds to a connected geometric covering of \bar{X} . Assume that it can be defined over a finite extension L of \mathbb{Q} . We can assume L/\mathbb{Q} is Galois. By the description above, it means that there is a compatible action of groups $\mathbb{Z}(1)_i, \mathbb{Z}^{\ast 2}$ and Gal_L on S . By increasing L , we can assume moreover that Gal_L fixes $[U]$.

Choose $p \neq \ell$ that splits completely in Gal_L , fix a prime \mathfrak{p} of L dividing p and let $\phi_{\mathfrak{p}} \in \text{Gal}_L$ denote a fixed Frobenius element. Let $t_3^{u_1}$ denote the unique element of $\psi_{|\pi_1^{\text{ét}}(\bar{X}_3, \bar{x}_3)}^{-1} \left(\begin{pmatrix} 1 & u_1 \\ & 1 \end{pmatrix} \right)$. Let $n \gg 0$. An easy calculation shows that $\psi(t_4^{-n} t_1 t_3 t_1^{-1} t_3^{-u_1} t_4^n) = 1_{\text{GL}_2(\mathbb{Q}_{\ell})} \in U$. Then $\phi_{\mathfrak{p}} \cdot [U] = \phi_{\mathfrak{p}} \cdot (t_4^{-n} t_1 t_3 t_1^{-1} t_3^{-u_1} t_4^n \cdot [U]) =$

$\phi_p(t_4^{-n}t_1t_3t_1^{-1}t_3^{-u_1}t_4^n) \cdot (\phi_p \cdot [U]) = t_4^{-n}t_1^p t_3^p t_1^{-p} t_3^{-pu_1} t_4^n \cdot [U]$. But

$$\begin{aligned} \psi(t_4^{-n}t_1^p t_3^p t_1^{-p} t_3^{-pu_1} t_4^n) &= \begin{pmatrix} \ell^{-n} & & & \\ & 1 & & \\ & & u_1^p & \\ & & & 1 \end{pmatrix} \begin{pmatrix} 1 & p \\ & 1 \end{pmatrix} \begin{pmatrix} u_1^{-p} & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & -pu_1 \\ & 1 \end{pmatrix} \begin{pmatrix} \ell^n & \\ & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & \ell^{-n} p(u_1^p - u_1) & & \\ & & & \\ & & & \\ & & & 1 \end{pmatrix} \notin U. \end{aligned}$$

As $n \gg 0$ and $u_1^p \neq u_1$, it follows that $\phi_p \cdot [U] \neq [U]$ — a contradiction.

It is important to note, that the above (counter)examples are possible only when considering the geometric coverings that are not trivialized by an étale cover (but one really needs to use the pro-étale cover to trivialize them). In [Bhatt and Scholze 2015], the category of geometric coverings trivialized by an étale cover on X is denoted by $\text{Loc}_{X_{\text{ét}}}$ and the authors prove the following

Fact 4.7 [Bhatt and Scholze 2015, Lemma 7.4.5]. Under $\text{Loc}_X \simeq \pi_1^{\text{proét}}(X) - \text{Sets}$, the full subcategory $\text{Loc}_{X_{\text{ét}}} \subset \text{Loc}_X$ corresponds to the full subcategory of those $\pi_1^{\text{proét}}(X) - \text{Sets}$ where an open subgroup acts trivially.

We are now going to prove:

Proposition 4.8. *Let X be a geometrically connected separated scheme of finite type over a field k . Let $Y \in \text{Cov}_{X_{\bar{k}}}$ be such that $Y \in \text{Loc}_{(X_{\bar{k}})_{\text{ét}}}$. Then there exists a finite extension l/k such and $Y_0 \in \text{Cov}_{X_l}$ such that $Y \simeq Y_0 \times_{X_l} X_{\bar{k}}$.*

Proof. By the topological invariance (Proposition 2.17), we can replace \bar{k} by k^{sep} if desired. By the assumption $Y \in \text{Loc}_{(X_{\bar{k}})_{\text{ét}}}$, there exists an étale cover of finite type that trivializes Y . Being of finite type, it is a base-change $X'_{\bar{k}} = X' \times_{\text{Spec}(l)} \text{Spec}(\bar{k}) \rightarrow X_{\bar{k}}$ of an étale cover $X' \rightarrow X_l$ for some finite extension l/k . Thus, $Y|_{X'_{\bar{k}}}$ is constant (i.e., $\simeq \bigsqcup_{s \in S} X' = \underline{S}$) and the isomorphism between the pull-backs of $Y|_{X'_{\bar{k}}}$ via the two projections $X'_{\bar{k}} \times_{X_{\bar{k}}} X'_{\bar{k}} \rightrightarrows X'_{\bar{k}}$ is expressed by an element of a constant sheaf $\underline{\text{Aut}}(\underline{S})(X'_{\bar{k}} \times_{X_{\bar{k}}} X'_{\bar{k}}) = \text{Aut}(\underline{S})(X'_{\bar{k}} \times_{X_{\bar{k}}} X'_{\bar{k}})$ (we use the fact that $X'_{\bar{k}}$ is étale over $X_{\bar{k}}$, and thus $\pi_0(X'_{\bar{k}} \times_{X_{\bar{k}}} X'_{\bar{k}})$ is discrete, in this case even finite). By enlarging l , we can assume that the connected components of the schemes involved: X' , $X' \times_{X_l} X'$ etc. are geometrically connected over l . Define $Y'_0 = \bigsqcup_{s \in S} X'$. The discussion above shows that the descent datum on $Y|_{X'_{\bar{k}}}$ with respect to $X'_{\bar{k}} \rightarrow X_{\bar{k}}$ is in fact the pull-back of a descent datum on Y'_0 with respect to $X' \rightarrow X_l$. As étale covers are morphisms of effective descent for geometric coverings (this follows from the fpqc descent for fpqc sheaves and the equivalence $\text{Cov}_{X_l} \simeq \text{Loc}_{X_l}$ of [Bhatt and Scholze 2015, Lemma 7.3.9]), the proof is finished. \square

Remark 4.9. Over a scheme with a nondiscrete set of connected components, $\underline{\text{Aut}}(\underline{S})$ might not be equal to $\text{Aut}(\underline{S})$.

Proposition 4.8 shows that our main theorem is significantly easier for π_1^{SGA3} .

Corollary 4.10. *Let X be a geometrically connected separated scheme of finite type over a field k . Fix an algebraic closure \bar{k} of k . Then*

$$\pi_1^{\text{SGA3}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{SGA3}}(X)$$

is a topological embedding.

4B. Preparation for the proof of Theorem 4.13. We are going to use the following proposition.

Proposition 4.11. *Let X be a scheme of finite type over a field k with a k -rational point x_0 and assume that $X_{\bar{k}}$ is connected. Let Y_1, \dots, Y_N be a set of connected finite étale coverings of $X_{\bar{k}}$. Then there exists a finite Galois covering Y of $X_{\bar{k}}$ that dominates each Y_i and such that the corresponding normal subgroup of $\pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$ is normalized by $\text{Gal}_k = \pi_1^{\text{ét}}(x_0, \bar{x}_0)$ in $\pi_1^{\text{ét}}(X, \bar{x}_0)$.*

Proof. There is a finite connected Galois covering of $X_{\bar{k}}$ dominating Y_1, \dots, Y_N . Thus, we can assume $N = 1$ and Y_1 is Galois. Fix a geometric point \bar{x}_0 over x_0 . The splitting $s : \text{Gal}_k = \pi_1^{\text{ét}}(x_0, \bar{x}_0) \rightarrow \pi_1^{\text{ét}}(X, \bar{x}_0)$ allows us to write $\pi_1^{\text{ét}}(X, \bar{x}_0) \simeq \pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0) \rtimes \text{Gal}_k$. Fix a geometric point \bar{y} on Y_1 over \bar{x}_0 . The group $U = \pi_1^{\text{ét}}(Y_1, \bar{y})$ is a normal open subgroup of $\pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$. As the pair (Y_1, \bar{y}) is defined over a finite Galois field extension l/k (contained in \bar{k}), it is easy to check that $\text{Gal}_l \subset \text{Gal}_k$ fixes U , i.e., ${}^\sigma U = U$ for $\sigma \in \text{Gal}_l$. It follows that the set of conjugates ${}^\sigma U$ is finite, of cardinality bounded by $[l : k]$. Define $V = \bigcap_{\sigma \in \text{Gal}_k} {}^\sigma U$. It follows that this is an open subgroup of $\pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$ fixed by the action of Gal_k . Moreover, it is normal in $\pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$, as for any $g \in \pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$, there is $g(\bigcap_{\sigma \in \text{Gal}_k} {}^\sigma U)g^{-1} = \bigcap_{\sigma \in \text{Gal}_k} g {}^\sigma U g^{-1} = \bigcap_{\sigma \in \text{Gal}_k} {}^\sigma ({}^{\sigma^{-1}} g)U({}^{\sigma^{-1}} g^{-1}) = \bigcap_{\sigma \in \text{Gal}_k} {}^\sigma U$, due to normality of U . This open subgroup $V < \pi_1^{\text{ét}}(X_{\bar{k}}, \bar{x}_0)$ corresponds to a covering with the desired properties. \square

Before starting the proof, we need to collect some facts about the Galois action on the geometric $\pi_1^{\text{ét}}$. They are discussed, for example, in [Stix 2013, Chapter 2]. The existence, functoriality and compatibility with compositions of the action can be readily seen to generalize to $\pi_1^{\text{proét}}$ as well, but *note (see the last point below) that one has to be careful when discussing continuity*. For a connected topologically noetherian scheme W and geometric points \bar{w}_1, \bar{w}_2 , let $\pi_1^{\text{proét}}(W, \bar{w}_1, \bar{w}_2) = \text{Isom}_{\text{Cov}_{W_{\bar{k}}}}(F_{\bar{w}_1}, F_{\bar{w}_2})$ denote the set of isomorphisms of the two fiber functors, topologized in a way completely analogous to the case when $\bar{w}_1 = \bar{w}_2$. By Corollary 3.18, it is a bitorsor under $\pi_1^{\text{proét}}(W, \bar{w}_1)$ and $\pi_1^{\text{proét}}(W, \bar{w}_2)$. The bitorsors under profinite groups $\pi_1^{\text{ét}}(W, \bar{w}_1, \bar{w}_2)$ are defined similarly and are rather standard. For a geometrically unibranch W , the two notions match.

Lemma 4.12. *For a scheme W of finite type over k and two geometric points \bar{w}_1, \bar{w}_2 on W lying over k -points, there is an abstract Gal_k -action on $\pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$ and $\pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$ such that:*

- (a) *It is given by $\psi_\sigma = \pi_1^{\text{ét}}(\text{id}_W \times_{\text{Spec}(k)} \text{Spec}(\sigma^{-1}), \bar{w}_1, \bar{w}_2)$ or an analogously defined automorphism of $\pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$. This makes sense as \bar{w}_1, \bar{w}_2 are Gal_k -invariant.*
- (b) *The morphism $\pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2) \rightarrow \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$ is Gal_k -equivariant. Similarly, maps of schemes $(W, \bar{w}_1, \bar{w}_2) \rightarrow (W', \bar{w}_1, \bar{w}_2)$ induce Gal_k -equivariant maps on $\pi_1^{\text{ét}}$ and $\pi_1^{\text{proét}}$.*

(c) For three geometric points $\bar{w}_1, \bar{w}_2, \bar{w}_3$, the Galois action is compatible with the composition maps, i.e., for any $\sigma \in \text{Gal}_k$, the following diagram commutes:

$$\begin{array}{ccc} \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_2, \bar{w}_3) \times \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2) & \xrightarrow{(-) \circ (-)} & \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_3) \\ \downarrow (\psi_\sigma, \psi_\sigma) & & \downarrow \psi_\sigma \\ \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_2, \bar{w}_3) \times \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2) & \xrightarrow{(-) \circ (-)} & \pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_3) \end{array}$$

Similarly for $\pi_1^{\text{ét}}$. Inductively, this also holds for arbitrary (finite) composition maps.

(d) Let s_{w_1}, s_{w_2} be the sections of the maps $\pi_1^{\text{ét}}(W, \bar{w}_i) \rightarrow \text{Gal}_k$ coming from rational points w_1, w_2 . Then $\psi_\sigma(\gamma) = s_{w_2}(\sigma) \circ \gamma \circ s_{w_1}(\sigma^{-1})$ for $\gamma \in \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2) \subset \pi_1^{\text{ét}}(W, \bar{w}_1, \bar{w}_2)$. Note that, while ψ_σ is defined for $\pi_1^{\text{proét}}$, the right-hand side of this formula only makes sense thanks to the fundamental exact sequence for $\pi_1^{\text{ét}}$ (and its version for the sets of paths, see [Stix 2013, Proposition 18]). Thus, at this stage, we cannot make an analogous statement for $\pi_1^{\text{proét}}$.

In terms of continuity of ψ_σ , there is a priori a huge difference in how much we can say about $\pi_1^{\text{ét}}$ and $\pi_1^{\text{proét}}$:

- (e) For each σ , the map ψ_σ is continuous as an automorphism of $\pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$ and $\pi_1^{\text{proét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$.
- (f) The action $\text{Gal}_k \times \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2) \rightarrow \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{w}_1, \bar{w}_2)$ is continuous. Note, however, that **at this stage of the proof we do not know whether this is true for $\pi_1^{\text{proét}}$** . In fact, this is closely related to the main result we need to prove.

4C. Proof that $\pi_1^{\text{proét}}(X_{\bar{k}}) \rightarrow \pi_1^{\text{proét}}(X)$ is a topological embedding. In this subsection we finally prove our main result.

Theorem 4.13. Let k be a field and fix an algebraic closure \bar{k} of k . Let X be a scheme of finite type over k such that the base-change $X_{\bar{k}}$ is connected. Let \bar{x} be a $\text{Spec}(\bar{k})$ -point on $X_{\bar{k}}$. Then the induced map

$$\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$$

is a topological embedding.

Then, we will derive the final form of the fundamental exact sequence.

Theorem 4.14. With the assumptions as in Theorem 4.13, the sequence of abstract groups

$$1 \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x}) \rightarrow \text{Gal}_k \rightarrow 1$$

is exact.

Moreover, the map $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$ is a topological embedding and the map $\pi_1^{\text{proét}}(X, \bar{x}) \rightarrow \text{Gal}_k$ is a quotient map of topological groups.

In the proof, after some preparatory steps (e.g., extending the field k), we define the set of *regular loops* in $\pi_1^{\text{proét}}(X_{\bar{k}})$ with respect to a fixed open subgroup $U < \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ and use it to construct an Galois invariant open subgroup V inside of U (see Steps II and III below). *There is also an alternative approach* to proving the existence of V that avoids the direct construction involving regular loops. *We sketch it in Remark 4.27.* While this latter approach is quicker, it is less instructive: as explained in Remark 4.26 below, the notion of a regular loop provides an insight of what goes wrong in the counterexample Example 4.5. Still, it might be worth having a look at, as our main approach is rather lengthy.

Step I: Setting things up and applying van Kampen. For any finite extension $\bar{k}/l/k$ of k , the map $\pi_1^{\text{proét}}(X_l, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$ is an embedding of an open subgroup and we have a factorization $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X_l, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$. Here, we have tacitly lifted \bar{x} to X_l . Thus, we can start by replacing k by a finite extension. Considering the normalization $X^\nu \rightarrow X$, base-changing the whole problem to a finite extension l of k , considering the factorization $l/l'/k$ into separable and purely inseparable extension of fields, and using first that the base-change along a separable field extension of a normal scheme is normal and then the topological invariance of $\pi_1^{\text{proét}}$, we can assume that we have a surjective finite morphism $h : \tilde{X} \rightarrow X$ such that the connected components of \tilde{X} , $\tilde{X} \times_X \tilde{X}$, $\tilde{X} \times_X \tilde{X} \times_X \tilde{X}$ are geometrically connected, have rational points and for each $W \in \pi_0(\tilde{X})$, there is $\pi_1^{\text{proét}}(W) = \pi_1^{\text{ét}}(W)$ and $\pi_1^{\text{proét}}(W_{\bar{k}}) = \pi_1^{\text{ét}}(W_{\bar{k}})$.

Let $\tilde{X} = \bigsqcup_{v \in \text{Vert}} \tilde{X}_v$ be the decomposition into connected components. Note that the indexing set Vert is finite. For each $t \in \pi_0(\tilde{X}) \cup \pi_0(\tilde{X} \times_X \tilde{X}) \cup \pi_0(\tilde{X} \times_X \tilde{X} \times_X \tilde{X})$, we fix a k -rational point $x(t)$ on t and a \bar{k} -point $\bar{x}(t)$ on $\bar{t} = t_{\bar{k}}$ lying over $x(t)$. We will often write \bar{x}_t to mean $\bar{x}(t)$. Let us fix $v_{\bar{x}} \in \text{Vert}$ for the rest of the text and say that the image of $\bar{x}(\tilde{X}_{v_{\bar{x}}, \bar{k}})$ in $X_{\bar{k}}$ will be the fixed geometric point \bar{x} of $X_{\bar{k}}$ and its image in X the fixed geometric point of X . For any $W_{\bar{k}}, W'_{\bar{k}} \in \pi_0(S_{\bullet}(\bar{h}))$ and every boundary map $\bar{\partial} : W_{\bar{k}} \rightarrow W'_{\bar{k}}$, we fix paths $\gamma_{W'_{\bar{k}}, W_{\bar{k}}} \in \pi_1^{\text{proét}}(W'_{\bar{k}}, \bar{x}_{W'_{\bar{k}}}, \bar{\partial}(\bar{x}_{W_{\bar{k}}}))$ between the chosen geometric points, as in Corollary 3.19. This is possible thanks to Lemma 3.17. We define $\gamma_{W', W}$ to be the image of this path.

Let $\bar{h} : \tilde{X}_{\bar{k}} \rightarrow X_{\bar{k}}$ be the base-change of h . We choose a maximal tree T (resp. T') in the graph $\Gamma = \pi_0(S_{\bullet}(h))_{\leq 1}$ (resp. $\Gamma' = \pi_0(S_{\bullet}(\bar{h}))_{\leq 1}$). After making these choices, we can apply Corollary 3.19 with Remark 3.21 to write the fundamental groups of (X, \bar{x}) and $(X_{\bar{k}}, \bar{x})$. This way we get a diagram

$$\begin{CD} ((\ast_v^{\text{top}} \pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v)) \ast^{\text{top}} \pi_1(\Gamma', T')) / \overline{\langle R'_1, R'_2 \rangle^{nc}} \text{Noohi} @>\cong>> \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \\ @VVV @VVV \\ ((\ast_v^{\text{top}} \pi_1^{\text{ét}}(\tilde{X}_v, \bar{x}_v)) \ast^{\text{top}} \pi_1(\Gamma, T)) / \overline{\langle R_1, R_2 \rangle^{nc}} \text{Noohi} @>\cong>> \pi_1^{\text{proét}}(X, \bar{x}) \end{CD}$$

where $\overline{(\cdot)}$ denotes the topological closure, $\langle R \rangle^{nc}$ denotes the normal subgroup generated by the set R , and R_1, R'_1, R_2, R'_2 are as in Remark 3.21.

Note that, while the (connected components of the) fiber products $\tilde{X} \times_X \tilde{X}$, $\tilde{X} \times_X \tilde{X} \times_X \tilde{X}$ are not necessarily normal nor satisfy $\pi_1^{\text{proét}}(W) = \pi_1^{\text{ét}}(W)$, we can effectively work as if this was the case, see Remark 3.21.

Observation 4.15. The maps and groups above enjoy the following properties:

(a) By Lemma 3.27, the left vertical map is the Noohi completion of the obvious map of the underlying quotients of free topological products.

(b) By geometrical connectedness of the schemes in sight, we can (and do) identify

$$\pi_0(\mathcal{S}_\bullet(h)) = \pi_0(\mathcal{S}_\bullet(\bar{h})), \quad \Gamma' = \Gamma \quad \text{and} \quad T' = T.$$

(c) As the $\gamma_{W,W}$ are chosen to be the images of the $\gamma_{W'_k, W_k}$, we see that $\alpha_{abc}^{(f)}$'s appearing in R_2 , and so a priori elements of the $\pi_1^{\text{ét}}(\tilde{X}_v, \bar{x}_v)$, are in fact in $\pi_1^{\text{ét}}(\tilde{X}_{v,\bar{k}}, \bar{x}_v)$. It follows that

$$R'_2 = R_2.$$

(d) The k -rational points $x(W)$ give identification

$$\pi_1^{\text{ét}}(W, \bar{x}_W) \simeq \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{x}_W) \rtimes \text{Gal}_k.$$

When $W = \tilde{X}_v$ for $v \in \text{Vert}$, we will write $\text{Gal}_{k,v}$ in the identification above to distinguish between different copies of Gal_k in the van Kampen presentation of $\pi_1^{\text{proét}}(X, \bar{x})$.

(e) As $\gamma_{W',W}$ is the image of the path $\gamma_{W'_k, W_k}$ on W'_k , it maps to the trivial element of $\text{Gal}_k = \pi_1^{\text{ét}}(\text{Spec}(k), \bar{x}(W), \bar{x}(W'))$. It implies, that the following diagram commutes:

$$\begin{array}{ccc} \pi_1^{\text{ét}}(W_{\bar{k}}, \bar{x}(W)) & \longrightarrow & \pi_1^{\text{ét}}(W, \bar{x}(W)) \\ \downarrow & & \downarrow \\ \pi_1^{\text{ét}}(W'_{\bar{k}}, \bar{x}(W')) & \longrightarrow & \pi_1^{\text{ét}}(W', \bar{x}(W')) \end{array} \quad \begin{array}{c} \nearrow \\ \searrow \end{array} \text{Gal}_k$$

Let P be a walk in Γ , i.e., a sequence of consecutive edges (with possible repetitions) e_1, \dots, e_m in Γ with an orientation such that the terminal vertex of e_i is the initial vertex of e_{i+1} . Using the orientation of Γ , it can be written as $\epsilon_1 e_1 \dots \epsilon_m e_m$ with $\epsilon_i \in \{\pm\}$ indicating whether the orientation agrees or not. This will come handy as follows: define $\partial_0^+ = \partial_0, \partial_0^- = \partial_1, \partial_1^+ = \partial_1, \partial_1^- = \partial_0$.

For each P as above with a vertex sequence $(v_1, v_2, \dots, v_{m+1})$, there is a map

$$\pi_1^{\text{ét}}(\tilde{X}_{v_{m+1},\bar{k}}, \partial_1^{\epsilon_m}(\bar{x}_{e_m}), \bar{x}_{v_{m+1}}) \times \pi_1^{\text{ét}}(\tilde{X}_{v_m,\bar{k}}, \bar{x}_{v_m}, \partial_0^{\epsilon_m}(\bar{x}_{e_m})) \times \dots \times \pi_1^{\text{ét}}(\tilde{X}_{v_2,\bar{k}}, \partial_1^{\epsilon_1}(\bar{x}_{e_1}), \bar{x}_{v_2}) \times \pi_1^{\text{ét}}(\tilde{X}_{v_1,\bar{k}}, \bar{x}_{v_1}, \partial_0^{\epsilon_1}(\bar{x}_{e_1})) \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}_{v_1}, \bar{x}_{v_{m+1}})$$

where

$$(\gamma_{2m}, \dots, \gamma_1) \mapsto \gamma_{2m} \circ \dots \circ \gamma_1.$$

In the following, we will use $\circ_?$ to denote the “composition of étale paths” and $\bullet_?$ to denote the multiplication in some group(oid) $?$. When $? = \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ or $\pi_1^{\text{proét}}(X, \bar{x})$, we will skip the subscript. While we could just use $\circ_?$ everywhere, it is sometimes convenient to keep track of when some paths “have been closed” by using $\bullet_?$.

Step II: Defining regular loops in $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$.

Definition 4.16. An element $\gamma \in \text{Isom}_{\text{Cov}_{X_{\bar{k}}}}(F_{\bar{x}_w}, F_{\bar{x}_v})$ is called an *étale path of special form supported on P* if it lies in the image of the composition map above for some walk P starting in w and ending in v .

Any element $(\gamma_{2m}, \dots, \gamma_1)$ in the preimage of such γ will be called a *presentation* of γ with respect to P .

For a walk P , denote by $l(P)$ the length of P , i.e., the number of consecutive edges (not necessarily different) it is composed of.

Observation 4.17. A useful example of a path of special form is the following. In the van Kampen presentation, the maps $\pi_1^{\text{ét}}(\tilde{X}_{v,\bar{k}}, \bar{x}_v) = \pi_1^{\text{proét}}(\tilde{X}_{v,\bar{k}}, \bar{x}_v) \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ are given by

$$\rho_v(-) = \gamma_v^{-1} \circ (-) \circ \gamma_v$$

where $\gamma_v \in \pi_1^{\text{proét}}(\tilde{X}_{v,\bar{k}}, \bar{x}, \bar{x}_v)$ is defined as follows: if $P_{v_{\bar{x}},v} \subset T$ denotes the unique shortest path in the tree $T \subset \Gamma$ (forgetting the orientation) from $v_{\bar{x}}$ to v , then the choices of paths $\gamma_{W_{\bar{k}}, W_{\bar{k}}}$ made when applying the van Kampen theorem give a unique étale path of special form γ_v supported on $P_{v_{\bar{x}},v}$.

Before introducing the main objects of the proof, we note a simple result.

Lemma 4.18. For a fixed path $\gamma \in \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}, \bar{y})$ of special form, the map $\text{Gal}_k \ni \sigma \mapsto \psi_\sigma(\gamma) \in \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}, \bar{y})$ is continuous.

Proof. This follows from the continuity of the composition maps of paths and the fact that the statement is true for $\pi_1^{\text{ét}}$. □

To prove [Theorem 4.13](#), it is enough to prove the following statement: any connected geometric covering Y of $X_{\bar{k}}$ can be dominated by a covering defined over a finite separable extension l/k .

Indeed, let $Y' \in \text{Cov}_{X_l}$ be a connected covering that dominates Y after base-change to \bar{k} . By looking at the separable closure of k in l and using the topological invariance of $\pi_1^{\text{proét}}$, we can assume l/k is separable. The composition $Y'' = Y' \rightarrow X_l \rightarrow X$ is an element of Cov_X and there is a diagonal embedding $Y' \times_{\text{Spec}(l)} \text{Spec}(\bar{k}) \rightarrow Y'' \times_{\text{Spec}(k)} \text{Spec}(\bar{k})$. By [Proposition 2.37\(5\)](#), the proof will be finished.

Let us fix a connected $Y \in \text{Cov}$ till the end of the proof and denote by $S = Y_{\bar{x}}$ the corresponding $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ -set. Fix some point $s_0 \in S$ and let $U = \text{Stab}_{\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})}(s_0)$.

Definition 4.19. For each $v \in \text{Vert}$, define

$$O_v^N = \{s \in F_{\bar{x}_v}(Y) \mid \exists_{\substack{\text{walk } P, \\ l(P) \leq N}} \exists_{\substack{\gamma \text{ of sp. form,} \\ \text{supp. on } P}} s = \gamma \cdot s_0\}$$

and call it the set of “elements at v reachable in at most N steps”.

The following is a crucial observation regarding O_v^N .

Lemma. For any v and N , the set O_v^N is finite.

Proof. We proceed by induction on N . For $N = 1$, the walks of length not greater than N starting in v_0 (are either trivial or) consist of a single edge whose initial vertex is necessarily $v_{\bar{x}}$. As Γ is finite, there are only finitely many such edges. Let us fix one, named e , with vertices v_0, w . We need to show that the set

$$\{(\theta \circ \delta) \cdot s_0 \in F_{\bar{x}_w}(Y) \mid \delta \in \pi_1^{\text{ét}}(\tilde{X}_{v_{\bar{x}}, \bar{k}}, \bar{x}, \partial_1^{\epsilon(e)}(\bar{x}_e)), \theta \in \pi_1^{\text{ét}}(\tilde{X}_{w, \bar{k}}, \partial_0^{\epsilon(e)}(\bar{x}_e), \bar{x}_w)\}$$

is finite. However, as in general the sets $\pi_1^{\text{ét}}(W, \bar{x}_1, \bar{x}_2)$ are (bi)torsors under profinite groups (namely $\pi_1^{\text{ét}}(W, \bar{x}_1)$ and $\pi_1^{\text{ét}}(W, \bar{x}_2)$) and the maps and actions in sight are continuous, we see that the finiteness of this last set follows directly from finiteness of orbits of points in discrete sets under an action by a profinite group.

Now, to see the inductive step, assume that the claim is true for N . To prove it for $N + 1$, note that any element in O_v^{N+1} can be connected by a single edge to an element of O_w^N (for some vertex w). As O_w^N is finite and as we have just explained that, starting from a fixed point, one can only reach finitely many points by applying étale paths of special forms supported on a single edge, the result follows. \square

Now, for each $v \in \text{Vert}$ and $N \in \mathbb{N}$, define $C_v^N \in \pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v) - \text{FSets}$ so that:

- (1) It is Galois.
- (2) It dominates each of the $\pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v)$ -orbits of elements of O_v^N .
- (3) The corresponding open normal subgroup $\ker(\pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v) \rightarrow \text{Aut}_{\text{Sets}}(C_v^N))$ is normalized by $\text{Gal}_{k, v}$, where we use the action $\text{Gal}_{k, v} \ni \sigma \mapsto \psi_\sigma \in \text{Aut}(\pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v))$ or, equivalently by [Lemma 4.12\(d\)](#), conjugation by $s_{\bar{x}_v}(\sigma)$ in $\pi_1^{\text{ét}}(\tilde{X}_v, \bar{x}_v)$.
- (4) There is a surjection $C_v^{N+1} \rightarrow C_v^N$ of $\pi_1^{\text{ét}}(\tilde{X}_v, \bar{x}_v)$ -sets.

We can find sets satisfying the first three conditions by applying [Proposition 4.11](#), and the last condition can be guaranteed by choosing the C_v^N 's inductively (for a given v).

We now proceed to define a subgroup of $\pi_1^{\text{proét}}(X, \bar{x})$ that will lead to the desired $\pi_1^{\text{proét}}(X, \bar{x})$ -set. For that we need to find a suitably large subgroup of elements of U that are well behaved under the Galois action.

Definition 4.20. We will call an element $g \in \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ a *regular loop* (with respect to U) if there exists v, m , a walk P of length m from $v_1 = v_{\bar{x}}$ to $v_{m+1} = v$, étale paths γ, γ' of special form supported on P and P^{-1} , respectively, and $\beta \in \pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v)$ such that:

- $g = \gamma' \circ \beta \circ \gamma$.
- β acts trivially on C_v^m , i.e.,

$$\beta \in \ker(\pi_1^{\text{ét}}(\tilde{X}_{v, \bar{k}}, \bar{x}_v) \rightarrow \text{Aut}(C_v^m)).$$


- There exist presentations $(\gamma_{2m}, \dots, \gamma_1)$ and $(\gamma'_1, \dots, \gamma'_{2m})$ of γ and γ' such that the following condition is satisfied. For any $1 \leq i \leq m$, there is

$$\gamma'_{2i-1} \circ \gamma_{2i-1} \in \ker(\pi_1^{\text{ét}}(\tilde{X}_{v_i, \bar{k}}, \bar{x}_{v_i}) \rightarrow \text{Aut}(C_{v_i}^i))$$

and

$$\gamma_{2i} \circ \gamma'_{2i} \in \ker(\pi_1^{\text{ét}}(\tilde{X}_{v_{i+1}, \bar{k}}, \bar{x}_{v_{i+1}}) \rightarrow \text{Aut}(C_{v_{i+1}}^i)).$$

The following picture might be useful to visualize the definition:



Here, the larger bullets correspond to the \bar{x}_{v_i} and the smaller ones to $\partial_{0 \text{ or } 1}^\epsilon(\bar{x}(e_i))$.

Remark 4.21. We find the definition involving the C_v^N quite convenient. One could, however, avoid introducing the C_v^N and make a slightly different definition. Define $O_v^{N,+}$ to be the set of (isomorphism classes of) Gal_k -conjugates of the $\pi_1^{\acute{e}t}(\tilde{X}_{v,\bar{k}}, \bar{x}_v)$ -sets in O_v^N . Proposition 4.11 then implies that $O_v^{N,+}$ are finite as well. Moreover, for each v , both O_v^N and $O_v^{N,+}$ are increasing with N . We could then require the β and the $(\gamma'_{2i-1} \circ \gamma_{2i-1})$ as above to act trivially on each element of $O_v^{m,+}$ and $O_{v_i}^{i,+}$, correspondingly.

Step III: Defining the desired open subgroup V , checking its properties and finishing the proof. We make the following central definition:

Let $V_0 < \pi_1^{\text{pro}\acute{e}t}(X_{\bar{k}}, \bar{x})$ denote the subgroup generated by the set of regular loops and let V be its topological closure.

Let $G = ((*_v^{\text{top}} \pi_1^{\acute{e}t}(\tilde{X}_{v,\bar{k}}, \bar{x}_v)) *^{\text{top}} \pi_1(\Gamma', T')) / \langle R'_1, R'_2 \rangle^{nc}$ denote the topological group appearing in the van Kampen presentation above. We have that $G^{\text{Noohi}} = \pi_1^{\text{pro}\acute{e}t}(X_{\bar{k}}, \bar{x})$. Let $\tilde{G} \subset \pi_1^{\text{pro}\acute{e}t}(X_{\bar{k}}, \bar{x})$ denote the subgroup of all étale paths (or “loops”, rather) of special form, supported on walks from $v_{\bar{x}}$ to $v_{\bar{x}}$.

Observation 4.22. By Observation 4.17, the map $G \rightarrow \pi_1^{\text{pro}\acute{e}t}(X_{\bar{k}}, \bar{x}) = G^{\text{Noohi}}$ factorizes through \tilde{G} . Directly from the definitions, there is $V_0 < \tilde{G}$. We are thus in the situation of Lemma 2.39. We will use it below.

For brevity, let us denote $\bar{G}_v = \pi_1^{\acute{e}t}(\tilde{X}_{v,\bar{k}}, \bar{x}_v)$ and $G_v = \pi_1^{\acute{e}t}(\tilde{X}_v, \bar{x}_v) \simeq \bar{G}_v \rtimes \text{Gal}_k$ in the proofs below.

Proposition 4.23. The following statements about the subgroup V hold:

- (1) There is a containment $V < U$.
- (2) It is an open subgroup.
- (3) The groups V_0 and V are invariant under the Galois action, i.e., $\psi_\sigma(V_0) = V_0$ and $\psi_\sigma(V) = V$ for all $\sigma \in \text{Gal}_k$.

Proof. (1) As any open subgroup is automatically closed, it is enough to show that any regular loop lies in U . Let g be a regular loop and write $g = \gamma' \circ \beta \circ \gamma$ with γ, γ' étale paths of special form supported on some walk (and its inverse) from $v_{\bar{x}}$ to v of length m , with presentations $(\gamma_1, \dots, \gamma_{2m})$ and $(\gamma'_{2m}, \dots, \gamma'_1)$ of γ and γ' , and $\beta \in \ker(\pi_1^{\acute{e}t}(\tilde{X}_{v,\bar{k}}, \bar{x}_v) \rightarrow C_v^m)$, as in the definition of a regular loop. Let us introduce the following notation (and analogously for γ')

$$\gamma_{i \leftarrow 1} = \gamma_i \circ \dots \circ \gamma_2 \circ \gamma_1.$$

By definition, there is

$$\gamma_{2i \leftarrow 1} \cdot s_0 \in O_{v_i}^i.$$

For $i = m$, it follows from the condition on β that $(\beta \circ \gamma) \cdot s_0 = \gamma \cdot s_0$. Similarly, the condition on $\gamma_{2m} \circ \gamma'_{2m}$ implies that $(\gamma'^{-1}_{2m} \circ \gamma_{2m}) \circ \gamma_{2m \leftarrow 1} \cdot s_0 = \gamma_{2m \leftarrow 1} \cdot s_0$, and thus

$$(\gamma' \circ \beta \circ \gamma) \cdot s_0 = (\gamma' \circ \gamma) \cdot s_0 = ((\gamma'_{1 \leftarrow 2m-1} \circ \gamma'_{2m} \circ \gamma_{2m \leftarrow 1})) \cdot s_0 = \gamma'_{1 \leftarrow 2m-1} \circ \gamma_{2m-1 \leftarrow 1} \cdot s_0.$$

The process continues in a similar fashion to show that g stabilizes s_0 , and thus belongs to U .

(2) By Lemma 2.39, it is enough to check that the map $G \rightarrow \text{Aut}(\tilde{G}/V_0)$ is continuous when \tilde{G}/V_0 is considered with the discrete topology.

Using the universal property of free topological products, continuity can be checked separately for \bar{G}_v and D . For D , this is automatic, as D is discrete. To see the result for \bar{G}_v 's, we need to show that the stabilizers of the action of \bar{G}_v on \tilde{G}/V_0 induced by $\bar{G}_v \rightarrow G$ are open. Fix $[gV_0] \in \tilde{G}/V_0$ and $g \in \tilde{G}$ representing it. The element g is represented by some étale path (or a “loop”, in fact) of special form ρ supported on a walk P_ρ of length $l(P_\rho)$. By Observation 4.17, the morphism $\bar{G}_v \rightarrow \tilde{G} \subset \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is also defined using an étale path of special form γ_v supported on a walk $P_{\bar{x}_v, v}$ in the tree $T \subset \Gamma$. Let $H_v = \ker(\bar{G}_v \rightarrow \text{Aut}(C_v^{l(P_\rho)+l(P_v)})) < \bar{G}_v$. Then H_v is open in \bar{G}_v and its image in \tilde{G} can be written as $\{\gamma_v^{-1} \circ \beta \circ \gamma_v \mid \beta \in H_v\}$. It follows from the setup that for $\beta \in H_v$,

$$g^{-1} \circ \gamma_v^{-1} \circ \beta \circ \gamma_v \circ g \in V_0$$

and so any element in the image of H_v fixes $[gV_0]$ in G/V . Thus, the stabilizer of $[gV_0]$ in \bar{G}_v is also open, as desired.

(3) For each σ , the map ψ_σ is continuous. As $V = \bar{V}_0^{G^{\text{Noohi}}}$, it is thus enough to prove that V_0 is Gal_k -invariant. By Lemma 4.12, it follows that under the action of Gal_k , an étale path of special form supported on a walk P is mapped again to an étale path of special form supported on P . Consequently, checking that the action of $\sigma \in \text{Gal}_k$ maps a regular loop g to another regular loop boils down to checking the following fact. If g has a presentation $g = \gamma' \circ \beta \circ \gamma$ as in the definition of a regular loop, then

- $\psi_\sigma(\beta)$ still acts trivially on $C_v^{l(P)}$;
- $\psi_\sigma(\gamma'_i \circ \gamma_i)$ or $\psi_\sigma(\gamma_i \circ \gamma'_i)$, depending on parity, still acts trivially on $C_v^{[l]_{l+1}}$ for every i .

However, this follows from property (3) in the definition of the C_v^j . □

Denote by S' the quotient $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})/V$ considered as a $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ -set.

Let $\rho_v : \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}_v) \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ be the isomorphism defined using the fixed (étale) path γ_v between \bar{x}_v and \bar{x} , as in Observation 4.17. We have an action given by $\sigma \mapsto \psi_\sigma$ on both $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ and $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}_v)$. We already know that $V < \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is invariant under this action, but this is not necessarily true for $\rho_v^{-1}(V) < \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}_v)$. This holds after a finite base field extension.

Lemma 4.24. For each $v \in \text{Vert}$, define an (abstract) $\text{Gal}_{k,v}$ -action on $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ to be

$$\sigma_v g = \rho_v(\psi_\sigma(\rho_v^{-1}(g))).$$

Then there exists a finite extension l/k , such that for all $v \in \text{Vert}$, there is:

(a) $\text{Gal}_{l,v}$ fixes V .

(b) The obtained induced $\text{Gal}_{l,v}$ -action on S' can be written as

$$\sigma_v \cdot [gV] = [(\gamma_v^{-1} \circ \psi_\sigma(\gamma_v)) \bullet \psi_\sigma(g)V].$$

(c) The induced $\text{Gal}_{l,v}$ action on S' is continuous and compatible with the \overline{G}_v -action.

Proof. As there are finitely many vertices v , it is enough to prove the statements for a single fixed v . Let $g \in V$. By definition of ρ_v , there is

$$\sigma_v g = \gamma_v^{-1} \circ (\psi_\sigma(\gamma_v \circ g \circ \gamma_v^{-1})) \circ \gamma_v = (\gamma_v^{-1} \circ \psi_\sigma(\gamma_v)) \bullet \psi_\sigma(g) \bullet (\psi_\sigma(\gamma_v^{-1}) \circ \gamma_v).$$

By [Proposition 4.23](#), we have $\psi_\sigma(g) \in V$ and we only need to show that $\gamma_v^{-1} \circ \psi_\sigma(\gamma_v) \in V$. By [Lemma 4.18](#) and [Observation 4.17](#), the map $\text{Gal}_k \ni \sigma \mapsto \psi_\sigma(\gamma_v) \in \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}, \bar{x}_v)$ is continuous, and we conclude that for an open subgroup of $\sigma \in \text{Gal}_k$ we have the desired containment.

It follows from the previous point that we get an induced action of $\text{Gal}_{l,v}$ on S' . Using that $\gamma_v^{-1} \circ \psi_\sigma(\gamma_v) \in V$, the alternative formula in the statement follows from the computation

$$\sigma_v \cdot [gV] = [\rho_v(\psi_\sigma(\rho_v^{-1}(g)))V] = [(\gamma_v^{-1} \circ \psi_\sigma(\gamma_v)) \bullet \psi_\sigma(g) \bullet (\psi_\sigma(\gamma_v)^{-1} \circ \gamma_v)V] = [(\gamma_v^{-1} \circ \psi_\sigma(\gamma_v)) \bullet \psi_\sigma(g)V].$$

Let us move to the last point. Compatibility with the \overline{G}_v -action follows from [Lemma 4.12\(d\)](#) and the fact that the map $\overline{G}_v \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is defined by postcomposing with ρ_v . To check continuity, fix $[gV]$. By [Lemma 2.39](#), this class is represented by a path (loop) of special form, and so we can assume this about g . Checking that the stabilizer of $[gV]$ is open boils down to checking that for an open subgroup of the σ in $\text{Gal}_{l,v}$, one has $g^{-1} \bullet (\gamma_v^{-1} \circ \psi_\sigma(\gamma_v)) \bullet \psi_\sigma(g) \in V$. However, this follows from the openness of V and [Lemma 4.18](#). \square

Proposition 4.25. *There is a (continuous) $\pi_1^{\text{proét}}(X_l, \bar{x})$ -action on S' that extends the $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ -action.*

Proof. By the van Kampen theorem for $\pi_1^{\text{proét}}(X_l, \bar{x})$, it is enough to show that there are continuous actions of the $\overline{G}_v \rtimes \text{Gal}_{l,v}$ and D compatible with the \overline{G}_v and D actions that S' is already equipped with, and such that the van Kampen relations are satisfied. We already have a continuous action by D on S' , and by [Lemma 4.24](#), we get an action of $\overline{G}_v \rtimes \text{Gal}_{l,v}$.

Let us now check that the van Kampen relations are preserved. In the case of relation R_2 , this is automatic by [Observation 4.15\(c\)](#). This is because we have left the \overline{G}_v -actions intact. To check that relation R_1 is respected, it suffices to check that $\pi_1^{\text{ét}}(\partial_1)(\sigma)\vec{E} = \vec{E}\pi_1^{\text{ét}}(\partial_0)(\sigma)$ for $\sigma \in \text{Gal}_{l,E}$ and E an edge in Γ with vertices v_-, v_+ . Let $\delta_{W',W} = \gamma_{W',W}$ denote the fixed paths from the van Kampen setup in the computation below to make the distinction from the γ_v clearer. Using [Lemma 4.12\(d\)](#), we compute that

$$\pi_1^{\text{ét}}(\partial_0)(\sigma) = \delta_{v_+,E}^{-1} \circ_{G_{v_+}} \sigma_E \circ_{G_{v_+}} \delta_{v_+,E} = \delta_{v_+,E}^{-1} \circ_{G_{v_+}} \psi_\sigma(\delta_{v_+,E}) \circ_{G_{v_+}} \sigma_{v_+} = (\delta_{v_+,E}^{-1} \circ_{G_{v_+}} \psi_\sigma(\delta_{v_+,E})) \bullet_{G_{v_+}} \sigma_{v_+}.$$

The image of $(\delta_{v_+,E}^{-1} \circ_{G_{v_+}} \psi_\sigma(\delta_{v_+,E}))$ in $\pi_1^{\text{proét}}(X, \bar{x})$ via ρ_{v_+} is $\gamma_{v_+}^{-1} \circ \delta_{v_+,E}^{-1} \circ \psi_\sigma(\delta_{v_+,E}) \circ \gamma_{v_+}$.

By definition, $\vec{E} \in \pi_1^{\text{proét}}(X, \bar{x})$ can be written as $\vec{E} = \gamma_{v-}^{-1} \circ \delta_{v-,E}^{-1} \circ \delta_{v+,E} \circ \gamma_{v+}$. Putting this together and using the formula of [Lemma 4.24](#), we have that $\vec{E} \cdot \pi_1^{\text{ét}}(\partial_0)(\sigma) \cdot [hV]$ equals

$$\begin{aligned} (\gamma_{v-}^{-1} \circ \delta_{v-,E}^{-1} \circ \delta_{v+,E} \circ \gamma_{v+}) \circ (\gamma_{v+}^{-1} \circ \delta_{v+,E}^{-1} \circ \psi_{\sigma}(\delta_{v+,E}) \circ \gamma_{v+}) \cdot [(\gamma_{v+}^{-1} \circ \psi_{\sigma}(\gamma_{v+})) \cdot \psi_{\sigma}(h)V] \\ = [(\gamma_{v-}^{-1} \circ \delta_{v-,E}^{-1} \circ \psi_{\sigma}(\delta_{v+,E} \circ \gamma_{v+})) \cdot \psi_{\sigma}(h)V]. \end{aligned}$$

A similar computation (left to the reader) shows that

$$\pi_1^{\text{ét}}(\partial_1)(\sigma) \cdot \vec{E} \cdot [hV] = [(\gamma_{v-}^{-1} \circ \delta_{v-,E}^{-1} \circ \psi_{\sigma}(\delta_{v+,E} \circ \gamma_{v+})) \cdot \psi_{\sigma}(h)V]$$

as well. This finishes the proof of the proposition. □

End of the proof of Theorem 4.13. We have proven that for a transitive $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ -set S there exists a finite extension l/k and a transitive $\pi_1^{\text{proét}}(X_l, \bar{x})$ -set S' that dominates S as $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ -sets. As explained above, this finishes the proof. □

We have finished our main proof, and thus the most difficult part of the exact sequence is now proven. We now obtain the final form of the fundamental exact sequence.

End of the proof of Theorem 4.14. We already know the statements of the “moreover” part and the near exactness in the middle of the sequence. All we have to prove is that $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is thickly closed in $\pi_1^{\text{proét}}(X, \bar{x})$. As $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$ is a topological embedding of Raïkov complete groups, $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is a closed subgroup of $\pi_1^{\text{proét}}(X, \bar{x})$; see, e.g., [\[Dikranjan 2013, Proposition 6.2.7.\]](#) By [Lemma 2.33](#), the proof will be finished if we show that $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ is normal in $\pi_1^{\text{proét}}(X, \bar{x})$. Observe that checking whether $\overline{\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})} = \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ can be performed after replacing $\pi_1^{\text{proét}}(X, \bar{x})$ by any open subgroup U such that $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) < U <^{\circ} \pi_1^{\text{proét}}(X, \bar{x})$. Choosing a suitably large finite field extension l/k and looking at $U = \pi_1^{\text{proét}}(X_l, \bar{x})$, we are reduced to the situation as in the proof of [Theorem 4.13](#), i.e., we have enough rational points on the connected components we are interested in when applying van Kampen. Let $\tilde{G} < \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$ be the dense subgroup defined above [Proposition 4.23](#). Note that by the van Kampen theorem applied to $\pi_1^{\text{proét}}(X, \bar{x})$ together with the observations in [Observation 4.15](#), it follows that the subgroup generated by \tilde{G} and the $\text{Gal}_{k,v}$ is dense in $\pi_1^{\text{proét}}(X, \bar{x})$. Putting this together, it follows that it is enough to check that, for each v , conjugation by elements of $\text{Gal}_{k,v}$ fixes \tilde{G} in $\pi_1^{\text{proét}}(X, \bar{x})$. This, however, follows from [Lemma 4.12\(c\), \(d\)](#) and the fact that $\text{Gal}_{k,v} \rightarrow \pi_1^{\text{proét}}(X, \bar{x})$ is defined as the composition $\text{Gal}_{k,v} \rightarrow \pi_1^{\text{proét}}(X, \bar{x}_v) \xrightarrow{\rho_v} \pi_1^{\text{proét}}(X, \bar{x})$, where $\rho_v = \gamma_v^{-1} \circ (-) \circ \gamma_v$ with $\gamma_v \in \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}, \bar{x}_v)$ of special form. □

Remark 4.26. Let us revisit the counterexample of [Example 4.5](#) from the point of view of the proof above. We will freely use the notation set there. In this example, we have started from the fixed point s_0 , and used the group elements to reach the point $s_1 = g_m h_{m-1} \cdots h_3 g_2 h_1 \cdot s_0$. We have then concluded that $s_2 = \zeta_{\ell^{m+1}} \cdot s_1 = g \cdot s_1$ and justified that the setup forces that this equality contradicts the possibility of extending the Galois action to the set S . The problem here is caused by the fact that, denoting




Figure 1. Graphical interpretation of S' .

$\gamma = g_m h_{m-1} \cdots h_3 g_2 h_1$, the element

$$\gamma^{-1} \circ g^{-1} \circ \zeta_{\ell^{m+1}} \circ \gamma$$

stabilizes s_0 , but it is not a “regular loop” in the language introduced above. Of course, this only means that this particular “obvious” presentation is not as in the definition of a regular loop. But, by now, we know that it provably cannot be a regular loop with any presentation.

Let us now apply the construction of our main proof in the context of this Example. Let X, \bar{x} and the $\pi_1^{\text{proét}}(\bar{X}, \bar{x})$ -set S be as in Example 4.5. Recall that S decomposes $S = \bigsqcup_{i \geq 1} a_{2i-1}$ (resp. $S = \bigsqcup_{i \geq 1} b_{2i}$) as $\pi_1^{\text{ét}}(\bar{X}_1, \bar{x}_1)$ -set (resp. $\pi_1^{\text{ét}}(\bar{X}_2, \bar{x}_2)$ -set), where we have fixed identifications $\pi_1^{\text{ét}}(\bar{X}_i, \bar{x}_i) \simeq \widehat{\mathbb{Z}}(1)$, $a_j = \mu_{\ell^j}$, $b_j = \mu_{\ell^j}$.

The 2-simplex obtained from the normalization $\tilde{X} = X_1 \sqcup X_2$ of X has two vertices. Let O_1^N, O_2^N denote the corresponding sets of “elements reachable in at most N steps”, as in Definition 4.19. Then one checks that

$$O_1^N = a_1 \cup \cdots \cup a_{2N+1}, \quad O_2^N = b_2 \cup \cdots \cup b_{2N}.$$

Let us denote $a'_i := a_{2i-1} \setminus \{1\}$, $b'_i := b_{2i} \setminus \{1\}$ (using the fixed identifications with the μ_j). Let $\Sigma = \bigcup_{i \geq 1} (a'_i \cup b'_i)$ be the alphabet consisting of all the elements of all the a'_j and b'_j . Let $S' \subset \text{words}(\Sigma)$ be the subset of words on Σ of the following form:

$$S' = \{\emptyset\} \cup \{\beta_m \alpha_{m-1} \cdots \beta_3 \alpha_2 \beta_2 \alpha_1\} \cup \{\alpha_m \beta_m \cdots \beta_3 \alpha_2 \beta_2 \alpha_1\} \cup \{\alpha_m \beta_{m-1} \cdots \alpha_3 \beta_2 \alpha_2 \beta_1\} \cup \{\beta_m \alpha_m \cdots \alpha_2 \beta_2 \alpha_1\},$$

where m runs over $\mathbb{N}_{\geq 1}$ and $\alpha_j \in a'_j$, $\beta_j \in b'_j$ for each j . Geometrically, S' can be thought of as an infinite tree: at the element \emptyset we glue copies of a_1 and b_2 so that $1 \in a_1$ and $1 \in b_2$ are identified at \emptyset . Now, to each element of $a_1 \setminus \{1\}$ we glue a copy of b_2 at $1 \in b_2$ and to each element of the initial $b_2 \setminus \{1\}$, we glue a copy of a_3 at $1 \in a_3$. Now, to each copy of the recently glued b_2 's, we glue a copy of a_3 , and to each copy of previously glued a_3 's, we glue b_4 . The procedure continues; see Figure 1.

There is an obvious action on such a tree by Gal_k , compatible with the $\pi_1^{\text{proét}}(\bar{X}, \bar{x})$ -action; via the description of S' in terms of words on Σ , it corresponds to applying the Gal_k -action to each letter via the identifications with the μ_j . There is moreover a $\pi_1^{\text{proét}}(\bar{X}, \bar{x})$ -equivariant surjective map $S' \rightarrow S$; Indeed, an

element of the form, for example, $\alpha_m \beta_m \cdots \beta_3 \alpha_2 \beta_2 \alpha_1 \in S'$ is mapped to the element $\alpha_m \beta_m \cdots \beta_3 \alpha_2 \beta_2 \alpha_1 \cdot s_0 \in S$, where the multiplication makes sense thanks to how the a_i, b_j grow with i and j . The constructed set S' thus satisfies the desired properties of the set sought in the proof of [Theorem 4.13](#). Up to some minor tweaking, it will correspond to the set obtained by following the proof of the theorem. We will not, however, try to give a precise proof of that last claim here.

While this “tree construction” example is much more enlightening in the simple cases of schemes glued at one point, it proved to be rather difficult to turn this intuition into a formal proof that would work for arbitrary schemes (i.e., where the normalization might no longer have such a pleasant form). For that reason, we have opted for a proof that is less geometric in nature.

Remark 4.27. We sketch a slightly different approach to the central part of the main proof. It is a bit quicker, but less constructive, i.e., does not “explicitly” construct the desired Galois invariant open subgroup in terms of regular loops. We will freely use the fact that a surjective map from a compact space onto a Hausdorff space is a quotient map.

Assume that we have already done the preparatory steps of the main proof, i.e., we have increased the base field to have many rational points and applied the van Kampen theorem. We want to prove that the action

$$\text{Gal}_k \times \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$$

given by ψ_σ is continuous. Let G, \tilde{G} be as introduced above [Observation 4.22](#).

Firstly, one checks that any element of \tilde{G} , so a path of special form, can be in fact rewritten with a presentation that makes it visibly an image of an element of G , at the expense of the presentation possibly getting longer. Another words, the map $G \rightarrow \tilde{G}$ is surjective. By default, \tilde{G} is considered with the subspace topology from $\pi_1^{\text{proét}}(X_{\bar{k}}, \bar{x})$. Let us denote (\tilde{G}, quot) the same group but considered with the quotient topology from G . We thus have a continuous bijection $(\tilde{G}, \text{quot}) \rightarrow \tilde{G}$.

The group G is a topological quotient of the free topological product of finitely many compact groups G_v and a finitely generated free group $D \simeq \mathbb{Z}^{*r}$. One checks from the universal properties that this free product can be written as a quotient of the free topological group $F(Z)$ (see [\[Arhangel'skii and Tkachenko 2008, Chapter 7\]](#)) on a compact space of generators $Z = \bigsqcup_v G_v \bigsqcup_{\{1, \dots, r\}} *$, i.e., the disjoint union of the G_v and r singletons.

By [\[Arhangel'skii and Tkachenko 2008, Theorem 7.4.1\]](#), $F(Z)$ is, as a topological space, a colimit of an increasing union $\cdots \subset B_n \subset B_{n+1} \subset \cdots$ of compact subspaces. These spaces are explicitly described as words of bounded length in $F(Z)$ (this makes sense, as the underlying group of $F(Z)$ is the abstract free group on Z). From this, it follows that (as a topological space) $(\tilde{G}, \text{quot}) = \text{colim } K_n$, with $K_n = \text{im}(B_n)$.

Working directly with the K_n is inconvenient for our purposes, as these sets are not necessarily preserved by the Galois action. The reason is that the van Kampen presentation as a quotient of a free product uses fixed paths, while applying Galois action will usually move the paths. One then has to conjugate by a suitable element to “return” to the paths fixed in van Kampen, possibly increasing the length of the word.

Instead, we can consider subsets $K'_n \subset \tilde{G}$ of elements that are paths of special form of length $\leq n$, i.e., possessing a presentation as a path of special form of length $\leq n$ (see [Definition 4.16](#)). By a reasonably simple combinatorics, one can cook up “brute force” bounds $f(n, d), g(n, d) \in \mathbb{N}$ in terms of n and the diameter $d = \text{diam}(\Gamma)$ of Γ such that there is

$$K_n \subset K'_{f(n,d)} \quad \text{and} \quad K'_n \subset K_{g(n,d)}.$$

In conclusion, $(\tilde{G}, \text{quot}) = \text{colim } K'_n$ in Top.

By [Lemma 4.12](#), the Gal_k -action preserves the sets K'_n and $\text{Gal}_k \times K'_n \rightarrow K'_n$ is continuous. As Gal_k is compact, $\text{Gal}_k \times (-)$ has a right adjoint $\text{Maps}_{\text{cts}}(\text{Gal}_k, -)$ in Top and so $\text{Gal}_k \times (\text{colim}_{n \in \mathbb{N}} K'_n) = \text{colim}_{n \in \mathbb{N}} (\text{Gal}_k \times K'_n)$. From this, we immediately get that $\text{Gal}_k \times (\tilde{G}, \text{quot}) \rightarrow (\tilde{G}, \text{quot})$ is continuous. As Gal_k -action respects the group action of \tilde{G} , it quickly follows that the action is still continuous when (\tilde{G}, quot) is equipped with the weakened topology τ making open subgroups a base at 1, as in [Lemma 2.25](#). By (the easier part of) [Lemma 2.39](#), this weakened topology on (\tilde{G}, quot) matches that of \tilde{G} . It follows that $\text{Gal}_k \times \tilde{G} \rightarrow \tilde{G}$ is continuous.

By [Lemma 2.25](#) again, one has to check that the continuity is not lost when passing to the Raïkov completion of the maximal Hausdorff quotient of (G, τ) . This in turn can be justified by similar arguments as in the proof of [Lemma 2.39](#). This finishes the sketch. See also [[Bhatt and Scholze 2015](#), Proposition 4.3.3].

Acknowledgements

The main ideas and results contained in this article are a part of my PhD thesis. I express my gratitude to my advisor H el ene Esnault for introducing me to the topic and her constant encouragement. I would like to thank my coadvisor Vasudevan Srinivas for his support and suggestions. I am thankful to Peter Scholze for explaining some parts of his work to me via e-mail. I thank Jo ao Pedro dos Santos for his comments and feedback. I owe special thanks to Fabio Tonini, Lei Zhang and Marco D’Addezio from our group in Berlin for many inspiring mathematical discussions. I thank Piotr Achinger and Jakob Stix for their support. I would also like to thank the referee for careful reading, valuable remarks and urging me to write a more streamlined version of the main proof.

My PhD was funded by the Einstein Foundation. This work is a part of the project “Kapibara” supported by the funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 802787).

The major revision was prepared at JU Krak ow and GU Frankfurt. I was supported by the Priority Research Area SciMat under the program Excellence Initiative — Research University at the Jagiellonian University in Krak ow. This research was also funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) TRR 326 Geometry and Arithmetic of Uniformized Structures, project number 444845124.

References

- [Achinger et al. 2022] P. Achinger, M. Lara, and A. Youcis, “Specialization for the pro-étale fundamental group”, *Compositio Math.* **158**:8 (2022), 1713–1745. [MR](#) [Zbl](#)
- [Arhangel’skii and Tkachenko 2008] A. Arhangel’skii and M. Tkachenko, *Topological groups and related structures: an introduction to topological algebra*, Atlantis Studies in Mathematics **1**, Atlantis, Paris, 2008. [MR](#)
- [Bhatt and Scholze 2015] B. Bhatt and P. Scholze, “The pro-étale topology for schemes”, pp. 99–201 in *De la géométrie algébrique aux formes automorphes, I*, edited by J.-B. Bost et al., Astérisque **369**, Soc. Math. France, Paris, 2015. [MR](#) [Zbl](#)
- [Bourbaki 1966] N. Bourbaki, *Elements of mathematics: general topology*, part 2, Hermann, Paris, 1966. [MR](#) [Zbl](#)
- [Çakar 2014] A. C. Çakar, “A category theoretical approach to classification of covering spaces”, senior project II, Bilkent University, 2014.
- [Dikranjan 2013] D. Dikranjan, “Introduction to topological groups”, preprint, Università degli Studi di Udine, 2013, <http://users.dimi.uniud.it/~dikran.dikranjan/ITG.pdf>.
- [Fulton 1995] W. Fulton, *Algebraic topology: a first course*, Graduate Texts in Mathematics **153**, Springer, 1995. [MR](#) [Zbl](#)
- [Graev 1948] M. I. Graev, “Free topological groups”, *Izv. Akad. Nauk SSSR Ser. Mat.* **12**:3 (1948), 279–324. In Russian; translated in *Amer. Math. Soc. Transl.* **8**:1 (1962), 305–364. [MR](#)
- [Grothendieck 1960] A. Grothendieck, “Technique de descente et théorèmes d’existence en géométrie algébriques, II: Le théorème d’existence en théorie formelle des modules”, exposé 195 in *Séminaire Bourbaki 1959/1960*, W. A. Benjamin, Amsterdam, 1960. Reprinted as pp. 369–390 in *Séminaire Bourbaki 5*, Soc. Math. France, Paris, 1995. [MR](#) [Zbl](#)
- [Hatcher 2002] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002. [MR](#) [Zbl](#)
- [de Jong 1995] A. J. de Jong, “Étale fundamental groups of non-archimedean analytic spaces”, *Compositio Math.* **97**:1-2 (1995), 89–118. [MR](#) [Zbl](#)
- [Lara 2022] M. Lara, “Homotopy exact sequence for the pro-étale fundamental group”, *Int. Math. Res. Not.* **2022**:19 (2022), 15355–15389. [MR](#) [Zbl](#)
- [Lavanda 2018] E. Lavanda, “Specialization map between stratified bundles and pro-étale fundamental group”, *Adv. Math.* **335** (2018), 27–59. [MR](#) [Zbl](#)
- [Lepage 2010] E. Lepage, *Géométrie anabélienne tempérée*, Ph.D. thesis, Université Paris-Diderot, 2010. [arXiv 1004.2150](https://arxiv.org/abs/1004.2150)
- [Noochi 2008] B. Noochi, “Fundamental groups of topological stacks with the slice property”, *Algebr. Geom. Topol.* **8**:3 (2008), 1333–1370. [MR](#) [Zbl](#)
- [Rydh 2010] D. Rydh, “Submersions and effective descent of étale morphisms”, *Bull. Soc. Math. France* **138**:2 (2010), 181–230. [MR](#) [Zbl](#)
- [Schwede 2005] K. Schwede, “Gluing schemes and a scheme without closed points”, pp. 157–172 in *Recent progress in arithmetic and algebraic geometry* (Knoxville, TN, 2002), *Contemp. Math.* **386**, Amer. Math. Soc., Providence, RI, 2005. [MR](#) [Zbl](#)
- [SGA 1 1971] A. Grothendieck, *Revêtements étales et groupe fondamental* (Séminaire de Géométrie Algébrique du Bois Marie 1960–1961), *Lecture Notes in Math.* **224**, Springer, 1971. [MR](#) [Zbl](#)
- [SGA 3 II 1970] M. Demazure and A. Grothendieck, *Schémas en groupes, Tome II: Groupes de type multiplicatif, et structure des schémas en groupes généraux, Exposés VIII–XVIII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), *Lecture Notes in Math.* **152**, Springer, 1970. [MR](#) [Zbl](#)
- [Stacks 2020] “The Stacks project”, electronic reference, 2020, <http://stacks.math.columbia.edu>.
- [Stix 2006] J. Stix, “A general Seifert–Van Kampen theorem for algebraic fundamental groups”, *Publ. Res. Inst. Math. Sci.* **42**:3 (2006), 763–786. [MR](#) [Zbl](#)
- [Stix 2013] J. Stix, *Rational points and arithmetic of fundamental groups: evidence for the section conjecture*, *Lecture Notes in Math.* **2054**, Springer, 2013. [MR](#) [Zbl](#)

Communicated by Bhargav Bhatt

Received 2020-01-27 Revised 2023-02-13 Accepted 2023-05-13

lara@math.uni-frankfurt.de, marcin.lara@mailbox.org

Institute of Mathematics, Goethe University Frankfurt, Frankfurt, Germany

*Institute of Mathematics, Faculty of Mathematics and Computer Science,
Jagiellonian University, Kraków, Poland*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Jonathan Pila	University of Oxford, UK
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Bjorn Poonen	Massachusetts Institute of Technology, USA
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	Rutgers University, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Melanie Matchett Wood	Harvard University, USA
János Kollár	Princeton University, USA	Shou-Wu Zhang	Princeton University, USA
Michael J. Larsen	Indiana University Bloomington, USA		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2024 is US \$525/year for the electronic version, and \$770/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 18 No. 4 2024

Fundamental exact sequence for the pro-étale fundamental group	631
MARCIN LARA	
Infinitesimal dilogarithm on curves over truncated polynomial rings	685
SINAN ÜNVER	
Wide moments of L -functions I: Twists by class group characters of imaginary quadratic fields	735
ASBJØRN CHRISTIAN NORDENTOFT	
On Ozaki's theorem realizing prescribed p -groups as p -class tower groups	771
FARSHID HAJIR, CHRISTIAN MAIRE and RAVI RAMAKRISHNA	
Supersolvable descent for rational points	787
YONATAN HARPAZ and OLIVIER WITTENBERG	
On Kato and Kuzumaki's properties for the Milnor K_2 of function fields of p -adic curves	815
DIEGO IZQUIERDO and GIANCARLO LUCCHINI ARTECHE	