

ANALYSIS & PDE

Volume 1

No. 2

2008

**THE MASS-CRITICAL NONLINEAR SCHRÖDINGER EQUATION
WITH RADIAL DATA IN DIMENSIONS THREE AND HIGHER**

ROWAN KILLIP, MONICA VISAN AND XIAOYI ZHANG



mathematical sciences publishers

THE MASS-CRITICAL NONLINEAR SCHRÖDINGER EQUATION WITH RADIAL DATA IN DIMENSIONS THREE AND HIGHER

ROWAN KILLIP, MONICA VISAN AND XIAOYI ZHANG

We establish global well-posedness and scattering for solutions to the mass-critical nonlinear Schrödinger equation $iu_t + \Delta u = \pm |u|^{4/d}u$ for large spherically symmetric $L^2_x(\mathbb{R}^d)$ initial data in dimensions $d \geq 3$. In the focusing case we require that the mass is strictly less than that of the ground state. As a consequence, we obtain that in the focusing case, any spherically symmetric blowup solution must concentrate at least the mass of the ground state at the blowup time.

1. Introduction

The d -dimensional mass-critical nonlinear Schrödinger equation is given by

$$iu_t + \Delta u = F(u) \quad \text{with } F(u) := \mu |u|^{\frac{4}{d}} u \tag{1-1}$$

where u is a complex-valued function of spacetime $\mathbb{R} \times \mathbb{R}^d$. Here $\mu = \pm 1$, with $\mu = 1$ known as the defocusing equation and $\mu = -1$ as the focusing equation.

The name “mass-critical” refers to the fact that the scaling symmetry

$$u(t, x) \mapsto u_\lambda(t, x) := \lambda^{-\frac{d}{2}} u(\lambda^{-2}t, \lambda^{-1}x)$$

leaves both the equation and the mass invariant. The mass of a solution is

$$M(u(t)) := \int_{\mathbb{R}^d} |u(t, x)|^2 dx$$

and is conserved under the flow.

In this paper, we investigate the Cauchy problem for (1-1) for spherically symmetric $L^2_x(\mathbb{R}^d)$ initial data in dimensions $d \geq 3$ by adapting the recent argument from [Killip et al. 2007], which treated the case $d = 2$. Before describing our results, we need to review some background material. We begin by making the notion of a solution more precise:

MSC2000: 35Q55.

Keywords: Nonlinear Schrödinger equation, mass-critical, focusing.

Killip was supported by NSF grants DMS-0701085 and DMS-0401277 and by a Sloan Foundation Fellowship. He is also grateful to the Institute for Advanced Study (Princeton) for its hospitality. Zhang was supported by NSF-10601060 and Project 973 in China. Visan was supported under NSF grant DMS-0111298. Any opinions, findings and conclusions or recommendations expressed are those of the authors and do not reflect the views of the National Science Foundation.

Definition 1.1 (Solution). A function $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ on a nonempty time interval $I \subset \mathbb{R}$ is a *solution* (more precisely, a strong $L_x^2(\mathbb{R}^d)$ solution) to (1-1) if it lies in the class

$$C_t^0 L_x^2(K \times \mathbb{R}^d) \cap L_{t,x}^{2(d+2)/d}(K \times \mathbb{R}^d)$$

for all compact $K \subset I$, and obeys the Duhamel formula

$$u(t_1) = e^{i(t_1-t_0)\Delta} u(t_0) - i \int_{t_0}^{t_1} e^{i(t_1-t)\Delta} F(u(t)) dt \tag{1-2}$$

for all $t_0, t_1 \in I$. Note that by Lemma 2.7 below, the condition $u \in L_{t,x}^{2(d+2)/d}$ locally in time guarantees that the integral converges, at least in a weak- L_x^2 sense.

Remark. The condition that u is in $L_{t,x}^{2(d+2)/d}$ locally in time is natural. This space appears in the Strichartz inequality (Lemma 2.7); consequently, all solutions to the linear problem lie in this space. Existence of solutions to (1-1) in this space is guaranteed by the local theory discussed below; it is also necessary in order to ensure uniqueness of solutions in this local theory. Solutions to (1-1) in this class have been intensively studied; see for example [Bégout and Vargas 2007; Bourgain 1998; Carles and Keraani 2007; Cazenave and Weissler 1989; Cazenave 2003; Keraani 2006; Merle and Vega 1998; Tao 2006; Tao et al. 2006; 2007; Tsutsumi 1985].

Associated to this notion of solution is a corresponding notion of blowup. As we will see in Theorem 1.3 below, this precisely corresponds to the impossibility of continuing the solution.

Definition 1.2 (Blowup). We say that a solution u to (1-1) *blows up forward in time* if there exists a time $t_0 \in I$ such that

$$\int_{t_0}^{\sup I} \int_{\mathbb{R}^d} |u(t, x)|^{2(d+2)/d} dx dt = \infty$$

and that u *blows up backward in time* if there exists a time $t_0 \in I$ such that

$$\int_{\inf I}^{t_0} \int_{\mathbb{R}^d} |u(t, x)|^{2(d+2)/d} dx dt = \infty.$$

The local theory for (1-1) was worked out by Cazenave and Weissler [1989]. They constructed local-in-time solutions for arbitrary initial data in $L_x^2(\mathbb{R}^d)$; however, due to the critical nature of the equation, the resulting time of existence depends on the profile of the initial data and not merely on its L_x^2 -norm. Cazenave and Weissler also constructed global solutions for small initial data. We summarize their results in the theorem below.

Theorem 1.3 (Local well-posedness [Cazenave and Weissler 1989; Cazenave 2003]). *Given $t_0 \in \mathbb{R}$ and $u_0 \in L_x^2(\mathbb{R}^d)$, there exists a unique maximal-lifespan solution u to (1-1) with $u(t_0) = u_0$. We will write I for the maximal lifespan. This solution also has the following properties:*

- (Local existence) I is an open neighbourhood of t_0 .
- (Mass conservation) The solution u obeys mass conservation: $M(u(t)) = M(u_0)$ for all $t \in I$.
- (Blowup criterion) If $\sup(I)$ or $\inf(I)$ is finite, then u blows up in the corresponding time direction.

- (Continuous dependence) *The map that takes initial data to the corresponding strong solution is uniformly continuous on compact time intervals for bounded sets of initial data.*
- (Scattering) *If $\sup(I) = +\infty$ and u does not blow up forward in time, then u scatters forward in time, that is, there exists a unique $u_+ \in L_x^2(\mathbb{R}^d)$ such that*

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u_+\|_{L_x^2(\mathbb{R}^d)} = 0.$$

Similarly, if $\inf(I) = -\infty$ and u does not blow up backward in time, then u scatters backward in time, that is, there is a unique $u_- \in L_x^2(\mathbb{R}^d)$ such that

$$\lim_{t \rightarrow -\infty} \|u(t) - e^{it\Delta} u_-\|_{L_x^2(\mathbb{R}^d)} = 0.$$

- (Small data global existence) *If $M(u_0)$ is sufficiently small depending on d , then u is a global solution with finite $L_{t,x}^{2(d+2)/d}$ norm.*

It is widely believed that in the defocusing case, all L_x^2 initial data lead to a global solution with finite $L_{t,x}^{2(d+2)/d}$ spacetime norm (and hence also scattering).

In the focusing case, the general consensus is more subtle. Let Q denote the *ground state*, that is, the unique positive radial solution to

$$\Delta Q + Q^{1+4/d} = Q.$$

(The existence and uniqueness of Q were established in [Berestycki and Lions 1979] and [Kwong 1989] respectively.) Then

$$u(t, x) := e^{it\Delta} Q(x)$$

is a solution to (1-1), which is global but blows up both forward and backward in time (in the sense of Definition 1.2). More dramatically, by applying the pseudoconformal transformation to u , we obtain a solution

$$v(t, x) := |t|^{-d/2} e^{i\frac{|x|^2-4}{4t}} Q\left(\frac{x}{t}\right)$$

with the same mass that blows up in finite time. It is widely believed that this ground state example is the minimal-mass obstruction to global well-posedness and scattering in the focusing case.

To summarize, we subscribe to:

Conjecture 1.4 (Global existence and scattering). *Let $d \geq 1$ and $\mu = \pm 1$. In the defocusing case $\mu = +1$, all maximal-lifespan solutions to (1-1) are global and do not blow up either forward or backward in time. In the focusing case $\mu = -1$, all maximal-lifespan solutions u to (1-1) with $M(u) < M(Q)$ are global and do not blow up either forward or backward in time.*

Remark. While this conjecture is phrased for $L_x^2(\mathbb{R}^d)$ solutions, it is equivalent to a scattering claim for smooth solutions; see [Bégout and Vargas 2007; Carles 2002; Keraani 2006; Tao 2006]. In [Blue and Colliander 2006; Tao 2006], it is also shown that the global existence and the scattering claims are equivalent in the $L_x^2(\mathbb{R}^d)$ category.

The contribution of this paper toward settling this conjecture is:

Theorem 1.5. *Let $d \geq 3$. Then Conjecture 1.4 is true in the class of spherically symmetric initial data (for either choice of sign μ).*

Conjecture 1.4 has been the focus of much intensive study and several partial results for various choices of d, μ , and sometimes with the additional assumption of spherical symmetry. The most compelling evidence in favour of this conjecture stems from results obtained under the assumption that u_0 has additional regularity. For the defocusing equation, it is easy to prove global well-posedness for initial data in H_x^1 ; this follows from the usual contraction mapping argument combined with the conservation of mass and energy; see, for example, [Cazenave 2003]. Recall that the energy is given by

$$E(u(t)) := \int_{\mathbb{R}^d} \left(\frac{1}{2} |\nabla u(t, x)|^2 + \mu \frac{d}{2(d+2)} |u(t, x)|^{\frac{2(d+2)}{d}} \right) dx. \tag{1-3}$$

Note that for general L_x^2 initial data, the energy need not be finite.

The focusing equation with data in H_x^1 was treated by Weinstein. A key ingredient was his proof of the following result:

Theorem 1.6 (Sharp Gagliardo–Nirenberg inequality [Weinstein 1983]).

$$\int_{\mathbb{R}^d} |f(x)|^{\frac{2(d+2)}{d}} dx \leq \frac{d+2}{d} \left(\frac{\|f\|_{L^2}^2}{\|Q\|_{L^2}^2} \right)^{\frac{2}{d}} \int_{\mathbb{R}^d} |\nabla f(x)|^2 dx.$$

As noticed by Weinstein, this inequality implies that the energy (1-3) is positive once $M(u_0) < M(Q)$; indeed, it gives an upper bound on the \dot{H}_x^1 -norm of the solution at all times of existence. Combining this with a contraction mapping argument and the conservation of mass and energy, Weinstein proved global well-posedness for the focusing equation with initial data in H_x^1 and mass smaller than that of the ground state.

Note that the iterative procedure used to obtain a global solution both for the defocusing and the focusing equations with initial data in H_x^1 does not yield finite spacetime norms; in particular, scattering does not follow even for more regular initial data.

In dimensions one and two, there has been much work [Bourgain 1998; Colliander et al. 2002; 2008b; Colliander et al. 2005; Colliander et al. 2007; De Silva et al. 2007a; Fang and Grillakis 2007; Tzirakis 2005] devoted to lowering the regularity of the initial data from H_x^1 toward $L_x^2(\mathbb{R}^d)$ and thus toward establishing the conjecture. For analogous results in higher dimensions, see [De Silva et al. 2007b; Visan and Zhang 2007].

In the case of spherically symmetric solutions, Conjecture 1.4 was recently settled in the high-dimensional defocusing case $\mu = +1, d \geq 3$ in [Tao et al. 2007]; thus, only the $\mu = -1$ case of Theorem 1.5 is new. However, the techniques used in that reference do not seem to be applicable to the focusing problem, primarily because the Morawetz inequality is no longer coercive in that case. Instead, our argument is based on the recent preprint by Killip, Tao and Visan [Killip et al. 2007], which resolved the conjecture for $\mu = \pm 1, d = 2$, and spherically symmetric data. That work, in turn, used techniques developed to treat the analogous conjecture for the energy-critical problem, such as [Bourgain 1999b; Colliander et al. 2008a; Ryckman and Visan 2007; Tao 2005; Visan 2006; 2007] and particularly [Kenig and Merle 2006a]. We will give a more thorough discussion of the relation of the current work to these predecessors later, when we outline the argument.

Mass concentration in the focusing problem. Neither Theorem 1.5 nor Conjecture 1.4 addresses the focusing problem for masses greater than or equal to that of the ground state. In this case, blowup

solutions exist and attention has been focused on describing their properties. For instance, finite-time blowup solutions with finite energy and mass equal to that of the ground state have been completely characterized by Merle [1993]; they are precisely the ground state solution up to symmetries of the equation.

Several works have shown that finite-time blowup solutions must concentrate a positive amount of mass around the blowup time T^* . For finite energy data, see [Merle and Tsutsumi 1990; Nawa 1999; Weinstein 1989] where it is shown that there exists $x(t) \in \mathbb{R}^d$ so that

$$\liminf_{t \nearrow T^*} \int_{|x-x(t)| \leq R} |u(t, x)|^2 dx \geq M(Q)$$

for any $R > 0$. For merely $L^2_x(\mathbb{R}^2)$ initial data, Bourgain [1998] proved that some small amount of mass must concentrate in parabolic windows (at least along a subsequence):

$$\limsup_{t \nearrow T^*} \sup_{x_0 \in \mathbb{R}^2} \int_{|x-x_0| \leq (T^*-t)^{1/2}} |u(t, x)|^2 dx \geq c,$$

where c is a small constant depending on the mass of u . This result was extended to other dimensions in [Bégout and Vargas 2007; Keraani 2006].

Combining Theorem 1.5 with the argument in [Killip et al. 2007, §10], one obtains the following concentration result.

Corollary 1.7 (Blowup solutions concentrate the mass of the ground state). *Let $d \geq 3$ and $\mu = -1$. Let u be a spherically symmetric solution to (1-1) that blows up at time $0 < T^* \leq \infty$. If $T^* < \infty$, there exists a sequence $t_n \nearrow T^*$ such that for any sequence $R_n \in (0, \infty)$ obeying $(T^* - t_n)^{-1/2} R_n \rightarrow \infty$,*

$$\limsup_{n \rightarrow \infty} \int_{|x| \leq R_n} |u(t_n, x)|^2 dx \geq M(Q).$$

If $T^ = \infty$, there exists a sequence $t_n \rightarrow \infty$ such that for any sequence R_n with $t_n^{-1/2} R_n \rightarrow \infty$ in $(0, \infty)$*

$$\limsup_{n \rightarrow \infty} \int_{|x| \leq R_n} |u(t_n, x)|^2 dx \geq M(Q).$$

The analogous statement holds in the negative time direction.

Outline of the proof. Beginning with Bourgain’s seminal work [1999b] on the energy-critical NLS, it has become apparent that in order to prove spacetime bounds for general solutions, it is sufficient to treat a special class of solutions, namely, those that are simultaneously localized in both frequency and space. For further developments, see [Colliander et al. 2008a; Ryckman and Visan 2007; Tao 2005; Visan 2006; 2007].

A new and much more efficient alternative to Bourgain’s induction on mass (or energy) method has recently been developed. It uses a (concentration) compactness technique to isolate minimal-mass/energy blowup solutions as opposed to the almost-blowup solutions of the induction method. Building on earlier developments in [Bégout and Vargas 2007; Bourgain 1998; Keraani 2001; 2006; Merle and Vega 1998], Kenig and Merle [2006a] introduced this method to treat the energy-critical focusing problem with radial data in dimensions three, four, and five; see also [Kenig and Merle 2006b; Killip et al. 2007; Killip and

Visan 2008; Tao 2008a; 2008b; 2008c; Tao et al. 2007] for subsequent applications/developments of this method.

To explain what the concentration compactness argument gives in our context, we need to introduce the following important notion:

Definition 1.8 (Almost periodicity modulo scaling). Given $d \geq 1$ and $\mu = \pm 1$, a solution u with lifespan I is said to be *almost periodic modulo scaling* if there exists a (possibly discontinuous) function $N : I \rightarrow \mathbb{R}^+$ and a function $C : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that

$$\int_{|x| \geq C(\eta)/N(t)} |u(t, x)|^2 dx \leq \eta \quad \text{and} \quad \int_{|\xi| \geq C(\eta)N(t)} |\hat{u}(t, \xi)|^2 d\xi \leq \eta$$

for all $t \in I$ and $\eta > 0$. We refer to the function N as the *frequency scale function* and to C as the *compactness modulus function*.

Remarks. (1) The parameter $N(t)$ measures the frequency scale of the solution at time t , and $1/N(t)$ measures the spatial scale; see [Tao et al. 2006; 2007] for further discussion. We have the freedom to modify $N(t)$ by any bounded function of t , provided that we also modify the compactness modulus function C accordingly. In particular, one could restrict $N(t)$ to be a power of 2 if one wished, although we will not do so here. Alternatively, the fact that the solution trajectory $t \mapsto u(t)$ is continuous in $L^2_x(\mathbb{R}^d)$ can be used to show that the function N may be chosen to depend continuously on t .

(2) By the Ascoli–Arzelà Theorem, a family of functions is precompact in $L^2_x(\mathbb{R}^d)$ if and only if it is norm-bounded and there exists a compactness modulus function C so that

$$\int_{|x| \geq C(\eta)} |f(x)|^2 dx + \int_{|\xi| \geq C(\eta)} |\hat{f}(\xi)|^2 d\xi \leq \eta$$

for all functions f in the family. Thus, an equivalent formulation of Definition 1.8 is as follows: u is almost periodic modulo scaling if and only if

$$\{u(t) : t \in I\} \subseteq \{\lambda^{-d/2} f(x/\lambda) : \lambda \in (0, \infty) \text{ and } f \in K\}.$$

for some compact subset K of $L^2_x(\mathbb{R}^d)$.

In [Tao et al. 2006, Theorems 1.13 and 7.2] the following result was established (see also [Bégout and Vargas 2007; Keraani 2006]), showing that any failure of Conjecture 1.4 must be “caused” by a very special type of solution. For simplicity we state it only in the spherically symmetric case.

Theorem 1.9 (Reduction to almost periodic solutions). *Fix μ and $d \geq 2$. Suppose that Conjecture 1.4 fails for spherically symmetric data. Then there exists a spherically symmetric maximal-lifespan solution u which is almost periodic modulo scaling and which blows up both forward and backward in time, and in the focusing case we also have $M(u) < M(Q)$.*

In [Killip et al. 2007], this result was further refined so as to identify three specific enemies. Once again, we state it only in the spherically symmetric case.

Theorem 1.10 (Three special scenarios for blowup [Killip et al. 2007]). *Fix μ and $d \geq 2$ and suppose that Conjecture 1.4 fails for spherically symmetric data. Then there exists a spherically symmetric maximal-lifespan solution u which is almost periodic modulo scaling, blows up both forward and backward in*

time, and in the focusing case also obeys $M(u) < M(Q)$. Moreover, the solution u may be chosen to match one of the following three scenarios:

- (Soliton-like solution) We have $I = \mathbb{R}$ and $N(t) = 1$ for all $t \in \mathbb{R}$ (thus the solution stays in a bounded space/frequency range for all time).
- (Double high-to-low frequency cascade) We have $I = \mathbb{R}$, $\liminf_{t \rightarrow -\infty} N(t) = \liminf_{t \rightarrow +\infty} N(t) = 0$, and $\sup_{t \in \mathbb{R}} N(t) < \infty$ for all $t \in I$.
- (Self-similar solution) We have $I = (0, +\infty)$ and

$$N(t) = t^{-1/2} \tag{1-4}$$

for all $t \in I$.

In light of this result, the proof of Theorem 1.5 is reduced to showing that none of these three scenarios can occur. In doing this, we follow the model set forth in [Killip et al. 2007]. In all cases, the key step is to prove that u has additional regularity. Indeed, to treat the first two scenarios, we need more than one derivative in L_x^2 ; for the self-similar scenario, H_x^1 suffices. The possibility of showing such additional regularity stems from the fact that u is both frequency and space localized; this in turn is an expression of the fact that u has minimal mass among all blowup solutions.

A further manifestation of this minimality is the absence of a scattered wave at the endpoints of the lifespan I . More formally:

Lemma 1.11 [Tao et al. 2006, Section 6]. *Let u be a solution to (1-1) which is almost periodic modulo scaling on its maximal-lifespan I . Then, for all $t \in I$,*

$$u(t) = \lim_{T \nearrow \sup I} i \int_t^T e^{i(t-t')\Delta} F(u(t')) dt' = - \lim_{T \searrow \inf I} i \int_T^t e^{i(t-t')\Delta} F(u(t')) dt', \tag{1-5}$$

as weak limits in L_x^2 .

Another important property of solutions that are almost periodic modulo scaling is that the behaviour of the spacetime norm is governed by that of $N(t)$. More precisely:

Lemma 1.12 (Spacetime bound [Killip et al. 2007]). *Let u be a nonzero solution to (1-1) with lifespan I , which is almost periodic modulo scaling with frequency scale function $N : I \rightarrow \mathbb{R}^+$. If J is any subinterval of I , then*

$$\int_J N(t)^2 dt \lesssim_u \int_J \int_{\mathbb{R}^d} |u(t, x)|^{\frac{2(d+2)}{d}} dx dt \lesssim_u 1 + \int_J N(t)^2 dt.$$

The nonexistence of self-similar solutions is proved in Section 3. We first prove that any such solution would belong to $C_t^0 H_x^1$ and then observe that H_x^1 solutions are global (see the discussion after Theorem 1.5), while self-similar solutions are not.

For the remaining two cases, higher regularity is proved in Section 5. In order to best take advantage of Lemma 1.11, we exploit a decomposition of spherically symmetric functions into incoming and outgoing waves; this is discussed in Section 4.

In Section 6, we use the additional regularity together with the conservation of energy to preclude the double high-to-low frequency cascade. In Section 7, we disprove the existence of soliton-like solutions using a truncated virial identity in much the same manner as [Kenig and Merle 2006a].

As noted earlier, the argument just described is closely modelled on [Killip et al. 2007], which treated the same equation in two dimensions. The main obstacle in extending that work to higher dimensions is the fractional power appearing in the nonlinearity. This problem presents itself when we prove additional regularity, which is already the most demanding part of [Killip et al. 2007]. Additional regularity is proved via a bootstrap argument using Duhamel’s formula. However, fractional powers can downgrade regularity (a fractional power of a smooth function need not be smooth); in particular, they preclude the simple Littlewood–Paley arithmetic that is usually used in the case of polynomial nonlinearities.

The remedy is twofold: first we use fractional chain rules (see Lemmas 2.3 and 2.4) that allow us to take more than one derivative of a nonlinearity that is merely $C^{1+4/d}$ in u . Secondly, we push through the resulting complexities in the bootstrap argument. An important role is played by Lemma 2.1 (a Gronwall-type result), which we use to untangle the intricate relationship between frequencies in u and those in $|u|^{4/d}u$.

2. Notation and linear estimates

This section contains the basic linear estimates we use repeatedly in the paper.

Some notation. We use $X \lesssim Y$ or $Y \gtrsim X$ whenever $X \leq CY$ for some constant $C > 0$. We use $O(Y)$ to denote any quantity X such that $|X| \lesssim Y$. We use the notation $X \sim Y$ whenever $X \lesssim Y \lesssim X$. The fact that these constants depend upon the dimension d will be suppressed. If C depends upon some additional parameters, we will indicate this with subscripts; for example, $X \lesssim_u Y$ denotes the assertion that $X \leq C_u Y$ for some C_u depending on u .

We use the “Japanese bracket” convention

$$\langle x \rangle := (1 + |x|^2)^{1/2}.$$

We write $L_t^q L_x^r$ to denote the Banach space with norm

$$\|u\|_{L_t^q L_x^r(\mathbb{R} \times \mathbb{R}^d)} := \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}^d} |u(t, x)|^r dx \right)^{q/r} dt \right)^{1/q},$$

with the usual modifications when q or r is equal to infinity, or when the domain $\mathbb{R} \times \mathbb{R}^d$ is replaced by a smaller region of spacetime such as $I \times \mathbb{R}^d$. When $q = r$ we abbreviate $L_t^q L_x^q$ as $L_{t,x}^q$.

The next lemma is a variant of Gronwall’s inequality that we will use to handle some bootstrap arguments below. The proof given is a standard application of techniques from the theories of Volterra and Toeplitz operators.

Lemma 2.1 (A Gronwall inequality). *Fix $r \in (0, 1)$ and $K \geq 4$. Let b_k be a bounded sequence of nonnegative numbers and x_k a sequence obeying $0 \leq x_k \leq b_k$ for $0 \leq k < K$ and*

$$0 \leq x_k \leq b_k + \sum_{l=0}^{k-K} r^{k-l} x_l, \tag{2-1}$$

for all $k \geq K$. Then

$$0 \leq x_k \lesssim \sum_{l=0}^k r^{k-l} \exp\left(\frac{\log(K-1)}{K-1} (k-l)\right) b_l \tag{2-2}$$

for all $k \geq 0$. In particular, if $b_k = O(2^{-k\sigma})$ and $2^\sigma r (K-1)^{1/(K-1)} < 1$, then $x_k = O(2^{-k\sigma})$.

Proof. Elementary monotonicity arguments show that we need only obtain the bound for the case of equality, namely, where

$$(1 - A)x = b. \tag{2-3}$$

Here x and b denote the semiinfinite vectors built from the corresponding sequences, while A is the matrix with entries

$$A_{k,l} = \begin{cases} r^{k-l} & \text{if } k-l \geq K, \\ 0 & \text{otherwise.} \end{cases}$$

The triangular structure of A guarantees that (2-3) can be solved (though not *a priori* in ℓ^∞); more precisely, it guarantees that the geometric series for $(1 - A)^{-1}$ converges entry-wise. To obtain bounds for the entries of this inverse matrix, it is simplest to use a functional model: under the mapping of sequences to functions

$$x_k \mapsto \sum_{k=0}^\infty x_k z^k \quad \text{and} \quad b_k \mapsto \sum_{k=0}^\infty b_k z^k,$$

the matrix A becomes multiplication by $r^K z^K (1 - rz)^{-1}$. In the same way, the entries of $(1 - A)^{-1}$ come from the Taylor coefficients of

$$a(z) := \frac{1 - rz}{1 - rz - r^K z^K}.$$

Using $e^x \geq 1 + x$ with $x = -\log |rz|$, we see that

$$|1 - rz| \geq \left(\frac{1}{r|z|} - 1\right) r|z| \geq \frac{\log(K-1)}{K-1} r|z| \geq \log(K-1) r^K |z|^K$$

on the disk $|z| \leq r^{-1} (K-1)^{-1/(K-1)}$. This shows that $a(z)$ is bounded and analytic on this disk. (Note that the hypothesis $K \geq 4$ implies that $\log(K-1) > 1$.) The inequality (2-2) now follows from the standard Cauchy estimates. □

Basic harmonic analysis. Consider a radial bump function $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$\varphi(\xi) = 1 \quad \text{if } |\xi| \leq 1 \quad \text{and} \quad \varphi(\xi) = 0 \quad \text{if } |\xi| \geq \frac{11}{10}. \tag{2-4}$$

For each number $N > 0$, define the Fourier multipliers

$$\begin{aligned} \widehat{P_{\leq N} f}(\xi) &:= \varphi(\xi/N) \hat{f}(\xi), & \widehat{P_{> N} f}(\xi) &:= (1 - \varphi(\xi/N)) \hat{f}(\xi), \\ \widehat{P_N f}(\xi) &:= \psi(\xi/N) \hat{f}(\xi) := (\varphi(\xi/N) - \varphi(2\xi/N)) \hat{f}(\xi), \end{aligned}$$

and similarly $P_{< N}$ and $P_{\geq N}$. We also define

$$P_{M < \cdot \leq N} := P_{\leq N} - P_{\leq M} = \sum_{M < N' \leq N} P_{N'}$$

whenever $M < N$. We will usually use these multipliers when M and N are *dyadic numbers* (that is, of the form 2^n for some integer n); in particular, all summations over N or M are understood to be over dyadic numbers. Nevertheless, it will occasionally be convenient to allow M and N to not be a power of 2. Note that P_N is not truly a projection; to get around this, we will occasionally need to use fattened Littlewood–Paley operators:

$$\tilde{P}_N := P_{N/2} + P_N + P_{2N}. \tag{2-5}$$

These obey $P_N \tilde{P}_N = \tilde{P}_N P_N = P_N$.

As with all Fourier multipliers, the Littlewood–Paley operators commute with the propagator $e^{it\Delta}$, as well as with differential operators such as $i\partial_t + \Delta$. We will use basic properties of these operators many times, including

Lemma 2.2 (Bernstein estimates). *For $1 \leq p \leq q \leq \infty$,*

$$\begin{aligned} \||\nabla|^{\pm s} P_N f\|_{L_x^p(\mathbb{R}^d)} &\sim N^{\pm s} \|P_N f\|_{L_x^p(\mathbb{R}^d)}, \\ \|P_{\leq N} f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{p} - \frac{d}{q}} \|P_{\leq N} f\|_{L_x^p(\mathbb{R}^d)}, \\ \|P_N f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{p} - \frac{d}{q}} \|P_N f\|_{L_x^p(\mathbb{R}^d)}. \end{aligned}$$

The next few results provide important tools for dealing with the fractional power appearing in the nonlinearity.

Lemma 2.3 (Fractional chain rule for a C^1 function [Christ and Weinstein 1991]). *Suppose $G \in C^1(\mathbb{C})$, $s \in (0, 1]$, and $1 < p, p_1, p_2 < \infty$ such that $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$. Then*

$$\||\nabla|^s G(u)\|_p \lesssim \|G'(u)\|_{p_1} \||\nabla|^s u\|_{p_2}.$$

When the function G is no longer C^1 , but merely Hölder continuous, we have the following useful chain rule:

Lemma 2.4 (Fractional chain rule for a Hölder continuous function [Visan 2007]). *Let G be a Hölder continuous function of order $0 < \alpha < 1$. Then, for every $0 < s < \alpha$, $1 < p < \infty$, and $\frac{s}{\alpha} < \sigma < 1$ we have*

$$\||\nabla|^s G(u)\|_p \lesssim \| |u|^{\alpha - \frac{s}{\sigma}} \|_{p_1} \||\nabla|^\sigma u\|_{\frac{p}{\sigma}}^{\frac{s}{\sigma}},$$

provided $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ and $(1 - s/\alpha\sigma)p_1 > 1$. The implicit constant depends upon s .

Corollary 2.5. *Let $0 \leq s < 1 + \frac{4}{d}$. Then on any spacetime slab $I \times \mathbb{R}^d$ we have*

$$\begin{aligned} \||\nabla|^s F(u)\|_{L_{t,x}^{2(d+2)/(d+4)}} &\lesssim \||\nabla|^s u\|_{L_{t,x}^{2(d+2)/d}} \|u\|_{L_{t,x}^{2(d+2)/d}}^{\frac{4}{d}}, \\ \||\nabla|^s F(u)\|_{L_t^\infty L_x^{\frac{2r}{r+4}}} &\lesssim \||\nabla|^s u\|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{\frac{2r}{d}}}^{\frac{4}{d}}, \end{aligned}$$

for any $\max\{d, 4\} \leq r \leq \infty$. The implicit constants depend upon s .

Proof. Fix a compact interval I . Throughout the proof, all spacetime estimates will be on $I \times \mathbb{R}^d$.

For $0 < s \leq 1$, both claims are easy consequences of Lemma 2.3. We now address the case $1 < s < 1 + \frac{4}{d}$ for $d \geq 5$; a few remarks on $d = 3, 4$ are given at the end of the proof. We start with the first claim.

By the chain rule and the fractional product rule, we estimate

$$\begin{aligned} & \left\| |\nabla|^s F(u) \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}} \\ & \lesssim \left\| |\nabla|^{s-1} (\nabla u F_z(u) + \nabla \bar{u} F_{\bar{z}}(u)) \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}} \\ & \lesssim \left\| |\nabla|^s u \right\|_{L_{t,x}^{2(d+2)/d}} \|u\|_{L_{t,x}^{\frac{4}{2(d+2)/d}}} + \left\| \nabla u \right\|_{L_{t,x}^{2(d+2)/d}} \left(\left\| |\nabla|^{s-1} F_z(u) \right\|_{L_{t,x}^{(d+2)/2}} + \left\| |\nabla|^{s-1} F_{\bar{z}}(u) \right\|_{L_{t,x}^{(d+2)/2}} \right). \end{aligned}$$

The claim will follow from this, once we establish

$$\left\| |\nabla|^{s-1} F_z(u) \right\|_{L_{t,x}^{(d+2)/2}} + \left\| |\nabla|^{s-1} F_{\bar{z}}(u) \right\|_{L_{t,x}^{(d+2)/2}} \lesssim \left\| |\nabla|^\sigma u \right\|_{L_{t,x}^{\frac{2(d+2)}{d}}} \|u\|_{L_{t,x}^{\frac{4-d-s-1}{d}}} \tag{2-6}$$

for some σ such that $\frac{1}{4}d(s-1) < \sigma < 1$. Indeed, one simply has to note that by interpolation we have

$$\begin{aligned} \left\| |\nabla|^\sigma u \right\|_{L_{t,x}^{2(d+2)/d}} & \lesssim \left\| |\nabla|^s u \right\|_{L_{t,x}^{2(d+2)/d}}^{\frac{\sigma}{s}} \|u\|_{L_{t,x}^{2(d+2)/d}}^{1-\frac{\sigma}{s}}, \\ \left\| \nabla u \right\|_{L_{t,x}^{2(d+2)/d}} & \lesssim \left\| |\nabla|^s u \right\|_{L_{t,x}^{2(d+2)/d}}^{\frac{1}{s}} \|u\|_{L_{t,x}^{2(d+2)/d}}^{1-\frac{1}{s}}. \end{aligned}$$

To derive (2-6), we remark that F_z and $F_{\bar{z}}$ are Hölder continuous functions of order $4/d$ and use Corollary 2.5 (with $\alpha := 4/d$ and $s := s-1$).

We now turn to the second claim. Note that the condition $r \geq 4$ simply insures that $\frac{2r}{r+4} \geq 1$. By the chain rule and the fractional product rule,

$$\begin{aligned} & \left\| |\nabla|^s F(u) \right\|_{L_t^\infty L_x^{\frac{2r}{r+4}}} \\ & \lesssim \left\| |\nabla|^{s-1} (\nabla u F_z(u) + \nabla \bar{u} F_{\bar{z}}(u)) \right\|_{L_t^\infty L_x^{\frac{2r}{r+4}}} \\ & \lesssim \left\| |\nabla|^s u \right\|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{\frac{2r}{d}}} + \left\| \nabla u \right\|_{L_t^\infty L_x^{\frac{2rs}{r+(s-1)d}}} \left\| |\nabla|^{s-1} O(|u|^{\frac{4}{d}}) \right\|_{L_t^\infty L_x^{\frac{2rs}{(r-d)(s-1)+4s}}}. \end{aligned}$$

By interpolation,

$$\left\| \nabla u \right\|_{L_t^\infty L_x^{\frac{2rs}{r+(s-1)d}}} \lesssim \left\| |\nabla|^s u \right\|_{L_t^\infty L_x^2}^{\frac{1}{s}} \|u\|_{L_t^\infty L_x^{\frac{2r}{d}}}^{1-\frac{1}{s}}.$$

Thus, the claim will follow once we establish that

$$\left\| |\nabla|^{s-1} O(|u|^{\frac{4}{d}}) \right\|_{L_t^\infty L_x^{\frac{2rs}{(r-d)(s-1)+4s}}} \lesssim \left\| |\nabla|^s u \right\|_{L_t^\infty L_x^2}^{1-\frac{1}{s}} \|u\|_{L_t^\infty L_x^{\frac{2r}{d}}}^{\frac{4}{d}+\frac{1}{s}-1}. \tag{2-7}$$

Applying Lemma 2.4, we obtain

$$\left\| |\nabla|^{s-1} O(|u|^{\frac{4}{d}}) \right\|_{L_t^\infty L_x^{\frac{2rs}{(r-d)(s-1)+4s}}} \lesssim \left\| |\nabla|^\sigma u \right\|_{L_t^\infty L_x^{\frac{2(d+2)}{d} + \frac{\sigma}{s-1}(r-d)}}^{\frac{s-1}{\sigma}} \|u\|_{L_t^\infty L_x^{\frac{4-d-s-1}{d} + \frac{2r}{d}}}$$

for any σ such that $\frac{1}{4}d(s - 1) < \sigma < 1$. The inequality (2-7) now follows from

$$\| |\nabla|^\sigma u \|_{L_t^\infty L_x^{\frac{2rs}{sd+\sigma(r-d)}}} \lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2}^{\frac{\sigma}{s}} \| u \|_{L_t^\infty L_x^{\frac{2r}{d}}}^{1-\frac{\sigma}{s}},$$

which is a consequence of interpolation.

Note that the restriction $r \geq d$ guarantees that certain Lebesgue exponents appearing above lie in the range $[1, \infty]$. In fact, one may relax this restriction a little, but we will not need this here.

The treatment of the two claims in the case $d = 4$ requires the bound

$$\| |\nabla|^\sigma |u| \|_{L_x^p} \lesssim \| |\nabla|^\sigma u \|_{L_x^p},$$

which holds for $1 < p < \infty$ and $0 < \sigma < 1$, in place of Lemma 2.4. Proofs of this slight variant of Lemma 2.3 can be found in [Kato 1995; Staffilani 1997; Taylor 2000].

We now discuss the case $d = 3$. When $1 < s < 2$, one may use the argument presented above. For $2 \leq s < \frac{7}{3}$, one first takes the Laplacian of $F(u)$ and then applies Lemma 2.4 to deal with the remaining fractional derivatives. Terms where the derivatives distribute themselves between several copies of u are dealt with by interpolation, as above. □

Strichartz estimates. Naturally, everything that we do for the nonlinear Schrödinger equation builds on basic properties of the linear propagator $e^{it\Delta}$.

From the explicit formula

$$e^{it\Delta} f(x) = \frac{1}{(4\pi i t)^{d/2}} \int_{\mathbb{R}^d} e^{i|x-y|^2/4t} f(y) dy,$$

we deduce the standard dispersive inequality

$$\| e^{it\Delta} f \|_{L^\infty(\mathbb{R}^d)} \lesssim \frac{1}{|t|^{d/2}} \| f \|_{L^1(\mathbb{R}^d)}$$

for all $t \neq 0$. Interpolating between this and the conservation of mass gives

$$\| e^{it\Delta} f \|_{L^p(\mathbb{R}^d)} \lesssim |t|^{\frac{d}{p}-\frac{d}{2}} \| f \|_{L^{p'}(\mathbb{R}^d)} \tag{2-8}$$

for all $t \neq 0$ and $2 \leq p \leq \infty$. Here p' is the dual of p , that is, $\frac{1}{p} + \frac{1}{p'} = 1$.

Finer bounds on the (frequency localized) linear propagator can be derived using stationary phase:

Lemma 2.6 (Kernel estimates). *For any $m \geq 0$, the kernel of the linear propagator obeys the estimates*

$$|(P_N e^{it\Delta})(x, y)| \lesssim_m \begin{cases} N^d \langle N|x-y\rangle^{-m} & \text{if } |t| \leq N^{-2}, \\ |t|^{-d/2} & \text{if } |t| \geq N^{-2} \text{ and } |x-y| \sim N|t|, \\ \frac{N^d}{|N^2 t|^m \langle N|x-y\rangle^m} & \text{otherwise.} \end{cases}$$

We also record the following standard Strichartz estimates:

Lemma 2.7 (Strichartz). *Let I be an interval, let $t_0 \in I$, and let*

$$u_0 \in L_x^2(\mathbb{R}^d) \quad \text{and} \quad f \in L_{t,x}^{2(d+2)/(d+4)}(I \times \mathbb{R}^d),$$

with $d \geq 3$. The function u defined by

$$u(t) := e^{i(t-t_0)\Delta} u_0 - i \int_{t_0}^t e^{i(t-t')\Delta} f(t') dt'$$

obeys the estimate

$$\|u\|_{C_t^0 L_x^2} + \|u\|_{L_{t,x}^2} \frac{2(d+2)}{d} + \|u\|_{L_t^2 L_x^{\frac{2d}{d-2}}} \lesssim \|u_0\|_{L_x^2} + \|f\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}}$$

where all spacetime norms are over $I \times \mathbb{R}^d$.

Proof. See, for example, [Ginibre and Velo 1992; Strichartz 1977]. For the endpoint see [Keel and Tao 1998]. □

We will also need three variants of the Strichartz inequality. First, we observe a weighted Strichartz estimate, which exploits the spherical symmetry heavily in order to obtain spatial decay. It is very useful in regions of space far from the origin $x = 0$.

Lemma 2.8 (Weighted Strichartz). *Let I be an interval, let $t_0 \in I$, and let*

$$u_0 \in L_x^2(\mathbb{R}^d) \quad \text{and} \quad f \in L_{t,x}^{2(d+2)/(d+4)}(I \times \mathbb{R}^d)$$

be spherically symmetric. The function u defined by

$$u(t) := e^{i(t-t_0)\Delta} u_0 - i \int_{t_0}^t e^{i(t-t')\Delta} f(t') dt'$$

obeys the estimate

$$\left\| |x|^{\frac{2(d-1)}{q}} u \right\|_{L_t^q L_x^{\frac{2q}{q-4}}(I \times \mathbb{R}^d)} \lesssim \|u_0\|_{L_x^2(\mathbb{R}^d)} + \|f\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I \times \mathbb{R}^d)}$$

if $4 \leq q \leq \infty$.

Proof. For $q = \infty$, this corresponds to the trivial endpoint in Strichartz inequality. We will only prove the result for the $q = 4$ endpoint, since the remaining cases then follow by interpolation.

As in the usual proof of Strichartz inequality, the method of TT^* together with the Christ–Kiselev lemma and Hardy–Littlewood–Sobolev inequality reduce matters to proving that

$$\left\| |x|^{\frac{d-1}{2}} e^{it\Delta} |x|^{\frac{d-1}{2}} g \right\|_{L_x^\infty(\mathbb{R}^d)} \lesssim |t|^{-\frac{1}{2}} \|g\|_{L_x^1(\mathbb{R}^d)} \tag{2-9}$$

for all radial functions g .

Let P_{rad} denote the projection onto radial functions. Then

$$[e^{it\Delta} P_{\text{rad}}](x, y) = (4\pi i t)^{-\frac{d}{2}} e^{i\frac{|x|^2+|y|^2}{4t}} \int_{S^{d-1}} e^{-i\frac{|y|\omega \cdot x}{2t}} d\sigma(\omega),$$

where $d\sigma$ denotes the uniform probability measure on the unit sphere S^{d-1} . This integral can be evaluated exactly in terms the Bessel function $J_{\frac{d-2}{2}}$. Using this, or simple stationary phase arguments, one sees that

$$|[e^{it\Delta} P_{\text{rad}}](x, y)| \lesssim |t|^{-\frac{d}{2}} \left(\frac{|y||x|}{|t|} \right)^{-\frac{d-1}{2}} \lesssim |t|^{-\frac{1}{2}} |x|^{-\frac{d-1}{2}} |y|^{-\frac{d-1}{2}}.$$

The radial dispersive estimate (2-9) now follows easily. □

We will rely crucially on a slightly different type of improvement to the Strichartz inequality in the spherically symmetric case due to Shao [2007], which improves the spacetime decay of the solution after localizing in frequency. The important thing for us will be the fact that this estimate can give decay as $N \rightarrow \infty$; this is not possible without the radial assumption.

Lemma 2.9 (Shao’s Strichartz estimate [Shao 2007, Corollary 6.2]). *For $f \in L^2_{\text{rad}}(\mathbb{R}^d)$ we have*

$$\|P_N e^{it\Delta} f\|_{L^q_{t,x}(\mathbb{R} \times \mathbb{R}^d)} \lesssim_q N^{\frac{d}{2} - \frac{d+2}{q}} \|f\|_{L^2_x(\mathbb{R}^d)},$$

provided $q > \frac{4d+2}{2d-1}$.

The last result is a bilinear estimate from [Visan 2006], which builds on earlier versions in [Bourgain 1999a; Colliander et al. 2008a]. It will be useful for controlling interactions between widely separated frequencies.

Lemma 2.10 (Bilinear Strichartz [Visan 2006, Lemma 2.5]). *For any spacetime slab $I \times \mathbb{R}^d$, any $t_0 \in I$, and any $M, N > 0$, we have*

$$\begin{aligned} \|(P_{\geq N}u)(P_{\leq M}v)\|_{L^2_{t,x}(I \times \mathbb{R}^d)} &\lesssim N^{-\frac{1}{2}} M^{\frac{d-1}{2}} \left(\|P_{\geq N}u(t_0)\|_{L^2} + \|(i\partial_t + \Delta)P_{\geq N}u\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I \times \mathbb{R}^d)} \right) \\ &\quad \times \left(\|P_{\leq M}v(t_0)\|_{L^2} + \|(i\partial_t + \Delta)P_{\leq M}v\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I \times \mathbb{R}^d)} \right), \end{aligned}$$

for all functions u, v on I .

3. The self-similar solution

In this section we preclude self-similar solutions. As mentioned in Section 1, the key ingredient is additional regularity.

Theorem 3.1 (Regularity in the self-similar case). *Let $d \geq 3$ and let u be a spherically symmetric solution to (1-1) that is almost periodic modulo scaling and that is self-similar in the sense of Theorem 1.10. Then $u(t) \in H^s_x(\mathbb{R}^d)$ for all $t \in (0, \infty)$ and all $0 \leq s < 1 + \frac{4}{d}$.*

Corollary 3.2 (Absence of self-similar solutions). *For $d \geq 3$ there are no nonzero spherically symmetric solutions to (1-1) that are self-similar in the sense of Theorem 1.10.*

Proof. By Theorem 3.1, any such solution would obey $u(t) \in H^1_x(\mathbb{R}^d)$ for all $t \in (0, \infty)$. Then, by the H^1_x global well-posedness theory described after Theorem 1.5, there exists a global solution with initial data $u(t_0)$ at any time $t_0 \in (0, \infty)$; recall that we assume $M(u) < M(Q)$ in the focusing case. On the other hand, self-similar solutions blow up at time $t = 0$. These two facts (combined with the uniqueness statement in Theorem 1.3) yield a contradiction. □

The remainder of this section is devoted to proving Theorem 3.1. We will regard s as fixed and will allow constants to implicitly depend on s .

Let u be as in Theorem 3.1. For any $A > 0$, we define

$$\begin{aligned} \mathcal{M}(A) &:= \sup_{T>0} \|u_{>AT^{-\frac{1}{2}}}(T)\|_{L_x^2(\mathbb{R}^d)}, \\ \mathcal{S}(A) &:= \sup_{T>0} \|u_{>AT^{-\frac{1}{2}}}\|_{L_{t,x}^{2(d+2)/d}([T,2T]\times\mathbb{R}^d)}, \\ \mathcal{N}(A) &:= \sup_{T>0} \|P_{>AT^{-\frac{1}{2}}}F(u)\|_{L_{t,x}^{2(d+2)/(d+4)}([T,2T]\times\mathbb{R}^d)}. \end{aligned} \tag{3-1}$$

The notation chosen indicates the quantity being measured, namely, the mass, the symmetric Strichartz norm, and the nonlinearity in the adjoint Strichartz norm, respectively. Since u is self-similar, $N(t)$ is comparable to $T^{-\frac{1}{2}}$ for t in the interval $[T, 2T]$. Thus, the Littlewood–Paley projections are adapted to the natural frequency scale on each dyadic time interval.

To prove Theorem 3.1 it suffices to show that for every $0 < s < 1 + \frac{4}{d}$ we have

$$\mathcal{M}(A) \lesssim_{s,u} A^{-s},$$

whenever A is sufficiently large depending on u and s . To establish this, we need a variety of estimates linking \mathcal{M} , \mathcal{S} , and \mathcal{N} . From mass conservation, Lemma 1.12, self-similarity, and Hölder’s inequality, we see that

$$\mathcal{M}(A) + \mathcal{S}(A) + \mathcal{N}(A) \lesssim_u 1 \tag{3-2}$$

for all $A > 0$. From the Strichartz inequality (Lemma 2.7), we also see that

$$\mathcal{S}(A) \lesssim \mathcal{M}(A) + \mathcal{N}(A) \tag{3-3}$$

for all $A > 0$. Another application of Strichartz combined with Lemma 1.12 and (1-4) shows that

$$\|u\|_{L_t^2 L_x^{\frac{2d}{d-2}}([T,2T]\times\mathbb{R}^d)} \lesssim_u 1. \tag{3-4}$$

Next, we obtain a deeper connection between these quantities.

Lemma 3.3 (Nonlinear estimate). *Let $\eta > 0$ and $0 < s < 1 + \frac{4}{d}$. If $A > 100$ and $0 < \beta \leq 1$, we have*

$$\mathcal{N}(A) \lesssim_u \sum_{N \leq \eta A^\beta} \left(\frac{N}{A}\right)^s \mathcal{S}(N) + (\mathcal{S}(\eta A^{\frac{\beta}{2(d-1)}}) + \mathcal{S}(\eta A^\beta))^{\frac{4}{d}} \mathcal{S}(\eta A^\beta) + A^{-\frac{2\beta}{d^2}} (\mathcal{M}(\eta A^\beta) + \mathcal{N}(\eta A^\beta)). \tag{3-5}$$

Proof. Fix $\eta > 0$ and $0 < s < 1 + \frac{4}{d}$. It suffices to bound

$$\|P_{>AT^{-\frac{1}{2}}}F(u)\|_{L_{t,x}^{2(d+2)/(d+4)}([T,2T]\times\mathbb{R}^d)}$$

by the right-hand side of (3-5) for arbitrary $T > 0$ and all $A > 100$ and $0 < \beta \leq 1$.

To achieve this, we decompose

$$\begin{aligned} F(u) &= F(u_{\leq \eta A^\beta T^{-\frac{1}{2}}}) + O(|u_{\leq \eta A^\beta T^{-\frac{1}{2}}}|^{\frac{4}{d}} |u_{> \eta A^\beta T^{-\frac{1}{2}}}|) \\ &\quad + O(|u_{\eta A^\beta T^{-\frac{1}{2}} < \cdot \leq \eta A^\beta T^{-\frac{1}{2}}}|^{\frac{4}{d}} |u_{> \eta A^\beta T^{-\frac{1}{2}}}|) + O(|u_{> \eta A^\beta T^{-\frac{1}{2}}}|^{1+\frac{4}{d}}), \end{aligned} \tag{3-6}$$

where

$$\alpha = \frac{\beta}{2(d-1)}.$$

To estimate the contribution from the last two terms in the expansion above, we discard the projection to high frequencies and then use Hölder’s inequality and (3-1):

$$\begin{aligned} \left\| |u_{\eta A^\alpha T^{-\frac{1}{2}} < \cdot \leq \eta A^\beta T^{-\frac{1}{2}}} |^{\frac{4}{d}} u_{> \eta A^\beta T^{-\frac{1}{2}}} \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T, 2T] \times \mathbb{R}^d)} &\lesssim \mathcal{G}(\eta A^\alpha)^{\frac{4}{d}} \mathcal{G}(\eta A^\beta), \\ \left\| |u_{> \eta A^\beta T^{-\frac{1}{2}}} |^{1+\frac{4}{d}} \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T, 2T] \times \mathbb{R}^d)} &\lesssim \mathcal{G}(\eta A^\beta)^{1+\frac{4}{d}}. \end{aligned}$$

To estimate the contribution coming from second term on the right-hand side of (3-6), we discard the projection to high frequencies and then use Hölder’s inequality, Lemmas 2.2 and 2.10, and (3-3):

$$\begin{aligned} &\left\| P_{> AT^{-\frac{1}{2}}} O\left(|u_{\leq \eta A^\alpha T^{-\frac{1}{2}}} |^{\frac{4}{d}} |u_{> \eta A^\beta T^{-\frac{1}{2}}} |\right) \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T, 2T] \times \mathbb{R}^d)} \\ &\lesssim \left\| u_{\leq \eta A^\alpha T^{-\frac{1}{2}}} u_{> \eta A^\beta T^{-\frac{1}{2}}} \right\|_{L_{t,x}^2([T, 2T] \times \mathbb{R}^d)}^{\frac{8}{d^2}} \\ &\quad \times \left\| u_{> \eta A^\beta T^{-\frac{1}{2}}} \right\|_{L_{t,x}^{2(d+2)/d}([T, 2T] \times \mathbb{R}^d)}^{1-\frac{8}{d^2}} \left\| u_{\leq \eta A^\alpha T^{-\frac{1}{2}}} \right\|_{L_{t,x}^2([T, 2T] \times \mathbb{R}^d)}^{\frac{4}{d}-\frac{8}{d^2}} \\ &\lesssim u \left((\eta A^\beta T^{-\frac{1}{2}})^{-\frac{1}{2}} (\eta A^\alpha T^{-\frac{1}{2}})^{\frac{d-1}{2}} \right)^{\frac{8}{d^2}} (\mathcal{M}(\eta A^\beta) + \mathcal{N}(\eta A^\beta))^{\frac{8}{d^2}} \mathcal{G}(\eta A^\beta)^{1-\frac{8}{d^2}} T^{\frac{2}{d}-\frac{4}{d^2}} \\ &\lesssim u A^{-\frac{2\beta}{d^2}} (\mathcal{M}(\eta A^\beta) + \mathcal{N}(\eta A^\beta)). \end{aligned}$$

We now turn to the first term on the right-hand side of (3-6). By Lemma 2.2 and Corollary 2.5 combined with (3-2), we estimate

$$\begin{aligned} \left\| P_{> AT^{-\frac{1}{2}}} F(u_{\leq \eta A^\beta T^{-\frac{1}{2}}}) \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T, 2T] \times \mathbb{R}^d)} &\lesssim (AT^{-\frac{1}{2}})^{-s} \left\| |\nabla|^s F(u_{\leq \eta A^\beta T^{-\frac{1}{2}}}) \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T, 2T] \times \mathbb{R}^d)} \\ &\lesssim u (AT^{-\frac{1}{2}})^{-s} \left\| |\nabla|^s u_{\leq \eta A^\beta T^{-\frac{1}{2}}} \right\|_{L_{t,x}^{\frac{2(d+2)}{d}}([T, 2T] \times \mathbb{R}^d)} \\ &\lesssim u \sum_{N \leq \eta A^\beta} \left(\frac{N}{A}\right)^s \mathcal{G}(N), \end{aligned}$$

which is acceptable. This finishes the proof of the lemma. □

We have some decay as $A \rightarrow \infty$:

Lemma 3.4 (Qualitative decay).

$$\lim_{A \rightarrow \infty} \mathcal{M}(A) = \lim_{A \rightarrow \infty} \mathcal{G}(A) = \lim_{A \rightarrow \infty} \mathcal{N}(A) = 0.$$

Proof. The vanishing of the first limit follows from Definition 1.8, self-similarity, and (3-1). By interpolation, (3-1), and (3-4),

$$\mathcal{G}(A) \lesssim \mathcal{M}(A)^{\frac{2}{d+2}} \left\| u_{\geq AT^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d-2}}([T, 2T] \times \mathbb{R}^d)}^{\frac{d}{d+2}} \lesssim u \mathcal{M}(A)^{\frac{2}{d+2}}.$$

Thus, as the first limit in Lemma 3.4 vanishes, we obtain that the second limit vanishes. The vanishing of the third limit follows from that of the second and Lemma 3.3. \square

We have now gathered enough tools to prove some regularity, albeit in the symmetric Strichartz space. As such, the next result is the crux of this section.

Proposition 3.5 (Quantitative decay estimate). *Let $0 < \eta < 1$ and $0 < s < 1 + \frac{4}{d}$. If η is sufficiently small depending on u and s , and A is sufficiently large depending on u , s , and η , then*

$$\mathcal{G}(A) \leq \sum_{N \leq \eta A} \left(\frac{N}{A}\right)^s \mathcal{G}(N) + A^{-\frac{1}{d^2}}. \tag{3-7}$$

In particular,

$$\mathcal{G}(A) \lesssim_u A^{-\frac{1}{d^2}}, \tag{3-8}$$

for all $A > 0$.

Proof. Fix $\eta \in (0, 1)$ and $0 < s < 1 + \frac{4}{d}$. To establish (3-7), it suffices to show

$$\|u_{>AT^{-\frac{1}{2}}}\|_{L_{t,x}^{\frac{2(d+2)}{d}}([T,2T] \times \mathbb{R}^d)} \lesssim_{u,\varepsilon} \sum_{N \leq \eta A} \left(\frac{N}{A}\right)^{s+\varepsilon} \mathcal{G}(N) + A^{-\frac{3}{2d^2}} \tag{3-9}$$

for all $T > 0$ and some small $\varepsilon = \varepsilon(d, s) > 0$, since then (3-7) follows by requiring η to be small and A to be large, both depending upon u and also ε .

Fix $T > 0$. By writing the Duhamel formula (1-2) beginning at $T/2$ and then using Lemma 2.7, we obtain

$$\begin{aligned} & \|u_{>AT^{-\frac{1}{2}}}\|_{L_{t,x}^{\frac{2(d+2)}{d}}([T,2T] \times \mathbb{R}^d)} \\ & \lesssim \|P_{>AT^{-\frac{1}{2}}} e^{i(t-T/2)\Delta} u(T/2)\|_{L_{t,x}^{\frac{2(d+2)}{d}}([T,2T] \times \mathbb{R}^d)} + \|P_{>AT^{-\frac{1}{2}}} F(u)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T,2T] \times \mathbb{R}^d)}. \end{aligned}$$

We first consider the second term. By (3-1), we have

$$\|P_{>AT^{-\frac{1}{2}}} F(u)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T,2T] \times \mathbb{R}^d)} \lesssim \mathcal{N}(A/2).$$

Using Lemma 3.3 (with $\beta = 1$ and s replaced by $s + \varepsilon$ for some $0 < \varepsilon < 1 + \frac{4}{d} - s$) combined with Lemma 3.4 (choosing A sufficiently large depending on u , s , and η), and (3-2), we derive

$$\|P_{>AT^{-\frac{1}{2}}} F(u)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}([T,2T] \times \mathbb{R}^d)} \lesssim_{u,\varepsilon} \sum_{N \leq \eta A} \left(\frac{N}{A}\right)^{s+\varepsilon} \mathcal{G}(N) + A^{-\frac{3}{2d^2}},$$

whose right-hand side is that of (3-9). Thus, the second term is acceptable.

We now consider the first term. It suffices to show

$$\|P_{>AT^{-\frac{1}{2}}} e^{i(t-T/2)\Delta} u(T/2)\|_{L_{t,x}^{\frac{2(d+2)}{d}}([T,2T] \times \mathbb{R}^d)} \lesssim_u A^{-\frac{3}{2d^2}}, \tag{3-10}$$

which we will deduce by first proving two estimates at a single frequency scale, interpolating between them, and then summing.

From Lemma 2.9 and mass conservation, we have

$$\| P_{BT^{-\frac{1}{2}}} e^{i(t-T/2)\Delta} u(T/2) \|_{L_{t,x}^q([T,2T] \times \mathbb{R}^d)} \lesssim_{u,q} (BT^{-\frac{1}{2}})^{\frac{d}{2} - \frac{d+2}{q}} \tag{3-11}$$

whenever

$$\frac{4d+2}{2d-1} < q \leq \frac{2(d+2)}{d}$$

and $B > 0$. This is our first estimate.

Using the Duhamel formula (1-2), we write

$$P_{BT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) = P_{BT^{-\frac{1}{2}}} e^{i(t-\varepsilon)\Delta} u(\varepsilon) - i \int_{\varepsilon}^{\frac{T}{2}} P_{BT^{-\frac{1}{2}}} e^{i(t-t')\Delta} F(u(t')) dt'$$

for any $\varepsilon > 0$. By self-similarity, the former term converges strongly to zero in L_x^2 as $\varepsilon \rightarrow 0$. Convergence to zero in $L_x^{2d/(d-2)}$ then follows from Lemma 2.2. Thus, using Hölder’s inequality followed by the dispersive estimate (2-8), and then (3-4), we estimate

$$\begin{aligned} & \| P_{BT^{-\frac{1}{2}}} e^{i(t-T/2)\Delta} u(T/2) \|_{L_{t,x}^{\frac{2d}{d-2}}([T,2T] \times \mathbb{R}^d)} \\ & \lesssim T^{\frac{d-2}{2d}} \left\| \int_0^{\frac{T}{2}} \frac{1}{t-t'} \| F(u(t')) \|_{L_x^{\frac{2d}{d+2}}} dt' \right\|_{L_t^\infty([T,2T])} \lesssim T^{-\frac{d+2}{2d}} \| F(u) \|_{L_t^1 L_x^{\frac{2d}{d+2}}((0, \frac{T}{2}] \times \mathbb{R}^d)} \\ & \lesssim T^{-\frac{d+2}{2d}} \sum_{0 < \tau \leq \frac{T}{4}} \| F(u) \|_{L_t^1 L_x^{\frac{2d}{d+2}}([\tau, 2\tau] \times \mathbb{R}^d)} \\ & \lesssim T^{-\frac{d+2}{2d}} \sum_{0 < \tau \leq \frac{T}{4}} \tau^{1/2} \| u \|_{L_t^2 L_x^{\frac{2d}{d-2}}([\tau, 2\tau] \times \mathbb{R}^d)} \| u \|_{L_t^\infty L_x^2([\tau, 2\tau] \times \mathbb{R}^d)}^{\frac{4}{d}} \\ & \lesssim_u T^{-1/d}. \end{aligned}$$

Interpolating between the estimate just proved and (3-11) with

$$q = \frac{2d(d+2)(4d-3)}{4d^3 - 3d^2 + 12},$$

we obtain

$$\| P_{BT^{-\frac{1}{2}}} e^{i(t-T/2)\Delta} u(T/2) \|_{L_{t,x}^{\frac{2(d+2)}{d}}([T,2T] \times \mathbb{R}^d)} \lesssim_u B^{-\frac{3}{2d^2}}.$$

Summing this over dyadic $B \geq A$ yields (3-10) and hence (3-9).

We now justify (3-8). Given an integer $K \geq 4$, we set $\eta = 2^{-K}$. Then there exists A_0 , depending on u and K , so that (3-7) holds for $A \geq A_0$. By (3-2), we need only bound $\mathcal{S}(A)$ for $A \geq A_0$.

Let $k \geq 0$ and set $A = 2^k A_0$ in (3-7). Then, writing $N = 2^l A_0$ and using (3-2), we have

$$\mathcal{G}(2^k A_0) \leq \sum_{l \leq k-K} 2^{-(k-l)s} \mathcal{G}(2^l A_0) + (2^k A_0)^{-\beta} \leq \sum_{l=0}^{k-K} 2^{-(k-l)s} \mathcal{G}(2^l A_0) + \frac{2^{-ks}}{1-2^{-s}} \mathcal{G}(0) + 2^{-k\beta} A_0^{-\beta},$$

where $\beta := d^{-2}$. Setting $s = 1$ and applying Lemma 2.1 with $x_k = \mathcal{G}(2^k A_0)$ and $b_k = O_u(2^{-k\beta})$, we deduce

$$\mathcal{G}(2^k A_0) \lesssim_u 2^{-k/d^2},$$

provided K is chosen sufficiently large. This gives the necessary bound on \mathcal{G} . □

Corollary 3.6. *For any $A > 0$ we have*

$$\mathcal{M}(A) + \mathcal{G}(A) + \mathcal{N}(A) \lesssim_u A^{-1/d^2}.$$

Proof. The bound on \mathcal{G} was proved in the previous proposition. The bound on \mathcal{N} follows from this, Lemma 3.3 with $\beta = 1$, and (3-2).

We now turn to the bound on \mathcal{M} . By Lemma 1.11,

$$\|P_{>AT^{-\frac{1}{2}}} u(T)\|_2 \lesssim \sum_{k=0}^{\infty} \left\| \int_{2^k T}^{2^{k+1} T} e^{i(T-t')\Delta} P_{>AT^{-\frac{1}{2}}} F(u(t')) dt' \right\|_2, \tag{3-12}$$

where weak convergence has become strong convergence because of the frequency projection and the fact that $N(t) = t^{-1/2} \rightarrow 0$ as $t \rightarrow \infty$. Intuitively, the reason for using (1-5) forward in time is that the solution becomes smoother as $N(t) \rightarrow 0$.

Combining (3-12) with Lemma 2.7 and (3-1), we get

$$\mathcal{M}(A) = \sup_{T>0} \|P_{>AT^{-\frac{1}{2}}} u(T)\|_2 \lesssim \sum_{k=0}^{\infty} \mathcal{N}(2^{k/2} A). \tag{3-13}$$

The desired bound on \mathcal{M} now follows from that on \mathcal{N} . □

Proof of Theorem 3.1. Let $0 < s < 1 + \frac{4}{d}$. Combining Lemma 3.3 (with $\beta = 1 - \frac{1}{2d^2}$), (3-3), and (3-13), we deduce that if

$$\mathcal{G}(A) + \mathcal{M}(A) + \mathcal{N}(A) \lesssim_u A^{-\sigma}$$

for some $0 < \sigma < s$, then

$$\mathcal{G}(A) + \mathcal{M}(A) + \mathcal{N}(A) \lesssim_u A^{-\sigma} \left(A^{-\frac{s-\sigma}{2d^2}} + A^{-\frac{(d+1)(3d-2)\sigma}{2d^3(d-1)}} + A^{-\frac{3-\sigma}{2d^2} - \frac{d^2-2}{2d^4}} \right).$$

More precisely, Lemma 3.3 provides the bound on $\mathcal{N}(A)$, then (3-13) gives the bound on $\mathcal{M}(A)$ and then finally (3-3) gives the bound on $\mathcal{G}(A)$.

Iterating this statement shows that $u(t) \in H_x^s(\mathbb{R}^d)$ for all $0 < s < 1 + \frac{4}{d}$. Note that Corollary 3.6 allows us to begin the iteration with $\sigma = d^{-2}$. □

4. An in/out decomposition

In this section, we will often write radial functions on \mathbb{R}^d just in terms of the radial variable. With this convention,

$$f(r) = r^{\frac{2-d}{2}} \int_0^\infty J_{\frac{d-2}{2}}(kr) \hat{f}(k) k^{\frac{d}{2}} dk \quad \text{and} \quad \hat{f}(k) = k^{\frac{2-d}{2}} \int_0^\infty J_{\frac{d-2}{2}}(kr) f(r) r^{\frac{d}{2}} dr,$$

as can be seen from [Stein and Weiss 1971, Theorem IV.3.3]. Here J_ν denotes the Bessel function of order ν . In particular,

$$g(k, r) := r^{\frac{2-d}{2}} J_{\frac{d-2}{2}}(kr)$$

solves the radial Helmholtz equation

$$-g_{rr} - \frac{d-1}{r} g_r = k^2 g, \tag{4-1}$$

which corresponds to the fact that $g(k, r)$ represents a spherical standing wave of frequency $k^2/(2\pi)$. Incoming and outgoing spherical waves are represented by two further solutions of (4-1), namely,

$$g_-(k, r) := r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(2)}(kr) \quad \text{and} \quad g_+(k, r) := r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(1)}(kr),$$

respectively. Note that

$$g = \frac{1}{2}g_+ + \frac{1}{2}g_-.$$

This leads us to define the projection onto outgoing spherical waves by

$$\begin{aligned} [P^+ f](r) &= \frac{1}{2} \int_0^\infty r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(1)}(kr) \hat{f}(k) k^{\frac{d}{2}} dk \\ &= \frac{1}{2} r^{\frac{2-d}{2}} \int_0^\infty \left(\int_0^\infty H_{\frac{d-2}{2}}^{(1)}(kr) J_{\frac{d-2}{2}}(k\rho) k dk \right) f(\rho) \rho^{\frac{d}{2}} d\rho \\ &= \frac{1}{2} f(r) + \frac{i}{\pi} \int_0^\infty \frac{r^{2-d} f(\rho) \rho^{d-1} d\rho}{r^2 - \rho^2}. \end{aligned} \tag{4-2}$$

In order to derive the last equality we used [Gradshteyn and Ryzhik 2000, §6.521.2] together with analytic continuation. Similarly, we define the projection onto incoming waves by

$$[P^- f](r) = \frac{1}{2} \int_0^\infty r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(2)}(kr) \hat{f}(k) k^{\frac{d}{2}} dk = \frac{1}{2} f(r) - \frac{i}{\pi} \int_0^\infty \frac{r^{2-d} f(\rho) \rho^{d-1} d\rho}{r^2 - \rho^2}.$$

Note that the kernel of P^- is the complex conjugate of that belonging to P^+ , as is required by time-reversal symmetry.

We will write P_N^\pm for the product $P^\pm P_N$.

Remark. For $f(\rho) \in L^2(\rho^{d-1} d\rho)$,

$$\int_0^\infty |f(\rho)|^2 \rho^{d-1} d\rho = \frac{1}{2} \int |s^{\frac{d-2}{4}} f(\sqrt{s})|^2 ds$$

and with $t = r^2$,

$$\int_0^\infty \frac{r^{2-d} f(\rho) \rho^{d-1} d\rho}{r^2 - \rho^2} = \frac{1}{2} t^{-\frac{d-2}{4}} \int_0^\infty \left(\frac{s}{t}\right)^{\frac{d-2}{4}} \frac{s^{\frac{d-2}{4}} f(\sqrt{s}) ds}{t-s}.$$

Thus $P^+ : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ is bounded if and only if the Hilbert transform is bounded in the weighted space $L^2([0, \infty), t^{-(d-2)/2} dt)$. Thus P^+ is unbounded on $L^2(\mathbb{R}^d)$ for $d \geq 4$.

Lemma 4.1 (Kernel estimates). *For $|x| \gtrsim N^{-1}$ and $t \gtrsim N^{-2}$, the integral kernel obeys*

$$|[P_N^\pm e^{\mp it\Delta}](x, y)| \lesssim \begin{cases} (|x||y|)^{-\frac{d-1}{2}} |t|^{-\frac{1}{2}} & \text{if } |y| - |x| \sim Nt, \\ \frac{N^d}{(N|x|)^{\frac{d-1}{2}} \langle N|y| \rangle^{\frac{d-1}{2}}} \langle N^2t + N|x| - N|y| \rangle^{-m} & \text{otherwise,} \end{cases}$$

for any $m \geq 0$. For $|x| \gtrsim N^{-1}$ and $|t| \lesssim N^{-2}$, the integral kernel obeys

$$|[P_N^\pm e^{\mp it\Delta}](x, y)| \lesssim \frac{N^d}{(N|x|)^{\frac{d-1}{2}} \langle N|y| \rangle^{\frac{d-1}{2}}} \langle N|x| - N|y| \rangle^{-m}$$

for any $m \geq 0$.

Proof. The proof is an exercise in stationary phase. We will only provide the details for $P_N^+ e^{-it\Delta}$, the other kernel being its complex conjugate. By (4-2) we have the following formula for the kernel:

$$[P_N^+ e^{-it\Delta}](x, y) = \frac{1}{2} (|x||y|)^{-\frac{d-2}{2}} \int_0^\infty H_{\frac{d-2}{2}}^{(1)}(k|x|) J_{\frac{d-2}{2}}(k|y|) e^{itk^2} \psi\left(\frac{k}{N}\right) k dk \quad (4-3)$$

where ψ is the multiplier from the Littlewood–Paley projection. To proceed, we use the following information about Bessel/Hankel functions:

$$J_{\frac{d-2}{2}}(r) = \frac{a(r)e^{ir}}{\langle r \rangle^{1/2}} + \frac{\bar{a}(r)e^{-ir}}{\langle r \rangle^{1/2}}, \quad (4-4)$$

where $a(r)$ obeys the symbol estimates

$$\left| \frac{\partial^m a(r)}{\partial r^m} \right| \lesssim_m \langle r \rangle^{-m} \quad \text{for all } m \geq 0. \quad (4-5)$$

The Hankel function $H_{\frac{d-2}{2}}^{(1)}(r)$ has a singularity at $r = 0$; however, for $r \gtrsim 1$,

$$H_{\frac{d-2}{2}}^{(1)}(r) = \frac{b(r)e^{ir}}{r^{1/2}} \quad (4-6)$$

for a smooth function $b(r)$ obeying (4-5). Since we assumed $|x| \gtrsim N^{-1}$, the singularity does not enter into our considerations.

Substituting (4-4) and (4-6) into (4-3), we see that a stationary phase point can only occur in the term containing $\bar{a}(r)$ and even then only if $|y| - |x| \sim Nt$. In this case, stationary phase yields the first estimate. In all other cases, integration by parts yields the second estimate.

The short-time estimate is also a consequence of (4-3) and stationary phase techniques. Since t is so small, e^{ik^2t} shows no appreciable oscillation and can be incorporated into $\psi(\frac{k}{N})$. For $||y|-|x|| \leq N^{-1}$, the result follows from the naive L^1 estimate. For larger $|x|-|y|$, one integrates by parts m times. \square

Lemma 4.2 (Properties of P^\pm). (i) $P^+ + P^-$ acts as the identity on $L^2_{\text{rad}}(\mathbb{R}^d)$.

(ii) Fix $N > 0$. For any spherically symmetric function $f \in L^2_x(\mathbb{R}^d)$,

$$\|P^\pm P_{\geq N} f\|_{L^2_x(|x| \geq \frac{1}{100} N^{-1})} \lesssim \|f\|_{L^2_x(\mathbb{R}^d)}$$

with an N -independent constant.

Proof. Part (i) is immediate from the definition.

We now turn to part (ii). We only prove the inequality for P^+ , as the result for P^- can be deduced from this. Let χ be a nonnegative smooth function on \mathbb{R}^+ vanishing in a neighborhood of the origin and obeying $\chi(r) = 1$ for $r \geq \frac{1}{100}$. With this definition and (4-2),

$$\begin{aligned} \|P^\pm P_{\geq N} f\|_{L^2_x(|x| \geq N^{-1})}^2 &\leq \|\chi(N|x|)P^\pm P_{\geq N} f\|_{L^2_x(\mathbb{R}^d)}^2 \\ &= \int_0^\infty \left| \int_0^\infty H_{\frac{d-2}{2}}^{(1)}(kr) \hat{f}(k) k^{\frac{d}{2}} \left(1 - \varphi\left(\frac{k}{N}\right)\right) dk \right|^2 \chi(Nr)^2 r \, dr, \end{aligned}$$

where φ is a cutoff function as in (2-4). Note that, by scaling, it suffices to treat the case $N = 1$. Because of the cutoffs, the only nonzero contribution comes from the region $kr \gtrsim 1$. This allows us to use the following information about Hankel functions: for $\rho \gtrsim 1$,

$$H_{\frac{d-2}{2}}^{(1)}(\rho) = \left(\frac{2}{\pi\rho}\right)^{\frac{1}{2}} (1 + b(\rho)) e^{i\rho - i(d-1)\frac{\pi}{4}}$$

where b is a symbol of order -1 , that is,

$$\left| \frac{\partial^m b(\rho)}{\partial \rho^m} \right| \lesssim_m \langle \rho \rangle^{-m-1},$$

for all $m \geq 0$; see for example [Gradshteyn and Ryzhik 2000]. Note that this is more refined than formula (4-6) used in the previous proof. With these observations, our goal has been reduced to showing that

$$\int_0^\infty \left| \int_0^\infty e^{ikr} (1 + b(kr))(1 - \varphi(k))g(k) \, dk \right|^2 \chi(r)^2 \, dr \lesssim \int_0^\infty |g(k)|^2 \, dk$$

or, equivalently, that

$$K(k, k') := (1 - \varphi(k))(1 - \varphi(k')) \int_0^\infty e^{i(k-k')r} (1 + b(kr))(1 + \bar{b}(k'r)) \chi(r)^2 \, dr$$

is the kernel of a bounded operator on $L^2_k([0, \infty))$. To this end, we will decompose K as the sum of two kernels, each of which we can estimate.

First, we consider

$$K_1(k, k') := (1 - \varphi(k))(1 - \varphi(k')) \int_0^\infty e^{i(k-k')r} \chi(r)^2 \, dr.$$

Without the prefactors, the integral is the kernel of a bounded Fourier multiplier and so a bounded operator on L^2_k . As φ is a bounded function, we may then deduce that K_1 is itself the kernel of a bounded operator.

Our second kernel is

$$K_2(k, k') := (1 - \varphi(k))(1 - \varphi(k')) \int_0^\infty e^{i(k-k')r} (b(kr) + \bar{b}(k'r) + b(kr)\bar{b}(k'r)) \chi(r)^2 dr,$$

which we will show to be bounded using Schur’s test. Note that the factors in front of the integral ensure that the kernel is zero unless $k \gtrsim 1$ and $k' \gtrsim 1$. By integration by parts, we see that

$$K_2(k, k') \lesssim_m |k - k'|^{-m}$$

for any $m \geq 1$, which offers ample control away from the diagonal. To obtain a good estimate near the diagonal, we need to break the integral into two pieces. We do this by writing

$$1 = \chi\left(\frac{r}{R}\right) + \left(1 - \chi\left(\frac{r}{R}\right)\right),$$

with $R \gg 1$. Integrating by parts once when r is large and not at all when r is small leads to

$$\begin{aligned} K_2(k, k') &\lesssim \frac{1}{|k-k'|} \int \left(\left(\frac{1}{kr^2} + \frac{1}{k'r^2} + \frac{1}{kk'r^3} \right) \chi\left(\frac{r}{R}\right) + \left(\frac{1}{kr} + \frac{1}{k'r} + \frac{1}{kk'r^2} \right) \frac{1}{R} \chi'\left(\frac{r}{R}\right) \right) dr \\ &\quad + \int \left(\frac{1}{kr} + \frac{1}{k'r} + \frac{1}{kk'r^2} \right) \chi(r)^2 \left(1 - \chi\left(\frac{r}{R}\right)\right) dr \\ &\lesssim \frac{1}{R|k-k'|} + \log R. \end{aligned}$$

Choosing $R = |k - k'|^{-1}$ provides sufficient control near the diagonal to complete the application of Schur’s test. □

5. Additional regularity

This section is devoted to prove:

Theorem 5.1 (Regularity in the global case). *Let $d \geq 3$ and let u be a global spherically symmetric solution to (1-1) that is almost periodic modulo scaling. Suppose also that $N(t) \lesssim 1$ for all $t \in \mathbb{R}$. Then $u \in L_t^\infty H_x^s(\mathbb{R} \times \mathbb{R}^d)$ for all $0 \leq s < 1 + \frac{4}{d}$.*

The argument mimics that in [Killip et al. 2007], though the nonpolynomial nature of the nonlinearity introduces several technical complications. That $u(t)$ is moderately smooth will follow from a careful study of the Duhamel formulae (1-5). Near t , we use the fact that there is little mass at high frequencies, as is implied by the definition of almost periodicity and the boundedness of the frequency scale function $N(t)$. Far from t , we use the spherical symmetry of the solution. As this symmetry is only valuable at large radii, we are only able to exploit it by using the in/out decomposition described in Section 4.

Let us now begin the proof. For the remainder of the section, u will denote a solution to (1-1) that obeys the hypotheses of Theorem 5.1. Once again, we will regard s as fixed and suppress the dependence of implicit constants upon this parameter.

We first record some basic local estimates. From mass conservation we have

$$\|u\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} \lesssim u \cdot 1,$$

while from Definition 1.8 and the fact that $N(t)$ is bounded we have

$$\lim_{N \rightarrow \infty} \|u_{\geq N}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} = 0.$$

From Lemma 1.12 and $N(t) \lesssim 1$, we have

$$\|u\|_{L_{t,x}^{\frac{2(d+2)}{d}}(J \times \mathbb{R}^d)} \lesssim u \langle |J| \rangle^{\frac{d}{2(d+2)}} \tag{5-1}$$

for all intervals $J \subset \mathbb{R}$. By Hölder’s inequality, this implies

$$\|F(u)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} \lesssim u \langle |J| \rangle^{\frac{d+4}{2(d+2)}}$$

and then, by the (endpoint) Strichartz inequality (Lemma 2.7),

$$\|u\|_{L_t^2 L_x^{\frac{2d}{d-2}}(J \times \mathbb{R}^d)} \lesssim u \langle |J| \rangle^{\frac{1}{2}}. \tag{5-2}$$

More precisely, one first treats the case $|J| = O(1)$ using (5-1) and then larger intervals by subdivision. Similarly, from the weighted Strichartz inequality (Lemma 2.8),

$$\| |x|^{\frac{d-1}{2}} u_{N_1 \leq \cdot \leq N_2} \|_{L_t^4 L_x^\infty(J \times \mathbb{R}^d)} \lesssim u \langle |J| \rangle^{\frac{1}{4}} \tag{5-3}$$

uniformly in $0 < N_1 \leq N_2 < \infty$.

Now, for any dyadic number N , define

$$\mathcal{M}(N) := \|u_{\geq N}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)}.$$

From the discussion above, we see that $\mathcal{M}(N) \lesssim u \cdot 1$ and

$$\lim_{N \rightarrow \infty} \mathcal{M}(N) = 0. \tag{5-4}$$

To prove Theorem 5.1, it suffices to show $\mathcal{M}(N) \lesssim_{u,s} N^{-s}$ for any $0 < s < 1 + \frac{4}{d}$ and all N sufficiently large depending on u and s . As we will explain momentarily, this will follow from Lemma 2.1 and the following

Proposition 5.2 (Regularity). *Let u be as in Theorem 5.1, let $0 < s < 1 + \frac{4}{d}$, and let $\eta > 0$ be a small number. Then*

$$\mathcal{M}(N) \leq N^{-s} + \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^s \mathcal{M}(M),$$

whenever N is sufficiently large depending on u, s , and η .

Indeed, given $\varepsilon > 0$, set $\eta = 2^{-K}$, where K is so large that

$$2 \log(K - 1) < \varepsilon(K - 1).$$

Let N_0 be sufficiently large depending on u, s , and K so that the inequality in Proposition 5.2 holds for $N \geq N_0$. If we write $r = 2^{-s}$, $x_k = \mathcal{M}(2^k N_0)$, and

$$b_k = 2^{-ks} N_0^{-s} + \sum_{l \leq -1} 2^{-s(k-l)} \mathcal{M}(2^l N_0) \lesssim_u 2^{-ks} \lesssim_u 2^{-k(s-\varepsilon)},$$

then (2-1) holds. Therefore, $\mathcal{M}(N) \lesssim_{u,s} N^{\varepsilon-s}$ by the last sentence in Lemma 2.1.

The rest of this section is devoted to proving Proposition 5.2. Fix $0 < s < 1 + \frac{4}{d}$ and $\eta > 0$. Our task is to show that

$$\|u_{\geq N}(t_0)\|_{L^2_x(\mathbb{R}^d)} \leq N^{-s} + \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^s \mathcal{M}(M)$$

for all times t_0 and all N sufficiently large (depending on u, s , and η). By time translation symmetry, we may assume $t_0 = 0$. As noted above, one of the keys to obtaining additional regularity is Lemma 1.11. Specifically, we have

$$\begin{aligned} u_{\geq N}(0) &= (P^+ + P^-)u_{\geq N}(0) \\ &= \lim_{T \rightarrow \infty} i \int_0^T P^+ e^{-it\Delta} P_{\geq N} F(u(t)) dt - \lim_{T \rightarrow \infty} i \int_{-T}^0 P^- e^{-it\Delta} P_{\geq N} F(u(t)) dt, \end{aligned} \tag{5-5}$$

where the limit is to be interpreted as a weak limit in L^2 . However, this representation is not useful for $|x|$ small because the kernels of P^\pm have a strong singularity at $x = 0$. To this end, we introduce the cutoff

$$\chi_N(x) := \chi(N|x|),$$

where χ is the characteristic function of $[1, \infty)$. As short times and large times will be treated differently, we rewrite (5-5) as

$$\begin{aligned} \chi_N(x)u_{\geq N}(0, x) &= i \int_0^\delta \chi_N(x) P^+ e^{-it\Delta} P_{\geq N} F(u(t)) dt - i \int_{-\delta}^0 \chi_N(x) P^- e^{-it\Delta} P_{\geq N} F(u(t)) dt \\ &\quad + \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_\delta^T \int_{\mathbb{R}^d} \chi_N(x) [P_M^+ e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt \\ &\quad - \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_{-T}^{-\delta} \int_{\mathbb{R}^d} \chi_N(x) [P_M^- e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt, \end{aligned} \tag{5-6}$$

as weak limits in L^2_x . We have used the identity

$$P_{\geq N} = \sum_{M \geq N} P_M \tilde{P}_M,$$

where $\tilde{P}_M := P_{M/2} + P_M + P_{2M}$, because of the way we will estimate the large-time integrals.

The analogous representation for treating small x is

$$\begin{aligned}
 & (1 - \chi_N(x))u_{\geq N}(0, x) \\
 &= \lim_{T \rightarrow \infty} i \int_0^T (1 - \chi_N(x))e^{-it\Delta} P_{\geq N} F(u(t)) dt \\
 &= i \int_0^\delta (1 - \chi_N(x))e^{-it\Delta} P_{\geq N} F(u(t)) dt \\
 &\quad + \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_\delta^T \int_{\mathbb{R}^d} (1 - \chi_N(x)) [P_M e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt, \tag{5-7}
 \end{aligned}$$

also as weak limits.

To deal with the poor nature of the limits in (5-6) and (5-7), we note that

$$f_T \rightarrow f \text{ weakly} \implies \|f\| \leq \limsup_{T \rightarrow \infty} \|f_T\|, \tag{5-8}$$

or equivalently, that the unit ball is weakly closed.

Despite the fact that different representations will be used depending on the size of $|x|$, some estimates can be dealt with in a uniform manner. The first such example is a bound on integrals over short times.

Lemma 5.3 (Local estimate). *Let $0 < s < 1 + \frac{4}{d}$. For any sufficiently small $\eta > 0$, there exists $\delta = \delta(u, \eta) > 0$ such that*

$$\left\| \int_0^\delta e^{-it\Delta} P_{\geq N} F(u(t)) dt \right\|_{L_x^2} \leq N^{-s} + \frac{1}{10} \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^s \mathcal{M}(M),$$

provided N is sufficiently large depending on u, s , and η . An analogous estimate holds for integration over $[-\delta, 0]$ and after premultiplication by $\chi_N P^\pm$.

Proof. By Lemma 2.7, it suffices to prove

$$\mathcal{N}(N) := \|P_{\geq N} F(u)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} \lesssim_u N^{-s-\varepsilon} + \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^{s+\varepsilon} \mathcal{M}(M) \tag{5-9}$$

for some small $\varepsilon = \varepsilon(d, s) > 0$, any interval J of length $|J| \leq \delta$, and all sufficiently large N depending on u, s , and η , since the claim would follow by requiring η small and N large, both depending on u .

From (5-4), there exists $N_0 = N_0(u, \eta)$ such that

$$\|u_{\geq N_0}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} \leq \eta^{100d^2}. \tag{5-10}$$

Let $N > N_1 := \eta^{-1} N_0$. We decompose

$$F(u) = F(u_{\leq \eta N}) + O(|u_{\leq N_0}|^{\frac{4}{d}} |u_{> \eta N}|) + O(|u_{N_0 \leq \cdot \leq \eta N}|^{\frac{4}{d}} |u_{> \eta N}|) + O(|u_{> \eta N}|^{1+\frac{4}{d}}). \tag{5-11}$$

Using Lemma 2.2, Corollary 2.5 together with (5-1), and Lemma 2.7, we estimate the contribution of the first term on the right-hand side of (5-11) as follows:

$$\begin{aligned} \|P_{\geq N} F(u_{\leq \eta N})\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} &\lesssim N^{-s-3\varepsilon} \||\nabla|^{s+3\varepsilon} F(u_{\leq \eta N})\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} \\ &\lesssim_u \langle \delta \rangle^{\frac{2}{d+2}} N^{-s-3\varepsilon} \||\nabla|^{s+3\varepsilon} u_{\leq \eta N}\|_{L_{t,x}^{\frac{2(d+2)}{d}}(J \times \mathbb{R}^d)} \\ &\lesssim_u \langle \delta \rangle^{\frac{2}{d+2}} \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^{s+3\varepsilon} (\mathcal{M}(M) + \mathcal{N}(M)) \\ &\lesssim_u \eta^\varepsilon \langle \delta \rangle^{\frac{2}{d+2}} \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^{s+2\varepsilon} (\mathcal{M}(M) + \mathcal{N}(M)), \end{aligned}$$

for any positive $\varepsilon < \frac{1}{3}(1 + \frac{4}{d} - s)$.

To estimate the contribution of the second term on the right-hand side of (5-11), we use Hölder’s inequality, Lemma 2.2, and (5-1):

$$\begin{aligned} \|O(|u_{\leq N_0}|^{\frac{4}{d}} |u_{> \eta N}|)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} &\lesssim \delta^{\frac{1}{2}} \|u_{\leq N_0}\|_{L_{t,x}^{2(d+2)/d}(J \times \mathbb{R}^d)}^{\frac{2}{d}} \|u_{\leq N_0}\|_{L_{t,x}^\infty(J \times \mathbb{R}^d)}^{\frac{2}{d}} \|u_{> \eta N}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \\ &\lesssim_u \delta^{\frac{1}{2}} \langle \delta \rangle^{\frac{1}{d+2}} N_0 \mathcal{M}(\eta N). \end{aligned}$$

Finally, to estimate the contribution of the last two terms on the right-hand side of (5-11), we use Hölder’s inequality, interpolation combined with (5-2) and (5-10), and then Lemma 2.7 to obtain

$$\begin{aligned} \|O(|u_{N_0 \leq \cdot \leq \eta N}|^{\frac{4}{d}} |u_{> \eta N}|)\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} &\lesssim \|u_{N_0 \leq \cdot \leq \eta N}\|_{L_{t,x}^{2(d+2)/d}(J \times \mathbb{R}^d)}^{\frac{4}{d}} \|u_{> \eta N}\|_{L_{t,x}^{2(d+2)/d}(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{N_0 \leq \cdot \leq \eta N}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)}^{\frac{8}{d(d+2)}} \|u_{N_0 \leq \cdot \leq \eta N}\|_{L_t^2 L_x^{2d/(d-2)}(J \times \mathbb{R}^d)}^{\frac{4}{d+2}} (\mathcal{M}(\eta N) + \mathcal{N}(\eta N)) \\ &\lesssim_u \eta^8 \langle \delta \rangle^{\frac{2}{d+2}} (\mathcal{M}(\eta N) + \mathcal{N}(\eta N)); \end{aligned}$$

similarly,

$$\|O(|u_{> \eta N}|^{1+\frac{4}{d}})\|_{L^{\frac{2(d+2)}{d+4}}(J \times \mathbb{R}^d)} \lesssim_u \eta^8 \langle \delta \rangle^{\frac{2}{d+2}} (\mathcal{M}(\eta N) + \mathcal{N}(\eta N)).$$

Putting everything together and taking η sufficiently small depending on u and s , then δ sufficiently small depending upon N_0 and η , we derive

$$\mathcal{N}(N) \leq \sum_{M \leq \eta N} \left(\frac{M}{N}\right)^{s+2\varepsilon} (\mathcal{M}(M) + \mathcal{N}(M)) \tag{5-12}$$

for all $N > N_1$ and $\varepsilon > 0$ as above. The claim (5-9) follows from this and Lemma 2.1. More precisely, let $\eta = 2^{-K}$ where K is sufficiently large so that $2 \log(K - 1) < \varepsilon(K - 1)$. If we write $r = 2^{-s-2\varepsilon}$, $x_k = \mathcal{N}(2^k N_1)$, and

$$b_k = \sum_{l \leq k-K} 2^{-(s+2\varepsilon)(k-l)} \mathcal{M}(2^l N_1) + \sum_{l \leq -1} 2^{-(s+2\varepsilon)(k-l)} \mathcal{N}(2^l N_1) \\ \lesssim_u \sum_{l \leq k-K} 2^{-(s+2\varepsilon)(k-l)} \mathcal{M}(2^l N_1) + 2^{-(s+2\varepsilon)k},$$

then (5-12) implies (2-1). With a few elementary manipulations, (2-2) implies (5-9).

The last claim follows from Lemma 4.2 after employing $P_{\geq N} = P_{\geq N/2} P_{\geq N}$. □

To estimate the integrals where $|t| \geq \delta$, we break the region of (t, y) integration into two pieces, namely, where $|y| \gtrsim M|t|$ and $|y| \ll M|t|$. The former is the more significant region; it contains the points where the integral kernels $P_M e^{-it\Delta}(x, y)$ and $P_M^\pm e^{-it\Delta}(x, y)$ are large (see Lemmas 2.6 and 4.1). More precisely, when $|x| \leq N^{-1}$, we use (5-7); in this case $|y - x| \sim M|t|$ implies $|y| \gtrsim M|t|$ for $|t| \geq \delta \geq N^{-2}$. (This last condition can be subsumed under our hypothesis N sufficiently large depending on u and η .) When $|x| \geq N^{-1}$, we use (5-6); in this case $|y| - |x| \sim M|t|$ implies $|y| \gtrsim M|t|$.

The next lemma bounds the integrals over the significant region $|y| \gtrsim M|t|$. Let χ_k denote the characteristic function of the set

$$\{(t, y) : 2^k \delta \leq |t| \leq 2^{k+1} \delta, |y| \gtrsim M|t|\}.$$

Lemma 5.4 (Main contribution). *Let $0 < s < 1 + \frac{4}{d}$, let $\eta > 0$ be a small number, and let δ be as in Lemma 5.3. Then*

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \leq \frac{1}{10} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L)$$

for all N sufficiently large depending on u, s , and η . An analogous estimate holds with P_M replaced by $\chi_N P_M^+$ or $\chi_N P_M^-$; moreover, the time integrals may be taken over $[-2^{k+1}\delta, -2^k\delta]$.

Proof. We decompose

$$F(u) = F(u_{\leq \eta M}) + O(|u_{> \eta M}|^{1+\frac{4}{d}}) + O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|). \tag{5-13}$$

We first consider the contribution coming from the last two terms in the decomposition above. By the adjoint Strichartz inequality and Hölder’s inequality,

$$\left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) \tilde{P}_M [O(|u_{> \eta M}|^{\frac{d+4}{d}}) + O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|)](y) dy dt \right\|_{L_x^2} \\ \lesssim \|\chi_k \tilde{P}_M [O(|u_{> \eta M}|^{\frac{d+4}{d}}) + O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|)]\|_{L_t^1 L_y^2} \\ \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{d}} (2^k \delta)^{\frac{d-1}{d}} \left(\left\| |y|^{\frac{2(d-1)}{d}} \tilde{P}_M O(|u_{> \eta M}|^{\frac{d+4}{d}}) \right\|_{L_t^d L_y^2([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right. \\ \left. + \left\| |y|^{\frac{2(d-1)}{d}} \tilde{P}_M O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|) \right\|_{L_t^d L_y^2([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right).$$

As \tilde{P}_M is a Mihlin multiplier and $|y|^{\frac{4(d-1)}{d}}$ is an A_2 weight, \tilde{P}_M is bounded on $L^2(|y|^{\frac{4(d-1)}{d}} dy)$; see [Stein 1993, Chapter V]. Thus, by Hölder's inequality and (5-3),

$$\begin{aligned} & \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) \tilde{P}_M [O(|u_{>\eta M}|^{\frac{d+4}{d}}) + O(|u_{\leq\eta M}|^{\frac{4}{d}} |u_{>\eta M}|)](y) dy dt \right\|_{L_x^2} \\ & \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{d}} (2^k \delta)^{\frac{d-1}{d}} \left(\left\| |y|^{\frac{2(d-1)}{d}} |u_{>\eta M}|^{\frac{d+4}{d}} \right\|_{L_t^d L_y^2([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right. \\ & \qquad \qquad \qquad \left. + \left\| |y|^{\frac{2(d-1)}{d}} |u_{\leq\eta M}|^{\frac{4}{d}} |u_{>\eta M}| \right\|_{L_t^d L_y^2([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right) \\ & \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{d}} (2^k \delta)^{\frac{d-1}{d}} \|u_{>\eta M}\|_{L_t^\infty L_y^2} \left(\left\| |y|^{\frac{d-1}{2}} u_{>\eta M} \right\|_{L_t^4 L_y^\infty([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right. \\ & \qquad \qquad \qquad \left. + \left\| |y|^{\frac{d-1}{2}} u_{\leq\eta M} \right\|_{L_t^4 L_y^\infty([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \right) \\ & \lesssim_u (M 2^k \delta)^{-\frac{2(d-1)}{d}} (2^k \delta)^{\frac{d-1}{d}} \mathcal{M}(\eta N) (2^k \delta)^{\frac{1}{d}}. \end{aligned}$$

Summing first in $k \geq 0$ and then in $M \geq N$, we estimate the contribution of the last two terms on the right-hand side of (5-13) by

$$(N^2 \delta)^{-1 + \frac{1}{d}} \mathcal{M}(\eta N).$$

Next we consider the contribution coming from the first term on the right-hand side of (5-13). By the adjoint of the weighted Strichartz inequality in Lemma 2.8, Hölder's inequality, Corollary 2.5, and Lemma 2.2,

$$\begin{aligned} & \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u_{\leq\eta M}(t))](y) dy dt \right\|_{L_x^2} \\ & \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{q}} \left\| \chi_k \tilde{P}_M F(u_{\leq\eta M}) \right\|_{L_t^{\frac{q}{q-1}} L_y^{\frac{2q}{q+4}}} \\ & \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{q}} (2^k \delta)^{\frac{q-1}{q}} M^{-s} \left\| |\nabla|^s F(u_{\leq\eta M}) \right\|_{L_t^\infty L_y^{\frac{2q}{q+4}}} \\ & \lesssim (M 2^k \delta)^{-\frac{2(d-1)}{q}} (2^k \delta)^{\frac{q-1}{q}} M^{-s} \|u_{\leq\eta M}\|_{L_t^\infty L_y^{\frac{2q}{d}}}^{\frac{4}{d}} \left\| |\nabla|^s u_{\leq\eta M} \right\|_{L_t^\infty L_y^2} \\ & \lesssim_u (M 2^k \delta)^{-\frac{2(d-1)}{q}} (2^k \delta)^{\frac{q-1}{q}} (\eta M)^{\frac{2(q-d)}{q}} \sum_{L \leq \eta M} \left(\frac{L}{M}\right)^s \mathcal{M}(L) \\ & \lesssim_u (M^2 2^k \delta)^{-\frac{2d-q-1}{q}} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L) \end{aligned}$$

provided $q \geq \max\{d, 4\}$ and $M \geq N$. In order to deduce the last inequality, we used the fact that, for $M \geq N$,

$$\begin{aligned} \sum_{L \leq \eta M} \left(\frac{L}{M}\right)^s \mathcal{M}(L) &\leq \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L) + \sum_{\eta N \leq L \leq \eta M} \left(\frac{L}{M}\right)^s \mathcal{M}(L) \\ &\lesssim \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L) + \eta^s \mathcal{M}(\eta N) \lesssim \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L). \end{aligned} \tag{5-14}$$

Therefore, choosing $q = d + 1$,

$$\begin{aligned} \sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u_{\leq \eta M}(t))](y) dy dt \right\|_{L_x^2} \\ \lesssim (N^2 \delta)^{-\frac{d-2}{d+1}} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L). \end{aligned}$$

Putting everything together we obtain

$$\begin{aligned} \sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \\ \lesssim_u ((N^2 \delta)^{-1+\frac{2}{d}} + (N^2 \delta)^{-1+\frac{1}{d}} + (N^2 \delta)^{-\frac{d-2}{d+1}}) \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L). \end{aligned}$$

Choosing N sufficiently large depending on u, δ , and s (and hence only on u, η , and s), we obtain the desired bound.

The last claim follows from the L_x^2 -boundedness of $\chi_N P^\pm P_M$ (see Lemma 4.2) and the time-reversal symmetry of the argument just presented. \square

We turn now to the region of (t, y) integration where $|y| \ll M|t|$. First, we describe the bounds that we will use for the kernels of the propagators. For $|x| \leq N^{-1}$, $|y| \ll M|t|$, and $|t| \geq \delta \gg N^{-2}$, we have

$$|P_M e^{-it\Delta}(x, y)| \lesssim \frac{1}{(M^2|t|)^{50d}} \frac{M^d}{\langle M(x-y) \rangle^{50d}}; \tag{5-15}$$

this follows from Lemma 2.6 since under these constraints, $|y-x| \ll M|t|$. For $|x| \geq N^{-1}$ and y and t as above,

$$|P_M^\pm e^{-it\Delta}(x, y)| \lesssim \frac{1}{(M^2|t|)^{50d}} \frac{M^d}{\langle Mx \rangle^{\frac{d-1}{2}} \langle My \rangle^{\frac{d-1}{2}} \langle M|x-M|y \rangle^{50d}}, \tag{5-16}$$

by Lemma 4.1. To simplify the bound in (5-16) we used the inequalities $|y-x| \ll M|t|$ and

$$\langle M^2|t| + M|x-M|y \rangle^{-100d} \lesssim (M^2|t|)^{-50d} \langle M|x-M|y \rangle^{-50d}.$$

From (5-15) and (5-16) we see that under the hypotheses set out above,

$$|P_M e^{-it\Delta}(x, y)| + |P_M^\pm e^{-it\Delta}(x, y)| \lesssim \frac{1}{(M^2|t|)^{50d}} K_M(x, y), \tag{5-17}$$

where

$$K_M(x, y) := \frac{M^d}{\langle M(x-y) \rangle^{50d}} + \frac{M^d}{\langle Mx \rangle^{\frac{d-1}{2}} \langle My \rangle^{\frac{d-1}{2}} \langle M|x| - M|y| \rangle^{50d}}.$$

Note that by Schur’s test, this is the kernel of a bounded operator on $L_x^2(\mathbb{R}^d)$.

Let $\tilde{\chi}_k$ denote the characteristic function of the set

$$\{(t, y) : 2^k \delta \leq |t| \leq 2^{k+1} \delta, |y| \ll M|t|\}.$$

Lemma 5.5 (The tail). *Let $0 < s < 1 + \frac{4}{d}$, let $\eta > 0$ be a small number, and let δ be as in Lemma 5.3. Then*

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2|t|)^{50d}} \tilde{\chi}_k(t, y) |\tilde{P}_M F(u(t))|(y) dy dt \right\|_{L_x^2} \leq \frac{1}{10} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L)$$

for all N sufficiently large depending on u, s , and η (in particular, we require $N \gg \delta^{-1/2}$).

Proof. Using Hölder’s inequality, the L^2 -boundedness of the operator with kernel K_M , and Lemma 2.2,

$$\begin{aligned} & \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2|t|)^{50d}} \tilde{\chi}_k(t, y) |\tilde{P}_M F(u(t))|(y) dy dt \right\|_{L_x^2} \\ & \lesssim (M^2 2^k \delta)^{-50d} (2^k \delta)^{\frac{d-2}{d}} M^{\frac{2(d-2)}{d}} \left\| \tilde{P}_M F(u) \right\|_{L_t^{\frac{d}{2}} L_x^{d^2+4d-8} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \\ & \lesssim (M^2 2^k \delta)^{-49d} \left\| \tilde{P}_M F(u) \right\|_{L_t^{\frac{d}{2}} L_x^{d^2+4d-8} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)}. \end{aligned}$$

We decompose

$$F(u) = F(u_{\leq \eta M}) + O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|) + O(|u_{> \eta M}|^{1+\frac{4}{d}}). \tag{5-18}$$

Discarding the projection \tilde{P}_M , we use Hölder and (5-2) to estimate

$$\begin{aligned} & \left\| \tilde{P}_M O(|u_{\leq \eta M}|^{\frac{4}{d}} |u_{> \eta M}|) \right\|_{L_t^{\frac{d}{2}} L_x^{d^2+4d-8} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \\ & \lesssim \left\| u_{\leq \eta M} \right\|_{L_t^2 L_x^{\frac{2d}{d-2}} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)}^{\frac{4}{d}} \left\| u_{> \eta M} \right\|_{L_t^\infty L_x^2} \lesssim_u \langle 2^k \delta \rangle^{\frac{2}{d}} \mathcal{M}(\eta N), \\ & \left\| \tilde{P}_M O(|u_{> \eta M}|^{1+\frac{4}{d}}) \right\|_{L_t^{\frac{d}{2}} L_x^{d^2+4d-8} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \\ & \lesssim \left\| u_{> \eta M} \right\|_{L_t^2 L_x^{\frac{2d}{d-2}} ([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)}^{\frac{4}{d}} \left\| u_{> \eta M} \right\|_{L_t^\infty L_x^2} \lesssim_u \langle 2^k \delta \rangle^{\frac{2}{d}} \mathcal{M}(\eta N). \end{aligned}$$

To estimate the contribution coming from the first term on the right-hand side of (5-18), we use Lemma 2.2, Corollary 2.5 (with $r = d^2/(d-2)$) combined with Hölder’s inequality in the time variable, (5-2),

and (5-14), to estimate

$$\begin{aligned} \|\tilde{P}_M F(u_{\leq \eta M})\|_{L_t^{\frac{d}{2}} L_x^{\frac{2d^2}{d^2+4d-8}}([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} &\lesssim M^{-s} \|\nabla|^s F(u_{\leq \eta M})\|_{L_t^{\frac{d}{2}} L_x^{\frac{2d^2}{d^2+4d-8}}([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \\ &\lesssim M^{-s} \|\nabla|^s u_{\leq \eta M}\|_{L_t^\infty L_x^2} \|u_{\leq \eta M}\|_{L_t^2 L_x^{\frac{2d}{d-2}}([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)} \\ &\lesssim u \langle 2^k \delta \rangle^{\frac{2}{d}} \sum_{L \leq \eta M} \left(\frac{L}{M}\right)^s \mathcal{M}(L) \lesssim u \langle 2^k \delta \rangle^{\frac{2}{d}} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L) \end{aligned}$$

for any $M \geq N$.

Putting everything together, we deduce

$$\left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2 |t|)^{50d}} \tilde{\chi}_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \lesssim u (M^2 2^k \delta)^{-49d} \langle 2^k \delta \rangle^{\frac{2}{d}} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L).$$

Summing over $k \geq 0$ and $M \geq N$, we obtain

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2 |t|)^{50d}} \tilde{\chi}_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \lesssim u (N^2 \delta)^{-49d} \sum_{L \leq \eta N} \left(\frac{L}{N}\right)^s \mathcal{M}(L).$$

The claim follows by choosing N sufficiently large depending on δ, η , and s (and hence only on u, s , and η). □

Combining it all together:

Proof of Proposition 5.2. Naturally, we may bound $\|u_{\geq N}\|_{L^2}$ by separately bounding the L^2 norm on the ball $\{|x| \leq N^{-1}\}$ and on its complement. On the ball, we use (5-7), while outside the ball we use (5-6). Invoking (5-8) and the triangle inequality, we reduce the proof to bounding certain integrals. The integrals over short times were estimated in Lemma 5.3. For $|t| \geq \delta$, we further partition the region of integration into two pieces. The first piece, where $|y| \gtrsim M|t|$, was dealt with in Lemma 5.4. To estimate the remaining piece, $|y| \ll M|t|$, one combines (5-17) and Lemma 5.5. □

6. The double high-to-low frequency cascade

In this section, we use the additional regularity provided by Theorem 5.1 to preclude double high-to-low frequency cascade solutions. We argue as in [Killip et al. 2007].

Proposition 6.1 (Absence of double cascades). *Let $d \geq 3$. There are no nonzero global spherically symmetric solutions to (1-1) that are double high-to-low frequency cascades in the sense of Theorem 1.10.*

Proof. Suppose to the contrary that there is such a solution u . By Theorem 5.1, u lies in $C_t^0 H_x^1(\mathbb{R} \times \mathbb{R}^d)$. Hence the energy

$$E(u) = E(u(t)) = \int_{\mathbb{R}^d} \frac{1}{2} |\nabla u(t, x)|^2 + \mu \frac{d}{2(d+2)} |u(t, x)|^{2(d+2)/d} dx$$

is finite and conserved (see [Cazenave 2003], for example). Since $M(u) < M(Q)$ in the focusing case, the sharp Gagliardo–Nirenberg inequality (Theorem 1.6) gives

$$\|\nabla u(t)\|_{L^2_x(\mathbb{R}^d)}^2 \sim_u E(u) \sim_u 1 \tag{6-1}$$

for all $t \in \mathbb{R}$. We will now reach a contradiction by proving that $\|\nabla u(t)\|_2 \rightarrow 0$ along any sequence where $N(t) \rightarrow 0$. The existence of two such time sequences is guaranteed by the fact that u is a double high-to-low frequency cascade.

Let $\eta > 0$ be arbitrary. By Definition 1.8, we can find $C = C(\eta, u) > 0$ such that

$$\int_{|\xi| \geq CN(t)} |\hat{u}(t, \xi)|^2 d\xi \leq \eta^2$$

for all t . Meanwhile, by Theorem 5.1, $u \in C_t^0 H_x^s(\mathbb{R} \times \mathbb{R}^d)$ for some $s > 1$. Thus,

$$\int_{|\xi| \geq CN(t)} |\xi|^{2s} |\hat{u}(t, \xi)|^2 d\xi \lesssim_u 1$$

for all t and some $s > 1$. Thus, by Hölder’s inequality,

$$\int_{|\xi| \geq CN(t)} |\xi|^2 |\hat{u}(t, \xi)|^2 d\xi \lesssim_u \eta^{2(s-1)/s}.$$

On the other hand, from mass conservation and Plancherel’s theorem we have

$$\int_{|\xi| \leq CN(t)} |\xi|^2 |\hat{u}(t, \xi)|^2 d\xi \lesssim_u C^2 N(t)^2.$$

Summing these last two bounds and using Plancherel’s theorem again, we obtain

$$\|\nabla u(t)\|_{L^2_x(\mathbb{R}^d)} \lesssim_u \eta^{(s-1)/s} + CN(t)$$

for all t . As $\eta > 0$ is arbitrary and there exists a sequence of times $t_n \rightarrow \infty$ such that $N(t_n) \rightarrow 0$ (u is a double high-to-low frequency cascade), we conclude $\|\nabla u(t_n)\|_2 \rightarrow 0$. This contradicts (6-1). \square

Remark. As mentioned in [Killip et al. 2007], the argument presented can be used to rule out nonradial single-sided cascade solutions that lie in $C_t^0 H_x^s$ for some $s > 1$. (By a single-sided cascade we mean a solution with $N(t)$ bounded on a semiinfinite interval, say $[T, \infty)$, with $\liminf_{t \rightarrow \infty} N(t) = 0$.) For such regular solutions u , we may define the total momentum $\int_{\mathbb{R}^d} \text{Im}(\bar{u} \nabla u)$, which is conserved. By a Galilean transformation, we can set this momentum equal to zero; thus

$$\int_{\mathbb{R}^d} \xi |\hat{u}(t, \xi)|^2 d\xi = 0.$$

From this, mass conservation, and the uniform H_x^s bound for some $s > 1$, one can show that $\xi(t) \rightarrow 0$ whenever $N(t) \rightarrow 0$. On the other hand, a modification of the above argument gives

$$1 \sim_u \|\nabla u(t)\|_2 \lesssim \eta^{(s-1)/s} + C(N(t) + |\xi(t)|),$$

which is absurd.

7. Death of a soliton

In this section, we use the additional regularity proved in Theorem 5.1 to rule out the third and final enemy, the soliton-like solution. Once again, we follow [Killip et al. 2007]; the method is similar to that in [Kenig and Merle 2006a]. Let

$$M_R(t) := 2 \operatorname{Im} \int_{\mathbb{R}^d} \psi\left(\frac{|x|}{R}\right) \bar{u}(t, x) x \cdot \nabla u(t, x) \, dx,$$

where ψ is a smooth function obeying

$$\psi(r) = \begin{cases} 1 & \text{if } r \leq 1, \\ 0 & \text{if } r \geq 2, \end{cases}$$

and R denotes a radius to be chosen momentarily. For solutions u to (1-1) belonging to $C_t^0 H_x^1$, $M_R(t)$ is a well-defined function. Indeed,

$$|M_R(t)| \lesssim R \|u(t)\|_2 \|\nabla u(t)\|_2 \lesssim_u R.$$

An oft-repeated calculation (essentially that in the derivation of the Morawetz and virial identities) gives:

Lemma 7.1.

$$\partial_t M_R(t) = 8E(u(t))$$

$$- \int_{\mathbb{R}^d} \left(\frac{d^2-1}{R|x|} \psi'\left(\frac{|x|}{R}\right) + \frac{2d+1}{R^2} \psi''\left(\frac{|x|}{R}\right) + \frac{|x|}{R^3} \psi'''\left(\frac{|x|}{R}\right) \right) |u(t, x)|^2 \, dx \tag{7-1}$$

$$+ 4 \int_{\mathbb{R}^d} \left(\psi\left(\frac{|x|}{R}\right) - 1 + \frac{|x|}{R} \psi'\left(\frac{|x|}{R}\right) \right) |\nabla u(t, x)|^2 \, dx \tag{7-2}$$

$$+ \frac{4\mu}{d+2} \int_{\mathbb{R}^d} \left(d \left(\psi\left(\frac{|x|}{R}\right) - 1 \right) + \frac{|x|}{R} \psi'\left(\frac{|x|}{R}\right) \right) |u(t, x)|^{\frac{2(d+2)}{d}} \, dx, \tag{7-3}$$

where $E(u)$ is the energy of u as defined in (1-3).

Proposition 7.2 (Absence of solitons). *Let $d \geq 3$. There are no nonzero global spherically symmetric solutions to (1-1) that are soliton-like in the sense of Theorem 1.10.*

Proof. Assume to the contrary that there is such a solution u . Then, by Theorem 5.1, $u \in C_t^0 H_x^s$ for some $s > 1$. In particular,

$$|M_R(t)| \lesssim_u R. \tag{7-4}$$

Recall that in the focusing case, $M(u) < M(Q)$. As a consequence, the sharp Gagliardo–Nirenberg inequality (Theorem 1.6) implies that the energy is a positive quantity in the focusing case as well as in the defocusing case. Indeed,

$$E(u) \gtrsim_u \int_{\mathbb{R}^d} |\nabla u(t, x)|^2 \, dx > 0.$$

We will show that for R sufficiently large, (7-1) through (7-3) are small terms compared with $E(u)$. Combining this fact with Lemma 7.1, we conclude $\partial_t M_R(t) \gtrsim E(u) > 0$, which contradicts (7-4).

We first turn our attention to (7-1). This is trivially bounded as

$$|(7-1)| \lesssim_u R^{-2}. \tag{7-5}$$

We now study (7-2) and (7-3). Let $\eta > 0$ be a small number to be chosen later. By Definition 1.8 and the fact that $N(t) = 1$ for all $t \in \mathbb{R}$, if R is sufficiently large depending on u and η , then

$$\int_{|x| \geq \frac{R}{4}} |u(t, x)|^2 dx \leq \eta \quad (7-6)$$

for all $t \in \mathbb{R}$. Let χ denote a smooth cutoff to the region $|x| \geq R/2$, chosen so that $\nabla \chi$ is bounded by R^{-1} and supported where $|x| \sim R$. As $u \in C_t^0 H_x^s$ for some $s > 1$, using interpolation and (7-6), we estimate

$$|(7-2)| \lesssim \|\chi \nabla u(t)\|_2^2 \lesssim \|\nabla(\chi u(t))\|_2^2 + \|u(t) \nabla \chi\|_2^2 \lesssim \|\chi u(t)\|_2^{\frac{2(s-1)}{s}} \|u(t)\|_{H_x^s}^{\frac{2}{s}} + \eta \lesssim u \eta^{\frac{s-1}{s}} + \eta. \quad (7-7)$$

Finally, we are left to consider (7-3). Using the same χ as above together with the Gagliardo–Nirenberg inequality and (7-6),

$$|(7-3)| \lesssim \|\chi u(t)\|_{\frac{2(d+2)}{d}}^{\frac{2(d+2)}{d}} \lesssim \|\chi u(t)\|_2^{\frac{4}{d}} \|\nabla(\chi u(t))\|_2^2 \lesssim u \eta^{\frac{2}{d}}. \quad (7-8)$$

Combining (7-5), (7-7), and (7-8) and choosing η sufficiently small depending on u and R sufficiently large depending on u and η , we obtain

$$|(7-1)| + |(7-2)| + |(7-3)| \leq \frac{1}{100} E(u).$$

This completes the proof of the proposition for the reasons explained in the third paragraph. \square

Acknowledgements

We are grateful to Shuanglin Shao for access to preliminary drafts of his work.

References

- [Bégout and Vargas 2007] P. Bégout and A. Vargas, “Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation”, *Trans. Amer. Math. Soc.* **359**:11 (2007), 5257–5282. MR 2008g:35190 Zbl 05178492
- [Berestycki and Lions 1979] H. Berestycki and P.-L. Lions, “Existence d’ondes solitaires dans des problèmes nonlinéaires du type Klein–Gordon”, *C. R. Acad. Sci. Paris Sér. A-B* **288**:7 (1979), A395–A398. MR 80i:35076 Zbl 0397.35024
- [Blue and Colliander 2006] P. Blue and J. Colliander, “Global well-posedness in Sobolev space implies global existence for weighted L^2 initial data for L^2 -critical NLS”, *Commun. Pure Appl. Anal.* **5**:4 (2006), 691–708. MR 2007f:35265 Zbl 1137.35066
- [Bourgain 1998] J. Bourgain, “Refinements of Strichartz’ inequality and applications to 2D-NLS with critical nonlinearity”, *Internat. Math. Res. Not.* **1998**:5 (1998), 253–283. MR 99f:35184 Zbl 0917.35126
- [Bourgain 1999a] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, American Mathematical Society Colloquium Publications **46**, American Mathematical Society, Providence, RI, 1999. MR 2000h:35147 Zbl 0933.35178
- [Bourgain 1999b] J. Bourgain, “Global wellposedness of defocusing critical nonlinear Schrödinger equation in the radial case”, *J. Amer. Math. Soc.* **12**:1 (1999), 145–171. MR 99e:35208 Zbl 0958.35126
- [Carles 2002] R. Carles, “Critical nonlinear Schrödinger equations with and without harmonic potential”, *Math. Models Methods Appl. Sci.* **12**:10 (2002), 1513–1523. MR 2004b:35298 Zbl 1029.35208
- [Carles and Keraani 2007] R. Carles and S. Keraani, “On the role of quadratic oscillations in nonlinear Schrödinger equations. II. The L^2 -critical case”, *Trans. Amer. Math. Soc.* **359**:1 (2007), 33–62. MR 2008a:35260 Zbl 1115.35119

- [Cazenave 2003] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics **10**, New York University Courant Institute of Mathematical Sciences, New York, 2003. MR 2004j:35266 Zbl 1055.35003
- [Cazenave and Weissler 1989] T. Cazenave and F. B. Weissler, “Some remarks on the nonlinear Schrödinger equation in the subcritical case”, pp. 59–69 in *New methods and results in nonlinear field equations* (Bielefeld, 1987), edited by P. Blanchard et al., Lecture Notes in Phys. **347**, Springer, Berlin, 1989. MR 91c:35136 Zbl 0699.35217
- [Christ and Weinstein 1991] F. M. Christ and M. I. Weinstein, “Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation”, *J. Funct. Anal.* **100**:1 (1991), 87–109. MR 92h:35203 Zbl 0743.35067
- [Colliander et al. 2002] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, “Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation”, *Math. Res. Lett.* **9**:5-6 (2002), 659–682. MR 2003j:35294 Zbl 01930988
- [Colliander et al. 2005] J. Colliander, S. Raynor, C. Sulem, and J. D. Wright, “Ground state mass concentration in the L^2 -critical nonlinear Schrödinger equation below H^1 ”, *Math. Res. Lett.* **12**:2-3 (2005), 357–375. MR 2006e:35300 Zbl 1084.35088
- [Colliander et al. 2007] J. Colliander, M. Grillakis, and N. Tzirakis, “Improved interaction Morawetz inequalities for the cubic nonlinear Schrödinger equation on \mathbb{R}^2 ”, *Int. Math. Res. Not.* **2007**:23 (2007), Art. ID rnm090. Zbl 1142.35085 arXiv math/0703606
- [Colliander et al. 2008a] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, “Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in \mathbb{R}^3 ”, *Ann. of Math. (2)* **167**:3 (2008), 767–865.
- [Colliander et al. 2008b] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, “Resonant decompositions and the I -method for the cubic nonlinear Schrödinger equation on \mathbb{R}^2 ”, *Discrete Contin. Dyn. Syst.* **21**:3 (2008), 665–686. MR 2399431 Zbl 05304859
- [De Silva et al. 2007a] D. De Silva, N. Pavlović, G. Staffilani, and N. Tzirakis, “Global well-posedness and polynomial bounds for the defocusing L^2 -critical nonlinear Schrödinger equation in \mathbb{R} ”, preprint, 2007. arXiv math/0702707
- [De Silva et al. 2007b] D. De Silva, N. Pavlović, G. Staffilani, and N. Tzirakis, “Global well-posedness for the L^2 critical nonlinear Schrödinger equation in higher dimensions”, *Commun. Pure Appl. Anal.* **6**:4 (2007), 1023–1041. MR 2341818 Zbl 05257774
- [Fang and Grillakis 2007] Y. F. Fang and M. G. Grillakis, “On the global existence of rough solutions of the cubic defocusing Schrödinger equation in \mathbf{R}^{2+1} ”, *J. Hyperbolic Differ. Equ.* **4**:2 (2007), 233–257. MR 2008c:35303 Zbl 1122.35132
- [Ginibre and Velo 1992] J. Ginibre and G. Velo, “Smoothing properties and retarded estimates for some dispersive evolution equations”, *Comm. Math. Phys.* **144**:1 (1992), 163–188. MR 93a:35065 Zbl 0762.35008
- [Gradshteyn and Ryzhik 2000] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, 6th ed., Academic Press, San Diego, CA, 2000. Translated from the Russian. MR 2001c:00002 Zbl 0981.65001
- [Kato 1995] T. Kato, “On nonlinear Schrödinger equations. II. H^s -solutions and unconditional well-posedness”, *J. Anal. Math.* **67** (1995), 281–306. MR 98a:35124a Zbl 0848.35124
- [Keel and Tao 1998] M. Keel and T. Tao, “Endpoint Strichartz estimates”, *Amer. J. Math.* **120**:5 (1998), 955–980. MR 2000d:35018 Zbl 0922.35028
- [Kenig and Merle 2006a] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case”, *Invent. Math.* **166**:3 (2006), 645–675. MR 2007g:35232 Zbl 1115.35125
- [Kenig and Merle 2006b] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy critical focusing non-linear wave equation”, preprint, 2006. arXiv math/0610801
- [Keraani 2001] S. Keraani, “On the defect of compactness for the Strichartz estimates of the Schrödinger equations”, *J. Differential Equations* **175**:2 (2001), 353–392. MR 2002j:35281 Zbl 1038.35119
- [Keraani 2006] S. Keraani, “On the blow up phenomenon of the critical nonlinear Schrödinger equation”, *J. Funct. Anal.* **235**:1 (2006), 171–192. MR 2007e:35260 Zbl 1099.35132
- [Killip and Visan 2008] R. Killip and M. Visan, “The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher”, preprint, 2008. arXiv 0804.1018

- [Killip et al. 2007] R. Killip, T. Tao, and M. Visan, “The cubic nonlinear Schrödinger equation in two dimensions with radial data”, preprint, 2007. arXiv 0707.3188
- [Kwong 1989] M. K. Kwong, “Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbf{R}^n ”, *Arch. Rational Mech. Anal.* **105**:3 (1989), 243–266. MR 90d:35015 Zbl 0676.35032
- [Merle 1993] F. Merle, “Determination of blow-up solutions with minimal mass for nonlinear Schrödinger equations with critical power”, *Duke Math. J.* **69**:2 (1993), 427–454. MR 94b:35262 Zbl 0808.35141
- [Merle and Tsutsumi 1990] F. Merle and Y. Tsutsumi, “ L^2 concentration of blow-up solutions for the nonlinear Schrödinger equation with critical power nonlinearity”, *J. Differential Equations* **84**:2 (1990), 205–214. MR 91e:35194 Zbl 0722.35047
- [Merle and Vega 1998] F. Merle and L. Vega, “Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D”, *Internat. Math. Res. Not.* **1998**:8 (1998), 399–425. MR 99d:35156 Zbl 0913.35126
- [Nawa 1999] H. Nawa, “Asymptotic and limiting profiles of blowup solutions of the nonlinear Schrödinger equation with critical power”, *Comm. Pure Appl. Math.* **52**:2 (1999), 193–270. MR 99m:35235 Zbl 0964.37014
- [Ryckman and Visan 2007] E. Ryckman and M. Visan, “Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in \mathbb{R}^{1+4} ”, *Amer. J. Math.* **129**:1 (2007), 1–60. MR 2007k:35474
- [Shao 2007] S. Shao, “Sharp linear and bilinear restriction estimates for paraboloid in the cylindrically symmetric case”, preprint, 2007. arXiv 0706.3759
- [Staffilani 1997] G. Staffilani, “On the generalized Korteweg-de Vries-type equations”, *Differential Integral Equations* **10**:4 (1997), 777–796. MR 2001a:35005 Zbl 0891.35135
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, Princeton, NJ, 1993. MR 95c:42002 Zbl 0821.42001
- [Stein and Weiss 1971] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Mathematical Series **32**, Princeton University Press, Princeton, N.J., 1971. MR 46 #4102 Zbl 0232.42007
- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. MR 58 #23577 Zbl 0372.35001
- [Tao 2005] T. Tao, “Global well-posedness and scattering for the higher-dimensional energy-critical nonlinear Schrödinger equation for radial data”, *New York J. Math.* **11** (2005), 57–80. MR 2006e:35308 Zbl 1119.35092
- [Tao 2006] T. Tao, “A pseudoconformal compactification of the nonlinear Schrödinger equation and applications”, preprint, 2006. arXiv math/0606254
- [Tao 2008a] T. Tao, “Global regularity of wave maps III”, preprint, 2008. arXiv 0805.4666
- [Tao 2008b] T. Tao, “Global regularity of wave maps IV”, preprint, 2008. arXiv 0806.3592
- [Tao 2008c] T. Tao, “Global regularity of wave maps V”, preprint, 2008. arXiv 0808.0368
- [Tao et al. 2006] T. Tao, M. Visan, and X. Zhang, “Minimal-mass blowup solutions of the mass-critical NLS”, preprint, 2006. To appear in Forum Math. arXiv math/0609690
- [Tao et al. 2007] T. Tao, M. Visan, and X. Zhang, “Global well-posedness and scattering for the defocusing mass-critical nonlinear Schrödinger equation for radial data in high dimensions”, *Duke Math. J.* **140**:1 (2007), 165–202. MR 2355070 Zbl 05208403
- [Taylor 2000] M. E. Taylor, *Tools for PDE. Pseudodifferential operators, paradifferential operators, and layer potentials*, Mathematical Surveys and Monographs **81**, American Mathematical Society, Providence, RI, 2000. MR 2001g:35004 Zbl 0963.35211
- [Tsutsumi 1985] Y. Tsutsumi, “Scattering problem for nonlinear Schrödinger equations”, *Ann. Inst. H. Poincaré Phys. Théor.* **43**:3 (1985), 321–347. MR 87g:35221 Zbl 0612.35104
- [Tzirakis 2005] N. Tzirakis, “The Cauchy problem for the semilinear quintic Schrödinger equation in one dimension”, *Differential Integral Equations* **18**:8 (2005), 947–960. MR 2006b:35306 Zbl 1075.35082
- [Visan 2006] M. Visan, *The defocusing energy-critical nonlinear Schrödinger equation in dimensions five and higher*, PhD Thesis, University of California, Los Angeles, 2006.
- [Visan 2007] M. Visan, “The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions”, *Duke Math. J.* **138**:2 (2007), 281–374. MR 2008f:35387 Zbl 1131.35081

- [Visan and Zhang 2007] M. Visan and X. Zhang, “On the blowup for the L^2 -critical focusing nonlinear Schrödinger equation in higher dimensions below the energy class”, *SIAM J. Math. Anal.* **39**:1 (2007), 34–56. MR 2008b:35284 Zbl 1135.35079
- [Weinstein 1983] M. I. Weinstein, “Nonlinear Schrödinger equations and sharp interpolation estimates”, *Comm. Math. Phys.* **87**:4 (1983), 567–576. MR 84d:35140 Zbl 0527.35023
- [Weinstein 1989] M. I. Weinstein, “The nonlinear Schrödinger equation—singularity formation, stability and dispersion”, pp. 213–232 in *The connection between infinite-dimensional and finite-dimensional dynamical systems* (Boulder, CO, 1987), edited by B. Nicolaenko et al., *Contemp. Math.* **99**, Amer. Math. Soc., Providence, RI, 1989. MR 90m:35188 Zbl 0703.35159

Received 11 Aug 2008. Revised 20 Aug 2008. Accepted 23 Sep 2008.

ROWAN KILLIP: killip@math.ucla.edu

520 Portola Plaza, Math Sciences Building 6363, Mailcode: 155505, Los Angeles, CA 90095, United States

<http://www.math.ucla.edu/~killip/>

MONICA VISAN: visan.m@gmail.com

Department of Mathematics, University of Chicago, 5734 University Avenue, Chicago, IL 60637-1514, United States

<http://www.math.uchicago.edu/~mvisan/>

XIAOYI ZHANG: zh.xiaoyi@gmail.com

School of Mathematics, Institute for Advanced Study, One Einstein Drive, Princeton, NJ 08540, United States