

# ANALYSIS & PDE

Volume 2

No. 3

2009

TERENCE TAO

**GLOBAL REGULARITY FOR A LOGARITHMICALLY  
SUPERCRITICAL HYPERDISSIPATIVE NAVIER-STOKES  
EQUATION**



mathematical sciences publishers

# GLOBAL REGULARITY FOR A LOGARITHMICALLY SUPERCRITICAL HYPERDISSIPATIVE NAVIER–STOKES EQUATION

TERENCE TAO

Let  $d \geq 3$ . We consider the global Cauchy problem for the generalized Navier–Stokes system

$$\partial_t u + (u \cdot \nabla)u = -D^2 u - \nabla p, \quad \nabla \cdot u = 0, \quad u(0, x) = u_0(x)$$

for  $u : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  and  $p : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ , where  $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is smooth and divergence free, and  $D$  is a Fourier multiplier whose symbol  $m : \mathbb{R}^d \rightarrow \mathbb{R}^+$  is nonnegative; the case  $m(\xi) = |\xi|$  is essentially Navier–Stokes. It is folklore that one has global regularity in the critical and subcritical hyperdissipation regimes  $m(\xi) = |\xi|^\alpha$  for  $\alpha \geq (d+2)/4$ . We improve this slightly by establishing global regularity under the slightly weaker condition that  $m(\xi) \geq |\xi|^{(d+2)/4}/g(|\xi|)$  for all sufficiently large  $\xi$  and some nondecreasing function  $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\int_1^\infty ds/(sg(s)^4) = +\infty$ . In particular, the results apply for the logarithmically supercritical dissipation  $m(\xi) := |\xi|^{(d+2)/4}/\log(2 + |\xi|^2)^{1/4}$ .

## 1. Introduction

Let  $d \geq 3$ . This note is concerned with solutions to the generalised Navier–Stokes system

$$\begin{aligned} \partial_t u + (u \cdot \nabla)u &= -D^2 u - \nabla p, \\ \nabla \cdot u &= 0, \\ u(0, x) &= u_0(x), \end{aligned} \tag{1}$$

where  $u : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,  $p : \mathbb{R}^+ \times \mathbb{R}^d \rightarrow \mathbb{R}$  are smooth, and  $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is smooth, compactly supported, and divergence-free, and  $D$  is a Fourier multiplier<sup>1</sup> whose symbol  $m : \mathbb{R}^d \rightarrow \mathbb{R}^+$  is nonnegative; the case  $m(\xi) = |\xi|$  is essentially the Navier–Stokes system, while the case  $m = 0$  is the Euler system.

For  $d \geq 3$ , the global regularity of the Navier–Stokes system is of course a notoriously difficult unsolved problem, due in large part to the supercritical nature of the equation with respect to the energy  $E(u(t)) := \int_{\mathbb{R}^d} |u(t, x)|^2 dx$ . This supercriticality can be avoided by strengthening the dissipative symbol  $m(\xi)$ , for instance setting  $m(\xi) := |\xi|^\alpha$  for some  $\alpha > 1$ . This *hyper-dissipative* variant of the Navier–Stokes equation becomes subcritical for  $\alpha > (d+2)/4$  (and critical for  $\alpha = (d+2)/4$ ), and it is known that global regularity can be recovered in these cases; see [Katz and Pavlović 2002] for further discussion. For  $1 \leq \alpha < (d+2)/4$ , only partial regularity results are known; see [Caffarelli et al. 1982] for the case  $\alpha = 1$  and [Katz and Pavlović 2002] for the case  $\alpha > 1$ .

MSC2000: 35Q30.

Keywords: Navier–Stokes, energy method.

The author is supported by NSF Research Award DMS-0649473, the NSF Waterman award and a grant from the MacArthur Foundation.

<sup>1</sup>The exact definition of the Fourier transform is inessential here, but for concreteness, take  $\hat{f}(\xi) := \int_{\mathbb{R}^d} f(x)e^{-ix \cdot \xi} dx$ .



The purpose of this note is to extend the global regularity result very slightly into the supercritical regime.

**Theorem 1.1.** *Suppose that  $m$  obeys the lower bound*

$$m(\xi) \geq |\xi|^{(d+2)/4} / g(|\xi|) \tag{2}$$

for all sufficiently large  $|\xi|$ , where  $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a nondecreasing function such that

$$\int_1^\infty \frac{ds}{sg(s)^4} = \infty. \tag{3}$$

Then for any smooth, compactly supported initial data  $u_0$ , one has a global smooth solution to (1).

Note that the hypotheses are for instance satisfied when

$$m(\xi) := |\xi|^{(d+2)/4} / \log^{1/4}(2 + |\xi|^2); \tag{4}$$

thus

$$|D|^2 = \frac{(-\Delta)^{(d+2)/4}}{\log^{1/2}(2 - \Delta)}.$$

Analogous “barely supercritical” global regularity results were established for the nonlinear wave equation recently [Tao 2007; Roy 2008; 2009].

The argument is quite simple, being based on the classical energy method and Sobolev embedding. The basic point is that whereas in the critical and subcritical cases one can get an energy inequality of the form

$$\partial_t \|u(t)\|_{H^k(\mathbb{R}^d)}^2 \leq Ca(t) \|u(t)\|_{H^k(\mathbb{R}^d)}^2$$

for some locally integrable function  $a(t)$  of time, a constant  $C$ , and some large  $k$ , which by Gronwall’s inequality is sufficient to establish a suitable *a priori* bound, in the logarithmically supercritical case (4) one instead obtains the slightly weaker inequality

$$\partial_t \|u(t)\|_{H^k(\mathbb{R}^d)}^2 \leq Ca(t) \|u(t)\|_{H^k(\mathbb{R}^d)}^2 \log(2 + \|u(t)\|_{H^k(\mathbb{R}^d)})$$

(thanks to an endpoint version of Sobolev embedding, closely related to an inequality of Brézis and Wainger [1980]), which is still sufficient to obtain an *a priori* bound (though one which is now double-exponential rather than single-exponential; compare [Beale et al. 1984]).

**Remark 1.2.** It may well be that the condition (3) can be relaxed further by a more sophisticated argument. Indeed, the following heuristic suggests that one should be able to weaken (3) to

$$\int_1^\infty \frac{ds}{sg(s)^2} = \infty,$$

thus allowing one to increase the  $\frac{1}{4}$  exponent in (4) to  $\frac{1}{2}$ . Consider a blowup scenario in which the solution blows up at some finite time  $T_*$ , and is concentrated on a ball of radius  $1/N(t)$  for times  $0 < t < T_*$ , where  $N(t) \rightarrow \infty$  as  $t \rightarrow T_*$ . As the energy of the fluid must stay bounded, we obtain the heuristic

bound  $u(t) = O(N(t)^{d/2})$  for times  $0 < t < T_*$ . In particular, we expect the fluid to propagate at speeds  $O(N(t)^{d/2})$ , leading to the heuristic ODE

$$\frac{d}{dt} \frac{1}{N(t)} = O(N(t)^{d/2})$$

for the radius  $1/N(t)$  of the fluid. Solving this ODE, we are led to a heuristic upper bound  $N(t) = O((T_* - t)^{2/(d+2)})$  on the blowup rate. On the other hand, from the energy inequality

$$2 \int_0^{T_*} \int_{\mathbb{R}^d} |Du(t, x)|^2 dx dt \leq \int_{\mathbb{R}^d} |u_0(x)|^2 dx$$

one is led to the heuristic bound

$$\int_0^{T_*} \frac{1}{N(t)^{(d+2)/2} g(N(t))^2} dt < \infty.$$

This is incompatible with the upper bound  $N(t) = O((T_* - t)^{2/(d+2)})$  if  $\int_1^{T_*} ds/(sg(s)^2) = \infty$ . Unfortunately the author was not able to make this argument precise, as there appear to be multiple and inequivalent ways to rigorously define an analogue of the “frequency scale”  $N(t)$ , and all attempts of the author to equate different versions of these analogues lost one or more powers of  $g(s)$ .

To go beyond the barrier  $\int_1^\infty ds/(sg(s)^2) = \infty$  (with the aim of getting closer to the Navier–Stokes regime, in which  $g(s) = s^{1/4}$  in three dimensions), the heuristic analysis above suggests that one would need to force the energy to not concentrate into small balls, but instead to exhibit turbulent behaviour.

## 2. Proof of theorem

We now prove [Theorem 1.1](#). Let  $k$  be a large integer (for example,  $k := 100d$  will suffice).

Standard energy method arguments (see, for example, [\[Kato 1985\]](#)) show that if the initial data is smooth and compactly supported, then either a smooth  $H^\infty$  solution exists for all time, or there exists a smooth solution up to some blowup time  $0 < T_* < \infty$ , and  $\|u(t)\|_{H^k(\mathbb{R}^d)} \rightarrow \infty$  as  $t \rightarrow T_*$ . Thus, to establish global regularity, it suffices to prove an *a priori bound* of the form

$$\|u(t)\|_{H^k(\mathbb{R}^d)} \leq C(k, d, \|u_0\|_{H^k(\mathbb{R}^d)}, T, g)$$

for all  $0 \leq t \leq T < \infty$  and all smooth  $H^\infty$  solutions  $u : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  to [\(1\)](#), where the quantity  $C(k, d, \|u_0\|_{H^k(\mathbb{R}^d)}, T, g)$  only depends on  $k, d, \|u_0\|_{H^k(\mathbb{R}^d)}, T$ , and  $g$ .

We now fix  $u_0, u, T$ , and let  $C$  denote any constant depending on  $k, d, \|u_0\|_{H^k(\mathbb{R}^d)}, T$ , and  $g$  (whose value can vary from line to line). Multiplying the Navier–Stokes equation by  $u$  and integrating by parts, we obtain the well-known energy identity

$$\partial_t \int_{\mathbb{R}^d} |u(t, x)|^2 dx = -2a(t),$$

where

$$a(t) := \|Du\|_{L^2(\mathbb{R}^d)}^2 \tag{5}$$

(note that the pressure term  $\nabla p$  disappears thanks to the divergence free condition); integrating this in time, we obtain the energy dissipation bound

$$\int_0^T a(t) dt \leq C. \tag{6}$$

Now, we consider the higher energy

$$E_k(t) := \sum_{j=0}^k \int_{\mathbb{R}^d} |\nabla^j u(t, x)|^2 dx. \tag{7}$$

Differentiating (7) in time and integrating by parts, we obtain

$$\partial_t E_k(t) = -2 \sum_{j=0}^k \|\nabla^j Du(t)\|_{L^2(\mathbb{R}^d)}^2 - 2 \sum_{j=0}^k \int_{\mathbb{R}^d} \nabla^j u(t, x) \cdot \nabla^j ((u \cdot \nabla)u)(t, x) dx;$$

again, the pressure term disappears thanks to the divergence-free condition. For brevity we shall now drop explicit mention of the  $t$  and  $x$  variables.

We apply the Leibniz rule to  $\nabla^j((u \cdot \nabla)u)$ . There is one term involving  $(j+1)$ -st derivatives of  $u$ , but the contribution of that term vanishes by integration by parts and the divergence free property. The remaining terms give contributions of the form

$$\sum_{j=0}^k \sum_{\substack{1 \leq j_1, j_2 \leq j \\ j_1 + j_2 = j+1}} \int_{\mathbb{R}^d} \mathbb{O}(\nabla^j u \nabla^{j_1} u \nabla^{j_2} u) dx,$$

where  $\mathbb{O}(\nabla^j u \nabla^{j_1} u \nabla^{j_2} u)$  denotes some constant-coefficient trilinear combination of the components of  $\nabla^j u$ ,  $\nabla^{j_1} u$ , and  $\nabla^{j_2} u$  whose explicit form is easily computed, but is not of importance to our argument. We can integrate by parts using  $D$  and  $D^{-1}$  and then use Cauchy–Schwarz to obtain the bound

$$\int_{\mathbb{R}^d} \mathbb{O}(\nabla^j u \nabla^{j_1} u \nabla^{j_2} u) dx \leq \|(1 + D)\nabla^j u\|_{L^2(\mathbb{R}^d)} \|(1 + D)^{-1}(\mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u))\|_{L^2(\mathbb{R}^d)}.$$

By the arithmetic mean-geometric mean inequality we then have

$$\int_{\mathbb{R}^d} \mathbb{O}(\nabla^j u \nabla^{j_1} u \nabla^{j_2} u) dx \leq c \|(1 + D)\nabla^j u\|_{L^2(\mathbb{R}^d)}^2 + \frac{1}{c} \|(1 + D)^{-1}(\mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u))\|_{L^2(\mathbb{R}^d)}^2$$

for any  $c > 0$ . Finally, from the triangle inequality, (7), and the fact that  $D$  commutes with  $\nabla^j$ , we have

$$\|(1 + D)\nabla^j u\|_{L^2(\mathbb{R}^d)}^2 \leq C (\|\nabla^j Du\|_{L^2(\mathbb{R}^d)}^2 + E_k).$$

Putting this all together and choosing  $c$  small enough, we conclude that

$$\partial_t E_k \leq C E_k + C \sum_{\substack{1 \leq j_1 \leq j_2 \leq k \\ j_1 + j_2 \leq k+1}} \|(1 + D)^{-1}(\mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u))\|_{L^2(\mathbb{R}^d)}^2. \tag{8}$$

To estimate this expression, we introduce a parameter  $N > 1$  (depending on  $t$ ) to be optimised later, and write

$$(1 + D)^{-1} = (1 + D)^{-1} P_{\leq N} + (1 + D)^{-1} P_{> N},$$

where  $P_{\leq N}$  and  $P_{> N}$  are the Fourier projections to the regions  $\{\zeta : |\zeta| \leq N\}$  and  $\{\zeta : |\zeta| > N\}$ .

We first deal with the low-frequency contribution to (8). From Plancherel’s theorem and (2) we obtain

$$\|(1 + D)^{-1} P_{\leq N}(\mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u))\|_{L^2(\mathbb{R}^d)} \leq Cg(N) \|\langle \nabla \rangle^{-(d+2)/4} \mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u)\|_{L^2(\mathbb{R}^d)},$$

where  $\langle \nabla \rangle^{-(d+2)/4}$  is the Fourier multiplier with symbol  $\langle \zeta \rangle^{-(d+2)/4}$ , where  $\langle \zeta \rangle := (1 + |\zeta|^2)^{1/2}$ . Applying Sobolev embedding, we can bound the right-hand side by

$$\leq Cg(N) \|\nabla^{j_1} u\|_{L^{4d/(3d+2)}(\mathbb{R}^d)} \|\nabla^{j_2} u\|_{L^{4d/(3d+2)}(\mathbb{R}^d)}.$$

By Hölder’s inequality and the Gagliardo–Nirenberg inequality, we can bound this by

$$\leq Cg(N) \|\nabla u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} \|\nabla^{j_1+j_2-1} u\|_{L^2(\mathbb{R}^d)},$$

which by (7) is bounded by

$$\leq Cg(N) \|\nabla u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} E_k^{1/2}.$$

Next, we partition

$$\|\nabla u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} \leq \|\nabla P_{\leq N} u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} + \|\nabla P_{> N} u\|_{L^{4d/(d+2)}(\mathbb{R}^d)}.$$

From Sobolev embedding and Plancherel, together with (2) and (5), we have

$$\|\nabla P_{\leq N} u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} \leq C \|\langle \nabla \rangle^{(d+2)/4} P_{\leq N} u\|_{L^2(\mathbb{R}^d)} \leq Cg(N)(1 + a(t))^{1/2}.$$

Meanwhile, from Sobolev embedding we have

$$\|\nabla P_{> N} u\|_{L^{4d/(d+2)}(\mathbb{R}^d)} \leq \frac{1}{N} E_k^{1/2},$$

(say) if  $k$  is large enough. Putting this all together, we see that the low-frequency contribution to (8) is

$$\leq Cg(N)^2 E_k \left[ g(N)^2 (1 + a(t)) + \frac{1}{N^2} E_k \right].$$

Next, we turn to the high-frequency contribution to (8). From Plancherel, Hölder’s inequality, and (7) we have

$$\begin{aligned} \|(1 + D)^{-1} P_{\geq N}(\mathbb{O}(\nabla^{j_1} u \nabla^{j_2} u))\|_{L^2(\mathbb{R}^d)} &\leq Cg(N) N^{-(d+2)/4} \|\nabla^{j_1} u\|_{L^2(\mathbb{R}^d)} \|\nabla^{j_2} u\|_{L^2(\mathbb{R}^d)} \\ &\leq Cg(N) N^{-(d+2)/4} \|\nabla^{j_1} u\|_{L^\infty(\mathbb{R}^d)} E_k^{1/2}, \end{aligned}$$

while from Sobolev embedding and (7) we see (for  $k$  large enough) that

$$\|\nabla^{j_1} u\|_{L^\infty(\mathbb{R}^d)} \leq C E_k^{1/2}.$$

Thus the high-frequency contribution to (8) is at most  $Cg(N)^2 N^{-(d+2)/2} E_k^2$ .

Putting this all together, we conclude that

$$\partial_t E_k \leq Cg(N)^2 E_k \left[ g(N)^2 (1 + a(t)) + \frac{1}{N} E_k \right].$$

We now optimize in  $N$ , setting  $N := 1 + E_k$ , to obtain

$$\partial_t E_k \leq Cg(1 + E_k)^4 E_k (1 + a(t)).$$

From (6), (3) and separation of variables we see that the ODE

$$\partial_t E = Cg(1 + E)^4 E(1 + a(t))$$

with initial data  $E(0) \geq 0$  does not blow up in time. Also, from (7) we have  $E_k(0) \leq C$ . A standard ODE comparison (or continuity) argument then shows that  $E_k(t) \leq C(T)$  for all  $0 \leq t \leq T$ , and the claim follows.

**Remark 2.1.** It should be clear to the experts that the domain  $\mathbb{R}^d$  here could be replaced by any other sufficiently smooth domain, for example, the torus  $\mathbb{R}^d/\mathbb{Z}^d$ , using standard substitutes for the Littlewood–Paley type operators  $P_{\leq N}$ ,  $P_{> N}$  (one could use spectral projections of the Laplacian). We omit the details.

### References

- [Beale et al. 1984] J. T. Beale, T. Kato, and A. Majda, “Remarks on the breakdown of smooth solutions for the 3-D Euler equations”, *Comm. Math. Phys.* **94**:1 (1984), 61–66. [MR 85j:35154](#) [Zbl 0573.76029](#)
- [Brézis and Wainger 1980] H. Brézis and S. Wainger, “A note on limiting cases of Sobolev embeddings and convolution inequalities”, *Comm. Partial Differential Equations* **5**:7 (1980), 773–789. [MR 81k:46028](#) [Zbl 0437.35071](#)
- [Caffarelli et al. 1982] L. Caffarelli, R. Kohn, and L. Nirenberg, “Partial regularity of suitable weak solutions of the Navier–Stokes equations”, *Comm. Pure Appl. Math.* **35**:6 (1982), 771–831. [MR 84m:35097](#) [Zbl 0509.35067](#)
- [Kato 1985] T. Kato, *Abstract differential equations and nonlinear mixed problems*, Scuola Normale Superiore, Pisa, 1985. [MR 88m:34058](#)
- [Katz and Pavlović 2002] N. H. Katz and N. Pavlović, “A cheap Caffarelli–Kohn–Nirenberg inequality for the Navier–Stokes equation with hyper-dissipation”, *Geom. Funct. Anal.* **12**:2 (2002), 355–379. [MR 2003e:35243](#) [Zbl 0999.35069](#)
- [Roy 2008] T. Roy, “Global existence of smooth solutions of a 3D loglog energy-supercritical wave equation”, preprint, 2008. [arXiv 0810.5175](#)
- [Roy 2009] T. Roy, “One remark on barely  $\dot{H}^{s_p}$  supercritical wave equations”, preprint, 2009. [arXiv 0906.0044](#)
- [Tao 2007] T. Tao, “Global regularity for a logarithmically supercritical defocusing nonlinear wave equation for spherically symmetric data”, *J. Hyperbolic Differ. Equ.* **4**:2 (2007), 259–265. [MR 2009b:35294](#) [Zbl 1124.35043](#)

Received 16 Jun 2009. Revised 22 Sep 2009. Accepted 23 Oct 2009.

TERENCE TAO: [tao@math.ucla.edu](mailto:tao@math.ucla.edu)

University of California, Department of Mathematics, Los Angeles, CA 90095-1555, United States

<http://www.math.ucla.edu/~tao/>

# Analysis & PDE

[pjm.math.berkeley.edu/apde](http://pjm.math.berkeley.edu/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
University of California  
Berkeley, USA

BOARD OF EDITORS

Michael Aizenman	Princeton University, USA <a href="mailto:aizenman@math.princeton.edu">aizenman@math.princeton.edu</a>	Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>
Luis A. Caffarelli	University of Texas, USA <a href="mailto:caffarel@math.utexas.edu">caffarel@math.utexas.edu</a>	Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Nigel Higson	Pennsylvania State University, USA <a href="mailto:higson@math.psu.edu">higson@math.psu.edu</a>
Vaughan Jones	University of California, Berkeley, USA <a href="mailto:vfr@math.berkeley.edu">vfr@math.berkeley.edu</a>	Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>	Steven Zelditch	Johns Hopkins University, USA <a href="mailto:szelditch@math.jhu.edu">szelditch@math.jhu.edu</a>

## PRODUCTION

[apde@mathscipub.org](mailto:apde@mathscipub.org)

Paulo Ney de Souza, Production Manager

Sheila Newbery, Production Editor

Silvio Levy, Senior Production Editor

---

See inside back cover or [pjm.math.berkeley.edu/apde](http://pjm.math.berkeley.edu/apde) for submission instructions.

---


The subscription price for 2009 is US \$120/year for the electronic version, and \$180/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

---

Analysis & PDE, at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer-review and production is managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY  
 **mathematical sciences publishers**  
<http://www.mathscipub.org>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2009 by Mathematical Sciences Publishers



# ANALYSIS & PDE

Volume 2    No. 3    2009

---

Global existence of smooth solutions of a 3D log-log energy-supercritical wave equation	261
TRISTAN ROY	
Periodic stochastic Korteweg–de Vries equation with additive space-time white noise	281
TADAHIRO OH	
Stability for strongly coupled critical elliptic systems in a fully inhomogeneous medium	305
OLIVIER DRUET and EMMANUEL HEBEY	
Global regularity for a logarithmically supercritical hyperdissipative Navier–Stokes equation	361
TERENCE TAO	