

ANALYSIS & PDE

Volume 6

No. 4

2013

THIERRY GALLAY AND YASUNORI MAEKAWA

**LONG-TIME ASYMPTOTICS FOR TWO-DIMENSIONAL
EXTERIOR FLOWS WITH SMALL CIRCULATION AT INFINITY**

LONG-TIME ASYMPTOTICS FOR TWO-DIMENSIONAL EXTERIOR FLOWS WITH SMALL CIRCULATION AT INFINITY

THIERRY GALLAY AND YASUNORI MAEKAWA

We consider the incompressible Navier–Stokes equations in a two-dimensional exterior domain Ω , with no-slip boundary conditions. Our initial data are of the form $u_0 = \alpha\Theta_0 + v_0$, where Θ_0 is the Oseen vortex with unit circulation at infinity and v_0 is a solenoidal perturbation belonging to $L^2(\Omega)^2 \cap L^q(\Omega)^2$ for some $q \in (1, 2)$. If $\alpha \in \mathbb{R}$ is sufficiently small, we show that the solution behaves asymptotically in time like the self-similar Oseen vortex with circulation α . This is a global stability result, in the sense that the perturbation v_0 can be arbitrarily large, and our smallness assumption on the circulation α is independent of the domain Ω .

1. Introduction

Let $\Omega \subset \mathbb{R}^2$ be a smooth exterior domain, namely an unbounded connected open subset of the Euclidean plane with a smooth compact boundary $\partial\Omega$. We consider the free motion of an incompressible viscous fluid in Ω , with no-slip boundary conditions on $\partial\Omega$. The evolution is governed by the Navier–Stokes equations

$$\begin{cases} \partial_t u + (u \cdot \nabla)u = \Delta u - \nabla p, & \operatorname{div} u = 0 & \text{for } x \in \Omega, \quad t > 0, \\ u(x, t) = 0 & & \text{for } x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0(x) & & \text{for } x \in \Omega, \end{cases} \quad (1)$$

where $u(x, t) \in \mathbb{R}^2$ denotes the velocity of a fluid particle at point $x \in \Omega$ and time $t > 0$, and $p(x, t)$ is the pressure in the fluid at the same point. For simplicity, both the kinematic viscosity and the density of the fluid have been normalized to 1. The initial velocity field $u_0 : \Omega \rightarrow \mathbb{R}^2$ is assumed to be divergence-free and tangent to the boundary on $\partial\Omega$.

If the initial velocity u_0 belongs to the energy space

$$L^2_\sigma(\Omega) = \{u \in L^2(\Omega)^2 \mid \operatorname{div} u = 0 \text{ in } \Omega, \quad u \cdot n = 0 \text{ on } \partial\Omega\},$$

where n denotes the unit normal on $\partial\Omega$, then it is known that system (1) has a unique global solution $u \in C^0([0, \infty); L^2_\sigma(\Omega)) \cap C^1((0, \infty); L^2_\sigma(\Omega)) \cap C^0((0, \infty); H^1_0(\Omega)^2 \cap H^2(\Omega)^2)$, which satisfies the energy equality

$$\frac{1}{2} \|u(\cdot, t)\|_{L^2(\Omega)}^2 + \int_0^t \|\nabla u(\cdot, s)\|_{L^2(\Omega)}^2 \, ds = \frac{1}{2} \|u_0\|_{L^2(\Omega)}^2 \quad \text{for all } t > 0.$$

The authors are partially supported by the ANR project PREFERED (Gallay) and by the Grant-in-Aid for Young Scientists (B) 22740090 (Maekawa).

MSC2010: 35B35, 35Q30, 76D05, 76D17.

Keywords: Navier–Stokes equations, exterior domains, long-time asymptotics, Oseen vortices.

This global well-posedness result was first established by Leray [1933] in the particular case where $\Omega = \mathbb{R}^2$, and subsequently extended to more general domains, including exterior domains, by various authors [Leray 1934; Ladyženskaja 1959; Lions and Prodi 1959; Kato and Fujita 1962; Fujita and Kato 1964; Kozono and Ogawa 1993b]. It is also known that the kinetic energy $\frac{1}{2} \|u(\cdot, t)\|_{L^2(\Omega)}^2$ converges to zero as $t \rightarrow \infty$ [Masuda 1984; Borchers and Miyakawa 1992; Kozono and Ogawa 1993b], and precise decay rates can be obtained under additional assumptions on the initial data [Kozono and Ogawa 1993a; He and Miyakawa 2006; Bae and Jin 2006].

In two-dimensional fluid mechanics, however, the assumption that the velocity field u be square integrable is quite restrictive, because it implies (if $u = 0$ on $\partial\Omega$) that the associated vorticity field $\omega = \partial_1 u_2 - \partial_2 u_1$ has zero mean over Ω ; see [Majda and Bertozzi 2002, Section 3.1.3]. In many important examples, this condition is not satisfied and the kinetic energy of the flow is therefore infinite. For instance, when $\Omega = \mathbb{R}^2$, the Navier–Stokes equations (1) have a family of explicit self-similar solutions of the form $u(x, t) = \alpha \Theta(x, t)$, $p(x, t) = \alpha^2 \Pi(x, t)$, where $\alpha \in \mathbb{R}$ is a parameter and

$$\Theta(x, t) = \frac{1}{2\pi} \frac{x^\perp}{|x|^2} \left(1 - e^{-\frac{|x|^2}{4(1+t)}}\right), \quad \nabla \Pi(x, t) = \frac{x}{|x|^2} |\Theta(x, t)|^2. \tag{2}$$

Here and in the sequel, if $x = (x_1, x_2) \in \mathbb{R}^2$, we define $x^\perp = (-x_2, x_1)$ and $|x|^2 = x_1^2 + x_2^2$. The solution (2) is called the *Lamb–Oseen vortex* with circulation α . Remark that $|\Theta(x, t)| = \mathcal{O}(|x|^{-1})$ as $|x| \rightarrow \infty$, so that $\Theta(\cdot, t) \notin L^2(\mathbb{R}^2)^2$, and that the circulation at infinity of the vector field Θ is equal to 1, in the sense that $\oint_{|x|=R} \Theta_1 dx_1 + \Theta_2 dx_2 \rightarrow 1$ as $R \rightarrow \infty$. The corresponding vorticity distribution

$$\Xi(x, t) = \partial_1 \Theta_2(x, t) - \partial_2 \Theta_1(x, t) = \frac{1}{4\pi(1+t)} e^{-\frac{|x|^2}{4(1+t)}} \tag{3}$$

has a constant sign and satisfies $\int_{\mathbb{R}^2} \Xi(x, t) dx = 1$ for all $t \geq 0$. Oseen’s vortex plays an important role in the dynamics of the Navier–Stokes equations in \mathbb{R}^2 , because it describes the long-time asymptotics of all solutions whose vorticity distribution is integrable. This result was first proved in [Giga and Kambe 1988] for small solutions, and subsequently in [Carpio 1994] for large solutions with small circulation. The general case was finally settled in [Gallay and Wayne 2005]. It is worth mentioning that all these results were obtained using the vorticity formulation of the Navier–Stokes equations.

In the case of an exterior domain $\Omega \subset \mathbb{R}^2$, much less is known about infinite-energy solutions, mainly because the vorticity formulation is not convenient anymore due to the boundary conditions. A general existence result was established in [Kozono and Yamazaki 1995], who proved that system (1) is globally well-posed for initial data u_0 in the weak L^2 space $L_\sigma^{2,\infty}(\Omega)$, provided that the local singularity of u_0 in $L^{2,\infty}$ is sufficiently small. In what follows, we consider initial data of the form

$$u_0 = \alpha \chi \Theta_0 + v_0, \tag{4}$$

where $\Theta_0(x) = \Theta(x, 0)$ is Oseen’s vortex at time $t = 0$, and $\chi : \mathbb{R}^2 \rightarrow [0, 1]$ is a smooth, radially symmetric cut-off function such that $\chi = 0$ on a neighborhood of $\mathbb{R}^2 \setminus \Omega$ and $\chi(x) = 1$ when $|x|$ is sufficiently large. For any $\alpha \in \mathbb{R}$ and any $v_0 \in L_\sigma^2(\Omega)$, Theorem 4 in [Kozono and Yamazaki 1995] asserts that the Navier–Stokes equation (1) has a global solution with initial data (4), which is unique in an appropriate

class. However, little is known about the long-time behavior of this solution, and in particular there is no a priori estimate which guarantees that the $L^{2,\infty}$ norm of u remains bounded for all times.

Very recently, a first result concerning the long-time behavior of solutions of (1) with initial data of the form (4) was obtained by Ifitimie, Karch, and Lacave:

Theorem 1.1 [Ifitimie et al. 2011]. *Let $\Omega \subset \mathbb{R}^2$ be a smooth exterior domain whose complement $\mathbb{R}^2 \setminus \Omega$ is a connected set in \mathbb{R}^2 . For any $v_0 \in L^2_\sigma(\Omega)$, there exists a constant $\epsilon = \epsilon(v_0, \Omega) > 0$ such that, for all $\alpha \in [-\epsilon, \epsilon]$, the solution of (1) with initial data (4) satisfies*

$$\lim_{t \rightarrow \infty} t^{\frac{1}{2} - \frac{1}{p}} \|u(\cdot, t) - \alpha \Theta(\cdot, t)\|_{L^p(\Omega)} = 0 \quad \text{for all } p \in (2, \infty). \tag{5}$$

Moreover, there exists $\epsilon_0 = \epsilon_0(\Omega) > 0$ such that $\epsilon \geq \epsilon_0$ if $\|v_0\|_{L^2} \leq \epsilon_0$.

Theorem 1.1 shows that solutions of (1) which are finite-energy perturbations of Oseen’s vortex $\alpha \Theta_0$ behave asymptotically in time like the self-similar Oseen vortex $\alpha \Theta(x, t)$, provided that the circulation at infinity, α , is sufficiently small, depending on the size of the initial perturbation. The conclusion holds in particular when both the circulation α and the finite-energy perturbation v_0 are small, so that Theorem 1.1 extends to exterior domains the result of [Giga and Kambe 1988]. For large solutions, however, the assumption that α be small depending on v_0 is very restrictive. The goal of the present paper is to prove the following result, which reaches a conclusion similar to that of Theorem 1.1 under different assumptions on the initial data:

Theorem 1.2. *Fix $q \in (1, 2)$, and let $\mu = 1/q - 1/2$. There exists a constant $\epsilon = \epsilon(q) > 0$ such that, for any smooth exterior domain $\Omega \subset \mathbb{R}^2$ and for all initial data of the form (4) with $|\alpha| \leq \epsilon$ and $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$, the solution of the Navier–Stokes equations (1) satisfies*

$$\|u(\cdot, t) - \alpha \Theta(\cdot, t)\|_{L^2(\Omega)} + t^{1/2} \|\nabla u(\cdot, t) - \alpha \nabla \Theta(\cdot, t)\|_{L^2(\Omega)} = \mathcal{O}(t^{-\mu}), \tag{6}$$

as $t \rightarrow +\infty$.

Here, we also suppose that the circulation at infinity is small, and we assume in addition that the initial perturbation belongs to $L^2_\sigma(\Omega) \cap L^q(\Omega)^2$ for some $q < 2$. Unlike in Theorem 1.1, the limiting case $q = 2$ is not included, and the proof shows that $\epsilon(q) = \mathcal{O}(\sqrt{2 - q})$ as $q \rightarrow 2$. However, there is absolutely no restriction on the size of the perturbation v_0 ; hence Theorem 1.2 establishes a *global stability property* for the Lamb–Oseen vortices (with small circulation) in two-dimensional exterior domains. In this sense, our result can be considered as a generalization to exterior domains of the work of Carpio [1994], although our proof relies on completely different ideas. On the other hand, since our perturbations decay faster at infinity (in space) than those considered by Ifitimie, Karch, and Lacave, we are able to show that the difference $u(x, t) - \alpha \Theta(x, t)$ converges rapidly to zero, like an inverse power of time, as $t \rightarrow \infty$. In particular, using (6) and elementary interpolation, we obtain the estimate

$$\sup_{t > 0} t^{\frac{1}{q} - \frac{1}{p}} \|u(\cdot, t) - \alpha \Theta(\cdot, t)\|_{L^p(\Omega)} < \infty \quad \text{for all } p \in [2, \infty),$$

which improves (5) since $q < 2$.

At this point, it is useful to mention that the assumption that u_0 can be decomposed as in (4) for some $\alpha \in \mathbb{R}$ and some $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$ is automatically satisfied if we suppose that the initial vorticity $\omega_0 = \text{curl } u_0$ is sufficiently localized. Indeed, let us assume for simplicity that u_0 vanishes on the boundary $\partial\Omega$. For $1 \leq p < \infty$, we denote by $\dot{W}^{1,p}_{0,\sigma}(\Omega)$ the completion with respect to the norm $u \mapsto \|\nabla u\|_{L^p}$ of the space of all smooth, divergence-free vector fields with compact support in Ω . Using this notation, we have the following result:

Proposition 1.3. *Fix $q \in (1, 2)$. Assume that u_0 belongs to $\dot{W}^{1,p}_{0,\sigma}(\Omega)$ for some $p \in [1, 2)$, and that the associated vorticity $\omega_0 = \text{curl } u_0$ satisfies*

$$\int_{\Omega} (1 + |x|^2)^m |\omega_0(x)|^2 \, dx < \infty \tag{7}$$

for some $m > 2/q$. If we define $\alpha = \int_{\Omega} \omega_0(x) \, dx$, then u_0 can be decomposed as in (4) for some $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$. In particular, if $|\alpha| \leq \epsilon$, the conclusion of Theorem 1.2 holds.

For completeness, we give a short proof of Proposition 1.3 in the Appendix. Returning to the discussion of Theorem 1.2, we emphasize that the smallness condition on the circulation α is independent of the domain Ω , which can be an arbitrary multiply connected exterior domain. In fact, the proof will show that the optimal constant $\epsilon(q)$ is entirely determined by quantities that appear in the evolution equation for the perturbation of Oseen’s vortex in the whole plane \mathbb{R}^2 . Note that Oseen vortices are known to be globally stable for all values of the circulation α when $\Omega = \mathbb{R}^2$ [Gallay and Wayne 2005], but in that particular case one can use the vorticity equation to obtain precise information on the solutions of (1). The reader who is not interested in precise convergence rates could consider the following variant of Theorem 1.2, where the condition on the circulation is totally explicit:

Corollary 1.4. *There exists a universal constant $\epsilon_* \geq 4.956$ such that, if $|\alpha| < \epsilon_*$ and if*

$$v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$$

for all $q \in (1, 2)$, the solution of the Navier–Stokes equations (1) with initial data (4) satisfies

$$\|u(\cdot, t) - \alpha\Theta(\cdot, t)\|_{L^2(\Omega)} \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

The rest of this paper is devoted to the proof of Theorem 1.2, which is quite different from that of Theorem 1.1 in [Iftimie et al. 2011]. In the preliminary section (Section 2), we collect various estimates on the truncated Oseen vortex χ^Θ , which can be verified by direct calculations. In Section 3, following the classical approach of [Fujita and Kato 1964], we prove the existence of a unique global solution of (1) for small initial data of the form (4), and we obtain the asymptotics (6) for small solutions. To deal with large solutions, we derive in Section 4 a “logarithmic energy estimate”, which shows that the energy norm of the perturbation v has at most a logarithmic growth as $t \rightarrow \infty$. This is the key new ingredient, which we use as a substitute for the classical energy inequality when $\alpha \neq 0$. Exploiting this estimate and our assumption that $v_0 \in L^q(\Omega)^2$, we control in Section 5 the evolution of a fractional primitive of v , and we deduce that the perturbation $v(\cdot, t)$ converges to zero in energy norm, at least along a sequence of times. Thus we can eventually use the results of Section 3, and the conclusion follows.

2. The truncated Oseen vortex

Fix $\rho \geq 1$ large enough so that $\{x \in \mathbb{R}^2 \mid |x| \geq \rho\} \subset \Omega$. Let $\chi(x) = \tilde{\chi}(x/\rho)$, where $\tilde{\chi} \in C^\infty(\mathbb{R}^2)$ is a radially symmetric cut-off function satisfying $\tilde{\chi}(x) = 0$ when $|x| \leq 1$, $\tilde{\chi}(x) = 1$ when $|x| \geq 2$, and $0 \leq \tilde{\chi}(x) \leq 1$ for all $x \in \mathbb{R}^2$. We define the *truncated Oseen vortex* (with unit circulation) as follows:

$$u^\chi(x, t) = \chi(x)\Theta(x, t) = \frac{1}{2\pi} \frac{x^\perp}{|x|^2} \left(1 - e^{-\frac{|x|^2}{4(1+t)}}\right) \chi(x), \quad x \in \mathbb{R}^2, t \geq 0. \tag{8}$$

Since χ is radially symmetric and $\text{supp } \chi \subset \{x \in \mathbb{R}^2 \mid |x| \geq \rho\} \subset \Omega$, it is clear that $u^\chi(x, t)$ is a smooth divergence-free vector field which vanishes in a neighborhood of $\mathbb{R}^2 \setminus \Omega$. Let $\omega^\chi = \partial_1 u_2^\chi - \partial_2 u_1^\chi$ be the corresponding vorticity field, namely

$$\omega^\chi(x, t) = \chi(x)\Xi(x, t) + \frac{1}{2\pi} \frac{1}{|x|^2} \left(1 - e^{-\frac{|x|^2}{4(1+t)}}\right) x \cdot \nabla \chi(x), \tag{9}$$

where $\Xi(x, t)$ is defined in (3). Since $u^\chi(x, t) = \Theta(x, t)$ whenever $|x| \geq 2\rho$, the circulation of u^χ at infinity is equal to 1, so that $\int_{\mathbb{R}^2} \omega^\chi \, dx = 1$. Moreover, a direct calculation shows that

$$(u^\chi \cdot \nabla)u^\chi = \frac{1}{2} \nabla |u^\chi|^2 + (u^\chi)^\perp \omega^\chi = -\frac{x}{|x|^2} |u^\chi|^2; \tag{10}$$

hence there exists a radially symmetric function $p^\chi(x, t)$ such that $-\nabla p^\chi = (u^\chi \cdot \nabla)u^\chi$. This shows that $P(u^\chi \cdot \nabla)u^\chi = 0$, where P denotes the Leray–Hopf projection in Ω , namely the orthogonal projection in $L^2(\Omega)^2$ onto the subspace $L^2_\sigma(\Omega)$.

The following elementary estimates will be useful:

Lemma 2.1. (i) *For any $p \in (2, \infty]$, there exists a constant $a_p > 0$ such that*

$$\|u^\chi(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \frac{a_p}{(1+t)^{\frac{1}{2} - \frac{1}{p}}}, \quad t \geq 0. \tag{11}$$

(ii) *For any $p \in (1, \infty]$, there exists a constant $b_p > 0$ such that*

$$\|\nabla u^\chi(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \frac{b_p}{(1+t)^{1 - \frac{1}{p}}}, \quad t \geq 0. \tag{12}$$

(iii) *For all $t, s \geq 0$, we have*

$$\|u^\chi(\cdot, t) - u^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 \leq \frac{1}{4\pi} \left| \log \frac{1+t}{1+s} \right|. \tag{13}$$

(iv) *There exists a constant $\kappa_1 > 0$ such that, for all $t, s \geq 0$,*

$$\|\nabla u^\chi(\cdot, t) - \nabla u^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 \leq \kappa_1 \left| \frac{1}{1+t} - \frac{1}{1+s} \right|. \tag{14}$$

Moreover all constants a_p, b_p , and κ_1 are independent of ρ , and hence of the domain Ω .

Proof. By (8) we have

$$u^\chi(x, t) = \chi(x)\Theta(x, t) = \frac{\chi(x)}{\sqrt{1+t}}\Theta_0\left(\frac{x}{\sqrt{1+t}}\right),$$

where $\Theta_0(x) = \Theta(x, 0)$. Since $0 \leq \chi \leq 1$ and $\Theta_0 \in L^p(\mathbb{R}^2)^2$ for all $p > 2$, we find

$$\|u^\chi(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \frac{1}{\sqrt{1+t}} \left\| \Theta_0\left(\frac{\cdot}{\sqrt{1+t}}\right) \right\|_{L^p(\mathbb{R}^2)} = \frac{\|\Theta_0\|_{L^p(\mathbb{R}^2)}}{(1+t)^{\frac{1}{2}-\frac{1}{p}}}, \quad t \geq 0.$$

This proves (11).

Similarly, we have $\partial_i u^\chi = \chi \partial_i \Theta + (\partial_i \chi) \Theta$ for $i = 1, 2$. As $\partial_i \Theta_0 \in L^p(\mathbb{R}^2)^2$ for all $p > 1$, we obtain as before

$$\|\chi \partial_i \Theta(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \frac{1}{1+t} \left\| \partial_i \Theta_0\left(\frac{\cdot}{\sqrt{1+t}}\right) \right\|_{L^p(\mathbb{R}^2)} = \frac{\|\partial_i \Theta_0\|_{L^p(\mathbb{R}^2)}}{(1+t)^{1-\frac{1}{p}}}, \quad t \geq 0. \tag{15}$$

On the other hand, the function $\partial_i \chi$ is supported in the annulus

$$D = \{x \in \mathbb{R}^2 \mid \rho \leq |x| \leq 2\rho\},$$

and satisfies $|\partial_i \chi(x)| \leq C\rho^{-1}$ for some $C > 0$ independent of ρ . Moreover, it follows from (2) that

$$|\Theta(x, t)| \leq \frac{1}{2\pi} \min\left(\frac{1}{|x|}, \frac{|x|}{4(1+t)}\right), \quad x \in \mathbb{R}^2, t \geq 0;$$

hence

$$|(\partial_i \chi(x))\Theta(x, t)| \leq C \min\left(\frac{1}{\rho^2}, \frac{1}{1+t}\right) \mathbf{1}_D(x), \quad x \in \mathbb{R}^2, t \geq 0,$$

where $\mathbf{1}_D$ is the characteristic function of D . Taking the L^p norm of both sides, we thus obtain

$$\|(\partial_i \chi)\Theta(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq C\rho^{2/p} \min\left(\frac{1}{\rho^2}, \frac{1}{1+t}\right) \leq \frac{C}{(1+t)^{1-\frac{1}{p}}}, \quad t \geq 0. \tag{16}$$

Combining (15) and (16), we arrive at (12).

To prove (13), we observe that

$$\begin{aligned} \|u^\chi(\cdot, t) - u^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 &\leq \frac{1}{4\pi^2} \int_{\mathbb{R}^2} \frac{1}{|x|^2} \left(e^{-\frac{|x|^2}{4(1+t)}} - e^{-\frac{|x|^2}{4(1+s)}} \right)^2 dx \\ &= \frac{1}{2\pi} \log \left\{ \frac{1}{2} \sqrt{\frac{1+t}{1+s}} + \frac{1}{2} \sqrt{\frac{1+s}{1+t}} \right\} \leq \frac{1}{4\pi} \left| \log \frac{1+t}{1+s} \right| \end{aligned}$$

for all $t, s \geq 0$. Finally, using (9), we find

$$\omega^\chi(x, t) - \omega^\chi(x, s) = \chi(x)(\Xi(x, t) - \Xi(x, s)) - \frac{x \cdot \nabla \chi(x)}{2\pi|x|^2} \left(e^{-\frac{|x|^2}{4(1+t)}} - e^{-\frac{|x|^2}{4(1+s)}} \right).$$

Thus $\|\nabla u^\chi(\cdot, t) - \nabla u^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 = \|\omega^\chi(\cdot, t) - \omega^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 \leq (J_1(t, s)^{1/2} + J_2(t, s)^{1/2})^2$, where

$$\begin{aligned} J_1(t, s) &= \int_{\mathbb{R}^2} \chi(x)^2 (\Xi(x, t) - \Xi(x, s))^2 dx \leq \int_{\mathbb{R}^2} (\Xi(x, t) - \Xi(x, s))^2 dx \\ &= \frac{1}{8\pi} \left\{ \frac{1}{1+t} + \frac{1}{1+s} - \frac{4}{t+s+2} \right\} \leq \frac{1}{8\pi} \left| \frac{1}{1+t} - \frac{1}{1+s} \right|, \end{aligned}$$

and

$$\begin{aligned} J_2(t, s) &= \int_{\mathbb{R}^2} \frac{|\nabla \chi(x)|^2}{4\pi^2|x|^2} \left(e^{-\frac{|x|^2}{4(1+t)}} - e^{-\frac{|x|^2}{4(1+s)}} \right)^2 dx \leq C\rho^{-4} \int_D \left(e^{-\frac{|x|^2}{4(1+t)}} - e^{-\frac{|x|^2}{4(1+s)}} \right)^2 dx \\ &\leq C\rho^{-2} \sup_{x \in D} \left| e^{-\frac{|x|^2}{4(1+t)}} - e^{-\frac{|x|^2}{4(1+s)}} \right| \leq C \left| \frac{1}{1+t} - \frac{1}{1+s} \right|. \end{aligned}$$

We thus obtain (14), which is the desired estimate. For later use, we also observe that $J_2(t, s)$ can be bounded by $C\rho^2 \left(\frac{1}{1+t} - \frac{1}{1+s}\right)^2$, for some $C > 0$ independent of ρ . Since $\rho \geq 1$, this gives the alternative estimate

$$\|\nabla u^\chi(\cdot, t) - \nabla u^\chi(\cdot, s)\|_{L^2(\mathbb{R}^2)}^2 \leq \frac{1}{8\pi} \left| \frac{1}{1+t} - \frac{1}{1+s} \right| + C\rho^2 \left| \frac{1}{1+t} - \frac{1}{1+s} \right|^{3/2}, \tag{17}$$

which will be used in Section 4. This concludes the proof of Lemma 2.1. □

The truncated Oseen vortex is not a solution of the Navier–Stokes equation, and therefore we need to control the remainder term $R^\chi = \Delta u^\chi - \partial_t u^\chi = (\Delta \chi)\Theta + 2(\nabla \chi \cdot \nabla)\Theta$, which has the explicit expression

$$R^\chi(x, t) = \Theta(x, t)\Delta \chi(x) + 2\frac{x \cdot \nabla \chi(x)}{|x|^2} (x^\perp \Xi(x, t) - \Theta(x, t)). \tag{18}$$

Lemma 2.2. *There exists a constant $\kappa_2 > 0$ (independent of ρ) such that, for any $p \in [1, \infty]$,*

$$\|R^\chi(\cdot, t)\|_{L^p(\mathbb{R}^2)} \leq \frac{\kappa_2 \rho^{\frac{2}{p}-1}}{1+t}, \quad t \geq 0. \tag{19}$$

Moreover, for any vector field $u \in H_{\text{loc}}^1(\mathbb{R}^2)^2$, we have

$$\left| \int_{\mathbb{R}^2} R^\chi(x, t) \cdot u(x) dx \right| \leq \frac{\kappa_2 \rho}{1+t} \|\nabla u\|_{L^2(D)}, \quad t \geq 0, \tag{20}$$

where $D = \{x \in \mathbb{R}^2 \mid \rho \leq |x| \leq 2\rho\}$.

Proof. It is clear from (18) that $|R^\chi(x, t)| \leq C\rho^{-1}(1+t)^{-1}\mathbf{1}_D(x)$ for all $x \in \mathbb{R}^2$ and all $t \geq 0$, and (19) follows immediately. Moreover, we have $R^\chi(x, t) = x^\perp Q^\chi(x, t)$ for some radially symmetric scalar function $Q^\chi(x, t)$; hence $R^\chi(\cdot, t)$ has zero mean over the annulus D . If $u \in H_{\text{loc}}^1(\mathbb{R}^2)^2$ and if we denote by \bar{u} the average of u over D , the Poincaré–Wirtinger inequality implies

$$\left| \int_{\mathbb{R}^2} R^\chi(x, t) \cdot u(x) dx \right| = \left| \int_D R^\chi(x, t) \cdot (u(x) - \bar{u}) dx \right| \leq C\rho \|R^\chi(\cdot, t)\|_{L^2(\mathbb{R}^2)} \|\nabla u\|_{L^2(D)},$$

and using (19) with $p = 2$ we obtain (20). □

3. Asymptotic behavior of small solutions

Given $\alpha \in \mathbb{R}$, we consider solutions of (1) of the form

$$u(x, t) = \alpha u^\chi(x, t) + v(x, t), \quad p(x, t) = \alpha^2 p^\chi(x, t) + q(x, t), \tag{21}$$

where $u^\chi(x, t)$ is the truncated Oseen vortex (8) and p^χ is the associated pressure. The perturbation $v(x, t)$ satisfies the no-slip boundary condition and the equation

$$\partial_t v + \alpha(u^\chi \cdot \nabla)v + \alpha(v \cdot \nabla)u^\chi + (v \cdot \nabla)v = \Delta v + \alpha R^\chi - \nabla q, \quad \operatorname{div} v = 0, \tag{22}$$

where R^χ is given by (18). If we apply the Leray–Hopf projection P and use the fact that $PR^\chi = R^\chi$, we obtain the equivalent system

$$\partial_t v + \alpha P((u^\chi \cdot \nabla)v + (v \cdot \nabla)u^\chi) + P(v \cdot \nabla)v = -Av + \alpha R^\chi, \tag{23}$$

where $A = -P\Delta$ is the Stokes operator, which is selfadjoint and nonnegative in $L^2_\sigma(\Omega)$ with domain $D(A) = L^2_\sigma(\Omega) \cap H^1_0(\Omega)^2 \cap H^2(\Omega)^2$; see [Constantin and Foias 1988].

In this section, we fix some initial time $t_0 \geq 0$ and prove the existence of global solutions to (23) with small initial data $v_0 = v(\cdot, t_0)$ in the energy space. The integral equation associated with (23) is

$$v(t) = S(t - t_0)v_0 + \int_{t_0}^t S(t - s) \{ \alpha R^\chi(s) - P(v(s) \cdot \nabla)v(s) - \alpha P((u^\chi(s) \cdot \nabla)v(s) + (v(s) \cdot \nabla)u^\chi(s)) \} ds, \tag{24}$$

where $v(t) \equiv v(\cdot, t)$ and $S(t) = \exp(-tA)$ is the Stokes semigroup. For $p \in (1, \infty)$, we denote by $L^p_\sigma(\Omega)$ the closure in $L^p(\Omega)^2$ of the set of all smooth divergence-free vector fields with compact support in Ω . We then have the following standard estimates:

Proposition 3.1. *The Stokes operator $-A$ generates an analytic semigroup of contractions in $L^2_\sigma(\Omega)$. Moreover, for each $t > 0$ the operator $S(t) = \exp(-tA)$ extends to a bounded linear operator from $L^q_\sigma(\Omega)$ into $L^2_\sigma(\Omega)$ for $1 < q \leq 2$, and there exists a constant $C = C(q) > 0$ (independent of Ω) such that*

$$t^{\frac{1}{q}-\frac{1}{2}} \|S(t)v_0\|_{L^2(\Omega)} + t^{\frac{1}{q}} \|\nabla S(t)v_0\|_{L^2(\Omega)} \leq C \|v_0\|_{L^q(\Omega)}, \quad t > 0, \tag{25}$$

for all $v_0 \in L^q_\sigma(\Omega)$. In particular, we can take $C = 2$ in (25) if $q = 2$.

Since A is selfadjoint and nonnegative, it is clear that $\{S(t)\}_{t \geq 0}$ is an analytic semigroup of contractions in $L^2_\sigma(\Omega)$. In particular, we have $\|S(t)v_0\|_{L^2} \leq \|v_0\|_{L^2}$ and $t^{1/2} \|\nabla S(t)v_0\|_{L^2} = t^{1/2} \|A^{1/2}S(t)v_0\|_{L^2} \leq \|v_0\|_{L^2}$ for all $t > 0$ if $v_0 \in L^2_\sigma(\Omega)$. On the other hand, general $L^q - L^p$ estimates for $S(t)$ were established in [Borchers and Varnhorn 1993; Dan and Shibata 1999a; 1999b; Kozono and Yamazaki 1995; Maremonti and Solonnikov 1997], but the corresponding constants depend a priori on the domain Ω . The fact that (25) holds with C independent of Ω was already observed in [Borchers and Miyakawa 1992; Kozono and Ogawa 1993a]. For the reader’s convenience, we reproduce the proof of (25) in Section 5 below.

The main result of this section is this:

Proposition 3.2. Fix $\mu \in (0, 1/2)$. There exist positive constants K_0, δ, V_Ω , and T_Ω such that, if $t_0 \geq T_\Omega$, if $|\alpha| \leq \delta$, and if $\|v_0\|_{L^2(\Omega)} \leq V_\Omega$, then the perturbation equation (23) has a unique global solution $v \in C^0([t_0, \infty); L^2_\sigma(\Omega))$ such that

$$\sup_{t \geq t_0} \|v(t)\|_{L^2(\Omega)} + \sup_{t > t_0} (t - t_0)^{\frac{1}{2}} \|\nabla v(t)\|_{L^2(\Omega)} \leq 4\|v_0\|_{L^2(\Omega)} + K_0 \rho^{\frac{1}{2}} |\alpha| (1 + t_0)^{-\frac{1}{4}}. \tag{26}$$

Here K_0 and δ are independent of Ω . In addition, if

$$M := \sup_{\tau > 0} \tau^\mu \|S(\tau)v_0\|_{L^2(\Omega)} + \sup_{\tau > 0} \tau^{\mu + \frac{1}{2}} \|\nabla S(\tau)v_0\|_{L^2(\Omega)} < \infty, \tag{27}$$

then

$$\sup_{t > t_0} (t - t_0)^\mu \|v(t)\|_{L^2(\Omega)} + \sup_{t > t_0} (t - t_0)^{\mu + \frac{1}{2}} \|\nabla v(t)\|_{L^2(\Omega)} \leq 2M + C_\Omega |\alpha|, \tag{28}$$

for some $C_\Omega > 0$ depending on Ω .

Proof. We follow the classical approach of [Fujita and Kato 1964]. Given $t_0 \geq 0$, we introduce the Banach space $X = \{v \in C^0([t_0, \infty); L^2_\sigma(\Omega)) \cap C^0((t_0, \infty); H^1_0(\Omega)^2) \mid \|v\|_X < \infty\}$, equipped with the norm

$$\|v\|_X = \sup_{t \geq t_0} \|v(t)\|_{L^2} + \sup_{t > t_0} (t - t_0)^{\frac{1}{2}} \|\nabla v(t)\|_{L^2}.$$

If $v_0 \in L^2_\sigma(\Omega)$, we define $\bar{v}(t) = S(t - t_0)v_0$ for $t \geq t_0$. In view of (25), we have $\bar{v} \in X$ and $\|\bar{v}\|_X \leq 2\|v_0\|_{L^2}$. On the other hand, given any $v \in X$ we define, for $t \geq t_0$,

$$(Fv)(t) = \int_{t_0}^t S(t - s) (\alpha R^\chi(s) + \alpha G_1^v(s) + G_2^v(s)) ds = \alpha F_0(t) + \alpha (F_1v)(t) + (F_2v)(t),$$

where $G_1^v(s) = -P(u^\chi(s) \cdot \nabla)v(s) - P(v(s) \cdot \nabla)u^\chi(s)$ and $G_2^v(s) = -P(v(s) \cdot \nabla)v(s)$. We shall show that F maps X into X , and that there exist positive constants $C_1, C_2, C_{3,\Omega}$ (independent of t_0) such that

$$\|Fv\|_X \leq C_1 \rho^{\frac{1}{2}} |\alpha| (1 + t_0)^{-\frac{1}{4}} + |\alpha| C_2 \|v\|_X + C_{3,\Omega} \|v\|_X^2, \tag{29}$$

$$\|Fv - F\tilde{v}\|_X \leq |\alpha| C_2 \|v - \tilde{v}\|_X + C_{3,\Omega} (\|v\|_X + \|\tilde{v}\|_X) \|v - \tilde{v}\|_X, \tag{30}$$

for all $v, \tilde{v} \in X$.

To prove (29), we estimate separately the contributions of F_0, F_1 , and F_2 . First, using (25) with $q = 4/3$, we obtain for $t > t_0$:

$$\|F_0(t)\|_{L^2} + (t - t_0)^{\frac{1}{2}} \|\nabla F_0(t)\|_{L^2} \leq C \int_{t_0}^t \left(\frac{1}{(t - s)^{\frac{1}{4}}} + \frac{(t - t_0)^{\frac{1}{2}}}{(t - s)^{\frac{3}{4}}} \right) \|R^\chi(s)\|_{L^{\frac{4}{3}}} ds, \tag{31}$$

and from Lemma 2.2 we know that $\|R^\chi(s)\|_{L^{4/3}} \leq C\rho^{1/2}(1 + s)^{-1}$ for all $s \geq 0$. It follows that $\|F_0\|_X \leq C_1 \rho^{1/2}(1 + t_0)^{-1/4}$ for some $C_1 > 0$ independent of t_0 and Ω . In a similar way, we find

$$\|(F_2v)(t)\|_{L^2} + (t - t_0)^{\frac{1}{2}} \|\nabla(F_2v)(t)\|_{L^2} \leq C \int_{t_0}^t \left(\frac{1}{(t - s)^{\frac{1}{4}}} + \frac{(t - t_0)^{\frac{1}{2}}}{(t - s)^{\frac{3}{4}}} \right) \|G_2^v(s)\|_{L^{\frac{4}{3}}} ds. \tag{32}$$

Using the fact that the Leray–Hopf projection is a bounded operator in $L^{4/3}(\Omega)^2$, whose norm depends a priori on Ω , we estimate

$$\|G_2^v(s)\|_{L^{\frac{4}{3}}} \leq C_\Omega \|v(s)\|_{L^4} \|\nabla v(s)\|_{L^2} \leq C_\Omega \|v(s)\|_{L^2}^{\frac{1}{2}} \|\nabla v(s)\|_{L^2}^{\frac{3}{2}} \leq \frac{C_\Omega \|v\|_X^2}{(s - t_0)^{\frac{3}{4}}},$$

for all $s > t_0$. It follows that $\|F_2 v\|_X \leq C_{3,\Omega} \|v\|_X^2$, where $C_{3,\Omega} > 0$ is independent of t_0 . Finally, to bound F_1 , we proceed in a slightly different way in order to obtain a constant C_2 that does not depend on Ω . Observing that $G_1^v(s) = -A^{1/2} A^{-1/2} P \operatorname{div}(u^\chi \otimes v + v \otimes u^\chi)(s)$, and that $\|A^{1/2} v\|_{L^2} = \|\nabla v\|_{L^2}$ for all $v \in L_\sigma^2(\Omega) \cap H_0^1(\Omega)^2$, we can use (25) with $q = 2$ to obtain

$$\|(F_1 v)(t)\|_{L^2} \leq \int_{t_0}^t (t-s)^{-\frac{1}{2}} \|A^{-1/2} P \operatorname{div}(u^\chi \otimes v + v \otimes u^\chi)(s)\|_{L^2} ds. \tag{33}$$

Similarly, the quantity $(t - t_0)^{\frac{1}{2}} \|\nabla(F_1 v)(t)\|_{L^2}$ can be bounded by

$$\int_{t_0}^{\frac{t+t_0}{2}} \frac{(t-t_0)^{\frac{1}{2}}}{t-s} \|A^{-1/2} P \operatorname{div}(u^\chi \otimes v + v \otimes u^\chi)(s)\|_{L^2} ds + \int_{\frac{t+t_0}{2}}^t \frac{(t-t_0)^{\frac{1}{2}}}{(t-s)^{\frac{1}{2}}} \|G_1^v(s)\|_{L^2} ds. \tag{34}$$

Since $A^{-1/2} P \operatorname{div}$ defines a bounded operator from $L^2(\Omega)^4$ into $L_\sigma^2(\Omega)$ whose norm is less than or equal to 1 (see [Sohr 2001, Lemma III-2-6-1]), we have from (11)

$$\|A^{-1/2} P \operatorname{div}(u^\chi \otimes v + v \otimes u^\chi)(s)\|_{L^2} \leq 2\|u^\chi(s)v(s)\|_{L^2} \leq 2a_\infty(1+s)^{-\frac{1}{2}} \|v\|_X.$$

Moreover, using (11) and (12) we find

$$\|G_1^v(s)\|_{L^2} \leq \|u^\chi(s)\nabla v(s)\|_{L^2} + \|v(s)\nabla u^\chi(s)\|_{L^2} \leq \frac{a_\infty \|v\|_X}{(1+s)^{\frac{1}{2}}(s-t_0)^{\frac{1}{2}}} + \frac{b_\infty \|v\|_X}{1+s}.$$

Inserting these estimates into (33) and (34), we obtain $\|F_1 v\|_X \leq C_2 \|v\|_X$ for some $C_2 > 0$ independent of t_0 and Ω . Since $Fv = \alpha F_0 + \alpha F_1 v + F_2 v$, this concludes the proof of (29), and the Lipschitz bound (30) is established in exactly the same way.

Now let $B_r = \{v \in X \mid \|v\|_X \leq r\}$, where $r > 0$ is small enough so that $4rC_{3,\Omega} \leq 1$. If we assume that $4|\alpha|C_2 \leq 1$, $8\|v_0\|_{L^2} \leq r$, and $4C_1\rho^{1/2}|\alpha|(1+t_0)^{-1/4} \leq r$, the estimates above imply that the map $v \mapsto \bar{v} + Fv$ leaves the closed ball B_r invariant and is a strict contraction in B_r . By construction, the unique fixed point of that map in B_r is the desired solution of (24). This proves the existence part of Proposition 3.2 with

$$K_0 = 2C_1, \quad \delta = \frac{1}{4C_2}, \quad V_\Omega = \frac{1}{32C_{3,\Omega}}, \quad T_\Omega = \left(\frac{4C_1C_{3,\Omega}\rho^{\frac{1}{2}}}{C_2}\right)^4.$$

In a second step, we assume that (27) holds for some $\mu \in (0, 1/2)$. Given any $T > t_0$, we define

$$\mathcal{E}_T = \sup_{t_0 \leq t \leq T} (t - t_0)^\mu \|v(t)\|_{L^2} + \sup_{t_0 < t \leq T} (t - t_0)^{\mu + \frac{1}{2}} \|\nabla v(t)\|_{L^2},$$

where v is the solution of (24) constructed in the previous step. Our goal is to show that \mathcal{E}_T is uniformly bounded by a constant which does not depend on T . Since $v(t) = S(t - t_0)v_0 + (Fv)(t)$, we have

$$\mathcal{E}_T \leq M + \sup_{t_0 \leq t \leq T} (t - t_0)^\mu \|(Fv)(t)\|_{L^2} + \sup_{t_0 < t \leq T} (t - t_0)^{\mu + \frac{1}{2}} \|\nabla(Fv)(t)\|_{L^2}, \tag{35}$$

where M is defined in (27). To estimate the last two terms, we proceed as above. Let $p \in (1, 2)$ be such that $1/p > \mu + 1/2$, and define $q \in (2, \infty)$ by the relation $1/q = 1/p - 1/2$. As in (31) and (32), we have

$$(t - t_0)^\mu \|F_0(t)\|_{L^2} + (t - t_0)^{\mu + \frac{1}{2}} \|\nabla F_0(t)\|_{L^2} \leq C \int_{t_0}^t \left(\frac{(t - t_0)^\mu}{(t - s)^{\frac{1}{q}}} + \frac{(t - t_0)^{\mu + \frac{1}{2}}}{(t - s)^{\frac{1}{p}}} \right) \|R^\chi(s)\|_{L^p} ds,$$

$$(t - t_0)^\mu \|(F_2v)(t)\|_{L^2} + (t - t_0)^{\mu + \frac{1}{2}} \|\nabla(F_2v)(t)\|_{L^2} \leq C \int_{t_0}^t \left(\frac{(t - t_0)^\mu}{(t - s)^{\frac{1}{q}}} + \frac{(t - t_0)^{\mu + \frac{1}{2}}}{(t - s)^{\frac{1}{p}}} \right) \|G_2^v(s)\|_{L^p} ds,$$

for $t \in (t_0, T]$. Moreover $\|R^\chi(s)\|_{L^p} \leq C\rho^{\frac{2}{p}-1}(1 + s)^{-1}$ and

$$\|P(v(s) \cdot \nabla)v(s)\|_{L^p} \leq C_\Omega \|v(s)\|_{L^q} \|\nabla v(s)\|_{L^2} \leq C_\Omega \|v(s)\|_{L^2}^{\frac{2}{q}} \|\nabla v(s)\|_{L^2}^{2 - \frac{2}{q}} \leq \frac{C_\Omega \|v\|_X \mathcal{E}_T}{(s - t_0)^{\mu + 1 - \frac{1}{q}}}$$

for all $s \in (t_0, T]$. The term involving F_1v is estimated as in (33) and (34), and we find

$$(t - t_0)^\mu \|(F_1v)(t)\|_{L^2} \leq C \int_{t_0}^t \frac{(t - t_0)^\mu \mathcal{E}_T}{(t - s)^{\frac{1}{2}}(1 + s)^{\frac{1}{2}}(s - t_0)^\mu} ds,$$

$$(t - t_0)^{\mu + \frac{1}{2}} \|\nabla(F_1v)(t)\|_{L^2} \leq C \int_{t_0}^{\frac{t+t_0}{2}} \frac{(t - t_0)^{\mu + \frac{1}{2}} \mathcal{E}_T}{(t - s)(1 + s)^{\frac{1}{2}}(s - t_0)^\mu} ds$$

$$+ C \int_{\frac{t+t_0}{2}}^t \frac{(t - t_0)^{\mu + \frac{1}{2}}}{(t - s)^{\frac{1}{2}}} \left(\frac{\mathcal{E}_T}{(1 + s)^{\frac{1}{2}}(s - t_0)^{\mu + \frac{1}{2}}} + \frac{\mathcal{E}_T}{(1 + s)(s - t_0)^\mu} \right) ds.$$

If we insert these estimates into (35), we obtain after elementary calculations

$$\mathcal{E}_T \leq M + \tilde{C}_1 \rho^{\frac{2}{p}-1} |\alpha| (1 + t_0)^{-\frac{1}{p} + \mu + \frac{1}{2}} + \tilde{C}_2 |\alpha| \mathcal{E}_T + \tilde{C}_{3,\Omega} \|v\|_X \mathcal{E}_T, \tag{36}$$

for some positive constants $\tilde{C}_1, \tilde{C}_2, \tilde{C}_{3,\Omega}$ independent of T and t_0 . Now, taking δ and V_Ω smaller and T_Ω larger if needed, we can ensure that $\tilde{C}_2 |\alpha| + \tilde{C}_{3,\Omega} \|v\|_X \leq 1/2$. Then (36) implies that

$$\mathcal{E}_T \leq 2M + 2 \frac{\tilde{C}_1 \rho^{\frac{2}{p}-1} |\alpha|}{(1 + t_0)^{\frac{1}{p} - \mu - \frac{1}{2}}}$$

for all $T > t_0$, and (28) follows. This concludes the proof. □

Remark 3.3. The proof of Proposition 3.2 can be modified in a classical way [Fujita and Kato 1964; Brezis 1994] to yield the following local existence result. For any $\alpha \in \mathbb{R}$, any $t_0 \geq 0$, and any $v_0 \in L^2_\sigma(\Omega)$, there exists $T = T(\alpha, v_0, \Omega) > 0$ such that (23) has a unique solution $v \in C^0([t_0, t_0 + T]; L^2_\sigma(\Omega)) \cap C^0((t_0, t_0 + T]; H^1_0(\Omega)^2)$ satisfying $v(t_0) = v_0$; moreover, any upper bound on $|\alpha| + \|v_0\|_{H^1}$ gives a lower bound on the local existence time T . In our formulation of Proposition 3.2, smallness conditions

were imposed on α and v_0 to ensure global existence, and the assumption on the initial time t_0 guarantees that the smallness condition on α is independent of the domain Ω .

4. A logarithmic energy estimate

In this section, we establish our key estimate for large solutions of (23) in the energy space. Fix $\alpha \in \mathbb{R}$, $v_0 \in L^2_\sigma(\Omega)$, and let $v \in C^0([0, T]; L^2_\sigma(\Omega)) \cap C^0((0, T]; H^1_0(\Omega)^2)$ be a solution of (23) with initial data $v(0) = v_0$; see Remark 3.3. We first derive a crude bound on v using a classical energy estimate. Multiplying both sides of (23) by v and integrating by parts over Ω , we find

$$\frac{1}{2} \frac{d}{dt} \|v(t)\|_{L^2}^2 + \|\nabla v(t)\|_{L^2}^2 = \alpha \langle v(t), R^\chi(t) \rangle - \alpha \langle v(t), (v(t) \cdot \nabla) u^\chi(t) \rangle, \tag{37}$$

where $\langle \cdot, \cdot \rangle$ denotes the usual scalar product in $L^2_\sigma(\Omega)$, so that $\|\cdot\|_{L^2} = \langle \cdot, \cdot \rangle^{1/2}$. Using (20), we easily obtain

$$|\alpha \langle v(t), R^\chi(t) \rangle| \leq \frac{\kappa_2 \rho |\alpha|}{1+t} \|\nabla v(t)\|_{L^2} \leq \frac{\eta}{2} \|\nabla v(t)\|_{L^2}^2 + \frac{\kappa_2^2 \rho^2 \alpha^2}{2\eta(1+t)^2},$$

for any $\eta \in (0, 1]$. Moreover, applying (12) with $p = \infty$, we see that

$$|\langle v(t), (v(t) \cdot \nabla) u^\chi(t) \rangle| \leq \frac{b_\infty}{1+t} \|v(t)\|_{L^2}^2.$$

We thus obtain the energy inequality

$$\frac{d}{dt} \|v(t)\|_{L^2}^2 + (2 - \eta) \|\nabla v(t)\|_{L^2}^2 \leq \frac{2b_\infty |\alpha|}{1+t} \|v(t)\|_{L^2}^2 + \frac{\kappa_2^2 \rho^2 \alpha^2}{\eta(1+t)^2}, \quad 0 < t \leq T.$$

Using Gronwall’s lemma, we deduce that

$$\|v(t)\|_{L^2}^2 + (2 - \eta) \int_{t_0}^t \|\nabla v(s)\|_{L^2}^2 ds \leq \left(\frac{1+t}{1+t_0} \right)^{2b_\infty |\alpha|} \left(\|v(t_0)\|_{L^2}^2 + \frac{\kappa_2^2 \rho^2 \alpha^2}{\eta(1+t_0)} \right), \tag{38}$$

for $0 \leq t_0 < t \leq T$.

We shall see that estimate (38) is pessimistic for large times, but it already implies that the solutions of (23) in the energy space $L^2_\sigma(\Omega)$ are global. Indeed, (38) shows that the norm $\|v(t)\|_{L^2}$ grows at most polynomially in time, and it is then straightforward to establish a similar result for $\|\nabla v(t)\|_{L^2}$. In particular, the H^1 norm of $v(t)$ cannot blow up in finite time, and using Remark 3.3 we conclude that all solutions of (23) in $L^2_\sigma(\Omega)$ are global.

The aim of this section is to establish the following “logarithmic energy estimate”, which improves (38) for large times.

Proposition 4.1. *There exists a constant $K_1 > 0$ (independent of Ω) such that, for any $\alpha \in \mathbb{R}$ and any $v_0 \in L^2_\sigma(\Omega)$, the solution of (23) with initial data v_0 satisfies, for all $t \geq 1$,*

$$\|v(t)\|_{L^2(\Omega)}^2 + \int_0^t \|\nabla v(s)\|_{L^2(\Omega)}^2 ds \leq K_1 (\|v_0\|_{L^2(\Omega)}^2 + \alpha^2 \log(1+t) + D_{\alpha,\rho}), \tag{39}$$

where $D_{\alpha,\rho} = \alpha^2 \log(1 + |\alpha|) + \alpha^2 \rho^2$.

Proof. As in (38), we introduce here a parameter $\eta \in (0, 1]$, which will be used in Section 5 below to specify the optimal smallness condition on the circulation α and prove Corollary 1.4. The reader who is not interested in optimal constants should set $\eta = 1$ everywhere.

Given any $\tau \geq 0$, we define

$$\tilde{v}(x, t) = u(x, t) - \alpha u^X(x, t + \tau) = v(x, t) + \alpha(u^X(x, t) - u^X(x, t + \tau)), \tag{40}$$

for all $x \in \Omega$ and all $t > 0$. Then \tilde{v} satisfies (23) where $u^X(x, t)$ and $R^X(x, t)$ are replaced by $u^X(x, t + \tau)$ and $R^X(x, t + \tau)$, respectively. Proceeding exactly as above, we thus obtain the following energy estimate:

$$\|\tilde{v}(t)\|_{L^2}^2 + (2 - \eta) \int_0^t \|\nabla \tilde{v}(s)\|_{L^2}^2 ds \leq \left(\frac{1+t+\tau}{1+\tau}\right)^{2b_\infty|\alpha|} \left(\|\tilde{v}(0)\|_{L^2}^2 + \frac{\kappa_2^2 \rho^2 \alpha^2}{\eta(1+\tau)}\right), \tag{41}$$

for all $t > 0$. Now, we fix $t \geq 1$ and choose $\tau = Nt - 1$, where

$$N = N_{\alpha, \eta} = \max\left(1, \frac{2b_\infty|\alpha|}{\log(1+\eta)}\right).$$

This choice implies that

$$\left(\frac{1+t+\tau}{1+\tau}\right)^{2b_\infty|\alpha|} = \left(1 + \frac{1}{N}\right)^{2b_\infty|\alpha|} \leq 1 + \eta.$$

On the other hand, using (13), (40), we find

$$\|v(t)\|_{L^2}^2 \leq (1+\eta)\|\tilde{v}(t)\|_{L^2}^2 + \frac{1+\eta}{\eta}\alpha^2\|u^X(t) - u^X(t+\tau)\|_{L^2}^2 \leq (1+\eta)\|\tilde{v}(t)\|_{L^2}^2 + \frac{\alpha^2}{2\pi\eta} \log(N+1),$$

$$\|\tilde{v}(0)\|_{L^2}^2 \leq \frac{1+\eta}{\eta}\|v_0\|_{L^2}^2 + (1+\eta)\alpha^2\|u^X(0) - u^X(\tau)\|_{L^2}^2 \leq \frac{2}{\eta}\|v_0\|_{L^2}^2 + \frac{(1+\eta)\alpha^2}{4\pi} \log(Nt).$$

Similarly, using (17), we find

$$\begin{aligned} \int_0^t \|\nabla v(s)\|_{L^2}^2 ds &\leq 2 \int_0^t \|\nabla \tilde{v}(s)\|_{L^2}^2 ds + 2\alpha^2 \int_0^t \|\nabla u^X(s) - \nabla u^X(s+\tau)\|_{L^2}^2 ds \\ &\leq 2 \int_0^t \|\nabla \tilde{v}(s)\|_{L^2}^2 ds + \frac{\alpha^2}{4\pi} \log(1+t) + C\rho^2\alpha^2. \end{aligned}$$

Thus, it follows from (41) that

$$\|v(t)\|_{L^2}^2 \leq \frac{(1+\eta)^3\alpha^2}{4\pi} \log t + \frac{C}{\eta} (\|v_0\|_{L^2}^2 + \alpha^2 \log(N+1) + \alpha^2 \rho^2), \tag{42}$$

$$\int_0^t \|\nabla v(s)\|_{L^2}^2 ds \leq \frac{(1+\eta)^3\alpha^2}{2\pi} \log(1+t) + \frac{C}{\eta} (\|v_0\|_{L^2}^2 + \alpha^2 \rho^2) + C\alpha^2 \log N, \tag{43}$$

for some universal constant $C > 0$. Setting $\eta = 1$ and using the definition of N , we see that (39) follows from (42), (43). □

5. Estimate for a fractional primitive of the velocity field

In this final section, we consider the solution of (23) with initial data $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$, for some fixed $q \in (1, 2)$, and we define $\mu = 1/q - 1/2 \in (0, 1/2)$. If A is the Stokes operator in $L^2_\sigma(\Omega)$, we recall that A is selfadjoint and nonnegative in $L^2_\sigma(\Omega)$, so that the fractional power A^β can be defined for all $\beta > 0$. The following result shows that the range of A^μ contains the (dense) subspace $L^2_\sigma(\Omega) \cap L^q(\Omega)^2$.

Lemma 5.1 [Borchers and Miyakawa 1992; Kozono and Ogawa 1993a]. *Let $q \in (1, 2)$ and $\mu = 1/q - 1/2$. For all $v \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$, there exists a unique $w \in D(A^\mu) \subset L^2_\sigma(\Omega)$ such that $v = A^\mu w$. Moreover, there exists a constant $C = C(q) > 0$ (independent of v and Ω) such that $\|w\|_{L^2(\Omega)} \leq C \|v\|_{L^q(\Omega)}$.*

Remark 5.2. If v, w are as in Lemma 5.1, we define $w = A^{-\mu}v$. The fact that inequality $\|w\|_{L^2(\Omega)} \leq C \|v\|_{L^q(\Omega)}$ holds with a constant C independent of the domain Ω follows directly from the proof given in [Kozono and Ogawa 1993a, Lemmas 2.1 and 2.2].

As a first application of Lemma 5.1, we give a short proof of inequality (25), which was used in Section 3.

Proof of Proposition 3.1. It is sufficient to prove (25) for $1 < q < 2$. Let $\mu = 1/q - 1/2$, and let $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$. By Lemma 5.1, there exists a unique $w_0 \in D(A^\mu)$ such that $v_0 = A^\mu w_0$. Thus

$$\|S(t)v_0\|_{L^2(\Omega)} = \|A^\mu S(t)w_0\|_{L^2(\Omega)} \leq t^{-\mu} \|w_0\|_{L^2(\Omega)} \leq C t^{-\mu} \|v_0\|_{L^q(\Omega)},$$

with C depending only on q . The estimate for the first derivative is proved in the same way, since $\|\nabla S(t)v_0\|_{L^2(\Omega)} = \|A^{\mu+1/2}S(t)w_0\|_{L^2(\Omega)}$. This proves (25) for all $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$, and the general case follows by a density argument. \square

Let $v \in C^0([0, \infty); L^2_\sigma(\Omega)) \cap C^0((0, \infty); H^1_0(\Omega)^2)$ be the solution of (23) with initial data v_0 , which was constructed in Sections 3 and 4. Since $v_0 \in L^q_\sigma(\Omega)$ by assumption, it is rather straightforward to verify that $v(t) \in L^q_\sigma(\Omega)$ for all $t > 0$. Thus, by Lemma 5.1, we can define $w(t) = A^{-\mu}v(t)$ for all $t > 0$. This quantity solves the equation

$$\partial_t w + Aw + \alpha F_\mu(u^\chi, v) + \alpha F_\mu(v, u^\chi) + F_\mu(v, v) = \alpha A^{-\mu} R^\chi, \tag{44}$$

where $F_\mu(u, v)$ is the bilinear term formally defined by

$$F_\mu(u, v) = A^{-\mu} P(u \cdot \nabla)v. \tag{45}$$

We refer to [Kozono and Ogawa 1993a, Section 2] for a rigorous definition and a list of properties of the bilinear map F_μ . Our goal here is to establish the following estimate:

Proposition 5.3. *There exist positive constants K_2 and c (independent of Ω) such that, for any $\alpha \in \mathbb{R}$ and any solution v of (23) with initial data $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$, the function $w(t) = A^{-\mu}v(t)$ satisfies, for all $t \geq 1$,*

$$\|w(t)\|_{L^2}^2 + \int_0^t \|\nabla w(s)\|_{L^2}^2 ds \leq K_2(1+t)^{c\alpha^2} \exp(K_2(\|v_0\|_{L^2}^2 + D_{\alpha,\rho}))(\|v_0\|_{L^q}^2 + \rho^2\alpha^2), \tag{46}$$

where $D_{\alpha,\rho} = \alpha^2 \log(1 + |\alpha|) + \alpha^2 \rho^2$.

Proof. Taking the scalar product of both sides of (44) with w , we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w(t)\|_{L^2}^2 + \|A^{1/2} w(t)\|_{L^2}^2 + \alpha \langle F_\mu(u^\chi(t), v(t)), w(t) \rangle + \alpha \langle F_\mu(v(t), u^\chi(t)), w(t) \rangle \\ + \langle F_\mu(v(t), v(t)), w(t) \rangle = \alpha \langle A^{-\mu} R^\chi(t), w(t) \rangle. \end{aligned} \quad (47)$$

We recall that $\|A^{1/2} w\|_{L^2} = \|\nabla w\|_{L^2}$ for all $w \in D(A^{1/2}) = L^2_\sigma(\Omega) \cap H^1_0(\Omega)^2$. To bound the other terms, we observe that

$$\begin{aligned} |\langle F_\mu(u^\chi, v), w \rangle| &= |\langle (u^\chi \cdot \nabla)v, A^{-\mu} w \rangle| = |\langle (u^\chi \cdot \nabla)A^{-\mu} w, v \rangle| \leq \|u^\chi\|_{L^\infty} \|A^{\frac{1}{2}-\mu} w\|_{L^2} \|v\|_{L^2} \\ &= \|u^\chi\|_{L^\infty} \|A^{\frac{1}{2}-\mu} w\|_{L^2} \|A^\mu w\|_{L^2} \leq \|u^\chi\|_{L^\infty} \|A^{1/2} w\|_{L^2} \|w\|_{L^2}, \end{aligned}$$

where in the last inequality we used the interpolation inequality for fractional powers of A . The same argument shows that $|\langle F_\mu(v, u^\chi), w \rangle| \leq \|u^\chi\|_{L^\infty} \|A^{1/2} w\|_{L^2} \|w\|_{L^2}$. In a similar way, we find

$$\begin{aligned} |\langle F_\mu(v, v), w \rangle| &= |\langle (v \cdot \nabla)v, A^{-\mu} w \rangle| = |\langle (v \cdot \nabla)A^{-\mu} w, v \rangle| \leq \|v\|_{L^4}^2 \|A^{\frac{1}{2}-\mu} w\|_{L^2} \\ &\leq C_*^2 \|\nabla v\|_{L^2} \|v\|_{L^2} \|A^{\frac{1}{2}-\mu} w\|_{L^2} \leq C_*^2 \|\nabla v\|_{L^2} \|A^{1/2} w\|_{L^2} \|w\|_{L^2}, \end{aligned}$$

where $C_* > 0$ is the best constant of Gagliardo–Nirenberg’s inequality

$$\|f\|_{L^4(\mathbb{R}^2)} \leq C_* \|f\|_{L^2(\mathbb{R}^2)}^{\frac{1}{2}} \|\nabla f\|_{L^2(\mathbb{R}^2)}^{\frac{1}{2}}. \quad (48)$$

Finally, since $|\langle A^{-\mu} R^\chi, w \rangle| = |\langle R^\chi, A^{-\mu} w \rangle| \leq \kappa_2 \rho (1+t)^{-1} \|A^{\frac{1}{2}-\mu} w\|_{L^2}$ by (20), we can use interpolation and Young’s inequality to obtain

$$|\alpha \langle A^{-\mu} R^\chi, w \rangle| \leq \frac{\kappa_2 \rho |\alpha|}{1+t} \|A^{1/2} w\|_{L^2}^{1-2\mu} \|w\|_{L^2}^{2\mu} \leq \frac{\eta}{4} \|A^{1/2} w\|_{L^2}^2 + \frac{\|w\|_{L^2}^2}{2(1+t)^{\gamma_1}} + \frac{C_\eta \rho^2 \alpha^2}{2(1+t)^{\gamma_2}},$$

for some exponents $\gamma_1, \gamma_2 > 1$ satisfying $\gamma_2 + 2\mu\gamma_1 = 2$. Here $\eta \in (0, 1]$ is as in the proof of Proposition 4.1, and $C_\eta > 0$ denotes a constant depending only on η . Inserting all these estimates into (47), we arrive at

$$\frac{d}{dt} \|w\|_{L^2}^2 + 2\|\nabla w\|_{L^2}^2 \leq 2H \|\nabla w\|_{L^2} \|w\|_{L^2} + \frac{\eta}{2} \|\nabla w\|_{L^2}^2 + \frac{\|w\|_{L^2}^2}{(1+t)^{\gamma_1}} + \frac{C_\eta \rho^2 \alpha^2}{(1+t)^{\gamma_2}}, \quad (49)$$

where $H = 2|\alpha| \|u^\chi\|_{L^\infty} + C_*^2 \|\nabla v\|_{L^2}$.

To exploit (49), we apply Young’s inequality again and obtain the differential inequality

$$\frac{d}{dt} \|w\|_{L^2}^2 + \eta \|\nabla w\|_{L^2}^2 \leq \left(\frac{H^2}{2-3\eta/2} + \frac{1}{(1+t)^{\gamma_1}} \right) \|w\|_{L^2}^2 + \frac{C_\eta \rho^2 \alpha^2}{(1+t)^{\gamma_2}},$$

which can be integrated using Gronwall’s lemma. The result is

$$\|w(t)\|_{L^2}^2 + \eta \int_0^t \|\nabla w(s)\|_{L^2}^2 ds \leq C \exp\left(\frac{\Phi(t)}{1-3\eta/4}\right) (\|w_0\|_{L^2}^2 + C_\eta \rho^2 \alpha^2), \quad t \geq 0, \quad (50)$$

where $\Phi(t) = \frac{1}{2} \int_0^t H(s)^2 ds$ and C is a positive constant depending only on γ_1, γ_2 . It remains to estimate the quantity $\Phi(t)$ in (50). Using (11) with $p = \infty$, the logarithmic energy estimate (43), and Minkowski’s

inequality, we find

$$\begin{aligned}
 2\Phi(t) &= \int_0^t H(s)^2 \, ds \leq \int_0^t \left\{ \frac{2|\alpha|a_\infty}{(1+s)^{1/2}} + C_*^2 \|\nabla v(s)\|_{L^2} \right\}^2 \, ds \\
 &\leq \left\{ |\alpha| \log(1+t)^{1/2} \left(2a_\infty + \frac{C_*^2(1+\eta)^{\frac{3}{2}}}{\sqrt{2\pi}} \right) + C_\eta (\|v_0\|_{L^2} + D_{\alpha,\rho}^{1/2}) \right\}^2 \\
 &\leq 2C_0(1+\eta)^4 \alpha^2 \log(1+t) + C_\eta (\|v_0\|_{L^2}^2 + D_{\alpha,\rho}), \quad t \geq 1,
 \end{aligned} \tag{51}$$

where $D_{\alpha,\rho} = \alpha^2 \log(1 + |\alpha|) + \alpha^2 \rho^2$ and

$$C_0 = \frac{1}{2} \left(2a_\infty + \frac{C_*^2}{\sqrt{2\pi}} \right)^2. \tag{52}$$

If we now replace (51) into (50) and set $\eta = 1$, we obtain (46) since $\|w_0\|_{L^2} \leq C \|v_0\|_{L^q}$ by Lemma 5.1. This concludes the proof. \square

Corollary 5.4. *Under the assumptions of Proposition 5.3, there exists a positive constant K depending on Ω , α , and $\|v_0\|_{L^2 \cap L^q}$ such that, for any $T \geq 2$, there exists a time $t \in [T/2, T]$ for which*

$$\|v(t)\|_{L^2(\Omega)}^2 \leq K(1+t)^{c\alpha^2-2\mu}. \tag{53}$$

Proof. Fix $T \geq 2$. In view of (46), there exists a time $t \in [T/2, T]$ such that

$$\|\nabla w(t)\|_{L^2}^2 \leq \frac{2}{T} \int_{T/2}^T \|\nabla w(s)\|_{L^2}^2 \, ds \leq \frac{2}{T} C(1+T)^{c\alpha^2} \leq 2^{c\alpha^2+2} C(1+t)^{c\alpha^2-1},$$

where C depends on ρ , α , and $\|v_0\|_{L^2 \cap L^q}$. Moreover, $\|w(t)\|_{L^2}^2 \leq C(1+t)^{c\alpha^2}$ by (46). Thus, using the interpolation inequality $\|v(t)\|_{L^2} = \|A^\mu w(t)\|_{L^2} \leq \|\nabla w(t)\|_{L^2}^{2\mu} \|w(t)\|_{L^2}^{1-2\mu}$, we obtain (53). \square

Proof of Theorem 1.2. Fix $q \in (1, 2)$, and assume that $\epsilon > 0$ is small enough so that $c\epsilon^2 < 2\mu$, where $\mu = 1/q - 1/2$ and c is as in Proposition 5.3. We also suppose that $\epsilon \leq \delta$, where $\delta > 0$ is as in Proposition 3.2. Given $\alpha \in [-\epsilon, \epsilon]$ and $v_0 \in L_\sigma^2(\Omega) \cap L^q(\Omega)^2$, let $v \in C^0([0, \infty); L_\sigma^2(\Omega)) \cap C^0((0, \infty); H_0^1(\Omega)^2)$ be the solution of (23) with initial data $v(0) = v_0$, which was constructed in Sections 3 and 4. In view of (53), since $c\alpha^2 < 2\mu$, we can take $t_0 > 0$ large enough (depending on Ω , α , and v_0) so that $\|v(t_0)\|_{L^2} \leq V_\Omega$, where V_Ω is as in Proposition 3.2. Moreover, since $v(t_0) = A^\mu w(t_0)$ for some $w(t_0) \in L_\sigma^2(\Omega)$, we have

$$\sup_{\tau>0} \tau^\mu \|S(\tau)v_0\|_{L^2} + \sup_{\tau>0} \tau^{\mu+\frac{1}{2}} \|\nabla S(\tau)v_0\|_{L^2} \leq C \|w(t_0)\|_{L^2} < \infty.$$

Applying Proposition 3.2, we conclude that the solution v of (23) satisfies (28), namely

$$\|u(\cdot, t) - \alpha u^\chi(\cdot, t)\|_{L^2} + t^{1/2} \|\nabla u(\cdot, t) - \alpha \nabla u^\chi(\cdot, t)\|_{L^2(\Omega)} = \mathcal{O}(t^{-\mu}), \tag{54}$$

as $t \rightarrow \infty$. But $\|u^\chi - \Theta\|_{L^2} + \|\nabla u^\chi - \nabla \Theta\|_{L^2} \leq C(1+t)^{-1}$ for all $t \geq 0$; hence (6) follows from (54). \square

Proof of Corollary 1.4. The proof of Proposition 5.3 shows that the constant c in (46), (53) satisfies $c \leq C_0(1 + \mathcal{O}(\eta))$, where C_0 is defined in (52) and $\eta \in (0, 1]$ can be chosen arbitrarily small. On the other hand, since by assumption $v_0 \in L_\sigma^2(\Omega) \cap L^q(\Omega)^2$ for all $q \in (1, 2)$, we can take $\mu = 1/q - 1/2$

arbitrarily close to $1/2$. Thus, if we assume that $|\alpha| < \epsilon_* = C_0^{-1/2}$, we see that the condition $c\alpha^2 < 2\mu$ can be fulfilled by an appropriate choice of η and μ . Now, take $t \geq 2$ and let $t_0 \in [t/2, t]$ be the time defined in Corollary 5.4, for which $\|v(t_0)\|_{L^2}^2 \leq K(1+t_0)^{c\alpha^2-2\mu}$. Using (38) with $\eta = 1$, we conclude

$$\|v(t)\|_{L^2}^2 \leq C \left(\frac{1+t}{1+t_0} \right)^{2b_\infty|\alpha|} (\|v(t_0)\|_{L^2}^2 + (1+t_0)^{-1}) \leq C(1+t)^{c\alpha^2-2\mu} \xrightarrow{t \rightarrow \infty} 0,$$

which is the desired result. Here the constant $C > 0$ depends on α, ρ , and v_0 , but not on t . To estimate ϵ_* , we use (52) and observe that $a_\infty = \|\Theta_0\|_{L^\infty} \approx 0.050784$. Moreover, the optimal constant in the Gagliardo–Nirenberg inequality (48) satisfies $C_*^4 \leq 2/(3\pi)$; see [Del Pino and Dolbeault 2002]. Using these values, we find $C_0 \leq 0.0407108$; hence $\epsilon_* = C_0^{-1/2} \geq 4.95616$. Finally, it was kindly pointed out to us by Jean Dolbeault that the optimal constant C_* can be computed numerically: $C_* \approx 0.6430$. This yields the approximate value $\epsilon_* \approx 5.306$. □

Appendix: Proof of Proposition 1.3

We recall the following characterization of the space $\dot{W}_{0,\sigma}^{1,p}(\Omega)$ for $1 \leq p < 2$:

$$\dot{W}_{0,\sigma}^{1,p}(\Omega) = \left\{ u \in L^{\frac{2p}{2-p}}(\Omega)^2 \mid \|\nabla u\|_{L^p} < \infty, u = 0 \text{ on } \partial\Omega, \operatorname{div} u = 0 \text{ in } \Omega \right\} \tag{A-1}$$

(see, e.g., [Galdi 1994, Chapter III.5]). Here ∇u and $\operatorname{div} u$ denote weak derivatives of u , and the condition “ $u = 0$ on $\partial\Omega$ ” means that the boundary trace of u , which is well defined because $\nabla u \in L^p(\Omega)^4$, vanishes.

Given $u_0 \in \dot{W}_{0,\sigma}^{1,p}(\Omega)$ satisfying (7), we define $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $\omega : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by

$$u(x) = \begin{cases} u_0(x) & \text{if } x \in \Omega, \\ 0 & \text{if } x \notin \Omega, \end{cases} \quad \omega(x) = \begin{cases} \omega_0(x) & \text{if } x \in \Omega, \\ 0 & \text{if } x \notin \Omega. \end{cases}$$

Since $u = 0$ on $\partial\Omega$, we have $\nabla u \in L^p(\mathbb{R}^2)^4$ and $\partial_1 u_2 - \partial_2 u_1 = \omega \in L^p(\mathbb{R}^2)$. Moreover (7) implies that $\omega \in L^2(m)$ for some $m > 2/q > 1$, where

$$L^2(m) = \left\{ \omega \in L^2(\mathbb{R}^2) \mid \int_{\mathbb{R}^2} (1+|x|^2)^m |\omega(x)|^2 dx < \infty \right\}.$$

Thus, using Hölder’s inequality, it is easy to verify that $\omega \in L^1(\mathbb{R}^2)$, so that we can define

$$\alpha = \int_{\mathbb{R}^2} \omega(x) dx = \int_{\Omega} \omega_0(x) dx.$$

Moreover, using the Biot–Savart formula in \mathbb{R}^2 and the fact that $u \in L^{2p/(2-p)}(\mathbb{R}^2)^2$, we obtain the equality

$$u(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(x-y)^\perp}{|x-y|^2} \omega(y) dy = \frac{1}{2\pi} \int_{\Omega} \frac{(x-y)^\perp}{|x-y|^2} \omega_0(y) dy, \tag{A-2}$$

for almost all $x \in \mathbb{R}^2$. We emphasize at this point that the representation (A-2) is *not* what is usually called the Biot–Savart law in the domain Ω , because the velocity field defined by (A-2) for an arbitrary vorticity $\omega_0 \in L^1(\Omega)$ will not, in general, be tangent to the boundary on $\partial\Omega$. However, if we start from a

velocity field u_0 that vanishes on $\partial\Omega$, the argument above shows that (A-2) holds with $\omega_0 = \text{curl } u_0$. We refer to [Ifimie et al. 2003] for a more detailed discussion of the Biot–Savart law in a two-dimensional exterior domain.

Now, we decompose

$$u(x) = \alpha u^\chi(x, 0) + v(x), \quad \omega(x) = \alpha \omega^\chi(x, 0) + w(x), \quad x \in \mathbb{R}^2,$$

where u^χ, ω^χ are defined in (8), (9). By construction, we have $w \in L^2(m)$ and $\int_{\mathbb{R}^2} w \, dx = 0$. Applying [Gallay and Wayne 2002, Proposition B.1], we deduce that the corresponding velocity field v , which is obtained from w via the Biot–Savart law in \mathbb{R}^2 , satisfies

$$\int_{\mathbb{R}^2} (1 + |x|^2)^{\frac{mr}{2}-1} |v(x)|^r \, dx < \infty,$$

for all $r > 2$. Using Hölder’s inequality again, we conclude that $v \in L^s(\mathbb{R}^2)^2$ for all $s > 2/m$; hence in particular $v \in L^2(\mathbb{R}^2)^2 \cap L^q(\mathbb{R}^2)^2$. Clearly $v(x) = 0$ for all $x \notin \Omega$; hence denoting by v_0 the restriction of v to Ω we obtain (4) with $v_0 \in L^2_\sigma(\Omega) \cap L^q(\Omega)^2$. \square

Acknowledgements

This work was initiated during a visit of Y. M. at Université Joseph Fourier (Grenoble I), whose hospitality is gratefully acknowledged.

References

- [Bae and Jin 2006] H.-O. Bae and B. J. Jin, “Asymptotic behavior for the Navier–Stokes equations in 2D exterior domains”, *J. Funct. Anal.* **240**:2 (2006), 508–529. MR 2008e:35145 Zbl 1115.35095
- [Borchers and Miyakawa 1992] W. Borchers and T. Miyakawa, “ L^2 -decay for Navier–Stokes flows in unbounded domains, with application to exterior stationary flows”, *Arch. Rational Mech. Anal.* **118**:3 (1992), 273–295. MR 93i:35100 Zbl 0756.76018
- [Borchers and Varnhorn 1993] W. Borchers and W. Varnhorn, “On the boundedness of the Stokes semigroup in two-dimensional exterior domains”, *Math. Z.* **213**:2 (1993), 275–299. MR 94i:35160 Zbl 0794.35070
- [Brezis 1994] H. Brezis, “Remarks on the preceding paper by M. Ben-Artzi: “Global solutions of two-dimensional Navier–Stokes and Euler equations””, *Arch. Rational Mech. Anal.* **128**:4 (1994), 359–360. MR 96h:35149 Zbl 0837.35112
- [Carpio 1994] A. Carpio, “Asymptotic behavior for the vorticity equations in dimensions two and three”, *Comm. Partial Differential Equations* **19**:5-6 (1994), 827–872. MR 95f:35193 Zbl 0816.35108
- [Constantin and Foias 1988] P. Constantin and C. Foias, *Navier–Stokes equations*, University of Chicago Press, Chicago, IL, 1988. MR 90b:35190 Zbl 0687.35071
- [Dan and Shibata 1999a] W. Dan and Y. Shibata, “On the L_q – L_r estimates of the Stokes semigroup in a two-dimensional exterior domain”, *J. Math. Soc. Japan* **51**:1 (1999), 181–207. MR 99m:35185 Zbl 0924.35094
- [Dan and Shibata 1999b] W. Dan and Y. Shibata, “Remark on the L_q – L_∞ estimate of the Stokes semigroup in a 2-dimensional exterior domain”, *Pacific J. Math.* **189**:2 (1999), 223–239. MR 2000g:35167 Zbl 0931.35021
- [Del Pino and Dolbeault 2002] M. Del Pino and J. Dolbeault, “Best constants for Gagliardo–Nirenberg inequalities and applications to nonlinear diffusions”, *J. Math. Pures Appl.* (9) **81**:9 (2002), 847–875. MR 2003h:35051 Zbl 1112.35310
- [Fujita and Kato 1964] H. Fujita and T. Kato, “On the Navier–Stokes initial value problem, I”, *Arch. Rational Mech. Anal.* **16** (1964), 269–315. MR 29 #3774 Zbl 0126.42301
- [Galdi 1994] G. P. Galdi, *An introduction to the mathematical theory of the Navier–Stokes equations, I: Linearized steady problems*, Springer Tracts in Natural Philosophy **38**, Springer, New York, 1994. MR 95i:35216a Zbl 0949.35004

- [Gallay and Wayne 2002] T. Gallay and C. E. Wayne, “Invariant manifolds and the long-time asymptotics of the Navier–Stokes and vorticity equations on \mathbb{R}^2 ”, *Arch. Ration. Mech. Anal.* **163**:3 (2002), 209–258. MR 2003c:37123 Zbl 1042.37058
- [Gallay and Wayne 2005] T. Gallay and C. E. Wayne, “Global stability of vortex solutions of the two-dimensional Navier–Stokes equation”, *Comm. Math. Phys.* **255**:1 (2005), 97–129. MR 2005m:35224 Zbl 1139.35084
- [Giga and Kambe 1988] Y. Giga and T. Kambe, “Large time behavior of the vorticity of two-dimensional viscous flow and its application to vortex formation”, *Comm. Math. Phys.* **117**:4 (1988), 549–568. MR 89k:35182 Zbl 0661.76018
- [He and Miyakawa 2006] C. He and T. Miyakawa, “Nonstationary Navier–Stokes flows in a two-dimensional exterior domain with rotational symmetries”, *Indiana Univ. Math. J.* **55**:5 (2006), 1483–1555. MR 2008a:35210 Zbl 1122.35095
- [Iftimie et al. 2003] D. Iftimie, M. C. Lopes Filho, and H. J. Nussenzweig Lopes, “Two dimensional incompressible ideal flow around a small obstacle”, *Comm. Partial Differential Equations* **28**:1-2 (2003), 349–379. MR 2004d:76009 Zbl 1094.76007
- [Iftimie et al. 2011] D. Iftimie, G. Karch, and C. Lacave, “Self-similar asymptotics of solutions to the Navier–Stokes system in two dimensional exterior domain”, preprint, 2011. arXiv 1107.205
- [Kato and Fujita 1962] T. Kato and H. Fujita, “On the nonstationary Navier–Stokes system”, *Rend. Sem. Mat. Univ. Padova* **32** (1962), 243–260. MR 26 #495 Zbl 0114.05002
- [Kozono and Ogawa 1993a] H. Kozono and T. Ogawa, “Decay properties of strong solutions for the Navier–Stokes equations in two-dimensional unbounded domains”, *Arch. Rational Mech. Anal.* **122**:1 (1993), 1–17. MR 94d:35130 Zbl 0781.35053
- [Kozono and Ogawa 1993b] H. Kozono and T. Ogawa, “Two-dimensional Navier–Stokes flow in unbounded domains”, *Math. Ann.* **297**:1 (1993), 1–31. MR 94h:35207 Zbl 0796.35129
- [Kozono and Yamazaki 1995] H. Kozono and M. Yamazaki, “Local and global unique solvability of the Navier–Stokes exterior problem with Cauchy data in the space $L^{p,\infty}$ ”, *Houston J. Math.* **21**:4 (1995), 755–799. MR 97a:35175 Zbl 0848.35099
- [Ladyženskaja 1959] O. A. Ladyženskaja, “Solution “in the large” of the nonstationary boundary value problem for the Navier–Stokes system with two space variables”, *Comm. Pure Appl. Math.* **12** (1959), 427–433. MR 21 #7674 Zbl 0103.19502
- [Leray 1933] J. Leray, “Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l’hydrodynamique”, *J. Math. Pures Appl. (9)* **12** (1933), 1–82. Zbl 0006.16702
- [Leray 1934] J. Leray, “Essai sur les mouvements plans d’une liquide visqueux que limitent des parois”, *J. Math. Pures Appl. (9)* **13** (1934), 331–418. JFM 60.0727.01
- [Lions and Prodi 1959] J.-L. Lions and G. Prodi, “Un théorème d’existence et unicité dans les équations de Navier–Stokes en dimension 2”, *C. R. Acad. Sci. Paris* **248** (1959), 3519–3521. MR 21 #7676 Zbl 0091.42105
- [Majda and Bertozzi 2002] A. J. Majda and A. L. Bertozzi, *Vorticity and incompressible flow*, Cambridge Texts in Applied Mathematics **27**, Cambridge University Press, 2002. MR 2003a:76002 Zbl 0983.76001
- [Maremonti and Solonnikov 1997] P. Maremonti and V. A. Solonnikov, “On nonstationary Stokes problem in exterior domains”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **24**:3 (1997), 395–449. MR 99c:35092 Zbl 0958.35103
- [Masuda 1984] K. Masuda, “Weak solutions of Navier–Stokes equations”, *Tohoku Math. J. (2)* **36**:4 (1984), 623–646. MR 86a:35117 Zbl 0568.35077
- [Sohr 2001] H. Sohr, *The Navier–Stokes equations: An elementary functional analytic approach*, Birkhäuser, Basel, 2001. MR 2004b:35265 Zbl 0983.35004

Received 28 Feb 2012. Accepted 6 Aug 2012.

THIERRY GALLAY: Thierry.Gallay@ujf-grenoble.fr

Institut Fourier, Université de Grenoble I, 100 rue des Maths, B.P. 74, 38402 Saint-Martin-d’Hères, France

<http://www-fourier.ujf-grenoble.fr/~gallay>

YASUNORI MAEKAWA: yasunori@math.kobe-u.ac.jp

Department of Mathematics, Graduate School of Science, Kobe University, 1-1 Rokkodai, Nada-ku, Kobe 657-8501, Japan

<http://www.math.kobe-u.ac.jp/home-j/yasunori.html>

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Michael Aizenman	Princeton University, USA aizenman@math.princeton.edu	Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr
Luis A. Caffarelli	University of Texas, USA caffarel@math.utexas.edu	Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Charles Fefferman	Princeton University, USA cf@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Nigel Higson	Pennsylvania State University, USA higson@math.psu.edu
Vaughan Jones	University of California, Berkeley, USA vfr@math.berkeley.edu	Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr
László Lempert	Purdue University, USA lempert@math.purdue.edu	Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Igor Rodnianski	Princeton University, USA irod@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 6 No. 4 2013

Cauchy problem for ultrasound-modulated EIT GUILLAUME BAL	751
Sharp weighted bounds involving A_∞ TUOMAS HYTÖNEN and CARLOS PÉREZ	777
Periodicity of the spectrum in dimension one ALEX IOSEVICH and MIHAL N. KOLOUNTZAKIS	819
A codimension-two stable manifold of near soliton equivariant wave maps IOAN BEJENARU, JOACHIM KRIEGER and DANIEL TATARU	829
Discrete Fourier restriction associated with KdV equations YI HU and XIAOCHUN LI	859
Restriction and spectral multiplier theorems on asymptotically conic manifolds COLIN GUILLARMOU, ANDREW HASSELL and ADAM SIKORA	893
Homogenization of Neumann boundary data with fully nonlinear operator SUNHI CHOI, INWON C. KIM and KI-AHM LEE	951
Long-time asymptotics for two-dimensional exterior flows with small circulation at infinity THIERRY GALLAY and YASUNORI MAEKAWA	973
Second order stability for the Monge–Ampère equation and strong Sobolev convergence of optimal transport maps GUIDO DE PHILIPPIS and ALESSIO FIGALLI	993



2157-5045(2013)6:4;1-E