

ANALYSIS & PDE

Volume 7

No. 3

2014

EDUARDO V. TEIXEIRA AND JOSÉ MIGUEL URBANO

**A GEOMETRIC TANGENTIAL APPROACH TO SHARP
REGULARITY
FOR DEGENERATE EVOLUTION EQUATIONS**

A GEOMETRIC TANGENTIAL APPROACH TO SHARP REGULARITY FOR DEGENERATE EVOLUTION EQUATIONS

EDUARDO V. TEIXEIRA AND JOSÉ MIGUEL URBANO

That the weak solutions of degenerate parabolic PDEs modelled on the inhomogeneous p -Laplace equation

$$u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = f \in L^{q,r}, \quad p > 2$$

are $C^{0,\alpha}$, for some $\alpha \in (0, 1)$, has been known for almost 30 years. What was hitherto missing from the literature was a precise and sharp knowledge of the Hölder exponent α in terms of p, q, r and the space dimension n . We show in this paper that

$$\alpha = \frac{(pq - n)r - pq}{q[(p - 1)r - (p - 2)]}$$

using a method based on the notion of geometric tangential equations and the intrinsic scaling of the p -parabolic operator. The proofs are flexible enough to be of use in a number of other nonlinear evolution problems.

1. Introduction

The understanding of the local behaviour of solutions to singular and degenerate parabolic equations has witnessed an impressive progress in the last three decades. At the heart of most developments lies a single unifying idea, namely that regularity results have to be interpreted in an intrinsic geometric configuration, a sort of signature to each particular PDE. The pioneering work of DiBenedetto [1993] was the starting point to a theory that has, in many aspects, reached its maturity (see [DiBenedetto et al. 2012] and [Urbano 2008] for recent accounts).

A central aspect in this endeavour has always been the Hölder continuity of bounded weak solutions, which ultimately follows from Harnack-type inequalities. Although powerful, this approach only provides qualitative estimates that depend solely on the structure of the equations and thus hold in a very general setting. The quest for precise, quantitative derivations of the Hölder exponent has hitherto eluded the community, the only exception being the two-dimensional result in [Iwaniec and Manfredi 1989] concerning p -harmonic functions. This type of quantitative information, apart from its own intrinsic value, plays an important role in the analysis of a number of qualitative issues for parabolic PDEs, such as blow-up analysis, Liouville type results, free boundary problems, and so forth.

MSC2010: 35K55, 35K65, 35B65.

Keywords: degenerate parabolic equations, sharp Hölder regularity, tangential equations, intrinsic scaling.

The main goal of this paper is to fill this gap, bringing the theory to a new level of understanding. We show that weak solutions of degenerate p -parabolic equations whose prototype is

$$u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = f \in L^{q,r}, \quad p \geq 2, \quad (1)$$

are locally of class $C^{0,\alpha}$ in space, with

$$\alpha := \frac{(pq - n)r - pq}{q[(p - 1)r - (p - 2)]}$$

a precise and sharp expression for the Hölder exponent in terms of p , the integrability of the source and the space dimension n . We also show that u is of class $C^{0,\alpha/\theta}$ in time, where θ is the α -interpolation between 2 and p . What makes the parabolic case more delicate to analyse is the inhomogeneity in the equation, the fact that it scales differently with respect to space and time. It is worth stressing that the integrability in time (respectively, in space) of the source affects the regularity in space (respectively, in time) of the solution.

To highlight the extent to which our result is sharp, we project it into the state of the art of the theory. For the linear case $p = 2$, we obtain

$$\alpha = 1 - \left(\frac{2}{r} + \frac{n}{q} - 1 \right),$$

which is the optimal Hölder exponent for the nonhomogeneous heat equation, and is in accordance with estimates obtained by energy considerations. When $p \rightarrow \infty$, we have $\alpha \rightarrow 1^-$, which gives an indication of the expected locally Lipschitz regularity for the case of the parabolic infinity-Laplacian. When the source f is independent of time, or else bounded in time, that is $r = \infty$, we obtain

$$\alpha = \frac{pq - n}{q(p - 1)} = \frac{p}{p - 1} \cdot \frac{q - n/p}{q},$$

which is exactly the optimal exponent obtained in [Teixeira 2013] for the elliptic case. It might also be interesting to compare our optimal result with the estimates from [Misawa 2013, Section 4], and also with the continuity estimates on p -parabolic obstacle problems from [Kuusi et al. 2014].

Within the general theory of p -parabolic equations, our result reveals a surprising feature. From the applied point of view, it is relevant to know what is the effect on the diffusion properties of the model as we dim the exponent p . Naïve physical interpretations could indicate that the higher the value of p , the less efficient should the diffusion properties of the p -parabolic operator turn out to be, *i.e.*, one should expect a less efficient smoothness effect of the operator. For instance, this is verified in the sharp regularity estimate for p -harmonic functions in 2D [Iwaniec and Manfredi 1989]. On the contrary, our estimate implies that for p -parabolic inhomogeneous equations, the Hölder regularity theory improves as p increases. In fact, a direct computation shows

$$\operatorname{sign}(\partial_p \alpha(p, n, q, r)) = \operatorname{sign}(q(2 - r) + nr) = +1,$$

in view of standard assumptions on the integrability exponents of the source term.

Although regularity estimates for degenerate evolution equations have been successfully obtained in great generality (see [Kinnunen and Lewis 2000; Acerbi and Mingione 2007]), explicit expressions for the Hölder exponent of continuity for weak solutions have only been known in the linear setting. For nonlinear equations, the classical tools from harmonic analysis, such as singular integrals, are precluded from being used and an entirely new approach is needed. The new estimates we obtain are striking in their simplicity but perhaps the most relevant contribution we offer is the technique employed. We develop a method based on the notion of geometric tangential equations, which explores the intrinsic scaling of the p -parabolic operator and the integrability of the forcing term. By means of appropriate scaled iterative arguments, we show that at each inhomogeneous equation there is a universal tangential space formed by $C^{0,1}$ in space and $C^{0,1/2}$ in time functions. The method then imports such regularity back to the original equation, properly corrected through the scaling used to access the tangential space. The method is new to the field and robust enough to be adapted to other evolutionary problems, as well as to a number of other issues in the theory.

2. Preliminary tools

Let $U \subset \mathbb{R}^n$ be open and bounded, and $T > 0$. We consider the space-time domain $U_T = U \times (0, T)$. We work with the prototype inhomogeneous equation

$$u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = f \quad \text{in } U_T, \quad (2)$$

with a source term $f \in L^{q,r}(U_T) \equiv L^r(0, T; L^q(U))$ satisfying

$$\frac{1}{r} + \frac{n}{pq} < 1 \quad (3)$$

and

$$\frac{2}{r} + \frac{n}{q} > 1. \quad (4)$$

The first assumption is the standard minimal integrability condition that guarantees the existence of bounded weak solutions, while (4) defines the borderline setting for optimal Hölder type estimates. For instance, when $r = \infty$, conditions (3) and (4) enforce

$$\frac{n}{p} < q < n,$$

which corresponds to the known range of integrability required in the elliptic theory for local $C^{0,\alpha}$ estimates to be available.

We start with the definition of weak solution to (2).

Definition 2.1. We say a function

$$u \in C_{\text{loc}}(0, T; L_{\text{loc}}^2(U)) \cap L_{\text{loc}}^p(0, T; W_{\text{loc}}^{1,p}(U))$$

is a weak solution to (2) if, for every compact $K \subset U$ and every subinterval $[t_1, t_2] \subset (0, T]$, there holds

$$\int_K u\varphi dx \Big|_{t_1}^{t_2} + \int_{t_1}^{t_2} \int_K \{-u\varphi_t + |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi\} dx dt = \int_{t_1}^{t_2} \int_K f\varphi dx dt,$$

for all $\varphi \in H_{loc}^1(0, T; L^2(K)) \cap L_{loc}^p(0, T; W_0^{1,p}(K))$.

The following alternative definition makes use of the Steklov average of a function $v \in L^1(U_T)$, defined for $0 < h < T$ by

$$v_h := \begin{cases} \frac{1}{h} \int_t^{t+h} v(\cdot, \tau) d\tau & \text{if } t \in (0, T - h], \\ 0 & \text{if } t \in (T - h, T], \end{cases}$$

and circumvents the difficulties related to the low regularity in time. In fact, these difficulties are more of a technical nature since the time derivative u_t is shown in [Lindqvist 2008] to be an element of a certain Lebesgue space.

Definition 2.2. We say a function

$$u \in C_{loc}(0, T; L_{loc}^2(U)) \cap L_{loc}^p(0, T; W_{loc}^{1,p}(U))$$

is a weak solution to (2) if, for every compact $K \subset U$ and every $0 < t < T - h$, there holds

$$\int_{K \times \{t\}} \{(u_h)_t \varphi + (|\nabla u|^{p-2} \nabla u)_h \cdot \nabla \varphi\} dx = \int_{K \times \{t\}} f_h \varphi dx, \tag{5}$$

for all $\varphi \in W_0^{1,p}(K)$.

One key ingredient in our analysis is the following Caccioppoli-type energy estimate enjoyed by weak solutions of (2).

Lemma 2.3 (Caccioppoli estimate). *Let u be a weak solution to (2). Given $K \times [t_1, t_2] \subset U \times (0, T]$, there exists a constant C , depending only on $n, p, K \times [t_1, t_2]$ and $\|f\|_{L^{q,r}}$, such that*

$$\begin{aligned} \sup_{t_1 < t < t_2} \int_K u^2 \xi^p dx + \int_{t_1}^{t_2} \int_K |\nabla u|^p \xi^p dx dt \\ \leq \int_{t_1}^{t_2} \int_K |u|^p (\xi^p + |\nabla \xi|^p) dx dt + \int_{t_1}^{t_2} \int_K u^2 \xi^{p-1} |\xi_t| dx dt + \|f\|_{q,r} \end{aligned} \tag{6}$$

for every $\xi \in \mathcal{C}_0^\infty(K \times (t_1, t_2))$ such that $\xi \in [0, 1]$.

Proof. Choose $\varphi = u_h \xi^p$ as a test function in (5) and perform the usual combination of integrating in time, passing to the limit in $h \rightarrow 0$ and applying Young’s inequality to derive the estimate. \square

We finally recall that, if v is a function belonging to $L^p(Q)$, its averaged norm is

$$\|v\|_{p, \text{avg}, Q} := \left(\int_Q |v|^p dx dt \right)^{1/p} = |Q|^{-1/p} \|v\|_{p, Q},$$

where, as usual, the integral average is defined by

$$\int_A \psi = \frac{1}{|A|} \int_A \psi.$$

3. Sharp Hölder estimate

We start by fixing universal constants, that depend only on the data. The intrinsic exponent to (2), with $f \in L^{q,r}$, is

$$\alpha := \frac{(pq - n)r - pq}{q[(p - 1)r - (p - 2)]} = \frac{p\left(1 - \frac{1}{r} - \frac{n}{pq}\right)}{\left(\frac{2}{r} + \frac{n}{q} - 1\right) + p\left(1 - \frac{1}{r} - \frac{n}{pq}\right)}, \tag{7}$$

which, in view of (3) and (4), satisfies $0 < \alpha < 1$. Next, let

$$\theta := \alpha + p - (p - 1)\alpha = p - (p - 2)\alpha = \alpha 2 + (1 - \alpha)p. \tag{8}$$

Clearly $2 < \theta < p$, since $0 < \alpha < 1$. For such θ , we define the intrinsic θ -parabolic cylinder

$$G_\tau := (-\tau^\theta, 0) \times B_\tau(0), \quad \tau > 0.$$

We first establish a key compactness result that states that if the source term f has a small norm in $L^{q,r}$, then a solution u to (2) is close to a p -caloric function in an inner subdomain. It is worth comparing such a result with the A -caloric approximation lemma obtained in [Duzaar and Mingione 2005, Lemma 4.1].

Lemma 3.1 (approximation to p -caloric functions). *For every $\delta > 0$, there exists $0 < \epsilon \ll 1$, such that if $\|f\|_{L^{q,r}(G_1)} \leq \epsilon$ and u is a local weak solution of (2) in G_1 , with $\|u\|_{p,\text{avg},G_1} \leq 1$, then there exists a ϕ that is p -caloric in $G_{1/2}$ in the sense that*

$$\phi_t - \text{div}(|\nabla\phi|^{p-2}\nabla\phi) = 0 \quad \text{in } G_{1/2}, \tag{9}$$

and moreover satisfies

$$\|u - \phi\|_{p,\text{avg},G_{1/2}} \leq \delta. \tag{10}$$

Proof. Suppose, for the sake of contradiction, that the thesis of the lemma fails. That is, assume, for some $\delta_0 > 0$, that there exists a sequence

$$(u^j)_j \in C_{\text{loc}}(-1, 0; L^2_{\text{loc}}(B_1)) \cap L^p_{\text{loc}}(-1, 0; W^{1,p}_{\text{loc}}(B_1))$$

and a sequence $(f^j)_j \in L^{q,r}(G_1)$ such that

$$u^j_t - \text{div}(|\nabla u^j|^{p-2}\nabla u^j) = f^j \quad \text{in } G_1, \tag{11}$$

$$\|u^j\|_{p,\text{avg},G_1} \leq 1, \tag{12}$$

$$\|f^j\|_{L^{q,r}(G_1)} \leq 1/j, \tag{13}$$

but still, for any j and any p -caloric function ϕ in $G_{1/2}$,

$$\|u^j - \phi\|_{p,\text{avg},G_{1/2}} > \delta_0. \tag{14}$$

Fix a cutoff function $\xi \in C_0^\infty(G_1)$, such that $\xi \in [0, 1]$, $\xi \equiv 1$ in $G_{1/2}$ and $\xi \equiv 0$ near $\partial_p G_1$. From the Caccioppoli estimate, using the notation

$$V(I \times U) = L^\infty(I; L^2(U)) \cap L^p(I; W^{1,p}(U)),$$

we obtain

$$\begin{aligned} \|u^j\|_{V(G_{1/2})} &\leq \sup_{-1 < t < 0} \int_{B_1} (u^j)^2 \xi^p dx + \int_{-1}^0 \int_{B_1} |\nabla u^j|^p \xi^p dx dt \\ &\leq \int_{-1}^0 \int_{B_1} \{|u^j|^p (\xi^p + |\nabla \xi|^p) + (u^j)^2 \xi^{p-1} |\xi_t|\} dx dt + \|f^j\|_{L^{q,r}(G_1)} \\ &\leq c \|u^j\|_{p,\text{avg},G_1}^p + c' \|u^j\|_{2,\text{avg},G_1}^2 + \frac{1}{j} \\ &\leq c. \end{aligned}$$

A control of the time derivative, along the lines of [Lindqvist 2008] (see also [Acerbi et al. 2004]), gives

$$\|u_t^j\|_{L^{s,1}(G_{1/2})} \leq c,$$

with $s = \min\{q, p/(p-1)\} < p$. We now use a classical compactness result (see [Simon 1987, Corollary 4]), with

$$W^{1,p} \hookrightarrow L^p \subset L^s,$$

to conclude that

$$u^j \rightarrow \psi,$$

strongly in $L^p(G_{1/2})$, in addition to the weak convergence in $V(G_{1/2})$.

Passing to the limit in (11), we find that

$$\psi_t - \operatorname{div}(|\nabla \psi|^{p-2} \nabla \psi) = 0 \quad \text{in } G_{1/2},$$

which contradicts (14), for $j \gg 1$. The proof is complete. □

Next, by means of geometric iteration, we shall establish the optimal Hölder continuity for solutions to the heterogeneous p -parabolic equation (2). Our approach explores the approximation by p -caloric functions, given by Lemma 3.1, and the fact that p -caloric functions are *universally* Lipschitz continuous in space and $C^{0,1/2}$ in time. The following is the crucial first iterative step.

Lemma 3.2. *Let $0 < \alpha < 1$ be fixed. There exists $\epsilon > 0$ and $0 < \lambda \ll 1/2$, depending only on p, n and α , such that if $\|f\|_{L^{q,r}(G_1)} \leq \epsilon$ and u is a local weak solution of (2) in G_1 , with $\|u\|_{p,\text{avg},G_1} \leq 1$, then there exists a universally bounded constant c_0 such that*

$$\|u - c_0\|_{p,\text{avg},G_\lambda} \leq \lambda^\alpha. \tag{15}$$

Proof. Take $0 < \delta < 1$, to be chosen later, and apply Lemma 3.1 to obtain $0 < \epsilon \ll 1$ and a p -caloric function ϕ in $G_{1/2}$, such that

$$\|u - \phi\|_{p,\text{avg},G_{1/2}} \leq \delta.$$

Observe that

$$\|\phi\|_{p,\text{avg},G_{1/2}} \leq \|u - \phi\|_{p,\text{avg},G_{1/2}} + \|u\|_{p,\text{avg},G_1} \leq \delta + 2^{(\theta+n)/p} \leq C. \tag{16}$$

Since ϕ is p -caloric, it follows from standard theory that ϕ is universally $C_{\text{loc}}^{0,1/2}$ in time and $C_{\text{loc}}^{0,1}$ in space. That is, for $\lambda \ll 1$, to be chosen soon, we have

$$\sup_{(x,t) \in G_\lambda} |\phi(x,t) - \phi(0,0)| \leq C \lambda,$$

for $C > 1$ universal. In fact, for $(x,t) \in G_\lambda$,

$$\begin{aligned} |\phi(x,t) - \phi(0,0)| &\leq |\phi(x,t) - \phi(0,t)| + |\phi(0,t) - \phi(0,0)| \\ &\leq C' |x - 0| + C'' |t - 0|^{1/2} \\ &\leq C' \lambda + C'' \lambda^{\theta/2} \leq C \lambda, \end{aligned}$$

since $\theta > 2$. We can therefore estimate

$$\begin{aligned} \|u(x,t) - \phi(0,0)\|_{p,\text{avg},G_\lambda} &\leq \|u(x,t) - \phi(x,t)\|_{p,\text{avg},G_\lambda} + \|\phi(x,t) - \phi(0,0)\|_{p,\text{avg},G_\lambda} \\ &\leq \left(\frac{1}{2\lambda}\right)^{\frac{\theta+n}{p}} \delta + C \lambda. \end{aligned} \tag{17}$$

Note that we will choose $\lambda \ll 1/2$ and thus

$$G_\lambda = (-\lambda^\theta, 0) \times B_\lambda \subset (-(1/2)^\theta, 0) \times B_{1/2} = G_{1/2}.$$

We put $c_0 := \phi(0,0)$, observing that, due to (16) and the fact that ϕ is p -caloric, c_0 is universally bounded. The next step is to fix the constants. We choose $\lambda \ll \frac{1}{2}$ so small that

$$C \lambda \leq \frac{1}{2} \lambda^\alpha,$$

and then we define

$$\delta = \frac{1}{2} \lambda^\alpha (2\lambda)^{(\theta+n)/p},$$

thus fixing, via Lemma 3.1, also $\epsilon > 0$. The lemma now follows from estimate (17) with the indicated choices. □

Our next step involves iterating Lemma 3.2 in the appropriate geometric scaling.

Theorem 3.3. *Under the conditions of the previous lemma, there exists a convergent sequence of real numbers $\{c_k\}_{k \geq 1}$, with*

$$|c_k - c_{k+1}| \leq c(n,p)(\lambda^\alpha)^k, \tag{18}$$

such that

$$\|u - c_k\|_{p,\text{avg},G_{\lambda^k}} \leq (\lambda^k)^\alpha. \tag{19}$$

Proof. The proof is by induction on $k \in \mathbb{N}$. For $k = 1$, (19) holds due to Lemma 3.2, with $c_1 = c_0$. Suppose the conclusion holds for k and let's show it also holds for $k + 1$. We start by defining the function $v: G_1 \rightarrow \mathbb{R}$ by

$$v(x, t) = \frac{u(\lambda^k x, \lambda^{k\theta} t) - c_k}{\lambda^{\alpha k}}. \quad (20)$$

We compute

$$v_t(x, t) = \lambda^{k\theta - \alpha k} u_t(\lambda^k x, \lambda^{k\theta} t)$$

and

$$\operatorname{div}(|\nabla v(x, t)|^{p-2} \nabla v(x, t)) = \lambda^{pk - (p-1)\alpha k} \operatorname{div}(|\nabla u(\lambda^k x, \lambda^{k\theta} t)|^{p-2} \nabla u(\lambda^k x, \lambda^{k\theta} t))$$

to conclude, recalling (8), that

$$v_t - \operatorname{div}(|\nabla v|^{p-2} \nabla v) = \lambda^{pk - (p-1)\alpha k} f(\lambda^k x, \lambda^{k\theta} t) = \tilde{f}(x, t).$$

We now compute

$$\begin{aligned} \|\tilde{f}\|_{L^{q,r}(G_1)}^r &= \int_{-1}^0 \left(\int_{B_1} |\tilde{f}(x, t)|^q dx \right)^{r/q} dt \\ &= \int_{-1}^0 \left(\int_{B_1} \lambda^{(pk - (p-1)\alpha k)q} |f(\lambda^k x, \lambda^{k\theta} t)|^q dx \right)^{r/q} dt \\ &= \int_{-1}^0 \left(\int_{B_{\lambda^k}} \lambda^{(pk - (p-1)\alpha k)q - kn} |f(x, \lambda^{k\theta} t)|^q dx \right)^{r/q} dt \\ &= \lambda^{((pk - (p-1)\alpha k)q - kn)\frac{r}{q}} \int_{-1}^0 \left(\int_{B_{\lambda^k}} |f(x, \lambda^{k\theta} t)|^q dx \right)^{r/q} dt \\ &= \lambda^{((pk - (p-1)\alpha k)q - kn)\frac{r}{q} - k\theta} \int_{-\lambda^{k\theta}}^0 \left(\int_{B_{\lambda^k}} |f(x, t)|^q dx \right)^{r/q} dt. \end{aligned} \quad (21)$$

Due to the crucial and sharp choice (7) of α , we have, recalling again (8),

$$((pk - (p-1)\alpha k)q - kn)\frac{r}{q} - k\theta = 0.$$

We go back to (21) to conclude

$$\|\tilde{f}\|_{L^{q,r}(G_1)} = \|f\|_{L^{q,r}((-\lambda^{k\theta}, 0) \times B_{\lambda^k})} \leq \|f\|_{L^{q,r}(G_1)} \leq \epsilon,$$

which entitles v to Lemma 3.2 (note that $\|v\|_{p,\text{avg},G_1} \leq 1$, due to the induction hypothesis).

It then follows that there exists a constant \tilde{c}_0 , with $|\tilde{c}_0| \leq c(n, p)$, such that

$$\|v - \tilde{c}_0\|_{p,\text{avg},G_\lambda} \leq \lambda^\alpha,$$

which is the same as

$$\|u - c_{k+1}\|_{p,\text{avg},G_{\lambda^{k+1}}} \leq \lambda^{\alpha(k+1)},$$

for $c_{k+1} := c_k + \tilde{c}_0 \lambda^{\alpha k}$; the induction is complete. We readily observe that

$$|c_{k+1} - c_k| \leq c(n, p)(\lambda^\alpha)^k,$$

thus obtaining also (18). □

Theorem 3.4. *A locally bounded weak solution of (2), with $f \in L^{q,r}$, satisfying (3)–(4), is locally Hölder continuous in the space variables, with exponent*

$$\alpha = \frac{(pq - n)r - pq}{q[(p - 1)r - (p - 2)]}$$

and locally Hölder continuous in time with exponent α/θ . In addition, there exists a constant C , that depends only on $p, n, \|f\|_{q,r}$ and $\|u\|_{p,\text{avg},G_1}$, such that

$$\|u\|_{C^{0;\alpha,\alpha/\theta}(G_{1/2})} \leq C.$$

Proof. We start by observing (see also [Araújo et al. 2013, Section 7]) that the smallness regime required in the assumptions of Theorem 3.3 is not restrictive since we can fall into that framework by scaling and contraction. Indeed, given a solution u , let

$$v(x, t) = \varrho u(\varrho^a x, \varrho^{(p-2)+ap} t)$$

(ϱ, a to be fixed), which is a solution of (2) with

$$\tilde{f}(x, t) = \varrho^{(p-1)+ap} f(\varrho^a x, \varrho^{(p-2)+ap} t).$$

We choose $a > 0$ such that

$$a < \frac{2}{n+p} \quad \text{and} \quad [(p-1)+ap]r - a(n+p) - (p-2) > 0,$$

which is always possible (observe that the second condition holds for $a = 0$ and use its continuity with respect to a), and then $0 < \varrho < 1$ such that

$$\|v\|_{p,\text{avg},G_1}^p \leq \varrho^{2-a(n+p)} \|u\|_{p,\text{avg},G_1}^p \leq 1$$

and

$$\|\tilde{f}\|_{L^{q,r}(G_1)}^r = \varrho^{[(p-1)+ap]r - a(n+p) - (p-2)} \|f\|_{L^{q,r}(G_1)}^r \leq \epsilon^r.$$

Due to (18), the sequence $\{c_k\}_{k \geq 1}$ is convergent and we put

$$\bar{c} := \lim_{k \rightarrow \infty} c_k.$$

It follows from (19) that, for arbitrary $0 < r < 1/2$,

$$\int_{G_r} |u - \bar{c}|^p dx dt \leq Cr^{p\alpha}.$$

Standard covering arguments, a remark in [Teixeira 2013, Lemma 3.2] and the characterisation of Hölder continuity of Campanato–Da Prato give the local $C^{0;\alpha,\alpha/\theta}$ -continuity and thus the result. □

4. Generalisations and beyond

The ideas and methods employed in this paper only explore the degenerate p -structure of the operator. The underlying heuristics is to interpret the homogeneous problem as the geometric tangential equation of its inhomogeneous counterpart, for small perturbations $f \in L^{r,q}$, $\|f\|_{r,q} \ll 1$. The proofs adapt to more general degenerate parabolic equations

$$u_t - \operatorname{div} \mathcal{A}(x, t, Du) = f \in L^{r,q} \quad (22)$$

satisfying the usual structure assumptions for $p \geq 2$.

We briefly comment on the modifications required. Lemma 2.3 is based on pure energy considerations, thus the very same proof works in the general case. Lemma 3.1 can be carried out universally in the structural class of operators, provided integrability bounds for the time-derivative are available (cf. [Acerbi et al. 2004, Section 7], where a more general version of the result in [Lindqvist 2008] on this issue is proved). As for Lemma 3.2, the very same proof works since solutions to the general homogeneous equation are also Lipschitz in space and $C^{0;1/2}$ in time. The only modification occurs when we iterate Lemma 3.2. The rescaled function v defined in (20) now solves the equation

$$v_t - \operatorname{div} \mathcal{A}_k(x, t, Dv) = \lambda^{p k - (p-1)\alpha k} f(\lambda^k x, \lambda^{k\theta} t),$$

where

$$\mathcal{A}_k(x, t, \xi) := (\lambda^{-\alpha k})^{1-p} \mathcal{A}(\lambda^k x, \lambda^{\theta k} t, \lambda^{-\alpha k} \xi)$$

belongs to the same structural class of \mathcal{A} . In particular, v is entitled to the conditions of Lemma 3.2 and the proof then follows exactly as before.

We would like to conclude by explaining how the idea of finding geometrical tangential equations can be employed to derive analytical tools for p -parabolic operators, continuously on p . For instance, one can access regularity estimates for degenerate parabolic equations by interpreting the heat operator as the tangential equation obtained when we differentiate the family of p -parabolic operators with respect to the exponent p , near $p = 2$.

It is possible to obtain a universal compactness device. Let $Q_\tau := I_\tau \times B_\tau = (-\tau, \tau) \times B_\tau$. We fix $M_0 \gg 2$ and work within the range $p \in [2, M_0]$.

Lemma 4.1 (uniform in p compactness). *Given $\delta > 0$, there exists $\epsilon > 0$, depending only on n , M_0 and δ , such that if $q \in [2, M_0]$, u is a q -caloric function in Q_1 , with $|u| \leq 1$, and $|q - p| < \epsilon$, then we can find a p -caloric function w in $Q_{1/2}$, with $|w| \leq 1$, such that*

$$\sup_{Q_{1/2}} |w - u| \leq \delta. \quad (23)$$

Proof. Suppose, for the sake of contradiction, that the thesis of the lemma does not hold true. This means that for a certain $\delta_0 > 0$, there exist sequences $(q_j)_j$, $(u_j)_j$ and $(p_j)_j$, with

$$\begin{cases} q_j \in [2, M_0], \\ (u_j)_t - \operatorname{div}(|\nabla u_j|^{q_j-2} \nabla u_j) = 0 & \text{in } Q_1, \\ |u_j| \leq 1, \\ |p_j - q_j| \leq \frac{1}{j}, \end{cases} \quad (24)$$

but such that, for every p_j -caloric function w in $Q_{1/2}$,

$$\sup_{Q_{1/2}} |u_j - w| > \delta_0. \quad (25)$$

By compactness, we have, up to subsequences,

$$q_j \rightarrow q_\infty \in [2, M_0] \quad (26)$$

and, from the last assertion in (24), also $p_j \rightarrow q_\infty$. As in the proof of Lemma 3.1, up to a subsequence, $u_j \rightarrow u_\infty$ in the appropriate space. Since $q_j \rightarrow q_\infty$, by stability (see [Kinnunen and Parviainen 2010]), we can pass to the limit in the equation satisfied by the u_j to conclude that u_∞ is q_∞ -caloric in $Q_{2/3}$.

We now solve, for each p_j , the boundary value problem

$$\begin{cases} (w_j)_t - \operatorname{div}(|\nabla w_j|^{p_j-2} \nabla w_j) = 0 & \text{in } Q_{2/3}, \\ w_j = u_\infty & \text{on } \partial Q_{2/3}, \end{cases} \quad (27)$$

and pass to the limit in j , concluding that also $w_j \rightarrow u_\infty$ uniformly in $Q_{1/2}$.

Finally, choosing j sufficiently large, we obtain

$$|u_j - w_j| \leq |u_j - u_\infty| + |w_j - u_\infty| \leq \frac{\delta_0}{2} + \frac{\delta_0}{2} = \delta_0 \quad \text{in } Q_{1/2},$$

which is a contradiction to (25). \square

Heuristically, Lemma 4.1 implies the continuity of the underlying regularity theory for p -parabolic operators with respect to p . In particular, improved sharp Hölder estimates can be derived by these methods for problems governed by p -parabolic operators, near the heat equation, *i.e.*, for p close to 2. We leave the development of these heuristics for a future work.

Acknowledgements

This work was developed in the framework of the Brazilian Program “Ciência sem Fronteiras”. The authors thank Giuseppe Mingione and Rico Zacher for interesting conversations on the topic of the paper, and acknowledge the warm hospitality of Universidade Federal do Ceará and Universidade de Coimbra.

The research of Teixeira was partially supported by CNPq–Brazil. The research of Urbano was supported by projects UTAustin/MAT/0035/2008, PTDC/MAT/098060/2008, UTA-CMU/MAT/0007/2009 and PTDC/MAT-CAL/0749/2012, by FCT grant SFRH/BSAB/1273/2012, and by CMUC, funded by the European Regional Development Fund through the program COMPETE and by FCT, under the project PEst-C/MAT/UI0324/2011.

References

- [Acerbi and Mingione 2007] E. Acerbi and G. Mingione, “Gradient estimates for a class of parabolic systems”, *Duke Math. J.* **136**:2 (2007), 285–320. MR 2007k:35211 Zbl 1113.35105
- [Acerbi et al. 2004] E. Acerbi, G. Mingione, and G. A. Seregin, “Regularity results for parabolic systems related to a class of non-Newtonian fluids”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **21**:1 (2004), 25–60. MR 2005a:35118 Zbl 1052.76004
- [Araújo et al. 2013] D. Araújo, G. Ricarte, and E. V. Teixeira, “Optimal gradient continuity for degenerate elliptic equations”, preprint, 2013. Accepted for publication in *Calc. Var. Partial Differential Equations* under the title “Geometric gradient estimates for solutions to degenerate elliptic equation”. arXiv 1206.4089
- [DiBenedetto 1993] E. DiBenedetto, *Degenerate parabolic equations*, Springer, New York, 1993. MR 94h:35130 Zbl 0794.35090
- [DiBenedetto et al. 2012] E. DiBenedetto, U. Gianazza, and V. Vespi, *Harnack’s inequality for degenerate and singular parabolic equations*, Springer, New York, 2012. MR 2865434 Zbl 1237.35004
- [Duzaar and Mingione 2005] F. Duzaar and G. Mingione, “Second order parabolic systems, optimal regularity, and singular sets of solutions”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **22**:6 (2005), 705–751. MR 2008h:35139 Zbl 1099.35042
- [Iwaniec and Manfredi 1989] T. Iwaniec and J. J. Manfredi, “Regularity of p -harmonic functions on the plane”, *Rev. Mat. Iberoamericana* **5**:1-2 (1989), 1–19. MR 91i:35071 Zbl 0805.31003
- [Kinnunen and Lewis 2000] J. Kinnunen and J. L. Lewis, “Higher integrability for parabolic systems of p -Laplacian type”, *Duke Math. J.* **102**:2 (2000), 253–271. MR 2001b:35152 Zbl 0994.35036
- [Kinnunen and Parviainen 2010] J. Kinnunen and M. Parviainen, “Stability for degenerate parabolic equations”, *Adv. Calc. Var.* **3**:1 (2010), 29–48. MR 2011b:35222 Zbl 1185.35123
- [Kuusi et al. 2014] T. Kuusi, G. Mingione, and K. Nyström, “Sharp regularity for evolutionary obstacle problems, interpolative geometries and removable sets”, *J. Math. Pures Appl.* (9) **101**:2 (2014), 119–151. MR 3158698 Zbl 06251653
- [Lindqvist 2008] P. Lindqvist, “On the time derivative in a quasilinear equation”, *Skr. K. Nor. Vidensk. Selsk.* **2008**:2 (2008), 1–7. MR 2010d:35207 Zbl 1165.35402
- [Misawa 2013] M. Misawa, “A Hölder estimate for nonlinear parabolic systems of p -Laplacian type”, *J. Differential Equations* **254**:2 (2013), 847–878. MR 2990053 Zbl 1273.35173
- [Simon 1987] J. Simon, “Compact sets in the space $L^p(0, T; B)$ ”, *Ann. Mat. Pura Appl.* (4) **146** (1987), 65–96. MR 89c:46055 Zbl 0629.46031
- [Teixeira 2013] E. V. Teixeira, “Sharp regularity for general Poisson equations with borderline sources”, *J. Math. Pures Appl.* (9) **99**:2 (2013), 150–164. MR 3007841 Zbl 1263.35071
- [Urbano 2008] J. M. Urbano, *The method of intrinsic scaling: a systematic approach to regularity for degenerate and singular PDEs*, Lecture Notes in Math. **1930**, Springer, Berlin, 2008. MR 2011a:35245 Zbl 1158.35003

Received 12 Aug 2013. Revised 20 Jan 2014. Accepted 17 Feb 2014.

EDUARDO V. TEIXEIRA: teixeira@mat.ufc.br

Department of Mathematics, Universidade Federal do Ceará, Campus of Pici, Bloco 914, 60455–760 Fortaleza-CE, Brazil

JOSÉ MIGUEL URBANO: jmurub@mat.uc.pt

CMUC, Department of Mathematics, University of Coimbra, 3001–501 Coimbra, Portugal

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2014 is US \$180/year for the electronic version, and \$355/year (+\$50, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 7 No. 3 2014

Prescription du spectre de Steklov dans une classe conforme PIERRE JAMMES	529
Semilinear geometric optics with boundary amplification JEAN-FRANCOIS COULOMBEL, OLIVIER GUÈS and MARK WILLIAMS	551
The 1-harmonic flow with values in a hyperoctant of the N -sphere LORENZO GIACOMELLI, JOSE M. MAZÓN and SALVADOR MOLL	627
Decomposition rank of \mathcal{L} -stable C^* -algebras AARON TIKUISIS and WILHELM WINTER	673
Scattering for a massless critical nonlinear wave equation in two space dimensions MARTIN SACK	701
Large-time blowup for a perturbation of the cubic Szegő equation HAIYAN XU	717
A geometric tangential approach to sharp regularity for degenerate evolution equations EDUARDO V. TEIXEIRA and JOSÉ MIGUEL URBANO	733
The theory of Hahn-meromorphic functions, a holomorphic Fredholm theorem, and its applications JÖRN MÜLLER and ALEXANDER STROHMAIER	745



2157-5045(2014)7:3;1-D