

ANALYSIS & PDE

Volume 8

No. 1

2015

SEIICHIRO KUSUOKA

HÖLDER CONTINUITY AND BOUNDS FOR FUNDAMENTAL
SOLUTIONS
TO NONDIVERGENCE FORM PARABOLIC EQUATIONS

HÖLDER CONTINUITY AND BOUNDS FOR FUNDAMENTAL SOLUTIONS TO NONDIVERGENCE FORM PARABOLIC EQUATIONS

SEIICHIRO KUSUOKA

We consider nondegenerate second-order parabolic partial differential equations in nondivergence form with bounded measurable coefficients (not necessary continuous). Under certain assumptions weaker than the Hölder continuity of the coefficients, we obtain Gaussian bounds and Hölder continuity of the fundamental solution with respect to the initial point. Our proofs employ pinned diffusion processes for the probabilistic representation of fundamental solutions and the coupling method.

1. Introduction and main result

Let $a(t, x) = (a_{ij}(t, x))$ be a symmetric $d \times d$ -matrix-valued bounded measurable function on $[0, \infty) \times \mathbb{R}^d$ which is uniformly positive-definite, i.e.,

$$\Lambda^{-1}I \leq a(t, x) \leq \Lambda I, \quad (1-1)$$

where Λ is a positive constant and I is the unit matrix. Let $b(t, x) = (b_i(t, x))$ be an \mathbb{R}^d -valued bounded measurable function on $[0, \infty) \times \mathbb{R}^d$ and $c(t, x)$ a bounded measurable function on $[0, \infty) \times \mathbb{R}^d$. Consider the parabolic partial differential equation

$$\begin{cases} \frac{\partial}{\partial t} u(t, x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(t, x) \frac{\partial^2}{\partial x_i \partial x_j} u(t, x) + \sum_{i=1}^d b_i(t, x) \frac{\partial}{\partial x_i} u(t, x) + c(t, x) u(t, x), \\ u(0, x) = f(x). \end{cases} \quad (1-2)$$

Generally, (1-2) does not have a unique solution. We will assume the continuity of a in spatial components uniformly in t , and this implies the uniqueness of the weak solution; see [Stroock and Varadhan 1979]. In the present paper, we always consider cases where the uniqueness of the weak solution holds. Set

$$L_t f(x) := \frac{1}{2} \sum_{i,j=1}^d a_{ij}(t, x) \frac{\partial^2}{\partial x_i \partial x_j} f(x) + \sum_{i=1}^d b_i(t, x) \frac{\partial}{\partial x_i} f(x) + c(t, x) f(x), \quad f \in C_b^2(\mathbb{R}^d)$$

and denote the fundamental solution to (1-2) by $p(s, x; t, y)$, i.e., $p(s, x; t, y)$ is a measurable function defined for $s, t \in [0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$ which satisfies

$$\frac{\partial}{\partial t} \int_{\mathbb{R}^d} f(y) p(s, \cdot; t, y) dy = L_t \left(\int_{\mathbb{R}^d} f(y) p(s, \cdot; t, y) dy \right) \quad \text{and} \quad \lim_{r \downarrow s} \int_{\mathbb{R}^d} f(y) p(s, \cdot; r, y) dy = f$$

MSC2010: primary 35B65, 35K10, 60H30; secondary 60H10, 60J60.

Keywords: parabolic partial differential equation, diffusion, fundamental solution, Hölder continuity, Gaussian estimate, stochastic differential equation, coupling method.

for $s, t \in [0, \infty)$ such that $s < t$ and a continuous function f with a compact support. In the present paper, we consider the existence and the regularity of $p(0, x; t, y)$.

The problem of regularity of the fundamental solutions to parabolic partial differential equations with bounded measurable coefficients has a long history. Parabolic equations in divergence form has been investigated more thoroughly than that those in nondivergence form, because the variational method is applicable to them. The Hölder continuity of the fundamental solution to $\partial_t u = \nabla \cdot a \nabla u$ for a matrix-valued bounded measurable function a with ellipticity condition $\Lambda^{-1}I \leq a \leq \Lambda I$ was originally obtained by De Giorgi [1957] and Nash [1958] independently. Precisely speaking, in their results the α -Hölder continuity of the fundamental solution, with some positive number $\alpha \in (0, 1]$, is obtained. The index α depends on many constants appearing in the Harnack inequality and so on. These results have been extended to the case of the more general equations $\partial_t u = \nabla \cdot a \nabla u + b \cdot \nabla u - cu$, where b, c are bounded measurable; see [Aronson 1967; Stroock 1988]. The case for unbounded coefficients is also studied; see, for example, [Metafuno et al. 2009; Porper and Eidelman 1984; 1992]. An analogy to the case of a type of nonlocal generators (the associated stochastic processes are called stable-like processes) is given by Chen and Kumagai [2003]. In the results above, the index of the Hölder continuity of the fundamental solution depends on many constants appearing in the estimates, and it is difficult to calculate its exact value. Moreover, it is difficult to obtain even a lower bound for the index.

The fundamental solutions to parabolic equations in nondivergence form with low-regular coefficients have been studied mainly in the case of Hölder-continuous coefficients. One of the most powerful tools for the problem is the parametrix method, and it yields the existence, uniqueness and Hölder continuity of the fundamental solution; see [Friedman 1964; Ladyženskaja et al. 1967; Porper and Eidelman 1984, Chapter I]. Furthermore, an a priori estimate (the so-called Schauder estimate) is known for the solutions, and twice-continuous differentiability in x of the fundamental solution $p(s, x; t, y)$ to (1-2) is obtained; see, for example, [Ladyženskaja et al. 1967; Krylov 1996; Bogachev et al. 2009; Bogachev et al. 2005]. We remark that all the coefficients a, b, c need to be Hölder-continuous to apply the Schauder estimate. Even in the case that a is the unit matrix and b is not continuous, we cannot expect the continuous differentiability of the fundamental solution; see [Karatzas and Shreve 1991, Remark 5.2, Chapter 6].

In the present paper, we consider the Gaussian estimate and the lower bound of the index for the Hölder continuity in x of the fundamental solution $p(s, x; t, y)$ to (1-2) by a probabilistic approach.

Now we fix some assumptions. Let $B(x, R)$ be the open ball in \mathbb{R}^d centered at x with radius R for $x \in \mathbb{R}^d$ and $R > 0$. We assume that

$$\sum_{i,j=1}^d \sup_{t \in [0, \infty)} \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial x_j} a_{ij}(s, x) \right|^\theta e^{-m|x|} dx \leq M, \quad (1-3)$$

where the derivatives are in the weak sense, θ is a constant in $[d, \infty) \cap (2, \infty)$, and m and M are nonnegative constants. We also assume the continuity of a in spatial component uniformly in t , i.e., for any $R > 0$ there exists a continuous and nondecreasing function ρ_R on $[0, \infty)$ such that $\rho_R(0) = 0$ and

$$\sup_{t \in [0, \infty)} \sup_{i,j} |a_{ij}(t, x) - a_{ij}(t, y)| \leq \rho_R(|x - y|), \quad x, y \in B(0; R). \quad (1-4)$$

We remark that under the assumptions (1-1) and (1-4), the equation (1-2) under consideration is well-posed (see [Stroock and Varadhan 1979, Chapter 7]), and for fixed $s \in [0, \infty)$ the fundamental solution $p(s, \cdot; t, \cdot)$ exists for almost all $t \in (s, \infty)$ (see [Stroock and Varadhan 1979, Theorem 9.1.9]). However, the fundamental solution does not always exist for all $t \in (s, \infty)$ under assumptions (1-1) and (1-4); see [Fabes and Kenig 1981]. We remark that under assumptions (1-1), (1-3) and (1-4), neither existence of the fundamental solutions nor examples where the fundamental solution does not exist are known. In the case that a does not depend on the time t , the fundamental solution exists for all t ; see [Stroock and Varadhan 1979, Theorem 9.2.6]. We also remark that (1-3) and (1-4) do not imply the local Hölder continuity of a in the spatial component.

Let $p^X(s, x; t, y)$ be the fundamental solution to the parabolic equation

$$\frac{\partial}{\partial t} u(t, x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(t, x) \frac{\partial^2}{\partial x_i \partial x_j} u(t, x),$$

and let

$$L_t^X = \frac{1}{2} \sum_{i,j=1}^d a_{ij}(t, x) \frac{\partial^2}{\partial x_i \partial x_j}. \quad (1-5)$$

Suppose that $\{a^{(n)}(t, x)\}$ is a sequence of symmetric $d \times d$ -matrix-valued functions with components in $C_b^\infty([0, \infty) \times \mathbb{R}^d)$ such that $a^{(n)}(t, x)$ converges to $a(t, x)$ for each $(t, x) \in [0, \infty) \times \mathbb{R}^d$. We also assume that (1-1), (1-3) and (1-4) hold for $a^{(n)}$ instead of a , with the same constants m, M, θ, R and Λ , and the same function ρ_R . Denote the fundamental solution to the parabolic equation associated with the generator

$$\frac{1}{2} \sum_{i,j=1}^d a_{ij}^{(n)}(t, x) \frac{\partial^2}{\partial x_i \partial x_j}$$

by $p^{X,(n)}$. We assume the uniform Gaussian estimates for the fundamental solutions to $p^{X,(n)}$, i.e., there exist positive constants $\gamma_G^-, \gamma_G^+, C_G^-$ and C_G^+ such that

$$\frac{C_G^-}{(t-s)^{d/2}} \exp\left(-\frac{\gamma_G^- |x-y|^2}{t-s}\right) \leq p^{X,(n)}(s, x; t, y) \leq \frac{C_G^+}{(t-s)^{d/2}} \exp\left(-\frac{\gamma_G^+ |x-y|^2}{t-s}\right) \quad (1-6)$$

for $s, t \in [0, \infty)$ such that $s < t$, $x, y \in \mathbb{R}^d$, and $n \in \mathbb{N}$. The Gaussian estimates for the fundamental solutions to parabolic equations in divergence form have been well investigated; see [Aronson 1967; Karrass 2001; Porper and Eidelman 1984; 1992]. However, there are not many known results in the case of nondivergence form. A sufficient condition for the Gaussian estimate is obtained in [Porper and Eidelman 1992, Theorem 19] by means of Dini's continuity condition. The result includes the case of Hölder-continuous coefficients. We remark that two-sided estimates similar to the Gaussian estimates for the equations with general coefficients are obtained in [Escauriaza 2000].

Now we state the main theorem of this paper:

Theorem 1.1. *Assume (1-1), (1-3), (1-4) and (1-6). Then, there exist constants C_1, C_2, γ_1 and γ_2 depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that*

$$\frac{C_1 e^{-C_1(t-s)}}{(t-s)^{d/2}} \exp\left(-\frac{\gamma_1|x-y|^2}{t-s}\right) \leq p(s, x; t, y) \leq \frac{C_2 e^{C_2(t-s)}}{(t-s)^{d/2}} \exp\left(-\frac{\gamma_2|x-y|^2}{t-s}\right)$$

for $s, t \in [0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$. Moreover, for any $R > 0$ and sufficiently small $\varepsilon > 0$, there exists a constant C depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that

$$|p(0, x; t, y) - p(0, z; t, y)| \leq C t^{-d/2-1} e^{Ct} |x-z|^{1-\varepsilon}$$

for $t \in (0, \infty), x, z \in B(0; R/2)$ and $y \in \mathbb{R}^d$.

The first assertion of Theorem 1.1 is the Gaussian estimate for p . The advantage of the result is obtaining the Gaussian estimate of the fundamental solution to the parabolic equation in nondivergence form without the continuity of b and c . Such a result seems difficult to obtain via the parametrix method. The second assertion of Theorem 1.1 implies that $p(0, x; t, y)$ is $(1-\varepsilon)$ -Hölder continuous in x , and this is a clear lower bound for the continuity. The approach in this paper is mainly probabilistic. The key method to prove Theorem 1.1 is the coupling method introduced by Lindvall and Rogers [1986]. This method enables us to discuss the Hölder continuity of $p(0, x; t, y)$ in x from the oscillation of the diffusion processes without regularity of the coefficients.

If a is uniformly continuous in the spatial component, our proof below follows without restriction on x, z , and the following corollary holds:

Corollary 1.2. *Assume (1-1), (1-3), (1-6) and that there exists a continuous and nondecreasing function ρ on $[0, \infty)$ such that $\rho(0) = 0$ and*

$$\sup_{t \in [0, \infty)} \sup_{i, j} |a_{ij}(t, x) - a_{ij}(t, y)| \leq \rho(|x-y|), \quad x, y \in \mathbb{R}^d.$$

Then, for sufficiently small $\varepsilon > 0$, there exists a constant C such that

$$|p(0, x; t, y) - p(0, z; s, y)| \leq C t^{-d/2-1} e^{Ct} |x-z|^{1-\varepsilon}$$

for $t \in (0, \infty)$ and $x, y, z \in \mathbb{R}^d$.

The assumption (1-6) may seem strict. However, as mentioned above, Theorem 19 of [Porper and Eidelman 1992] gives a Gaussian estimate for the parabolic equations with coefficients which satisfy a version of Dini's continuity condition. From this sufficient condition and Theorem 1.1, we have the following corollary:

Corollary 1.3. *Assume (1-1), (1-3), and that there exists a continuous and nondecreasing function ρ on $[0, \infty)$ such that $\rho(0) = 0$,*

$$\int_0^1 \frac{1}{r_2} \left(\int_0^{r_2} \frac{\rho(r_1)}{r_1} dr_1 \right) dr_2 < \infty \quad \text{and} \quad \sup_{t \in [0, \infty)} \sup_{i, j} |a_{ij}(t, x) - a_{ij}(t, y)| \leq \rho(|x-y|), \quad x, y \in \mathbb{R}^d. \quad (1-7)$$

Then, for sufficiently small $\varepsilon > 0$, there exists a constant C such that

$$|p(0, x; t, y) - p(0, z; s, y)| \leq Ct^{-d/2-1} e^{Ct} |x - z|^{1-\varepsilon}$$

for $t \in (0, \infty)$ and $x, y, z \in \mathbb{R}^d$.

We remark that, for $\alpha \in (0, 1]$ and a positive constant C , $\rho(r) = Cr^\alpha$ satisfies (1-7). Furthermore, $\rho(r) = C \min\{1, (-\log r)^{-\alpha}\}$ satisfies (1-7) for $\alpha \in (2, \infty)$. We also remark that continuity of b and c are not assumed in Corollary 1.3.

The organization of the paper is as follows:

In Section 2, we prepare the probabilistic representation of the fundamental solution to (1-2). It should be remarked that we consider the case where a is smooth in Sections 2–4, and the general case is considered only in Section 5. The representation enable us to consider the Hölder continuity of the fundamental solution by a probabilistic way, and actually in Section 4 we prove the constant appearing in the Hölder continuity of $p(0, x; t, y)$ in x depends only on the suitable constants. The representation is obtained by the Feynman–Kac formula and the Girsanov transformation, and in the end of this section $p(s, x; t, y)$ is represented by the functional of the pinned diffusion process.

In Section 3, we prepare some estimates. The goal of this section is Lemma 3.5, which concerns the integrability of a functional of the pinned diffusion process. Generally speaking, it is much harder to see the integrability with respect to conditional probability measures than with respect to the original probability measure. In our case, conditioning generates a singularity, and this fact makes the estimate difficult. To overcome the difficulty, we begin with Lemma 3.1, which is an estimate of the derivative of $p(s, x; t, y)$. The proof of this lemma is analytic, and (1-3) is assumed for the lemma. In this section, we also have the Gaussian estimate for $p(s, x; t, y)$.

In Section 4, we prove that the constant appearing in the $(1 - \varepsilon)$ -Hölder continuity of $p(0, x; t, y)$ in x depends only on the suitable constants. This section is the main part of our argument. To show this, we apply the coupling method to diffusion processes. By virtue of the coupling method, the continuity problem of the fundamental solution is reduced to the problem of the local behavior of the pinned diffusion processes. To see the local behavior, (1-4) is needed. Finally, by showing an estimate of the coupling time, we obtain the $(1 - \varepsilon)$ -Hölder continuity of $p(0, x; t, y)$ in x and the suitable dependence of the constant appearing in the Hölder continuity.

In Section 5, we consider the case of general a and prove Theorem 1.1. Our approach is just smoothly approximating a and using the result obtained in Section 4.

Throughout this paper, we denote the inner product in the Euclidean space \mathbb{R}^d by $\langle \cdot, \cdot \rangle$, and all random variables are considered on a probability space (Ω, \mathcal{F}, P) . We denote the expectation of random variables by $E[\cdot]$ and the expectation on the event $A \in \mathcal{F}$ (i.e., $\int_A \cdot dP$) by $E[\cdot; A]$. We denote the smooth functions with bounded derivatives on S by $C_b^\infty(S)$ and the smooth functions on S with compact support by $C_0^\infty(S)$.

2. Probabilistic representation of the fundamental solution

In this section, we assume that $a_{ij}(t, x) \in C_b^{0,\infty}([0, \infty) \times \mathbb{R}^d)$. Define a $d \times d$ -matrix-valued function $\sigma(t, x)$ as the square root of $a(t, x)$. Then, (1-1) implies that $\sigma_{ij}(t, x) \in C_b^{0,\infty}([0, \infty) \times \mathbb{R}^d)$, $a(t, x) =$

$\sigma(t, x)\sigma(t, x)^T$ and

$$\sup_{t \in [0, \infty)} \sup_{i, j} |\sigma_{ij}(t, x) - \sigma_{ij}(t, y)| \leq C\rho_R(|x - y|), \quad x, y \in B(0; R), \quad (2-1)$$

where C is a constant depending on Λ . Note that (1-1) implies that

$$\Lambda^{-1/2}I \leq \sigma(t, x) \leq \Lambda^{1/2}I. \quad (2-2)$$

Consider the stochastic differential equation:

$$\begin{cases} dX_t^x = \sigma(t, X_t^x) dB_t, \\ X_0^x = x. \end{cases} \quad (2-3)$$

Lipschitz continuity of σ implies the existence of a solution and its pathwise uniqueness. Let (\mathcal{F}_t) be the σ -field generated by $(B_s : s \in [0, t])$. Then, pathwise uniqueness implies that the solution X_t^x is \mathcal{F}_t -measurable. All stopping times appearing in this paper are associated with (\mathcal{F}_t) . We remark that the generator of (X_t^x) is given by (1-5), and therefore the transition probability density of (X_t^x) coincides with the fundamental solution p^X of the parabolic equation generated by (L_t^X) . The smoothness of σ implies the smoothness of $p^X(s, x; t, y)$ on $(0, \infty) \times \mathbb{R}^d \times (0, \infty) \times \mathbb{R}^d$; see, for example, [Kusuoka and Stroock 1985] for the probabilistic proof and [Lax and Milgram 1954] for the analytic proof.

There is a relation between the fundamental solution and the generator, as follows. Since p^X is smooth, by the definition of p^X we have

$$\frac{\partial}{\partial t} p^X(s, x; t, y) = [L_t^X p^X(s, \cdot; t, y)](x) \quad (2-4)$$

for $s, t \in [0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$. Let $(L_t^X)^*$ be the dual operator of L_t^X on $L^2(\mathbb{R}^d)$. Define $T_{s,t}^X$ and $(T_{s,t}^X)^*$ as the semigroups generated by L_t^X and $(L_t^X)^*$, respectively. Since

$$\int_{\mathbb{R}^d} \phi(x) (T_{s,t}^X \psi)(x) dx = \int_{\mathbb{R}^d} \psi(x) [(T_{s,t}^X)^* \phi](x) dx,$$

we have

$$\int_{\mathbb{R}^d} \phi(x) \left(\int_{\mathbb{R}^d} \psi(y) p^X(s, x; t, y) dy \right) dx = \int_{\mathbb{R}^d} \psi(x) \left(\int_{\mathbb{R}^d} \phi(y) (p^X)^*(s, x; t, y) dy \right) dx,$$

where $(p^X)^*(s, x; t, y)$ is the fundamental solution associated with $(L_t^X)^*$. Hence, it holds that

$$p^X(s, x; t, y) = (p^X)^*(s, y; t, x)$$

for $s, t \in (0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$. Differentiating both sides of this equation with respect to t , we obtain

$$[L_t^X p^X(s, \cdot; t, y)](x) = [(L_t^X)^* (p^X)^*(s, \cdot; t, x)](y) = [(L_t^X)^* p^X(s, x; t, \cdot)](y) \quad (2-5)$$

for $s, t \in [0, \infty)$ such that $s < t$, and $x, y \in \mathbb{R}^d$. By the Chapman–Kolmogorov equation, we have, for $s, t, u \in [0, \infty)$ such that $u < s < t$ and $x, y \in \mathbb{R}^d$, that

$$p^X(u, x; t, y) = \int_{\mathbb{R}^d} p^X(u, x; s, \xi) p^X(s, \xi; t, y) d\xi.$$

Differentiating both sides of this equation with respect to s , we have

$$0 = \int_{\mathbb{R}^d} \left(\frac{\partial}{\partial s} p^X(u, x; s, \xi) \right) p^X(s, \xi; t, y) d\xi + \int_{\mathbb{R}^d} p^X(u, x; s, \xi) \left(\frac{\partial}{\partial s} p^X(s, \xi; t, y) \right) d\xi$$

for $s, u \in [0, \infty)$ such that $u < s$ and $x, y \in \mathbb{R}^d$. Since (2-4) and (2-5) imply that

$$\frac{\partial}{\partial s} p^X(u, x; s, \xi) = [L_s^X p^X(u, \cdot; s, \xi)](x) = [(L_s^X)^* p^X(u, x; s, \cdot)](\xi),$$

we have, for $s, t, u \in [0, \infty)$ such that $u < s < t$ and $x, y \in \mathbb{R}^d$, that

$$\begin{aligned} \int_{\mathbb{R}^d} p^X(u, x; s, \xi) \left(\frac{\partial}{\partial s} p^X(s, \xi; t, y) \right) d\xi &= - \int_{\mathbb{R}^d} [(L_s^X)^* p^X(u, x; s, \cdot)](\xi) p^X(s, \xi; t, y) d\xi \\ &= - \int_{\mathbb{R}^d} p^X(u, x; s, \xi) [L_s^X p^X(s, \cdot; t, y)](\xi) d\xi. \end{aligned}$$

Noting that $p^X(u, x; s, \xi)$ converges to $\delta_x(\xi)$ as $u \uparrow s$ in the sense of Schwartz distributions, we obtain

$$\frac{\partial}{\partial s} p^X(s, x; t, y) = -[L_s^X p^X(s, \cdot; t, y)](x) \quad (2-6)$$

for $s, t \in (0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$.

Next we study the probabilistic representation of $p(s, x; t, y)$ by $p^X(s, x; t, y)$. By the Feynman–Kac formula (see, for example, [Revuz and Yor 1999, Proposition 3.10, Chapter VIII]) and the Girsanov transformation (see, for example, [Ikeda and Watanabe 1989, Theorem 4.2, Chapter IV]), we have the following representation of $u(t, x)$ by X_t^x :

$$u(t, x) = E \left[f(X_t^x) \exp \left(\int_0^t \langle b_\sigma(s, X_s^x), dB_s \rangle - \frac{1}{2} \int_0^t |b_\sigma(s, X_s^x)|^2 ds + \int_0^t c(s, X_s^x) ds \right) \right], \quad (2-7)$$

where $b_\sigma(t, x) := \sigma(t, x)^{-1} b(t, x)$. For $s \leq t$ and $x \in \mathbb{R}^d$, let

$$\mathcal{E}(s, t; X^x) := \exp \left(\int_s^t \langle b_\sigma(u, X_u^x), dB_u \rangle - \frac{1}{2} \int_s^t |b_\sigma(u, X_u^x)|^2 du + \int_s^t c(u, X_u^x) du \right).$$

Then, by the definition of the fundamental solution and (2-7), we obtain the probabilistic representation of the fundamental solution:

$$p(0, x; t, y) = p^X(0, x; t, y) E^{X_t^x=y} [\mathcal{E}(0, t; X^x)], \quad (2-8)$$

where $P^{X_t^x=y}$ is the conditional probability measure of P on $X_t^x = y$ and $E^{X_t^x=y}[\cdot]$ is the expectation with respect to $P^{X_t^x=y}$. Hence, to see the regularity of $p(0, x; t, y)$ in x , it is sufficient to see the regularity of the function $x \mapsto p^X(0, x; t, y)E^{X_t^x=y}[\mathcal{E}(0, t; X^x)]$. We prove Theorem 1.1 by studying the regularity of this function. The definition of \mathcal{E} implies that

$$\mathcal{E}(0, t; X^x) - \mathcal{E}(\tau \wedge t, t; X^x) = \mathcal{E}(\tau \wedge t, t; X^x)(\mathcal{E}(0, \tau \wedge t; X^x) - 1) \quad (2-9)$$

for any stopping time τ and $t \in [0, \infty)$, and by Itô's formula we have

$$\mathcal{E}(s, t; X^x) - 1 = \int_s^t \mathcal{E}(s, u; X^x) \langle b_\sigma(u, X_u^x), dB_u \rangle + \int_s^t \mathcal{E}(s, u; X^x) c(u, X_u^x) du \quad (2-10)$$

for $s, t \in [0, \infty)$ such that $s \leq t$. We use these equations in the proof.

Now we consider the diffusion process X^x pinned at y at time t . Let $s, t \in [0, \infty)$ such that $s < t$, $x, y \in \mathbb{R}^d$ and $\varepsilon > 0$. By the Markov property of X , we have for $A \in \mathcal{F}_s$ that

$$P(A \cap \{X_t^x \in B(y; \varepsilon)\})_{B(y; \varepsilon)} \left(\int_{\mathbb{R}^d} p^X(s, \xi; t, \xi') P(A \cap \{X_s^x \in d\xi\}) d\xi' \right)$$

Hence, we obtain

$$P^{X_t^x=y}(A) = \frac{1}{p^X(0, x; t, y)} \int_{\mathbb{R}^d} p^X(s, \xi; t, y) P(A \cap \{X_s^x \in d\xi\}) \quad (2-11)$$

for $s, t \in (0, \infty)$ such that $s < t$, $A \in \mathcal{F}_s$ and $x, y \in \mathbb{R}^d$. This formula enables us to see the generator of the pinned diffusion process. By Itô's formula, (2-6) and (2-11) we have, for $f \in C_b^2(\mathbb{R}^d)$, $s, t \in [0, \infty)$ such that $s < t$ and $x, y \in \mathbb{R}^d$,

$$\begin{aligned} & p^X(0, x; t, y) E^{X_t^x=y}[f(X_s^x)] - p^X(0, x; t, y) f(x) \\ &= E[f(X_s^x) p^X(s, X_s^x; t, y)] - E[f(X_0^x) p^X(0, X_0^x; t, y)] \\ &= E \left[\int_0^s (L_u^X f)(X_u^x) p^X(u, X_u^x; t, y) du \right] + E \left[\int_0^s f(X_u^x) \left(\frac{\partial}{\partial u} p^X(u, \xi; t, y) \right) \Big|_{\xi=X_u^x} du \right] \\ &\quad + E \left[\int_0^s f(X_u^x) (L_u^X p^X(u, \cdot; t, y))(X_u^x) du \right] \\ &\quad + \frac{1}{2} E \left[\int_0^s \langle \sigma(u, X_u^x)^T \nabla f(X_u^x), \sigma(u, X_u^x)^T (\nabla p^X(u, \cdot; t, y))(X_u^x) \rangle du \right] \\ &= p^X(0, x; t, y) \int_0^s E^{X_t^x=y}[(L_u^X f)(X_u^x)] du \\ &\quad + \frac{1}{2} p^X(0, x; t, y) \int_0^s E^{X_t^x=y} \left[\left\langle \nabla f(X_u^x), a(u, X_u^x) \frac{(\nabla p^X(u, \cdot; t, y))(X_u^x)}{p^X(u, X_u^x; t, y)} \right\rangle \right] du. \end{aligned}$$

Hence, the generator of X pinned at y at time t is

$$\frac{1}{2} \sum_{i,j=1}^d a_{ij}(s, x) \frac{\partial^2}{\partial x_i \partial x_j} + \left\langle \frac{1}{2} a(s, x) \frac{\nabla p^X(s, \cdot; t, y)(x)}{p^X(s, x; t, y)}, \nabla \right\rangle$$

for $s \in [0, t)$ and $x \in \mathbb{R}^d$. Of course, pinned Brownian motion is an example of pinned diffusion processes; see [Ikeda and Watanabe 1989, Example 8.5, Chapter IV].

3. Estimates

In this section we prepare some estimates for the proof of the main theorem. Assume that a is smooth and fix notation as in Section 2.

Lemma 3.1. *Let $t \in (0, \infty)$ and ϕ be a nonnegative continuous function on $(0, t) \times \mathbb{R}^d$ such that $\phi(\cdot, x) \in W_{\text{loc}}^{1,1}((0, t), ds)$ for $x \in \mathbb{R}^d$ and $\phi(s, \cdot) \in W_{\text{loc}}^{1,2}(\mathbb{R}^d, dx)$ for $s \in (0, t)$. Then, for $s_1, s_2 \in (0, t)$ such that $s_1 \leq s_2$,*

$$\begin{aligned} & \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_{\xi} p^X(u, \xi; t, y), \nabla_{\xi} p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi du \\ & \leq C(1 + |\log(t - s_1)|) \int_{\mathbb{R}^d} \phi(s_1, \xi) d\xi + C(t - s_1)^{-1} \int_{\mathbb{R}^d} |y - \xi|^2 \phi(s_1, \xi) d\xi \\ & \quad + C(1 + |\log(t - s_2)|) \int_{\mathbb{R}^d} \phi(s_2, \xi) d\xi + C(t - s_2)^{-1} \int_{\mathbb{R}^d} |y - \xi|^2 \phi(s_2, \xi) d\xi + C \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \phi(u, \xi) d\xi du \\ & \quad + C \int_{s_1}^{s_2} \int_{\text{supp } \phi} \frac{|\nabla_{\xi} \phi(u, \xi)|^2}{\phi(u, \xi)} d\xi du + C \int_{s_1}^{s_2} (1 + |\log(t - u)|) \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du \\ & \quad + C \int_{s_1}^{s_2} (t - u)^{-1} \int_{\mathbb{R}^d} |y - \xi|^2 \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du + C \sum_{i,j=1}^d \int_0^t \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial \xi_j} a_{ij}(u, \xi) \right|^2 \phi(u, \xi) d\xi du, \end{aligned}$$

where C is a constant depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+$, and Λ , and $\text{supp } \phi$ is the support of ϕ .

Remark 3.2. If ϕ is a continuous function on \mathbb{R}^d , the Lebesgue measure of $\text{supp } \phi \setminus \{x \in \mathbb{R}^d : \phi(x) > 0\}$ is zero.

Proof of Lemma 3.1. It is sufficient to show the theorem for $\phi \in C_0^\infty([0, t] \times \mathbb{R}^d)$, because the general case is obtained by approximation. Let $u \in (0, t)$. Recall that the components of the coefficient σ of (2-3) are in $C_b^{0,\infty}([0, \infty) \times \mathbb{R}^d)$ and σ is uniformly positive-definite. Hence, the associated transition probability density $p^X(\cdot, \cdot; t, \cdot)$ is smooth on $(0, t) \times \mathbb{R}^d \times \mathbb{R}^d$, and $p^X(s, x; t, y) > 0$ for $s \in (0, t)$ and $x, y \in \mathbb{R}^d$; see, for example, [Aronson 1967]. By the Leibniz rule, we have

$$\begin{aligned} & \frac{\partial}{\partial u} \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \phi(u, \xi) d\xi \\ & = \int_{\mathbb{R}^d} \frac{(\partial/\partial u) p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi + \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \frac{\partial}{\partial u} \phi(u, \xi) d\xi. \quad (3-1) \end{aligned}$$

The equality (2-6) and integration by parts imply

$$\begin{aligned}
& \int_{\mathbb{R}^d} \frac{(\partial/\partial u)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi \\
&= -\frac{1}{2} \sum_{i,j=1}^d \int_{\mathbb{R}^d} \frac{a_{ij}(u, \xi)(\partial^2/\partial\xi_i\partial\xi_j)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi \\
&= -\frac{1}{2} \sum_{i,j=1}^d \int_{\mathbb{R}^d} \frac{(\partial/\partial\xi_i)(a_{ij}(u, \xi)(\partial/\partial\xi_j)p^X(u, \xi; t, y))}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi \\
&\quad + \frac{1}{2} \sum_{i,j=1}^d \int_{\mathbb{R}^d} \frac{((\partial/\partial\xi_i)a_{ij}(u, \xi))(\partial/\partial\xi_j)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi \\
&= -\frac{1}{2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_\xi p^X(u, \xi; t, y), \nabla_\xi p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi \\
&\quad + \frac{1}{2} \int_{\mathbb{R}^d} \left\langle \frac{a(u, \xi) \nabla_\xi p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)}, \nabla_\xi \phi(u, \xi) \right\rangle d\xi \\
&\quad + \frac{1}{2} \sum_{i,j=1}^d \int_{\mathbb{R}^d} \frac{((\partial/\partial\xi_i)a_{ij}(u, \xi))(\partial/\partial\xi_j)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi.
\end{aligned}$$

Hence, by (3-1) we have

$$\begin{aligned}
\frac{\partial}{\partial u} \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \phi(u, \xi) d\xi &= -\frac{1}{2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_\xi p^X(u, \xi; t, y), \nabla_\xi p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi \\
&\quad + \frac{1}{2} \int_{\mathbb{R}^d} \left\langle \frac{a(u, \xi) \nabla_\xi p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)}, \nabla_\xi \phi(u, \xi) \right\rangle d\xi \\
&\quad + \frac{1}{2} \sum_{i,j=1}^d \int_{\mathbb{R}^d} \frac{((\partial/\partial\xi_i)a_{ij}(u, \xi))(\partial/\partial\xi_j)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi \\
&\quad + \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \frac{\partial}{\partial u} \phi(u, \xi) d\xi.
\end{aligned}$$

Integrating both sides from s_1 to s_2 with respect to u , we obtain

$$\begin{aligned}
& \frac{1}{2} \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_\xi p^X(u, \xi; t, y), \nabla_\xi p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi du \\
&= \int_{\mathbb{R}^d} (\log p^X(s_1, \xi; t, y)) \phi(s_1, \xi) d\xi - \int_{\mathbb{R}^d} (\log p^X(s_2, \xi; t, y)) \phi(s_2, \xi) d\xi \\
&\quad + \frac{1}{2} \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left\langle \frac{a(u, \xi) \nabla_\xi p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)}, \nabla_\xi \phi(u, \xi) \right\rangle d\xi du \\
&\quad + \frac{1}{2} \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{((\partial/\partial\xi_i)a_{ij}(u, \xi))(\partial/\partial\xi_j)p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi du \\
&\quad + \int_{s_1}^{s_2} \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \frac{\partial}{\partial u} \phi(u, \xi) d\xi du. \tag{3-2}
\end{aligned}$$

Now we consider the estimates for the terms on the right-hand side of this equation. By (1-6), we have for $s \in (0, t)$ that

$$\left| \int_{\mathbb{R}^d} (\log p^X(s, \xi; t, y)) \phi(s, \xi) d\xi \right| \leq \int_{\mathbb{R}^d} \left(|\log C_G^+| + |\log C_G^-| + \frac{d}{2} |\log(t-s)| + \frac{\gamma_G^- |y - \xi|^2}{t-s} \right) \phi(s, \xi) d\xi.$$

Hence, there exists a constant C depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+$, and Λ such that, for $s \in (0, t)$,

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} (\log p^X(s, \xi; t, y)) \phi(s, \xi) d\xi \right| \\ & \leq C(1 + |\log(t-s)|) \int_{\mathbb{R}^d} \phi(s, \xi) d\xi + C(t-s)^{-1} \int_{\mathbb{R}^d} |y - \xi|^2 \phi(s, \xi) d\xi. \end{aligned} \quad (3-3)$$

The third term of the right-hand side of (3-2) can be estimated as follows:

$$\begin{aligned} & \frac{1}{2} \left| \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left\langle \frac{a(u, \xi) \nabla_z p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)}, \nabla_\xi \phi(u, \xi) \right\rangle d\xi du \right| \\ & \leq \frac{1}{8} \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_\xi p^X(u, \xi; t, y), \nabla_\xi p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi du \\ & \quad + 8 \int_{s_1}^{s_2} \int_{\text{supp } \phi} \frac{\langle a(u, \xi) \nabla_\xi \phi(u, \xi), \nabla_\xi \phi(u, \xi) \rangle}{\phi(u, \xi)} d\xi du. \end{aligned} \quad (3-4)$$

To estimate the fourth term of the right-hand side of (3-2), we observe that

$$\begin{aligned} & \frac{1}{2} \left| \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{((\partial/\partial \xi_i) a_{ij}(u, \xi)) (\partial/\partial \xi_j) p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi du \right| \\ & \leq \frac{1}{8\Lambda} \sum_{j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{|(\partial/\partial \xi_j) p^X(u, \xi; t, y)|^2}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi du \\ & \quad + 8d\Lambda \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial \xi_j} a_{ij}(u, \xi) \right|^2 \phi(u, \xi) d\xi du. \end{aligned}$$

Hence, by (1-1) we have

$$\begin{aligned} & \frac{1}{2} \left| \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{(\frac{\partial}{\partial \xi_i} a_{ij}(t, \xi)) \frac{\partial}{\partial \xi_j} p^X(u, \xi; t, y)}{p^X(u, \xi; t, y)} \phi(u, \xi) d\xi du \right| \\ & \leq \frac{1}{8} \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \frac{\langle a(u, \xi) \nabla_\xi p^X(u, \xi; t, y), \nabla_\xi p^X(u, \xi; t, y) \rangle}{p^X(u, \xi; t, y)^2} \phi(u, \xi) d\xi du \\ & \quad + C \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial \xi_j} a_{ij}(u, \xi) \right|^2 \phi(u, \xi) d\xi du, \end{aligned} \quad (3-5)$$

where C depends on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+$ and Λ . By using (1-6), we estimate the final term of the right-hand side of (3-2) as follows:

$$\begin{aligned} & \left| \int_{s_1}^{s_2} \int_{\mathbb{R}^d} (\log p^X(u, \xi; t, y)) \frac{\partial}{\partial u} \phi(u, \xi) d\xi du \right| \\ & \leq \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left(|\log C_G^+| + |\log C_G^-| + \frac{d}{2} |\log(t-u)| + \frac{\gamma_G^- |y-\xi|^2}{t-u} \right) \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du. \end{aligned}$$

Hence, there exists a constant C depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+$ and Λ such that

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} (\log p^X(s, \xi; t, y)) \phi(s_1, \xi) d\xi \right| \\ & \leq C \int_{s_1}^{s_2} (1 + |\log(t-u)|) \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du + C \int_{s_1}^{s_2} (t-u)^{-1} \int_{\mathbb{R}^d} |y-\xi|^2 \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du. \quad (3-6) \end{aligned}$$

Therefore, by (3-2), (3-3), (3-4), (3-5) and (3-6) we obtain the lemma. \square

Next, we state the fact on the integrability of \mathcal{E} as a lemma. The proof is obtained by the standard argument; see, for example, [Stroock and Varadhan 1979, Theorem 4.2.1]. So, we omit it.

Lemma 3.3. *Let τ_1, τ_2 be stopping times such that $0 \leq \tau_1 \leq \tau_2$ almost surely. It holds that, for any $q \in \mathbb{R}$,*

$$E[\mathcal{E}(t \wedge \tau_1, t \wedge \tau_2; X^x)^q] \leq e^{C(1+q^2)t}, \quad t > 0, \quad x, y \in \mathbb{R}^d,$$

where C is a constant depending on $d, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$.

Lemma 3.4. *Let τ_1, τ_2 be stopping times such that $0 \leq \tau_1 \leq \tau_2 \leq t$ almost surely. It holds that*

$$p^X(0, x; t, y) E^{X_t^y} \left[\int_0^t \mathcal{E}(u \wedge \tau_1, u \wedge \tau_2; X^x)^q du \right] \leq C t^{-d/2+1-\varepsilon} e^{C(1+q^2)t} \exp\left(-\frac{\gamma|x-y|^2}{t}\right)$$

for $t \in (0, \infty)$, $x, y \in \mathbb{R}^d$, $q \in \mathbb{R}$ and sufficiently small $\varepsilon > 0$, where C and γ are positive constants depending on $d, \varepsilon, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$.

Proof. In view of Fubini's theorem and (2-11), it is sufficient to show that there exist positive constants C and γ , depending on $d, \varepsilon, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$, such that

$$\int_0^t E[\mathcal{E}(u \wedge \tau_1, u \wedge \tau_2; X^x)^q p^X(u, X_u; t, y)] du \leq C t^{-d/2+1-\varepsilon} e^{C(1+q^2)t} \exp\left(-\frac{\gamma|x-y|^2}{t}\right) \quad (3-7)$$

for $t \in (0, \infty)$ and $x, y \in \mathbb{R}^d$. By (1-6) and Hölder's inequality, we have

$$\begin{aligned} & \int_0^t E[\mathcal{E}(u \wedge \tau_1, u \wedge \tau_2; X^x)^q p^X(u, X_u; t, y)] du \\ & \leq C_G^+ \int_0^t E \left[\mathcal{E}(u \wedge \tau_1, u \wedge \tau_2; X^x)^q (t-u)^{-\frac{d}{2}} \exp\left(-\frac{\gamma_G^+ |X_u - y|^2}{t-u}\right) \right] du \\ & \leq C_G^+ \left(\int_0^t E[\mathcal{E}(u \wedge \tau_1, u \wedge \tau_2; X^x)^{(d+\varepsilon)q/\varepsilon}] du \right)^{\frac{\varepsilon}{d+\varepsilon}} \\ & \quad \times \left(\int_0^t E \left[(t-u)^{-(d+\varepsilon)/2} \exp\left(-\frac{(d+\varepsilon)\gamma_G^+ |X_u - y|^2}{d(t-u)}\right) \right] du \right)^{\frac{d}{d+\varepsilon}}. \end{aligned}$$

Hence, in view of Lemma 3.3, to show (3-7) it is sufficient to prove that

$$\int_0^t E \left[(t-u)^{-(d+\varepsilon)/2} \exp\left(-\frac{(d+\varepsilon)\gamma_G^+ |X_u^x - y|^2}{d(t-u)}\right) \right] du \leq C t^{-(d+\varepsilon)/2+1} \exp\left(-\gamma \frac{|x-y|^2}{t}\right) \quad (3-8)$$

for $t \in (0, \infty)$ and $x, y \in \mathbb{R}^d$, where C and γ are constants depending on $d, \varepsilon, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$.

Let $\tilde{\gamma} := (1 + \varepsilon/d)\gamma_G^+$. By (1-6) again, we have for $u \in (0, t)$ that

$$\begin{aligned} & E \left[(t-u)^{-d/2} \exp\left(-\frac{\tilde{\gamma} |X_u^x - y|^2}{t-u}\right) \right] \\ & \leq C_G^+ u^{-d/2} (t-u)^{-d/2} \int_{\mathbb{R}^d} \exp\left(-\frac{\tilde{\gamma} |\xi - y|^2}{t-u}\right) \exp\left(-\frac{\gamma_G^+ |\xi - x|^2}{u}\right) d\xi \\ & = C_G^+ u^{-d/2} (t-u)^{-d/2} \exp\left(-\frac{\gamma_G^+ \tilde{\gamma}}{u\tilde{\gamma} + (t-u)\gamma_G^+} |x-y|^2\right) \\ & \quad \times \int_{\mathbb{R}^d} \exp\left(-\frac{u\tilde{\gamma} + (t-u)\gamma_G^+}{u(t-u)} \left| \xi - \frac{\gamma_G^+(t-u)x + \tilde{\gamma}uy}{u\tilde{\gamma} + (t-u)\gamma_G^+} \right|^2\right) d\xi \\ & = (2\pi)^{d/2} C_G^+ (u\tilde{\gamma} + (t-u)\gamma_G^+)^{-d/2} \exp\left(-\frac{\gamma_G^+ \tilde{\gamma}}{u\tilde{\gamma} + (t-u)\gamma_G^+} |x-y|^2\right) \\ & \leq (2\pi)^{d/2} C_G^+ \gamma_G^{+d/2} t^{-d/2} \exp\left(-\frac{\gamma_G^+ |x-y|^2}{t}\right). \end{aligned}$$

Hence, there exists a positive constant C depending on $d, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$, such that

$$E \left[(t-u)^{-d/2} \exp\left(-\frac{\tilde{\gamma} |X_u^x - y|^2}{t-u}\right) \right] \leq C t^{-d/2} \exp\left(-\frac{\gamma_G^+ |x-y|^2}{t}\right), \quad u \in (0, t).$$

Thus, we obtain (3-8). \square

Lemma 3.5. *Let $t \in (0, \infty)$ and τ_1, τ_2 be stopping times such that $0 \leq \tau_1 \leq \tau_2 \leq t$ almost surely. Then, it holds that*

$$p^X(0, x; t, y) E^{X_t^x=y}[\mathcal{E}(s \wedge \tau_1, s \wedge \tau_2; X^x)^q] \leq C t^{-d/2} e^{C(1+q^2)t} \exp\left(-\frac{\gamma |x-y|^2}{t}\right)$$

for $t \in (0, \infty)$, $x, y \in \mathbb{R}^d$, $q \in \mathbb{R}$ and $s \in [0, t)$, where C and γ are positive constants depending on d , γ_G^- , γ_G^+ , C_G^- , C_G^+ , m , M , θ , Λ , $\|b\|_\infty$ and $\|c\|_\infty$.

Proof. Let $s_1, s_2 \in (0, t)$ such that $s_1 \leq s_2$. In view of (2-10), by (2-11) and Itô's formula we have

$$\begin{aligned}
& p^X(0, x; t, y) E^{X_t^x=y} [\mathcal{E}((s \wedge \tau_1) \vee s_1, (s \wedge \tau_2) \wedge s_2; X^x)^q] \\
&= E[p^X(s_2, X_{s_2}^x; t, y) \mathcal{E}((s \wedge \tau_1) \vee s_1, (s \wedge \tau_2) \wedge s_2; X^x)^q] \\
&= p^X(0, x; t, y) + E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} \left(\frac{\partial}{\partial u} p^X(u, \xi; t, y) \right) \Big|_{\xi=X_u^x} \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q du \right] \\
&+ E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} (L_u^X p^X(u, \cdot; t, y))(X_u^x) \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q du \right] \\
&+ \frac{q}{2} E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q \times \langle \sigma(u, X_u^x)^T \nabla_z p^X(u, z; t, y) \Big|_{z=X_u^x}, b_\sigma(u, X_u^x) \rangle du \right] \\
&+ \frac{q^2}{2} E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} p^X(u, X_u^x; t, y) \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q |b_\sigma(u, X_u^x)|^2 du \right] \\
&+ q E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} p^X(u, X_u^x; t, y) \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q c(u, X_u^x) du \right].
\end{aligned}$$

Hence, by (2-6) we obtain

$$\begin{aligned}
& p^X(0, x; t, y) E^{X_t^x=y} [\mathcal{E}((s \wedge \tau_1) \vee s_1, (s \wedge \tau_2) \wedge s_2; X^x)^q] \\
&= p^X(0, x; t, y) \\
&+ \frac{q}{2} E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q \times \langle \sigma(u, X_u^x)^T \nabla_z p^X(u, z; t, y) \Big|_{z=X_u^x}, b_\sigma(u, X_u^x) \rangle du \right] \\
&+ \frac{q^2}{2} E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} p^X(u, X_u^x; t, y) \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q |b_\sigma(u, X_u^x)|^2 du \right] \\
&+ q E \left[\int_{(s \wedge \tau_1) \vee s_1}^{(s \wedge \tau_2) \wedge s_2} p^X(u, X_u^x; t, y) \mathcal{E}((s \wedge \tau_1) \vee s_1, u; X^x)^q c(u, X_u^x) du \right].
\end{aligned}$$

In view of the boundedness of $\det \sigma$, b and c , the desired estimate is obtained, once we show the estimates

$$\begin{aligned}
& E \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q p^X(u, X_u^x; t, y) du \right] \\
&\leq C t^{-d/2+1-\varepsilon} e^{C(1+q^2)t} \exp\left(-\gamma \frac{|x-y|^2}{t}\right), \quad (3-9)
\end{aligned}$$

$$\begin{aligned}
& E \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q |\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x} du \right] \\
&\leq C t^{-d/2+(1-\varepsilon)/2} e^{C(1+q^2)t} \exp\left(-\frac{\gamma|x-y|^2}{t}\right) \quad (3-10)
\end{aligned}$$

for sufficiently small $\varepsilon > 0$, where C and γ are positive constants depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. The first estimate (3-9) follows, because by (2-11) and Lemma 3.4 we have

$$\begin{aligned} E \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q p^X(u, X_u^x; t, y) du \right] \\ = p^X(0, x; t, y) E^{X_t^x=y} \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q du \right] \\ \leq C t^{-d/2+1-\varepsilon} e^{C(1+q^2)t} \exp\left(-\gamma \frac{|x-y|^2}{t}\right), \end{aligned}$$

where C and γ are positive constants depending on $d, \varepsilon, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Now we show (3-10). By (2-11) and Hölder's inequality, we have

$$\begin{aligned} E \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q |\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x} du \right] \\ = p^X(0, x; t, y) E^{X_t^x=y} \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q \frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} du \right] \\ \leq p^X(0, x; t, y) \left(\int_{s_1}^{s_2} E^{X_t^x=y} \left[\mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^{2q/(1-2\varepsilon)} du \right] \right)^{1/2-\varepsilon} \\ \times \left(\int_{s_1}^{s_2} [u(t-u)]^{-1/2} du \right)^\varepsilon \left(\int_{s_1}^{s_2} [u(t-u)]^\varepsilon E^{X_t^x=y} \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 du \right] \right)^{1/2}. \end{aligned}$$

Lemma 3.4 and (1-6) imply that

$$\begin{aligned} E \left[\int_{s_1}^{s_2} \mathcal{E}(u \wedge [(s \wedge \tau_1) \vee s_1], u; X^x)^q |\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x} du \right] \\ \leq C t^{-d/4+1/2-\varepsilon} e^{C(1+q^2)t} \exp\left(-\gamma \frac{|x-y|^2}{t}\right) \\ \times \left(p^X(0, x; t, y) \int_{s_1}^{s_2} [u(t-u)]^\varepsilon E^{X_t^x=y} \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 du \right] \right)^{1/2}, \end{aligned}$$

where C and γ are positive constants depending on $d, \varepsilon, \gamma_G^+, C_G^+, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Hence, to show (3-10), it is sufficient to prove

$$\begin{aligned} p^X(0, x; t, y) \int_0^t [u(t-u)]^{\varepsilon/2} E^{X_t^x=y} \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 du \right] \\ \leq C (t^{-d/2+\varepsilon} + t^{-d/2+\varepsilon} |\log t|), \quad (3-11) \end{aligned}$$

where C is a constant depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta$ and Λ . Equation (2-11) implies that

$$\begin{aligned} p^X(0, x; t, y) & \int_0^t [u(t-u)]^{\varepsilon/2} E^{X_t^x=y} \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 \right] du \\ & = \int_0^t [u(t-u)]^{\varepsilon/2} E \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 p^X(u, X_u^x; t, y) \right] du \\ & = \int_0^t [u(t-u)]^{\varepsilon/2} \int_{\mathbb{R}^d} \left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|}{p^X(u, \xi; t, y)} \right)^2 p^X(u, \xi; t, y) p^X(0, x; u, \xi) d\xi du. \end{aligned}$$

By (1-6), we have

$$\begin{aligned} p^X(0, x; t, y) & \int_0^t [u(t-u)]^{\varepsilon/2} E^{X_t^x=y} \left[\left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|_{\xi=X_u^x}}{p^X(u, X_u^x; t, y)} \right)^2 \right] du \\ & \leq (C_G^+)^2 \int_0^t \int_{\mathbb{R}^d} \left(\frac{|\nabla_\xi p^X(u, \xi; t, y)|}{p^X(u, \xi; t, y)} \right)^2 [u(t-u)]^{-(d-\varepsilon)/2} \\ & \quad \times \exp \left[-\gamma_G^+ \left(\frac{|\xi-x|^2}{u} + \frac{|y-\xi|^2}{t-u} \right) \right] d\xi du. \quad (3-12) \end{aligned}$$

For fixed t, x and y , let

$$\phi(u, \xi) := [u(t-u)]^{-(d-\varepsilon)/2} \exp \left[-\gamma_G^+ \left(\frac{|\xi-x|^2}{u} + \frac{|y-\xi|^2}{t-u} \right) \right].$$

Denote the surface area of the unit sphere in \mathbb{R}^d by ω_d for $d \geq 2$. In the case $d = 1$, let $\omega_d = 2$. Explicit calculation implies that

$$\begin{aligned} & \int_0^t \left(\int_{\mathbb{R}^d} e^{2m|\xi|/(\theta-2)} \phi(u, \xi)^{\theta/(\theta-2)} d\xi \right)^{(\theta-2)/\theta} du \\ & = \int_0^t [u(t-u)]^{-(d-\varepsilon)/2} \times \left(\int_{\mathbb{R}^d} e^{2m|\xi|/(\theta-2)} \exp \left[-\frac{\gamma_G^+ \theta}{\theta-2} \left(\frac{|\xi-x|^2}{u} + \frac{|y-\xi|^2}{t-u} \right) \right] d\xi \right)^{(\theta-2)/\theta} du \\ & = \exp \left(-\frac{\gamma_G^+ |x-y|^2}{t} \right) \int_0^t [u(t-u)]^{-(d-\varepsilon)/2} \\ & \quad \times \left(\int_{\mathbb{R}^d} e^{2m|\xi|/(\theta-2)} \exp \left[-\frac{\gamma_G^+ \theta t}{(\theta-2)u(t-u)} \left| \xi - \frac{(t-u)x + uy}{t} \right|^2 \right] d\xi \right)^{(\theta-2)/\theta} du. \end{aligned}$$

Hence, noting that for $\mu_1 \in [0, \infty)$, $\mu_2 \in (0, \infty)$ and $v \in \mathbb{R}^d$

$$\begin{aligned}
& \int_{\mathbb{R}^d} e^{\mu_1|\xi|} \exp(-\mu_2|\xi - v|^2) d\xi \\
&= \int_{\mathbb{R}^d} e^{\mu_1|\xi+v|} \exp(-\mu_2|\xi|^2) d\xi \\
&\leq e^{\mu_1|v|} \int_{\mathbb{R}^d} \exp(\mu_1|\xi| - \mu_2|\xi|^2) d\xi \\
&= \omega_d e^{\mu_1|v|} \int_{(0,\infty)} r^{d-1} \exp(\mu_1 r - \mu_2 r^2) dr \\
&= \omega_d \mu_2^{-d/2} e^{\mu_1|v|} \int_{(0,\infty)} r^{d-1} \exp\left(\frac{\mu_1}{\sqrt{\mu_2}} r - r^2\right) dr \\
&= \omega_d \mu_2^{-d/2} \int_0^{1+\mu_1/\sqrt{\mu_2}} r^{d-1} \exp\left(\frac{\mu_1}{\sqrt{\mu_2}} r - r^2\right) dr + \omega_d \mu_2^{-d/2} \int_{1+\mu_1/\sqrt{\mu_2}}^\infty r^{d-1} \exp\left(\frac{\mu_1}{\sqrt{\mu_2}} r - r^2\right) dr \\
&\leq \frac{\omega_d \mu_2^{-d/2}}{d} \left(1 + \frac{\mu_1}{\sqrt{\mu_2}}\right)^d \exp\left[\frac{\mu_1}{\sqrt{\mu_2}} \left(1 + \frac{\mu_1}{\sqrt{\mu_2}}\right)\right] + \omega_d \mu_2^{-d/2} \int_{1+\mu_1/\sqrt{\mu_2}}^\infty r^{d-1} e^{-r} dr \\
&\leq C \mu_2^{-d/2} \exp\left[C \frac{\mu_1}{\sqrt{\mu_2}} \left(1 + \frac{\mu_1}{\sqrt{\mu_2}}\right)\right],
\end{aligned}$$

where C is a constant depending on d , we have

$$\begin{aligned}
& \int_0^t \left(\int_{\mathbb{R}^d} e^{2m|\xi|/(\theta-2)} \phi(u, \xi)^{\theta/(\theta-2)} d\xi \right)^{(\theta-2)/\theta} du \\
&\leq C_1 t^{-d/2+d/\theta} \exp\left(-\frac{\gamma_G^+ |x-y|^2}{t}\right) \\
&\quad \times \int_0^t [u(t-u)]^{-d/\theta+\varepsilon/2} \exp\left[C_1 \sqrt{u\left(1-\frac{u}{t}\right)} \left(1 + \sqrt{u\left(1-\frac{u}{t}\right)}\right)\right] du \\
&\leq C_2 t^{-d/2+d/\theta} e^{C_2 t} \int_0^t [u(t-u)]^{-d/\theta+\varepsilon/2} du \\
&\leq C_3 t^{-d/2+1-d/\theta+\varepsilon/2} e^{C_2 t},
\end{aligned}$$

where C_1, C_2, C_3 are constants depending on $d, \varepsilon, \gamma_G^+, m$ and θ . Hence, by Hölder's inequality and (1-3), we have

$$\begin{aligned}
& \sum_{i,j=1}^d \int_{s_1}^{s_2} \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial \xi_j} a_{ij}(u, \xi) \right|^2 \phi(u, \xi) d\xi du \\
&\leq \int_0^t \left(\int_{\mathbb{R}^d} e^{2m|\xi|/(\theta-2)} \phi(u, \xi)^{\theta/(\theta-2)} d\xi \right)^{(\theta-2)/\theta} du \sum_{i,j=1}^d \left(\sup_{u \in [0,t]} \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial \xi_j} a_{ij}(u, \xi) \right|^\theta e^{-m|\xi|} d\xi \right)^{2/\theta} \\
&\leq C t^{-d/2+1-d/\theta+\varepsilon/2} e^{Ct},
\end{aligned}$$

where C is a constant depending on $d, \varepsilon, \gamma_G^+, m, M$ and θ . On the other hand, by explicit calculation, we have

$$\begin{aligned}
\lim_{s \downarrow 0} (1 + |\log(t-s)|) \int_{\mathbb{R}^d} \phi(s, \xi) d\xi &= 0, \\
\lim_{s \downarrow 0} (t-s)^{-1} \int_{\mathbb{R}^d} |y-\xi|^2 \phi(s, \xi) d\xi &= 0, \\
\lim_{s \uparrow t} (1 + |\log(t-s)|) \int_{\mathbb{R}^d} \phi(s, \xi) d\xi &= 0, \\
\lim_{s \uparrow t} (t-s)^{-1} \int_{\mathbb{R}^d} |y-\xi|^2 \phi(s, \xi) d\xi &= 0, \\
\int_0^t \int_{\mathbb{R}^d} \frac{|\nabla_\xi \phi(u, \xi)|^2}{\phi(u, \xi)} d\xi du &\leq C t^{-d/2+\varepsilon}, \\
\int_0^t (1 + |\log(t-u)|) \int_{\mathbb{R}^d} \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du &\leq C t^{-d/2+\varepsilon} |\log t|, \\
\int_0^t (t-u)^{-1} \int_{\mathbb{R}^d} |y-\xi|^2 \left| \frac{\partial}{\partial u} \phi(u, \xi) \right| d\xi du &\leq C t^{-d/2+\varepsilon},
\end{aligned}$$

where C is a constant depending on d, ε and γ_G^+ . In view of these results, applying Lemma 3.1 to (3-12), we obtain (3-11). \square

From Lemma 3.5 we can easily show the Gaussian estimate for p with constants depending on the suitable constants.

Proposition 3.6. *For $s, t \in [0, \infty)$ such that $s < t$, and $x, y \in \mathbb{R}^d$,*

$$\frac{C_1 e^{-C_1(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_1 |x-y|^2}{t-s}\right) \leq p(s, x; t, y) \leq \frac{C_2 e^{C_2(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_2 |x-y|^2}{t-s}\right),$$

where γ_1, γ_2, C_1 and C_2 are positive constants depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$.

Proof. Since the argument follows even if a, b and c are replaced by $a(\cdot - s, \cdot)$, $b(\cdot - s, \cdot)$ and $c(\cdot - s, \cdot)$ respectively, it is sufficient to show that there exist positive constants γ_1, γ_2, C_1 and C_2 depending on $d, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that

$$C_1 t^{-d/2} e^{-C_1 t} \exp\left(-\frac{\gamma_1 |x-y|^2}{t}\right) \leq p(0, x; t, y) \leq C_2 t^{-d/2} e^{C_2 t} \exp\left(-\frac{\gamma_2 |x-y|^2}{t}\right) \quad (3-13)$$

for $t \in (0, \infty)$ and $x, y \in \mathbb{R}^d$. The upper estimate in (3-13) follows immediately from (2-8) and Lemma 3.5.

Now we prove the lower estimate in (3-13). From Hölder's inequality, it follows that

$$1 \leq E^{X_t^i=y} [\mathcal{E}(s \wedge \tau_1, s \wedge \tau_2; X^x)^{-1}] E^{X_t^i=y} [\mathcal{E}(s \wedge \tau_1, s \wedge \tau_2; X^x)].$$

Hence, by Lemma 3.5, we have

$$\begin{aligned} p^X(0, x; t, y) E^{X_t^x=y}[\mathcal{E}(s \wedge \tau_1, s \wedge \tau_2; X^x)] &\geq \frac{p^X(0, x; t, y)^2}{p^X(0, x; t, y) E^{X_t^x=y}[\mathcal{E}(s \wedge \tau_1, s \wedge \tau_2; X^x)^{-1}]} \\ &\geq C t^{d/2} e^{-C't} p^X(0, x; t, y)^2, \end{aligned}$$

where C and C' are positive constants depending on $d, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. This inequality, (2-8) and (1-6) imply the lower bound in (3-13). \square

4. The regularity of $p(0, x; t, y)$ in x

Assume that a is smooth and set notation as in Section 2. In this section, we prove the Hölder continuity of $p(0, x; t, y)$ in x , with constant depending only on suitable ones. The precise statement is as follows:

Proposition 4.1. *For any $R > 0$ and sufficiently small $\varepsilon > 0$, there exists a constant C depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that*

$$|p(0, x; t, y) - p(0, z; t, y)| \leq C t^{-d/2-1} e^{Ct} |x - z|^{1-\varepsilon}$$

for $t \in (0, \infty), x, z \in B(0; R/2)$ and $y \in \mathbb{R}^d$.

We use the coupling method; see, for example, [Lindvall and Rogers 1986; Cranston 1991]. Let $x, z \in \mathbb{R}^d$. Given (X^x, B) defined by (2-3), we consider the stochastic process Z^z defined by

$$\begin{cases} Z_t^z = z + \int_0^{t \wedge \tau} \sigma(s, Z_s^z) d\tilde{B}_s + \int_{t \wedge \tau}^t \sigma(s, Z_s^z) dB_s, \\ \tilde{B}_t = \int_0^{t \wedge \tau} \left(I - \frac{2(\sigma(s, Z_s^z)^{-1}(X_s^x - Z_s^z)) \otimes (\sigma(s, Z_s^z)^{-1}(X_s^x - Z_s^z))}{|\sigma(s, Z_s^z)^{-1}(X_s^x - Z_s^z)|^2} \right) dB_s, \end{cases} \quad (4-1)$$

where τ is the stopping time defined by $\tau := \inf\{t \geq 0 : X_t^x = Z_t^z\}$.

To see the existence and uniqueness of Z^z , for each $n \in \mathbb{N}$ consider the following stochastic differential equation for $\check{Z}^{n,z}$:

$$\begin{cases} d\check{Z}_t^{n,z} = \sigma(t, \check{Z}_t^{n,z}) \times \left(I - \frac{2(\sigma(t, \check{Z}_t^{n,z})^{-1}(X_t^x - \check{Z}_t^{n,z})) \otimes (\sigma(t, \check{Z}_t^{n,z})^{-1}(X_t^x - \check{Z}_t^{n,z}))}{|\sigma(t, \check{Z}_t^{n,z})^{-1}(X_t^x - \check{Z}_t^{n,z})|^2} \right) \mathbb{1}_{\{t < \check{\tau}_n\}} dB_t \\ \check{Z}_0^{n,z} = z, \end{cases}$$

where $\check{\tau}_n := \inf\{t \geq 0 : |X_t^x - \check{Z}_t^{n,z}| < 1/n\}$. Note that the equations have random coefficients, since we are considering equations where X^x is given. Now we see the existence and the uniqueness of $\check{Z}^{n,z}$. Let

$$G_n(t, \xi) := \begin{cases} \sigma(t, \xi) \left(I - \frac{2(\sigma(t, \xi)^{-1}(X_t^x - \xi)) \otimes (\sigma(t, \xi)^{-1}(X_t^x - \xi))}{|\sigma(t, \xi)^{-1}(X_t^x - \xi)|^2} \right) & \text{if } (t, \xi) \in [0, \infty) \times \mathbb{R}^d \\ & \text{and } |X_t^x - \xi| \geq 1/n, \\ 0 & \text{otherwise.} \end{cases}$$

Then, there exists a constant C_n such that $|G_n(t, \xi) - G_n(t, \eta)| \leq C_n$ for $t \in [0, \infty)$ and $\xi, \eta \in \mathbb{R}^d$. Note that C_n is nonrandom. Let Y, W be stochastic processes satisfying

$$\begin{cases} dY_t = G_n(t, Y_t) \mathbb{1}_{\{t < \tau_n^Y\}} dB_t, \\ Y_0 = z, \end{cases} \quad \begin{cases} dW_t = G_n(t, W_t) \mathbb{1}_{\{t < \tau_n^W\}} dB_t, \\ W_0 = z, \end{cases} \quad (4-2)$$

where $\tau_n^Y := \inf\{t \geq 0 : |X_t^x - Y_t| < 1/n\}$ and $\tau_n^W := \inf\{t \geq 0 : |X_t^x - W_t| < 1/n\}$. Then, by Proposition 1.1(iv), Chapter II and (6.16) of Theorem 6.10, Chapter I of [Ikeda and Watanabe 1989], we have

$$\begin{aligned} E \left[\sup_{s \in [0, t]} |Y_{s \wedge \tau_n^Y \wedge \tau_n^W} - W_{s \wedge \tau_n^Y \wedge \tau_n^W}|^2 \right] &= E \left[\sup_{s \in [0, t]} \left| \int_0^{s \wedge \tau_n^Y \wedge \tau_n^W} (G_n(v, Y_v) \mathbb{1}_{\{v < \tau_n^Y\}} - G_n(v, W_v) \mathbb{1}_{\{v < \tau_n^W\}}) dB_v \right|^2 \right] \\ &= E \left[\sup_{s \in [0, t]} \left| \int_0^s (G_n(v, Y_v) - G_n(v, W_v)) \mathbb{1}_{\{v < \tau_n^Y\}} \mathbb{1}_{\{v < \tau_n^W\}} dB_v \right|^2 \right] \\ &\leq 4E \left[\int_0^t |G_n(v, Y_v) - G_n(v, W_v)|^2 \mathbb{1}_{\{v < \tau_n^Y\}} \mathbb{1}_{\{v < \tau_n^W\}} dv \right] \\ &\leq 4C_n^2 \int_0^t E \left[|Y_v - W_v|^2 \mathbb{1}_{\{v < \tau_n^Y\}} \mathbb{1}_{\{v < \tau_n^W\}} \right] dv \\ &\leq 4C_n^2 \int_0^t E \left[\sup_{s \in [0, v]} |Y_{s \wedge \tau_n^Y \wedge \tau_n^W} - W_{s \wedge \tau_n^Y \wedge \tau_n^W}|^2 \right] dv. \end{aligned}$$

Hence, by Gronwall's inequality, we have

$$Y_{t \wedge \tau_n^Y \wedge \tau_n^W} = W_{t \wedge \tau_n^Y \wedge \tau_n^W}, \quad t \in [0, \infty) \quad (4-3)$$

almost surely. If $\tau_n^Y \leq \tau_n^W$ and $\tau_n^Y < \infty$ for some events, then by letting $t \rightarrow \infty$ in (4-3) we have $Y_{\tau_n^Y} = W_{\tau_n^Y}$. Hence, $\tau_n^Y = \tau_n^W$ for these events. Similarly, if $\tau_n^Y \geq \tau_n^W$ and $\tau_n^W < \infty$ for some events, then we have $\tau_n^Y = \tau_n^W$ for these events. Therefore, we obtain

$$\tau_n^Y = \tau_n^W \quad (4-4)$$

almost surely. On the other hand, (4-2) implies that $Y_{\tau_n^Y} = Y_{t \vee \tau_n^Y}$ and $W_{\tau_n^W} = W_{t \vee \tau_n^W}$ for $t \in [0, \infty)$. Hence, by (4-3) and (4-4) we obtain that $Y_t = W_t$ for $t \in [0, \infty)$ almost surely. Thus, we have uniqueness. To see existence, let

$$\widehat{G}_n(t, \xi) := \sigma(t, \xi) \left(I - \frac{2(\sigma(t, \xi))^{-1}(X_t^x - \xi) \otimes (\sigma(t, \xi))^{-1}(X_t^x - \xi)}{(|\sigma(t, \xi))^{-1}(X_t^x - \xi)| \vee (\Lambda^{-(1/2)} n^{-1})^2} \right)$$

for $t \in [0, \infty)$ and $\xi \in \mathbb{R}^d$. Then, there exists a constant C_n such that $|\widehat{G}_n(t, \xi) - \widehat{G}_n(t, \eta)| \leq C_n$ for $t \in [0, \infty)$ and $\xi, \eta \in \mathbb{R}^d$. Define a sequence of stochastic processes $\{Y^m : m \in \mathbb{N} \cup \{0\}\}$ by $Y_t^0 = z$ for $t \in [0, \infty)$ and

$$Y_t^m := z + \int_0^t \widehat{G}_n(s, Y_s^{m-1}) dB_s \quad (4-5)$$

for $t \in [0, \infty)$ and $m \in \mathbb{N}$ by iteration. Then, by a similar calculation as above, we have for $m \in \mathbb{N}$ and $t \in [0, \infty)$ that

$$E \left[\sup_{s \in [0, t]} |Y_s^{m+1} - Y_s^m|^2 \right] \leq 4C_n^2 \int_0^t E \left[\sup_{s \in [0, v]} |Y_s^m - Y_s^{m-1}|^2 \right] dv.$$

Applying this inequality iteratively, for $m \in \mathbb{N}$ and $t \in [0, \infty)$ we obtain

$$\begin{aligned} E \left[\sup_{s \in [0, t]} |Y_s^{m+1} - Y_s^m|^2 \right] &\leq (4C_n^2)^m \int_0^t \int_0^{v_m} \cdots \int_0^{v_2} E \left[\sup_{s \in [0, v_1]} |Y_s^1 - Y_s^0|^2 \right] dv_1 dv_2 \cdots dv_m \\ &= (4C_n^2)^m \int_0^t \int_0^{v_m} \cdots \int_0^{v_2} E \left[\sup_{s \in [0, v_1]} \left| \int_0^s \widehat{G}_n(w, z) dB_w \right|^2 \right] dv_1 dv_2 \cdots dv_m \\ &\leq \frac{(4C_n^2)^m t^m}{(m+1)!} E \left[\sup_{s \in [0, t]} \left| \int_0^s \widehat{G}_n(w, z) dB_w \right|^2 \right]. \end{aligned}$$

Since [Ikeda and Watanabe 1989, (6.16) of Theorem 6.10, Chapter I] implies

$$E \left[\sup_{s \in [0, t]} \left| \int_0^s \widehat{G}_n(w, z) dB_w \right|^2 \right] \leq 4E \left[\int_0^t |\widehat{G}_n(w, z)|^2 dw \right] < \infty$$

for $t \in [0, \infty)$, we have

$$\sum_{m=1}^{\infty} E \left[\sup_{s \in [0, t]} |Y_s^{m+1} - Y_s^m|^2 \right] < \infty$$

for $t \in [0, \infty)$. Hence, $\{Y^m\}$ is a Cauchy sequence in $L^2(\Omega; \mathcal{C})$, where \mathcal{C} is the complete metric space $C([0, \infty); \mathbb{R}^d)$ with distance function given by

$$D(w, w') := \sum_{k=1}^{\infty} 2^{-k} \left(\sup_{t \in [0, k]} |w(t) - w'(t)| \right) \wedge 1, \quad w, w' \in C([0, \infty); \mathbb{R}^d).$$

Therefore, there exists a stochastic process Y in $L^2(\Omega; \mathcal{C})$ which satisfies

$$\lim_{m \rightarrow \infty} E \left[\sup_{s \in [0, t]} |Y_s - Y_s^m|^2 \right] = 0$$

for $t \in [0, \infty)$. By taking the limit in (4-5) as $m \rightarrow \infty$, we have

$$Y_t = z + \int_0^t \widehat{G}_n(s, Y_s) dB_s, \quad t \in [0, \infty) \quad (4-6)$$

almost surely. Let $\tau_n^Y := \inf\{t \geq 0 : |X_t^x - Y_t| < 1/n\}$. Note that (2-2) implies that

$$|\sigma(t, Y_t)^{-1}(X_t^x - Y_t)| \vee (\Lambda^{-1/2})n^{-1} = |\sigma(t, Y_t)^{-1}(X_t^x - Y_t)|, \quad t \in [0, \tau_n^Y)$$

almost surely. Applying [Ikeda and Watanabe 1989, Proposition 1.1(iv), Chapter II] to (4-6), we see that the $Y_{\cdot \wedge \tau_n^Y}$ satisfy the stochastic differential equation for $\check{Z}^{n, z}$. Thus, we obtain existence.

We remark that $\{\check{Z}^{n,z} : n \in \mathbb{N}\}$ is consistent; i.e., $\check{Z}_{t \wedge \check{\tau}_n}^{m,z} = \check{Z}_{t \wedge \check{\tau}_n}^{n,z}$ for $m > n$ almost surely. This fact is immediately obtained by [Ikeda and Watanabe 1989, Proposition 1.1(iv), Chapter II] and uniqueness. Define the stochastic processes $(Z_t^z, \tilde{B}_t; t \in [0, \tau])$ by

$$\begin{aligned} Z_t^z &= \check{Z}_t^{n,z}, \\ \tilde{B}_t &= \int_0^t \left(I - \frac{2(\sigma(s, \check{Z}_s^{n,z})^{-1}(X_s^x - \check{Z}_s^{n,z})) \otimes (\sigma(s, \check{Z}_s^{n,z})^{-1}(X_s^x - \check{Z}_s^{n,z}))}{|\sigma(s, \check{Z}_s^{n,z})^{-1}(X_s^x - \check{Z}_s^{n,z})|^2} \right) dB_s \end{aligned}$$

for $t \in [0, \tau_n^Z)$ and $n \in \mathbb{N}$, where $\tau_n^Z := \inf\{t \geq 0 : |X_t^x - Z_t^z| < 1/n\}$. Then, (4-1) holds for $t \in [0, \tau)$. On the other hand, by applying [Ikeda and Watanabe 1989, Proposition 1.1(iv), Chapter II], we have that $Z_{\cdot \wedge \tau_n^Z}^z$ solves the stochastic differential equation of $\check{Z}^{n,z}$ for $n \in \mathbb{N}$. Hence, $(Z_t^z, \tilde{B}_t; t \in [0, \tau])$ are determined almost surely and uniquely. Let

$$H_t := I - \frac{2[\sigma(t, Z_t^z)^{-1}(X_t^x - Z_t^z)] \otimes [\sigma(t, Z_t^z)^{-1}(X_t^x - Z_t^z)]}{|\sigma(t, Z_t^z)^{-1}(X_t^x - Z_t^z)|^2}$$

for $t \in [0, \tau)$. Then, H_t is an orthogonal matrix for all $t \in [0, \tau)$, and hence \tilde{B}_t is a d -dimensional Brownian motion for $t \in [0, \tau)$. Hence, $(Z_t^z, \tilde{B}_t; t \in [0, \tau))$ are extended to $(Z_t^z, \tilde{B}_t; t \in [0, \tau])$ almost surely and uniquely. By the Lipschitz continuity of σ , (4-1) is solved almost surely and uniquely for $t \in [\tau, \infty)$, and thus we obtain $(Z_t^z; t \in [0, \infty))$ almost surely and uniquely; see [Stroock and Varadhan 1979, Section 6.6]. From this fact we have that Z_t^z is \mathcal{F}_t -measurable for $t \in [0, \infty)$. Hence, if $x = z$, X^x and Z^z have the same law. Moreover, $X_t^x = Z_t^z$ for $t \in [\tau, \infty)$ almost surely.

Lemma 4.2. *For $R > 0$ and sufficiently small $\varepsilon > 0$, there exist positive constants C and c_0 depending on $d, \varepsilon, R, \rho_R$ and Λ such that*

$$E[t \wedge \tau] \leq C(1 + t^2)|x - z|^{1-\varepsilon} \quad (4-7)$$

for $t \in [0, \infty)$ and $x, z \in B(0; R/2)$ such that $|x - z| \leq c_0$.

Proof. Let $R > 0$ and $x, z \in B(0; R/2)$. Define

$$\xi_t := X_t^x - Z_t^z \quad \text{and} \quad \alpha_t := \sigma(t, X_t^x) - \sigma(t, Z_t^z)H_t.$$

Then, by Itô's formula we have, for $t \in [0, \tau)$,

$$d(|\xi_t|) = \left\langle \frac{\xi_t}{|\xi_t|}, \alpha_t dB_t \right\rangle + \frac{1}{2|\xi_t|} \left(\text{tr}(\alpha_t \alpha_t^T) - \frac{|\alpha_t^T \xi_t|^2}{|\xi_t|^2} \right) dt, \quad (4-8)$$

where $\text{tr}(A)$ is the trace of the matrix A . Now we follow the argument in [Lindvall and Rogers 1986, Section 3]. Since

$$\begin{aligned} \alpha_t &= \sigma(t, X_t^x) - \sigma(t, Z_t^z) + \frac{2\xi_t \otimes (\sigma(t, Z_t^z)^{-1}\xi_t)}{|\sigma(t, Z_t^z)^{-1}\xi_t|^2} \\ &= \sigma(t, X_t^x) - \sigma(t, Z_t^z) + \frac{2\xi_t \xi_t^T (\sigma(t, Z_t^z)^{-1})^T}{|\sigma(t, Z_t^z)^{-1}\xi_t|^2}, \end{aligned}$$

it holds that

$$\operatorname{tr}(\alpha_t \alpha_t^T) - \frac{|\alpha_t^T \xi_t|^2}{|\xi_t|^2} = \operatorname{tr}([\sigma(t, X_t^x) - \sigma(t, Z_t^z)][\sigma(t, X_t^x) - \sigma(t, Z_t^z)]^T) - \frac{[[\sigma(t, X_t^x) - \sigma(t, Z_t^z)]^T \xi_t]^2}{|\xi_t|^2}.$$

Hence, in view of (2-1), there exists a positive constant γ_1 depending on d and Λ such that

$$\left| \operatorname{tr}(\alpha_t \alpha_t^T) - \frac{|\alpha_t^T \xi_t|^2}{|\xi_t|^2} \right| \leq \gamma_1 \rho_R(|\xi_t|) \quad \text{for } t \in [0, \tau) \text{ such that } X_t^x, Z_t^z \in B(0; R). \quad (4-9)$$

On the other hand, following the argument in [Lindvall and Rogers 1986, Section 3], we have a positive constant γ_2 depending on d and Λ such that

$$\frac{|\alpha_t^T \xi_t|}{|\xi_t|} \geq \gamma_2^{-1} \quad \text{for } t \in [0, \tau) \text{ such that } |\sigma(t, X_t^x) - \sigma(t, Z_t^z)| \leq 2\Lambda^{-1}. \quad (4-10)$$

Note that if $\rho_R(|\xi_t|) \leq 2\Lambda^{-1}$ and $X_t^x, Z_t^z \in B(0; R)$, then $|\sigma(t, X_t^x) - \sigma(t, Z_t^z)| \leq 2\Lambda^{-1}$. Let $\gamma := \gamma_1 \vee \gamma_2$. Define stopping times τ_n by $\tau_n := \inf\{t > 0 : |X_t^x - Z_t^z| \leq 1/n\}$ for $n \in \mathbb{N}$. For a given $\varepsilon > 0$, let

$$\begin{aligned} \tilde{\tau} &:= \tau \wedge \inf \left\{ t \in [0, \infty) : \rho_R(|\xi_t|) > \frac{\varepsilon}{2\gamma^3} \wedge 2\Lambda^{-1}, X_t^x \notin B(0; R) \text{ or } Z_t^z \notin B(0; R) \right\}, \\ \tilde{\tau}_n &:= \tau_n \wedge \inf \left\{ t \in [0, \infty) : \rho_R(|\xi_t|) > \frac{\varepsilon}{2\gamma^3} \wedge 2\Lambda^{-1}, X_t^x \notin B(0; R) \text{ or } Z_t^z \notin B(0; R) \right\} \end{aligned}$$

for $n \in \mathbb{N}$. Then, it holds that $\tilde{\tau}_n \uparrow \tilde{\tau}$ almost surely as $n \rightarrow \infty$. By Itô's formula, (4-8), (4-9) and (4-10), we have for $t \in [0, \infty)$ that

$$\begin{aligned} E[|\xi_{t \wedge \tilde{\tau}_n}|^{1-\varepsilon}] &= |x - z|^{1-\varepsilon} + (1-\varepsilon)E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-\varepsilon} \frac{1}{2|\xi_s|} \left(\operatorname{tr}(\alpha_s \alpha_s^T) - \frac{|\alpha_s^T \xi_s|^2}{|\xi_s|^2} \right) ds \right] \\ &\quad - \frac{\varepsilon(1-\varepsilon)}{2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} \frac{|\alpha_s^T \xi_s|^2}{|\xi_s|^2} ds \right] \\ &\leq |x - z|^{1-\varepsilon} + \frac{(1-\varepsilon)\gamma}{2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} \rho_R(|\xi_s|) ds \right] - \frac{\varepsilon(1-\varepsilon)}{2\gamma^2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} ds \right] \\ &\leq |x - z|^{1-\varepsilon} + \frac{\varepsilon(1-\varepsilon)}{4\gamma^2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} ds \right] - \frac{\varepsilon(1-\varepsilon)}{2\gamma^2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} ds \right] \\ &\leq |x - z|^{1-\varepsilon} - \frac{\varepsilon(1-\varepsilon)}{4\gamma^2} E \left[\int_0^{t \wedge \tilde{\tau}_n} |\xi_s|^{-1-\varepsilon} ds \right] \\ &\leq |x - z|^{1-\varepsilon} - \frac{\varepsilon(1-\varepsilon)}{2^{3+\varepsilon}\gamma^2 R^{1+\varepsilon}} E[t \wedge \tilde{\tau}_n]. \end{aligned}$$

Hence, it holds that

$$E[t \wedge \tilde{\tau}] \leq C|x - z|^{1-\varepsilon} \quad \text{for } t \in [0, \infty), \quad (4-11)$$

where C is a constant depending on d, ε, R and Λ .

Now we consider the estimate of the expectation of τ by using that of $\tilde{\tau}$. To simplify the notation, let

$$\delta_0 := \frac{1}{3} \rho_R^{-1} \left(\frac{\varepsilon}{2\gamma^3} \wedge 2\Lambda^{-1} \right).$$

Since

$$|\xi_t| > 3\delta_0 \text{ implies } |X_t^x - x| > \delta_0, |Z_t^z - z| > \delta_0, \text{ or } |x - z| > \delta_0,$$

$$X_t^x \notin B(0; R) \text{ or } Z_t^z \notin B(0; R) \text{ implies } |X_t^x - x| > \frac{R}{2} \text{ or } |Z_t^z - z| > \frac{R}{2},$$

we have, for $x, z \in B(0; R/2)$ such that $|x - z| \leq \delta_0$,

$$P(\tau \geq t) \leq P(\tilde{\tau} \geq t) + P\left(\sup_{s \in [0, t]} |X_s^x - x| > \delta_0 \wedge \frac{R}{2}\right) + P\left(\sup_{s \in [0, t]} |Z_s^z - z| > \delta_0 \wedge \frac{R}{2}\right). \quad (4-12)$$

Let $\eta = x$ or z . By Chebyshev's inequality and Burkholder's inequality we have

$$\begin{aligned} P\left(\sup_{s \in [0, t]} |X_s^\eta - \eta| > \delta_0 \wedge \frac{R}{2}\right) &\leq \left(\delta_0 \wedge \frac{R}{2}\right)^{-2/\varepsilon} E\left[\sup_{s \in [0, t]} |X_s^\eta - \eta|^{2/\varepsilon}\right] \\ &\leq \left(\delta_0 \wedge \frac{R}{2}\right)^{-2/\varepsilon} E\left[\sup_{s \in [0, t]} \left|\int_0^s \sigma(u, X_u^\eta) dB_u\right|^{2/\varepsilon}\right] \\ &\leq \left(\delta_0 \wedge \frac{R}{2}\right)^{-2/\varepsilon} CE\left[\left(\sum_{i,j=1}^d \int_0^t \sigma_{ij}(u, X_u^\eta) \sigma_{ji}(u, X_u^\eta) du\right)^{1/\varepsilon}\right] \\ &\leq d^{1/\varepsilon} \left(\delta_0 \wedge \frac{R}{2}\right)^{-2/\varepsilon} C\Lambda^{1/\varepsilon} t^{1/\varepsilon}, \end{aligned}$$

where C is a constant depending on ε . Hence, there exists a constant C depending on $d, \varepsilon, R, \rho_R$ and Λ such that

$$P\left(\sup_{s \in [0, t]} |X_s^\eta - \eta| > \delta_0 \wedge \frac{R}{2}\right) \leq C|x - z| \quad (4-13)$$

for $\eta = x, z$ and $t \in [0, |x - z|^\varepsilon]$. By (4-11), (4-12) and (4-13) we have, for $x, z \in B(0; R/2)$ such that $|x - z| \leq \delta_0$, and $t \in [0, |x - z|^\varepsilon]$,

$$\begin{aligned} E[t \wedge \tau] &\leq \int_0^t P(\tau \geq s) ds \\ &\leq \int_0^t P(\tilde{\tau} \geq s) ds + t \left[P\left(\sup_{s \in [0, t]} |X_s^x - x| > \delta_0 \wedge \frac{R}{2}\right) + P\left(\sup_{s \in [0, t]} |Z_s^z - z| > \delta_0 \wedge \frac{R}{2}\right) \right] \\ &\leq C(1+t)|x - z|^{1-\varepsilon}, \end{aligned}$$

where C is a constant depending on $d, \varepsilon, R, \rho_R$ and Λ . Therefore, we obtain

$$E[t \wedge \tau] \leq C(1+t)|x - z|^{1-\varepsilon} \quad (4-14)$$

for $x, z \in B(0; R/2)$ such that $|x - z| \leq \delta_0$ and $t \in [0, |x - z|^\varepsilon]$. By using Chebyshev's inequality, we calculate $E[t \wedge \tau]$ as

$$\begin{aligned} E[t \wedge \tau] &= \int_0^{|x-z|^\varepsilon} P(\tau \geq s) ds + \int_{|x-z|^\varepsilon}^t P(\tau \geq s) ds \leq E[|x - z|^\varepsilon \wedge \tau] + tP(\tau \geq |x - z|^\varepsilon) \\ &\leq E[|x - z|^\varepsilon \wedge \tau] + \frac{t}{|x - z|^\varepsilon} E[\tau \wedge |x - z|^\varepsilon] \\ &\leq (1 + t|x - z|^{-\varepsilon})E[|x - z|^\varepsilon \wedge \tau]. \end{aligned}$$

Thus, applying (4-14) with $t = |x - z|^\varepsilon$ and choosing another small ε , we obtain (4-7) for all $t \in [0, \infty)$. \square

Lemma 4.3. *For $R > 0$ and sufficiently small $\varepsilon > 0$, there exist positive constants C and c_0 depending on $d, \varepsilon, C_G^+, R, \rho_R$ and Λ such that*

$$\begin{aligned} p^X(0, x; t, y)E^{X_t^x=y}[t \wedge \tau] &\leq Ct^{-d/2}(1+t^2)|x - z|^{1-\varepsilon}, \\ p^X(0, z; t, y)E^{Z_t^z=y}[t \wedge \tau] &\leq Ct^{-d/2}(1+t^2)|x - z|^{1-\varepsilon} \end{aligned}$$

for $t \in (0, \infty)$, $x, z \in B(0; R/2)$ such that $|x - z| \leq c_0$, and $y \in \mathbb{R}^d$.

Proof. It holds that

$$E^{X_t^x=y}[t \wedge \tau] = E^{X_t^x=y}[(t \wedge \tau)\mathbb{1}_{[0, t/2]}(\tau)] + E^{X_t^x=y}[(t \wedge \tau)\mathbb{1}_{[t/2, \infty)}(\tau)]. \quad (4-15)$$

By (2-11) and (1-6), we have

$$p^X(0, x; t, y)E^{X_t^x=y}[(t \wedge \tau)\mathbb{1}_{[0, t/2]}(\tau)] = E\left[(t \wedge \tau)\mathbb{1}_{[0, t/2]}(\tau) p^X\left(\frac{t}{2}, X_{t/2}^x; t, y\right)\right] \leq 2^{d/2}C_G^+ t^{-d/2} E[t \wedge \tau].$$

Hence, in view of Lemma 4.2, there exists positive constants C and c_0 depending on $d, \varepsilon, C_G^+, R, \rho_R$ and Λ such that

$$p^X(0, x; t, y)E^{X_t^x=y}[(t \wedge \tau)\mathbb{1}_{[0, t/2]}(\tau)] \leq Ct^{-d/2}(1+t^2)|x - z|^{1-\varepsilon} \quad (4-16)$$

for $x, z \in B(0; R/2)$ such that $|x - z| \leq c_0$ and $y \in \mathbb{R}^d$.

On the other hand, by (2-11) and (1-6), we have

$$\begin{aligned} p^X(0, x; t, y)E^{X_t^x=y}[(t \wedge \tau)\mathbb{1}_{[t/2, \infty)}(\tau)] &\leq tp^X(0, x; t, y)P^{X_t^x=y}\left(\tau > \frac{t}{2}\right) \\ &= t \int_{\mathbb{R}^d} p^X\left(\frac{t}{2}, z; t, y\right)P\left(\tau > \frac{t}{2}, X_{t/2}^x \in dz\right) \\ &\leq 2^{d/2}C_G^+ t^{-d/2+1}P\left(\tau > \frac{t}{2}\right). \end{aligned}$$

Hence, by applying Chebyshev's inequality we have

$$p^X(0, x; t, y) E^{X_t^x=y}[(t \wedge \tau)_{[t/2, \infty)}(\tau)] \leq C t^{-d/2} E[t \wedge \tau],$$

where C is a constant depending on d and C_G^+ . Thus, Lemma 4.2 implies that

$$p^X(0, x; t, y) E^{X_t^x=y}[(t \wedge \tau)_{[t/2, \infty)}(\tau)] \leq C t^{-d/2} (1+t^2) |x-z|^{1-\varepsilon} \quad (4-17)$$

for $x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, where C and c_0 are positive constants depending on $d, \varepsilon, C_G^+, R, \rho_R$ and Λ . Therefore, we obtain the assertion for x by (4-15), (4-16) and (4-17). Similar argument yields the assertion for z . \square

Lemma 4.4. *For $q \geq 1, R > 0$ and sufficiently small $\varepsilon > 0$, there exist positive constants C and c_0 depending on $q, d, \varepsilon, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$, such that*

$$\begin{aligned} E \left[\sup_{s \in [0, t]} |\mathcal{E}(0, \tau \wedge s; X^x) - 1|^q \right] &\leq C e^{Ct} |x-z|^{2/(q\sqrt{2})-\varepsilon}, \\ E \left[\sup_{s \in [0, t]} |\mathcal{E}(0, \tau \wedge s; Z^z) - 1|^q \right] &\leq C e^{Ct} |x-z|^{2/(q\sqrt{2})-\varepsilon} \end{aligned}$$

for $t \in [0, \infty), x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, and $y \in \mathbb{R}^d$.

Proof. By (2-10) we have

$$\begin{aligned} &E \left[\sup_{v \in [0, \tau \wedge t]} |\mathcal{E}(0, v; X^x) - 1|^q \right] \\ &= E \left[\sup_{v \in [0, \tau \wedge t]} \left| \int_0^v \mathcal{E}(0, u; X^x) \langle b_\sigma(u, X_u^x), dB_u \rangle + \int_0^v \mathcal{E}(0, u; X^x) c(u, X_u^x) du \right|^q \right] \\ &\leq C E \left[\sup_{v \in [0, \tau \wedge t]} \left| \int_0^v \mathcal{E}(0, u; X^x) \langle b_\sigma(u, X_u^x), dB_u \rangle \right|^q \right] + C E \left[\sup_{v \in [0, \tau \wedge t]} \left| \int_0^v \mathcal{E}(0, u; X^x) c(u, X_u^x) du \right|^q \right], \end{aligned}$$

where C is a constant depending on q . The terms of the right-hand side of this inequality are dominated as follows: By Burkholder's inequality and Hölder's inequality we have

$$\begin{aligned} &E \left[\sup_{v \in [0, \tau \wedge t]} \left| \int_0^v \mathcal{E}(0, u; X^x) \langle b_\sigma(u, X_u^x), dB_u \rangle \right|^q \right] \\ &\leq C E \left[\left(\int_0^{\tau \wedge t} \mathcal{E}(0, u; X^x)^2 |b_\sigma(u, X_u^x)|^2 du \right)^{q/2} \right] \\ &\leq C \Lambda^q \|b\|_\infty^q t^{1-1/(q/2) \vee 1} E \left[\int_0^{\tau \wedge t} \mathcal{E}(0, u; X^x)^{2[(q/2) \vee 1]} du \right]^{1/[(q/2) \vee 1]} \\ &\leq C \Lambda^q \|b\|_\infty^q t^{1-2/[q\sqrt{2}]} E[\tau \wedge t]^{2(1-\varepsilon)/(q\sqrt{2})} E \left[\left(\int_0^t \mathcal{E}(0, u; X^x)^{(q\sqrt{2})/\varepsilon} du \right)^{2\varepsilon/(q\sqrt{2})} \right], \end{aligned}$$

where C is a constant depending on q , and by Hölder's inequality we have

$$\begin{aligned} E \left[\sup_{v \in [0, \tau \wedge t]} \left| \int_0^v \mathcal{E}(0, u; X^x) c(u, X_u^x) du \right|^q \right] &\leq \|c\|_\infty^q t^{1-1/q} E \left[\int_0^{\tau \wedge t} \mathcal{E}(0, u; X^x)^q du \right] \\ &\leq \|c\|_\infty^q t^{1-1/q} E[\tau \wedge t]^{1-\varepsilon} E \left[\int_0^t \mathcal{E}(0, u; X^x)^{q/\varepsilon} du \right]^\varepsilon. \end{aligned}$$

Thus, applying by Lemmas 3.3 and 4.2 to these inequalities and choosing another small ε , we obtain

$$E \left[\sup_{v \in [0, \tau \wedge t]} |\mathcal{E}(0, v; X^x) - 1|^q \right] \leq C e^{Ct} |x - z|^{2/(q\sqrt{2})-\varepsilon},$$

where C is a constant depending on $q, d, \varepsilon, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Similar argument yields the same estimate for Z^z . \square

Now we start the proof of Proposition 4.1. Let $t \in (0, \infty)$, $x, z \in B(0; R/2)$ such that $x \neq z$, $y \in \mathbb{R}^d$ and $s \in (t/2, t)$. Recall that X^z and Z^z have the same law. By (2-11) and (2-9) we have

$$\begin{aligned} &\left| p^X(0, x; t, y) E^{X_t^x=y} \left[\mathcal{E}(0, s; X^x); \tau \leq \frac{t}{2} \right] - p^X(0, z; t, y) E^{Z_t^z=y} \left[\mathcal{E}(0, s; Z^z); \tau \leq \frac{t}{2} \right] \right| \\ &= \left| E \left[\mathcal{E}(0, s; X^x) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] - E \left[\mathcal{E}(0, s; Z^z) p^X(s, Z_s^z; t, y); \tau \leq \frac{t}{2} \right] \right| \\ &\leq E \left[\mathcal{E}(0, s; Z^z) |p^X(s, X_s^x; t, y) - p^X(s, Z_s^z; t, y)|; \tau \leq \frac{t}{2} \right] \\ &\quad + E \left[|\mathcal{E}(0, \tau \wedge s; X^x) - \mathcal{E}(0, \tau \wedge s; Z^z)| \mathcal{E}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \\ &\quad + E \left[\mathcal{E}(0, \tau \wedge s; X^x) |\mathcal{E}(\tau \wedge s, s; X^x) - \mathcal{E}(\tau \wedge s, s; Z^z)| p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \end{aligned}$$

Noting that

$$X_s^x = Z_s^z \quad \text{for } s \geq \tau,$$

we obtain

$$\begin{aligned} &\left| p^X(0, x; t, y) E^{X_t^x=y} \left[\mathcal{E}(0, s; X^x); \tau \leq \frac{t}{2} \right] - p^X(0, z; t, y) E^{Z_t^z=y} \left[\mathcal{E}(0, s; Z^z); \tau \leq \frac{t}{2} \right] \right| \\ &\leq E \left[|\mathcal{E}(0, \tau \wedge s; X^x) - \mathcal{E}(0, \tau \wedge s; Z^z)| \mathcal{E}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right]. \quad (4-18) \end{aligned}$$

By the triangle inequality and Hölder's inequality, we obtain

$$\begin{aligned}
& E \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - \mathfrak{G}(0, \tau \wedge s; Z^z)| \mathfrak{G}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \\
& \leq E \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - 1| \mathfrak{G}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \\
& \quad + E \left[|\mathfrak{G}(0, \tau \wedge s; Z^z) - 1| \mathfrak{G}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \\
& \leq \left(E \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - 1|^{2/(2-\varepsilon)} p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right. \\
& \quad \left. + E \left[|\mathfrak{G}(0, \tau \wedge s; Z^z) - 1|^{2/(2-\varepsilon)} p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right) \\
& \quad \times E \left[\mathfrak{G}(\tau \wedge s, s; Z^z)^{2/\varepsilon} p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right]^{\varepsilon/2}.
\end{aligned}$$

Hence, by (2-11) and (1-6), we have

$$\begin{aligned}
& E \left[\mathfrak{G}(0, \tau \wedge s; X^x) - \mathfrak{G}(0, \tau \wedge s; Z^z) | \mathfrak{G}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \\
& \leq \left(E^{X_t^x=y} \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - 1|^{2/(2-\varepsilon)}; \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right. \\
& \quad \left. + E^{X_t^x=y} \left[|\mathfrak{G}(0, \tau \wedge s; Z^z) - 1|^{2/(2-\varepsilon)}; \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right) \\
& \quad \times p^X(0, x; t, y) E^{X_t^x=y} \left[\mathfrak{G}(\tau \wedge s, s; Z^z)^{2/\varepsilon}; \tau \leq \frac{t}{2} \right]^{\varepsilon/2} \\
& \leq \left(E \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - 1|^{2/(2-\varepsilon)} p^X(t/2, X_{t/2}^x; t, y); \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right. \\
& \quad \left. + E \left[|\mathfrak{G}(0, \tau \wedge s; Z^z) - 1|^{2/(2-\varepsilon)} p^X(t/2, X_{t/2}^z; t, y); \tau \leq \frac{t}{2} \right]^{1-\varepsilon/2} \right) \\
& \quad \times \left(p^X(0, x; t, y) E^{X_t^x=y} \left[\mathfrak{G}(\tau \wedge s, s; Z^z)^{2/\varepsilon}; \tau \leq \frac{t}{2} \right] \right)^{\varepsilon/2} \\
& \leq (C_G^+)^{1-\varepsilon/2} t^{-d/2+d\varepsilon/4} (p^X(0, x; t, y) E^{X_t^x=y} [\mathfrak{G}(\tau \wedge s, s; Z^z)^{2/\varepsilon}])^{\varepsilon/2} \\
& \quad \times (E[|\mathfrak{G}(0, \tau \wedge s; X^x) - 1|^{2/(2-\varepsilon)}]^{1-\varepsilon/2} + E[|\mathfrak{G}(0, \tau \wedge s; Z^z) - 1|^{2/(2-\varepsilon)}]^{1-\varepsilon/2}).
\end{aligned}$$

Applying Lemmas 3.3 and 4.4 to this inequality, we obtain

$$E \left[|\mathfrak{G}(0, \tau \wedge s; X^x) - \mathfrak{G}(0, \tau \wedge s; Z^z)| \mathfrak{G}(\tau \wedge s, s; Z^z) p^X(s, X_s^x; t, y); \tau \leq \frac{t}{2} \right] \leq C t^{-d/2} e^{Ct} |x-z|^{1-\varepsilon} \quad (4-19)$$

for $x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, where C and c_0 are constants depending on $d, \varepsilon, C_G^+, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$.

Hölder's inequality and Chebyshev's inequality imply

$$\begin{aligned} E^{X_t^x=y} \left[\mathcal{E}(0, s; X^x); \tau \geq \frac{t}{2} \right] &\leq P^{X_t^x=y} \left(\tau \geq \frac{t}{2} \right)^{1-\varepsilon/2} E^{X_t^x=y} [\mathcal{E}(0, s; X^x)^{2/\varepsilon}]^{\varepsilon/2} \\ &\leq \frac{2^{1-\varepsilon/2}}{t^{1-\varepsilon/2}} E^{X_t^x=y} [\tau \wedge t]^{1-\varepsilon/2} E^{X_t^x=y} [\mathcal{E}(0, s; X^x)^{2/\varepsilon}]^{\varepsilon/2}. \end{aligned}$$

Hence, by Lemmas 3.5 and 4.3, we obtain

$$p^X(0, x; t, y) E^{X_t^x=y} \left[\mathcal{E}(0, s; X^x); \tau \geq \frac{t}{2} \right] \leq C t^{-d/2-1} e^{Ct} |x-z|^{1-\varepsilon} \quad (4-20)$$

for $x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, where C and c_0 are constants depending on $d, \varepsilon, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Similarly we have

$$p^X(0, z; t, y) E^{Z_t^z=y} \left[\mathcal{E}(0, s; Z^z); \tau \geq \frac{t}{2} \right] \leq C t^{-d/2-1} e^{Ct} |x-z|^{1-\varepsilon} \quad (4-21)$$

for $x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, where C and c_0 are constants depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Thus, (2-8), (4-18), (4-19), (4-20) and (4-21) imply

$$|p(0, x; t, y) - p(0, z; t, y)| \leq C t^{-d/2-1+\varepsilon/2} e^{Ct} |x-z|^{1-\varepsilon}$$

for $t \in (0, \infty)$, $x, z \in B(0; R/2)$ such that $|x-z| \leq c_0$, and $y \in \mathbb{R}^d$, with constants C and c_0 depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. By (1-6) we can remove the restriction on $|x-z|$, and, therefore, we obtain Proposition 4.1.

5. The case of general a (proof of the main theorem)

Let $a^{(n)}(t, x) = (a_{ij}^{(n)}(t, x))$ be symmetric $d \times d$ -matrix-valued bounded measurable functions on $[0, \infty) \times \mathbb{R}^d$ which converge to $a(t, x)$ for each $(t, x) \in [0, \infty) \times \mathbb{R}^d$ and satisfy (1-1), (1-3) and (1-4). Consider the parabolic partial differential equation

$$\begin{cases} \frac{\partial}{\partial t} u(t, x) = \frac{1}{2} \sum_{i,j=1}^d a_{ij}^{(n)}(t, x) \frac{\partial^2}{\partial x_i \partial x_j} u(t, x) + \sum_{i=1}^d b_i(t, x) \frac{\partial}{\partial x_i} u(t, x) + c(t, x) u(t, x), \\ u(0, x) = f(x). \end{cases} \quad (5-1)$$

Denote the fundamental solution to (5-1) by $p^{(n)}(s, x; t, y)$. From (1-6) and Proposition 3.6 we have positive constants γ_1, γ_2, C_1 and C_2 depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that

$$\frac{C_1 e^{-C_1(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_1 |x-y|^2}{t-s}\right) \leq p^{(n)}(s, x; t, y) \leq \frac{C_2 e^{C_2(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_2 |x-y|^2}{t-s}\right) \quad (5-2)$$

for $s, t \in [0, \infty)$ such that $s < t$, $x, y \in \mathbb{R}^d$ and $n \in \mathbb{N}$.

It is known that local Hölder continuity of the fundamental solution follows, with index and constant depending only on the constants appearing in the Gaussian estimate; see [Stroock 1988]. This fact and (5-2)

imply that the Arzelà–Ascoli theorem is applicable to $p^{(n)}$. Moreover, in view of Proposition 4.1, there exists a constant C depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that

$$|p^{(n)}(0, x; t, y) - p^{(n)}(0, z; t, y)| \leq Ct^{-d/2-1} e^{Ct} |x - z|^{1-\varepsilon}$$

for $t \in (0, \infty)$, $y \in \mathbb{R}^d$ and $x, z \in B(0; R/2)$. Hence, there exists a continuous function $p^{(\infty)}(0, \cdot; \cdot, \cdot)$ on $\mathbb{R}^d \times (0, \infty) \times \mathbb{R}^d$ such that

$$\lim_{n \rightarrow \infty} \sup_{|x| \leq R/2} |p^{(n)}(0, x; t, y) - p^{(\infty)}(0, x; t, y)| = 0, \quad (5-3)$$

$$|p^{(\infty)}(0, x; t, y) - p^{(\infty)}(0, z; t, y)| \leq Ct^{-d/2-1} e^{Ct} |x - z|^{1-\varepsilon}, \quad x, z \in B(0; R/2), \quad (5-4)$$

for $t \in (0, \infty)$ and $y \in \mathbb{R}^d$, where C is a constant depending on $d, \varepsilon, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, R, \rho_R, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$. Moreover, we have positive constants C_1, C_2, γ_1 and γ_2 depending on $d, \gamma_G^-, \gamma_G^+, C_G^-, C_G^+, m, M, \theta, \Lambda, \|b\|_\infty$ and $\|c\|_\infty$ such that

$$\frac{C_1 e^{-C_1(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_1 |x-y|^2}{t-s}\right) \leq p^{(\infty)}(s, x; t, y) \leq \frac{C_2 e^{C_2(t-s)}}{(t-s)^{\frac{d}{2}}} \exp\left(-\frac{\gamma_2 |x-y|^2}{t-s}\right)$$

for $s, t \in [0, \infty)$ such that $s < t$, and $x, y \in \mathbb{R}^d$. To prove Theorem 1.1, we show that $p^{(\infty)}(0, \cdot; \cdot, \cdot)$ coincides with the fundamental solution $p(0, \cdot; \cdot, \cdot)$ of the original parabolic partial differential equation (1-2). Let $\phi, \psi \in C_0^\infty(\mathbb{R}^d)$, and set

$$P_t^{(n)} g(x) := \int_{\mathbb{R}^d} g(y) p^{(n)}(0, x; t, y) dy \quad \text{for } g \in C_b(\mathbb{R}^d),$$

$$L_t^{(n)} := \frac{1}{2} \sum_{i,j=1}^d a_{ij}^{(n)}(t, x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^d b_i(t, x) \frac{\partial}{\partial x_i} + c(t, x).$$

Noting that $p^{(n)}(s, x; t, y)$ is smooth in (s, x, t, y) , we have $P_t^{(n)} L_t^{(n)} \phi(x) = (\partial/\partial t) P_t^{(n)} \phi(x)$. Hence,

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \phi(y) p^{(n)}(0, x; t, y) dy \right) \psi(x) dx - \int_{\mathbb{R}^d} \phi(x) \psi(x) dx \\ &= \int_{\mathbb{R}^d} [P_t^{(n)} \phi(x)] \psi(x) dx - \int_{\mathbb{R}^d} \phi(x) \psi(x) dx \\ &= \int_0^t \int_{\mathbb{R}^d} \left[\frac{\partial}{\partial s} P_s^{(n)} \phi(x) \right] \psi(x) dx ds \\ &= \int_0^t \int_{\mathbb{R}^d} [P_s^{(n)} L_s^{(n)} \phi(x)] \psi(x) dx ds \\ &= \int_0^t \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \left[\frac{1}{2} \sum_{i,j=1}^d a_{ij}^{(n)}(s, y) \frac{\partial^2}{\partial y_i \partial y_j} + \sum_{i=1}^d b_i(s, y) \frac{\partial}{\partial y_i} + c(s, y) \right] \phi(y) p^{(n)}(0, x; s, y) dy \right) \psi(x) dx ds. \end{aligned}$$

Taking the limit as $n \rightarrow \infty$, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \phi(y) p^{(\infty)}(0, x; t, y) dy \right) \psi(x) dx - \int_{\mathbb{R}^d} \phi(x) \psi(x) dx \\ &= \int_0^t \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \left[\frac{1}{2} \sum_{i,j=1}^d a_{ij}(s, y) \frac{\partial^2}{\partial y_i \partial y_j} + \sum_{i=1}^d b_i(s, y) \frac{\partial}{\partial y_i} + c(s, y) \right] \phi(y) p^{(\infty)}(0, x; s, y) dy \right) \psi(x) dx ds. \end{aligned}$$

This equality implies that $p^{(\infty)}(0, x; t, y)$ is also the fundamental solution to the parabolic partial differential equation (1-2). Since the weak solution to (1-2) is unique, $p^{(\infty)}(0, x; t, y)$ coincides with $p(0, x; t, y)$. Therefore, we obtain Theorem 1.1.

Acknowledgments

The author is grateful to Professor Felix Otto for finding a mistake and giving the author the information on the Schauder estimate. The author also thank to the anonymous referee for careful reading, indicating mistakes in the previous version and giving the author information on references. This work was supported by JSPS KAKENHI grant number 25800054.

References

- [Aronson 1967] D. G. Aronson, “Bounds for the fundamental solution of a parabolic equation”, *Bull. Amer. Math. Soc.* **73** (1967), 890–896. MR 36 #534 Zbl 0153.42002
- [Bogachev et al. 2005] V. I. Bogachev, M. Röckner, and S. V. Shaposhnikov, “Глобальная регулярность и оценки решений параболических уравнений”, *Teor. Veroyatn. Primen.* **50:4** (2005), 652–674. Translated as “Global regularity and bounds for solutions of parabolic equations for probability measures” in *Theory Probab. Appl.* **50:4** (2006), 561–581. MR 2008f:60073 Zbl 1203.60095
- [Bogachev et al. 2009] V. I. Bogachev, N. V. Krylov, and M. Röckner, “Эллиптические и параболические уравнения для мер”, *Uspekhi Mat. Nauk* **64:6(390)** (2009), 5–116. Translated as “Elliptic and parabolic equations for measures” in *Russian Math. Surveys* **64:6** (2009), 973–1078. MR 2011c:35592 Zbl 1194.35481
- [Chen and Kumagai 2003] Z.-Q. Chen and T. Kumagai, “Heat kernel estimates for stable-like processes on d -sets”, *Stochastic Process. Appl.* **108:1** (2003), 27–62. MR 2005d:60135 Zbl 1075.60556
- [Cranston 1991] M. Cranston, “Gradient estimates on manifolds using coupling”, *J. Funct. Anal.* **99:1** (1991), 110–124. MR 93a:58175 Zbl 0770.58038
- [Escauriaza 2000] L. Escauriaza, “Bounds for the fundamental solution of elliptic and parabolic equations in nondivergence form”, *Comm. Partial Differential Equations* **25:5-6** (2000), 821–845. MR 2001i:35009 Zbl 0946.35004
- [Fabes and Kenig 1981] E. B. Fabes and C. E. Kenig, “Examples of singular parabolic measures and singular transition probability densities”, *Duke Math. J.* **48:4** (1981), 845–856. MR 86j:35081 Zbl 0482.35021
- [Friedman 1964] A. Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, Englewood Cliffs, NJ, 1964. MR 31 #6062 Zbl 0144.34903
- [Giorgi 1957] E. De Giorgi, “Sulla differenziabilità e l’analiticità delle estremali degli integrali multipli regolari”, *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat.* (3) **3** (1957), 25–43. Reprinted, pp. 167–184 in *Selected papers*, edited by G. Dal Maso et al., Springer, New York, 2006; translated as “On the differentiability and the analyticity of extremals of regular multiple integrals”, *ibid.*, 149–166. MR 20 #172 Zbl 0084.31901
- [Ikeda and Watanabe 1989] N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*, 2nd ed., North-Holland Mathematical Library **24**, North-Holland, Amsterdam, 1989. MR 90m:60069 Zbl 0684.60040

- [Karatzas and Shreve 1991] I. Karatzas and S. E. Shreve, *Brownian motion and stochastic calculus*, 2nd ed., Graduate Texts in Mathematics **113**, Springer, New York, 1991. MR 92h:60127 Zbl 0734.60060
- [Karrmann 2001] S. Karrmann, “Gaussian estimates for second-order operators with unbounded coefficients”, *J. Math. Anal. Appl.* **258**:1 (2001), 320–348. MR 2002d:35085 Zbl 0983.35053
- [Krylov 1996] N. V. Krylov, *Lectures on elliptic and parabolic equations in Hölder spaces*, Graduate Studies in Mathematics **12**, American Mathematical Society, Providence, RI, 1996. MR 97i:35001 Zbl 0865.35001
- [Kusuoka and Stroock 1985] S. Kusuoka and D. W. Stroock, “Applications of the Malliavin calculus, II”, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **32**:1 (1985), 1–76. MR 86k:60100b Zbl 0568.60059
- [Ladyženskaja et al. 1967] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural’ceva, *Линейные и квазилинейные уравнения параболического типа*, Nauka, Moscow, 1967. Translated as *Linear and quasilinear equations of parabolic type*, Transl. Math. Monogr. **23**, American Mathematical Society, Providence, RI, 1968. MR 39 #3159b Zbl 0164.12302
- [Lax and Milgram 1954] P. D. Lax and A. N. Milgram, “Parabolic equations”, pp. 167–190 in *Contributions to the theory of partial differential equations*, edited by L. Bers et al., Annals of Mathematics Studies **33**, Princeton University Press, 1954. MR 16,709b Zbl 0058.08703
- [Lindvall and Rogers 1986] T. Lindvall and L. C. G. Rogers, “Coupling of multidimensional diffusions by reflection”, *Ann. Probab.* **14**:3 (1986), 860–872. MR 88b:60179 Zbl 0593.60076
- [Metafune et al. 2009] G. Metafune, D. Pallara, and A. Rhandi, “Global properties of transition probabilities of singular diffusions”, *Teor. Veroyatn. Primen.* **54**:1 (2009), 116–148. MR 2011m:35144 Zbl 1206.60072
- [Nash 1958] J. Nash, “Continuity of solutions of parabolic and elliptic equations”, *Amer. J. Math.* **80** (1958), 931–954. MR 20 #6592 Zbl 0096.06902
- [Porper and Eidelman 1984] F. O. Porper and S. D. Eidelman, “Двусторонние оценки фундаментальных решений параболических уравнений второго порядка и некоторые их приложения”, *Uspekhi Mat. Nauk* **39**:3(237) (1984), 107–156. Translated as “Two-sided estimates of fundamental solutions of second-order parabolic equations, and some applications” in *Russian Math. Surveys* **39**:3 (1984), 119–178. MR 86b:35078 Zbl 0582.35052
- [Porper and Eidelman 1992] F. O. Porper and S. D. Eidelman, “Свойства решений параболических уравнений второго порядка с младшими членами”, *Trudy Moskov. Mat. Obshch.* **54** (1992), 118–159. Translated as “Properties of solutions of second-order parabolic equations with lower-order terms” in *Trans. Moscow Math. Soc.* **54** (1993), 101–137. MR 95b:35084 Zbl 0783.35023
- [Revuz and Yor 1999] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, 3rd ed., Grundlehren der Mathematischen Wissenschaften **293**, Springer, Berlin, 1999. MR 2000h:60050 Zbl 0917.60006
- [Stroock 1988] D. W. Stroock, “Diffusion semigroups corresponding to uniformly elliptic divergence form operators”, pp. 316–347 in *Séminaire de Probabilités XXII*, edited by J. Azéma et al., Lecture Notes in Math. **1321**, Springer, Berlin, 1988. MR 90b:35071 Zbl 0651.47031
- [Stroock and Varadhan 1979] D. W. Stroock and S. R. S. Varadhan, *Multidimensional diffusion processes*, Grundlehren der Mathematischen Wissenschaften **233**, Springer, Berlin, 1979. MR 81f:60108 Zbl 0426.60069

Received 16 Oct 2013. Revised 6 Nov 2014. Accepted 21 Dec 2014.

SEIICHIRO KUSUOKA: kusuoka@math.tohoku.ac.jp

Graduate School of Science, Tohoku University, 6-3 Aramaki Aza-Aoba, Aoba-ku, Sendai 980-8578, Japan

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 1 2015

Hölder continuity and bounds for fundamental solutions to nondivergence form parabolic equations SEIICHIRO KUSUOKA	1
Eigenvalue distribution of optimal transportation BO'AZ B. KLARTAG and ALEXANDER V. KOLESNIKOV	33
Nonlocal self-improving properties TUOMO KUUSI, GIUSEPPE MINGIONE and YANNICK SIRE	57
Symbol calculus for operators of layer potential type on Lipschitz surfaces with VMO normals, and related pseudodifferential operator calculus STEVE HOFMANN, MARIUS MITREA and MICHAEL E. TAYLOR	115
Criteria for Hankel operators to be sign-definite DIMITRI R. YAFAEV	183
Nodal sets and growth exponents of Laplace eigenfunctions on surfaces GUILLAUME ROY-FORTIN	223



2157-5045(2015)8:1;1-D