

ANALYSIS & PDE

Volume 8

No. 2

2015

JONATHAN ZINSL AND DANIEL MATTHES

**EXPONENTIAL CONVERGENCE TO EQUILIBRIUM
IN A COUPLED GRADIENT FLOW SYSTEM MODELING
CHEMOTAXIS**

EXPONENTIAL CONVERGENCE TO EQUILIBRIUM IN A COUPLED GRADIENT FLOW SYSTEM MODELING CHEMOTAXIS

JONATHAN ZINSL AND DANIEL MATTHES

We study a system of two coupled nonlinear parabolic equations. It constitutes a variant of the Keller–Segel model for chemotaxis; i.e., it models the behavior of a population of bacteria that interact by means of a signaling substance. We assume an external confinement for the bacteria and a nonlinear dependency of the chemotactic drift on the signaling substance concentration.

We perform an analysis of existence and long-time behavior of solutions based on the underlying gradient flow structure of the system. The result is that, for a wide class of initial conditions, weak solutions exist globally in time and converge exponentially fast to the unique stationary state under suitable assumptions on the convexity of the confinement and the strength of the coupling.

1. Introduction

1A. The equations and their variational structure. This paper is concerned with existence and long-time behavior of weak nonnegative solutions to the initial value problem

$$\partial_t u(t, x) = \operatorname{div}(u(t, x)D[u(t, x) + W(x) + \varepsilon\phi(v(t, x))]), \quad (1)$$

$$\partial_t v(t, x) = \Delta v(t, x) - \kappa v(t, x) - \varepsilon u(t, x)\phi'(v(t, x)), \quad (2)$$

$$u(0, x) = u_0(x) \geq 0, \quad v(0, x) = v_0(x) \geq 0, \quad (3)$$

where the sought functions u and v are defined for $(t, x) \in [0, \infty) \times \mathbb{R}^3$. Below, we comment in detail on the origin of (1)–(2) from mathematical biology. In brief, u is the spatial density of bacteria that interact with each other by means of a signaling substance of local concentration v .

In (1)–(2), ε and κ are given positive constants; we are mainly concerned with the case where the *coupling strength* ε is sufficiently small. Strict positivity of κ is essential for our approach, as explained below. The *response function* $\phi \in C^2([0, \infty))$ is assumed to be convex and strictly decreasing, with

$$0 < -\phi'(w) \leq -\phi'(0) < \infty, \quad 0 \leq \phi''(w) \leq \overline{\phi''} < \infty \quad \text{for all } w \geq 0 \quad (4)$$

This research was supported by the German Research Foundation (DFG) Collaborative Research Center SFB-TR 109.

MSC2010: primary 35K45; secondary 35A15, 35B40, 35D30, 35Q92.

Keywords: gradient flow, Wasserstein metric, chemotaxis.

for an appropriate constant $\overline{\phi}'' \geq 0$, the paradigmatic examples being

$$\phi(w) = -w \quad (\text{classical Keller–Segel model}), \quad (5)$$

$$\phi(w) = -\log(1+w) \quad (\text{weak saturation effect}), \quad (6)$$

$$\phi(w) = \frac{1}{1+w} \quad (\text{strong saturation effect}). \quad (7)$$

The *external potential* $W \in C^2(\mathbb{R}^3)$ is assumed to grow quadratically: it has globally bounded second-order partial derivatives and is uniformly convex with a constant $\lambda_0 > 0$, that is,

$$D^2W(x) \geq \lambda_0 \mathbb{1} \quad \text{for all } x \in \mathbb{R}^3 \text{ in the sense of symmetric matrices.} \quad (8)$$

Without loss of generality, we may assume that $W \geq 0$.

Equations (1)–(2) possess a variational structure. Formally, they can be written as a gradient flow of the entropy functional

$$\mathcal{H}(u, v) := \int_{\mathbb{R}^3} \left(\frac{1}{2}u^2 + uW + \frac{1}{2}|Dv|^2 + \frac{1}{2}\kappa v^2 + \varepsilon u\phi(v) \right) dx$$

with respect to a metric **dist**, defined on the space $X := \mathcal{P}_2(\mathbb{R}^3) \times L^2_+(\mathbb{R}^3)$ by

$$\mathbf{dist}((u_1, v_1), (u_2, v_2)) := \sqrt{W_2^2(u_1, u_2) + \|v_1 - v_2\|_{L^2(\mathbb{R}^3)}^2} \quad \text{for } (u_1, v_1), (u_2, v_2) \in X. \quad (9)$$

Here W_2 is the L^2 -Wasserstein metric on the space $\mathcal{P}_2(\mathbb{R}^3)$ of probability measures on \mathbb{R}^3 with finite second moment; see Section 2A for the definition. This gradient flow structure is at the basis of our proof for global existence of weak solutions to (1)–(3), and it is also the key element for our analysis of long-time behavior. We remark that, even with this variational structure at hand, the analysis is far from trivial since \mathcal{H} is *not* convex along geodesics. Therefore, the established general theory on λ -contractive gradient flows in metric spaces [Ambrosio et al. 2008] is not directly applicable.

1B. Statement of the main results. In the first part of this work, we show that a weak solution to (1)–(2) can be obtained by means of the time-discrete implicit Euler approximation (also known as *minimizing movement* or the *JKO scheme*). More precisely, for each sufficiently small time step $\tau > 0$, let $(u_\tau^0, v_\tau^0) := (u_0, v_0)$, and then define inductively for each $n \in \mathbb{N}$

$$(u_\tau^n, v_\tau^n) \in \operatorname{argmin}_{(u, v) \in \mathcal{P}_2(\mathbb{R}^3) \times L^2(\mathbb{R}^3)} \left(\frac{1}{2\tau} \mathbf{dist}((u, v), (u_\tau^{n-1}, v_\tau^{n-1}))^2 + \mathcal{H}(u, v) \right). \quad (10)$$

We will prove in Section 4A that this construction is well defined, i.e., that a minimizer exists for every $n \in \mathbb{N}$. Further, introduce the piecewise-constant interpolation $(u_\tau, v_\tau) : \mathbb{R}_+ \rightarrow \mathcal{P}_2(\mathbb{R}^3) \times L^2(\mathbb{R}^3)$ by

$$u_\tau(t) = u_\tau^n, \quad v_\tau(t) = v_\tau^n \quad \text{for all } t \in ((n-1)\tau, n\tau]. \quad (11)$$

Our existence result — which does not require a small coupling strength ε — reads as follows.

Theorem 1.1 (existence of weak solutions to (1)–(2)). *Let $\kappa > 0$ and $\varepsilon > 0$ be given, and assume that the response function ϕ satisfies (4) and that the convex confinement potential W grows quadratically.*

Let further initial conditions $u_0 \in \mathcal{P}_2(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ and $v_0 \in W^{1,2}(\mathbb{R}^3)$ be given, with $v_0 \geq 0$, and define for each $\tau > 0$ a function (u_τ, v_τ) by means of the scheme (10) and (11). Then there is a sequence $(\tau_k)_{k \in \mathbb{N}}$ with $\tau_k \downarrow 0$ such that (u_{τ_k}, v_{τ_k}) converges to a weak solution $(u, v) : [0, \infty) \times \mathbb{R}^3 \rightarrow [0, \infty]^2$ of (1)–(3) in the following sense:

$$u_{\tau_k}(t) \rightarrow u(t) \quad \text{narrowly in } \mathcal{P}(\mathbb{R}^3), \text{ pointwise with respect to } t \in [0, T],$$

$$v_{\tau_k}(t) \rightarrow v(t) \quad \text{in } L^2(\mathbb{R}^3), \text{ uniformly with respect to } t \in [0, T],$$

$$u \in C^{1/2}([0, T], \mathcal{P}_2(\mathbb{R}^3)) \cap L^\infty([0, T], L^2(\mathbb{R}^3)) \cap L^2([0, T], W^{1,2}(\mathbb{R}^3)),$$

$$v \in C^{1/2}([0, T], L^2(\mathbb{R}^3)) \cap L^\infty([0, T], W^{1,2}(\mathbb{R}^3)) \cap L^2([0, T], W^{2,2}(\mathbb{R}^3)) \cap W^{1,2}([0, T], L^2(\mathbb{R}^3))$$

for all $T > 0$, and (u, v) satisfies

$$\partial_t u = \operatorname{div}(uD[u + W + \varepsilon\phi(v)]) \quad \text{in the sense of distributions,} \tag{12}$$

$$\partial_t v = \Delta v - \kappa v - \varepsilon u \phi'(v) \quad \text{a.e. in } (0, +\infty) \times \mathbb{R}^3, \tag{13}$$

$$u(0) = u_0, \quad v(0) = v_0. \tag{14}$$

The convergence of (u_{τ_k}, v_{τ_k}) is actually much stronger; see Proposition 4.7 for details.

The key a priori estimate yielding sufficient compactness of (u_τ, v_τ) follows from a dissipation estimate, which formally amounts to

$$-\frac{d}{dt} \int_{\mathbb{R}^3} \left(u \log u + \frac{1}{2} |Dv|^2 + \frac{1}{2} \kappa v^2 \right) dx \geq \frac{1}{2} \int_{\mathbb{R}^3} (|Du|^2 + (\Delta v - \kappa v)^2) dx - C \left(\|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{W^{1,2}(\mathbb{R}^3)}^2 + \|\Delta W\|_{L^\infty(\mathbb{R}^3)} \right).$$

Related existence results have been proved recently for similar systems of equations, using essentially the same technique, in [Laurençot and Matioc 2013; Blanchet and Laurençot 2013; Zinsl 2014; Blanchet et al. 2014]. Therefore, we keep the technical details to a minimum. Note that our method of proof yields neither contractivity of the flow nor uniqueness of weak solutions due to the lack of convexity of the entropy functional.

Our main result is the following on the long-time behavior of solutions:

Theorem 1.2 (exponential convergence to equilibrium). *Let κ, ϕ and W be as in Theorem 1.1 above. Then there are constants $\bar{\varepsilon} > 0, L > 0$ and $C > 0$ such that, for every $\varepsilon \in (0, \bar{\varepsilon})$ and with $\Lambda_\varepsilon := \min(\kappa, \lambda_0) - L\varepsilon$, the following is true.*

Let initial conditions $u_0 \in \mathcal{P}_2(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$ and $v_0 \in W^{1,2}(\mathbb{R}^3)$ be given, with $v_0 \geq 0$, and assume in addition that $v_0 \in L^{6/5}(\mathbb{R}^3)$. Let further (u, v) be a weak solution to (1)–(3) obtained as a limit of the scheme (10) and (11). Then (u, v) converges to the unique nonnegative stationary solution $(u_\infty, v_\infty) \in (\mathcal{P}_2 \cap L^2)(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3)$ of (1)–(2) exponentially fast with rate Λ_ε in the sense

$$\begin{aligned} & \mathbf{W}_2(u(t, \cdot), u_\infty) + \|u(t, \cdot) - u_\infty\|_{L^2(\mathbb{R}^3)} + \|v(t, \cdot) - v_\infty\|_{W^{1,2}(\mathbb{R}^3)} \\ & \leq C(1 + \|v_0\|_{L^{6/5}(\mathbb{R}^3)}) (\mathcal{H}(u_0, v_0) - \mathcal{H}(u_\infty, v_\infty) + 1) e^{-\Lambda_\varepsilon t} \quad \text{for all } t \geq 0. \end{aligned} \tag{15}$$

We give a brief and formal indication of the main idea for the proof of Theorem 1.2. First, we decompose the entropy in the form

$$\mathcal{H}(u, v) - \mathcal{H}(u_\infty, v_\infty) = \mathcal{L}_u(u) + \mathcal{L}_v(v) + \varepsilon \mathcal{L}_*(u, v), \quad (16)$$

where

$$\begin{aligned} \mathcal{L}_u(u) &:= \int_{\mathbb{R}^3} \left(\frac{1}{2}(u^2 - u_\infty^2) + [W + \varepsilon \phi(v_\infty)](u - u_\infty) \right) dx, \\ \mathcal{L}_v(v) &:= \int_{\mathbb{R}^3} \frac{1}{2} (|\mathbf{D}(v - v_\infty)|^2 + \kappa(v - v_\infty)^2) dx, \\ \mathcal{L}_*(u, v) &:= \int_{\mathbb{R}^3} (u[\phi(v) - \phi(v_\infty)] - u_\infty \phi'(v_\infty)[v - v_\infty]) dx. \end{aligned}$$

There, \mathcal{L}_u and \mathcal{L}_v are λ_ε -convex and κ -convex functionals — with $\lambda_\varepsilon = \lambda_0 - C\varepsilon > 0$ — in $(\mathcal{P}_2, \mathbf{W}_2)$ and in L^2 , respectively, which are minimized by the stationary solution (u_∞, v_∞) ; the functional \mathcal{L}_* has no useful convexity properties. On a very formal level — pretending that \mathcal{L}_u , \mathcal{L}_v and \mathcal{L}_* are smooth functionals on Euclidean spaces and denoting their “gradients” by ∇_u and ∇_v — the dissipation of the *principal entropy* $\mathcal{L}_u + \mathcal{L}_v$ amounts to

$$\begin{aligned} -\frac{d}{dt}(\mathcal{L}_u + \mathcal{L}_v) &= \nabla_u \mathcal{L}_u \cdot \nabla_u \mathcal{H} + \nabla_v \mathcal{L}_v \cdot \nabla_v \mathcal{H} \\ &= \|\nabla_u \mathcal{L}_u\|^2 + \|\nabla_v \mathcal{L}_v\|^2 + \varepsilon \nabla_u \mathcal{L}_u \cdot \nabla_u \mathcal{L}_* + \varepsilon \nabla_v \mathcal{L}_v \cdot \nabla_v \mathcal{L}_* \\ &\geq (1 - \varepsilon)\|\nabla_u \mathcal{L}_u\|^2 + (1 - \varepsilon)\|\nabla_v \mathcal{L}_v\|^2 - \frac{1}{2}\varepsilon(\|\nabla_u \mathcal{L}_*\|^2 + \|\nabla_v \mathcal{L}_*\|^2). \end{aligned} \quad (17)$$

By convexity of \mathcal{L}_u and \mathcal{L}_v , one has the inequalities

$$\|\nabla_u \mathcal{L}_u\|^2 \geq 2\lambda_\varepsilon \mathcal{L}_u, \quad \|\nabla_v \mathcal{L}_v\|^2 \geq 2\kappa \mathcal{L}_v,$$

and so we are *almost* in the situation to apply the Gronwall estimate to (17) and conclude convergence to equilibrium with an exponential rate of $\min(\lambda_\varepsilon, \kappa) > 0$. However, it remains to estimate the terms involving the “gradients” of \mathcal{L}_* . This is relatively straightforward if the entropy $\mathcal{H}(u, v)$ is sufficiently close to its minimal value $\mathcal{H}(u_\infty, v_\infty)$ but is rather difficult for (u, v) far from equilibrium. Moreover, rigorous estimates have to be carried out on the time-discrete level (with subsequent passage to continuous time) since our notion of solution is too weak to carry out the respective estimates in continuous time.

In the language of gradient flows, our results can be interpreted as follows. For $\varepsilon = 0$, the functional \mathcal{H} is Λ_0 -convex along geodesics in (X, \mathbf{dist}) with $\Lambda_0 = \min(\lambda_0, \kappa) > 0$. Consequently, there is an associated Λ_0 -contractive gradient flow defined on all of X that satisfies (1)–(2), and in particular, all solutions converge with the exponential rate Λ_0 to the unique equilibrium. For every $\varepsilon > 0$, the convexity of \mathcal{H} is lost; see [Zinsl 2014] for a discussion of (non)convexity in a similar situation. By Theorem 1.1, Equations (1)–(2) still define a continuous flow on the proper domain of \mathcal{H} , which is $X \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$. Further, we show that, on the (almost exhaustive) subset of those (u, v) with $v \in L^{6/5}(\mathbb{R}^3)$, this flow still converges to an equilibrium with an exponential rate $\Lambda_\varepsilon \geq \Lambda_0 - L\varepsilon > 0$ (Theorem 1.2) for all $\varepsilon > 0$ sufficiently small.

From this point of view, our result is perturbative: the uncoupled system ($\varepsilon = 0$) exhibiting a strictly contractive flow is perturbed in such a way that the perturbed system ($\varepsilon > 0$) still yields exponential convergence towards the unique equilibrium — with a slightly slower convergence rate than in the unperturbed case. For this approach to work, we obviously need to require $\kappa > 0$ and $\lambda_0 > 0$. On the other hand, this theorem is stronger than a usual perturbation result: the crucial point is that we do *not* require the initial condition (u_0, v_0) to be close to equilibrium, apart from the rather harmless additional hypothesis that $v_0 \in L^{6/5}(\mathbb{R}^3)$, which could be weakened further with additional technical effort.

Our result on global existence of weak solutions (Theorem 1.1), however, can be generalized to the case of $\kappa = 0$ and no convexity assumption on the confinement potential (see, e.g., [Zinsl 2014]). Further generalization of Theorem 1.1 to the case of nonlinear, but nonquadratic, diffusion can be achieved with similar techniques as in [Zinsl 2014]. However, in our analysis of the long-time behavior, the right entropy dissipation estimates are not at hand to the best of our knowledge when dealing with nonquadratic diffusion. To keep technicalities to a minimum, we consider the quadratic case throughout this work.

We expect that similar results can be proved for system (1)–(2) on a bounded domain $\Omega \subset \mathbb{R}^3$, even with vanishing confinement $W \equiv 0$. The role of the confinement will then be played by Poincaré’s inequality. Our setup with a convex confinement on \mathbb{R}^3 fits much more naturally into the variational framework.

1C. Modeling background. The system of equations (1)–(2) is a variant of the so-called *Keller–Segel model for chemotaxis* describing the time-dependent distribution of biological cells or microorganisms in response to gradients of chemical substances (*chemotaxis*). The original model — corresponding to the linear response function from (5) — has been developed by Keller and Segel [1970] to describe slime mold aggregation. However, chemotactic processes occur in many (and highly different) biological systems; for the biological details, we refer to the book by Eisenbach [Eisenbach 2004]. For example, many bacteria like *Escherichia coli* possess *flagella* driven by small motors that respond to gradients of signaling molecules in the environment. Chemotaxis also plays an important role in embryonal development, e.g., in the development of blood vessels (*angiogenesis*), which is also a crucial step in tumor growth. Starting from the basic Keller–Segel model, many different model extensions are conceivable. A broad range of those is summarized in the review articles by Hillen and Painter [2009] and Horstmann [2003]. Details on the modeling aspects can be found, e.g., in the books by Murray [2003] and Perthame [2007].

In the model (1)–(2) under consideration here, u is the time-dependent spatial density of the cells and v is the time-dependent concentration of the signaling substance. Equation (1) describes the temporal change in cell density due to the directed drift of cells towards regions with a higher concentration of the substance and due to undirected diffusion. Equation (2) models the degradation of the signaling substance as well as its production by the cells. Two special aspects are included in this particular model: nonlinear diffusion, i.e., the use of a nonconstant, u -dependent mobility coefficient for the diffusive motion of the bacteria, and signal-dependent chemotactic sensitivity, i.e., the use of the — in general nonlinear — response $\phi(v)$ instead of the concentration v itself. For the first, we refer to [Hillen and Painter 2009] and the references therein for biological motivation. The second is motivated by the fact that the conversion of an external signal into a reaction of the considered microorganism (*signal transduction*) often occurs by binding and dissociation of molecules to certain receptors. The movement of the cell is then caused

by gradients in the number of receptors occupied by signaling molecules rather than by concentration gradients of signaling molecules themselves. For growing concentrations, the number of bound receptors can exhibit a saturation such that the gradient vanishes. In [Hillen and Painter 2009; Segel 1977; Lapidus and Schiller 1976], this was incorporated into the model by the *chemotactic sensitivity function*

$$\phi'(v) = -\frac{1}{(1+v)^2},$$

which fits into our model with the response function ϕ defined in (7). Finally, an external background potential W is included in order to generate a spatial confinement of the bacterial population.

For the dynamics of the signaling substance, we assume linear diffusion according to Fick's laws and degradation with a constant, exponential rate κ . The nonnegative term $-\varepsilon u \phi'(v)$ models the production of signaling substance by the microorganisms; here it is taken into account that the cells might be less active in producing additional substance the higher its local concentration already is. This is consistent with the models presented in [Horstmann 2003, §6]. Often, the two processes of chemotactic response and production of the chemoattractant are modeled with different response functions. Here we require them to be equal (or scalar multiples of each other) to ensure that the system has a gradient flow structure.

By definition, u and v are density/concentration functions and thus should be nonnegative. Note that it is part of our results that, for given nonnegative initial data (of sufficient regularity), there exists a weak solution that is nonnegative for all times $t > 0$. On a formal level, nonnegativity is an easy consequence of the particular structure of the system (1)–(2).

1D. Relation to the existing literature. The rapidly growing mathematical literature about the Keller–Segel model and its manifold variants is devoted primarily to the dichotomy *global existence versus finite-time blow-up* of (weak, possibly measure-valued) solutions, but the long-time behavior of global solutions has been intensively investigated as well.

Global existence and blow-up in the classical parabolic-parabolic Keller–Segel model, which is (1)–(2) with $\phi(v) = -v$, $W \equiv 0$ and *linear* diffusion, has been thoroughly studied by Calvez and Corrias [2008] in space dimension $d = 2$ and by Corrias and Perthame [2008] in higher space dimensions $d > 2$; see also [Biler et al. 2011; Kozono and Sugiyama 2009; Mizoguchi 2013; Nagai et al. 2003; Senba and Suzuki 2006; Sugiyama and Kunii 2006; Yamada 2011]. Recently, in [Carrapatoso and Mischler 2014], uniqueness and long-time behavior of solutions of the parabolic-parabolic Keller–Segel system was studied by means of a perturbation of the parabolic-elliptic framework.

Variants with nonlinear diffusion and drift have been studied for instance by Sugiyama [2006; 2007]. The results from [Sugiyama 2006] already indicate that, in the model (1)–(2) under consideration, blow-up *never* occurs, in accordance with Theorem 1.1.

In the aforementioned works [Corrias and Perthame 2008; Nagai et al. 2003], the intermediate asymptotics of global solutions have been studied as well: it is proved that the cell density converges to the self-similar solution of the heat equation at an algebraic rate, i.e., in a properly scaled frame, the density approaches a Gaussian. See also [Di Francesco and Rosado 2008] for an extension of this result to a model with size-exclusion. Similar asymptotic behavior has been proved in models with

nonlinear, homogeneous diffusion, e.g., by Luckhaus and Sugiyama [2006; 2007]. There, the intermediate asymptotics are that of a porous medium equation with the respective homogeneous nonlinearity; i.e., the rescaled bacterial density converges to a Barenblatt profile. These intermediate asymptotics are — at least morally — related to Theorem 1.2: recall that algebraic convergence to self-similarity for the unconfined porous medium equation is comparable to exponential convergence to an equilibrium for the equation with λ -convex confinement.

The fully parabolic model (1)–(2) with a nonlinear response ϕ has not been rigorously analyzed so far, with the following exception: in her thesis, Post [1999] proves existence and uniqueness of solutions to a similar system with linear diffusion and vanishing confinement on a bounded domain by nonvariational methods and obtains convergence to the (spatially homogeneous) stationary solution from compactness arguments. Variants of the classical parabolic-parabolic or parabolic-elliptic Keller–Segel models with a nonlinear chemotactic sensitivity coefficient have also been studied, e.g., in [Nagai and Senba 1998; Winkler 2010].

Despite the fact that energy/entropy methods are one of the key tools for the analysis of Keller–Segel-type systems, the use of genuine variational methods is relatively recent in that context. The variational machinery of gradient flows in transportation metrics, originally developed by Jordan, Kinderlehrer and Otto [Jordan et al. 1998] for the linear Fokker–Planck equation, has been applied to a variety of dynamical systems: mainly to nonlinear diffusion [Carrillo and Toscani 2000; Otto 2001; Carrillo et al. 2006a; Agueh 2008] but also to aggregation [Carrillo et al. 2003; 2006b; 2011] and fourth-order equations [Giacomelli and Otto 2001; Gianazza et al. 2009; Matthes et al. 2009].

For the parabolic-elliptic Keller–Segel model, which can be reduced to a single nonlocal scalar equation, the variational framework was established by Blanchet, Calvez and Carrillo [Blanchet et al. 2008], who represented the evolution as a gradient flow of an appropriate potential with respect to the Wasserstein distance and constructed a numerical scheme on these grounds. Later, the gradient flow structure has been used for a detailed analysis of the basin of attraction in the critical mass case by Blanchet, Carlen and Carrillo [Blanchet et al. 2012] (see also, e.g., [Blanchet et al. 2009; Calvez and Carrillo 2012; López-Gómez et al. 2013]).

The parabolic-parabolic Keller–Segel model was somewhat harder to fit into the framework since the two equations are (formally) gradient flows with respect to *different* metrics: Wasserstein and L^2 . The first rigorous analytical result on grounds of this structure was given by Blanchet and Laurençot [2013], where they constructed weak solutions for the system with critical exponents of nonlinear diffusion. Later their result was generalized to other, noncritical parameter situations in [Zinsl 2014]. In the recent work by Blanchet et al. [2014], a similar strategy was used to reprove the result in [Calvez and Corrias 2008] about the global existence of weak solutions to the classical Keller–Segel system in two spatial dimensions. To the best of our knowledge, our approach taken here to prove long-time asymptotics by gradient flow techniques in a combined Wasserstein- L^2 -metric is novel.

1E. Plan of the paper. First, we summarize common facts and definitions on gradient flows in metric spaces in Section 2. After that, various properties of the entropy functional are derived in Section 3. On grounds of these properties, we construct a weak solution by means of the minimizing movement

scheme in Section 4, proving Theorem 1.1. Existence, uniqueness and regularity of stationary solutions are studied in Section 5, and the proof of Theorem 1.2 is completed in Section 6.

2. Preliminaries

In this section, we recall the relevant definitions and properties related to gradient flows in metric spaces (X, \mathbf{d}) , following [Ambrosio et al. 2008]. The two metric spaces of interest here are $L^2(\mathbb{R}^d)$ with the metric induced by the norm and the space $\mathcal{P}_2(\mathbb{R}^d)$ of probability measures, endowed with the L^2 -Wasserstein distance \mathbf{W}_2 . We also discuss the compound metric **dist** from (9).

2A. Spaces of probability measures and the Wasserstein distance. We denote by $\mathcal{P}(\mathbb{R}^d)$ the space of probability measures on \mathbb{R}^d . By abuse of notation, we will frequently identify absolutely continuous measures $\mu \in \mathcal{P}(\mathbb{R}^d)$ with their respective (Lebesgue) density functions $u = \mathrm{d}\mu/\mathrm{d}x \in L^1_+(\mathbb{R}^d)$, where $L^p_+(\mathbb{R}^d)$ for $p \geq 1$ denotes the subspace of those $L^p(\mathbb{R}^d)$ functions with nonnegative values.

A sequence $(\mu_n)_{n \in \mathbb{N}}$ in $\mathcal{P}(\mathbb{R}^d)$ is called *narrowly convergent* to its limit $\mu \in \mathcal{P}(\mathbb{R}^d)$ if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \varphi(x) \mathrm{d}\mu_n(x) = \int_{\mathbb{R}^d} \varphi(x) \mathrm{d}\mu(x)$$

for every bounded, continuous function $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$. By $\mathcal{P}_2(\mathbb{R}^d)$, we denote the subspace of those $\mu \in \mathcal{P}(\mathbb{R}^d)$ with finite second moment

$$m_2(\mu) := \int_{\mathbb{R}^d} |x|^2 \mathrm{d}\mu(x).$$

$\mathcal{P}_2(\mathbb{R}^d)$ turns into a complete metric space when endowed with the L^2 -Wasserstein distance \mathbf{W}_2 . We do not recall the general definition of \mathbf{W}_2 here. Instead, since we are concerned with absolutely continuous measures in $\mathcal{P}_2(\mathbb{R}^d)$ only, we remark that, for probability density functions $u_1, u_2 \in L^1_+(\mathbb{R}^d)$, the Wasserstein distance is given by the infimum

$$\mathbf{W}_2^2(u_1, u_2) = \inf \left\{ \int_{\mathbb{R}^d} |t(x) - x|^2 u_1(x) \mathrm{d}x \mid t : \mathbb{R}^d \rightarrow \mathbb{R}^d \text{ Borel-measurable and } t \# u_1 = u_2 \right\},$$

where $t \# u$ denotes the *push-forward* with respect to the map t . In this case, the infimum above is attained by an optimal transport map [Villani 2003, Theorem 2.32]. Convergence in the metric space $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$ is equivalent to narrow convergence and convergence of the second moment. Further, \mathbf{W}_2 is lower semicontinuous in both components with respect to narrow convergence.

2B. Geodesic convexity and gradient flows in metric spaces. A functional $\mathfrak{A} : X \rightarrow \mathbb{R} \cup \{\infty\}$ defined on the metric space (X, \mathbf{d}) is called *geodesically λ -convex* for some $\lambda \in \mathbb{R}$ if, for every $w_0, w_1 \in X$ and $s \in [0, 1]$, one has

$$\mathfrak{A}(w_s) \leq (1-s)\mathfrak{A}(w_0) + s\mathfrak{A}(w_1) - \frac{1}{2}\lambda s(1-s)\mathbf{d}^2(w_0, w_1),$$

where $w_s : [0, 1] \rightarrow X$, $s \mapsto w_s$ is a *geodesic connecting* w_0 and w_1 .

On $L^2(\mathbb{R}^d)$, the (unique up to rescaling) geodesic from w_0 to w_1 is given by linear interpolation, i.e., $w_s = (1 - s)w_0 + sw_1$. Hence, a functional $\mathfrak{F} : L^2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\infty\}$ of the form

$$\mathfrak{F}(w) = \int_{\mathbb{R}^d} f(w(x), Dw(x), D^2w(x)) \, dx$$

with a given continuous function $f : \mathbb{R} \times \mathbb{R}^d \times \mathbb{R}^{d \times d} \rightarrow \mathbb{R}$ is λ -convex if and only if $(z, p, Q) \mapsto f(z, p, Q) - \frac{1}{2}\lambda z^2$ is (jointly) convex.

In the metric space $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$, geodesic λ -convexity is a much more complicated concept. We recall two important classes of λ -convex functionals (see, e.g., [Ambrosio et al. 2008, Chapter 9.3; Villani 2003, Theorem 5.15]).

Theorem 2.1 (criteria for geodesic convexity in $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$). *The following statements are true:*

- (a) *Let a function $h \in C^0([0, \infty))$ be given, and define the functional \mathfrak{H} on $\mathcal{P}_2(\mathbb{R}^d)$ by $\mathfrak{H}(u) := \int_{\mathbb{R}^d} h(u(x)) \, dx$ for $u \in (\mathcal{P}_2 \cap L^1)(\mathbb{R}^d)$. If $h(0) = 0$ and $r \mapsto r^d h(r^{-d})$ is convex and nonincreasing on $(0, \infty)$, then \mathfrak{H} is 0-geodesically convex and lower semicontinuous in $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$.*
- (b) *Let a function $W \in C^0(\mathbb{R}^d)$ be given, and define the functional $\mathfrak{H}(\mu) := \int_{\mathbb{R}^d} W(x) \, d\mu(x)$ for all $\mu \in \mathcal{P}_2(\mathbb{R}^d)$. If W is λ -convex (as a functional on the metric space \mathbb{R}^d with the Euclidean distance) for some $\lambda \in \mathbb{R}$, then \mathfrak{H} is λ -geodesically convex in $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$.*

Next, we introduce a notion of *gradient flow*. There are various possible characterizations. For our purposes here, we need the following very strong one:

Definition 2.2. Let $\mathfrak{A} : X \rightarrow \mathbb{R} \cup \{\infty\}$ be a lower semicontinuous functional on the metric space (X, \mathbf{d}) . A continuous semigroup $S_{(\cdot)}^{\mathfrak{A}}$ on (X, \mathbf{d}) is called κ -flow for some $\kappa \in \mathbb{R}$ if the *evolution variational inequality*

$$\frac{1}{2} \frac{d^+}{dt} \mathbf{d}^2(S_t^{\mathfrak{A}}(w), \tilde{w}) + \frac{\kappa}{2} \mathbf{d}^2(S_t^{\mathfrak{A}}(w), \tilde{w}) + \mathfrak{A}(S_t^{\mathfrak{A}}(w)) \leq \mathfrak{A}(\tilde{w}) \tag{18}$$

holds for arbitrary w and \tilde{w} in the domain of \mathfrak{A} and for all $t \geq 0$.

If $S_{(\cdot)}^{\mathfrak{A}}$ is a κ -flow for the λ -convex functional \mathfrak{A} , then $S_{(\cdot)}^{\mathfrak{A}}$ is also a *gradient flow* for \mathfrak{A} in essentially all possible interpretations of that notion. For the metric spaces $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$ and $L^2(\mathbb{R}^d)$, it can be proved that every lower semicontinuous and geodesically λ -convex functional possesses a unique κ -flow, with $\kappa := \lambda$ (see [Ambrosio et al. 2008, Theorem 11.1.4, Corollary 4.3.3], respectively).

In these metric spaces, λ -geodesic convexity with $\lambda > 0$ implies existence and uniqueness of a minimizer w_{\min} of \mathfrak{A} , for which the following holds (see, e.g., [Ambrosio et al. 2008, Lemma 2.4.8, Theorem 4.0.4]):

$$\frac{\lambda}{2} \mathbf{d}^2(w, w_{\min}) \leq \mathfrak{A}(w) - \mathfrak{A}(w_{\min}) \leq \frac{1}{2\lambda} \lim_{h \downarrow 0} \frac{\mathfrak{A}(w) - \mathfrak{A}(S_h^{\mathfrak{A}}(w))}{h}. \tag{19}$$

Remark 2.3 (formal calculation of evolution equations associated to gradient flows). In the metric spaces of interest here, one can explicitly write an evolution equation for the flow $S_{(\cdot)}^{\mathfrak{A}}$ of a sufficiently regular

functional \mathfrak{A} ; see, e.g., [Villani 2003, §8.2]. On $(\mathcal{P}_2(\mathbb{R}^d), \mathbf{W}_2)$, one has

$$\partial_t S_t^{\mathfrak{A}}(w) = \operatorname{div} \left(S_t^{\mathfrak{A}}(w) D \left(\frac{\delta \mathfrak{A}}{\delta w} (S_t^{\mathfrak{A}}(w)) \right) \right),$$

and on $L^2(\mathbb{R}^d)$, one has

$$\partial_t S_t^{\mathfrak{A}}(w) = - \frac{\delta \mathfrak{A}}{\delta w} (S_t^{\mathfrak{A}}(w)).$$

Here, $\delta \mathfrak{A} / \delta w$ stands for the usual first variation of the functional \mathfrak{A} on L^2 .

2C. The metric \mathbf{dist} . It is easily verified that $\mathbf{X} := \mathcal{P}_2(\mathbb{R}^3) \times L_+^2(\mathbb{R}^3)$ becomes a complete metric space when endowed with the compound metric \mathbf{dist} defined in (9). The topology on \mathbf{X} induced by \mathbf{dist} is that of the cartesian product. Moreover:

Lemma 2.4. *The distance \mathbf{dist} is weakly lower semicontinuous on \mathbf{X} in the following sense: if $(u_n, v_n)_{n \in \mathbb{N}}$ is a sequence in \mathbf{X} such that u_n converges to $u \in \mathcal{P}_2(\mathbb{R}^3)$ narrowly and v_n converges to $v \in L^2(\mathbb{R}^3)$ weakly in $L^2(\mathbb{R}^3)$, then*

$$\mathbf{dist}((u, v), (\tilde{u}, \tilde{v})) \leq \liminf_{n \rightarrow \infty} \mathbf{dist}((u_n, v_n), (\tilde{u}, \tilde{v}))$$

holds, for every $(\tilde{u}, \tilde{v}) \in \mathbf{X}$.

For our purposes, it suffices to discuss convexity and gradient flows for functionals $\Phi : \mathbf{X} \rightarrow \mathbb{R} \cup \{\infty\}$ of the separable form $\Phi(u, v) = \Phi_1(u) + \Phi_2(v)$. One immediately verifies:

Lemma 2.5. *Assume that Φ_1 and Φ_2 are λ -convex and lower semicontinuous functionals on the respective spaces $(\mathcal{P}_2(\mathbb{R}^3), \mathbf{W}_2)$ and $L^2(\mathbb{R}^3)$, and denote their respective gradient flows by $S_{(\cdot)}^1$ and $S_{(\cdot)}^2$. Then $\Phi : \mathbf{X} \rightarrow \mathbb{R} \cup \{\infty\}$ with $\Phi(u, v) = \Phi_1(u) + \Phi_2(v)$ is a λ -convex and lower semicontinuous functional on $(\mathbf{X}, \mathbf{dist})$, and the semigroup $S_{(\cdot)}^\Phi$ given by $S_t^\Phi(u, v) = (S_t^1(u), S_t^2(v))$ is a λ -flow for Φ .*

3. Properties of the entropy functional

Recall the definition of the metric space $(\mathbf{X}, \mathbf{dist})$. We define the *entropy functional* $\mathcal{H} : \mathbf{X} \rightarrow \mathbb{R} \cup \{\infty\}$ as follows. For all $(u, v) \in \mathbf{X} \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$, set

$$\mathcal{H}(u, v) := \int_{\mathbb{R}^3} \left(\frac{1}{2} u^2 + u W + \frac{1}{2} |\operatorname{D}v|^2 + \frac{1}{2} \kappa v^2 + \varepsilon u \phi(v) \right) dx, \quad (20)$$

which is a finite value by our assumptions on ϕ and W . For all other $(u, v) \in \mathcal{P}_2(\mathbb{R}^3) \times L^2(\mathbb{R}^3)$, we set $\mathcal{H}(u, v) = +\infty$.

Proposition 3.1 (properties of the entropy functional \mathcal{H}). *The functional \mathcal{H} defined in (20) has the following properties:*

(a) *There exist $C_0, C_1 > 0$ such that*

$$\mathcal{H}(u, v) \geq C_0 \left[\|u\|_{L^2(\mathbb{R}^3)}^2 + \mathbf{m}_2(u) + \|v\|_{W^{1,2}(\mathbb{R}^3)}^2 - C_1 \right]. \quad (21)$$

In particular, \mathcal{H} is bounded from below.

(b) \mathcal{H} is weakly lower semicontinuous in the following sense: for every sequence $(u_n, v_n)_{n \in \mathbb{N}}$ in \mathbf{X} , where $(u_n)_{n \in \mathbb{N}}$ converges narrowly to some $u \in \mathcal{P}_2(\mathbb{R}^3)$ and where $(v_n)_{n \in \mathbb{N}}$ converges weakly in $L^2(\mathbb{R}^3)$ to some $v \in L^2(\mathbb{R}^3)$, one has

$$\mathcal{H}(u, v) \leq \liminf_{n \rightarrow \infty} \mathcal{H}(u_n, v_n).$$

(c) For sufficiently small $\varepsilon > 0$, \mathcal{H} is λ' -geodesically convex for some $\lambda' > 0$ with respect to the distance induced by the norm $\|(\tilde{u}, \tilde{v})\|_{L^2 \times L^2} := \sqrt{\|\tilde{u}\|_{L^2(\mathbb{R}^3)}^2 + \|\tilde{v}\|_{L^2(\mathbb{R}^3)}^2}$.

Proof. For part (a), we observe that, due to λ_0 -convexity of W , one has $W(x) \geq \frac{1}{4}\lambda_0|x|^2 - \frac{1}{2}\lambda_0|x_{\min}|^2$, where $x_{\min} \in \mathbb{R}^3$ is the unique minimizer of W . Moreover, with convexity of ϕ , we deduce

$$\int_{\mathbb{R}^3} u\phi(v) \, dx \geq \phi(0) + \phi'(0)\|uv\|_{L^1(\mathbb{R}^3)} \geq \phi(0) + C\phi'(0)\|Dv\|_{L^2(\mathbb{R}^3)}\|u\|_{L^2(\mathbb{R}^3)}^{1/3},$$

using that $\|u\|_{L^1(\mathbb{R}^3)} = 1$ and the chain of inequalities

$$\|uv\|_{L^1(\mathbb{R}^3)} \leq \|u\|_{L^{6/5}(\mathbb{R}^3)}\|v\|_{L^6(\mathbb{R}^3)} \leq C\|Dv\|_{L^2(\mathbb{R}^3)}\|u\|_{L^1(\mathbb{R}^3)}^{2/3}\|u\|_{L^2(\mathbb{R}^3)}^{1/3}. \tag{22}$$

All in all, we arrive at

$$\begin{aligned} \mathcal{H}(u, v) \geq & \frac{1}{2}\|u\|_{L^2(\mathbb{R}^3)}^2 + \frac{1}{4}\lambda_0\mathbf{m}_2(u) - \lambda_0|x_{\min}|^2 + \frac{1}{2}\|Dv\|_{L^2(\mathbb{R}^3)}^2 + \frac{1}{2}\kappa\|v\|_{L^2(\mathbb{R}^3)}^2 \\ & - \varepsilon|\phi(0)| - \varepsilon C|\phi'(0)|\|Dv\|_{L^2(\mathbb{R}^3)}\|u\|_{L^2(\mathbb{R}^3)}^{1/3}. \end{aligned}$$

From this, the desired estimate follows by Young's inequality.

In (b), the claimed lower semicontinuity of the integral with $\varepsilon = 0$ follows from joint convexity of the map

$$\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}^3 \ni (r, z, p) \mapsto \frac{1}{2}r^2 + W(x)r + \frac{1}{2}|p|^2 + \frac{1}{2}\kappa z^2,$$

for every $x \in \mathbb{R}^3$. It thus remains to prove semicontinuity of the integral of $u\phi(v)$. Let a sequence $(u_n, v_n)_{n \in \mathbb{N}}$ with the mentioned properties be given, and assume — without loss of generality — that $\mathcal{H}(u_n, v_n) \rightarrow H < \infty$. With these prerequisites at hand, we are even able to prove the continuity of the integral of $u\phi(v)$: it follows by (21) that $(u_n)_{n \in \mathbb{N}}$ and $(v_n)_{n \in \mathbb{N}}$ are bounded sequences in $L^2(\mathbb{R}^3)$ and in $W^{1,2}(\mathbb{R}^3)$, respectively. Moreover, the sequence of second moments $(\mathbf{m}_2(u_n))_{n \in \mathbb{N}}$ is bounded. Hence, $(u_n)_{n \in \mathbb{N}}$ converges to u weakly in $L^2(\mathbb{R}^3)$, and $(v_n)_{n \in \mathbb{N}}$ converges to v weakly in $W^{1,2}(\mathbb{R}^3)$ and strongly in $L^2(B_R(0))$, for every ball $B_R(0) \subset \mathbb{R}^3$. Recalling our assumptions (4) on ϕ , we conclude that

$$|\phi(v_n) - \phi(v)|^2 \leq \phi'(0)^2|v_n - v|^2,$$

and thus, $(\phi(v_n))_{n \in \mathbb{N}}$ converges to $\phi(v)$ strongly in $L^2(B_R(0))$. We proceed by a truncation argument. Let therefore $R > 0$, and choose $\beta_R \in C^\infty(\mathbb{R}^3)$ with

$$0 \leq \beta_R \leq 1, \quad \beta_R \equiv 1 \quad \text{on } B_R(0), \quad \beta_R \equiv 0 \quad \text{on } \mathbb{R}^3 \setminus B_{2R}(0).$$

Using the triangle inequality, we see

$$\begin{aligned} & \left| \int_{\mathbb{R}^3} (u_n \phi(v_n) - u \phi(v)) \, dx \right| \\ & \leq \left| \int_{\mathbb{R}^3} \phi(v)(u_n - u) \, dx \right| + \left| \int_{\mathbb{R}^3} \beta_R u_n (\phi(v_n) - \phi(v)) \, dx \right| + \left| \int_{\mathbb{R}^3} (1 - \beta_R) u_n (\phi(v_n) - \phi(v)) \, dx \right|. \end{aligned} \quad (23)$$

Since $u_n \rightharpoonup u$ weakly on $L^2(\mathbb{R}^3)$ and $\phi(v) \in L^2(\mathbb{R}^3)$, the first term in (23) converges to zero. The same holds for the second one due to strong convergence of $\phi(v_n)$ to $\phi(v)$ on $L^2(B_{2R}(0))$ and boundedness of $\|u_n\|_{L^2(\mathbb{R}^3)}$. The third term in (23) can be estimated using (22):

$$\left| \int_{\mathbb{R}^3} (1 - \beta_R) u_n (\phi(v_n) - \phi(v)) \, dx \right| \leq \|\phi(v_n) - \phi(v)\|_{L^6(\mathbb{R}^3)} \|u_n\|_{L^{6/5}(\mathbb{R}^3 \setminus B_R(0))}.$$

Consequently,

$$\begin{aligned} \left| \int_{\mathbb{R}^3} (1 - \beta_R) u_n (\phi(v_n) - \phi(v)) \, dx \right| & \leq C \|\phi(v_n) - \phi(v)\|_{W^{1,2}(\mathbb{R}^3)} \|u_n\|_{L^2(\mathbb{R}^3)}^{1/3} \left(\int_{\mathbb{R}^3 \setminus B_R(0)} \frac{|x|^2}{R^2} u_n(x) \, dx \right)^{2/3} \\ & \leq C R^{-4/3} (\|\phi(v_n)\|_{W^{1,2}(\mathbb{R}^3)} + \|\phi(v)\|_{W^{1,2}(\mathbb{R}^3)}) \|u_n\|_{L^2(\mathbb{R}^3)}^{1/3} (\mathbf{m}_2(u_n))^{2/3} \\ & \leq 2\tilde{C} R^{-4/3}. \end{aligned}$$

Hence, for all $R > 0$,

$$\limsup_{n \rightarrow \infty} \left| \int_{\mathbb{R}^3} (u_n \phi(v_n) - u \phi(v)) \, dx \right| \leq 2\tilde{C} R^{-4/3},$$

proving the claim.

Finally, to prove (c), consider a geodesic $w_s = (u_s, v_s)$ with respect to the flat metric induced by $\|\cdot\|_{L^2 \times L^2}$; that is, $u_s = (1-s)u_0 + su_1$ and $v_s = (1-s)v_0 + sv_1$ for given $u_0, u_1 \in (\mathcal{P}_2 \cap L^2)(\mathbb{R}^3)$ and $v_0, v_1 \in W^{1,2}(\mathbb{R}^3)$. It then follows that

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{H}(u_s, v_s) & = \int_{\mathbb{R}^3} ((u_1 - u_0)^2 + |D(v_1 - v_0)|^2 + \kappa(v_1 - v_0)^2 \\ & \quad + 2\varepsilon \phi'(v_s)(u_1 - u_0)(v_1 - v_0) + \varepsilon u_s \phi''(v_s)(v_1 - v_0)^2) \, dx \\ & \geq \int_{\mathbb{R}^3} \begin{pmatrix} u_1 - u_0 \\ v_1 - v_0 \end{pmatrix}^\top A_s \begin{pmatrix} u_1 - u_0 \\ v_1 - v_0 \end{pmatrix} \, dx \quad \text{with } A_s := \begin{pmatrix} 1 & \varepsilon \phi'(v_s) \\ \varepsilon \phi'(v_s) & \kappa \end{pmatrix}, \end{aligned}$$

where we have used that ϕ is convex. Thus, \mathcal{H} is λ' -convex with respect to the flat distance above if $A_s \geq \lambda' \mathbb{1}$ for all $s \in [0, 1]$. Recalling that $0 < -\phi'(v_s) \leq -\phi'(0)$ by hypothesis (4), it follows from elementary linear algebra that $\varepsilon^2 \phi'(0)^2 < \kappa$ is sufficient to find a suitable $\lambda' > 0$ with $A_s \geq \lambda' \mathbb{1}$. \square

4. Existence of weak solutions

In this section, we prove Theorem 1.1 by construction of a weak solution using the minimizing movement scheme.

4A. Time discretization. Recall the discretization scheme from (10). We introduce the step size $\tau > 0$ and define the associated Yosida penalization \mathcal{H}_τ of the entropy by

$$\mathcal{H}_\tau(u, v \mid \tilde{u}, \tilde{v}) := \frac{1}{2\tau} \mathbf{dist}^2((u, v), (\tilde{u}, \tilde{v})) + \mathcal{H}(u, v) \quad (24)$$

for all $(u, v), (\tilde{u}, \tilde{v}) \in X$. Set $(u_\tau^0, v_\tau^0) := (u_0, v_0)$, and define the sequence $(u_\tau^n, v_\tau^n)_{n \in \mathbb{N}}$ inductively by choosing

$$(u_\tau^n, v_\tau^n) \in \underset{(u,v) \in X}{\operatorname{argmin}} \mathcal{H}_\tau(u, v \mid u_\tau^{n-1}, v_\tau^{n-1}). \quad (25)$$

Lemma 4.1. *For every $(\tilde{u}, \tilde{v}) \in X$, there exists at least one minimizer $(u, v) \in X$ of $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$ that satisfies $u \in L^2(\mathbb{R}^3)$ and $v \in W^{1,2}(\mathbb{R}^3)$.*

Proof. The proof is an application of the direct methods from the calculus of variations to the functional $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$.

First, observe that, on any given sublevel S of $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$, both $\mathbf{W}_2(u, \tilde{u})$ and $\|v\|_{L^2(\mathbb{R}^3)}$ are uniformly bounded. The first bound implies that also the second moment $\mathbf{m}_2(u)$ is uniformly bounded, and thus, the u -components in S belong to a subset of $\mathcal{P}_2(\mathbb{R}^3)$ that is relatively compact in the narrow topology by Prokhorov's theorem. The other bound implies via Alaoglu's theorem that the v -components belong to a weakly relatively compact subset of $L^2(\mathbb{R}^3)$.

Next, recall the properties of \mathcal{H} and of \mathbf{dist} given in Proposition 3.1 and Lemma 2.4. From these, it follows that $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$ is lower semicontinuous with respect to narrow convergence in the first and L^2 -weak convergence in the second components.

Combining these properties with the fact that $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$ is bounded from below (e.g., by zero), the existence of a minimizer follows. The additional regularity is a consequence of the fact that the proper domain of \mathcal{H} is a subset of $L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3)$. \square

Given the sequence $(u_\tau^n, v_\tau^n)_{n \in \mathbb{N}}$, define the *discrete solution* $(u_\tau, v_\tau) : [0, \infty) \rightarrow X$ as in (11) by piecewise constant interpolation:

$$(u_\tau, v_\tau)(t) := (u_\tau^n, v_\tau^n) \quad \text{for } t \in ((n-1)\tau, n\tau] \text{ and } n \geq 1. \quad (26)$$

We start by recalling a collection of estimates on (u_τ, v_τ) that follows immediately from the construction by minimizing movements.

Proposition 4.2 (classical estimates). *The following hold for $T > 0$:*

$$\mathcal{H}(u_\tau^n, v_\tau^n) \leq \mathcal{H}(u_0, v_0) < \infty \quad \text{for all } n \geq 0, \quad (27)$$

$$\sum_{n=1}^{\infty} (\mathbf{W}_2^2(u_\tau^n, u_\tau^{n-1}) + \|v_\tau^n - v_\tau^{n-1}\|_{L^2(\mathbb{R}^3)}^2) \leq 2\tau(\mathcal{H}(u_0, v_0) - \inf \mathcal{H}), \quad (28)$$

$$\mathbf{W}_2(u_\tau(s), u_\tau(t)) + \|v_\tau(s) - v_\tau(t)\|_{L^2(\mathbb{R}^3)} \leq 2[2(\mathcal{H}(u_0, v_0) - \inf \mathcal{H}) \max(\tau, |t-s|)]^{1/2} \quad \text{for all } 0 \leq s, t \leq T, \quad (29)$$

the infimum $\inf \mathcal{H}$ of \mathcal{H} on X being finite.

By the well known *JKO method* [Jordan et al. 1998], we derive an approximate weak formulation satisfied by (u_τ, v_τ) . The idea is to choose test functions $\eta, \gamma \in C_c^\infty(\mathbb{R}^3)$ and perturb the minimizer (u_τ^n, v_τ^n) of the functional $\mathcal{H}_\tau(\cdot \mid u_\tau^{n-1}, v_\tau^{n-1})$ over an auxiliary time $s \geq 0$ as follows:

$$u_\tau^n \rightsquigarrow S_s^{D\eta} \# u_\tau^n, \quad v_\tau^n \rightsquigarrow v + s\gamma.$$

Here $S_{(\cdot)}^{D\eta}$ is the flow on \mathbb{R}^3 generated by the gradient vector field $D\eta$. Since the calculations are very similar to the ones performed in [Zinsl 2014], we skip the details and directly state the result.

Lemma 4.3. *For all $n \in \mathbb{N}$ and all test functions $\eta, \gamma \in C_c^\infty(\mathbb{R}^3)$ and $\psi \in C_c^\infty((0, \infty)) \cap C([0, \infty))$, the following discrete weak formulation holds:*

$$\begin{aligned} 0 = & \int_0^\infty \int_{\mathbb{R}^3} [u_\tau(t, x)\eta(x) - v_\tau(t, x)\gamma(x)] \frac{\psi(\lfloor t/\tau \rfloor \tau) - \psi(\lfloor t/\tau \rfloor \tau + \tau)}{\tau} \, dx \, dt + O(\tau) \\ & + \int_0^\infty \int_{\mathbb{R}^3} \psi(\lfloor t/\tau \rfloor \tau) \left(-\frac{1}{2} u_\tau(t, x)^2 \Delta \eta(x) + u_\tau(t, x) DW(x) \cdot D\eta(x) \right. \\ & \quad \left. + Dv_\tau(t, x) \cdot D\gamma(x) + \kappa v_\tau(t, x)\gamma(x) \right. \\ & \quad \left. + \varepsilon u_\tau(t, x)\phi'(v_\tau(t, x))[\gamma(x) + Dv_\tau(t, x) \cdot D\eta(x)] \right) \, dx \, dt. \end{aligned} \quad (30)$$

Our goal for the rest of this section is to pass to the limit $\tau \downarrow 0$ in (30) and obtain the (time-continuous) weak formulation (12)–(13).

4B. Regularity of the discrete solution. Since the discrete weak formulation (30) contains nonlinear terms with respect to u_τ and v_τ , further compactness estimates are needed to pass to the continuous time limit $\tau \rightarrow 0$. As a preparation, we state:

Lemma 4.4 (flow interchange lemma [Matthes et al. 2009, Theorem 3.2]). *Let \mathfrak{A} be a proper, lower semi-continuous and λ -geodesically convex functional on (X, \mathbf{dist}) , which is defined on $X \cap L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3)$ at least. Further, assume that $S_{(\cdot)}^{\mathfrak{A}}$ is a λ -flow for \mathfrak{A} . Then, the following holds for every $n \in \mathbb{N}$:*

$$\mathfrak{A}(u_\tau^n, v_\tau^n) + \tau D^{\mathfrak{A}}\mathcal{H}(u_\tau^n, v_\tau^n) + \frac{1}{2}\lambda \mathbf{dist}^2((u_\tau^n, v_\tau^n), (u_\tau^{n-1}, v_\tau^{n-1})) \leq \mathfrak{A}(u_\tau^{n-1}, v_\tau^{n-1}).$$

There, $D^{\mathfrak{A}}\mathcal{H}(w)$ denotes the dissipation of the entropy \mathcal{H} along $S_{(\cdot)}^{\mathfrak{A}}$, i.e.,

$$D^{\mathfrak{A}}\mathcal{H}(w) := \limsup_{h \downarrow 0} \frac{\mathcal{H}(w) - \mathcal{H}(S_h^{\mathfrak{A}}(w))}{h}.$$

The necessary additional regularity is provided by the following estimate on the minimizers of \mathcal{H}_τ :

Proposition 4.5 (additional regularity). *Let $(u, v), (\tilde{u}, \tilde{v}) \in X \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$ with $(u, v) \in \operatorname{argmin} \mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$. Denoting $\mathcal{E}(u) := \int_{\mathbb{R}^3} u \log(u) \, dx$ and $\mathcal{F}(v) := \int_{\mathbb{R}^3} (\frac{1}{2}|Dv|^2 + \frac{1}{2}\kappa v^2) \, dx$, the following estimate holds for some constant $K > 0$:*

$$\begin{aligned} & \|Du\|_{L^2(\mathbb{R}^3)}^2 + \|\Delta v - \kappa v\|_{L^2(\mathbb{R}^3)}^2 \\ & \leq K \left(\|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{W^{1,2}(\mathbb{R}^3)}^2 + \|D\tilde{u}\|_{L^\infty(\mathbb{R}^3)} + \frac{1}{\tau} (\mathcal{E}(\tilde{u}) - \mathcal{E}(u) + \mathcal{F}(\tilde{v}) - \mathcal{F}(v)) \right). \end{aligned} \quad (31)$$

Proof. The method of proof used here is based on the flow interchange lemma (Lemma 4.4). The idea is to calculate the dissipation of \mathcal{H} along the gradient flow of an auxiliary functional, namely the heat flow and the heat flow with decay, respectively.

Therefore, we recall that the functional $\mathcal{E}(u) := \int_{\mathbb{R}^3} u \log(u) \, dx$ is 0-geodesically convex on $\mathcal{P}_2(\mathbb{R}^3)$ and its gradient flow $S_{(\cdot)}^{\mathcal{E}}$ is the heat flow satisfying

$$\partial_s S_s^{\mathcal{E}}(u) = \Delta S_s^{\mathcal{E}}(u).$$

Moreover, with the evolution variational inequality (18), we deduce as in [Blanchet and Laurençot 2013; Zinsl 2014] by integration over time using that \mathcal{E} is a Lyapunov functional along $S_{(\cdot)}^{\mathcal{E}}$

$$\frac{1}{2}(\mathbf{W}_2^2(S_s^{\mathcal{E}}(u), \tilde{u}) - \mathbf{W}_2^2(u, \tilde{u})) \leq \int_0^s (\mathcal{E}(\tilde{u}) - \mathcal{E}(S_\sigma^{\mathcal{E}}(u))) \, d\sigma \leq s[\mathcal{E}(\tilde{u}) - \mathcal{E}(S_s^{\mathcal{E}}(u))]. \quad (32)$$

Analogous to that, $\mathcal{F}(v) := \int_{\mathbb{R}^3} (\frac{1}{2}|Dv|^2 + \frac{1}{2}\kappa v^2) \, dx$ is κ -geodesically convex on $L^2(\mathbb{R}^3)$ and its gradient flow $S_{(\cdot)}^{\mathcal{F}}$ is given by

$$\partial_s S_s^{\mathcal{F}}(v) = \Delta S_s^{\mathcal{F}}(v) - \kappa S_s^{\mathcal{F}}(v).$$

The application of the evolution variational inequality (18) then shows

$$\frac{1}{2}(\|S_s^{\mathcal{F}}(v) - \tilde{v}\|_{L^2(\mathbb{R}^3)}^2 - \|v - \tilde{v}\|_{L^2(\mathbb{R}^3)}^2) \leq \int_0^s (\mathcal{F}(\tilde{v}) - \mathcal{F}(S_\sigma^{\mathcal{F}}(v))) \, d\sigma \leq s[\mathcal{F}(\tilde{v}) - \mathcal{F}(S_s^{\mathcal{F}}(v))]. \quad (33)$$

Well known results of parabolic theory ensure that $(S_s^{\mathcal{E}}(u), S_s^{\mathcal{F}}(v)) \in X \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$ if $(u, v) \in X \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$. For the sake of clarity, we introduce the notation $(\mathcal{U}_s, \mathcal{V}_s) := (S_s^{\mathcal{E}}(u), S_s^{\mathcal{F}}(v))$ and calculate for $s > 0$

$$\begin{aligned} & \frac{d}{ds} \mathcal{H}(\mathcal{U}_s, \mathcal{V}_s) \\ &= \int_{\mathbb{R}^3} ([\mathcal{U}_s + W + \varepsilon \phi(\mathcal{V}_s)] \Delta \mathcal{U}_s + [-\Delta \mathcal{V}_s + \kappa \mathcal{V}_s + \varepsilon \mathcal{U}_s \phi'(\mathcal{V}_s)] [\Delta \mathcal{V}_s - \kappa \mathcal{V}_s]) \, dx \\ &= \int_{\mathbb{R}^3} (-|D\mathcal{U}_s|^2 - \mathcal{U}_s \Delta W - (\Delta \mathcal{V}_s - \kappa \mathcal{V}_s)^2 - \varepsilon \phi'(\mathcal{V}_s) D\mathcal{V}_s \cdot D\mathcal{U}_s + \varepsilon \mathcal{U}_s \phi'(\mathcal{V}_s) [\Delta \mathcal{V}_s - \kappa \mathcal{V}_s]) \, dx, \quad (34) \end{aligned}$$

where the last line follows by integration by parts. An application of Young's inequality yields

$$\begin{aligned} & \int_{\mathbb{R}^3} (-|D\mathcal{U}_s|^2 - \mathcal{U}_s \Delta W - (\Delta \mathcal{V}_s - \kappa \mathcal{V}_s)^2 - \varepsilon \phi'(\mathcal{V}_s) D\mathcal{V}_s \cdot D\mathcal{U}_s + \varepsilon \mathcal{U}_s \phi'(\mathcal{V}_s) [\Delta \mathcal{V}_s - \kappa \mathcal{V}_s]) \, dx \\ & \leq \int_{\mathbb{R}^3} (-\frac{1}{2}|D\mathcal{U}_s|^2 - \mathcal{U}_s \Delta W - \frac{1}{2}(\Delta \mathcal{V}_s - \kappa \mathcal{V}_s)^2 + \frac{1}{2}\varepsilon^2 \phi'(0)^2 (|D\mathcal{V}_s|^2 + \mathcal{U}_s^2)) \, dx. \end{aligned}$$

Exploiting the monotonicity of the L^2 norm along $S_{(\cdot)}^{\mathcal{E}}$ and of the $W^{1,2}$ norm along $S_{(\cdot)}^{\mathcal{F}}$, one gets

$$\begin{aligned} & \int_{\mathbb{R}^3} (-\frac{1}{2}|D\mathcal{U}_s|^2 - \mathcal{U}_s \Delta W - \frac{1}{2}(\Delta \mathcal{V}_s - \kappa \mathcal{V}_s)^2 + \frac{1}{2}\varepsilon^2 \phi'(0)^2 (|D\mathcal{V}_s|^2 + \mathcal{U}_s^2)) \, dx \\ & \leq -\frac{1}{2} \|D\mathcal{U}_s\|_{L^2(\mathbb{R}^3)}^2 - \frac{1}{2} \|\Delta \mathcal{V}_s - \kappa \mathcal{V}_s\|_{L^2(\mathbb{R}^3)}^2 + \|\Delta W\|_{L^\infty(\mathbb{R}^3)} + \frac{1}{2} \varepsilon^2 \phi'(0)^2 (\|Dv\|_{L^2(\mathbb{R}^3)}^2 + \|u\|_{L^2(\mathbb{R}^3)}^2). \end{aligned}$$

All in all, we have estimated the dissipation of \mathcal{H} along $S_{(\cdot)}^{\mathcal{E}}$ and $S_{(\cdot)}^{\mathcal{F}}$:

$$\frac{d}{ds} \mathcal{H}(\mathcal{U}_s, \mathcal{V}_s) \leq -\frac{1}{2}(\|\mathbf{D}\mathcal{U}_s\|_2^2 + \|\Delta\mathcal{V}_s - \kappa\mathcal{V}_s\|_2^2) + C\|u\|_2^2 + C\|v\|_{W^{1,2}(\mathbb{R}^3)}^2 + \|\Delta W\|_\infty. \quad (35)$$

As a final step of the proof of Proposition 4.5, we use the minimizing property of (u, v) . Clearly,

$$0 \leq \mathcal{H}_\tau(\mathcal{U}_s, \mathcal{V}_s \mid \tilde{u}, \tilde{v}) - \mathcal{H}_\tau(u, v \mid \tilde{u}, \tilde{v}).$$

We insert (32), (33) and (35) and obtain

$$\begin{aligned} & \frac{1}{s} \int_0^s (\|\mathbf{D}\mathcal{U}_\sigma\|_{L^2(\mathbb{R}^3)}^2 + \|\Delta\mathcal{V}_\sigma - \kappa\mathcal{V}_\sigma\|_{L^2(\mathbb{R}^3)}^2) d\sigma \\ & \leq K \left(\|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{W^{1,2}(\mathbb{R}^3)}^2 + \|\Delta W\|_{L^\infty(\mathbb{R}^3)} + \frac{1}{\tau} (\mathcal{E}(\tilde{u}) - \mathcal{E}(\mathcal{U}_s) + \mathcal{F}(\tilde{v}) - \mathcal{F}(\mathcal{V}_s)) \right), \end{aligned}$$

for some constant $K > 0$. Similar to [Blanchet and Laurençot 2013; Zinsl 2014; Blanchet 2013], passing to the lim inf as $s \rightarrow 0$ yields (31) by lower semicontinuity of norms and continuity of the entropies \mathcal{E} and \mathcal{F} along their respective gradient flows. \square

4C. Compactness estimates and passage to continuous time. The following compactness estimates in addition to the results of Proposition 4.2 are needed to pass to the limit $\tau \rightarrow 0$ in the nonlinear terms of the discrete weak formulation (30) afterwards. The method of proof is essentially the same as in [Zinsl 2014, §7]. For the sake of brevity, the details are omitted here.

Proposition 4.6 (additional a priori estimates). *Let (u_τ, v_τ) be the discrete solution obtained by the minimizing movement scheme (25). Then the following hold for $T > 0$:*

$$\mathbf{m}_2(u_\tau^n) \leq C_1 < \infty \quad \text{for all } n \leq \lfloor T/\tau \rfloor, \quad (36)$$

$$\|u_\tau^n\|_{L^2(\mathbb{R}^3)} \leq C_3 < \infty \quad \text{for all } n \geq 0, \quad (37)$$

$$\|v_\tau^n\|_{W^{1,2}(\mathbb{R}^3)} \leq C_5 < \infty \quad \text{for all } n \geq 0, \quad (38)$$

$$\int_0^T \|u_\tau(t)\|_{W^{1,2}(\mathbb{R}^3)}^2 dt \leq C_6 < \infty, \quad (39)$$

$$\int_0^T \|v_\tau(t)\|_{W^{2,2}(\mathbb{R}^3)}^2 dt \leq C_7 < \infty, \quad (40)$$

with constants $C_j > 0$ only depending on T and the initial condition (u_0, v_0) .

The estimates (36)–(38) are a consequence of those in Proposition 4.2. Also employing Proposition 4.5 yields (39)–(40).

The estimates of Propositions 4.2 and 4.6 enable us to prove the existence of the continuous-time limit of the discrete solution.

Proposition 4.7 (continuous-time limit). *Let $(\tau_k)_{k \geq 0}$ be a vanishing sequence of step sizes, i.e., $\tau_k \rightarrow 0$ as $k \rightarrow \infty$, and let $(u_{\tau_k}, v_{\tau_k})_{k \geq 0}$ be the corresponding sequence of discrete solutions obtained by the minimizing movement scheme.*

Then there exists a subsequence (nonreabeled) such that, for fixed $t \in [0, T]$, $u_{\tau_k}(t)$ converges to a limit $u(t)$ narrowly in $\mathcal{P}(\mathbb{R}^3)$ and $v_{\tau_k}(t)$ converges to a limit $v(t)$ strongly in $L^2(\mathbb{R}^3)$. The second convergence is uniform with respect to $t \in [0, T]$. The limit curves satisfy $u \in C^{1/2}([0, T], \mathcal{P}_2(\mathbb{R}^3))$ and $v \in C^{1/2}([0, T], L^2_+(\mathbb{R}^3))$. Furthermore, the following additional convergence properties hold for $k \rightarrow \infty$:

- (a) $u_{\tau_k} \rightharpoonup u$ weakly in $L^2([0, T], W^{1,2}(\mathbb{R}^3))$,
- (b) $v_{\tau_k} \rightharpoonup v$ weakly in $L^2([0, T], W^{2,2}(\mathbb{R}^3))$,
- (c) $u_{\tau_k} \rightarrow u$ strongly in $L^2([0, T], L^2(\Omega))$ for all bounded domains $\Omega \subset \mathbb{R}^3$,
- (d) $v_{\tau_k} \rightarrow v$ strongly in $L^2([0, T], W^{1,2}(\Omega))$ for all bounded domains $\Omega \subset \mathbb{R}^3$.

Now, to complete the proof of Theorem 1.1, one needs to verify that the obtained limit curve (u, v) indeed satisfies the weak formulation (12)–(13). This will be omitted here for the sake of brevity.

5. The stationary solution

In this section, we provide the characterization of a stationary state of system (1)–(2) and prove some relevant properties.

5A. Existence and uniqueness. At first, we show existence and uniqueness of the stationary solution to system (1)–(2).

Proposition 5.1. *For each sufficiently small $\varepsilon > 0$, there exists a unique minimizer $(u_\infty, v_\infty) \in X \cap (W^{1,2}(\mathbb{R}^3) \times W^{2,2}(\mathbb{R}^3))$ of \mathcal{H} , for which the following holds: (u_∞, v_∞) is a stationary solution to (1)–(2) and to the Euler–Lagrange system*

$$\Delta v_\infty - \kappa v_\infty = \varepsilon u_\infty \phi'(v_\infty), \tag{41}$$

$$u_\infty = [U_\varepsilon - W - \varepsilon \phi(v_\infty)]_+, \tag{42}$$

where $U_\varepsilon \in \mathbb{R}$ is chosen such that $\|u_\infty\|_{L^1(\mathbb{R}^3)} = 1$ and $[\cdot]_+$ denotes the positive part.

Moreover, $v_\infty \in C^0(\mathbb{R}^3)$ and there exists $V > 0$ independent of $\varepsilon > 0$ such that $\|v_\infty\|_{L^\infty(\mathbb{R}^3)} \leq V$.

Proof. We prove that \mathcal{H} possesses a unique minimizer (u_∞, v_∞) . Let a minimizing sequence $(u_n, v_n)_{n \in \mathbb{N}}$ be given such that $\lim_{n \rightarrow \infty} \mathcal{H}(u_n, v_n) = \inf \mathcal{H} > -\infty$. As the sequence $(\mathcal{H}(u_n, v_n))_{n \in \mathbb{N}}$ is bounded, we can, by the same argument as in the proof of Proposition 3.1, extract a (nonreabeled) subsequence, on which $(u_n)_{n \in \mathbb{N}}$ converges weakly in $L^2(\mathbb{R}^3)$ to some $u_\infty \in L^2_+(\mathbb{R}^3)$ and $(v_n)_{n \in \mathbb{N}}$ converges weakly in $W^{1,2}(\mathbb{R}^3)$ to some $v_\infty \in L^2_+(\mathbb{R}^3) \cap W^{1,2}(\mathbb{R}^3)$ as $n \rightarrow \infty$. By the same argument as in the proof of Proposition 3.1(b), (u_∞, v_∞) is indeed a minimizer of \mathcal{H} and hence an element of $X \cap (L^2(\mathbb{R}^3) \times W^{1,2}(\mathbb{R}^3))$.

Since $(u_\infty, v_\infty) \in \operatorname{argmin} \mathcal{H}_\tau(\cdot | u_\infty, v_\infty)$ for arbitrary $\tau > 0$, Proposition 4.5 immediately yields

$$\|u_\infty\|_{W^{1,2}(\mathbb{R}^3)}^2 + \|v_\infty\|_{W^{2,2}(\mathbb{R}^3)}^2 \leq V_0(\mathcal{H}(u_\infty, v_\infty) + \|\Delta W\|_{L^\infty(\mathbb{R}^3)} + V_1)$$

for some constants $V_0, V_1 > 0$. Because of the continuous embedding of $W^{2,2}(\mathbb{R}^3)$ into $C^0(\mathbb{R}^3)$, it follows that $\|v_\infty\|_{L^\infty(\mathbb{R}^3)} \leq V$ for some $V > 0$.

Uniqueness of the minimizer is, by [Villani 2003, Theorem 5.32], a consequence of λ' -geodesic convexity of \mathcal{H} with respect to the distance induced by $\|\cdot\|_{L^2 \times L^2}$ for some $\lambda' > 0$ as proved in Proposition 3.1(c).

It remains to show that there is a set of Euler–Lagrange equations characterizing (u_∞, v_∞) .

The following variational inequality holds thanks to the minimizing property of (u_∞, v_∞) :

$$0 \leq \left. \frac{d^+}{ds} \right|_{s=0} \mathcal{H}(u_\infty + s\tilde{u}, v_\infty + s\tilde{v}) \\ = \int_{\mathbb{R}^3} (u_\infty + W + \varepsilon\phi(v_\infty))\tilde{u} \, dx + \int_{\mathbb{R}^3} (-\Delta v_\infty + \kappa v_\infty + \varepsilon u_\infty \phi'(v_\infty))\tilde{v} \, dx \quad (43)$$

for arbitrary maps \tilde{u} and \tilde{v} such that $u_\infty + \tilde{u} \geq 0$ on \mathbb{R}^3 and $\int_{\mathbb{R}^3} \tilde{u} \, dx = 0$.

First, we consider the second component and thus set $\tilde{u} = 0$ in (43). As there are no constraints on v_∞ , it is allowed to replace \tilde{v} by $-\tilde{v}$ in (43), yielding equality and hence (41).

Second, we consider the first component and set $\tilde{v} = 0$ in (43). For arbitrary ψ such that $\int_{\mathbb{R}^3} \psi \, dx \leq 1$ and $\psi + u_\infty \geq 0$ on \mathbb{R}^3 , we put

$$\tilde{u}_\psi := \frac{1}{2}\psi - \frac{1}{2}u_\infty \int_{\mathbb{R}^3} \psi \, dx$$

and observe that $u_\infty + \tilde{u}_\psi \geq 0$ on \mathbb{R}^3 and $\int_{\mathbb{R}^3} \tilde{u}_\psi \, dx = 0$ since u_∞ has mass equal to 1. By straightforward calculation, we obtain

$$0 \leq \int_{\mathbb{R}^3} (u_\infty + W + \varepsilon\phi(v_\infty) - U_\varepsilon)\psi \, dx \quad (44)$$

for all ψ as above and the constant

$$U_\varepsilon := \int_{\mathbb{R}^3} (u_\infty^2 + Wu_\infty + \varepsilon u_\infty \phi(v_\infty)) \, dx \in \mathbb{R}.$$

Fix $x \in \mathbb{R}^3$. If $u_\infty(x) > 0$, choosing ψ supported on a small neighborhood of x and replacing by $-\psi$ in (44) eventually yields

$$u_\infty(x) = U_\varepsilon - W(x) - \varepsilon\phi(v_\infty(x)).$$

If $u_\infty(x) = 0$, we obtain

$$U_\varepsilon - W(x) - \varepsilon\phi(v_\infty(x)) \leq 0.$$

Hence, for all $x \in \mathbb{R}^3$,

$$u_\infty(x) = [U_\varepsilon - W(x) - \varepsilon\phi(v_\infty(x))]_+. \quad \square$$

5B. Properties. As a preparation to prove some crucial regularity estimates on the stationary solution (u_∞, v_∞) , several properties of solutions to the elliptic partial differential equation $-\Delta h + \kappa h = f$ are needed.

Therefore, we introduce for $\kappa > 0$ the *Yukawa potential* (also called *screened Coulomb* or *Bessel potential*) \mathbf{G}_κ by

$$\mathbf{G}_\kappa(x) := \frac{1}{4\pi|x|} \exp(-\sqrt{\kappa}|x|) \quad \text{for all } x \in \mathbb{R}^3 \setminus \{0\}. \quad (45)$$

Additionally, we define for $\sigma > 0$ the kernel Y_σ by

$$Y_\sigma := \frac{1}{\sigma} \mathbf{G}_{1/\sigma}.$$

In subsequent parts of this work, we will need the iterates Y_σ^k for $k \in \mathbb{N}$ defined inductively by

$$Y_\sigma^1 := Y_\sigma, \quad Y_\sigma^{k+1} := Y_\sigma * Y_\sigma^k.$$

The relevant properties of \mathbf{G}_κ and Y_σ are summarized in Lemma 5.2 below. For the proof, we refer to Appendix A.

Lemma 5.2 (Yukawa potential). *The following statements hold for all $\kappa > 0$, $\sigma > 0$ and $k \in \mathbb{N}$:*

- (a) \mathbf{G}_κ and Y_σ are the fundamental solutions to $-\Delta h + \kappa h = f$ and $-\sigma \Delta h + h = f$ on \mathbb{R}^3 , respectively.
- (b) Let $p > 1$. If $f \in L^p(\mathbb{R}^3)$, then $\mathbf{G}_\kappa * f \in W^{2,p}(\mathbb{R}^3)$ and

$$\kappa \|\mathbf{G}_\kappa * f\|_{L^p(\mathbb{R}^3)} + \sqrt{\kappa} \|\mathbf{D}(\mathbf{G}_\kappa * f)\|_{L^p(\mathbb{R}^3)} + \|\mathbf{D}^2(\mathbf{G}_\kappa * f)\|_{L^p(\mathbb{R}^3)} \leq C_p \|f\|_{L^p(\mathbb{R}^3)} \quad (46)$$

for some p -dependent constant $C_p > 0$. (Note that this fact is not obvious as $\mathbf{D}^2(\mathbf{G}_\kappa) \notin L^1(\mathbb{R}^3, \mathbb{R}^{3 \times 3})$.)

- (c) For all $x \in \mathbb{R}^3 \setminus \{0\}$,

$$Y_\sigma(x) = \int_0^\infty \mathbf{H}_{\sigma t}(x) e^{-t} dt,$$

where \mathbf{H}_t is the heat kernel on \mathbb{R}^3 at time $t > 0$, i.e.,

$$\mathbf{H}_t(\xi) = t^{-3/2} \mathbf{H}_1(t^{-1/2} \xi) \quad \text{with } \mathbf{H}_1(\zeta) = (4\pi)^{-3/2} \exp(-\frac{1}{4} |\zeta|^2).$$

Additionally, one has

$$Y_\sigma^k = \int_0^\infty \mathbf{H}_{\sigma r} \frac{r^{k-1} e^{-r}}{\Gamma(k)} dr. \quad (47)$$

Moreover, $Y_\sigma^k \in W^{1,q}(\mathbb{R}^3)$ for each $q \in [1, \frac{3}{2})$, and there are universal constants Y_q such that

$$\|\mathbf{D}Y_\sigma^k\|_{L^q(\mathbb{R}^3)} \leq Y_q (\sigma k)^{-Q}, \quad \text{where } Q := 2 - \frac{3}{2q} \in [\frac{1}{2}, 1). \quad (48)$$

Now, we are in position to prove several estimates on the stationary solution.

Proposition 5.3 (estimates on the stationary solution). *The following uniform estimates hold for all $x \in \mathbb{R}^3$:*

- (a) $u_\infty(x) \leq U_0 - \varepsilon V \phi'(0)$, where $U_0 \in \mathbb{R}$ is chosen in such a way that $\int_{\mathbb{R}^3} [U_0 - W]_+ dx = 1$ and $V > 0$ is the constant from Proposition 5.1.
- (b) $|\mathbf{D}v_\infty(x)| \leq C\varepsilon$ for some constant $C > 0$.
- (c) $-C'\varepsilon \mathbb{1} \leq \mathbf{D}^2 v_\infty(x) \leq C'\varepsilon \mathbb{1}$ in the sense of symmetric matrices, for some constant $C' > 0$.

Proof. (a) We first prove that $U_\varepsilon \leq U_0 + \varepsilon \phi(0)$, which in turn follows if

$$\int_{\mathbb{R}^3} [U_0 + \varepsilon \phi(0) - W - \varepsilon \phi(v_\infty)]_+ dx \geq 1.$$

One has

$$\begin{aligned} \int_{\mathbb{R}^3} [U_0 + \varepsilon\phi(0) - W - \varepsilon\phi(v_\infty)]_+ dx &= \int_{\{U_0 - W \geq 0\}} [U_0 - W + \varepsilon(\phi(0) - \phi(v_\infty))] dx \\ &+ \int_{\{0 > U_0 - W \geq \varepsilon(\phi(v_\infty) - \phi(0))\}} [U_0 - W + \varepsilon(\phi(0) - \phi(v_\infty))] dx. \end{aligned} \quad (49)$$

From $\phi(0) - \phi(v_\infty) \geq 0$ and the definition of U_0 , we deduce that the first term on the right-hand side of (49) is larger than or equal to 1. The second term on the right-hand side of (49) is nonnegative because the integrand is nonnegative on the domain of integration.

Now, if $u_\infty(x) > 0$ for some $x \in \mathbb{R}^3$, we also have due to convexity of ϕ

$$u_\infty(x) \leq U_\varepsilon - W(x) - \varepsilon\phi(0) - \varepsilon v_\infty(x)\phi'(0) \leq U_0 + \varepsilon\phi(0) - \varepsilon\phi(0) - \varepsilon V\phi'(0),$$

from which the desired estimate follows.

(b) Define

$$f_v : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad f_v(x) := \varepsilon[U_\varepsilon - W(x) - \varepsilon\phi(v(x))]_+\phi'(v(x)).$$

Then, $f_v \in L^\infty(\mathbb{R}^3)$ with compact support $\text{supp}(f_v) \subset B_R(0)$ where $R > 0$ can be chosen independently of $\varepsilon \in (0, 1)$. Moreover, by Lemma 5.2(a), (u_∞, v_∞) is the solution to the integral equation

$$v = -(\mathbf{G}_\kappa * f_v)$$

with the Yukawa potential \mathbf{G}_κ defined in (45). Since $W^{2,4}(\mathbb{R}^3)$ is continuously embedded in $C^1(\mathbb{R}^3)$ [Zeidler 1990, Appendix, §(45) et seq.] and $f_v \in L^4(\mathbb{R}^3)$, we deduce from Lemma 5.2(b) that

$$\|v\|_{C^1(\mathbb{R}^3)} \leq \tilde{C}\|f_v\|_{L^4(\mathbb{R}^3)},$$

for some constant $\tilde{C} > 0$. Hence, we obtain (b) by using (a):

$$\|Dv_\infty\|_{L^\infty(\mathbb{R}^3)} \leq \tilde{C}\|f_{v_\infty}\|_{L^4(\mathbb{R}^3)} \leq \tilde{C}\varepsilon(U_0 - \varepsilon V\phi'(0))|\phi'(0)||B_R(0)|^{1/4} =: C\varepsilon.$$

(c) First, consider $x \in \mathbb{R}^3 \setminus B_{R+1}(0)$, where $R > 0$ is such that $\text{supp}(f_{v_\infty}) \subset B_R(0)$. Smoothness of \mathbf{G}_κ on $\mathbb{R}^3 \setminus \{0\}$ yields for all $i, j \in \{1, 2, 3\}$

$$\begin{aligned} |\partial_i \partial_j v_\infty(x)| &= \left| \int_{B_R(0)} (\partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x-y)) f_{v_\infty}(y) dy \right| \\ &= \left| \int_{x+B_R(0)} (\partial_i \partial_j \mathbf{G}_\kappa(z)) f_{v_\infty}(x-z) dz \right|, \end{aligned}$$

where the last equality follows by the transformation $z := x - y$. Obviously, we obtain the estimate

$$|\partial_i \partial_j v_\infty(x)| \leq \|f_{v_\infty}\|_{L^\infty(\mathbb{R}^3)} \int_{\mathbb{R}^3 \setminus B_1(0)} |\partial_i \partial_j \mathbf{G}_\kappa(z)| dz.$$

Since, for $|z| \geq 1$, one has (see Appendix B for the derivatives of \mathbf{G}_κ)

$$|\partial_i \partial_j \mathbf{G}_\kappa(z)| \leq \frac{C(\kappa) \exp(-\sqrt{\kappa}|z|)}{4\pi|z|},$$

we arrive at

$$|\partial_i \partial_j v_\infty(x)| \leq C(\kappa) \|f_{v_\infty}\|_{L^\infty(\mathbb{R}^3)} \int_1^\infty \exp(-\sqrt{\kappa}r) r \, dr,$$

the last integral obviously being finite.

Consider now the case $|x| \leq R + 1$, and set $y := (R + 2)e_1 \neq x$. By the triangular inequality, we have for $\alpha \in (0, 1)$ that

$$|\partial_i \partial_j v_\infty(x)| \leq |\partial_i \partial_j v_\infty(y)| + \frac{|\partial_i \partial_j v_\infty(x) - \partial_i \partial_j v_\infty(y)|}{|x - y|^\alpha} |x - y|^\alpha.$$

By the arguments above, f_{v_∞} is α -Hölder-continuous for some $\alpha \in (0, 1)$ since u_∞ is Lipschitz-continuous and of compact support. By Lemma B.1 in Appendix B, we know that there exists $C > 0$ such that

$$[\partial_i \partial_j v_\infty]_{C^{0,\alpha}(\mathbb{R}^3)} \leq C [f_{v_\infty}]_{C^{0,\alpha}(\mathbb{R}^3)}.$$

Hence, since $|x - y| \leq 2R + 3$, one has

$$|\partial_i \partial_j v_\infty(x)| \leq |\partial_i \partial_j v_\infty(y)| + C(2R + 3)^\alpha [f_{v_\infty}]_{C^{0,\alpha}(\mathbb{R}^3)}.$$

Combining both cases yields

$$\begin{aligned} |\partial_i \partial_j v_\infty(x)| &\leq |\partial_i \partial_j v_\infty((R + 2)e_1)| + C(2R + 3)^\alpha [f_{v_\infty}]_{C^{0,\alpha}(\mathbb{R}^3)} \\ &\leq C_0 \|f_{v_\infty}\|_{L^\infty(\mathbb{R}^3)} + C_1 \|f_{v_\infty}\|_{W^{1,\infty}(\mathbb{R}^3)} \end{aligned}$$

for some $C_0, C_1 > 0$ and *all* $x \in \mathbb{R}^3$. Using (a) and (b), it is straightforward to conclude that there exists $C_2 > 0$ with

$$(\|Df\|_{C^0(\mathbb{R}^3)} + \|f\|_{L^\infty(\mathbb{R}^3)}) \leq C_2 \varepsilon.$$

All in all, we proved the existence of $C_3 > 0$ such that for all $x \in \mathbb{R}^3$ and all $i, j \in \{1, 2, 3\}$

$$|\partial_i \partial_j v_\infty(x)| \leq C_3 \varepsilon.$$

Obviously, this estimate yields the assertion (for a different constant $C' > 0$). □

6. Convergence to equilibrium

In this section, we prove Theorem 1.2. The strategy of proof is as follows. We first show that the entropy $\mathcal{H}(u, v) - \mathcal{H}_\infty$ can indeed be decomposed as in (16). Furthermore, the second component v_τ^n of the discrete solution admits a control estimate enabling us to prove boundedness of the auxiliary entropy $\mathcal{L}_u(u) + \mathcal{L}_v(v)$ in (16) for large times. From that, we can deduce an explicit temporal bound such that exponential decay to zero of this entropy occurs for sufficiently large times. The previous two steps essentially comprise a rigorous version of (17) from the introduction. Finally, these estimates are converted into the desired estimate for the continuous weak solution, completing the proof of Theorem 1.2.

Since our claim only concerns the solutions (u, v) to (1)–(3) that are constructed as in the proof of Theorem 1.1, i.e., by the minimizing movement scheme, we assume in the following that we are given a family of time-discrete approximations $(u_\tau^n, v_\tau^n)_{n \in \mathbb{N}}$ that converge to the weak solution (u, v) in the

sense discussed in Section 4 as $\tau \downarrow 0$. Therefore, we may assume without loss of generality that $\tau > 0$ is sufficiently small.

Throughout this section, we shall use the abbreviation $[a]_\tau := (1/\tau) \log(1 + a\tau)$, where $a > 0$. Note that, for every $\tau > 0$ and an index $m_\tau \in \mathbb{N}$ given such that $m_\tau \tau \geq T$ with a fixed $T \geq 0$,

$$(1 + a\tau)^{-m_\tau} \leq e^{-[a]_\tau T} \downarrow e^{-aT} \quad \text{as } \tau \downarrow 0. \quad (50)$$

In order to keep track of the dependencies of certain quantities on ε , we are going to define several positive numbers ε_j such that the estimates in a certain proof are uniform with respect to $\varepsilon \in (0, \varepsilon_j)$. When we want to emphasize that a quantity is independent of $\varepsilon \in (0, \varepsilon_j)$ — and also of τ and the initial condition (u_0, v_0) — we call it a *system constant*. System constants are (in principle) expressible as a function of λ_0, κ and ϕ and truly universal constants. Finally, we write $\mathcal{H}_\infty := \mathcal{H}(u_\infty, v_\infty)$.

6A. Decomposition of the entropy. The key element in the proof of Theorem 1.2 is the decomposition of the entropy functional as announced in (16). Introduce the *perturbed potential* W_ε by

$$W_\varepsilon(x) := W(x) + \varepsilon\phi(v_\infty(x)). \quad (51)$$

Recall that (u_∞, v_∞) is the minimizer of \mathcal{H} on X , and define

$$\begin{aligned} \mathcal{L}_u(u) &:= \int_{\mathbb{R}^3} \left(\frac{1}{2}(u^2 - u_\infty^2) + W_\varepsilon(u - u_\infty) \right) dx, \\ \mathcal{L}_v(v) &:= \int_{\mathbb{R}^3} \frac{1}{2} (|D(v - v_\infty)|^2 + \kappa(v - v_\infty)^2) dx, \\ \mathcal{L}_*(u, v) &:= \int_{\mathbb{R}^3} (u[\phi(v) - \phi(v_\infty)] - u_\infty\phi'(v_\infty)[v - v_\infty]) dx. \end{aligned}$$

Finally, let $\mathcal{L}(u, v) := \mathcal{L}_u(u) + \mathcal{L}_v(v)$ denote the *auxiliary entropy*.

Lemma 6.1. *The decomposition (16) holds:*

$$\mathcal{H}(u, v) - \mathcal{H}_\infty = \mathcal{L}(u, v) + \varepsilon\mathcal{L}_*(u, v).$$

Proof. By the properties of ϕ and the fact that u_∞ has compact support, \mathcal{L}_* is well defined on all of X while \mathcal{L}_u and \mathcal{L}_v are finite precisely on $(\mathcal{P}_2 \cap L^2)(\mathbb{R}^3)$ and $W^{1,2}(\mathbb{R}^3)$, respectively. Thus, both sides in (16) are finite on the same subset of X . Now, for every such pair (u, v) , we have on the one hand that

$$\mathcal{L}_u(u) = \int_{\mathbb{R}^3} \left(\frac{1}{2}u^2 + uW + \varepsilon u\phi(v_\infty) \right) dx - \int_{\mathbb{R}^3} \left(\frac{1}{2}u_\infty^2 + u_\infty W + \varepsilon u_\infty\phi(v_\infty) \right) dx \quad (52)$$

and on the other hand that

$$\mathcal{L}_v(v) = \int_{\mathbb{R}^3} \left(\frac{1}{2}|Dv|^2 + \frac{1}{2}\kappa v^2 \right) dx + \int_{\mathbb{R}^3} \left(\frac{1}{2}|Dv_\infty|^2 + \frac{1}{2}\kappa v_\infty^2 \right) dx - \int_{\mathbb{R}^3} (Dv \cdot Dv_\infty + \kappa v v_\infty) dx.$$

Integration by parts in the last integral yields, recalling the defining equation (41) for v_∞ , that

$$- \int_{\mathbb{R}^3} (Dv \cdot Dv_\infty + \kappa v v_\infty) dx = \int_{\mathbb{R}^3} (\Delta v_\infty - \kappa v_\infty)v dx = \varepsilon \int_{\mathbb{R}^3} u_\infty\phi'(v_\infty)v dx.$$

Similarly, integration by parts in the middle integral leads to

$$\int_{\mathbb{R}^3} \left(\frac{1}{2} |\mathbf{D}v_\infty|^2 + \frac{1}{2} \kappa v_\infty^2 \right) dx = -\frac{\varepsilon}{2} \int_{\mathbb{R}^3} u_\infty \phi'(v_\infty) v_\infty dx.$$

And so,

$$\mathcal{L}_v(v) = \int_{\mathbb{R}^3} \left(\frac{1}{2} |\mathbf{D}v|^2 + \frac{1}{2} \kappa v^2 \right) dx - \int_{\mathbb{R}^3} \left(\frac{1}{2} |\mathbf{D}v_\infty|^2 + \frac{1}{2} \kappa v_\infty^2 \right) dx + \varepsilon \int_{\mathbb{R}^3} u_\infty \phi'(v_\infty) (v - v_\infty) dx. \quad (53)$$

Combining (52) and (53) with the definition of \mathcal{L}_* yields (16). \square

We summarize some useful properties of the auxiliary entropy \mathcal{L} in the following:

Proposition 6.2 (properties of \mathcal{L}). *There are constants $K, L > 0$ and some $\varepsilon_0 > 0$ such that the following are true for every $\varepsilon \in (0, \varepsilon_0)$:*

- (a) $W_\varepsilon \in C^2(\mathbb{R}^3)$ is λ_ε -convex with $\lambda_\varepsilon := \lambda_0 - L\varepsilon > 0$.
- (b) \mathcal{L}_u is λ_ε -convex in $(\mathcal{P}_2(\mathbb{R}^3), \mathbf{W}_2)$, and for every $u \in (\mathcal{P}_2 \cap W^{1,2})(\mathbb{R}^3)$, one has

$$\frac{1}{2} \|u - u_\infty\|_{L^2(\mathbb{R}^3)}^2 \leq \mathcal{L}_u(u) \leq \frac{1}{2\lambda_\varepsilon} \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx. \quad (54)$$

- (c) \mathcal{L}_v is κ -convex in $L^2(\mathbb{R}^3)$, and for every $v \in W^{2,2}(\mathbb{R}^3)$, one has

$$\frac{1}{2} \kappa \|v - v_\infty\|_{L^2(\mathbb{R}^3)}^2 \leq \mathcal{L}_v(v) \leq \frac{1}{2\kappa} \int_{\mathbb{R}^3} (\Delta(v - v_\infty) - \kappa(v - v_\infty))^2 dx. \quad (55)$$

- (d) For every $(u, v) \in \mathbf{X}$,

$$\mathcal{L}(u, v) \leq (1 + K\varepsilon)(\mathcal{H}(u, v) - \mathcal{H}_\infty). \quad (56)$$

Proof. (a) Since $W_\varepsilon = W + \varepsilon\phi(v_\infty)$, the chain rule yields

$$\mathbf{D}^2 W_\varepsilon = \mathbf{D}^2 W + \varepsilon \phi''(v_\infty) \mathbf{D}v_\infty \otimes \mathbf{D}v_\infty + \varepsilon \phi'(v_\infty) \mathbf{D}^2 v_\infty.$$

Using our assumptions on ϕ and by Proposition 5.3, there are some $L > 0$ and some ε_0 such that

$$\phi''(v_\infty) \mathbf{D}v_\infty \otimes \mathbf{D}v_\infty + \phi'(v_\infty) \mathbf{D}^2 v_\infty \geq -L\mathbb{1}$$

holds uniformly with respect to $\varepsilon \in (0, \varepsilon_0)$. And thus also $\mathbf{D}^2 W_\varepsilon \geq \lambda_\varepsilon \mathbb{1}$ with the indicated definition of λ_ε .

(b) Since W_ε is λ_ε -convex, \mathcal{L}_u is also λ_ε -geodesically convex in \mathbf{W}_2 because it is the sum of a 0-geodesically convex functional and a λ_ε -geodesically convex functional; see Theorem 2.1.

The Wasserstein subdifferential of \mathcal{L}_u has been calculated in [Ambrosio et al. 2008, Lemma 10.4.1]. Together with (19), this shows the second inequality in (54). For the first inequality, observe that

$$\mathcal{L}_u(u) = \frac{1}{2} \int_{\mathbb{R}^3} (u - u_\infty)^2 dx + \int_{\mathbb{R}^3} (W_\varepsilon + u_\infty)(u - u_\infty) dx.$$

It thus suffices to prove nonnegativity of the second integral term for all $u \in \mathcal{P}_2(\mathbb{R}^3)$. First, as u and u_∞ have equal mass, and by the definition of u_∞ ,

$$0 = \int_{\mathbb{R}^3} (u_\infty - u) dx = \int_{\{u_\varepsilon - W_\varepsilon > 0\}} u_\infty dx - \int_{\mathbb{R}^3} u dx,$$

and consequently,

$$\int_{\{U_\varepsilon - W_\varepsilon > 0\}} (u - u_\infty) \, dx = - \int_{\{U_\varepsilon - W_\varepsilon \leq 0\}} u \, dx. \quad (57)$$

Also, by the definition of u_∞ ,

$$\int_{\mathbb{R}^3} (W_\varepsilon + u_\infty)(u - u_\infty) \, dx = \int_{\{U_\varepsilon - W_\varepsilon > 0\}} U_\varepsilon (u - u_\infty) \, dx + \int_{\{U_\varepsilon - W_\varepsilon \leq 0\}} W_\varepsilon u \, dx.$$

Combining this with (57) yields

$$\int_{\{U_\varepsilon - W_\varepsilon > 0\}} U_\varepsilon (u - u_\infty) \, dx + \int_{\{U_\varepsilon - W_\varepsilon \leq 0\}} W_\varepsilon u \, dx = \int_{\{U_\varepsilon - W_\varepsilon \leq 0\}} (W_\varepsilon - U_\varepsilon) u \, dx \geq 0$$

as the integrand is nonnegative on the domain of integration.

(c) This is an immediate consequence of (19) for the L^2 subdifferential of \mathcal{L}_v .

(d) Since ϕ is convex, we have

$$\phi(v) - \phi(v_\infty) - \phi'(v_\infty)[v - v_\infty] \geq 0,$$

and so we can estimate \mathcal{L}_* from below by

$$\begin{aligned} \mathcal{L}_*(u, v) &= \int_{\mathbb{R}^3} (u - u_\infty)[\phi(v) - \phi(v_\infty)] \, dx + \int_{\mathbb{R}^3} (\phi(v) - \phi(v_\infty) - \phi'(v_\infty)[v - v_\infty]) \, dx \\ &\geq -\frac{1}{2} \int_{\mathbb{R}^3} (u - u_\infty)^2 \, dx - \frac{\phi'(0)^2}{2} \int_{\mathbb{R}^3} (v - v_\infty)^2 \, dx \\ &\geq -\mathcal{L}_u(u) - \frac{\phi'(0)^2}{\kappa} \mathcal{L}_v(v), \end{aligned}$$

using the properties (b) and (c) above. By (16), we conclude

$$(1 - K'\varepsilon)\mathcal{L}(u, v) = \mathcal{H}(u, v) - \mathcal{H}_\infty \quad \text{with } K' := \max\left(1, \frac{\phi'(0)^2}{\kappa}\right),$$

which clearly implies (56) for all $\varepsilon \in (0, \varepsilon_0)$, possibly after diminishing ε_0 . \square

6B. Dissipation. We can now formulate the main a priori estimate for the time-discrete solution.

Proposition 6.3. *Given $(\tilde{u}, \tilde{v}) \in X$ with $\mathcal{H}(\tilde{u}, \tilde{v}) < \infty$, let $(u, v) \in X$ be a minimizer of the functional $\mathcal{H}_\tau(\cdot \mid \tilde{u}, \tilde{v})$ introduced in (24). Then*

$$\mathcal{L}_u(u) + \tau \mathcal{D}_u(u, v) \leq \mathcal{L}_u(\tilde{u}), \quad \mathcal{L}_v(v) + \tau \mathcal{D}_v(u, v) \leq \mathcal{L}_v(\tilde{v}), \quad (58)$$

where the dissipation terms are given by

$$\mathcal{D}_u(u, v) = \left(1 - \frac{\varepsilon}{2}\right) \int_{\mathbb{R}^3} u |D(u + W_\varepsilon)|^2 \, dx - \frac{\varepsilon}{2} \int_{\mathbb{R}^3} u |D(\phi(v) - \phi(v_\infty))|^2 \, dx, \quad (59)$$

$$\mathcal{D}_v(u, v) = \left(1 - \frac{\varepsilon}{2}\right) \int_{\mathbb{R}^3} (\Delta(v - v_\infty) - \kappa(v - v_\infty))^2 \, dx - \frac{\varepsilon}{2} \int_{\mathbb{R}^3} (u\phi'(v) - u_\infty\phi'(v_\infty))^2 \, dx. \quad (60)$$

Proof. Naturally, these estimates are derived by means of the flow interchange lemma (Lemma 4.4).

For given $\nu > 0$, introduce the regularized functional $\mathcal{L}_u^\nu = \mathcal{L}_u + \nu \mathcal{E}$, where

$$\mathcal{E}(u) = \int_{\mathbb{R}^3} u \log u \, dx.$$

Note that \mathcal{E} is finite on $(\mathcal{P}_2 \cap L^2)(\mathbb{R}^3)$; see, e.g., [Zinsl 2014, Lemma 5.3]. Moreover, \mathcal{L}_u^ν is λ_ε -convex in W_2 by Theorem 2.1. We claim that the λ_ε -flow associated to \mathcal{L}_u^ν satisfies the evolution equation

$$\partial_s \mathcal{U} = \nu \Delta \mathcal{U} + \frac{1}{2} \Delta \mathcal{U}^2 + \operatorname{div}(\mathcal{U} D W_\varepsilon). \tag{61}$$

Since $\nu > 0$, this equation is strictly parabolic. Therefore, for every initial condition $\mathcal{U}_0 \in (\mathcal{P}_2 \cap L^2)(\mathbb{R}^3)$, there exists a smooth and positive solution $\mathcal{U} : \mathbb{R}_+ \times \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\mathcal{U}(s, \cdot) \rightarrow \mathcal{U}_0$ both in W_2 and in $L^2(\mathbb{R}^3)$ as $s \downarrow 0$. By [Ambrosio et al. 2008, Theorem 11.2.8], the solution operator to (61) can be identified with the λ_ε -flow of \mathcal{L}_u^ν .

Now, let \mathcal{U} be the smooth solution to (61) with initial condition $\mathcal{U}_0 = u$. By smoothness of \mathcal{U} , the equation (61) is satisfied in the classical sense at every time $s > 0$, and the following integration by parts is justified:

$$\begin{aligned} -\frac{d}{ds} \mathcal{H}(\mathcal{U}, \nu) &= - \int_{\mathbb{R}^3} [\mathcal{U} + W_\varepsilon + \varepsilon(\phi(\nu) - \phi(\nu_\infty))] \operatorname{div}[\mathcal{U} D(\mathcal{U} + W_\varepsilon) + \nu D\mathcal{U}] \, dx \\ &= \int_{\mathbb{R}^3} \mathcal{U} |D(\mathcal{U} + W_\varepsilon)|^2 \, dx + \varepsilon \int_{\mathbb{R}^3} \mathcal{U} D(\phi(\nu) - \phi(\nu_\infty)) \cdot D(\mathcal{U} + W_\varepsilon) \, dx \\ &\quad + \nu \int_{\mathbb{R}^3} D[\mathcal{U} + W + \varepsilon\phi(\nu)] \cdot D\mathcal{U} \, dx. \end{aligned}$$

The very last integral has already been estimated in the proof of Proposition 4.5 (see (34) and following). Rewriting the middle integral by means of the elementary inequality

$$2ab \leq a^2 + b^2, \tag{62}$$

we arrive at

$$\begin{aligned} -\frac{d}{ds} \mathcal{H}(\mathcal{U}, \nu) &\geq \left(1 - \frac{\varepsilon}{2}\right) \int_{\mathbb{R}^3} \mathcal{U} |D(\mathcal{U} + W_\varepsilon)|^2 \, dx - \frac{\varepsilon}{2} \int_{\mathbb{R}^3} \mathcal{U} |D(\phi(\nu) - \phi(\nu_\infty))|^2 \, dx \\ &\quad - \nu K \left(\|\mathcal{U}\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{W^{1,2}(\mathbb{R}^3)}^2 \right). \end{aligned}$$

We pass to the limit $s \downarrow 0$. Recall that \mathcal{U} converges (strongly) to its initial datum $\mathcal{U}_0 = u$ in $L^2(\mathbb{R}^3)$, and observe that the expressions on the right-hand side are lower semicontinuous with respect to that convergence. In fact, this is clear except perhaps for the first integral, which however can be rewritten, using integration by parts, in the form

$$\int_{\mathbb{R}^3} \mathcal{U} |D(\mathcal{U} + W_\varepsilon)|^2 \, dx = \frac{4}{9} \int_{\mathbb{R}^3} |D\mathcal{U}^{3/2}|^2 \, dx - \int_{\mathbb{R}^3} \mathcal{U}^2 \Delta W_\varepsilon \, dx + \int_{\mathbb{R}^3} \mathcal{U} |\nabla W_\varepsilon|^2 \, dx,$$

in which the lower semicontinuity is obvious since $\Delta W_\varepsilon \in L^\infty(\mathbb{R}^3)$. Applying now Lemma 4.4, we arrive at

$$\begin{aligned} \mathcal{L}_u^v(u) + (1 - \varepsilon) \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx - \frac{\varepsilon}{2} \int_{\mathbb{R}^3} u |\mathbf{D}(\phi(v) - \phi(v_\infty))|^2 dx \\ \leq \mathcal{L}_u^v(\tilde{u}) + \nu K (\|u\|_{L^2(\mathbb{R}^3)}^2 + \|u\|_{W^{1,2}(\mathbb{R}^3)}^2). \end{aligned}$$

Finally, passage to the limit $\nu \downarrow 0$ yields the dissipation (59).

The dissipation (60) is easier to obtain. It is immediate that the κ -flow in $L^2(\mathbb{R}^3)$ of \mathcal{L}_v satisfies the linear parabolic evolution equation

$$\partial_s \mathcal{V} = \Delta(\mathcal{V} - v_\infty) - \kappa(\mathcal{V} - v_\infty). \quad (63)$$

Solutions \mathcal{V} to (63) exist for arbitrary initial conditions $\mathcal{V}_0 \in L^2(\mathbb{R}^3)$, and they have at least the spatial regularity of v_∞ . Hence, with $\mathcal{V}_0 := v$, we have, also recalling the defining equation (41) for v_∞ ,

$$-\frac{d}{ds} \mathcal{H}(u, \mathcal{V}) = \int_{\mathbb{R}^3} [\Delta(\mathcal{V} - v_\infty) - \kappa(\mathcal{V} - v_\infty) - \varepsilon(u\phi'(\mathcal{V}) - u_\infty\phi'(v_\infty))] \cdot [\Delta(\mathcal{V} - v_\infty) - \kappa(\mathcal{V} - v_\infty)] dx.$$

Another application of the elementary inequality (62) yields

$$-\frac{d}{ds} \mathcal{H}(u, \mathcal{V}) \geq \left(1 - \frac{\varepsilon}{2}\right) \int_{\mathbb{R}^3} [\Delta(\mathcal{V} - v_\infty) - \kappa(\mathcal{V} - v_\infty)]^2 dx - \frac{\varepsilon}{2} \int_{\mathbb{R}^3} (u\phi'(\mathcal{V}) - u_\infty\phi'(v_\infty))^2 dx.$$

We pass to the limit $s \downarrow 0$ so that \mathcal{V} converges to v in $L^2(\mathbb{R}^3)$. The first integral is obviously lower semicontinuous. Concerning the second integral, note that the integrand converges pointwise a.e. on \mathbb{R}^3 on a subsequence and that it is pointwise a.e. bounded by the integrable function $2\phi'(0)^2(u^2 + u_\infty^2)$. Hence, we can pass to the limit using the dominated convergence theorem. Now another application of Lemma 4.4 yields the desired result. \square

We will need below two further estimates for the dissipation terms from (59)–(60).

Lemma 6.4. *There is a constant $\theta > 0$ such that, for every $\varepsilon \in (0, \varepsilon_0)$ and every $u \in (\mathcal{P}_2 \cap W^{1,2})(\mathbb{R}^3)$,*

$$\|u\|_{L^3(\mathbb{R}^3)}^4 \leq \theta \left(1 + \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx\right). \quad (64)$$

Proof. Integrating by parts, it is easily seen that

$$\frac{4}{9} \int_{\mathbb{R}^3} |\mathbf{D}u^{3/2}|^2 dx + \int_{\mathbb{R}^3} u |\mathbf{D}W_\varepsilon|^2 dx = \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx + \int_{\mathbb{R}^3} u^2 \Delta W_\varepsilon dx.$$

By Proposition 5.3 on the regularity of u_∞ and v_∞ , there exists a constant C such that

$$\Delta W_\varepsilon = \Delta W + \varepsilon\phi'(v_\infty)\Delta v_\infty + \varepsilon\phi''(v_\infty)|\mathbf{D}v_\infty|^2 \leq C \quad \text{on } \mathbb{R}^3$$

for all $\varepsilon \in (0, \varepsilon_1)$. Moreover,

$$\frac{1}{2} \int_{\mathbb{R}^3} u^2 dx \leq \int_{\mathbb{R}^3} u_\infty^2 dx + \frac{1}{\lambda_\varepsilon} \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx$$

by (54). Invoking again Proposition 5.3, it follows that there exists an ε -uniform constant C' such that

$$\|Du^{3/2}\|_{L^2(\mathbb{R}^3)}^2 \leq C' \left(1 + \int_{\mathbb{R}^3} u |D(u + W_\varepsilon)|^2 dx \right)$$

holds for all $u \in \mathcal{P}_2(\mathbb{R}^3)$. On the other hand, Hölder's and Sobolev's inequalities provide

$$\|u\|_{L^3(\mathbb{R}^3)} \leq \|u^{3/2}\|_{L^6(\mathbb{R}^3)}^{1/2} \|u\|_{L^1(\mathbb{R}^3)}^{1/4} \leq C'' \|Du^{3/2}\|_{L^2(\mathbb{R}^3)}^{1/2},$$

where we have used that u is of unit mass. Together, this yields (64). \square

Lemma 6.5. *For every $v \in W^{2,2}(\mathbb{R}^3)$,*

$$\min(1, 2\kappa, \kappa^2) \|v - v_\infty\|_{W^{2,2}(\mathbb{R}^3)}^2 \leq \int_{\mathbb{R}^3} (\Delta(v - v_\infty) - \kappa(v - v_\infty))^2 dx. \quad (65)$$

Proof. Set $\hat{v} := v - v_\infty$ for brevity. Integration by parts yields

$$\begin{aligned} \int_{\mathbb{R}^3} (\Delta\hat{v} - \kappa\hat{v})^2 dx &= \int_{\mathbb{R}^3} (\Delta\hat{v})^2 dx - 2\kappa \int_{\mathbb{R}^3} \hat{v} \Delta\hat{v} dx + \kappa^2 \int_{\mathbb{R}^3} \hat{v}^2 dx \\ &= \int_{\mathbb{R}^3} \|D^2\hat{v}\|^2 dx + 2\kappa \int_{\mathbb{R}^3} |D\hat{v}|^2 dx + \kappa^2 \int_{\mathbb{R}^3} \hat{v}^2 dx, \end{aligned}$$

which clearly implies (65). \square

6C. Control of the v component. For our estimates below, we need some preliminaries concerning solutions to the time-discrete heat equation. Here, we use the iterates Y_σ^k defined in (47) to write a semiexplicit representation of the components v_τ^n for a particular choice of σ .

Lemma 6.6. *For every $n \in \mathbb{N}$,*

$$v_\tau^n = (1 + \kappa\tau)^{-n} Y_\sigma^n * v_0 + \tau \sum_{m=1}^n (1 + \kappa\tau)^{-m} Y_\sigma^m * f_\tau^{n+1-m}, \quad (66)$$

where we have set

$$f_\tau^k := -\varepsilon u_\tau^k \phi'(v_\tau^k), \quad \sigma := \frac{\tau}{1 + \kappa\tau}.$$

Proof. We proceed by induction on n . By the flow interchange lemma (Lemma 4.4), using the auxiliary functional $\mathfrak{A}(u, v) := \int_{\mathbb{R}^3} \gamma v dx$ for an arbitrary test function $\gamma \in C_c^\infty(\mathbb{R}^3)$, one sees by analogous (but easier) arguments as in the proof of (60) that v_τ^n is the — unique in $L^2(\mathbb{R}^3)$ — distributional solution to

$$v_\tau^n - \sigma \Delta v_\tau^n = (1 + \kappa\tau)^{-1} v_\tau^{n-1} + \tau(1 + \kappa\tau)^{-1} f_\tau^n.$$

Hence, it can be written as

$$v_\tau^n = (1 + \kappa\tau)^{-1} Y_\sigma * v_\tau^{n-1} + \tau(1 + \kappa\tau)^{-1} Y_\sigma * f_\tau^n.$$

For $n = 1$, this is (66) because $v_\tau^0 = v_0$. Now, if $n > 1$ and (66) holds with $n - 1$ in place of n , then

$$v_\tau^n = (1 + \kappa\tau)^{-n} Y_\sigma * (Y_\sigma^{n-1} * v_0) + \tau \sum_{m=1}^{n-1} (1 + \kappa\tau)^{-(m+1)} Y_\sigma * (Y_\sigma^m * f_\tau^{n-m}) + \tau(1 + \kappa\tau)^{-1} Y_\sigma * f_\tau^n.$$

Using that $Y_\sigma * (Y_\sigma^k * f) = Y_\sigma^{k+1} * f$, we obtain (66). \square

We are now able to prove the main result of this section.

Proposition 6.7. *Provided that $v_0 \in L^{6/5}(\mathbb{R}^3)$, then $Dv_\tau^n \in L^{6/5}(\mathbb{R}^3)$ for every $n \in \mathbb{N}$, and the following estimate holds:*

$$\|Dv_\tau^n\|_{L^{6/5}(\mathbb{R}^3)} \leq a \|v_0\|_{L^{6/5}(\mathbb{R}^3)} e^{-[\kappa]_\tau n \tau} (n \tau)^{-1/2} + \varepsilon M_1 \quad (67)$$

with the system constants

$$a := (1 + \kappa) Y_1, \quad M_1 := |\phi'(0)| Y_{6/5} (1 + \kappa)^{3/4} \int_0^\infty (1 + \kappa)^{-s} s^{-3/4} ds. \quad (68)$$

Proof. From the representation formula (66), it follows that

$$\|Dv_\tau^n\|_{L^{6/5}(\mathbb{R}^3)} \leq (1 + \kappa \tau)^{-n} \|DY_\sigma^n\|_{L^1(\mathbb{R}^3)} \|v_0\|_{L^{6/5}(\mathbb{R}^3)} + \tau \sum_{m=1}^n (1 + \kappa \tau)^{-m} \|DY_\sigma^m\|_{L^{6/5}(\mathbb{R}^3)} \|f_\tau^{n+1-m}\|_{L^1(\mathbb{R}^3)}.$$

Now apply estimate (48), once with $q := 1$ and $Q := \frac{1}{2}$ to the first term and once with $q := \frac{6}{5}$ and $Q := \frac{3}{4}$ to the second term on the right-hand side. Further, since u_τ^n is of unit mass, one has

$$\|f_\tau^k\|_{L^1(\mathbb{R}^3)} = \varepsilon \|u_\tau^k \phi'(v_\tau^k)\|_{L^1(\mathbb{R}^3)} \leq \varepsilon |\phi'(0)|.$$

This yields

$$\|Dv_\tau^n\|_{L^{6/5}(\mathbb{R}^3)} \leq Y_1 \|v_0\|_{L^{6/5}(\mathbb{R}^3)} (1 + \kappa \tau)^{-n} (\sigma n)^{-1/2} + \varepsilon |\phi'(0)| Y_{6/5} \tau \sum_{m=1}^n (1 + \kappa \tau)^{-m} (\sigma m)^{-3/4}. \quad (69)$$

The sum in (69) is bounded uniformly in n and τ because

$$\tau \sum_{m=1}^\infty (1 + \kappa \tau)^{-m} (\sigma m)^{-3/4} \leq (1 + \kappa \tau)^{3/4} \int_0^\infty e^{-[\kappa]_\tau t} t^{-3/4} dt.$$

Without loss of generality, we assume that $\tau \leq 1$. By the monotone convergence $e^{-[\kappa]_\tau t} \downarrow e^{-\kappa t}$ as $\tau \downarrow 0$, we can estimate the sum in (69) as

$$\tau \sum_{m=1}^\infty (1 + \kappa \tau)^{-m} (\sigma m)^{-3/4} \leq (1 + \tau)^{3/4} \int_0^\infty (1 + \kappa)^{-t} t^{-3/4} dt,$$

and the right-hand side is finite. Thus, (69) implies (67) with the given constants. \square

In view of (50), we can draw the following conclusion from (67) with $\varepsilon_1 := \min(\varepsilon_0, \frac{1}{2})$, where $\varepsilon_0 > 0$ was implicitly characterized in Proposition 6.2:

Corollary 6.8. *Assume that $v_0 \in L^{6/5}(\mathbb{R}^3)$, and define*

$$T_1 := \max\left(0, \frac{1}{\kappa} \log \frac{a \|v_0\|_{L^{6/5}(\mathbb{R}^3)}}{M_1}\right) \quad (70)$$

with the system constants a and M_1 from (68). Then for every $\varepsilon \in (0, \varepsilon_1)$, for every sufficiently small τ and for every n such that $n \tau \geq T_1$, one has

$$\|Dv_\tau^n\|_{L^{6/5}(\mathbb{R}^3)} \leq 2M_1. \quad (71)$$

6D. Bounds on the auxiliary entropy. We are now in position to prove the main estimate leading towards exponential decay and boundedness of the auxiliary entropy \mathcal{L} along the discrete solution.

Lemma 6.9. *There are system constants L' and M' and an $\varepsilon_2 \in (0, \varepsilon_1)$ such that, for every $\varepsilon \in (0, \varepsilon_2)$, for every sufficiently small $\tau > 0$ and for every n with $n\tau > T_1$, we have that*

$$(1 + 2\Lambda'_\varepsilon \tau) \mathcal{L}(u_\tau^n, v_\tau^n) \leq \mathcal{L}(u_\tau^{n-1}, v_\tau^{n-1}) + \tau \varepsilon M' \tag{72}$$

with $\Lambda'_\varepsilon := \min(\kappa, \lambda_0) - L'\varepsilon$.

Proof. For brevity, we simply write u and v in place of u_τ^n and v_τ^n , respectively, and we introduce $\hat{v} := v - v_\infty$. Since $n\tau > T_1$ by hypothesis, Corollary 6.8 implies that

$$\|\mathbf{D}\hat{v}\|_{L^{6/5}(\mathbb{R}^3)} \leq \|\mathbf{D}v\|_{L^{6/5}(\mathbb{R}^3)} + \|\mathbf{D}v_\infty\|_{L^{6/5}(\mathbb{R}^3)} \leq Z := 2M_1 + \sup_{0 < \varepsilon < \varepsilon_1} \|\mathbf{D}v_\infty\|_{L^{6/5}(\mathbb{R}^3)} < \infty.$$

Now, since

$$|\mathbf{D}(\phi(v) - \phi(v_\infty))|^2 \leq 2\phi'(v)^2 |\mathbf{D}\hat{v}|^2 + 2(\phi'(v) - \phi'(v_\infty))^2 |\mathbf{D}v_\infty|^2 \leq \alpha |\mathbf{D}\hat{v}|^2 + \beta \hat{v}^2,$$

with the system constants

$$\alpha := 2\phi'(0)^2, \quad \beta := 2\overline{\phi}''^2 \sup_{0 < \varepsilon < \varepsilon_1} \|\mathbf{D}v_\infty\|_{L^\infty(\mathbb{R}^3)}^2, \tag{73}$$

we conclude that

$$\begin{aligned} \int_{\mathbb{R}^3} u |\mathbf{D}(\phi(v) - \phi(v_\infty))|^2 dx &\leq \alpha \int_{\mathbb{R}^3} u |\mathbf{D}\hat{v}|^2 dx + \beta \int_{\mathbb{R}^3} u \hat{v}^2 dx \\ &\leq \alpha \|u\|_{L^3(\mathbb{R}^3)} \|\mathbf{D}\hat{v}\|_{L^3(\mathbb{R}^3)}^2 + \beta \|u\|_{L^1(\mathbb{R}^3)} \|\hat{v}\|_{L^\infty(\mathbb{R}^3)}^2 \\ &\leq \|u\|_{L^3(\mathbb{R}^3)}^4 + \alpha^{4/3} \|\mathbf{D}\hat{v}\|_{L^3(\mathbb{R}^3)}^{8/3} + \beta \|\hat{v}\|_{L^\infty(\mathbb{R}^3)}^2 \\ &\leq \|u\|_{L^3(\mathbb{R}^3)}^4 + \alpha^{4/3} (S_1 \|\hat{v}\|_{W^{2,2}(\mathbb{R}^3)}^{3/4} \|\mathbf{D}\hat{v}\|_{L^{6/5}(\mathbb{R}^3)}^{1/4})^{8/3} + \beta S_2 \|\hat{v}\|_{W^{2,2}(\mathbb{R}^3)}^2 \\ &\leq \theta \left(1 + \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx \right) + \frac{\alpha^{4/3} S_1^{8/3} Z^{2/3} + \beta S_2}{\min(1, 2\kappa, \kappa^2)} \int_{\mathbb{R}^3} (\Delta \hat{v} - \kappa \hat{v})^2 dx, \end{aligned} \tag{74}$$

where θ is the constant from (64) and S_1 and S_2 are Sobolev constants. Next, observe that

$$(u\phi'(v) - u_\infty\phi'(v_\infty))^2 \leq 2(u - u_\infty)^2 \phi'(v)^2 + 2u_\infty^2 (\phi'(v) - \phi'(v_\infty))^2 \leq \alpha(u - u_\infty)^2 + \beta \|u_\infty\|_{L^\infty(\mathbb{R}^3)}^2 \hat{v}^2$$

with the same constants as in (73). Therefore, using (54), (55) and Proposition 5.3(a),

$$\begin{aligned} \int_{\mathbb{R}^3} (u\phi'(v) - u_\infty\phi'(v_\infty))^2 dx &\leq \alpha \|u - u_\infty\|_{L^2(\mathbb{R}^3)}^2 + \beta (U_0 - \varepsilon V\phi'(0))^2 \|\hat{v}\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq 2\alpha \mathcal{L}_u(u) + \frac{2\beta}{\kappa} (U_0 - \varepsilon V\phi'(0))^2 \mathcal{L}_v(v). \end{aligned}$$

Altogether, we have shown that there is a system constant M' such that (recall the dissipation terms $\mathcal{D}_u(u, v)$ and $\mathcal{D}_v(u, v)$ from (59)–(60))

$$\begin{aligned} \mathcal{D}_u(u, v) + \mathcal{D}_v(u, v) &\geq (1 - M'\varepsilon) \int_{\mathbb{R}^3} u |\mathbf{D}(u + W_\varepsilon)|^2 dx + (1 - M'\varepsilon) \int_{\mathbb{R}^3} (\Delta \hat{v} - \kappa \hat{v})^2 dx \\ &\quad - M'\varepsilon \mathcal{L}_u(u) - M'\varepsilon \mathcal{L}_v(v) - M'\varepsilon \end{aligned} \quad (75)$$

for all $\varepsilon \in (0, \varepsilon_1)$. Provided that $M'\varepsilon < 1$, we can apply (54) and (55) to estimate further:

$$\mathcal{D}_u(u, v) + \mathcal{D}_v(u, v) \geq (2\lambda_\varepsilon(1 - M'\varepsilon) - M'\varepsilon) \mathcal{L}_u(u) + (2\kappa(1 - M'\varepsilon) - M'\varepsilon) \mathcal{L}_v(v) - M'\varepsilon.$$

Finally, we can choose $\varepsilon_2 \in (0, \varepsilon_1)$ so small that the coefficients of \mathcal{L}_u and \mathcal{L}_v above are nonnegative for every $\varepsilon \in (0, \varepsilon_2)$, and thus, we arrive at the final estimate

$$\mathcal{D}_u(u, v) + \mathcal{D}_v(u, v) \geq 2(\min(\kappa, \lambda_\varepsilon) - L'\varepsilon) \mathcal{L}(u, v) - \varepsilon M'$$

with a suitable choice of L' . Now estimate (58) implies (72) with Λ'_ε given as above. \square

Diminishing ε_2 such that the constant $1 + K\varepsilon_2$ in (56) is less than or equal to 2, we derive the following explicit estimate:

Proposition 6.10. *Assume that $v_0 \in L^{6/5}(\mathbb{R}^3)$, and let T_1 be defined as in (70). Then, for every $\varepsilon \in (0, \varepsilon_2)$, for every sufficiently small τ and for every n with $n\tau > T_1$, the following estimate holds:*

$$\mathcal{L}(u_\tau^n, v_\tau^n) \leq 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty) e^{-2[\Lambda'_\varepsilon]_1(n\tau - T_1)} + \varepsilon M_2 \quad (76)$$

with the system constant

$$M_2 := \frac{M'}{2 \inf_{0 < \varepsilon < \varepsilon_2} \Lambda'_\varepsilon}.$$

Proof. We prove a slightly refined estimate: given $\bar{n} \in \mathbb{N}$ with $\bar{n}\tau \geq T_1$, we conclude by induction on $n \geq \bar{n}$ that

$$\mathcal{L}(u_\tau^n, v_\tau^n) \leq 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)(1 + 2\Lambda'_\varepsilon\tau)^{-(n-\bar{n})} + \frac{M'\varepsilon}{2\Lambda'_\varepsilon}(1 - (1 + 2\Lambda'_\varepsilon\tau)^{-(n-\bar{n})}), \quad (77)$$

which clearly implies (76). For $n = \bar{n}$, (77) is a consequence of (56) and the energy estimate (27). Now assume (77) for some $n \geq \bar{n}$, and apply the iterative estimate (72):

$$\begin{aligned} \mathcal{L}(u_\tau^{n+1}, v_\tau^{n+1}) &\leq (1 - 2\Lambda'_\varepsilon\tau)^{-1} \mathcal{L}(u_\tau^n, v_\tau^n) + (1 + 2\Lambda'_\varepsilon\tau)^{-1} \tau M'\varepsilon \\ &\leq 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)(1 + 2\Lambda'_\varepsilon\tau)^{-((n+1)-\bar{n})} \\ &\quad + \frac{M'\varepsilon}{2\Lambda'_\varepsilon} \left((1 + 2\Lambda'_\varepsilon\tau)^{-1} - (1 + 2\Lambda'_\varepsilon\tau)^{-((n+1)-\bar{n})} \right) + (1 + 2\Lambda'_\varepsilon\tau)^{-1} \tau M'\varepsilon. \end{aligned}$$

Elementary calculations show that the last expression above equals the right-hand side of (77) with $n + 1$ in place of n . \square

Invoking again (50), we obtain the following analog to Corollary 6.8:

Corollary 6.11. *Assume that $v_0 \in L^{6/5}(\mathbb{R}^3)$, and define*

$$T_2 := T_1 + \max\left(0, \frac{1}{2\Lambda'_\varepsilon} \log \frac{2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)}{M_2}\right). \tag{78}$$

Then for every $\varepsilon \in (0, \varepsilon_2)$, for every sufficiently small τ and for every n such that $n\tau \geq T_2$, one has

$$\mathcal{L}(u_\tau^n, v_\tau^n) \leq 2M_2. \tag{79}$$

We have thus proved that, for $t \geq T_2$, the auxiliary entropy \mathcal{L} is bounded by a system constant. Next, we prove that \mathcal{L} is not only bounded but actually convergent to zero exponentially fast.

6E. Exponential decay for large times.

Lemma 6.12. *There is a constant $L'' > L'$ and some $\varepsilon_3 \in (0, \varepsilon_2)$ such that, for every $\varepsilon \in (0, \varepsilon_3)$, for every sufficiently small $\tau > 0$ and for every n such that $n\tau > T_2$, we have*

$$(1 + 2\Lambda''_\varepsilon \tau) \mathcal{L}(u_\tau^n, v_\tau^n) \leq \mathcal{L}(u_\tau^{n-1}, v_\tau^{n-1}) \tag{80}$$

with $\Lambda''_\varepsilon := \min(\lambda_0, \kappa) - L''\varepsilon$.

Proof. We proceed like in the proof of Lemma 6.9 with the following modification. By Corollary 6.11, we know that

$$\mathcal{L}_u(u_\tau^n) \leq \mathcal{L}(u_\tau^n, v_\tau^n) \leq 2M_2.$$

Using the first inequality in (54), we can estimate the L^2 -norm of u_τ^n by a system constant:

$$\|u\|_{L^2(\mathbb{R}^3)} \leq \|u_\infty\|_{L^2(\mathbb{R}^3)} + \|u - u_\infty\|_{L^2(\mathbb{R}^3)} \leq Z := \sup_{0 < \varepsilon < \varepsilon_2} \|u_\infty\|_{L^2(\mathbb{R}^3)} + 2\sqrt{M_2}.$$

This allows us to replace the chain of estimates (74) by a simpler one:

$$\int_{\mathbb{R}^3} u |D(\phi(v) - \phi(v_\infty))|^2 dx \leq \|u\|_{L^2(\mathbb{R}^3)} (\alpha \|D\hat{v}\|_{L^4(\mathbb{R}^3)}^2 + \beta \|\hat{v}\|_{L^4(\mathbb{R}^3)}^2)$$

with the constants from (73). Using the Sobolev inequalities

$$\|D\hat{v}\|_{L^4(\mathbb{R}^3)} \leq S \|\hat{v}\|_{W^{2,2}(\mathbb{R}^3)}, \quad \|\hat{v}\|_{L^4(\mathbb{R}^3)} \leq S \|\hat{v}\|_{W^{1,2}(\mathbb{R}^3)}$$

in combination with (65) and (55), respectively, we arrive at

$$\int_{\mathbb{R}^3} u |D(\phi(v) - \phi(v_\infty))|^2 dx \leq \frac{\alpha Z S^2}{\min(1, 2\kappa, \kappa^2)} \int_{\mathbb{R}^3} (\Delta \hat{v} - \kappa \hat{v})^2 dx + \frac{2\beta Z S^2}{\min(1, \kappa)} \mathcal{L}_v(v).$$

This eventually leads to the dissipation estimate (75) again, with a different constant M' , but *without the constant term $-\varepsilon M'$* . By means of (58), this implies (80) for appropriate choices of L'' and ε_3 . \square

By iteration of (80), starting from (79), one immediately obtains

Proposition 6.13. *For all sufficiently small τ and every n such that $n\tau \geq T_2$, we have*

$$\mathcal{L}(u_\tau^n, v_\tau^n) \leq 2M_2 e^{-2[\Lambda''_\varepsilon]_\tau (n\tau - T_2)}. \tag{81}$$

6F. Passage to continuous time. To complete the proof of Theorem 1.2, we consider the limit $\tau \downarrow 0$ of the estimates obtained above. Here $\tau \downarrow 0$ means that we consider a vanishing sequence $(\tau_k)_{k \in \mathbb{N}}$ such that the corresponding sequence of discrete solutions $(u_{\tau_k}, v_{\tau_k})_{k \in \mathbb{N}}$ converges in the sense of Section 4 to a weak solution (u, v) to (1)–(3). Since the convergence of $(u_{\tau_k}, v_{\tau_k})_{k \in \mathbb{N}}$ in X is locally uniform on each compact time interval, the lower semicontinuity of \mathcal{L} in X allows one to conclude that

$$\mathcal{L}(t) := \mathcal{L}(u(t), v(t)) \leq \liminf_{\tau \downarrow 0} \mathcal{L}(u_\tau(t), v_\tau(t)) \quad \text{for every } t \geq 0.$$

We prove that

$$\mathcal{L}(t) \leq C(1 + \|v_0\|_{L^{6/5}(\mathbb{R}^3)})^2(1 + \mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)^2 e^{-2\Lambda_\varepsilon'' t} \quad \text{for all } t \geq 0. \quad (82)$$

From this, claim (15) in Theorem 1.2 follows with $\Lambda_\varepsilon := \Lambda_\varepsilon''$.

Recalling (50), we conclude from (81) that

$$\mathcal{L}(t) \leq 2M_2 e^{-2\Lambda_\varepsilon''(t-T_2)} \quad \text{for all } t \geq T_2. \quad (83)$$

Moreover, from (56) and the energy estimate (27), we obtain

$$\mathcal{L}(t) \leq 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty) \quad \text{for all } t \geq 0. \quad (84)$$

We distinguish:

Case 1 ($\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty \leq \frac{1}{2}M_2$). Then, from the definition of T_2 in (78), one has $T_2 = T_1$, and in consequence of (83),

$$\mathcal{L}(t) \leq 2M_2 e^{-2\Lambda_\varepsilon''(t-T_1)} \quad \text{for all } t \geq T_2 = T_1.$$

Since further $\mathcal{L}(t) \leq M_2$ for all $t \geq 0$ by (84), the first inequality extends to all times $t \geq 0$:

$$\mathcal{L}(t) \leq 2M_2 e^{-2\Lambda_\varepsilon''(t-T_1)} \quad \text{for all } t \geq 0. \quad (85)$$

Case 2 ($\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty > \frac{1}{2}M_2$). Substitute

$$T_2 = T_1 + \frac{1}{2\Lambda_\varepsilon'} \log\left(\frac{2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)}{M_2}\right)$$

in (83) to find

$$\mathcal{L}(t) \leq 2M_2 e^{-2\Lambda_\varepsilon''(t-T_1)} \left(\frac{2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)}{M_2}\right)^{\Lambda_\varepsilon''/\Lambda_\varepsilon'} \quad \text{for all } t \geq T_2 > T_1. \quad (86)$$

Using $\Lambda_\varepsilon'' \leq \Lambda_\varepsilon'$, we conclude from (86)

$$\mathcal{L}(t) \leq 4(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty) e^{-2\Lambda_\varepsilon''(t-T_1)} \quad \text{for all } t \geq T_2 > T_1. \quad (87)$$

Define $A := 4(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty) \max(1, (\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)/M_2)$. Then, from (87) and the fact that $Ae^{-2\Lambda_\varepsilon''(T_2-T_1)} \geq 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)$, we deduce

$$\mathcal{L}(t) \leq Ae^{-2\Lambda_\varepsilon''(t-T_1)} \quad \text{for all } t \geq 0. \quad (88)$$

Together, (85) and (88) yield

$$\begin{aligned} \mathcal{L}(t) &\leq \max(2M_2, 4(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)) \max\left(1, \frac{1}{M_2}(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)\right) e^{-2\Lambda''_\varepsilon(t-T_1)} \\ &\leq \frac{2}{M_2} e^{2\Lambda''_\varepsilon T_1} \max(M_2, 2(\mathcal{H}(u_0, v_0) - \mathcal{H}_\infty))^2 e^{-2\Lambda''_\varepsilon t} \quad \text{for all } t \geq 0. \end{aligned}$$

Since $\kappa \geq \Lambda''_\varepsilon$, we have

$$e^{2\Lambda''_\varepsilon T_1} \leq e^{2\kappa T_1} \leq \max\left(1, \left[\frac{a}{M_1} \|v_0\|_{L^{6/5}(\mathbb{R}^3)}\right]^2\right)$$

and consequently (82):

$$\mathcal{L}(t) \leq C(1 + \|v_0\|_{L^{6/5}(\mathbb{R}^3)})^2 (1 + \mathcal{H}(u_0, v_0) - \mathcal{H}_\infty)^2 e^{-2\Lambda''_\varepsilon t} \quad \text{for all } t \geq 0.$$

Appendix A: Proof of Lemma 5.2

(a) The proof of the first assertion can be found in [Lieb and Loss 2001, Theorem 6.23]. From that, the second one follows by elementary calculations.

(b) According to [Stein 1970, Chapter V, §3.3, Theorem 3], one has for $p > 1$

$$\|\mathbf{G}_1 * f\|_{W^{2,p}(\mathbb{R}^3)} \leq C_p \|f\|_{L^p(\mathbb{R}^3)}. \quad (89)$$

To prove assertion (b), we use a rescaling of the equation $-\Delta h + \kappa h = f$ by $\tilde{x} := \sqrt{\kappa}x$. Consequently, $h(\tilde{x}) = (\mathbf{G}_\kappa * f)(\tilde{x}/\sqrt{\kappa})$ is a solution to $-\Delta_{\tilde{x}} h + h = f/\kappa$, i.e., $h(\tilde{x}) = (\mathbf{G}_1 * (f/\kappa))(\tilde{x})$. By the transformation theorem, we obtain

$$\begin{aligned} \left(\int_{\mathbb{R}^3} \left|\frac{f(\tilde{x})}{\kappa}\right|^p d\tilde{x}\right)^{1/p} &= \kappa^{3/2-1} \|f\|_{L^p(\mathbb{R}^3)}, \\ \left(\int_{\mathbb{R}^3} \left|(\mathbf{G}_1 * \frac{f}{\kappa})(\tilde{x})\right|^p d\tilde{x}\right)^{1/p} &= \kappa^{3/2} \|\mathbf{G}_\kappa * f\|_{L^p(\mathbb{R}^3)}, \\ \left(\int_{\mathbb{R}^3} \left|D_{\tilde{x}}(\mathbf{G}_1 * \frac{f}{\kappa})(\tilde{x})\right|^p d\tilde{x}\right)^{1/p} &= \kappa^{3/2-1/2} \|D_x(\mathbf{G}_\kappa * f)\|_{L^p(\mathbb{R}^3)}, \\ \left(\int_{\mathbb{R}^3} \left|D_{\tilde{x}}^2(\mathbf{G}_1 * \frac{f}{\kappa})(\tilde{x})\right|^p d\tilde{x}\right)^{1/p} &= \kappa^{3/2-1} \|D_x^2(\mathbf{G}_\kappa * f)\|_{L^p(\mathbb{R}^3)}, \end{aligned}$$

which yields (46) after insertion into (89) and simplification.

(c) The first statement is a straightforward consequence of the integral-type representation of \mathbf{G}_κ in [Lieb and Loss 2001, Theorem 6.23]. To prove the first claim of the second statement, we proceed by induction. For $k = 1$, Equation (47) is just the definition of \mathbf{Y}_σ . Now assume that (47) holds for some $k \in \mathbb{N}$. Using

the semigroup property $\mathbf{H}_{t_1+t_2} = \mathbf{H}_{t_1} * \mathbf{H}_{t_2}$ of the heat kernel, we find that

$$\begin{aligned} \mathbf{Y}_\sigma^{k+1} &= \int_0^\infty \int_0^\infty \mathbf{H}_{\sigma r_1} * \mathbf{H}_{\sigma r_2} e^{-r_1} r_2^{k-1} e^{-r_2} \frac{dr_1 dr_2}{\Gamma(k)} \\ &= \int_0^\infty \int_0^\infty \mathbf{H}_{\sigma(r_1+r_2)} e^{-(r_1+r_2)} r_2^{k-1} \frac{dr_1 dr_2}{\Gamma(k)}. \end{aligned}$$

Now perform a change of variables

$$r := r_1 + r_2, \quad s := r_2,$$

which is of determinant 1 and leads to

$$\mathbf{Y}_\sigma^{k+1} = \int_0^\infty \mathbf{H}_{\sigma r} e^{-r} \left(\int_0^r s^{k-1} ds \right) \frac{dr}{\Gamma(k)} = \int_0^\infty \mathbf{H}_{\sigma r} \frac{e^{-r} r^k dr}{k\Gamma(k)},$$

which is (47) with $k+1$ in place of k , using that $k\Gamma(k) = \Gamma(k+1)$.

For (48), first observe that $r \mapsto r^{k-1} e^{-r} / \Gamma(k)$ defines a probability density on \mathbb{R}_+ . We can thus apply Jensen's inequality to obtain

$$\|\mathbf{D}\mathbf{Y}_\sigma^k\|_{L^q(\mathbb{R}^3)} \leq \int_0^\infty \|\mathbf{D}\mathbf{H}_{\sigma r}\|_{L^q(\mathbb{R}^3)} \frac{r^{k-1} e^{-r} dr}{\Gamma(k)}. \quad (90)$$

The L^q -norm of $\mathbf{D}\mathbf{H}_{\sigma r}$ is easily evaluated using its definition,

$$\begin{aligned} \|\mathbf{D}\mathbf{H}_{\sigma r}\|_{L^q(\mathbb{R}^3)} &= (\sigma r)^{-3/2} \left(\int_{\mathbb{R}^3} |\mathbf{D}_\xi \mathbf{H}_1((\sigma r)^{-1/2} \xi)|^q d\xi \right)^{1/q} \\ &= (\sigma r)^{-3/2} \left(\int_{\mathbb{R}^3} |(\sigma r)^{-1/2} \mathbf{D}_\zeta \mathbf{H}_1(\zeta)|^q (\sigma r)^{3/2} d\zeta \right)^{1/q} = (\sigma r)^{-Q} \|\mathbf{D}\mathbf{H}_1\|_{L^q(\mathbb{R}^3)}. \end{aligned}$$

By definition of the gamma function, we thus obtain from (90) that

$$\|\mathbf{D}\mathbf{Y}_\sigma^k\|_{L^q(\mathbb{R}^3)} \leq \|\mathbf{D}\mathbf{H}_1\|_{L^q(\mathbb{R}^3)} \frac{\Gamma(k-Q)}{\Gamma(k)} \sigma^{-Q}.$$

For further estimation, observe that the sequence $(a_k)_{k \in \mathbb{N}}$ with $a_k = k^Q \Gamma(k-Q) / \Gamma(k)$ is monotonically decreasing (to zero). Indeed,

$$\frac{a_{k+1}}{a_k} = \frac{(k+1)k^Q (k-Q)\Gamma(k-Q)\Gamma(k)}{k^Q k\Gamma(k)\Gamma(k-Q)} = \left(1 + \frac{1}{k}\right)^Q \left(1 - \frac{Q}{k}\right)$$

is always smaller than 1 since $\xi \mapsto (1+\xi)^{-Q}$ is convex. Therefore, $a_k \leq a_1$ for all $k \in \mathbb{N}$, and so (48) follows with $Y_q := \Gamma(1-Q) \|\mathbf{D}\mathbf{H}_1\|_{L^q(\mathbb{R}^3)}$. \square

Appendix B: Hölder estimate for the kernel G_κ

As a preparation, we calculate the derivatives of G_κ in $\mathbb{R}^3 \setminus \{0\}$. For all $i, j, k \in \{1, 2, 3\}$, one has

$$\begin{aligned} \partial_i G_\kappa(x) &= -\frac{1}{4\pi} \frac{\exp(-\sqrt{\kappa}|x|)}{|x|^3} (\sqrt{\kappa}|x| + 1)x_i, \\ \partial_i \partial_j G_\kappa(x) &= -\frac{1}{4\pi} \exp(-\sqrt{\kappa}|x|) \left[\left(\frac{\kappa}{|x|^3} + \frac{3\sqrt{\kappa}}{|x|^4} + \frac{3}{|x|^5} \right) x_i x_j - \left(\frac{\sqrt{\kappa}}{|x|^2} + \frac{1}{|x|^3} \right) \delta_{ij} \right], \\ \partial_i \partial_j \partial_k G_\kappa(x) &= -\frac{1}{4\pi} \exp(-\sqrt{\kappa}|x|) \frac{-\sqrt{\kappa}x_k}{|x|} \left[\left(\frac{\kappa}{|x|^3} + \frac{3\sqrt{\kappa}}{|x|^4} + \frac{3}{|x|^5} \right) x_i x_j - \left(\frac{\sqrt{\kappa}}{|x|^2} + \frac{1}{|x|^3} \right) \delta_{ij} \right] \\ &\quad - \frac{1}{4\pi} \exp(-\sqrt{\kappa}|x|) \left[\left(-\frac{3\kappa}{|x|^4} - \frac{12\sqrt{\kappa}}{|x|^5} - \frac{15}{|x|^6} \right) \frac{x_i x_j x_k}{|x|} \right. \\ &\quad \left. + \delta_{ij} \left(\frac{2\sqrt{\kappa}}{|x|^3} + \frac{3}{|x|^4} \right) \frac{x_k}{|x|} + \left(\frac{\kappa}{|x|^3} + \frac{3\sqrt{\kappa}}{|x|^4} + \frac{3}{|x|^5} \right) (\delta_{ik} x_j + \delta_{jk} x_i) \right], \end{aligned}$$

where δ_{ij} denotes Kronecker's delta.

We prove the following:

Lemma B.1 (Hölder estimate for second derivative). *Let $f \in C^{0,\alpha}(\mathbb{R}^3)$ for some $\alpha \in (0, 1)$, and assume that it is of compact support. Then, there exists $C > 0$ such that for all $i, j \in \{1, 2, 3\}$ the following estimate holds:*

$$[\partial_i \partial_j (G_\kappa * f)]_{C^{0,\alpha}(\mathbb{R}^3)} \leq C[f]_{C^{0,\alpha}(\mathbb{R}^3)}.$$

Here,

$$[g]_{C^{0,\alpha}(\mathbb{R}^3)} := \sup_{x,y \in \mathbb{R}^3, x \neq y} \frac{|g(x) - g(y)|}{|x - y|}$$

denotes the Hölder seminorm of $g : \mathbb{R}^3 \rightarrow \mathbb{R}$.

Proof. This result is an extension of the respective result for Poisson's equation (corresponding to $\kappa = 0$) proved by Lieb and Loss [2001, Theorem 10.3]. Their method of proof is adapted here. In the following, C and \tilde{C} denote generic nonnegative constants.

The following holds for arbitrary test functions $\psi \in C_c^\infty(\mathbb{R}^3)$:

$$-\int_{\mathbb{R}^3} (\partial_j \psi)(x) (\partial_i v)(x) dx = \int_{\mathbb{R}^3} f(y) \int_{\mathbb{R}^3} (\partial_j \psi)(x) \partial_{x_i} G_\kappa(x - y) dx dy,$$

which can be rewritten by the dominated convergence theorem and integration by parts as

$$\begin{aligned} &\int_{\mathbb{R}^3} f(y) \int_{\mathbb{R}^3} (\partial_j \psi)(x) \partial_{x_i} G_\kappa(x - y) dx dy \\ &= \lim_{\delta \rightarrow 0} \int_{\mathbb{R}^3} f(y) \int_{\mathbb{R}^3 \setminus B_\delta(y)} (\partial_j \psi)(x) \partial_{x_i} G_\kappa(x - y) dx dy \\ &= \lim_{\delta \rightarrow 0} \int_{\mathbb{R}^3} f(y) \left[- \int_{\partial B_\delta(y)} \psi(x) \partial_{x_i} G_\kappa(x - y) e_j \cdot \nu_{y,\delta}(x) dS(x) - \int_{\mathbb{R}^3 \setminus B_\delta(y)} \psi(x) \partial_{x_i} \partial_{x_j} G_\kappa(x - y) dx \right] dy, \end{aligned}$$

where e_j is the j -th unit vector and $\nu_{y,\delta}(x) = (x - y)/\delta$ is the unit outward normal vector in x on the sphere $\partial B_\delta(y)$.

The first part can be simplified explicitly by the transformation $z := (x - y)/\delta$:

$$-\int_{\partial B_\delta(y)} \psi(x) \partial_{x_i} \mathbf{G}_\kappa(x - y) e_j \cdot \nu_{y,\delta}(x) \, dS(x) = \frac{1}{4\pi} \int_{\partial B_1(0)} \psi(\delta z + y) \exp(-\sqrt{\kappa}\delta)(\sqrt{\kappa}\delta + 1) z_i z_j \, dS(z),$$

which converges as $\delta \rightarrow 0$ to $\psi(y)\delta_{ij}/3$.

For the second part, we split the domain of integration $\mathbb{R}^3 \setminus B_\delta(y)$ into two parts:

$$\begin{aligned} \int_{\mathbb{R}^3 \setminus B_\delta(y)} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx \\ = \int_{\mathbb{R}^3 \setminus B_\delta(1)} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx + \int_{\{1 \geq |x-y| \geq \delta\}} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx. \end{aligned}$$

We use integration by parts to insert convenient additional terms:

$$\begin{aligned} \int_{\mathbb{R}^3 \setminus B_\delta(1)} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx + \int_{\{1 \geq |x-y| \geq \delta\}} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx \\ = \int_{\mathbb{R}^3 \setminus B_\delta(1)} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx + \int_{\{1 \geq |x-y| \geq \delta\}} \psi(x) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx \\ - \int_{\{1 \geq |x-y| \geq \delta\}} \psi(y) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dx \\ + \int_{\partial B_1(y)} \psi(y) \partial_{x_j} \mathbf{G}_\kappa(x - y) e_i \cdot \nu_{y,1}(x) \, dS(x) - \int_{\partial B_\delta(y)} \psi(y) \partial_{x_j} \mathbf{G}_\kappa(x - y) e_i \cdot \nu_{y,\delta}(x) \, dS(x). \end{aligned}$$

Now we calculate again explicitly and obtain in the limit $\delta \rightarrow 0$

$$\begin{aligned} \int_{\partial B_1(y)} \psi(y) \partial_{x_j} \mathbf{G}_\kappa(x - y) e_i \cdot \nu_{y,1}(x) \, dS(x) - \int_{\partial B_\delta(y)} \psi(y) \partial_{x_j} \mathbf{G}_\kappa(x - y) e_i \cdot \nu_{y,\delta}(x) \, dS(x) \\ \rightarrow -\frac{1}{3} \delta_{ij} [\exp(-\sqrt{\kappa})(\sqrt{\kappa} + 1) - 1] \psi(y). \end{aligned}$$

In summary, one gets

$$\begin{aligned} -\int_{\mathbb{R}^3} (\partial_j \psi)(x) (\partial_i v)(x) \, dx = \int_{\mathbb{R}^3} \psi(x) \left[\frac{1}{3} \delta_{ij} f(x) \exp(-\sqrt{\kappa})(\sqrt{\kappa} + 1) + \int_{\mathbb{R}^3 \setminus B_1(x)} f(y) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dy \right. \\ \left. + \lim_{\delta \rightarrow 0} \int_{\{1 \geq |x-y| \geq \delta\}} (f(x) - f(y)) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) \, dy \right] \, dx. \end{aligned}$$

From α -Hölder continuity of f , we conclude that, independent of δ ,

$$\mathbb{1}_{\{1 \geq |x-y| \geq \delta\}}(y) |[f(x) - f(y)] \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y)| \leq C|x - y|^{\alpha-3},$$

which is integrable as $\alpha - 3 + 2 > -1$. So using again the dominated convergence theorem, we have, with [Lieb and Loss 2001, Theorem 6.10],

$$(\partial_i \partial_j v)(x) = \frac{1}{3} \delta_{ij} \exp(-\sqrt{\kappa})(\sqrt{\kappa} + 1) + \int_{\mathbb{R}^3 \setminus B_1(x)} f(y) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) dy + \int_{B_1(x)} [f(x) - f(y)] \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) dy. \quad (91)$$

Obviously, the first term in (91) is Hölder-continuous. For the second term in (91), we obtain for all $x, z \in \mathbb{R}^3$, $x \neq z$,

$$\left| \int_{\mathbb{R}^3 \setminus B_1(x)} f(y) \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) dy - \int_{\mathbb{R}^3 \setminus B_1(z)} f(y) \partial_{z_i} \partial_{z_j} \mathbf{G}_\kappa(z - y) dy \right| = \left| \int_{B_1(0)} [f(z - a) - f(x - a)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) da \right|$$

by the transformations $a := x - y$ in the first and $a := z - y$ in the second integrals. From α -Hölder continuity of f , we get the estimate

$$\left| \int_{B_1(0)} [f(z - a) - f(x - a)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) da \right| \leq C|x - z|^\alpha \int_{\mathbb{R}^3 \setminus B_1(x)} |\partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a)| da,$$

where the integral on the right-hand side is finite because $\partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a)$ behaves as $r^{-1} \exp(-r)$ for $r \rightarrow \infty$, which is integrable.

The same integral transformation yields for the third term in (91)

$$\left| \int_{B_1(x)} [f(x) - f(y)] \partial_{x_i} \partial_{x_j} \mathbf{G}_\kappa(x - y) dy - \int_{B_1(z)} [f(z) - f(y)] \partial_{z_i} \partial_{z_j} \mathbf{G}_\kappa(z - y) dy \right| = \left| \int_{B_1(0)} [f(z) - f(z - a) - f(x) + f(x - a)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) da \right|.$$

We now proceed as in [Lieb and Loss 2001] and write $B_1(0) = A \cup B$ with

$$A := \{a : 0 \leq |a| < 4|x - z|\},$$

$$B := \{a : 4|x - z| < |a| < 1\},$$

where $B = \emptyset$ for $|x - z| \geq \frac{1}{4}$, and calculate, using that $|\partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a)| \leq C|a|^{-3}$,

$$\left| \int_A [f(z) - f(z - a) - f(x) + f(x - a)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) da \right| \leq \int_A 2C|a|^{\alpha-3} da = \tilde{C}|x - z|^\alpha.$$

It remains to consider the case $|x - z| < \frac{1}{4}$. One has

$$\left| \int_B [f(z) - f(x)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) da \right| = \left| \int_{\partial B} [f(z) - f(x)] \partial_{a_j} \mathbf{G}_\kappa(a) e_i \cdot \nu(a) dS(a) \right|,$$

and by similar arguments as above,

$$\begin{aligned} & \left| \int_{\partial B} [f(z) - f(x)] \partial_{a_j} \mathbf{G}_\kappa(a) e_i \cdot \nu(a) \, dS(a) \right| \\ &= \frac{1}{3} \delta_{ij} |f(z) - f(x)| \left| \exp(-\sqrt{\kappa})(\sqrt{\kappa} + 1) - \exp(-4\sqrt{\kappa}|x - z|)(4\sqrt{\kappa}|x - z| + 1) \right|. \end{aligned}$$

Note that the real-valued map $[0, \infty) \ni r \mapsto \exp(-\sqrt{\kappa}r)(\sqrt{\kappa}r + 1)$ is monotonically decreasing. This yields

$$\left| \int_B [f(z) - f(x)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) \, da \right| \leq C |z - x|^\alpha.$$

By the transformations $b := x - a - z$ and $b := -a$, we get

$$\begin{aligned} & \left| \int_B [f(x - a) - f(z - a)] \partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a) \, da \right| \\ &= \left| \int_B f(z + b) \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) \, db - \int_D f(b + z) \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b - x + z) \, db \right| \quad (92) \end{aligned}$$

with $D := \{b : 4|x - z| < |b - x + z| < 1\}$.

Note that

$$\int_B \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) \, db = \int_D \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b - x + z) \, db.$$

This enables us to rewrite (92) as

$$\begin{aligned} & \left| \int_B f(z + b) \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) \, db - \int_D f(b + z) \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b - x + z) \, db \right| \\ &= \left| \int_B [f(z + b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) \, db - \int_D [f(z + b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b - x + z) \, db \right|. \quad (93) \end{aligned}$$

We consider (93) separately on the sets $B \cap D$, $B \setminus D$ and $D \setminus B$.

Note that, by the triangular inequality, $B \cap D \subset \{b : 3|x - z| < |b| < 1 + |x - z|\}$ and by Taylor's theorem

$$(\partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa)(b) - (\partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa)(b - x + z) = \sum_{k=1}^3 (\partial_k \partial_i \partial_j \mathbf{G}_\kappa)(b^*) (x_k - z_k)$$

for some $b^* = b - \beta(x - z)$ with $\beta \in (0, 1)$. Therefore, one has, by the triangular inequality, $|b^*| \geq |b| - \beta|x - z| \geq (1 - \frac{1}{3}\beta)|b| \geq \frac{2}{3}|b|$ on $B \cap D$ and consequently

$$|(\partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa)(b) - (\partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa)(b - x + z)| \leq C |b^*|^{-4} |x - z| \leq \tilde{C} |b|^{-4} |x - z|.$$

This allows us to estimate

$$\begin{aligned} \left| \int_{B \cap D} [f(z+b) - f(z)] [\partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) - \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b-x+z)] db \right| \\ \leq C|x-z| \int_{3|x-z|}^{1+|x-z|} r^{-4+\alpha+2} dr \\ \leq \frac{C|x-z|}{1-\alpha} [(3|x-z|)^{\alpha-1} - (1+|x-z|)^{\alpha-1}] \frac{\tilde{C}}{1-\alpha} |x-z|^\alpha. \end{aligned}$$

For the remaining terms, we split up as in [Lieb and Loss 2001]:

$$\begin{aligned} B \setminus D &\subset E \cup G, \\ D \setminus B &\subset E' \cup G', \end{aligned}$$

where

$$\begin{aligned} E &:= \{b : 4|x-z| < |b| \leq 5|x-z|\}, \\ G &:= \{b : 1 - |x-z| \leq |b| < 1\}, \\ E' &:= \{b : 4|x-z| < |b-x+z| \leq 5|x-z|\}, \\ G' &:= \{b : 1 - |x-z| \leq |b-x+z| < 1\}. \end{aligned}$$

Consider at first the real-valued map $[0, \frac{1}{4}] \ni s \mapsto (1-s)^\beta$ for arbitrary $\beta > 0$. Obviously, it is continuously differentiable and therefore α -Hölder continuous because its domain of definition is compact. Hence, the following holds for all $0 \leq s \leq \frac{1}{4}$:

$$1 - (1-s)^\beta = (1-0)^\beta - (1-s)^\beta \leq Cs^\alpha. \tag{94}$$

Now, we estimate the integral on $B \setminus D$, where we use again the estimate $|\partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a)| \leq C|a|^{-3}$:

$$\begin{aligned} \left| \int_{B \setminus D} [f(z+b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b) db \right| &\leq C \left(\int_{4|x-z|}^{5|x-z|} r^{\alpha-3+2} dr + \int_{1-|x-z|}^1 r^{\alpha-3+2} dr \right) \\ &= \frac{C}{\alpha} [(5|x-z|)^\alpha - (4|x-z|)^\alpha + 1 - (1-|x-z|)^\alpha] \leq \frac{C}{\alpha} (5^\alpha + \tilde{C}) |x-z|^\alpha, \end{aligned}$$

where we have used (94) for $\beta := \alpha$ in the last step.

For the remaining integral on $D \setminus B$, we consider the domains E' and G' separately and note at first that, using the triangular inequality, $E' \subset \{0 < |b| \leq 6|x-z|\}$. Subsequently, this yields that $|b-x+z|^{-3} < (4|x-z|)^{-3} \leq C|b|^{-3}$ on E' . Hence, by the estimate $|\partial_{a_i} \partial_{a_j} \mathbf{G}_\kappa(a)| \leq C|a|^{-3}$,

$$\int_{E'} |[f(z+b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b-x+z)| db \leq C \int_0^{6|x-z|} r^{\alpha-3+2} dr = \tilde{C} |x-z|^\alpha.$$

On G' , one has $|b-x+z| \geq 1 - |x-z| > \frac{3}{4}$. Consequently, it holds that

$$\begin{aligned} \int_{G'} |[f(z+b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b-x+z)| db &\leq C \left(\frac{3}{4}\right)^{-3} \int_{1-|x-z|}^1 r^{\alpha+2} dr \\ &= \tilde{C} [1 - (1-|x-z|)^{3+\alpha}] \leq \tilde{C} |x-z|^\alpha, \end{aligned}$$

where we have used (94) for $\beta := 3 + \alpha$ in the last step. Together,

$$\left| \int_{D \setminus B} [f(z+b) - f(z)] \partial_{b_i} \partial_{b_j} \mathbf{G}_\kappa(b-x+z) db \right| \leq C|x-z|^\alpha,$$

and the assertion is proved. \square

References

- [Agueh 2008] M. Agueh, “Rates of decay to equilibria for p -Laplacian type equations”, *Nonlinear Anal.* **68**:7 (2008), 1909–1927. MR 2009e:35117 Zbl 1185.35017
- [Ambrosio et al. 2008] L. Ambrosio, N. Gigli, and G. Savaré, *Gradient flows in metric spaces and in the space of probability measures*, 2nd ed., Birkhäuser, Basel, 2008. MR 2009h:49002 Zbl 1145.35001
- [Biler et al. 2011] P. Biler, L. Corrias, and J. Dolbeault, “Large mass self-similar solutions of the parabolic-parabolic Keller–Segel model of chemotaxis”, *J. Math. Biol.* **63**:1 (2011), 1–32. MR 2012c:35431 Zbl 1230.92011
- [Blanchet 2013] A. Blanchet, “A gradient flow approach to the Keller–Segel systems”, preprint, 2013, Available at http://www.tse-fr.eu/sites/default/files/medias/doc/by/blanchet/dec_2010/b4.pdf.
- [Blanchet and Laurençot 2013] A. Blanchet and P. Laurençot, “The parabolic-parabolic Keller–Segel system with critical diffusion as a gradient flow in \mathbb{R}^d , $d \geq 3$ ”, *Comm. Partial Differential Equations* **38**:4 (2013), 658–686. MR 3040679 Zbl 1282.35202
- [Blanchet et al. 2008] A. Blanchet, V. Calvez, and J. A. Carrillo, “Convergence of the mass-transport steepest descent scheme for the subcritical Patlak–Keller–Segel model”, *SIAM J. Numer. Anal.* **46**:2 (2008), 691–721. MR 2009a:35113 Zbl 1205.65332
- [Blanchet et al. 2009] A. Blanchet, J. A. Carrillo, and P. Laurençot, “Critical mass for a Patlak–Keller–Segel model with degenerate diffusion in higher dimensions”, *Calc. Var. Partial Differential Equations* **35**:2 (2009), 133–168. MR 2010e:35149 Zbl 1172.35035
- [Blanchet et al. 2012] A. Blanchet, E. A. Carlen, and J. A. Carrillo, “Functional inequalities, thick tails and asymptotics for the critical mass Patlak–Keller–Segel model”, *J. Funct. Anal.* **262**:5 (2012), 2142–2230. MR 2876403 Zbl 1237.35155
- [Blanchet et al. 2014] A. Blanchet, J. A. Carrillo, D. Kinderlehrer, M. Kowalczyk, P. Laurençot, and S. Lisini, “A hybrid variational principle for the Keller–Segel system in \mathbb{R}^2 ”, preprint, 2014. arXiv 1407.5562v1
- [Calvez and Carrillo 2012] V. Calvez and J. A. Carrillo, “Refined asymptotics for the subcritical Keller–Segel system and related functional inequalities”, *Proc. Amer. Math. Soc.* **140**:10 (2012), 3515–3530. MR 2929020 Zbl 1277.35051
- [Calvez and Corrias 2008] V. Calvez and L. Corrias, “The parabolic-parabolic Keller–Segel model in \mathbb{R}^2 ”, *Commun. Math. Sci.* **6**:2 (2008), 417–447. MR 2009m:35518 Zbl 1149.35360
- [Carrapatoso and Mischler 2014] K. Carrapatoso and S. Mischler, “Uniqueness and long time asymptotic for the parabolic-parabolic Keller–Segel equation”, preprint, 2014. arXiv 1406.6006v2
- [Carrillo and Toscani 2000] J. A. Carrillo and G. Toscani, “Asymptotic L^1 -decay of solutions of the porous medium equation to self-similarity”, *Indiana Univ. Math. J.* **49**:1 (2000), 113–142. MR 2001j:35155 Zbl 0963.35098
- [Carrillo et al. 2003] J. A. Carrillo, R. J. McCann, and C. Villani, “Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates”, *Rev. Mat. Iberoamericana* **19**:3 (2003), 971–1018. MR 2005a:35126 Zbl 1073.35127
- [Carrillo et al. 2006a] J. A. Carrillo, M. Di Francesco, and G. Toscani, “Intermediate asymptotics beyond homogeneity and self-similarity: long time behavior for $u_t = \Delta \phi(u)$ ”, *Arch. Ration. Mech. Anal.* **180**:1 (2006), 127–149. MR 2006k:35131 Zbl 1096.35015
- [Carrillo et al. 2006b] J. A. Carrillo, R. J. McCann, and C. Villani, “Contractions in the 2-Wasserstein length space and thermalization of granular media”, *Arch. Ration. Mech. Anal.* **179**:2 (2006), 217–263. MR 2006j:76121 Zbl 1082.76105
- [Carrillo et al. 2011] J. A. Carrillo, M. Di Francesco, A. Figalli, T. Laurent, and D. Slepčev, “Global-in-time weak measure solutions and finite-time aggregation for nonlocal interaction equations”, *Duke Math. J.* **156**:2 (2011), 229–271. MR 2012c:35447 Zbl 1215.35045

- [Corrias and Perthame 2008] L. Corrias and B. Perthame, “Asymptotic decay for the solutions of the parabolic-parabolic Keller–Segel chemotaxis system in critical spaces”, *Math. Comput. Modelling* **47**:7–8 (2008), 755–764. MR 2009c:35200 Zbl 1134.92006
- [Di Francesco and Rosado 2008] M. Di Francesco and J. Rosado, “Fully parabolic Keller–Segel model for chemotaxis with prevention of overcrowding”, *Nonlinearity* **21**:11 (2008), 2715–2730. MR 2009m:35196 Zbl 1157.35398
- [Eisenbach 2004] M. Eisenbach, *Chemotaxis*, Imperial College, London, 2004.
- [Giacomelli and Otto 2001] L. Giacomelli and F. Otto, “Variational formulation for the lubrication approximation of the Hele–Shaw flow”, *Calc. Var. Partial Differential Equations* **13**:3 (2001), 377–403. MR 2003a:76046 Zbl 1086.35004
- [Gianazza et al. 2009] U. Gianazza, G. Savaré, and G. Toscani, “The Wasserstein gradient flow of the Fisher information and the quantum drift-diffusion equation”, *Arch. Ration. Mech. Anal.* **194**:1 (2009), 133–220. MR 2010i:35371 Zbl 1223.35264
- [Hillen and Painter 2009] T. Hillen and K. J. Painter, “A user’s guide to PDE models for chemotaxis”, *J. Math. Biol.* **58**:1–2 (2009), 183–217. MR 2009m:92017 Zbl 1161.92003
- [Horstmann 2003] D. Horstmann, “From 1970 until present: the Keller–Segel model in chemotaxis and its consequences, I”, *Jahresber. Deutsch. Math.-Verein.* **105**:3 (2003), 103–165. MR 2005f:35163 Zbl 1071.35001
- [Jordan et al. 1998] R. Jordan, D. Kinderlehrer, and F. Otto, “The variational formulation of the Fokker–Planck equation”, *SIAM J. Math. Anal.* **29**:1 (1998), 1–17. MR 2000b:35258 Zbl 0915.35120
- [Keller and Segel 1970] E. F. Keller and L. A. Segel, “Initiation of slime mold aggregation viewed as an instability”, *J. Theor. Biol.* **26**:3 (1970), 399–415. Zbl 1170.92306
- [Kozono and Sugiyama 2009] H. Kozono and Y. Sugiyama, “Global strong solution to the semi-linear Keller–Segel system of parabolic-parabolic type with small data in scale invariant spaces”, *J. Differential Equations* **247**:1 (2009), 1–32. MR 2010i:35147 Zbl 1207.35178
- [Lapidus and Schiller 1976] I. R. Lapidus and R. Schiller, “Model for the chemotactic response of a bacterial population”, *Biophys. J.* **16**:7 (1976), 779–789.
- [Laurençot and Matioc 2013] P. Laurençot and B.-V. Matioc, “A gradient flow approach to a thin film approximation of the Muskat problem”, *Calc. Var. Partial Differential Equations* **47**:1–2 (2013), 319–341. MR 3044141 Zbl 1264.35129
- [Lieb and Loss 2001] E. H. Lieb and M. Loss, *Analysis*, 2nd ed., Graduate Studies in Mathematics **14**, Amer. Math. Soc., Providence, RI, 2001. MR 2001i:00001 Zbl 0966.26002
- [López-Gómez et al. 2013] J. López-Gómez, T. Nagai, and T. Yamada, “The basin of attraction of the steady-states for a chemotaxis model in \mathbb{R}^2 with critical mass”, *Arch. Ration. Mech. Anal.* **207**:1 (2013), 159–184. MR 3004770 Zbl 06148937
- [Luckhaus and Sugiyama 2006] S. Luckhaus and Y. Sugiyama, “Large time behavior of solutions in super-critical cases to degenerate Keller–Segel systems”, *M2AN Math. Model. Numer. Anal.* **40**:3 (2006), 597–621. MR 2007e:35127 Zbl 1113.35028
- [Luckhaus and Sugiyama 2007] S. Luckhaus and Y. Sugiyama, “Asymptotic profile with the optimal convergence rate for a parabolic equation of chemotaxis in super-critical cases”, *Indiana Univ. Math. J.* **56**:3 (2007), 1279–1297. MR 2008k:35272 Zbl 1118.35006
- [Matthes et al. 2009] D. Matthes, R. J. McCann, and G. Savaré, “A family of nonlinear fourth order equations of gradient flow type”, *Comm. Partial Differential Equations* **34**:10–12 (2009), 1352–1397. MR 2011e:35149 Zbl 1187.35131
- [Mizoguchi 2013] N. Mizoguchi, “Global existence for the Cauchy problem of the parabolic-parabolic Keller–Segel system on the plane”, *Calc. Var. Partial Differential Equations* **48**:3–4 (2013), 491–505. MR 3116019 Zbl 1288.35286
- [Murray 2003] J. D. Murray, *Mathematical biology, II: Spatial models and biomedical applications*, 3rd ed., Interdisciplinary Applied Mathematics **18**, Springer, New York, 2003. MR 2004b:92001 Zbl 1006.92002
- [Nagai and Senba 1998] T. Nagai and T. Senba, “Global existence and blow-up of radial solutions to a parabolic-elliptic system of chemotaxis”, *Adv. Math. Sci. Appl.* **8**:1 (1998), 145–156. MR 2000a:35241 Zbl 0902.35010
- [Nagai et al. 2003] T. Nagai, R. Syukuinn, and M. Umesako, “Decay properties and asymptotic profiles of bounded solutions to a parabolic system of chemotaxis in \mathbb{R}^n ”, *Funkcial. Ekvac.* **46**:3 (2003), 383–407. MR 2005h:35144 Zbl 02112808
- [Otto 2001] F. Otto, “The geometry of dissipative evolution equations: the porous medium equation”, *Comm. Partial Differential Equations* **26**:1–2 (2001), 101–174. MR 2002j:35180 Zbl 0984.35089

- [Perthame 2007] B. Perthame, *Transport equations in biology*, Birkhäuser, Basel, 2007. MR 2007j:35004 Zbl 1185.92006
- [Post 1999] K. Post, *A non-linear parabolic system modeling chemotaxis with sensitivity functions*, Ph.D. thesis, Humboldt-Universität zu Berlin, 1999, Available at <http://edoc.hu-berlin.de/docviews/abstract.php?id=26314>.
- [Segel 1977] L. Segel, “A theoretical study of receptor mechanisms in bacterial chemotaxis”, *SIAM J. Appl. Math.* **32**:3 (1977), 653–665. Zbl 0356.92009
- [Senba and Suzuki 2006] T. Senba and T. Suzuki, “A quasi-linear parabolic system of chemotaxis”, *Abstr. Appl. Anal.* **2006** (2006), 23061. MR 2006j:35121 Zbl 1134.35059
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series **30**, Princeton University, 1970. MR 44 #7280 Zbl 0207.13501
- [Sugiyama 2006] Y. Sugiyama, “Global existence in sub-critical cases and finite time blow-up in super-critical cases to degenerate Keller–Segel systems”, *Differential Integral Equations* **19**:8 (2006), 841–876. MR 2008h:35134 Zbl 1212.35240
- [Sugiyama 2007] Y. Sugiyama, “Time global existence and asymptotic behavior of solutions to degenerate quasi-linear parabolic systems of chemotaxis”, *Differential Integral Equations* **20**:2 (2007), 133–180. MR 2008a:35169 Zbl 1212.35241
- [Sugiyama and Kunii 2006] Y. Sugiyama and H. Kunii, “Global existence and decay properties for a degenerate Keller–Segel model with a power factor in drift term”, *J. Differential Equations* **227**:1 (2006), 333–364. MR 2008j:35103 Zbl 1102.35046
- [Villani 2003] C. Villani, *Topics in optimal transportation*, Graduate Studies in Mathematics **58**, Amer. Math. Soc., Providence, RI, 2003. MR 2004e:90003 Zbl 1106.90001
- [Winkler 2010] M. Winkler, “Absence of collapse in a parabolic chemotaxis system with signal-dependent sensitivity”, *Math. Nachr.* **283**:11 (2010), 1664–1673. MR 2011k:35090 Zbl 1205.35037
- [Yamada 2011] T. Yamada, “Improvement of convergence rates for a parabolic system of chemotaxis in the whole space”, *Math. Methods Appl. Sci.* **34**:17 (2011), 2103–2124. MR 2012k:35217 Zbl 1242.35054
- [Zeidler 1990] E. Zeidler, *Nonlinear functional analysis and its applications, II/B: Nonlinear monotone operators*, Springer, New York, 1990. MR 91b:47002 Zbl 0684.47029
- [Zinsl 2014] J. Zinsl, “Existence of solutions for a nonlinear system of parabolic equations with gradient flow structure”, *Monats. Math.* **174**:4 (2014), 653–679. MR 3233116 Zbl 1302.35202

Received 12 May 2014. Revised 25 Nov 2014. Accepted 9 Jan 2015.

JONATHAN ZINSL: zinsl@ma.tum.de

Zentrum für Mathematik, Technische Universität München, Boltzmannstraße 3, 85747 Garching, Germany

DANIEL MATTHES: matthes@ma.tum.de

Zentrum für Mathematik, Technische Universität München, Boltzmannstraße 3, 85747 Garching, Germany

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 2 2015

Smooth parametric dependence of asymptotics of the semiclassical focusing NLS SERGEY BELOV and STEPHANOS VENAKIDES	257
Tunnel effect for semiclassical random walks JEAN-FRANÇOIS BONY, FRÉDÉRIC HÉRAU and LAURENT MICHEL	289
Traveling wave solutions in a half-space for boundary reactions XAVIER CABRÉ, NEUS CÓNSUL and JOSÉ V. MANDÉ	333
Locally conformally flat ancient Ricci flows GIOVANNI CATINO, CARLO MANTEGAZZA and LORENZO MAZZIERI	365
Motion of three-dimensional elastic films by anisotropic surface diffusion with curvature regularization IRENE FONSECA, NICOLA FUSCO, GIOVANNI LEONI and MASSIMILIANO MORINI	373
Exponential convergence to equilibrium in a coupled gradient flow system modeling chemotaxis JONATHAN ZINSL and DANIEL MATTHES	425
Scattering for the radial 3D cubic wave equation BENJAMIN DODSON and ANDREW LAWRIE	467
Counterexamples to the well posedness of the Cauchy problem for hyperbolic systems FERRUCCIO COLOMBINI and GUY MÉTIVIER	499



2157-5045(2015)8:2;1-C