

ANALYSIS & PDE

Volume 8

No. 8

2015

GIULIO GALISE, CYRIL IMBERT AND RÉGIS MONNEAU

**A JUNCTION CONDITION BY SPECIFIED HOMOGENIZATION
AND APPLICATION TO TRAFFIC LIGHTS**



A JUNCTION CONDITION BY SPECIFIED HOMOGENIZATION AND APPLICATION TO TRAFFIC LIGHTS

GIULIO GALISE, CYRIL IMBERT AND RÉGIS MONNEAU

Given a coercive Hamiltonian which is quasiconvex with respect to the gradient variable and periodic with respect to time and space, at least “far away from the origin”, we consider the solution of the Cauchy problem of the corresponding Hamilton–Jacobi equation posed on the real line. Compact perturbations of coercive periodic quasiconvex Hamiltonians enter into this framework, for example. We prove that the rescaled solution converges towards the solution of the expected effective Hamilton–Jacobi equation, but whose “flux” at the origin is “limited” in a sense made precise by Imbert and Monneau. In other words, the homogenization of such a Hamilton–Jacobi equation yields to supplement the expected homogenized Hamilton–Jacobi equation with a junction condition at the single discontinuous point of the effective Hamiltonian. We also illustrate possible applications of such a result by deriving, for a traffic flow problem, the effective flux limiter generated by the presence of a finite number of traffic lights on an ideal road. We also provide meaningful qualitative properties of the effective limiter.

1. Introduction	1891
2. Proof of convergence	1899
3. Homogenized Hamiltonians	1901
4. Truncated cell problems	1906
5. Proof of Theorem 1.12	1912
Appendix: Proofs of some technical results	1924
Acknowledgements	1928
References	1928

1. Introduction

Setting of the general problem. This article is concerned with the study of the limit of the solution $u^\varepsilon(t, x)$ of the equation

$$u_t^\varepsilon + H\left(\frac{t}{\varepsilon}, \frac{x}{\varepsilon}, u_x^\varepsilon\right) = 0 \quad \text{for } (t, x) \in (0, T) \times \mathbb{R} \tag{1}$$

subject to the initial condition

$$u^\varepsilon(0, x) = u_0(x) \quad \text{for } x \in \mathbb{R} \tag{2}$$

for a Hamiltonian H satisfying the following assumptions:

MSC2010: 35B27, 49L25, 35F21.

Keywords: Hamilton–Jacobi equations, quasiconvex Hamiltonians, homogenization, junction condition, flux-limited solution, viscosity solution.

(A0) Continuity: $H : \mathbb{R}^3 \rightarrow \mathbb{R}$ is continuous.

(A1) Time periodicity: For all $k \in \mathbb{Z}$ and $(t, x, p) \in \mathbb{R}^3$,

$$H(t+k, x, p) = H(t, x, p).$$

(A2) Uniform modulus of continuity in time: There exists a modulus of continuity ω such that, for all $t, s, x, p \in \mathbb{R}$,

$$H(t, x, p) - H(s, x, p) \leq \omega(|t-s|(1 + \max(H(s, x, p), 0))).$$

(A3) Uniform coercivity: $\lim_{|q| \rightarrow +\infty} H(t, x, q) = +\infty$

uniformly with respect to (t, x) .

(A4) Quasiconvexity of H for large x : There exists some $\rho_0 > 0$ such that, for all $x \in \mathbb{R} \setminus (-\rho_0, \rho_0)$, there exists a continuous map $t \mapsto p^0(t, x)$ such that

$$\begin{cases} H(t, x, \cdot) & \text{is nonincreasing in } (-\infty, p^0(t, x)), \\ H(t, x, \cdot) & \text{is nondecreasing in } (p^0(t, x), +\infty). \end{cases}$$

(A5) Left and right Hamiltonians: There exist two Hamiltonians $H_\alpha(t, x, p)$, $\alpha = L, R$, such that

$$\begin{cases} H(t, x+k, p) - H_L(t, x, p) \rightarrow 0 & \text{as } \mathbb{Z} \ni k \rightarrow -\infty, \\ H(t, x+k, p) - H_R(t, x, p) \rightarrow 0 & \text{as } \mathbb{Z} \ni k \rightarrow +\infty, \end{cases}$$

uniformly with respect to $(t, x, p) \in [0, 1]^2 \times \mathbb{R}$ and, for all $k, j \in \mathbb{Z}$, $(t, x, p) \in \mathbb{R}^3$ and $\alpha \in \{L, R\}$,

$$H_\alpha(t+k, x+j, p) = H_\alpha(t, x, p).$$

We have to impose some condition in order to ensure that effective Hamiltonians \bar{H}_α are quasiconvex; indeed, we will see that the effective equation should be solved with *flux-limited solutions*, recently introduced by Imbert and Monneau [2013]; such a theory relies on the quasiconvexity of the Hamiltonians.

(B-i) Quasiconvexity of the left and right Hamiltonians: H_α , $\alpha = L, R$, does not depend on time and there exists p_α^0 (independent of (t, x)) such that

$$\begin{cases} H_\alpha(x, \cdot) & \text{is nonincreasing on } (-\infty, p_\alpha^0), \\ H_\alpha(x, \cdot) & \text{is nondecreasing on } (p_\alpha^0, +\infty). \end{cases}$$

(B-ii) Convexity of the left and right Hamiltonians: For each $\alpha = L, R$ and for all $(t, x) \in \mathbb{R} \times \mathbb{R}$, the map $p \mapsto H_\alpha(t, x, p)$ is convex.

Example 1.1. A simple example of such a Hamiltonian is

$$H(t, x, p) = |p| - f(t, x)$$

with a continuous function f satisfying $f(t+1, x) = f(t, x)$ and $f(t, x) \rightarrow 0$ as $|x| \rightarrow +\infty$ uniformly with respect to $t \in \mathbb{R}$.

Main results. Our main result is concerned with the limit of the solution u^ε of (1)–(2). It joins part of the huge literature dealing with homogenization of Hamilton–Jacobi equation, starting with the pioneering work of Lions, Papanicolaou and Varadhan [Lions et al. 1986]. In particular, we need to use the perturbed test function introduced by Evans [1989]. As pointed out to us by the referee, there are few papers dealing with Hamiltonians that depend on time, which implies in particular that so-called correctors also depend on time. The reader is referred to [Barles and Souganidis 2000; Bernard and Roquejoffre 2004] for the large time behaviour and to [Forcadel et al. 2009a; 2009b; 2012] for homogenization results. This limit satisfies an effective Hamilton–Jacobi equation posed on the real line whose Hamiltonian is discontinuous. More precisely, the effective Hamiltonian equals the one which is expected (see (A5)) in $(-\infty; 0)$ and $(0; +\infty)$; in particular, it is discontinuous in the space variable (piecewise constant, in fact). In order to get a unique solution, a flux limiter should be identified [Imbert and Monneau 2013], henceforth abbreviated [IM].

Homogenized Hamiltonians and effective flux limiter. The homogenized left and right Hamiltonians are classically determined by the study of some “cell problems”.

Proposition 1.2 (homogenized left and right Hamiltonians). *Assume (A0)–(A5) and either (B-i) or (B-ii). Then, for every $p \in \mathbb{R}$ and $\alpha = L, R$, there exists a unique $\lambda \in \mathbb{R}$ such that there exists a bounded solution v^α of*

$$\begin{cases} v_t^\alpha + H_\alpha(t, x, p + v_x^\alpha) = \lambda & \text{in } \mathbb{R} \times \mathbb{R}, \\ v^\alpha \text{ is } \mathbb{Z}^2\text{-periodic.} \end{cases} \tag{3}$$

If $\bar{H}_\alpha(p)$ denotes such a λ , then the map $p \mapsto \bar{H}_\alpha(p)$ is continuous, coercive and quasiconvex.

Remark 1.3. We recall that a function \bar{H}_α is quasiconvex if the sets $\{\bar{H}_\alpha \leq \lambda\}$ are convex for all $\lambda \in \mathbb{R}$. If \bar{H}_α is also coercive, then \bar{p}_α^0 denotes in proofs some $p \in \operatorname{argmin} \bar{H}_\alpha$.

The effective flux limiter \bar{A} is the smallest $\lambda \in \mathbb{R}$ for which there exists a solution w of the global-in-time Hamilton–Jacobi equation

$$\begin{cases} w_t + H(t, x, w_x) = \lambda, & (t, x) \in \mathbb{R} \times \mathbb{R}, \\ w \text{ is 1-periodic in } t. \end{cases} \tag{4}$$

Theorem 1.4 (effective flux limiter). *Assume (A0)–(A5) and either (B-i) or (B-ii). The set*

$$E = \{\lambda \in \mathbb{R} : \text{there is a subsolution } w \text{ of (4)}\}$$

is nonempty and bounded from below. Moreover, if \bar{A} denotes the infimum of E , then

$$\bar{A} \geq A_0 := \max_{\alpha=L,R} (\min \bar{H}_\alpha). \tag{5}$$

Remark 1.5. We will see below (Theorem 4.6) that the infimum is in fact a minimum: there exists a global corrector which, in particular, can be rescaled properly.

We can now define the effective junction condition:

Definition 1.6. The effective junction function $F_{\bar{A}}$ is defined by

$$F_{\bar{A}}(p_L, p_R) := \max(\bar{A}, \bar{H}_L^+(p_L), \bar{H}_R^-(p_R)),$$

where

$$\bar{H}_\alpha^-(p) = \begin{cases} \bar{H}_\alpha(p) & \text{if } p < \bar{p}_\alpha^0, \\ \bar{H}_\alpha(\bar{p}_\alpha^0) & \text{if } p \geq \bar{p}_\alpha^0, \end{cases} \quad \text{and} \quad \bar{H}_\alpha^+(p) = \begin{cases} \bar{H}_\alpha(\bar{p}_\alpha^0) & \text{if } p \leq \bar{p}_\alpha^0, \\ \bar{H}_\alpha(p) & \text{if } p > \bar{p}_\alpha^0, \end{cases}$$

where $\bar{p}_\alpha^0 \in \operatorname{argmin} \bar{H}_\alpha$.

The convergence result. Our main result is the following theorem:

Theorem 1.7 (junction condition by homogenization). *Assume (A0)–(A5) and either (B-i) or (B-ii). Assume that the initial datum u_0 is Lipschitz continuous and, for $\varepsilon > 0$, let u^ε be the solution of (1)–(2). Then u^ε converges locally uniformly to the unique flux-limited solution u^0 of*

$$\begin{cases} u_t^0 + \bar{H}_L(u_x^0) = 0, & t > 0, x < 0, \\ u_t^0 + \bar{H}_R(u_x^0) = 0, & t > 0, x > 0, \\ u_t^0 + F_{\bar{A}}(u_x^0(t, 0^-), u_x^0(t, 0^+)) = 0, & t > 0, x = 0, \end{cases} \tag{6}$$

subject to the initial condition (2).

Remark 1.8. The notion of flux-limited solution for (6) was introduced in [IM].

This theorem asserts in particular that the slopes of the limit solution at the origin are characterized by the effective flux limiter \bar{A} . Its proof relies on the construction of a global “corrector”, i.e., a solution of (4) which is close to an appropriate V-shaped function after rescaling. This latter condition is necessary so that the slopes at infinity of the corrector fit the expected slopes of the solution of the limit problem at the origin. Here is a precise statement:

Theorem 1.9 (existence of a global corrector for the junction). *Assume (A0)–(A5) and either (B-i) or (B-ii). There exists a solution w of (4) with $\lambda = \bar{A}$ such that the function*

$$w^\varepsilon(t, x) = \varepsilon w(\varepsilon^{-1}t, \varepsilon^{-1}x)$$

converges locally uniformly (along a subsequence $\varepsilon_n \rightarrow 0$) towards a function $W = W(x)$ which satisfies $W(0) = 0$ and

$$\hat{p}_R x 1_{\{x>0\}} + \hat{p}_L x 1_{\{x<0\}} \geq W(x) \geq \bar{p}_R x 1_{\{x>0\}} + \bar{p}_L x 1_{\{x<0\}}, \tag{7}$$

where

$$\begin{cases} \bar{p}_R = \min E_R, \\ \hat{p}_R = \max E_R, \end{cases} \quad \text{with} \quad E_R := \{p \in \mathbb{R} : \bar{H}_R^+(p) = \bar{H}_R(p) = \bar{A}\}, \tag{8}$$

$$\begin{cases} \bar{p}_L = \max E_L, \\ \hat{p}_L = \min E_L, \end{cases} \quad \text{with} \quad E_L := \{p \in \mathbb{R} : \bar{H}_L^-(p) = \bar{H}_L(p) = \bar{A}\}. \tag{9}$$

The construction of this global corrector is the reason why homogenization is referred to as being “specified”; see also related results on p. 1897. As a matter of fact, we will prove a stronger result; see Theorem 4.6.

Extension: application to traffic lights. The techniques developed to prove Theorem 1.7 allow us to deal with a different situation inspired by traffic flow problems. As explained in [Imbert et al. 2013], such problems are related to the study of some Hamilton–Jacobi equations. Theorem 1.12 below is motivated by aiming to figuring out how the traffic flow on an ideal (infinite, straight) road is modified by the presence of a finite number of traffic lights.

We can consider a Hamilton–Jacobi equation whose Hamiltonian does not depend on (t, x) for x outside a (small) interval of the form $N_\varepsilon = (b_1\varepsilon, b_N\varepsilon)$, and is piecewise constant with respect to x in $(b_1\varepsilon, b_N\varepsilon)$. At space discontinuities, junction conditions are imposed with ε -time-periodic flux limiters. The limit solution satisfies the equation after the “neighbourhood” N_ε disappears. We will see that the equation keeps memory of what happened there through a flux limiter at the origin $x = 0$.

Let us be more precise now. We are given, for $N \geq 1$ and $K \in \mathbb{N}$, a finite number of junction points $-\infty = b_0 < b_1 < b_2 < \dots < b_N < b_{N+1} = +\infty$ and times $0 = \tau_0 < \tau_1 < \dots < \tau_K < 1 = \tau_{K+1}$. For $\alpha \in \{0, \dots, N\}$, ℓ_α denotes $b_{\alpha+1} - b_\alpha$. Note that $\ell_\alpha = +\infty$ for $\alpha = 0, N$.

We then consider the solution u^ε of (1) where the Hamiltonian H satisfies the following conditions:

(C1) The Hamiltonian is given by

$$H(t, x, p) = \begin{cases} \bar{H}_\alpha(p) & \text{if } b_\alpha < x < b_{\alpha+1}, \\ \max(\bar{H}_{\alpha-1}^+(p^-), \bar{H}_\alpha^-(p^+), a_\alpha(t)) & \text{if } x = b_\alpha, \alpha \neq 0. \end{cases}$$

(C2) The Hamiltonians \bar{H}_α for $\alpha = 0, \dots, N$ are continuous, coercive and quasiconvex.

(C3) The flux limiters a_α for $\alpha = 1, \dots, N$, and $i = 0, \dots, K$, satisfy

$$a_\alpha(s + 1) = a_\alpha(s) \quad \text{with} \quad a_\alpha(s) = A_\alpha^i \quad \text{for all } s \in [\tau_i, \tau_{i+1})$$

with $(A_\alpha^i)_{\alpha=1, \dots, N}^{i=0, \dots, K}$ satisfying $A_\alpha^i \geq \max_{\beta=\alpha-1, \alpha}(\min \bar{H}_\beta)$.

Remark 1.10. The Hamiltonians outside N_ε are denoted by \bar{H}_α instead of H_α in order to emphasize that they do not depend on time and space.

Remark 1.11. In view of the literature in traffic modelling, the Hamiltonians could be assumed to be convex. But we prefer to stick to the quasiconvex framework since it seems to us that it is the natural one (in view of [IM]).

The equation is supplemented with the initial condition

$$u^\varepsilon(0, x) = U_0^\varepsilon(x) \quad \text{for } x \in \mathbb{R} \tag{10}$$

with

$$U_0^\varepsilon \text{ equi-Lipschitz continuous} \quad \text{and} \quad U_0^\varepsilon \rightarrow u_0 \text{ locally uniformly.} \tag{11}$$

Then the following convergence result holds true:

Theorem 1.12 (time homogenization of traffic lights). *Assume (C1)–(C3) and (11). Let u^ε be the solution of (1) and (10) for all $\varepsilon > 0$. Then:*

- (i) *Homogenization: There exists some $\bar{A} \in \mathbb{R}$ such that u^ε converges locally uniformly as ε tends to zero towards the unique viscosity solution u^0 of (6) and (2) with*

$$\bar{H}_L := \bar{H}_0, \quad \bar{H}_R := \bar{H}_N.$$

- (ii) *Qualitative properties of \bar{A} : For $\alpha = 1, \dots, N$, $\langle a_\alpha \rangle$ denotes $\int_0^1 a_\alpha(s) ds$. The effective limiter \bar{A} satisfies the following properties:*

- For all α , \bar{A} is nonincreasing with respect to ℓ_α .
- For $N = 1$,

$$\bar{A} = \langle a_1 \rangle. \tag{12}$$

- For $N \geq 1$,

$$\bar{A} \geq \max_{\alpha=1, \dots, N} \langle a_\alpha \rangle. \tag{13}$$

- For $N \geq 2$, there exists a critical distance $d_0 \geq 0$ such that

$$\bar{A} = \max_{\alpha=1, \dots, N} \langle a_\alpha \rangle \quad \text{if} \quad \min_{\alpha} \ell_\alpha \geq d_0; \tag{14}$$

this distance d_0 only depends on $\max_{\alpha=1, \dots, N} \|a_\alpha\|_\infty$, $\max_{\alpha=1, \dots, N} \langle a_\alpha \rangle$ and the \bar{H}_α .

- We have

$$\bar{A} \rightarrow \langle \bar{a} \rangle \quad \text{as} \quad (\ell_1, \dots, \ell_{N-1}) \rightarrow (0, \dots, 0), \tag{15}$$

where $\bar{a}(\tau) = \max_{\alpha=1, \dots, N} a_\alpha(\tau)$.

Remark 1.13. Since the function $a(t)$ is piecewise constant, the way u^ε satisfies (1) has to be made precise. An L^1 theory in time (following for instance the approach of [Bourgoing 2008a; 2008b]) could probably be developed for such a problem, but we will use here a different, elementary approach. The Cauchy problem is understood as the solution of successive Cauchy problems. This is the reason why we will first prove a global Lipschitz bound on the solution, so that there indeed exists such a solution.

Remark 1.14. The result of Theorem 1.4 still holds for (1) under assumptions (C1)–(C3), with the set E defined for subsolutions which are moreover assumed to be globally Lipschitz (without fixed bound on the Lipschitz constant). The reader can check that the proof is unchanged.

Remark 1.15. It is somewhat easy to get (12) when the Hamiltonians \bar{H}_α are convex by using the optimal control interpretation of the problem. In the more general case of quasiconvex Hamiltonians, the result still holds true but the proof is more involved.

Remark 1.16. We may have $\bar{A} > \max_{\alpha=1, \dots, N} \langle a_\alpha \rangle$. It is possible to deduce it from (15) in the case $N = 2$ by using the traffic light interpretation of the problem. If we have two traffic lights very close to each other (let us say that the distance in between is at most the space for only one car) and if the traffic lights have common period and are exactly in opposite phases (with, for instance, one minute for the green phase and one minute for the red phase), then the effect of the two traffic lights together gives a very low

flux which is much lower than the effect of a single traffic light alone (i.e., here at most one car every two minutes will go through the two traffic lights).

Traffic flow interpretation of Theorem 1.12. We mentioned above that there are some connections between our problem and traffic flows.

Inequality (13) has a natural traffic interpretation, saying that the average limitation on the traffic flow created by several traffic lights on a single road is greater than or equal to the one created by the traffic light which creates the highest limitation. Moreover, this average limitation is smaller if the distances between traffic lights are bigger, as says the monotonicity of \bar{A} with respect to the distances ℓ_α .

Property (14) says that the minimal limitation is reached if the distances between the traffic lights are bigger than a critical distance d_0 . The proof of this result is quite involved and is reflected in the fact that the bounds that we have on d_0 are not continuous on the data (namely $\max_{\alpha=1,\dots,N} \|a_\alpha\|_\infty$, $\max_{\alpha=1,\dots,N} \langle a_\alpha \rangle$ and the \bar{H}_α).

Finally, property (15) is very natural from the point of view of traffic, since the limit corresponds to the case where all the traffic lights would be at the same position.

Related results. Achdou and Tchou [2015] studied a singular perturbation problem which has the same flavour as the one we are looking at in the present paper. More precisely, they consider the simplest network (a so-called junction) embedded in a star-shaped domain. They prove that the value function of an infinite horizon control problem converges, as the star-shaped domain “shrinks” to the junction, to the value function of a control problem posed on the junction. We borrow from them the idea of studying the cell problem on truncated domains with state constraints. We provide a different approach, which is in some sense more general because it can be applied to problems outside the framework of optimal control theory. Our approach relies in an essential way on the general theory developed in [IM].

The general theme of the lectures by P.-L. Lions [2013–2014] at the Collège de France was “Elliptic or parabolic equations and specified homogenization”. As far as first-order Hamilton–Jacobi equations are concerned, the term “specified homogenization” refers to the problem of constructing correctors to cell problems associated with Hamiltonians that are typically the sum of a periodic one, H , and a compactly supported function f depending only on x , say. Lions exhibits sufficient conditions on f such that the effective Hamilton–Jacobi equation is not perturbed. In terms of flux limiters [IM], it corresponds to looking for sufficient conditions such that the effective flux limiter \bar{A} given by Theorem 1.4 is (less than or) equal to $A_0 = \min H$.

Barles, Briani and Chasseigne [Barles et al. 2013, Theorem 6.1] considered the case

$$H(x, p) = \varphi\left(\frac{x}{\varepsilon}\right)H_R(p) + \left(1 - \varphi\left(\frac{x}{\varepsilon}\right)\right)H_L(p)$$

for some continuous increasing function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\lim_{s \rightarrow -\infty} \varphi(s) = 0 \quad \text{and} \quad \lim_{s \rightarrow +\infty} \varphi(s) = 1.$$

They prove that u^ε converges towards a value function denoted by U^- , which they characterize as the solution to a particular optimal control problem. It is proved in [IM] that U^- is the solution of (6) with

$\bar{H}_\alpha = H_\alpha$ and \bar{A} replaced by $A_J^+ = \max(A_0, A^*)$ with

$$A_0 = \max(\min H_R, \min H_L) \quad \text{and} \quad A^* = \max_{q \in [\min(p_R^0, p_L^0), \max(p_R^0, p_L^0)]} (\min(H_R(q), H_L(q))).$$

Giga and Hamamuki [2013] develop a theory which allows them in particular to prove existence and uniqueness for the following Hamilton–Jacobi equation (changing u to $-u$) in \mathbb{R}^d :

$$\begin{cases} \partial_t u + |\nabla u| = 0 & \text{for } x \neq 0, \\ \partial_t u + |\nabla u| + c = 0 & \text{at } x = 0. \end{cases}$$

The solutions of [Giga and Hamamuki 2013] are constructed as limits of the equation

$$\partial_t u^\varepsilon + |\nabla u^\varepsilon| + c \left(1 - \frac{|x|}{\varepsilon}\right)^+ = 0.$$

In the monodimensional case ($d = 1$), Theorem 1.7 implies that u^ε converges towards

$$\begin{cases} \partial_t u + |\nabla u| = 0 & \text{for } x \neq 0, \\ \partial_t u + \max(A, |\nabla u|) = 0 & \text{at } x = 0, \end{cases}$$

for some $A \in \mathbb{R}$. In view of Theorem 1.4, it is not difficult to prove that $A = \max(0, c)$. The Hamiltonian $\max(c, |\nabla u|)$ is identified in [Giga and Hamamuki 2013] and is referred to as the *relaxed* one.

It is known that homogenization of Hamilton–Jacobi equations is closely related to the study of the large time behaviour of solutions. Hamamuki [2013] discusses the large time behaviour of Hamilton–Jacobi equations with discontinuous source terms in two cases: for compactly supported ones and periodic ones. In our setting, we can address both, and even the sum of a periodic source term and a compactly supported one. It would be interesting to address such a problem in the case of traffic lights. Jin and Yu [2015] study the large time behaviour of the solutions of a Hamilton–Jacobi equations with an x -periodic Hamiltonian and what can be interpreted as a flux limiter depending periodically on time.

Further extensions. It is also possible to address the time homogenization problem of Theorem 1.12 with any finite number of junctions (with limiter functions $a_\alpha(t)$ that are piecewise constants — or continuous — and 1-periodic), either separated with distance of order $O(1)$ or with distance of order $O(\varepsilon)$, or mixing both, and even on a complicated network. See also [Jin and Yu 2015] for other connections between Hamilton–Jacobi equations and traffic light problems, and [Andreianov et al. 2010] for green waves modelling.

Note that the method presented in this paper can be readily applied (without modifying proofs) to the study of homogenization on a finite number of branches and not only two branches; the theory developed in [IM] should also be used for the limit problem.

Similar questions in higher dimensions with point defects of other codimensions will be addressed in future works.

Organization of the article. Section 2 is devoted to the proof of the convergence result (Theorem 1.7). Section 3 is devoted to the construction of correctors far from the junction point (Proposition 1.2), while the junction case, i.e., the proof of Theorem 4.6, is addressed in Section 4. We recall that Theorem 1.9 is

a straightforward corollary of this stronger result. The proof of Theorem 4.6 makes use of a comparison principle which is expected but not completely standard. This is the reason why a proof is sketched in the Appendix, together with two others that are rather standard but included for the reader’s convenience.

Notation. A ball centred at x of radius r is denoted by $B_r(x)$. If $\{u^\varepsilon\}_\varepsilon$ is locally bounded, the upper and lower relaxed limits are defined as

$$\begin{cases} \limsup_\varepsilon *u^\varepsilon(X) = \limsup_{Y \rightarrow X, \varepsilon \rightarrow 0} u^\varepsilon(Y), \\ \liminf_\varepsilon *u^\varepsilon(X) = \liminf_{Y \rightarrow X, \varepsilon \rightarrow 0} u^\varepsilon(Y). \end{cases}$$

In our proofs, constants may change from line to line.

2. Proof of convergence

This section is devoted to the proof of Theorem 1.7. We first construct barriers.

Lemma 2.1 (barriers). *There exists a nonnegative constant C such that, for any $\varepsilon > 0$,*

$$|u^\varepsilon(t, x) - u_0(x)| \leq Ct \quad \text{for } (t, x) \in (0, T) \times \mathbb{R}. \tag{16}$$

Proof. Let L_0 be the Lipschitz constant of the initial datum u_0 . Taking

$$C = \sup_{\substack{(t,x) \in \mathbb{R} \times \mathbb{R} \\ |p| \leq L_0}} |H(t, x, p)| < +\infty,$$

owing to (A0) and (A5), the functions $u^\pm(t, x) = u_0(x) \pm Ct$ are a super- and a sub-solution, respectively, of (1)–(2) and (16) follows via comparison principle. □

We can now prove the convergence theorem.

Proof of Theorem 1.7. We classically consider the upper and lower relaxed semilimits

$$\begin{cases} \bar{u} = \limsup_\varepsilon *u^\varepsilon, \\ \underline{u} = \liminf_\varepsilon *u^\varepsilon. \end{cases}$$

Notice that these functions are well defined because of Lemma 2.1. In order to prove convergence of u^ε towards u^0 , it is sufficient to prove that \bar{u} and \underline{u} are a sub- and a super-solution, respectively, of (6) and (2). The initial condition follows immediately from (16). We focus our attention on the subsolution case, since the supersolution one can be handled similarly.

We first check that

$$\bar{u}(t, 0) = \limsup_{(s,y) \rightarrow (t,0), y>0} \bar{u}(s, y) = \limsup_{(s,y) \rightarrow (t,0), y<0} \bar{u}(s, y). \tag{17}$$

This is a consequence of the stability of such a “weak continuity” condition; see [IM]. Indeed, it is shown in [IM] that classical viscosity solution can be viewed as a flux-limited one; in particular, u^ε solves

$$u_t^\varepsilon + H^-\left(\frac{t}{\varepsilon}, \frac{0}{\varepsilon}, u_x^\varepsilon(t, 0^+)\right) \vee H^+\left(\frac{t}{\varepsilon}, \frac{0}{\varepsilon}, u_x^\varepsilon(t, 0^-)\right) = 0 \quad \text{for } t > 0.$$

Since these ε -Hamiltonians are uniformly coercive and u^ε is continuous, we conclude that (17) holds true.

Let φ be a test function such that

$$(\bar{u} - \varphi)(t, x) < (\bar{u} - \varphi)(\bar{t}, \bar{x}) = 0 \quad \text{for all } (t, x) \in B_{\bar{r}}(\bar{t}, \bar{x}) \setminus \{(\bar{t}, \bar{x})\}. \tag{18}$$

We argue by contradiction, by assuming that

$$\varphi_t(\bar{t}, \bar{x}) + \bar{H}(\bar{x}, \varphi_x(\bar{t}, \bar{x})) = \theta > 0, \tag{19}$$

where

$$\bar{H}(\bar{x}, \varphi_x(\bar{t}, \bar{x})) := \begin{cases} \bar{H}_R(\varphi_x(\bar{t}, \bar{x})) & \text{if } \bar{x} > 0, \\ \bar{H}_L(\varphi_x(\bar{t}, \bar{x})) & \text{if } \bar{x} < 0, \\ F_{\bar{A}}(\varphi_x(\bar{t}, 0^-), \varphi_x(\bar{t}, 0^+)) & \text{if } \bar{x} = 0. \end{cases}$$

We only treat the case where $\bar{x} = 0$, since the case $\bar{x} \neq 0$ is somewhat classical. This latter case is detailed in Section A in the Appendix for the reader's convenience. Using [IM, Proposition 2.5], we may suppose that

$$\varphi(t, x) = \phi(t) + \bar{p}_L x 1_{\{x < 0\}} + \bar{p}_R x 1_{\{x > 0\}}, \tag{20}$$

where ϕ is a C^1 function defined in $(0, +\infty)$. In this case, (19) becomes

$$\phi'(\bar{t}) + F_{\bar{A}}(\bar{p}_L, \bar{p}_R) = \phi'(\bar{t}) + \bar{A} = \theta > 0. \tag{21}$$

Let us consider a solution w of the equation

$$w_t + H(t, x, w_x) = \bar{A}, \tag{22}$$

provided by Theorem 1.9, which is in particular 1-periodic with respect to time. We recall that the function W is the limit of $w^\varepsilon = \varepsilon w(\cdot/\varepsilon)$ as $\varepsilon \rightarrow 0$. We claim that, if $\varepsilon > 0$ is small enough, the perturbed test function $\varphi^\varepsilon(t, x) = \phi(t) + w^\varepsilon(t, x)$ [Evans 1989] is a viscosity supersolution of

$$\varphi_t^\varepsilon + H\left(\frac{t}{\varepsilon}, \frac{x}{\varepsilon}, \varphi_x^\varepsilon\right) = \frac{\theta}{2} \quad \text{in } B_r(\bar{t}, 0)$$

for some sufficiently small $r > 0$. In order to justify this fact, let $\psi(t, x)$ be a test function touching φ^ε from below at $(t_1, x_1) \in B_r(\bar{t}, 0)$. In this way,

$$w\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}\right) = \frac{1}{\varepsilon}(\psi(t_1, x_1) - \phi(t_1))$$

and

$$w(s, y) \geq \frac{1}{\varepsilon}(\psi(\varepsilon s, \varepsilon y) - \phi(\varepsilon s))$$

for (s, y) in a neighbourhood of $(t_1/\varepsilon, x_1/\varepsilon)$. Hence, from (21)–(22),

$$\psi_t(t_1, x_1) + H\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \psi_x(t_1, x_1)\right) \geq \bar{A} + \phi'(t_1) \geq \bar{A} + \phi'(\bar{t}) - \frac{\theta}{2} \geq \frac{\theta}{2}$$

provided r is small enough. Hence, the claim is proved.

Combining (7) from Theorem 1.9 with (18) and (20), we can fix $\kappa_r > 0$ and $\varepsilon > 0$ small enough so that

$$u^\varepsilon + \kappa_r \leq \varphi^\varepsilon \quad \text{on } \partial B_r(\bar{t}, 0).$$

By the comparison principle the previous inequality holds in $B_r(\bar{t}, 0)$. Passing to the limit as $\varepsilon \rightarrow 0$ and $(t, x) \rightarrow (\bar{t}, \bar{x})$, we get the contradiction

$$\bar{u}(\bar{t}, 0) + \kappa_r \leq \varphi(\bar{t}, 0) = \bar{u}(\bar{t}, 0).$$

The proof of convergence is now complete. \square

Remark 2.2. For the supersolution property, φ in (20) should be replaced with

$$\varphi(t, x) = \phi(t) + \hat{p}_L x 1_{\{x < 0\}} + \hat{p}_R x 1_{\{x > 0\}}.$$

3. Homogenized Hamiltonians

In order to prove Proposition 1.2, we first prove the following lemma. Even if the proof is standard, we give it in full detail since we will adapt it when constructing global correctors for the junction.

Lemma 3.1 (existence of a corrector). *There exists $\lambda \in \mathbb{R}$ for which there is a bounded (discontinuous) viscosity solution of (3).*

Remark 3.2. If H_α does not depend on t , then it is possible to construct a corrector which does not depend on time either. We leave the details to the reader.

Proof. For any $\delta > 0$, it is possible to construct a (possibly discontinuous) viscosity solution v^δ of

$$\begin{cases} \delta v^\delta + v_t^\delta + H_\alpha(t, x, p + v_x^\delta) = 0 & \text{in } \mathbb{R} \times \mathbb{R}, \\ v^\delta \text{ is } \mathbb{Z}^2\text{-periodic.} \end{cases}$$

First, the comparison principle implies

$$|\delta v^\delta| \leq C_\alpha, \tag{23}$$

where

$$C_\alpha = \sup_{(t,x) \in [0,1]^2} |H_\alpha(t, x, p)|.$$

Second, the function

$$m^\delta(x) = \sup_{t \in \mathbb{R}} (v^\delta)^*(t, x)$$

is a subsolution of

$$H_\alpha(t(x), x, p + m_x^\delta) \leq C_\alpha$$

(for some function $t(x)$). Assumptions (A3) and (A5) imply that there exists $C > 0$ independent of δ such that

$$|m_x^\delta| \leq C \quad \text{and} \quad v_t^\delta \leq C.$$

In particular, the comparison principle implies that, for all $t \in \mathbb{R}$, $x \in \mathbb{R}$ and $h \geq 0$,

$$v^\delta(t + h, x) \leq v^\delta(t, x) + Ch.$$

Combining this inequality with the time-periodicity of v^δ yields

$$|v^\delta(t, x) - m^\delta(x)| \leq C;$$

in particular,

$$|v^\delta(t, x) - v^\delta(0, 0)| \leq C. \quad (24)$$

Hence, the half-relaxed limits

$$\bar{v} = \limsup_{\delta \rightarrow 0}^*(v^\delta - v^\delta(0, 0)) \quad \text{and} \quad \underline{v} = \liminf_{\delta \rightarrow 0}^*(v^\delta - v^\delta(0, 0))$$

are finite. Moreover, (23) implies that $\delta v^\delta(0, 0) \rightarrow -\lambda$ (at least along a subsequence). Hence, the discontinuous stability of viscosity solutions implies that \bar{v} is a \mathbb{Z}^2 -periodic subsolution of (3) and \underline{v} is a \mathbb{Z}^2 -periodic supersolution of the same equation. Perron's method then allows us to construct a corrector between \bar{v} and $\underline{v} + C$ with $C = \sup(\bar{v} - \underline{v})$. The proof of the lemma is now complete. \square

The following lemma is completely standard; the proof is given in Section B in the Appendix for the reader's convenience.

Lemma 3.3 (uniqueness of λ). *The real number λ given by Lemma 3.1 is unique. If $\bar{H}_\alpha(p)$ denotes this real number, the function \bar{H}_α is continuous.*

Lemma 3.4 (coercivity of \bar{H}_α). *The continuous function \bar{H}_α is coercive:*

$$\lim_{|p| \rightarrow +\infty} \bar{H}_\alpha(p) = +\infty.$$

Proof. In view of the uniform coercivity in p of H_α with respect to (t, x) (see (A3)), for any $R > 0$ there exists a positive constant C_R such that

$$|p| \geq C_R \quad \implies \quad H_\alpha(t, x, p) \geq R \quad \text{for all } (t, x) \in \mathbb{R} \times \mathbb{R}. \quad (25)$$

Let v^α be the discontinuous corrector given by Lemma 3.1 and (\bar{t}, \bar{x}) the point of supremum of its upper semicontinuous envelope $(v^\alpha)^*$. Then we have

$$H_\alpha(\bar{t}, \bar{x}, p) \leq \bar{H}_\alpha(p),$$

which implies

$$\bar{H}_\alpha(p) \geq R \quad \text{for } |p| \geq C_R. \quad (26)$$

The proof of the lemma is now complete. \square

We first prove the quasiconvexity of \bar{H}_α under assumption (B-ii). We in fact prove more: the effective Hamiltonian is convex in this case.

Lemma 3.5 (convexity of \bar{H}_α under (B-ii)). *Assume (A0)–(A5) and (B-ii). Then the function \bar{H}_α is convex.*

Proof. For $p, q \in \mathbb{R}$, let v_p and v_q be solutions of (3) with $\lambda = \bar{H}_\alpha(p)$ and $\bar{H}_\alpha(q)$, respectively. We also set

$$u_p(t, x) = v_p(t, x) + px - t\bar{H}_\alpha(p)$$

and define u_q similarly.

Step 1: u_p and u_q are locally Lipschitz continuous. In this case, we have, almost everywhere,

$$\begin{cases} (u_p)_t + H_\alpha(t, x, (u_p)_x) = 0, \\ (u_q)_t + H_\alpha(t, x, (u_q)_x) = 0. \end{cases}$$

For $\mu \in [0, 1]$, let

$$\bar{u} = \mu u_p + (1 - \mu)u_q.$$

By convexity, we get, almost everywhere,

$$\bar{u}_t + H_\alpha(t, x, \bar{u}_x) \leq 0. \tag{27}$$

We claim that the convexity of H_α (in the gradient variable) implies that \bar{u} is a viscosity subsolution. To see this, we use an argument of [Bardi and Capuzzo-Dolcetta 1997, Proposition 5.1]. For $P = (t, x)$, we define a mollifier $\rho_\delta(P) = \delta^{-2}\rho(\delta^{-1}P)$ and set

$$\bar{u}_\delta = \bar{u} \star \rho_\delta$$

Then, by convexity, we get, with $Q = (s, y)$,

$$(\bar{u}_\delta)_t + H_\alpha(P, (\bar{u}_\delta)_x) \leq \int dQ \{H_\alpha(P, \bar{u}_x(Q)) - H_\alpha(Q, \bar{u}_x(Q))\} \rho_\delta(P - Q).$$

The fact that \bar{u}_x is locally bounded and the fact that H_α is continuous imply that the right-hand side goes to zero as $\delta \rightarrow 0$. We deduce (by stability of viscosity subsolutions) that (27) holds true in the viscosity sense. Then the comparison principle implies that

$$\mu \bar{H}_\alpha(p) + (1 - \mu)\bar{H}_\alpha(q) \geq \bar{H}_\alpha(\mu p + (1 - \mu)q). \tag{28}$$

Step 2: u_p and u_q are continuous. We proceed in two substeps:

Step 2.1: the case of a single function u . We first want to show that if $u = u_p$ is continuous and satisfies (27) almost everywhere, then u is a viscosity subsolution. To this end, we will use the structural assumptions satisfied by the Hamiltonian. The ones that were useful to prove the comparison principle will be also useful to prove the result we want, so we will revisit that proof. We also use the fact that

$$u(t, x) - px + t\bar{H}_\alpha(p) \text{ is bounded.} \tag{29}$$

For $\nu > 0$, we set

$$u^\nu(t, x) = \sup_{s \in \mathbb{R}} \left(u(s, x) - \frac{(t - s)^2}{2\nu} \right) = u(s_\nu, x) - \frac{(t - s_\nu)^2}{2\nu}.$$

As usual, we get from (29) that

$$|t - s_\nu| \leq C\sqrt{\nu} \text{ with } C = C(p, T) \tag{30}$$

for $t \in (-T, T)$. In particular $s_\nu \rightarrow t$ locally uniformly. If a test function φ touches u^ν from above at some point (t, x) , then we have $\varphi_t(t, x) = -(t - s_\nu)/\nu$ and

$$\begin{aligned} \varphi_t(t, x) + H_\alpha(t, x, \varphi_x(t, x)) &\leq H_\alpha(t, x, \varphi_x(t, x)) - H_\alpha(s_\nu, x, \varphi_x(t, x)) \\ &\leq \omega(|t - s_\nu|(1 + \max(0, H_\alpha(s_\nu, x, \varphi_x(t, x)))) \\ &\leq \omega\left(\frac{(t - s_\nu)^2}{\nu} + |t - s_\nu|\right), \end{aligned} \tag{31}$$

where we have used (A2) in the third line. The right-hand side goes to zero as ν goes to zero since

$$\frac{(t - s_\nu)^2}{\nu} \rightarrow 0 \quad \text{locally uniformly with respect to } (t, x)$$

(recall u is continuous). Indeed, this can be checked for (t, x) replaced by (t_ν, x_ν) because, for any sequence $(t_\nu, s_\nu, x_\nu) \rightarrow (t, t, x)$, we have

$$u(t_\nu, x_\nu) \leq u^\nu(t_\nu, x_\nu) = u(s_\nu, x_\nu) - \frac{(t_\nu - s_\nu)^2}{2\nu},$$

where the continuity of u implies the result. For a given $\nu > 0$, we see that (30) and (31) imply that

$$|\varphi_t|, |\varphi_x| \leq C_{\nu,p}.$$

This implies in particular that u^ν is Lipschitz continuous, and then

$$u_t^\nu + H(t, x, u_x^\nu) \leq o_\nu(1) \quad \text{a.e.},$$

where $o_\nu(1)$ is locally uniform with respect to (t, x) .

Step 2.2: application. Applying Step 2.1, we get, for $z = p, q$,

$$(u_z^\nu)_t + H(t, x, (u_z^\nu)_x) \leq o_\nu(1) \quad \text{a.e.},$$

where $o_\nu(1)$ is locally uniform with respect to (t, x) . Step 1 implies that

$$\bar{u}^\nu := \mu u_p^\nu + (1 - \mu)u_q^\nu$$

is a viscosity subsolution of

$$(\bar{u}^\nu)_t + H_\alpha(t, x, (\bar{u}^\nu)_x) \leq o_\nu(1),$$

where $o_\nu(1)$ is locally uniform with respect to (t, x) . In the limit $\nu \rightarrow 0$, we recover (by stability of subsolutions) that \bar{u} is a viscosity subsolution, i.e., satisfies (27) in the viscosity sense. This then gives the same conclusion as in Step 1.

Step 3: the general case. To cover the general case, we simply replace u_p by \tilde{u}_p , the solution to the Cauchy problem

$$\begin{cases} (\tilde{u}_p)_t + H_\alpha(t, x, (\tilde{u}_p)_x) = 0 & \text{for } (t, x) \in (0, +\infty) \times \mathbb{R}, \\ \tilde{u}_p(0, x) = px. \end{cases}$$

Then \tilde{u}_p is continuous and satisfies $|\tilde{u}_p - u_p| \leq C$. Proceeding similarly with \tilde{u}_q and using Step 2, we deduce the desired inequality (28). The proof is now complete. \square

We finally prove the quasiconvexity of \bar{H}_α under assumption (B-i).

Lemma 3.6 (quasiconvexity of \bar{H}_α under (B-i)). *Assume (A0)–(A5) and (B-i). Then the function \bar{H}_α is quasiconvex.*

Proof. We reduce quasiconvexity to convexity by composing with an increasing function γ ; note that such a reduction was already used in optimization and in partial differential equations; see, for instance, [Lions 1981; Kawohl 1985].

We first assume that H_α satisfies

$$\begin{cases} H_\alpha \in C^2, \\ D_{pp}^2 H_\alpha(x, p_\alpha^0) > 0, \\ D_p H_\alpha(x, p) < 0 & \text{for } p \in (-\infty, p_\alpha^0), \\ D_p H_\alpha(x, p) > 0 & \text{for } p \in (p_\alpha^0, +\infty), \\ H_\alpha(x, p) \rightarrow +\infty & \text{as } |p| \rightarrow +\infty \text{ uniformly with respect to } x \in \mathbb{R}. \end{cases} \quad (32)$$

For a function γ such that

$$\gamma \text{ is convex, } \gamma \in C^2(\mathbb{R}) \text{ and } \gamma' \geq \delta_0 > 0,$$

we have

$$D_{pp}^2(\gamma \circ H_\alpha) > 0$$

if and only if

$$(\ln \gamma')'(\lambda) > -\frac{D_{pp}^2 H_\alpha(x, p)}{(D_p H_\alpha(x, p))^2} \text{ for } p = \pi_\alpha^\pm(x, \lambda) \text{ and } \lambda \geq H_\alpha(x, p), \quad (33)$$

where $\pi_\alpha^\pm(x, \lambda)$ is the only real number r such that $\pm r \geq 0$ and $H_\alpha(x, r) = \lambda$. Because $D_{pp}^2 H_\alpha(x, p_\alpha^0) > 0$, we see that the right-hand side is negative for λ close enough to $H_\alpha(x, p_\alpha^0)$ and it is indeed possible to construct such a function γ .

In view of Remark 3.2, we can construct a solution of $\delta v^\delta + \gamma \circ H_\alpha(x, p + v_x^\delta) = 0$ with $-\delta v^\delta \rightarrow \overline{\gamma \circ H_\alpha}(p)$ as $\delta \rightarrow 0$, and a solution of

$$\gamma \circ H_\alpha(x, p + v_x) = \overline{\gamma \circ H_\alpha}(p).$$

This shows that

$$\bar{H}_\alpha = \gamma^{-1} \circ \overline{\gamma \circ H_\alpha}.$$

Thanks to Lemmas 3.4 and 3.5, we know that $\overline{\gamma \circ H_\alpha}$ is coercive and convex. Hence, \bar{H}_α is quasiconvex.

If now H_α does not satisfies (32) then, for all $\varepsilon > 0$, there exists $H_\alpha^\varepsilon \in C^2$ such that

$$\begin{cases} (D_{pp}^2 H_\alpha^\varepsilon)(x, p_\alpha^0) > 0, \\ D_p H_\alpha^\varepsilon(x, p) < 0 & \text{for } p \in (-\infty, p_\alpha^0), \\ D_p H_\alpha^\varepsilon(x, p) > 0 & \text{for } p \in (p_\alpha^0, +\infty), \\ |H_\alpha^\varepsilon - H_\alpha| < \varepsilon. \end{cases}$$

Then we can argue as in the proof of continuity of \bar{H}_α and deduce that

$$\bar{H}_\alpha(\rho) = \lim_{\varepsilon \rightarrow 0} \bar{H}_\alpha^\varepsilon(\rho).$$

Moreover, the previous case implies that $\bar{H}_\alpha^\varepsilon$ is quasiconvex. Hence, so is \bar{H}_α . The proof of the lemma is now complete. □

Proof of Proposition 1.2. Combine Lemmas 3.1, 3.3, 3.4, 3.5 and 3.6. □

4. Truncated cell problems

We consider the following problem: find $\lambda_\rho \in \mathbb{R}$ and w such that

$$\begin{cases} w_t + H(t, x, w_x) = \lambda_\rho & \text{for } (t, x) \in \mathbb{R} \times (-\rho, \rho), \\ w_t + H^-(t, x, w_x) = \lambda_\rho & \text{for } (t, x) \in \mathbb{R} \times \{-\rho\}, \\ w_t + H^+(t, x, w_x) = \lambda_\rho & \text{for } (t, x) \in \mathbb{R} \times \{\rho\}, \\ w \text{ is 1-periodic in } t. \end{cases} \tag{34}$$

Even if our approach is different, we borrow here an idea from [Achdou and Tchou 2015] by truncating the domain and considering correctors in $[-\rho, \rho]$ with $\rho \rightarrow +\infty$.

A comparison principle.

Proposition 4.1 (comparison principle for a mixed boundary value problem). *Let $\rho_2 > \rho_1 > \rho_0$ and $\lambda \in \mathbb{R}$ and v be a supersolution of the boundary value problem*

$$\begin{cases} v_t + H(t, x, v_x) \geq \lambda & \text{for } (t, x) \in \mathbb{R} \times (\rho_1, \rho_2), \\ v_t + H^+(t, x, v_x) \geq \lambda & \text{for } (t, x) \in \mathbb{R} \times \{\rho_2\}, \\ v(t, x) \geq U_0(t) & \text{for } (t, x) \in \mathbb{R} \times \{\rho_1\}, \\ v \text{ is 1-periodic in } t, \end{cases} \tag{35}$$

where U_0 is continuous and, for $\varepsilon_0 > 0$, let u be a subsolution of

$$\begin{cases} u_t + H(t, x, u_x) \leq \lambda - \varepsilon_0 & \text{for } (t, x) \in \mathbb{R} \times (\rho_1, \rho_2), \\ u_t + H^+(t, x, u_x) \leq \lambda - \varepsilon_0 & \text{for } (t, x) \in \mathbb{R} \times \{\rho_2\}, \\ u(t, x) \leq U_0(t) & \text{for } (t, x) \in \mathbb{R} \times \{\rho_1\}, \\ u \text{ is 1-periodic in } t. \end{cases} \tag{36}$$

Then $u \leq v$ in $\mathbb{R} \times [\rho_1, \rho_2]$.

Remark 4.2. A similar result holds true if the Dirichlet condition is imposed at $x = \rho_2$ and junction conditions

$$\begin{aligned} v_t + H^-(t, x, v_x) &\geq \lambda && \text{at } x = \rho_1, \\ u_t + H^-(t, x, u_x) &\leq \lambda - \varepsilon_0 && \text{at } x = \rho_1, \end{aligned}$$

are imposed at $x = \rho_1$.

The proof of Proposition 4.1 is very similar to (in fact simpler than) the proof of the comparison principle for Hamilton–Jacobi equations on networks contained in [IM]. The main difference lies in the fact that, in our case, u and v are global in time and the space domain is bounded. A sketch of the proof is provided in Section C in the Appendix, shedding some light on the main differences. Here, the parameter $\varepsilon_0 > 0$ in (36) is used in place of the standard correction term $-\eta/(T - t)$ for a Cauchy problem.

Correctors on truncated domains.

Proposition 4.3 (existence and properties of a corrector on a truncated domain). *There exists a unique $\lambda_\rho \in \mathbb{R}$ such that there exists a solution $w^\rho = w$ of (34). Moreover, there exists a constant $C > 0$ independent of $\rho \in (\rho_0, +\infty)$ and a function $m^\rho: [-\rho, \rho] \rightarrow \mathbb{R}$ such that*

$$\begin{cases} |\lambda_\rho| \leq C, \\ |m^\rho(x) - m^\rho(y)| \leq C|x - y| & \text{for } x, y \in [-\rho, \rho], \\ |w^\rho(t, x) - m^\rho(x)| \leq C & \text{for } (t, x) \in \mathbb{R} \times [-\rho, \rho]. \end{cases} \tag{37}$$

Proof. In order to construct a corrector on the truncated domain, we proceed classically by considering

$$\begin{cases} \delta w^\delta + w_t^\delta + H(t, x, w_x^\delta) = 0 & \text{for } (t, x) \in \mathbb{R} \times (-\rho, \rho), \\ \delta w^\delta + w_t^\delta + H^-(t, x, w_x^\delta) = 0 & \text{for } (t, x) \in \mathbb{R} \times \{-\rho\}, \\ \delta w^\delta + w_t^\delta + H^+(t, x, w_x^\delta) = 0 & \text{for } (t, x) \in \mathbb{R} \times \{\rho\}, \\ w^\delta \text{ is 1-periodic in } t. \end{cases} \tag{38}$$

A discontinuous viscosity solution of (38) is constructed by Perron’s method (in the class of 1-periodic functions in time) since $\pm\delta^{-1}C$ are trivial super- and sub-solutions if C is chosen to be

$$C = \sup_{\substack{t \in \mathbb{R} \\ x \in \mathbb{R}}} |H(t, x, 0)|.$$

In particular, the solution w^δ satisfies, by construction,

$$|w^\delta| \leq \frac{C}{\delta}. \tag{39}$$

We next consider

$$m^\delta(x) = \sup_{t \in \mathbb{R}} (w^\delta)^*(t, x).$$

We remark that the supremum is reached since w^δ is periodic in time; we also remark that m^δ is a viscosity subsolution of

$$H(t(x), x, m_x^\delta) \leq C, \quad x \in (-\rho, \rho),$$

(for some function $t(x)$). In view of (A3), we conclude that m^δ is globally Lipschitz continuous and

$$|m_x^\delta| \leq C \tag{40}$$

for some constant C which still only depends on H . Assumption (A3) also implies that

$$w_t^\delta \leq C$$

(with C only depending on H). In particular, the comparison principle implies that, for all $t \in \mathbb{R}$, $x \in (-\rho, \rho)$ and $h \geq 0$,

$$w^\delta(t+h, x) \leq w^\delta(t, x) + Ch.$$

Combining this information with the periodicity of w^δ in t , we conclude that, for $t \in \mathbb{R}$ and $x \in (-\rho, \rho)$,

$$|w^\delta(t, x) - m^\delta(x)| \leq C.$$

In particular,

$$|w^\delta(t, x) - w^\delta(0, 0)| \leq C.$$

We then consider

$$\bar{w} = \limsup_{\delta}^*(w^\delta - w^\delta(0, 0)) \quad \text{and} \quad \underline{w} = \liminf_{\delta}^*(w^\delta - w^\delta(0, 0)).$$

We next remark that (39) and (40) imply that there exists $\delta_n \rightarrow 0$ such that

$$\begin{aligned} m^{\delta_n} - m^{\delta_n}(0) &\rightarrow m^\rho \quad \text{as } n \rightarrow +\infty, \\ \delta_n w^{\delta_n}(0, 0) &\rightarrow -\lambda_\rho \quad \text{as } n \rightarrow +\infty, \end{aligned}$$

(the first convergence being locally uniform). In particular, λ , \bar{w} , \underline{w} and m^ρ satisfy

$$\begin{aligned} |\lambda_\rho| &\leq C, \\ |\bar{w} - m^\rho| &\leq C, \\ |\underline{w} - m^\rho| &\leq C, \\ |m_x^\rho| &\leq C. \end{aligned}$$

Discontinuous stability of viscosity solutions of Hamilton–Jacobi equations implies that $\bar{w} - 2C$ and \underline{w} are a sub- and a super-solution, respectively, of (34), and

$$\bar{w} - 2C \leq \underline{w}.$$

Perron’s method is used once again in order to construct a solution w^ρ of (34) which is 1-periodic in time. In view of the previous estimates, λ_ρ , m^ρ and w^ρ satisfy (37). Proving the uniqueness of λ_ρ is classical, so we skip it. The proof of the proposition is now complete. \square

Proposition 4.4 (first definition of the effective flux limiter). *The map $\rho \mapsto \lambda_\rho$ is nondecreasing and bounded in $(0, +\infty)$. In particular,*

$$\bar{A} = \lim_{\rho \rightarrow +\infty} \lambda_\rho$$

exists and $\bar{A} \geq \lambda_\rho$ for all $\rho > 0$.

Proof. For $\rho' > \rho > 0$, we see that the restriction of $w^{\rho'}$ to $[-\rho, \rho]$ is a subsolution, as a consequence of [IM, Proposition 2.15]. The boundedness of the map follows from Proposition 4.3. \square

We next prove that we can control w^ρ from below under appropriate assumptions on \bar{A} .

Proposition 4.5 (control of slopes on a truncated domain). *Assume first that $\bar{A} > \min \bar{H}_R$. Then, for all $\delta > 0$, there exists $\rho_\delta > 0$ and $C_\delta > 0$ (independent of ρ) such that, for $x \geq \rho_\delta$ and $h \geq 0$,*

$$w^\rho(t, x+h) - w^\rho(t, x) \geq (\bar{p}_R - \delta)h - C_\delta. \quad (41)$$

If we now assume that $\bar{A} > \min \bar{H}_L$ then, for $x \leq -\rho_\delta$ and $h \geq 0$,

$$w^\rho(t, x-h) - w^\rho(t, x) \geq (-\bar{p}_L - \delta)h - C_\delta \quad (42)$$

for some $\rho_\delta > 0$ and $C_\delta > 0$ as above.

Proof. We only prove (41), since the proof of (42) follows along the same lines. Let $\delta > 0$. In view of (A5), we know that there exists ρ_δ such that

$$|H(t, x, p) - H_R(t, x, p)| \leq \delta \quad \text{for } x \geq \rho_\delta. \quad (43)$$

Assume that $\bar{A} > \min \bar{H}_R$. Then Proposition 1.2 implies that we can pick p_R^δ such that

$$\bar{H}_R(p_R^\delta) = \bar{H}_R^+(p_R^\delta) = \lambda_\rho - 2\delta$$

for $\rho \geq \rho_0$ and $\delta \leq \delta_0$, by choosing ρ_0 large enough and δ_0 small enough.

We now fix $\rho \geq \rho_\delta$ and $x_0 \in [\rho_\delta, \rho]$. In view of Proposition 1.2 applied to $p = p_R^\delta$, we know that there exists a corrector v_R solving (3) with $\alpha = R$. Since it is \mathbb{Z}^2 -periodic, it is bounded and $w_R = p_R^\delta x + v_R(t, x)$ solves

$$(w_R)_t + H_R(t, x, (w_R)_x) = \lambda_\rho - 2\delta \quad \text{for } (t, x) \in \mathbb{R} \times \mathbb{R}.$$

In particular, the restriction of w_R to $[\rho_\delta, \rho]$ satisfies (see [IM, Proposition 2.15])

$$\begin{cases} (w_R)_t + H_R(t, x, (w_R)_x) \leq \lambda_\rho - 2\delta & \text{for } (t, x) \in \mathbb{R} \times (\rho_\delta, \rho), \\ (w_R)_t + H_R^+(t, x, (w_R)_x) \leq \lambda_\rho - 2\delta & \text{for } (t, x) \in \mathbb{R} \times \{\rho\}. \end{cases}$$

In view of (43), this implies

$$\begin{cases} (w_R)_t + H(t, x, (w_R)_x) \leq \lambda_\rho - \delta & \text{for } (t, x) \in \mathbb{R} \times (\rho_\delta, \rho), \\ (w_R)_t + H^+(t, x, (w_R)_x) \leq \lambda_\rho - \delta & \text{for } (t, x) \in \mathbb{R} \times \{\rho\}. \end{cases}$$

Now we remark that $v = w^\rho - w^\rho(0, x_0)$ and $u = w_R - w_R(0, x_0) - 2C - 2\|v_R\|_\infty$ satisfy

$$v(t, x_0) \geq -2C \geq u(t, x_0),$$

where C is given by (37). Thanks to the comparison principle from Proposition 4.1, we thus get, for $x \in [x_0, \rho]$,

$$w^\rho(t, x) - w^\rho(t, x_0) \geq p_R^\delta(x - x_0) - C_\delta,$$

where C_δ is a large constant which does not depend on ρ . In particular, we get (41), reducing δ if necessary. \square

Construction of global correctors. We now state and prove a result which implies Theorem 1.9, stated in the introduction.

Theorem 4.6 (existence of a global corrector for the junction). *Assume (A0)–(A5) and either (B-i) or (B-ii).*

(i) General properties: *There exists a solution w of (4) with $\lambda = \bar{A}$ such that, for all $(t, x) \in \mathbb{R}^2$,*

$$|w(t, x) - m(x)| \leq C \tag{44}$$

for some globally Lipschitz continuous function m , and

$$\bar{A} \geq A_0.$$

(ii) Bound from below at infinity: *If $\bar{A} > \max_{\alpha=L,R}(\min \bar{H}_\alpha)$ then there exists $\delta_0 > 0$ such that, for every $\delta \in (0, \delta_0)$, there exists $\rho_\delta > \rho_0$ such that w satisfies*

$$\begin{cases} w(t, x+h) - w(t, x) \geq (\bar{p}_R - \delta)h - C_\delta & \text{for } x \geq \rho_\delta \text{ and } h \geq 0, \\ w(t, x-h) - w(t, x) \geq (-\bar{p}_L - \delta)h - C_\delta & \text{for } x \leq -\rho_\delta \text{ and } h \geq 0. \end{cases} \tag{45}$$

The first line of (45) also holds if we have only $\bar{A} > \min \bar{H}_R$, while the second line of (45) also holds if we have only $\bar{A} > \min \bar{H}_L$.

(iii) Rescaling w : *For $\varepsilon > 0$, we set*

$$w^\varepsilon(t, x) = \varepsilon w(\varepsilon^{-1}t, \varepsilon^{-1}x).$$

Then (along a subsequence $\varepsilon_n \rightarrow 0$) we have that w^ε converges locally uniformly towards a function $W = W(x)$ which satisfies

$$\begin{cases} |W(x) - W(y)| \leq C|x - y| & \text{for all } x, y \in \mathbb{R}, \\ \bar{H}_R(W_x) = \bar{A} \quad \text{and} \quad \hat{p}_R \geq W_x \geq \bar{p}_R & \text{for } x \in (0, +\infty), \\ \bar{H}_L(W_x) = \bar{A} \quad \text{and} \quad \hat{p}_L \leq W_x \leq \bar{p}_L & \text{for } x \in (-\infty, 0). \end{cases} \tag{46}$$

In particular, we have $W(0) = 0$ and

$$\hat{p}_R x 1_{\{x>0\}} + \hat{p}_L x 1_{\{x<0\}} \geq W(x) \geq \bar{p}_R x 1_{\{x>0\}} + \bar{p}_L x 1_{\{x<0\}}. \tag{47}$$

Proof. We consider (up to some subsequence)

$$\bar{w} = \limsup_{\rho \rightarrow +\infty}^*(w^\rho - w^\rho(0, 0)), \quad \underline{w} = \liminf_{\rho \rightarrow +\infty}^*(w^\rho - w^\rho(0, 0)) \quad \text{and} \quad m = \lim_{\rho \rightarrow +\infty} (m^\rho - m^\rho(0)).$$

We derive from (37) that \underline{w} and \bar{w} are finite and

$$m - C \leq \underline{w} \leq \bar{w} \leq m + C.$$

Moreover, discontinuous stability of viscosity solutions implies that $\bar{w} - 2C$ and \underline{w} are a sub- and a super-solution, respectively, of (4) with $\lambda = \bar{A}$ (recall Proposition 4.4). Hence, a discontinuous viscosity solution w of (4) can be constructed by Perron’s method (in the class of functions that are 1-periodic in time).

Using (37) again, w and m satisfy (44). We also get (45) from Proposition 4.5 (use (37) and pass to the limit with m instead of w if necessary).

We now study $w^\varepsilon(t, x) = \varepsilon w(\varepsilon^{-1}t, \varepsilon^{-1}x)$. Note that (37) implies in particular that

$$w^\varepsilon(t, x) = \varepsilon m(\varepsilon^{-1}x) + O(\varepsilon).$$

In particular, we can find a sequence $\varepsilon_n \rightarrow 0$ such that

$$w^{\varepsilon_n}(t, x) \rightarrow W(x) \quad \text{locally uniformly as } n \rightarrow +\infty,$$

with $W(0) = 0$. Arguing as in the proof of convergence away from the junction point (see the case $\bar{x} \neq 0$ in Section A in the Appendix), we deduce that W satisfies

$$\begin{aligned} \bar{H}_R(W_x) &= \bar{A} \quad \text{for } x > 0, \\ \bar{H}_L(W_x) &= \bar{A} \quad \text{for } x < 0. \end{aligned}$$

We also deduce from (45) that, for all $\delta > 0$ and $x > 0$,

$$W_x \geq \bar{p}_R - \delta$$

in the case where $\bar{A} > \min \bar{H}_R$. Assume now that $\bar{A} = \min \bar{H}_R$. This implies that

$$\bar{p}_R \leq W_x \leq \hat{p}_R$$

and, in all cases, we thus get (47) for $x > 0$.

Similarly, we can prove for $x < 0$ that

$$\hat{p}_L \leq W_x \leq \bar{p}_L$$

and the proof of (46) of is achieved. This implies (47). The proof of Theorem 4.6 is now complete. \square

Proof of Theorem 1.4. Let \bar{A} denote the limit of A_ρ (see Proposition 4.4). We want to prove that $\bar{A} = \inf E$, where we recall that

$$E = \{\lambda \in \mathbb{R} : \text{there exists a subsolution } w \text{ of (4)}\}.$$

In view of (4), subsolutions are assumed to be periodic in time; we will see that they also automatically satisfy some growth conditions at infinity, see (48) below.

We argue by contradiction, by assuming that there exist $\lambda < \bar{A}$ and a subsolution w_λ of (4). The function

$$m_\lambda(x) = \sup_{t \in \mathbb{R}} (w_\lambda)^*(t, x)$$

satisfies

$$H(t(x), x, (m_\lambda)_x) \leq C$$

(for some function $t(x)$). Assumption (A3) implies that m_λ is globally Lipschitz continuous. Moreover, since w_λ is 1-periodic in time and $(w_\lambda)_t \leq C$,

$$|w_\lambda(t, x) - m_\lambda(x)| \leq C.$$

Hence,

$$w_\lambda^\varepsilon(t, x) = \varepsilon w_\lambda(\varepsilon^{-1}t, \varepsilon^{-1}x)$$

has a limit W^λ which satisfies

$$\bar{H}_R(W_x^\lambda) \leq \lambda \quad \text{for } x > 0.$$

In particular, for $x > 0$,

$$W_x^\lambda \leq \hat{p}_R^\lambda := \max\{p \in \mathbb{R} : \bar{H}_R(p) = \lambda\} < \bar{p}_R,$$

where \bar{p}_R is as defined in (8). Similarly,

$$W_x^\lambda \geq \hat{p}_L^\lambda := \min\{p \in \mathbb{R} : \bar{H}_L(p) = \lambda\} > \bar{p}_L$$

with \bar{p}_L as defined in (9). These two inequalities imply in particular that, for all $\delta > 0$, there exists \tilde{C}_δ such that

$$w_\lambda(t, x) \leq \begin{cases} (\hat{p}_R^\lambda + \delta)x + \tilde{C}_\delta & \text{for } x > 0, \\ (\hat{p}_L^\lambda + \delta)x + \tilde{C}_\delta & \text{for } x < 0. \end{cases} \quad (48)$$

In particular,

$$w_\lambda < w \quad \text{for } |x| \geq R$$

if δ is small enough and R is large enough. Hence,

$$w_\lambda < w + C_R \quad \text{for } x \in \mathbb{R}.$$

Note finally that $u(t, x) = w(t, x) + C_R - \bar{A}t$ is a solution and $u_\lambda(t, x) = w_\lambda(t, x) - \lambda t$ is a subsolution of (1) with $\varepsilon = 1$ and $u_\lambda(0, x) \leq u(0, x)$. Hence, the comparison principle implies that

$$w_\lambda(t, x) - \lambda t \leq w(t, x) - \bar{A}t + C_R.$$

Dividing by t and letting $t \rightarrow +\infty$, we get the contradiction

$$\bar{A} \leq \lambda.$$

The proof is now complete. □

5. Proof of Theorem 1.12

This section is devoted to the proof of Theorem 1.12. As pointed out in Remark 1.13 above, the notion of solutions for (1) has to first be made precise, because the Hamiltonian is discontinuous with respect to time.

Notion of solutions for (1). For $\varepsilon = 1$, a function u is a *solution* of (1) if it is globally Lipschitz continuous (in space and time) and it solves successively the Cauchy problems on time intervals $[\tau_i + k, \tau_{i+1} + k)$ for $i = 0, \dots, K$ and $k \in \mathbb{N}$.

Because of this definition and approach, we have to show that, if the initial datum u_0 is globally Lipschitz continuous, then the solution to the successive Cauchy problems is also globally Lipschitz continuous (which of course ensures its uniqueness from the classical comparison principle). See Lemma 5.1 below.

Proof of Theorem 1.12(i). In view of the proof of Theorem 1.7, the reader can check that it is enough to get a global Lipschitz bound on the solution u^ε and to construct a global corrector in this new framework. The proof of these two facts is postponed; see Lemmas 5.1 and 5.2 following this proof. Notice that half-relaxed limits are not necessary anymore and that the reasoning can be completed by considering locally converging subsequences of $\{u^\varepsilon\}$. Notice also that the perturbed test function method of [Evans 1989] still works. As usual, if the viscosity subsolution inequality is not satisfied at the limit, this implies that the perturbed test function is a supersolution except at times $\varepsilon(\mathbb{Z} + \{\tau_0, \dots, \tau_K\})$. Still, a localized comparison principle in each slice of times for each Cauchy problem is sufficient to conclude. \square

Lemma 5.1 (global Lipschitz bound). *The function u^ε is equi-Lipschitz continuous with respect to time and space.*

Proof. It is enough to get the result for $\varepsilon = 1$, since $u(t, x) = \varepsilon^{-1}u^\varepsilon(\varepsilon t, \varepsilon x)$ satisfies the equation with $\varepsilon = 1$ and the initial condition

$$u_0^\varepsilon(x) = \varepsilon^{-1}U_0^\varepsilon(\varepsilon x)$$

is equi-Lipschitz continuous. For the sake of clarity, we drop the ε superscript in u_0^ε and simply write u_0 .

We first derive bounds on the time interval $[\tau_0, \tau_1) = [0, \tau_1)$. In order to do so, we assume that the initial data satisfies $|(u_0)_x| \leq L$. Then, as usual, there is a constant $C > 0$ such that

$$u^\pm(t, x) = u_0(x) \pm Ct$$

are super- and sub-solutions of (1) and (10) with H given by (C1) with, for instance,

$$C := \max\left(\max_{\alpha=1, \dots, N} \|a_\alpha\|_\infty, \max_{\alpha=0, \dots, N} \left(\max_{|p| \leq L} |\bar{H}_\alpha(p)|\right)\right). \tag{49}$$

Let u be the standard (continuous) viscosity solution of (1) on the time interval $(0, \tau_1)$ with initial data given by u_0 (recall that $\varepsilon = 1$). Then, for any $h > 0$ small enough, we have $-Ch \leq u(h, x) - u(0, x) \leq Ch$. The comparison principle implies, for $t \in (0, \tau_1 - h)$,

$$-Ch \leq u(t + h, x) - u(t, x) \leq Ch,$$

which shows the Lipschitz bound in time, on the time interval $[0, \tau_1)$,

$$|u_t| \leq C. \tag{50}$$

From the Hamilton–Jacobi equation, we now deduce the following Lipschitz bound in space on the time interval $(0, \tau_1)$:

$$|\bar{H}_\alpha(u_x(t, \cdot))|_{L^\infty(b_\alpha, b_{\alpha+1})} \leq C \quad \text{for } \alpha = 0, \dots, N. \tag{51}$$

We can now derive bounds on the time interval $[\tau_1, \tau_2)$ as follows. We deduce first that (51) still holds true at time $t = \tau_1$. Combined with our definition (49) of the constant C , we also deduce that

$$v^\pm(t, x) = u(\tau_1, x) \pm C(t - \tau_1)$$

are sub- and super-solutions of (6) for $t \in (\tau_1, \tau_2)$, where H is given by (C1). Reasoning as above, we get bounds (50) and (51) on the time interval $[\tau_1, \tau_2)$.

Such reasoning can be used iteratively to get the Lipschitz bounds (50) and (51) for $t \in [0, +\infty)$. The proof of the lemma is now complete. \square

Lemma 5.2. *The conclusion of Theorem 4.6 still holds true in this new framework.*

Proof. The proof proceeds in several steps.

Step 1: construction of a time-periodic corrector w^ρ on $[-\rho, \rho]$. We first construct a Lipschitz corrector on a truncated domain. This, too, requires several steps.

Step 1.1: first Cauchy problem on $(0, +\infty)$. The method presented in the proof of Proposition 4.3, using a term δw^δ , has the inconvenience that it would not clearly provide a Lipschitz solution. In order to stick to our notion of globally Lipschitz solutions, we simply solve the Cauchy problem for $\rho > \rho_0 := \max_{\alpha=1, \dots, N} |b_\alpha|$,

$$\begin{cases} w_t^\rho + H(t, x, w_x^\rho) = 0 & \text{on } (0, +\infty) \times (-\rho, \rho), \\ w_t^\rho + \bar{H}_N^-(w_x^\rho) = 0 & \text{on } (0, +\infty) \times \{-\rho\}, \\ w_t^\rho + \bar{H}_0^+(w_x^\rho) = 0 & \text{on } (0, +\infty) \times \{\rho\}, \\ w^\rho(0, x) = 0 & \text{for } x \in [-\rho, \rho]. \end{cases} \quad (52)$$

As in the proof of the previous lemma, we get global Lipschitz bounds with a constant C (independent of $\rho > 0$ and the distances $\ell_\alpha = b_{\alpha+1} - b_\alpha$):

$$|w_t^\rho|, |\bar{H}_\alpha(w_x^\rho(t, \cdot))|_{L^\infty((b_\alpha, b_{\alpha+1}) \cap (-\rho, \rho))} \leq C \quad \text{for } \alpha = 0, \dots, N. \quad (53)$$

Arguing as in [Forcadel et al. 2009a], for instance, we deduce that there exists a real number λ_ρ with

$$|\lambda_\rho| \leq C$$

and a constant C_0 (that depends on ρ) such that

$$|w^\rho(t, x) + \lambda_\rho t| \leq C_0. \quad (54)$$

Details are given in Section D in the Appendix for the reader's convenience.

Step 1.2: getting global sub- and super-solutions. Let us now define the following function (up to some subsequence $k_n \rightarrow +\infty$):

$$w_\infty^\rho(t, x) = \lim_{k_n \rightarrow +\infty} (w^\rho(t + k_n, x) + \lambda_\rho k_n),$$

which still satisfies (53) and (54). Then we also define the two functions

$$\bar{w}_\infty^\rho(t, x) = \inf_{k \in \mathbb{Z}} (w_\infty^\rho(t + k, x) + k\lambda_\rho) \quad \text{and} \quad \underline{w}_\infty^\rho(t, x) = \sup_{k \in \mathbb{Z}} (w_\infty^\rho(t + k, x) + k\lambda_\rho).$$

They still satisfy (53) and (54) and are a super- and a sub-solution, respectively, of the problem in $\mathbb{R} \times [-\rho, \rho]$. They moreover satisfy that $\bar{w}_\infty^\rho(t, x) + \lambda_\rho t$ and $\underline{w}_\infty^\rho(t, x) + \lambda_\rho t$ are 1-periodic in time, which implies the bounds

$$|\bar{w}_\infty^\rho(t, x) - \bar{w}_\infty^\rho(0, x) + \lambda_\rho t| \leq C \quad \text{and} \quad |\underline{w}_\infty^\rho(t, x) - \underline{w}_\infty^\rho(0, x) + \lambda_\rho t| \leq C.$$

Step 1.3: a new Cauchy problem on $(0, +\infty)$ and construction of a time-periodic solution. We note that $\bar{w}_\infty^\rho + 2C_0 \geq \underline{w}_\infty^\rho$ and we now solve the Cauchy problem with new initial data $\underline{w}_\infty^\rho(0, x)$ instead of the zero initial data and call \tilde{w}^ρ the solution of this new Cauchy problem. From the comparison principle, we get

$$\underline{w}_\infty^\rho \leq \tilde{w}^\rho \leq \bar{w}_\infty^\rho + 2C_0.$$

In particular,

$$\tilde{w}^\rho(1, x) \geq \underline{w}_\infty^\rho(1, x) \geq \tilde{w}^\rho(0, x) - \lambda_\rho.$$

This implies, by comparison, that

$$\tilde{w}^\rho(k + 1, x) \geq \tilde{w}^\rho(k, x) - \lambda_\rho. \tag{55}$$

Moreover \tilde{w}^ρ still satisfies (53) (indeed with the same constant because, by construction, this is also the case for $\underline{w}_\infty^\rho$). We now define (up to some subsequence $k_n \rightarrow +\infty$)

$$\tilde{w}_\infty^\rho(t, x) = \lim_{k_n \rightarrow +\infty} (\tilde{w}^\rho(t + k_n, x) + \lambda_\rho k_n),$$

which, because of (55) and the fact that $\tilde{w}^\rho(t, x) + \lambda_\rho t$ is bounded, satisfies

$$\tilde{w}_\infty^\rho(k + 1, x) + \lambda = \tilde{w}_\infty^\rho(k, x)$$

and then $\tilde{w}_\infty^\rho(t, x) + \lambda_\rho t$ is 1-periodic in time. Moreover \tilde{w}_∞^ρ is still a solution of the Cauchy problem and satisfies (53). We define

$$w^\rho := \tilde{w}_\infty^\rho + \lambda_\rho t,$$

which satisfies (37) and then provides the analogue of the function given in Proposition 4.3.

Step 2: construction of w on \mathbb{R} . The result of Theorem 4.6 still holds true for

$$w = \lim_{\rho \rightarrow +\infty} (w^\rho - w^\rho(0, 0)),$$

which is globally Lipschitz continuous in space and time and satisfies (53) with $\rho = +\infty$, and

$$\bar{A} = \lim_{\rho \rightarrow +\infty} \lambda_\rho. \tag{□}$$

Proof of (12) from Theorem 1.12. We recall that $\bar{H}_L = \bar{H}_0$ and $\bar{H}_R = \bar{H}_1$ and set $a = a_1$ and (up to translation) $b_1 = 0$.

Step 1: the convex case; identification of \bar{A} .

Step 1.1: a convex subcase. We first work in the particular case where both \bar{H}_α for $\alpha = L, R$ are convex and given by the Legendre–Fenchel transform of convex Lagrangians L_α which satisfy, for some compact interval I_α ,

$$L_\alpha(p) = \begin{cases} \text{finite} & \text{if } q \in I_\alpha, \\ +\infty & \text{if } q \notin I_\alpha. \end{cases} \tag{56}$$

Then it is known (see for instance the section on optimal control in [IM]) that the solution of (1) on the time interval $[0, \varepsilon \tau_1)$ is given by

$$u^\varepsilon(t, x) = \inf_{y \in \mathbb{R}} \left(\inf_{X \in \mathcal{S}_{0,y;t,x}} \left\{ u^\varepsilon(0, X(0)) + \int_0^t L_\varepsilon(s, X(s), \dot{X}(s)) ds \right\} \right) \tag{57}$$

with

$$L_\varepsilon(s, x, p) = \begin{cases} \bar{H}_L^*(p) & \text{if } x < 0, \\ \bar{H}_R^*(p) & \text{if } x > 0, \\ \min(-a(\varepsilon^{-1}s), \min_{\alpha=L,R} L_\alpha(0)) & \text{if } x = 0, \end{cases}$$

and, for $s < t$, the set of trajectories

$$\mathcal{S}_{s,y;t,x} = \{X \in \text{Lip}((s, t); \mathbb{R}) : X(s) = y, X(t) = x\}.$$

Combining this formula with the other one on the time interval $[\varepsilon \tau_1, \varepsilon \tau_2)$, and iterating on all necessary intervals, we get that (57) is a representation formula of the solution u^ε of (1) for all $t > 0$. We also know (see the section on optimal control in [IM]), that the optimal trajectories from $(0, y)$ to (t_0, x_0) intersect the axis $x = 0$ at most on a time interval $[t_1^\varepsilon, t_2^\varepsilon]$ with $0 \leq t_1^\varepsilon \leq t_2^\varepsilon \leq t_0$. If this interval is not empty, then we have $t_i^\varepsilon \rightarrow t_i^0$ for $i = 1, 2$ and we can easily pass to the limit in (57). In general, u^ε converges to u^0 given by the formula

$$u^0(t, x) = \inf_{y \in \mathbb{R}} \left(\inf_{X \in \mathcal{S}_{0,y;t,x}} \left\{ u^0(0, X(0)) + \int_x^t L_0(s, X(s), \dot{X}(s)) ds \right\} \right)$$

with

$$L_0(s, x, p) = \begin{cases} \bar{H}_L^*(p) & \text{if } x < 0, \\ \bar{H}_R^*(p) & \text{if } x > 0, \\ \min(-\langle a \rangle, \min_{\alpha=L,R} L_\alpha(0)) & \text{if } x = 0, \end{cases}$$

and, from [IM], we see that u^0 is the unique solution of (6) and (2) with $\bar{A} = \langle a \rangle$.

Step 1.2: the general convex case. The general case of convex Hamiltonians is recovered, because, for Lipschitz continuous initial data u_0 , we know that the solution is globally Lipschitz continuous. Therefore, we can always modify the Hamiltonians \bar{H}_α outside some compact intervals so that the modified Hamiltonians satisfy (56).

Step 2: general quasiconvex Hamiltonians; identification of \bar{A} .

Step 2.1: subsolution inequality. From Theorem 2.10 in [IM], we know that $w(t, 0)$, as a function of time only, satisfies, in the viscosity sense,

$$w_t(t, 0) + a(t) \leq \bar{A} \quad \text{for all } t \notin \bigcup_{i=1, \dots, K+1} \tau_i + \mathbb{Z}.$$

Using the 1-periodicity in time of w , we see that the integration in time on one period implies

$$\langle a \rangle \leq \bar{A}. \tag{58}$$

Step 2.2: supersolution inequality. Recall that $\bar{A} \geq \langle a \rangle \geq A_0 := \max_{\alpha=L,R} \min(\bar{H}_\alpha)$. If $\bar{A} = A_0$, then obviously we get $\bar{A} = \langle a \rangle$. Hence, it remains to treat the case $\bar{A} > A_0$.

Step 2.3: construction of a supersolution for $x \neq 0$. Recall that \bar{p}_R and \bar{p}_L are defined in (8) and (9) and the minimum of \bar{H}_α is reached for \bar{p}_α^0 , $\alpha = R, L$. Since $\bar{A} > A_0$, there exists some $\delta > 0$ such that

$$\bar{p}_L + 2\delta < \bar{p}_L^0 \quad \text{and} \quad \bar{p}_R^0 < \bar{p}_R - 2\delta. \quad (59)$$

If w denotes a global corrector given by Lemma 5.2 (or Theorem 4.6), let us define

$$\underline{w}_R(t, x) = \inf_{h \geq 0} (w(t, x+h) - \bar{p}_R^0 h) \quad \text{for } x \geq 0$$

and similarly

$$\underline{w}_L(t, x) = \inf_{h \geq 0} (w(t, x-h) + \bar{p}_L^0 h) \quad \text{for } x \leq 0.$$

From (45) with $\rho_\delta = 0$, we deduce that we have, for some $\bar{h} \geq 0$,

$$w(t, x) \geq \underline{w}_R(t, x) = w(t, x + \bar{h}) - \bar{p}_R^0 \bar{h} \geq w(t, x) + (\bar{p}_R - \delta - \bar{p}_R^0) \bar{h} - C_\delta.$$

From (59), this implies

$$0 \leq \bar{h} \leq \frac{C_\delta}{\delta} \quad (60)$$

and, using the fact that w is globally Lipschitz continuous, we deduce that, for $\alpha = R$,

$$w \geq \underline{w}_\alpha \geq w - C_1. \quad (61)$$

Moreover, by construction — as an infimum of (globally Lipschitz continuous) supersolutions — \underline{w}_R is a (globally Lipschitz continuous) supersolution of the problem in $\mathbb{R} \times (0, +\infty)$. We also have, for $x = y + z$ with $z \geq 0$,

$$\begin{aligned} \underline{w}_R(t, x) - \underline{w}_R(t, y) &= w(t, x + \bar{h}) - \bar{p}_R^0 \bar{h} - \underline{w}_R(t, y) \\ &\geq w(t, x + \bar{h}) - \bar{p}_R^0 \bar{h} - (w(t, y + \bar{h} + z) - \bar{p}_R^0 (\bar{h} + z)) \\ &\geq \bar{p}_R^0 z = \bar{p}_R^0 (x - y), \end{aligned}$$

which shows that

$$(\underline{w}_R)_x \geq \bar{p}_R^0. \quad (62)$$

Similarly (and we can also use a symmetry argument to see it), we get that \underline{w}_L is a (globally Lipschitz continuous) supersolution in $\mathbb{R} \times (-\infty, 0)$, it satisfies (61) with $\alpha = L$, and

$$(\underline{w}_L)_x \leq \bar{p}_L^0. \quad (63)$$

We now define

$$\underline{w}(t, x) = \begin{cases} \underline{w}_R(t, x) & \text{if } x > 0, \\ \underline{w}_L(t, x) & \text{if } x < 0, \\ \min(\underline{w}_L(t, 0), \underline{w}_R(t, 0)) & \text{if } x = 0, \end{cases} \quad (64)$$

which, by construction is lower semicontinuous and satisfies (61), and is a supersolution for $x \neq 0$.

Step 2.4: checking the supersolution property at $x = 0$. Let φ be a test function touching \underline{w} from below at $(t_0, 0)$ with $t_0 \notin \bigcup_{i=1, \dots, K+1} \tau_i + \mathbb{Z}$. We want to check that

$$\varphi_t(t_0, 0) + F_{a(t_0)}(\varphi_x(t_0, 0^-), \varphi_x(t_0, 0^+)) \geq \bar{A}. \tag{65}$$

We may assume that

$$\underline{w}(t_0, 0) = \underline{w}_R(t_0, 0),$$

since the case $\underline{w}(t_0, 0) = \underline{w}_L(t_0, 0)$ is completely similar. Let $\bar{h} \geq 0$ be such that

$$\underline{w}_R(t_0, 0) = w(t_0, 0 + \bar{h}) - \bar{p}_R^0 \bar{h}.$$

We distinguish two cases. Assume first that $\bar{h} > 0$. Then we have, for all $h \geq 0$,

$$\varphi(t, 0) \leq w(t, 0 + h) - \bar{p}_R^0 h,$$

with equality for $(t, h) = (t_0, \bar{h})$. This implies the viscosity inequality

$$\varphi_t(t_0, 0) + \bar{H}_R(\bar{p}_R^0) \geq \bar{A},$$

which implies (65), because $F_{a(t_0)}(\varphi_x(t_0, 0^-), \varphi_x(t_0, 0^+)) \geq a(t_0) \geq A_0 \geq \min \bar{H}_R = \bar{H}_R(\bar{p}_R^0)$.

Assume now that $\bar{h} = 0$. Then we have $\varphi \leq \underline{w} \leq w$, with equality at $(t_0, 0)$. This immediately implies (65).

Step 2.5: conclusion. We deduce that \underline{w} is a supersolution on $\mathbb{R} \times \mathbb{R}$. Now let us consider a C^1 function $\psi(t)$ such that

$$\psi(t) \leq \underline{w}(t, 0),$$

with equality at $t = t_0$. Because of (62) and (63), we see that

$$\varphi(t, x) = \psi(t) + \bar{p}_L^0 x 1_{\{x < 0\}} + \bar{p}_R^0 x 1_{\{x > 0\}}$$

satisfies

$$\varphi \leq \underline{w},$$

with equality at $(t_0, 0)$. This implies (65) and, at almost every point t_0 where the Lipschitz continuous function $\underline{w}(t, 0)$ is differentiable, we have

$$\underline{w}_t(t_0, 0) + a(t_0) \geq \bar{A}.$$

Because w is 1-periodic in time, we get, after an integration on one period,

$$\langle a \rangle \geq \bar{A}. \tag{66}$$

Together with (58), we deduce that $\langle a \rangle = \bar{A}$, which is the desired result, for $N = 1$. □

Proof of (13) in Theorem 1.12. . We simply remark, using the subsolution viscosity inequality at each junction condition, that, for $\alpha = 1, \dots, N$,

$$\bar{A} \geq \langle a_\alpha \rangle,$$

which is the desired result. This achieves the proof of (12) and (13). \square

Proof of the monotonicity of \bar{A} in Theorem 1.12. Let $N \geq 2$ and, for $i = c, d$, let us assume some given $b_1^i < \dots < b_N^i$. and let us call w^i a global corrector given by Lemma 5.2 (or Theorem 4.6) with $\lambda = \bar{A}^i$ and $H = H^i$ for $i = c, d$, respectively.

We let $\ell_\alpha^i = b_{\alpha+1}^i - b_\alpha^i > 0$ and assume that

$$0 < \ell_{\alpha_0}^d - \ell_{\alpha_0}^c =: \delta_{\alpha_0} \quad \text{for some } \alpha_0 \in \{1, \dots, N - 1\}$$

and

$$\ell_\alpha^d = \ell_\alpha^c \quad \text{for all } \alpha \in \{1, \dots, N - 1\} \setminus \{\alpha_0\}.$$

Calling $\bar{p}_{\alpha_0}^0$ a point of global minimum of \bar{H}_{α_0} , we define

$$\tilde{w}^d(t, x) = \begin{cases} w^c(t, x - b_{\alpha_0}^d + b_{\alpha_0}^c) & \text{if } x \leq b_{\alpha_0}^d + \ell_{\alpha_0}^c/2 =: x_-, \\ w^c(t, x - b_{\alpha_0}^d + b_{\alpha_0}^c) + \bar{p}_{\alpha_0}^0(x - x_-) & \text{if } x_- \leq x \leq x_+, \\ w^c(t, x - b_{\alpha_0+1}^d + b_{\alpha_0+1}^c) + \bar{p}_{\alpha_0}^0(x_+ - x_-) & \text{if } x \geq b_{\alpha_0+1}^d - \ell_{\alpha_0}^c/2 =: x_+. \end{cases}$$

Recall that $w^i, i = c, d$, are globally Lipschitz continuous in space and time. This shows that \tilde{w}^d is also Lipschitz continuous in space and time by construction, because it is continuous at $x = x_-, x_+$. Moreover, \tilde{w}^d is 1-periodic in time. We now want to check that \tilde{w}^d is a subsolution of the equation satisfied by w^d with \bar{A}^c on the right-hand side instead of \bar{A}^d . We only have to check it for all times $\bar{t} \notin \{\tau_0, \dots, \tau_K\}$ and $\bar{x} \in [x_-, x_+]$, i.e., we have to show that

$$\tilde{w}_t^d(\bar{t}, \bar{x}) + \bar{H}_{\alpha_0}(\tilde{w}_x^d(\bar{t}, \bar{x})) \leq \bar{A}^c \quad \text{for all } \bar{x} \in [x_-, x_+]. \tag{67}$$

Assume that φ is a test function touching \tilde{w}^d from above at such a point (\bar{t}, \bar{x}) with $\bar{x} \in [x_-, x_+]$. Then this implies in particular that $\psi(t, x) = \varphi(t, x) - \bar{p}_{\alpha_0}^0(x - x_-)$ touches $\tilde{w}^d(\cdot, x_-) = w^c(\cdot, x_0)$ from above at time \bar{t} with $x_0 = b_{\alpha_0}^c + \ell_{\alpha_0}^c/2$. Recall that w^c is a solution of

$$w_t^c + \bar{H}_{\alpha_0}(w_x^c) = \bar{A}^c \quad \text{on } (b_{\alpha_0}^c, b_{\alpha_0+1}^c).$$

From the characterization of subsolutions (see Theorem 2.10 in [IM]), we then deduce that

$$\psi_t(\bar{t}) + \bar{H}_{\alpha_0}(\bar{p}_{\alpha_0}^0) \leq \bar{A}^c.$$

If $\bar{x} \in (x_-, x_+)$, then we have $\varphi_x(\bar{t}, \bar{x}) = \bar{p}_{\alpha_0}^0$. This means, in particular,

$$\varphi_t + \bar{H}_{\alpha_0}(\varphi_x) \leq \bar{A}^c \quad \text{at } (\bar{t}, \bar{x}) \text{ if } \bar{x} \in (x_-, x_+). \tag{68}$$

Now, using (68), and Theorem 2.10 in [IM] again, we deduce that we have, in the viscosity sense,

$$\tilde{w}_t^d(\bar{t}, \bar{x}) + \max(\bar{H}_{\alpha_0}^-(\tilde{w}_x^d(\bar{t}, \bar{x}^+)), \bar{H}_{\alpha_0}^+(\tilde{w}_x^d(\bar{t}, \bar{x}^-))) \leq \bar{A}^c \quad \text{for } \bar{x} = x_\pm. \tag{69}$$

Therefore, (68) and (69) imply (67).

Let us now call H^d the Hamiltonian in assumption (C1) constructed with the points $\{b_\alpha^d\}_{\alpha=1, \dots, N}$. Then we have

$$\tilde{w}_t^d + H^d(t, x, \tilde{w}_x^d) \leq \bar{A}^c \quad \text{for all } t \notin \{\tau_0, \dots, \tau_K\}.$$

Note that the proof of Theorem 1.4 is unchanged for the present problem and then Theorem 1.4 still holds true. This shows that

$$\bar{A}^d \leq \bar{A}^c, \tag{70}$$

which is the expected monotonicity. The proof is now complete. □

Remark 5.3. In the previous proof, it would also be possible to compare the subsolution given by the restriction of \tilde{w}^d on some interval $[-\rho, \rho]$ for $\rho > 0$ large enough (see Proposition 2.16 in [IM]) with the approximation $w^{d,\rho}$ of w^d on $[-\rho, \rho]$ with $\bar{A}^d \geq \bar{A}_\rho^d \rightarrow \bar{A}^d$ as $\rho \rightarrow +\infty$. The comparison for large times would imply $\bar{A}_\rho^d \leq \bar{A}^c$. As $\rho \rightarrow +\infty$, this would give the same conclusion (70).

Proof of (14) in Theorem 1.12. Let w be a global corrector associated to \bar{A} .

Recall that

$$\bar{A} \geq \bar{A}_0 := \max_{\alpha=1,\dots,N} \langle a_\alpha \rangle \geq A_0 := \max_{\alpha=1,\dots,N} A_0^\alpha \quad \text{with} \quad A_0^\alpha = \max_{\beta=\alpha-1,\alpha} (\min \bar{H}_\beta). \tag{71}$$

Our goal is to prove (14), i.e., that $\bar{A} = \bar{A}_0$ when all the distances ℓ_α are large enough. Let us assume that

$$\bar{A} > \bar{A}_0.$$

Step 1: considering another corrector with the same $\langle \hat{a}_\alpha \rangle = \bar{A}_0$. Let $\mu_\alpha \geq 0$ be such that

$$\hat{a}_\alpha = \mu_\alpha + a_\alpha \quad \text{with} \quad \langle \hat{a}_\alpha \rangle = \bar{A}_0 \quad \text{for all } \alpha = 1, \dots, N.$$

Let us call \hat{w} the corresponding corrector with associated constant \hat{A} . Then Theorem 1.4 (still valid here) implies that

$$\hat{A} \geq \bar{A} > \bar{A}_0.$$

We also split the set $\{1, \dots, N\}$ into two disjoint sets,

$$I_0 = \{\alpha \in \{1, \dots, N\} : \bar{A}_0 = A_0^\alpha\}$$

and

$$I_1 = \{\alpha \in \{1, \dots, N\} : \bar{A}_0 > A_0^\alpha\}.$$

Note that, by (71), if $\alpha \in I_0$ then $\langle a_\alpha \rangle = A_0^\alpha$ and then, by (C3), we have $a_\alpha(t) = \text{const} = A_0^\alpha$ for all $t \in \mathbb{R}$. For later use, we then claim that \hat{w} satisfies

$$\hat{w}_t(t, x) + \max(\bar{H}_\alpha^-(\hat{w}_x(t, x^+)), \bar{H}_{\alpha-1}^+(\hat{w}_x(t, x^-))) = \hat{A} \quad \text{for all } (t, x) \in \mathbb{R} \times \{b_\alpha\} \tag{72}$$

and not only for $t \in \mathbb{R} \setminus (\mathbb{Z} + \{\tau_0, \dots, \tau_K\})$. Let us show this for subsolutions (the proof being similar for supersolutions). Let φ be a test function touching \hat{w} from above at some point $(\bar{t}, \bar{x}) = (j + \tau_k, b_\alpha)$ for some $j \in \mathbb{Z}, k \in \{0, \dots, K\}$. Assume also that the contact between φ and \hat{w} only holds at that point (\bar{t}, \bar{x}) . The proof is a variant of a standard argument. For $\eta > 0$, let us consider the test function

$$\varphi_\eta(t, x) = \varphi(t, x) + \frac{\eta}{\bar{t} - t} \quad \text{for } t \in (-\infty, \bar{t}).$$

Then, for $r > 0$ fixed, we have

$$\inf_{\substack{(t,x) \in \overline{B_r(\bar{t}, \bar{x})} \\ t < \bar{t}}} (\varphi_\eta - \hat{w})(t, x) = (\varphi_\eta - \hat{w})(t_\eta, x_\eta)$$

with

$$\begin{cases} P_\eta = (t_\eta, x_\eta) \rightarrow (\bar{t}, \bar{x}) = \bar{P} & \text{as } \eta \rightarrow 0, \\ \varphi_t(\bar{P}) \leq \limsup_{\eta \rightarrow 0} (\varphi_\eta)_t(P_\eta). \end{cases}$$

This implies that \hat{w} is a relaxed viscosity subsolution at (\bar{t}, \bar{x}) in the sense of Definition 2.2 in [IM]. By Proposition 2.5 in [IM], we deduce that \hat{w} is also a standard (i.e., not relaxed) viscosity subsolution at (\bar{t}, \bar{x}) . Finally, we get (72).

Step 2: defining a space supersolution. Let us define the function

$$M(x) = \inf_{t \in \mathbb{R}} \hat{w}(t, x).$$

Because \hat{w} is globally Lipschitz continuous, we deduce that M is also globally Lipschitz continuous. Moreover, we have the viscosity supersolution inequality

$$\bar{H}_\alpha(M_x(x)) \geq \hat{A} > \bar{A}_0 \quad \text{for all } x \in (b_\alpha, b_{\alpha+1}), \alpha = 0, \dots, N.$$

Let us call, for $\alpha = 0, \dots, N$,

$$\begin{aligned} \bar{p}_{\alpha,R} &= \min E_{\alpha,R} & \text{with } E_{\alpha,R} &= \{p \in \mathbb{R} : \bar{H}_\alpha^+(p) = \bar{H}_\alpha(p) = \bar{A}_0\}, \\ \bar{p}_{\alpha,L} &= \max E_{\alpha,L} & \text{with } E_{\alpha,L} &= \{p \in \mathbb{R} : \bar{H}_\alpha^-(p) = \bar{H}_\alpha(p) = \bar{A}_0\}. \end{aligned}$$

Let us now consider $\alpha = 0, \dots, N$ and two points $x_- < x_+$ with $x_\pm \in (b_\alpha, b_{\alpha+1})$. Let us assume that there is a test function φ^\pm touching M from below at x_\pm . Then we have

$$\bar{H}_\alpha(\varphi_x^\pm(x_\pm)) \geq \hat{A} > \bar{A}_0$$

with

$$\varphi_x^\pm(x_\pm) \geq \bar{p}_{\alpha,R} \quad \text{or} \quad \varphi_x^\pm(x_\pm) \leq \bar{p}_{\alpha,L}.$$

Moreover, if $\bar{A}_0 > \min \bar{H}_\alpha$, then we have

$$\bar{p}_{\alpha,L} < \bar{p}_\alpha^0 < \bar{p}_{\alpha,R}$$

for any \bar{p}_α^0 which is a point of global minimum of \bar{H}_α .

Step 3: a property of the space supersolution. We now claim that the following case is impossible:

$$p^- := \varphi_x^-(x_-) < \varphi_x^+(x_+) =: p^+ \quad \text{and} \quad \inf_{[p^-, p^+]} \bar{H}_\alpha < \hat{A}.$$

Indeed, if $\bar{p} \in (p^-, p^+)$ is such that $\bar{H}_\alpha(\bar{p}) < \hat{A}$, then the geometry of the graph of the function M implies that

$$\inf_{x \in [x_-, x_+]} (M(x) - x \bar{p}) = M(\bar{x}) - \bar{x} \bar{p} \quad \text{for some } \bar{x} \in (x_-, x_+)$$

and then we have the viscosity supersolution inequality, at \bar{x} ,

$$\bar{H}_\alpha(\bar{p}) \geq \hat{A},$$

which leads to a contradiction. Therefore (in either case, $\bar{A}_0 > \min \bar{H}_\alpha$ or $\bar{A}_0 = \min \bar{H}_\alpha$), it is possible to check that there is a point $\bar{x}_\alpha \in [b_\alpha, b_{\alpha+1}]$ such that the Lipschitz continuous function M satisfies, in the viscosity sense,

$$\begin{cases} M_x \geq \bar{p}_{\alpha,R} & \text{in } (b_\alpha, \bar{x}_\alpha), \\ -M_x \geq -\bar{p}_{\alpha,L} & \text{in } (\bar{x}_\alpha, b_{\alpha+1}). \end{cases}$$

Moreover, from Theorem 4.6(ii) (see Lemma 5.2), we deduce from $\hat{A} > \max(\min H_N, \min H_0)$ that

$$\bar{x}_N = +\infty \quad \text{and} \quad \bar{x}_0 = -\infty.$$

In particular, we deduce that there exists at least one $\alpha_0 \in \{1, \dots, N\}$ such that

$$\bar{x}_{\alpha_0} - b_{\alpha_0} \geq \frac{1}{2}\ell_{\alpha_0} \quad \text{and} \quad b_{\alpha_0} - \bar{x}_{\alpha_0-1} \geq \frac{1}{2}\ell_{\alpha_0-1}. \tag{73}$$

Step 4: the case $\alpha_0 \in I_0$. In this case, we see that there exists a time \bar{t} such that the test function

$$\varphi(t, x) = \begin{cases} \bar{p}_{\alpha_0,R}(x - b_{\alpha_0}) & \text{for } x \geq b_{\alpha_0}, \\ \bar{p}_{\alpha_0-1,L}(x - b_{\alpha_0}) & \text{for } x \leq b_{\alpha_0}, \end{cases}$$

is a test function touching (up to some additive constant) \hat{w} from below at (\bar{t}, b_{α_0}) . By (72), this implies

$$\bar{A}_0 = \max(\bar{H}_{\alpha_0}(\bar{p}_{\alpha_0,R}), \bar{H}_{\alpha_0-1}(\bar{p}_{\alpha_0-1,L})) \geq \hat{A} \geq \bar{A}.$$

This is a contradiction.

Step 5: consequences on \hat{w} . From the fact that \hat{w} is 1-periodic in time and C -Lipschitz continuous in time (with a constant C depending only on $\max_{\alpha=1,\dots,N} \|\hat{a}_\alpha\|_\infty$ and the \bar{H}_α ; see (49)), we deduce that we have

$$\begin{cases} \hat{w}(t, x+h) - \hat{w}(t, x) \geq \bar{p}_{\alpha,R}h - 2C & \text{for } x, x+h \in (b_\alpha, \bar{x}_\alpha), \\ \hat{w}(t, x-h) - \hat{w}(t, x) \geq -\bar{p}_{\alpha,R}h - 2C & \text{for } x, x+h \in (\bar{x}_\alpha, b_{\alpha+1}). \end{cases} \tag{74}$$

Step 6: the case $\alpha_0 \in I_1$; definition of a spacetime supersolution. Proceeding similarly to Step 3 of the proof of (12), we define

$$\underline{\hat{w}}_{\alpha_0,R}(t, x) = \inf_{\ell_{\alpha_0}/4 \geq h \geq 0} (\hat{w}(t, x+h) - \bar{p}_{\alpha_0}^0 h) \quad \text{for } b_{\alpha_0} \leq x \leq b_{\alpha_0} + \frac{1}{4}\ell_{\alpha_0}$$

and

$$\underline{\hat{w}}_{(\alpha_0-1),L}(t, x) = \inf_{\ell_{(\alpha_0-1)}/4 \geq h \geq 0} (\hat{w}(t, x-h) + \bar{p}_{\alpha_0-1}^0 h) \quad \text{for } b_{\alpha_0} - \frac{1}{4}\ell_{\alpha_0-1} \leq x \leq b_{\alpha_0}.$$

From (74), we deduce that we have, for some $\bar{h} \in [0, \frac{1}{4}\ell_{\alpha_0}]$,

$$\hat{w}(t, x) \geq \underline{\hat{w}}_{\alpha_0,R}(t, x) = \hat{w}(t, x + \bar{h}) - \bar{p}_{\alpha_0}^0 \bar{h} \geq \hat{w}(t, x) + (\bar{p}_{\alpha_0,R} - \bar{p}_{\alpha_0}^0)\bar{h} - 2C,$$

which implies

$$0 \leq \bar{h} \leq \frac{2C}{\bar{p}_{\alpha_0,R} - \bar{p}_{\alpha_0}^0}.$$

As in Step 3 of the proof of (12), if

$$\frac{\ell_{\alpha_0}}{4} > \frac{2C}{\bar{p}_{\alpha_0,R} - \bar{p}_{\alpha_0}^0}, \tag{75}$$

this implies that $\hat{w}_{\alpha_0,R}$ is a supersolution for $x \in (b_{\alpha_0}, b_{\alpha_0} + \frac{1}{4}\ell_{\alpha_0})$. Similarly, if

$$\frac{\ell_{\alpha_0-1}}{4} > \frac{2C}{\bar{p}_{\alpha_0-1}^0 - \bar{p}_{\alpha_0-1,L}}, \tag{76}$$

then $\hat{w}_{\alpha_0-1,L}$ is a supersolution for $x \in (b_{\alpha_0} - \frac{1}{4}\ell_{\alpha_0-1}, b_{\alpha_0})$. We now define

$$\hat{w}(t, x) = \begin{cases} \hat{w}_{\alpha_0,R}(t, x) & \text{if } x \in (b_{\alpha_0}, b_{\alpha_0} + \frac{1}{4}\ell_{\alpha_0}), \\ \hat{w}_{\alpha_0-1,L}(t, x) & \text{if } x \in (b_{\alpha_0} - \frac{1}{4}\ell_{\alpha_0-1}, b_{\alpha_0}), \\ \min(\hat{w}_{\alpha_0-1,L}(t, b_{\alpha_0}), \hat{w}_{\alpha_0,R}(t, b_{\alpha_0})) & \text{if } x = b_{\alpha_0}. \end{cases}$$

Then, as in Steps 4 and 5 of the proof of (12), we deduce that \hat{w} is a supersolution up to the junction point $x = b_{\alpha_0}$ and that

$$\bar{A}_0 = \langle \hat{a}_{\alpha_0} \rangle \geq \hat{A} \geq \bar{A}.$$

This is a contradiction.

Step 7: conclusion. If (75) and (76) hold true for any $\alpha_0 \in I_1$, then we deduce that $\bar{A} \leq \bar{A}_0$, which implies $\bar{A} = \bar{A}_0$. This ends the proof of (14) in Theorem 1.12. \square

Proof of (15) in Theorem 1.12. Let us consider

$$\bar{a}(t) = \max_{\alpha=1, \dots, N} a_\alpha(t)$$

and (w, \bar{A}) a solution (given by Theorem 4.6 (see also Lemma 5.2)) of

$$\begin{cases} w_t + \bar{H}_0(w_x) = \bar{A} & \text{if } x < 0, \\ w_t + \bar{H}_N(w_x) = \bar{A} & \text{if } x > 0, \\ w_t(t, 0) + \max(\bar{a}(t), \bar{H}_N^-(w_x(t, 0^+)), \bar{H}_0^+(w_x(t, 0^-))) = \bar{A} & \text{if } x = 0, \\ w \text{ is 1-periodic in } t. \end{cases}$$

From Theorem 1.12, we also know that

$$\bar{A} = \langle \bar{a} \rangle.$$

For $N \geq 2$, we set $\ell = (\ell_1, \dots, \ell_{N-1}) \in (0, +\infty)^{N-1}$ and consider $b_0 = -\infty < b_1 < \dots < b_N < b_{N+1} = +\infty$ with

$$\ell_\alpha = b_{\alpha+1} - b_\alpha \quad \text{for } \alpha = 1, \dots, N - 1.$$

We now call (w^ℓ, \bar{A}^ℓ) a global corrector, given by Theorem 4.6 (see also Lemma 5.2). The remainder of the proof is divided into several steps.

Step 1: bound from above on \bar{A}^ℓ . We define

$$\tilde{w}(t, x) = \begin{cases} w(t, x - b_1) & \text{if } x \leq b_1, \\ w(t, 0) + \bar{p}_\alpha^0(x - b_\alpha) + \sum_{\beta=1, \dots, \alpha-1} \bar{p}_\beta^0(b_{\beta+1} - b_\beta) & \text{if } b_\alpha \leq x \leq b_{\alpha+1}, \alpha \in \{1, \dots, N-1\}, \\ w(t, x - b_N) + \sum_{\beta=1, \dots, N-1} \bar{p}_\beta^0(b_{\beta+1} - b_\beta) & \text{if } x \geq b_N. \end{cases}$$

Proceeding as in Step 1 of the proof of Theorem 1.12(ii), it is easy to check that \tilde{w} is a subsolution of the equation satisfied by w^ℓ with \bar{A} on the right-hand side instead of \bar{A}^ℓ . Then Theorem 1.4 implies that

$$\bar{A}^\ell \leq \bar{A} = \langle \bar{a} \rangle. \tag{77}$$

Step 2: bound from below on \bar{A}^ℓ . From Theorem 2.10 in [IM], we deduce that we have, in the viscosity sense (in time only),

$$w_t^\ell(t, b_\alpha) + a_\alpha(t) \leq \bar{A}^\ell \quad \text{for all } t \notin \bigcup_{k=0}^K \{\tau_k + \mathbb{Z}\}.$$

Let us call

$$\underline{A} = \liminf_{\ell \rightarrow 0} \bar{A}^\ell.$$

We also know that w^ℓ is 1-periodic and globally Lipschitz continuous with a constant which is independent of ℓ . Therefore, there exists a 1-periodic and Lipschitz continuous function $g = g(t)$ such that

$$w^\ell(t, b_\alpha) \rightarrow g(t) \quad \text{as } \ell \rightarrow 0 \quad \text{for all } \alpha = 1, \dots, N.$$

The stability of viscosity solutions implies, in the viscosity sense,

$$g'(t) + a_\alpha(t) \leq \underline{A} \quad \text{for all } \alpha = 1, \dots, N \text{ and } t \notin \bigcup_{k=0}^K \{\tau_k + \mathbb{Z}\}.$$

Because g is Lipschitz continuous, this inequality also holds for almost every $t \in \mathbb{R}$. This implies

$$g'(t) + \bar{a}(t) \leq \underline{A} \quad \text{for a.e. } t \in \mathbb{R}.$$

An integration on one period gives

$$\langle \bar{a} \rangle \leq \underline{A}. \tag{78}$$

Step 3: conclusion. Combining (77) with (78) finally yields that $\bar{A}^\ell \rightarrow \langle \bar{a} \rangle$ as $\ell \rightarrow 0$. The proof of (15) in Theorem 1.12 is now complete. □

Appendix: Proofs of some technical results

A. The case $\bar{x} \neq 0$ in the proof of convergence. We only deal with the subcase $\bar{x} > 0$, since the subcase $\bar{x} < 0$ is treated in the same way. Reducing \bar{r} if necessary, we may assume that $B_{\bar{r}}(\bar{t}, \bar{x})$ is compactly embedded in the set $\{(t, x) \in (0, +\infty) \times (0, +\infty) : x > 0\}$, because there exists a positive constant $c_{\bar{r}}$ such that

$$(t, x) \in B_{\bar{r}}(\bar{t}, \bar{x}) \implies x > c_{\bar{r}}. \tag{79}$$

Let $p = \varphi_x(\bar{t}, \bar{x})$ and let $v^R = v^R(t, x)$ be a solution of the cell problem

$$v_t^R + H_R(t, x, p + v_x^R) = \bar{H}_R(p) \quad \text{in } \mathbb{R} \times \mathbb{R}. \tag{80}$$

We claim that, if $\varepsilon > 0$ is small enough, the perturbed test function [Evans 1989]

$$\varphi^\varepsilon(t, x) = \varphi\left(t, x\right) + \varepsilon v^R\left(\frac{t}{\varepsilon}, \frac{x}{\varepsilon}\right)$$

satisfies, in the viscosity sense, the inequality

$$\varphi_t^\varepsilon + H\left(\frac{t}{\varepsilon}, \frac{x}{\varepsilon}, \varphi_x^\varepsilon\right) \geq \frac{\theta}{2} \quad \text{in } B_r(\bar{t}, \bar{x}) \tag{81}$$

for sufficiently small $r > 0$. To see this, let ψ be a test function touching φ^ε from below at (t_1, x_1) in $B_r(\bar{t}, \bar{x}) \subseteq B_{\bar{r}}(\bar{t}, \bar{x})$. In this way, the function

$$\eta(s, y) = \frac{1}{\varepsilon}(\psi(\varepsilon s, \varepsilon y) - \varphi(\varepsilon s, \varepsilon y))$$

touches v^R from below at $(s_1, y_1) = (t_1/\varepsilon, x_1/\varepsilon)$ and (80) yields

$$\psi_t(t_1, x_1) - \varphi_t(t_1, x_1) + H_R\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, p + \psi_x(t_1, x_1) - \varphi_x(t_1, x_1)\right) \geq \bar{H}_R(p). \tag{82}$$

Since (79) implies that $x/\varepsilon \rightarrow +\infty$, as $\varepsilon \rightarrow 0$, uniformly with respect to $(t, x) \in B_{\bar{r}}(\bar{t}, \bar{x})$, we can find, owing to (A5), an $\varepsilon_0 > 0$ independent of ψ and (t_1, x_1) such that the inequality

$$H\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \psi_x(t_1, x_1)\right) \geq H_R\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \psi_x(t_1, x_1)\right) - \frac{\theta}{4} \tag{83}$$

holds true for $\varepsilon < \varepsilon_0$. Combining (19), (82) and (83) and using the continuity of φ_x and φ_t , we have

$$\begin{aligned} & \psi_t(t_1, x_1) + H\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \psi_x(t_1, x_1)\right) \\ & \geq \psi_t(t_1, x_1) + H_R\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, p + \psi_x(t_1, x_1) - \varphi_x(t_1, x_1)\right) \\ & \quad + H_R\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \psi_x(t_1, x_1)\right) - H_R\left(\frac{t_1}{\varepsilon}, \frac{x_1}{\varepsilon}, \varphi_x(\bar{t}, \bar{x}) + \psi_x(t_1, x_1) - \varphi_x(t_1, x_1)\right) - \frac{\theta}{4} \\ & \geq \frac{\theta}{2} \end{aligned}$$

if r is sufficiently close to 0. The claim (81) is proved.

Since φ is strictly above \bar{u} , if ε and r are small enough then

$$u^\varepsilon + \kappa_r \leq \varphi^\varepsilon \quad \text{on } \partial B_r(\bar{t}, \bar{x})$$

for a suitable positive constant κ_r . By the comparison principle we deduce

$$u^\varepsilon + \kappa_r \leq \varphi^\varepsilon \quad \text{in } B_r(\bar{t}, \bar{x})$$

and, passing to the limit as $\varepsilon \rightarrow 0$ and $(t, x) \rightarrow (\bar{t}, \bar{x})$ on both sides of the previous inequality, we produce the contradiction

$$\bar{u}(\bar{t}, \bar{x}) < \bar{u}(\bar{t}, \bar{x}) + \kappa_r \leq \varphi(\bar{t}, \bar{x}) = \bar{u}(\bar{t}, \bar{x}).$$

B. Proof of Lemma 3.3. We first address uniqueness. Let us assume that we have two solutions of (3), (v^i, λ^i) for $i = 1, 2$. Let

$$u^i(t, x) = v^i(t, x) + px - \lambda^i t.$$

Then u^i solves

$$u_t^i + H_\alpha(t, x, u_x^i) = 0$$

with

$$u^1(0, x) \leq u^2(0, x) + C.$$

The comparison principle implies

$$u^1 \leq u^2 + C \quad \text{for all } t > 0$$

and then $\lambda^1 \geq \lambda^2$. Similarly, we get the reverse inequality and then $\lambda^1 = \lambda^2$.

We now turn to the continuity of the map $p \mapsto \bar{H}_\alpha(p)$. It follows from the stability of viscosity sub- and super-solutions, from the fact that the constant C in (24) is bounded for bounded p and from the comparison principle. This achieves the proof of the lemma.

C. Sketch of the proof of Proposition 4.1. Consider

$$M_\nu = \sup_{\substack{x \in [\rho_1, \rho_2] \\ s, t \in \mathbb{R}}} \left\{ u(t, x) - v(s, x) - \frac{(t-s)^2}{2\nu} \right\}.$$

We want to prove that

$$M = \lim_{\nu \rightarrow 0} M_\nu \leq 0.$$

We argue by contradiction by assuming that $M > 0$. The supremum defining M_ν is reached; let s_ν, t_ν and x_ν denote a maximizer. Choose ν small enough so that $M_\nu \geq \frac{1}{2}M > 0$. We classically get

$$|t_\nu - s_\nu| \leq C\sqrt{\nu}.$$

If there exists $\nu_n \rightarrow 0$ such that $x_{\nu_n} = \rho_1$ for all $n \in \mathbb{N}$, then

$$\frac{1}{2}M \leq M_{\nu_n} \leq U_0(t_{\nu_n}) - U_0(s_{\nu_n}) \leq \omega_0(t_{\nu_n} - s_{\nu_n}) \leq \omega_0(C\sqrt{\nu_n}),$$

where ω_0 denotes the modulus of continuity of U_0 . The contradiction $M \leq 0$ is obtained by letting n go to $+\infty$.

Hence, we can assume that, for ν small enough, $x_\nu > \rho_1$. Reasoning as in [IM, Theorem 7.8], we can easily reduce to the case where $H(t_\nu, x_\nu, \cdot)$ reaches its minimum for $p = p_0 = 0$. We can also consider the vertex test function G^γ associated with the single Hamiltonian H (using the notation of [IM], it corresponds to the case $N = 1$) and the free parameter γ . If $x_\nu < \rho_2$, then $G^\gamma(x, y)$ reduces to the standard test function $\frac{1}{2}(x - y)^2$.

We next consider

$$M_{v,\varepsilon} = \sup_{\substack{x,y \in [\rho_1, \rho_2] \cap \overline{B_r(x_v)} \\ s,t \in \mathbb{R}}} \left\{ u(t, x) - v(s, y) - \frac{(t-s)^2}{2\nu} - \varepsilon G^\gamma(\varepsilon^{-1}x, \varepsilon^{-1}y) - \varphi^v(t, s, x) \right\},$$

where $r = r_v$ is chosen so that $\rho_1 \notin \overline{B_r(x_v)}$ and φ^v is the localization function

$$\varphi^v(t, s, x) = \frac{1}{2}((t - t_v)^2 + (s - s_v)^2 + (x - x_v)^2).$$

The supremum defining $M_{v,\varepsilon}$ is reached and, if (t, s, x, y) denotes a maximizer, then

$$(t, s, x, y) \rightarrow (t_v, s_v, x_v, x_v) \quad \text{as } (\varepsilon, \gamma) \rightarrow 0.$$

In particular, $x, y \in B_r(x_v)$ for ε and γ small enough. The remaining of the proof is completely analogous (in fact much simpler).

D. Construction of λ_ρ in the proof of Lemma 5.2. In order to get λ_ρ , it is enough to apply the following lemma:

Lemma D.1. *Let u be the solution of a Hamilton–Jacobi equation of evolution type subject to the initial condition $u(0, x) = 0$ and posed on a compact set K . Assume that:*

- *the comparison principle holds true;*
- *u is L -globally Lipschitz continuous in time and space;*
- *$u(k + \cdot, \cdot) + C$ is a solution for all $k \in \mathbb{N}$ and $C \in \mathbb{R}$.*

There then exists $\lambda \in \mathbb{R}$ such that

$$|u(t, x) - \lambda t| \leq C_0$$

and

$$|\lambda| \leq L,$$

where $C_0 = L(2 + 3\rho)$ if ρ denotes the diameter of K .

Proof. Define

$$\lambda^+(T) = \sup_{\tau \geq 0} \frac{u(\tau + T, 0) - u(\tau, 0)}{T} \quad \text{and} \quad \lambda^-(T) = \inf_{\tau \geq 0} \frac{u(\tau + T, 0) - u(\tau, 0)}{T}.$$

Note that $T \mapsto \pm T\lambda^\pm(T)$ is subadditive. The fact that u is L -Lipschitz continuous with respect to time implies that $\lambda^\pm(T)$ are both finite:

$$|\lambda^\pm(T)| \leq L.$$

The ergodic theorem implies that $\lambda^\pm(T)$ converges towards λ^\pm and

$$\lambda^+ = \inf_{T>0} \lambda^+(T) \quad \text{and} \quad \lambda^- = \sup_{T>0} \lambda^-(T).$$

If, moreover,

$$|\lambda^+(T) - \lambda^-(T)| \leq \frac{C}{T}, \tag{84}$$

then the proof of the lemma is complete. Indeed, (84) implies in particular that $\lambda^+ = \lambda^-$ and

$$-\frac{C}{T} \leq \lambda^-(T) - \lambda \leq \lambda^+(T) - \lambda \leq \frac{C}{T}.$$

This implies that $|u(t, 0) - \lambda t| \leq C$. Finally, we get

$$|u(t, x) - \lambda t| \leq C + L\rho.$$

It remains to prove (84). There exists $k \in \mathbb{Z}$ and $\beta \in [0, 1)$ such that $\tau^+ = k + \tau^- + \beta$. Moreover,

$$u(\tau^+, x) \leq u(\tau^- + \beta, x) + u(\tau^+, 0) - u(\tau^- + \beta, 0) + 2L\rho,$$

where $\rho = \text{diam } K$. Now note that $u(\tau^- + \beta + t, x) + D$ is a solution in $[\tau^+, +\infty)$ for all constant D . Hence, we get by comparison that, for all $t > 0$ and $x \in K$,

$$u(\tau^+ + t, x) \leq u(\tau^- + \beta + t, x) + u(\tau^+, 0) - u(\tau^- + \beta, 0) + 2L\rho.$$

In particular,

$$u(\tau^+ + T, 0) - u(\tau^+, 0) \leq u(\tau^- + \beta + T, 0) - u(\tau^- + \beta, 0) + 2L\rho \leq u(\tau^- + T, 0) - u(\tau^-, 0) + 2L(1 + \rho).$$

Finally, we get (after letting $\varepsilon \rightarrow 0$)

$$\lambda^+(T) \leq \lambda^-(T) + \frac{2L(1 + \rho)}{T}.$$

Similarly, we can get

$$\lambda^+(T) \geq \lambda^-(T) - \frac{2L(1 + \rho)}{T}.$$

This implies (84) with $C = 2L(1 + \rho)$. The proof of the lemma is now complete. \square

Acknowledgements

The authors thank the referees for their valuable comments. The authors thank Y. Achdou, K. Han and N. Tchou for stimulating discussions. The authors thank N. Seguin for interesting discussions on green waves. Imbert thanks Giga for the interesting discussions they had together and for drawing his attention towards papers such as [Hamamuki 2013]. Monneau thanks G. Costeseque for his comments on traffic lights modelling and his simulations, which inspired certain complementary results. Imbert and Monneau are partially supported by ANR-12-BS01-0008-01 HJnet project.

References

- [Achdou and Tchou 2015] Y. Achdou and N. Tchou, “Hamilton–Jacobi equations on networks as limits of singularly perturbed problems in optimal control: dimension reduction”, *Comm. Partial Differential Equations* **40**:4 (2015), 652–693. MR 3299352 Zbl 06448676
- [Andreianov et al. 2010] B. Andreianov, P. Goatin, and N. Seguin, “Finite volume schemes for locally constrained conservation laws”, *Numer. Math.* **115**:4 (2010), 609–645. MR 2011e:35212 Zbl 1196.65151
- [Bardi and Capuzzo-Dolcetta 1997] M. Bardi and I. Capuzzo-Dolcetta, *Optimal control and viscosity solutions of Hamilton–Jacobi–Bellman equations*, Birkhäuser, Boston, 1997. MR 99e:49001 Zbl 0890.49011

- [Barles and Souganidis 2000] G. Barles and P. E. Souganidis, “On the large time behavior of solutions of Hamilton–Jacobi equations”, *SIAM J. Math. Anal.* **31**:4 (2000), 925–939. MR 2002b:49056 Zbl 0960.70015
- [Barles et al. 2013] G. Barles, A. Briani, and E. Chasseigne, “A Bellman approach for two-domains optimal control problems in \mathbb{R}^N ”, *ESAIM Control Optim. Calc. Var.* **19**:3 (2013), 710–739. MR 3092359 Zbl 1287.49028
- [Bernard and Roquejoffre 2004] P. Bernard and J.-M. Roquejoffre, “Convergence to time-periodic solutions in time-periodic Hamilton–Jacobi equations on the circle”, *Comm. Partial Differential Equations* **29**:3–4 (2004), 457–469. MR 2005d:35028 Zbl 1072.35055
- [Bourgoing 2008a] M. Bourgoing, “Viscosity solutions of fully nonlinear second order parabolic equations with L^1 dependence in time and Neumann boundary conditions”, *Discrete Contin. Dyn. Syst.* **21**:3 (2008), 763–800. MR 2009h:35204 Zbl 1152.35411
- [Bourgoing 2008b] M. Bourgoing, “Viscosity solutions of fully nonlinear second order parabolic equations with L^1 dependence in time and Neumann boundary conditions: existence and applications to the level-set approach”, *Discrete Contin. Dyn. Syst.* **21**:4 (2008), 1047–1069. MR 2010b:35214 Zbl 1165.35401
- [Evans 1989] L. C. Evans, “The perturbed test function method for viscosity solutions of nonlinear PDE”, *Proc. Roy. Soc. Edinburgh Sect. A* **111**:3–4 (1989), 359–375. MR 91c:35017 Zbl 0679.35001
- [Forcadel et al. 2009a] N. Forcadel, C. Imbert, and R. Monneau, “Homogenization of fully overdamped Frenkel–Kontorova models”, *J. Differential Equations* **246**:3 (2009), 1057–1097. MR 2009j:35022 Zbl 1171.49023
- [Forcadel et al. 2009b] N. Forcadel, C. Imbert, and R. Monneau, “Homogenization of some particle systems with two-body interactions and of the dislocation dynamics”, *Discrete Contin. Dyn. Syst.* **23**:3 (2009), 785–826. MR 2010h:35021 Zbl 1154.35306
- [Forcadel et al. 2012] N. Forcadel, C. Imbert, and R. Monneau, “Homogenization of accelerated Frenkel–Kontorova models with n types of particles”, *Trans. Amer. Math. Soc.* **364**:12 (2012), 6187–6227. MR 2958933 Zbl 1276.35024
- [Giga and Hamamuki 2013] Y. Giga and N. Hamamuki, “Hamilton–Jacobi equations with discontinuous source terms”, *Comm. Partial Differential Equations* **38**:2 (2013), 199–243. MR 3009078 Zbl 1263.35066
- [Hamamuki 2013] N. Hamamuki, “On large time behavior of Hamilton–Jacobi equations with discontinuous source terms”, pp. 83–112 in *Nonlinear analysis in interdisciplinary sciences—modellings, theory and simulations*, edited by T. Aiki et al., GAKUTO Internat. Ser. Math. Sci. Appl. **36**, Gakkōtoshō, Tokyo, 2013. MR 3205346
- [Imbert and Monneau 2013] C. Imbert and R. Monneau, “Flux-limited solutions for quasi-convex Hamilton–Jacobi equations on networks”, preprint, 2013. arXiv 1306.2428
- [Imbert et al. 2013] C. Imbert, R. Monneau, and H. Zidani, “A Hamilton–Jacobi approach to junction problems and application to traffic flows”, *ESAIM Control Optim. Calc. Var.* **19**:1 (2013), 129–166. MR 3023064 Zbl 1262.35080
- [Jin and Yu 2015] W.-L. Jin and Y. Yu, “Asymptotic solution and effective Hamiltonian of a Hamilton–Jacobi equation in the modeling of traffic flow on a homogeneous signalized road”, *J. de Math. Pures et Appliquées* (online publication July 2015).
- [Kawohl 1985] B. Kawohl, *Rearrangements and convexity of level sets in PDE*, Lecture Notes in Mathematics **1150**, Springer, Berlin, 1985. MR 87a:35001 Zbl 0593.35002
- [Lions 1981] P.-L. Lions, “Two geometrical properties of solutions of semilinear problems”, *Applicable Anal.* **12**:4 (1981), 267–272. MR 83e:35051 Zbl 0445.35043
- [Lions 2013–2014] P.-L. Lions, “Elliptic or parabolic equations, and improved homogenization”, lectures, Collège de France, 2013–2014, <http://www.college-de-france.fr/site/en-pierre-louis-lions/course-2013-2014.htm>.
- [Lions et al. 1986] P.-L. Lions, G. Papanicolaou, and S. R. S. Varadhan, “Homogeneization of Hamilton–Jacobi equations”, 1986, http://www.math.unipd.it/~bardi/didattica/Nonlinear_PDE_%20homogenization_Dott._%202011/LPV87.pdf. Unpublished.

Received 8 Sep 2014. Revised 16 Mar 2015. Accepted 11 May 2015.

GIULIO GALISE: ggalise@unisa.it

Department of Mathematics, University of Salerno, Via Giovanni Paolo II, 132, I-84084 Fisciano (SA), Italy

CYRIL IMBERT: cyril.imbert@math.cnrs.fr

CNRS, UMR 7580, CNRS, Université Paris-Est Créteil, 61 avenue du Général de Gaulle, 94010 Paris Créteil, France

RÉGIS MONNEAU: monneau@cermics.enpc.fr

CERMICS (ENPC), Université Paris-Est, 6–8 Avenue Blaise Pascal, Cité Descartes, F-77455 Champs-sur-Marne Marne-la-Vallée Cedex 2, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Maciej Zworski
zworski@math.berkeley.edu
University of California
Berkeley, USA

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Institute of Technology, USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2015 is US \$205/year for the electronic version, and \$390/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 8 No. 8 2015

- Semilinear wave equations on asymptotically de Sitter, Kerr–de Sitter and Minkowski space- 1807
times
PETER HINTZ and ANDRÁS VASY
- A junction condition by specified homogenization and application to traffic lights 1891
GIULIO GALISE, CYRIL IMBERT and RÉGIS MONNEAU
- Existence and classification of singular solutions to nonlinear elliptic equations with a gradient 1931
term
JOSHUA CHING and FLORICA CÎRSTEĂ
- A topological join construction and the Toda system on compact surfaces of arbitrary genus 1963
ALEKS JEVIKAR, SADOK KALLEL and ANDREA MALCHIODI
- Well-posedness and scattering for the Zakharov system in four dimensions 2029
IOAN BEJENARU, ZIHUA GUO, SEBASTIAN HERR and KENJI NAKANISHI



2157-5045(2015)8:8;1-6