

ANALYSIS & PDE

Volume 9

No. 4

2016

MARIUS BECEANU

**DISPERSIVE ESTIMATES IN \mathbb{R}^3
WITH THRESHOLD EIGENSTATES AND RESONANCES**



DISPERSIVE ESTIMATES IN \mathbb{R}^3 WITH THRESHOLD EIGENSTATES AND RESONANCES

MARIUS BECEANU

We prove dispersive estimates in \mathbb{R}^3 for the Schrödinger evolution generated by the Hamiltonian $H = -\Delta + V$, under optimal decay conditions on V , in the presence of zero-energy eigenstates and resonances.

1. Introduction	813
1A. Classification of exceptional Hamiltonians	813
1B. Main result	814
1C. History of the problem	815
2. Proof of the statements	817
2A. Notations	817
2B. Auxiliary results	818
2C. Wiener spaces	822
2D. Regular points and regular Hamiltonians	824
2E. Exceptional Hamiltonians of the first kind	828
2F. Exceptional Hamiltonians of the third kind	840
Acknowledgments	857
References	857

1. Introduction

1A. Classification of exceptional Hamiltonians. Consider a Hamiltonian of the form $H = -\Delta + V$, where V is a real-valued scalar potential on \mathbb{R}^3 .

We assume $V \in L^{\frac{3}{2},1} \subset L^{\frac{3}{2}}$, which is the predual of weak- L^3 and a Lorentz space, $L^{\frac{3}{2},1} \subset L^{\frac{3}{2}-\epsilon} \cap L^{\frac{3}{2}+\epsilon}$; for its definition and properties, see [Bergh and Löfström 1976]. By [Simon 1982], this is sufficient to guarantee the self-adjointness of $H = -\Delta + V$.

Let $R_0(\lambda) := (-\Delta - \lambda)^{-1}$ be the free resolvent corresponding to the free evolution $e^{-it\Delta}$ and let $R_V(\lambda) := (-\Delta + V - \lambda)^{-1}$ be the perturbed resolvent corresponding to the perturbed evolution e^{itH} . Explicitly, in three dimensions and for $\text{Im } \lambda \geq 0$,

$$R_0((\lambda + i0)^2)(x, y) = \frac{1}{4\pi} \frac{e^{i\lambda|x-y|}}{|x-y|}. \tag{1-1}$$

The author was partially supported by a Rutgers Research Council grant.

MSC2010: primary 35J10; secondary 47D08.

Keywords: pointwise decay estimates, resonances, zero-energy eigenfunctions.

It will be shown below that, under reasonable assumptions, H has only finitely many negative eigenvalues. Then the Schrödinger evolution restricted to the continuous spectrum $[0, \infty)$ has the representation formula

$$e^{itH} P_c = \lim_{\epsilon \rightarrow 0} \frac{1}{2\pi i} \int_0^\infty e^{it\eta} (R_V(\eta + i\epsilon) - R_V(\eta - i\epsilon)) d\eta.$$

By the work of Ionescu and Jerison [2003] and Goldberg and Schlag [2004b], it is known that, when $V \in L^{\frac{3}{2}}$, the perturbed resolvent $R_V(\lambda \pm i0)$ is uniformly bounded in $\mathcal{B}(L^{\frac{6}{5}}, L^6)$ on any interval $\lambda \in [\epsilon_0, \infty)$, where $\epsilon_0 > 0$, and has no singularities in $[0, \infty)$ except potentially at $\lambda = 0$.

Observe that $R_V = (I + R_0 V)^{-1} R_0$, so R_V has a singularity at zero precisely when $I + R_0(0)V$, which is a compact perturbation of the identity, is not invertible.

We denote the null space of $I + R_0(0)V$ by

$$\mathcal{M} := \{\phi \in L^\infty \mid \phi + R_0(0)V\phi = 0\}.$$

If $\mathcal{M} \neq \emptyset$, we say that H is of *exceptional type*, while if $\mathcal{M} = \emptyset$, we say that H is of *generic type*.

The sesquilinear form $-\langle u, Vv \rangle$ is an inner product on \mathcal{M} ; see Lemma 2.2. This pairing is well-defined when $V \in L^{\frac{3}{2},1}$ because $u, v \in L^{3,\infty} \cap L^\infty$ by Lemma 2.1.

Let $\mathcal{E} := \mathcal{M} \cap L^2$ and P_0 be the orthogonal L^2 projection onto \mathcal{E} . In Lemma 2.3, we provide a characterization of \mathcal{E} and show that $\text{codim}_{\mathcal{M}} \mathcal{E} \leq 1$.

The set $\mathcal{E}_1 := \mathcal{E} \cap L^1$ also plays a special part in the proof. In Lemma 2.5, we give a characterization of \mathcal{E}_1 and prove that $\text{codim}_{\mathcal{E}} \mathcal{E}_1 \leq 12$.

A function $\phi \in \mathcal{M} \setminus \mathcal{E}$ is called a zero-energy resonance of H . Following [Jensen and Kato 1979; Yajima 2005], we classify exceptional Hamiltonians H as follows:

- (1) H is of exceptional type of the first kind if it has a zero-energy resonance, but no zero-energy eigenfunctions: $\{0\} = \mathcal{E} \subsetneq \mathcal{M}$.
- (2) H is of exceptional type of the second kind if it has zero-energy eigenfunctions, but no zero-energy resonance: $\{0\} \subsetneq \mathcal{E} = \mathcal{M}$.
- (3) H is of exceptional type of the third kind if it has both resonances and eigenfunctions at zero energy: $\{0\} \subsetneq \mathcal{E} \subsetneq \mathcal{M}$.

1B. Main result. When H is of exceptional type of the first kind, we let the *canonical resonance* be $\phi \in \mathcal{M}$ such that $\langle V, \phi \rangle > 0$ and $-\langle \phi, V\phi \rangle = 1$ (one can make these choices by Lemmas 2.3 and 2.2, respectively).

Using the canonical resonance $\phi(x)$, we define a constant a and a function $\zeta_t(x)$ by

$$a = \frac{4\pi i}{|\langle V, \phi \rangle|^2}, \quad \zeta_t(x) = e^{\frac{i|x|^2}{4t}} \phi(x).$$

We also define a function $\mu_t(x)$ by

$$\mu_t(x) := \frac{i}{|x|} \int_0^1 (e^{\frac{i|x|^2}{4t}} - e^{\frac{i\theta x|^2}{4t}}) d\theta.$$

Let the operators $R(t)$ and $S(t)$ be given by

$$\begin{aligned} R(t) &:= \frac{ae^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} \zeta_t(x) \otimes \zeta_t(y), \\ S(t) &:= \frac{e^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} \left(-iP_0V \frac{|x-y|^2}{24\pi} VP_0 + \mu_t(x) \frac{|x-y|}{8\pi} VP_0 + P_0V \frac{|x-y|}{8\pi} \mu_t(y) \right). \end{aligned} \quad (1-2)$$

Note that

$$\|R(t)u\|_{L^{3,\infty}} + \|S(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

Proposition 1.1 (main result). *Assume that $\langle x \rangle^2 V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is exceptional of the first kind. Then, for $1 \leq p < \frac{3}{2}$ and any $u \in L^2 \cap L^p$,*

$$e^{-itH} P_c u = Z(t)u + R(t)u, \quad \|Z(t)u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|f\|_{L^p},$$

where p' is the dual exponent, that is, $\frac{1}{p} + \frac{1}{p'} = 1$. Furthermore, assuming only that $V \in L^{\frac{3}{2},1}$, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH} P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|u\|_{L^p}, \quad \|e^{-itH} P_c u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

Assume that $\langle x \rangle^4 V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is exceptional of the second or third kind. Then, for $1 \leq p < \frac{3}{2}$ and any $u \in L^2 \cap L^p$,

$$e^{-itH} P_c u = Z(t)u + R(t)u + S(t)u, \quad \|Z(t)u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|u\|_{L^p}, \quad (1-3)$$

where $R(t)$ is missing if H is an exceptional Hamiltonian of the second kind.

In the case when all the zero-energy eigenfunctions of H are in L^1 , one can omit $S(t)$ from (1-3).

Assume that $\langle x \rangle^2 V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is exceptional of the second or third kind. Then, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH} P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|u\|_{L^p}, \quad \|e^{-itH} P_c u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

Note that, in terms of powers of x , the decay conditions on the potential correspond to $|V| \lesssim \langle x \rangle^{-2-}$, $|V| \lesssim \langle x \rangle^{-4-}$, and $|V| \lesssim \langle x \rangle^{-6-}$.

Additionally, note that these decay estimates also imply a certain range of Strichartz estimates.

The rest of the paper is dedicated to proving this main result, which is a combination of Propositions 2.13, 2.15, 2.18, and 2.19. For brevity, we omit the proof in the case when H is an exceptional Hamiltonian of the second kind, which is similar to the case when H is exceptional of the third kind.

1C. History of the problem. We study solutions to the linear Schrödinger equation in \mathbb{R}^3 with potential

$$i \partial_t u + \Delta u - V u = 0, \quad u(0) \text{ given.}$$

By the RAGE theorem, every solution is the sum of a bound and a scattering component. The quantitative study of scattering states began with Rauch [1978], who proved that if $H = -\Delta + gV$, where $g \in \mathbb{C}$,

with exponentially decaying V , then $e^{itH} P_c$ has a local decay rate of $t^{-\frac{3}{2}}$, with at most a discrete set of exceptional g for which the decay rate is $t^{-\frac{1}{2}}$. Here P_c is the projection on the space of scattering solutions.

Threshold estimates in the presence of eigenvalues and resonances go back to the work of Jensen and Kato [1979], who obtained an asymptotic expansion of the resolvent $R(\zeta) = (H - \zeta)^{-1}$ into

$$R(\zeta) = -\zeta^{-1} B_{-2} - i\zeta^{-\frac{1}{2}} B_{-1} + B_0 + i\zeta^{\frac{1}{2}} B_1 + \dots$$

and similar ones for the spectral density and the S -matrix. The condition imposed on the potential was polynomial decay at infinity of the form $(1 + |x|^\beta)V(x) \in L^{\frac{3}{2}}(\mathbb{R}^3)$, where $\beta > 2$.

The possible singularities in this expansion are due to the presence of resonances or eigenstates at zero. B_{-2} is the L^2 orthogonal projection on the zero eigenspace, while B_{-1} is given by

$$B_{-1} = P_0 V \frac{|x - y|^2}{24\pi} V P_0 - \phi \otimes \phi,$$

where ϕ is the canonical zero resonance; see above.

Jensen and Kato also obtained an asymptotic expansion for the evolution $e^{itH} P_c$ in two cases: if zero is a regular point, then

$$e^{itH} P_c = -(4\pi i)^{-\frac{1}{2}} t^{-\frac{3}{2}} B_0 + o(t^{-\frac{3}{2}}),$$

and if there is only a resonance ϕ at zero then

$$e^{itH} P_c = (\pi i)^{-\frac{1}{2}} t^{-\frac{1}{2}} \phi \otimes \phi + o(t^{-\frac{1}{2}}).$$

Murata [1982] extended these results by obtaining an asymptotic expansion to any order, for a more general evolution, with or without singular points, and then proving that each term in the expansion is degenerate. Murata’s expansion and proof are valid in weighted L^2 spaces.

Erdoğ an and Schlag [2004] obtained an asymptotic expansion of the evolution $e^{itH} P_c$ in the pointwise L^1 -to- L^∞ setting using the Jensen–Nenciu lemma [2001]. The condition assumed for the potential was that $|V(x)| \lesssim \langle x \rangle^{-12-\epsilon}$. The same method works in the case of nonselfadjoint Hamiltonians (see [Erdoğ an and Schlag 2006]) of the form

$$\mathcal{H} = \begin{pmatrix} -\Delta + \mu + V_1 & V_2 \\ -V_2 & \Delta - \mu - V_1 \end{pmatrix},$$

assuming that $|V_1(x)| + |V_2(x)| \lesssim \langle x \rangle^{-10-\epsilon}$.

At the same time, Yajima [2005] proved a similar expansion for generic Hamiltonians $H = -\Delta + V$ when $|V(x)| \leq \langle x \rangle^{-\frac{5}{2}-\epsilon}$, for singular Hamiltonians of the first kind when $|V(x)| \leq \langle x \rangle^{-\frac{9}{2}-\epsilon}$, and of the second and third kind when $|V(x)| \leq \langle x \rangle^{-\frac{11}{2}-\epsilon}$. His main result stated the following:

Theorem 1.2 [Yajima 2005, Theorem 1.3]. (1) *Let V satisfy $|V(x)| \leq C \langle x \rangle^{-\beta}$ for some $\beta > \frac{5}{2}$. Suppose that H is of generic type. Then, for any $1 \leq q \leq 2 \leq p \leq \infty$ such that $\frac{1}{p} + \frac{1}{q} = 1$,*

$$\|e^{-itH} P_c u\|_p \leq C_p t^{-3(\frac{1}{2}-\frac{1}{p})} \|u\|_q, \quad \text{where } u \in L^2 \cap L^q. \tag{1-4}$$

(2) Let V satisfy $|V(x)| \leq C\langle x \rangle^{-\beta}$ for some $\beta > \frac{11}{2}$. Suppose that H is of exceptional type. Then the following statements are satisfied:

- (a) Estimate (1-4) holds when p and q are restricted to $\frac{3}{2} < q \leq 2 \leq p < 3$ and $\frac{1}{p} + \frac{1}{q} = 1$.
- (b) Estimate (1-4) holds when $p = 3$ and $q = \frac{3}{2}$ provided that L^3 and $L^{\frac{3}{2}}$ are respectively replaced by Lorentz spaces $L^{3,\infty}$ and $L^{\frac{3}{2},1}$.
- (c) When $3 < p \leq \infty$ and $1 \leq q < \frac{3}{2}$ are such that $\frac{1}{p} + \frac{1}{q} = 1$, there exists a constant C_{pq} such that, for any $u \in L^2 \cap L^q$,

$$\|(e^{-itH} P_c - R(t) - S(t))u\|_p \lesssim C_{pq} t^{-3\left(\frac{1}{2} - \frac{1}{p}\right)} \|u\|_q.$$

If H is of exceptional type of the first kind, statement (2) holds under a weaker decay condition $|V(x)| \leq C\langle x \rangle^{-\beta}$ with $\beta > \frac{9}{2}$.

However, note that, due to a mistake in the proof, the requirement $\beta > \frac{11}{2}$ should be replaced by $\beta > 8$.

When the zero-energy eigenfunctions ϕ_k of H have enough decay, both $R(t)$ and $S(t)$ can be taken to be zero. Indeed, Goldberg [2010] showed that if $V \in L^{\frac{3}{2}-\epsilon} \cap L^{\frac{3}{2}+\epsilon}$ and the zero-energy eigenfunctions are in L^1 then $\|e^{-itH} P_c u\|_{L^\infty} \lesssim t^{-\frac{3}{2}} \|u\|_{L^1}$. We retrieve a similar result in our context.

Some of our results for exceptional potentials of the first kind hold under the same decay assumption as those for generic potentials: $V \in L^{\frac{3}{2},1}$. A similar fact was also recently noticed by Egorova, Kopylova, Marchenko and Teschl [Egorova et al. 2014] in dimension one.

Several results [Journé et al. 1991; Goldberg and Schlag 2004a; Goldberg 2006; Beceanu and Goldberg 2012] address the issue of pointwise decay in the case of generic Hamiltonians — for $L^{\frac{3}{2}-\epsilon} \cap L^{\frac{3}{2}+\epsilon}$ potentials in [Goldberg 2006] and Kato-class potentials in [Beceanu and Goldberg 2012].

Results obtained in other dimensions include [Cardoso et al. 2009; Egorova et al. 2014; Erdoğan et al. 2014; Erdoğan and Green 2010; 2013a; 2013b; 2013c; Goldberg 2007; Goldberg and Green 2014; 2015; Green 2012; Schlag 2005].

The current result, Proposition 1.1, represents an improvement on [Yajima 2005] by half a power of potential decay for exceptional Hamiltonians of the first kind. We expect the rate of potential decay from Proposition 1.1 to be optimal for this sort of result.

The same considerations apply in the case of exceptional Hamiltonians of the second and third kind, also leading to similar improved results. These will constitute the subject of a separate paper.

Below we mostly follow the scheme of Yajima's proof [2005], making the changes from Hölder spaces to Wiener spaces needed to improve the result. The proof method that we use here is the same as in [Beceanu 2011; Beceanu and Goldberg 2012].

2. Proof of the statements

2A. Notations. We denote the usual Lebesgue spaces by L^p and the Lorentz spaces by $L^{p,q}$, where $1 \leq p, q \leq \infty$. Note here that $L^{p,p} = L^p$, $L^{p,\infty}$ is weak- L^p , and $L^{p,q_1} \subset L^{p,q_2}$ for $q_1 \leq q_2$. For the definition and further properties, see [Bergh and Löfström 1976].

Let Sobolev spaces be $W^{s,p}$, where $s \in \mathbb{R}$ and $1 \leq p \leq \infty$, and denote weighted Lebesgue spaces by $f(x)L^p = \{f(x)g(x) \mid g \in L^p\}$.

Fix the Fourier transform to

$$\hat{f}(\xi) = \int_{\mathbb{R}^d} e^{-ix\xi} f(x) dx, \quad \check{f}(x) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{i\xi x} f(\xi) d\xi.$$

Let $R_0(\lambda) := (-\Delta - \lambda)^{-1}$ and for $\lambda \in \mathbb{R}$,

$$R_{0a}(\lambda) := \frac{1}{i} (R_0(\lambda + i0) - R_0(\lambda - i0)).$$

Concerning the Fourier transform, resolvents, and the free evolution, note that with our definitions

$$\begin{aligned} e^{itH_0} &= (R_{0a}(\lambda))^\vee(t), \\ R_{0a}(\lambda) &= (e^{itH_0})^\wedge \quad \text{for } \lambda \in \mathbb{R}, \\ iR_0(\lambda) &= (\chi_{[0,\infty)}(t)e^{itH_0})^\wedge(\lambda) \quad \text{for } \text{Im } \lambda < 0. \end{aligned}$$

Likewise let $R_V(\lambda) := (-\Delta + V - \lambda)^{-1}$.

Also, let

- χ_A be the characteristic function of the set A ;
- \mathcal{M} be the space of finite-mass Borel measures on \mathbb{R} ;
- δ_x denote Dirac's measure at x ;
- $\langle x \rangle = (1 + |x|^2)^{\frac{1}{2}}$;
- $\mathcal{B}(X, Y)$ be the Banach space of bounded operators from X to Y and $\mathcal{B}(X)$ be the Banach space of bounded operators from X to itself;
- C be any constant (not always the same throughout the paper);
- $a \lesssim b$ mean $|a| \leq C|b|$;
- \mathcal{S} be the Schwartz space;
- $u \otimes v$ mean the rank-one operator $\langle \cdot, v \rangle u$;
- $K(x, y)$ denote the operator having $K(x, y)$ as its integral kernel.

For a potential V , let $V_1 = |V|^{\frac{1}{2}}$ and $V_2 = |V|^{\frac{1}{2}} \text{sgn } V$.

2B. Auxiliary results. Recall that \mathcal{M} is the kernel of $I + R_0(0)V$ in L^∞ .

Lemma 2.1. *Let $V \in L^{\frac{3}{2},1}$; then $\mathcal{M} \subset L^{3,\infty}$. Conversely, any $\phi \in L^{3,\infty}$ that satisfies the equation $\phi + R_0(0)V\phi = 0$ must be in L^∞ , hence in \mathcal{M} .*

Proof of Lemma 2.1. Let $V = V^1 + V^2$, where V^1 is smooth of compact support and $\|V^2\|_{L^{3/2,1}} \ll 1$. Then, if ϕ solves the equation

$$\begin{aligned} \phi &= -(I + R_0(0)V^2)^{-1} R_0(0)V^1\phi \\ &= -\left(\sum_{k=0}^{\infty} (-1)^k (R_0(0)V^2)^k \right) R_0(0)V^1\phi, \end{aligned}$$

where the inverse is the sum of a Neumann series, and thus is bounded on $L^{3,\infty}$ and on L^∞ .

If $\phi \in L^\infty$, then $V^1\phi \in L^1$; hence $R_0(0)V^1\phi \in L^{3,\infty}$, so $\phi \in L^{3,\infty}$.

If $\phi \in L^{3,\infty}$, then $V^1\phi \in L^{\frac{3}{2},1}$; hence $R_0(0)V^1\phi \in L^\infty$, so $\phi \in L^\infty$. \square

Lemma 2.2. *The quadratic form $-\langle u, Vv \rangle$ is an inner product on \mathcal{M} .*

Proof. Suppose that $u, v \in \mathcal{M}$. By the definition of \mathcal{M} , observe that $-\langle u, Vv \rangle = \langle u, -\Delta v \rangle$, where $u \in L^{3,\infty} \cap L^\infty$ by Lemma 2.1 and $-\Delta v = Vv \in L^1 \cap L^{\frac{3}{2},1}$. Thus the pairing is well-defined.

Furthermore, $\nabla u = \nabla R_0(0)Vu \in L^{\frac{3}{2},\infty} \cap L^{3,\infty} \subset L^2$ and the same holds for ∇v , so their pairing is also well-defined and we can write $(u, -\Delta v) = (\nabla u, \nabla v)$.

This expression is positively defined because, setting $u = v$, the equation $\langle \nabla u, \nabla u \rangle = 0$ implies that u is constant; hence, in view of the fact that $u \in L^{3,\infty}$ by Lemma 2.1, $u = 0$. \square

Recall that $\mathcal{E} = \mathcal{M} \cap L^2$.

Lemma 2.3. *Assume that $V \in L^{\frac{3}{2},1}$. Then, for any $\phi \in \mathcal{M}$, we have $\phi(x) \in \langle x \rangle^{-1}L^\infty$.*

Assume that $V \in L^1 \cap L^{\frac{3}{2},1}$. Then, for any $\phi \in \mathcal{M}$, we have

$$\phi(x) - \frac{\langle \phi, V \rangle}{4\pi|x|} \in |x|^{-1}L^{3,\infty} \cap |x|^{-1}L^\infty \subset L^2.$$

Thus $\phi \in \mathcal{M}$ is in \mathcal{E} if and only if $\langle \phi, V \rangle = 0$; thus $\text{codim}_{\mathcal{M}} \mathcal{E} \leq 1$. Also, $\mathcal{E} \subset \langle x \rangle^{-2}L^\infty$.

Proof of Lemma 2.3. First, assume that $V \in L^{\frac{3}{2},1}$. Rewrite the eigenfunction equation

$$\phi(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{1}{|x-y|} V(y)\phi(y) dy$$

as

$$\begin{aligned} |x|\phi(x) + \frac{1}{4\pi} \int_{|y| \geq R} \frac{|x| - |x-y|}{|x-y||y|} V(y)|y|\phi(y) dy \\ = -\frac{1}{4\pi} \int_{\mathbb{R}^3} V(y)\phi(y) dy - \frac{1}{4\pi} \int_{|y| \leq R} \frac{|x| - |x-y|}{|x-y|} V(y)\phi(y) dy. \end{aligned}$$

Note that $||x| - |x-y|| \leq |y|$ and $\lim_{R \rightarrow \infty} \|\chi_{|x| \geq R}(x)V(x)\|_{L^{3/2,1}} = 0$. Then, for sufficiently large R , we can invert

$$(T_0\phi)(x) = \phi(x) + \frac{1}{4\pi} \int_{|y| \geq R} \frac{|x| - |x-y|}{|x-y||y|} V(y)\phi(y) dy$$

as an operator in $\mathcal{B}(L^\infty)$. Since $\phi(y) \in L^{3,\infty} \cap L^\infty$, the right-hand side is in L^∞ , so we obtain that $|x|\phi(x) \in L^\infty$.

Next, assume that $V \in L^1 \cap L^{\frac{3}{2},1}$. Start from

$$\begin{aligned} \phi(x) - \frac{\langle \phi, V \rangle}{4\pi|x|} &= -\frac{1}{4\pi} \int_{\mathbb{R}^3} \left(\frac{1}{|x-y|} - \frac{1}{|x|} \right) V(y)\phi(y) dy \\ &= -\frac{1}{4\pi|x|} \int_{\mathbb{R}^3} \frac{|x| - |x-y|}{|x-y|} V(y)\phi(y) dy, \end{aligned}$$

which is bounded in absolute value by

$$\frac{1}{4\pi|x|} \int_{\mathbb{R}^3} \frac{|y||V(y)||\phi(y)|}{|x-y|} dy.$$

Since $\phi \in \langle x \rangle^{-1} L^\infty$ and $V \in L^1 \cap L^{\frac{3}{2},1}$, this expression is in $|x|^{-1} L^\infty \cap |x|^{-1} L^{3,\infty} \subset \langle x \rangle^{-1} L^{3,\infty} \subset L^2$.

Since whenever $\langle \phi, V \rangle \neq 0$ we have $\langle \phi, V \rangle / (4\pi|x|) \notin L^2$, it follows that for ϕ to be in L^2 it is necessary and sufficient that $\langle \phi, V \rangle = 0$.

The space \mathcal{E} is then the kernel of the rank-one map $\phi \mapsto \langle \phi, V \rangle$ from \mathcal{M} to \mathbb{C} , so it has codimension at most 1.

Finally, we already know that $\mathcal{E} \subset \mathcal{M} \subset \langle x \rangle^{-1} L^\infty$. The eigenfunction equation for a function $\phi \in \mathcal{E}$ for which $\langle \phi, V \rangle = 0$, can be written as

$$\phi(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{|x| - |x-y|}{|x-y||x|} V(y) \phi(y) dy.$$

We further rewrite it as

$$\begin{aligned} |x|^2 \phi(x) + \frac{1}{4\pi} \int_{|y| \geq R} \frac{(|x| - |x-y|)^2}{|x-y||y|^2} V(y) |y|^2 \phi(y) dy \\ = -\frac{1}{4\pi} \int_{\mathbb{R}^3} (|x| - |x-y|) V(y) \phi(y) dy - \frac{1}{4\pi} \int_{|y| \leq R} \frac{(|x| - |x-y|)^2}{|x-y|} V(y) \phi(y) dy. \end{aligned}$$

The right-hand side is in L^∞ and, for sufficiently large R , the left-hand side is invertible, as above. This shows that $|x|^2 \phi(x) \in L^\infty$. \square

We can continue the asymptotic expansion of eigenfunctions to any order, but first we need the following lemma.

Lemma 2.4. For $x, y \in \mathbb{R}^3$,

$$\left| \frac{1}{|x-y|} - \left(\frac{1}{|x|} + \frac{xy}{|x|^3} \right) \right| \lesssim \frac{|y|^2}{|x|^2|x-y|} \quad (2-1)$$

and

$$\left| \frac{1}{|x-y|} - \left(\frac{1}{|x|} + \frac{xy}{|x|^3} + \frac{|y|^2}{2|x|^3} - \frac{3(xy)^2}{2|x|^5} \right) \right| \lesssim \frac{|y|^3}{|x|^3|x-y|}. \quad (2-2)$$

More generally, it seems to be the case (one can prove by induction) that

$$\left| \frac{1}{|x+y|} - \sum_{k=0}^N d^k \frac{1}{|x-\cdot|} (y, \dots, y) \right| \lesssim \frac{|y|^{N+1}}{|x|^{N+1}|x-y|}.$$

Proof of Lemma 2.4. Indeed, we start from

$$(|x|^2 + 2xy + |y|^2)^{\frac{1}{2}} - (|x|^2)^{\frac{1}{2}} = \frac{2xy}{|x+y|+|x|} + \frac{|y|^2}{|x+y|+|x|}. \quad (2-3)$$

Then

$$\left| \frac{2xy}{|x+y|+|x|} - \frac{xy}{|x|} \right| = \left| \frac{xy(|x|-|x+y|)}{(|x+y|+|x|)|x|} \right| \lesssim \frac{|y|^2}{|x|}.$$

Therefore,

$$\left| |x+y| - |x| - \frac{xy}{|x|} \right| \lesssim \frac{|y|^2}{|x|}. \quad (2-4)$$

Consequently,

$$|x|^2(|x| - |x - y|) - xy|x - y| \leq |x|^2 \left| |x - y| - |x| + \frac{xy}{|x|} \right| + |xy(|x| - |x - y|)| \lesssim |y|^2|x|.$$

Dividing by $|x|^3|x - y|$, we obtain (2-1).

We next perform a more detailed analysis of the same inequality. In (2-3), by (2-4) we have

$$\begin{aligned} & \left| \frac{xy(|x| - |x + y|)}{(|x + y| + |x|)|x|} + \frac{(xy)^2}{2|x|^3} \right| \\ & \lesssim \left| \frac{xy(|x| - |x + y|)}{(|x + y| + |x|)|x|} - \frac{xy(|x| - |x + y|)}{2|x|^2} \right| + \left| \frac{xy\left(\frac{xy}{|x|} + |x| - |x + y|\right)}{2|x|^2} \right| \lesssim \frac{|y|^3}{|x|^2}. \end{aligned}$$

Furthermore, also in (2-3),

$$\frac{|y|^2}{|x + y| + |x|} - \frac{|y|^2}{2|x|} \lesssim \frac{|y|^3}{|x|^2}.$$

Therefore,

$$\left| |x + y| - |x| - \frac{xy}{|x|} - \frac{|y|^2}{2|x|} + \frac{(xy)^2}{2|x|^3} \right| \lesssim \frac{|y|^3}{|x|^2}. \quad (2-5)$$

By (2-4) and (2-5), we then obtain (2-2). \square

We can now establish the asymptotic expansion of eigenfunctions.

Lemma 2.5. *Assume that $V \in L^1 \cap L^{\frac{3}{2},1}$. Let $\phi \in \mathcal{E}$ be a zero-energy eigenfunction of H . Then*

$$\phi(x) - \sum_{k=1}^3 \langle V\phi, y_k \rangle \frac{x_k}{|x|^3} \in |x|^{-2}(L^{3,\infty} \cap L^\infty).$$

Further assume that $V \in \langle x \rangle^{-1}L^1 \cap L^{\frac{3}{2},1}$. Then

$$\phi(x) - \sum_{k=1}^3 \langle V\phi, y_k \rangle \frac{x_k}{|x|^3} - \sum_{k,\ell=1}^3 \langle \phi V, y_k y_\ell \rangle \left(\frac{\delta_{k\ell}}{2|x|^3} - \frac{3x_k x_\ell}{2|x|^5} \right) \in |x|^{-3}(L^{3,\infty} \cap L^\infty).$$

In particular, $\phi \in \mathcal{E}$ is in L^1 if and only if $\langle V\phi, y_k \rangle = 0$ and $\langle \phi V, y_k y_\ell \rangle = 0$ for $1 \leq k, \ell \leq 3$.

Let $\mathcal{E}_1 := \mathcal{E} \cap L^1$. Then $\text{codim}_{\mathcal{E}} \mathcal{E}_1 \leq 12$.

Proof of Lemma 2.5. We start from the eigenfunction equation

$$\phi(x) = -\frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{1}{|x - y|} V(y)\phi(y) dy.$$

Recall that $\langle \phi, V \rangle = 0$. Using (2-1), we obtain that

$$\left| \phi(x) - \sum_{k=1}^3 \langle V\phi, y_k \rangle \frac{x_k}{|x|^3} \right| \lesssim \frac{1}{|x|^2} \int_{\mathbb{R}^3} \frac{|y|^2 |V(y)| |\phi(y)| dy}{|x - y|}.$$

Since $\phi \in \langle x \rangle^{-2}L^\infty$ and $V \in L^1 \cap L^{\frac{3}{2},1}$, the right-hand side is in $|x|^{-2}(L^{3,\infty} \cap L^\infty)$.

Using (2-2), we obtain instead that

$$\left| \phi(x) - \sum_{k=1}^3 \langle \phi V, y_k \rangle \frac{x_k}{|x|^3} - \sum_{k,\ell=1}^3 \langle \phi V, y_k y_\ell \rangle \left(\frac{\delta_{k\ell}}{2|x|^3} - \frac{3x_k x_\ell}{2|x|^5} \right) \right| \lesssim \frac{1}{|x|^3} \int_{\mathbb{R}^3} \frac{|y|^3 |V(y)| |\phi(y)| dy}{|x-y|}.$$

Since $\phi \in \langle x \rangle^{-2} L^\infty$ and $V \in \langle x \rangle^{-1} L^1 \cap L^{\frac{3}{2},1}$, the right-hand side is in $|x|^{-3} (L^{3,\infty} \cap L^\infty)$.

These estimates matter only in the region $\{x : |x| \geq 1\}$, since near zero, $\phi \in L^\infty \subset L^1(\{|x| \leq 1\})$. As $|x|^{-3} L^{3,\infty} \subset L^1(\{|x| \geq 1\})$ and

$$\frac{x_k}{|x|^3}, \frac{\delta_{k\ell}}{2|x|^3} - \frac{3x_k x_\ell}{2|x|^5} \notin L^1$$

are linearly independent, it follows that $\phi \in \mathcal{E}$ is in L^1 if and only if all the coefficients $\langle V\phi, y_k \rangle$ and $\langle V\phi, y_k y_\ell \rangle$ are zero.

Then \mathcal{E}_1 is the kernel of a rank-12 map $\phi \mapsto (\langle \phi V, y_k \rangle, \langle \phi V, y_k y_\ell \rangle)$ from \mathcal{E} to \mathbb{C}^{12} , so $\text{codim}_{\mathcal{E}} \mathcal{E}_1 \leq 12$. \square

2C. Wiener spaces.

Definition. For a Banach lattice X , let the space \mathcal{V}_X consist of kernels $T(x, y, \sigma)$ such that, for each pair (x, y) , we have that $T(x, y, \sigma)$ is a finite measure in σ on \mathbb{R} and

$$M(T)(x, y) := \int_{\mathbb{R}} d|T(x, y, \sigma)|$$

is an X -bounded operator.

\mathcal{V}_X is an algebra under

$$(T_1 * T_2)(x, z, \sigma) := \int T_1(x, y, \rho) T_2(y, z, \sigma - \rho) dy ds.$$

Elements of \mathcal{V}_X have Fourier transforms

$$\widehat{T}(x, y, \lambda) := \int_{\mathbb{R}} e^{-i\sigma\lambda} dT(x, y, \sigma),$$

which are uniformly X -bounded operators, $\widehat{T}(\lambda) \in L^\infty_\lambda \mathcal{B}(X)$, and, for every $\lambda \in \mathbb{R}$, we have $\widehat{T}_1(\lambda) \widehat{T}_2(\lambda) = (T_1 * T_2)^\wedge(\lambda)$.

The space \mathcal{V}_X contains elements of the form $\delta_0(\sigma)T(x, y)$, whose Fourier transform is constantly the operator $T(x, y) \in \mathcal{B}(X)$. In particular, rank-one operators $\delta_0(\sigma)\phi(x) \otimes \psi(y)$ are in \mathcal{V}_X when $\psi \in X^*$ and $\phi \in X$. More generally, $f(\sigma)T(x, y) \in \mathcal{V}_X$ if $f \in L^1$ and $T \in \mathcal{B}(X)$.

Moreover, for two Banach lattices X and Y of functions on \mathbb{R}^3 , we also define the space $\mathcal{V}_{X,Y}$ of kernels $T(x, y, \sigma)$ such that $M(T)(x, y)$ is a bounded operator from X to Y . The category of such operators forms an abroid, in the sense that

$$\|T_1 * T_2\|_{\mathcal{V}_{X,Z}} \leq \|T_1\|_{\mathcal{V}_{Y,Z}} \|T_2\|_{\mathcal{V}_{X,Y}}.$$

For example, note that

$$(R_0((\lambda + i0)^2))^\wedge \in \mathcal{V}_{L^{3/2,1}, L^\infty} \cap \mathcal{V}_{L^1, L^{3,\infty}} \quad \text{and} \quad (\partial_\lambda R_0((\lambda + i0)^2))^\wedge \in \mathcal{V}_{L^1, L^\infty}.$$

Indeed, the Fourier transform in λ is

$$(R_0((\lambda + i0)^2))^{\wedge}(\sigma)(x, y) = (4\pi\sigma)^{-1}\delta_{|x-y|}(\sigma),$$

so we have

$$M((R_0((\lambda + i0)^2))^{\wedge}) = \frac{1}{4\pi|x-y|}.$$

Clearly $1/(4\pi|x-y|)$ is in $\mathcal{B}(L^{\frac{3}{2},1}, L^{\infty}) \cap \mathcal{B}(L^1, L^{3,\infty})$.

Likewise,

$$(\partial_{\lambda} R_0((\lambda + i0)^2))^{\wedge}(\sigma)(x, y) = (4\pi)^{-1}i\delta_{|x-y|}(\sigma),$$

so we have

$$M((\partial_{\lambda} R_0((\lambda + i0)^2))^{\wedge}) = (4\pi)^{-1}1 \otimes 1,$$

which is in $\mathcal{B}(L^1, L^{\infty})$.

A space that will repeatedly intervene in computations is

$$\mathcal{W} = \{L \mid L^{\vee} \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}, (\partial_{\lambda} L)^{\vee} \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}\}.$$

This space has the algebra property that $L_1, L_2 \in \mathcal{W} \implies L_1(\lambda)L_2(\lambda) \in \mathcal{W}$.

The following technical lemma will be useful:

Lemma 2.6 (Fourier transforms).

$$\begin{aligned} M\left(\left(\frac{e^{is|x-y|}}{4\pi|x-y|}\right)^{\wedge}\right) &= \frac{1}{4\pi|x-y|}, \\ M\left(\left(\partial_s \frac{e^{is|x-y|}}{4\pi|x-y|}\right)^{\wedge}\right) &= \frac{1 \otimes 1}{4\pi}, \\ M\left(\left(\frac{R_0((s+i0)^2) - R_0(0)}{s}\right)^{\wedge}\right) &= \frac{1 \otimes 1}{4\pi}, \\ M\left(\left(\partial_s \frac{R_0((s+i0)^2) - R_0(0)}{s}\right)^{\wedge}\right) &= \frac{|x-y|}{8\pi}, \\ M\left(\left(\frac{R_0((s+i0)^2) - R_0(0) - is\frac{1 \otimes 1}{4\pi}}{s^2}\right)^{\wedge}\right) &= \frac{|x-y|}{8\pi}, \\ M\left(\left(\partial_s \frac{R_0((s+i0)^2) - R_0(0) - is\frac{1 \otimes 1}{4\pi}}{s^2}\right)^{\wedge}\right) &= \frac{|x-y|^2}{24\pi}. \end{aligned}$$

Proof. Let $a > 0$. Observe that the Fourier transform of $e^{i\lambda a}$ in λ is $\delta_a(t)$. Then

$$\frac{e^{i\lambda a} - 1}{i\lambda} = \int_0^a e^{i\lambda b} db,$$

so $((e^{i\lambda a} - 1)/(i\lambda))^{\wedge} = \chi_{[0,a]}(\lambda)$. Also

$$\frac{e^{i\lambda a} - 1 - i\lambda a}{i\lambda^2} = \int_0^a \frac{e^{i\lambda b} - 1}{\lambda} db,$$

so $((e^{i\lambda a} - 1 - i\lambda a)/(i\lambda^2))^\wedge = (a - t)\chi_{[0,a]}(t)$.

Note that

$$R_0((s + i0)^2) = \frac{e^{is|x-y|}}{4\pi|x-y|}$$

has the Fourier transform $\delta_{|x-y|}(\sigma)/(4\pi|x-y|)$. Thus

$$R_0((s + i0)^2)^\wedge = \left(\frac{e^{is|x-y|}}{4\pi|x-y|}\right)^\wedge = \frac{\delta_{|x-y|}(\sigma)}{4\pi|x-y|}.$$

Integrating the absolute value in σ , we obtain $1/(4\pi|x-y|)$.

Likewise,

$$\left(\frac{R_0((s + i0)^2) - R_0(0)}{s}\right)^\wedge = \frac{i\chi_{[0,|x-y|]}(\sigma)}{4\pi|x-y|}.$$

Integrating the absolute value in σ , we get $1/(4\pi) = (1 \otimes 1)/(4\pi)$.

The Fourier transform of the derivative is

$$\left(\partial_s \frac{R_0((s + i0)^2) - R_0(0)}{s}\right)^\wedge = \frac{i\sigma\chi_{[0,|x-y|]}(\sigma)}{4\pi|x-y|}.$$

Integrating in σ , we obtain $|x-y|/(8\pi)$.

Next,

$$\begin{aligned} \left(\frac{R_0((s + i0)^2) - R_0(0) - is1 \otimes 1}{s^2}\right)^\wedge &= \left(\frac{e^{is|x-y|} - 1 - is|x-y|}{4\pi s^2|x-y|}\right)^\wedge \\ &= \frac{(|x-y| - \sigma)\chi_{[0,|x-y|]}(\sigma)}{4\pi|x-y|}. \end{aligned} \tag{2-6}$$

Integrating in σ , we obtain $|x-y|/(8\pi)$.

The Fourier transform of the derivative is

$$\left(\partial_s \frac{R_0((s + i0)^2) - R_0(0) - is1 \otimes 1}{s^2}\right)^\wedge = \frac{\sigma(|x-y| - \sigma)\chi_{[0,|x-y|]}(\sigma)}{4\pi|x-y|}.$$

Integrating in σ , we obtain $|x-y|^2/(24\pi)$. □

2D. Regular points and regular Hamiltonians. Before examining the possible singularity at zero, we study what happens at regular points in the spectrum.

Recall the notation $V_1 = |V|^{1/2}$ and $V_2 = |V|^{1/2} \operatorname{sgn} V$. The following two properties play an important part in the study:

Lemma 2.7. *Let*

$$T(x, y, \rho) := \frac{V_2(x)V_1(y)}{4\pi|x-y|} \delta_{-|x-y|}(\rho),$$

so $\widehat{T}(\lambda) = V_2 R_0((\lambda + i0)^2) V_1$. Then:

(C1) $\lim_{R \rightarrow \infty} \|\chi_{\rho \geq R}(\rho) T(\rho)\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} = 0$.

(C2) For some $N \geq 1$, we have $\lim_{\epsilon \rightarrow 0} \|T^N(\rho + \epsilon) - T^N(\rho)\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} = 0$.

Here the powers of T mean repeated convolution. We refer the reader to similar properties that appear in the proof of [Beceanu and Goldberg 2012, Theorem 5].

Proof of Lemma 2.7. Suppose V_1 and V_2 are bounded functions with compact support in $B(0, D)$. It follows that for $R > 2D$, we have $\chi(\frac{t}{R})T(t) = 0$, so in particular

$$\|\chi_{t \geq R} T\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} \rightarrow 0$$

as $R \rightarrow \infty$, and property (C1) is preserved by taking the limits of V_1 and V_2 in $L^{3,2}$.

Next, fix $p \in (1, \frac{4}{3}]$ and assume that V_1 and V_2 are bounded and of compact support.

Since V_1 and V_2 are bounded and of compact support, T also has the local and distal properties

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \left\| \chi_{<\epsilon}(|x-y|) \frac{V_2(x)V_1(y)}{|x-y|} \right\|_{\mathcal{B}(L^{3/2,2}) \cap \mathcal{B}(L^{3,2})} &= 0, \\ \lim_{R \rightarrow \infty} \left\| \chi_{>R}(|x-y|) \frac{V_2(x)V_1(y)}{|x-y|} \right\|_{\mathcal{B}(L^{3/2,2}) \cap \mathcal{B}(L^{3,2})} &= 0. \end{aligned}$$

Combined with condition (C1), this implies that for any $\epsilon > 0$ there exists a cutoff function χ compactly supported in $(0, \infty)$ such that

$$\|\chi(\rho)T(\rho) - T(\rho)\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} < \epsilon.$$

Thus, it suffices to show that condition (C2) holds for $\chi(\rho)T(\rho)$.

The Fourier transform of $\chi(\rho)T(\rho)$ has the form

$$(\chi(\rho)T(\rho))^\wedge(\lambda) = V_2(x) \frac{e^{i\lambda|x-y|}}{4\pi|x-y|} \chi(|x-y|) V_1(y). \quad (2-7)$$

Such oscillating kernels have decay in the L^p operator norm for $p > 1$. By [Stein 1993, Lemma on p. 392], with p' being the dual exponent, that is, we have $\frac{1}{p} + \frac{1}{p'} = 1$,

$$\|(\chi(\rho)T(\rho))^\wedge(\lambda) f\|_{L^p} \lesssim \lambda^{-\frac{3}{p'}} \|f\|_{L^{p'}}. \quad (2-8)$$

Taking into account the fact that $(\chi(\rho)T(\rho))^\wedge(\lambda)$ has a kernel bounded in absolute value by

$$\frac{|V(x)|^{\frac{1}{2}} |V(y)|^{\frac{1}{2}}}{4\pi|x-y|}$$

(where $|V|^{\frac{1}{2}} = V_1$ is bounded and has compact support by assumption), it follows that $(\chi(\rho)T(\rho))^\wedge(\lambda)$ is uniformly bounded in $\mathcal{B}(X, L^p)$, $\mathcal{B}(L^p, X)$, and $\mathcal{B}(L^p)$ for all λ , where X is $L^{\frac{3}{2},2}$ or $L^{3,2}$. Therefore, by also using (2-8) for the middle factors,

$$\|((\chi(\rho)T(\rho))^\wedge(\lambda))^N f\|_X \lesssim \langle \lambda \rangle^{-\frac{3(N-2)}{p'}} \|f\|_X.$$

For $N > 2 + \frac{2p'}{3}$, this shows that $\partial_\rho(\chi(\rho)T(\rho))^N$ are uniformly bounded operators in $\mathcal{B}(X)$, where X is either $L^{\frac{3}{2},2}$ or $L^{3,2}$. Since $(\chi(\rho)T(\rho))^N$ has compact support in ρ , this in turn implies (C2).

For general $V \in L^{\frac{3}{2},1}$, choose a sequence of bounded compactly supported approximations for which (C2) holds, as shown above. By a limiting process, we obtain that (C2) also holds for V . \square

Lemma 2.8. *Let $\widehat{T}(\lambda) = V_2 R_0((\lambda + i0)^2) V_1$. Assume that $V \in L^{\frac{3}{2},1}$ and let $\lambda_0 \neq 0$. Consider a cutoff function χ . Then, for $\epsilon \ll 1$, we have*

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}.$$

The same holds for $\lambda_0 = 0$ if V is a generic potential.

Infinity has the same property: for $R \gg 1$, we have

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}.$$

Proof of Lemma 2.8. Note that $I + \widehat{T}(\lambda_0)$ is invertible in $\mathcal{B}(L^{\frac{3}{2},2})$ and in $\mathcal{B}(L^{3,2})$ for all $\lambda_0 \neq 0$, the only issue being at zero.

Indeed, assume that $I + \widehat{T}(\lambda_0)$ is not invertible in $\mathcal{B}(L^{\frac{3}{2},2})$; then, by Fredholm’s alternative, there exists a nonzero $f \in L^{\frac{3}{2},2}$ such that

$$f = -V_2 R_0((\lambda_0 + i0)^2) V_1 f.$$

Let $V_1 = V_1^1 + V_1^2$ and $V_2 = V_2^1 + V_2^2$, where V_1^1 and V_2^1 have compact support and are bounded with $\|V_1^1\|_{L^{3,2}}, \|V_2^1\|_{L^{3,2}} \ll 1$. Then

$$f = -(I + V_2 R_0((\lambda_0 + i0)^2) V_1^2 + V_2^2 R_0((\lambda_0 + i0)^2) V_1^1)^{-1} V_2^1 R_0((\lambda_0 + i0)^2) V_1^1 f,$$

which implies that $f \in L^2$. Letting $g = R_0((\lambda_0 + i0)^2) V_1 f$, we obtain a nonzero $L^{6,\infty}$ solution g of the equation

$$g = -R_0((\lambda_0 + i0)^2) V g.$$

However, this is impossible for $\lambda_0 \neq 0$ due to the results of Ionescu and Jerison [2003] and Goldberg and Schlag [2004b].

When $\lambda_0 = 0$, we have that g is a zero-energy eigenfunction or resonance for $H = -\Delta + V$, which cannot happen if V is a generic potential.

Let

$$S_\epsilon(\lambda) = \chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right)(\widehat{T}(\lambda) - \widehat{T}(\lambda_0)).$$

A simple argument based on condition (C1) shows that $\lim_{\epsilon \rightarrow 0} \|S_\epsilon^\vee\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} = 0$. Then

$$\begin{aligned} \chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right)(I + \widehat{T}(\lambda))^{-1} &= \chi\left(\frac{\lambda}{\epsilon}\right)\left(I + \widehat{T}(\lambda_0) + \chi\left(\frac{\lambda - \lambda_0}{2\epsilon}\right)(\widehat{T}(\lambda) - \widehat{T}(\lambda_0))\right)^{-1} \\ &= \chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right)(I + \widehat{T}(\lambda_0))^{-1} \sum_{k=0}^\infty (-1)^k (S_{2\epsilon}(\lambda)(I + \widehat{T}(\lambda_0))^{-1})^k. \end{aligned}$$

The Fourier transform of the series above converges for sufficiently small ϵ , showing that

$$\left(\chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right)(I + \widehat{T}(\lambda))^{-1}\right)^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}.$$

Concerning the derivative,

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right) \partial_\lambda (I + \widehat{T}(\lambda))^{-1} = -\chi\left(\frac{\lambda - \lambda_0}{\epsilon}\right) (I + \widehat{T}(\lambda))^{-1} \partial_\lambda \widehat{T}(\lambda) \chi\left(\frac{\lambda - \lambda_0}{2\epsilon}\right) (I + \widehat{T}(\lambda))^{-1}.$$

Here

$$\left(\chi\left(\frac{\lambda-\lambda_0}{2\epsilon}\right)(I+\widehat{T}(\lambda))^{-1}\right)^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}} \quad \text{and} \quad (\partial_\lambda \widehat{T}(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2},L^{3,2}}$$

since

$$M((\partial_\lambda T(\lambda))^\vee) = \frac{|V_2(x)| \otimes |V_1(y)|}{4\pi}.$$

Then

$$\left(\chi\left(\frac{\lambda-\lambda_0}{\epsilon}\right)\partial_\lambda(I+\widehat{T}(\lambda))^{-1}\right)^\vee \in \mathcal{V}_{L^{3/2,2},L^{3,2}}.$$

At infinity, for any real number L , one can express $(1-\chi(\frac{\lambda}{R}))\widehat{T}(\lambda)$ as the Fourier transform of

$$S_R(\rho) = (T - R\check{\chi}(R\cdot) * T)(\rho) = \int_{\mathbb{R}} R\check{\chi}(R\sigma)[T(\rho) - T(\rho - \sigma)] d\sigma.$$

Thanks to condition (C2), the norm of the right-hand side integral vanishes as $L \rightarrow \infty$. This makes it possible to construct an inverse Fourier transform for

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1} = \left(1 - \chi\left(\frac{\lambda}{R}\right)\right) \sum_{k=0}^{\infty} (-1)^k \left(\left(1 - \chi\left(\frac{2\lambda}{R}\right)\right)\widehat{T}(\lambda)\right)^k$$

via this power series expansion, which converges for sufficiently large R .

If only T^N satisfies (C2) then one constructs an inverse Fourier transform for

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I - (-\widehat{T})^N(\lambda))^{-1}$$

in this manner and observes that

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1} = \left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I - (-\widehat{T}(\lambda))^N)^{-1} \sum_{k=0}^{N-1} (-1)^k \widehat{T}^k(\lambda).$$

Finally, concerning the derivative in a neighborhood of infinity, we note that

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)\partial_\lambda(I + \widehat{T}(\lambda))^{-1} = -\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1}\partial_\lambda \widehat{T}(\lambda)\left(1 - \chi\left(\frac{2\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1}.$$

Here

$$\left(\left(1 - \chi\left(\frac{2\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1}\right)^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}} \quad \text{and} \quad (\partial_\lambda \widehat{T}(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2},L^{3,2}}.$$

Therefore,

$$\left(\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)\partial_\lambda(I + \widehat{T}(\lambda))^{-1}\right)^\vee \in \mathcal{V}_{L^{3/2,2},L^{3,2}}. \quad \square$$

In the case when H is generic, we can cover the whole spectrum $[0, \infty)$ by open neighborhoods of regular points, plus an open neighborhood of infinity, and choose a subordinate partition of unity. We retrieve a form of [Beceanu and Goldberg 2012, Theorem 2]:

Theorem 2.9. *Let $V \in L^{\frac{3}{2},1}$ be a real-valued potential for which the Schrödinger operator $H = -\Delta + V$ has no resonances or eigenvalues at zero energy. Then*

$$\|e^{-itH} P_c f\|_\infty \lesssim |t|^{-\frac{3}{2}} \|f\|_1. \quad (2-9)$$

In the context of the wave equation, again if the Hamiltonian H is generic, we retrieve the results of [Beceanu and Goldberg 2014].

Proof of Theorem 2.9. Consider a sufficiently large R such that

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$$

by Lemma 2.8. Also by Lemma 2.8, for every $\lambda_0 \in [-4R, 4R]$ (including zero, since V is a generic potential), there exists $\epsilon(\lambda_0) > 0$ such that

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon(\lambda_0)}\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}.$$

Since $[-4R, 4R]$ is a compact set, there exists a finite covering

$$[-4R, 4R] \subset \bigcup_{k=1}^N (\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)).$$

Then we construct a finite partition of unity on \mathbb{R} by smooth functions $1 = \sum_{k=1}^N \chi_k(\lambda) + \chi_\infty(\lambda)$, where $\text{supp } \chi_\infty \subset \mathbb{R} \setminus (-2R, 2R)$ and $\text{supp } \chi_k \subset [\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)]$. By our construction, for each $1 \leq k \leq N$ and for $k = \infty$, we have $\chi_k(\lambda)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$, so summing up we obtain that $(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$.

By spectral calculus, we express the perturbed evolution as

$$\begin{aligned} e^{itH} P_c f &= \frac{1}{2\pi i} \int_0^\infty e^{it\lambda} (R_V(\lambda+i0) - R_V(\lambda-i0)) f \, d\lambda \\ &= \frac{1}{\pi i} \int_{-\infty}^\infty e^{it\lambda^2} R_V((\lambda+i0)^2) f \lambda \, d\lambda \\ &= \frac{1}{\pi i} \int_{-\infty}^\infty e^{it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 (I + \widehat{T}(\lambda))^{-1} V_2 R_0((\lambda+i0)^2)) f \lambda \, d\lambda \\ &= \frac{1}{2\pi t} \int_{-\infty}^\infty e^{it\lambda^2} \partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 (I + \widehat{T}(\lambda))^{-1} V_2 R_0((\lambda+i0)^2)) f \, d\lambda \\ &= \frac{C}{t^{\frac{3}{2}}} \int_{-\infty}^\infty e^{i\frac{\rho^2}{4t}} (\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 (I + \widehat{T}(\lambda))^{-1} V_2 R_0((\lambda+i0)^2)))^\vee(\rho) f \, d\rho. \end{aligned} \tag{2-10}$$

Since $(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$, it follows that $(\partial_\lambda (I + \widehat{T}(\lambda))^{-1})^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$. Taking into account that $R_0((\lambda + i0)^2) V_1 \in \mathcal{V}_{L^1, L^{3/2,2}}$ and $V_2 R_0((\lambda + i0)^2) \in \mathcal{V}_{L^{3,2}, L^\infty}$, we obtain that

$$R_0((\lambda + i0)^2) V_1 (I + \widehat{T}(\lambda))^{-1} V_2 R_0((\lambda + i0)^2) \in \mathcal{V}_{L^1, L^\infty}.$$

By definition, this ensures a bound of $|t|^{-\frac{3}{2}}$ for this expression's contribution to (2-10). The other terms are handled similarly. □

We next consider the effect of singularities at zero.

2E. Exceptional Hamiltonians of the first kind. Let

$$Q = -\frac{1}{2\pi i} \int_{|z+1|=\delta} (V_2 R_0(0) V_1 - zI)^{-1} dz$$

and $\bar{Q} = 1 - Q$. Assuming that $H = -\Delta + V$ has only a resonance ϕ at zero, then (recalling that $-\langle \phi, V\phi \rangle = 1$), by the analytic Fredholm theorem,

$$Q = -V_2\phi \otimes V_1\phi.$$

The resonance $\phi \in \mathcal{M}$ satisfies the equation $\phi = -R_0(0)V\phi$. Since $\phi \in L^{3,\infty} \cap L^\infty$, we have that Q is bounded on $L^{\frac{3}{2},2}$ and on $L^{3,2}$, so the constant family of operators Q is in \mathcal{W} . Moreover, Q is in $\mathcal{B}(L^{\frac{3}{2},2}, L^{3,2})$ and in $\mathcal{B}(L^{3,2}, L^{\frac{3}{2},2})$.

Note that, since

$$e^{i\lambda|x-y|} - 1 \lesssim \min(1, \lambda|x-y|) \implies e^{i\lambda|x-y|} - 1 \lesssim \lambda^\delta |x-y|^\delta,$$

one has

$$V_2(x) \left(\frac{e^{i\lambda|x-y|}}{|x-y|} - \frac{1}{|x-y|} \right) V_1(y) \lesssim |V_2(x)| \lambda |V_1(y)|. \quad (2-11)$$

Thus, when $V \in \langle x \rangle^{-1} L^{\frac{3}{2},1}$,

$$I + \hat{T}(\lambda) = I + V_2 R_0((\lambda + i0)^2) V_1$$

is Lipschitz continuous in $\mathcal{B}(L^2)$. This implies that, more generally, when $V \in L^{\frac{3}{2},1}$, we have that $\hat{T}(\lambda)$ is continuous in $\mathcal{B}(L^2)$ (the proof is by approximation).

In a similar manner, by approximating $V \in L^{\frac{3}{2},1}$ with $\langle x \rangle^{-2} L^{\frac{3}{2},1}$ potentials, we obtain that $\hat{T}(\lambda)$ is continuous in $\mathcal{B}(L^{\frac{3}{2},2}) \cap \mathcal{B}(L^{3,2})$.

Let

$$K = (I + V_2 R_0(0) V_1 + Q)^{-1} \bar{Q}.$$

Then K is the inverse of $\bar{Q}(I + \hat{T}(0))\bar{Q} = \bar{Q}(I + V_2 R_0(0) V_1)\bar{Q}$ in $\mathcal{B}(\bar{Q}L^{\frac{3}{2},2} \cap \bar{Q}L^{3,2})$, in the sense that

$$K\bar{Q}(I + V_2 R_0(0) V_1)\bar{Q} = \bar{Q}(I + V_2 R_0(0) V_1)\bar{Q}K = \bar{Q}. \quad (2-12)$$

By continuity, $\bar{Q}(I + V_2 R_0((\lambda + i0)^2) V_1)\bar{Q}$ is also invertible for $|\lambda| \ll 1$.

The following lemma, also known as the Feshbach lemma, is extremely useful in studying the singularity at zero.

Lemma 2.10 (see [Yajima 2005, Lemma 4.7]). *Let $X = X_0 + X_1$ be a direct sum decomposition of a vector space X . Suppose that a linear operator $L \in \mathcal{B}(X)$ is written in the form*

$$L = \begin{pmatrix} L_{00} & L_{01} \\ L_{10} & L_{11} \end{pmatrix}$$

with respect to this decomposition and that L_{00}^{-1} exists. Set $C = L_{11} - L_{10}L_{00}^{-1}L_{01}$. Then, L^{-1} exists if and only if C^{-1} exists. In this case,

$$L^{-1} = \begin{pmatrix} L_{00}^{-1} + L_{00}^{-1}L_{01}C^{-1}L_{10}L_{00}^{-1} & -L_{00}^{-1}L_{01}C^{-1} \\ -C^{-1}L_{10}L_{00}^{-1} & C^{-1} \end{pmatrix}. \quad (2-13)$$

By definition, an exceptional point $\lambda \in \mathbb{C}$ is one where $I + V_2 R_0(\lambda) V_1$ is not L^2 -invertible.

Lemma 2.11. *Assume that $V \in \langle x \rangle^{-2} L^{\frac{3}{2},1} \subset \langle x \rangle^{-1} L^1 \cap L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is exceptional of the first type, with a resonance ϕ at zero. Let χ be a fixed cutoff function. Then, for some $\epsilon > 0$,*

$$\chi\left(\frac{\lambda}{\epsilon}\right)(I + \hat{T}(\lambda))^{-1} = L(\lambda) - \lambda^{-1} \chi\left(\frac{\lambda}{\epsilon}\right) \frac{4\pi i}{|\langle V, \phi \rangle|^2} V_2 \phi \otimes V_1 \phi,$$

where $L \in \mathcal{W}$.

Moreover, zero is an isolated exceptional point, so $H = -\Delta + V$ has finitely many negative eigenvalues.

The computations in the proof of this lemma parallel those in [Yajima 2005, Section 4.3]. The main difference is using \hat{L}^1 -related spaces instead of Hölder spaces.

Proof of Lemma 2.11. We apply Lemma 2.10 to

$$I + \hat{T}(\lambda) := \begin{pmatrix} \bar{Q}(I + \hat{T}(\lambda))\bar{Q} & \bar{Q}\hat{T}(\lambda)\mathcal{Q} \\ \mathcal{Q}\hat{T}(\lambda)\bar{Q} & \mathcal{Q}(I + \hat{T}(\lambda))\mathcal{Q} \end{pmatrix} = \begin{pmatrix} T_{00}(\lambda) & T_{01}(\lambda) \\ T_{10}(\lambda) & T_{11}(\lambda) \end{pmatrix}.$$

Note that $T_{00}(\lambda) := \bar{Q}(I + V_2 R_0((\lambda + i0)^2) V_1) \bar{Q}$ is invertible in $\mathcal{B}(\bar{Q}L^{\frac{3}{2},2}) \cap \mathcal{B}(\bar{Q}L^{3,2})$ for $|\lambda| \ll 1$ because

$$T_{00}(0) = \bar{Q}(I + \hat{T}(0))\bar{Q} = \bar{Q}(I + V_2 R_0(0) V_1) \bar{Q}$$

is invertible on $\bar{Q}L^{\frac{3}{2},2}$ and on $\bar{Q}L^{3,2}$ with inverse K (see (2-12)), and $T_{00}(\lambda)$ is continuous in the norm of $\mathcal{B}(\bar{Q}L^{\frac{3}{2},2}) \cap \mathcal{B}(\bar{Q}L^{3,2})$ (see (2-11) above).

Furthermore, start from

$$(R_0((\lambda + i0)^2))^{\wedge} \in \mathcal{V}_{L^{3/2,1}, L^\infty} \cap \mathcal{V}_{L^1, L^{3,\infty}} \quad \text{and} \quad (\partial_\lambda R_0((\lambda + i0)^2))^{\wedge} \in \mathcal{V}_{L^1, L^\infty}.$$

We know that

$$|V|^{\frac{1}{2}} \in \mathcal{B}(L^{\frac{3}{2},2}, L^1) \cap \mathcal{B}(L^\infty, L^{3,2}) \cap \mathcal{B}(L^{3,\infty}, L^{\frac{3}{2},2}) \cap \mathcal{B}(L^{3,2}, L^{\frac{3}{2},1}).$$

Thus $V_2 R_0((\lambda + i0)^2) V_1 \in \mathcal{W}$ and \bar{Q} preserves that. Then $T_{00}(\lambda) \in \mathcal{W}$ as well.

Next, since $T_{00}(0)$ is invertible, for small ϵ we have $\chi(\frac{\lambda}{\epsilon}) T_{00}^{-1}(\lambda) \in \mathcal{W}$. The proof is as follows: Let

$$S_\epsilon(\lambda) := \chi\left(\frac{\lambda}{\epsilon}\right) \bar{Q}(\hat{T}(\lambda) - \hat{T}(0))\bar{Q}.$$

A simple argument based on condition (C1) shows that $\lim_{\epsilon \rightarrow 0} \|S_\epsilon^\vee\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} = 0$. Then

$$\begin{aligned} \chi\left(\frac{\lambda}{\epsilon}\right) T_{00}^{-1}(\lambda) &= \chi\left(\frac{\lambda}{\epsilon}\right) \left(T_{00}(0) + \chi\left(\frac{\lambda}{2\epsilon}\right) \bar{Q}(\hat{T}(\lambda) - \hat{T}(0))\bar{Q} \right)^{-1} \\ &= \chi\left(\frac{\lambda}{\epsilon}\right) T_{00}^{-1}(0) \sum_{k=0}^{\infty} (-1)^k (S_{2\epsilon}(\lambda) T_{00}^{-1}(0))^k. \end{aligned}$$

The series above converges for sufficiently small ϵ , showing that $\chi(\frac{\lambda}{\epsilon}) T_{00}^{-1}(\lambda) \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}$.

Concerning the derivative,

$$\chi\left(\frac{\lambda}{\epsilon}\right) \partial_\lambda T_{00}^{-1}(\lambda) = -\chi\left(\frac{\lambda}{\epsilon}\right) T_{00}^{-1}(\lambda) \partial_\lambda T_{00}(\lambda) \chi\left(\frac{\lambda}{2\epsilon}\right) T_{00}^{-1}(\lambda).$$

In this expression, $(\chi(\frac{\lambda}{2\varepsilon})T_{00}^{-1}(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}$ and $(\partial_\lambda T_{00}(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$. Thus

$$\left(\chi\left(\frac{\lambda}{\varepsilon}\right)\partial_\lambda T_{00}^{-1}(\lambda)\right)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}.$$

This computation shows that $\chi(\frac{\lambda}{\varepsilon})T_{00}^{-1}(\lambda) \in \mathcal{W}$.

Let

$$\begin{aligned} J(\lambda) &:= \frac{\widehat{T}(\lambda) - (V_2 R_0(0)V_1 + i\lambda(4\pi)^{-1}V_2 \otimes V_1)}{\lambda^2} \\ &= \frac{V_2 R_0((\lambda + i0)^2)V_1 - V_2 R_0(0)V_1 - i\lambda(4\pi)^{-1}V_2 \otimes V_1}{\lambda^2}. \end{aligned}$$

Then (recall that $Q = -V_2\phi \otimes V_1\phi$),

$$\begin{aligned} T_{11}(\lambda) &= Q(I + \widehat{T}(\lambda))Q = Q(I + V_2 R_0((\lambda + i0)^2)V_1)Q \\ &= Q(V_2 R_0((\lambda + i0)^2)V_1 - V_2 R_0(0)V_1)Q \\ &= V_2\phi \otimes V\phi(R_0((\lambda + i0)^2) - R_0(0))V\phi \otimes V_1\phi \\ &= \left(\lambda \frac{|\langle V, \phi \rangle|^2}{4i\pi} - \lambda^2 \langle V_1\phi, J(\lambda)V_2\phi \rangle\right)Q \\ &= (\lambda a^{-1} - \lambda^2 \langle V_1\phi, J(\lambda)V_2\phi \rangle)Q \\ &=: \lambda c_0(\lambda)Q. \end{aligned} \tag{2-14}$$

Note that $c_0(0) = a^{-1} \neq 0$. Recall that $a = 4i\pi/|\langle V, \phi \rangle|^2$.

By the third line of (2-14), $c_0(\lambda) \in \widehat{L}^1$ if

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x)\phi(x)V(y)\phi(y) \left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} dx dy < \infty.$$

For every x and y , by Lemma 2.6,

$$\left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} = \left\| \frac{\chi_{[0,|x-y|]}(t)}{|x-y|} \right\|_{L_t^1} = 1,$$

so it is enough to assume that $V\phi \in L^1$, i.e., that $V \in L^{\frac{3}{2},1}$, to prove that $c_0(\lambda) \in \widehat{L}^1$.

In order for $\partial_\lambda c_0(\lambda)$ to be in \widehat{L}^1 , it suffices that

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x)\phi(x)V(y)\phi(y) \left\| \partial_\lambda \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} dx dy < \infty.$$

For every x and y , by Lemma 2.6,

$$\left\| \partial_\lambda \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} = \left\| \frac{t\chi_{[0,|x-y|]}(t)}{|x-y|} \right\|_{L_t^1} = \frac{|x-y|}{2},$$

so $\partial_\lambda c_0(\lambda) \in \widehat{L}^1$ when $V\phi \in \langle x \rangle^{-1}L^1$, i.e., when $V \in L^1$.

Regarding $J(\lambda)$, if $V \in L^1$ then

$$\langle J(\lambda)V_2\phi, V_1\phi \rangle = \left\langle \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda(4\pi)^{-1}1 \otimes 1}{\lambda^2} V\phi, V\phi \right\rangle \in \widehat{L}_\lambda^1. \tag{2-15}$$

Moreover, when $V \in \langle x \rangle^{-1}L^1$, we know $\langle \partial_\lambda J(\lambda)V_2\phi, V_1\phi \rangle \in \widehat{L}_\lambda^1$.

Furthermore, considering the fact that $\phi + R_0(0)V\phi = 0$, let us define

$$\begin{aligned} \lambda\tilde{\psi}(\lambda) &:= (I + \widehat{T}(\lambda))V_2\phi = (V_2R_0((\lambda + i0)^2)V - V_2R_0(0)V)\phi \\ &= \lambda \left(i \frac{V_2 \otimes V_1}{4\pi} + \lambda J(\lambda) \right) V_2\phi \end{aligned}$$

and

$$\begin{aligned} \lambda\tilde{\psi}^*(\lambda) &:= (I + \widehat{T}(\lambda)^*)V_1\phi = (V_1R_0^*((\lambda + i0)^2)V - V_1R_0(0)V)\phi \\ &= \lambda \left(-i \frac{V_1 \otimes V_2}{4\pi} + \lambda J^*(\lambda) \right) V_1\phi. \end{aligned}$$

Note that

$$M(J(\lambda)^\vee) = |V_2(x)| \frac{|x-y|}{8\pi} |V_1(y)|$$

is a bounded operator from $L^{\frac{3}{2},2}$ to $L^{3,2}$, assuming that $V \in \langle x \rangle^{-2}L^{\frac{3}{2},1}$. Thus $J(\lambda)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$ and the same goes for $\lambda\partial_\lambda J(\lambda)$.

Moreover,

$$M((\lambda J(\lambda))^\vee) = \frac{|V_2| \otimes |V_1|}{2\pi}.$$

Thus $(\lambda J(\lambda))^\vee \in \mathcal{V}_{L^2}$ for $V \in L^1$ and $(\lambda J(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}$ when $V \in \langle x \rangle^{-2}L^{\frac{3}{2},1}$. Further note that $(\partial_\lambda(\lambda J(\lambda)))^\vee = (J(\lambda) + \lambda\partial_\lambda J(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$. It follows that $\lambda J(\lambda) \in \mathcal{W}$.

Then (recalling that $Q = -V_2\phi \otimes V_1\phi$),

$$\begin{aligned} T_{01}(\lambda) &:= \overline{Q}\widehat{T}(\lambda)Q = \overline{Q}(I + \widehat{T}(\lambda))Q = (I + \widehat{T}(\lambda))Q - Q(I + \widehat{T}(\lambda))Q \\ &= -\lambda\tilde{\psi}(\lambda) \otimes V_1\phi - \lambda c_0(\lambda)Q \\ &= -\lambda(\tilde{\psi}(\lambda) + c_0(\lambda)V_2\phi) \otimes V_1\phi. \end{aligned}$$

Likewise,

$$T_{10}(\lambda) = -\lambda V_2\phi \otimes (\tilde{\psi}^*(\lambda) + \overline{c_0(\lambda)}V_1\phi).$$

By our above computations, it follows that $T_{01}(\lambda) = \lambda E_1(\lambda)$ and $T_{10}(\lambda) = \lambda E_2(\lambda)$ with $E_1, E_2 \in \mathcal{W}$.

Then $-T_{10}(\lambda)T_{00}^{-1}(\lambda)T_{01}(\lambda) = \lambda^2 c_1(\lambda)Q$, where

$$\begin{aligned} c_1(\lambda) &:= -\langle \tilde{\psi}^*(\lambda) + \overline{c_0(\lambda)}V_1\phi, T_{00}^{-1}(\lambda)(\tilde{\psi}(\lambda) + c_0(\lambda)V_2\phi) \rangle \\ &= -\left\langle \left(-i \frac{V_1 \otimes V_2}{4\pi} + \lambda J^*(\lambda) \right) V_1\phi + \overline{c_0(\lambda)}V_1\phi, T_{00}^{-1}(\lambda) \left(\left(i \frac{V_2 \otimes V_1}{4\pi} + \lambda J(\lambda) \right) V_2\phi + c_0(\lambda)V_2\phi \right) \right\rangle. \end{aligned} \tag{2-16}$$

For example, one of the terms in (2-16) has the form

$$\langle \lambda J^*(\lambda)V_1\phi, T_{00}^{-1}(\lambda)\lambda J(\lambda)V_2\phi \rangle. \tag{2-17}$$

Since $\lambda J(\lambda) \in \mathcal{W}$ and $\chi\left(\frac{\lambda}{\epsilon}\right)T_{00}^{-1}(\lambda) \in \mathcal{W}$ and since $V_1\phi, V_2\phi \in L^{\frac{3}{2},2} \cap L^{3,2}$, it immediately follows that $\chi\left(\frac{\lambda}{\epsilon}\right)(2-17)$ is in \widehat{L}^1 and its derivative is also in \widehat{L}^1 .

We then recognize from formula (2-16) that, for a cutoff function χ ,

$$\chi\left(\frac{\lambda}{\epsilon}\right)c_1(\lambda) \in \widehat{L}^1 \quad \text{and} \quad \chi\left(\frac{\lambda}{\epsilon}\right)\partial_\lambda c_1(\lambda) \in \widehat{L}^1$$

when $V \in \langle x \rangle^{-2}L^{\frac{3}{2},1}$.

Let

$$C(\lambda) := T_{11}(\lambda) - T_{10}(\lambda)T_{00}^{-1}(\lambda)T_{01}(\lambda).$$

Then

$$C(\lambda) = (\lambda a^{-1} - \lambda^2 \langle V_1\phi, J(\lambda)V_2\phi \rangle + \lambda^2 c_1(\lambda))Q =: \lambda a^{-1}Q + \lambda^2 c_2(\lambda)Q.$$

Thus $C(\lambda)/\lambda$ is invertible for $|\lambda| \ll 1$, and when $V \in \langle x \rangle^{-2}L^{\frac{3}{2},1}$ one has that

$$\begin{aligned} C^{-1}(\lambda) &= \frac{1}{\lambda a^{-1} + \lambda^2 c_2(\lambda)}Q \\ &= \left(\frac{1}{\lambda a^{-1}} + \frac{1}{\lambda a^{-1} + \lambda^2 c_2(\lambda)} - \frac{1}{\lambda a^{-1}} \right)Q \\ &= \left(\frac{a}{\lambda} - \frac{c_2(\lambda)}{(a^{-1} + \lambda c_2(\lambda))a^{-1}} \right)Q \\ &=: a\lambda^{-1}Q + E(\lambda). \end{aligned}$$

By our computations, such as (2-15), $\chi\left(\frac{\lambda}{\epsilon}\right)c_2(\lambda) \in \widehat{L}^1$ and $\chi\left(\frac{\lambda}{\epsilon}\right)\partial_\lambda c_2(\lambda) \in \widehat{L}^1$. Therefore for sufficiently small ϵ , as $Q \in \mathcal{B}(L^{\frac{3}{2},2}) \cap \mathcal{B}(L^{3,2}) \cap \mathcal{B}(L^{\frac{3}{2},2}, L^{3,2})$, it follows that $\chi\left(\frac{\lambda}{\epsilon}\right)E(\lambda) \in \mathcal{W}$.

The inverse of $I + \widehat{T}(\lambda)$ is then given for small λ by formula (2-13):

$$(I + \widehat{T})^{-1} = \begin{pmatrix} T_{00}^{-1} + T_{00}^{-1}T_{01}C^{-1}T_{10}T_{00}^{-1} & -T_{00}^{-1}T_{01}C^{-1} \\ -C^{-1}T_{10}T_{00}^{-1} & C^{-1} \end{pmatrix}.$$

Three of the matrix elements belong to \mathcal{W} when localized by $\chi\left(\frac{\lambda}{\epsilon}\right)$. Indeed, recall that $\chi\left(\frac{\lambda}{\epsilon}\right)T_{00}^{-1}(\lambda) \in \mathcal{W}$, $T_{10}(\lambda) = \lambda E_1(\lambda)$ and $T_{01}(\lambda) = \lambda E_2(\lambda)$, while $C^{-1} = \lambda^{-1}E_3(\lambda)$, with $E_1, E_2, \chi\left(\frac{\lambda}{\epsilon}\right)E_3 \in \mathcal{W}$.

The fourth matrix element is C^{-1} in the lower-right corner, which is the sum of the regular term $\chi\left(\frac{\lambda}{\epsilon}\right)E(\lambda) \in \mathcal{W}$ and the singular term

$$a\lambda^{-1}\chi\left(\frac{\lambda}{\epsilon}\right)Q = -a\lambda^{-1}\chi\left(\frac{\lambda}{\epsilon}\right)V_2\phi \otimes V_1\phi.$$

As an aside, note that $\lambda^{-1}(1 - \chi\left(\frac{\lambda}{\epsilon}\right)) \in \widehat{L}^1$ and the same holds for its derivative. Thus we can also write the singular term as $a\lambda^{-1}Q$.

Further note that $(I + \widehat{T})^{-1}$ is well-defined on a whole cut neighborhood of zero by formula (2-13) above. Thus zero is an isolated exceptional point, so there are finitely many negative eigenvalues. \square

The next lemma shows what happens in the case when the potential has the critical rate of decay.

Lemma 2.12. *Assume that $V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is exceptional of the first kind. Let χ be a standard cutoff function. Then*

$$\chi\left(\frac{\lambda}{\epsilon}\right)(I + \widehat{T}(\lambda))^{-1} = L(\lambda) + \lambda^{-1}S(\lambda),$$

with $L(\lambda) \in \mathcal{W}$ and $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$ for sufficiently small $\epsilon > 0$.

Furthermore, 0 is an isolated exceptional point, so H has finitely many negative eigenvalues.

Proof of Lemma 2.12. We again apply Lemma 2.10 to

$$I + \widehat{T}(\lambda) := \begin{pmatrix} \overline{Q}(I + \widehat{T}(\lambda))\overline{Q} & \overline{Q}\widehat{T}(\lambda)Q \\ Q\widehat{T}(\lambda)\overline{Q} & Q(I + \widehat{T}(\lambda))Q \end{pmatrix} \equiv \begin{pmatrix} T_{00}(\lambda) & T_{01}(\lambda) \\ T_{10}(\lambda) & T_{11}(\lambda) \end{pmatrix}.$$

The proof of the fact that $\chi(\frac{\lambda}{\epsilon})T_{00}^{-1}(\lambda) \in \mathcal{W}$ is the same as in Lemma 2.11.

Then note that

$$\begin{aligned} T_{11}(\lambda) &= Q(I + \widehat{T}(\lambda))Q = Q(I + V_2R_0((\lambda + i0)^2)V_1)Q \\ &= Q(V_2R_0((\lambda + i0)^2)V_1 - V_2R_0(0)V_1)Q \\ &= V_2\phi \otimes V\phi(R_0((\lambda + i0)^2) - R_0(0))V\phi \otimes V_1\phi \\ &=: \lambda c_0(\lambda)Q. \end{aligned}$$

Observe that $c_0(0) = a^{-1} \neq 0$. Recall that $a = 4i\pi/|\langle V, \phi \rangle|^2$.

Note that $c_0(\lambda) \in \widehat{L}^1$ if

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x)\phi(x)V(y)\phi(y) \left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} dx dy < \infty.$$

For every x and y , by Lemma 2.6,

$$\left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\widehat{L}_\lambda^1} = \left\| \frac{\chi_{[0,|x-y|]}(t)}{|x-y|} \right\|_{L_t^1} = 1,$$

so it is enough to assume that $V\phi \in L^1$, i.e., that $V \in L^{\frac{3}{2},1}$, to prove that $c_0(\lambda) \in \widehat{L}^1$.

Furthermore, recalling that $Q = -V_2\phi \otimes V_1\phi$,

$$\begin{aligned} T_{01}(\lambda) &:= \overline{Q}\widehat{T}(\lambda)Q = \overline{Q}(I + \widehat{T}(\lambda))Q = (I + \widehat{T}(\lambda))Q - Q(I + \widehat{T}(\lambda))Q \\ &= -(V_2(R_0((\lambda + i0)^2) - R_0(0))V\phi + \lambda c_0(\lambda)V_2\phi) \otimes V_1\phi \\ &= -\lambda \left(V_2 \frac{R_0((\lambda + i0)^2) - R_0(0)}{\lambda} V\phi + c_0(\lambda)V_2\phi \right) \otimes V_1\phi. \end{aligned} \tag{2-18}$$

Likewise,

$$\begin{aligned} T_{10}(\lambda) &= -V_2\phi \otimes (V_1(R_0^*((\lambda + i0)^2) - R_0(0))V\phi + \overline{\lambda c_0(\lambda)}V_1\phi) \\ &= -\lambda V_2\phi \otimes \left(V_1 \frac{R_0^*((\lambda + i0)^2) - R_0(0)}{\lambda} V\phi + \overline{c_0(\lambda)}V_1\phi \right). \end{aligned} \tag{2-19}$$

Thus T_{10}^\vee and T_{01}^\vee are both in $\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}} - T_{10}^\vee$ by the second line of (2-18) and T_{01}^\vee by the first line of (2-19) — when $V \in L^{\frac{3}{2},1}$. Indeed, following the definition, this reduces to

$$\int_{\mathbb{R}^3} \frac{|V_2(x)||V(y)||\phi(y)|}{4\pi|x-y|} dy \in L_x^{\frac{3}{2},2} \cap L_x^{3,2}.$$

Next, $-T_{10}(\lambda)T_{00}^{-1}(\lambda)T_{01}(\lambda) = \lambda c_1(\lambda)Q$, where

$$c_1(\lambda) = -\left\langle V_1(R_0^*((\lambda+i0)^2) - R_0(0))V\phi + \overline{\lambda c_0(\lambda)}V_1\phi, T_{00}^{-1}(\lambda)\left(V_2\frac{R_0((\lambda+i0)^2) - R_0(0)}{\lambda}V\phi + c_0(\lambda)V_2\phi\right)\right\rangle. \quad (2-20)$$

For example, one term from formula (2-20) has the form

$$\left\langle V_1(R_0^*((\lambda+i0)^2) - R_0(0))V_2V_1\phi, T_{00}^{-1}(\lambda)V_2\frac{R_0((\lambda+i0)^2) - R_0(0)}{\lambda}V_1V_2\phi\right\rangle. \quad (2-21)$$

Note that $V_1(R_0^*((\lambda+i0)^2) - R_0(0))V_2$ and $\chi(\frac{\lambda}{\epsilon})T_{00}^{-1}(\lambda)$ are in \mathcal{W} , while

$$M\left(V_2\frac{R_0((\lambda+i0)^2) - R_0(0)}{\lambda}V_1\right) = \frac{|V_2| \otimes |V_1|}{4\pi} \in \mathcal{B}(L^{\frac{3}{2},2}, L^{3,2}),$$

so

$$V_2\frac{R_0((\lambda+i0)^2) - R_0(0)}{\lambda}V_1 \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}.$$

Taking into account the fact that $V_1\phi, V_2\phi \in L^{\frac{3}{2},2}$, it follows that (2-21) is in \widehat{L}^1 .

Thus we recognize from (2-20) that $c_1(\lambda) \in \widehat{L}^1$ when $V \in L^{\frac{3}{2},1}$.

Further note that, since $R_0^*((\lambda+i0)^2) - R_0(0) = 0$ when $\lambda = 0$, we have $c_1(0) = 0$.

Let

$$C(\lambda) := T_{11}(\lambda) - T_{10}(\lambda)T_{00}^{-1}(\lambda)T_{01}(\lambda).$$

Then

$$C(\lambda) = \lambda(c_0(\lambda) + c_1(\lambda))Q.$$

Thus $C(\lambda)/\lambda$ is invertible for $|\lambda| \ll 1$ and $C^{-1}(\lambda) = \lambda^{-1}c_2(\lambda)Q$, with c_2 locally in \widehat{L}^1 . Consequently, for small ϵ , we have $(\chi(\frac{\lambda}{\epsilon})\lambda C^{-1}(\lambda))^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$.

The inverse of $I + \widehat{T}(\lambda)$ is then given for small λ by formula (2-13):

$$(I + \widehat{T})^{-1} = \begin{pmatrix} T_{00}^{-1} + T_{00}^{-1}T_{01}C^{-1}T_{10}T_{00}^{-1} & -T_{00}^{-1}T_{01}C^{-1} \\ -C^{-1}T_{10}T_{00}^{-1} & C^{-1} \end{pmatrix}.$$

Since $T_{00}^{-1} \in \mathcal{W}$ and $T_{01}^\vee, T_{10}^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}$, while

$$\left(\chi\left(\frac{\lambda}{\epsilon}\right)\lambda C^{-1}(\lambda)\right)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}},$$

it immediately follows that

$$\lambda((I + \widehat{T}(\lambda))^{-1} - T_{00}^{-1}(\lambda)) \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$$

and that $(I + \widehat{T})^{-1}$, given by formula (2-13), exists on a whole cut neighborhood of zero. \square

Recall that by (1-2)

$$R(t) := \frac{ae^{-i\frac{3\pi}{4}}}{\sqrt{i\pi t}} \zeta_t(x) \otimes \zeta_t(y), \quad \zeta_t(x) := e^{i\frac{|x|^2}{4t}} \phi(x).$$

Proposition 2.13. *Assume that $\langle x \rangle^2 V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is an exceptional Hamiltonian of the first kind with canonical resonance ϕ at zero. Then, for $1 \leq p < \frac{3}{2}$ and $R(t)$ as above,*

$$e^{-itH} P_c u = Z(t)u + R(t)u,$$

$$\|Z(t)u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|f\|_{L^p}, \quad \|Z(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|f\|_{L^{3/2,1}}.$$

Furthermore, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH} P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|u\|_{L^p}.$$

Here $\frac{1}{p} + \frac{1}{p'} = 1$.

Proof of Proposition 2.13. Write the evolution as

$$e^{-itH} P_c f = \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda + i0)^2) - R_0((\lambda + i0)^2) V_1 \widehat{T}(\lambda)^{-1} V_2 R_0((\lambda + i0)^2)) f \lambda \, d\lambda.$$

We consider a partition of unity subordinated to the neighborhoods of Lemmas 2.8 and 2.11. First, take a sufficiently large R such that $(1 - \chi(\frac{\lambda}{R}))(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$. Then for every $\lambda_0 \in [-4R, 4R]$ there exists $\epsilon(\lambda_0) > 0$ such that

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon(\lambda_0)}\right) (I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$$

if $\lambda_0 \neq 0$, while the conclusion of Lemma 2.11 holds when $\lambda_0 = 0$.

Since $[-4R, 4R]$ is a compact set, there exists a finite covering

$$[-4R, 4R] \subset \bigcup_{k=1}^N (\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)).$$

Then we construct a finite partition of unity on \mathbb{R} by smooth functions $1 = \chi_0(\lambda) + \sum_{k=1}^N \chi_k(\lambda) + \chi_\infty(\lambda)$, where $\text{supp } \chi_\infty \subset \mathbb{R} \setminus (-2R, 2R)$, $\text{supp } \chi_0 \subset [-\epsilon(0), \epsilon(0)]$, and $\text{supp } \chi_k \subset [\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)]$.

By Lemma 2.8, for any $k \neq 0$, we have $\chi_k(\lambda)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$, so $(1 - \chi_0(\lambda))(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$. By Lemma 2.11, $\chi_0(\lambda)\widehat{T}(\lambda)$ also decomposes into a regular term $L \in \mathcal{W}$ and a singular term $-\lambda^{-1}\chi_0(\lambda)aV_2\phi \otimes V_1\phi$.

Let Z_1 be given by the sum of all the regular terms in the decomposition:

$$Z_1(t) := \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda + i0)^2) - R_0((\lambda + i0)^2) V_1 L(\lambda) V_2 R_0((\lambda + i0)^2) - (1 - \chi_0(\lambda)) R_0((\lambda + i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda + i0)^2)) \lambda \, d\lambda$$

$$= \frac{1}{2\pi t} \int_{\mathbb{R}} e^{-it\lambda^2} \partial_\lambda (R_0((\lambda + i0)^2) - R_0((\lambda + i0)^2) V_1 L(\lambda) V_2 R_0((\lambda + i0)^2) - (1 - \chi_0(\lambda)) R_0((\lambda + i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda + i0)^2)) \, d\lambda$$

$$= \frac{C}{t^{\frac{3}{2}}} \int_{\mathbb{R}} e^{-i\frac{\rho^2}{4t}} (\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee(\rho) d\rho.$$

The fact that $\|Z_1(t)u\|_{L^\infty} \lesssim |t|^{-\frac{3}{2}}\|u\|_{L^1}$ follows by knowing that

$$(\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee \in \mathcal{V}_{L^1, L^\infty}.$$

The fact that $\|Z_1(t)u\|_{L^2} \lesssim \|u\|_{L^2}$ follows by smoothing estimates. Indeed, the first term is bounded since it represents the free evolution, and note that

$$\begin{aligned} \|V_2R_0(\lambda \pm i0)f\|_{L^2_{\lambda,x}} &\lesssim \|f\|_{L^2_x}, \\ \|e^{-it\lambda}(L(\pm\sqrt{\lambda}) + (1-\chi_0(\pm\sqrt{\lambda}))\widehat{T}(\pm\sqrt{\lambda}))\|_{L^\infty_{\lambda}B(L^2)} &< \infty, \\ \left\| \int_{\mathbb{R}} R_0(\lambda \pm i0)V_1F(x, \lambda) d\lambda \right\|_{L^2_x} &\lesssim \|F\|_{L^2_{\lambda,x}}. \end{aligned}$$

Combining these three estimates, we obtain the L^2 boundedness of Z_1 .

By interpolation between the two bounds, we obtain that, for $\frac{1}{p} + \frac{1}{p'} = 1$, with $1 \leq p \leq 2$,

$$\|Z_1(t)u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p} - \frac{1}{p'})} \|u\|_{L^p},$$

as well as

$$\|Z_1(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

Let Z_2 be the term corresponding to the singular part of the decomposition from [Lemma 2.11](#), given by

$$\begin{aligned} Z_2(t) &:= \frac{a}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} \chi_0(\lambda) R_0((\lambda+i0)^2) V\phi \otimes V\phi R_0((\lambda+i0)^2) d\lambda \\ &= \frac{a}{i\pi} \int_{\mathbb{R}} \int_{(\mathbb{R}^3)^2} e^{-it\lambda^2} \chi_0(\lambda) \frac{e^{i\lambda|x-z_1|}}{4\pi|x-z_1|} V(z_1)\phi(z_1) V(z_2)\phi(z_2) \frac{e^{i\lambda|z_2-y|}}{4\pi|z_2-y|} dz_1 dz_2 d\lambda. \end{aligned}$$

The subsequent [Lemma 2.14](#) is the same as [[Yajima 2005](#), Lemma 4.10], the only difference being the space of potentials for which the result holds. For the sake of completeness, we repeat the proof given in [[Yajima 2005](#)].

Lemma 2.14. For $V \in \langle x \rangle^{-1} L^{\frac{3}{2},1}$,

$$\|(Z_2(t) - R(t))u\|_{L^\infty} \lesssim t^{-\frac{3}{2}} \|u\|_{L^1}, \quad \|Z_2(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}. \quad (2-22)$$

Proof of Lemma 2.14. Let $b = |x - z_1| + |z_2 - y|$ and

$$C(t, b) = \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2 + i\lambda b} \chi_0(\lambda) d\lambda.$$

We express $Z_2(t)$ as

$$Z_2(t) = \int_{(\mathbb{R}^3)^2} C(t, b) a \frac{V(z_1)\phi(z_1)V(z_2)\phi(z_2)}{|x - z_1||z_2 - y|} dz_1 dz_2.$$

Note that

$$C(t, b) = \frac{e^{-i\frac{3\pi}{4}} e^{i\frac{b^2}{4t}}}{\sqrt{\pi t}} (e^{i\frac{s^2}{4t}} \chi_0^\vee(s))^\wedge \left(\frac{b}{2t}\right).$$

Then $C(t, b) \lesssim t^{-\frac{1}{2}}$ and

$$|Z_2(t)(x, y)| \lesssim t^{-\frac{1}{2}} \int_{(\mathbb{R}^3)^2} \frac{|V(z_1)\phi(z_1)V(z_2)\phi(z_2)|}{|z_1 - x||z_2 - y|} dz_1 dz_2.$$

Clearly

$$\int_{\mathbb{R}^3} \frac{|V(z_1)\phi(z_1)|}{|z_1 - x|} dz_1 \in L_x^{3,\infty} \quad \text{and} \quad \int_{\mathbb{R}^3} \frac{|V(z_2)\phi(z_2)|}{|z_2 - y|} dz_2 \in L_y^{3,\infty},$$

implying the second half of (2-22):

$$\|Z_2(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

We also have

$$\left| \mathcal{F}(e^{i\frac{s^2}{4t}} \chi_0^\vee(s)) \left(\frac{b}{2t}\right) - 1 \right| \lesssim t^{-1} (\|s^2 \chi_0^\vee\|_1 + |b|).$$

It is easy to see, for

$$B = 2(|x - z_1||z_1| + |z_2 - y||z_2| + |x - z_1||z_2 - y|) + |z_1|^2 + |z_2|^2,$$

that

$$|e^{ib^2/4t} - e^{i(x^2+y^2)/4t}| = |e^{i(|x-z_1|+|z_2-y|)^2/4t} - e^{i(x^2+y^2)/4t}| \leq \frac{B}{4t}.$$

It follows that

$$C(t, b) - \frac{e^{-i\frac{3\pi}{4}} e^{i(x^2+y^2)/4t}}{\sqrt{\pi t}} \lesssim (1 + b + B)t^{-\frac{3}{2}}.$$

Then

$$\begin{aligned} \left| Z_2(t) - \int_{(\mathbb{R}^3)^2} \frac{e^{i\frac{3\pi}{4}} e^{i(x^2+y^2)/4t}}{\sqrt{\pi t}} a \frac{V(z_1)\phi(z_1)V(z_2)\phi(z_2)}{|x - z_1||y - z_2|} dz_1 dz_2 \right| \\ \lesssim t^{-\frac{3}{2}} \int_{(\mathbb{R}^3)^2} \frac{(1 + b + B) |V(z_1)\phi(z_1)V(z_2)\phi(z_2)|}{|x - z_1||z_2 - y|} dz_1 dz_2. \end{aligned}$$

Now note that, for $V \in \langle x \rangle^{-1} L^{\frac{3}{2},1}$ and $\phi(x) \lesssim |x|^{-1}$,

$$\sup_{x,y} \int_{(\mathbb{R}^3)^2} \frac{(1 + b + B) |V(z_1)\phi(z_1)V(z_2)\phi(z_2)|}{|x - z_1||z_2 - y|} dz_1 dz_2 < \infty$$

and

$$\int_{\mathbb{R}^3} \frac{V(z_1)\phi(z_1)}{|x - z_1|} dz_1 = \phi(x).$$

The first part of conclusion (2-22) follows. □

Note that $R(t)$ also satisfies $\|R(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$, so the same holds for the difference:

$$\|(Z_2(t) - R(t))u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}.$$

By interpolation with the L^1 -to- L^∞ estimate of [Lemma 2.14](#), we obtain that, for $1 \leq p < \frac{3}{2}$,

$$\|(Z_2(t) - R(t))u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}\left(\frac{1}{p} - \frac{1}{p'}\right)}\|u\|_{L^p}.$$

Since the same is true for Z_1 , we obtain for $1 \leq p < \frac{3}{2}$ that

$$\|(Z(t)u)\|_{L^{p'}} = \|(Z_1(t) + Z_2(t) - R(t))u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}\left(\frac{1}{p} - \frac{1}{p'}\right)}\|u\|_{L^p},$$

where $e^{-itH}P_c u = Z_1(t)u + Z_2(t)u = Z(t)u + R(t)u$.

Knowing that $\|Z_i(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$ leads to the conclusion that $\|e^{-itH}P_c u\|_{L^{3,\infty}} \lesssim \|u\|_{L^{3/2,1}}$. Combining this with the L^2 estimate $\|e^{-itH}P_c u\|_{L^2} \lesssim \|u\|_{L^2}$, we obtain that, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH}P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}\left(\frac{1}{p} - \frac{1}{p'}\right)}\|u\|_{L^p}.$$

Thus we have proved all the conclusions of [Proposition 2.13](#). □

Proposition 2.15. *Assume that $V \in L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is an exceptional Hamiltonian of the first kind. Then*

$$\|e^{-itH}P_c u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}},$$

and, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH}P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}\left(\frac{1}{p} - \frac{1}{p'}\right)}\|u\|_{L^p}.$$

Here $\frac{1}{p} + \frac{1}{p'} = 1$.

Proof of Proposition 2.15. Write the evolution as

$$e^{-itH}P_c f = \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda + i0)^2) - R_0((\lambda + i0)^2)V_1 \hat{T}(\lambda)^{-1}V_2 R_0((\lambda + i0)^2)) f \lambda d\lambda.$$

We consider a partition of unity subordinated to the neighborhoods of [Lemmas 2.8](#) and [2.12](#). First, take a sufficiently large R such that $(1 - \chi(\frac{\lambda}{R}))(I + \hat{T}(\lambda))^{-1} \in \mathcal{W}$. Then, for every $\lambda_0 \in [-4R, 4R]$, there exists $\epsilon(\lambda_0) > 0$ such that

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon(\lambda_0)}\right)(I + \hat{T}(\lambda))^{-1} \in \mathcal{W}$$

if $\lambda_0 \neq 0$, while the conclusion of [Lemma 2.12](#) holds when $\lambda_0 = 0$.

Since $[-4R, 4R]$ is a compact set, there exists a finite covering

$$[-4R, 4R] \subset \bigcup_{k=1}^N (\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)).$$

Then we construct a finite partition of unity on \mathbb{R} by smooth functions $1 = \chi_0(\lambda) + \sum_{k=1}^N \chi_k(\lambda) + \chi_\infty(\lambda)$, where $\text{supp } \chi_\infty \subset \mathbb{R} \setminus (-2R, 2R)$, $\text{supp } \chi_0 \subset [-\epsilon(0), \epsilon(0)]$, and $\text{supp } \chi_k \subset [\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)]$.

By Lemma 2.8, for any $k \neq 0$, we have $\chi_k(\lambda)(I + \hat{T}(\lambda))^{-1} \in \mathcal{W}$, so $(1 - \chi_0(\lambda))(I + \hat{T}(\lambda))^{-1} \in \mathcal{W}$. By Lemma 2.12, $\chi_0(\lambda)(I + \hat{T}(\lambda))^{-1}$ also decomposes into a regular term $L \in \mathcal{W}$ and a singular term $\lambda^{-1}S$, with the property that $S^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$.

Let Z_1 be given by the sum of all the regular terms of the decomposition:

$$\begin{aligned} Z_1(t) &:= \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\hat{T}(\lambda)V_2R_0((\lambda+i0)^2))\lambda d\lambda \\ &= \frac{1}{2\pi t} \int_{\mathbb{R}} e^{-it\lambda^2} \partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\hat{T}(\lambda)V_2R_0((\lambda+i0)^2)) d\lambda \\ &= \frac{C}{t^{\frac{3}{2}}} \int_{\mathbb{R}} e^{-i\frac{\rho^2}{4t}} (\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\hat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee(\rho) d\rho. \end{aligned}$$

The fact that $\|Z_1(t)u\|_{L^\infty} \lesssim |t|^{-\frac{3}{2}}\|u\|_{L^1}$ follows by knowing that

$$\begin{aligned} &(\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\hat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee \in \mathcal{V}_{L^1, L^\infty}. \end{aligned}$$

Using smoothing estimates, it immediately follows that $Z_1(t)$ is L^2 -bounded; see the proof of Proposition 2.13. Interpolating, we obtain the desired $\|Z_1(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3,1}}$ estimate.

Let Z_2 be the singular part of the decomposition from Lemma 2.12, given by

$$Z_2(t) := \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} R_0((\lambda+i0)^2)V_1S(\lambda)V_2R_0((\lambda+i0)^2) d\lambda. \tag{2-23}$$

Note that $(R_0((\lambda+i0)^2)V_1)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,\infty}}$, $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$, and $(V_2R_0((\lambda+i0)^2))^\vee \in \mathcal{V}_{L^{3/2,1}, L^{3,2}}$. Thus

$$R_0((\lambda+i0)^2)V_1(\lambda S(\lambda))V_2R_0((\lambda+i0)^2) \in \mathcal{V}_{L^{3/2,1}, L^{3,\infty}}.$$

By taking the Fourier transform in (2-23), this immediately implies the conclusion that $\|Z_2(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$.

Putting the two estimates for Z_1 and Z_2 together, we obtain that $\|e^{-itH}P_cu\|_{L^{3,\infty}} \lesssim \|u\|_{L^{3/2,1}}$. Interpolating with the obvious L^2 bound $\|e^{-itH}P_cu\|_{L^2} \lesssim \|u\|_{L^2}$, we obtain the stated conclusion. \square

2F. Exceptional Hamiltonians of the third kind. We next consider the case in which H is exceptional of the third kind; that is, there are both zero eigenvectors and zero resonances. Recall that $\hat{T}(\lambda) = V_2R_0((\lambda+i0)^2)V_1$.

Lemma 2.16. *Suppose that $V \in \langle x \rangle^{-4}L^{\frac{3}{2},1}$ and $H = -\Delta + V$ has both eigenvectors and resonances at zero. Let χ be a standard cutoff function. Then, for sufficiently small ϵ ,*

$$\chi\left(\frac{\lambda}{\epsilon}\right)(I + \hat{T}(\lambda))^{-1} = L(\lambda) + \chi\left(\frac{\lambda}{\epsilon}\right)\left(\frac{V_2P_0V_1}{\lambda^2} + \frac{iV_2P_0V|x-y|^2VP_0V_1}{\lambda} - \frac{aV_2\phi \otimes V_1\phi}{\lambda}\right),$$

where $L(\lambda) \in \mathcal{W}$ and ϕ is a certain resonance for $H = -\Delta + V$.

Furthermore, 0 is an isolated exceptional point for H , meaning that H has finitely many negative eigenvalues.

The computations in the proof of this lemma parallel those in [Yajima 2005, Section 4.5]. The main difference is in using the space \mathcal{W} instead of Hölder spaces.

Proof of Lemma 2.16. We study $(I + \hat{T}(\lambda))^{-1} := (I + V_2 R_0((\lambda + i0)^2) V_1)^{-1}$ near $\lambda = 0$.

Let

$$Q = -\frac{1}{2\pi i} \int_{|z+1|=\delta} (V_2 R_0(0) V_1 - zI)^{-1} dz.$$

Take the orthonormal basis $\{\phi_1, \dots, \phi_N\}$ with respect to the inner product $-(Vu, v)$ for \mathcal{M} so that $\{\phi_2, \dots, \phi_N\}$ is a basis of \mathcal{E} and $\langle \phi_1, V \rangle > 0$. This condition determines ϕ_1 uniquely.

Define the orthogonal projections π_1 onto $\mathbb{C}V_1\phi_1$ and π_2 onto $V_1 P_0 L^2$ with respect to the inner product $-(\text{sgn } Vu, v)$, i.e., $\pi_1 = -V_2\phi_1 \otimes V_1\phi_1$ and $\pi_2 = -\sum_{j=2}^N V_2\phi_j \otimes V_1\phi_j$, and let

$$Q_0 = \bar{Q} := 1 - Q, \quad Q_1 := Q\pi_1 Q, \quad Q_2 := Q\pi_2 Q.$$

The following identities hold in L^2 :

$$\begin{aligned} Q_j Q_k &= \delta_{jk} I \quad \text{for } j, k = 0, 1, 2, & Q_0 + Q_1 + Q_2 &= I, \\ (I + V_2 R_0(0) V_1) Q_1 &= Q_1 (I + V_2 R_0(0) V_1) = 0, \\ (I + V_2 R_0(0) V_1) Q_2 &= Q_2 (I + V_2 R_0(0) V_1) = 0, \\ Q_2 (V_2 \otimes V_1) Q_0 &= 0, \quad Q_2 (V_2 \otimes V_1) Q_1 = 0, \quad Q_2 (V_2 \otimes V_1) Q_2 = 0, \\ Q_0 (V_2 \otimes V_1) Q_2 &= 0, \quad Q_1 (V_2 \otimes V_1) Q_2 = 0. \end{aligned}$$

These identities follow from $Q_2 V_2 = 0$ and $Q_2^* V_1 = 0$, which in turn follow from the fact that eigenvectors ϕ_k are orthogonal to V , that is, $\langle \phi_k, V \rangle = 0$ for $2 \leq k \leq N$.

We first apply Lemma 2.10 to invert $Q(I + \hat{T}(\lambda))Q$ in QL^2 for small λ , after writing it in matrix form with respect to the decomposition $QL^2 = Q_1 L^2 + Q_2 L^2$:

$$Q(I + \hat{T}(\lambda))Q = \begin{pmatrix} Q_1(I + \hat{T}(\lambda))Q_1 & Q_1 \hat{T}(\lambda) Q_2 \\ Q_2 \hat{T}(\lambda) Q_1 & Q_2(I + \hat{T}(\lambda))Q_2 \end{pmatrix} =: \begin{pmatrix} T_{11}(\lambda) & T_{12}(\lambda) \\ T_{21}(\lambda) & T_{22}(\lambda) \end{pmatrix}.$$

The inverse will be given by formula (2-13); that is,

$$(Q(I + \hat{T}(\lambda))Q)^{-1} = \begin{pmatrix} T_{11}^{-1} + T_{11}^{-1} T_{12} C_{22}^{-1} T_{21} T_{11}^{-1} & -T_{11}^{-1} T_{12} C_{22}^{-1} \\ -C_{22}^{-1} T_{21} T_{11}^{-1} & C_{22}^{-1} \end{pmatrix}, \quad (2-24)$$

where

$$C_{22} = T_{22} - T_{21} T_{11}^{-1} T_{12}.$$

As in the case of exceptional Hamiltonians of the first kind, let

$$J(\lambda) := \frac{\hat{T}(\lambda) - (V_2 R_0(0) V_1 + i\lambda(4\pi)^{-1} V_2 \otimes V_1)}{\lambda^2}.$$

Then (recall that $Q_1 = -V_2\phi_1 \otimes V_1\phi_1$),

$$\begin{aligned} T_{11}(\lambda) &= Q_1(I + \widehat{T}(\lambda))Q_1 = Q_1(I + V_2R_0((\lambda + i0)^2)V_1)Q_1 \\ &= Q_1(V_2R_0((\lambda + i0)^2)V_1 - V_2R_0(0)V_1)Q_1 \\ &= V_2\phi_1 \otimes V\phi_1(R_0((\lambda + i0)^2) - R_0(0))V\phi_1 \otimes V_1\phi_1 \\ &= \left(\lambda \frac{|\langle V, \phi_1 \rangle|^2}{4i\pi} - \lambda^2 \langle V_1\phi_1, J(\lambda)V_2\phi_1 \rangle \right) Q_1 \\ &=: (\lambda a^{-1} + \lambda^2 c_1(\lambda))Q_1. \end{aligned}$$

Here $a = 4i\pi/|\langle V, \phi_1 \rangle|^2 \neq 0$. As in the proof of [Lemma 2.11](#), note that $c_1(\lambda) \in \widehat{L}^1$ when $V \in L^1$ and $\partial_\lambda c_1(\lambda) \in \widehat{L}^1$ when $V \in \langle x \rangle^{-1}L^1$.

It follows that $T_{11}(\lambda)$ is invertible for $|\lambda| \ll 1$ in Q_1L^2 and

$$\begin{aligned} T_{11}^{-1}(\lambda) &= \frac{1}{\lambda a^{-1} + \lambda^2 c_1(\lambda)} Q_1 \\ &= \left(\frac{a}{\lambda} - \frac{c_1(\lambda)}{(a^{-1} + \lambda c_1(\lambda))a^{-1}} \right) Q_1 \\ &= \lambda^{-1}aQ_1 + E(\lambda). \end{aligned}$$

Here and below we denote various regular terms by $E(\lambda)$, i.e., terms with the property that $\chi(\frac{\lambda}{\epsilon})E(\lambda) \in \mathcal{W}$ for sufficiently small ϵ .

Likewise, since $Q_2(V_2 \otimes V_1) = (V_2 \otimes V_1)Q_2 = 0$,

$$\begin{aligned} T_{12}(\lambda) &= Q_1(I + V_2R_0((\lambda + i0)^2)V_1)Q_2 \\ &= Q_1V_2 \left(R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi} \right) V_1Q_2 \\ &= -\lambda^2 Q_1 \left(V_2 \frac{|x-y|}{8\pi} V_1 + \lambda V_2 e_1(\lambda) V_1 \right) Q_2 \\ &= -\lambda^2 Q_1 V_2 \frac{|x-y|}{8\pi} V_1 Q_2 + \lambda^3 E(\lambda), \end{aligned}$$

where

$$e_1(\lambda) := \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi} + \lambda^2 \frac{|x-y|}{8\pi}}{-\lambda^3}.$$

By [Lemma 2.6](#),

$$M((e_1(\lambda))^\wedge) = \frac{|x-y|^2}{24\pi} \quad \text{and} \quad M((\partial_\lambda e_1(\lambda))^\wedge) = \frac{|x-y|^3}{96\pi}.$$

Thus $E(\lambda) := Q_1V_2e_1(\lambda)V_1Q_2 \in \mathcal{W}$ when

$$\int_{(\mathbb{R}^3)^2} V(x)\phi_1(x)|x-y|^3V(y)\phi_k(y) dx dy < \infty,$$

which takes place when $V \in \langle x \rangle^{-2}L^1$ (recall that $|\phi_1(y)| \lesssim \langle y \rangle^{-1}$).

Likewise we obtain

$$T_{21}(\lambda) = -\lambda^2 Q_2 V_2 \frac{|x-y|}{8\pi} V_1 Q_1 + \lambda^3 E(\lambda);$$

hence, combining the previous results,

$$T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) = \lambda^3 a Q_2 V_2 \frac{|x-y|}{8\pi} V_1 Q_1 V_2 \frac{|x-y|}{8\pi} V_1 Q_2 + E(\lambda).$$

Furthermore,

$$\begin{aligned} T_{22}(\lambda) &= Q_2 (I + V_2 R_0((\lambda + i0)^2) V_1) Q_2 \\ &= Q_2 V_2 \left(R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi} \right) V_1 Q_2 \\ &= -\lambda^2 Q_2 \left(V_2 \frac{|x-y|}{8\pi} V_1 + i\lambda V_2 \frac{|x-y|^2}{24\pi} V_1 - \lambda^2 V_2 e_2(\lambda) V_1 \right) Q_2. \end{aligned}$$

Here

$$e_2(\lambda) := \lambda^{-4} \left(R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi} - i\lambda^2 \frac{|x-y|}{8\pi} - \lambda^3 \frac{|x-y|^2}{24\pi} \right).$$

By [Lemma 2.6](#),

$$M(e_2(\lambda)^\wedge) = \frac{|x-y|^3}{96\pi} \quad \text{and} \quad M((\partial_\lambda e_2(\lambda))^\wedge) = \frac{|x-y|^4}{480\pi}.$$

Thus $E(\lambda) := Q_2 V_2 e_2(\lambda) V_1 Q_2 \in \mathcal{W}$ when

$$\int_{(\mathbb{R}^3)^2} V(x) \phi_k(x) |x-y|^4 V(y) \phi_k(y) dx dy < \infty,$$

which holds true when $V \in \langle x \rangle^{-2} L^1$ (recall that $|\phi_k(y)| \lesssim \langle y \rangle^{-2}$). Then

$$T_{22}(\lambda) = -\lambda^2 Q_2 \left(V_2 \frac{|x-y|}{8\pi} V_1 + i\lambda V_2 \frac{|x-y|^2}{24\pi} V_1 \right) Q_2 + \lambda^4 E(\lambda). \quad (2-25)$$

Let P_0 be the L^2 orthogonal projection onto the set \mathcal{E} spanned by ϕ_2, \dots, ϕ_N . By relation (4.38) of [\[Yajima 2005\]](#),

$$\left(Q_2 V_2 \frac{|x-y|}{8\pi} V_1 Q_2 \right)^{-1} = -V_2 P_0 V_1.$$

Also note that

$$V_2 P_0 V_1 Q_2 = Q_2 V_2 P_0 V_1 = V_2 P_0 V_1.$$

By (2-25),

$$\begin{aligned} T_{22}^{-1}(\lambda) &= -\lambda^{-2} \left(Q_2 V_2 \frac{|x-y|}{8\pi} V_1 Q_2 \right)^{-1} \\ &\quad \sum_{k=0}^{\infty} (-1)^k \left(\left(i\lambda Q_2 V_2 \frac{|x-y|^2}{24\pi} V_1 Q_2 - \lambda^2 E(\lambda) \right) \left(Q_2 V_2 \frac{|x-y|}{8\pi} V_1 Q_2 \right)^{-1} \right)^k \\ &= \lambda^{-2} V_2 P_0 V_1 \sum_{k=0}^{\infty} \left(i\lambda V_2 \frac{|x-y|^2}{24\pi} V_1 - \lambda^2 E(\lambda) \right) V_2 P_0 V_1. \end{aligned}$$

Therefore, by grouping the terms by the powers of λ , for $|\lambda| \ll 1$,

$$T_{22}^{-1}(\lambda) = \lambda^{-2} V_2 P_0 V_1 + i \lambda^{-1} V_2 P_0 V \frac{|x-y|^2}{24\pi} V P_0 V_1 + E(\lambda).$$

Then we write

$$\begin{aligned} C_{22}(\lambda) &= T_{22}(\lambda) - T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) \\ &= (I - T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) T_{22}^{-1}(\lambda)) T_{22}(\lambda). \end{aligned}$$

By our previous estimates, $T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) T_{22}^{-1}(\lambda) = \lambda E(\lambda)$, where $E(\lambda) \in \mathcal{W}$. Then, by means of a Neumann series expansion, we retrieve that

$$\begin{aligned} C_{22}^{-1}(\lambda) &= T_{22}^{-1}(\lambda) \sum_{k=0}^{\infty} (T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) T_{22}^{-1}(\lambda))^k \\ &= T_{22}^{-1}(\lambda) + T_{22}^{-1}(\lambda) T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) T_{22}^{-1}(\lambda) + E(\lambda), \end{aligned}$$

so

$$\begin{aligned} C_{22}^{-1}(\lambda) &= \lambda^{-2} V_2 P_0 V_1 + i \lambda^{-1} V_2 P_0 V \frac{|x-y|^2}{24\pi} V P_0 V_1 \\ &\quad + a \lambda^{-1} V_2 P_0 V \frac{|x-y|}{8\pi} V_1 Q_1 V_2 \frac{|x-y|}{8\pi} V P_0 V_1 + E(\lambda). \end{aligned}$$

If we set

$$\tilde{\phi}_1 = P_0 V \frac{|x-y|}{8\pi} V \phi_1 \in \mathcal{E},$$

then

$$V_2 P_0 V \frac{|x-y|}{8\pi} V_1 Q_1 V_2 \frac{|x-y|}{8\pi} V P_0 V_1 = -V_2 \tilde{\phi}_1 \otimes \tilde{\phi}_1 V_1.$$

Then we get that

$$C_{22}^{-1}(\lambda) = \lambda^{-2} V_2 P_0 V_1 + i \lambda^{-1} V_2 P_0 V \frac{|x-y|^2}{24\pi} V P_0 V_1 - \lambda^{-1} a V_2 \tilde{\phi}_1 \otimes \tilde{\phi}_1 V_1 + E(\lambda).$$

Furthermore,

$$\begin{aligned} -T_{11}^{-1}(\lambda) T_{12}(\lambda) C_{22}^{-1}(\lambda) &= (\lambda^{-1} a Q_1 + E(\lambda)) \lambda^2 Q_1 \left(V_2 \frac{|x-y|}{8\pi} V_1 + \lambda E(\lambda) \right) Q_2 (\lambda^{-2} V_2 P_0 V_1 + i \lambda^{-1} E(\lambda)) \\ &= \lambda^{-1} a (-V_2 \phi_1 \otimes V_1 \phi_1) V_2 \frac{|x-y|}{8\pi} V P_0 V_1 + E(\lambda) \\ &= -a \lambda^{-1} V_2 \phi_1 \otimes \tilde{\phi}_1 V_1 + E(\lambda). \end{aligned}$$

Likewise we obtain

$$\begin{aligned} -C_{22}^{-1}(\lambda) T_{21}(\lambda) T_{11}^{-1}(\lambda) &= -a \lambda^{-1} V_2 \tilde{\phi}_1 \otimes \phi_1 V_1 + E(\lambda), \\ T_{11}^{-1}(\lambda) T_{12}(\lambda) C_{22}^{-1}(\lambda) T_{21}(\lambda) T_{11}^{-1}(\lambda) &= E(\lambda). \end{aligned}$$

By (2-24), we have that $(Q(I + \hat{T}(\lambda))Q)^{-1}$ is given in matrix form modulo $E(\lambda) \in \mathcal{W}$ by

$$\begin{pmatrix} -a \lambda^{-1} V_2 \phi_1 \otimes V_1 \phi_1 & -a \lambda^{-1} V_2 \phi_1 \otimes V_1 \tilde{\phi}_1 \\ -a \lambda^{-1} V_2 \tilde{\phi}_1 \otimes V_1 \phi_1 & \lambda^{-2} V_2 P_0 V_1 + i \lambda^{-1} V_2 P_0 V \frac{|x-y|^2}{24\pi} V P_0 V_1 - \lambda^{-1} a V_2 \tilde{\phi}_1 \otimes V_1 \tilde{\phi}_1 \end{pmatrix}. \quad (2-26)$$

Therefore, if we define the canonical resonance as $\phi = \phi_1 - \tilde{\phi}_1$, we have that ϕ satisfies $\phi \in \mathcal{M}$ and $\langle \phi, V \rangle = 1$ and

$$(Q(I + \hat{T}(\lambda))Q)^{-1} = \frac{V_2 P_0 V_1}{\lambda^2} + \frac{i V_2 P_0 V \frac{|x-y|^2}{24\pi} V P_0 V_1}{\lambda} - \frac{a V_2 \phi \otimes V_1 \phi}{\lambda} + E(\lambda). \quad (2-27)$$

We apply [Lemma 2.10](#) again after writing $I + \hat{T}(\lambda)$ in matrix form with respect to the decomposition $L^2 = \bar{Q}L^2 + QL^2$, where $QL^2 = V_2\mathcal{M}$:

$$I + \hat{T}(\lambda) = \begin{pmatrix} \bar{Q}(I + \hat{T}(\lambda))\bar{Q} & \bar{Q}\hat{T}(\lambda)Q \\ Q\hat{T}(\lambda)\bar{Q} & Q(I + \hat{T}(\lambda))Q \end{pmatrix} := \begin{pmatrix} S_{00}(\lambda) & S_{01}(\lambda) \\ S_{10}(\lambda) & S_{11}(\lambda) \end{pmatrix}.$$

Next, let $A(\lambda) := S_{00}(\lambda)^{-1}$. Then $\chi\left(\frac{\lambda}{\epsilon}\right)A(\lambda) \in \mathcal{W}$ for sufficiently small ϵ . Indeed, it is easy to see that $S_{00}(\lambda) \in \mathcal{W}$. Furthermore, $S_{00}(0)$ is invertible on $\bar{Q}L^{\frac{3}{2},2} \cap \bar{Q}L^{3,2}$ of inverse K ; see [\(2-12\)](#).

As in the proof of [Lemma 2.11](#), let

$$S_\epsilon(\lambda) = \chi\left(\frac{\lambda}{\epsilon}\right)\bar{Q}(\hat{T}(\lambda) - \hat{T}(0))\bar{Q}.$$

A simple argument based on condition [\(C1\)](#) shows that $\lim_{\epsilon \rightarrow 0} \|S_\epsilon(\lambda)\|_{\mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}} = 0$. Then

$$\begin{aligned} \chi\left(\frac{\lambda}{\epsilon}\right)S_{00}^{-1}(\lambda) &= \chi\left(\frac{\lambda}{\epsilon}\right)\left(S_{00}(0) + \chi\left(\frac{\lambda}{2\epsilon}\right)\bar{Q}(\hat{T}(\lambda) - \hat{T}(0))\bar{Q}\right)^{-1} \\ &= \chi\left(\frac{\lambda}{\epsilon}\right)S_{00}^{-1}(0) \sum_{k=0}^{\infty} (-1)^k (S_{2\epsilon}(\lambda)S_{00}^{-1}(0))^k. \end{aligned}$$

This series converges for sufficiently small ϵ , showing that $(\chi\left(\frac{\lambda}{\epsilon}\right)S_{00}^{-1}(\lambda))^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}}$.

Concerning the derivative,

$$\chi\left(\frac{\lambda}{\epsilon}\right)\partial_\lambda S_{00}^{-1}(\lambda) = -\chi\left(\frac{\lambda}{\epsilon}\right)S_{00}^{-1}(\lambda)\partial_\lambda S_{00}(\lambda)\chi\left(\frac{\lambda}{2\epsilon}\right)S_{00}^{-1}(\lambda).$$

In this expression,

$$\left(\chi\left(\frac{\lambda}{\epsilon}\right)S_{00}^{-1}(\lambda)\right)^\vee \in \mathcal{V}_{L^{3/2,2}} \cap \mathcal{V}_{L^{3,2}} \quad \text{and} \quad \left(\chi\left(\frac{\lambda}{2\epsilon}\right)\partial_\lambda S_{00}(\lambda)\right)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$$

since $M((\partial_\lambda T_{00}(\lambda))^\vee) = (|V_2| \otimes |V_1|)/(4\pi)$. Thus

$$\left(\chi\left(\frac{\lambda}{\epsilon}\right)\partial_\lambda S_{00}^{-1}(\lambda)\right)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,2}}.$$

From this we infer that $\chi\left(\frac{\lambda}{\epsilon}\right)A(\lambda) \in \mathcal{W}$, so A is a regular term.

We compute the inverse of $I + \hat{T}(\lambda)$ by finding each of its matrix elements:

$$(I + \hat{T}(\lambda))^{-1} = \begin{pmatrix} A + AS_{01}C^{-1}S_{10}A & AS_{01}C^{-1} \\ -C^{-1}S_{10}A & C^{-1} \end{pmatrix}. \quad (2-28)$$

Here

$$C(\lambda) = S_{11}(\lambda) - S_{10}(\lambda)A(\lambda)S_{01}(\lambda).$$

$S_{10}(\lambda)A(\lambda)S_{01}(\lambda) = Q\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q$ may be written as

$$\begin{pmatrix} Q_1\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_1 & Q_1\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 \\ Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_1 & Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 \end{pmatrix} = \begin{pmatrix} \lambda^2 E_{11}(\lambda) & \lambda^3 E_{12}(\lambda) \\ \lambda^3 E_{21}(\lambda) & \lambda^4 E_{22}(\lambda) \end{pmatrix}. \tag{2-29}$$

Indeed, consider, for example, $Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2$. It can be reexpressed as

$$Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 = \lambda^4 Q_2 V_2 \frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1 A(\lambda) V_2 \frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1 Q_2. \tag{2-30}$$

For this computation, we assume that $V \in \langle x \rangle^{-4} L^{\frac{3}{2},1}$. Taking a derivative of (2-30), we obtain terms such as

$$Q_2 V_2 \partial_\lambda \left(\frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} \right) V_1 A(\lambda) V_2 \frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1 Q_2. \tag{2-31}$$

Note that the range of Q_2 is spanned by functions $V_2 \phi_k$, with $2 \leq k \leq N$, such that $|\phi_k(y)| \lesssim \langle y \rangle^{-2}$ and $V_2 \in \langle x \rangle^{-2} L^{3,2}$, so $V_2 \phi \in \langle y \rangle^{-4} L^{3,2}$. Also

$$M \left(\left(V_2 \partial_\lambda \left(\frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} \right) V_1 \right)^\wedge \right) = |V_2| \frac{|x-y|^2}{24\pi} |V_1| \in \mathcal{B}(L^{\frac{3}{2},2}, L^{3,2}).$$

Likewise

$$M \left(\left(V_2 \frac{R_0((\lambda+i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1 \right)^\wedge \right) = |V_2| \frac{|x-y|}{8\pi} |V_1| \in \mathcal{B}(L^{\frac{3}{2},2}, L^{\frac{3}{2},2}).$$

This shows that (2-31) $\in \mathcal{V}_{L^{3/2,2}, L^{3,2}}$. By such computations, we obtain that $Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 = \lambda^4 E_{22}(\lambda)$, where $\chi(\frac{\lambda}{\epsilon}) E_{22}(\lambda) \in \mathcal{W}$ for sufficiently small ϵ . In this manner, we prove (2-29).

By (2-26), we have $S_{11}^{-1}(\lambda) = (Q\hat{T}(\lambda)Q)^{-1}$ is of the form

$$S_{11}^{-1}(\lambda) = \begin{pmatrix} \lambda^{-1} E(\lambda) & \lambda^{-1} E(\lambda) \\ \lambda^{-1} E(\lambda) & \lambda^{-2} E(\lambda) \end{pmatrix}.$$

Then, letting $N(\lambda) := S_{11}^{-1}(\lambda)S_{10}(\lambda)A(\lambda)S_{01}(\lambda)$, by (2-29),

$$\begin{aligned} N(\lambda) &:= S_{11}^{-1}(\lambda)S_{10}(\lambda)S_{00}^{-1}(\lambda)S_{01}(\lambda) \\ &= \begin{pmatrix} \lambda^{-1} E(\lambda) & \lambda^{-1} E(\lambda) \\ \lambda^{-1} E(\lambda) & \lambda^{-2} E(\lambda) \end{pmatrix} \begin{pmatrix} \lambda^2 E_{11}(\lambda) & \lambda^3 E_{12}(\lambda) \\ \lambda^3 E_{21}(\lambda) & \lambda^4 E_{22}(\lambda) \end{pmatrix} \\ &= \begin{pmatrix} \lambda E(\lambda) & \lambda^2 E(\lambda) \\ \lambda E(\lambda) & \lambda^2 E(\lambda) \end{pmatrix}. \end{aligned}$$

This shows that $C(\lambda)$ is invertible for $\lambda \ll 1$:

$$C(\lambda) = S_{11}(\lambda) - S_{10}(\lambda)A(\lambda)S_{01}(\lambda) = S_{11}(\lambda)(1 - N(\lambda)),$$

so

$$\begin{aligned} C^{-1}(\lambda) &= (I - N(\lambda))^{-1} S_{11}^{-1}(\lambda) \\ &= S_{11}^{-1}(\lambda) + (I - N(\lambda))^{-1} N(\lambda) S_{11}^{-1}(\lambda). \end{aligned} \tag{2-32}$$

A computation shows that $(I - N(\lambda))^{-1}N(\lambda)S_{11}^{-1}(\lambda)$ is a regular term:

$$\begin{aligned} (1 - N(\lambda))^{-1}N(\lambda)S_{11}^{-1}(\lambda) &= E(\lambda) \begin{pmatrix} \lambda E(\lambda) & \lambda^2 E(\lambda) \\ \lambda E(\lambda) & \lambda^2 E(\lambda) \end{pmatrix} \begin{pmatrix} \lambda^{-1} E(\lambda) & \lambda^{-1} E(\lambda) \\ \lambda^{-1} E(\lambda) & \lambda^{-2} E(\lambda) \end{pmatrix} \\ &= E(\lambda). \end{aligned}$$

By (2-32) and (2-27),

$$\begin{aligned} C^{-1}(\lambda) &= S_{11}^{-1}(\lambda) + E(\lambda) \\ &= \lambda^{-2}V_2P_0V_1 + i\lambda^{-1}V_2P_0V\frac{|x-y|^2}{24}VP_0V_1 - a\lambda^{-1}V_2\phi \otimes V_1\phi + E(\lambda). \end{aligned}$$

One can then also write C^{-1} as

$$C^{-1}(\lambda) = \begin{pmatrix} \lambda^{-1}E(\lambda) & \lambda^{-1}E(\lambda) \\ \lambda^{-1}E(\lambda) & \lambda^{-2}E(\lambda) \end{pmatrix}.$$

We also have

$$S_{01}(\lambda) = \bar{Q}(I + \hat{T}(\lambda))Q = \lambda E_1(\lambda)Q_1 + \lambda^2 E_2(\lambda)Q_2$$

with regular terms $E_1, E_2 \in \mathcal{W}$:

$$\begin{aligned} E_1(\lambda) &:= \bar{Q}V_2 \frac{R_0((\lambda + i0)^2) - R_0(0)}{\lambda} V_1Q_1, \\ E_2(\lambda) &:= \bar{Q}V_2 \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda 1 \otimes 1}{\lambda^2} V_1Q_2. \end{aligned}$$

Showing that $E_1, E_2 \in \mathcal{W}$ requires assuming that $V \in \langle x \rangle^{-4}L^{\frac{3}{2},1}$.

Therefore, the following matrix element of (2-28) is regular near zero:

$$A(\lambda)S_{01}(\lambda)C^{-1}(\lambda) = (\lambda A(\lambda)E_1(\lambda) \lambda^2 A(\lambda)E_2(\lambda)) \begin{pmatrix} \lambda^{-1}E(\lambda) & \lambda^{-1}E(\lambda) \\ \lambda^{-1}E(\lambda) & \lambda^{-2}E(\lambda) \end{pmatrix} = E(\lambda).$$

One shows in the same manner that the matrix element $C^{-1}(\lambda)S_{10}(\lambda)A(\lambda)$ of (2-28) is regular near zero.

Finally, the last remaining matrix element $A + AS_{01}C^{-1}S_{10}A$ of (2-28) consists of the regular part A and

$$\begin{aligned} AS_{01}C^{-1}S_{10}A &= E(\lambda) (\lambda E(\lambda) \lambda^2 E(\lambda)) \begin{pmatrix} \lambda^{-1}E(\lambda) & \lambda^{-1}E(\lambda) \\ \lambda^{-1}E(\lambda) & \lambda^{-2}E(\lambda) \end{pmatrix} \begin{pmatrix} \lambda E(\lambda) \\ \lambda^2 E(\lambda) \end{pmatrix} E(\lambda) \\ &= \lambda E(\lambda). \end{aligned}$$

Thus this is also a regular term. It follows by (2-28) that $\hat{T}(\lambda)^{-1}$ is up to regular terms given by

$$\lambda^{-2}V_2P_0V_1 + i\lambda^{-1}V_2P_0V\frac{|x-y|^2}{24}VP_0V_1 - a\lambda^{-1}V_2\phi \otimes V_1\phi,$$

which was to be shown. \square

We next prove a corresponding statement in the case when V has an almost minimal amount of decay. One can also obtain a resolvent expansion when $V \in \langle x \rangle^{-1}L^{\frac{3}{2},1}$, but it does not lead to decay estimates.

Lemma 2.17. *Suppose that $V \in \langle x \rangle^{-2} L^{\frac{3}{2},1}$ and $H = -\Delta + V$ is an exceptional Hamiltonian of the third kind. Let χ be a standard cutoff function. Then, for sufficiently small ϵ ,*

$$\chi\left(\frac{\lambda}{\epsilon}\right)(I + \hat{T}(\lambda))^{-1} = L(\lambda) + \lambda^{-1} S(\lambda) + \lambda^{-2} V_2 P_0 V_1,$$

where $L(\lambda) \in \mathcal{W}$, $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$, and P_0 is the L^2 orthogonal projection on \mathcal{E} .

Furthermore, 0 is an isolated exceptional point, so H has finitely many negative eigenvalues.

Proof of Lemma 2.17. We study $(I + \hat{T}(\lambda))^{-1} := (I + V_2 R_0((\lambda + i0)^2) V_1)^{-1}$ near $\lambda = 0$.

Let $Q = Q_1 + Q_2$, $Q_0 = \bar{Q}$, and Q_1 and Q_2 be as in the proof of Lemma 2.16.

Also take again the orthonormal basis $\{\phi_1, \dots, \phi_N\}$ with respect to the inner product $-(Vu, v)$ for \mathcal{M} so that $\{\phi_2, \dots, \phi_N\}$ is a basis of \mathcal{E} and $\langle \phi_1, V \rangle > 0$.

We apply Lemma 2.10 to invert $Q(I + \hat{T}(\lambda))Q$ in QL^2 for small λ , after writing it in matrix form with respect to the decomposition $QL^2 = Q_1 L^2 + Q_2 L^2$:

$$Q(I + \hat{T}(\lambda))Q = \begin{pmatrix} Q_1(I + \hat{T}(\lambda))Q_1 & Q_1 \hat{T}(\lambda) Q_2 \\ Q_2 \hat{T}(\lambda) Q_1 & Q_2(I + \hat{T}(\lambda))Q_2 \end{pmatrix} := \begin{pmatrix} T_{11}(\lambda) & T_{12}(\lambda) \\ T_{21}(\lambda) & T_{22}(\lambda) \end{pmatrix}.$$

The inverse will be given by formula (2-13), that is,

$$(Q(I + \hat{T}(\lambda))Q)^{-1} = \begin{pmatrix} T_{11}^{-1} + T_{11}^{-1} T_{12} C_{22}^{-1} T_{21} T_{11}^{-1} & -T_{11}^{-1} T_{12} C_{22}^{-1} \\ -C_{22}^{-1} T_{21} T_{11}^{-1} & C_{22}^{-1} \end{pmatrix}, \tag{2-33}$$

where

$$C_{22} = T_{22} - T_{21} T_{11}^{-1} T_{12}.$$

Then (recall that $Q_1 = -V_2 \phi_1 \otimes V_1 \phi_1$),

$$\begin{aligned} T_{11}(\lambda) &= Q_1(I + \hat{T}(\lambda))Q_1 = Q_1(I + V_2 R_0((\lambda + i0)^2) V_1)Q_1 \\ &= Q_1(V_2 R_0((\lambda + i0)^2) V_1 - V_2 R_0(0) V_1)Q_1 \\ &= V_2 \phi_1 \otimes V \phi_1 (R_0((\lambda + i0)^2) - R_0(0)) V \phi_1 \otimes V_1 \phi_1 \\ &=: \lambda c_0(\lambda) Q_1. \end{aligned}$$

Here $c_0(0) = a = 4i\pi/|\langle V, \phi_1 \rangle|^2 \neq 0$. Note that $c_0(\lambda) \in \hat{L}^1$ when

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x) \phi_1(x) V(y) \phi_1(y) \left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\hat{L}_\lambda^1} dx dy < \infty.$$

Since

$$\left\| \frac{e^{i\lambda|x-y|} - 1}{\lambda|x-y|} \right\|_{\hat{L}_\lambda^1} = 1,$$

it is enough to assume that $V \phi_1 \in L^1$, i.e., that $V \in L^{\frac{3}{2},1}$, in view of the fact that $\phi_1 \in \langle x \rangle^{-1} L^\infty$.

It follows that $T_{11}(\lambda)$ is invertible for $|\lambda| \ll 1$ in $Q_1 L^2$ and

$$T_{11}^{-1}(\lambda) = \lambda^{-1} c_0^{-1}(\lambda) Q_1 = \lambda^{-1} E(\lambda).$$

Here $\chi(\frac{\lambda}{\epsilon})c_0^{-1}(\lambda) \in \widehat{L}^1$ for sufficiently small ϵ .

Likewise, since $Q_2(V_2 \otimes V_1) = (V_2 \otimes V_1)Q_2 = 0$,

$$\begin{aligned} T_{12}(\lambda) &= Q_1(I + V_2 R_0((\lambda + i0)^2) V_1) Q_2 \\ &= \lambda^2 Q_1 V_2 \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda(4\pi)^{-1} 1 \otimes 1}{\lambda^2} V_1 Q_2 \\ &= \lambda^2 Q_1 e(\lambda) Q_2. \end{aligned}$$

Since by [Lemma 2.6](#)

$$M\left(\left(\frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda(4\pi)^{-1} 1 \otimes 1}{\lambda^2}\right)^\wedge\right) = \frac{|x - y|}{8\pi},$$

it follows that $e(\lambda) \in \widehat{L}^1$ if

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x) \phi_1(x) V(y) \phi_k(y) |x - y| < \infty,$$

that is, if $V \in L^1$.

Likewise we obtain $T_{21}(\lambda) = \lambda^2 Q_2 e(\lambda) Q_1$; hence, combining the previous results,

$$T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) = \lambda^3 Q_2 e(\lambda) Q_2.$$

Furthermore,

$$\begin{aligned} T_{22}(\lambda) &= Q_2(I + V_2 R_0((\lambda + i0)^2) V_1) Q_2 \\ &= \lambda^2 Q_2 V_2 \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1 Q_2 \\ &= -\lambda^2 \left(Q_2 V_2 \frac{|x - y|}{8\pi} V_1 Q_2 + \lambda Q_2 e(\lambda) Q_2 \right). \end{aligned}$$

Again by [Lemma 2.6](#), $e(\lambda) \in \widehat{L}^1$ if

$$\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} V(x) \phi_k(x) V(y) \phi_\ell(y) |x - y|^2 < \infty,$$

that is (taking into account that $\phi_k, \phi_\ell \lesssim \langle x \rangle^{-2}$), if $V \in L^1$.

Let P_0 be the L^2 orthogonal projection onto the set \mathcal{E} spanned by ϕ_2, \dots, ϕ_N . By relation (4.38) of [\[Yajima 2005\]](#),

$$\left(Q_2 V_2 \frac{|x - y|}{8\pi} V_1 Q_2 \right)^{-1} = -V_2 P_0 V_1.$$

Then

$$\begin{aligned} C_{22}(\lambda) &= T_{22}(\lambda) - T_{21}(\lambda) T_{11}^{-1}(\lambda) T_{12}(\lambda) \\ &= -\lambda^2 Q_2 V_2 \frac{|x - y|}{8\pi} V_1 Q_2 + \lambda^3 Q_2 e(\lambda) Q_2. \end{aligned}$$

Therefore,

$$C_{22}^{-1}(\lambda) = \lambda^{-2} V_2 P_0 V_1 + \lambda^{-1} Q_2 e(\lambda) Q_2.$$

Furthermore, we then obtain that

$$\begin{aligned} -T_{11}^{-1}(\lambda)T_{12}(\lambda)C_{22}^{-1}(\lambda) &= \lambda^{-1}Q_1e(\lambda)Q_1\lambda^2Q_1e(\lambda)Q_2\lambda^{-2}Q_2e(\lambda)Q_2 \\ &= \lambda^{-1}Q_1e(\lambda)Q_2. \end{aligned}$$

Likewise we obtain

$$\begin{aligned} -C_{22}^{-1}(\lambda)T_{21}(\lambda)T_{11}^{-1}(\lambda) &= \lambda^{-1}Q_2e(\lambda)Q_1, \\ T_{11}^{-1}(\lambda)T_{12}(\lambda)C_{22}^{-1}(\lambda)T_{21}(\lambda)T_{11}^{-1}(\lambda) &= Q_1e(\lambda)Q_1. \end{aligned}$$

By (2-33), we know that $(Q(I + \hat{T}(\lambda))Q)^{-1}$ is given in matrix form by

$$(Q(I + \hat{T}(\lambda))Q)^{-1} = \begin{pmatrix} \lambda^{-1}Q_1e(\lambda)Q_1 & \lambda^{-1}Q_1e(\lambda)Q_2 \\ \lambda^{-1}Q_2e(\lambda)Q_1 & \lambda^{-2}V_2P_0V_1 + \lambda^{-1}Q_2e(\lambda)Q_2 \\ & \lambda^{-1}Qe(\lambda)Q + \lambda^{-2}V_2P_0V_1 \end{pmatrix}, \tag{2-34}$$

where $\chi(\frac{\lambda}{\epsilon})e(\lambda) \in \hat{L}^1$ for sufficiently small ϵ .

We apply Lemma 2.10 again after writing $I + \hat{T}(\lambda)$ in matrix form with respect to the decomposition $L^2 = \bar{Q}L^2 + QL^2$, where $QL^2 = V_2\mathcal{M}$:

$$I + \hat{T}(\lambda) = \begin{pmatrix} \bar{Q}(I + \hat{T}(\lambda))\bar{Q} & \bar{Q}\hat{T}(\lambda)Q \\ Q\hat{T}(\lambda)\bar{Q} & Q(I + \hat{T}(\lambda))Q \end{pmatrix} := \begin{pmatrix} S_{00}(\lambda) & S_{01}(\lambda) \\ S_{10}(\lambda) & S_{11}(\lambda) \end{pmatrix}.$$

Next, as in the proof of Lemma 2.16, let $A(\lambda) = S_{00}^{-1}(\lambda)$. Then $\chi(\frac{\lambda}{\epsilon})A(\lambda) \in \mathcal{W}$ for sufficiently small ϵ .

We compute the inverse of $I + \hat{T}(\lambda)$ by finding each of its matrix elements:

$$(I + \hat{T}(\lambda))^{-1} = \begin{pmatrix} A + AS_{01}C^{-1}S_{10}A & AS_{01}C^{-1} \\ -C^{-1}S_{10}A & C^{-1} \end{pmatrix}. \tag{2-35}$$

Here

$$C(\lambda) = S_{11}(\lambda) - S_{10}(\lambda)A(\lambda)S_{01}(\lambda).$$

$S_{10}(\lambda)A(\lambda)S_{01}(\lambda) = Q\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q$ may be written as

$$\begin{pmatrix} Q_1\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_1 & Q_1\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 \\ Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_1 & Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 \end{pmatrix} = \begin{pmatrix} \lambda^2Q_1e(\lambda)Q_1 & \lambda^3Q_1e(\lambda)Q_2 \\ \lambda^3Q_2e(\lambda)Q_1 & \lambda^3Q_2e(\lambda)Q_2 \end{pmatrix}, \tag{2-36}$$

where $e(\lambda) \in \hat{L}^1$.

Indeed, consider, for example, $Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2$. It can be rewritten as

$$\begin{aligned} &Q_2\hat{T}(\lambda)A(\lambda)\hat{T}(\lambda)Q_2 \\ &= \lambda^3Q_2V_2\frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda 1 \otimes 1}{\lambda}V_1A(\lambda)V_2\frac{R_0((\lambda + i0)^2) - R_0(0)}{\lambda}V_1Q_2. \end{aligned} \tag{2-37}$$

Assuming that $V \in \langle x \rangle^{-2}L^{\frac{3}{2},1}$,

$$M\left(\left(V_2\frac{R_0((\lambda + i0)^2) - R_0(0)}{\lambda}V_1\right)^\wedge\right) = \frac{|V_2| \otimes |V_1|}{4\pi} \in \mathcal{B}(L^{\frac{3}{2},2}).$$

Likewise

$$M\left(\left(V_2 \frac{R_0((\lambda + i0)^2) - R_0(0) - i\lambda \frac{1 \otimes 1}{4\pi}}{\lambda^2} V_1\right)^\wedge\right) = |V_2| \frac{|x-y|}{8\pi} |V_1| \in \mathcal{B}(L^{\frac{3}{2},2}, L^{3,2}).$$

This implies that (2-37) $= \lambda^3 Q_2 e_0(\lambda) Q_2$ and $e_0(\lambda) \in \hat{L}^1$. In this manner, we prove (2-36).

By (2-34), we know that $S_{11}^{-1}(\lambda) = (Q\hat{T}(\lambda)Q)^{-1}$ is of the form

$$S_{11}^{-1}(\lambda) = \begin{pmatrix} \lambda^{-1} Q_1 e(\lambda) Q_1 & \lambda^{-1} Q_1 e(\lambda) Q_2 \\ \lambda^{-1} Q_2 e(\lambda) Q_1 & \lambda^{-2} Q_2 e(\lambda) Q_2 \end{pmatrix}.$$

Then, letting $N(\lambda) := S_{11}^{-1}(\lambda) S_{10}(\lambda) A(\lambda) S_{01}(\lambda)$, by (2-36),

$$\begin{aligned} N(\lambda) &:= S_{11}^{-1}(\lambda) S_{10}(\lambda) S_{00}^{-1}(\lambda) S_{01}(\lambda) \\ &= \begin{pmatrix} \lambda^{-1} Q_1 e(\lambda) Q_1 & \lambda^{-1} Q_1 e(\lambda) Q_2 \\ \lambda^{-1} Q_2 e(\lambda) Q_1 & \lambda^{-2} Q_2 e(\lambda) Q_2 \end{pmatrix} \begin{pmatrix} \lambda^2 Q_1 e(\lambda) Q_1 & \lambda^3 Q_1 e(\lambda) Q_2 \\ \lambda^3 Q_2 e(\lambda) Q_1 & \lambda^3 Q_2 e(\lambda) Q_2 \end{pmatrix} \\ &= \begin{pmatrix} \lambda Q_1 e(\lambda) Q_1 & \lambda^2 Q_1 e(\lambda) Q_2 \\ \lambda Q_2 e(\lambda) Q_1 & \lambda Q_2 e(\lambda) Q_2 \end{pmatrix}. \end{aligned}$$

Therefore $N(0) = 0$. This shows that $C(\lambda)$ is invertible for $\lambda \ll 1$:

$$C(\lambda) = S_{11}(\lambda) - S_{10}(\lambda) A(\lambda) S_{01}(\lambda) = S_{11}(\lambda)(I - N(\lambda)),$$

so

$$\begin{aligned} C^{-1}(\lambda) &= (I - N(\lambda))^{-1} S_{11}^{-1}(\lambda) \\ &= S_{11}^{-1}(\lambda) + (I - N(\lambda))^{-1} N(\lambda) S_{11}^{-1}(\lambda). \end{aligned} \tag{2-38}$$

A computation shows that

$$\begin{aligned} (I - N(\lambda))^{-1} N(\lambda) S_{11}^{-1}(\lambda) &= Qe(\lambda)Q \begin{pmatrix} \lambda Q_1 e(\lambda) Q_1 & \lambda^2 Q_1 e(\lambda) Q_2 \\ \lambda Q_2 e(\lambda) Q_1 & \lambda Q_2 e(\lambda) Q_2 \end{pmatrix} \begin{pmatrix} \lambda^{-1} Q_1 e(\lambda) Q_1 & \lambda^{-1} Q_1 e(\lambda) Q_2 \\ \lambda^{-1} Q_2 e(\lambda) Q_1 & \lambda^{-2} Q_2 e(\lambda) Q_2 \end{pmatrix} \\ &= \begin{pmatrix} Q_1 e(\lambda) Q_1 & Q_1 e(\lambda) Q_2 \\ Q_2 e(\lambda) Q_1 & \lambda^{-1} Q_2 e(\lambda) Q_2 \end{pmatrix}. \end{aligned}$$

By (2-38) and (2-34),

$$\begin{aligned} C^{-1}(\lambda) &= S_{11}^{-1}(\lambda) + \lambda^{-1} Qe(\lambda)Q \\ &= \lambda^{-2} V_2 P_0 V_1 + \lambda^{-1} Qe(\lambda)Q. \end{aligned}$$

Note that

$$\begin{aligned} S_{01}(\lambda) &= \bar{Q}\hat{T}(\lambda)Q = \bar{Q}(I + \hat{T}(\lambda))Q \\ &= \lambda \bar{Q} V_2 \frac{R_0((\lambda + i0)^2) - R_0(0)}{\lambda} V_1 Q = \lambda E_1(\lambda), \end{aligned}$$

where $E_1(\lambda)^\vee \in \mathcal{V}_{L^{3/2,2}}$ when $V \in \langle x \rangle^{-1} L^1$. Therefore,

$$A(\lambda) S_{01}(\lambda) C^{-1}(\lambda) = A(\lambda) \lambda E_1(\lambda) \lambda^{-2} Qe(\lambda)Q = \lambda^{-1} S(\lambda),$$

where $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$. Likewise $S_{10}(\lambda) = \lambda E_2(\lambda)$, where $E_2(\lambda)^\vee \in \mathcal{V}_{L^{3,2}}$. Then

$$C^{-1}(\lambda)S_{10}(\lambda)A(\lambda) = \lambda^{-1}S(\lambda),$$

where $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$.

Finally, for the last remaining matrix element $A + AS_{01}C^{-1}S_{10}A$ of (2-35), we use the fact that

$$AS_{01}C^{-1}S_{10}A = A(\lambda)\lambda E_1(\lambda)\lambda^{-2}Qe(\lambda)Q\lambda E_2(\lambda)A(\lambda) = S(\lambda),$$

where $S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$. Also recall that $A(\lambda) \in \mathcal{W}$.

We have thus analyzed all the terms in (2-35) and the conclusion follows. □

Recall that

$$R(t) := \frac{ae^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} \zeta_t(x) \otimes \zeta_t(y), \quad \zeta_t(x) := e^{i\frac{|x|^2}{4t}} \phi(x),$$

$$S(t) := \frac{e^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} \left(-iP_0V \frac{|x-y|^2}{24\pi} VP_0 + \mu_t(x) \frac{|x-y|}{8\pi} VP_0 + P_0V \frac{|x-y|}{8\pi} \mu_t(y) \right),$$

where

$$\mu_t(x) := \frac{i}{|x|} \int_0^1 (e^{i\frac{|x|^2}{4t}} - e^{i\frac{|\theta x|^2}{4t}}) d\theta.$$

Although it is not immediately obvious, it is also true that

$$\|S(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}. \tag{2-39}$$

Indeed, note that since $\langle \phi_k, V \rangle = 0$ for the eigenvectors ϕ_k , with $2 \leq k \leq N$ (recall that ϕ_1 is the resonance),

$$\mu_t(x)|x-y|VP_0 = \mu_t(x)(|x-y|-|x|)VP_0,$$

which is bounded in absolute value by

$$\sum_{k=2}^N |\mu_t(x)| \int_{\mathbb{R}^3} |y| |V(y)| |\phi_k(y)| dy \otimes |\phi_k(z)|.$$

By definition, $|\mu_t(x)| \lesssim |x|^{-1}$. This leads to (2-39), since $\phi_k \in \langle x \rangle^{-2} L^\infty$ and $V \in L^{\frac{3}{2},1}$.

We use Lemma 2.16 as the basis for the following decay estimate:

Proposition 2.18. *Let V satisfy $\langle x \rangle^4 V(x) \in L^{\frac{3}{2},1}$. Suppose that H is of exceptional type of the third kind. Then, for $1 \leq p < \frac{3}{2}$ and $u \in L^2 \cap L^p$,*

$$e^{-itH} P_c u = Z(t)u + R(t)u + S(t)u, \quad \|Z(t)u\|_{L^{p'}} \lesssim t^{-\frac{3}{2}(\frac{1}{p}-\frac{1}{p'})} \|u\|_{L^p}. \tag{2-40}$$

Here $\frac{1}{p} + \frac{1}{p'} = 1$. If in addition all the zero-energy eigenfunctions ϕ_k , with $2 \leq k \leq N$, are in L^1 , then we can take $S(t) = 0$.

Proof of Proposition 2.18. Write the dispersive component of the evolution as

$$e^{itH} P_c f = \frac{1}{i\pi} \int_{\mathbb{R}} e^{it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 \widehat{T}(\lambda)^{-1} V_2 R_0((\lambda+i0)^2)) f \lambda d\lambda.$$

We use the same method as in the proofs of Propositions 2.13 and 2.15. Consider a partition of unity subordinated to the neighborhoods of Lemmas 2.8 and 2.16. First, following Lemma 2.8, take a sufficiently large R such that

$$\left(1 - \chi\left(\frac{\lambda}{R}\right)\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}.$$

Then, again by Lemma 2.8, for every $\lambda_0 \in [-4R, 4R]$, there exists $\epsilon(\lambda_0) > 0$ such that

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon(\lambda_0)}\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$$

if $\lambda_0 \neq 0$ while the conclusion of Lemma 2.16 holds when $\lambda_0 = 0$.

Since $[-4R, 4R]$ is a compact set, there exists a finite covering

$$[-4R, 4R] \subset \bigcup_{k=1}^N (\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)).$$

Then we construct a finite partition of unity on \mathbb{R} by smooth functions $1 = \chi_0(\lambda) + \sum_{k=1}^N \chi_k(\lambda) + \chi_\infty(\lambda)$, where $\text{supp } \chi_\infty \subset \mathbb{R} \setminus (-2R, 2R)$, $\text{supp } \chi_0 \subset [-\epsilon(0), \epsilon(0)]$, and $\text{supp } \chi_k \subset [\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)]$.

By Lemma 2.8, for any $k \neq 0$, we have $\chi_k(\lambda)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$, so $(1 - \chi_0(\lambda))(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$. By Lemma 2.16, for $L \in \mathcal{W}$,

$$\chi_0(\lambda)(I + \widehat{T}(\lambda))^{-1} = L(\lambda) + \chi_0(\lambda) \left(\frac{V_2 P_0 V_1}{\lambda^2} + \frac{i V_2 P_0 V |x-y|^2 V P_0 V_1}{\lambda} - \frac{a}{\lambda} V_2 \phi \otimes V_1 \phi \right).$$

Let Z_1 be the contribution of all the regular terms in this decomposition, such as the free resolvent, $(1 - \chi_0(\lambda))(I + \widehat{T}(\lambda))^{-1}$, and $L(\lambda)$:

$$\begin{aligned} Z_1(t) &:= \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 L(\lambda) V_2 R_0((\lambda+i0)^2) \\ &\quad - (1 - \chi_0(\lambda)) R_0((\lambda+i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda+i0)^2)) \lambda d\lambda \\ &= \frac{1}{2\pi t} \int_{\mathbb{R}} e^{-it\lambda^2} \partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 L(\lambda) V_2 R_0((\lambda+i0)^2) \\ &\quad - (1 - \chi_0(\lambda)) R_0((\lambda+i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda+i0)^2)) d\lambda \\ &= \frac{C}{t^{\frac{3}{2}}} \int_{\mathbb{R}} e^{-i\frac{\rho^2}{4t}} (\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 L(\lambda) V_2 R_0((\lambda+i0)^2) \\ &\quad - (1 - \chi_0(\lambda)) R_0((\lambda+i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda+i0)^2)))^\vee(\rho) d\rho. \end{aligned}$$

The fact that $\|Z_1(t)u\|_{L^1} \lesssim |t|^{-\frac{3}{2}} \|u\|_{L^\infty}$ follows by knowing that

$$\begin{aligned} &(\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 L(\lambda) V_2 R_0((\lambda+i0)^2) \\ &\quad - (1 - \chi_0(\lambda)) R_0((\lambda+i0)^2) V_1 \widehat{T}(\lambda) V_2 R_0((\lambda+i0)^2)))^\vee \in \mathcal{V}_{L^1, L^\infty}. \end{aligned}$$

By smoothing estimates, it also follows that $Z_1(t)$ is L^2 -bounded; see the proof of [Proposition 2.13](#). By interpolation, we also obtain the estimate $\|Z_1(t)u\|_{L^{3,\infty}} \lesssim \|u\|_{L^{3/2,1}}$.

Let $Z_2(t)$ be the contribution of the term $a\lambda^{-1}\chi_0(\lambda)V_2\phi \otimes V_1\phi$:

$$Z_2(t) := \frac{a}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} \chi_0(\lambda) R_0((\lambda + i0)^2) V\phi \otimes V\phi R_0((\lambda + i0)^2) d\lambda.$$

By [Lemma 2.14](#),

$$\|(Z_2(t) - R(t))u\|_{L^\infty} \leq t^{-\frac{3}{2}} \|u\|_{L^1}, \quad \|Z_2(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}.$$

We are left with the terms

$$\lambda^{-2} R_0((\lambda + i0)^2) V P_0 V R_0((\lambda + i0)^2) \quad \text{and} \quad i\lambda^{-1} R_0((\lambda + i0)^2) V P_0 V \frac{|x - y|^2}{24\pi} V P_0 V R_0((\lambda + i0)^2).$$

Let their contributions be

$$X_2(t) := \frac{-1}{\pi} \int_{\mathbb{R}} e^{-it\lambda^2} R_0((\lambda + i0)^2) V P_0 V \frac{|x - y|^2}{24\pi} V P_0 V R_0((\lambda + i0)^2) d\lambda,$$

$$X_3(t) := \frac{-1}{i\pi} \lim_{\delta \rightarrow 0} \int_{|\lambda| > \delta} e^{-it\lambda^2} R_0((\lambda + i0)^2) V P_0 V R_0((\lambda + i0)^2) \lambda^{-1} d\lambda.$$

By [\[Yajima 2005, Lemma 4.12\]](#),

$$\begin{aligned} \|X_2(t)u\|_{L^{3,\infty}} &\lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}, \\ \left\| X_2(t)u + i \frac{e^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} P_0 V \frac{|x - y|^2}{24\pi} V P_0 \right\|_{L^\infty} &\lesssim t^{-\frac{3}{2}} \|u\|_{L^1}. \end{aligned} \tag{2-41}$$

This lemma has a proof similar to [Lemma 2.14](#). It requires, in addition, that $|\phi_j(x)| \lesssim |x|^{-2}$ for every eigenfunction $\phi_j \in \mathcal{E}$, with $2 \leq j \leq N$, which is guaranteed by [Lemma 2.3](#).

By [\[Yajima 2005, Lemma 4.14\]](#),

$$\begin{aligned} \|X_3(t)u\|_{L^{3,\infty}} &\lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}, \\ \left\| X_3(t)u - \frac{e^{-i\frac{3\pi}{4}}}{\sqrt{\pi t}} \left(\mu_t(x) \frac{|x - y|}{8\pi} V P_0 + P_0 V \frac{|x - y|}{8\pi} \mu_t(y) \right) \right\|_{L^\infty} &\lesssim t^{-\frac{3}{2}} \|u\|_{L^1}. \end{aligned} \tag{2-42}$$

The proof of [\[Yajima 2005, Lemma 4.14\]](#) depends on $\langle y \rangle^3 V(y)\phi(y)$ being integrable, which is also true here since $|\phi(y)| \lesssim \langle y \rangle^{-1}$ and $\langle y \rangle^2 V(y) \in \langle y \rangle^{-2} L^{\frac{3}{2},1} \subset L^1$.

Combining the two results (2-41) and (2-42) and knowing that $\|S(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}$ by (2-39), we obtain that

$$\|(X_2(t) + X_3(t) - S(t))u\|_{L^\infty} \lesssim t^{-\frac{3}{2}} \|u\|_{L^1}, \quad \|(X_2(t) + X_3(t) - S(t))u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}} \|u\|_{L^{3/2,1}}. \tag{2-43}$$

Recall that

$$e^{-itH} P_c = Z_1(t) + Z_2(t) + X_2(t) + X_3(t) = Z(t) + R(t) + S(t).$$

We obtain for $Z(t) = Z_1(t) + (Z_2(t) - R(t)) + (X_2(t) + X_3(t) - S(t))$ that

$$\|Z(t)u\|_{L^\infty} \lesssim t^{-\frac{3}{2}}\|u\|_{L^1}, \quad \|Z(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}.$$

Conclusion (2-40) follows by interpolation.

Finally, assume that all the eigenfunctions ϕ_k are in L^1 for $2 \leq k \leq N$ (recall that ϕ_1 is the resonance). Then, by Lemma 2.5, it follows that $\langle V\phi_k, y_\ell \rangle = \langle V\phi_k, y_\ell y_m \rangle = 0$ for all ℓ and m and all $2 \leq k \leq N$. As a consequence, we immediately see that

$$P_0 V|x-y|^2 V P_0 = P_0 V(|x|^2 + |y|^2) P_0 - 2 \sum_{k=1}^3 P_0 V x_k y_k V P_0 = 0.$$

Since $\langle \phi_k, V \rangle = 0$ and $\langle V\phi_k, y_\ell \rangle = 0$, we can also rewrite

$$\mu_t(x)|x-y|VP_0 = \mu_t(x) \left(|x-y| - |x| + \frac{xy}{|x|} \right) VP_0.$$

Then note that $|x| \left(|x-y| - |x| + \frac{xy}{|x|} \right) VP_0$ is bounded in absolute value by

$$\sum_{k=2}^N \int_{\mathbb{R}^3} |y|^2 |V(y)| |\phi_k(y)| dy \otimes |\phi_k(z)|,$$

which is bounded from L^1 to L^∞ since $\phi_k \in \langle x \rangle^{-2} L^\infty$ and $V \in \langle x \rangle^{-1} L^{\frac{3}{2},1}$. Having gained a power of decay in x , we use it by $|\mu_t(x)|x|^{-1}| \lesssim t^{-1}$. Therefore,

$$\|t^{-\frac{1}{2}}\mu_t(x)|x-y|VP_0u\|_{L^\infty} \lesssim t^{-\frac{3}{2}}\|u\|_{L^1}.$$

Consequently, when $\phi_k \in L^1$ for $2 \leq k \leq N$, we can remove $S(t)$ from (2-43). Hence we retrieve conclusion (2-40) without S , as claimed. \square

Proposition 2.19. *Assume that $V \in \langle x \rangle^{-2} L^{\frac{3}{2},1}$ and that $H = -\Delta + V$ is an exceptional Hamiltonian of the third kind. Then*

$$\|e^{-itH} P_c u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$$

and, for $\frac{3}{2} < p \leq 2$,

$$\|e^{-itH} P_c u\|_{L^{p'}} \lesssim t^{-\frac{3}{2} \left(\frac{1}{p} - \frac{1}{p'} \right)} \|u\|_{L^p}.$$

Here $\frac{1}{p} + \frac{1}{p'} = 1$.

The proof of this proposition parallels the proof of Proposition 2.15.

Proof of Proposition 2.19. Write the evolution as

$$e^{-itH} P_c f = \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2) V_1 \hat{T}(\lambda)^{-1} V_2 R_0((\lambda+i0)^2)) f \lambda d\lambda.$$

We consider a partition of unity subordinated to the neighborhoods of Lemmas 2.8 and 2.17. First, take a sufficiently large R such that

$$\left(1 - \chi\left(\frac{\lambda}{R}\right) \right) (I + \hat{T}(\lambda))^{-1} \in \mathcal{W}.$$

Then, for every $\lambda_0 \in [-4R, 4R]$, there exists $\epsilon(\lambda_0) > 0$ such that

$$\chi\left(\frac{\lambda - \lambda_0}{\epsilon(\lambda_0)}\right)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$$

if $\lambda_0 \neq 0$, while the conclusion of [Lemma 2.12](#) holds when $\lambda_0 = 0$.

Since $[-4R, 4R]$ is a compact set, there exists a finite covering

$$[-4R, 4R] \subset \bigcup_{k=1}^N (\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)).$$

Then we construct a finite partition of unity on \mathbb{R} by smooth functions $1 = \chi_0(\lambda) + \sum_{k=1}^N \chi_k(\lambda) + \chi_\infty(\lambda)$, where $\text{supp } \chi_\infty \subset \mathbb{R} \setminus (-2R, 2R)$, $\text{supp } \chi_0 \subset [-\epsilon(0), \epsilon(0)]$, and $\text{supp } \chi_k \subset [\lambda_k - \epsilon(\lambda_k), \lambda_k + \epsilon(\lambda_k)]$.

By [Lemma 2.8](#), for any $k \neq 0$, we have $\chi_k(\lambda)(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$, so $(1 - \chi_0(\lambda))(I + \widehat{T}(\lambda))^{-1} \in \mathcal{W}$. By [Lemma 2.17](#),

$$\chi_0(\lambda)(I + \widehat{T}(\lambda))^{-1} = L(\lambda) + \lambda^{-1}S(\lambda) + \lambda^{-2}V_2P_0V_1,$$

where $L \in \mathcal{W}$ and $S^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}}$.

Let Z_1 be given by the sum of all the regular terms of the decomposition:

$$\begin{aligned} Z_1(t) &:= \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2))\lambda \, d\lambda \\ &= \frac{1}{2\pi t} \int_{\mathbb{R}} e^{-it\lambda^2} \partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2)) \, d\lambda \\ &= \frac{C}{t^{\frac{3}{2}}} \int_{\mathbb{R}} e^{-i\frac{\rho^2}{4t}} (\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee(\rho) \, d\rho. \end{aligned}$$

The fact that $\|Z_1(t)u\|_{L^\infty} \lesssim |t|^{-\frac{3}{2}}\|u\|_{L^1}$ follows by knowing that

$$\begin{aligned} &(\partial_\lambda (R_0((\lambda+i0)^2) - R_0((\lambda+i0)^2)V_1L(\lambda)V_2R_0((\lambda+i0)^2) \\ &\quad - (1-\chi_0(\lambda))R_0((\lambda+i0)^2)V_1\widehat{T}(\lambda)V_2R_0((\lambda+i0)^2)))^\vee \in \mathcal{V}_{L^1, L^\infty}. \end{aligned}$$

Using smoothing estimates, it immediately follows that $Z_1(t)$ is L^2 -bounded; see the proof of [Proposition 2.13](#). Interpolating, we obtain that $\|Z_1(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3,1}}$.

Let Z_2 be the following singular term in the decomposition of [Lemma 2.17](#):

$$\begin{aligned} Z_2(t) &:= \frac{1}{i\pi} \int_{\mathbb{R}} e^{-it\lambda^2} R_0((\lambda+i0)^2)V_1S(\lambda)V_2R_0((\lambda+i0)^2) \, d\lambda \\ &= \frac{C}{t^{\frac{1}{2}}} \int_{\mathbb{R}} e^{-i\frac{\rho^2}{4t}} (R_0((\lambda+i0)^2)V_1S(\lambda)V_2R_0((\lambda+i0)^2))^\vee(\rho) \, d\rho. \end{aligned}$$

Note that

$$(R_0((\lambda + i0)^2)V_1)^\vee \in \mathcal{V}_{L^{3/2,2}, L^{3,\infty}}, \quad S(\lambda)^\vee \in \mathcal{V}_{L^{3,2}, L^{3/2,2}} \quad \text{and} \quad (V_2 R_0((\lambda + i0)^2))^\vee \in \mathcal{V}_{L^{3/2,1}, L^{3,2}}.$$

Thus

$$R_0((\lambda + i0)^2)V_1(\lambda S(\lambda))V_2 R_0((\lambda + i0)^2) \in \mathcal{V}_{L^{3/2,1}, L^{3,\infty}}.$$

This immediately implies that $\|Z_2(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$.

We are left with the contribution of the term $\lambda^{-2}V_2 P_0 V_1$. This is the same as the term X_3 from the proof of Proposition 2.18. By (2-42), we have $\|X_3(t)u\|_{L^{3,\infty}} \lesssim t^{-\frac{1}{2}}\|u\|_{L^{3/2,1}}$.

Putting the three estimates for Z_1 , Z_2 , and X_3 together, we obtain that $\|e^{-itH} P_c u\|_{L^{3,\infty}} \lesssim \|u\|_{L^{3/2,1}}$. Interpolating with the obvious L^2 bound $\|e^{-itH} P_c u\|_{L^2} \lesssim \|u\|_{L^2}$, we obtain the stated conclusion. \square

Acknowledgments

This paper is indebted to Professor Kenji Yajima's article [2005], from which it borrows several lemmas, as well as the general plan of the proof. I would like to thank Professor Avy Soffer for referring me to that article and Professor Wilhelm Schlag for introducing me to this problem. I would also like to thank the anonymous referee for the many useful comments.

The author was partially supported by the NSF grant DMS-1128155 and by an AMS-Simons Foundation travel grant.

References

- [Beceanu 2011] M. Beceanu, "New estimates for a time-dependent Schrödinger equation", *Duke Math. J.* **159**:3 (2011), 417–477. [MR 2831875](#) [Zbl 1229.35224](#)
- [Beceanu and Goldberg 2012] M. Beceanu and M. Goldberg, "Schrödinger dispersive estimates for a scaling-critical class of potentials", *Comm. Math. Phys.* **314**:2 (2012), 471–481. [MR 2958960](#) [Zbl 1250.35047](#)
- [Beceanu and Goldberg 2014] M. Beceanu and M. Goldberg, "Strichartz estimates and maximal operators for the wave equation in \mathbb{R}^3 ", *J. Funct. Anal.* **266**:3 (2014), 1476–1510. [MR 3146823](#) [Zbl 1292.35063](#)
- [Bergh and Löfström 1976] J. Bergh and J. Löfström, *Interpolation spaces: An introduction*, Grundlehren der Mathematischen Wissenschaften **223**, Springer, Berlin, 1976. [MR 0482275](#) [Zbl 0344.46071](#)
- [Cardoso et al. 2009] F. Cardoso, C. Cuevas, and G. Vodev, "Dispersive estimates for the Schrödinger equation in dimensions four and five", *Asymptot. Anal.* **62**:3-4 (2009), 125–145. [MR 2521760](#) [Zbl 1163.35482](#)
- [Egorova et al. 2014] I. Egorova, E. Kopylova, V. Marchenko, and G. Teschl, "Dispersion estimates for one-dimensional Schrödinger and Klein-Gordon equations revisited", preprint, 2014. To appear in *Russian Math. Surveys*. [arXiv 1411.0021](#)
- [Erdoğan and Green 2010] M. B. Erdoğan and W. R. Green, "Dispersive estimates for the Schrödinger equation for $C^{\frac{n-3}{2}}$ potentials in odd dimensions", *Int. Math. Res. Not.* **2010**:13 (2010), 2532–2565. [MR 2669658](#) [Zbl 1200.35036](#)
- [Erdoğan and Green 2013a] M. B. Erdoğan and W. R. Green, "Dispersive estimates for matrix Schrödinger operators in dimension two", *Discrete Contin. Dyn. Syst.* **33**:10 (2013), 4473–4495. [MR 3049087](#) [Zbl 1277.35294](#)
- [Erdoğan and Green 2013b] M. B. Erdoğan and W. R. Green, "Dispersive estimates for Schrödinger operators in dimension two with obstructions at zero energy", *Trans. Amer. Math. Soc.* **365**:12 (2013), 6403–6440. [MR 3105757](#) [Zbl 1282.35143](#)
- [Erdoğan and Green 2013c] M. B. Erdoğan and W. R. Green, "A weighted dispersive estimate for Schrödinger operators in dimension two", *Comm. Math. Phys.* **319**:3 (2013), 791–811. [MR 3040376](#) [Zbl 1272.35053](#)
- [Erdoğan and Schlag 2004] M. B. Erdoğan and W. Schlag, "Dispersive estimates for Schrödinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three, I", *Dyn. Partial Differ. Equ.* **1**:4 (2004), 359–379. [MR 2127577](#) [Zbl 1080.35102](#)

- [Erdoğan and Schlag 2006] M. B. Erdoğan and W. Schlag, “Dispersive estimates for Schrödinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three, II”, *J. Anal. Math.* **99** (2006), 199–248. [MR 2279551](#) [Zbl 1146.35324](#)
- [Erdoğan et al. 2014] M. B. Erdoğan, M. Goldberg, and W. R. Green, “Dispersive estimates for four dimensional Schrödinger and wave equations with obstructions at zero energy”, *Comm. Partial Differential Equations* **39**:10 (2014), 1936–1964. [MR 3250981](#) [Zbl 1325.35017](#)
- [Goldberg 2006] M. Goldberg, “Dispersive bounds for the three-dimensional Schrödinger equation with almost critical potentials”, *Geom. Funct. Anal.* **16**:3 (2006), 517–536. [MR 2238943](#) [Zbl 1158.35408](#)
- [Goldberg 2007] M. Goldberg, “Transport in the one-dimensional Schrödinger equation”, *Proc. Amer. Math. Soc.* **135**:10 (2007), 3171–3179. [MR 2322747](#) [Zbl 1123.35046](#)
- [Goldberg 2010] M. Goldberg, “A dispersive bound for three-dimensional Schrödinger operators with zero energy eigenvalues”, *Comm. Partial Differential Equations* **35**:9 (2010), 1610–1634. [MR 2754057](#) [Zbl 1223.35265](#)
- [Goldberg and Green 2014] M. Goldberg and W. R. Green, “Dispersive estimates for higher dimensional Schrödinger operators with threshold eigenvalues, II: The even dimensional case”, preprint, 2014. To appear in *J. Spectr. Theory*. [arXiv 1409.6328](#)
- [Goldberg and Green 2015] M. Goldberg and W. R. Green, “Dispersive estimates for higher dimensional Schrödinger operators with threshold eigenvalues, I: The odd dimensional case”, *J. Funct. Anal.* **269**:3 (2015), 633–682. [MR 3350725](#) [Zbl 1317.35216](#)
- [Goldberg and Schlag 2004a] M. Goldberg and W. Schlag, “Dispersive estimates for Schrödinger operators in dimensions one and three”, *Comm. Math. Phys.* **251**:1 (2004), 157–178. [MR 2096737](#) [Zbl 1086.81077](#)
- [Goldberg and Schlag 2004b] M. Goldberg and W. Schlag, “A limiting absorption principle for the three-dimensional Schrödinger equation with L^p potentials”, *Int. Math. Res. Not.* **2004**:75 (2004), 4049–4071. [MR 2112327](#) [Zbl 1069.35063](#)
- [Green 2012] W. R. Green, “Dispersive estimates for matrix and scalar Schrödinger operators in dimension five”, *Illinois J. Math.* **56**:2 (2012), 307–341. [MR 3161326](#) [Zbl 06233916](#)
- [Ionescu and Jerison 2003] A. D. Ionescu and D. Jerison, “On the absence of positive eigenvalues of Schrödinger operators with rough potentials”, *Geom. Funct. Anal.* **13**:5 (2003), 1029–1081. [MR 2024415](#) [Zbl 1055.35098](#)
- [Jensen and Kato 1979] A. Jensen and T. Kato, “Spectral properties of Schrödinger operators and time-decay of the wave functions”, *Duke Math. J.* **46**:3 (1979), 583–611. [MR 544248](#) [Zbl 0448.35080](#)
- [Jensen and Nenciu 2001] A. Jensen and G. Nenciu, “A unified approach to resolvent expansions at thresholds”, *Rev. Math. Phys.* **13**:6 (2001), 717–754. [MR 1841744](#) [Zbl 1029.81067](#)
- [Journé et al. 1991] J.-L. Journé, A. Soffer, and C. D. Sogge, “Decay estimates for Schrödinger operators”, *Comm. Pure Appl. Math.* **44**:5 (1991), 573–604. [MR 1105875](#) [Zbl 0743.35008](#)
- [Murata 1982] M. Murata, “Asymptotic expansions in time for solutions of Schrödinger-type equations”, *J. Funct. Anal.* **49**:1 (1982), 10–56. [MR 680855](#) [Zbl 0499.35019](#)
- [Rauch 1978] J. Rauch, “Local decay of scattering solutions to Schrödinger’s equation”, *Comm. Math. Phys.* **61**:2 (1978), 149–168. [MR 0495958](#) [Zbl 0381.35023](#)
- [Schlag 2005] W. Schlag, “Dispersive estimates for Schrödinger operators in dimension two”, *Comm. Math. Phys.* **257**:1 (2005), 87–117. [MR 2163570](#) [Zbl 1134.35321](#)
- [Simon 1982] B. Simon, “Schrödinger semigroups”, *Bull. Amer. Math. Soc. (N.S.)* **7**:3 (1982), 447–526. [MR 670130](#) [Zbl 0524.35002](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. [MR 1232192](#) [Zbl 0821.42001](#)
- [Yajima 2005] K. Yajima, “Dispersive estimates for Schrödinger equations with threshold resonance and eigenvalue”, *Comm. Math. Phys.* **259**:2 (2005), 475–509. [MR 2172692](#) [Zbl 1079.81021](#)

Received 2 Mar 2015. Revised 17 Dec 2015. Accepted 26 Feb 2016.

MARIUS BECEANU: mbeceanu@ias.edu

Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
László Lempert	Purdue University, USA lempert@math.purdue.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 4 2016

Peierls substitution for magnetic Bloch bands	773
SILVIA FREUND and STEFAN TEUFEL	
Dispersive estimates in \mathbb{R}^3 with threshold eigenstates and resonances	813
MARIUS BECEANU	
Interior nodal sets of Steklov eigenfunctions on surfaces	859
JIUYI ZHU	
Some counterexamples to Sobolev regularity for degenerate Monge–Ampère equations	881
CONNOR MOONEY	
Mean ergodic theorem for amenable discrete quantum groups and a Wiener-type theorem for compact metrizable groups	893
HUICHI HUANG	
Resonance free regions for nontrapping manifolds with cusps	907
KIRIL DATCHEV	
Characterizing regularity of domains via the Riesz transforms on their boundaries	955
DORINA MITREA, MARIUS MITREA and JOAN VERDERA	