

ANALYSIS & PDE

Volume 9

No. 8

2016

TERENCE TAO

**FINITE-TIME BLOWUP FOR
A SUPERCRITICAL DEFOCUSING NONLINEAR WAVE SYSTEM**



FINITE-TIME BLOWUP FOR A SUPERCRITICAL DEFOCUSING NONLINEAR WAVE SYSTEM

TERENCE TAO

We consider the global regularity problem for defocusing nonlinear wave systems

$$\square u = (\nabla_{\mathbb{R}^m} F)(u)$$

on Minkowski spacetime \mathbb{R}^{1+d} with d'Alembertian $\square := -\partial_t^2 + \sum_{i=1}^d \partial_{x_i}^2$, where the field $u : \mathbb{R}^{1+d} \rightarrow \mathbb{R}^m$ is vector-valued, and $F : \mathbb{R}^m \rightarrow \mathbb{R}$ is a smooth potential which is positive and homogeneous of order $p + 1$ outside of the unit ball for some $p > 1$. This generalises the scalar defocusing nonlinear wave (NLW) equation, in which $m = 1$ and $F(v) = 1/(p + 1)|v|^{p+1}$. It is well known that in the energy-subcritical and energy-critical cases when $d \leq 2$ or $d \geq 3$ and $p \leq 1 + 4/(d - 2)$, one has global existence of smooth solutions from arbitrary smooth initial data $u(0), \partial_t u(0)$, at least for dimensions $d \leq 7$. We study the supercritical case where $d = 3$ and $p > 5$. We show that in this case, there exists a smooth potential F for some sufficiently large m (in fact we can take $m = 40$), positive and homogeneous of order $p + 1$ outside of the unit ball, and a smooth choice of initial data $u(0), \partial_t u(0)$ for which the solution develops a finite-time singularity. In fact the solution is discretely self-similar in a backwards light cone. The basic strategy is to first select the mass and energy densities of u , then u itself, and then finally design the potential F in order to solve the required equation. The Nash embedding theorem is used in the second step, explaining the need to take m relatively large.

1. Introduction

Let \mathbb{R}^m be a Euclidean space, with the usual Euclidean norm $v \mapsto \|v\|_{\mathbb{R}^m}$ and Euclidean inner product $v, w \mapsto \langle v, w \rangle_{\mathbb{R}^m}$. A function $F : \mathbb{R}^m \rightarrow \mathbb{R}$ is said to be *homogeneous of order α* for some real α if we have

$$F(\lambda v) = \lambda^\alpha F(v) \tag{1-1}$$

for all $\lambda > 0$ and $v \in \mathbb{R}^m$. In particular, differentiating this at $\lambda = 1$ we obtain *Euler's identity*

$$\langle v, (\nabla_{\mathbb{R}^m} F)(v) \rangle_{\mathbb{R}^m} = \alpha F(v), \tag{1-2}$$

where $\nabla_{\mathbb{R}^m}$ denotes the gradient in \mathbb{R}^m , assuming of course that the gradient $\nabla_{\mathbb{R}^m} F$ of F exists at v . When α is not an integer, it is not possible for such homogeneous functions to be smooth at the origin unless they are identically zero (this can be seen by performing a Taylor expansion of F around the origin). To avoid this technical issue, we also introduce the notion of F being *homogeneous of order α outside of the unit ball*, by which we mean that (1-1) holds for $\lambda \geq 1$ and $v \in \mathbb{R}^m$ with $\|v\|_{\mathbb{R}^m} \geq 1$.

MSC2010: primary 35Q30, 35L71; secondary 35L67.

Keywords: nonlinear wave equation, Nash embedding theorem.

Define a *potential* to be a function $F : \mathbb{R}^m \rightarrow \mathbb{R}$ that is smooth away from the origin; if F is also smooth at the origin, we call it a *smooth potential*. We say that the potential is *defocusing* if F is positive away from the origin, and *focusing* if F is negative away from the origin. In this paper we consider nonlinear wave systems of the form

$$\square u = (\nabla_{\mathbb{R}^m} F)(u), \tag{1-3}$$

where the unknown field $u : \mathbb{R}^{1+d} \rightarrow \mathbb{R}^m$ is assumed to be smooth, $\square = \partial^\alpha \partial_\alpha = -\partial_t^2 + \sum_{i=1}^d \partial_{x_i}^2$ is the d’Alembertian operator on Minkowski spacetime

$$\mathbb{R}^{1+d} := \{(t, x_1, \dots, x_d) : t, x_1, \dots, x_d \in \mathbb{R}\} = \{(t, x) : t \in \mathbb{R}, x \in \mathbb{R}^d\}$$

with the usual Minkowski metric

$$\eta_{\alpha\beta} x^\alpha x^\beta = -t^2 + x_1^2 + \dots + x_d^2$$

and the usual Einstein summation, raising, and lowering conventions, $m, d \geq 1$ are integers, and $F : \mathbb{R}^m \rightarrow \mathbb{R}$ is a smooth potential. This is a Lagrangian field equation, in the sense that (1-3) is (formally, at least) the Euler–Lagrange equations for the Lagrangian

$$\int_{\mathbb{R}^{1+d}} \frac{1}{2} \langle \partial^\alpha u, \partial_\alpha u \rangle_{\mathbb{R}^m} + F(u) \, d\eta.$$

We will restrict attention to potentials F which are homogeneous outside of the unit ball of order $p + 1$ for some exponent $p > 1$. The well-studied *nonlinear wave equation* (NLW) corresponds to the case when $m = 1$ and $F(v) = |v|^{p+1}/(p + 1)$ (for the defocusing NLW) or $F(v) = -|v|^{p+1}/(p + 1)$ (for the focusing NLW), with the caveat that one needs to restrict p to be an odd integer if one wants these potentials to be smooth. Later in the paper we will restrict attention to the physical case $d = 3$, basically to take advantage of a form of the sharp Huygens’ principle.

The natural initial value problem to study here is the Cauchy initial value problem, in which one specifies a smooth initial position $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}^m$ and initial velocity $u_1 : \mathbb{R}^d \rightarrow \mathbb{R}^m$, and asks for a smooth solution u to (1-3) with $u(0, x) = u_0(x)$ and $\partial_t u(0, x) = u_1(x)$. Standard energy methods (see, e.g., [Shatah and Struwe 1998]) show that for any choice of smooth initial data $u_0, u_1 : \mathbb{R}^d \rightarrow \mathbb{R}^m$, one can construct a solution u to (1-3) in an open neighbourhood Ω in \mathbb{R}^{1+d} of the initial time slice $\{(0, x) : x \in \mathbb{R}^d\}$ with this initial data. Furthermore, either such a solution can be extended to be globally defined in \mathbb{R}^{1+d} , or else there is a solution u defined on some open neighbourhood Ω of $\{(0, x) : x \in \mathbb{R}^d\}$ that “blows up” in the sense that it cannot be smoothly continued to some boundary point (t_*, x_*) of Ω . The *global regularity problem* for a given choice of potential F asks if the latter situation does not occur, that is to say that for every choice of smooth initial data there is a smooth global solution. Note that as the equation (1-3) enjoys finite speed of propagation, there is no need to specify any decay hypotheses on the initial data as this will not affect the answer to the global regularity problem.

For focusing potentials F , there are well-known blowup examples that show that global regularity fails. For instance, if $m = 1$ and $F : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$F(v) := -\frac{2}{(p-1)^2} |v|^{p+1} \tag{1-4}$$

for all $|v| \geq 1$ (and extended arbitrarily in some smooth fashion to the region $|v| < 1$ while remaining negative away from the origin), then F is a focusing potential that is homogeneous of order $p + 1$ outside of the unit ball, and the function $u : \{(t, x) \in \mathbb{R}^{1+d} : 0 < t \leq 1\} \rightarrow \mathbb{R}$ defined by

$$u(t, x) := t^{-\frac{2}{p-1}} \quad (1-5)$$

solves (1-3) but blows up at the boundary $t = 0$; applying the time reversal symmetry $(t, x) \mapsto (1 - t, x)$, we obtain a counterexample to global regularity for this choice of F . We will thus henceforth restrict attention to defocusing potentials F , which excludes ODE-type blowup examples (1-5) in which $u(t, x)$ depends only on t .

The *energy* (or *Hamiltonian*)

$$E[u(t)] := \int_{\mathbb{R}^d} \frac{1}{2} \|\partial_t u(t, x)\|_{\mathbb{R}^m}^2 + \frac{1}{2} \|\nabla_x u(t, x)\|_{\mathbb{R}^m \otimes \mathbb{R}^d}^2 + F(u(t, x)) \, dx \quad (1-6)$$

is (formally, at least) conserved by the flow (1-3). A dimensional analysis of this quantity then naturally splits the range of parameters (d, p) into three cases:

- The *energy-subcritical* case when $d \leq 2$, or when $d \geq 3$ and $p < 1 + \frac{4}{d-2}$.
- The *energy-critical* case when $d \geq 3$ and $p = 1 + \frac{4}{d-2}$.
- The *energy-supercritical* case when $d \geq 3$ and $p > 1 + \frac{4}{d-2}$.

In the energy-subcritical and energy-critical cases one has global regularity for any defocusing NLW system, at least when $d \leq 7$; see¹ [Jörgens 1961] for the subcritical case, and [Grillakis 1990; 1992; Struwe 1988; Shatah and Struwe 1998] for the critical case. These results were also extended to the logarithmically supercritical case (in which the potential F grows faster than the energy-critical potential by a logarithmic factor) in [Tao 2007; Roy 2009]. A major ingredient in the proof of global regularity in these cases is the conservation of the energy (1-6), which is nonnegative in the defocusing case. In the energy-critical (and logarithmically supercritical) case, one also takes advantage of Morawetz inequalities such as

$$\int_0^T \int_{\mathbb{R}^d} \frac{F(u(t, x))}{|x|} \, dx \, dt \leq CE[u(0)] \quad (1-7)$$

for any time interval $[0, T]$ on which the solution exists. These bounds can be deduced from the properties of the stress-energy tensor

$$T_{\alpha\beta} := \langle \partial_\alpha u, \partial_\beta u \rangle - \frac{1}{2} \eta_{\alpha\beta} (\langle \partial^\gamma u, \partial_\gamma u \rangle_{\mathbb{R}^m} + F(u))$$

and in particular in the divergence-free nature $\partial^\beta T_{\alpha\beta} = 0$ of this tensor.

It thus remains to address the energy-supercritical case for defocusing smooth potentials F . In this case it is known that the Cauchy problem is ill-posed in various technical senses at low regularities [Lebeau 2001; 2005; Christ et al. 2003; Brenner and Kumlin 2000; Burq et al. 2007; Ibrahim et al. 2011],

¹Several of these references restrict attention to the scalar NLW or to three spatial dimensions, but the arguments extend without difficulty to the energy-critical NLW systems considered here in the range $3 \leq d \leq 7$. There are technical difficulties establishing global regularity in extremely high dimension, even when the potential F and all of its derivatives are bounded; see, e.g., [Brenner and von Wahl 1981].

despite the existence of global weak solutions [Segal 1963; Strauss 1989], as well as global smooth solutions from sufficiently small initial data [Lindblad and Sogge 1996] (assuming that F vanishes to sufficiently high order at the origin); see also [Zheng 1991] for a partial regularity result. However, to the author’s knowledge, finite-time blowup of smooth solutions has not actually been demonstrated for such equations. The main result of this paper is to establish such a finite-time blowup for at least some choices of defocusing potential F and parameters d, p, m :

Theorem 1.1 (finite-time blowup). *Let $d = 3$, let $p > 1 + \frac{4}{d-2}$, and let*

$$m \geq 2 \max\left(\frac{(d+1)(d+6)}{2}, \frac{(d+1)(d+4)}{2} + 5\right) + 2$$

be an integer. Then there exists a defocusing smooth potential $F : \mathbb{R}^m \rightarrow \mathbb{R}$ that is homogeneous of order $p + 1$ outside of the unit ball, and a smooth choice of initial data $u_0, u_1 : \mathbb{R}^d \rightarrow \mathbb{R}^m$ such that there is no global smooth solution $u : \mathbb{R}^{1+d} \rightarrow \mathbb{R}^m$ to the nonlinear wave system (1-3) with initial data $u(0) = u_0, \partial_t u(0) = u_1$.

Of course, since d is set equal to 3, the conditions on p and m reduce to $p > 5$ and $m \geq 40$ respectively. However, our restriction to the $d = 3$ case is largely for technical reasons (basically in order to exploit the strong Huygens principle), and we believe the results should extend to higher values of d , with the indicated constraints on d and p , though we will not pursue this matter here. The rather large value of m is due to our use of the Nash embedding theorem (!) at one stage of the argument. It would of course be greatly desirable to lower the number m of degrees of freedom down to 1, in order to establish blowup for the scalar defocusing supercritical NLW, but our methods crucially need a large value of m in order to ensure that a certain map from a $(1+d)$ -dimensional space into the sphere S^{m-1} is embedded, which is where the Nash embedding theorem comes in. Nevertheless, even though Theorem 1.1 does not directly show that the scalar defocusing supercritical NLW exhibits finite-time blowup, it does demonstrate a significant *barrier* to any attempt to prove global regularity for this equation, as such an attempt must necessarily use some special property of the scalar equation that is not shared by the more general system (1-3).

We briefly discuss the methods used to prove Theorem 1.1. The singularity constructed is a discretely self-similar blowup in a backwards light cone; see the reduction to Theorem 2.1 below. In particular, the blowup is “locally of type II” in the sense that scale-invariant norms inside the light cone stay bounded, but not “globally of type II”, as a significant amount of energy (as measured using scale-invariant norms) radiates out of the backwards light cone at all scales. This is compatible with the results in [Kenig and Merle 2008; Killip and Visan 2011a; 2011b], which rule out “global” type II blowup, but not “local” type II blowup. It would be natural to seek a *continuously* self-similar smooth blowup solution, but it turns out² that these are ruled out; see Proposition 2.2 below. Hence we will not restrict attention to

²On the other hand, it is possible to use perturbative methods to create *rough* solutions to (1-3) that are continuously self-similar: see [Planchon 2000; Ribaud and Youssfi 2002]. However, these methods do not seem to be adaptable to generate smooth solutions, and indeed Proposition 2.2 suggests that there are strong obstacles in trying to create such an adaptation. The negative result here also stands in contrast to the situation of high-dimensional wave maps into negatively curved targets, where ODE methods were used in [Cazenave et al. 1998] to construct continuously self-similar blowup examples in seven and higher spatial dimensions.

continuously self-similar solutions. It also turns out to be convenient not to initially restrict attention to spherically symmetric solutions, although we will eventually do so later in the argument.

Traditionally, one thinks of the potential F as being prescribed in advance, and the field u as the unknown to be solved for. However, as we have the freedom to *select* F in Theorem 1.1, it turns out to be more convenient to prescribe u first, and only then design an F for which the equation (1-3) is obeyed. This turns out to be possible as long as the map

$$\theta : (t, x) \mapsto \frac{u(t, x)}{\|u(t, x)\|_{\mathbb{R}^m}}$$

has certain nondegeneracy properties, and if the stress-energy tensor $T_{\alpha\beta}$ (which can be defined purely in terms of u) is divergence-free; see the reduction to Theorem 3.2 below. The stress-energy tensor $T_{\alpha\beta}$ (or more precisely, some related fields which we call the *mass density* M and the *energy tensor* $E_{\alpha\beta}$) can be viewed as prescribing the metric geometry of the map θ , and the Nash embedding theorem can then be used to locate a choice of θ with the desired nondegeneracy properties and the prescribed metric, so long as the fields M and $E_{\alpha\beta}$ obey a number of conditions (one of which relates to the divergence-free nature of the stress-energy tensor, and another to the positive definiteness of the Gram matrix of u). This reduces the problem to a certain “semidefinite program” (see Theorem 4.1), in which one now only needs to specify the fields M and $E_{\alpha\beta}$, rather than the original field u or the potential F .

It is at this point (after some additional technical reductions in which certain fields are allowed to degenerate to zero) that it finally becomes convenient to make symmetry reductions, working with fields M , $E_{\alpha\beta}$ that are both continuously self-similar and spherically symmetric, and assuming that there are no angular components to the energy tensor. In three spatial dimensions, this reduces the divergence-free nature of the stress-energy tensor to a single transport equation for the null energy e_+ (which, in terms of the original field u , is given in polar coordinates by $e_+ = \frac{1}{2}\|(\partial_t + \partial_r)(ru)\|_H^2$), in terms of a certain “potential energy density” V (which, in terms of the original data u and F , is given by $V = rF(u)$); see Theorem 5.4 for a precise statement. The strategy is then to solve for these fields e_+ , V first, and then choose all the remaining unknown fields in such a way that the remaining requirements of the semidefinite program are satisfied. This turns out to be possible if the fields e_+ , V are chosen to concentrate close to the boundary of the light cone.

2. Reduction to discretely self-similar solution

We begin the proof of Theorem 1.1.

We first observe that from finite speed of propagation and the symmetries of the equation, Theorem 1.1 follows from the claim below, in which the solution is restricted to a truncated light cone and is discretely self-similar and the potential is now homogeneous everywhere (not just outside of the unit ball), but no longer required to be smooth. This reduction does not use any of the hypotheses on m , d , p .

Theorem 2.1 (first reduction). *Let $d = 3$, let $p > 1 + \frac{4}{d-2}$, and let*

$$m \geq 2 \max\left(\frac{(d+1)(d+6)}{2}, \frac{(d+1)(d+4)}{2} + 5\right) + 2$$

be an integer. Then there exists a defocusing potential $F : \mathbb{R}^m \rightarrow \mathbb{R}$ which is homogeneous of order $p + 1$ and a smooth function $u : \Gamma_d \rightarrow \mathbb{R}^m \setminus \{0\}$ on the light cone $\Gamma_d := \{(t, x) \in \mathbb{R}^{1+d} : t > 0; |x| \leq t\}$ that solves (1-3) on its domain and is nowhere vanishing, and also discretely self-similar in the sense that there exists $S > 0$ such that

$$u(e^S t, e^S x) = e^{-\frac{2}{p-1}S} u(t, x) \tag{2-1}$$

for all $(t, x) \in \Gamma_d$.

A key point here is that u is smooth all the way up to the boundary of the light cone Γ_d , rather than merely being smooth in the interior. The exponent $-\frac{2}{p-1}$ is mandated by dimensional analysis considerations. It would be natural to consider solutions that are continuously self-similar in the sense that (2-1) holds for *all* $S \in \mathbb{R}$, but as we shall shortly see, it will not be possible to generate such solutions in the three-dimensional defocusing setting.

Let us assume Theorem 2.1 for the moment, and show how it implies Theorem 1.1. Let F, S, u be as in Theorem 2.1. Since u is smooth and nonzero on the compact region $\{(t, x) \in \Gamma_d : e^{-S} \leq t \leq 1\}$, it is bounded from below in this region. By replacing u with Cu and F with $v \mapsto C^2 F(v/C)$ for some large constant C , we may thus assume that

$$\|u(t, x)\|_{\mathbb{R}^m} \geq 1$$

whenever $(t, x) \in \Gamma_d$ with $e^{-S} \leq t \leq 1$. Using the discrete self-similarity property (2-1), we then have this bound for all $0 < t \leq 1$; in fact we have a lower bound on $\|u(t, x)\|_{\mathbb{R}^m}$ that goes to infinity as $t \rightarrow 0$, ensuring in particular that u has no smooth extension to $(0, 0)$.

Using a smooth cutoff function, one can find a smooth defocusing potential $\tilde{F} : \mathbb{R}^m \rightarrow \mathbb{R}$ that agrees with F in the region $\{v \in \mathbb{R}^m : \|v\|_{\mathbb{R}^m} \geq 1\}$. Then u solves (1-3) with this potential in the truncated light cone $\{(t, x) \in \mathbb{R}^{1+d} : 0 < t \leq 1; |x| \leq t\}$ with F replaced by \tilde{F} . Choose smooth initial data $v_0, v_1 : \mathbb{R}^d \rightarrow \mathbb{R}^m$ such that

$$v_0(x) = u(1, x)$$

and

$$v_1(x) = -\partial_t u(1, x)$$

for all $|x| \leq 1$ (where we use $|x| := \|x\|_{\mathbb{R}^d}$ to denote the magnitude of $x \in \mathbb{R}^d$); such data exists from standard smooth extension theorems (see, e.g., [Seeley 1964]) since the functions $u(1, x), \partial_t u(1, x)$ are smooth on the closed ball $\{x : |x| \leq 1\}$. Suppose for contradiction that Theorem 1.1 failed (with F replaced by \tilde{F}); then we have a global smooth solution $v : \mathbb{R}^{1+d} \rightarrow \mathbb{R}^m$ to (1-3) (for \tilde{F}) with initial data $v(0) = v_0, \partial_t v(0) = v_1$. The function $\tilde{u} : (t, x) \mapsto v(1-t, x)$ is then another global smooth solution to (1-3) (for \tilde{F}) such that $\tilde{u}(1, x) = u(1, x)$ and $\partial_t \tilde{u}(1, x) = \partial_t u(1, x)$ for all $|x| \leq 1$. Finite speed of propagation (see, e.g., [Tao 2006, Proposition 3.3]) then shows that \tilde{u} and u agree in the region $\{(t, x) \in \mathbb{R}^{1+d} : 0 < t \leq 1; |x| \leq t\}$; as \tilde{u} is smoothly extendible to $(0, 0)$, we know u is also, giving the desired contradiction. This concludes the derivation of Theorem 1.1 from Theorem 2.1.

It remains to prove Theorem 2.1. This will be the focus of the remaining sections of the paper. For now, let us show why continuously self-similar solutions are not available in the defocusing case, at

least for some choices of parameters d, p . The point will be that continuous self-similarity gives a new monotonicity formula for a certain quantity $f(t, r)$ (measuring a sort of “equipartition of energy”) that can be used to derive a contradiction.

Proposition 2.2 (no self-similar defocusing solutions). *Let $d \geq 3$ and $p > 1$ be such that $\frac{d-3}{2} - \frac{2}{p-1} < 0$, let m be a natural number, and let $F : \mathbb{R}^m \rightarrow \mathbb{R}$ be a defocusing potential that is homogeneous of order $p + 1$. Then there does not exist a smooth solution $u : \Gamma_d \rightarrow \mathbb{R}^m \setminus \{0\}$ to (1-3) that is homogeneous of order $-\frac{2}{p-1}$.*

Note in particular that in the physical case $d = 3$, the condition $\frac{d-3}{2} - \frac{2}{p-1} < 0$ is automatic, and so no self-similar defocusing solutions exist in this case. We do not know if this condition is necessary in the above proposition.

Proof. Suppose for contradiction that such a u exists. Equation (1-3) in polar coordinates (t, r, ω) reads

$$-\partial_{tt}u + \partial_{rr}u + \frac{d-1}{r}\partial_r u + \frac{1}{r^2}\Delta_\omega u = (\nabla F)(u),$$

where Δ_ω is the Laplace–Beltrami operator on the sphere S^{d-1} . Making the substitution

$$\phi(t, r, \omega) := r^{\frac{d-1}{2}}u(t, r, \omega), \quad (2-2)$$

this becomes

$$-\partial_{tt}\phi + \partial_{rr}\phi - \frac{1}{r^2}\left(-\Delta_\omega + \frac{(d-1)(d-3)}{4}\right)\phi = r^{\frac{d-1}{2}}(\nabla F)(r^{-\frac{d-1}{2}}\phi) \quad (2-3)$$

for $r > 0$.

We introduce the scaling vector field $S := t\partial_t + r\partial_r$ and the Lorentz boost $L := r\partial_t + t\partial_r$. Observe that L and S commute with

$$-S^2 + L^2 = (t^2 - r^2)(-\partial_{tt} + \partial_{rr}) \quad (2-4)$$

and thus

$$-\langle S^2\phi, L\phi \rangle_{\mathbb{R}^m} + \langle L^2\phi, L\phi \rangle_{\mathbb{R}^m} = (t^2 - r^2)\langle -\partial_{tt}\phi + \partial_{rr}\phi, L\phi \rangle_{\mathbb{R}^m}.$$

As u is assumed homogeneous of order $-\frac{2}{p-1}$, we know ϕ is homogeneous of order $\frac{d-1}{2} - \frac{2}{p-1}$. From Euler’s identity (1-2) we thus have ϕ an eigenfunction of S ,

$$S\phi = \left(\frac{d-1}{2} - \frac{2}{p-1}\right)\phi,$$

and thus (by the commutativity of L and S)

$$\langle L\phi, S^2\phi \rangle_{\mathbb{R}^m} = \langle LS\phi, S\phi \rangle_{\mathbb{R}^m} = \frac{1}{2}L\|S\phi\|_{\mathbb{R}^m}^2.$$

We also have

$$\langle L\phi, L^2\phi \rangle_{\mathbb{R}^m} = \frac{1}{2}L\|L\phi\|_{\mathbb{R}^m}^2.$$

Putting all of these facts together, we conclude that

$$L\left(-\frac{1}{2}\|S\phi\|_{\mathbb{R}^m}^2 + \frac{1}{2}\|L\phi\|_{\mathbb{R}^m}^2\right) = (t^2 - r^2)\langle -\partial_{tt}\phi + \partial_{rr}\phi, L\phi \rangle_{\mathbb{R}^m}.$$

A computation similar to (2-4) shows that

$$-\|S\phi\|_{\mathbb{R}^m}^2 + \|L\phi\|_{\mathbb{R}^m} = (t^2 - r^2)(-\|\partial_t\phi\|_{\mathbb{R}^m}^2 + \|\partial_r\phi\|_{\mathbb{R}^m}^2).$$

Since $t^2 - r^2$ is annihilated by L , we conclude that

$$L(-\frac{1}{2}\|\partial_t\phi\|_{\mathbb{R}^m}^2 + \frac{1}{2}\|\partial_r\phi\|_{\mathbb{R}^m}^2) = \langle -\partial_{tt}\phi + \partial_{rr}\phi, \phi \rangle_{\mathbb{R}^m}.$$

By (2-3), the right-hand side is equal to

$$\frac{1}{r^2} \langle -\Delta_\omega\phi, L\phi \rangle_{\mathbb{R}^m} + \frac{(d-1)(d-3)}{4r^2} \langle \phi, L\phi \rangle_{\mathbb{R}^m} + r^{\frac{d-1}{2}} \langle (\nabla F)(r^{-\frac{d-1}{2}}\phi), L\phi \rangle_{\mathbb{R}^m}.$$

To deal with the angular Laplacian, we integrate over S^{d-1} and then integrate by parts to conclude that

$$\begin{aligned} &L \int_{S^{d-1}} (-\frac{1}{2}\|\partial_t\phi\|_{\mathbb{R}^m}^2 + \frac{1}{2}\|\partial_r\phi\|_{\mathbb{R}^m}^2) d\omega \\ &= \int_{S^{d-1}} \frac{1}{2r^2} L\|\nabla_\omega\phi\|_{\mathbb{R}^m \otimes \mathbb{R}^d}^2 + \frac{(d-1)(d-3)}{8r^2} L\|\phi\|_{\mathbb{R}^m}^2 + r^{\frac{d-1}{2}} \langle (\nabla F)(r^{-\frac{d-1}{2}}\phi), L\phi \rangle_{\mathbb{R}^m} d\omega, \end{aligned}$$

where we use the fact that the Lorentz boost L commutes with angular derivatives, and where $d\omega$ denotes surface measure on S^{d-1} .

From the chain and product rules, noting that $Lr = t$, we have

$$L\phi = r^{\frac{d-1}{2}} L(r^{-\frac{d-1}{2}}\phi) + \frac{d-1}{2} \frac{t}{r} \phi$$

and thus (using (1-2))

$$\begin{aligned} \langle (\nabla F)(r^{-\frac{d-1}{2}}\phi), L\phi \rangle_{\mathbb{R}^m} &= r^{\frac{d-1}{2}} \left(LF(r^{-\frac{d-1}{2}}\phi) + \frac{d-1}{2} \frac{t}{r} \langle r^{-\frac{d-1}{2}}\phi, (\nabla F)(r^{-\frac{d-1}{2}}\phi) \rangle_{\mathbb{R}^m} \right) \\ &= r^{\frac{d-1}{2}} \left(LF(r^{-\frac{d-1}{2}}\phi) + \frac{(d-1)(p+1)}{2} \frac{t}{r} F(r^{-\frac{d-1}{2}}\phi) \right). \end{aligned}$$

Putting all this together, we see that if we introduce the quantity

$$\begin{aligned} f(t, r) := \int_{S^{d-1}} &-\frac{1}{2}\|\partial_t\phi\|_{\mathbb{R}^m}^2 + \frac{1}{2}\|\partial_r\phi\|_{\mathbb{R}^m}^2 - \frac{1}{2r^2}\|\nabla_\omega\phi\|_{\mathbb{R}^m \otimes \mathbb{R}^d}^2 \\ &- \frac{(d-1)(d-3)}{8r^2} \|\phi\|_{\mathbb{R}^m}^2 - r^{d-1} F(r^{-\frac{d-1}{2}}\phi) d\omega \end{aligned}$$

then we have the formula

$$Lf = \int_{S^{d-1}} \frac{t}{r^3} \|\nabla_\omega\phi\|_{\mathbb{R}^m \otimes \mathbb{R}^d}^2 + \frac{(d-1)(d-3)t}{4r^3} \|\phi\|_{\mathbb{R}^m}^2 + \frac{(d-1)(p-1)}{2} \frac{t}{r} r^{d-1} F(r^{-\frac{d-1}{2}}\phi) d\omega$$

for any $r > 0$. In particular, $f(\cosh y, \sinh y)$ is a strictly function of y for $y > 0$, since

$$\frac{d}{dy} f(\cosh y, \sinh y) = (Lf)(\cosh y, \sinh y) > 0$$

with the strict positivity coming from the defocusing nature of F . On the other hand, when $y \rightarrow 0^+$, we see from (2-2) that all the negative integrands in the definition of $f(\cosh y, \sinh y)$ go to zero, and thus

$$\lim_{y \rightarrow 0^+} f(\cosh y, \sinh y) \geq 0.$$

Combining these two facts, we conclude in particular that

$$\lim_{y \rightarrow +\infty} f(\cosh y, \sinh y) > 0. \quad (2-5)$$

On the other hand, as ϕ is homogeneous of order $\frac{d-1}{2} - \frac{2}{p-1}$ and F is homogeneous of order $p+1$, we see that the integrand in the definition of $f(t, r)$ is homogeneous of order $2\left(\frac{d-3}{2} - \frac{2}{p-1}\right)$, which is negative by hypothesis. This implies that $f(\cosh y, \sinh y)$ goes to zero as $y \rightarrow +\infty$, contradicting (2-5). \square

3. Eliminating the potential

We now exploit the freedom to select the defocusing potential F by eliminating it from the equations of motion. To motivate this elimination, let us temporarily make the a priori assumption that we have a solution u to (1-3) in the light cone Γ_d from Theorem 2.1 that is nowhere vanishing. Taking the inner product of (1-3) with u and using (1-2) then gives an equation for $F(u)$:

$$F(u) = \frac{1}{p+1} \langle u, \square u \rangle_{\mathbb{R}^m}. \quad (3-1)$$

In particular, since F is defocusing and u is nowhere vanishing, we have the defocusing property

$$\langle u, \square u \rangle_{\mathbb{R}^m} > 0 \quad (3-2)$$

throughout Γ_d . Next, if ∂_α denotes one of the $d+1$ derivative operators $\partial_t, \partial_{x_1}, \dots, \partial_{x_d}$, we have from the chain rule that

$$\partial_\alpha F(u) = \langle \partial_\alpha u, (\nabla F)(u) \rangle_{\mathbb{R}^m}$$

and hence from (1-3) and (3-1) we have the equation

$$\partial_\alpha \langle u, \square u \rangle_{\mathbb{R}^m} = (p+1) \langle \partial_\alpha u, \square u \rangle_{\mathbb{R}^m}. \quad (3-3)$$

Remark 3.1. One can rewrite the equation (3-3) in the more familiar form

$$\partial^\beta T_{\alpha\beta} = 0,$$

where $T_{\alpha\beta}$ is the stress-energy tensor

$$T_{\alpha\beta} = \langle \partial_\alpha u, \partial_\beta u \rangle_{\mathbb{R}^m} - \eta_{\alpha\beta} \left(\frac{1}{2} \langle \partial^\gamma u, \partial_\gamma u \rangle_{\mathbb{R}^m} + \frac{1}{p+1} \langle u, \square u \rangle \right).$$

Now assume that u obeys the discrete self-similarity hypothesis (2-1). Let $\theta := u/\|u\|_{\mathbb{R}^m}$ denote the direction vector of u ; then θ is smooth map from Γ_d to the unit sphere $S^{m-1} := \{v \in \mathbb{R}^m : \|v\|_{\mathbb{R}^m} = 1\}$ of \mathbb{R}^m . From the discrete self-similarity (2-1) we see that θ is invariant under the dilation action of the multiplicative group $e^{S\mathbb{Z}} := \{e^{nS} : n \in \mathbb{Z}\}$ on Γ_d . Thus θ descends to a smooth map $\tilde{\theta} : \Gamma_d/e^{S\mathbb{Z}} \rightarrow S^{m-1}$

on the compact quotient $\Gamma_d/e^{S\mathbb{Z}}$, which is a smooth surface with boundary (diffeomorphic to the product of a d -dimensional closed ball and a circle). Under some nondegeneracy hypotheses on this map, we can now eliminate the potential F , reducing Theorem 2.1 to the following claim:

Theorem 3.2 (second reduction). *Let $d = 3$, let $p > 1 + \frac{4}{d-2}$, and let*

$$m \geq 2 \max\left(\frac{(d+1)(d+6)}{2}, \frac{(d+1)(d+4)}{2} + 5\right) + 2$$

be an integer. Then there exists $S > 0$ and a smooth nowhere vanishing function $u : \Gamma_d \rightarrow \mathbb{R}^m \setminus \{0\}$ which is discretely self-similar in the sense of (2-1) and obeys the defocusing property (3-2) and the equations (3-3) throughout Γ_d . Furthermore, the map $\tilde{\theta} : \Gamma_d/\lambda^{\mathbb{Z}} \rightarrow S^{m-1}$ defined as above is injective, and immersed in the sense that the $d + 1$ derivatives $\partial_\alpha \tilde{\theta}(t, x)$ for $\alpha = 0, \dots, d$ are linearly independent in \mathbb{R}^m for each $(t, x) \in \Gamma_d$.

Let us assume Theorem 3.2 for now and see how it implies Theorem 2.1. As in the previous section, our arguments here will not depend on our hypotheses on m, d, p .

Since the map $\tilde{\theta} : \Gamma_d/e^{S\mathbb{Z}} \rightarrow S^{m-1}$ is assumed to be injective and immersed, it is a smooth embedding of the set $\Gamma_d/e^{S\mathbb{Z}}$ to S^{m-1} , so that $\tilde{\theta}(\Gamma_d/e^{S\mathbb{Z}}) = \theta(\Gamma_d)$ is a smooth manifold with boundary contained in S^{m-1} . We define a function $F_0 : \theta(\Gamma_d) \rightarrow \mathbb{R}$ by the formula

$$F_0\left(\frac{u(t, x)}{\|u(t, x)\|_{\mathbb{R}^m}}\right) := \frac{1}{(p+1)\|u(t, x)\|_{\mathbb{R}^m}^{p+1}} \langle u(t, x), \square u(t, x) \rangle_{\mathbb{R}^m} \tag{3-4}$$

for any $(t, x) \in \Gamma_d$. As θ is injective and u is nowhere vanishing and discretely self-similar, one verifies that F_0 is well-defined. As the map θ is immersed, we also see that F_0 is smooth. From (3-2) we see that F_0 is positive on $\theta(\Gamma_d)$. Intuitively, F_0 is going to be our choice for F on the set $\theta(\Gamma_d)$ (this choice is forced upon us by (3-1) and homogeneity).

We define an auxiliary function $T : \theta(\Gamma_d) \rightarrow \mathbb{R}^m$ by the formula

$$T\left(\frac{u(t, x)}{\|u(t, x)\|_{\mathbb{R}^m}}\right) := \frac{1}{\|u(t, x)\|_{\mathbb{R}^m}^p} \square u(t, x) - \frac{1}{\|u(t, x)\|_{\mathbb{R}^m}^{p+2}} \langle u(t, x), \square u(t, x) \rangle_{\mathbb{R}^m} u(t, x) \tag{3-5}$$

for all $(t, x) \in \Gamma_d$; geometrically, this is the orthogonal projection of $(1/\|u\|_{\mathbb{R}^m}^p) \square u$ to the tangent plane of S^m at $u/\|u\|_{\mathbb{R}^m}$, and will be our choice for the S^{m-1} gradient

$$(\nabla_{S^{m-1}} F)\left(\frac{u}{\|u\|}\right) = (\nabla_{\mathbb{R}^m} F)\left(\frac{u}{\|u\|_{\mathbb{R}^m}}\right) - \left\langle \frac{u}{\|u\|_{\mathbb{R}^m}}, (\nabla_{\mathbb{R}^m} F)\left(\frac{u}{\|u\|}\right) \right\rangle_{\mathbb{R}^m} \frac{u}{\|u\|_{\mathbb{R}^m}}$$

of F at $u/\|u\|_{\mathbb{R}^m}$.

As θ is injective and u is nowhere vanishing and discretely self-similar, one verifies as before that T is well-defined, and from the immersed nature of θ we see that T is smooth. Clearly $T(\omega)$ is also orthogonal to ω for any $\omega \in \theta(\Gamma_d)$. We also claim that T is an extension of the gradient $\nabla_{\theta(\Gamma_d)} F_0$ of F_0 on $\theta(\Gamma_d)$, in the sense that

$$\langle v, \nabla_{\theta(\Gamma_d)} F_0(\omega) \rangle_{\mathbb{R}^m} = \langle v, T(\omega) \rangle_{\mathbb{R}^m} \tag{3-6}$$

for any $\omega \in \theta(\Gamma_d)$ and tangent vectors $v \in T_\omega\theta(\Gamma_d)$ to $\theta(\Gamma_d)$ at ω . To verify (3-6), we write

$$\omega = \frac{u(t, x)}{\|u(t, x)\|_{\mathbb{R}^m}} = \frac{u}{\|u\|}$$

for some $(t, x) \in \Gamma_d$; henceforth we suppress the explicit dependence on (t, x) for brevity. The tangent space to $\theta(\Gamma_d)$ at ω is spanned by $\partial_\alpha(u/\|u\|)$ for $\partial_\alpha = \partial_t, \partial_{x_1}, \dots, \partial_{x_d}$, so it suffices to show that

$$\left\langle \partial_\alpha \frac{u}{\|u\|}, \nabla_{\theta(\Gamma_d)} F_0(\omega) \right\rangle_{\mathbb{R}^m} = \left\langle \partial_\alpha \frac{u}{\|u\|}, T(\omega) \right\rangle_{\mathbb{R}^m}$$

for each ∂_α . But from the chain and product rules and (3-4), (3-3) and (3-5) we have

$$\begin{aligned} \left\langle \partial_\alpha \frac{u}{\|u\|}, \nabla_{\theta(\Gamma_d)} F_0(\omega) \right\rangle_{\mathbb{R}^m} &= \partial_\alpha F_0 \left(\frac{u}{\|u\|} \right) \\ &= \frac{1}{p+1} \partial_\alpha \left(\frac{1}{\|u\|_{\mathbb{R}^m}^{p+1}} \langle u, \square u \rangle_{\mathbb{R}^m} \right) \\ &= -\frac{\langle u, \partial_\alpha u \rangle_{\mathbb{R}^m}}{\|u\|_{\mathbb{R}^m}^{p+3}} \langle u, \square u \rangle_{\mathbb{R}^m} + \frac{\partial_\alpha \langle u, \square u \rangle_{\mathbb{R}^m}}{(p+1)\|u\|_{\mathbb{R}^m}^{p+1}} \\ &= -\frac{\langle u, \partial_\alpha u \rangle_{\mathbb{R}^m}}{\|u\|_{\mathbb{R}^m}^{p+3}} \langle u, \square u \rangle_{\mathbb{R}^m} + \frac{\langle \partial_\alpha u, \square u \rangle_{\mathbb{R}^m}}{\|u\|_{\mathbb{R}^m}^{p+1}} \\ &= \left\langle \frac{1}{\|u\|_{\mathbb{R}^m}} \partial_\alpha u, T \left(\frac{u}{\|u\|_{\mathbb{R}^m}} \right) \right\rangle_{\mathbb{R}^m} \\ &= \left\langle \partial_\alpha \frac{u}{\|u\|_{\mathbb{R}^m}}, T \left(\frac{u}{\|u\|_{\mathbb{R}^m}} \right) \right\rangle_{\mathbb{R}^m} \end{aligned}$$

as desired, where in the final line comes from the orthogonality of $T(u/\|u\|_{\mathbb{R}^m})$ with scalar multiples of u .

We now claim that we may find an open neighbourhood U of $\theta(\Gamma_d)$ in S^{m-1} and a smooth extension $F_1 : U \rightarrow \mathbb{R}$ of F_0 , with the property that

$$\nabla_{S^{m-1}} F_1(\omega) = T(\omega) \tag{3-7}$$

for all $\omega \in \theta(\Gamma_d)$. Indeed, we can define

$$F_1(\omega + v) := F_0(\omega) + \langle v, T(\omega) \rangle_{\mathbb{R}^m}$$

for all $\omega \in \theta(\Gamma_d)$ and sufficiently small $v \in \mathbb{R}^m$ orthogonal to the tangent space $T_\omega\theta(\Gamma_d/e^{S\mathbb{Z}})$ with $\omega + v \in S^{m-1}$; one can verify that this is well-defined as a smooth extension of F_0 to a sufficiently small normal neighbourhood of $\theta(\Gamma_d)$ with the desired gradient property (3-7) (here we use (3-6) to deal with tangential components of the gradient), and one may smoothly extend this to an open neighbourhood of $\theta(\Gamma_d)$ by Seeley's theorem [1964].

Next, if we extend F_1 by zero to all of S^{m-1} and define $F_2 : S^{m-1} \rightarrow \mathbb{R}$ to be the function $F_2 := \psi F_1 + (1-\psi)$ for some smooth function $\psi : S^{m-1} \rightarrow [0, 1]$ supported in U that equals 1 on a neighbourhood of $\theta(\Gamma_d)$, then F_2 is a smooth extension of F_0 to S^{m-1} that is strictly positive, and which also obeys

(3-7). If we then set $F : \mathbb{R}^m \rightarrow \mathbb{R}$ to be the function

$$F(\lambda\omega) := \lambda^{p+1} F_2(\omega)$$

for all $\lambda \geq 0$ and $\omega \in S^{m-1}$, then F is a defocusing potential, homogeneous of order $p + 1$, which extends F_0 , and such that

$$\nabla_{S^{m-1}} F(\omega) = T(\omega)$$

for $\omega \in \theta(\Gamma_d)$. By homogeneity (1-1), the radial derivative $\langle \omega, \nabla_{\mathbb{R}^m} F(\omega) \rangle_{\mathbb{R}^m}$ is

$$(p + 1)F(\omega) = (p + 1)F_0(\omega)$$

for such ω , and hence for $\omega = u/\|u\|$ by (3-5) and (3-4) we have

$$\begin{aligned} \nabla_{\mathbb{R}^m} F(\omega) &= T(\omega) + (p + 1)F_0(\omega)\omega \\ &= \frac{1}{\|u\|^p} \square u - \frac{\langle u, \square u \rangle}{\|u\|^{p+2}} u + \frac{p + 1}{(p + 1)\|u\|^{p+1}} \langle u, \square u \rangle \frac{u}{\|u\|} \\ &= \frac{1}{\|u\|^p} \square u; \end{aligned}$$

since $\nabla_{\mathbb{R}^m} F$ is homogeneous of order p , this gives (1-3) as required.

It remains to establish Theorem 3.2. This will be the focus of the remaining sections of the paper.

4. Eliminating the field

Having eliminated the potential F from the problem, the next step is (perhaps surprisingly) to eliminate the unknown field u , replacing it with quadratic data such as the mass density

$$M(t, x) := \|u(t, x)\|_{\mathbb{R}^m}^2 \tag{4-1}$$

and the energy tensor

$$E_{\alpha\beta}(t, x) := \langle \partial_\alpha u(t, x), \partial_\beta u(t, x) \rangle. \tag{4-2}$$

If u has the discrete self-similarity property (2-1), then M and E similarly obey the discrete self-similarity properties

$$M(e^S t, e^S x) = e^{-\frac{4}{p-1}S} M(t, x) \tag{4-3}$$

and

$$E_{\alpha\beta}(e^S t, e^S x) = e^{-\frac{2(p+1)}{p-1}S} E_{\alpha\beta}(t, x). \tag{4-4}$$

Next, observe from the product rule that

$$\langle u, \square u \rangle_{\mathbb{R}^m} = \frac{1}{2} \square M - \eta^{\beta\gamma} E_{\beta\gamma}, \tag{4-5}$$

where η is the Minkowski metric. Thus, the defocusing property (3-2) can be rewritten as

$$\frac{1}{2} \square M - \eta^{\alpha\beta} E_{\alpha\beta} > 0. \tag{4-6}$$

In a similar spirit, we have

$$\langle \partial_\alpha u, \square u \rangle = \partial^\beta E_{\alpha\beta} - \frac{1}{2} \partial_\alpha (\eta^{\beta\gamma} E_{\beta\gamma})$$

and hence the equation (3-3) can be expressed in terms of M and E as

$$\partial_\alpha \left(\frac{1}{2} \square M - \eta^{\beta\gamma} E_{\beta\gamma} \right) = (p+1) \left(\partial^\beta E_{\alpha\beta} - \frac{1}{2} \partial_\alpha (\eta^{\beta\gamma} E_{\beta\gamma}) \right). \quad (4-7)$$

Finally, observe that the $(2+d) \times (2+d)$ Gram matrix

$$\begin{pmatrix} \langle u(t, x), u(t, x) \rangle_{\mathbb{R}^m} & \langle u(t, x), \partial_t u(t, x) \rangle_{\mathbb{R}^m} & \cdots & \langle u(t, x), \partial_{x_d} u(t, x) \rangle_{\mathbb{R}^m} \\ \langle \partial_t u(t, x), u(t, x) \rangle_{\mathbb{R}^m} & \langle \partial_t u(t, x), \partial_t u(t, x) \rangle_{\mathbb{R}^m} & \cdots & \langle \partial_t u(t, x), \partial_{x_d} u(t, x) \rangle_{\mathbb{R}^m} \\ \vdots & \vdots & \ddots & \vdots \\ \langle \partial_{x_d} u(t, x), u(t, x) \rangle_{\mathbb{R}^m} & \langle \partial_{x_d} u(t, x), \partial_t u(t, x) \rangle_{\mathbb{R}^m} & \cdots & \langle \partial_{x_d} u(t, x), \partial_{x_d} u(t, x) \rangle_{\mathbb{R}^m} \end{pmatrix} \quad (4-8)$$

can be expressed in terms of E, M as

$$\begin{pmatrix} M(t, x) & \frac{1}{2} \partial_t M(t, x) & \cdots & \frac{1}{2} \partial_{x_d} M(t, x) \\ \frac{1}{2} \partial_t M(t, x) & E_{00}(t, x) & \cdots & E_{0d}(t, x) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{2} \partial_{x_d} M(t, x) & E_{d0}(t, x) & \cdots & E_{dd}(t, x) \end{pmatrix}. \quad (4-9)$$

In particular, the matrix (4-9) is positive semidefinite for every t, x .

It turns out that with the aid of the Nash embedding theorem and our hypothesis that m is large, we can largely reverse the above observations, reducing Theorem 3.2 to the following claim that no longer directly involves the field u (or the range dimension m).

Theorem 4.1 (third reduction). *Let $d = 3$, and let $p > 1 + \frac{4}{d-2}$. Then there exists $S > 0$ and smooth functions $M : \Gamma_d \rightarrow \mathbb{R}$ and $E_{\alpha\beta} : \Gamma_d \rightarrow \mathbb{R}$ for $\alpha, \beta = 0, \dots, d$ which are discretely self-similar in the sense of (4-3) and (4-4), obey the defocusing property (4-6) and the equation (4-7) on Γ_d for all $\alpha = 0, \dots, d$, and such that the matrix (4-9) is strictly positive definite on Γ_d (in particular, this forces M to be strictly positive).*

Let us assume Theorem 4.1 for the moment and show Theorem 3.2. Let $d, p, S, M, E_{\alpha\beta}$ be as in Theorem 4.1, and let m be as in Theorem 3.2. Our task is to obtain a function $u : \Gamma_d \rightarrow \mathbb{R}^m \setminus \{0\}$ obeying all the properties claimed in Theorem 3.2.

The idea is to build u in such a fashion that (4-1) and (4-2) are obeyed. Accordingly, we will use an ansatz

$$u(t, x) := M(t, x)^{\frac{1}{2}} \theta(t, x) \quad (4-10)$$

for some smooth $\theta : \Gamma_d \rightarrow S^{m-1}$ to be constructed shortly. As M is strictly positive, such a function u will be smooth on Γ and obey (4-1); differentiating, we see that

$$\langle u, \partial_\alpha u \rangle_{\mathbb{R}^m} = \frac{1}{2} \partial_\alpha M \quad (4-11)$$

for $\alpha = 0, \dots, d$. If θ obeys the discrete self-similarity property

$$\theta(e^S t, e^S x) = \theta(t, x) \quad (4-12)$$

then u will obey (2-1). Thus we shall impose (4-12); that is to say we assume that θ is lifted from a smooth map $\tilde{\theta} : \Gamma_d/e^{S\mathbb{Z}} \rightarrow S^{m-1}$.

From the product rule, (4-1) and (4-11) we have (after some calculation)

$$\langle \partial_\alpha \theta, \partial_\beta \theta \rangle_{\mathbb{R}^m} = M^{-1} \langle \partial_\alpha u, \partial_\beta u \rangle_{\mathbb{R}^m} - M^{-2} (\partial_\alpha M) \partial_\beta M.$$

Thus, if we wish for (4-2) to be obeyed, then the $(1+d) \times (1+d)$ Gram matrix

$$(\langle \partial_\alpha \theta, \partial_\beta \theta \rangle_{\mathbb{R}^m})_{\alpha, \beta=0, \dots, d}$$

must be equal to

$$M(E_{\alpha\beta} - (\partial_\alpha M)M^{-1}\partial_\beta M)_{\alpha, \beta=0, \dots, d}. \tag{4-13}$$

The matrix in (4-13) is a Schur complement of the matrix in (4-9). Since the matrix in (4-9) is assumed to be strictly positive definite, we conclude that (4-13) is also.

If we denote the matrix in (4-13) by $g(t, x)$, then from (4-3) and (4-4) we have the discrete self-similarity property

$$g(e^S t, e^S x) = e^{-2S} g(t, x). \tag{4-14}$$

As g is a positive definite and symmetric $(1+d) \times (1+d)$ matrix, we can view g as a smooth Riemannian metric on Γ_d . Given that the dilation operator $(t, x) \mapsto (e^S t, e^S x)$ dilates tangent vectors to Γ_d by a factor of e^S , we see that the metric g is lifted from a smooth Riemannian metric \tilde{g} on the quotient space $\Gamma_d/e^{S\mathbb{Z}}$.

The space $(\Gamma_d/e^{S\mathbb{Z}}, \tilde{g})$ is a smooth compact $(1+d)$ -dimensional Riemannian manifold with boundary; it is easy to embed it in a smooth compact $(1+d)$ -dimensional Riemannian manifold without boundary (for instance by using the theorems in [Seeley 1964]). Applying the Nash embedding theorem (for instance in the form in [Günther 1991]), we can thus isometrically embed $(\Gamma_d/e^{S\mathbb{Z}}, \tilde{g})$ in a Euclidean space \mathbb{R}^D with

$$D := \max\left(\frac{(d+1)(d+6)}{2}, \frac{(d+1)(d+4)}{2} + 5\right).$$

The embedded copy of $(\Gamma_d/e^{S\mathbb{Z}}, \tilde{g})$ is compact and is thus contained in a cube $[-R, R]^D$ for some finite R . We use a generic³ linear isometry from \mathbb{R}^D to \mathbb{R}^{D+1} to embed $[-R, R]^D$ to some compact subset of \mathbb{R}^{D+1} . The image of this isometry is a generic hyperplane, which can be chosen to avoid the lattice $(1/\sqrt{2D+2})\mathbb{Z}^{D+1}$, and thus we can embed $[-R, R]^D$ isometrically into the torus $\mathbb{R}^{D+1}/((1/\sqrt{D+1})\mathbb{Z}^{D+1})$, which is isometric to $(1/\sqrt{D+1})(S^1)^{D+1}$. But from Pythagoras' theorem, $(1/\sqrt{D+1})(S^1)^{D+1}$ is contained in S^{2D+1} , which is in turn contained in S^{m-1} by the largeness hypothesis on m . Thus we have an isometric embedding $\tilde{\theta} : \Gamma_d/e^{S\mathbb{Z}} \rightarrow S^{m-1}$ from $(\Gamma_d/e^{S\mathbb{Z}}, \tilde{g})$ into the round sphere S^{m-1} . In particular, $\tilde{\theta}$ is injective and immersed, and lifting $\tilde{\theta}$ back to Γ_d , we obtain a smooth map $\theta : \Gamma_d \rightarrow S^{m-1}$ with Gram matrix (4-13) that is discretely self-similar in the sense of (4-12), so that the function u defined by (4-10) obeys (2-1). Reversing the calculations that led to (4-13), we

³We thank Marc Nardmann for this argument, which improved the value of m from our previous argument by a factor of approximately two.

see that the Gram matrix (4-8) of u is given by (4-9). In particular, (4-2) holds. Reversing the derivation of (4-6), we now obtain (3-2), while from reversing the derivation of (4-7), we obtain (3-3). We have now obtained all the required properties claimed by Theorem 3.2, as desired.

It remains to establish Theorem 4.1. This will be the focus of the remaining sections of the paper.

5. Reduction to a self-similar (1+1)-dimensional problem

In reducing Theorem 1.1 to Theorem 4.1, we have achieved the somewhat remarkable feat of converting a nonlinear PDE problem to a convex (or positive semidefinite) PDE problem, in that all of the constraints⁴ on the remaining unknowns M , $E_{\alpha\beta}$ are linear equalities and inequalities, or assertions that certain matrices are positive definite. Among other things, this shows that if one has a given solution M , $E_{\alpha\beta}$ to Theorem 4.1, and then one averages that solution over some compact symmetry group that acts on the space of such solutions, then the average will also be a solution to Theorem 4.1. In particular, one can then reduce without any loss of generality to considering solutions that are invariant with respect to that symmetry.

For instance, given that M , $E_{\alpha\beta}$ are already discretely self-similar by (4-3) and (4-4), the space of solutions has an action of the compact dilation group $\mathbb{R}^+/e^{S\mathbb{Z}}$, with (the quotient representative of) any real number $\lambda > 0$ acting on M , $E_{\alpha\beta}$ by the action

$$(\lambda \cdot M)(t, x) := \frac{1}{\lambda^{\frac{4}{p-1}}} M\left(\frac{t}{\lambda}, \frac{x}{\lambda}\right)$$

and

$$(\lambda \cdot E_{\alpha\beta})(t, x) := \frac{1}{\lambda^{\frac{2(p+1)}{p-1}}} E_{\alpha\beta}\left(\frac{t}{\lambda}, \frac{x}{\lambda}\right);$$

this is initially an action of the multiplicative group \mathbb{R}^+ , but descends to an action of $\mathbb{R}^+/e^{S\mathbb{Z}}$ thanks to (4-3) and (4-4). By the preceding discussion, we may restrict without loss of generality to the case when M , $E_{\alpha\beta}$ are invariant with respect to this $\mathbb{R}^+/e^{S\mathbb{Z}}$, or equivalently that M , $E_{\alpha\beta}$ are homogeneous of order $-\frac{4}{p-1}$ and $\frac{2(p+1)}{p-1}$ respectively. With this restriction, the parameter S no longer plays a role and may be discarded.

Remark 5.1. This reduction may seem at first glance to be in conflict with the negative result in Proposition 2.2. However, the requirement that the mass density M and the energy tensor $E_{\alpha\beta}$ be homogeneous is strictly weaker than the hypothesis that the field u itself is homogeneous. For instance, one could imagine a “twisted self-similar” solution in which the homogeneity condition (1-1) on u is replaced with a more general condition of the form

$$u(\lambda t, \lambda x) = \lambda^{-\frac{2}{p-1}} \exp(J \log \lambda) u(t, x)$$

for all $(t, x) \in \Gamma_d$ and $\lambda > 0$, where $J : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a fixed skew-adjoint linear transformation. (To be compatible with (1-3), one would also wish to require that the potential F is invariant with respect to the orthogonal transformations $\exp(sJ)$ for $s \in \mathbb{R}$.) Such solutions u would not be homogeneous, but the associated densities M , $E_{\alpha\beta}$ would still be homogeneous of the order specified above.

⁴Compare with the “kernel trick” in machine learning, or with semidefinite relaxation in optimization.

We may similarly apply the above reductions to the orthogonal group $O(d)$, which acts on the scalar field M and on the 2-tensor $E_{\alpha\beta}$ in the usual fashion; thus

$$(UM)(t, x) := M(t, U^{-1}x)$$

and

$$(UE)_{\alpha\beta}(t, x)(Uv)^\alpha(Uv)^\beta = E_{\alpha\beta}(t, U^{-1}x)v^\alpha v^\beta$$

for all $(t, x) \in \Gamma_d$, $U \in O(d)$, and $v \in \mathbb{R}^{1+d}$, where U acts on \mathbb{R}^{1+d} by $(t, x) \mapsto (t, Ux)$. This allows us to reduce to fields $M, E_{\alpha\beta}$ which are $O(d)$ -invariant; thus M is spherically symmetric, and $E_{\alpha\beta}$ takes the form⁵

$$E_{00} = E_{tt}, \tag{5-1}$$

$$E_{0i} = E_{i0} = \frac{x_i}{r} E_{tr}, \tag{5-2}$$

$$E_{ij} = \frac{x_i x_j}{r^2} (E_{rr} - E_{\omega\omega}) + \delta_{ij} E_{\omega\omega} \tag{5-3}$$

for $i, j = 1, \dots, d$ and some spherically symmetric scalar functions $E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$, where $r := |x|$ is the radial variable and δ_{ij} is the Kronecker delta. Observe that if $E_{tt}, E_{rr}, E_{\omega\omega} : \Gamma_1 \rightarrow \mathbb{R}$ are smooth even functions and $E_{tr} : \Gamma_1 \rightarrow \mathbb{R}$ is a smooth odd function on the $(1+1)$ -dimensional light cone

$$\Gamma_1 := \{(t, r) \in \mathbb{R}^{1+1} : t > 0; -t \leq r \leq t\}$$

with $E_{rr} - E_{\omega\omega}$ vanishing to second order at $r = 0$, then the above equations define a smooth field $E_{\alpha\beta}$ on Γ_d , which will be homogeneous of order $-\frac{2(p+1)}{p-1}$ if $E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$ are.

Using polar coordinates, we have

$$\frac{1}{2} \square M - \eta^{\beta\gamma} E_{\beta\gamma} = \frac{1}{2} \left(-\partial_{tt} M + \partial_{rr} M + \frac{d-1}{r} M \right) - (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega});$$

thus the condition (4-6) is now

$$\frac{1}{2} \left(-\partial_{tt} M + \partial_{rr} M + \frac{d-1}{r} M \right) - (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) > 0. \tag{5-4}$$

By rotating x to be of the form $x = r e_1$, we see that the matrix (4-9) is conjugate to

$$\begin{pmatrix} M & \frac{1}{2} \partial_t M & \frac{1}{2} \partial_r M & 0 & \cdots & 0 \\ \frac{1}{2} \partial_t M & E_{tt} & E_{tr} & 0 & \cdots & 0 \\ \frac{1}{2} \partial_r M & E_{tr} & E_{rr} & 0 & \cdots & 0 \\ 0 & 0 & 0 & E_{\omega\omega} & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & E_{\omega\omega} \end{pmatrix}$$

⁵To see that $E_{\alpha\beta}$ must be of this form, rotate the spatial variable x to equal $x = r e_1$, then use the orthogonal transformation $(x_1, x_2, \dots, x_d) \mapsto (x_1, -x_2, \dots, -x_d)$, which preserves $r e_1$, to see that $E_{0i} = E_{1i} = 0$ for all $i = 2, \dots, d$; further use of orthogonal transformations preserving $r e_1$ can be then used to show that $E_{ij} = 0$ and $E_{ii} = E_{jj}$ for $2 \leq i < j \leq d$ (basically because the only matrices that commute with all orthogonal transformations are scalar multiples of the identity). This places $E_{\alpha\beta}$ in the desired form in the $x = r e_1$ case, and the general case follows from rotation.

so the positive definiteness of (4-9) is equivalent to the positive definiteness of the 3×3 matrix

$$\begin{pmatrix} M & \frac{1}{2}\partial_t M & \frac{1}{2}\partial_r M \\ \frac{1}{2}\partial_t M & E_{tt} & E_{tr} \\ \frac{1}{2}\partial_r M & E_{tr} & E_{rr} \end{pmatrix} \quad (5-5)$$

together with the positivity of $E_{\omega\omega}$. It will be convenient to isolate the $r = 0$ case of this condition (in order to degenerate E_{rr} to zero at $r = 0$ later in the argument). In this case, the odd functions $\partial_r M$ and E_{tr} vanish, and E_{rr} is equal to $E_{\omega\omega}$, so the condition reduces to the positive definiteness of the 2×2 matrix

$$\begin{pmatrix} M & \frac{1}{2}\partial_t M \\ \frac{1}{2}\partial_t M & E_{tt} \end{pmatrix} \quad (5-6)$$

together with the aforementioned positivity of $E_{\omega\omega}$.

Finally, we turn to the condition (4-7). Again, we can rotate the position x to be of the form $x = r e_1$. In the angular cases $\alpha = 2, \dots, d$, both sides of (4-7) automatically vanish, basically because $\partial_\alpha f(r e_1) = 0$ for any spherically symmetric f (and because $E_{\alpha\beta}$ vanishes to second order for any $\beta \neq \alpha$). So the only nontrivial cases of (4-7) are $\alpha = 0$ and $\alpha = 1$. Applying (5-1), (5-2), and (5-3), we can write these cases of (4-7) as

$$\begin{aligned} \partial_t \left[\frac{1}{2} \left(-\partial_{tt} M + \partial_{rr} M + \frac{d-1}{r} M \right) - (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) \right] \\ = (p+1) \left[-\partial_t E_{tt} + \partial_r E_{tr} + \frac{d-1}{r} E_{tr} - \frac{1}{2} \partial_t (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) \right] \end{aligned} \quad (5-7)$$

and

$$\begin{aligned} \partial_r \left[\frac{1}{2} \left(-\partial_{tt} M + \partial_{rr} M + \frac{d-1}{r} M \right) - (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) \right] \\ = (p+1) \left[-\partial_t E_{tr} + \partial_r E_{rr} + \frac{d-1}{r} (E_{rr} - E_{\omega\omega}) - \frac{1}{2} \partial_r (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) \right] \end{aligned} \quad (5-8)$$

respectively.

To summarise, we have reduced Theorem 4.1 to

Theorem 5.2 (fourth reduction). *Let $d = 3$, and let $p > 1 + \frac{4}{d-2}$. Then there exist smooth even functions $M, E_{tt}, E_{rr}, E_{\omega\omega} : \Gamma_1 \rightarrow \mathbb{R}$ and a smooth odd function $E_{tr} : \Gamma_1 \rightarrow \mathbb{R}$, with M homogeneous of order $-\frac{4}{p-1}$ and $E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$ homogeneous of order $-\frac{2(p+1)}{p-1}$, and with $E_{rr} - E_{\omega\omega}$ vanishes to second order at $r = 0$, obeying the defocusing property (5-4) and the equations (5-7) and (5-8) on Γ_1 , such that*

$$E_{\omega\omega} > 0 \quad (5-9)$$

and the 3×3 matrix (5-5) is strictly positive definite on Γ_1 with $r \neq 0$, and the 2×2 matrix (5-6) is positive definite when $r = 0$.

It remains to prove Theorem 5.2. To do so, we make a few technical relaxations. Firstly, we claim that we may relax the strict conditions (5-4) and (5-9) to their nonstrict counterparts

$$\frac{1}{2} \left(-\partial_{tt} M + \partial_{rr} M + \frac{d-1}{r} M \right) - (-E_{tt} + E_{rr} + (d-1)E_{\omega\omega}) \geq 0 \tag{5-10}$$

and

$$E_{\omega\omega} \geq 0. \tag{5-11}$$

To see this, suppose that $M, E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$ obey the conclusions of Theorem 5.2 with the conditions (5-4), (5-9) replaced by (5-10), (5-11). We let $\varepsilon > 0$ be a small quantity to be chosen later, and define new fields $M^\varepsilon, E_{tt}^\varepsilon, E_{tr}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$ by the formulae⁶

$$\begin{aligned} M^\varepsilon &:= M - c\varepsilon t^{-\frac{4}{p-1}}, \\ E_{tt}^\varepsilon &:= E_{tt} - (d+1)\varepsilon t^{-\frac{2(p+1)}{p-1}}, \\ E_{tr}^\varepsilon &:= E_{tr}, \\ E_{rr}^\varepsilon &:= E_{rr} + \varepsilon t^{-\frac{2(p+1)}{p-1}}, \\ E_{\omega\omega}^\varepsilon &:= E_{\omega\omega} + \varepsilon t^{-\frac{2(p+1)}{p-1}}, \end{aligned}$$

where c is the constant such that

$$\frac{c}{2} \frac{4}{p-1} \frac{p+3}{p-1} - (2d+1) = \frac{p+1}{2}.$$

Clearly these new fields $M^\varepsilon, E_{tt}^\varepsilon, E_{tr}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$ are still smooth, with $M^\varepsilon, E_{tt}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$ even and E_{tr}^ε odd, with M^ε homogeneous of order $-\frac{4}{p-1}$ and $E_{tt}^\varepsilon, E_{tr}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$ homogeneous of order $-\frac{2(p+1)}{p-1}$, with $E_{rr}^\varepsilon - E_{\omega\omega}^\varepsilon$ vanishing to second order at $r = 0$. A calculation using the definition of c shows that the equations (5-7) and (5-8) continue to be obeyed when the fields $M, E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$ are replaced by $M^\varepsilon, E_{tt}^\varepsilon, E_{tr}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$. With this replacement, the left-hand side of (5-10) increases by

$$\frac{p+1}{2} \varepsilon t^{-\frac{2(p+1)}{p-1}},$$

and so (5-4) now holds. The remaining task is to show that with these new fields $M^\varepsilon, E_{tt}^\varepsilon, E_{tr}^\varepsilon, E_{rr}^\varepsilon, E_{\omega\omega}^\varepsilon$, (5-5) is positive definite when $r \neq 0$ and (5-6) is positive definite when $r = 0$. By the scale invariance it suffices to verify these latter properties when $t = 1$. The positive definiteness of (5-6) when $r = 0$ then follows by continuity for ε small enough. For (5-5), we have to take a little care because the condition $r \neq 0$ is noncompact. We need to ensure the positive definiteness of

$$\begin{pmatrix} M - c\varepsilon & \frac{1}{2}\partial_t M + \frac{2c}{p-1}\varepsilon & \frac{1}{2}\partial_r M \\ \frac{1}{2}\partial_t M + \frac{2c}{p-1}\varepsilon & E_{tt} - (d+1)\varepsilon & E_{tr} \\ \frac{1}{2}\partial_r M & E_{tr} & E_{rr} + \varepsilon \end{pmatrix}$$

⁶The ability to freely manipulate the fields $M, E_{tt}, E_{tr}, E_{rr}, E_{\omega\omega}$ in this fashion is a major advantage of the formulation of Theorem 5.2. It would be very difficult to perform analogous manipulations if the original field u or the potential F were still present.

when $t = 1$ and $r \neq 0$ for ε small enough. Continuity will ensure this if $|r|$ is bounded away from zero (independently of ε), so we may assume that r is in a small neighbourhood of the origin (independent of ε). Given that the above matrix is already positive definite when $\varepsilon = 0$, it suffices by a continuity argument to show that the above matrix has positive determinant for sufficiently small ε ; by the hypothesis (5-5) and the fundamental theorem of calculus, it thus suffices to show that

$$\frac{d}{d\varepsilon} \det \begin{pmatrix} M - c\varepsilon & \frac{1}{2}\partial_t M + \frac{2c}{p-1}\varepsilon & \frac{1}{2}\partial_r M \\ \frac{1}{2}\partial_t M + \frac{2c}{p-1}\varepsilon & E_{tt} - (d+1)\varepsilon & E_{tr} \\ \frac{1}{2}\partial_r M & E_{tr} & E_{rr} + \varepsilon \end{pmatrix} > 0$$

for r near zero and sufficiently small ε . But since $\partial_r M, E_{tr}, E_{rr}$ vanish at $r = 0$, we can use cofactor expansion to write the left-hand side as

$$\begin{pmatrix} M(1, 0) & \frac{1}{2}\partial_t M(1, 0) \\ \frac{1}{2}\partial_t M(1, 0) & E_{tt}(1, 0) \end{pmatrix} + O(|r|) + O(\varepsilon)$$

and the claim then follows from the hypothesis (5-5). This concludes the relaxation of the conditions (5-4), (5-9) to (5-10), (5-11).

Now that we allow equality in (5-11), we sacrifice some generality by restricting to the special case $E_{\omega\omega} = 0$ (which basically corresponds to considering spherically symmetric blowup solutions). While this gives up some flexibility, this will simplify our calculations a bit as we now only have four fields $M, E_{tt}, E_{tr}, E_{rr}$ to deal with, rather than five.

Until now we have avoided using the hypothesis $d = 3$. Now we will embrace this hypothesis. In Proposition 2.2 it was convenient to make the change of variables $\phi = r^{\frac{d-1}{2}} u = ru$ to eliminate lower-order terms such as $(\frac{d-1}{r})\partial_r u$; this change of variables is particularly pleasant in the three-dimensional case as the lower-order term involving the coefficient $\frac{1}{4}(d-1)(d-3)$ vanishes completely (this vanishing is closely tied to the strong Huygens principle in three dimensions). The corresponding change of variables in this setting, aimed at eliminating the lower-order terms $(\frac{d-1}{r})E_{tr}$ and $(\frac{d-1}{r})E_{rr}$ in (5-7) and (5-8), is to replace the fields $M, E_{tt}, E_{tr}, E_{rr}$ by the fields $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{tr}, \tilde{E}_{rr} : \Gamma_1 \rightarrow \mathbb{R}^+$ defined by

$$\begin{aligned} \tilde{M} &:= r^2 M, \\ \tilde{E}_{tt} &:= r^2 E_{tt}, \\ \tilde{E}_{tr} &:= r^2 E_{tr} + \frac{1}{2}r\partial_t M = r^2 E_{tr} + \frac{1}{2r}\partial_t \tilde{M}, \\ \tilde{E}_{rr} &:= r^2 E_{rr} + r\partial_r M + M = r^2 E_{rr} + \frac{1}{r}\partial_r \tilde{M} - \frac{1}{r^2}\tilde{M}. \end{aligned}$$

Observe that if $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{rr}$ are smooth and even, and \tilde{E}_{tr} is odd, with $\tilde{M}, \tilde{E}_{tt}$ vanishing to second order at $r = 0$,

$$\tilde{E}_{tr} - \frac{1}{2r}\partial_t \tilde{M}$$

vanishing to third order, and

$$\tilde{E}_{rr} - \frac{1}{r}\partial_r \tilde{M} + \frac{1}{r^2}\tilde{M}$$

to fourth order, then these fields determine smooth fields $M, E_{tt}, E_{tr}, E_{rr}$ with M, E_{tt}, E_{rr} even, E_{tr} odd, and E_{rr} vanishing to second order at $r = 0$. Furthermore, if \tilde{M} is homogeneous of order $\frac{2p-6}{p-1}$ and $\tilde{E}_{tt}, \tilde{E}_{tr}, \tilde{E}_{rr}$ are homogeneous of order $-\frac{4}{p-1}$, then we know M will be homogeneous of order $-\frac{4}{p-1}$ and E_{tt}, E_{tr}, E_{rr} will be homogeneous of order $-\frac{2(p+1)}{p-1}$.

If we introduce the quantity

$$V := \frac{1}{p+1} \left(\frac{1}{2} (-\partial_{tt} \tilde{M} + \partial_{rr} \tilde{M}) + \tilde{E}_{tt} - \tilde{E}_{rr} \right) \quad (5-12)$$

then a brief calculation shows that

$$V = \frac{r^2}{2(p+1)} \left((-\partial_{tt} M + \partial_{rr} M + \frac{2}{r} M) - (-E_{tt} + E_{rr}) \right)$$

and so the condition (5-10) is equivalent to

$$V \geq 0. \quad (5-13)$$

The equations (5-7) and (5-8) can now be expressed as

$$\partial_t \left[\frac{1}{r^2} V \right] = -\partial_t E_{tt} + \partial_r E_{tr} + \frac{2}{r} E_{tr} - \frac{1}{2} \partial_t (-E_{tt} + E_{rr})$$

and

$$\partial_r \left[\frac{1}{r^2} V \right] = -\partial_t E_{tr} + \partial_r E_{rr} + \frac{2}{r} E_{rr} - \frac{1}{2} \partial_r (-E_{tt} + E_{rr}),$$

which rearrange as an energy conservation law

$$\partial_t \left(\frac{1}{2} E_{tt} + \frac{1}{2} E_{rr} + \frac{1}{r^2} V \right) = \partial_r E_{tr} + \frac{2}{r} E_{tr}$$

and a momentum conservation law

$$\partial_t E_{tr} = \partial_r \left(\frac{1}{2} E_{tt} + \frac{1}{2} E_{rr} - \frac{1}{r^2} V \right) + \frac{2}{r} E_{rr};$$

multiplying these equations by r^2 and writing E_{tt}, E_{tr}, E_{rr} in terms of $\tilde{E}_{tt}, \tilde{E}_{tr}, \tilde{E}_{rr}$ and \tilde{M} one obtains (after some calculation, as well as (5-12) in the case of (5-15)) the slightly simpler equations

$$\partial_t \left(\frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} + V \right) = \partial_r \tilde{E}_{tr} \quad (5-14)$$

and

$$\partial_t \tilde{E}_{tr} = \partial_r \left(\frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} - V \right) - \frac{p-1}{r} V. \quad (5-15)$$

The expressions in (5-14) are even, while the expressions in (5-15) are odd. Thus we may combine these equations into a single equation by adding them together, which after some rearranging becomes the transport-type equation

$$(\partial_t - \partial_r) e_+ + (\partial_t + \partial_r) V = -\frac{p-1}{r} V, \quad (5-16)$$

where e_+ is the null energy density

$$e_+ := \frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} + \tilde{E}_{tr}. \quad (5-17)$$

Remark 5.3. It may be instructive to derive these equations in the specific context of a solution $u : \Gamma_3 \rightarrow \mathbb{R}$ to the scalar defocusing NLW

$$\square u = |u|^{p-1}u,$$

which in polar coordinates becomes

$$-\partial_{tt}u + \partial_{rr}u + \frac{2}{r}u = |u|^{p-1}u.$$

Making the change of variables $\phi = ru$, this becomes

$$-\partial_{tt}\phi + \partial_{rr}\phi = \frac{|\phi|^{p-1}\phi}{r^{p-1}}.$$

Introducing the null energy

$$e_+ := \frac{1}{2}|\partial_t\phi + \partial_r\phi|^2$$

and the potential energy

$$V := \frac{1}{p+1} \frac{|\phi|^{p+1}}{r^{p-1}}$$

as well as the additional densities

$$\tilde{M} := |\phi|^2, \quad \tilde{E}_{tt} := |\partial_t\phi|^2, \quad \tilde{E}_{rr} := |\partial_r\phi|^2, \quad \tilde{E}_{tr} := \partial_t\phi\partial_r\phi,$$

one can readily verify the identities (5-12), (5-16), and (5-17). It is similar for the other properties of \tilde{M} , \tilde{E}_{tt} , \tilde{E}_{rr} , \tilde{E}_{tr} identified in this section.

Finally, we translate the positive definiteness of (5-5) (when $r \neq 0$) and (5-6) (when $r = 0$) into conditions involving the fields \tilde{M} , \tilde{E}_{tt} , \tilde{E}_{rr} , \tilde{E}_{tr} . From the identity

$$\begin{pmatrix} \tilde{M} & \frac{1}{2}\partial_t\tilde{M} & \frac{1}{2}\partial_r\tilde{M} \\ \frac{1}{2}\partial_t\tilde{M} & \tilde{E}_{tt} & \tilde{E}_{tr} \\ \frac{1}{2}\partial_r\tilde{M} & \tilde{E}_{tr} & \tilde{E}_{rr} \end{pmatrix} = r^2 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{r} & 0 & 1 \end{pmatrix} \begin{pmatrix} M & \frac{1}{2}\partial_t M & \frac{1}{2}\partial_r M \\ \frac{1}{2}\partial_t M & E_{tt} & E_{tr} \\ \frac{1}{2}\partial_r M & E_{tr} & E_{rr} \end{pmatrix} \begin{pmatrix} 1 & 0 & \frac{1}{r} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

we see (for $r \neq 0$) that (5-5) is strictly positive definite if and only if the matrix

$$\begin{pmatrix} \tilde{M} & \frac{1}{2}\partial_t\tilde{M} & \frac{1}{2}\partial_r\tilde{M} \\ \frac{1}{2}\partial_t\tilde{M} & \tilde{E}_{tt} & \tilde{E}_{tr} \\ \frac{1}{2}\partial_r\tilde{M} & \tilde{E}_{tr} & \tilde{E}_{rr} \end{pmatrix} \quad (5-18)$$

is strictly positive definite. Now we turn to (5-6) when $r = 0$. By homogeneity, it suffices to verify this condition when $(t, r) = (1, 0)$. From (1-1), we have $\partial_t M(1, 0) = -\frac{4}{p-1}M(1, 0)$, so the positive definiteness of (5-6) is equivalent to the condition

$$E_{tt}(1, 0) > \left(\frac{2}{p-1}\right)^2 M(1, 0) > 0,$$

which in terms of $\tilde{E}_{tt}, \tilde{M}$ becomes

$$\partial_{rr} \tilde{E}_{tt}(1, 0) > \left(\frac{2}{p-1}\right)^2 \partial_{rr} \tilde{M}(1, 0) > 0. \quad (5-19)$$

Summarising the above discussion, we now see that Theorem 5.2 is a consequence of the following:

Theorem 5.4 (fifth reduction). *Let $p > 5$. Then there exist smooth even functions $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{rr} : \Gamma_1 \rightarrow \mathbb{R}$ and a smooth odd function $\tilde{E}_{tr} : \Gamma_1 \rightarrow \mathbb{R}$, with \tilde{M} homogeneous of order $\frac{2p-6}{p-1}$ and $\tilde{E}_{tt}, \tilde{E}_{tr}, \tilde{E}_{rr}$ homogeneous of order $-\frac{4}{p-1}$, with $\tilde{M}, \tilde{E}_{tt}$ vanishing to second order at $r = 0$,*

$$\tilde{E}_{tr} - \frac{1}{2r} \partial_t \tilde{M}$$

vanishing to third order, and

$$\tilde{E}_{rr} - \frac{1}{r} \partial_r \tilde{M} + \frac{1}{r^2} \tilde{M}$$

to fourth order. Furthermore, if one defines the fields $V, e_+ : \Gamma_1 \rightarrow \mathbb{R}$ by (5-12) and (5-17), we have the weak defocusing property (5-13) and the null transport equation (5-16). Finally, the matrix (5-18) is strictly positive definite for $r \neq 0$, and for $r = 0$ one has the condition (5-19).

It remains to establish Theorem 5.4. This will be the focus of the final section of the paper.

6. Constructing the mass and energy fields

Fix $p > 5$. We will need a large constant $A > 1$ depending only on p , and then sufficiently small parameter $\delta > 0$ (depending on p, A) to be chosen later. We use the notation $X \lesssim Y$, $Y \gtrsim X$, or $X = O(Y)$ to denote an estimate of the form $|X| \leq CY$, where C can depend on p but is independent of δ, A .

We need to construct smooth fields $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{rr}, \tilde{E}_{tr} : \Gamma_1 \rightarrow \mathbb{R}$ which generate some further fields $V, e_+ : \Gamma_1 \rightarrow \mathbb{R}$, which are all required to obey a certain number of constraints. The problem is rather underdetermined, and so there will be some flexibility in selecting these fields; most of these fields will end up being concentrated in the region $\{(t, r) \in \Gamma_1 : r = (\pm 1 + O(\delta))t\}$ near the boundary of the light cone. Given that the constraint (5-16) only involves the two fields V and e_+ , it is natural to proceed by constructing V and e_+ first. In fact we will proceed as follows.

Selection of e_+ in the left half of the cone. We begin by making a choice for the function $e_+ : \Gamma_1 \rightarrow \mathbb{R}$ in the left half $\Gamma_1^l := \{(t, r) \in \Gamma_1 : r \leq 0\}$ of the cone. When $t = 1$, we choose $e_+(1, r)$ to be a smooth function with the following properties:

- One has

$$e_+(1, r) = (1+r)^{-\frac{4}{p-1}} \quad (6-1)$$

for $-1 + \delta \leq r \leq 0$.

- One has

$$e_+(1, r) \geq (1+r)^{-\frac{4}{p-1}} \quad (6-2)$$

for $-1 + \frac{1}{2}\delta \leq r \leq -1 + \delta$. Furthermore, one has

$$\int_{-1+\frac{1}{2}\delta}^{-1+\delta} e_+(1, r) dr \geq A\delta^{1-\frac{4}{p-1}}. \quad (6-3)$$

- One has

$$\delta^{-\frac{4}{p-1}} \lesssim e_+(1, r) \lesssim A\delta^{-\frac{4}{p-1}} \quad (6-4)$$

and

$$\left| \frac{d}{dr} e_+(1, r) \right| \lesssim A\delta^{-\frac{p+3}{p-1}} \quad (6-5)$$

for $-1 \leq r \leq -1 + \delta$.

Clearly we can find a smooth function $r \rightarrow e_+(1, r)$ on $[-1, 0]$ with these properties. We then extend e_+ to the entire left half Γ_1^I of the cone by requiring it to be homogeneous of order $-\frac{4}{p-1}$; thus

$$e_+(t, r) := t^{-\frac{4}{p-1}} e_+\left(1, \frac{r}{t}\right). \quad (6-6)$$

In particular, e_+ is smooth on this half of the cone, and we have

$$e_+(t, r) = (t+r)^{-\frac{4}{p-1}}$$

for $-(1-\delta)t \leq r \leq 0$.

The properties (6-1)–(6-5) are largely used to ensure that the potential energy V that we will construct below is nonnegative.

Selection of V in the left half of the cone. Once e_+ has been selected on Γ_1^I , we construct V on Γ_1^I by solving (5-16), or more explicitly by the formula

$$V(t, r) := \frac{1}{2|r|^{p-1}} \int_r^0 |s|^{p-1} ((\partial_t - \partial_r)e_+)(t-r+s, s) ds \quad (6-7)$$

for $-t \leq r < 0$. Note that as $(\partial_t - \partial_r)e_+$ vanishes for $-(1-\delta)t < r < 0$, the potential energy V vanishes on this region also, and so one can smoothly extend V to all of Γ_1^I . It is easy to see that V is homogeneous of order $-\frac{4}{p-1}$. From the fundamental theorem of calculus and the chain rule, we have

$$(\partial_t + \partial_r)(|r|^{p-1}V) = |r|^{p-1}(\partial_t - \partial_r)e_+$$

for $-t \leq r < 0$, and hence by the product rule we see that (5-16) is obeyed for $-t \leq r < 0$, and hence to all of Γ_1^I by smoothness. We have already seen that V vanishes in the region $-(1-\delta)t < r \leq 0$. In the region $-t \leq r \leq -(1-\delta)t$, we have the following estimate and nonnegativity property:

Proposition 6.1. *For $-t \leq r \leq -(1-\delta)t$, we have*

$$0 \leq V(t, r) \lesssim At^{-\frac{4}{p-1}} \delta^{\frac{p-5}{p-1}}.$$

We remark that to get the lower bound $V(t, r)$, the supercriticality hypothesis $p > 5$ will be crucial.

Proof. By homogeneity we may assume that $t-r=2$, so that $t=1-O(\delta)$ and $r=-1+O(\delta)$, and it will suffice to show that

$$0 \leq V(t, r) \lesssim A\delta^{\frac{p-5}{p-1}}. \quad (6-8)$$

Write $e_+(t, r) = (t+r)^{-\frac{4}{p-1}} + f(t, r)$; then from (6-7) we have

$$V(t, r) = \frac{1}{2|r|^{p-1}} \int_r^0 |s|^{p-1} ((\partial_t - \partial_r)f)(2+s, s) ds. \quad (6-9)$$

The function f is homogeneous of order $-\frac{4}{p-1}$; hence by (1-2)

$$(t\partial_t + t\partial_r)f = -\frac{4}{p-1}f.$$

From the identity

$$\partial_t - \partial_r = -\frac{t+r}{t-r}(\partial_t + \partial_r) + \frac{2}{t-r}(t\partial_t + t\partial_r)$$

and the chain rule, we thus have

$$((\partial_t - \partial_r)f)(2+s, s) = -(1+s)\frac{d}{ds}f(2+s, s) - \frac{4}{p-1}f(2+s, s).$$

Inserting this into (6-9) and integrating by parts, we conclude that

$$V(t, r) = \frac{1+r}{2}f(t, r) + \frac{1}{2|r|^{p-1}} \int_r^0 \frac{d}{ds}(|s|^{p-1}(1+s))f(2+s, s) - |s|^{p-1} \frac{4}{p-1}f(2+s, s) ds,$$

which by the product rule is equal to

$$V(t, r) = \frac{1+r}{2}f(t, r) + \frac{1}{2|r|^{p-1}} \int_r^0 |s|^{p-1} \left[\frac{p-5}{p-1} + \frac{(p-1)(1+s)}{s} \right] f(2+s, s) ds. \quad (6-10)$$

Note that $f(2+s, s)$ is only nonzero when $s = -1 + O(\delta)$, in which case it is of size $O(A\delta^{-\frac{4}{p-1}})$ thanks to (6-2) and (6-4). This gives the upper bound in (6-8). Now we turn to the lower bound. First suppose that $-(1 - \frac{1}{2}\delta)t \leq r$; then f is nonnegative in all of its appearances in (6-10). As we are in the supercritical case $p > 5$, the factor

$$\frac{p-5}{p-1} + \frac{(p-1)(1+s)}{s}$$

is positive (indeed it is $\gtrsim 1$) for δ small enough, and the claim follows in this case.

It remains to consider the case when $-t \leq r \leq -(1 - \frac{1}{2}\delta)t$. In this case we can use the lower bound

$$f(t, r) \geq -(t+r)^{-\frac{4}{\delta}}$$

and conclude that the term $\frac{1}{2}(1+r)f(t, r)$ is at least $-O(\delta^{\frac{p-5}{p-1}})$. A similar argument shows that the contribution to (6-10) coming from those s with

$$-(2+s) \leq s \leq -(1 - \frac{1}{2}\delta)(2+s)$$

is at least $-O(\delta^{\frac{p-5}{p-1}})$. On the other hand, from (6-3), the contribution of those s with

$$s > -(1 - \frac{1}{2}\delta)(2+s)$$

is $\gtrsim (A - O(1))\delta^{\frac{p-5}{p-1}}$. As A is assumed to be large, the claim follows. \square

On the support of V in Γ_1^l , we see from (6-5) and (6-6) that

$$(\partial_t - \partial_r)e_+ = O(At^{-\frac{p+3}{p-1}}\delta^{-\frac{4}{p-1}})$$

and hence by (5-16) and Proposition 6.1,

$$(\partial_t + \partial_r)V = O(At^{-\frac{p+3}{p-1}}\delta^{-\frac{4}{p-1}}). \quad (6-11)$$

Selection of V in the right half of the cone. Once V has been constructed in the left half Γ_1^l of the light cone, we extend it to the right half $\Gamma_1^r := \{(t, r) \in \Gamma_1 : r \geq 0\}$ by even extension; thus

$$V(t, r) := V(t, -r)$$

for all $(t, r) \in \Gamma_1^r$. Since V vanished for $-(1-\delta)t \leq r \leq 0$, we see that V is smooth on all of Γ_1 , and vanishing in the interior cone $\{(t, r) \in \Gamma_1 : |r| \leq (1-\delta)t\}$. It also obeys the nonnegativity property (5-13). From reflecting (6-11) and Proposition 6.1 we have the bounds

$$V = O(At^{-\frac{4}{p-1}}\delta^{\frac{p-5}{p-1}}) \quad (6-12)$$

and

$$(\partial_t - \partial_r)V = O(At^{-\frac{p+3}{p-1}}\delta^{-\frac{4}{p-1}}) \quad (6-13)$$

when $(1-\delta)t \leq r \leq t$.

Selection of e_+ in the right half of the cone. Thus far, V has been defined on all of Γ_1 , and e_+ defined on Γ_1^l . We now extend e_+ to Γ_1^r by solving (5-16), or more precisely by setting

$$e_+(t, r) := e_+(t+r, 0) + \int_0^r ((\partial_t + \partial_r)V)(t+r-s, s) + \frac{p-1}{s}V(t+r-s, s) ds \quad (6-14)$$

for $0 < r \leq t$; note that the integral is well-defined since V vanishes near the time axis. One easily checks that $e_+(t, r) = (t+r)^{-\frac{4}{p-1}}$ for $0 \leq r \leq (1-\delta)t$, and so e_+ extends smoothly to all of Γ_1 and is equal to $(t+r)^{-\frac{4}{p-1}}$ in the interior cone $\{(t, r) \in \Gamma_1 : |r| \leq (1-\delta)t\}$. It is also clear from construction that e_+ is homogeneous of order $-\frac{4}{p-1}$. From the fundamental theorem of calculus we see that e_+ and V obey (5-16) on Γ_1^r , and hence on all of Γ_1 . From (6-12) and (6-13) we see that the integrand is of size $O(At^{-\frac{p+3}{p-1}}\delta^{-\frac{4}{p-1}})$ when $r = (1-O(\delta))t$, and vanishes otherwise, which leads (for δ small enough) to the crude upper and lower bounds

$$t^{-\frac{p+3}{p-1}} \lesssim e_+(t, r) \lesssim t^{-\frac{p+3}{p-1}} \quad (6-15)$$

throughout Γ_1^r .

Selection of e_- and \tilde{E}_{tr} . We reflect the function e_+ around the time axis to create a new function $e_- : \Gamma_1 \rightarrow \mathbb{R}$:

$$e_-(t, r) := e_+(t, r).$$

Like e_+ , the function e_- is smooth and homogeneous of order $-\frac{4}{p-1}$. It equals $(t-r)^{-\frac{4}{p-1}}$ in the interior cone $\{(t, r) \in \Gamma_1 : |r| \leq (1-\delta)t\}$. On Γ_1^l it obeys the crude upper and lower bounds

$$t^{-\frac{p+3}{p-1}} \lesssim e_-(t, r) \lesssim t^{-\frac{p+3}{p-1}} \quad (6-16)$$

and in the region $(1 - \delta)t \leq r \leq t$ we have the bounds

$$(\delta t)^{-\frac{4}{p-1}} \lesssim e_-(t, r) \lesssim A(\delta t)^{-\frac{4}{p-1}} \quad (6-17)$$

thanks to (6-4).

Recall from (5-17) that the field e_+ is intended to ultimately be of the form $\frac{1}{2}\tilde{E}_{tt} + \frac{1}{2}\tilde{E}_{rr} + \tilde{E}_{tr}$. Similarly, e_- is intended to be of the form

$$e_- = \frac{1}{2}\tilde{E}_{tt} + \frac{1}{2}\tilde{E}_{rr} - \tilde{E}_{tr}. \quad (6-18)$$

Accordingly, we may now define \tilde{E}_{tr} as

$$\tilde{E}_{tr} := \frac{e_+ - e_-}{2}. \quad (6-19)$$

This is clearly smooth, odd, and homogeneous of order $-\frac{4}{p-1}$. We also see that the quantity $\tilde{E}_{tt} + \tilde{E}_{rr}$ is now specified:

$$\tilde{E}_{tt} + \tilde{E}_{rr} = e_+ + e_-. \quad (6-20)$$

We are left with two remaining unknown scalar fields to specify: the mass density \tilde{M} and the energy equipartition $-\tilde{E}_{tt} + \tilde{E}_{rr}$, which determines the fields \tilde{E}_{tt} and \tilde{E}_{rr} by (6-20). The requirements needed for Theorem 5.4 that have not already been verified are as follows:

- \tilde{M} is smooth, even, and homogeneous of order $\frac{2p-6}{p-1}$, and $-\tilde{E}_{tt} + \tilde{E}_{rr}$ is smooth, even, and homogeneous of order $-\frac{4}{p-1}$.
- $\tilde{M}, \tilde{E}_{tt}$ vanish to second order at $r = 0$, $\tilde{E}_{tr} - \frac{1}{2r}\partial_t\tilde{M}$ vanishes to third order, and $\tilde{E}_{rr} - \frac{1}{r}\partial_r\tilde{M} + \frac{1}{r^2}\tilde{M}$ to fourth order.
- One has the equations (5-12) and (5-17) (and hence also (6-18)).
- The matrix (5-18) is strictly positive definite for $r \neq 0$, and for $r = 0$ one has the condition (5-19).

As there is only one equation (beyond homogeneity and reflection symmetry) constraining \tilde{M} and $-\tilde{E}_{tt} + \tilde{E}_{rr}$ — namely, (5-12) — the problem of selecting these two fields is underdetermined, and thus subject to a certain amount of arbitrary choices. We will select these fields first in the exterior region $\{(t, r) \in \Gamma : |r| \geq \frac{t}{2}\}$, and then fill in the interior using a different method.

Selection of $\tilde{M}, -\tilde{E}_{tt} + \tilde{E}_{rr}$ away from the time axis. In the exterior region $\{(t, r) \in \Gamma : |r| \geq \frac{t}{2}\}$, we shall simply select the field \tilde{M} to be a small but otherwise rather arbitrary field, and then use (5-12) to determine $-\tilde{E}_{tt} + \tilde{E}_{rr}$.

More precisely, let $\tilde{M}(1, r)$ be a smooth even function on the region $\{r : \frac{1}{2} \leq |r| \leq 1\}$ obeying the following properties:

- For $\frac{1}{2} \leq |r| \leq \frac{3}{4}$, one has

$$\tilde{M}(1, r) = \delta\left((1+r)^{\frac{2p-6}{p-1}} + (1-r)^{\frac{2p-6}{p-1}}\right). \quad (6-21)$$

(This condition will not be used directly in this part of the construction, but is needed for compatibility with the next part.)

- For $\frac{1}{2} \leq |r| \leq 1$, one has the bounds

$$\delta \lesssim \tilde{M}(1, r) \lesssim \delta \quad (6-22)$$

and

$$\frac{d}{dr} \tilde{M}(1, r), \frac{d^2}{dr^2} \tilde{M}(1, r) = O(\delta). \quad (6-23)$$

It is clear that one can select such a function. We then extend \tilde{M} to $\{(t, r) \in \Gamma_1 : |r| \geq \frac{t}{2}\}$ by requiring that \tilde{M} be homogeneous of order $\frac{2p-6}{p-1}$. Then \tilde{M} is smooth and even, and one has the bounds

$$\delta t^{\frac{2p-6}{p-1}} \lesssim \tilde{M}(t, r) \lesssim \delta t^{\frac{2p-6}{p-1}}, \quad (6-24)$$

$$\frac{d}{dr} \tilde{M}(t, r), \frac{d}{dt} \tilde{M}(t, r) = O(\delta t^{\frac{p-5}{p-1}}), \quad (6-25)$$

$$\frac{d^2}{dr^2} \tilde{M}(t, r), \frac{d^2}{dt^2} \tilde{M}(t, r) = O(\delta t^{-\frac{4}{p-1}}) \quad (6-26)$$

in the region $\{(t, r) \in \Gamma_1 : |r| \geq \frac{t}{2}\}$.

We then define $-\tilde{E}_{tt} + \tilde{E}_{rr}$ on this region by enforcing (5-12); thus

$$-\tilde{E}_{tt} + \tilde{E}_{rr} := \frac{1}{2}(-\partial_{tt} \tilde{M} + \partial_{rr} \tilde{M}) - (p+1)V. \quad (6-27)$$

Combining this with (6-20), this defines \tilde{E}_{tt} and \tilde{E}_{rr} . It is easy to see that these fields are smooth, even and homogeneous of order $-\frac{4}{p-1}$ on $\{(t, r) \in \Gamma_1 : |r| \geq \frac{t}{2}\}$.

We now claim that the matrix (5-18) is strictly positive definite in the region $\{(t, r) \in \Gamma_1 : |r| \geq \frac{t}{2}\}$. By homogeneity and reflection symmetry, it suffices to verify this when $t = 1$ and $\frac{1}{2} \leq r \leq 1$. Using the identity

$$\begin{pmatrix} \tilde{M} & \frac{1}{2}(\partial_t + \partial_r)\tilde{M} & \frac{1}{2}(\partial_t - \partial_r)\tilde{M} \\ \frac{1}{2}(\partial_t + \partial_r)\tilde{M} & 2e_+ & -\tilde{E}_{tt} + \tilde{E}_{rr} \\ \frac{1}{2}(\partial_t - \partial_r)\tilde{M} & -\tilde{E}_{tt} + \tilde{E}_{rr} & 2e_- \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} \tilde{M} & \frac{1}{2}\partial_t \tilde{M} & \frac{1}{2}\partial_r \tilde{M} \\ \frac{1}{2}\partial_t \tilde{M} & \tilde{E}_{tt} & \tilde{E}_{tr} \\ \frac{1}{2}\partial_r \tilde{M} & \tilde{E}_{tr} & \tilde{E}_{rr} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & 1 \end{pmatrix},$$

it suffices to show that the matrix

$$\begin{pmatrix} \tilde{M} & \frac{1}{2}(\partial_t + \partial_r)\tilde{M} & \frac{1}{2}(\partial_t - \partial_r)\tilde{M} \\ \frac{1}{2}(\partial_t + \partial_r)\tilde{M} & 2e_+ & -\tilde{E}_{tt} + \tilde{E}_{rr} \\ \frac{1}{2}(\partial_t - \partial_r)\tilde{M} & -\tilde{E}_{tt} + \tilde{E}_{rr} & 2e_- \end{pmatrix}$$

is strictly positive definite.

If $r \leq 1 - \delta$, then all off-diagonal terms are $O(\delta)$ thanks to (6-23) and (6-27), while the diagonal terms are $\gtrsim \delta$, $\gtrsim 1$, and $\gtrsim 1$ respectively, and the positive definiteness is easily verified, since the associated quadratic form is at least

$$\gtrsim \delta x_1^2 + x_2^2 + x_3^2 - O(\delta|x_1||x_2|) - O(\delta|x_1||x_3|) - O(\delta|x_2||x_3|),$$

which is easily seen to be positive for δ small enough. If $r < 1 - \delta$, then the off-diagonal terms are $O(\delta)$ in the top row and left column, and $O(A\delta^{\frac{p-5}{p-1}})$ in the bottom right minor by (6-12), while the diagonal

terms are $\gtrsim \delta$, $\gtrsim 1$, and $\gtrsim \delta^{-\frac{4}{p-1}}$ by (6-22), (6-15) and (6-17), so the associated quadratic form is

$$\gtrsim \delta x_1^2 + x_2^2 + \delta^{-\frac{4}{p-1}} x_3^2 - O(\delta|x_1||x_2|) - O(\delta|x_1||x_3|) - O(A\delta^{\frac{p-5}{p-1}}|x_2||x_3|),$$

which is again positive definite (note that $A\delta^{\frac{p-5}{p-1}}$ can be chosen to be much smaller than the geometric mean of δ and $\delta^{-\frac{4}{p-1}}$).

Selection of $M, \tilde{E}_{tt}, \tilde{E}_{rr}$ near the time axis. Now we restrict attention to the interior region $\Gamma_1^i := \{(t, r) \in \Gamma_1 : |r| \leq \frac{t}{2}\}$; all identities and estimates here are understood to be on this region unless otherwise specified.

We will now reverse the Gram matrix reduction from previous sections, and construct $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{rr}$ in Γ_1^i from an (infinite-dimensional) vector-valued solution to the (free, (1+1)-dimensional) wave equation. Let H be a Hilbert space and let $t \mapsto f(t)$ be a family of vectors $f(t)$ in H smoothly parameterised by a parameter $t \in (0, +\infty)$ (so that all derivatives in t exist in the strong sense and are continuous); we will select this family more precisely later. We introduce the smooth vector-valued field $\phi : \Gamma_1^i \rightarrow H$ by the formula

$$\phi(t, r) := f(t + r) - f(t - r)$$

and we will define $\tilde{M}, \tilde{E}_{tt}, \tilde{E}_{rr} : \Gamma_1^i \rightarrow \mathbb{R}$ by the formulae

$$\begin{aligned} \tilde{M}(t, r) &:= \langle \phi(t, r), \phi(t, r) \rangle_H, \\ \tilde{E}_{tt}(t, r) &:= \langle \partial_t \phi(t, r), \partial_t \phi(t, r) \rangle_H, \\ \tilde{E}_{rr}(t, r) &:= \langle \partial_r \phi(t, r), \partial_r \phi(t, r) \rangle_H. \end{aligned}$$

Since ϕ is smooth and odd in r , these functions are smooth and even in r . If we impose the additional hypothesis that the Gram matrix $\langle f(s), f(t) \rangle_H$ has the scaling symmetry

$$\langle f(\lambda s), f(\lambda t) \rangle_H = \lambda^{\frac{2p-6}{p-1}} \langle f(s), f(t) \rangle_H \tag{6-28}$$

for $s, t, \lambda > 0$, then M will be homogeneous of order $\frac{2p-6}{p-1}$; furthermore, by differentiating (6-28) with respect to both s and t we see that

$$\langle f'(\lambda s), f'(\lambda t) \rangle_H = \lambda^{-\frac{4}{p-1}} \langle f'(s), f'(t) \rangle_H \tag{6-29}$$

(where f' denotes the derivative of f) and so $\tilde{E}_{tt}, \tilde{E}_{rr}$ will be homogeneous of order $-\frac{4}{p-1}$.

Observe that

$$\begin{aligned} \frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} + \langle \partial_t \phi, \partial_r \phi \rangle_H &= \frac{1}{2} \|(\partial_t + \partial_r)\phi\|_H^2 \\ &= 2 \|f'(t+r)\|_H^2 \end{aligned}$$

and similarly

$$\frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} - \langle \partial_t \phi, \partial_r \phi \rangle_H = 2 \|f'(t-r)\|_H^2.$$

Thus, if we impose the additional normalisation

$$\|f'(1)\|_H = \frac{1}{\sqrt{2}}, \tag{6-30}$$

and hence by (6-29),

$$\|f'(t)\|_H = \frac{1}{\sqrt{2}} t^{-\frac{2}{p-1}}, \quad (6-31)$$

we see from the identities $e_{\pm}(t, r) = (t \pm r)^{-\frac{4}{p-1}}$ in Γ_1^i that

$$\frac{1}{2} \tilde{E}_{tt} + \frac{1}{2} \tilde{E}_{rr} \pm \langle \partial_t \phi, \partial_r \phi \rangle_H = e_{\pm}.$$

In particular, (6-20) holds, and from (6-19) one has

$$\tilde{E}_{tr} = \langle \partial_t \phi, \partial_r \phi \rangle_H.$$

We also obtain the equations (5-17) and (6-18).

Next, it is clear that ϕ solves the wave equation

$$-\partial_{tt} \phi + \partial_{rr} \phi = 0,$$

so in particular

$$\langle \phi, -\partial_{tt} \phi + \partial_{rr} \phi \rangle_H = 0,$$

which implies in particular (cf. (4-5)) that

$$\frac{1}{2} (-\partial_{tt} \tilde{M} + \partial_{rr} \tilde{M}) + \tilde{E}_{tt} - \tilde{E}_{rr} = 0.$$

Since V vanishes on Γ_1 , we conclude that (5-12) holds.

Next, from differentiating the formula for \tilde{M} , one has

$$\frac{1}{2} \partial_t \tilde{M} = \langle \phi, \partial_t \phi \rangle_H$$

and

$$\frac{1}{2} \partial_r \tilde{M} = \langle \phi, \partial_r \phi \rangle_H$$

and so the quadratic form associated with (5-18) factorises as

$$\|x_1 \phi + x_2 \partial_t \phi + x_3 \partial_r \phi\|_H^2.$$

This is clearly positive semidefinite at least; to make it positive definite for $r \neq 0$, it will suffice to enforce the condition

$$f(s), f(t), f'(s), f'(t) \text{ linearly independent} \quad (6-32)$$

for all distinct $s, t > 0$.

Suppose we assume the long-range orthogonality condition

$$\langle f(s), f(t) \rangle_H = 0 \quad (6-33)$$

whenever $\frac{t}{s} > 1.1$ or $\frac{s}{t} > 1.1$. Then in the region $\{(t, r) \in \Gamma_1^i : |r| \geq \frac{t}{4}\}$ away from the time axis, we have from Pythagoras' theorem that

$$\tilde{M}(t, r) = \|f(t+r)\|_H^2 + \|f(t-r)\|_H^2.$$

In particular, if we also impose the normalisation

$$\|f(1)\|_H = \sqrt{\delta} \quad (6-34)$$

then (from (6-28)) we have

$$\tilde{M}(t, r) = \delta \left((t+r)^{\frac{2p-6}{p-1}} + (t-r)^{\frac{2p-6}{p-1}} \right)$$

in the region $\{(t, r) \in \Gamma_1^i : |r| \geq \frac{t}{4}\}$. In particular from (6-21) and homogeneity we see that \tilde{M} on Γ_1^i joins up smoothly with its counterpart in the exterior region $\{(t, r) \in \Gamma_1 : |r| \geq \frac{t}{2}\}$; by (5-12) we see that $-\tilde{E}_{tt} + \tilde{E}_{rr}$ does too. By (6-20) and (6-19) we now see that all of the fields M , \tilde{E}_{tt} , \tilde{E}_{rr} , \tilde{E}_{tr} are smooth on all of Γ_1 .

Now we study the vanishing properties of the various fields constructed at $r = 0$ for a fixed value of t . From Taylor expansion we have

$$\phi(t, r) = 2rf'(t) + \frac{1}{3}r^3 f'''(t) + O(|r|^5)$$

as $r \rightarrow 0$ (where the error term denotes a quantity in H of norm $O(|r|^5)$, and the implied constant can depend on t and ϕ). Furthermore, these asymptotics behave in the expected fashion with respect to differentiation in time or space; thus for instance

$$\begin{aligned} \partial_r \phi(t, r) &= 2f'(t) + r^2 f'''(t) + O(|r|^4), \\ \partial_t \phi(t, r) &= 2rf''(t) + \frac{1}{3}r^3 f^{(4)}(t) + O(|r|^5). \end{aligned}$$

Taking inner products, we conclude the asymptotics

$$\begin{aligned} \tilde{M}(t, r) &= 4r^2 \|f'(t)\|_H^2 + \frac{4}{3}r^4 \langle f'(t), f'''(t) \rangle_H + O(|r|^6), \\ \tilde{E}_{tt}(t, r) &= 4r^2 \|f''(t)\|_H^2 + O(|r|^4), \\ \tilde{E}_{tr}(t, r) &= 4r \langle f'(t), f''(t) \rangle_H + O(|r|^3), \\ \tilde{E}_{rr} &= 4 \|f'(t)\|_H^2 + 4r^2 \langle f'(t), f'''(t) \rangle_H + O(|r|^4). \end{aligned}$$

The asymptotic for \tilde{M} behaves well with respect to derivatives; thus for instance

$$\begin{aligned} \partial_t \tilde{M}(t, r) &= 8r^2 \langle f'(t), f''(t) \rangle_H + O(|r|^4), \\ \partial_r \tilde{M}(t, r) &= 8r \|f'(t)\|_H^2 + \frac{16}{3}r^3 \langle f'(t), f'''(t) \rangle_H + O(|r|^5). \end{aligned}$$

Among other things, this shows (using (6-30)) that the condition (5-19) reduces to

$$\|f''(1)\|_H > \frac{2}{p-1} \frac{1}{\sqrt{2}}. \quad (6-35)$$

It is also clear from these asymptotics that \tilde{E} and \tilde{E}_{tt} vanish to second order, and $\tilde{E}_{tr} - \frac{1}{2r} \partial_t \tilde{M}$ vanishes to third order; a brief calculation also shows that $\tilde{E}_{rr} - \frac{1}{r} \partial_r \tilde{M} + \frac{1}{r^2} \tilde{M}$ vanishes to fourth order.

To summarise: in order to conclude all the required properties for Theorem 5.4, it suffices to locate a smooth curve $t \mapsto f(t)$ in a Hilbert space H which obeys the hypotheses (6-28), (6-30), (6-32), (6-33), (6-34) and (6-35).

We take the Hilbert space H to be the space $L^2(\mathbb{R})$ of square-integrable real-valued functions on \mathbb{R} with Lebesgue measure. The functions $f(t) \in H$ will take the form

$$f(t)(x) := t^{\frac{p-3}{p-1}} \psi(x - \log t),$$

where $\psi : \mathbb{R} \rightarrow \mathbb{R}$ is a bump function whose (closed) support is precisely $[0, 0.01]$ (that is to say, the set $\{\psi \neq 0\}$ is a dense subset of $[0, 0.01]$) depending on δ and p to be chosen shortly. It is clear from construction that (6-28) and (6-33) hold. The condition (6-34) becomes

$$\int_{\mathbb{R}} \psi(x)^2 dx = \delta,$$

while the condition (6-30) becomes

$$\int_{\mathbb{R}} \psi'(x)^2 dx = \frac{1}{2}.$$

It is easy to see that we can select ψ with closed support precisely $[0, 0.01]$ with both of these normalisations, basically because the Dirichlet form $\langle \phi', \psi' \rangle$ is unbounded on $L^2([0, 0.01])$.

Now we verify the linear independence claim (6-32). We may assume without loss of generality that $s = 1$ and $t > 1$. Then we have a linear dependence between ψ and ψ' in a neighbourhood of 0; since ψ, ψ' vanish to the left of 0, the Picard uniqueness theorem for ODEs then implies that ψ vanishes a little to the right of 0 also, contradicting the hypothesis that ψ has closed support containing 0. This gives (6-32).

A similar argument shows that $f'(1)$ and $f''(1)$ are linearly independent. Squaring and differentiating (6-31) at $t = 1$ gives

$$\langle f'(1), f''(1) \rangle_H = -\frac{2}{p-1} \frac{1}{2}$$

and (6-35) then follows from (6-30) and the Cauchy–Schwarz inequality, using the linear independence to get the strict inequality. This (finally) completes the proof of Theorem 5.4 and hence Theorem 1.1.

Acknowledgments

The author is supported by NSF grant DMS-1266164 and by a Simons Investigator Award. The author is also indebted to Sergiu Klainerman for suggesting this question, and Yi Jin Gao for corrections.

References

- [Brenner and Kumlin 2000] P. Brenner and P. Kumlin, “On wave equations with supercritical nonlinearities”, *Arch. Math. (Basel)* **74**:2 (2000), 129–147. MR 1735230 Zbl 0971.35051
- [Brenner and von Wahl 1981] P. Brenner and W. von Wahl, “Global classical solutions of nonlinear wave equations”, *Math. Z.* **176**:1 (1981), 87–121. MR 606174 Zbl 0457.35059
- [Burq et al. 2007] N. Burq, S. Ibrahim, and P. Gérard, “Instability results for nonlinear Schrödinger and wave equations”, preprint, 2007.
- [Cazenave et al. 1998] T. Cazenave, J. Shatah, and A. S. Tahvildar-Zadeh, “Harmonic maps of the hyperbolic space and development of singularities in wave maps and Yang–Mills fields”, *Ann. Inst. H. Poincaré Phys. Théor.* **68**:3 (1998), 315–349. MR 1622539 Zbl 0918.58074
- [Christ et al. 2003] M. Christ, J. Colliander, and T. Tao, “Ill-posedness for nonlinear Schrödinger and wave equations”, preprint, 2003. arXiv math/0311048

- [Grillakis 1990] M. G. Grillakis, “Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity”, *Ann. of Math.* (2) **132**:3 (1990), 485–509. MR 1078267 Zbl 0736.35067
- [Grillakis 1992] M. G. Grillakis, “Regularity for the wave equation with a critical nonlinearity”, *Comm. Pure Appl. Math.* **45**:6 (1992), 749–774. MR 1162370 Zbl 0785.35065
- [Günther 1991] M. Günther, “Isometric embeddings of Riemannian manifolds”, pp. 1137–1143 in *Proceedings of the International Congress of Mathematicians, Vol. II* (Kyoto, 1990), Math. Soc. Japan, Tokyo, 1991. MR 1159298 Zbl 0745.53031
- [Ibrahim et al. 2011] S. Ibrahim, M. Majdoub, and N. Masmoudi, “Well- and ill-posedness issues for energy supercritical waves”, *Anal. PDE* **4**:2 (2011), 341–367. MR 2859857 Zbl 1267.35131
- [Jörgens 1961] K. Jörgens, “Das Anfangswertproblem im Grossen für eine Klasse nichtlinearer Wellengleichungen”, *Math. Z.* **77** (1961), 295–308. MR 0130462 Zbl 0111.09105
- [Kenig and Merle 2008] C. E. Kenig and F. Merle, “Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation”, *Acta Math.* **201**:2 (2008), 147–212. MR 2461508 Zbl 1183.35202
- [Killip and Visan 2011a] R. Killip and M. Visan, “The defocusing energy-supercritical nonlinear wave equation in three space dimensions”, *Trans. Amer. Math. Soc.* **363**:7 (2011), 3893–3934. MR 2775831 Zbl 1230.35068
- [Killip and Visan 2011b] R. Killip and M. Visan, “The radial defocusing energy-supercritical nonlinear wave equation in all space dimensions”, *Proc. Amer. Math. Soc.* **139**:5 (2011), 1805–1817. MR 2763767 Zbl 1218.35148
- [Lebeau 2001] G. Lebeau, “Non linear optic and supercritical wave equation”, *Bull. Soc. Roy. Sci. Liège* **70**:4–6 (2001), 267–306. MR 1904059 Zbl 1034.35137
- [Lebeau 2005] G. Lebeau, “Perte de régularité pour les équations d’ondes sur-critiques”, *Bull. Soc. Math. France* **133**:1 (2005), 145–157. MR 2145023 Zbl 1071.35020
- [Lindblad and Sogge 1996] H. Lindblad and C. D. Sogge, “Long-time existence for small amplitude semilinear wave equations”, *Amer. J. Math.* **118**:5 (1996), 1047–1135. MR 1408499 Zbl 0869.35064
- [Planchon 2000] F. Planchon, “Self-similar solutions and semi-linear wave equations in Besov spaces”, *J. Math. Pures Appl.* (9) **79**:8 (2000), 809–820. MR 1782103 Zbl 0979.35106
- [Ribaud and Youssfi 2002] F. Ribaud and A. Youssfi, “Global solutions and self-similar solutions of semilinear wave equation”, *Math. Z.* **239**:2 (2002), 231–262. MR 1888223 Zbl 1005.35068
- [Roy 2009] T. Roy, “Global existence of smooth solutions of a 3D log-log energy-supercritical wave equation”, *Anal. PDE* **2**:3 (2009), 261–280. MR 2603799 Zbl 1195.35222
- [Seeley 1964] R. T. Seeley, “Extension of C^∞ functions defined in a half space”, *Proc. Amer. Math. Soc.* **15**:4 (1964), 625–626. MR 0165392 Zbl 0127.28403
- [Segal 1963] I. E. Segal, “The global Cauchy problem for a relativistic scalar field with power interaction”, *Bull. Soc. Math. France* **91** (1963), 129–135. MR 0153967 Zbl 0178.45403
- [Shatah and Struwe 1998] J. Shatah and M. Struwe, *Geometric wave equations*, Courant Lecture Notes in Mathematics **2**, New York University, Courant Institute of Mathematical Sciences, New York, 1998. MR 1674843 Zbl 0993.35001
- [Strauss 1989] W. A. Strauss, *Nonlinear wave equations*, CBMS Regional Conference Series in Mathematics **73**, American Mathematical Society, Providence, RI, 1989. MR 1032250 Zbl 0714.35003
- [Struwe 1988] M. Struwe, “Globally regular solutions to the u^5 Klein–Gordon equation”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **15**:3 (1988), 495–513. MR 1015805 Zbl 0728.35072
- [Tao 2006] T. Tao, *Nonlinear dispersive equations: local and global analysis*, CBMS Regional Conference Series in Mathematics **106**, American Mathematical Society, Providence, RI, 2006. MR 2233925 Zbl 1106.35001
- [Tao 2007] T. Tao, “Global regularity for a logarithmically supercritical defocusing nonlinear wave equation for spherically symmetric data”, *J. Hyperbolic Differ. Equ.* **4**:2 (2007), 259–265. MR 2329385 Zbl 1124.35043
- [Zheng 1991] Y. X. Zheng, “Concentration in sequences of solutions to the nonlinear Klein–Gordon equation”, *Indiana Univ. Math. J.* **40**:1 (1991), 201–235. MR 1101227 Zbl 0714.35079

Received 24 Feb 2016. Revised 6 Aug 2016. Accepted 29 Sep 2016.

TERENCE TAO: tao@math.ucla.edu

Department of Mathematics, UCLA, Los Angeles, CA 90095-1555, United States

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 8 2016

Estimates for radial solutions of the homogeneous Landau equation with Coulomb potential	1773
MARIA PIA GUALDANI and NESTOR GUILLEN	
Forward self-similar solutions of the Navier–Stokes equations in the half space	1811
MIKHAIL KOROBKOV and TAI-PENG TSAI	
Decay of solutions of Maxwell–Klein–Gordon equations with arbitrary Maxwell field	1829
SHIWU YANG	
Invariant distributions and the geodesic ray transform	1903
GABRIEL P. PATERNAIN and HANMING ZHOU	
Multiple vector-valued inequalities via the helicoidal method	1931
CRISTINA BENEÀ and CAMIL MUSCALU	
Structure of modular invariant subalgebras in free Araki–Woods factors	1989
RÉMI BOUTONNET and CYRIL HOUDAYER	
Finite-time blowup for a supercritical defocusing nonlinear wave system	1999
TERENCE TAO	
A long \mathbb{C}^2 without holomorphic functions	2031
LUKA BOČ THALER and FRANC FORSTNERIČ	



2157-5045(2016)9:8;1-4