

ANALYSIS & PDE

Volume 10

No. 4

2017

MAXIME INGREMEAU

DISTORTED PLANE WAVES IN CHAOTIC SCATTERING

DISTORTED PLANE WAVES IN CHAOTIC SCATTERING

MAXIME INGREMEAU

We provide a precise description of distorted plane waves for semiclassical Schrödinger operators under the assumption that the classical trapped set is hyperbolic and that a certain topological pressure (a quantity defined using thermodynamical formalism) is negative. Distorted plane waves are generalized eigenfunctions of the Schrödinger operator which differ from free plane waves, $e^{i(x \cdot \xi)/h}$, by an outgoing term. Under our assumptions we show that they can be written as a convergent sum of Lagrangian states. That provides a description of their semiclassical defect measures in the spirit of quantum ergodicity and extends results of Guillarmou and Naud obtained for hyperbolic quotients to our setting.

1. Introduction

In this paper, we will consider on \mathbb{R}^d a semiclassical Hamiltonian of the form

$$P_h = -h^2 \Delta + V(x), \quad V \in C_c^\infty(\mathbb{R}^d).$$

We will study the “distorted plane waves”, or “scattering states” associated to P_h . They are a family of functions $E_h^\xi \in C^\infty(\mathbb{R}^d)$ with parameter $\xi \in \mathbb{S}^d$ (the direction of propagation of the incoming wave) which are generalized eigenfunctions of P_h ; that is to say, they satisfy the differential equation

$$(P_h - 1)E_h^\xi = 0, \tag{1}$$

but are not in $L^2(\mathbb{R}^d)$ (since P_h has no embedded eigenvalues in \mathbb{R}^+).

These distorted plane waves resemble the actual plane waves, in the sense that we may write

$$E_h^\xi(x) = e^{i/h x \cdot \xi} + E_{\text{out}}^\xi, \tag{2}$$

where E_{out} is outgoing in the sense that it satisfies the Sommerfeld radiation condition:

$$\lim_{|x| \rightarrow \infty} |x|^{(d-1)/2} \left(\frac{\partial}{\partial |x|} - \frac{i}{h} \right) E_{\text{out}}^\xi(x) = 0. \tag{3}$$

One can show (see, for instance, [Melrose 1995, §2; Dyatlov and Zworski 2017, §4]) that for any $\xi \in \mathbb{S}^{d-1}$ and $h > 0$, there exists a unique function E_h^ξ satisfying conditions (1), (2) and (3).

Condition (3) may be equivalently stated by asking that E_{out}^ξ is the image of a function in $C_c^\infty(\mathbb{R}^d)$ by the outgoing resolvent $(P_h - (1 + i0)^2)^{-1}$, or by asking that E_{out}^ξ be of the form

$$E_{\text{out}}^\xi(x) = e^{i|x|/h} |x|^{-(d-1)/2} \left(a_h^\xi(\omega) + O\left(\frac{1}{|x|}\right) \right),$$

MSC2010: 35P20, 35P25, 81Q50.

Keywords: scattering theory, quantum chaos, semiclassical measures, distorted plane waves.

where $\omega = x/|x|$. The function $a_h(\xi, \omega) := a_h^\xi(\omega)$ is called the *scattering amplitude*, and is the integral kernel of the scattering matrix minus identity. The scattering amplitude, and hence the distorted plane waves, are central objects in scattering theory.

The aim of this paper is to discuss the behaviour of distorted plane waves in the semiclassical limit $h \rightarrow 0$. Distorted plane waves can be seen as an analogue, on manifolds of infinite volume, of the eigenfunctions of a Schrödinger operator on a compact manifold. It is therefore natural to ask questions similar to those in the compact case: what can be said about the semiclassical measures of distorted plane waves, about the behaviour of their L^p norms as $h \rightarrow 0$, and about their nodal sets and nodal domains?

The answer to these questions will depend in a drastic way on the properties of the underlying *classical dynamics*. Let us define the classical Hamiltonian by

$$p(x, \xi) = |\xi|^2 + V(x),$$

and the layer of energy 1 as

$$\mathcal{E} = \{\rho \in T^*\mathbb{R}^d : p(\rho) = 1\}.$$

Note that this is a noncompact set, but its intersection with any fibre T_x^*X is compact.

We also denote, for each $t \in \mathbb{R}$, the Hamiltonian flow generated by p by $\Phi^t : T^*\mathbb{R}^d \rightarrow T^*\mathbb{R}^d$. For $\rho \in \mathcal{E}$, we will say that $\rho \in \Gamma^\pm$ if $\{\Phi^t(\rho) : \pm t \leq 0\}$ is a bounded subset of $T^*\mathbb{R}^d$; that is to say, ρ does not “go to infinity”, respectively in the past or in the future. The sets Γ^\pm are called respectively the *outgoing* and *incoming* tails (at energy 1).

The *trapped set* is defined as

$$K := \Gamma^+ \cap \Gamma^-. \tag{4}$$

It is a flow-invariant set, and it is compact, because V is compactly supported.

If the trapped set is empty, then we can easily describe the distorted plane waves in the semiclassical limit. Namely, one can show (see [Dyatlov and Guillarmou 2014, §5.1]) that E_h^ξ is a *Lagrangian (WKB) state*. Furthermore, for any $\chi \in C_c^\infty(\mathbb{R}^d)$, the norm $\|\chi E_h^\xi\|_{L^2}$ is bounded independently of h .

However, if the trapped set is nonempty, the distorted plane waves may not be bounded uniformly in L^2_{loc} as $h \rightarrow 0$. Actually, $\|\chi E_h^\xi\|_{L^2}$ could grow exponentially fast as $h \rightarrow 0$. If we want this quantity to remain bounded uniformly in h , we must therefore make some additional assumptions on the classical dynamics. Let us now detail these assumptions.

Hypotheses on the classical dynamics.

- *Hyperbolicity assumption:* In the sequel, we will suppose that the potential V is such that the trapped set contains no fixed point, and is a *hyperbolic set*. We refer to Section 2.1.2 for the definition of a hyperbolic set. The potential in Figure 1 is an example of such a potential.
- *Topological pressure assumption:* For our result on distorted plane waves to hold, we must also make the assumption (Hypothesis 46) that the topological pressure associated to half the logarithm of the unstable Jacobian of the flow on K is negative. The definition of the topological pressure will be recalled in Section 3.4. Hypothesis 46 roughly says that the system is “very open”. One should note that in dimension 2, this condition is equivalent to the fact that the Hausdorff dimension of K is strictly smaller




Figure 1. An example of a potential on $(\mathbb{R}^2, g_{\text{flat}})$ such that the dynamics is hyperbolic on the trapped set in some energy range. (See [Sjöstrand 1990, Appendix C] for details.)

than 2. In the three-bumps potential of Figure 1, this condition is satisfied if the three bumps are far enough from each other, but it is not satisfied if the bumps are close to each other.

- *Transversality assumption:* Our last assumption does not concern directly the classical dynamics, but the Lagrangian manifold¹

$$\Lambda_\xi := \{(x, \xi) : x \in \mathbb{R}^d\}. \tag{5}$$

Note that the plane wave $e^{i/\hbar x \cdot \xi}$ is a Lagrangian state associated with the Lagrangian manifold Λ_ξ .

We need to make a *transversality assumption* on Λ_ξ . This assumption roughly says that the direction ξ defining Λ_ξ is such that the incoming tail Γ^- and Λ_ξ intersect transversally. We postpone the precise statement of this assumption to Hypothesis 16 in Section 2.1.4. This assumption is probably generic in ξ , although we don't know how to prove it. In [Ingremeau 2017], we show that it is always satisfied for every ξ , when we consider geometric scattering on a manifold of nonpositive curvature.

Statement of the results. In Theorem 47, we will give a precise description of E_h^ξ as a sum of WKB states, under the assumptions above. Since the precise statement of the theorem is a bit technical, we postpone it to Section 3.5, and only state two important consequences of this result.

The first one is a bound analogous to what we would get in the nontrapping case.

Theorem 1. *Suppose that Hypothesis 10 on hyperbolicity holds, that Hypothesis 46 concerning the topological pressure is satisfied, and that $\xi \in \mathbb{S}^{d-1}$ is such that Λ_ξ satisfies Hypothesis 16 of transversality.*

Let $\chi \in C_c^\infty(X)$. Then there exists a constant $C_{\xi, \chi}$ independent of h such that for any $h > 0$, we have

$$\|\chi E_h^\xi\|_{L^2} \leq C_{\xi, \chi}. \tag{6}$$

Remark 2. The bound (6) could not be obtained directly from resolvent estimates. Indeed, as we will see in Section 3.3.2, the term E_{out} in (2) can be written as the outgoing resolvent $(P_h - (1 + i0)^2)^{-1}$

¹By a Lagrangian manifold, we mean a d -dimensional submanifold of a $2d$ -dimensional symplectic manifold, on which the symplectic form vanishes. We will allow Lagrangian manifolds to have boundaries, and to be disconnected.

applied to a term which is compactly supported, and whose L^2 norm is $O(h)$. Therefore, we have a priori that $\|\chi E_h^\xi\|_{L^2} \leq O(h)\|\chi(P_h - (1 + i0)^2)^{-1}\chi\|_{L^2 \rightarrow L^2}$, as least if the support of χ is large enough. But under Hypotheses 10 and 46, it is known from [Nonnenmacher and Zworski 2009] (see Theorem 45) that

$$\|\chi(P_h - (1 + i0)^2)^{-1}\chi\|_{L^2 \rightarrow L^2} \leq C \frac{|\log h|}{h},$$

and such estimates are sharp in the presence of trapping (see [Bony et al. 2010]). Such a priori estimates would therefore only give $\|\chi E_h^\xi\|_{L^2} \leq C|\log h|$.

Our next result concerns the *semiclassical measure* of E_h^ξ . Consider on $T^*\mathbb{R}^d$ the measure μ_0^ξ given by

$$d\mu_0^\xi(x, v) = dx \delta_{v=\xi}.$$

The measure μ_0^ξ is the semiclassical measure associated to $e^{\frac{i}{h}x \cdot \xi}$, in the sense that for any $\psi \in C_c^\infty(T^*\mathbb{R}^d)$ and any $\chi \in C_c^\infty(\mathbb{R}^d)$, we have

$$\lim_{h \rightarrow 0} \langle \text{Op}_h(\psi) \chi e^{\frac{i}{h}x \cdot \xi}, \chi e^{\frac{i}{h}x \cdot \xi} \rangle = \int_{T^*\mathbb{R}^d} \chi^2(x) \psi(x, v) d\mu_0^\xi(x, v).$$

For the definition and properties of the Weyl quantization Op_h , we refer the reader to Section 3.1.1.

We then define a measure μ^ξ on $T^*\mathbb{R}^d$ by

$$\int_{T^*\mathbb{R}^d} a d\mu^\xi := \lim_{t \rightarrow \infty} \int_{T^*\mathbb{R}^d} a \circ \Phi^t d\mu_0^\xi$$

for any $a \in C_c^0(T^*\mathbb{R}^d)$.

We will show in Section 6.3 that this limit exists under our above assumptions. Actually, the proof will not use Hypothesis 46 that the topological pressure of half the unstable jacobian is negative, but the much weaker assumption that the topological pressure of the unstable jacobian is negative.

The following theorem tells us that, under our hypotheses, μ^ξ is the semiclassical measure associated to E_h^ξ , and it gives us a precise description of μ^ξ close to the trapped set.

Theorem 3. *Suppose that Hypothesis 10 on hyperbolicity holds, that Hypothesis 46 concerning the topological pressure is satisfied, and that $\xi \in \mathbb{S}^{d-1}$ is such that Λ_ξ satisfies Hypothesis 16 of transversality.*

Then for any $\psi \in C_c^\infty(T^\mathbb{R}^d)$ and any $\chi \in C_c^\infty(\mathbb{R}^d)$, we have*

$$\langle \text{Op}_h(\psi) \chi E_h^\xi, \chi E_h^\xi \rangle = \int_{T^*\mathbb{R}^d} \psi(x, v) d\mu^\xi(x, v) + O(h^c).$$

Furthermore, for any $\rho \in K$, there exists a small neighbourhood $U_\rho \subset T^*\mathbb{R}^d$ of ρ , and a local change of symplectic coordinates $\kappa_\rho : U_\rho \rightarrow T^*\mathbb{R}^d$ with $\kappa_\rho(\rho) = 0$ such that the following holds. There exists a constant $c > 0$ and two sequences of functions $f_n, \phi_n \in C_c^\infty(\mathbb{R}^d)$ for $n \in \mathbb{N}$ such that for any $(y, \eta) \in \kappa_\rho(U_\rho)$, we have

$$d\mu^\xi(\kappa_\rho^{-1}(y, \eta)) = \sum_{n=0}^\infty f_n(y) \delta_{\{\eta = \partial \phi_n(y)\}} dy,$$

and where the functions f_n satisfy

$$\sum_{n=0}^{\infty} \|f_n\|_{C^0} < \infty. \quad (7)$$

Remark 4. [Theorem 3](#) tells us that the distorted plane waves E_h^ξ have a unique semiclassical measure. This result is therefore analogous to the quantum unique ergodicity conjecture for eigenfunctions of the Laplace–Beltrami operator on manifolds of negative curvature. However, on compact manifolds of negative curvature, the semiclassical measure we expect is the Liouville measure. Here, the semiclassical measure given by [Theorem 3](#) is very different from the Liouville measure, since, close to the trapped set, it is concentrated on a countable union of Lagrangian submanifolds of T^*X . There is therefore a deep difference between compact and noncompact manifolds concerning the semiclassical measure of eigenfunctions, a fact which was already noted in [\[Guillarmou and Naud 2014\]](#).

Idea of proof. [Theorems 1](#) and [3](#) will be deduced from a precise description of the distorted plane waves E_h^ξ microlocally near the trapped set. In [Theorem 47](#), we will show that, microlocally near the trapped set, E_h^ξ can be written as a convergent sum of WKB states. Let us now explain how this result is obtained.

By definition, the distorted plane waves E_h^ξ are generalized eigenfunctions of the operator P_h . Therefore, if we write $U(t) = e^{-\frac{i}{h}P_h}$ for the Schrödinger propagator associated to P_h , we would like to write formally that $U(t)E_h^\xi = e^{-\frac{it}{h}}E_h^\xi$. Of course, this expression can only be formal, since $E_h^\xi \notin L^2$, but we will give it a precise meaning by truncating it by some cut-off functions.

By equation (2), E_h^ξ may be decomposed into two terms, which we will write as E_h^0 and E_h^1 in the sequel. E_h^0 is a Lagrangian state associated to the Lagrangian manifold Λ_ξ , while E_h^1 is the image of a smooth compactly supported function by the resolvent $(P_h - (1 + i0)^2)^{-1}$.

Using some resolvent estimates and hyperbolic dispersion estimates, we will show in the sequel that, for any compactly supported function χ , we have $\lim_{t \rightarrow \infty} \|\chi U(t)E_h^1\| = 0$.

Therefore, in order to describe E_h^ξ , we only have to study $U(t)E_h^0$ for some very long times. Since E_h^0 is a Lagrangian state, its evolution can be described using the WKB method. To do this, we will have to understand the classical evolution of the Lagrangian manifold Λ_ξ for large times. We will show that for any $t > 0$, the restriction of $\Phi^t(\Lambda_\xi)$ to a region close to the trapped set consists of finitely many Lagrangian manifolds, most of which are very close to the “outgoing tail” of the trapped set (see [Theorem 17](#) for more details).

Relation to other works. The study of the high frequency behaviour of eigenfunctions of Schrödinger operators, and of their semiclassical measures, in the case where the associated classical dynamics has a chaotic behaviour, has a long story. It goes back to the classical works [\[Shnirelman 1974; Zelditch 1987; Colin de Verdière 1985\]](#) dealing with quantum ergodicity on compact manifolds.

Analogous results on manifolds of infinite volume are much more recent. In [\[Dyatlov and Guillarmou 2014\]](#), the authors studied the semiclassical measures associated to distorted plane waves in a very general framework, with very mild assumptions on the classical dynamics. The counterpart of this generality is that the authors have to average on directions ξ and on an energy interval of size h to be able to define the semiclassical measure of distorted plane waves. Their result can be seen as a form of quantum ergodicity result on noncompact manifolds, although no “ergodicity” assumption is made.

In [Guillarmou and Naud 2014], the authors considered the case where $X = \Gamma \backslash \mathbb{H}^d$ is a manifold of infinite volume, with sectional curvature constant equal to -1 (convex cocompact hyperbolic manifold), and with the assumption that the Hausdorff dimension of the limit set of Γ is smaller than $(d-1)/2$. In this setting, distorted plane waves are often called *Eisenstein series*. The authors prove that there is a unique semiclassical measure for the Eisenstein series with a given incoming direction, and they give a very explicit formula for it. This result can hence be seen as a quantum unique ergodicity result in infinite volume.

Our result is a generalization of those of [Guillarmou and Naud 2014]. Indeed, we also obtain a unique semiclassical measure for the distorted plane waves with a given incoming direction. Our assumption on the topological pressure is a natural generalization of the assumption on the Hausdorff dimension of the limit set of Γ to the case of nonconstant curvature. As in [Guillarmou and Naud 2014], the main ingredient of the proof is a decomposition of the distorted plane waves as a sum of WKB states. Although our description of the distorted plane waves and of their semiclassical measure is slightly less explicit than that of [Guillarmou and Naud 2014], our methods are much more versatile, since they rely on the properties of the Hamiltonian flow close to the trapped set, instead of relying on the global quotient structure.

In [Dyatlov 2012], the author was able to obtain semiclassical convergence of distorted plane waves on manifolds of finite volume (with cusps), by working at complex energies; see also [Bonthonneau 2014] for more precise results. The main argument of [Dyatlov 2012], [Bonthonneau 2014] and [Dyatlov and Guillarmou 2014], which is to describe the distorted plane waves as plane waves propagated during a long time by the Schrödinger flow, is the starting point of our proof. However, the reason for the convergence in the long-time limit is very different in the papers above than in the present paper.

Many of the tools used in this paper were inspired by [Nonnenmacher and Zworski 2009]. We will use the notations and methods of this paper a lot.

Most of the results of the present paper can be made more precise if we suppose that we work on a manifold of nonpositive sectional curvature, without a potential. This has been studied in [Ingremeau 2017], where the author is able to show, by using the methods developed in the present paper, that distorted plane waves are bounded in L_{loc}^∞ independently of h , and to give sharp bounds on the Hausdorff measure of nodal sets of the real part of distorted plane waves restricted to a compact set.

Organization of the paper. In Section 2, we will state and prove a result concerning the propagation by the Hamiltonian flow of Lagrangian manifolds similar to Λ_ξ near the trapped set, under general assumptions. In Section 3, we will state Theorem 47, which is our main theorem, giving a description of distorted plane waves as a sum of WKB states. We will deduce Theorem 1 as an easy corollary. In Section 4, we will recall various tools which were introduced in [Nonnenmacher and Zworski 2009], and which will play a role in the proof of Theorem 47. We shall then prove Theorem 47 in Section 5. Section 6 will be devoted to the proof of the Theorem 3.

The main reason why we want to state Theorem 47 for generalized eigenfunctions that are more general than distorted plane waves on \mathbb{R}^d is that our results do also apply if the manifold is hyperbolic near infinity (which allows us to recover some of the results of [Guillarmou and Naud 2014]), as is shown in [Ingremeau 2017, Appendix B]. Our results do probably also apply if the manifold is asymptotically hyperbolic; this shall be pursued elsewhere.

2. Propagation of Lagrangian manifolds

2.1. General assumptions for propagation of Lagrangian manifolds. Let (X, g) be a noncompact complete Riemannian manifold of dimension d , and let $V : X \rightarrow \mathbb{R}$ be a smooth compactly supported potential.

We denote by $p(x, \xi) = p(\rho) : T^*X \rightarrow \mathbb{R}$, $p(x, \xi) = \|\xi\|^2 + V(x)$, the classical Hamiltonian.

For each $t \in \mathbb{R}$, we denote by $\Phi^t : T^*X \rightarrow T^*X$ the Hamiltonian flow at time t for the Hamiltonian p .

Given any smooth function $f : X \rightarrow \mathbb{R}$, it may be lifted to a function $f : T^*X \rightarrow \mathbb{R}$, which we denote by the same letter. We may then define $\dot{f}, \ddot{f} \in C^\infty(T^*X)$ to be the derivatives of f with respect to the Hamiltonian flow:

$$\dot{f}(x, \xi) := \left. \frac{df(\Phi^t(x, \xi))}{dt} \right|_{t=0}, \quad \ddot{f}(x, \xi) := \left. \frac{d^2f(\Phi^t(x, \xi))}{dt^2} \right|_{t=0}.$$

2.1.1. Hypotheses near infinity. We suppose the following conditions are fulfilled.

Hypothesis 5 (structure of X near infinity). *We suppose the manifold (X, g) is such that the following holds:*

- (1) *There exists a compactification \bar{X} of X , that is, a compact manifold with boundaries \bar{X} such that X is diffeomorphic to the interior of \bar{X} . The boundary $\partial\bar{X}$ is called the boundary at infinity.*
- (2) *There exists a boundary-defining function b on X , that is, a smooth function $b : \bar{X} \rightarrow [0, \infty)$ such that $b > 0$ on X , and b vanishes to first order on $\partial\bar{X}$.*
- (3) *There exists a constant $\epsilon_0 > 0$ such that for any point $(x, \xi) \in \mathcal{E}$,*

$$\text{if } b(x, \xi) \leq \epsilon_0 \text{ and } \dot{b}(x, \xi) = 0 \text{ then } \ddot{b}(x, \xi) < 0.$$

Note that, although part (3) of the hypothesis makes reference to the Hamiltonian flow, it is only an assumption on the manifold (X, g) and not on the potential V , because V is assumed to be compactly supported.

Example 6. \mathbb{R}^d fulfills **Hypothesis 5** by taking the boundary-defining function $b(x) = (1 + |x|^2)^{-1/2}$. We then have $\bar{X} \equiv B(0, 1)$.

Example 7. The Poincaré space \mathbb{H}^d also fulfills **Hypothesis 5**. Indeed, in the ball model $B_0(1) = \{x \in \mathbb{R}^d : |x| < 1\}$, where $|\cdot|$ denotes the Euclidean norm, \mathbb{H}^d compactifies to the closed unit ball, and the boundary-defining function $b(x) = 2(1 - |x|)/(1 + |x|)$ fulfills conditions (2) and (3).

We will write $X_0 := \{x \in X : b(x) \geq \epsilon_0/2\}$. By possibly taking ϵ_0 smaller, we can assume that $\text{supp}(V) \subset \{x \in X : b(x) > \epsilon_0\}$. We will call X_0 the *interaction region*. We will also write

$$W_0 := T^*(X \setminus X_0) = \{\rho \in T^*X : b(\rho) < \epsilon_0/2\}, \quad \mathcal{W}_0 = W_0 \cap \mathcal{E}. \tag{8}$$

By possibly taking ϵ_0 even smaller, we may ask that

$$\forall \rho \in \mathcal{W}_0, \quad b(\Phi^1(\rho)) < \epsilon_0. \tag{9}$$

Definition 8. If $\rho = (x, \xi) \in \mathcal{E}$, we say that ρ escapes directly in the forward direction, denoted $\rho \in \mathcal{DE}_+$, if $b(x) < \epsilon_0/2$ and $\dot{b}(x, \xi) \leq 0$.

If $\rho = (x, \xi) \in \mathcal{E}$, we say that ρ escapes directly in the backward direction, denoted $\rho \in \mathcal{DE}_-$, if $b(x) < \epsilon_0/2$ and $\dot{b}(x, \xi) \geq 0$.

Note that we have

$$\mathcal{W}_0 = \mathcal{DE}_- \cup \mathcal{DE}_+.$$

Part (3) of [Hypothesis 5](#) implies the following *geodesic convexity* result, which reflects the fact that once a trajectory has left the interaction region, it cannot come back to it.

Lemma 9. For any $t \geq 0$, we have

$$\Phi^t(\mathcal{E} \cap T^*X_0) \cap \mathcal{DE}_- = \emptyset.$$

Proof. Suppose that there exists a $\rho \in \Phi^t(\mathcal{E} \cap T^*X_0) \cap \mathcal{DE}_-$ for some $t \geq 0$. Then there exists $\rho' \in \mathcal{E} \cap T^*X_0$ such that $\rho = \Phi^t(\rho')$. Let us consider $f(s) := b(\Phi^s(\rho'))$. We have $f(0) > \epsilon_0/2$, $f(t) < \epsilon_0/2$ and $f'(t) \geq 0$ by hypothesis. This is impossible, because by [Hypothesis 5](#), point (3), whenever $f(s) \leq \epsilon_0$ and $f'(s) = 0$, we have $f''(s) < 0$. □

2.1.2. Hyperbolicity. Recall that the *trapped set* was defined in (4). In the sequel, we will always suppose that the trapped set is a *hyperbolic set*, as follows.

Hypothesis 10 (hyperbolicity of the trapped set). We assume that K is a hyperbolic set for the flow $\Phi^t|_{\mathcal{E}}$. That is to say, there exists a metric g_{ad} on a neighbourhood of K included in \mathcal{E} , and $\lambda > 0$, such that the following holds. For each $\rho \in K$, there is a decomposition

$$T_\rho \mathcal{E} = \mathbb{R}H_\rho(\rho) \oplus E_\rho^+ \oplus E_\rho^-$$

such that

$$\|d\Phi_\rho^t(v)\|_{g_{\text{ad}}} \leq e^{-\lambda|t|} \|v\|_{g_{\text{ad}}} \quad \text{for all } v \in E_\rho^\mp, \pm t \geq 0.$$

We will call E^\pm the *unstable* (resp. *stable*) subspaces at the point ρ .

We may extend g_{ad} to a metric on the whole energy layer, so that outside of the interaction region, it coincides with the metric on T^*X induced from the Riemannian metric on X . From now on, d will denote the Riemannian distance associated to this metric on \mathcal{E} .

Let us recall a few properties of hyperbolic dynamics (see [[Katok and Hasselblatt 1995](#), Chapter 6] for the proofs of the statements).

(i) The hyperbolic set is *structurally stable*, in the following sense. For $E > 0$, define the layer of energy E as

$$\mathcal{E}_E := \{\rho \in T^*X : p(\rho) = E\}, \tag{10}$$

and the trapped set at energy E as

$$K_E := \{\rho \in \mathcal{E}_E : \Phi^t(\rho) \text{ remains in a compact set for all } t \in \mathbb{R}\}. \tag{11}$$




Figure 2. A surface which has negative curvature close to the trapped set of the geodesic flow, and which is isometric to two copies of $\mathbb{R}^2 \setminus B(0, R_0)$ outside of a compact set. It satisfies [Hypothesis 10](#) near the trapped set and [Hypothesis 5](#) at infinity.

If K is a hyperbolic set for $\Phi^t|_E$, then

$$\exists \delta > 0, \forall E \in (1 - \delta, 1 + \delta), \quad K_E \text{ is a hyperbolic set for } \Phi^t|_{\mathcal{E}_E}. \tag{12}$$

(ii) $d\Phi^t_\rho(E_\rho^\pm) = E_{\Phi^t(\rho)}^\pm$.

(iii) $K \ni \rho \mapsto E_\rho^\pm \subset T_\rho(\mathcal{E})$ is Hölder-continuous.

(iv) Any $\rho \in K$ admits local strongly (un)stable manifolds $W_{\text{loc}}^\pm(\rho)$ tangent to E_ρ^\pm , defined by

$$W_{\text{loc}}^\pm(\rho) = \left\{ \rho' \in \mathcal{E} : d(\Phi^t(\rho), \Phi^t(\rho')) < \epsilon \text{ for all } \pm t \leq 0 \text{ and } \lim_{t \rightarrow \mp\infty} d(\Phi^t(\rho'), \Phi^t(\rho)) = 0 \right\},$$

where $\epsilon > 0$ is some small number.

We call

$$E_\rho^{+0} := E_\rho^+ \oplus \mathbb{R}H_p(\rho), \quad E_\rho^{-0} := E_\rho^- \oplus \mathbb{R}H_p(\rho)$$

the *weak unstable* and *weak stable* subspaces at the point ρ respectively.

2.1.3. Adapted coordinates. Let us now describe the construction of a local system of coordinates which is adapted to the stable and unstable directions near a point. In the sequel, these coordinates will be considered as fixed, and used to state [Theorem 17](#).

Lemma 11. *Let $\rho \in K$. There exists an adapted system of symplectic coordinates (y^ρ, η^ρ) on a neighbourhood of ρ in T^*X such that the following holds:*

- (i) $\rho \equiv (0, 0)$.
- (ii) $E_\rho^+ = \text{span}\{(\partial/\partial y_i^\rho)(\rho) : i = 2, \dots, d\}$.
- (iii) $E_\rho^- = \text{span}\{(\partial/\partial \eta_i^\rho)(\rho) : i = 2, \dots, d\}$.
- (iv) $\eta_1^\rho = p - 1$ is the energy coordinate.
- (v) $\langle (\partial/\partial y_i^\rho)(\rho), (\partial/\partial y_j^\rho)(\rho) \rangle_{g_{\text{ad}}(\rho)} = \delta_{i,j}, \quad i, j = 2, \dots, d$.

Proof. We may identify a neighbourhood of $\rho \in T^*X$ with a neighbourhood of $(0, 0) \in T^*\mathbb{R}^d$. Let us take $e_1^\rho = H_p(\rho)$, and complete it into a basis $(e_1^\rho, \dots, e_d^\rho)$ of E_ρ^{+0} such that $\langle e_i^\rho, e_j^\rho \rangle_{g^{\text{ad}}(\rho)} = 1$ for $2 \leq i, j \leq d$.

Since $E^{\pm 0}$ are Lagrangian subspaces (which follows from the hyperbolicity assumption), it is then possible to find vectors $(f_1^\rho, \dots, f_d^\rho)$ such that $E_\rho^- = \text{span}\{f_2^\rho, \dots, f_d^\rho\}$ and such that $\omega(f_j^\rho, e_k^\rho) = \delta_{j,k}$ for any $1 \leq j, k \leq d$. In particular, we have $\omega(f_1^\rho, e_1^\rho) = dp(f_1) = 1$.

From Darboux’s theorem, there exists a nonlinear symplectic chart $(y^{\text{flat}}, \eta^{\text{flat}})$ near the origin such that $\eta_1^{\text{flat}} = p - 1$. There also exists a linear symplectic transformation A such that the coordinates $(y, \eta) = A(y^{\text{flat}}, \eta^{\text{flat}})$ satisfy $\eta_1 = \eta_1^{\text{flat}}$ as well as

$$\eta_1 = p - 1, \quad \frac{\partial}{\partial y_j}(0, 0) = e_j \text{ and } \frac{\partial}{\partial \eta_j}(0, 0) = f_j, \quad j = 1, \dots, d. \quad \square$$

We will often write

$$y^\rho := (y_2^\rho, \dots, y_d^\rho) \quad \text{and} \quad \eta^\rho := (\eta_2^\rho, \dots, \eta_d^\rho). \quad (13)$$

For any $\epsilon > 0$, write $D_\epsilon = \{u \in \mathbb{R}^{d-1} : |u| < \epsilon\}$. We define the following polydisk centred at ρ :

$$U^\rho(\epsilon) \equiv \{(y^\rho, \eta^\rho) : |y_1^\rho| < \epsilon, |\eta_1^\rho| < \delta, y^\rho \in D_\epsilon, \eta \in D_\epsilon\}, \quad (14)$$

where δ comes from (12).

We also define *unstable Lagrangian manifolds*, which are needed in the statement of [Theorem 17](#).

Definition 12. Let $\Lambda \subset \mathcal{E}$ be an isoenergetic Lagrangian manifold (not necessarily connected) included in a small neighbourhood W of a point $\rho \in K$, and let $\gamma > 0$. We will say that Λ is a γ -unstable Lagrangian manifold (or that Λ is in the γ -unstable cone) in the coordinates (y^ρ, η^ρ) if it can be written in the form

$$\Lambda = \{(y^\rho; 0, F(y^\rho)) : y^\rho \in D\},$$

where $D \subset \mathbb{R}^d$ is an open subset with finitely many connected components, and with piecewise smooth boundary, and $F : \mathbb{R}^d \rightarrow \mathbb{R}^{d-1}$ is a smooth function with $\|dF\|_{C^0} \leq \gamma$.

Note that, since F is defined on \mathbb{R}^d , a γ -unstable manifold may always be seen as a submanifold of a *connected γ -unstable Lagrangian manifold*.

Let us also note that, since Λ is isoenergetic and is Lagrangian, an immediate computation shows that F does not depend on y_1^ρ , so that Λ can actually be put in the form

$$\Lambda = \{(y^\rho; 0, f(y^\rho)) : y^\rho \in D\},$$

where $f : \mathbb{R}^{d-1} \rightarrow \mathbb{R}^{d-1}$ is a smooth function with $\|df\|_{C^0} \leq \gamma$.

2.1.4. Hypotheses on the incoming Lagrangian manifold. Let us consider an isoenergetic Lagrangian manifold $\mathcal{L}_0 \subset \mathcal{E}$ of the form

$$\mathcal{L}_0 := \{(x, \phi(x)) : x \in X_1\},$$

where X_1 is a closed subset of $X \setminus X_0$ with finitely many connected components and piecewise smooth boundary, and $\phi : x \rightarrow \phi(x), X_2 \mapsto T_x^*X$, is a smooth covector field defined on some neighbourhood X_2 of X_1 .

We make the following additional hypothesis on \mathcal{L}_0 :

Hypothesis 13 (invariance hypothesis). *We suppose that \mathcal{L}_0 satisfies the following invariance hypotheses:*

$$\forall t \geq 0, \quad \Phi^t(\mathcal{L}_0) \cap \mathcal{DE}_- = \mathcal{L}_0 \cap \mathcal{DE}_-. \tag{15}$$

Example 14. Given a $\xi \in \mathbb{R}^d$ with $|\xi|^2 = 1$, the Lagrangian manifold Λ_ξ defined in the [Introduction](#) fulfills [Hypothesis 13](#).

Example 15. Suppose that $(X \setminus X_0, g) \cong (\mathbb{R}^d \setminus B(0, R), g_{\text{Eucl}})$ for some $R > 0$. Then the incoming spherical Lagrangian, defined by

$$\Lambda_{\text{sph}} := \left\{ \left(x, -\frac{x}{|x|} \right) : |x| > R \right\},$$

fulfills [Hypothesis 13](#).

We also make the following transversality assumption on the Lagrangian manifold \mathcal{L}_0 . It roughly says that \mathcal{L}_0 intersects the stable manifold transversally.

Hypothesis 16 (transversality hypothesis). *We suppose that \mathcal{L}_0 is such that, for any $\rho \in K$, for any $\rho' \in \mathcal{L}_0$, for any $t \geq 0$, we have*

$$\Phi^t(\rho') \in W_{\text{loc}}^-(\rho) \implies W_{\text{loc}}^-(\rho) \text{ and } \Phi^t(\mathcal{L}_0) \text{ intersect transversally at } \Phi^t(\rho'),$$

that is to say

$$T_{\Phi^t(\rho')}\mathcal{L}_0 \oplus T_{\Phi^t(\rho')}W_{\text{loc}}^-(\rho) = T_{\Phi^t(\rho')}\mathcal{E}. \tag{16}$$

Note that (16) is equivalent to $T_{\Phi^t(\rho')}\mathcal{L}_0 \cap T_{\Phi^t(\rho')}W_{\text{loc}}^-(\rho) = \{0\}$.

On $X = \mathbb{R}^d$, [Hypothesis 16](#) is likely to hold for almost every $\xi \in \mathbb{S}^{d-1}$, at least for a generic V . In [\[Ingremeau 2017\]](#), the author shows that this hypothesis is satisfied for every ξ on manifolds of nonpositive curvature which have several Euclidean ends (like the one in [Figure 2](#)), when there is no potential.

2.2. Statement of the result. Let us now state the main result of this section, which describes the “truncated evolution” of Lagrangian manifolds.

Truncated Lagrangians. Let $(W_a)_{a \in A}$ be a finite family of open sets in T^*X . Let $N \in \mathbb{N}$, and let $\alpha = \alpha_0, \alpha_1 \cdots \alpha_{N-1} \in A^N$. Let Λ be a Lagrangian manifold in T^*X . We define the sequence of (possibly empty) Lagrangian manifolds $(\Phi_\alpha^k(\Lambda))_{0 \leq k \leq N}$ by recurrence by

$$\Phi_\alpha^0(\Lambda) = \Lambda \cap W_{\alpha_0}, \quad \Phi_\alpha^{k+1}(\Lambda) = W_{\alpha_{k+1}} \cap \Phi^1(\Phi_\alpha^k(\Lambda)).$$

In the sequel, we will consider families with indices in $A = A_1 \sqcup A_2 \sqcup \{0\}$. For any $\alpha \in A^N$ such that $\alpha_{N-1} \neq 0$, we will define

$$\tau(\alpha) := \max\{1 \leq i \leq N - 1 : \alpha_i = 0\} \tag{17}$$

if there exists $1 \leq i \leq N - 1$ with $\alpha_i = 0$, and $\tau(\alpha) = 0$ otherwise.

Theorem 17. *Suppose that the manifold X satisfies [Hypothesis 5](#) at infinity, that the Hamiltonian flow (Φ^t) satisfies [Hypothesis 10](#), and that the Lagrangian manifold \mathcal{L}_0 satisfies [Hypothesis 13](#) on invariance as well as [Hypothesis 16](#) of transversality.*

*Fix $\gamma_{\text{uns}} > 0$ small enough. There exists $\varepsilon_0 > 0$ such that the following holds. Let $(W_a)_{a \in A_1}$ be any open cover of K in T^*X of diameter $< \varepsilon_0$ such that there exist points $\rho_a \in W_a \cap K$, and such that the adapted coordinates (y^a, η^a) centred on ρ^a are well defined on W_a for every $a \in A_1$. Then we may complete this cover into $(W_a)_{a \in A}$ an open cover of \mathcal{E} in T^*X where $A = A_1 \sqcup A_2 \sqcup \{0\}$ (with W_0 defined as in [\(8\)](#)) such that the following holds.*

There exists $N_{\text{uns}} \in \mathbb{N}$ such that for all $N \in \mathbb{N}$, for all $\alpha \in A^N$ and all $a \in A_1$, we have $W_a \cap \Phi_\alpha^N(\mathcal{L}_0)$ is either empty, or is a Lagrangian manifold in some unstable cone in the coordinates (y^a, η^a) .

Furthermore, if $N - \tau(\alpha) \geq N_{\text{uns}}$, then $W_a \cap \Phi_\alpha^N(\mathcal{L}_0)$ is a γ_{uns} -unstable Lagrangian manifold in the coordinates (y^a, η^a) .

Remark 18. For a sequence $\alpha \in A^N$, $N - \tau(\alpha)$ corresponds to the time spent in the interaction region. Our last statement therefore says that if a part of \mathcal{L}_0 stays in the interaction region for long enough when propagated, then its tangents will form a small angle with the unstable direction at ρ^a .

Remark 19. The constant ε_0 and the sets $(W_a)_{a \in A_2}$ depend on the Lagrangian manifold \mathcal{L}_0 . If we take a whole family of Lagrangian manifolds $(\mathcal{L}_z)_{z \in Z}$ satisfying [Hypotheses 13](#) and [16](#), then we will need some additional conditions on the whole family to be able to find a common choice of ε_0 and $(W_a)_{a \in A_2}$ independent of $z \in Z$. An example of such a condition will be provided by equations [\(36\)](#) and [\(37\)](#). Note that these equations are automatically satisfied if Z is finite.

2.3. Proof of [Theorem 17](#).

Proof. From now on, we will fix a $\gamma_{\text{uns}} > 0$.

Let $\rho_0 \in K$, and consider the system of adapted coordinates in a neighbourhood of ρ_0 constructed in [Section 2.1.3](#). Recall that the set $U^{\rho_0}(\epsilon)$ was defined in [\(14\)](#). We define a *Poincaré section* by

$$\Sigma^{\rho_0} = \Sigma^{\rho_0}(\epsilon) = \{(y^{\rho_0}, \eta^{\rho_0}) \in U^{\rho_0}(\epsilon) : y_1^{\rho_0} = \eta_1^{\rho_0} = 0\}.$$

Note that the spaces $E_{\rho_0}^\pm$ are tangent to Σ^{ρ_0} , and that the coordinates $(y^{\rho_0}, \eta^{\rho_0})$ introduced in [\(13\)](#) form a symplectic chart on Σ^{ρ_0} .

Actually, we will often need a nonsymplectic system of coordinates built from the coordinates (y^ρ, η^ρ) .

Before building this nonsymplectic system of coordinates, let us explain why it is a crucial ingredient of our argument. The main tool in the proof of [Theorem 17](#) is the so-called “inclination lemma”, which roughly says that a Lagrangian manifold which intersects the stable manifold transversally will get more and more unstable when propagated in the future. This is a very easy result in the case of linear hyperbolic diffeomorphisms, but we must add some quantifiers in the case of nonlinear dynamics to make it rigorous. Namely, one can say, as in [[Nonnenmacher and Zworski 2009](#), Proposition 5.1], that given a $\gamma > 0$, there exists $\epsilon_\gamma > 0$ such that if Λ is a γ -unstable Lagrangian manifold included in some $U^\rho(\epsilon_\gamma)$, then for any ρ' , $\Phi^1(\Lambda) \cap U^{\rho'}(\epsilon_\gamma)$ is still γ -unstable.

However, we may not use this result directly for the following reason. The smaller we take ϵ , the longer the points of the Lagrangian manifold \mathcal{L}_0 may spend in the part of the interaction region which is not affected by the hyperbolic dynamics before entering in some $U^\rho(\epsilon)$ for some $\rho \in K$. Yet the longer they spend in this “intermediate” region, the more stable the Lagrangian manifold may a priori become. To avoid such a circular reasoning, we should introduce another system of coordinates, in which the description of the propagation of the Lagrangian manifolds in the intermediate region is easier.

2.3.1. Alternative coordinates. In this section we will describe a system of “alternative”, or “twisted” coordinates built from the one we introduced in Section 2.1.3, but which may differ slightly from them.

Given a $\rho \in K$, we introduce a system of smooth coordinates $(\tilde{y}^\rho, \tilde{\eta}^\rho)$ as follows.

On Σ^ρ , these coordinates are such that

$$W_{\text{loc}}^{0+}(\rho) \cap \Sigma^\rho \equiv \{(\tilde{y}^\rho, 0) : \tilde{y}^\rho \in D_\epsilon\}, \quad W_{\text{loc}}^{0-}(\rho) \cap \Sigma^\rho \equiv \{(0, \tilde{\eta}^\rho) : \tilde{\eta}^\rho \in D_\epsilon\},$$

and if we denote by L_ρ the map

$$L_\rho : (y^\rho, \eta^\rho) \mapsto (\tilde{y}^\rho, \tilde{\eta}^\rho) \tag{18}$$

defined in a neighbourhood of $(0, 0)$, we have

$$dL_\rho(0, 0) = \text{Id}_{\mathbb{R}^{2d-2}}. \tag{19}$$

Now, if $\hat{\rho}$ has straight coordinates $(y^\rho(\hat{\rho}), \eta^\rho(\hat{\rho}))$, we let $\hat{\rho}' \in \Sigma^\rho$ be the point with straight coordinates $(0, y^\rho(\hat{\rho}), 0, \eta^\rho(\hat{\rho}))$. We then define the twisted coordinates of $\hat{\rho}$ by

$$\tilde{y}_1^\rho(\hat{\rho}) = y_1^\rho(\hat{\rho}), \quad \tilde{\eta}_1^\rho(\hat{\rho}) = \eta_1^\rho(\hat{\rho}), \quad \tilde{y}^\rho(\hat{\rho}) = \tilde{y}^\rho(\hat{\rho}'), \quad \tilde{\eta}^\rho(\hat{\rho}) = \tilde{\eta}^\rho(\hat{\rho}').$$

Note that this system of coordinates doesn't have to be symplectic.

We have

$$\frac{\partial y_j^\rho}{\partial \tilde{y}_1^\rho} = \frac{\partial \eta_j^\rho}{\partial \tilde{\eta}_1^\rho} = 0 \quad \text{for } j = 1, \dots, d-1, \quad \frac{\partial y_1^\rho}{\partial \tilde{y}_1^\rho} = 1. \tag{20}$$

Given a $\rho \in K$, and $\epsilon, \epsilon' > 0$, we define

$$\tilde{U}^\rho(\epsilon, \epsilon') \equiv \{(\tilde{y}^\rho, \tilde{\eta}^\rho) : |\tilde{y}_1^\rho| < \epsilon, |\tilde{\eta}_1^\rho| < \delta, \tilde{y}^\rho \in D_{\epsilon'}, \tilde{\eta}^\rho \in D_\epsilon\}, \tag{21}$$

where δ is an energy interval on which the dynamics remains uniformly hyperbolic.

Finally, the Poincaré section in the alternative coordinates is represented as

$$\tilde{\Sigma}^\rho(\epsilon, \epsilon') := \{(\tilde{y}^\rho, \tilde{\eta}^\rho) \in \tilde{U}^\rho(\epsilon, \epsilon') : \tilde{y}_1^\rho = \tilde{\eta}_1^\rho = 0\}.$$

In the sequel, we will be working most of the time in a situation where $\epsilon' \ll \epsilon$ (that is, with sets much thinner in the unstable direction than in the stable direction).

The main reason why we needed to introduce alternative coordinates is that they give a simpler expression for the *Poincaré map* (see Remark 20). Let us now define this map.

2.3.2. The Poincaré map. Let $\rho_0 \in K$, and let $\epsilon > 0$ be small enough so that the twisted coordinates around ρ_0 and $\Phi^1(\rho_0)$ are well defined in some neighbourhoods $\tilde{U}^{\rho_0}(\epsilon, \epsilon)$ and $\tilde{U}^{\Phi^1(\rho_0)}(\epsilon, \epsilon)$. The Poincaré map κ_{ρ_0} is defined, for $\rho \in \tilde{\Sigma}^{\rho_0}(\epsilon)$ near ρ_0 , by taking the intersection of the trajectory $(\Phi^s(\rho))_{|s-1| \leq \epsilon}$ with the section $\tilde{\Sigma}^{\Phi^1(\rho_0)}$ (this intersection consists of at most one point). In the sequel, we will sometimes omit the reference to ρ_0 and simply write the Poincaré map κ .

The map κ_{ρ_0} need not be symplectic, since it is defined in the twisted coordinates which need not be symplectic. However, if we had defined the Poincaré map in the straight coordinates, it would have been automatically symplectic. The linearisations of the two systems of coordinates are identical at ρ_0 by (19). Therefore, by using the hyperbolicity assumption, we see that the differential of κ at ρ_0 takes the form

$$d\kappa(\rho_0) \equiv \begin{pmatrix} A & 0 \\ 0 & {}^tA^{-1} \end{pmatrix},$$

and there exists

$$\nu = e^{-\lambda} < 1 \tag{22}$$

such that the matrix A satisfies

$$\|A^{-1}\| \leq \nu, \tag{23}$$

where $\|\cdot\|$ corresponds to the matrix norm. Hence, the Poincaré map κ_{ρ_0} takes the form

$$\kappa_{\rho_0}(\tilde{y}^{\rho_0}, \tilde{\eta}^{\rho_0}) = (A\tilde{y}^{\rho_0} + \tilde{\alpha}(\tilde{y}^{\rho_0}, \tilde{\eta}^{\rho_0}), {}^tA^{-1}\tilde{\eta}^{\rho_0} + \tilde{\beta}(\tilde{y}^{\rho_0}, \tilde{\eta}^{\rho_0})), \tag{24}$$

and the functions $\tilde{\alpha}$ and $\tilde{\beta}$ satisfy

$$\tilde{\alpha}(0, \tilde{\eta}^{\rho_0}) = \tilde{\beta}(\tilde{y}^{\rho_0}, 0) \equiv 0 \quad \text{and} \quad d\tilde{\alpha}(0, 0) = d\tilde{\beta}(0, 0) = 0. \tag{25}$$

We therefore have

$$\|\tilde{\alpha}\|_{C^1(V)} \leq C_0\epsilon, \quad \|\tilde{\beta}\|_{C^1(V)} \leq C_0\epsilon \tag{26}$$

for some constant C_0 , since κ is uniformly C^2 .

Remark 20. Equation (25) is the main reason why we needed to introduce alternative coordinates, and will play a key role in the proof of Lemma 31. If we had defined the Poincaré map in the straight coordinates, we wouldn't have had $\alpha(0, \eta^{\rho_0}) = 0$ or $\beta(y^{\rho_0}, 0) = 0$.

Remark 21. By compactness of the trapped set, the constants C_0 and ν may be chosen independent of the point ρ_0 . We may also find a $C > 1$ such that, independently of ρ_0 and ρ_1 in K , we have

$$\|A\| \leq C. \tag{27}$$

Finally, by possibly taking C_0 larger, we may assume that all the second derivatives of the map L_ρ defined in (18) are bounded by C_0 independently on $\rho \in K$.

2.3.3. Changes of coordinates and Lagrangian manifolds. Let us describe how a Lagrangian manifold is affected when we go from twisted coordinates to straight coordinates centred at the same point.

Lemma 22. Suppose that a Lagrangian manifold $\Lambda \subset \tilde{U}^\rho(\epsilon, \epsilon)$ may be written in the twisted coordinates centred on $\rho \in K$ as $\Lambda = \{(\tilde{y}_1^\rho, \tilde{y}^\rho; 0, \tilde{F}(\tilde{y}^\rho)) : \tilde{y}^\rho \in \rho\}$, where $\rho \subset \mathbb{R}^d$ is a small open set, and with $\|d\tilde{F}\|_{C^0} \leq \gamma$. Suppose furthermore that

$$C_0\epsilon\gamma < 1.$$

Then, in the straight coordinates, Λ may be written as

$$\Lambda = \{(y_1^\rho, \mathbf{y}^\rho; 0, f(\mathbf{y}^\rho)) : \mathbf{y}^\rho \in D_\rho\},$$

with $\|df\|_{C^0} \leq \gamma(1 - C_0\gamma\epsilon)^{-1}(1 + 2C_0\epsilon)$.

Proof. To lighten the notations, we will not write the indices ρ .

Points on Λ are parametrized by the coordinate \tilde{y} . We may hence see their straight coordinates u, s as functions of \tilde{y} .

By equations (19), (20) and Remark 21, we have

$$\frac{\partial y}{\partial \tilde{y}} = \frac{\partial y}{\partial \tilde{y}} + \frac{\partial y}{\partial \tilde{\eta}} \frac{\partial \tilde{F}(\tilde{y})}{\partial \tilde{y}} = I + R$$

with $\|R\| \leq C_0\gamma\epsilon < 1$.

Therefore, on Λ , we know $\tilde{y} \mapsto y$ is invertible. We may hence write η as a function of y , and we have

$$\frac{\partial \eta}{\partial y} = \frac{\partial \tilde{y}}{\partial y} \left[\frac{\partial \eta}{\partial \tilde{y}} + \frac{\tilde{F}(\tilde{y})}{\partial \tilde{y}} \frac{\partial \eta}{\partial \tilde{\eta}} \right] = (I + R)^{-1}(\gamma(I + R')),$$

with $\|R'\| \leq 2C_0\epsilon$. Hence $\|\partial\eta/\partial y\| \leq \gamma(1 - C_0\gamma\epsilon)^{-1}(1 + 2C_0\epsilon)$.

That η is actually independent of y_1 comes from the fact that Λ is an isoenergetic Lagrangian manifold, and that we are working in symplectic coordinates. □

Let us now describe the change between two systems of twisted coordinates. Let $\rho, \rho' \in K$. If they are close enough to each other, the map $L : (\tilde{y}^\rho, \tilde{\eta}^\rho) \mapsto (\tilde{y}^{\rho'}, \tilde{\eta}^{\rho'})$ is well defined on a set containing both ρ and ρ' , of diameter $d(\rho, \rho')$.

Combining the fact that the (un)stable subspaces E_ρ^\pm are Hölder continuous with respect to $\rho \in K^\delta$ with some Hölder exponent $p > 0$, and point (v) of Lemma 11, we get

$$dL_{(0,0)} = L + R_{\rho,\rho'}, \tag{28}$$

where

$$\|R_{\rho,\rho'}\| \leq Cd^p(\rho, \rho') \quad \text{for some } p > 0, \tag{29}$$

and where L is of the form

$$L = \begin{pmatrix} U_y & 0 \\ 0 & L_\eta \end{pmatrix}$$

for some unitary matrix U_y . Here, L_η might not be unitary, but it is invertible, and by compactness of K , $\|L_\eta\|^{-1}$ may be bounded independently on ρ .

Now, by compactness, the second derivatives of L may be bounded independently of ρ and ρ' . Therefore, for any ρ'' in a neighbourhood of ρ , we have

$$dL_{\rho''} = dL_{(0,0)} + R_{\rho''}, \tag{30}$$

with $R_{\rho''} \leq C'd(\rho, \rho'')$ and C' independent of ρ' .

By possibly enlarging C_0 , we may assume that $\|L_\eta\|^{-1} \leq C_0$. We may also assume that $C_0/2$ is larger than the constants C and C' appearing in the bounds on $R_{\rho,\rho'}$ and $R_{\rho''}$.

We will use the previous remarks in the form of the following lemma, which describes the effect of a change of twisted coordinates on a Lagrangian manifold.

Lemma 23. *Let $\rho, \rho' \in K$ be such that $d(\rho, \rho') < \epsilon$, and let Λ be a Lagrangian manifold which may be written in the twisted coordinates centred on ρ as $\Lambda = \{(\tilde{y}_1^\rho, \tilde{y}^\rho; 0, \tilde{F}^\rho(\tilde{y}^\rho)) : \tilde{y}^\rho \in \rho\}$, where $\rho \subset \mathbb{R}^d$ is a small open set, and with $\|d\tilde{F}^\rho\|_{C^0} \leq \gamma < 1/(4C_0\epsilon^p)$.*

Then, $\Lambda \cap \tilde{U}^{\rho'}(\epsilon, \epsilon)$ may be written in the coordinates centred at ρ' as

$$\Lambda \cap \tilde{U}^{\rho'}(\epsilon, \epsilon) = \{(\tilde{y}_1^{\rho'}, \tilde{y}^{\rho'}; 0, \tilde{F}^{\rho'}(\tilde{y}^{\rho'})) : \tilde{y}^{\rho'} \in \rho'\},$$

where $\rho' \subset \mathbb{R}^d$ is a small open set, and with

$$\|d\tilde{F}^{\rho'}\|_{C^0} \leq (\gamma(1 + C_0\epsilon^p) + C_0\epsilon^p)(1 - 2\gamma C_0\epsilon^p)^{-1} < \infty.$$

Proof. Consider points on Λ . By assumption, their $\tilde{\eta}^\rho$ -coordinate is a function of their \tilde{y}^ρ -coordinate. Therefore, using the map L , their coordinates $(\tilde{y}^{\rho'}, \tilde{\eta}^{\rho'})$ may be seen as functions of \tilde{y}^ρ .

Let us denote by L_y and L_η the two components of L . By definition, we have

$$\tilde{y}^{\rho'} = L_y(\tilde{y}^\rho, \tilde{\eta}^\rho) = L_y(\tilde{y}^\rho, \tilde{F}^\rho(\tilde{y}^\rho)),$$

where $\tilde{F}^\rho(\tilde{y}^\rho)$ satisfies $\|\partial\tilde{F}^\rho(\tilde{y}^\rho)/\partial\tilde{y}^\rho\| \leq \gamma$. Therefore, we have

$$\frac{\partial\tilde{y}^{\rho'}}{\partial\tilde{y}^\rho} = \frac{\partial L_y}{\partial\tilde{y}^\rho} + \frac{\partial\tilde{F}^\rho(\tilde{y}^\rho)}{\partial\tilde{y}^\rho} \frac{\partial L_y}{\partial\tilde{\eta}^\rho} = U + \tilde{R},$$

where U is unitary.

By equations (28) and (30), we have $\|\tilde{R}\| \leq 2\gamma C_0\epsilon^p < 1$ by assumption. Therefore, $\tilde{y}^\rho \mapsto \tilde{y}^{\rho'}$ is invertible, and we have $\|\partial\tilde{y}^\rho/\partial\tilde{y}^{\rho'}\| \leq (1 - 2\gamma C_0\epsilon^p)^{-1}$. We may see $\tilde{\eta}^{\rho'}$ as a function of $\tilde{y}^{\rho'}$, and we have

$$\left\| \frac{\partial\tilde{\eta}^{\rho'}}{\partial\tilde{y}^{\rho'}} \right\| = \left\| \frac{\partial\tilde{y}^\rho}{\partial\tilde{y}^{\rho'}} \frac{\partial\tilde{\eta}^{\rho'}}{\partial\tilde{y}^\rho} + \frac{\partial\tilde{y}^\rho}{\partial\tilde{y}^{\rho'}} \frac{\partial\tilde{\eta}^\rho}{\partial\tilde{y}^\rho} \frac{\partial\tilde{\eta}^{\rho'}}{\partial\tilde{\eta}^\rho} \right\| \leq (1 - 2\gamma C_0\epsilon^p)^{-1} (C_0\epsilon^p + \gamma(1 + C_0\epsilon^p)),$$

and the lemma follows. □

2.3.4. Propagation for bounded times. Let us fix a $\nu_1 \in (\nu, 1)$, where ν was defined in (22). Recall that p was defined in (29) as the Hölder exponent of the stable and unstable directions. From now on, we fix an $\epsilon > 0$ small enough so that

$$\frac{\nu + C_0\epsilon^p}{\nu^{-1} - C_0\epsilon^p} < \nu_1, \quad \text{and} \quad \frac{C_0\epsilon^p}{\nu^{-1} - 2C_0\epsilon^p} < \frac{\gamma_{\text{uns}}(1 - \nu_1)}{8}, \tag{31}$$

$$\left(1 - \frac{(1 + \nu_1)\gamma_{\text{uns}}}{1 + 2C_0\epsilon^p} C_0\epsilon^p\right)^{-1} \left(\gamma_{\text{uns}} \frac{(1 + \nu_1)(1 + C_0\epsilon^p)}{2 + 4C_0\epsilon^p} + C_0\epsilon^p\right) < \frac{\gamma_{\text{uns}}}{1 + 2C_0\epsilon^p}. \tag{32}$$

This is possible because $(1 + \nu_1)/2 < 1$. We also ask that $C_0\epsilon^p < 1/2$. Note that, although condition (32) looks horrible, it is designed to work well with Lemma 23.




Figure 3. A representation of some of the different sets we introduce in the proof of [Theorem 17](#), intersected with a Poincaré section.

Let us introduce a first decomposition of the energy layer. Recall that we defined \mathcal{W}_0 in (8) as the external part of the energy layer. We define $\mathcal{W}_1 := \{\rho \in \mathcal{E} \setminus \mathcal{W}_0 : d(\rho, K) < \epsilon/2\}$ for the part of the energy layer close to the trapped set, and $\mathcal{W}_2 := \{\rho \in \mathcal{E} \setminus \mathcal{W}_0 : d(\rho, K) \geq \epsilon/2\}$ for the intermediate region. See [Figure 3](#) for a representation of these different sets. Note that we will later introduce a finer open cover of the energy layer, using the sets W_a appearing in the statement of the theorem.

The following lemma tells us that the set \mathcal{W}_2 is a transient set, that is to say, points spend only a finite time inside it.

Lemma 24. *There exists $N_\epsilon \in \mathbb{N}$ an integer which depends on ϵ such that for all $\rho \in \mathcal{W}_2$, we have either $\Phi^{N_\epsilon}(\rho) \in \mathcal{W}_0$ or $\Phi^{-N_\epsilon}(\rho) \in \mathcal{W}_0$.*

Proof. This result comes from the uniform transversality of the stable and unstable manifolds (which is a direct consequence of the compactness of K).

It gives us the existence of a $d_1(\epsilon) > 0$ such that, for all $\rho \in \mathcal{W}_2 \cup \mathcal{W}_1$,

$$d(\rho, \Gamma^+) + d(\rho, \Gamma^-) \leq 2d_1 \implies d(\rho, K) \leq \epsilon/2.$$

We may therefore write

$$\mathcal{W}_2 = \{\rho \in \mathcal{W}_2 : d(\rho, \Gamma^+) > d_1\} \cup \{\rho \in \mathcal{W}_2 : d(\rho, \Gamma^-) > d_1\}.$$

A point in the first set will leave the interaction region in finite time in the future, while a point in the second set will leave it in finite time in the past. By compactness, we can find a uniform N_ϵ such as the one in the statement of the lemma. □

The following lemma is a consequence of the transversality assumption we made. It tells us that when we propagate \mathcal{L}_0 during a finite time N and restrict it to a small set $\tilde{U}^\rho(\epsilon, \varrho)$ close to the trapped set, we

obtain a finite union of Lagrangian manifolds in the alternative coordinates. Here, the size ϱ of the set in the unstable direction depends on N , but its size ϵ in the stable direction does not.

Lemma 25. *Let $N \in \mathbb{N}$. There exists $\mathcal{N}_N \in \mathbb{N}$, $\tilde{\varrho}_N > 0$ and $\tilde{\gamma}_N > 0$ such that for all $0 < \varrho \leq \tilde{\varrho}_N$, for all $\rho \in K$, and for all $1 \leq t \leq N$, the set $\Phi^t(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \varrho)$ can be written in the coordinates $(\tilde{y}^\rho, \tilde{\eta}^\rho)$ as the union of at most \mathcal{N}_N disjoint Lagrangian manifolds, which are all $\tilde{\gamma}_N$ -unstable:*

$$\Phi^t(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \varrho) \equiv \bigcup_{l=0}^{l(\varrho)} \hat{\Lambda}_l,$$

with $l(\varrho) \leq \mathcal{N}_N$ and

$$\hat{\Lambda}_l = \{(\tilde{y}_1^\rho, \tilde{y}^\rho; 0, f^l(\tilde{y}^\rho)) : \tilde{y}^\rho \in D_\varrho\}$$

for some smooth functions f^l with $\|df^l(\tilde{y}^\rho)\|_{C^0(D_\varrho)} \leq \tilde{\gamma}_N$.

Proof. Let us consider a $1 \leq t \leq N$. First of all, since Φ^t is a symplectomorphism, it sends Lagrangian manifolds to Lagrangian manifolds. The restriction of a Lagrangian manifold to a region of phase space is a union of Lagrangian manifolds.

We now have to prove that, if we take ϱ small enough, these Lagrangian manifolds are all $\tilde{\gamma}_N$ unstable for some $\tilde{\gamma}_N > 0$ which is independent of ρ .

Let $\rho \in K$. By hypothesis, $W_{\text{loc}}^-(\rho)$ and $\Phi^t(\mathcal{L}_0)$ are transverse when they intersect.

Therefore, in a small neighbourhood of the stable manifold $\{\tilde{y}^\rho = 0\}$, each connected component of $\Phi^t(\mathcal{L}_0)$ may be projected smoothly on the twisted unstable manifold $\{\tilde{\eta}^\rho = 0\}$. That is to say, there exists a $\varrho > 0$ and a $\gamma > 0$ such that each connected component of $\Phi^t(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \varrho)$ is γ -unstable in the twisted coordinates around ρ for some $\gamma > 0$.

Now, since the changes of coordinates between twisted coordinates are continuous, we may use the compactness of K to find uniform constants $\varrho > 0$ and $\gamma > 0$ such that each connected component of $\Phi^t(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \varrho)$ is γ -unstable in the twisted coordinates around ρ , independently of $\rho \in K$ and $1 \leq t \leq N$.

By compactness of $\tilde{U}^\rho(\epsilon, \varrho)$, the number of Lagrangian manifolds making up $\Phi^t(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \varrho)$ is finite. □

Applying this lemma to $N = N_\epsilon + 2$, we define the following constants, which we shall need later in the proof (recall that γ_{uns} has been fixed):

$$(\gamma_0, \varrho_0) := (\tilde{\gamma}_{N_\epsilon+2}, \tilde{\varrho}_{N_\epsilon+2}), \tag{33}$$

$$N_1 := \left\lfloor \frac{\log(\gamma_{\text{uns}}/4\gamma_0)}{\log((1 + \nu_1)/2)} \right\rfloor + 1, \quad N_{\text{uns}} := N_1 + N_\epsilon + 2, \tag{34}$$

$$\varrho_1 := \min\left(\frac{\epsilon}{2\gamma_0}, \varrho_0\right), \quad \varrho_2 := \min((C + C_0\epsilon^p)^{-N_{\text{uns}}} \varrho_1, \tilde{\delta}_{N_{\text{uns}}}), \tag{35}$$

where C comes from [Remark 21](#), and C_0 comes from equation [\(26\)](#).

Remark 26. As explained in [Lemma 24](#), N_ϵ is the maximal time spent by a trajectory in the intermediate region \mathcal{W}_2 . The time N_1 will be the time necessary to incline a γ_0 -unstable Lagrangian manifold to a γ_{uns} -unstable Lagrangian manifold, as explained in [Proposition 30](#). As for the constant ϱ_2 , it has been chosen

small enough so that at each step of the aforementioned propagation during a time N_1 , the Lagrangian manifolds we consider are contained in a single coordinate chart, as explained in [Proposition 30](#).

Remark 27. The constant ε_0 in [Theorem 17](#) will depend only on γ_0 and ϱ_0 . Therefore, the proof of [Lemma 25](#) tells us that if we consider a whole family of Lagrangian manifolds $(\mathcal{L}_z)_{z \in Z}$ satisfying [Hypotheses 13](#) and [16](#), we will be able to find an $\varepsilon_0 > 0$ uniform in $z \in Z$ provided we have the following uniform transversality condition:

$$\forall t \in \mathbb{N}, \forall \rho \in K, \exists \delta, \gamma > 0 \text{ such that } \forall z \in Z, \Phi^t(\mathcal{L}_z) \cap \tilde{U}^\rho(\varepsilon, \delta) \text{ is } \gamma\text{-unstable.} \tag{36}$$

Lemma 28. *There exists a neighbourhood \mathcal{W}_3 of $\Gamma^- \cap \mathcal{W}_1$ in \mathcal{E} , a finite set of points $(\rho_i)_{i \in I} \subset K$ and $0 < \varepsilon_1 < \varrho_1$, such that the following holds:*

- (i) *The sets $(\tilde{U}_i)_{i \in I} := (\tilde{U}^{\rho_i}(\varepsilon, \varrho_2))_{i \in I}$ form an open cover of a neighbourhood of \mathcal{W}_3 .*
- (ii) $\rho \in [\mathcal{W}_1 \setminus \mathcal{W}_3] \cup \{\rho' \in \mathcal{W}_2 : d(\rho', \Gamma^-) > d_1\} \implies \forall t \geq 0, d(\Phi^t(\rho), K) \geq \varepsilon_1$.
- (iii) *For any open set W of diameter $< \varepsilon_1$ included in \mathcal{W}_3 , there exists an $i \in I$ such that $W \subset \tilde{U}_i$.*

Proof. The sets $(\tilde{U}^\rho(\varepsilon, \varrho_2))_{\rho \in K}$ form an open cover of a neighbourhood of $(\Gamma^- \cap \mathcal{W}_1)$. Let us denote by \mathcal{W}_3 such a neighbourhood.

By compactness, we may extract from it a finite open cover $(\tilde{U}_i)_{i \in I} := (\tilde{U}^{\rho_i}(\varepsilon, \varrho_2))_{i \in I}$, which still satisfies (i).

Since \mathcal{W}_3 is a neighbourhood of $\Gamma^- \cap \mathcal{W}_1$, there exists a constant $\varrho'_2 > 0$ such that the following holds:

$$\forall \rho \in \mathcal{W}_1 \setminus \mathcal{W}_3, \quad d(\rho, \Gamma^-) > \varrho'_2.$$

Therefore, there exists $0 < \varepsilon_1 < \min(\varrho_1, \varepsilon)$ such that

$$\rho \in [\mathcal{W}_1 \setminus \mathcal{W}_3] \cup \{\rho' \in \mathcal{W}_2 : d(\rho', \Gamma^-) \geq d_1\} \implies \forall t \geq 0, d(\Phi^t(\rho), K) > \varepsilon_1,$$

which is (ii). Finally, since the set \tilde{U}_i are open, we may shrink ε_1 so that (iii) is satisfied. □

Remark 29. The constant ε_0 appearing in [Theorem 17](#) will be smaller than ε_1 (see [Lemma 33](#)); therefore each of the sets $(W_a)_{a \in A_1}$ will be contained in some \tilde{U}_i . Furthermore, we will have $W_a \subset \{\rho \in \mathcal{E} : d(\rho, K) < \varepsilon_0\}$. Hence, a point $\rho \in [\mathcal{W}_1 \setminus \mathcal{W}_3] \cup \{\rho' \in \mathcal{W}_2 : d(\rho', \Gamma^-) \geq d_1\}$ will not be contained in any of the sets $(W_a)_{a \in A_1}$ when propagated in the future.

[Lemma 25](#) tells us that $\Phi^{N\varepsilon}(\mathcal{L}_0) \cap \tilde{U}_i$ consists of finitely many γ_0 -unstable Lagrangian manifolds. Our aim will now be to take a Lagrangian manifold included in a \tilde{U}_{i_1} , to propagate it during some time $N \geq N_1$, then to restrict it to a \tilde{U}_{i_2} for $i_1, i_2 \in I$. The remaining part of the Lagrangian, which is in $\mathcal{W}_1 \setminus \mathcal{W}_3$, will not meet the sets $(W_a)_{a \in A_1}$ when propagated in the future, as explained in [Remark 29](#).

2.3.5. Propagation in the sets \tilde{U}_i . For $N \in \mathbb{N}$ and $\iota = (i_0 i_1 \cdots i_{N-1}) \in I^N$, we define

$$\Phi_\iota(\Lambda) := \Phi^1(\tilde{U}_{i_{N-1}} \cap \Phi^1(\cdots \Phi^1(\tilde{U}_{i_0} \cap \Lambda) \cdots)).$$

The propagation of Lagrangian manifolds in the sets \tilde{U}_i is described in the following proposition, which is the cornerstone of the proof of [Theorem 17](#). Recall that γ_{uns} was chosen arbitrarily at the beginning of the proof, and that N_1 was defined in [\(34\)](#).

Proposition 30. *Let $N \geq N_1$, $\iota = (i_0 i_1 \cdots i_{N-1}) \in I^N$ and $i \in I$. Let $\Lambda^0 \subset \tilde{U}_{i_0}$ be an isoenergetic Lagrangian manifold which is γ_0 -unstable in the twisted coordinates centred on ρ_{i_0} . Then $\tilde{U}_i \cap \Phi_\iota(\Lambda)$ is a Lagrangian manifold contained in \tilde{U}_i , and it is $(\gamma_{\text{uns}}/(1 + 2C_0\epsilon^p)^2)$ -unstable in the twisted coordinates centred on ρ_i .*

Proof. The first part of the proof consists in understanding how $\Phi^n(\Lambda^0)$ behaves for $n \leq N_1$, in the twisted coordinates centred on ρ_{i_0} . This is the content of the following lemma, which is an adaptation to our context of the ‘‘inclination lemma’’. (See [\[Katok and Hasselblatt 1995, Theorem 6.2.8\]](#); see also [\[Nonnenmacher and Zworski 2009, Proposition 5.1\]](#) for a statement closer to our context and notation.)

Lemma 31. $\Phi^{N_1}(\Lambda^0)$ is a Lagrangian manifold, which can be written in the chart $(\tilde{y}^{\Phi^{N_1}(\rho_{i_0})}, \tilde{\eta}^{\Phi^{N_1}(\rho_{i_0})})$ in the form

$$\Phi^{N_1}(\Lambda^0) \equiv \{(\tilde{y}_1^{\Phi^{N_1}(\rho_{i_0})}, \tilde{y}^{\Phi^{N_1}(\rho_{i_0})}; 0, f^{N_1}(\tilde{y}^{\Phi^{N_1}(\rho_{i_0})})) : \tilde{y}^{\Phi^{N_1}(\rho_{i_0})} \in D_{N_1}\},$$

with $D_{N_1} \subset B(0, \varrho_1)$ and $\|df^{N_1}\|_{C^0(D_k)} \leq (1 + \nu_1)\gamma_{\text{uns}}/4$.

Note that $\Phi^{N_1}(\Lambda^0)$ is a priori not contained in a single set \tilde{U}_i , but the lemma states that it is contained in the set $\tilde{U}^{\Phi^{N_1}(\rho_{i_0})}(\epsilon, \varrho_1)$, where the twisted coordinates are well defined.

Proof. By assumption, Λ^0 may be put in the form

$$\Lambda^0 \equiv \{(\tilde{y}_1^{\rho_{i_0}}, \tilde{y}^{\rho_{i_0}}; 0, f^0(\tilde{y}^{\rho_{i_0}})) : |\tilde{y}^{\rho_{i_0}}| < \varrho_2\}, \quad \text{with } \|df^0(\tilde{y}^{\rho_{i_0}})\|_{C^0} \leq \gamma_0.$$

We will consider restrictions of the Lagrangian manifolds at intermediate times to the Poincaré sections centred at $\Phi^k(\rho_{i_0})$:

$$\Lambda_{\text{sec}}^k := \Phi^k(\Lambda^0) \cap \Sigma^{\Phi^k(\rho_{i_0})}(\epsilon, \varrho_0).$$

We have $\Lambda_{\text{sec}}^{k+1} = \kappa^k(\Lambda_{\text{sec}}^k)$, where $\kappa^k := \kappa_{\Phi^k(\rho_{i_0}), \Phi^{k+1}(\rho_{i_0})}$ is of the form [\(24\)](#). From equation [\(24\)](#) and the definition of \mathcal{C} , we see that the maximal rate of expansion in the unstable direction is bounded by $(\mathcal{C} + C_0\epsilon^p)$. Therefore, the definition of ϱ_2 implies that for any $k \leq N_1$, the projection of Λ_{sec}^k on the unstable direction is supported in $B(0, \varrho_1)$.

To lighten the notations, we will write \tilde{y}^k and $\tilde{\eta}^k$ instead of $\tilde{y}^{\Phi^k(\rho_{i_0})}$ and $\tilde{\eta}^{\Phi^k(\rho_{i_0})}$.

Let $k \geq 0$, and suppose we may write

$$\Lambda_{\text{sec}}^k \equiv \{(\tilde{y}^k, f^k(\tilde{y}^k)) : \tilde{y}^k \in D_k\},$$

where $D_k \subset B(0, \varrho_1)$, and $\|df^k\|_{C^0} \leq \gamma_k$ for some $0 < \gamma_k \leq \gamma_0$.

Note that the key point in the following computations is that, since we have chosen ‘‘alternative’’ coordinates, we have $|\partial_{\tilde{\eta}} \tilde{\alpha}^k(\tilde{y}^k, \tilde{\eta}^k)| \leq C_0 \tilde{y}^k \leq C_0 \varrho_1$.

The projection of $\Phi^1_{|\Lambda_{\text{sec}}^k}$ on the horizontal subspace is given by

$$\tilde{y}^k \mapsto \tilde{y}^{k+1} = \pi \Phi^1(\tilde{y}^k, f^k(\tilde{y}^k)) = A_k \tilde{y}^k + \tilde{\alpha}^k(\tilde{y}^k, f^k(\tilde{y}^k)),$$

where for each k , we have A_k is a matrix as in (23).

By differentiating, we obtain

$$\frac{\partial \tilde{\mathbf{y}}^{k+1}}{\partial \tilde{\mathbf{y}}^k} = A_k + \frac{\partial \tilde{\alpha}^k}{\partial \tilde{\mathbf{y}}^k} + \frac{\partial \tilde{\alpha}^k}{\partial \tilde{\eta}^k} \frac{\partial f_k}{\partial \tilde{\mathbf{y}}^k} = A_k + r_k,$$

where r_k has entries bounded by $C_0 \varrho_1 \gamma_0 \leq C_0 \epsilon$.

Therefore, the map is invertible, and $\tilde{\mathbf{y}}^{k+1} \mapsto \tilde{\mathbf{y}}^k$ is contracting. This implies that $\Lambda_{\text{sec}}^{k+1}$ can be represented as a graph

$$\Lambda_{\text{sec}}^{k+1} \equiv \{(\tilde{\mathbf{y}}^{k+1}, f^{k+1}(\tilde{\mathbf{y}}^{k+1})) : \tilde{\mathbf{y}}^{k+1} \in D_{k+1}\},$$

with

$$f^{k+1}(\tilde{\mathbf{y}}^{k+1}) = {}^t A_k^{-1} f^k(\tilde{\mathbf{y}}^k) + \tilde{\beta}_k(\tilde{\mathbf{y}}^k, f^k(\tilde{\mathbf{y}}^k)).$$

Differentiating with respect to $\tilde{\mathbf{y}}^{k+1}$, we get

$$\frac{\partial f^{k+1}}{\partial \tilde{\mathbf{y}}^{k+1}} = \left(\frac{\partial \tilde{\mathbf{y}}^k}{\partial \tilde{\mathbf{y}}^{k+1}} \right) \left[({}^t A_k^{-1} + \partial_{\tilde{\eta}} \tilde{\beta}^k(\tilde{\mathbf{y}}^k, f^k(\tilde{\mathbf{y}}^k))) \frac{\partial f^k}{\partial \tilde{\mathbf{y}}^k}(\tilde{\mathbf{y}}^k) + \partial_{\tilde{\mathbf{y}}} \tilde{\beta}^k(\tilde{\mathbf{y}}^k, f^k(\tilde{\mathbf{y}}^k)) \right].$$

Therefore, we have

$$\begin{aligned} \left\| \frac{\partial f^{k+1}}{\partial \tilde{\mathbf{y}}^{k+1}} \right\| &\leq \frac{\|{}^t A_k^{-1}\| \gamma_k + |\partial_{\tilde{\mathbf{y}}} \tilde{\beta}| + |\partial_{\tilde{\eta}} \tilde{\beta}| \gamma_k}{\nu^{-1} - |\partial_{\tilde{\mathbf{y}}} \tilde{\alpha}| - |\partial_{\tilde{\eta}} \tilde{\beta}| \gamma_k} \leq \frac{\gamma_k \nu + C_0 \epsilon^p (1 + \gamma_k)}{\nu^{-1} - 2C_0 \epsilon^p} \\ &\leq \nu_1 \gamma_k + \frac{(1 - \nu_1) \gamma_{\text{uns}}}{8} = \gamma_k \left(\nu_1 + \frac{\gamma_{\text{uns}}(1 - \nu_1)}{8\gamma_k} \right), \end{aligned}$$

where the last inequality comes from (31). First of all, the fact that this slope is bounded uniformly on $\Lambda_{\text{sec}}^{k+1}$ implies that $\Lambda_{\text{sec}}^{k+1}$ can indeed be written in the form

$$\Lambda_{\text{sec}}^{k+1} \equiv \{(\tilde{\mathbf{y}}^{k+1}, f^{k+1}(\tilde{\mathbf{y}}^{k+1})) : \tilde{\mathbf{y}}^{k+1} \in D_{k+1}\},$$

where $D_{k+1} \subset B(0, \varrho_1)$, and $\|df^{k+1}\|_{C^0} \leq \gamma_{k+1}$, where

$$\gamma_{k+1} \leq \gamma_k \left(\nu_1 + \frac{\gamma_{\text{uns}}(1 - \nu_1)}{8\gamma_k} \right).$$

Now, if $\gamma_k > \gamma_{\text{uns}}/4$, then

$$\nu_1 + \frac{\gamma_{\text{uns}}(1 - \nu_1)}{8\gamma_k} < \frac{1 + \nu_1}{2} < 1,$$

so that γ_k decreases exponentially fast, while if $\gamma_k \leq (1 + \nu_1)\gamma_{\text{uns}}/4$, then $\gamma_{k+1} < (1 + \nu_1)\gamma_{\text{uns}}/4$.

The time N_1 has been chosen large enough so that $\gamma_{N_1} < (1 + \nu_1)\gamma_{\text{uns}}/4$, which concludes the proof of the lemma. \square

After times $N > N_1$, the Lagrangian manifold may not be included in $\tilde{U}^{\Phi^N(\rho_{i_0})}(\epsilon, \varrho_1)$. Therefore, we may have to use a change of coordinates. By Lemma 31, at time N_1 , our Lagrangian manifold $\Phi^{N_1}(\Lambda^0)$ is included in $\tilde{U}^{\Phi^{N_1}(\rho_{i_0})}(\epsilon, \varrho_1)$ and is $((1 + \nu_1)\gamma_{\text{uns}}/4)$ -unstable.

We want to study $\tilde{U}_j \cap \Phi^{N_1}(\Lambda^0)$ for $j \in I$ in the coordinates centred at ρ_j , and to apply the computations made in the proof of [Lemma 31](#) again. Let us see how all this works.

If, for some $j \in I$, we have $\tilde{U}_j \cap \Phi^{N_1}(\Lambda^0) \neq \emptyset$, then $d(\Phi^{N_1}(\rho_{i_0}), \rho_j) < \epsilon$. Now, by applying [Lemma 23](#) as well as equation (32), we obtain that $\Phi^{N_1}(\Lambda^0) \cap \tilde{U}_j$ is $(\gamma_{\text{uns}}/2)$ -unstable in the twisted coordinates centred at ρ_j .

We may continue this argument of changing coordinates and propagating to any time $N \geq N_1$: we always obtain a single Lagrangian manifold which is $((1 + \nu_1)\gamma_{\text{uns}}/4)$ -unstable. This concludes the proof of [Proposition 30](#), because we assumed that $C_0\epsilon^p < 1/2$. □

Remark 32. In [[Nonnenmacher and Zworski 2009](#), Proposition 5.1], the authors prove using the chain rule that for each $\ell \in \mathbb{N}$, there exists a constant C_ℓ large enough such that the following holds. If $i_1, i_2 \in I$ and if $\Lambda \subset \tilde{U}_{i_1}$ is a Lagrangian manifold in some unstable cone, generated by a function f in the coordinates $(\tilde{y}^{\rho_{i_1}}, \tilde{\eta}^{\rho_{i_1}})$ with $\|f\|_{C^\ell} \leq C_\ell$, then $\Phi^1(\Lambda) \cap \tilde{U}_{i_2}$ is a union of finitely many Lagrangian manifolds, all of which are in some unstable cone in the coordinates $(\tilde{y}^{\rho_{i_2}}, \tilde{\eta}^{\rho_{i_2}})$, and are generated by functions with a C^ℓ norm smaller than C_ℓ .

In particular, this shows that on the Lagrangian manifold $\Phi^N(\Lambda)$ described in [Proposition 30](#), the function $s^{\rho_i}(y^{\rho_i})$ has a C^ℓ norm smaller than C_ℓ , where C_ℓ is a constant independent of N .

2.3.6. Properties of the sets $(W_a)_{a \in A_1}$. The following lemma is an adaptation of [Lemma 25](#) to the “straight coordinates”. Note that the main reason why we want to use these straight coordinates is because they are symplectic, which will play a crucial role in the proof of [Theorem 47](#).

Lemma 33. *There exists $\epsilon_0 < \epsilon_1$ such that, if $(W_a)_{a \in A_1}$ is an adapted cover of K of diameter ϵ_0 such that for each $a \in A_1$, we have $W_a \cap \mathcal{W}_0 = \emptyset$, and there exists a point $\rho_a \in W_a \cap K \neq \emptyset$, then there exist $\mathcal{N}_{N_{\text{uns}}} \in \mathbb{N}$ and γ' such that the following holds.*

For each $a \in A_1$, for each $1 \leq N \leq N_{\text{uns}}$, the set $\Phi^N(\mathcal{L}_0) \cap W_a$ consists of at most $\mathcal{N}_{N_{\text{uns}}}$ Lagrangian manifolds, all of which are γ' -unstable in the straight coordinates centred on ρ_a .

Proof. Let us choose $\epsilon_0 > 0$ small enough so that $C_0\epsilon_0\tilde{\gamma}_{N_{\text{uns}}} < 1$ and such that each set of diameter smaller than ϵ_0 and which intersects K is contained in some $\tilde{U}^\rho(\epsilon, \delta)$, with $\delta < \tilde{\delta}_{N_{\text{uns}}}$. By applying [Lemma 25](#), we know that there exists $\mathcal{N}_{N_{\text{uns}}} \in \mathbb{N}$, $\tilde{\delta}_{N_{\text{uns}}} > 0$ and $\tilde{\gamma}_{N_{\text{uns}}} > 0$ such that for all $0 < \delta \leq \tilde{\delta}_{N_{\text{uns}}}$, for all $\rho \in K$ and for all $1 \leq N \leq N_{\text{uns}}$, the set $\Phi^N(\mathcal{L}_0) \cap \tilde{U}^\rho(\epsilon, \delta)$ can be written in the coordinates $(\tilde{y}^\rho, \tilde{\eta}^\rho)$ as the union of at most $\mathcal{N}_{N_{\text{uns}}}$ Lagrangian manifolds, which are all $\tilde{\gamma}_{N_{\text{uns}}}$ -unstable. This gives us the statement in the twisted coordinates. To go to the straight coordinates, we may simply use [Lemma 22](#) thanks to the assumption made on ϵ_0 . □

For any $a \in A_1$, $1 \leq k \leq N_{\text{uns}}$, the set $W_a \cap \Phi^k(\mathcal{L}_0)$ consists of finitely many Lagrangian manifolds. Let us define $d_{a,k}$ as the minimal distance (with respect to the distance d) between the Lagrangian manifolds which make up $W_a \cap \Phi^k(\mathcal{L}_0)$, with the convention that this quantity is equal to $+\infty$ if $W_a \cap \Phi^k(\mathcal{L}_0)$ consists of a single Lagrangian manifold or is empty. We then set

$$d := \min(\epsilon_0, \min_{\substack{a \in A_1 \\ 1 \leq k \leq N_{\text{uns}}}} \{d_{a,k}\}) > 0.$$

Remark 34. If we consider a whole family of Lagrangian manifolds $(\mathcal{L}_z)_{z \in Z}$ satisfying Hypotheses 13 and 16, we will be able to apply Theorem 17 to them with sets $(W_a)_{a \in A_2}$ independent of $z \in Z$ provided the constant d is well-defined, that is to say, provided we have

$$\inf_{\substack{a \in A_1, z \in Z \\ 1 \leq k \leq N_{\text{uns}}}} \{d_{a,k}^z\} > 0, \tag{37}$$

where $d_{a,k}^z$ is the minimal distance between the Lagrangian manifolds which make up $W_a \cap \Phi^k(\mathcal{L}_z)$, with the convention that this quantity is equal to $+\infty$ if $W_a \cap \Phi^k(\mathcal{L}_z)$ consists of a single Lagrangian manifold or is empty.

The flow (Φ^t) is C^1 with respect to time, and hence Lipschitz on $[0, N_{\text{uns}}]$. Therefore, there exists a constant $C > 0$ such that for all $t \in [0, N_{\text{uns}}]$ and for all $\rho_1, \rho_2 \in \mathcal{E}$, we have

$$d(\Phi^t(\rho_1), \Phi^t(\rho_2)) \leq C d(\rho_1, \rho_2).$$

We take

$$\varepsilon_2 := d/C.$$

We now complete $(W_a)_{a \in A_1}$ to cover the whole energy layer.

2.3.7. Construction and properties of the sets $(W_a)_{a \in A_2}$. Recall that $W_0 = T^*(X \setminus X_0)$, and that b is the boundary-defining function introduced in Hypothesis 5.

We build the sets $(W_a)_{a \in A_2}$ so that, if we set $A = A_1 \cup A_2 \cup \{0\}$, the following holds:

- Each of the sets $(W_a)_{a \in A_2}$ has a diameter smaller than ε_2 .
- For each $a \in A_2$, we have $d(W_a, K) > \varepsilon_2/2$.
- $(W_a)_{a \in A}$ is an open cover of \mathcal{E} .

Our next lemma is the first brick of the proof of the uniqueness of the Lagrangian manifold making up $\Phi_\alpha^N(\mathcal{L}_0)$. It relies on the fact that the sets $(W_a)_{a \in A_2}$ have been built small enough.

Lemma 35. *Let $k \leq N_{\text{uns}}$, $\alpha \in A^k$, and $a \in A_1$. Then the set $W_a \cap \Phi_\alpha^k(\mathcal{L}_0)$ is empty or consists of a single Lagrangian manifold.*

Proof. Let us suppose that $\Phi^k(\mathcal{L}_0) \cap W_a$ is nonempty. We have seen in Lemma 33 that it consists of finitely many Lagrangian manifolds, with a distance between them larger than d . Therefore, for any $1 \leq k' \leq k$, the sets $\Phi^{-k'}(\Phi^k(\mathcal{L}_0) \cap W_a)$ consist of Lagrangian manifolds which are at a distance larger than ε_2 from each other. Because of the assumption (9) we made, we have $\alpha_{k'} \in A_2$ for some $k' \leq k$. Since the sets $(W_a)_{a \in A_2}$ have a diameter smaller than ε_2 , they separate the Lagrangian manifolds which make up $\Phi^{-k'}(\Phi^k(\mathcal{L}_0) \cap W_a)$. We deduce from this the lemma. □

2.3.8. Structure of the admissible sequences. We will now state two of lemmas which put some constraints on the sequences $\alpha \in A^N$, with $\alpha_N \in A_1$ such that $\Phi_\alpha^N(\mathcal{L}_0) \neq \emptyset$.

The first of these lemmas tell us that we may restrict ourselves to sequences such that $\alpha_k \neq 0$ for $k \geq 1$.

Lemma 36. *Let $N \in \mathbb{N}$, and let $\alpha \in A^N$, and $a \in A_1$. Suppose that $\alpha_k = 0$ for some $1 \leq k \leq N - 1$, and that $W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \neq \emptyset$. Then*

$$W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \subset \Phi_{\alpha_{k+1} \dots \alpha_{N-1}}^{N-k}(\mathcal{L}_0).$$

Proof. By hypothesis, $\Phi_{\alpha_1 \dots \alpha_k}^k(\mathcal{L}_0) \subset \mathcal{W}_0$, and it intersects \mathcal{W}_1 in the future. We have $\mathcal{W}_0 = \mathcal{D}\mathcal{E}_- \cup \mathcal{D}\mathcal{E}_+$, and a point in $\mathcal{D}\mathcal{E}_+$ cannot intersect \mathcal{W}_1 in the future. Therefore, the points in $\Phi_{\alpha_1 \dots \alpha_k}^k(\mathcal{L}_0)$ which intersect \mathcal{W}_1 in the future are all in $\mathcal{D}\mathcal{E}_-$. But by Lemma 9, the point in $\mathcal{D}\mathcal{E}_-$ can only have preimages in W_0 . Therefore, we have

$$W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \subset W_a \cap \Phi_{0 \dots 0 \alpha_{k+1} \dots \alpha_{N-1}}^N(\mathcal{L}_0) \subset \Phi_{\alpha_{k+1} \dots \alpha_{N-1}}^{N-k}(\mathcal{L}_0),$$

where the second inclusion comes from Hypothesis 13. □

Let us now take advantage of Remark 29 to show that, from time $k \geq N_\epsilon + 2$, all the interesting dynamics takes place in \mathcal{W}_3 .

Lemma 37. *Let $N \geq N_\epsilon + 2$, and $\alpha \in A^N$ with $\alpha_i \neq 0$ for $i \geq 1$.*

Let $N_\epsilon + 2 \leq k \leq N$, and $\rho \in \Phi_{\alpha_1 \dots \alpha_k}^k(\mathcal{L}_0)$ be such that $\Phi^{N-k}(\rho) \in W_a$ for some $a \in A_1$. Then $\rho \in \mathcal{W}_3$.

Proof. If $\rho \in \mathcal{W}_1$, then the result follows from Remark 29. We must therefore check that we cannot have $\rho \in \mathcal{W}_2 \cup \mathcal{W}_0$. First of all, note that Lemma 9 implies that we cannot have $\rho \in \mathcal{W}_0$. This lemma also implies that for each $a' \in A_1 \cup A_2$, we have

$$\Phi^1(W_{a'} \setminus W_0) \cap \mathcal{D}\mathcal{E}_- = \emptyset. \tag{38}$$

Suppose now that $\rho \in \mathcal{W}_2$. Since $k \geq N_\epsilon + 2$, and $\alpha_i \neq 0$ for $i \geq 1$, we have $\Phi^{-N_\epsilon-1}(\rho) \in W_{a'}$ for some $a' \in A_1 \cup A_2$. Therefore, by equation (38), we have $\Phi^{-N_\epsilon}(\rho) \notin \mathcal{W}_0$.

By the proof of Lemma 24, this would imply that $d(\rho, \Gamma^-) \geq d_1$. By Remark 29, this implies that we cannot have $\Phi^{N-k}(\rho) \in W_a$ for some $a \in A_1$, a contradiction. □

2.3.9. End of the proof of Theorem 17. Let $N \geq 0$, $\alpha \in A^N$ and $a \in A_1$. If $N \leq N_{\text{uns}}$, the result of Theorem 17 is a consequence of Lemmas 33 and 35.

Consider now $N \geq N_{\text{uns}} > N_\epsilon + 2$. We will assume that $W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \neq \emptyset$. Thanks to Lemma 36 and to Hypothesis 13, we may assume that $\alpha_i \neq 0$ for all $i \geq 1$.

From Lemma 37, we deduce that

$$W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \subset \bigcup_{\substack{\iota \in I^{N-N_\epsilon-1} \\ \iota_{N-N_\epsilon} = i_\alpha}} \Phi_\iota(\Phi_{\alpha_1 \dots \alpha_{N_\epsilon+2}}^{N_\epsilon+2}(\mathcal{L}_0)), \tag{39}$$

where $i_\alpha \in I$ is such that $W_{\alpha_N} \subset \tilde{U}_{i_\alpha}$.

Let us define

$$\Lambda_k := \{\rho \in \Phi_\alpha^k(\mathcal{L}_0) : \forall k' \geq 0, \Phi^{k'}(\rho) \in W_{\alpha_{k+k'}}\}.$$

By Lemma 37, for each $k \geq N_\epsilon + 2$, we have $\Lambda_k \subset \mathcal{W}_3 \cap W_{\alpha_k}$. Therefore, by Lemma 28(iii), there exists a $i_k \in I$ such that $\Lambda_k \subset \tilde{U}_{i_k}$, and we obtain that

$$W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \subset \Phi_{i_{N_\epsilon+2} \dots i_N}^{N-N_\epsilon-2}(\Phi_{\alpha_1 \dots \alpha_{N_\epsilon+2}}^{N_\epsilon+2}(\mathcal{L}_0)).$$

We know from Lemmas 25 and 35 that $\Phi_{\alpha_1 \dots \alpha_{N_\epsilon+2}}^{N_\epsilon+2}(\mathcal{L}_0)$ consists of a single Lagrangian manifold, which is γ_0 -unstable in the coordinates centred on any point of K . Applying Proposition 30, we know that the right-hand side of (39) is a Lagrangian manifold which is $(\gamma_{\text{uns}}/(1 + 2C_a \epsilon^p)^2)$ -unstable in the twisted coordinates centred on ρ_{i_α} .

We first apply Lemma 23 to write this Lagrangian manifold in the twisted coordinates centred on ρ_a . Thanks to equation (32), it is $(\gamma_{\text{uns}}/(1 + 2C_a \epsilon^p))$ -unstable. We then use Lemma 22 to write this Lagrangian manifold in the straight coordinates centred on ρ_{α_N} , and we deduce that it is γ_{uns} -unstable. This concludes the proof of Theorem 17. □

Remark 38. Therefore, in the coordinates (y^a, η^a) , the set $W_a \cap \Phi_\alpha^N(\mathcal{L}_0)$ may be put in the form

$$W_a \cap \Phi_\alpha^N(\mathcal{L}_0) \equiv \{(y_1^a, y^a; 0, f_{N,\alpha,a}(y^a)) : y^a \in D_{N,\alpha,a}\}$$

for some open set $D_{N,\alpha,a} \subset \mathbb{R}^d$.

Remark 32 tells us that for any $\ell \in \mathbb{N}$, the functions $f_{N,\alpha,a}$ have C^ℓ norms which are bounded independently of N, α and a .

3. Generalized eigenfunctions

We shall state our results about generalized eigenfunctions under rather general assumptions. We shall then explain why these assumptions hold in the case of distorted plane waves on manifolds which are Euclidean near infinity.

In the sequel, we will consider a Riemannian manifold (X, g) with a real-valued potential $V \in C_c^\infty(X)$, and define the Schrödinger operator

$$P_h = -h^2 \Delta_g - c_0 h^2 + V(x).$$

Here $c_0 > 0$ is a constant, which will be 0 in the case of Euclidean-near-infinity manifolds (see Section 3.3 for the definition of such manifolds).

Before stating our assumptions, let us recall a few definitions and facts from semiclassical analysis.

3.1. Refresher on semiclassical analysis.

3.1.1. Pseudodifferential calculus. We shall use the class $S^{\text{comp}}(T^*X)$ of symbols $a \in C_c^\infty(T^*X)$, which may depend on h , but whose seminorms and supports are all bounded independently of h . We will sometimes write $S^{\text{comp}}(X)$ for the set of symbols in $S^{\text{comp}}(T^*X)$ which depend only on the base variable. If U is an open subset of T^*X , we will denote by $S^{\text{comp}}(U)$ the set of functions in $S^{\text{comp}}(T^*X)$ whose support is contained in U .

Definition 39. Let $a \in S^{\text{comp}}(T^*Y)$. We will say that a is a *classical symbol* if there exists a sequence of symbols $a_k \in S^{\text{comp}}(T^*Y)$ such that for any $n \in \mathbb{N}$,

$$a - \sum_{k=0}^n h^k a_k \in h^{n+1} S^{\text{comp}}(T^*Y).$$

We will then write

$$a^0(x, \xi) := \lim_{h \rightarrow 0} a(x, \xi; h)$$

for the *principal symbol* of a .

We associate to $S^{\text{comp}}(T^*X)$ the class of pseudodifferential operators $\Psi_h^{\text{comp}}(X)$, through a surjective quantization map

$$\text{Op}_h : S^{\text{comp}}(T^*X) \rightarrow \Psi_h^{\text{comp}}(X).$$

This quantization map is defined using coordinate charts, and the standard Weyl quantization on \mathbb{R}^d . It is therefore not intrinsic. However, the principal symbol map

$$\sigma_h : \Psi_h^{\text{comp}}(X) \rightarrow S^{\text{comp}}(T^*X)/hS^{\text{comp}}(T^*X)$$

is intrinsic, and we have

$$\sigma_h(A \circ B) = \sigma_h(A)\sigma_h(B)$$

and

$$\sigma_h \circ \text{Op} : S^{\text{comp}}(T^*X) \rightarrow S^{\text{comp}}(T^*X)/hS^{\text{comp}}(T^*X)$$

is the natural projection map.

For more details on all these maps and their construction, we refer the reader to [Zworski 2012, Chapter 14].

For $a \in S^{\text{comp}}(T^*X)$, we say its essential support is equal to a given compact $K \Subset T^*X$, denoted by

$$\text{ess supp}_h a = K \Subset T^*X,$$

if and only if, for all $\chi \in S(T^*X)$,

$$\text{supp } \chi \subset (T^*X \setminus K) \implies \chi a \in h^\infty S(T^*X).$$

For $A \in \Psi_h^{\text{comp}}(X)$, $A = \text{Op}_h(a)$, we define the wave front set of A as

$$\text{WF}_h(A) = \text{ess supp}_h a,$$

noting that this definition does not depend on the choice of the quantisation. When K is a compact subset of T^*X and $\text{WF}_h(A) \subset K$, we will sometimes say that A is *microsupported* inside K .

Let us now state a lemma which is a consequence of Egorov theorem [Zworski 2012, Theorem 11.1]. Recall that $U(t)$ is the Schrödinger propagator $U(t) = e^{-itP_h/h}$.

Lemma 40. Let $A, B \in \Psi_h^{\text{comp}}(X)$, and suppose that $\Phi^t(\text{WF}_h(A)) \cap \text{WF}_h(B) = \emptyset$. Then we have

$$AU(t)B = O_{L^2 \rightarrow L^2}(h^\infty).$$

If U, V are bounded open subsets of T^*X , and if $T, T' : L^2(X) \rightarrow L^2(X)$ are bounded operators, we shall say that $T \equiv T'$ *microlocally* near $U \times V$ if there exist bounded open sets $\tilde{U} \supset \bar{U}$ and $\tilde{V} \supset \bar{V}$ such that for any $A, B \in \Psi_h^{\text{comp}}(X)$ with $\text{WF}(A) \subset \tilde{U}$ and $\text{WF}(B) \subset \tilde{V}$, we have

$$A(T - T')B = O_{L^2 \rightarrow L^2}(h^\infty)$$

Tempered distributions. Let $u = (u(h))$ be an h -dependent family of distributions in $\mathcal{D}'(X)$. We say it is h -tempered if for any bounded open set $U \subset X$, there exists $C > 0$ and $N \in \mathbb{N}$ such that

$$\|u(h)\|_{H_h^{-N}(U)} \leq Ch^{-N},$$

where $\|\cdot\|_{H_h^{-N}(U)}$ is the semiclassical Sobolev norm.

For a tempered distribution $u = (u(h))$, we say that a point $\rho \in T^*X$ does not lie in the wave front set $\text{WF}(u)$ if there exists a neighbourhood V of ρ in T^*X such that for any $A \in \Psi_h^{\text{comp}}(X)$ with $\text{WF}(a) \subset V$, we have $Au = O(h^\infty)$.

3.1.2. Lagrangian distributions and Fourier integral operators.

Phase functions. Let $\phi(x, \theta)$ be a smooth real-valued function on some open subset U_ϕ of $X \times \mathbb{R}^L$ for some $L \in \mathbb{N}$. We call x the *base variable* and θ the *oscillatory variable*. We say that ϕ is a *nondegenerate phase function* if the differentials $d(\partial_{\theta_1}\phi) \cdots d(\partial_{\theta_L}\phi)$ are linearly independent on the *critical set*

$$C_\phi := \{(x, \theta) : \partial_\theta \phi = 0\} \subset U_\phi.$$

In this case

$$\Lambda_\phi := \{(x, \partial_x \phi(x, \theta)) : (x, \theta) \in C_\phi\} \subset T^*X$$

is an immersed Lagrangian manifold. By shrinking the domain of ϕ , we can make it an embedded Lagrangian manifold. We say that ϕ *generates* Λ_ϕ .

Lagrangian distributions. Given a phase function ϕ and a symbol $a \in S^{\text{comp}}(U_\phi)$, consider the h -dependent family of functions

$$u(x; h) = h^{-L/2} \int_{\mathbb{R}^L} e^{i\phi(x, \theta)/h} a(x, \theta; h) d\theta. \tag{40}$$

We call $u = (u(h))$ a *Lagrangian distribution*, (or a *Lagrangian state*) generated by ϕ . By the method of nonstationary phase, if $\text{supp } a$ is contained in some h -independent compact set $K \subset U_\phi$, then

$$\text{WF}_h(u) \subset \{(x, \partial_x \phi(x, \theta)) : (x, \theta) \in C_\phi \cap K\} \subset \Lambda_\phi.$$

Definition 41. Let $\Lambda \subset T^*X$ be an embedded Lagrangian submanifold. We say that an h -dependent family of functions $u(x; h) \in C_c^\infty(X)$ is a (compactly supported and compactly microlocalized) *Lagrangian distribution associated to* Λ , if it can be written as a sum of finitely many functions of the form (40), for different phase functions ϕ parametrizing open subsets of Λ , plus an $O(h^\infty)$ remainder. We will denote by $I^{\text{comp}}(\Lambda)$ the space of all such functions.

Fourier integral operators. Let X, X' be two manifolds of the same dimension d , and let κ be a symplectomorphism from an open subset of T^*X to an open subset of T^*X' . Consider the Lagrangian

$$\Lambda_\kappa = \{(x', -v'; x, v) : \kappa(x, v) = (x', v')\} \subset T^*X' \times T^*X = T^*(X' \times X).$$

A compactly supported operator $U : \mathcal{D}'(X) \rightarrow C_c^\infty(X')$ is called a (semiclassical) *Fourier integral operator* associated to κ if its Schwartz kernel $K_U(x', x)$ lies in $h^{-d/2}I^{\text{comp}}(\Lambda_\kappa)$. We write $U \in I^{\text{comp}}(\kappa)$. The $h^{-d/2}$ factor is explained as follows: the normalization for Lagrangian distributions is chosen so that $\|u\|_{L^2} \sim 1$, while the normalization for Fourier integral operators is chosen so that $\|U\|_{L^2(X) \rightarrow L^2(X')} \sim 1$.

Note that if $\kappa \circ \kappa'$ is well defined, and if $U \in I^{\text{comp}}(\kappa)$ and $U' \in I^{\text{comp}}(\kappa')$, then $U \circ U' \in I^{\text{comp}}(\kappa \circ \kappa')$.

If $U \in I^{\text{comp}}(\kappa)$ and $O \subset T^*X$ is an open bounded set, we shall say that U is *microlocally unitary* near O if $U^*U \equiv I_{L^2(X) \rightarrow L^2(X)}$ microlocally near $O \times \kappa(O)$.

3.1.3. Local properties of Fourier integral operators. In this section we shall see that, if we work locally, we may describe many Fourier integral operators without the help of oscillatory coordinates. In particular, following [Nonnenmacher and Zworski 2009, §4.1], we will recall the effect of a Fourier integral operator on a Lagrangian distribution which has no caustics. We will recall in Section 4.2 how this formalism may be applied to the study of the Schrödinger propagator.

Let $\kappa : T^*\mathbb{R}^d \rightarrow T^*\mathbb{R}^d$ be a local symplectic diffeomorphism. By performing phase-space translations, we may assume that κ is defined in a neighbourhood of $(0, 0)$ and that $\kappa(0, 0) = (0, 0)$.

Without loss of generality, we can find linear Lagrangian subspaces, $\Gamma_j, \Gamma_j^\perp \subset T^*\mathbb{R}^d$, $j = 0, 1$, with the following properties:

- Γ_j^\perp is transversal to Γ_j .
- If π_j (resp. π_j^\perp) is the projection $T^*\mathbb{R}^d \rightarrow \Gamma_j$ along Γ_j^\perp (resp. the projection $T^*\mathbb{R}^d \rightarrow \Gamma_j^\perp$ along Γ_j), then, for some neighbourhood U of ρ_0 , the map

$$\kappa(U) \times U \rightarrow \Gamma_1 \times \Gamma_0^\perp, \quad (\kappa(\rho), \rho) \mapsto \pi_1(\kappa(\rho)) \times \pi_0^\perp,$$

is a local diffeomorphism from the graph of $\kappa|_U$ to a neighbourhood of the origin in $\Gamma_1 \times \Gamma_0^\perp$.

Let $A_j, j = 0, 1$ be linear symplectic transformations with the properties

$$A_j(\Gamma_j) = \{(x, 0)\} \subset T^*\mathbb{R}^d \quad \text{and} \quad A_j(\Gamma_j^\perp) = \{(0, \xi)\} \subset T^*\mathbb{R}^d,$$

and let M_j be *metaplectic quantizations* of the A_j as defined in [Dimassi and Sjöstrand 1999, Appendix to Chapter 7]. Then the rotated diffeomorphism

$$\tilde{\kappa} := A_1 \circ \kappa \circ A_0^{-1}$$

is such that the projection from the graph of $\tilde{\kappa}$

$$T^*\mathbb{R}^d \times T^*\mathbb{R}^d \rightarrow \mathbb{R}^d \times \mathbb{R}^d, \quad (x^1, \xi^1; x^0, \xi^0) \mapsto (x^1, \xi^0), \quad (x^1, \xi^1) = \tilde{\kappa}(x^0, \xi^0), \quad (41)$$

is a diffeomorphism near the origin. It then follows that there exists a unique function $\tilde{\psi} \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ such that for (x^1, ξ^0) near $(0, 0)$,

$$\tilde{\kappa}(\tilde{\psi}'_\xi(x^1, \xi^0), \xi^0) = (x^1, \tilde{\psi}'_x(x^1, \xi^0)), \quad \det \tilde{\psi}''_{x\xi} \neq 0 \quad \text{and} \quad \tilde{\psi}(0, 0) = 0.$$

The function $\tilde{\psi}$ is said to *generate* the transformation $\tilde{\kappa}$ near $(0, 0)$.

Note that if $\tilde{T} \in I^{\text{comp}}(\tilde{\kappa})$, then

$$T := M_1^{-1} \circ \tilde{T} \circ M_0 \in I^{\text{comp}}(\kappa). \tag{42}$$

Thanks to assumption (41), a Fourier integral operator $\tilde{T} \in I^{\text{comp}}(\tilde{\kappa})$ may then be written in the form

$$\tilde{T}u(x^1) := \frac{1}{(2\pi h)^d} \iint_{\mathbb{R}^{2n}} e^{i(\tilde{\psi}(x^1, \xi^0) - \langle x^0, \xi^0 \rangle / h)} \alpha(x^1, \xi^0; h) u(x^0) dx^0 d\xi^0, \tag{43}$$

with $\alpha \in S^{\text{comp}}(\mathbb{R}^{2d})$.

Now, let us state a lemma which was proven in [Nonnenmacher and Zworski 2009, Lemma 4.1], and which describes the effect of a Fourier integral operator of the form (43) on a Lagrangian distribution which projects on the base manifold without caustics.

Lemma 42. *Consider a Lagrangian $\Lambda_0 = \{(x_0, \phi'_0(x_0)) : x \in \Omega_0\}$, $\phi_0 \in C_b^\infty(\Omega_0)$, contained in a small neighbourhood $V \subset T^*\mathbb{R}^d$ such that κ is generated by ψ near V . We assume that*

$$\kappa(\Lambda_0) = \Lambda_1 = \{(x, \phi'_1(x)) : x \in \Omega_1\}, \quad \phi_1 \in C_b^\infty(\Omega_1).$$

Then, for any symbol $a \in S^{\text{comp}}(\Omega_0)$, the application of a Fourier integral operator T of the form (43) to the Lagrangian state

$$a(x)e^{i\phi_0(x)/h}$$

associated with Λ_0 can be expanded, for any $L > 0$, into

$$T(ae^{i\phi_0/h})(x) = e^{i\phi_1(x)/h} \left(\sum_{j=0}^{L-1} b_j(x)h^j + h^L r_L(x, h) \right),$$

where $b_j \in S^{\text{comp}}$, and for any $\ell \in \mathbb{N}$, we have

$$\|b_j\|_{C^\ell(\Omega_1)} \leq C_{\ell,j} \|a\|_{C^{\ell+2j}(\Omega_0)}, \quad 0 \leq j \leq L-1,$$

$$\|r_L(\cdot, h)\|_{C^\ell(\Omega_1)} \leq C_{\ell,L} \|a\|_{C^{\ell+2L+n}(\Omega_0)}.$$

The constants $C_{\ell,j}$ depend only on κ , α and $\sup_{\Omega_0} |\partial^\beta \phi_0|$ for $0 < |\beta| \leq 2\ell + j$.

3.2. Assumptions on the generalized eigenfunctions. We consider generalized eigenfunctions of P_h at energy 1, that is to say, a family of smooth functions $E_h \in C^\infty(X)$ indexed by $h \in (0, 1]$ which satisfy

$$(P_h - 1)E_h = 0.$$

We will furthermore assume that these generalized eigenfunctions may be decomposed as follows.

Hypothesis 43. *We suppose that E_h can be put in the form*

$$E_h = E_h^0 + E_h^1, \tag{44}$$

where E_h^0 is a tempered distribution which is a Lagrangian state associated to a Lagrangian manifold which satisfies Hypothesis 13 of invariance, as well as Hypothesis 16 of transversality, and where E_h^1 is a tempered distribution such that for each $\rho \in \text{WF}_h(E_h^1)$, we have $\rho \in \mathcal{E}$.

Furthermore, we suppose that E_h^1 is **outgoing** in the sense that there exists $\epsilon_2 > 0$ such that for all $\chi, \chi' \in C_c^\infty$ such that $\chi \equiv 1$ on $\{x \in X : b(x) \geq \epsilon_2\}$, there exists $T_\chi > 0$ such that for all $t \geq T_\chi$, we have

$$\Phi^t(\text{WF}((1 - \chi)\chi' E_h^1)) \cap \text{spt}(\chi) = \emptyset. \tag{45}$$

The most natural example of such generalized eigenfunctions is given by distorted plane waves, which we are now going to define. Note that they depend on a parameter $\xi \in \partial\bar{X}$, so that they actually form a whole family of generalized eigenfunctions.

It is also possible to define generalized eigenfunctions which satisfy [Hypothesis 43](#) on manifolds which are hyperbolic near infinity. This is done in [[Ingremeau 2017](#), Appendix B]; the construction mainly follows [[Dyatlov and Guillarmou 2014](#), §6], but some work has to be done to check that E_h^1 is a tempered distribution.

3.3. Distorted plane waves on Euclidean-near-infinity manifolds.

Definition 44. We say that X is *Euclidean near infinity* if there exists a compact set $X_0 \subset X$ and a $R_0 > 0$ such that $X \setminus X_0$ has finitely many connected components, which we denote by X_1, \dots, X_l , such that for each $1 \leq i \leq l$, we have (X_i, g) is isometric to $(\mathbb{R}^d \setminus B(0, R_0), g_{\text{Eucl}})$.

The surface in [Figure 2](#) is an example of a Euclidean-near-infinity manifold. We may assume that $\text{supp } V \subset X_0$. Also, any Euclidean-near-infinity manifold fulfills [Hypothesis 5](#). Indeed, we may take a boundary-defining function b such that $b(x) = (1 + |x|^2)^{-1/2}$ if $x \in X_i$ which we identify with $\mathbb{R}^d \setminus B(0, R_0)$.

To define distorted plane waves, we will simply give a definition of each of the two terms which compose them as in [\(44\)](#).

3.3.1. Definition of E_h^0 . By definition of a Euclidean-near-infinity manifold, we have

$$X = X_0 \sqcup \left(\bigsqcup_{i=1}^N X_i \right)$$

with X_0 compact, and for each $1 \leq i \leq N$, there exists an isometric isomorphism

$$x_i : X_i \rightarrow \mathbb{R}^d \setminus B(0, R_0), \tag{46}$$

equipped with the Euclidean metric g_0 .

The boundary of \bar{X} may then be identified with a union of spheres:

$$\partial X \cong \bigsqcup_{i=1}^N S_i,$$

with $S_i \cong \mathbb{S}^n$.

Let $\xi \in \partial\bar{X}$. We have $\xi \in S_i$ for some $1 \leq i \leq m$. Take a smooth function $\tilde{\chi} : X \rightarrow [0, 1]$ which vanishes outside of X_i , and which is equal to 1 in a neighbourhood of S_i .

We define the incoming wave E_h^0 by $E_h^0(\xi, \cdot) : X \rightarrow \mathbb{C}$ by

$$E_h^0(\xi, x) = \begin{cases} \tilde{\chi}(x) e^{\frac{i}{h} x_i(x) \cdot \xi} & \text{if } x \in X_i, \\ 0 & \text{otherwise.} \end{cases}$$

If we write \mathcal{L}_0 for the Lagrangian submanifold (with boundaries) $X_i \times \{\xi\} \subset T^*X$, then E_h^0 is a Lagrangian distribution associated to \mathcal{L}_0 , which satisfies [Hypothesis 13](#) of invariance.

3.3.2. Definition of the distorted plane waves. Let us set

$$F_h := -[P_h, \tilde{\chi}]E_h^0(\xi).$$

Note that we have $F_h \in S^{\text{comp}}(X)$.

Recall that the outgoing resolvent $R_h(1)$ is defined as $R_h(1) := \lim_{\epsilon \rightarrow 0^+} (P_h - (1 + i\epsilon)^2)^{-1}$, the limit being taken in the topology of bounded operators from $L^2_{\text{comp}}(X)$ to $L^2_{\text{loc}}(X)$.

We shall use the following resolvent estimate, which was proved in [\[Nonnenmacher and Zworski 2009\]](#).

Theorem 45 (resolvent estimates for Euclidean-near-infinity manifolds). *Let X be a Euclidean-near-infinity manifold such that [Hypothesis 10](#) on hyperbolicity and [Hypothesis 46](#) on topological pressure hold. Then for any $\chi \in C_c^\infty(X)$, there exists $C > 0$ such that for all $0 < h < h_0$, we have*

$$\|\chi R_h(1)\chi\|_{L^2(X) \rightarrow L^2(X)} \leq C \frac{\log(1/h)}{h}. \tag{47}$$

We define

$$E_h^1 := R_h(1)F_h,$$

which is a tempered distribution thanks to [Theorem 45](#).

We then define the distorted plane wave as

$$E_h^\xi := E_h^0 + E_h^1.$$

To check the outgoing assumption on E_h^1 , we must explain why there exists $\epsilon_2 > 0$ such that for all $\chi, \chi' \in C_c^\infty$ with $\chi \equiv 1$ on $\{x \in X : b(x) \geq \epsilon_2\}$, there exists $T_\chi > 0$ such that for all $t > T_\chi$, we have

$$\Phi^t(\text{WF}((1 - \chi)\chi'E_h^1)) \cap \text{spt}(\chi) = \emptyset. \tag{48}$$

From [\[Dyatlov and Guillarmou 2014, §6.2\]](#), we know that for any $\rho \in \text{WF}_h(E_h^1)$, we have $\rho \in \mathcal{E}$, and either $\rho \in \Gamma^+$ or there exists a $t > 0$ such that $\Phi^{-t}(\rho) = (x, \xi)$ where $x \in \text{spt}(\partial\tilde{\chi})$, where $\tilde{\chi}$ is as in [Section 3.3.1](#).

We may take $\epsilon_2 < \epsilon_0$ small enough so that $\text{spt}(\tilde{\chi}) \subset \{x \in X : b(x) > \epsilon_2\}$. Suppose that $\rho = (x, \xi)$ is such that $x \in \text{spt}(1 - \chi)$ and $\pi_X(\Phi^t(\rho)) \in \text{spt}(\chi)$. Then, by geodesic convexity, $(x, -\xi) \in \mathcal{DE}_+$. Therefore, since $\text{spt}(\tilde{\chi}) \subset \{x \in X : b(x) > \epsilon_2\}$ and $\text{spt}(1 - \chi) \subset \{x \in X : b(x) < \epsilon_2\}$ and since b decreases in the future along the trajectory of $(x, -\xi)$, it is impossible that there exists $t > 0$ such that $\Phi^{-t}(\rho) = (x, \xi)$ where $x \in \text{spt}(\partial\tilde{\chi})$. Therefore, if $\rho \in \Phi^t(\text{WF}((1 - \chi)\chi'E_h^1)) \cap \text{spt}(\chi)$, we must have $\rho \in \mathcal{DE}_+$.

On the other hand, if $\rho \in \mathcal{DE}_+$, then [\(48\)](#) is always satisfied as long as T_χ is large enough so that $\Phi^{T_\chi}(\mathcal{DE}_+ \cap T^*(\text{spt}(1 - \chi))) \cap T^*\text{spt}(\chi) = \emptyset$. This shows that E_h^1 is outgoing.

Finally, one readily checks that we have, in the sense of PDEs,

$$(P_h - 1)E_h^\xi = 0.$$

We will sometimes simply write E_h instead of E_h^ξ , to avoid cumbersome notations.

The definition of E_h seems to depend on the choices of the cut-off functions we made. Actually, the distorted plane waves can be defined in a much more intrinsic fashion, using the structure of the resolvent at infinity. We don't want to enter into the details here (see [Dyatlov and Guillarmou 2014, §6; Melrose 1995, Chapter 2]).

3.4. Topological pressure. We shall now give a definition of topological pressure, so as to formulate [Hypothesis 46](#). Recall that the distance d was defined in [Section 2.1.2](#), and that it was associated to the adapted metric. We say that a set $\mathcal{S} \subset K$ is (ϵ, t) -separated if for $\rho_1, \rho_2 \in \mathcal{S}$, $\rho_1 \neq \rho_2$, we have $d(\Phi^{t'}(\rho_1), \Phi^{t'}(\rho_2)) > \epsilon$ for some $0 \leq t \leq t'$. (Such a set is necessarily finite.)

The metric g_{ad} induces a volume form Ω on any d -dimensional subspace of $T(T^*\mathbb{R}^d)$. Using this volume form, we will define the unstable Jacobian on K . For any $\rho \in K$, the determinant map

$$\Lambda^n d\Phi^t(\rho)|_{E_\rho^{+0}} : \Lambda^n E_\rho^{+0} \rightarrow \Lambda^n E_{\Phi^t(\rho)}^{+0}$$

can be identified with the real number

$$\det(d\Phi^t(\rho)|_{E_\rho^{+0}}) := \frac{\Omega_{\Phi^t(\rho)}(d\Phi^t v_1 \wedge d\Phi^t v_2 \wedge \dots \wedge d\Phi^t v_n)}{\Omega_\rho(v_1 \wedge v_2 \wedge \dots \wedge v_n)},$$

where (v_1, \dots, v_n) can be any basis of E_ρ^{+0} . This number defines the unstable Jacobian:

$$\exp \lambda_t^+(\rho) := \det(d\Phi^t(\rho)|_{E_\rho^{+0}}). \tag{49}$$

From there, we take

$$Z_t(\epsilon, s) := \sup_{\mathcal{S}} \sum_{\rho \in \mathcal{S}} \exp(-s\lambda_t^+(\rho)),$$

where the supremum is taken over all (ϵ, t) -separated sets. The pressure is then defined as

$$\mathcal{P}(s) := \lim_{\epsilon \rightarrow 0} \limsup_{t \rightarrow \infty} \frac{1}{t} \log Z_t(\epsilon, s).$$

This quantity is actually independent of the volume form Ω and of the metric chosen: after taking logarithms, a change in Ω or in the metric will produce a term $O(1)/t$, which is not relevant in the $t \rightarrow \infty$ limit.

Hypothesis 46. We assume the following inequality on the topological pressure associated with Φ^t on K :

$$\mathcal{P}\left(\frac{1}{2}\right) < 0. \tag{50}$$

We will give an equivalent definition of topological pressure in [Section 4.1](#), better suited to our purpose.

3.5. Statement of the results concerning distorted plane waves. We may now formulate our main result.

Theorem 47. Suppose that the manifold X satisfies [Hypothesis 5](#) at infinity, and that the Hamiltonian flow (Φ^t) satisfies [Hypothesis 10](#) on hyperbolicity and [Hypothesis 46](#) concerning the topological pressure. Let E_h be a generalized eigenfunction of the form described in [Hypothesis 43](#), where E_h^0 is associated to a Lagrangian manifold \mathcal{L}_0 which satisfies the invariance [Hypothesis 13](#) as well as the transversality [Hypothesis 16](#).

Then there exists a finite set of points $(\rho_b)_{b \in B_1} \subset K$ and a family $(\Pi_b)_{b \in B_1}$ of operators in $\Psi_h^{\text{comp}}(X)$ microsupported in a small neighbourhood of ρ_b such that $\sum_{b \in B_1} \Pi_b = I$ microlocally on a neighbourhood of K in T^*X such that the following holds.

Let $\mathcal{U}_b : L^2(X) \rightarrow L^2(\mathbb{R}^d)$ be a Fourier integral operator quantizing the symplectic change of local coordinates $\kappa_b : (x, \xi) \mapsto (y^{\rho_b}, \eta^{\rho_b})$, and which is microlocally unitary on the microsupport of Π_b .

For any $r > 0$, there exists $M_r > 0$ such that we have

$$\mathcal{U}_b \Pi_b E_h(y^{\rho_b}) = \sum_{n=0}^{\lfloor M_r \ell \rfloor \log h} \sum_{\beta \in \tilde{\mathcal{B}}_n} e^{i\phi_{n,\beta,b}(y^{\rho_b})/h} a_{n,\beta,b}(y^{\rho_b}; h) + R_r, \tag{51}$$

where $a_{n,\beta,b} \in S^{\text{comp}}(\mathbb{R}^d)$ are classical symbols, and each $\phi_{n,\beta,b}$ is a smooth function independent of h , and defined in a neighbourhood of the support of $a_{n,\beta,b}$. The set $\tilde{\mathcal{B}}_n$ will be defined in (85). Its cardinal behaves like some exponential of n .

We have the following estimate on the remainder

$$\|R_r\|_{L^2} = O(h^r).$$

For any $\ell \in \mathbb{N}$, $\epsilon > 0$, there exists $C_{\ell,\epsilon}$ such that for all $n \geq 0$ and for all $h \in (0, h_0]$, we have

$$\sum_{\beta \in \tilde{\mathcal{B}}_n} \|a_{n,\beta,b}\|_{C^\ell} \leq C_{\ell,\epsilon} e^{n(\mathcal{P}(1/2)+\epsilon)}. \tag{52}$$

Remark 48. This theorem can be considered as a quantum analogue of [Theorem 17](#). Indeed, as we explained in [Section 1](#), we will prove it by describing the evolution of the Schrödinger flow of Lagrangian states, while [Theorem 17](#) described the evolution by the Hamiltonian flow of associated Lagrangian manifolds. Actually, the sets containing the microsupports of the operators $(\Pi_b)_{b \in B_1}$ will be built from the sets $(W_a)_{a \in A_1}$ constructed in [Theorem 17](#), as explained in [Section 4.1](#).

Remark 49. The remainder R_r is compactly microlocalised, since the other two terms in the decomposition (51) are compactly microlocalised. Therefore, for any $\ell \in \mathbb{N}$, by possibly taking M_r larger, we may ask that

$$\|R_r\|_{C^\ell} = O(h^r).$$

[Theorem 47](#) may be used to identify the semiclassical measures associated to our generalized eigenfunctions, as in [Theorem 3](#). We shall do this only microlocally close to the trapped set, since the expression for the semiclassical measure on the whole manifold may become very complicated.

Let us denote by π_b the principal symbol of the operators Π_b introduced in the statement of [Theorem 47](#). The following corollary is a more precise version of (the second part of) [Theorem 3](#).

Corollary 50. *There exists a constant $0 < c \leq 1$ and functions $e_{n,\beta,b}$ for $n \in \mathbb{N}$, $\beta \in \tilde{\mathcal{B}}_n$ and $b \in B_1$ such that for any $a \in C_c^\infty(T^*X)$ and for any $\chi \in C_c^\infty(X)$, we have*

$$\langle \text{Op}_h(\pi_b^2 a) \chi E_h, \chi E_h \rangle = \int_{T^*X} a(x, v) d\mu_{b,\chi}(x, v) + O(h^c),$$

with

$$d\mu_{b,\chi}(\kappa_b^{-1}(y^{\rho_b}, \eta^{\rho_b})) = \sum_{n=0}^{\infty} \sum_{\beta \in \tilde{\mathcal{B}}_n} e_{n,\beta,b}(y^{\rho_b}) \delta_{\{\eta^{\rho_b} = \partial\phi_{j,n}(y^{\rho_b})\}} dy^{\rho_b}.$$

The functions $e_{n,\beta,b}$ satisfy an exponential decay estimate as in (52).

The functions $e_{n,\beta,b}$ will be closely related to $a_{n,\beta,b}^0(y^{\rho_b})$, the principal symbol of $a_{n,\beta,b}(y^{\rho_b})$ appearing in (51). Actually, $e_{n,\beta,b}(y^{\rho_b})$ will either be the square of the modulus of $a_{n,\beta,b}^0(y^{\rho_b})$, or the square of the modulus of the sum of a finite number of $a_{n,\beta,b}^0(y^{\rho_b})$ for different values of n and β . These different terms will come from the fact that a point may belong to $\Phi_{\beta}^{n,t_0}(\mathcal{L}_0)$ for several values of n, β .

3.6. Strategy of proof. To study the asymptotic behaviour of the distorted plane wave as h goes to zero, we would like to write that $\tilde{U}(t)E_h = E_h$, where $\tilde{U}(t) := e^{it/h}U(t)$. However, this equation can only be formal, because $E_h \notin L^2(X)$. Instead, we use [Dyatlov and Guillarmou 2014, Lemma 3.10]:

Lemma 51. *Let $\chi \in C_c^\infty(X)$. Take $t \in \mathbb{R}$, and a cut-off function $\chi_t \in C_c^\infty(X)$ supported in the interior of a compact set K_t , such that*

$$d_g(\text{supp } \chi, \text{supp}(1 - \chi_t)) > 2|t|,$$

where d_g denotes the Riemannian distance on M . Then, for any $\xi \in \mathbb{S}^d$, we have

$$\chi E_h = \chi \tilde{U}(t) \chi_t E_h + O(h^\infty \|E_h\|_{L^2(K_t)}). \tag{53}$$

Since E_h is a tempered distribution by assumption, we have, for any $t > 0$ and $\chi \in C_c^\infty(X)$,

$$\|\chi E_h - \chi \tilde{U}(t) \chi_t E_h\|_{L^2} = O(h^\infty),$$

where χ_t is as in Lemma 51.

We may then iterate this equation as follows: we write that $\chi_t = \chi + \chi_t(1 - \chi)$, and obtain

$$\chi E_h = \chi \tilde{U}(t)((1 - \chi)\chi_t) E_h + \chi \tilde{U}(t) \chi \tilde{U}(t) \chi_t E_h + O(h^\infty).$$

We may iterate this method to times $Nt \leq Mt|\log h|$ for any given $M > 0$. We obtain

$$\chi E_h = (\chi \tilde{U}(t))^N \chi_t E_h + \sum_{k=1}^N (\chi \tilde{U}(t))^k (1 - \chi)\chi_t E_h + O(h^\infty). \tag{54}$$

Now, choose $\chi \in C_c^\infty(X)$ as in Hypothesis 43, and take $t > T_\chi$.

Lemma 52. *Let $t > T_\chi$, $M > 0$, and $\chi \in C_c^\infty(X)$ be such that $\chi \equiv 1$ on $\{x \in X : b(b) > \epsilon_2\}$, where $\epsilon_2 < \epsilon_0$ is as in Hypothesis 43. For any $k \leq M|\log h|$, we have*

$$\|(\chi \tilde{U}(t))^k (1 - \chi)\chi_t E_h^1\|_{L^2} = O(h^\infty).$$

Proof. We only have to prove that $\|(\chi \tilde{U}(t))(1 - \chi)\chi_t E_h^1\|_{L^2} = O(h^\infty)$. This is a consequence of (45) in Hypothesis 43. □

Therefore, we have for any $\chi \in C_c^\infty(X)$ as in Lemma 52,

$$\chi E_h = (\chi \tilde{U}(t))^N \chi_t E_h^0 + (\chi \tilde{U}(t))^N \chi_t E_h^1 + \sum_{k=1}^N (\chi \tilde{U}(t))^k (1 - \chi) \chi_t E_h^0 + O(h^\infty). \tag{55}$$

Let us now introduce tools from [Nonnenmacher and Zworski 2009] to analyse these terms in more detail.

4. Tools for the proofs of Theorem 47

4.1. Another definition of topological pressure. Recall that \mathcal{E}_E and K_E were defined in (10) and (11) respectively. For any $\delta > 0$ small enough so that (12) holds, we define

$$\mathcal{E}^\delta := \bigcup_{|E-1| < \delta} \mathcal{E}_E, \quad K^\delta := \bigcup_{|E-1| < \delta} K_E.$$

Let $\mathcal{W} = (W_\alpha)_{\alpha \in A_1}$ be a finite open cover of $K^{\delta/2}$ such that the W_α are all strictly included in \mathcal{E}^δ and of diameter $< \varepsilon_0$, where ε_0 comes from Theorem 17. For any $T \in \mathbb{N}^*$, define $W(T) := (W_\alpha)_{\alpha \in A_1^T}$ by

$$W_\alpha := \bigcap_{k=0}^{T-1} \Phi^{-k}(W_{a_k}),$$

where $\alpha = a_0, \dots, a_{T-1}$. Let \mathcal{A}'_T be the set of $\alpha \in A_1^T$ such that $W_\alpha \cap K^\delta \neq \emptyset$. If $V \subset \mathcal{E}^\delta$, $V \cap K^{\delta/2} \neq \emptyset$, define

$$S_T(V) := - \inf_{\rho \in V \cap K^{\delta/2}} \lambda_T^+(\rho), \quad \text{with } \lambda_T^+ \text{ as in (49),}$$

$$Z_T(\mathcal{W}, s) := \inf \left\{ \sum_{\alpha \in \mathcal{A}_T} \exp\{s S_T(W_\alpha)\} : \mathcal{A}_T \subset \mathcal{A}'_T, K^{\delta/2} \subset \bigcup_{\alpha \in \mathcal{A}_T} W_\alpha \right\},$$

$$\mathcal{P}^\delta(s) := \lim_{\text{diam } \mathcal{W} \rightarrow 0} \lim_{T \rightarrow \infty} \frac{1}{T} \log Z_T(\mathcal{W}, s).$$

The topological pressure is then

$$\mathcal{P}(s) = \lim_{\delta \rightarrow 0} \mathcal{P}^\delta(s). \tag{56}$$

Recall that we assumed that

$$\mathcal{P}\left(\frac{1}{2}\right) < 0.$$

Let us fix $\varepsilon_0 > 0$ so that $\mathcal{P}\left(\frac{1}{2}\right) + 2\varepsilon_0 < 0$. Then there exists $t_0 > 0$, and $\widehat{\mathcal{W}}$ an open cover of K^δ with $\text{diam}(\widehat{\mathcal{W}}) < \varepsilon_0$ such that

$$\left| \frac{1}{t_0} \log Z_{t_0}(\widehat{\mathcal{W}}, s) - \mathcal{P}^\delta(s) \right| \leq \varepsilon_0. \tag{57}$$

We can find \mathcal{A}_{t_0} so that $\{W_\alpha : \alpha \in \mathcal{A}_{t_0}\}$ is an open cover of K^δ in \mathcal{E}^δ and such that

$$\sum_{\alpha \in \mathcal{A}_{t_0}} \exp\{s S_{t_0}(W_\alpha)\} \leq \exp\{t_0(\mathcal{P}^\delta(s) + \varepsilon_0)\}.$$

Therefore, if we take δ small enough, and if we rename $\{W_\alpha : \alpha \in \mathcal{A}_{t_0}\}$ as $\{V_b : b \in B_1\}$, we have

$$\sum_{b \in B_1} \exp\{\frac{1}{2}S_{t_0}(V_b)\} \leq \exp\{t_0(\mathcal{P}(\frac{1}{2}) + 2\epsilon_0)\}. \tag{58}$$

By taking t_0 large enough, we can assume that $\log(1 + \epsilon_0) + t_0(\mathcal{P}(1/2) + \epsilon_0) < 0$.

A new open cover of \mathcal{E} . By hypothesis, the diameter of $\widehat{\mathcal{W}}$ in (57) is smaller than ϵ_0 , so that we may apply Theorem 17 to it. We complete it into an open cover $(W_\alpha)_{\alpha \in A}$ as in Theorem 17, and if $\alpha \in A^N$ for some $N \geq 0$, we define as previously $W_\alpha := \bigcap_{k=0}^{N-1} \Phi^{-k}(W_{\alpha_k})$.

Let us rewrite as $(V_b)_{b \in B_2}$ the sets $(W_\alpha)_{\alpha \in A^{t_0}}$, where $\alpha \in A^{t_0} \setminus \mathcal{A}_{t_0}$ such that $\alpha_k \neq 0$ for some $0 \leq k \leq t_0 - 1$. We will also write V_0 for the set $W_{0,0,\dots,0}$.

If we write $B = B_1 \sqcup B_2 \sqcup \{0\}$, the sets $(V_b)_{b \in B}$ form an open cover of \mathcal{E} in T^*X .

Actually, by compactness of the interaction region, we may find a $\delta > \delta' > 0$ small enough so that (12) holds and such that, by replacing V_0 by $V_0 \cap \mathcal{E}^\delta$, the sets $(V_b)_{b \in B}$ form an open cover of $\mathcal{E}^{\delta'}$ included in \mathcal{E}^δ .

If $\beta = b_0 \cdots b_{N-1} \in B^N$ for some $N \in \mathbb{N}$, and if Λ is a Lagrangian manifold, we will define for each $0 \leq k \leq N - 1$ the set $\Phi_\beta^{k,t_0}(\Lambda)$ by

$$\begin{aligned} \Phi_\beta^{0,t_0}(\Lambda) &= \Lambda \cap V_{b_0}, \\ \Phi_\beta^{k,t_0}(\Lambda) &= \Phi^{t_0}(V_{b_k} \cap \Phi_\beta^{k-1,t_0}(\Lambda)) \quad \text{for } 1 \leq k \leq N - 1. \end{aligned}$$

By the definition of the sets $b \in B$, we have $\Phi_\beta^{N,t_0}(\Lambda) = \Phi_{\alpha_\beta}^{N,t_0}(\Lambda)$, where $\alpha_\beta \in A^{Nt_0}$ is the concatenation of all the sequences which make up the b_k , $0 \leq k \leq N - 1$.

Therefore, once we have fixed a point $\rho^b \in K \cap V_b$ for each $b \in B_1$, we have the following analogue of Theorem 17.

Corollary 53. *If there exists $N'_{\text{uns}} \in \mathbb{N}$ such that for all $N \in \mathbb{N}$, for all $\beta \in B^N$ and all $b \in B_1$, then $V_b \cap \Phi_\beta^{N,t_0}(\mathcal{L}_0)$ is either empty, or is a Lagrangian manifold in some unstable cone in the coordinates $(y^{\rho^b}, \eta^{\rho^b})$.*

Furthermore, if $N - \tau(\beta) \geq N'_{\text{uns}}$, then $V_b \cap \Phi_\beta^N(\mathcal{L}_0)$ is a γ_{uns} -unstable Lagrangian manifold in the coordinates $(y^{\rho^b}, \eta^{\rho^b})$.

Remark 54 (new definition of the sets $(V_b)_{b \in B_1}$). The sets $(V_b)_{b \in B_1}$ form an open cover of K . By compactness, they form an open cover of $\{\rho \in \mathcal{E} : d(\rho, K) \leq \epsilon_3\}$ for some $\epsilon_3 > 0$. Hence, if for each $b \in B_2$ we replace each V_b by $V_b \cap \{\rho \in \mathcal{E} : d(\rho, K) > \epsilon_3/2\}$, which we still denote by V_b , the sets $(V_b)_{b \in B}$ still form an open cover of \mathcal{E} , and the conclusions of Corollary 53 do still apply.

By adapting the proof of Lemma 24, we see that by possibly enlarging N'_{uns} , we may suppose that for all $b \in B_2$ and for all $\rho \in V_b$, we have $\Phi^{N'_{\text{uns}}t_0}(\rho) \in V_0 \setminus (\bigcup_{b \in B_2} V_b)$ or $\Phi^{-N'_{\text{uns}}t_0}(\rho) \in V_0 \setminus (\bigcup_{b \in B_2} V_b)$.

Note also that thanks to Lemma 9, for any $b \in B_1 \cup B_2$ and for any $k \geq 1$, we have $\Phi^{kt_0}(\mathcal{L}_0 \cap V_b) \cap \mathcal{W}_0 \cap \mathcal{D}\mathcal{E}_- = \emptyset$.

Remark 55. In [Nonnenmacher and Zworski 2009, Proposition 5.2] the authors proved the following statement. There exists a $\gamma_1 > 0$ such that the following holds. Let $b, b' \in B_1$, and let Λ be a Lagrangian

manifold contained in V_b , γ -unstable in the coordinates $(y^{\rho_b}, \eta^{\rho_b})$ for some $\gamma \leq \gamma_1$. Then $\Phi^{t_0}(\Lambda) \cap V_b$ is also a Lagrangian manifold which is γ -unstable in the coordinates $(y^{\rho_{b'}}, \eta^{\rho_{b'}})$.

Furthermore, the map $y^{\rho_b} \mapsto y^{\rho_{b'}}$ obtained by projecting $\Phi^{t_0}|_\Lambda$ onto the planes $\{(y^{\rho_b}, \eta^{\rho_b}) : \eta^{\rho_b} = 0\}$ and $\{(y^{\rho_{b'}}, \eta^{\rho_{b'}}) : \eta^{\rho_{b'}} = 0\}$ satisfies the following estimate on its domain of definition:

$$\det\left(\frac{\partial y^{\rho_{b'}}}{\partial y^{\rho_b}}\right) = (1 + O(\epsilon^p))e^{\lambda_{t_0}^+(\rho_b)},$$

where $\lambda_{t_0}^+(\rho_b)$ is the unstable Jacobian of ρ_b , defined in (49).

In the sequel, we will always suppose that $\gamma_{\text{uns}} < \gamma_1$.

For each $b \in B_1$, we will denote by \mathcal{U}_b a Fourier integral operator quantizing the local change of symplectic coordinates $(x, \xi) \mapsto (y^{\rho_b}, \eta^{\rho_b})$.

4.2. The Schrödinger propagator as a Fourier integral operator. Let us explain how the formalism of Section 3.1.3 may be used to describe the Schrödinger propagator $U(t)$ acting on $L^2(X)$. We shall state a lemma proven in [Nonnenmacher and Zworski 2009, Lemma 4.2]. Recall that for $0 < \delta < 1$, we defined \mathcal{E}^δ as $\bigcup_{|E-1|<\delta} \mathcal{E}$.

Lemma 56. *Let $V_0 \in \mathcal{E}^\delta$, $V_1 \subset \Phi^t(V_0)$ for some $t > 0$. Take some $\rho_0 \in V_0 \cap \mathcal{E}$ and set $\rho_1 = \Phi^t(\rho_0) \in V_1$. Let $f_j : \pi(V_j) \rightarrow \mathbb{R}^d$, $j = 0, 1$ be local coordinates such that $f_0(\pi(\rho_0)) = f_1(\pi(\rho_1)) = 0 \in \mathbb{R}^d$. They induce on V_0 and V_1 the symplectic coordinates*

$$F_j(x, \xi) := (f_j(x), (df_j(x)^t)^{-1}\xi - \xi^{(j)}), \quad j = 0, 1,$$

where $\xi^{(j)} \in \mathbb{R}^d$ is fixed by the condition $F_j(\rho_j) = (0, 0)$. Then the operator on $L^2(\mathbb{R}^d)$,

$$T(t) := e^{-i\langle x, \xi^{(1)} \rangle/h} (f_1^{-1})^* U(t) (f_0)^* e^{i\langle x, \xi^{(0)} \rangle/h},$$

is of the form (42) for some choice of the A_j microlocally near $(0, 0) \times (0, 0)$.

4.3. Iterations of Fourier integral operators. We recall here the main results from [Nonnenmacher and Zworski 2009, §4] concerning the iterations of semiclassical Fourier integral operators in $T^*\mathbb{R}^d$.

Let $V \subset T^*\mathbb{R}^d$ be an open neighbourhood of 0, and take a sequence of symplectomorphisms $(\kappa_i)_{i=1, \dots, N}$ from V to $T^*\mathbb{R}^d$ such that for all $i \in \{1, \dots, N\}$, we have $\kappa_i(0) \in V$, and the projection

$$(x_1, \xi_1; x_0, \xi_0) \mapsto (x_1, \xi_0), \quad \text{where } (x_1, \xi_1) = \kappa(x_0, \xi_0),$$

is a diffeomorphism close to the origin. We consider Fourier integral operators (T_i) which quantise κ_i and which are microlocally unitary near an open set $U \times U$, where $U \Subset V$, which contains the origin. Let $\Omega \subset \mathbb{R}^d$ be an open set such that $U \Subset T^*\Omega$, and, for all i , we have $\kappa_i(U) \Subset T^*\Omega$. For each i , we take a smooth cut-off function $\chi_i \in C_c^\infty(U; [0, 1])$, and let

$$S_i := \text{Op}_h(\chi_i) \circ T_i. \tag{59}$$

Let us consider a family of Lagrangian manifolds $\Lambda_k = \{(x, \phi'_k(x)) : x \in \Omega\} \subset T^*\mathbb{R}^d$, $k = 0, \dots, N$, such that

$$|\partial^\alpha \phi_k| \leq C_\alpha, \quad 0 \leq k \leq N, \quad \alpha \in \mathbb{N}^d. \tag{60}$$

We assume that there exists a sequence of integers $(i_k \in \{1, \dots, J\})_{k=1, \dots, N}$ such that

$$\kappa_{i_{k+1}}(\Lambda_k \cap U) \subset \Lambda_{k+1}, \quad k = 0, \dots, N - 1.$$

We define g_k by

$$g_k(x) = \pi \circ \kappa_{i_k}^{-1}(x, \phi'_k(x)).$$

That is to say, $\kappa_{i_k}^{-1}(x, \phi'_k(x)) = (g_k(x), \phi'_{k-1}(g_k(x)))$.

We will say that a point $x \in \Omega$ is N -admissible if we can define recursively a sequence by $x^N = x$, and, for $k = N, \dots, 1$, we have $x^{k-1} = g_k(x^k)$. This procedure is possible if, for any k , we have x^k is in the domain of definition of g_k .

Let us assume that, for any admissible sequence $(x^N \dots x^0)$, the Jacobian matrices are uniformly bounded from above:

$$\left\| \frac{\partial x^k}{\partial x^l} \right\| = \left\| \frac{\partial(g_{k+1} \circ g_{k+2} \circ \dots \circ g_l)}{\partial x^l}(x^l) \right\| \leq C_D, \quad 0 \leq k < l \leq N,$$

where C_D is independent of N . This assumption roughly says that the maps g_k are (weakly) contracting.

We will also use the notation

$$D_k := \sup_{x \in \Omega} |\det dg_k(x)|^{1/2}, \quad J_k := \prod_{k'=1}^k D_{k'},$$

and assume that the D_k are uniformly bounded: $1/C_D \leq D_k \leq C_D$.

The following result can be found in [Nonnenmacher and Zworski 2009, Proposition 4.1].

Proposition 57. *We use the above definitions and assumptions, and take N arbitrarily large, possibly varying with h . Take any $a \in S^{\text{comp}}$ and consider the Lagrangian state $u = ae^{i\phi_0/h}$ associated with the Lagrangian Λ_0 . Then we may write*

$$(S_{i_N} \circ \dots \circ S_{i_1})(ae^{i\phi_0/h})(x) = e^{i\phi_N(x)/h} \left(\sum_{j=0}^{L-1} h^j a_j^N(x) + h^L R_L^N(x, h) \right),$$

where each $a_j^N \in C_c^\infty(\Omega)$ depends on h only through N , and $R_L^N \in C^\infty((0, 1]_h, S(\mathbb{R}^d))$. If $x^N \in \Omega$ is N -admissible, and defines a sequence (x^k) , $k = N, \dots, 1$, then

$$|a_0^N(x^N)| = \left(\prod_{k=1}^N \chi_{i_k}(x^k, \phi'_k(x^k)) |\det dg_k(x^k)|^{1/2} \right) |a(x^0)|;$$

otherwise $a_j^N(x^N) = 0$, $j = 0, \dots, L-1$. We also have the bounds

$$\|a_j^N\|_{C^\ell(\Omega)} \leq C_{j,\ell} J_N (N+1)^{\ell+3j} \|a\|_{C^{\ell+2j}(\Omega)}, \quad j = 0, \dots, L-1, \quad \ell \in \mathbb{N}, \tag{61}$$

$$\|R_L^N\|_{L^2(\mathbb{R}^d)} \leq C_L \|a\|_{C^{2L+d}(\Omega)} (1 + C_0 h)^N \sum_{k=1}^N J_k k^{3L+d}, \tag{62}$$

$$\|R_L^N\|_{C^\ell(\mathbb{R}^d)} \leq C_{L,\ell} h^{-d/2-\ell} \|a\|_{C^{2L+d}(\Omega)} (1 + C_0 h)^N \sum_{k=1}^N J_k k^{3L+d}. \tag{63}$$

The constants $C_{j,\ell}$, C_0 and C_L depend on the constants in (60) and on the operators $\{S_j\}_{j=1}^J$.

We shall mainly be using this proposition in the case where for all k , we have $D_k \leq \nu < 1$. In this case, the estimates (61), (62) and (63) imply that for any $\ell \in \mathbb{N}$, there exists C_ℓ independent of N such that for any $N \in \mathbb{N}$, we have

$$\|a^N\|_{C^\ell} \leq \|a_0^N\|_{C^\ell} (1 + C_\ell h). \tag{64}$$

4.4. Microlocal partition. We take a partition of unity $\sum_{b \in B} \pi_b$ such that

$$\sum_{b \in B} \pi_b(x) \equiv 1 \quad \text{for all } x \in \mathcal{E}^{\delta'},$$

and $\text{supp}(\pi_b) \subset V_b \subset \mathcal{E}^\delta$ for all $b \in B$.

For $b \in B_1 \cup B_2$, we set $\Pi_b := \text{Op}_h(\pi_b)$. We have

$$\text{WF}_h(\Pi_b) \subset V_b \cap \mathcal{E}^\delta \quad \text{and} \quad \Pi_b = \Pi_b^*.$$

We then set

$$\Pi_0 := \text{Id} - \sum_{b \in B_1 \cup B_2} \Pi_b.$$

We can decompose the propagator at time t_0 into

$$\tilde{U}(t_0) = \sum_{b \in B} \tilde{U}_b, \quad \text{where } \tilde{U}_b := \Pi_b e^{it_0/h} U(t_0).$$

The propagator at time Nt_0 may then be decomposed as

$$\tilde{U}(Nt_0) = \sum_{\beta \in B^N} \tilde{U}_\beta, \tag{65}$$

where $\tilde{U}_\beta := \tilde{U}_{\beta_{N-1}} \circ \dots \circ \tilde{U}_{\beta_0}$.

4.5. Hyperbolic dispersion estimates. We will use the following hyperbolic dispersion estimate, coming from [Nonnenmacher and Zworski 2009, Proposition 6.3], the proof of which can be found in Section 7 of that paper.

Lemma 58 (hyperbolic dispersion estimate). *Let $M > 0$ be fixed. There exists an $h_0 > 0$ and a $C > 0$ such that for any $0 < h < h_0$, for any $N < M \log(1/h)$ and for any $\beta \in B_1^N$, we have*

$$\|\tilde{U}_\beta\|_{L^2 \rightarrow L^2} \leq C h^{-d/2} (1 + \epsilon_0)^N \prod_{j=1}^N \exp\left[\frac{1}{2} S_{t_0}(V_{\beta_j})\right]. \tag{66}$$

5. Proof of Theorem 47

Proof. Having introduced these different tools, we may now come back to the proof of [Theorem 47](#).

5.1. Decomposition of χE_h . Let $\chi \in C_c^\infty(X)$ be as in [Lemma 52](#). We may suppose $T_\chi \leq t_0$. Then, by equation (55), we have

$$\chi E_h = (\chi \tilde{U}(t_0))^N \chi_{t_0} E_h + \sum_{k=1}^N (\chi \tilde{U}(t_0))^k (1 - \chi) \chi_{t_0} E_h^0 + O(h^\infty), \tag{67}$$

where the cut-off function $\chi_{t_0} \in C_c^\infty(X)$ is such that

$$d_X(\text{supp } \chi, \text{supp}(1 - \chi_{t_0})) > 2|t_0|,$$

where d_X denotes the Riemannian distance on X .

We shall require the following lemma. The proof of (i) is the same as that of [Lemma 24](#), while the proof of (ii) essentially follows from point (3) of [Hypothesis 5](#).

Lemma 59. (i) *There exists $N_\chi \in \mathbb{N}$ such that for any $N \in \mathbb{N}$ if $\rho \in \text{supp}(\chi_{t_0})$ and $\Phi^N(\rho) \in \text{supp}(\chi)$, then for any $N_\chi \leq k \leq N - N_\chi$, we have $\Phi^{k t_0}(\rho) \in V_b$ for some $b \in B_1 \cup B_2$.*

(ii) *If $\rho \in \mathcal{E}$ is such that $\Phi^{k t_0}(\rho) \in V_0$ for some $k \in \mathbb{N}$, but $\Phi^{(k+1)t_0}(\rho) \in V_b$ for some $b \in B_1 \cup B_2$, then $\Phi^{k'}(\rho)$ is in \mathcal{DE}_- (and hence in V_0) for any $k' \leq k$.*

From [Lemma 59](#), we deduce that for any $k \geq 2N_\chi + 2$, we have

$$(\chi \tilde{U}(t_0))^k = \sum_{l=0}^{N_\chi+1} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in B^{k-2N_\chi-2+l}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^{N_\chi-l} + O_{L^2 \rightarrow L^2}(h^\infty). \tag{68}$$

For any $N \in \mathbb{N} \setminus \{0\}$, define $\mathcal{B}_N \subset (B_1 \cup B_2)^N$ by

$$\mathcal{B}_N := \begin{cases} (B_1 \cup B_2)^N & \text{if } N \leq 2N'_{\text{uns}} + 2, \\ (B_1 \cup B_2)^{N'_{\text{uns}}+1} B_1^{N-2N'_{\text{uns}}-2} (B_1 \cup B_2)^{N'_{\text{uns}}+1} & \text{otherwise.} \end{cases} \tag{69}$$

Lemma 60. *For any $N \geq 2N'_{\text{uns}} + 2$ and for any $\beta \in (B_1 \cup B_2)^N \setminus \mathcal{B}_N$, we have*

$$\|\tilde{U}_\beta\|_{L^2 \rightarrow L^2} = O(h^\infty).$$

Proof. Let $\beta \in (B_1 \cup B_2)^N \setminus \mathcal{B}_N$. Then there exists $N'_{\text{uns}} + 2 \leq k \leq N - N'_{\text{uns}} + 2$ such that $\beta_k \in B_2$. Recall from [Remark 54](#) that N'_{uns} is such that for any $\rho \in V_{\beta_k}$, we have $\Phi^{N'_{\text{uns}} t_0}(\rho) \in V_0 \setminus (\bigcup_{b \in B_2} V_b)$ or $\Phi^{-N'_{\text{uns}} t_0}(\rho) \in V_0 \setminus (\bigcup_{b \in B_2} V_b)$. The result then follows from [Lemma 40](#). \square

Equation (68) may then be rewritten as

$$(\chi \tilde{U}(t_0))^k = \sum_{l=0}^{N_\chi+1} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in B^{k-2N_\chi-2+l}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^{N_\chi-l} + O_{L^2 \rightarrow L^2}(h^\infty). \tag{70}$$

By summing over k and reordering the terms, we get, for any $K > 2N_\chi + 3N'_{\text{uns}} + 4$,

$$\begin{aligned} \sum_{k=0}^K (\chi \tilde{U}(t_0))^k &= \sum_{n=1}^{\mathfrak{N}} \sum_{l=0}^{N_\chi+1} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+R N'_{\text{uns}}+2}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^l \\ &\quad - \sum_{n=K-2N_\chi-2}^{\mathfrak{N}} \sum_{l=0}^{\mathfrak{N}-n} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^l \\ &\quad + \sum_{l=0}^{K-\mathfrak{N}-1} (\chi \tilde{U}(t_0))^l + O_{L^2 \rightarrow L^2}(h^\infty), \end{aligned} \tag{71}$$

where $\mathfrak{N} = K - 3N'_{\text{uns}} - N_\chi - 4$.

Let us note that from [Lemma 42](#) and [Hypothesis 13](#), for each $0 \leq l \leq N_\chi$, there exists $\chi_l \in S^{\text{comp}}(X)$ such that

$$(\chi \tilde{U}_0)^{N_\chi-l} (1 - \chi) \chi_{t_0} E_h^0 = \chi_l E_h^0 + O(h^\infty). \tag{72}$$

Let us introduce the notation

$$\bar{\chi} := \sum_{l=0}^{N_\chi+1} \chi_l. \tag{73}$$

Thanks to equation (71), we can study the different terms in equation (67). The first term in the right-hand side of (67) may be bounded by the following lemma.

Lemma 61. *Let $r > 0$. We may find a constant $M_r \geq 0$ such that for any $M > M_r$ and for any $M_r |\log h| \leq N \leq M |\log h|$, we have*

$$\|(\chi \tilde{U}(t_0))^N \chi_{t_0} E_h\|_{L^2} = O(h^r).$$

Proof. We use (70), [Lemma 58](#) and the topological pressure assumption to obtain

$$\begin{aligned} \|(\chi \tilde{U}(t_0))^N \chi_{t_0} E_h\|_{L^2} &\leq C \left\| \sum_{\beta \in \mathcal{B}_{N-2N_\chi-2}} \tilde{U}_\beta \chi_{t_0} E_h \right\| + O(h^\infty) \\ &\leq C \sum_{\beta \in \mathcal{B}_1^{N-2N_{\text{uns}}-2N_\chi-4}} \|\tilde{U}_\beta \chi_{t_0} E_h\| \\ &\leq C h^{-d/2} (1 + \epsilon_0)^N \sum_{\beta \in \mathcal{B}_1^{N-2N_{\text{uns}}-2N_\chi-4}} \prod_{j=1}^{N-2N_{\text{uns}}-2} \exp\left[\frac{1}{2} S_{t_0}(V_{\beta_j})\right] \|\chi_{t_0} E_h\| \\ &\leq C h^{-d/2} (1 + \epsilon_0)^N \left(\sum_{b \in \mathcal{B}_1} \exp\left[\frac{1}{2} S_{t_0}(V_b)\right] \right)^N \|\chi_{t_0} E_h\| \\ &\leq C h^{-d/2} (1 + \epsilon_0)^N \exp\{N t_0 (\mathcal{P}(\frac{1}{2}) + 2N\epsilon_0)\} \|\chi_{t_0} E_h\|. \end{aligned}$$

By assumption, E_h is a tempered distribution, so that $\|\chi_{t_0} E_h\|_{L^2} \leq C/h''$. Therefore

$$\|(\chi \tilde{U}(t_0))^N \chi_{t_0} E_h\|_{L^2} \leq C h^{-r''-d/2-\epsilon} \exp\{N t_0 (\mathcal{P}(\frac{1}{2}) + 2N\epsilon_0)\}$$

for some small ϵ . The lemma follows by taking M_r large enough. □

Using [Lemma 61](#), and equation (71), we may rewrite equation (67) as

$$\begin{aligned} \chi E_h &= \sum_{n=1}^{M_r |\log h|} \sum_{l=0}^{N_\chi+1} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^l (1-\chi) \chi_{t_0} E_h^0 \\ &\quad - \sum_{n=M_r |\log h| - N_\chi}^{M_r |\log h|} \sum_{l=0}^{M_r |\log h| - N_\chi - 2 - n} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}} \tilde{U}_\beta \right) (\chi \tilde{U}_0)^l (1-\chi) \chi_{t_0} E_h^0 \\ &\quad + \sum_{l=0}^{3N'_{\text{uns}}+N_\chi+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 + O_{L^2}(h^r). \end{aligned}$$

The second term may be bounded by $O(h^r)$ thanks to [Lemma 61](#). By using equations (72) and (73), we get

$$\chi E_h = \sum_{n=1}^{M_r |\log h|} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}} \tilde{U}_\beta \right) \bar{\chi} E_h^0 + \sum_{l=0}^{3N'_{\text{uns}}+N_\chi+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 + O_{L^2}(h^r). \tag{74}$$

5.2. Evolution of the WKB states.

5.2.1. Construction of $\tilde{\mathcal{B}}_0$. From now on, we fix $b \in B_1$ and $r > 1$. We may write

$$\mathcal{U}_b \Pi_b \sum_{l=0}^{3N'_{\text{uns}}+N_\chi+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 = \sum_{l=0}^{N_\chi+3N'_{\text{uns}}+3} \sum_{\beta \in B^l} \mathcal{U}_b \Pi_b U_\beta^\chi (1-\chi) \chi_{t_0} E_h^0, \tag{75}$$

where we have used the notation

$$U_\beta^\chi = \chi \tilde{U}_{\beta_l} \chi \cdots \chi \tilde{U}_{\beta_0}. \tag{76}$$

Note that each of the $\mathcal{U}_b \Pi_b U_\beta^\chi$ is a Fourier integral operator from $L^2(X)$ to $L^2(\mathbb{R}^d)$. Thanks to [Corollary 53](#), we may use [Lemma 42](#) to describe the action of each of these Fourier integral operators on the Lagrangian state $(1-\chi) \chi_{t_0} E_h^0$. If we denote by $\tilde{\mathcal{B}}_0$ the set $\bigcup_{l=0}^{N_\chi+3N'_{\text{uns}}+3} B^l$, we may write

$$\mathcal{U}_b \Pi_b \sum_{l=0}^{N_\chi+3N'_{\text{uns}}+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 = \sum_{\beta \in \tilde{\mathcal{B}}_0} e_{0,\beta,b}, \tag{77}$$

where $e_{0,\beta,b}(y^b) = e^{\phi_{0,\beta,b}(y^{\rho b})/h} a_{0,\beta,b}(y^{\rho b}; h)$, with $a_{0,\beta,b}$ and $\phi_{0,\beta,b}$ as in the statement of [Theorem 47](#).

Let us now consider the other terms on the right-hand side of equation (74), which will be indexed by $\tilde{\mathcal{B}}_n$, $n \geq 1$.

5.2.2. Evolution in the intermediate region. Let $n \geq 1$ and $\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}$. By the definition of $\mathcal{B}_{n+3N'_{\text{uns}}+2}$, for $N'_{\text{uns}} + 1 \leq i \leq n + 2N'_{\text{uns}} + 1$, we have $\beta_i \in B_1$.

According to [Theorem 17](#), $\Phi_{\beta}^{2N'_{\text{uns}}+1, t_0}(\mathcal{L}_0)$ consists of a single Lagrangian manifold, which is γ_{uns} -unstable in the symplectic coordinates in $V_{\beta_{2N'_{\text{uns}}+1}}$.

Thus, we may say that $\tilde{\mathcal{U}}_{\beta_0 \dots \beta_{2N'_{\text{uns}}+1}}(\bar{\chi} E_h^0)$ is a Lagrangian state associated to the Lagrangian manifold $\Phi_{\beta}^{2N'_{\text{uns}}+1, t_0}(\mathcal{L}_0)$. Thanks to [Lemma 56](#), we may use [Lemma 42](#) to write

$$(\mathcal{U}_{\beta_{2N'_{\text{uns}}+1}} \Pi_{\beta_{2N'_{\text{uns}}+1}} \tilde{\mathcal{U}}_{\beta_0 \dots \beta_{2N'_{\text{uns}}+1}}(\bar{\chi} E_h^0))(y^{\rho_{\beta_{2N'_{\text{uns}}+1}}}) = a(y^{\rho_{\beta_{2N'_{\text{uns}}+1}}}; h) e^{i\phi(y^{\rho_{\beta_{2N'_{\text{uns}}+1}}})/h}$$

for some $a \in S^{\text{comp}}(\mathbb{R}^d)$.

5.2.3. Propagation of Lagrangian states close to the trapped set. To lighten the notations, let us write $\hat{n} := n + 2N'_{\text{uns}} + 1$.

For each $2N'_{\text{uns}} + 1 \leq k \leq \hat{n}$, we write

$$T_{\beta_{k'+1}, \beta_{k'}} := \mathcal{U}_{\beta_{k'+1}} \tilde{\mathcal{U}}_{\beta_{k'+1}} \mathcal{U}_{\beta_{k'}}^*.$$

Now $T_{\beta_{k'+1}, \beta_{k'}}$ is an operator quantising the map $\kappa_{\beta_{k'+1}, \beta_{k'}}$ obtained by expressing Φ^{t_0} in the coordinates $(y^{\beta_{k'}}, \eta^{\beta_{k'}}) \mapsto (y^{\beta_{k'+1}}, \eta^{\beta_{k'+1}})$. It is of the form [\(59\)](#).

We will write

$$T_{\beta}^{2N'_{\text{uns}}+1, \hat{n}} := T_{\beta_{\hat{n}}, \beta_{\hat{n}}} \circ \dots \circ T_{\beta_{2N'_{\text{uns}}+2}, \beta_{2N'_{\text{uns}}+1}}.$$

Thanks to [Remark 55](#), we may apply [Proposition 57](#) to describe the action of $T_{\beta}^{2N'_{\text{uns}}+1, \hat{n}}$ on the Lagrangian state $\mathcal{U}_{\beta_{2N'_{\text{uns}}+1}} \tilde{\mathcal{U}}_{\beta_0 \dots \beta_{2N'_{\text{uns}}+1}}(\bar{\chi} E_h^0)$. Note that

$$T_{\beta}^{2N'_{\text{uns}}+1, \hat{n}} \mathcal{U}_{\beta_{2N'_{\text{uns}}+1}} \tilde{\mathcal{U}}_{\beta_0 \dots \beta_{2N'_{\text{uns}}+1}} = \mathcal{U}_{\beta_{\hat{n}}} \tilde{\mathcal{U}}_{\beta_0 \dots \beta_{\hat{n}}}.$$

We obtain that $\mathcal{U}_{\beta_{\hat{n}+1}} \Pi_{\beta_{\hat{n}+1}} \tilde{\mathcal{U}}_{\beta_0 \dots \beta_{\hat{n}}}(\bar{\chi} E_h^0) = e_{\hat{n}, \beta}$, with

$$e_{\hat{n}, \beta}(y) = a^{\hat{n}, \beta}(y) e^{i\phi_{\hat{n}, \beta}(y)/h}, \quad y \in \mathbb{R}^d. \tag{78}$$

In the notation of [Section 4.3](#), by [Remark 55](#) that for any $N'_{\text{uns}} + 1 \leq k' \leq \hat{n}$, we have

$$D_{k'} = S_T(V_{\beta_{k'}})(1 + O(\epsilon^p)) < 1.$$

We therefore set

$$J_{\beta_{N'_{\text{uns}}+1} \dots \beta_{\hat{n}}} := \prod_{k'=N'_{\text{uns}}+1}^{\hat{n}} (S_{t_0}(V_{\beta_{k'}})(1 + O(\epsilon^p))). \tag{79}$$

Thanks to equation [\(61\)](#) in [Proposition 57](#) and equation [\(64\)](#), we obtain for any $\ell \in \mathbb{N}$,

$$\|a^{\hat{n}, \beta}\|_{C^{\ell}} \leq (1 + C_{\ell} h) C'_{\ell} J_{\beta_{N'_{\text{uns}}+1} \dots \beta_{\hat{n}}} (\hat{n} + 1)^{\ell} \tag{80}$$

for some constants C_{ℓ}, C'_{ℓ} .

5.2.4. End of the propagation. Using equation (74) and the results of the previous subsection, we have

$$\begin{aligned} \chi E_h &= \sum_{n=1}^{M_r |\log h|} (\chi \tilde{U}(t_0))^{N_\chi+1} \left(\sum_{\beta \in \mathcal{B}_{n+3N'_{\text{uns}}+2}} \tilde{U}_{\beta_{\hat{n}} \dots \beta_n} \mathcal{U}_{\beta_{\hat{n}}}^* e_{\hat{n},\beta} \right) \\ &+ \sum_{l=0}^{N_\chi+3N'_{\text{uns}}+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 + O_{L^2}(h^r), \end{aligned} \tag{81}$$

with

$$\mathcal{U}_b \Pi_b \sum_{l=0}^{N_\chi+3N'_{\text{uns}}+3} (\chi \tilde{U}(t_0))^l (1-\chi) \chi_{t_0} E_h^0 = \sum_{\beta \in \tilde{\mathcal{B}}_0} e_{0,\beta,b}.$$

To finish the proof, we have to apply $\mathcal{U}_b \Pi_b (\chi \tilde{U}(t_0))^{N_\chi+1} \tilde{U}_{\beta_{\hat{n}} \dots \beta_n} \mathcal{U}_{\beta_{\hat{n}}}^*$ to $e_{\hat{n},\beta}$.

To do this, one should once again decompose the propagator, and study

$$\sum_{\beta' \in \mathcal{B}^{N_\chi+1}} \mathcal{U}_b \Pi_b U_{\beta'}^\chi \tilde{U}_{\beta_{\hat{n}} \dots \beta_n} \mathcal{U}_{\beta_{\hat{n}}}^* e_{\hat{n},\beta}, \tag{82}$$

with $U_{\beta'}^\chi$ as in (76). To analyse each of the terms on the right-hand side of (82), we use once again Lemma 42 (the lemma may be applied, thanks to Theorem 17 and to Lemma 56).

We obtain that

$$\mathcal{U}_b \Pi_b U_{\beta'}^\chi \tilde{U}_{\beta_{\hat{n}} \dots \beta_n} \mathcal{U}_{\beta_{\hat{n}}}^* e_{\hat{n},\beta}(y) = a^{n,\beta,\beta'}(y) e^{i\phi_{n,\beta,\beta'}(y)/h}, \quad y \in \mathbb{R}^d, \tag{83}$$

and thanks to equation (80), we get

$$\|a^{n,\beta,\beta'}\|_{C^\ell} \leq (1 + C_\ell h) C'_\ell J_{\beta_{N'_{\text{uns}}+1} \dots \beta_{\hat{n}}} (\hat{n} + 1)^\ell \tag{84}$$

for some constants C_ℓ, C'_ℓ .

For any $n \geq 1$, we write

$$\tilde{\mathcal{B}}_n = \mathcal{B}_{n+3N'_{\text{uns}}+2} \times \mathcal{B}^{N_\chi+1}. \tag{85}$$

As announced, the cardinal of $\tilde{\mathcal{B}}_n$ grows exponentially with n . If $\beta = (\beta', \beta'') \in \tilde{\mathcal{B}}_n$ with $\beta \in \mathcal{B}_{n+2N'_{\text{uns}}+1}$, we define

$$a_{n,\beta,b} = a^{N_n+2N_\chi+2,l,\beta,\beta'}, \quad \phi_{n,\beta,b} = \phi_{N_n+2N_\chi+2,l,\beta,\beta'}.$$

With these notations, combining (81) with (83) gives us the decomposition (51).

The key point to obtaining estimate (52) is to notice that for any $N \geq N'_{\text{uns}} + 1$, we have, thanks to (58),

$$\begin{aligned} \sum_{\beta_{N'_{\text{uns}}+1} \dots \beta_{\hat{N}} \in \mathcal{B}_1^{N-N'_{\text{uns}}-1}} J_{\beta_{N'_{\text{uns}}+1} \dots \beta_{\hat{N}}} &= \left(\sum_{b \in \mathcal{B}_1} S_{t_0}(V_b) (1 + O(\epsilon^p)) \right)^{N-N'_{\text{uns}}-1} \\ &\leq \exp\left[(N - N'_{\text{uns}} - 1) (t_0 \mathcal{P}(\frac{1}{2}) (1 + O(\epsilon^p))) \right]. \end{aligned} \tag{86}$$

By applying (86) for $N = N_{n+2N_\chi+2,l}$, and combining it with (84), we get (52). □

Note that, although the statement of [Theorem 47](#) describes the generalized eigenfunctions E_h only very close to the trapped set, [\(81\)](#) can be used to describe E_h in any compact set, though in a less explicit way.

Using the estimate [\(52\)](#) as well as the fact that $\|\chi\tilde{U}(t_0)\|_{L^2 \rightarrow L^2} \leq 1$ and $\|\mathcal{U}_b\|_{L^2 \rightarrow L^2} \leq 1$, we deduce [Theorem 1](#).

6. Semiclassical measures

The main ingredient in the proof of [Corollary 50](#) is nonstationary phase. Let us recall the estimate we will use, and which can be proven by integrating by parts.

Let $a, \phi \in S^{\text{comp}}(X)$. We consider the oscillatory integral

$$I_h(a, \phi) := \int_X a(x) e^{\frac{i\phi(x,h)}{h}} dx.$$

Proposition 62. *Let $\epsilon > 0$. Suppose that there exists $C > 0$ such that for all $x \in \text{spt}(a)$ and for all $0 < h < h_0$, we have $|\partial\phi(x, h)| \geq Ch^{1/2-\epsilon}$. Then*

$$I_h(a, \phi) = O(h^\infty).$$

We shall only give a sketch of proof here, and refer to [\[Hörmander 1983, §7.7\]](#) for more details.

Sketch of proof. To prove this result, we simply integrate by parts, noting that

$$I_h(a, \phi) = \frac{h}{i} \int_X \frac{a}{|\partial\phi|^2} \partial\phi \cdot \partial(e^{\frac{i\phi(x,h)}{h}}) dx.$$

Hence, when we integrate by parts, the worst term in the integrand will involve second derivatives of ϕ times $h/|\partial\phi|^2$, and will therefore be a $O(h^{2\epsilon})$ by assumption. By integrating by parts more times, we will gain a factor $h^{2\epsilon}$ every time, so that $I_h(a, \phi)$ is actually a $O(h^\infty)$. □

Note that the sketch of proof above tells us that, if we could say that when $\partial\phi(x, h)$ is small, then the higher derivatives of ϕ are small as well, i.e., if we had

$$\forall k \geq 2, \exists C_k \quad \text{such that} \quad |\partial^k \phi(x, h)| \leq C_k |\partial\phi(x, h)|,$$

then we would have $I_h(a, \phi) = O(h^\infty)$ provided $|\partial\phi(x, h)| \geq Ch^{1-\epsilon}$. However, it is not clear that we can estimate the higher derivatives of the phase functions which appear in this section.

6.1. Distance between the Lagrangian manifolds. To take advantage of [Proposition 62](#), we need a lower bound on the distance between the Lagrangian manifolds which make up $\Phi^{n,t_0}(\mathcal{L}_0) \cap V_b$. To prove such a lower bound, let us first state an elementary topological lemma.

Lemma 63. *There exists $c_0 > 0$ such that for any $\rho, \rho' \in T^*X_0 \cap \mathcal{E}$ such that $d(\rho, \rho') < c_0$, there exists $b \in B$ such that $\rho, \rho' \in V_b$.*

Proof. Suppose for contradiction that for any $\epsilon > 0$, there exists $\rho_\epsilon, \rho'_\epsilon$ such that $d(\rho_\epsilon, \rho'_\epsilon) < \epsilon$ and such that for all $b \in B$ such that $\rho_\epsilon \in V_b$, we have $\rho'_\epsilon \notin V_b$. By compactness of $T^*X_0 \cap \mathcal{E}$, we may suppose that ρ_ϵ converges to some ρ . We then have $\rho'_\epsilon \rightarrow x$, and if $b \in B$ is such that $\rho \in V_b$, then $\rho_\epsilon, \rho'_\epsilon \in V_b$ for ϵ small enough, a contradiction. □

We may now state our lower bound on the distance between the Lagrangian leaves which make up $\Phi^{n,t_0}(\mathcal{L}_0) \cap V_b$.

Let $N \in \mathbb{N}$, $\beta \in B^N$ and $b \in B_1$. The set $\Phi_\beta^{n,t_0}(\mathcal{L}_0) \cap V_b$ may be written in the form $\{(y^{\rho_b}, \partial\tilde{\phi}_{n,\beta,b}(y^{\rho_b}))\}$ for some smooth function $\tilde{\phi}_{n,\beta,b}$.

For any $\beta \in B^N$, $\beta' \in B^{N'}$, let us define

$$\sigma(\beta, \beta') := \max(N - \tau(\beta), N' - \tau(\beta')),$$

with $\tau(\beta)$ as in (17).

Proposition 64. *There exist constants $C'_1, C'_2 > 0$ such that for any $N, N' \in \mathbb{N}$, for any $\beta \in B^N, \beta' \in B^{N'}$, for any $b \in B_1$ and for any y^{ρ_b} , we have either $\partial\tilde{\phi}_{N,\beta,b}(y^{\rho_b}) = \partial\tilde{\phi}_{N',\beta',b}(y^{\rho_b})$ or*

$$|\partial\tilde{\phi}_{N,\beta,b}(y^{\rho_b}) - \partial\tilde{\phi}_{N',\beta',b}(y^{\rho_b})| \geq C'_1 e^{C'_2 \sigma(\beta, \beta')}.$$

Proof. Since $T^*X_0 \cap \mathcal{E}$ is compact, we may find a constant $C > 0$ such that for any $\rho, \rho' \in \mathcal{E} \cap T^*X_0$,

$$d(\Phi^t(\rho), \Phi^t(\rho')) \leq e^{Ct} d(\rho, \rho'), \quad (87)$$

where d is the distance on the energy layer which we introduced in Section 2.1.2.

Let $b \in B_1$, and $y^{\rho_b} \in D_{\beta,b} \cap D_{\beta',b}$ be such that

$$\partial\tilde{\phi}_{N,\beta,b}(y^{\rho_b}) \neq \partial\tilde{\phi}_{N',\beta',b}(y^{\rho_b}).$$

Let us denote by ρ the point $(y^{\rho_b}, \partial\tilde{\phi}_{N,\beta,b}(y^{\rho_b}))$ and by ρ' the point $(y^{\rho_b}, \partial\tilde{\phi}_{N',\beta',b}(y^{\rho_b}))$.

We claim that there exists $0 \leq k \leq \sigma(\beta, \beta')$ such that for each $b' \in B$, if $\Phi^{-kt_0}(\rho) \in V_{b'}$, then $\Phi^{-kt_0}(\rho') \notin V_b$. Indeed, if no such k existed, then for each k , there would exist $b_k \in B$ such that $\Phi^{-kt_0}(\rho) \in V_{b_k}$ and $\Phi^{-kt_0}(\rho') \in V_{b_k}$ for each $0 \leq k \leq \sigma(\beta, \beta')$. We would then have $\rho \in \Phi_{\beta''}^{\max(N, N'), t_0}(\mathcal{L}_0)$ and $\rho' \in \Phi_{\beta''}^{\max(N, N'), t_0}(\mathcal{L}_0)$ for some sequence β'' built by possibly adding some 0's in front of the sequences β and β' . This would contradict the statement of Corollary 53.

Thanks to Lemma 63, we deduce from this that there exists $0 \leq k \leq \sigma(\beta, \beta')$ such that

$$d(\Phi^{-kt_0}(\rho), \Phi^{-kt_0}(\rho')) \geq c_0.$$

Combining this fact with equation (87), we get

$$d(\rho, \rho') \geq c_0 e^{-C\sigma(\beta, \beta')}$$

Since all metrics are equivalent on a compact set, we may compare $d(\rho, \rho')$ with $|\partial\tilde{\phi}_{N,\beta,b}(y^{\rho_b}) - \partial\tilde{\phi}_{N',\beta',b}(y^{\rho_b})|$ and we deduce from this the proposition. \square

Using the definition of $\tilde{\mathcal{B}}_n$, we deduce the following result about the functions $\phi_{n,\beta,b}$ in the statement of Theorem 47.

Corollary 65. *There exist constants $C'_1, C'_2 > 0$ such that for any $n, n' \in \mathbb{N}$, for any $\beta \in \tilde{\mathcal{B}}_n, \beta' \in \tilde{\mathcal{B}}_{n'}$, for any $b \in B_1$ and for any y^{ρ_b} , we have either $\partial\phi_{n,\beta,b}(y^{\rho_b}) = \partial\phi_{n',\beta',b}(y^{\rho_b})$ or*

$$|\partial\phi_{n,\beta,b}(y^{\rho_b}) - \partial\phi_{n',\beta',b}(y^{\rho_b})| \geq C'_1 e^{C'_2 \min(n, n')}.$$

6.2. Proof of Corollary 50. We shall now prove Corollary 50, which we recall.

Corollary 50. *There exists a constant $0 < c \leq 1$ and functions $e_{n,\beta,b}$ for $n \in \mathbb{N}$, $\beta \in \tilde{\mathcal{B}}_n$ and $b \in B_1$ such that for any $a \in C_c^\infty(T^*X)$ and for any $\chi \in C_c^\infty(X)$, we have*

$$\langle \text{Op}_h(\pi_b^2 a) \chi E_h, \chi E_h \rangle = \int_{T^*X} a(x, v) \, d\mu_{b,\chi}(x, v) + O(h^c),$$

with

$$d\mu_{b,\chi}(\kappa_b^{-1}(y^{\rho_b}, \eta^{\rho_b})) = \sum_{n=0}^{\infty} \sum_{\beta \in \tilde{\mathcal{B}}_n} e_{n,\beta,b}(y^{\rho_b}) \delta_{\{\eta^{\rho_b} = \partial \phi_{j,n}(y^{\rho_b})\}} dy^{\rho_b},$$

The functions $e_{n,\beta,b}$ satisfy the estimate (52).

Proof. Take any small $\epsilon > 0$, and set

$$M := \frac{1}{2C'_2} - \epsilon, \quad c := (M - \epsilon)\mathcal{P}\left(\frac{1}{2}\right) = \frac{\mathcal{P}\left(\frac{1}{2}\right)}{2C'_2} - \epsilon,$$

where C'_2 comes from Corollary 65.

Let $a \in C_c^\infty(T^*X)$, $\chi \in C_c^\infty(X)$ and $b \in B_1$. Using the fact that $\text{Op}_h(ab) = \text{Op}_h(a)\text{Op}_h(b) + O_{L^2 \rightarrow L^2}(h)$ for any $a, b \in S^{\text{comp}}(X)$, the self-adjointness of Π_b , and the unitarity of \mathcal{U}_b on the microsupport of Π_b , we see that we have

$$\begin{aligned} \langle \text{Op}_h(\pi_b^2 a) \chi E_h, \chi E_h \rangle_{L^2(X)} &= \langle \text{Op}_h(a) \Pi_b \chi E_h, \Pi_b \chi E_h \rangle_{L^2(X)} + O(h) \\ &= \langle \mathcal{U}_b \text{Op}_h(a) \mathcal{U}_b^* \mathcal{U}_b \Pi_b E_h, \mathcal{U}_b \Pi_b \chi E_h \rangle_{L^2(X)} + O(h). \end{aligned}$$

Now, using Egorov's theorem ([Zworski 2012, Theorem 11.1]), we know that

$$\mathcal{U}_b \text{Op}_h(a) \mathcal{U}_b^* \mathcal{U}_b \Pi_b = \text{Op}_h(a_b) \mathcal{U}_b \Pi_b + O_{L^2(X) \rightarrow L^2(\mathbb{R}^d)}(h^\infty),$$

where $a_b = a \circ \kappa_b + O_{L^2}(h)$. Using decomposition (51), we have

$$\begin{aligned} &\langle \text{Op}_h(\pi_b^2 a) \chi E_h, \chi E_h \rangle_{L^2(X)} \\ &= \sum_{n=0}^{\lfloor M_c |\log h| \rfloor} \sum_{\beta \in \tilde{\mathcal{B}}_n} \left\langle \text{Op}_h(a_b) [e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M_c |\log h| \rfloor} \sum_{\beta' \in \tilde{\mathcal{B}}_{n'}} e^{i\phi_{n',\beta',b}/h} a_{n',\beta',b} \right\rangle + O(h^c). \end{aligned} \quad (88)$$

But thanks to estimate (52),

$$\sum_{n=0}^{\lfloor M_c |\log h| \rfloor} \sum_{\beta \in \tilde{\mathcal{B}}_n} e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b} = \sum_{n=0}^{\lfloor M |\log h| \rfloor} \sum_{\beta \in \tilde{\mathcal{B}}_n} e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b} + O_{L^2}(h^c),$$

so that

$$\begin{aligned} &\langle \text{Op}_h(\pi_b^2 a) \chi E_h, \chi E_h \rangle_{L^2(X)} \\ &= \sum_{n=0}^{\lfloor M |\log h| \rfloor} \sum_{\beta \in \tilde{\mathcal{B}}_n} \left\langle \text{Op}_h(a_b) [e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M |\log h| \rfloor} \sum_{\beta' \in \tilde{\mathcal{B}}_{n'}} e^{i\phi_{n',\beta',b}/h} a_{n',\beta',b} \right\rangle + O(h^c). \end{aligned} \quad (89)$$

We now want to fix a $n \leq M|\log h|$ and a $\beta \in \tilde{\mathcal{B}}_n$, and to analyse the behaviour of

$$\left\langle \text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M|\log h \rfloor} \sum_{\beta' \in \tilde{\mathcal{B}}_n} e^{i\phi_{n',\beta',b}/h}a_{n',\beta',b} \right\rangle.$$

Let us define $Y_{n',\beta'} = \{y^{\rho_b} \in \text{spt}(\phi_{n,\beta,b}) \cap \text{spt}(\phi_{n',\beta',b}) : \partial\phi_{n',\beta',b}(y^{\rho_b}) = \partial\phi_{n,\beta,b}(y^{\rho_b})\}$. We have

$$\begin{aligned} & \left\langle \text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}], e^{i\phi_{n',\beta',b}/h}a_{n',\beta',b} \right\rangle \\ &= \int_{Y_{n',\beta'}} (\text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}])(y^{\rho_b}) e^{i\phi_{n',\beta',b}(y^{\rho_b})/h} a_{n',\beta',b}(y^{\rho_b}; h) dy^{\rho_b} \\ &+ \int_{\mathbb{R}^d \setminus Y_{n',\beta'}} (\text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}])(y^{\rho_b}) e^{i\phi_{n',\beta',b}(y^{\rho_b})/h} a_{n',\beta',b}(y^{\rho_b}; h) dy^{\rho_b}. \quad (90) \end{aligned}$$

Recall that the integrals are well defined, because the phase functions are well defined in a neighbourhood of the functions $a_{n,\beta,b}$.

The second term on the right-hand side of (90) is a $O(h^\infty)$. Indeed, the image of a Lagrangian state by a pseudodifferential operator is still a Lagrangian state with the same phase. Therefore, we are computing scalar products between Lagrangian states with respective phases $\phi_{n,\beta,b}$ and $\phi_{n',\beta',b}$.

Now, by the choice of M , and by Corollary 65, we know that for each $y^{\rho_b} \in \mathbb{R}^d \setminus Y_{n',\beta'}$ we have $|\partial\phi_{n,\beta,b}(y^{\rho_b}) - \partial\phi_{n',\beta',b}(y^{\rho_b})| \geq Ch^{1/2+\epsilon}$ for some $C, \epsilon > 0$. Hence by Proposition 62, we deduce that the second term on the right-hand side of (90) is a $O(h^\infty)$.

We should now try to understand the properties of the set $Y_{n',\beta'}$.

First of all, $Y_{n',\beta'}$ is an open set. Indeed, if $y^{\rho_b} \in Y_{n',\beta'}$, then the point $\rho = (y^{\rho_b}, \partial\phi_{n,\beta,b}(y^{\rho_b}))$ (in the coordinates centred at ρ_b) belongs to $\Phi_\beta^{n,t_0}(\mathcal{L}_0)$ as well as to $\Phi_{\beta'}^{n',t_0}(\mathcal{L}_0)$ in the notation of Proposition 64. Suppose for simplicity that $n = n'$ (the general case works the same). Then the condition $y^{\rho_b} \in Y_{n',\beta'}$ simply means that $\Phi^{n-k}(\rho)$ was both in V_{β_k} and in $V_{\beta'_k}$ at each intermediate time k . This is clearly an open condition.

On the other hand, by continuity of the phase functions, $Y_{n',\beta'}$ is a closed set. Therefore, $Y_{n',\beta'}$ consists of a certain number of connected components of the support of $\phi_{n',\beta',b}$.

We know that the support of $a_{n',\beta',b}$ is included in the domain of definition of $\phi_{n',\beta',b}$. Therefore, some of the connected components of $\text{spt}(a_{n',\beta',b})$ may be included in $Y_{n',\beta'}$, while others are included in $\mathbb{R}^d \setminus Y_{n',\beta'}$, but none of them may intersect both sets. Therefore, if we set $a_{n',\beta',b}^{n,\beta}(y^{\rho_b}) = a_{n',\beta',b}(y^{\rho_b})$ if $y^{\rho_b} \in Y_{n',\beta'}$ and equal to 0 otherwise, then $a_{n',\beta',b}^{n,\beta} \in S$, and we have

$$\begin{aligned} & \left\langle \text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M|\log h \rfloor} \sum_{\beta' \in \mathcal{B}_n} e^{i\phi_{n',\beta',b}/h}a_{n',\beta',b} \right\rangle \\ &= \int_{\mathbb{R}^d} (\text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h}a_{n,\beta,b}])(y^{\rho_b}) e^{-i\phi_{n,\beta,b}(y^{\rho_b})/h} \left(\sum_{n'=0}^{\lfloor M|\log h \rfloor} \sum_{\beta' \in \mathcal{B}_n} a_{n',\beta',b}^{n,\beta}(y^{\rho_b}) \right) dy^{\rho_b}. \quad (91) \end{aligned}$$

Let us write

$$\tilde{a}_{n,\beta,b} := \sum_{n'=0}^{\lfloor M|\log h| \rfloor} \sum_{\beta' \in \mathcal{B}_n} a_{n',\beta',b}^{n,\beta}.$$

So $\tilde{a}_{n,\beta,b}(y^{\rho_b})$ is the sum of all the symbols in the expansion (51) having phase $\phi_{n,\beta,b}(y^{\rho_b})$. We see by the estimate (52) that $\tilde{a}_{n,\beta,b}$ satisfies (52) itself, and that

$$\begin{aligned} & \left\langle \text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M|\log h| \rfloor} \sum_{\beta' \in \mathcal{B}_n} e^{i\phi_{n',\beta',b}/h} a_{n',\beta',b} \right\rangle \\ &= \int_{\mathbb{R}^d} (\text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b}])(y^{\rho_b}) e^{-i\phi_{n,\beta,b}(y^{\rho_b})/h} \tilde{a}_{n,\beta,b}(y^{\rho_b}) dy^{\rho_b} + O(h^\infty). \end{aligned}$$

We may then compute this expression using stationary phase, just as to compute the semiclassical measure of a Lagrangian state (see [Zworski 2012, §5.1]). We obtain

$$\left\langle \text{Op}_h(a_b)[e^{i\phi_{n,\beta,b}/h} a_{n,\beta,b}], \sum_{n'=0}^{\lfloor M|\log h| \rfloor} \sum_{\beta' \in \mathcal{B}_n} e^{i\phi_{n',\beta',b}/h} a_{n',\beta',b} \right\rangle = \int_{\mathbb{R}^{2n}} a_b d\mu_{n,\beta,b},$$

where

$$d\mu_{n,\beta,b} = a_{n,\beta,b}(y^{\rho_b}) \overline{\tilde{a}_{n,\beta,b}(y^{\rho_b})} \delta_{\{\eta^{\rho_b} = \partial\phi_{n,\beta,b}(y^{\rho_b})\}} dy^{\rho_b}.$$

Summing over all n, β and using equation (89), we obtain indeed that

$$\langle \text{Op}_h(\pi_b^2 a) E_h, E_h \rangle = \int_{T^*X} a(x, \xi) d\mu_{b,\chi}(x, \xi) + O(h^c),$$

with $(\kappa_b)^* \mu_{b,\chi} = \sum_{n=0}^\infty \sum_{\beta \in \mathcal{B}_n} \mu_{n,\beta,b}$; that is to say

$$d\mu_{b,\chi}(\kappa_b^{-1}(y^{\rho_b}, \eta^{\rho_b})) = \sum_{n=0}^\infty \sum_{\beta \in \mathcal{B}_n} e_{n,\beta,b}(y^{\rho_b}) \delta_{\{\eta^{\rho_b} = \partial\phi_{n,\beta,b}(y^{\rho_b})\}} dy^{\rho_b},$$

where $e_{n,\beta,b}(y^{\rho_b}) := \lim_{h \rightarrow 0} (a_{n,\beta,b} \overline{\tilde{a}_{n,\beta,b}})(y^{\rho_b})$. This concludes the proof of Corollary 50. □

6.3. Construction of the measure μ^ξ . In the Introduction we defined the measure μ^ξ by

$$\int_{T^*\mathbb{R}^d} a d\mu^\xi := \lim_{t \rightarrow \infty} \int_{T^*\mathbb{R}^d} a \circ \Phi^t d\mu_0^\xi$$

for any $a \in C_c^0(T^*\mathbb{R}^d)$. We will now give a sketch of the proof of why the hyperbolicity and transversality hypotheses, along with the assumption that $\mathcal{P}(1) < 0$, imply that the above limit exists.

Note that the assumption $\mathcal{P}(1) < 0$ is really less restrictive than $\mathcal{P}(\frac{1}{2}) < 0$. For instance, if we assume that the flow (Φ^t) is *axiom A*, that is to say, that the periodic orbits are dense in K , then [Bowen 1975, §4.C] guarantees us that $\mathcal{P}(1) < 0$.

Note that, if a is nonnegative, then $t \mapsto \int_{T^*\mathbb{R}^d} a \circ \Phi^t d\mu_0^\xi$ is nondecreasing, so that we only have to show that this quantity is bounded.

If μ is a measure, we define $\Phi_*^t \mu$ by

$$\int_{T^*\mathbb{R}^d} a \, d(\Phi_*^t \mu) := \int_{T^*\mathbb{R}^d} a \circ \Phi^t \, d\mu.$$

If $\pi \in C^\infty(T^*\mathbb{R}^d; [0, 1])$ we define the measure $\pi\mu$ by

$$\int_{T^*\mathbb{R}^d} a \, d(\pi\mu) := \int_{T^*X} a\pi \, d\mu.$$

Remark 66. Note that if μ is the semiclassical measure associated to a Lagrangian state ϕ_h , then $\pi\mu$ is the semiclassical measure associated to $\sqrt{p_i}\phi_h$, and, by Egorov’s theorem, $\Phi_*^t \mu$ is the semiclassical measure associated to $U(t)\phi_h$.

We shall use the functions π_b from Section 4.4. If $\beta \in B^n$, we set

$$\Phi_\beta \mu := \pi_{\beta_n} \Phi_*^{t_0} (\cdots \pi_{\beta_2} \Phi_*^{t_0} (\pi_{\beta_1} \Phi_*^{t_0} \mu)).$$

Let ϕ_h be a Lagrangian state associated to a Lagrangian manifold which is γ -unstable in the coordinates (y^ρ, η^ρ) , and let μ be the semiclassical measure associated to ϕ_h . The propagation $U_\beta \phi_h$ can be described using the methods of Section 4.3 along with the results of Section 2. In particular, we obtain, like in [Nonnenmacher and Zworski 2009, (7.12)], that we may find $C, \epsilon > 0$ such that for all $N \in \mathbb{N}$ and all $\beta \in B_1^N$, we have

$$\|U_\beta \phi_h\|_{L^2} \leq C(1 + C\epsilon)^N \prod_{j=1}^N \exp\left[\frac{1}{2} S_{t_0}(V_{\beta_j})\right].$$

We may deduce from this the following bound for the measure $\Phi_\beta \mu$. Note that this could also be deduced directly from the transport equations for measures, without using Schrödinger propagators and Egorov’s theorem.

For any $a \in C_c^0(T^*X)$, if $\beta \in B_1^N$, we have that

$$\langle \Phi_\beta \mu, a \rangle \leq C(a)(1 + C\epsilon)^N \prod_{j=1}^N \exp[S_{t_0}(V_{\beta_j})].$$

By possibly taking the sets V_b smaller, we may ensure, just like in Section 4.1, that

$$\sum_{b \in B_1} \exp\{S_{t_0}(V_b)\} \leq \exp\{t_0(\mathcal{P}(1) + \epsilon)\}.$$

Therefore, we obtain that

$$\sum_{\beta \in B_1^N} \langle \Phi_\beta \mu, a \rangle \leq C(a) \exp[-Nt_0(\mathcal{P}(1) - \epsilon)]. \tag{92}$$

If we assume that the flow (Φ^t) is axiom A, that is to say, that the periodic orbits are dense in K , then [Bowen 1975, §4.C] guarantees us that $\mathcal{P}(1) < 0$.

Now, we have that

$$\Phi_*^{Nt_0} \mu^\xi = \sum_{\beta \in \tilde{\mathcal{B}}^N} \Phi_\beta \mu^\xi,$$

and we may use (92) along with the assumption that $\mathcal{P}(1) < 0$ to show that, if a is nonnegative, $t \mapsto \int_{T^*\mathbb{R}^d} a \circ \Phi^t d\mu_0^\xi$ is bounded.

Showing that μ^ξ is the semiclassical measure associated to E_h follows from [Dyatlov and Guillarmou 2014, §5.1] (which relies on Egorov’s theorem), along with estimate (47).

Acknowledgements

The author is partially supported by the Agence Nationale de la Recherche project GeRaSic (ANR-13-BS01-0007-01).

The author would like to thank Stéphane Nonnenmacher for suggesting this project, as well as for his advice during the redaction of this paper. He also thanks the anonymous referee for suggesting several clarifications and improvements in the paper.

References

- [Bonthonneau 2014] Y. Bonthonneau, “Long time quantum evolution of observables”, preprint, 2014. [arXiv](#)
- [Bony et al. 2010] J.-F. Bony, N. Burq, and T. Ramond, “Minoration de la résolvente dans le cas captif”, *C. R. Math. Acad. Sci. Paris* **348**:23–24 (2010), 1279–1282. [MR](#) [Zbl](#)
- [Bowen 1975] R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Mathematics **470**, Springer, 1975. [MR](#) [Zbl](#)
- [Dimassi and Sjöstrand 1999] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series **268**, Cambridge University Press, 1999. [MR](#) [Zbl](#)
- [Dyatlov 2012] S. Dyatlov, “Microlocal limits of Eisenstein functions away from the unitarity axis”, *J. Spectr. Theory* **2**:2 (2012), 181–202. [MR](#) [Zbl](#)
- [Dyatlov and Guillarmou 2014] S. Dyatlov and C. Guillarmou, “Microlocal limits of plane waves and Eisenstein functions”, *Ann. Sci. Éc. Norm. Supér. (4)* **47**:2 (2014), 371–448. [MR](#) [Zbl](#)
- [Dyatlov and Zworski 2017] S. Dyatlov and M. Zworski, “Mathematical theory of scattering resonances”, book in progress, 2017, available at http://math.mit.edu/~dyatlov/res/res_20170323.pdf.
- [Guillarmou and Naud 2014] C. Guillarmou and F. Naud, “Equidistribution of Eisenstein series on convex co-compact hyperbolic manifolds”, *Amer. J. Math.* **136**:2 (2014), 445–479. [MR](#) [Zbl](#)
- [Hörmander 1983] L. Hörmander, *The analysis of linear partial differential operators, I: Distribution theory and Fourier analysis*, Grundlehren der Mathematischen Wissenschaften **256**, Springer, 1983. [MR](#) [Zbl](#)
- [Ingreneau 2017] M. Ingreneau, “Distorted plane waves on manifolds of nonpositive curvature”, *Comm. Math. Phys.* **350**:2 (2017), 845–891. [MR](#) [Zbl](#)
- [Katok and Hasselblatt 1995] A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications **54**, Cambridge University Press, 1995. [MR](#) [Zbl](#)
- [Melrose 1995] R. B. Melrose, *Geometric scattering theory*, Cambridge University Press, 1995. [MR](#) [Zbl](#)
- [Nonnenmacher and Zworski 2009] S. Nonnenmacher and M. Zworski, “Quantum decay rates in chaotic scattering”, *Acta Math.* **203**:2 (2009), 149–233. [MR](#) [Zbl](#)
- [Shnirelman 1974] A. I. Shnirelman, “Ergodic properties of eigenfunctions”, *Uspehi Mat. Nauk* **29**:6(180) (1974), 181–182. In Russian. [MR](#) [Zbl](#)

- [Sjöstrand 1990] J. Sjöstrand, “Geometric bounds on the density of resonances for semiclassical problems”, *Duke Math. J.* **60**:1 (1990), 1–57. [MR](#) [Zbl](#)
- [Colin de Verdière 1985] Y. Colin de Verdière, “Ergodicité et fonctions propres du laplacien”, *Comm. Math. Phys.* **102**:3 (1985), 497–502. [MR](#) [Zbl](#)
- [Zelditch 1987] S. Zelditch, “Uniform distribution of eigenfunctions on compact hyperbolic surfaces”, *Duke Math. J.* **55**:4 (1987), 919–941. [MR](#) [Zbl](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, American Mathematical Society, Providence, RI, 2012. [MR](#) [Zbl](#)

Received 7 Jan 2016. Revised 19 Nov 2016. Accepted 7 Mar 2017.

MAXIME INGREMEAU: maxime.ingremeau@math.u-psud.fr
Université Paris-Sud, 91400 Orsay, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2017 is US \$265/year for the electronic version, and \$470/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 10 No. 4 2017

The Fuglede conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_p$	757
ALEX IOSEVICH, AZITA MAYELI and JONATHAN PAKIANATHAN	
Distorted plane waves in chaotic scattering	765
MAXIME INGREMEAU	
A Fourier restriction theorem for a two-dimensional surface of finite type	817
STEFAN BUSCHENHENKE, DETLEF MÜLLER and ANA VARGAS	
On the 3-dimensional water waves system above a flat bottom	893
XUECHENG WANG	
Improving Beckner's bound via Hermite functions	929
PAATA IVANISVILI and ALEXANDER VOLBERG	
Positivity for fourth-order semilinear problems related to the Kirchhoff–Love functional	943
GIULIO ROMANI	
Geometric control condition for the wave equation with a time-dependent observation domain	983
JÉRÔME LE ROUSSEAU, GILLES LEBEAU, PEPPINO TERPOLILLI and EMMANUEL TRÉLAT	