

ANALYSIS & PDE

Volume 11 No. 3 2018

DAVID CRUZ-URIBE, JOSÉ MARÍA MARTELL AND CRISTIAN RIOS

**ON THE KATO PROBLEM AND
EXTENSIONS FOR DEGENERATE ELLIPTIC OPERATORS**



ON THE KATO PROBLEM AND EXTENSIONS FOR DEGENERATE ELLIPTIC OPERATORS

DAVID CRUZ-URIBE, JOSÉ MARÍA MARTELL AND CRISTIAN RIOS

We study the Kato problem for divergence form operators whose ellipticity may be degenerate. The study of the Kato conjecture for degenerate elliptic equations was begun by Cruz-Uribe and Rios (2008, 2012, 2015). In these papers the authors proved that given an operator $L_w = -w^{-1} \operatorname{div}(A \nabla)$, where w is in the Muckenhoupt class A_2 and A is a w -degenerate elliptic measure (that is, $A = wB$ with $B(x)$ an $n \times n$ bounded, complex-valued, uniformly elliptic matrix), then L_w satisfies the weighted estimate $\|\sqrt{L_w} f\|_{L^2(w)} \approx \|\nabla f\|_{L^2(w)}$. In the present paper we solve the L^2 -Kato problem for a family of degenerate elliptic operators. We prove that under some additional conditions on the weight w , the following unweighted L^2 -Kato estimates hold:

$$\|L_w^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}.$$

This extends the celebrated solution to the Kato conjecture by Auscher, Hofmann, Lacey, McIntosh, and Tchamitchian, allowing the differential operator to have some degree of degeneracy in its ellipticity. For example, we consider the family of operators $L_\gamma = -|x|^\gamma \operatorname{div}(|x|^{-\gamma} B(x) \nabla)$, where B is any bounded, complex-valued, uniformly elliptic matrix. We prove that there exists $\varepsilon > 0$, depending only on dimension and the ellipticity constants, such that

$$\|L_\gamma^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}, \quad -\varepsilon < \gamma < \frac{2n}{n+2}.$$

The case $\gamma = 0$ corresponds to the case of uniformly elliptic matrices. Hence, our result gives a range of γ 's for which the classical Kato square root proved in Auscher et al. (2002) is an interior point.

Our main results are obtained as a consequence of a rich Calderón–Zygmund theory developed for certain operators naturally associated with L_w . These results, which are of independent interest, establish estimates on $L^p(w)$, and also on $L^p(v dw)$ with $v \in A_\infty(w)$, for the associated semigroup, its gradient, the functional calculus, the Riesz transform, and vertical square functions. As an application, we solve some unweighted L^2 -Dirichlet, regularity and Neumann boundary value problems for degenerate elliptic operators.

1.	Introduction	610
2.	Preliminaries	613
3.	Off-diagonal estimates for the semigroup e^{-tL_w}	621
4.	The functional calculus	624
5.	Square function estimates for the semigroup	628

MSC2010: primary 35B45, 35J15, 35J25, 35J70, 42B20; secondary 42B37, 47A07, 47B44, 47D06.

Keywords: Muckenhoupt weights, degenerate elliptic operators, Kato problem, semigroups, holomorphic functional calculus, square functions, square roots of elliptic operators, Riesz transforms, Dirichlet problem, regularity problem, Neumann problem.

6. Reverse inequalities	633
7. The gradient of the semigroup $\sqrt{t}\nabla e^{-tL_w}$	638
8. An upper bound for $\mathcal{K}(L_w)$	642
9. Riesz transform estimates	645
10. Square function estimates for the gradient of the semigroup	650
11. Unweighted L^2 Kato estimates	652
12. Applications to L^2 boundary value problems	656
Acknowledgements	659
References	659

1. Introduction

We study the degenerate elliptic operators $L_w = -w^{-1} \operatorname{div} A \nabla$, where w is in the Muckenhoupt class A_2 and $A(x)$ is an $n \times n$ complex-valued matrix that satisfies the degenerate ellipticity condition

$$\lambda w(x) |\xi|^2 \leq \operatorname{Re} \langle A(x) \xi, \xi \rangle, \quad |\langle A(x) \xi, \eta \rangle| \leq \Lambda w(x) |\xi| |\eta|, \quad \xi, \eta \in \mathbb{C}^n, \text{ a.e. } x \in \mathbb{R}^n.$$

Equivalently, $A(x) = w(x) B(x)$, where B is an $n \times n$ complex-valued matrix that satisfies the uniform ellipticity conditions

$$\lambda |\xi|^2 \leq \operatorname{Re} \langle B(x) \xi, \xi \rangle, \quad |\langle B(x) \xi, \eta \rangle| \leq \Lambda |\xi| |\eta|, \quad \xi, \eta \in \mathbb{C}^n, \text{ a.e. } x \in \mathbb{R}^n.$$

Such operators were first studied (with A a real symmetric matrix) by Fabes, Kenig and Serapioni [Fabes et al. 1982]. When A is complex-valued and uniformly elliptic (i.e., $w \equiv 1$), a landmark result was the proof by Auscher, Hofmann, Lacey, McIntosh, and Tchamitchian [Auscher et al. 2002] of the Kato conjecture, which states that for all $f \in H^1$,

$$\|L^{1/2} f\|_2 \approx \|\nabla f\|_2.$$

The proof of this long-standing conjecture led naturally to the study of the operators associated with L : the semigroup e^{-tL} , its gradient $\sqrt{t}\nabla e^{-tL}$, the Riesz transform $\nabla L^{-1/2}$, the H^∞ functional calculus and square functions; for details and complete references, see [Auscher 2007]. These estimates are interesting in themselves; moreover, it is well known that L^p estimates for these operators yield regularity results for boundary value problems for L ; for details, see the introduction to [Auscher and Tchamitchian 1998].

In [Cruz-Uribe and Rios 2015] (see also [Cruz-Uribe and Rios 2008; 2012; Auscher et al. 2015]), the first and third authors solved the Kato problem for degenerate elliptic operators: they showed that if $w \in A_2$ and A satisfies the degenerate ellipticity conditions, then for all $f \in H^1(w)$,

$$\|L_w^{1/2} f\|_{L^2(w)} \approx \|\nabla f\|_{L^2(w)}. \tag{1.1}$$

In this paper we consider the problem of determining those A_2 weights such that the classical Kato problem can be solved for L_w , that is, finding weights such that L_w satisfies the unweighted estimate

$$\|L_w^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}$$

for f in a class of nice functions (a posteriori, by standard density arguments, the estimate can be extended to all $f \in H^1(\mathbb{R}^n)$). We solve this problem in two steps. The first is to prove weighted L^p estimates for some operators associated with L_w (the semigroup, its gradient, the Riesz transform, the functional calculus, and square functions). These results, which are of interest in their own right, are analogous to those obtained in the uniformly elliptic case. However, a significant technical obstruction is that given a weight $w \in A_2$, while it is the case that there exists $\varepsilon > 0$ such that $w \in A_{2-\varepsilon}$, it is easy to construct examples to show that ε may be arbitrarily small. Therefore, our bounds in the range $1 < p < 2$ need to take this into account.

The second step is to find conditions on the weight w so that these operators satisfy *unweighted* L^2 estimates. Both steps are carried out simultaneously, and the proofs are intertwined. Our approach is to apply the theory of off-diagonal estimates on balls developed by Auscher and the second author [Auscher and Martell 2006; 2007a; 2007b; 2008]. We will in fact prove weighted estimates on $L^p(v \, dw)$, where v satisfies Muckenhoupt and reverse Hölder conditions with respect to the measure $dw = w \, dx$; $L^p(w)$ estimates are then obtained by taking $v = 1$, and unweighted estimates by taking $v = w^{-1}$.

The unweighted L^2 estimates are delicate, since they require a careful estimate of the constants that appear. Nevertheless, we are able to give useful sufficient conditions; e.g., $w \in A_1 \cap \text{RH}_{n/2+1}$. (For definitions of these classes, see Section 2 below.) For example, we have the following result that is a special case of one of our main results (cf. Theorem 11.11).

Theorem 1.2. *Let $L_w = -w^{-1} \operatorname{div} A \nabla$ be a degenerate elliptic operator as above. If $w \in A_1 \cap \text{RH}_{n/2+1}$, then the Kato problem can be solved for L_w : for every $f \in H^1(\mathbb{R}^n)$,*

$$\|L_w^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}.$$

The implicit constants depend only on the dimension, the ellipticity constants, and the A_1 and $\text{RH}_{n/2+1}$ constants of w .

Furthermore, if we define $L_\gamma = -|x|^\gamma \operatorname{div}(|x|^{-\gamma} B(x) \nabla)$, where B is an $n \times n$ complex-valued matrix that satisfies the uniform ellipticity condition, then there exists $0 < \varepsilon < \frac{1}{2}$ small enough (depending only on the dimension and the ratio Λ/λ) such that

$$\|L_\gamma^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}, \quad -\varepsilon < \gamma < \frac{2n}{n+2}.$$

Remark 1.3. In Theorem 1.2 the operator $L_w^{1/2}$ is a priori only defined on $H^1(w)$; however, this means that it is defined on $C_0^\infty(\mathbb{R}^n)$ and so by a standard density argument we can extend our results to all $f \in H^1(\mathbb{R}^n)$. Hereafter we will make this extension without further comment.

We emphasize that in Theorem 1.2, when $\gamma = 0$ we are back at the uniformly elliptic case, which is the celebrated solution to the Kato square root problem by Auscher, Hofmann, Lacey, McIntosh, and Tchamitchian [Auscher et al. 2002]. Here we are able to find a range of γ 's for which the same estimates hold and the classical Kato square root problem (i.e., $\gamma = 0$) is an interior point in that range.

These unweighted L^2 estimates have important applications to boundary value problems for degenerate elliptic operators. Consider, for example, the following Dirichlet problem on $\mathbb{R}_+^{n+1} = \mathbb{R}^n \times [0, \infty)$:

$$\begin{cases} \partial_t^2 u - L_w u = 0 & \text{on } \mathbb{R}_+^{n+1}, \\ u = f & \text{on } \partial\mathbb{R}_+^{n+1} = \mathbb{R}^n. \end{cases}$$

If $f \in L^2(\mathbb{R}^n)$, then $u(x, t) = e^{-tL_w^{1/2}} f(x)$ is a solution, and if L_w has a bounded H^∞ functional calculus on L^2 , then $\sup_{t>0} \|u(\cdot, t)\|_2 \lesssim \|f\|_2$. Similar results hold for the corresponding Neumann and regularity problems.

Our proofs are unavoidably technical, and the results for each operator considered build upon what was proved previously for other operators. We have organized the material as follows. In Section 2 we gather some essential definitions and results about weights, degenerate elliptic operators, and off-diagonal estimates. Central to all of our subsequent work are Theorems 2.35 and 2.39, which were proved in [Auscher and Martell 2006].

In Sections 3, 4, and 5 we prove estimates for the semigroup e^{-tL_w} , $t > 0$, the H^∞ functional calculus (i.e., operators $\varphi(L_w)$ where $\varphi \in \mathcal{H}^\infty$), the vertical square function associated to the semigroup,

$$g_{L_w} f(x) = \left(\int_0^\infty |(tL_w)^{1/2} e^{-tL_w} f(x)|^2 \frac{dt}{t} \right)^{1/2},$$

and its discrete analog. Here and in subsequent sections we prove both $L^p(w)$ estimates and weighted $L^p(v dw)$ estimates. In many cases these results are proved simultaneously, with the unweighted results (i.e., in $L^p(w)$) following from the weighted results (i.e., in $L^p(v dw)$) by taking $v = 1$.

In Section 6 we prove the so-called reverse inequality, $\|L_w^{1/2}\|_{L^p(w)} \lesssim \|\nabla f\|_{L^p(w)}$, that generalizes the $L^2(w)$ estimate in (1.1). We note that while the equivalence in (1.1) follows at once from the reverse inequality for $p = 2$ by duality, the two inequalities behave differently when $p \neq 2$.

In Sections 7 and 8 we prove estimates for the gradient of the semigroup, $\sqrt{t}\nabla e^{-tL_w}$. The proof that there exists $q_+ > 2$ such that this operator satisfies $L^p(w)$ estimates for $2 < p < q_+$ is quite involved as it requires preliminary estimates for the Riesz transform and the Hodge projection. We note that, as opposed to the nondegenerate case, here we cannot use “global” embeddings, nor can we rescale. Also we cannot expect to obtain that the gradient of the semigroup maps globally $L^2(w)$ into $L^p(w)$ for $p \neq 2$. All these difficulties arise naturally from the lack of isotropy of the natural underlying measure $w(x) dx$ and make the typical arguments used in the uniformly elliptic case (see [Auscher 2007, Chapter 4]) unusable. We also note that in some sense our result is the best possible: even in the nondegenerate case it is known [Auscher 2007] that given any $p > 2$ there exists a matrix A and operator L such that gradient of the semigroup is not bounded on L^p .

In Section 9 we prove $L^p(w)$ estimates for the Riesz transform $\nabla L^{-1/2}$, and in Section 10 we prove $L^p(w)$ estimates for the square function associated to the gradient of the semigroup,

$$G_{L_w} f(x) = \left(\int_0^\infty |t^{1/2} \nabla e^{-tL_w} f(x)|^2 \frac{dt}{t} \right)^{1/2}.$$

In Section 11 we prove unweighted L^2 inequalities for the operators we have considered in previous sections. These are a consequence of the weighted estimates and are obtained by taking $v = w^{-1}$. The main problem is determining conditions on w for these to hold. We essentially have two different kinds of estimates, one for operators that do not involve the gradient, and one for those that do. The latter are more delicate as they involve careful bounds for the parameter q_+ from Section 8 in terms of the weight w . We also show that we get unweighted L^p estimates for p very close to 2.

Finally, in Section 12 we describe in more detail the application of our results to L^2 boundary value problems for degenerate elliptic operators. The results in this section are the culmination of our work, as they depend on all the estimates derived in previous sections.

As we were completing this project, we learned that related results had been obtained independently by other authors. Le [2015] studied (among other things) the $L^p(w)$ theory for some of the operators considered here and proved estimates for values of p in the range $(2 - \varepsilon, 2 + \varepsilon)$. His proofs differ from ours in a number of details. Hofmann, Le and Morris [Hofmann et al. 2015] established some Carleson measure estimates and considered the Dirichlet problem for degenerate elliptic operators. Also, very recently we learned that Yang and Zhang [2017] proved Kato-type estimates in $L^p(w)$ for p in the range $(p_0, 2]$. Finally, we note that the paper [Chen et al. 2016] complements our work here as it considers the conical square functions associated to the operator L_w .

2. Preliminaries

Throughout, n will denote the dimension of the underlying space \mathbb{R}^n and we will always assume $n \geq 2$. If we write $A \lesssim B$ we mean that there exists a constant C such that $A \leq CB$. We write $A \approx B$ if $A \lesssim B$ and $B \lesssim A$. The constant C in these estimates may depend on the dimension n and other (fixed) parameters that should be clear from the context. All constants, explicit or implicit, may change at each appearance.

Given a ball B , let $r(B)$ denote the radius of B . Let λB denote the concentric ball with radius $r(\lambda B) = \lambda r(B)$.

Weights. By a weight w we mean a nonnegative, locally integrable function. For brevity, we will often write dw for $w dx$. We will use the following notation for averages: given a set E such that $0 < w(E) < \infty$,

$$\int_E f dw = \frac{1}{w(E)} \int_E f dw,$$

or, if $0 < |E| < \infty$,

$$\int_E f dx = \frac{1}{|E|} \int_E f dx.$$

We state some definitions and basic properties of Muckenhoupt weights. For further details, see [Duoandikoetxea 2001; García-Cuerva and Rubio de Francia 1985]. We say that $w \in A_p$, $1 < p < \infty$, if

$$[w]_{A_p} = \sup_Q \int_Q w(x) dx \left(\int_Q w(x)^{1-p'} dx \right)^{p-1} < \infty.$$

When $p = 1$, we say that $w \in A_1$ if

$$[w]_{A_1} = \sup_Q \int_Q w(x) dx \operatorname{ess\,sup}_{x \in Q} w(x)^{-1} < \infty.$$

We say that $w \in \operatorname{RH}_s$, $1 < s < \infty$, if

$$[w]_{\operatorname{RH}_s} = \sup_Q \left(\int_Q w(x) dx \right)^{-1} \left(\int_Q w(x)^s dx \right)^{1/s} < \infty,$$

and we say that $w \in \operatorname{RH}_\infty$ if

$$[w]_{\operatorname{RH}_\infty} = \sup_Q \left(\int_Q w(x) dx \right)^{-1} \operatorname{ess\,sup}_{x \in Q} w(x) < \infty.$$

Let

$$A_\infty = \bigcup_{1 \leq p < \infty} A_p = \bigcup_{1 < s \leq \infty} \operatorname{RH}_s.$$

Weights in the A_p and RH_s classes have a self-improving property: if $w \in A_p$, there exists $\varepsilon > 0$ such that $w \in A_{p-\varepsilon}$, and similarly if $w \in \operatorname{RH}_s$, then $w \in \operatorname{RH}_{s+\delta}$ for some $\delta > 0$. Hereafter, given $w \in A_p$, let

$$r_w = \inf\{p : w \in A_p\}, \quad s_w = \sup\{q : w \in \operatorname{RH}_q\}.$$

An important property of A_p weights is that they are doubling: given $w \in A_p$, for all $\tau \geq 1$ and any ball B ,

$$w(\tau B) \leq [w]_{A_p} \tau^{pn} w(B).$$

In particular, hereafter let $D \leq pn$ be the doubling order of w , that is, the smallest exponent such that this inequality holds.

As a consequence of this doubling property, we have that with the ordinary Euclidean distance $|\cdot|$, $(\mathbb{R}^n, dw, |\cdot|)$ is a space of homogeneous type. In this setting we can define the new weight classes $A_p(w)$ and $\operatorname{RH}_s(w)$ by replacing Lebesgue measure in the definitions above with dw ; e.g., $v \in A_p(w)$ if

$$[v]_{A_p(w)} = \sup_Q \int_Q v(x) dw \left(\int_Q v(x)^{1-p'} dw \right)^{p-1} < \infty.$$

It follows at once from these definitions that there is a “duality” relationship between the weighted and unweighted A_p and RH_s conditions: $v = w^{-1} \in A_p(w)$ if and only if $w \in \operatorname{RH}_{p'}$ and $v = w^{-1} \in \operatorname{RH}_s(w)$ if and only if $w \in A_{s'}$.

Weighted Poincaré–Sobolev inequalities were proved in [Fabes et al. 1982].

Theorem 2.1. *Given $w \in A_p$, $p \geq 1$, let $p_w^* = pnr_w/(nr_w - p)$ if $p < nr_w$, and $p_w^* = \infty$ otherwise. Then for every $p \leq q < p_w^*$, ball B and $f \in C_0^\infty(B)$,*

$$\left(\int_B |f(x)|^q dw(x) \right)^{1/q} \leq Cr(B) \left(\int_B |\nabla f(x)|^p dw \right)^{1/p}. \quad (2.2)$$

Moreover, if $f \in C^\infty(B)$, then

$$\left(\int_B |f(x) - f_{B,w}|^q dw(x) \right)^{1/q} \leq Cr(B) \left(\int_B |\nabla f(x)|^p dw \right)^{1/p}, \tag{2.3}$$

where $f_{B,w} = \int_B f dw$.

Remark 2.4. In the special case when $w \in A_1$ and $1 < p < n$ we can also take $q = p_w^* = p^*$, i.e., the regular Sobolev exponent. See [Pérez 1999, Theorem 2.5.2].

Remark 2.5. If we let $q = np/(n - 1) < p_w^*$, then we can get a sharp estimate for the constant C in (2.2) and (2.3): it is of the form $C(p, n)[w]_{A_p}^\kappa$, where $\kappa = (np - 1)/(np(p - 1))$. This follows from the sharp weighted estimates for the fractional integral operator due to Alberico, Cianchi and Sbordone [Alberico et al. 2009] and the standard pointwise estimates used to prove Poincaré–Sobolev inequalities; see [Fabes et al. 1982] for details.

Remark 2.6. By a standard density argument, once we know that (2.3) holds for smooth functions in B we can easily extend that estimate to any function $f \in L^q(w)$ with $\nabla f \in L^p(w)$. Details are left to the reader.

Degenerate elliptic operators. Given $w \in A_2$ and constants $0 < \lambda \leq \Lambda < \infty$, let $\mathcal{E}_n(w, \lambda, \Lambda)$ denote the class of $n \times n$ matrices $A = (A_{ij}(x))_{i,j=1}^n$ of complex-valued, measurable functions satisfying the degenerate ellipticity condition

$$\lambda w(x)|\xi|^2 \leq \operatorname{Re}\langle A\xi, \xi \rangle, \quad |\langle A\xi, \eta \rangle| \leq \Lambda w(x)|\xi||\eta|, \quad \xi, \eta \in \mathbb{C}^n. \tag{2.7}$$

Given $A \in \mathcal{E}_n(w, \lambda, \Lambda)$, we define the degenerate elliptic operator in divergence form

$$L_w = -w^{-1} \operatorname{div} A \nabla.$$

These operators were developed in [Cruz-Uribe and Rios 2008] and we refer the reader there for complete details. Here we sketch the key ideas.

Given a weight $w \in A_2$, the space $H^1(w)$ is the weighted Sobolev space that is the completion of C_c^∞ with respect to the norm

$$\|f\|_{H^1(w)} = \left(\int_{\mathbb{R}^n} (|f(x)|^2 + |\nabla f(x)|^2) dw \right)^{1/2}.$$

Note that the space defined above would usually be denoted by $H_0^1(w)$. The space $H^1(w)$ is defined as the set of distributions for which both f and $|\nabla f|$ belong to $L^2(w)$. However, since the underlying domain is \mathbb{R}^n , this definition implies that the “boundary” values vanish in the $L^2(w)$ -sense, and both definitions agree [Miller 1982].

Given a matrix $A \in \mathcal{E}_n(w, \lambda, \Lambda)$, define $\mathfrak{a}(f, g)$ to be the sesquilinear form

$$\mathfrak{a}(f, g) = \int_{\mathbb{R}^n} A(x) \nabla f(x) \cdot \overline{\nabla g(x)} dx. \tag{2.8}$$

Since $w \in A_2$ and A satisfies (2.7), α is a closed, maximally accretive, continuous sesquilinear form. Therefore, there exists an operator L_w whose domain $\mathcal{D}(L_w) \subset H^1(w)$ is dense in $L^2(w)$ and such that for every $f \in \mathcal{D}(L_w)$ and every $g \in H^1(w)$,

$$\alpha(f, g) = \langle L_w f, g \rangle_w = \int_{\mathbb{R}^n} L_w f(x) \overline{g(x)} dx. \quad (2.9)$$

We note that the operator L_w is one-to-one. Indeed, if $u, v \in \mathcal{D}(L_w)$ are such that $L_w u = L_w v$, then for all $g \in H^1(w)$

$$0 = \int_{\mathbb{R}^n} A(x) \nabla(u(x) - v(x)) \cdot \overline{\nabla g(x)} dx.$$

Taking $g = u - v$ implies $\nabla u(x) = \nabla v(x)$ and so $u = v$.

The properties of the sesquilinear form guarantee that on $L^2(w)$ there exists a bounded, strongly continuous semigroup e^{-tL_w} . Further, it has a holomorphic extension. Let

$$\Sigma_\omega = \{z \in \mathbb{C} : z \neq 0, |\arg(z)| < \omega\}$$

and define $\vartheta, \vartheta^* \in [0, \frac{\pi}{2})$ by

$$\vartheta = \sup\{|\arg\langle Lf, f \rangle_w| : f \in \mathcal{D}(L_w)\}, \quad \vartheta^* = \arctan \sqrt{\frac{\Lambda^2}{\lambda^2} - 1}.$$

Then there exists a complex semigroup e^{-zL_w} on $\Sigma_{\pi/2-\vartheta}$ of bounded operators on $L^2(w)$. By the weighted ellipticity condition (2.7), we have $0 \leq \vartheta \leq \vartheta^* < \frac{\pi}{2}$.

Holomorphic functional calculus. Our operator L_w is “an operator of type ω ” with $\omega = \vartheta$, as defined in [McIntosh 1986]. Indeed, the ellipticity conditions imply that L_w is closed and densely defined, its spectrum is contained in Σ_ϑ , and its resolvent satisfies standard decay estimates [Cruz-Uribe and Rios 2008]. Therefore, we can define an $L^2(w)$ functional calculus as in [McIntosh 1986].

Given $\mu \in (\vartheta, \pi)$, let $\mathcal{H}^\infty(\Sigma_\mu)$ be the collection of bounded holomorphic functions on Σ_μ . To define $\varphi(L_w)$ for $\varphi \in \mathcal{H}^\infty(\Sigma_\mu)$ we first consider a smaller class: we say that $\varphi \in \mathcal{H}_0^\infty(\Sigma_\mu)$ if for some $c, s > 0$ it satisfies

$$|\varphi(z)| \leq c|z|^s(1+|z|)^{-2s}, \quad z \in \Sigma_\mu.$$

We then have an integral representation of $\varphi(L_w)$. Let Γ_θ be the boundary of Σ_θ with positive orientation, and let $\vartheta < \theta < \nu < \min(\mu, \frac{\pi}{2})$; then

$$\varphi(L_w) = \int_{\Gamma_{\pi/2-\theta}} e^{-zL_w} \eta(z) dz, \quad (2.10)$$

where

$$\eta(z) = \frac{1}{2\pi i} \int_{\gamma_\nu(z)} e^{\xi z} \varphi(\xi) d\xi \quad (2.11)$$

and $\gamma_\nu(z) = \mathbb{R}^+ e^{i \operatorname{sign}(\operatorname{Im}(z))\nu}$. Note that

$$|\eta(z)| \lesssim \min\{1, |z|^{-s-1}\}, \quad z \in \Gamma_{\pi/2-\theta},$$

so the representation (2.10) converges in $L^2(w)$, and we have the bound

$$\|\varphi(L_w)f\|_{L^2(w)} \leq C \|\varphi\|_\infty \|f\|_{L^2(w)}, \quad f \in \mathcal{H}_0^\infty(\Sigma_\mu). \quad (2.12)$$

Now, since L_w is a one-to-one operator of type ω , it has dense range [Cowling et al. 1996, Theorem 2.3], and so the results in [McIntosh 1986] (see also [Cowling et al. 1996, Corollary 2.2]) imply that L_w has an H^∞ functional calculus and (2.12) extends to all of $\mathcal{H}^\infty(\Sigma_\mu)$. Moreover, in [McIntosh 1986, Section 8] the equivalence between the existence of this H^∞ functional calculus and square function estimates for L_w and L_w^* is established:

$$\left(\int_0^\infty \|\varphi(tL_w)\|_{L^2(w)}^2 \frac{dt}{t} \right)^{1/2} \leq C \|\varphi\|_\infty \|f\|_{L^2(w)}, \quad \varphi \in \mathcal{H}_0^\infty(\Sigma_\mu), \quad (2.13)$$

with similar estimates for L_w^* .

The operators $\varphi(L_w)$ also have the following properties:

- If φ and ψ are bounded holomorphic functions, then we have the operator identity $\varphi(L)\psi(L) = (\varphi\psi)(L)$.
- Given any sequence $\{\varphi_k\}$ of bounded holomorphic functions converging uniformly on compact subsets of Σ_μ to φ , we have that $\varphi_k(L_w)$ converges to $\varphi(L_w)$ in the strong operator topology (of operators on $L^2(w)$).

Remark 2.14. The H^∞ functional calculus can be extended to more general holomorphic functions, such as powers, for which the operators $\varphi(L_w)$ can be defined as unbounded operators; see [Haase 2006; McIntosh 1986].

Gaffney-type estimates. The semigroup and its gradient satisfy Gaffney-type estimates on $L^2(w)$. Below, we will see that these are a particular case of what we will call full off-diagonal estimates; see Definition 2.33.

Theorem 2.15. *Given $w \in A_2$ and $A \in \mathcal{E}_n(w, \lambda, \Lambda)$, for any closed sets E and F , for $f \in L^2(w)$ and for all $z \in \Sigma_\nu$, where $0 < \nu < \frac{\pi}{2} - \vartheta$,*

- (1) $\|e^{-zL_w}(f\chi_E)\chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)^2/|z|} \|f\chi_E\|_{L^2(w)},$
- (2) $\|\sqrt{z}\nabla e^{-zL_w}(f\chi_E)\chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)^2/|z|} \|f\chi_E\|_{L^2(w)},$
- (3) $\|zL_w e^{-zL_w}(f\chi_E)\chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)^2/|z|} \|f\chi_E\|_{L^2(w)}.$

Proof. The semigroup estimate (1) was proved in [Cruz-Uribe and Rios 2008, Theorem 1.6] for real z , but the same proof can be readily modified to prove the analytic version. Alternatively, estimates (1) and (2) follow from the resolvent bounds

$$\|(1 + z^2L_w)^{-1}(f\chi_E)\chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)/|z|} \|f\chi_E\|_{L^2(w)}, \quad (2.16)$$

$$\|z\nabla(1 + z^2L_w)^{-1}(f\chi_E)\chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)/|z|} \|f\chi_E\|_{L^2(w)}, \quad (2.17)$$

obtained in [Cruz-Uribe and Rios 2015, Lemma 2.10] for $z \in \Sigma_{\pi/2+\nu}$, together with the integral representation of the semigroup

$$e^{-zL_w} f = \frac{1}{2\pi} \int_{\Gamma} e^{z\xi} (\xi + L_w)^{-1} f d\xi,$$

where Γ is the boundary of Σ_{θ} with positive orientation and $\frac{\pi}{2} < \theta < \frac{\pi}{2} + \nu - \arg(z)$.

Finally, from (2.16) and (2.17) we obtain the estimate

$$\|z^2 L_w (1 + z^2 L_w)^{-1} (f \chi_E) \chi_F\|_{L^2(w)} \leq C e^{-cd(E,F)/|z|} \|f \chi_E\|_{L^2(w)},$$

and then by the same kind of argument we get (3). \square

The Kato estimate. The starting point for all of our estimates is the $L^2(w)$ Kato estimates for the square root operator $L_w^{1/2}$ proved in [Cruz-Uribe and Rios 2015] (see also [Auscher et al. 2015] for a different proof). This operator is the unique, maximal accretive operator such that $L_w^{1/2} L_w^{1/2} = L_w$. It has the integral representation

$$L_w^{1/2} = \frac{1}{\sqrt{\pi}} \int_0^{\infty} \sqrt{t} L_w e^{-tL_w} \frac{dt}{t}.$$

(For further details, see [Auscher and Tchamitchian 1998; McIntosh 1986].)

Theorem 2.18 [Cruz-Uribe and Rios 2015, Theorem 1.1]. *Given $w \in A_2$ and $A \in \mathcal{E}_n(w, \lambda, \Lambda)$, the domain of L_w is $H^1(w)$ and there exist constants c and C , depending on n , Λ/λ and $[w]_{A_2}$, such that for all $f \in H^1(w)$,*

$$c \|\nabla f\|_{L^2(w)} \leq \|L_w^{1/2} f\|_{L^2(w)} \leq C \|\nabla f\|_{L^2(w)}. \quad (2.19)$$

The Riesz transform associated to L_w is the operator $\nabla L_w^{-1/2}$. Formally, by (2.19) we have that the Riesz transform is a bounded operator on $L^2(w, \mathbb{C}^n)$. To legitimize this, we define

$$\nabla L_w^{-1/2} = \frac{1}{\sqrt{\pi}} \int_0^{\infty} \sqrt{t} \nabla e^{-tL_w} \frac{dt}{t}. \quad (2.20)$$

However, it is not immediate that this integral converges at 0 or ∞ . To rectify this, for $\varepsilon > 0$ define

$$S_{\varepsilon} = S_{\varepsilon}(L_w) = \frac{1}{\sqrt{\pi}} \int_{\varepsilon}^{1/\varepsilon} \sqrt{t} e^{-tL_w} \frac{dt}{t}. \quad (2.21)$$

Since $S_{\varepsilon}(z)$ is a uniformly bounded holomorphic function on the right half-plane for all $0 < \varepsilon < 1$, by the $L^2(w)$ functional calculus described above, $S_{\varepsilon}(L_w)$ is uniformly bounded on $L^2(w)$ for that range of ε . Further, for $f \in L_c^{\infty}$, we have $S_{\varepsilon} f \in \mathcal{D}(L_w) \subset \mathcal{D}(L_w^{1/2})$, and so by inequality (2.19) and the functional calculus,

$$\|\nabla S_{\varepsilon} f\|_{L^2(w)} \lesssim \|L_w^{1/2} S_{\varepsilon} f\|_{L^2(w)} = \|\varphi_{\varepsilon}(L_w) f\|_{L^2(w)}, \quad (2.22)$$

where

$$\varphi_{\varepsilon}(z) = \frac{1}{\sqrt{\pi}} \int_{\varepsilon}^{1/\varepsilon} \sqrt{t} \sqrt{z} e^{-tz} \frac{dt}{t}.$$

The sequence $\{\varphi_\varepsilon\}$ is uniformly bounded and converges uniformly to 1 on compact subsets of the sector Σ_μ , $0 < \mu < \frac{\pi}{2}$. Therefore, $L^{1/2}S_\varepsilon f \rightarrow f$ strongly in $L^2(w)$. If we combine this fact with (2.22) we see that $\{\nabla S_\varepsilon f\}$ is Cauchy and so it converges in $L^2(w)$. We therefore define

$$\nabla L^{-1/2} f = \lim_{\varepsilon \rightarrow 0} \nabla S_\varepsilon f,$$

where the limit is in $L^2(w)$.

Given this definition, hereafter, when we are proving $L^2(w)$ estimates for the Riesz transform, we should actually prove estimates for ∇S_ε that are independent of ε . These arguments will remain implicit unless there are details we need to emphasize.

Off-diagonal estimates. Off-diagonal estimates as we define them were introduced in [Auscher and Martell 2007b] and we will refer repeatedly to this paper for further information and results. Throughout this section we will assume that given a weight w , we have $w \in A_2$.

Given a ball B , for $j \geq 2$ we define the annuli $C_j(B) = 2^{j+1}B \setminus 2^jB$. We let $C_1(B) = 4B$. By a slight abuse of notation, we will define

$$\int_{C_j(B)} h \, dw = \frac{1}{w(2^{j+1}B)} \int_{C_j(B)} h \, dw.$$

If $w \in A_2$ (as it will be hereafter), then $w(2^{j+1}B) \approx w(C_j(B))$, so this definition is equivalent to the one given above up to a constant. Finally, for $s > 0$ we set $\Upsilon(s) = \max\{s, s^{-1}\}$.

Definition 2.23. Given $1 \leq p \leq q \leq \infty$, a family $\{T_t\}_{t>0}$ of sublinear operators satisfies $L^p(w) - L^q(w)$ off-diagonal estimates on balls, denoted by

$$T_t \in \mathcal{O}(L^p(w) \rightarrow L^q(w)),$$

if there exist constants $\theta_1, \theta_2 > 0$ and $c > 0$ such that for every $t > 0$ and for any ball B , setting $r = r(B)$,

$$\left(\int_B |T_t(\chi_B f)|^q \, dw \right)^{1/q} \lesssim \Upsilon\left(\frac{r}{\sqrt{t}}\right)^{\theta_2} \left(\int_B |f|^p \, dw \right)^{1/p}, \tag{2.24}$$

and for all $j \geq 2$,

$$\left(\int_B |T_t(\chi_{C_j(B)} f)|^q \, dw \right)^{1/q} \lesssim 2^{j\theta_1} \Upsilon\left(\frac{2^j r}{\sqrt{t}}\right)^{\theta_2} e^{-c4^j r^2/t} \left(\int_{C_j(B)} |f|^p \, dw \right)^{1/p}, \tag{2.25}$$

$$\left(\int_{C_j(B)} |T_t(\chi_B f)|^q \, dw \right)^{1/q} \lesssim 2^{j\theta_1} \Upsilon\left(\frac{2^j r}{\sqrt{t}}\right)^{\theta_2} e^{-c4^j r^2/t} \left(\int_B |f|^p \, dw \right)^{1/p}. \tag{2.26}$$

If the family of sublinear operators $\{T_z\}_{z \in \Sigma_\mu}$ is defined on a complex sector Σ_μ , we say that it satisfies $L^p(w) - L^q(w)$ off-diagonal estimates on balls in Σ_μ if (2.24)–(2.26) hold for $z \in \Sigma_\mu$ with t replaced by $|z|$ in the right-hand terms. We denote this by $T_z \in \mathcal{O}(L^p(w) \rightarrow L^q(w), \Sigma_\mu)$.

We give some basic properties of off-diagonal estimates on balls as a series of lemmas taken from [Auscher and Martell 2007b, Section 2.2]. The first follows immediately by real interpolation, the second by Hölder’s inequality, and the third by duality.

Lemma 2.27. *Given $1 \leq p_i \leq q_i \leq \infty$, $i = 1, 2$, if $T_t \in \mathcal{O}(L^{p_1}(w) \rightarrow L^{q_1}(w))$ and $T_t : L^{p_2}(w) \rightarrow L^{q_2}(w)$ is uniformly bounded, then $T_t \in \mathcal{O}(L^{p_\theta}(w) \rightarrow L^{q_\theta}(w))$, $0 < \theta < 1$, where*

$$\frac{1}{p_\theta} = \frac{\theta}{p_1} + \frac{1-\theta}{p_2}, \quad \frac{1}{q_\theta} = \frac{\theta}{q_1} + \frac{1-\theta}{q_2}.$$

Lemma 2.28. *If $1 \leq p \leq p_1 \leq q_1 \leq q \leq \infty$, then*

$$\mathcal{O}(L^p(w) \rightarrow L^q(w)) \subset \mathcal{O}(L^{p_1}(w) \rightarrow L^{q_1}(w)).$$

Lemma 2.29. *If for some $1 \leq p \leq q \leq \infty$, we have $T_t \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, and the operators T_t are linear, then $T_t^* \in \mathcal{O}(L^{q'}(w) \rightarrow L^{p'}(w))$. (Here T_t^* is the dual operator for the inner product $\int_{\mathbb{R}^n} fg \, dw$.)*

Lemma 2.30 [Auscher and Martell 2007b, Theorem 2.3]. (1) *If $T_t \in \mathcal{O}(L^p(w) \rightarrow L^p(w))$, $1 \leq p \leq \infty$, then $T_t : L^p(w) \rightarrow L^p(w)$ is uniformly bounded.*

(2) *If $1 \leq p \leq q \leq r \leq \infty$, $T_t \in \mathcal{O}(L^q(w) \rightarrow L^r(w))$, and $S_t \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, then $T_t \circ S_t \in \mathcal{O}(L^p(w) \rightarrow L^r(w))$.*

Remark 2.31. *If $p < q$, then $T_t \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$ does not guarantee that T_t is bounded from $L^p(w)$ to $L^q(w)$.*

Remark 2.32. *Since complex sectors Σ_μ , $0 \leq \mu < \pi$, are closed under addition, the proof of Lemma 2.30 extends to give off-diagonal estimates on complex sectors $\mathcal{O}(L^p(w) \rightarrow L^q(w), \Sigma_\mu)$.*

Definition 2.33. *Given $1 \leq p \leq q \leq \infty$, a family of operators $\{T_t\}$ satisfies full off-diagonal estimates from $L^p(w)$ to $L^q(w)$, denoted by*

$$T_t \in \mathcal{F}(L^p(w) \rightarrow L^q(w)),$$

if there exist constants $C, c, \theta > 0$ such that given any closed sets E, F ,

$$\|T_t(f \chi_E) \chi_F\|_{L^q(w)} \leq C t^{-\theta} e^{-cd^2(E,F)/t} \|f \chi_E\|_{L^p(w)}.$$

The connection between full off-diagonal estimates and off-diagonal estimates on balls is given in the following lemma from [Auscher and Martell 2007b, Section 3.1].

Lemma 2.34. *Given $1 \leq p \leq q \leq \infty$:*

- (1) *if $T_t \in \mathcal{F}(L^p(w) \rightarrow L^q(w))$, then $T_t : L^p(w) \rightarrow L^q(w)$ is uniformly bounded;*
- (2) *$T_t \in \mathcal{F}(L^p(w) \rightarrow L^p(w))$ if and only if $T_t \in \mathcal{O}(L^p(w) \rightarrow L^p(w))$.*

The importance of off-diagonal estimates is that they will let us prove weighted norm inequalities for the operators we are interested in. To do so we will make repeated use of two results first proved in [Auscher and Martell 2007a]; however, we will use special cases of these results as given in [Auscher and Martell 2006, Theorems 2.2 and 2.4].

Theorem 2.35. *Given $w \in A_2$ and $1 \leq p_0 < q_0 \leq \infty$, let T be a sublinear operator acting on $L^{p_0}(w)$, $\{\mathcal{A}_r\}_{r>0}$ a family of operators acting from a subspace \mathcal{D} of $L^{p_0}(w)$ into $L^{p_0}(w)$, and S an operator from \mathcal{D} into the space of measurable functions on \mathbb{R}^n . Suppose that for every $f \in \mathcal{D}$ and ball B with radius r ,*

$$\left(\int_B |T(I - \mathcal{A}_r)f|^{p_0} dw \right)^{1/p_0} \leq \sum_{j \geq 1} g(j) \left(\int_{2^{j+1}B} |Sf|^{p_0} dw \right)^{1/p_0}, \tag{2.36}$$

$$\left(\int_B |T\mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} \leq \sum_{j \geq 1} g(j) \left(\int_{2^{j+1}B} |Tf|^{p_0} dw \right)^{1/p_0}, \tag{2.37}$$

where $\sum g(j) < \infty$. Then for every p , $p_0 < p < q_0$, and weights

$$v \in A_{p/p_0}(w) \cap \text{RH}_{(q_0/p)'}(w),$$

there is a constant C such that for all $f \in \mathcal{D}$,

$$\|Tf\|_{L^p(v dw)} \leq C \|Sf\|_{L^p(v dw)}.$$

Remark 2.38. In Theorem 2.35 and Theorem 2.39 below, the case $q_0 = \infty$ is understood in the sense that the $L^{q_0}(w)$ -average is replaced by the essential supremum. Also in Theorem 2.35, if $q_0 = \infty$, then the condition on v becomes $v \in A_{p/p_0}$.

Theorem 2.39. *Given $w \in A_2$ with doubling order D , and $1 \leq p_0 < q_0 \leq \infty$, let $T : L^{q_0}(w) \rightarrow L^{q_0}(w)$ be a sublinear operator, and $\{\mathcal{A}_r\}_{r>0}$ a family of linear operators acting from L_c^∞ into $L^{q_0}(w)$. Suppose that for every ball B with radius r , $f \in L_c^\infty$ with $\text{supp}(f) \subset B$ and $j \geq 2$,*

$$\left(\int_{C_j(B)} |T(I - \mathcal{A}_r)f|^{p_0} dw \right)^{1/p_0} \leq g(j) \left(\int_B |f|^{p_0} dw \right)^{1/p_0}. \tag{2.40}$$

Suppose further that for every $j \geq 1$,

$$\left(\int_{C_j(B)} |\mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} \leq g(j) \left(\int_B |f|^{p_0} dw \right)^{1/p_0}, \tag{2.41}$$

where $\sum g(j)2^{Dj} < \infty$. Then for all p , $p_0 < p < q_0$, there exists a constant C such that for all $f \in L_c^\infty$,

$$\|Tf\|_{L^p(w)} \leq C \|f\|_{L^p(w)}.$$

3. Off-diagonal estimates for the semigroup e^{-tL_w}

In this section we consider off-diagonal estimates for the semigroup associated to L_w . Throughout this and subsequent sections, let $w \in A_2$ and $A \in \mathcal{E}_n(w, \Lambda, \lambda)$ be fixed. Our goal is to characterize the set of pairs (p, q) , $p \leq q$, such that these operators are in $\mathcal{O}(L^p(w) \rightarrow L^q(w))$. By Theorem 2.15 we have

$$e^{-tL_w} \in \mathcal{F}(L^2(w) \rightarrow L^2(w)) \subset \mathcal{O}(L^2(w) \rightarrow L^2(w)).$$

We will show that in the (p, q) -plane this set contains a right triangle; see Figure 1.




Figure 1. (p, q) such that $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$

Let $\tilde{\mathcal{J}}(L_w) \subset [1, \infty]$ be the set of all exponents p such that $e^{-tL_w} : L^p(w) \rightarrow L^p(w)$ is uniformly bounded for all $t > 0$. By Theorem 2.15 and Lemma 2.34, $2 \in \tilde{\mathcal{J}}(L_w)$, and if it contains more than one point, then by interpolation $\tilde{\mathcal{J}}(L_w)$ is an interval. The set of pairs (p, q) such that $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$ is completely characterized by the next result.

Proposition 3.1. *There exists an interval $\mathcal{J}(L_w) \subset [1, \infty]$ such that $p, q \in \mathcal{J}(L_w)$ if and only if $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$. Furthermore, $\mathcal{J}(L_w)$ has the following properties:*

- (1) $\mathcal{J}(L_w) \subset \tilde{\mathcal{J}}(L_w)$.
- (2) $\text{Int } \mathcal{J}(L_w) = \text{Int } \tilde{\mathcal{J}}(L_w)$.
- (3) If $p_-(L_w)$ and $p_+(L_w)$ are respectively the left and right endpoints of $\mathcal{J}(L_w)$, then $p_-(L_w) \leq (2_w^*)'$ and $p_+(L_w) \geq 2_w^*$, where 2_w^* is as in Theorem 2.1. In particular, $2 \in \text{Int}(\mathcal{J}(L_w))$.

Remark 3.2. The smaller the value of r_w , the better our bounds on the size of the set $\mathcal{J}(L_w)$. In the limiting case when $w \in A_1$, we have $p_-(L_w) \leq 2n/(n+2)$ and $p_+(L_w) \geq 2n/(n-2)$. These values should be compared to the estimates in [Auscher 2007, Corollary 4.6] for the nondegenerate case that corresponds to the case $w = 1$.

We get two corollaries to Proposition 3.1. The first gives us weighted off-diagonal estimates.

Corollary 3.3. *Let $p_-(L_w) < p \leq q < p_+(L_w)$. If $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/q)'(w)}$, then $e^{-tL_w} \in \mathcal{O}(L^p(v dw) \rightarrow L^q(v dw))$.*

Proof. By Proposition 3.1, if $p_-(L_w) < p \leq q < p_+(L_w)$, then $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$. Therefore, by [Auscher and Martell 2007b, Proposition 2.6], if $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/q)'(w)}$, then we have $e^{-tL_w} \in \mathcal{O}(L^p(v dw) \rightarrow L^q(v dw))$. \square

As our second corollary we get off-diagonal estimates for the holomorphic extension of the semigroup.

Corollary 3.4. *For any ν , $0 < \nu < \frac{\pi}{2} - \vartheta$, and for any $p \leq q$ such that $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, then for all $m \in \mathbb{N} \cup \{0\}$, $(zL_w)^m e^{-zL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w), \Sigma_\nu)$.*

Proof. This follows from [Auscher and Martell 2007b, Theorem 4.3] and the fact that, by Theorem 2.15, $e^{-zL_w} \in \mathcal{F}(L^2(w) \rightarrow L^2(w))$ for these values of z . \square

Proof of Proposition 3.1. Fix $2 < q < 2_w^*$. (If $w \in A_1$ we let $q = 2_w^* = 2^*$.) We will show that $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$. Given this, then we also have $e^{-tL_w} \in \mathcal{O}(L^{q'}(w) \rightarrow L^2(w))$. For if L_w^* is the adjoint of L_w (with respect to $L^2(w)$), then $L_w^* = -w^{-1} \operatorname{div}(A^* \nabla f)$ and the same estimates hold for L_w^* . Hence, $e^{-tL_w^*} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$, and so by Lemma 2.29, $e^{-tL_w} \in \mathcal{O}(L^{q'}(w) \rightarrow L^2(w))$. Since e^{-tL_w} is a semigroup, by Lemma 2.30 we have $e^{-tL_w} \in \mathcal{O}(L^{q'}(w) \rightarrow L^q(w))$. Therefore, by [Auscher and Martell 2007b, Proposition 4.1], we have that there exists an interval $\mathcal{J}(L_w)$ and properties (1) and (2) hold. Moreover, we have $[q', q] \subset \mathcal{J}(L_w)$, so if we let $q \rightarrow 2_w^*$, then we immediately get property (3).

It therefore remains to prove that $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$. We first show (2.24). Fix B and for brevity write $r = r(B)$ and $C_j = C_j(B)$. By our choice of q , the Poincaré inequality (2.3) holds. Moreover, as we noted above, $e^{-tL_w}, \sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$; we may assume that the same exponents θ_1, θ_2 hold for both operators. We thus get that

$$\begin{aligned} & \left(\int_B |e^{-tL_w}(\chi_B f)|^q dw \right)^{1/q} \\ & \leq |(e^{-tL_w}(\chi_B f))_{B,w}| + \left(\int_B |e^{-tL_w}(\chi_B f)(x) - (e^{-tL_w}(\chi_B f))_{B,w}|^q dw(x) \right)^{1/q} \\ & \lesssim \left(\int_B |e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2} + r \left(\int_B |\nabla e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2} \\ & \lesssim \left(1 + \frac{r}{\sqrt{t}} \right) \Upsilon \left(\frac{r}{\sqrt{t}} \right)^{\theta_2} \left(\int_B |f|^2 dw \right)^{1/2} \\ & \lesssim \Upsilon \left(\frac{r}{\sqrt{t}} \right)^{1+\theta_2} \left(\int_B |f|^2 dw \right)^{1/2}. \end{aligned}$$

The proof that (2.25) holds is gotten by nearly the same argument:

$$\begin{aligned} & \left(\int_B |e^{-tL_w}(\chi_{C_j} f)|^q dw \right)^{1/q} \\ & \leq |(e^{-tL_w}(\chi_{C_j} f))_{B,w}| + \left(\int_B |e^{-tL_w}(\chi_{C_j} f)(x) - (e^{-tL_w}(\chi_{C_j} f))_{B,w}|^q dw(x) \right)^{1/q} \\ & \lesssim \left(\int_B |e^{-tL_w}(\chi_{C_j} f)|^2 dw \right)^{1/2} + r \left(\int_B |\nabla e^{-tL_w}(\chi_{C_j} f)|^2 dw \right)^{1/2} \\ & \lesssim 2^{j\theta_1} \left(1 + \frac{r}{\sqrt{t}} \right) \Upsilon \left(\frac{2^j r}{\sqrt{t}} \right)^{\theta_2} e^{-c4^j r^2/t} \left(\int_{C_j} |f|^2 dw \right)^{1/2} \\ & \lesssim 2^{j\theta_1} \Upsilon \left(\frac{2^j r}{\sqrt{t}} \right)^{1+\theta_2} e^{-c4^j r^2/t} \left(\int_{C_j} |f|^2 dw \right)^{1/2}. \end{aligned}$$

Finally, to prove that (2.26) holds we use a covering argument. Fix $j \geq 2$; then we can cover the annulus C_j by a collection of balls $\{B_k\}_{k=1}^N$, $r(B_k) = 2^{j-2}r$, with centers $x_{B_k} \in C_j$. The number of balls required, N , depends only on the dimension. For any such ball, since dw is a doubling measure, we have

$$\begin{aligned} & \left(\int_{B_k} |e^{-tL_w}(\chi_B f)|^q dw \right)^{1/q} \\ & \leq |(e^{-tL_w}(\chi_B f))_{B_k, w}| + \left(\int_{B_k} |e^{-tL_w}(\chi_B f)(x) - (e^{-tL_w}(\chi_B f))_{B_k, w}|^q dw(x) \right)^{1/q} \\ & \lesssim \left(\int_{B_k} |e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2} + r(B_k) \left(\int_{B_k} |\nabla e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2} \\ & \lesssim \left(\int_{2^{j+2}B \setminus 2^{j-1}B} |e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2} + 2^j r \left(\int_{2^{j+2}B \setminus 2^{j-1}B} |\nabla e^{-tL_w}(\chi_B f)|^2 dw \right)^{1/2}. \end{aligned}$$

If $j \geq 3$, then $2^{j+2}B \setminus 2^{j-1}B = C_{j+1} \cup C_j \cup C_{j-1}$; then to estimate the last two terms we use the fact that $e^{-tL_w}, \sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$ and apply (2.26) with $p = q = 2$ in each annulus C_i , $j-1 \leq i \leq j+1$. (These annuli have comparable measure since dw is a doubling measure, so we can divide the average up into three averages). If $j = 2$, then $2^4B \setminus 2B = C_3 \cup C_2 \cup (4B \setminus 2B)$. On C_3 and C_2 we argue as before using (2.26). On $4B \setminus B$ we apply [Auscher and Martell 2007b, Lemma 6.1]. (We note that in the notation there, $\hat{C}_1(B) = 4B \setminus 2B$.)

If we combine all of these estimates, we get that for every $j \geq 2$,

$$\begin{aligned} \left(\int_{B_k} |e^{-tL_w}(\chi_B f)|^q dw \right)^{1/q} & \lesssim 2^{j\theta_1} \left(1 + \frac{2^j r}{\sqrt{t}} \right) \Gamma \left(\frac{2^j r}{\sqrt{t}} \right)^{\theta_2} \left(\int_B |f|^2 dw \right)^{1/2} \\ & \lesssim 2^{j\theta_1} \Gamma \left(\frac{2^j r}{\sqrt{t}} \right)^{1+\theta_2} e^{-c4^j r^2/t} \left(\int_B |f|^2 dw \right)^{1/2}. \end{aligned}$$

Since $C_j \subset \bigcup_k B_k$, we can sum in k to get

$$\begin{aligned} \left(\int_{C_j(B)} |e^{-tL_w}(\chi_B f)|^q dw \right)^{1/q} & \lesssim \sum_{k=1}^N \left(\int_{B_k} |e^{-tL_w}(\chi_B f)|^q dw \right)^{1/q} \\ & \lesssim 2^{j\theta_1} \Gamma \left(\frac{2^j r}{\sqrt{t}} \right)^{1+\theta_2} e^{-c4^j r^2/t} \left(\int_B |f|^2 dw \right)^{1/2}. \quad \square \end{aligned}$$

This completes the proof that $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$.

4. The functional calculus

In this section we show that the operator L_w has an $L^p(w)$ holomorphic functional calculus. As we discussed in Section 2 above, we know already that if φ is a bounded holomorphic function on Σ_μ , $\mu \in (\vartheta, \pi)$, then $\varphi(L_w)$ is a bounded operator on $L^2(w)$. Recall that for any $\mu \in (\vartheta, \pi)$, we say that $\varphi \in \mathcal{H}_0^\infty(\Sigma_\mu)$ if for some $c, s > 0$,

$$|\varphi(z)| \leq c|z|^s(1+|z|)^{-2s}, \quad z \in \Sigma_\mu. \quad (4.1)$$

We say that L_w has a bounded holomorphic functional calculus on $L^p(w)$ if for any such φ ,

$$\|\varphi(L_w)f\|_{L^p(w)} \leq C \|\varphi\|_\infty \|f\|_{L^p(w)}, \quad f \in L^p(w) \cap L^2(w), \tag{4.2}$$

where C depends only on p, w, ϑ and μ (but not on the decay of φ). By a standard density argument, (4.2) implies that $\varphi(L_w)$ extends to a bounded operator on all of $L^p(w)$. Furthermore, we then have this inequality holds if φ is any bounded holomorphic function. For the details of this extension, see [Haase 2006; McIntosh 1986].

Proposition 4.3. *Let $p_-(L_w) < p < p_+(L_w)$ and $\mu \in (\vartheta, \pi)$. Then for any $\varphi \in \mathcal{H}_0^\infty(\Sigma_\mu)$,*

$$\|\varphi(L_w)f\|_{L^p(w)} \leq C \|\varphi\|_\infty \|f\|_{L^p(w)}, \tag{4.4}$$

with C independent of φ and f . Hence, L_w has a bounded holomorphic functional calculus on $L^p(w)$. Moreover, if $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/p)'}(w)$ then L_w also has a bounded holomorphic functional calculus on $L^p(v dw)$:

$$\|\varphi(L_w)f\|_{L^p(v dw)} \leq C \|\varphi\|_\infty \|f\|_{L^p(v dw)}, \tag{4.5}$$

with C independent of φ and f .

Proof. For brevity, let $p_- = p_-(L_w)$ and $p_+ = p_+(L_w)$. By density it will suffice to assume that $f \in L_c^\infty$. Fix $\varphi \in \mathcal{H}_0^\infty(\Sigma_\mu)$; by linearity we may assume that $\|\varphi\|_\infty = 1$.

We divide the proof into two steps. We first obtain (4.4) for $p_- < p < 2$ by applying Theorem 2.39 and following the ideas in [Auscher 2007]. To do so, we will pick $q_0 = 2$ and $p_0 > p_-$ arbitrarily close to p_- . In the second step, using some ideas from [Auscher and Martell 2006], we will use Theorem 2.35 to get (4.5); in particular this yields (4.4) for every $2 < p < p_+$ by taking $v \equiv 1$. To apply Theorem 2.35 we will choose $p_0 > p_-$ arbitrarily close to p_- and $q_0 < p_+$ arbitrarily close to p_+ . We will also use the fact that $\varphi(L_w)$ is bounded on $L^{p_0}(w)$; this follows from the first step choosing $p_- < p_0 < 2$.

To apply Theorem 2.39, fix $p_- < p_0 < p < 2$ and let $q_0 = 2, T = \varphi(L_w)$, and

$$A_r f(x) = (I - (I - e^{-r^2 L_w})^m) f(x), \tag{4.6}$$

where m is a positive integer that will be chosen below. We first show that inequality (2.41) holds. By Proposition 3.1 we have $e^{-tL_w} \in \mathcal{O}(L^{p_0}(w) \rightarrow L^2(w))$. Using

$$A_r = \sum_{k=1}^m \binom{m}{k} (-1)^{k+1} e^{-kr^2 L_w}, \tag{4.7}$$

and that for each fixed m and $1 \leq k \leq m$

$$\Upsilon\left(\frac{r}{\sqrt{kt}}\right) \leq \sqrt{m} \Upsilon\left(\frac{r}{t}\right) \quad \text{and} \quad \exp\left(-\frac{c}{k} \frac{4^j r^2}{t^2}\right) \leq \exp\left(-\frac{c}{m} \frac{4^j r^2}{t^2}\right),$$

Proposition 3.1 implies

$$A_r \in \mathcal{O}(L^p(w) \rightarrow L^q(w)) \quad \text{for all } p_-(L_w) < p \leq q < p_+(L_w). \tag{4.8}$$

In particular, we have $\mathcal{A}_r \in \mathcal{O}(L^{p_0}(w) \rightarrow L^2(w))$. Thus, given any ball B with radius r , if $\text{supp}(f) \subset B$, then for all $j \geq 1$,

$$\left(\int_{C_j(B)} |\mathcal{A}_r f|^2 dw \right)^{1/2} \lesssim 2^{j\theta_1} \Upsilon(2^j)^{\theta_2} e^{-c4^j} \left(\int_B |f|^{p_0} dw \right)^{1/p_0}. \quad (4.9)$$

This establishes (2.41) with $g(j) = C 2^{j(\theta_1+\theta_2)} e^{-c4^j}$, for in this case we have

$$\sum_{j \geq 1} 2^{j(\theta_1+\theta_2+D)} e^{-c4^j} < \infty,$$

where D is the doubling constant of w .

We next prove that (2.40) holds. Since $\varphi(z)(1 - e^{-r^2 z})^m \in \mathcal{H}_0^\infty(\Sigma_{\{\min\{\mu, \pi/2\}\}})$, by the functional calculus representation (2.10) we have

$$\varphi(L_w)(I - \mathcal{A}_r)f = \int_{\Gamma} e^{-zL_w} f \eta(z) dz,$$

where $\Gamma = \partial\Sigma_{\pi/2-\theta}$, with $0 < \vartheta < \theta < \nu < \min\{\mu, \frac{\pi}{2}\}$, and we choose θ so that the hypotheses of Corollary 3.4 are satisfied for $z \in \Gamma$. Moreover, we have the estimate

$$|\eta(z)| \lesssim \frac{r^{2m}}{|z|^{m+1}};$$

see [Auscher 2007, Section 5.1] for details.

We can now argue as follows: given a ball B with radius r , for each $j \geq 2$, by Minkowski's inequality and Corollary 3.4 (since $p_0 \in \text{Int } \mathcal{J}(L_w)$),

$$\begin{aligned} & \left(\int_{C_j(B)} |\varphi(L_w)(I - \mathcal{A}_{r(B)})f|^{p_0} dw \right)^{1/p_0} \\ &= \left(\int_{C_j(B)} \left| \int_{\Gamma} e^{-zL_w} f \eta(z) dz \right|^{p_0} dw \right)^{1/p_0} \\ &\lesssim \int_{\Gamma} \left(\int_{C_j(B)} |e^{-zL_w} f|^{p_0} dw \right)^{1/p_0} \frac{r^{2m}}{|z|^{m+1}} |dz| \\ &\lesssim \left(\int_B |f|^{p_0} dw \right)^{1/p_0} \int_{\Gamma} \frac{r^{2m}}{|z|^{m+1}} 2^{j\theta_1} \Upsilon\left(\frac{2^j r}{\sqrt{|z|}}\right)^{\theta_2} e^{-c(r^2/|z|)4^j} |dz| \\ &= \left(\int_B |f|^{p_0} dw \right)^{1/p_0} 2^{j(\theta_1-2m)} \int_0^\infty \sigma^{2m} \Upsilon(\sigma)^{\theta_2} e^{-c\sigma^2} \frac{d\sigma}{\sigma} \\ &\lesssim 2^{j(\theta_1-2m)} \left(\int_B |f|^{p_0} dw \right)^{1/p_0}; \end{aligned} \quad (4.10)$$

the final inequality holds (i.e., the integral in σ converges) provided $2m > \theta_2$. Moreover, if we choose $2m > \theta_1 + D$, we have that (2.40) holds with $g(j) = C 2^{(j-1)(\theta_1-2m)}$ and

$$\sum_{j \geq 2} g(j) 2^{jD} \lesssim \sum_{j \geq 2} 2^{j(\theta_1+D-2m)} < \infty.$$

We have shown that inequalities (2.40) and (2.41) hold, and so by Theorem 2.39 inequality (4.4) holds for all p such that $p_- < p \leq 2$.

We will now apply Theorem 2.35 to show that (4.5) holds for $p_- < p < p_+$. (Inequality (4.4) then follows for $2 < p < p_+$ if we take $v \equiv 1$.) Fix p , $p_- < p < p_+$, and $v \in A_{p/p_-}(w) \cap \text{RH}_{(p_+/p)'}(w)$. By the openness properties of the A_q and RH_s classes there exist p_0, q_0 such that

$$p_- < p_0 < \min\{p, 2\} \leq p < q_0 < p_+, \quad v \in A_{p/p_0}(w) \cap \text{RH}_{(q_0/p)'}(w).$$

Let $T = \varphi(L_w)$, $A_r = I - (I - e^{-r^2 L_w})^m$, $S = I$, and fix the above values of p_0 and q_0 . By the previous argument we have that $\varphi(L_w)$ is bounded on $L^{p_0}(w)$.

We first show that (2.36) holds. Fix a ball B and decompose f as

$$f = \sum_{j \geq 1} f \chi_{C_j(B)} := \sum_{j \geq 1} f_j. \tag{4.11}$$

Then, by the same functional calculus argument as given above, we have that for each j ,

$$\begin{aligned} & \left(\int_B |\varphi(L_w)(I - A_r)f_j|^{p_0} dw \right)^{1/p_0} \\ &= \left(\int_B \left| \int_{\Gamma} e^{-zL_w} f_j \eta(z) dz \right|^{p_0} dw \right)^{1/p_0} \\ &\lesssim \int_{\Gamma} \left(\int_B |e^{-zL_w} f_j|^{p_0} dw \right)^{1/p_0} \frac{r^{2m}}{|z|^{m+1}} |dz| \\ &\lesssim \left(\int_{C_j(B)} |f|^{p_0} dw \right)^{1/p_0} 2^{j(\theta_1-2m)} \int_{\Gamma} \left(\frac{2^j r}{\sqrt{|z|}} \right)^{2m} \Upsilon \left(\frac{2^j r}{\sqrt{|z|}} \right)^{\theta_2} e^{-c4^j r^2/|z|} \frac{|dz|}{|z|} \\ &\lesssim 2^{j(\theta_1-2m)} \left(\int_{C_j(B)} |f|^{p_0} dw \right)^{1/p_0}; \end{aligned}$$

the last inequality holds provided $2m > \theta_2$. Hence, since $2^{j+1}B \supset C_j$, by Minkowski's inequality we have (since the sum $\sum f_j$ is finite for $f \in L_c^\infty$)

$$\begin{aligned} \left(\int_B |\varphi(L_w)(I - A_r)f|^{p_0} dw \right)^{1/p_0} &\leq \sum_{j \geq 1} \left(\int_B |\varphi(L_w)(I - A_r)f_j|^{p_0} dw \right)^{1/p_0} \\ &\lesssim \sum_{j \geq 1} 2^{j(\theta_1-2m)} \left(\int_{2^{j+1}B} |f|^{p_0} dw \right)^{1/p_0}. \end{aligned}$$

This establishes (2.36) with $g(j) = C 2^{j(\theta_1-2m)}$. If we take $2m > \max\{\theta_1, \theta_2\}$, then $\sum g(j) < \infty$.

We now show that (2.37) holds. Fix a ball B and $j \geq 1$. Since $A_r \in \mathcal{O}(L^{p_0}(w) \rightarrow L^{q_0}(w))$ (see (4.8)),

$$\left(\int_B |A_r(\chi_{C_j(B)}\varphi(L_w)f)|^{q_0} dw \right)^{1/q_0} \lesssim 2^{j\theta_1} \Upsilon(2^j)^{\theta_2} e^{-c4^j} \left(\int_{C_j(B)} |\varphi(L_w)f|^{p_0} d\mu \right)^{1/p_0}.$$

Therefore, since $\varphi(L_w)$ and \mathcal{A}_r commute, by Minkowski's inequality we obtain

$$\left(\int_B |\varphi(L_w) \mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} \lesssim \sum_{j \geq 1} 2^{j(\theta_1 + \theta_2)} e^{-c4^j} \left(\int_{C_j(B)} |\varphi(L_w) f|^{p_0} d\mu \right)^{1/p_0}.$$

This establishes (2.37) with $g(j) = C 2^{j(\theta_1 + \theta_2)} e^{-c4^j}$; again, $\sum g(j) < \infty$. Therefore, our proof is complete. \square

5. Square function estimates for the semigroup

In this section we prove $L^p(w)$ norm inequalities for the vertical square function associated to the semigroup e^{-tL_w} :

$$g_{L_w} f(x) = \left(\int_0^\infty |(tL_w)^{1/2} e^{-tL_w} f(x)|^2 \frac{dt}{t} \right)^{1/2}.$$

Proposition 5.1. *Let $p_-(L_w) < p < p_+(L_w)$. Then*

$$\|g_{L_w} f\|_{L^p(w)} \approx \|f\|_{L^p(w)}. \quad (5.2)$$

Conversely if for some p the equivalence (5.2) holds, then $p \in \tilde{\mathcal{J}}(L_w)$ — i.e., the interior of the interval on which (5.2) holds is $(p_-(L_w), p_+(L_w))$.

Moreover, if $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/p)'}(w)$, then

$$\|g_{L_w} f\|_{L^p(v dw)} \approx \|f\|_{L^p(v dw)}. \quad (5.3)$$

We note that the upper bounds in the previous result could be obtained by combining Proposition 4.3 with the operator theory methods developed in [Cowling et al. 1996]. To reach a wider audience we present a self-contained harmonic analysis proof. We will use an auxiliary Hilbert space related to square functions, following the approach in [Auscher and Martell 2006]. Let \mathbb{H} denote the Hilbert space $L^2((0, \infty), \frac{dt}{t})$ with norm

$$\|h\| = \left(\int_0^\infty |h(t)|^2 \frac{dt}{t} \right)^{1/2}.$$

In particular, we have

$$g_{L_w} f(x) = \|\varphi(L, \cdot) f(x)\|,$$

where $\varphi(z, t) = (tz)^{1/2} e^{-tz}$. Furthermore, we define $L_{\mathbb{H}}^p(w)$ to be the space of \mathbb{H} -valued functions with the norm

$$\|h\|_{L_{\mathbb{H}}^p(w)} = \left(\int_{\mathbb{R}^n} \|h(x, \cdot)\|^p dw(x) \right)^{1/p}.$$

The following lemma lets us extend scalar-valued inequalities to \mathbb{H} -valued inequalities. For a proof, see [Auscher and Martell 2006, Lemma 7.4].

Lemma 5.4. *Given a Borel measure μ on \mathbb{R}^n , let \mathcal{D} be a subspace of \mathcal{M} , the space of measurable functions in \mathbb{R}^n , and let S, T be linear operators from \mathcal{D} into \mathcal{M} . Fix $1 \leq p \leq q < \infty$ and suppose there exists $C_0 > 0$ such that for all $f \in \mathcal{D}$,*

$$\|Tf\|_{L^q(\mu)} \leq C_0 \sum_{j \geq 1} \alpha_j \|Sf\|_{L^p(F_j, \mu)},$$

where the F_j are measurable subsets of \mathbb{R}^n and $\alpha_j \geq 0$. Then there is an \mathbb{H} -valued inequality with the same constant: for all $f : \mathbb{R}^n \times (0, \infty) \rightarrow \mathbb{C}$ such that for almost all $t > 0$, $f(\cdot, t) \in \mathcal{D}$,

$$\|Tf\|_{L^q_{\mathbb{H}}(\mu)} \leq C_0 \sum_{j \geq 1} \alpha_j \|Sf\|_{L^p_{\mathbb{H}}(F_j, \mu)}.$$

The extension of a linear operator T on \mathbb{C} -valued functions to \mathbb{H} -valued functions is defined for $x \in \mathbb{R}^n$ and $t > 0$ by $(Th)(x, t) = T(h(\cdot, t))(x)$; that is, t can be considered as a parameter and T acts only on the variable in \mathbb{R}^n .

Proof of Proposition 5.1. We shall first prove the upper bound inequalities. We first claim that the upper bound inequality in (5.2) holds for $p = 2$. Indeed, since $\varphi(z) = z^{1/2}e^{-z} \in \mathcal{H}_0^\infty(\Sigma_\mu)$, it follows from (2.13) that we have the bound

$$\|g_{L_w} f\|_{L^2(w)} \lesssim \|f\|_{L^2(w)}.$$

For brevity, let $p_- = p_-(L_w)$ and $p_+ = p_+(L_w)$. As in previous proofs, we divide our proof into two steps. We will first prove the upper bound in (5.2) for $p_- < p < 2$ by applying Theorem 2.39. Fix $p_- < p < q_0 = 2$, and let $\mathcal{A}_r = I - (I - e^{-r^2 L_w})^m$, where m will be chosen below. Notice that, by (4.8), \mathcal{A}_r is bounded on $L^{q_0}(w)$ for each m . Fix $f \in L_c^\infty$; the result for general $f \in L^p(w)$ then follows by a density argument.

We have $(tL_w)^{1/2}e^{-tL_w}(I - \mathcal{A}_r)f = \varphi(L_w, t)f$, where

$$\varphi(z, t) = (tz)^{1/2}e^{-tz}(1 - e^{-r^2 z})^m.$$

Moreover, since $\varphi(\cdot, t) \in \mathcal{H}_0^\infty(\Sigma_{\{\min\{\mu, \pi/2\}\}})$, by the functional calculus representation (2.10) we have

$$(tL_w)^{1/2}e^{-tL_w}(I - \mathcal{A}_r)f = \int_{\Gamma} \eta(z, t)e^{-zL_w} f dz,$$

where $\Gamma = \partial\Sigma_{\pi/2-\theta}$, with $0 < \vartheta < \theta < \nu < \min\{\mu, \frac{\pi}{2}\}$, and we choose θ so that the hypotheses of Corollary 3.4 are satisfied for $z \in \Gamma$. Moreover, we have the estimate [Auscher 2007; Auscher and Martell 2006]

$$|\eta(z, t)| \lesssim \frac{t^{1/2}r^{2m}}{(|z| + t)^{m+3/2}}, \quad z \in \Gamma.$$

Therefore,

$$\|\eta(z, \cdot)\| = \left(\int_0^\infty |\eta(z, t)|^2 \frac{dt}{t} \right)^{1/2} \lesssim \frac{r^{2m}}{|z|^{m+1}}. \tag{5.5}$$

Now let $f \in L_c^\infty$ with $\text{supp}(f) \subset B$. For $j \geq 2$, we have

$$\begin{aligned}
 \left(\int_{C_j(B)} |g_{L_w}(I - \mathcal{A}_r)f|^p dw \right)^{1/p} &= \left(\int_{C_j(B)} \left| \left(\int_0^\infty \left| \int_{\Gamma_{\pi/2-\theta}} \eta(z,t) e^{-zL_w} f dz \right|^2 \frac{dt}{t} \right)^{1/2} \right|^p dw \right)^{1/p} \\
 &\leq \left(\int_{C_j(B)} \left| \int_{\Gamma_{\pi/2-\theta}} |e^{-zL_w} f| \|\eta(z, \cdot)\| d|z| \right|^p dw \right)^{1/p} \\
 &\lesssim \int_{\Gamma_{\pi/2-\theta}} \left(\int_{C_j(B)} |e^{-zL_w} f|^p dw \right)^{1/p} \frac{r^{2m}}{|z|^{m+1}} d|z| \\
 &\lesssim 2^{j\theta_1} \left(\int_B |f|^p dw \right)^{1/p} \int_{\Gamma_{\pi/2-\theta}} \Upsilon \left(\frac{2^j r}{\sqrt{|z|}} \right)^{\theta_2} e^{-c4^j r^2/|z|} \frac{r^{2m}}{|z|^m} \frac{d|z|}{|z|} \\
 &\lesssim 2^{j\theta_1} 4^{-mj} \left(\int_B |f|^p dw \right)^{1/p}; \tag{5.6}
 \end{aligned}$$

in the second inequality we applied (5.5) and the off-diagonal estimates for e^{-zL_w} from Corollary 3.4, and the last inequality holds provided $2m > \theta_2$. Thus, if we take $m > \theta_1 + D$, where D is the doubling order of w , the operator g_{L_w} satisfies (2.40) in Theorem 2.39 with $g(j) = C 2^{j(\theta_1 - 2m)}$. Since we already established (2.41) in (4.9) with $g(j) = C 2^{j(\theta_1 + \theta_2)} 4^{-mj}$, the hypotheses of Theorem 2.39 are satisfied if $m > \theta_1 + \theta_2 + D$. Therefore, for each $p_- < p < 2$ there exists a constant C such that

$$\|g_{L_w} f\|_{L^p(w)} \leq C \|f\|_{L^p(w)}. \tag{5.7}$$

In the second part of the proof we will show that if $p_- < p < p_+$ and $v \in A_{p/p_-}(w) \cap \text{RH}_{(p_+/p)'}(w)$, then the upper bound inequality in (5.3) holds. If we take $v \equiv 1$, then we immediately get (5.2). To do so, first note that if we fix p and v , then by the openness properties of weights there exist p_0, q_0 such that

$$p_- < p_0 < \min\{p, 2\} \leq \max\{p, 2\} < q_0 < p_+$$

and $v \in A_{p_0/p_-}(w) \cap \text{RH}_{(q_0/p)'}(w)$.

We will apply Theorem 2.35 with $T = g_{L_w}$, $S = I$ and $\mathcal{D} = L^{p_0}(w)$ (again, note that by (4.8), \mathcal{A}_r is bounded on $L^{p_0}(w)$). We first prove that inequality (2.36) holds. For each $j \geq 1$, let $f_j = f \chi_{C_j(B)}$; then we can argue exactly as we did in the proof of (5.6), exchanging the roles of B and $C_j(B)$, to get

$$\left(\int_B |g_{L_w}(I - \mathcal{A}_r)f_j|^p dw \right)^{1/p} \lesssim 2^{j\theta_1} 4^{-mj} \left(\int_{2^{j+1}B} |f|^p dw \right)^{1/p}.$$

Inequality (2.36) follows if we sum over all j and take $g(j) = 2^{j\theta_1} 4^{-mj}$.

We will now show that inequality (2.37) holds. To do so, we need to prove a vector-valued version of a key inequality. By Proposition 3.1, given a ball B with radius r , we have for all $j \geq 1$, g with $\text{supp}(g) \subset C_j(B)$, and $1 \leq k \leq m$,

$$\left(\int_B |e^{-kr^2 L_w} g|^{q_0} dw \right)^{1/q_0} \leq C_0 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j} \left(\int_{C_j(B)} |g|^{p_0} dw \right)^{1/p_0}. \tag{5.8}$$

We now apply Lemma 5.4 with $S = I$ and $T : L^{p_0}(w) \rightarrow L^{q_0}(w)$ given by

$$Tg = (C_0 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j})^{-1} \frac{w(2^{j+1}B)^{1/p_0}}{w(B)^{1/q_0}} \chi_B e^{-kr^2 L_w} (g \chi_{C_j(B)}).$$

This yields the \mathbb{H} -valued extension of (5.8): for all $g \in L_{\mathbb{H}}^{p_0}(w)$ with $\text{supp}(g(\cdot, t)) \subset C_j(B)$, $t > 0$, we have

$$\left(\int_B \| \| e^{-kr^2 L_w} g(x, \cdot) \| \|^{q_0} dw \right)^{1/q_0} \leq C_0 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j} \left(\int_{C_j(B)} \| \| g(x, \cdot) \| \|^{p_0} dw \right)^{1/p_0}. \tag{5.9}$$

Given an arbitrary $g \in L_{\mathbb{H}}^{p_0}(w)$, decompose it as

$$g(x, t) = \sum_{j \geq 1} g(x, t) \chi_{C_j(B)}(x) = \sum_{j \geq 1} g_j(x, t).$$

Then inequality (5.9) yields

$$\begin{aligned} \left(\int_B \| \| e^{-kr^2 L_w} g(x, \cdot) \| \|^{q_0} dw \right)^{1/q_0} &\leq \sum_{j \geq 1} \left(\int_B \| \| e^{-kr^2 L_w} g_j(x, \cdot) \| \|^{q_0} dw \right)^{1/q_0} \\ &\lesssim \sum_{j \geq 1} 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j} \left(\int_{2^{j+1}B} \| \| g(x, \cdot) \| \|^{p_0} dw \right)^{1/p_0}. \end{aligned} \tag{5.10}$$

Define $g(x, t) = (tL_w)^{1/2} e^{-tL_w} f(x)$. Then $g_{L_w} f(x) = \| \| g(x, \cdot) \| \|$; by our choice of p_0 and the first step of the proof we have $g \in L_{\mathbb{H}}^{p_0}(w)$. Moreover, since for each $t > 0$ we know that $(tL_w)^{1/2} e^{-tL_w}$ and $e^{-kr^2 L_w}$ commute,

$$g_{L_w} (e^{-kr^2 L_w} f)(x) = \| \| e^{-kr^2 L_w} g(x, \cdot) \| \|.$$

We can now use (4.7) and (5.10) to get

$$\begin{aligned} \left(\int_B |g_{L_w} \mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} &\lesssim \sum_{k=1}^m \left(\int_B \| \| e^{-kr^2 L_w} g(x, \cdot) \| \|^{q_0} dw \right)^{1/q_0} \\ &\lesssim \sum_{j \geq 1} 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j} \left(\int_{2^{j+1}B} |g_{L_w} f|^p dw \right)^{1/p_0}. \end{aligned}$$

This proves (2.37) with $g(j) = C 2^{j(\theta_1 + \theta_2)} e^{-c 4^j}$. Therefore, by Theorem 2.35 we get

$$\| \| g_{L_w} f \| \|_{L^p(v dw)} \lesssim \| \| f \| \|_{L^p(v dw)}.$$

It remains to show the reverse inequalities. We will prove the lower bound in (5.3); then the lower bound in (5.2) holds if we take $v \equiv 1$. Fix $p_- < p < p_+$ and $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/p)'}(w)$. By the duality properties of weights [Auscher and Martell 2007a, Lemma 4.4] and since $p_{\pm}(L_w)' = p_{\mp}(L_w^*)$, where L_w^* is the adjoint (on $L^2(w)$) of L_w ,

$$v^{1-p'} \in A_{p'/p_-(L^*)}(w) \cap \text{RH}_{(p_+(L^*)/p')'}(w). \tag{5.11}$$

We now proceed as in the proof of [Auscher and Martell 2006, Theorem 7.3]. Given $F \in L^p_{\mathbb{H}}(v \, dw) \cap L^2_{\mathbb{H}}(w)$ and $x \in \mathbb{R}^n$, we set

$$T_{L_w} F(x) = \int_0^\infty (t L_w)^{1/2} e^{-tL_w} F(x, t) \frac{dt}{t}. \tag{5.12}$$

Recall that $(t L_w)^{1/2} e^{-tL_w} F(x, t) = (t L_w)^{1/2} e^{-tL_w} (F(\cdot, t))(x)$. Hence, T_{L_w} maps \mathbb{H} -valued functions to \mathbb{C} -valued functions. For $h \in L^{p'}(v^{1-p'} \, dw) \cap L^2(w)$ with $\|h\|_{L^{p'}(v^{1-p'} \, dw)} = 1$, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^n} T_{L_w} F \bar{h} \, dw \right| &= \left| \int_{\mathbb{R}^n} \int_0^\infty F(x, t) \overline{(t L_w^*)^{1/2} e^{-tL_w^*} h(x)} \frac{dt}{t} \, dw(x) \right| \\ &\leq \int_{\mathbb{R}^n} \|F(x, \cdot)\| \|g_{L_w^*} h(x)\| \, dw(x) \\ &\lesssim \|F\|_{L^p_{\mathbb{H}}(v \, dw)} \|g_{L_w^*} h\|_{L^{p'}(v^{1-p'} \, dw)} \lesssim \|F\|_{L^p_{\mathbb{H}}(v \, dw)}, \end{aligned}$$

where the last estimate uses the fact that $g_{L_w^*}$ is bounded on $L^{p'}(v^{1-p'} \, dw)$. This follows from the upper bound in (5.3) (with L_w^* in place of L_w), which we proved above, and (5.11). Taking the supremum over all such functions h and using a standard density argument we have obtained that T_{L_w} is bounded from $L^p_{\mathbb{H}}(v \, dw)$ to $L^p(v \, dw)$.

Next, given $f \in L^p(v \, dw) \cap L^2(dw)$, if we define $F(x, t) = (tL_w)^{1/2} e^{-tL_w} f(x)$, then $F \in L^p_{\mathbb{H}}(v \, dw) \cap L^2_{\mathbb{H}}(w)$ since $\|F\|_{L^p_{\mathbb{H}}(v \, dw)} = \|g_{L_w} f\|_{L^p(v \, dw)}$ and analogously for $L^2(w)$. Also, by the $L^2(w)$ functional calculus we have

$$f(x) = 2 \int_0^\infty (tL_w)^{1/2} e^{-tL_w} F(x, t) \frac{dt}{t} = 2T_{L_w} F(x). \tag{5.13}$$

Therefore,

$$\|f\|_{L^p(v \, dw)} = 2\|T_{L_w} F\|_{L^p(v \, dw)} \lesssim \|F\|_{L^p_{\mathbb{H}}(v \, dw)} = \|g_{L_w} f\|_{L^p(v \, dw)},$$

and this completes the proof of (5.3).

To finish the proof of Proposition 5.1 we need to show that the equivalence of norms in (5.2) implies that the semigroup is uniformly bounded. However, this follows immediately from the definition of g_{L_w} and the semigroup property: for any $s > 0$,

$$g_{L_w}(e^{-sL_w} f)(x) = \left(\int_0^\infty |L_w^{1/2} e^{-(s+t)L_w} f(x)|^2 \, dt \right)^{1/2} \leq g_{L_w} f(x). \quad \square$$

We conclude this section by proving a version of Proposition 5.1 for the ‘‘adjoint’’ of a discrete square function. We will need this estimate in the proof of Proposition 6.1 below.

Proposition 5.14. *Define the holomorphic function ψ on the sector $\Sigma_{\pi/2}$ by*

$$\psi(z) = \frac{1}{\sqrt{\pi}} \int_1^\infty z e^{-tz} \frac{dt}{\sqrt{t}}. \tag{5.15}$$

If $p_-(L_w) < p < p_+(L_w)$, then for any sequence of functions $\{\beta_k\}_{k \in \mathbb{Z}}$,

$$\left\| \sum_{k \in \mathbb{Z}} \psi(4^k L_w) \beta_k \right\|_{L^p(w)} \lesssim \left\| \left(\sum_{k \in \mathbb{Z}} |\beta_k|^2 \right)^{1/2} \right\|_{L^p(w)}. \tag{5.16}$$

Proof. By duality and since $p_{\pm}(L_w)' = p_{\mp}(L_w^*)$, it will suffice to show that for every $p_-(L_w^*) < p < p_+(L_w^*)$,

$$\left\| \left(\sum_{k \in \mathbb{Z}} |\bar{\psi}(4^k L_w^*) h|^2 \right)^{1/2} \right\|_{L^p(w)} \lesssim \|h\|_{L^p(w)}. \tag{5.17}$$

The function ψ satisfies $|\psi(z)| \leq C|z|^{1/2}e^{-c|z|}$ uniformly on subsectors Σ_{μ} , $0 \leq \mu < \frac{\pi}{2}$. Thus the operator on the left-hand side of (5.17) is a discrete analog of the square function $g_{L_w^*}$, changing continuous times t to discrete times 4^k and $z^{1/2}e^{-z}$ to $\bar{\psi}(z)$. Since $\bar{\psi}(z)$ has the same quantitative properties as $z^{1/2}e^{-z}$ (decay at 0 and at infinity), we can repeat the previous argument and obtain the desired estimates as in the proof of Proposition 5.1. \square

Remark 5.18. In Proposition 5.14 we can also get $L^p(v dw)$ estimates, but in the proof of Proposition 6.1 below we will only need the unweighted estimates. Further details and the precise statements are left to the interested reader.

6. Reverse inequalities

In this section we will prove $L^p(w)$ estimates of the form $\|L_w^{1/2} f\|_{L^p(w)} \leq C \|\nabla f\|_{L^p(w)}$, which generalize the $L^2(w)$ Kato estimates in Theorem 2.18. These are referred to as reverse inequalities since if we replace f by $L_w^{-1/2} f$, then formally we get a reverse-type inequality for the Riesz transform: $\|f\|_{L^p(w)} \leq C \|\nabla L_w^{-1/2} f\|_{L^p(w)}$.

Since these estimates involve the gradient, in proving them we will rely (implicitly and explicitly) on the weighted Poincaré inequality (2.3). This will require an additional assumption on p when $p < 2$. To state it simply, define

$$(p_-(L_w))_{w,*} = \frac{nr_w p_-(L_w)}{nr_w + p_-(L_w)} < p_-(L_w).$$

Proposition 6.1. *Let $\max\{r_w, (p_-(L_w))_{w,*}\} < p < p_+(L_w)$. Then for all $f \in \mathcal{S}$,*

$$\|L_w^{1/2} f\|_{L^p(w)} \leq C \|\nabla f\|_{L^p(w)}, \tag{6.2}$$

with C independent of f . Furthermore, if

$$\max\{r_w, p_-(L_w)\} < p < p_+(L_w) \quad \text{and} \quad v \in A_{p/\max\{r_w, p_-(L_w)\}}(w) \cap \text{RH}_{(p_+(L_w)/p)'(w)},$$

then for all $f \in \mathcal{S}$,

$$\|L_w^{1/2} f\|_{L^p(v dw)} \leq C \|\nabla f\|_{L^p(v dw)}. \tag{6.3}$$

Remark 6.4. The quantity $\max\{r_w, (p_-(L_w))_{w,*}\}$ can be equal to either term. For instance, it equals r_w if $p_-(L_w) \leq n'r_w$. From Proposition 3.1 we know that $p_-(L_w) < (2_w^*)' = 2nr_w/(nr_w + 2)$, but this only implies the previous inequality for some values of n and r_w .

Proof. As before, let $p_- = p_-(L_w)$ and $p_+ = p_+(L_w)$. Fix p , $\max\{r_w, (p_-)_{w,*}\} < p < 2$, and $f \in \mathcal{S}$. We will first show

$$\|L_w^{1/2} f\|_{L^{p,\infty}(w)} \lesssim \|\nabla f\|_{L^p(w)}. \tag{6.5}$$

First note that since $p > r_w$, we have $w \in A_p$. Therefore, given $\alpha > 0$ we can form the Calderón–Zygmund decomposition given in [Auscher and Martell 2006, Lemma 6.6]. There exist a collection of balls $\{B_i\}_i$, smooth functions $\{b_i\}_i$ and a function $g \in L^1_{\text{loc}}(w)$ such that

$$f = g + \sum_i b_i \tag{6.6}$$

and the following properties hold:

$$|\nabla g(x)| \leq C\alpha \quad \text{for } w\text{-a.e. } x, \tag{6.7}$$

$$\text{supp}(b_i) \subset B_i \quad \text{and} \quad \int_{B_i} |\nabla b_i|^p dw \leq C\alpha^p w(B_i), \tag{6.8}$$

$$\sum_i w(B_i) \leq \frac{C}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw, \tag{6.9}$$

$$\sum_i \chi_{B_i} \leq N, \tag{6.10}$$

$$\left(\int_{B_i} |b_i|^q dw \right)^{1/q} \lesssim C\alpha r(B_i) \quad \text{for } 1 \leq q \leq p_w^*, \tag{6.11}$$

where C and N depend only on n , p , q and the doubling constant of w .

To prove (6.5) we will prove the corresponding weak-type estimates with f replaced by g and b_i . For g , we use the $L^2(w)$ Kato estimate (2.19), (6.7), and the fact that $p < 2$ to get

$$\begin{aligned} w\left(\left\{|L_w^{1/2} g| > \frac{\alpha}{3}\right\}\right) &\lesssim \frac{1}{\alpha^2} \int_{\mathbb{R}^n} |L_w^{1/2} g|^2 dw \\ &\lesssim \frac{1}{\alpha^2} \int_{\mathbb{R}^n} |\nabla g|^2 dw \\ &\lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla g|^p dw \\ &\lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw + \frac{1}{\alpha^p} \int_{\mathbb{R}^n} \left| \sum_i \nabla b_i \right|^p dw \\ &\lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw, \end{aligned}$$

where the last estimate follows from (6.10), (6.8), and (6.9).

To prove a weak-type estimate for $L_w^{1/2}(\sum_i b_i)$, let $r_i = 2^k$ if $2^k \leq r(B_i) < 2^{k+1}$. Then for all i , $r_i \sim r(B_i)$. Write

$$L_w^{1/2} = \frac{1}{\sqrt{\pi}} \int_0^{r_i^2} L_w e^{-tL_w} \frac{dt}{\sqrt{t}} + \frac{1}{\sqrt{\pi}} \int_{r_i^2}^\infty L_w e^{-tL_w} \frac{dt}{\sqrt{t}} = T_i + U_i;$$

then we have

$$\begin{aligned} & w\left(\left\{\left|\sum_i L_w^{1/2} b_i\right| > \frac{2\alpha}{3}\right\}\right) \\ & \leq w\left(\bigcup_i 4B_i\right) + w\left(\left\{\left|\sum_i U_i b_i\right| > \frac{\alpha}{3}\right\}\right) + w\left(\left(\mathbb{R}^n \setminus \bigcup_i 4B_i\right) \cap \left\{\left|\sum_i T_i b_i\right| > \frac{\alpha}{3}\right\}\right) \\ & \lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw + I_1 + I_2, \end{aligned}$$

where the last inequality follows from (6.9).

We first estimate I_2 . Since $p > (p-)_{w,*}$ we have $p_w^* > ((p-)_{w,*})_w^* = p-$, and we can choose $q \in \mathcal{J}(L_w)$ such that (6.11) is satisfied. By Corollary 3.4, $tL_w e^{-tL_w} \in \mathcal{O}(L^q(w) \rightarrow L^q(w))$, and so

$$\begin{aligned} I_2 & \lesssim \frac{1}{\alpha} \sum_i \sum_{j \geq 2} \int_{C_j(B_i)} |T_i b_i| dw \\ & \lesssim \frac{1}{\alpha} \sum_i \sum_{j \geq 2} w(2^j B_i) \int_0^{r_i^2} \int_{C_j(B_i)} |tL_w e^{-tL_w} b_i| dw \frac{dt}{t^{3/2}} \\ & \lesssim \frac{1}{\alpha} \sum_i \sum_{j \geq 2} 2^{jD} w(B_i) \int_0^{r_i^2} 2^{j\theta_1} \Upsilon\left(2^j r_i / \sqrt{t}\right)^{\theta_2} e^{-c4^j r_i^2/t} \frac{dt}{t^{3/2}} \left(\int_{B_i} |b_i|^q dw\right)^{1/q} \\ & \lesssim \sum_i \sum_{j \geq 2} 2^{jD} e^{-c4^j} w(B_i) \\ & \lesssim \sum_i w(B_i) \lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw, \end{aligned}$$

where we have used (6.11) and (6.9), and D is the doubling order of dw .

We will now estimate I_1 . For q as above, by Proposition 4.3 we have an $L^q(w)$ functional calculus for L_w . Therefore, we can write U_i as $r_i^{-1} \psi(r_i^2 L_w)$ with ψ defined by (5.15). Let $\beta_k = \sum_{i: r_i=2^k} b_i / r_i$; then,

$$\sum_i U_i b_i = \sum_{k \in \mathbb{Z}} \psi(4^k L_w) \left(\sum_{i: r_i=2^k} \frac{b_i}{r_i}\right) = \sum_{k \in \mathbb{Z}} \psi(4^k L_w) \beta_k.$$

Therefore, by Proposition 5.14, (6.10), (6.11), the fact that $r_i \sim r(B_i)$ and (6.9), we have

$$\begin{aligned} I_1 & \lesssim \frac{1}{\alpha^q} \left\| \sum_i U_i b_i \right\|_{L^q(w)}^q \lesssim \frac{1}{\alpha^q} \left\| \left(\sum_{k \in \mathbb{Z}} |\beta_k|^2\right)^{1/2} \right\|_{L^q(w)}^q \\ & \lesssim \frac{1}{\alpha^q} \int_{\mathbb{R}^n} \sum_i \frac{|b_i|^q}{r_i^q} dw \lesssim \sum_i w(B_i) \lesssim \frac{1}{\alpha^p} \int_{\mathbb{R}^n} |\nabla f|^p dw. \end{aligned}$$

If we combine all of the estimates we have obtained, we get (6.5) as desired.

To prove (6.2) from the weak-type estimate (6.5) we will use an interpolation argument from [Auscher and Martell 2006]. Fix p and r such that $\max\{r_w, (p-)_{w,*}\} < r < p < 2$. Then by (6.5) and (2.19) we

have that for every $f \in \mathcal{S}$,

$$\|L_w^{1/2} f\|_{L^{r,\infty}(w)} \lesssim \|\nabla f\|_{L^r(w)}, \quad \|L_w^{1/2} f\|_{L^2(w)} \lesssim \|\nabla f\|_{L^2(w)}. \quad (6.12)$$

Formally, to apply Marcinkiewicz interpolation, we let $g = \nabla f$ to get a weak (r, r) and strong $(2, 2)$ inequality; this would immediately yield a strong (p, p) inequality. To formalize this we must justify this substitution.

For every $q > r_w$, by [Auscher and Martell 2006, Lemma 6.7] we have that

$$\mathcal{E} = \{(-\Delta)^{1/2} f : f \in \mathcal{S}, \text{supp } \hat{f} \subset \mathbb{R}^n \setminus \{0\}\}$$

is dense in $L^q(w)$, where \hat{f} denotes the Fourier transform of f . Moreover, since $r > r_w$, we have $w \in A_r$ and the Riesz transforms, $R_j = \partial_j (-\Delta)^{-1/2}$, are bounded on $L^r(w)$ [García-Cuerva and Rubio de Francia 1985]. It follows from this and the identity $-I = R_1^2 + \dots + R_n^2$ that for $g \in L^r(w)$,

$$\|g\|_{L^r(w)} \sim \|\nabla (-\Delta)^{-1/2} g\|_{L^r(w)}.$$

Thus, for $g \in \mathcal{E}$, we know $L_w^{1/2} (-\Delta)^{-1/2} g = L_w^{1/2} f$ if $f = (-\Delta)^{-1/2} g$ and $\|\nabla f\|_{L^r(w)} \sim \|g\|_{L^r(w)}$ for $r > r_w$. Thus (6.12) becomes weighted weak (r, r) and strong $(2, 2)$ inequalities for $T = L_w^{1/2} (-\Delta)^{-1/2}$, and this operator is defined a priori on \mathcal{E} . Since \mathcal{E} is dense in each $L^q(w)$, we can extend T by density in both cases and their restrictions to the space of simple functions agree. Hence, we can apply Marcinkiewicz interpolation and conclude, again by density, that (6.2) holds for all p with $r < p < 2$. Since r is arbitrary, we get (6.2) in the range $\max\{r_w, (p_-)_{w,*}\} < p < 2$.

For the second step of the proof we will prove (6.3) using Theorem 2.35. Inequality (6.2) for its full range of exponents then follows by letting $v = 1$. Define $\tilde{p}_- = \max\{r_w, p_-\} < 2$, and fix $\tilde{p}_- < p < p_+$ and $v \in A_{p/\tilde{p}_-}(w) \cap \text{RH}_{(p_+/p)'}(w)$. By the openness properties of A_q and RH_s weights, there exist p_0, q_0 such that

$$\tilde{p}_- < p_0 < \min\{p, 2\} \leq p < q_0 < p_+, \quad v \in A_{p/p_0}(w) \cap \text{RH}_{(q_0/p)'}(w).$$

To apply Theorem 2.35, let $T = L_w^{1/2}$, $S = \nabla$, and $\mathcal{A}_r = I - (I - e^{-r^2 L_w})^m$, where the value of m will be fixed below. We will first show that (2.37) holds. By (4.8) we have $\mathcal{A}_r \in \mathcal{O}(L^{p_0}(w) \rightarrow L^{q_0}(w))$ since $p_0, q_0 \in \mathcal{J}(L_w)$. Let $h = L_w^{1/2} f$ and decompose h as we decomposed f in (4.11). Then, since $L_w^{1/2}$ and \mathcal{A}_r commute, it follows that

$$\begin{aligned} \left(\int_B |L_w^{1/2} \mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} &\lesssim \sum_{j \geq 1} \left(\int_B |\mathcal{A}_r h_j|^{q_0} dw \right)^{1/q_0} \\ &\lesssim \sum_{j \geq 1} 2^{j\theta_1} \Upsilon(2^j)^{\theta_2} e^{-c4^j} \left(\int_{C_j} |h|^{p_0} dw \right)^{1/p_0} \\ &\leq \sum_{j \geq 1} 2^{j(\theta_1 + \theta_2)} e^{-c4^j} \left(\int_{2^{j+1}B} |L_w^{1/2} f|^{p_0} dw \right)^{1/p_0}. \end{aligned}$$

This gives us (2.37) with $g(j) = C 2^{j(\theta_1 + \theta_2)} e^{-c4^j}$; clearly, $\sum g(j) < \infty$.

We now prove that (2.36) holds. Fix $f \in \mathcal{S}$ and let $\varphi(z) = z^{1/2}(1 - e^{-r^2z})^m$ so that

$$\varphi(L_w)f = L_w^{1/2}(I - e^{-r^2L_w})^m f.$$

By the conservation property [Cruz-Uribe and Rios 2015; Auscher 2007, Section 2.5],

$$\varphi(L_w)f = \varphi(L_w)(f - f_{4B,w}) = \sum_{j \geq 1} \varphi(L_w)h_j, \quad (6.13)$$

where $h_j = (f - f_{4B,w})\phi_j$, $\phi_j = \chi_{C_j(B)}$ for $j \geq 3$, ϕ_1 is a smooth function with support in $4B$, $0 \leq \phi_1 \leq 1$, $\phi_1 = 1$ in $2B$ and $\|\nabla\phi_1\|_\infty \leq C/r$, and ϕ_2 is chosen so that $\sum_{j \geq 1} \phi_j = 1$.

We estimate each term in the right-hand side of (6.13) separately. When $j = 1$, since $p_- < p_0 < p_+$, by the bounded holomorphic functional calculus on $L^{p_0}(w)$ (Proposition 4.3) and the fact that $\varphi(L_w)h_1 = (I - e^{-r^2L_w})^m L_w^{1/2}h_1$, we have

$$\|\varphi(L_w)h_1\|_{L^{p_0}(w)} \lesssim \|L_w^{1/2}h_1\|_{L^{p_0}(w)}$$

uniformly in r . By the above argument we have that (6.2) holds for $p = p_0$ since $\tilde{p}_- < p_0 < 2$. Further, since $f \in \mathcal{S}$, we have $h_1 \in \mathcal{S}$ by our choice of ϕ_1 . This, together with the $L^{p_0}(w)$ -Poincaré inequality (2.3) (since $p_0 > r_w$, $w \in A_{p_0}$) and the definition of h_1 yield

$$\begin{aligned} \|L_w^{1/2}h_1\|_{L^{p_0}(w)} &\lesssim \|\nabla h_1\|_{L^{p_0}(w)} \\ &\lesssim \|(\nabla f)\chi_{4B}\|_{L^{p_0}(w)} + r^{-1}\|(f - f_{4B,w})\chi_{4B}\|_{L^{p_0}(w)} \lesssim \|(\nabla f)\chi_{4B}\|_{L^{p_0}(w)}. \end{aligned}$$

Therefore,

$$\left(\int_B |\varphi(L_w)h_1|^{p_0} dw \right)^{1/p_0} \lesssim \left(\int_{4B} |\nabla f|^{p_0} dw \right)^{1/p_0}.$$

When $j \geq 3$, the functions η associated with φ by (2.11) satisfy

$$|\eta(z)| \lesssim \frac{r^{2m}}{|z|^{m+3/2}}, \quad z \in \Gamma_{\pi/2-\theta}.$$

Since $p_0 \in \mathcal{J}(L_w)$, by Corollary 3.4, $e^{-zL_w} \in \mathcal{O}(L^{p_0}(w) \rightarrow L^{p_0}(w), \Sigma_\mu)$. This, together with the representation (2.10), gives us that

$$\begin{aligned} \left(\int_B |\varphi(L_w)h_j|^{p_0} dw \right)^{1/p_0} &\leq \int_{\Gamma_{\pi/2-\theta}} \left(\int_B |e^{-zL}h_j|^{p_0} dw \right)^{1/p_0} |\eta(z)||dz| \\ &\lesssim 2^{j\theta_1} \int_{\Gamma_{\pi/2-\theta}} \Upsilon\left(\frac{2^j r}{\sqrt{|z|}}\right)^{\theta_2} e^{-\alpha 4^j r^2/|z|} \frac{r^{2m}}{|z|^{m+3/2}} |dz| \left(\int_{C_j(B)} |h_j|^{p_0} dw \right)^{1/p_0} \\ &\lesssim 2^{j(\theta_1-2m-1)} r^{-1} \left(\int_{2^{j+1}B} |f - f_{4B,w}|^{p_0} dw \right)^{1/p_0} \\ &\lesssim 2^{j(\theta_1-2m-1)} \sum_{l=1}^j 2^l \left(\int_{2^{l+1}B} |\nabla f|^{p_0} dx \right)^{1/p_0}, \end{aligned}$$

provided $2m + 1 > \theta_2$. The last estimate follows from the $L^{p_0}(w)$ -Poincaré inequality (2.3) (here we again use that $p_0 > r_w$ and so $w \in A_{p_0}$):

$$\begin{aligned} \left(\int_{2^{j+1}B} |f - f_{4B,w}|^{p_0} dw \right)^{1/p_0} &\leq \left(\int_{2^{j+1}B} |f - f_{2^{j+1}B,w}|^{p_0} dw \right)^{1/p_0} + \sum_{l=2}^j |f_{2^l B,w} - f_{2^{l+1}B,w}| \\ &\lesssim \sum_{l=1}^j \left(\int_{2^{l+1}B} |f - f_{2^{l+1}B}|^{p_0} dx \right)^{1/p_0} \\ &\lesssim r \sum_{l=1}^j 2^l \left(\int_{2^{l+1}B} |\nabla f|^{p_0} dx \right)^{1/p_0}. \end{aligned} \tag{6.14}$$

When $j = 2$ we can argue similarly, using the fact that

$$|h_2| \leq |f - f_{4B,w}| \chi_{8B \setminus 2B} \leq |f - f_{2B,w}| \chi_{8B \setminus 2B} + |f_{4B,w} - f_{2B,w}| \chi_{8B \setminus 2B}.$$

If we combine these estimates, then by (6.13) and Minkowski’s inequality we get

$$\left(\int_B |\varphi(L_w)h|^{p_0} dw \right)^{1/p_0} \lesssim \sum_{j \geq 1} \left(\int_B |\varphi(L_w)h_j|^{p_0} dw \right)^{1/p_0} \leq \sum_{j \geq 1} g(j) \left(\int_B |\nabla f|^{p_0} dw \right)^{1/p_0}$$

with $g(j) = C_m 2^{j(\theta_1 - 2m)}$ provided $2m + 1 > \theta_2$. If we further assume that $2m > \theta_1$, then $\sum_j g(j) < \infty$. This proves that (2.36) holds. Therefore, by Theorem 2.35 we get (6.3) as desired. \square

7. The gradient of the semigroup $\sqrt{t} \nabla e^{-tL_w}$

Let $\tilde{\mathcal{K}}(L_w) \subset [1, \infty]$ be the set of all exponents p such that $\sqrt{t} \nabla e^{-tL_w} : L^p(w) \rightarrow L^p(w)$ is uniformly bounded for all $t > 0$. By Theorem 2.15 and Lemma 2.34, $2 \in \tilde{\mathcal{K}}(L_w)$ and if it contains more than one point, then by interpolation $\tilde{\mathcal{K}}(L_w)$ is an interval. In this section we give a partial description of the set of (p, q) such that $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$.

Proposition 7.1. *There exists an interval $\mathcal{K}(L_w)$ such that if $p, q \in \mathcal{K}(L_w)$, $p \leq q$, then $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$. Moreover, $\mathcal{K}(L_w)$ has the following properties:*

- (1) $\mathcal{K}(L_w) \subset \tilde{\mathcal{K}}(L_w)$.
- (2) If $q_-(L_w)$ and $q_+(L_w)$ are the left and right endpoints of $\mathcal{K}(L_w)$, then $q_-(L_w) = p_-(L_w)$, $2 \leq q_+(L_w) \leq (q_+(L_w))_w^* \leq p_+(L_w)$. In particular, $2 \in \mathcal{K}(L_w)$ and $\mathcal{K}(L_w) \subset \mathcal{J}(L_w)$.
- (3) If $q \geq 2$ and $p < q$, and if $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, then $p, q \in \mathcal{K}(L_w)$.
- (4) $\sup \tilde{\mathcal{K}}(L_w) = q_+(L_w)$.

Remark 7.2. Unlike in the unweighted case [Auscher and Martell 2007b], we are unable to give a complete characterization of $\mathcal{K}(L_w)$. More precisely, if we have an off-diagonal estimate and $p < q < 2$, then we cannot prove that $p, q \in \mathcal{K}(L_w)$.

Remark 7.3. In Section 8 below we will show that $q_+(L_w) > 2$; in particular, this gives that $2 \in \text{Int } \mathcal{K}(L_w)$.

As an immediate consequence of Proposition 7.1 we get weighted inequalities for the gradient of the semigroup. The proof is identical to the proofs of Corollaries 3.3 and 3.4.

Corollary 7.4. *Let $q_-(L_w) < p \leq q < q_+(L_w)$. If $v \in A_{p/q_-(L_w)}(w) \cap \text{RH}_{(q_+(L_w)/q)'(w)}$, then $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(v \, dw) \rightarrow L^q(v \, dw))$ and $\sqrt{z} \nabla e^{-zL_w} \in \mathcal{O}(L^p(v \, dw) \rightarrow L^q(v \, dw), \Sigma_v)$ for all v , $0 < v < \frac{\pi}{2} - \vartheta$.*

The proof of Proposition 7.1 requires two lemmas.

Lemma 7.5. *Given $w \in A_\infty$ and a family of sublinear operators $\{T_t\}_{t>0}$ such that $T_t \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, with $1 \leq p < q \leq \infty$, there exist $\alpha, \beta > 0$ such that for any ball B with radius r and for any $t > 0$,*

$$\left(\int_B |T_t(\chi_B f)|^q \, dw \right)^{1/q} \lesssim \max \left\{ \left(\frac{r}{\sqrt{t}} \right)^\alpha, \left(\frac{r}{\sqrt{t}} \right)^\beta \right\} \left(\int_B |f|^p \, dw \right)^{1/p}. \tag{7.6}$$

Proof. This result is implicit in [Auscher and Martell 2007b, Proof of Proposition 2.4, p. 306]; here we reprove it with a small improvement in the constant. There it was shown that in Definition 2.23 it is sufficient to consider the case where $r \approx \sqrt{t}$. But in this case we get that $\Upsilon(r/\sqrt{t}) \approx 1$ and for all $j \geq 2$, $\Upsilon(2^j r/\sqrt{t}) \approx 2^j$. The argument in [loc. cit., p. 306] shows that if we assume that (2.24)–(2.26) hold when $r \approx \sqrt{t}$, then (2.24) holds in general with constant $\max\{1, (r/\sqrt{t})^\alpha\}$ for some $\alpha > 0$ depending on p, q and w . In this maximum the 1 occurs when $r \leq \sqrt{t}$; therefore, to prove (7.6) we need to show that if $r \leq \sqrt{t}$, then we can replace 1 by the better constant $(r/\sqrt{t})^\beta$ for some $\beta > 0$.

Fix $r \leq \sqrt{t}$. If $B = B(x, r)$, then $B \subset B_t = B(x, \sqrt{t})$. As in [loc. cit., p. 306] we apply (2.24) to T_t and B_t ; this yields

$$\begin{aligned} \left(\int_B |T_t(\chi_B f)|^q \, dw \right)^{1/q} &\leq \left(\frac{w(B_t)}{w(B)} \right)^{1/q} \left(\int_{B_t} |T_t(\chi_B f)|^q \, dw \right)^{1/q} \\ &\lesssim \left(\frac{w(B_t)}{w(B)} \right)^{1/q} \left(\int_{B_t} |\chi_B f|^p \, dw \right)^{1/p} \\ &\leq \left(\frac{w(B)}{w(B_t)} \right)^{1/p-1/q} \left(\int_B |f|^p \, dw \right)^{1/p}. \end{aligned}$$

Since $w \in A_\infty$, we have that for some $\theta > 0$,

$$\frac{w(B)}{w(B_t)} \lesssim \left(\frac{|B|}{|B_t|} \right)^\theta = \left(\frac{r}{\sqrt{t}} \right)^{\theta n}.$$

Since $p < q$ we have

$$\left(\int_B |T_t(\chi_B f)|^q \, dw \right)^{1/q} \lesssim \left(\frac{r}{\sqrt{t}} \right)^{(1/p-1/q)\theta n} \left(\int_B |f|^p \, dw \right)^{1/p}.$$

Therefore, if we combine this with the argument from [loc. cit., p. 306] described above, we get that (7.6) holds with $\beta = (1/p - 1/q) \theta n$. □

The second lemma gives the close connection between off-diagonal estimates for e^{-tL_w} and $\sqrt{t} \nabla e^{-tL_w}$ for $p < 2$.

Lemma 7.7. *Given $1 \leq p < 2$ the following are equivalent:*

- (1) $e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$.
- (2) $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$.
- (3) $tL_w e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$.

Proof. We follow the proof of [Auscher and Martell 2007b, Lemma 5.3]. To prove that (1) implies (2), note that by Theorem 2.15, $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$. If we compose this with (1), by Lemma 2.30 and the semigroup property, we get (2).

To prove that (2) implies (3), define $S_t \vec{f} = \sqrt{t} e^{-tL_w} (w^{-1} \operatorname{div}(A \vec{f}))$. By duality, we have

$$\begin{aligned} \langle S_t \vec{f}, g \rangle_{L^2(w)} &= \langle w^{-1} \operatorname{div}(A \vec{f}), \sqrt{t} e^{-tL_w^*} g \rangle_{L^2(w)} = \langle \operatorname{div}(A \vec{f}), \sqrt{t} e^{-tL_w^*} g \rangle_{L^2} \\ &= -\langle \vec{f}, A^* \sqrt{t} \nabla e^{-tL_w^*} g \rangle_{L^2} = \langle \vec{f}, w^{-1} A^* \sqrt{t} \nabla e^{-tL_w^*} g \rangle_{L^2(w)}. \end{aligned}$$

The matrix $w^{-1} A^*$ is uniformly elliptic, and so multiplication by it is bounded on $L^2(w)$. Furthermore, $\sqrt{t} \nabla e^{-tL_w^*} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$. Therefore, it follows that $S_t \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$. If we combine this with (2), we get that $-tL_w e^{-2tL_w} = S_t \circ \sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$. This proves (3).

Finally we show that (3) implies (1). We first prove (2.24). Fix B and f, g such that $(\int_B |f|^p dw)^{1/p} = (\int_B |g|^2 dw)^{1/2} = 1$, and assume also that $f \in L^2(B, dw)$. Define

$$h(t) = \int_B e^{-tL_w} (\chi_B f)(x) g(x) dw(x).$$

By duality it will suffice to show that $|h(t)| \lesssim \Upsilon(r/\sqrt{t})^\theta$. (Note that our assumption implies that $th'(t)$ satisfies such a bound.) First, we claim that

$$\lim_{t \rightarrow \infty} h(t) = 0.$$

To see this we use the fact (discussed in Section 2) that L_w has a bounded holomorphic functional calculus on $L^2(w)$. Given this, since $z \mapsto e^{-tz}$ converges to 0 uniformly on compact subsets of $\operatorname{Re} z > 0$, we get the desired limit.

Hence, we can write $h(t) = -\int_t^\infty h'(s) ds$. Notice that $|th'(t)| \lesssim \Upsilon(r/\sqrt{t})^{\theta_2}$ but this does not give a convergent integral. However, if we apply Lemma 7.5 to $tL_w e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$, we get that $|th'(t)| \lesssim \tilde{\Upsilon}(r/\sqrt{t})$ with $\tilde{\Upsilon}(s) = \max\{s^\alpha, s^\beta\}$. It follows from this estimate that

$$|h(t)| \leq \int_t^\infty |h'(s)| ds \lesssim \int_t^\infty \tilde{\Upsilon}\left(\frac{r}{\sqrt{s}}\right) \frac{ds}{s} \approx \int_0^{r/\sqrt{t}} \tilde{\Upsilon}(s) \frac{ds}{s} \lesssim \tilde{\Upsilon}\left(\frac{r}{\sqrt{t}}\right) \lesssim \Upsilon\left(\frac{r}{\sqrt{t}}\right)^{\alpha+\beta}.$$

To prove (2.25) we argue as before, but with $(\int_{C_j(B)} |f|^p dw)^{1/p} = (\int_B |g|^2 dw)^{1/2} = 1$ and

$$h(t) = \int_B e^{-tL_w} (\chi_{C_j(B)} f)(x) g(x) dw(x).$$

Since $d(B, C_j(B)) > 0$, by Theorem 2.15 and Hölder’s inequality, $h(t) \rightarrow 0$ as $t \rightarrow 0$. Therefore, $h(t) = \int_0^t h'(s) ds$. Since $tL_w e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$, we have

$$\begin{aligned} h(t) &\leq \int_0^t |h'(s)| ds \lesssim 2^{j\theta_1} \int_0^t \Upsilon\left(\frac{2^j r}{\sqrt{s}}\right)^{\theta_2} e^{-c4^j r^2/s} \frac{ds}{s} \\ &\approx 2^{j\theta_1} \int_{2^j r/\sqrt{t}}^\infty \Upsilon(s)^{\theta_2} e^{-cs^2} \frac{ds}{s} \lesssim 2^{j\theta_1} \Upsilon\left(\frac{2^j r}{\sqrt{t}}\right)^{\theta_2} e^{-c4^j r^2/t}. \end{aligned}$$

This is (2.25).

Finally, the proof of (2.26) is essentially the same and we omit the details. This completes the proof that (3) implies (1). \square

Proof of Proposition 7.1. Define the sets $\mathcal{K}_-(L_w)$ and $\mathcal{K}_+(L_w)$ to be

$$\begin{aligned} \mathcal{K}_-(L_w) &= \{p \in [1, 2] : \sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))\}, \\ \mathcal{K}_+(L_w) &= \{p \in [2, \infty] : \sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^p(w))\}, \end{aligned}$$

and let $\mathcal{K}(L_w) = \mathcal{K}_-(L_w) \cup \mathcal{K}_+(L_w)$. The set is nonempty, since $2 \in \mathcal{K}(L_w)$. By Lemma 2.28 it is an interval. Now fix $p, q \in \mathcal{K}(L_w)$ with $p < q$. If $p < q \leq 2$ or $2 \leq p < q$, then by Lemma 2.28, $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$ since $p, q \in \mathcal{K}_-(L_w)$ or $p, q \in \mathcal{K}_+(L_w)$. If $p \leq 2 < q$, then $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$ and by Lemma 7.7, $e^{-tL} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$. Hence, by Lemma 2.30 and the semigroup property, $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$. Thus, in every case we get the desired off-diagonal estimate.

We now prove (1)–(4). By Lemma 2.30, off-diagonal estimates on balls imply uniform boundedness, and so $\mathcal{K}(L_w) \subset \tilde{\mathcal{K}}(L_w)$. This proves (1).

To prove (2), we first note that if $p < 2$, then by Lemma 7.7, $p \in \mathcal{J}(L_w)$ if and only if $p \in \mathcal{K}_-(L_w)$. Thus $\mathcal{J}(L_w) \cap [1, 2] = \mathcal{K}_-(L_w)$ and so $q_-(L_w) = p_-(L_w)$. To show that $(q_+(L_w))_w^* \leq p_+(L_w)$, first note that if $q_+(L_w) = 2$, then by Proposition 3.1 we have $(q_+(L_w))_w^* = 2_w^* \leq p_+(L_w)$. If $q_+(L_w) > 2$, then we proceed as in the proof of this proposition. Let $2 < p < q_+(L_w)$ and $p < q < p_w^*$. Then by (2.3), and the facts that $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$ and $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^p(w))$, we get

$$\begin{aligned} \left(\int_B |e^{-tL_w}(\chi_B f)|^q dw\right)^{1/q} &\lesssim \left(\int_B |e^{-tL_w}(\chi_B f)|^2 dw\right)^{1/2} + r \left(\int_B |\nabla e^{-tL_w}(\chi_B f)|^p dw\right)^{1/p} \\ &\lesssim \Upsilon\left(\frac{r}{\sqrt{t}}\right)^{1+\theta_2} \left(\int_B |f|^2 dw\right)^{1/2}. \end{aligned}$$

This gives us inequality (2.24). The other two inequalities in Definition 2.23 can be proved in exactly the same way. Thus $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$, which implies $q \leq p_+(L_w)$. Letting $p \nearrow q_+(L_w)$ and $q \nearrow p_w^*$, we conclude that $(q_+(L_w))_w^* \leq p_+(L_w)$.

The last estimate implies in particular that $q_+(L_w) \leq p_+(L_w)$. If $q_+(L_w) < \infty$, we clearly have $q_+(L_w) < p_+(L_w)$ and so $\mathcal{K}_+(L_w) \subset \mathcal{J}(L_w)$. Otherwise, $p_+(L) = \infty$ and again we have $\mathcal{K}_+(L_w) \subset \mathcal{J}(L_w)$. This completes the proof of (2).

To prove (3), suppose first that $2 \leq p < q$ and $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$. We will show that $p, q \in \mathcal{K}(L_w)$. Since we also have $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$, by interpolation (Lemma 2.27), $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^{p_\theta}(w) \rightarrow L^{q_\theta}(w))$, where $1/p_\theta = (1 - \theta)/p + \theta/2$, $1/q_\theta = (1 - \theta)/q + \theta/2$ and $\theta \in (0, 1)$. If $p \notin \mathcal{K}_+(L_w)$, then $q > \sup \mathcal{K}_+(L_w)$. We can choose θ such that $p_\theta < \sup \mathcal{K}_+(L_w) < q_\theta$. Since $\mathcal{K}_+(L_w) \subset \mathcal{J}(L_w)$, we have $p_\theta \in \mathcal{J}(L_w)$; i.e., $e^{-tL} \in \mathcal{O}(L^2(w) \rightarrow L^{p_\theta}(w))$. By composition and the semigroup property, $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^{q_\theta}(w))$; hence, $q_\theta \in \mathcal{K}_+(L_w)$, a contradiction. Therefore, $p \in \mathcal{K}_+(L_w)$. As we have $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$ by assumption and $e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^p(w))$ since $p \in \mathcal{J}(L_w)$, by composition and the semigroup property, $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$. Hence, $q \in \mathcal{K}_+(L_w)$.

The case $p < 2 \leq q$ is straightforward. Since $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^q(w))$, by Lemma 2.28 we have $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^2(w) \rightarrow L^q(w))$ and $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^p(w) \rightarrow L^2(w))$. Hence, $p \in \mathcal{K}_-(L_w)$ and $q \in \mathcal{K}_+(L_w)$.

Finally, we prove (4). Suppose to the contrary that $\sup \tilde{\mathcal{K}}(L_w) > q_+(L_w)$. Then there exist p, q such that $q_+(L_w) < p < q < \sup \tilde{\mathcal{K}}(L_w)$. Fix r such that $p_-(L_w) = q_-(L_w) < r < 2$. Then we have that $\sqrt{t} \nabla e^{-tL_w}$ is uniformly bounded on $L^q(w)$ and in $\mathcal{O}(L^r(w) \rightarrow L^2(w))$. By Lemma 2.27 we can interpolate between these to get that $\sqrt{t} \nabla e^{-tL_w} \in \mathcal{O}(L^s(w) \rightarrow L^p(w))$ for some $s < p$. But then by the above converse, we have $p \in \mathcal{K}(L_w)$, which is a contradiction. \square

8. An upper bound for $\mathcal{K}(L_w)$

In this section we will prove that $q_+(L_w) > 2$; that is, the set $\mathcal{K}(L_w)$ contains 2 in its interior. In general, all we can say is that $q_+(L_w) > 2$; as noted in [Auscher 2007, Section 4.5], even in the unweighted case this is the best possible bound, since given any $\varepsilon > 0$ it is possible to find an operator L such that $q_+(L) < 2 + \varepsilon$. In Section 11 below we will give some estimates for $q_+(L_w)$ in terms of $[w]_{A_2}$.

We have broken the proof that $q_+(L_w) > 2$ into a series of discrete steps where we borrow some ideas from [Auscher and Coulhon 2005]. We first prove a reverse Hölder inequality and use Gehring’s inequality to get a higher-integrability estimate. We then prove that the Hodge projection is bounded on $L^q(w)$ for a range of $q > 2$ and use this to prove the Riesz transform is also bounded for exponents greater than 2. (In Section 9 we give a more complete discussion of the Riesz transform.) From this we deduce that $q_+(L_w) > 2$.

A reverse Hölder inequality. Fix a ball B_0 and let $u \in H^1(w)$ be any weak solution of $L_w u = 0$ in $4B_0$. Then for any ball B such that $3B \subset 4B_0$, we can again prove via a standard argument a Caccioppoli inequality:

$$\left(\int_B |\nabla u|^2 dw \right)^{1/2} \leq \frac{C_1}{r} \left(\int_{2B} |u - u_{2B,w}|^2 dw \right)^{1/2},$$

where $C_1 = C(n, \Lambda/\lambda)[w]_{A_2}^{1/2} \geq 1$. Fix q such that

$$\max \left\{ \frac{2(n-1)}{n}, r_w, \frac{2nr_w}{2+nr_w} \right\} < q < 2; \tag{8.1}$$

such a q exists since $r_w < 2$. Our choice of q guarantees that $2 < q_w^*$ and also that $2 < nq/(q - 1)$. Then, by the weighted Poincaré inequality, Theorem 2.1,

$$\frac{1}{r} \left(\int_{2B} |u - u_{2B,w}|^2 dw \right)^{1/2} \leq C_2 \left(\int_{2B} |\nabla u|^q dw \right)^{1/q}, \tag{8.2}$$

where $C_2 = C(n)[w]_{A_2}^\kappa \geq 1$ and $\kappa = (nq - 1)/(nq(q - 1))$. (By our choice of q we can get this sharp estimate; see Remark 2.5. Since $q < 2$ we could write $[w]_{A_q}$, but we use that $[w]_{A_q} \leq [w]_{A_2}$.) If we combine these inequalities, we get a reverse Hölder inequality:

$$\left(\int_B |\nabla u|^2 dw \right)^{1/2} \leq C_1 C_2 \left(\int_{2B} |\nabla u|^q dw \right)^{1/q}.$$

We now apply Gehring’s lemma in the setting of spaces of homogeneous type [Björn and Björn 2011, Theorem 3.22] to get that there exists $p_0 > 2$ such that for every such B ,

$$\left(\int_B |\nabla u|^{p_0} dw \right)^{1/p_0} \leq C_0 \left(\int_{2B} |\nabla u|^2 dw \right)^{1/2}. \tag{8.3}$$

Moreover, we can take the values $C_0 = 8C_1^2 C_2^2 [w]_{A_2}^{31}$ and

$$p_0 = 2 + \frac{2 - q}{2^{4/q+1} C_1^2 C_2^2 [w]_{A_2}^{6/q+17}}. \tag{8.4}$$

In Section 11 below we will need these precise values. Here, it suffices to note that in inequality (8.3) we have $p_0 > 2$.

The Hodge projection. Define the Hodge projection operator by

$$T = \nabla L_w^{-1/2} (\nabla (L_w^*)^{-1/2})^*,$$

where the adjoint operators are defined with respect to the inner product in $L^2(w)$. As we noted in Section 2, the Riesz transform is bounded on $L^2(w)$; hence, the Hodge projection is also bounded. By duality, $(\nabla (L_w^*)^{-1/2})^* \vec{f} = -L_w^{-1/2} (w^{-1} \operatorname{div}(w \vec{f}))$, and so

$$T \vec{f} = -\nabla L_w^{-1/2} L_w^{-1/2} (w^{-1} \operatorname{div}(w \vec{f})) = -\nabla L_w^{-1} (w^{-1} \operatorname{div}(w \vec{f})).$$

Now fix $\vec{f} \in L^2(w, \mathbb{C}^n) \cap L^{p_0}(w, \mathbb{C}^n)$ such that $\operatorname{supp}(\vec{f}) \subset \mathbb{R}^n \setminus 4B_0$. Let $u \in H^1(w)$ be a solution to the equation

$$L_w u = w^{-1} \operatorname{div}(w \vec{f});$$

by a standard Lax–Milgram argument because A satisfies (2.7) [Fabes et al. 1982, Theorem 2.2], we know u exists. Then

$$T \vec{f} = -\nabla L_w^{-1} L_w u = -\nabla u,$$

where equality is in the sense of distributions. In particular, since $f = 0$ on $4B_0$, we know $L_w u = 0$ on $4B_0$. Therefore, we can apply (8.3) to u : on any ball B such that $3B \subset 4B_0$,

$$\left(\int_B |T \vec{f}|^{p_0} dw \right)^{1/p_0} = \left(\int_B |\nabla u|^{p_0} dw \right)^{1/p_0} \leq C_0 \left(\int_{2B} |\nabla u|^2 dw \right)^{1/2} = \left(\int_{2B} |T \vec{f}|^2 dw \right)^{1/2}.$$

As a consequence of this inequality, by [Auscher and Martell 2007a, Theorem 3.14] (see also Section 5 of the same paper), for all q , $2 \leq q < p_0$, we have $T : L^q(w, \mathbb{C}^n) \rightarrow L^q(w, \mathbb{C}^n)$.

Boundedness of the Riesz transform. To show that the Riesz transform $\nabla L_w^{-1/2}$ is bounded, fix q such that

$$\max\{p_-(L_w^*), r_w, p'_0\} = \max\left\{p_-(L_w^*), r_w, p'_0, \frac{nr_w p_-(L_w^*)}{nr_w + p_-(L_w^*)}\right\} < q' < 2.$$

(The reason for including $p_-(L_w^*)$ will be made clear below.) By the above argument we have that T^* is bounded on $L^{q'}(w)$, where $T^* \vec{f} = -\nabla(L_w^*)^{-1}(w^{-1} \operatorname{div}(w \vec{f}))$. Furthermore, by Proposition 6.1, we have

$$\|(L_w^*)^{1/2} f\|_{L^{q'}(w)} \leq C \|\nabla f\|_{L^{q'}(w)}.$$

Therefore,

$$\begin{aligned} \|(\nabla L_w^{-1/2})^* \vec{f}\|_{L^{q'}(w)} &= \|(L_w^*)^{-1/2}(w^{-1} \operatorname{div}(w \vec{f}))\|_{L^{q'}(w)} \\ &= \|(L_w^*)^{1/2}(L_w^*)^{-1}(w^{-1} \operatorname{div}(w \vec{f}))\|_{L^{q'}(w)} \\ &\lesssim \|\nabla(L_w^*)^{-1}(w^{-1} \operatorname{div}(w \vec{f}))\|_{L^{q'}(w)} \\ &= \|T^* \vec{f}\|_{L^{q'}(w)} \lesssim \|\vec{f}\|_{L^{q'}(w)}. \end{aligned}$$

Hence, by duality we have $\nabla L_w^{-1/2} : L^q(w) \rightarrow L^q(w)$ for all q such that

$$2 < q < \min\{p_+(L_w), r'_w, p_0\} = q_w;$$

here we have used the fact that by duality, $p_-(L_w^*)' = p_+(L_w)$.

Boundedness of the gradient of the semigroup. Finally, we show that if $2 < q < q_w$, then $\sqrt{t} \nabla e^{-tL_w} : L^q(w) \rightarrow L^q(w)$. The desired estimate for $q_+(L_w)$ follows from this: by Proposition 7.1, part (4),

$$q_+(L_w) = \sup \tilde{\mathcal{K}}(L_w) \geq q_w > 2.$$

Fix such a q ; then by the above estimate for the Riesz transform,

$$\begin{aligned} \|\sqrt{t} \nabla e^{-tL_w} f\|_{L^q(w)} &= \|\nabla L_w^{-1/2}(tL_w)^{1/2} e^{-tL_w} f\|_{L^q(w)} \\ &\lesssim \|(tL_w)^{1/2} e^{-tL_w} f\|_{L^q(w)} = \|\varphi_t(L_w) f\|_{L^q(w)}, \end{aligned}$$

where $\varphi_t(z) = (tz)^{1/2} e^{-tz}$. For all $t > 0$, this is a uniformly bounded holomorphic function in the right half-plane. Therefore, since $2 < q < p_+(L_w)$, by Proposition 4.3 we have

$$\|\sqrt{t} \nabla e^{-tL_w} f\|_{L^q(w)} \lesssim \|\varphi_t\|_\infty \|f\|_{L^q(w)} \lesssim \|f\|_{L^q(w)}$$

and the bound is independent of t . This completes the proof that $q_+(L_w) > 2$.

9. Riesz transform estimates

In this section we prove $L^p(w)$ norm inequalities for the Riesz transform $\nabla L_w^{-1/2}$. We have already proved such inequalities for a small range of values $q > 2$ in Section 8. Here we prove the following result.

Proposition 9.1. *Let $q_-(L_w) < p < q_+(L_w)$. Then there exists a constant C such that*

$$\|\nabla L_w^{-1/2} f\|_{L^p(w)} \leq C \|f\|_{L^p(w)}. \tag{9.2}$$

Furthermore, if $v \in A_{p/q_-(L_w)}(w) \cap \text{RH}_{(q_+(L_w)/p)'}(w)$, then

$$\|\nabla L_w^{-1/2} f\|_{L^p(v dw)} \leq C \|f\|_{L^p(v dw)}. \tag{9.3}$$

To prove Proposition 9.1 we would like to follow the same outline as the proof of Proposition 4.3. The first step (i.e., proving (9.2) holds when $q_-(L_w) < p < 2$) does work with the appropriate changes. However, the second step (i.e., the proof that (9.3) holds) runs into difficulties since $\nabla L_w^{-1/2}$ and the auxiliary operators \mathcal{A}_r do not commute. One approach to overcoming this obstacle would be to adapt the proof in [Auscher and Martell 2006]; see also [Auscher 2007]. In this case we would need to use an $L^{p_0}(w)$ -Poincaré inequality, which may not hold unless we assume $w \in A_{p_0}$. This would yield estimates in the range $\max\{r_w, q_-(L_w)\} < p < q_+(L_w)$, analogous to those in Proposition 6.1.

There is, however, an alternative approach. In [Auscher and Martell 2008] the authors considered Riesz transforms associated with the Laplace–Beltrami operator of a complete, noncompact Riemannian manifold. Their proof avoids Poincaré inequalities for p close to 1 as these may not hold. Instead, they use a duality argument based on ideas in [Bernicot and Zhao 2008]; this requires that they first prove that the Riesz transform is bounded for $p > 2$ in the appropriate range of values. This reverses the order used in the proof of Proposition 4.3.

Proof of Proposition 9.1. For brevity, let $q_- = q_-(L_w)$ and $q_+ = q_+(L_w)$. To implement the approach sketched above, we divide the proof in two steps. First we will prove that (9.2) holds when $2 < p < q_+$. We do so using Theorem 2.35 and some ideas from [Auscher 2007; Auscher and Martell 2006]. We note that since the Riesz transform and \mathcal{A}_r do not commute, we will use an $L^2(w)$ -Poincaré inequality. This holds since $w \in A_2$; the problem with using the Poincaré inequality only occurs with exponents less than 2. The second step is to prove that (9.3) holds by adapting the proof in [Auscher and Martell 2008]. Here we will use duality and a result from [Auscher and Martell 2007a] that is based on good- λ inequalities. Inequality (9.2) then holds when $q_- < p < 2$ by taking $v \equiv 1$.

To apply Theorem 2.35, fix $2 < p < q_+$ and let $T = \nabla L_w^{-1/2}$, $S = I$ and $\mathcal{D} = L_c^\infty$. Let $p_0 = 2$ and fix q_0 such that $2 < p < q_0 < q_+$. As before we take $\mathcal{A}_r = I - (I - e^{-r^2 L_w})^m$, where m will be chosen below. We first show that (2.36) holds. Let $f \in L_c^\infty$ and decompose it as in (4.11); then we have

$$\left(\int_B |\nabla L_w^{-1/2} (I - e^{-r^2 L_w})^m f|^2 dw \right)^{1/2} \leq \sum_{j \geq 1} \left(\int_B |\nabla L_w^{-1/2} (I - e^{-r^2 L_w})^m f_j|^2 dw \right)^{1/2}.$$

To estimate the first term, note that $\nabla L_w^{-1/2}$ and $e^{-r^2 L_w}$ are bounded on $L^2(w)$ by Theorems 2.15 and 2.18. Hence,

$$\left(\int_B |\nabla L_w^{-1/2} (I - e^{-r^2 L_w})^m f_1|^2 dw \right)^{1/2} \lesssim \left(\int_{4B} |f|^2 dw \right)^{1/2}. \tag{9.4}$$

Fix $j \geq 2$; to get the desired L^2 estimates we will use the L^2 bounds for the gradient of the square function. If $h \in L^2(w)$, by (2.20)

$$\nabla L_w^{-1/2} (I - e^{-r^2 L_w})^m h = \frac{1}{\sqrt{\pi}} \int_0^\infty \sqrt{t} \nabla \varphi(L_w, t) h \frac{dt}{t}, \tag{9.5}$$

where $\varphi(z, t) = e^{-tz} (1 - e^{-r^2 z})^m \in \mathcal{H}_0^\infty(\Sigma_\mu)$. We can therefore use the integral representation (2.10) for $\varphi(\cdot, t)$. The function $\eta(\cdot, t)$ in this representation satisfies

$$|\eta(z, t)| \lesssim \frac{r^{2m}}{(|z| + t)^{m+1}}, \quad z \in \Gamma, t > 0.$$

By Theorem 2.15, $\sqrt{z} \nabla e^{-z L_w} \in \mathcal{O}(L^2(w) \rightarrow L^2(w))$; hence,

$$\begin{aligned} & \left(\int_B \left| \int_\Gamma \eta(z) \sqrt{t} \nabla e^{-z L_w} f_j dz \right|^2 dw \right)^{1/2} \\ & \leq \int_\Gamma \left(\int_B |\sqrt{z} \nabla e^{-z L_w} f_j|^2 dw \right)^{1/2} \frac{\sqrt{t}}{\sqrt{|z|}} |\eta(z)| |dz| \\ & \lesssim 2^{j\theta_1} \int_\Gamma \Upsilon \left(\frac{2^j r}{\sqrt{|z|}} \right)^{\theta_2} e^{-\alpha 4^j r^2 / |z|} \frac{\sqrt{t}}{\sqrt{|z|}} |\eta(z)| |dz| \left(\int_{C_j(B)} |f|^2 dw \right)^{1/2} \\ & \lesssim 2^{j\theta_1} \int_0^\infty \Upsilon \left(\frac{2^j r}{\sqrt{s}} \right)^{\theta_2} e^{-\alpha 4^j r^2 / s} \frac{\sqrt{t}}{\sqrt{s}} \frac{r^{2m}}{(s+t)^{m+1}} ds \left(\int_{C_j(B)} |f|^2 dw \right)^{1/2}. \end{aligned} \tag{9.6}$$

When $2m > \theta_2$,

$$\int_0^\infty \int_0^\infty \Upsilon \left(\frac{2^j r}{\sqrt{s}} \right)^{\theta_2} e^{-\alpha 4^j r^2 / s} \frac{\sqrt{t}}{\sqrt{s}} \frac{r^{2m}}{(s+t)^{m+1}} ds \frac{dt}{t} = C 4^{-jm}. \tag{9.7}$$

If we insert this into the representation (2.10) we get

$$\begin{aligned} \left(\int_B |\nabla e^{-t L_w} (I - e^{-r^2 L_w})^m f_j|^2 dw \right)^{1/2} & \lesssim \int_0^\infty \left(\int_B |\sqrt{t} \nabla \varphi(L_w, t) f_j|^2 dw \right)^{1/2} \frac{dt}{t} \\ & \lesssim 2^{j(\theta_1 - 2m)} \left(\int_{C_j(B)} |f|^2 dw \right)^{1/2}. \end{aligned} \tag{9.8}$$

If we now combine (9.4) and (9.8) we get (2.36) with $g(j) = C_m 2^{j(\theta_1 - 2m)}$; if we also fix $2m > \theta_1$, we get that $\sum g(j) < \infty$.

We now show that (2.37) holds. As we remarked above, the Riesz transform does not commute with \mathcal{A}_r . To overcome this obstacle, we will prove an off-diagonal estimate for the gradient of the semigroup (using the $L^2(w)$ -Poincaré inequality), and then use an approximation argument to get the desired estimate for the Riesz transform.

More precisely, we claim that for every $f \in H^1(w)$ and $1 \leq k \leq m$,

$$\left(\int_B |\nabla e^{-k r^2 L w} f|^{q_0} dw \right)^{1/q_0} \leq \sum_{j \geq 1} g(j) \left(\int_{2^{j+1} B} |\nabla f|^2 dw \right)^{1/2}, \tag{9.9}$$

where $g(j) = C_m 2^j \sum_{l \geq j} 2^{l \theta} e^{-\alpha 4^l}$. Assume for the moment that (9.9) holds. Then for every $\varepsilon > 0$ we can apply this estimate to $S_\varepsilon f$, defined by (2.21), since $S_\varepsilon f \in H^1(w)$. Moreover, we have that \mathcal{A}_r and S_ε commute, and so if we expand $\mathcal{A}_r = I - (I - e^{-r^2 L})^m$ and apply (9.9), we get

$$\left(\int_B |\nabla S_\varepsilon \mathcal{A}_r f|^{q_0} dw \right)^{1/q_0} \leq C_m \sum_{j \geq 1} g(j) \left(\int_{2^{j+1} B} |\nabla S_\varepsilon f|^2 dw \right)^{1/2}.$$

If we let ε go to 0, we obtain (2.37). (The justification of this uses the observations made in Section 2 after (2.21) and is left to the reader.) Moreover, we have $\sum_{j \geq 1} g(j) < \infty$, and so by Theorem 2.35 with $v \equiv 1$, which trivially satisfies $v \in A_{p/2}(w) \cap \text{RH}_{(q_0/p)'}(w)$, we have that (9.2) holds for $f \in L_c^\infty$ and for every $2 < p < q_+$.

To complete this step we need to prove (9.9). Fix $1 \leq k \leq m$ and $f \in H^1(w)$. Let $h = f - f_{4B,w}$, where $f_{4B,w} = \int_{4B} f dw$. Then by the conservation property (see [Cruz-Uribe and Rios 2015], or the proof in [Auscher 2007, Section 2.5]), $e^{-tL} 1 = 1$ for all $t > 0$, and so

$$\nabla e^{-k r^2 L w} f = \nabla e^{-k r^2 L w} (f - f_{4B,w}) = \nabla e^{-k r^2 L w} h = \sum_{j \geq 1} \nabla e^{-k r^2 L w} h_j,$$

where $h_j = h \chi_{C_j(B)}$. Hence,

$$\left(\int_B |\nabla e^{-k r^2 L w} f|^{q_0} dw \right)^{1/q_0} \leq \sum_{j \geq 1} \left(\int_B |\nabla e^{-k r^2 L w} h_j|^{q_0} dw \right)^{1/q_0}.$$

Since $2 < q_0 < q_+$, by Proposition 7.1, $\sqrt{t} \nabla e^{-tL} \in \mathcal{O}(L^2(w) \rightarrow L^{q_0}(w))$. If we apply this and the $L^2(w)$ -Poincaré inequality (see Remark 2.6 with $p = q = 2$), then for each $j \geq 1$ we get

$$\begin{aligned} & \left(\int_B |\nabla e^{-k r^2 L w} h_j|^{q_0} dw \right)^{1/q_0} \\ & \lesssim \frac{2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j}}{r} \left(\int_{C_j(B)} |h_j|^2 dw \right)^{1/2} \\ & \leq \frac{2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j}}{r} \left(\int_{2^{j+1} B} |f - f_{4B,w}|^2 dw \right)^{1/2} \\ & \leq \frac{2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j}}{r} \left(\left(\int_{2^{j+1} B} |f - f_{2^{j+1} B,w}|^2 dw \right)^{1/2} + \sum_{l=2}^j |f_{2^l B,w} - f_{2^{l+1} B,w}| \right) \\ & \lesssim \frac{2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j}}{r} \sum_{l=1}^j \left(\int_{2^{l+1} B} |f - f_{2^{l+1} B,w}|^2 dw \right)^{1/2} \\ & \lesssim 2^{j(\theta_1 + \theta_2)} e^{-\alpha 4^j} \sum_{l=1}^j 2^l \left(\int_{2^{l+1} B} |\nabla f|^2 dw \right)^{1/2}. \end{aligned}$$

If we combine these two estimates and exchange the order of summation we get (9.9) with $\theta = \theta_1 + \theta_2$. This completes the proof that (9.2) holds when $2 < p < q_+$.

For the second step of our proof we show that (9.3) holds for all p , $q_- < p < q_+$, and $v \in A_{p/q_-}(w) \cap \text{RH}_{(q_+/p)'}(w)$. Fix such a p and v ; then by the openness properties of A_q and RH_s weights, there exist p_0, q_0 such that

$$q_- < p_0 < \min\{p, 2\} \leq \max\{p, 2\} < q_0 < q_+ \quad \text{and} \quad v \in A_{p/p_0}(w) \cap \text{RH}_{(q_0/p)'}(w).$$

By the duality properties of weights [Auscher and Martell 2007a, Lemma 4.4],

$$u = v^{1-p'} \in A_{p'/q'_0}(w) \cap \text{RH}_{(p'_0/p)'}(w).$$

Let $T = \nabla L_w^{-1/2}$; then T is bounded from $L^p(\mathbb{R}^n, v \, dw)$ to $L^p(\mathbb{R}^n; \mathbb{C}^n, v \, dw)$ if and only if T^* is bounded from $L^{p'}(\mathbb{R}^n; \mathbb{C}^n, u \, dw)$ to $L^{p'}(\mathbb{R}^n; u \, dw)$. (Note that T takes scalar-valued functions to vector-valued functions and T^* does the opposite.)

Therefore, it will suffice to prove the boundedness of T^* . We will do so using a particular case of [Auscher and Martell 2007a, Theorem 3.1]. This result is stated there in the Euclidean setting but it extends to spaces of homogeneous type. Here we give the weighted version we need; see [loc. cit., Section 5].

Theorem 9.10. *Fix $1 < q < \infty$, $a \geq 1$ and $u \in \text{RH}_{s'}(w)$, $1 < s < \infty$. Then there exists $C > 1$ with the following property: suppose $F \in L^1(w)$ and G are nonnegative measurable functions such that for any ball B there exist nonnegative functions G_B and H_B with $F(x) \leq G_B(x) + H_B(x)$ for a.e. $x \in B$ and, for all $x \in B$,*

$$\left(\int_B H_B^q \, dw \right)^{1/q} \leq a M_w F(x), \quad \int_B G_B \, dw \leq G(x), \tag{9.11}$$

where M_w is the Hardy–Littlewood maximal function with respect to dw . Then for $1 < t < q/s$,

$$\|M_w F\|_{L^t(u \, dw)} \leq C \|G\|_{L^t(u \, dw)}. \tag{9.12}$$

To apply Theorem 9.10, fix $\vec{f} \in L_c^\infty(\mathbb{R}^n; \mathbb{C}^n)$, and let $h = T^* \vec{f}$ and $F = |h|^{q'_0}$. Then $F \in L^1(w)$; by the argument above, since $2 < q_0 < q_+$, we have that T is bounded from $L^{q_0}(\mathbb{R}^n, w)$ to $L^{q_0}(\mathbb{R}^n; \mathbb{C}^n, w)$, thus, T^* is bounded from $L^{q'_0}(\mathbb{R}^n; \mathbb{C}^n, w)$ to $L^{q'_0}(\mathbb{R}^n, w)$.

Now let $\mathcal{A}_r = I - (I - e^{-r^2 L_w})^m$, where $m > 0$ will be fixed below. Given a ball B with radius r , we define

$$F \leq 2^{q'_0-1} |(I - \mathcal{A}_r)^* h|^{q'_0} + 2^{q'_0-1} |\mathcal{A}_r^* h|^{q'_0} \equiv G_B + H_B,$$

where, as before, the adjoint is with respect to $L^2(w)$. To complete the proof, suppose for the moment that we could prove (9.11) with $q = p'_0/q'_0$ and $G = M_w(|\vec{f}|^{q'_0})$. Since $u \in \text{RH}_{(p'_0/p)'}(w)$, by the openness property of reverse Hölder weights, $u \in \text{RH}_{s'}(w)$ for some $s < p'_0/p'$. Then if we let $t = p'/q'_0 = (p'_0/q'_0)/(p'_0/p') < q/s$, we have $u \in A_t(w)$, and so M_w is bounded on $L^t(u \, dw)$. Therefore, by (9.12),

$$\|T^* \vec{f}\|_{L^{p'}(u \, dw)}^{q'_0} \leq \|M_w F\|_{L^t(u \, dw)} \leq C \|G\|_{L^t(u \, dw)} = C \|M_w(|\vec{f}|^{q'_0})\|_{L^t(u \, dw)} \lesssim \|\vec{f}\|_{L^{p'}(u \, dw)}^{q'_0}.$$

To complete the proof we need to show that (9.11) holds. We first estimate H_B . By duality there exists $g \in L^{p_0}(B, dw/w(B))$ with norm 1 such that for all $x \in B$,

$$\begin{aligned} \left(\int_B H_B^q dw\right)^{1/(q q'_0)} &\lesssim w(B)^{-1} \int_{\mathbb{R}^n} |h| |\mathcal{A}_r g| dw \\ &\lesssim \sum_{j=1}^{\infty} 2^{jD} \left(\int_{C_j(B)} |h|^{q'_0} dw\right)^{1/q'_0} \left(\int_{C_j(B)} |\mathcal{A}_r g|^{q_0} dw\right)^{1/q_0} \\ &\lesssim M_w F(x)^{1/q'_0} \sum_{j=1}^{\infty} 2^{j(D+\theta_1+\theta_2)} e^{-\alpha 4^j} \left(\int_B |g|^{p_0} dw\right)^{1/p_0} \lesssim M_w F(x)^{1/q'_0}, \end{aligned}$$

where in the second-to-last inequality we used the fact that by our choice of p_0, q_0 , we have $e^{-tLw} \in \mathcal{O}(L^{p_0}(w) \rightarrow L^{q_0}(w))$, and so \mathcal{A}_r is as well.

We now estimate G_B . Again by duality there exists $g \in L^{q_0}(B, dw/w(B))$ with norm 1 such that for all $x \in B$,

$$\begin{aligned} \left(\int_B G_B dw\right)^{1/q'_0} &\lesssim w(B)^{-1} \int_{\mathbb{R}^n} |\vec{f}| |T(I - \mathcal{A}_r)g| dw \\ &\lesssim \sum_{j=1}^{\infty} 2^{jD} \left(\int_{C_j(B)} |\vec{f}|^{q'_0} dw\right)^{1/q'_0} \left(\int_{C_j(B)} |T(I - \mathcal{A}_r)g|^{q_0} dw\right)^{1/q_0} \\ &\leq M_w (|\vec{f}|^{q'_0})(x)^{1/q'_0} \sum_{j=1}^{\infty} 2^{jD} \left(\int_{C_j(B)} |T(I - \mathcal{A}_r)g|^{q_0} d\mu\right)^{1/q_0}. \end{aligned} \tag{9.13}$$

To estimate each term in the sum, we argue as in the first half of the proof. When $j = 1$, we know that $\nabla L_w^{-1/2}$ and $e^{-r^2 Lw}$ are bounded on $L^{q_0}(w)$ by the first part of the proof and Theorem 2.15. Hence,

$$\left(\int_{4B} |\nabla L_w^{-1/2}(I - e^{-r^2 Lw})^m g|^{q_0} dw\right)^{1/q_0} \lesssim \left(\int_B |g|^{q_0} dw\right)^{1/q_0} = 1. \tag{9.14}$$

For $j \geq 2$ we use the integral representation (9.5). If we estimate as in (9.6), with the roles of B and $C_j(B)$ switched and using the fact that $\sqrt{z} \nabla e^{-zLw} \in \mathcal{O}(L^{q_0}(w) \rightarrow L^{q_0}(w))$ since $2 < q_0 < q_+$, we see that

$$\begin{aligned} \left(\int_{C_j(B)} \left| \int_{\Gamma} \eta(z) \sqrt{t} \nabla e^{-zLw} g dz \right|^{q_0} dw\right)^{1/q_0} &\leq \int_{\Gamma} \left(\int_{C_j(B)} |\sqrt{z} \nabla e^{-zLw} g|^{q_0} dw\right)^{1/q_0} \frac{\sqrt{t}}{\sqrt{|z|}} |\eta(z)| |dz| \\ &\lesssim 2^{j\theta_1} \int_{\Gamma} \Upsilon\left(\frac{2^j r}{\sqrt{|z|}}\right)^{\theta_2} e^{-\alpha 4^j r^2/|z|} \frac{\sqrt{t}}{\sqrt{|z|}} |\eta(z)| |dz| \left(\int_B |g|^{q_0} dw\right)^{1/2} \\ &\lesssim 2^{j\theta_1} \int_0^{\infty} \Upsilon\left(\frac{2^j r}{\sqrt{s}}\right)^{\theta_2} e^{-\alpha 4^j r^2/s} \frac{\sqrt{t}}{\sqrt{s}} \frac{r^{2m}}{(s+t)^{m+1}} ds. \end{aligned}$$

If we take $2m > \theta_2$, we can combine this with (9.7). We can then insert this estimate into the representation (2.10) to get that for every $j \geq 2$,

$$\begin{aligned} \left(\int_{C_j(B)} |\nabla e^{-tL_w} (I - e^{-r^2 L_w})^m g|^{q_0} dw \right)^{1/q_0} \\ \lesssim \int_0^\infty \left(\int_{C_j(B)} |\sqrt{t} \nabla \varphi(L_w, t) g|^{q_0} dw \right)^{1/q_0} \frac{dt}{t} \lesssim 2^{j(\theta_1 - 2m)}. \end{aligned} \quad (9.15)$$

Taken together, (9.13)–(9.15) yield

$$\left(\int_B G_B dw \right)^{1/q'_0} \lesssim M_w(|\vec{f}|^{q'_0})(x)^{1/q'_0} \sum_{j=1}^\infty 2^{j(D + \theta_1 - 2m)} \lesssim M_w(|\vec{f}|^{q'_0})(x)^{1/q'_0} = G(x)^{1/q'_0},$$

provided we take m large enough so that $D + \theta_1 - 2m < 0$. This completes the estimate of H_B and G_B and so completes our proof. □

10. Square function estimates for the gradient of the semigroup

In this section we prove $L^p(w)$ estimates for the vertical square function

$$G_{L_w} f(x) = \left(\int_0^\infty |t^{1/2} \nabla e^{-tL_w} f(x)|^2 \frac{dt}{t} \right)^{1/2}.$$

Proposition 10.1. *Let $q_-(L_w) < p < q_+(L_w)$. Then*

$$\|G_{L_w} f\|_{L^p(w)} \lesssim \|f\|_{L^p(w)}. \quad (10.2)$$

Furthermore, if $v \in A_{p/q_-(L_w)}(w) \cap \text{RH}_{(q_+(L_w)/p)'}(w)$, then

$$\|G_{L_w} f\|_{L^p(v dw)} \lesssim \|f\|_{L^p(v dw)}. \quad (10.3)$$

We can also prove a reverse inequality for G_{L_w} . To do so we need to introduce an auxiliary operator. Define the weighted Laplacian by $\Delta_w = -w^{-1} \text{div } w \nabla$; i.e., Δ_w is the operator L_w if we take the matrix A to be wI , where I is the identity matrix.

Proposition 10.4. *Let $q_+(\Delta_w)' < p < \infty$. Then*

$$\|f\|_{L^p(w)} \lesssim \|G_{L_w} f\|_{L^p(w)}. \quad (10.5)$$

Furthermore, if $v \in A_{p/q_+(\Delta_w)'}(w)$, then

$$\|f\|_{L^p(v dw)} \lesssim \|G_{L_w} f\|_{L^p(v dw)}. \quad (10.6)$$

Proof of Proposition 10.1. The proof could be done in a way similar to those for the square function g_{L_w} in Section 5. However, we will give a shorter proof that uses the Riesz transform estimates from Section 9.

Let $q_- = q_-(L_w)$ and $q_+ = q_+(L_w)$. Fix p ,

$$q_- = p_-(L_w) < p < q_+ \leq p_+(L_w),$$

and $v \in A_{p/q_-}(w) \cap \text{RH}_{(q_+/p)'}(w)$. Then by Proposition 9.1, the Riesz transform is bounded on $L^p(v dw)$, and so by Lemma 5.4 it has a bounded extension to $L^p_{\mathbb{H}}(v dw)$; i.e., if $g(x, t) \in L^p_{\mathbb{H}}(v dw)$, then $\|\nabla L_w^{-1/2} g\|_{L^p_{\mathbb{H}}(v dw)} \lesssim \|g\|_{L^p_{\mathbb{H}}(v w)}$, where the extension of $\nabla L_w^{-1/2}$ to \mathbb{H} -valued functions is defined for $x \in \mathbb{R}^n$ and $t > 0$ by $(\nabla L_w^{-1/2} g)(x, t) = \nabla L_w^{-1/2}(g(\cdot, t))(x)$.

Define $g_f(x, t) = (tL_w)^{1/2} e^{-tL_w} f(x)$ and $G_f(x, t) = t^{1/2} \nabla e^{-tL_w} f(x)$; then we clearly have $\|g_{L_w f}\|_{L^p(v dw)} = \|g_f\|_{L^p_{\mathbb{H}}(v dw)}$ and $\|G_{L_w f}\|_{L^p(v dw)} = \|G_f\|_{L^p_{\mathbb{H}}(v dw)}$. Furthermore, $G_f(x, t) = \nabla L_w^{-1/2}(g_f(\cdot, t))(x) = (\nabla L_w^{-1/2} g_f)(x, t)$. Hence,

$$\begin{aligned} \|G_{L_w f}\|_{L^p(v dw)} &= \|G_f\|_{L^p_{\mathbb{H}}(v dw)} = \|\nabla L_w^{-1/2} g_f\|_{L^p_{\mathbb{H}}(v dw)} \\ &\lesssim \|g_f\|_{L^p_{\mathbb{H}}(v dw)} = \|g_{L_w f}\|_{L^p(v dw)} \lesssim \|f\|_{L^p(v dw)}. \end{aligned}$$

To prove the last inequality, we used Proposition 5.1; we also used the fact that $q_- = p_-(L_w) < p < q_+ \leq p_+(L_w)$ and $v \in A_{p/q_-}(w) \cap \text{RH}_{(q_+/p)'}(w)$, which together imply $v \in A_{p/p_-(L_w)}(w) \cap \text{RH}_{(p_+(L_w)/p)'}(w)$. This proves (10.3). To prove inequality (10.2), we take $v \equiv 1$. \square

To prove Proposition 10.4 we need the following identity relating G_{L_w} and Δ_w . It is a straightforward extension of a similar unweighted result given in [Auscher 2007, Section 7.1]. For completeness we include the proof.

Lemma 10.7. *If $f, g \in L^\infty(w)$ then*

$$\left| \int_{\mathbb{R}^n} f(x) \overline{g(x)} dw \right| \leq (\Lambda + 1) \int_{\mathbb{R}^n} G_{L_w} f(x) \overline{G_{\Delta_w} g(x)} dw.$$

Proof. By the definition and properties of the operators L_w and Δ_w we have

$$\begin{aligned} \int_{\mathbb{R}^n} f(x) \overline{g(x)} dw &= \lim_{\varepsilon \downarrow 0} \int_{\mathbb{R}^n} e^{-\varepsilon L_w} f(x) \overline{e^{-\varepsilon \Delta_w} g(x)} dw - \lim_{R \uparrow \infty} \int_{\mathbb{R}^n} e^{-RL_w} f(x) \overline{e^{-R\Delta_w} g(x)} dw \\ &= - \int_0^\infty \frac{d}{dt} \int_{\mathbb{R}^n} e^{-tL_w} f(x) \overline{e^{-t\Delta_w} g(x)} dw dt \\ &= \int_0^\infty \int_{\mathbb{R}^n} (L_w e^{-tL_w} f(x) \overline{e^{-t\Delta_w} g(x)} + e^{-tL_w} f(x) \overline{\Delta_w e^{-t\Delta_w} g(x)}) dw dt \\ &= \int_0^\infty \int_{\mathbb{R}^n} (A(x)w(x)^{-1} + I)(\nabla e^{-tL_w} f(x) \overline{\nabla e^{-t\Delta_w} g(x)}) dw dt. \end{aligned}$$

Since $\|Aw^{-1}\|_\infty \leq \Lambda$, if we apply Hölder’s inequality in the t variable we get the desired result. \square

Proof of Proposition 10.4. As a consequence of the Gaussian estimate for weighted operators with real symmetric coefficients that were proved in [Cruz-Uribe and Rios 2008], we have that $\Delta_w \in \mathcal{O}(L^1(w) \rightarrow L^\infty(w))$. In particular, $q_-(\Delta_w) = p_-(L_{\Delta_w}) = 1$. Further, by the results in Section 8 we have $q_+(\Delta_w) > 2$.

Therefore, by Proposition 10.1, if $1 < p' < q_+(\Delta_w)$, and

$$u \in A_{p'}(w) \cap \text{RH}_{(q_+(\Delta_w)/p)'}(w), \tag{10.8}$$

then

$$\|G_{\Delta_w} f\|_{L^{p'}(u dw)} \lesssim \|f\|_{L^{p'}(u dw)}. \tag{10.9}$$

We want to apply inequality (10.9) with $u = v^{1-p'}$. By [Auscher and Martell 2007a, Lemma 4.4], the condition (10.8) is equivalent to $v \in A_{p/q_+(w)'}(w)$.

Now fix $f, g \in L_c^\infty$, and a weight $v \in A_{p/q_+(w)'}(w)$. Then by Lemma 10.7, for $q_+(\Delta_w)' < p < \infty$,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f(x)g(x) dw \right| &\leq (\Lambda + 1) \int_{\mathbb{R}^n} G_{L_w} f(x)G_{\Delta_w} g(x) dw \\ &= (\Lambda + 1) \int_{\mathbb{R}^n} G_{L_w} f(x)G_{\Delta_w} g(x)v^{1/p} v^{-1/p} dw \\ &\leq (\Lambda + 1) \|G_{L_w} f\|_{L^p(v dw)} \|G_{\Delta_w} g\|_{L^{p'}(v^{1-p'} dw)} \\ &\lesssim \|G_{L_w} f\|_{L^p(v dw)} \|g\|_{L^{p'}(v^{1-p'} dw)}; \end{aligned}$$

the last inequality follows from (10.9). If we take $g = \text{sign}(f)|f|^{p-1}v$, we get

$$\|f\|_{L^p(v dw)}^p \lesssim \|G_{L_w} f\|_{L^p(v dw)} \| |f|^{p-1}v \|_{L^{p'}(v^{1-p'} dw)} = \|G_{L_w} f\|_{L^p(v dw)} \|f\|_{L^p(v dw)}^{p/p'}.$$

This immediately gives us the desired inequality. □

11. Unweighted L^2 Kato estimates

In this section we prove unweighted L^2 estimates for the operators we have considered in the previous sections. These will all be consequences of the weighted $L^p(v dw)$ estimates we have already proved: it will only be necessary to find further conditions on $w \in A_2$ so that the weight $v = w^{-1}$ satisfies the requisite conditions.

We are particularly interested in power weights and we recall some well-known facts about them. Define $w_\alpha(x) = |x|^\alpha$, $\alpha > -n$; this restriction guarantees that w_α is locally integrable. We can exactly determine the Muckenhoupt A_p and reverse Hölder RH_s classes of these weights in terms of α : if $-n < \alpha \leq 0$, then $w \in A_1$; for $1 < p < \infty$, we have $w \in A_p$ if $-n < \alpha < n(p - 1)$. Furthermore, if $0 \leq \alpha < \infty$, then $w \in \text{RH}_\infty$; for $1 < q < \infty$, we have $w \in \text{RH}_q$ if $-n/q < \alpha < \infty$. Hence, we easily see that

$$r_{w_\alpha} = \max\{1, 1 + \alpha/n\}, \quad s_{w_\alpha} = (\max\{1, (1 + \alpha/n)^{-1}\})'. \tag{11.1}$$

We first consider the semigroup e^{-tL_w} , the functional calculus, and the square function g_{L_w} , since these estimates will depend on $p_-(L_w)$ and $p_+(L_w)$ and we have good estimates for these quantities.

Theorem 11.2. *Given a weight $w \in A_2$, suppose $1 \leq r_w < 1 + \frac{2}{n}$ and $s_w > \frac{n}{2}r_w + 1$. Then $e^{-tL_w} : L^2 \rightarrow L^2$ is uniformly bounded for all $t > 0$. Similarly, $\varphi(L_w) : L^2 \rightarrow L^2$, where φ is any bounded holomorphic function on Σ_μ , $\mu \in (\vartheta, \pi)$, and $g_{L_w} : L^2 \rightarrow L^2$.*

In particular, these L^2 estimates hold if we assume that $w \in A_1 \cap \text{RH}_{1+n/2}$, or more generally if $w \in A_r \cap \text{RH}_{(n/2)r+1}$ for $1 < r \leq 1 + \frac{2}{n}$, or if we take the power weights

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} < \alpha < 2.$$

Proof. Let $p = q = 2$, $p_0 = (2_w^*)'$, $q_0 = 2_w^*$, and let $v = w^{-1}$. Then by Proposition 3.1, Corollary 3.3 and the nesting properties of weights, $e^{-tL_w} \in \mathcal{O}(L^2 \rightarrow L^2)$ if $w^{-1} \in A_{2/p_0}(w) \cap \text{RH}_{(q_0/2)'}(w)$; in

particular, by Lemma 2.30, $e^{-tL_w} : L^2 \rightarrow L^2$ is uniformly bounded. However, this weight condition is equivalent to

$$w \in \text{RH}_{(2/p_0)'} \cap A_{q_0/2}.$$

A straightforward computation shows that

$$\frac{q_0}{2} = \frac{nr_w}{nr_w - 2}, \quad \left(\frac{2}{p_0}\right)' = \frac{n}{2}r_w + 1.$$

Since $r_w < 1 + \frac{2}{n}$, we have $r_w < nr_w/(nr_w - 2)$, so we automatically have $w \in A_{q_0/2}$. Therefore, the desired bounds hold if we have $s_w > \frac{n}{2}r_w + 1$. If $w \in A_r \cap \text{RH}_{(n/2)r+1}$ with $1 \leq r \leq 1 + \frac{2}{n}$, then $r_w \leq r$ and $s_w > \frac{n}{2}r + 1 \geq \frac{n}{2}r_w + 1$. The desired conclusion for power weights follows at once from (11.1).

The same argument holds for $\varphi(L_w)$ and g_{L_w} , using Proposition 4.3 or Proposition 5.1, respectively. \square

It is straightforward to construct weights more general than power weights that satisfy the conditions on r_w and s_w in the above theorems. For instance, $w \in A_{1+2/n} \cap \text{RH}_{2+n/2}$ (which corresponds to the choice $r = 1 + \frac{2}{n}$) if and only if there exist $u_1, u_2 \in A_1$ such that

$$w = u_1^{2/(n+4)} u_2^{-2/n}.$$

This follows from the Jones factorization theorem and the properties of A_1 weights; see [Cruz-Uribe and Neugebauer 1995].

Remark 11.3. We can modify the proof of Theorem 11.2 to get unweighted L^p estimates for values of p close to 2. We leave the details to the interested reader.

For the reverse inequalities we must take into account the slightly stronger hypotheses in Proposition 6.1; otherwise, the proof of the following result follows exactly as in the proof of Theorem 11.2.

Theorem 11.4. *Given a weight $w \in A_2$, suppose that*

$$1 \leq r_w < 1 + \frac{2}{n} \quad \text{and} \quad s_w > \max\left\{\left(\frac{2}{r_w}\right)', \frac{n}{2}r_w + 1\right\}.$$

Then

$$\|L_w^{1/2} f\|_{L^2} \leq C \|\nabla f\|_{L^2}, \quad f \in \mathcal{S}. \tag{11.5}$$

In particular, this is the case if we either assume that $w \in A_1 \cap \text{RH}_{1+n/2}$, or more generally that $w \in A_r \cap \text{RH}_{\max\{(2/r)', (n/2)r+1\}}$, with $1 < r \leq 1 + \frac{2}{n}$, or for power weights if we take

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} = -\min\left\{\frac{n}{2}, \frac{2n}{n+2}\right\} < \alpha < 2.$$

Remark 11.6. Note that $\max\left\{\left(\frac{2}{r}\right)', \frac{n}{2}r + 1\right\} = \frac{n}{2}r + 1$ provided $r \leq 2 - \frac{2}{n}$ and this always holds if $n \geq 4$ as $1 + \frac{2}{n} \leq 2 - \frac{2}{n}$. In this case, the conditions in the second part of Theorem 11.4 simplify to the same conditions as in Theorem 11.2.

Remark 11.7. We note that in Theorems 11.2 and 11.4 we can replace $1 \leq r_w < 1 + \frac{2}{n}$ with the possibly weaker condition $1 \leq r_w < p_+(L_w)/2$. The proof only requires us to take $q_0 = p_+(L_w)$.

For the gradient of the semigroup $\sqrt{t}\nabla e^{-tL_w}$, the Riesz transform $\nabla L_w^{-1/2}$, and the square function G_{L_w} , our estimates depend on $q_+(L_w)$.

Theorem 11.8. *Given a weight $w \in A_2$, suppose $1 \leq r_w < q_+(L_w)/2$ and $s_w > \frac{n}{2}r_w + 1$. Then $\sqrt{t}\nabla e^{-tL_w} : L^2 \rightarrow L^2$ is uniformly bounded for all $t > 0$. Similarly, we have $\nabla L_w^{-1/2} : L^2 \rightarrow L^2$ and $G_{L_w} : L^2 \rightarrow L^2$.*

In particular, this is the case if we assume that $w \in A_1 \cap \text{RH}_{n/2+1}$. Furthermore, these L^2 estimates hold if the following is true: given $\Theta \geq 1$ there exists $\varepsilon_0 = \varepsilon_0(\Theta, n, \Lambda/\lambda)$, $0 < \varepsilon_0 \leq \frac{1}{2n}$, such that $w \in A_{1+\varepsilon} \cap \text{RH}_{(n/2)(1+\varepsilon)+1}$, $0 \leq \varepsilon < \varepsilon_0$, and $[w]_{A_2} \leq \Theta$.

For power weights, there exists $\varepsilon_1 = \varepsilon_1(n, \Lambda/\lambda)$, $0 < \varepsilon_1 \leq \frac{1}{2}$, such that these estimate holds for

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} < \alpha < \varepsilon_1.$$

Proof. We will prove this result for $\sqrt{t}\nabla e^{-tL_w}$ using Proposition 7.1. The proof for $\nabla L_w^{-1/2}$ or G_{L_w} is exactly the same, using Proposition 9.1 or Proposition 10.1.

By Proposition 7.1, $\sqrt{t}\nabla e^{-tL_w} : L^2 \rightarrow L^2$ if $w^{-1} = v \in A_{2/q_-(L_w)}(w) \cap \text{RH}_{q_+(L_w)/2}'(w)$, which is equivalent to

$$w \in \text{RH}_{(2/q_-(L_w))}' \cap A_{q_+(L_w)/2}.$$

Therefore, we need $r_w < q_+(L_w)/2$. Furthermore, since we have $q_-(L_w) = p_-(L_w) \leq (2_w^*)'$, we can take

$$s_w > \left(\frac{2}{(2_w^*)'}\right)' = \frac{n}{2}r_w + 1.$$

To get the particular examples stated in the theorem, note first that if we let $r_w = 1$, then it clearly suffices to assume $w \in A_1 \cap \text{RH}_{n/2+1}$, since we showed in Section 8 that $q_+(L_w) > 2$ for every $w \in A_2$.

We now prove the condition for weights $w \in A_{1+\varepsilon}$. In this case it is more difficult to satisfy the condition $r_w < q_+(L_w)/2$ since the right-hand side can be very close to 1, depending on w . Assume then that $w \in A_{1+\varepsilon} \cap \text{RH}_{(n/2)(1+\varepsilon)+1}$, with $0 \leq \varepsilon < \varepsilon_0 \leq \frac{1}{2n}$, $[w]_{A_2} \leq \Theta$, and with $\varepsilon_0 > 0$ to be fixed below. Then we have

$$s_w > \frac{n}{2}(1 + \varepsilon) + 1 \geq \frac{n}{2}r_w + 1.$$

Therefore, in order to apply the first half of the theorem we need to show that we can choose ε_0 sufficiently small so that $r_w < q_+(L_w)/2$. To do so we will use the notation and computations from Section 8. There we showed that $q_+(L_w) \geq q_w$, and so it will suffice to show that

$$2r_w < q_w = \min\{r'_w, p_+(L_w), p_0\}. \tag{11.9}$$

We will compare r_w to each term in the minimum in turn.

The first two terms are straightforward. First, we have $r_w < 1 + \varepsilon < 1 + \frac{1}{2n} < \frac{3}{2}$ and so $2r_w < r'_w$. Second, $r_w < 1 + \frac{1}{2n} < 1 + \frac{2}{n}$, and it follows at once from this that $2r_w < 2_w^*$. By Proposition 3.1, $2_w^* \leq p_+(L_w)$ and so $2r_w < p_+(L_w)$.

Finally, we estimate p_0 , the exponent from the higher-integrability condition (8.3). We will use the formula (8.4). First, we need to fix the exponent q from the Poincaré inequality (8.2). Let $q = 2 - 1/n$; this value satisfies (8.1) since $r_w < 1 + \frac{1}{2n} < 1 + \frac{1}{n}$. With this choice of q (that only depends on n), we have

$$p_0 = 2 + \frac{2 - q}{2^{4/q+1} C_1^2 C_2^2 [w]_{A_2}^{6/q+17}} = 2 + \frac{1}{nC(n, \Lambda/\lambda) [w]_{A_2}^{\theta_n}},$$

where $C(n, \Lambda/\lambda) \geq 1$ depends only on n and the ratio Λ/λ of the ellipticity constants of the matrix A used to define L_w , and where $\theta_n \geq 1$ depends only on n . Then, since we also assumed that $[w]_{A_2} \leq \Theta$, we get that

$$p_0 = 2 + \frac{1}{nC(n, \Lambda/\lambda) [w]_{A_2}^{\theta_n}} \geq 2 + \frac{1}{nC(n, \Lambda/\lambda) \Theta^{\theta_n}} = 2 + 2\varepsilon_0,$$

and $\varepsilon_0 = (2nC(n, \Lambda/\lambda) \Theta^{\theta_n})^{-1}$ is such that $0 < \varepsilon_0 \leq \frac{1}{2n}$. Thus $2r_w < 2(1 + \varepsilon) < 2(1 + \varepsilon_0) \leq p_0$ and so $2r_w < p_0$. This completes the proof that (11.9) is satisfied, and so the L^2 estimates hold for weights that satisfy $w \in A_{1+\varepsilon} \cap \text{RH}_{(n/2)(1+\varepsilon)+1}$.

Finally, we consider power weights. First, it is easy to see that

$$w_\alpha(x) = |x|^\alpha, \quad \frac{-2n}{n+2} < \alpha \leq 0$$

yields the desired estimates, since in this case $r_w = 1$ and $s_w > \frac{n}{2} + 1 = \frac{n}{2}r_w + 1$.

Now consider the case $\alpha > 0$. If we assume that $\alpha < \frac{1}{2}$, then $w \in A_{1+1/(2n)} \cap \text{RH}_\infty$. Moreover, it is straightforward to show that for all such α , there exists Θ , depending only on n , such that $[w_\alpha]_{A_2} \leq \Theta$. Now apply the above argument to find $\varepsilon_0 \in (0, \frac{1}{2n}]$; this value will only depend on n and the ratio Λ/λ . If we let $\varepsilon_1 = n\varepsilon_0$ and assume that $0 < \alpha < \varepsilon_1$, then $\alpha < \frac{1}{2}$ and $w_\alpha \in A_{1+\varepsilon}$ for some $\varepsilon < \varepsilon_0$ as desired. \square

To find examples of weights other than power weights to which Theorem 11.8 apply, we argue as before. If $u_1 \in A_1$, then

$$w = u_1^{2/(n+2)} \in A_1 \cap \text{RH}_{n/2+1}.$$

To get weights that are not in A_1 , take $u \in A_2$ and let $w = u^\theta$. If θ is sufficiently small (depending on n , the ratio Λ/λ and $[u]_{A_2}$), we can show that w satisfies the final conditions given in Theorem 11.8. Details are left to the interested reader.

Remark 11.10. To get the unweighted lower estimate

$$\|f\|_{L^2} \leq C \|G_{L_w} f\|_{L^2},$$

we note that by (10.6) we need $w^{-1} \in A_{2/q_+(\Delta_w)'}(w)$, or equivalently, $w \in \text{RH}_{(2/q_+(\Delta_w))'}$. Hence, it suffices to assume

$$s_w > 1 + \frac{q_+(\Delta_w)}{q_+(\Delta_w) - 2}.$$

Arguing as above we can construct weights that satisfy this condition; details are left to the interested reader.

If we combine Theorems 11.4, 11.8, and Remark 11.7 we solve the Kato square root problem for degenerate elliptic operators.

Theorem 11.11. *Let $L_w = -w^{-1} \operatorname{div} A \nabla$ be a degenerate elliptic operator with $w \in A_2$. If*

$$1 \leq r_w < \frac{q_+(L_w)}{2} \quad \text{and} \quad s_w > \max \left\{ \left(\frac{2}{r_w} \right)', \frac{n}{2} r_w + 1 \right\},$$

then the Kato problem can be solved for L_w ; that is, for every $f \in H^1(\mathbb{R}^n)$,

$$\|L_w^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}, \tag{11.12}$$

where the implicit constants depend only on the dimension, the ellipticity constants λ, Λ , and w .

In particular, (11.12) holds if $w \in A_1 \cap \operatorname{RH}_{n/2+1}$. Further, (11.12) holds if the following is true: given $\Theta \geq 1$ there exists $\varepsilon_0 = \varepsilon_0(\Theta, n, \Lambda/\lambda)$, $0 < \varepsilon_0 \leq \frac{1}{2n}$, such that $w \in A_{1+\varepsilon} \cap \operatorname{RH}_{\max\{(2/(1+\varepsilon))', (n/2)(1+\varepsilon)+1\}}$, $0 \leq \varepsilon < \varepsilon_0$, and $[w]_{A_2} \leq \Theta$.

For power weights, there exists $\varepsilon_1 = \varepsilon_1(n, \Lambda/\lambda)$, $0 < \varepsilon_1 \leq \frac{1}{2}$, such that inequality (11.12) holds (with w_α in place of w) if

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} < \alpha < \varepsilon_1.$$

We can restate the final part of Theorem 11.11 as follows: consider the family of operators $L_\gamma = -|x|^\gamma \operatorname{div}(|x|^{-\gamma} B(x) \nabla)$, where B is an $n \times n$ complex-valued matrix that satisfies the uniform ellipticity condition

$$\lambda |\xi|^2 \leq \operatorname{Re} \langle B(x) \xi, \xi \rangle, \quad |\langle B(x) \xi, \eta \rangle| \leq \Lambda |\xi| |\eta|, \quad \xi, \eta \in \mathbb{C}^n, \text{ a.e. } x \in \mathbb{R}^n.$$

Then,

$$\|L_\gamma^{1/2} f\|_{L^2(\mathbb{R}^n)} \approx \|\nabla f\|_{L^2(\mathbb{R}^n)}, \quad -\varepsilon_1 < \gamma < \frac{2n}{n+2}. \tag{11.13}$$

When $\gamma = 0$ we get the classical Kato square root problem solved by Auscher, Hofmann, Lacey, McIntosh, and Tchamitchian [Auscher et al. 2002]. Inequality (11.13) shows that we can find an open interval containing 0 such that if γ is in this interval, the same estimate holds.

12. Applications to L^2 boundary value problems

In this section we apply the results from the previous section to some L^2 boundary value problems involving the degenerate elliptic operator L_w . We follow the ideas in [Auscher and Tchamitchian 1998] and consider semigroup solutions: for the Dirichlet or regularity problems we let $u(x, t) = e^{-tL_w^{1/2}} f(x)$; for the Neumann problem we let $u(x, t) = -L_w^{-1/2} e^{-tL_w^{1/2}} f(x)$. In each case, for $t > 0$ fixed, $L_w u(\cdot, t)$ makes sense in a weak sense since $u(\cdot, t)$ is in the domain of L_w . Further, derivatives in t are well-defined because of the semigroup properties. Finally, note that by the strong continuity of the semigroup and the off-diagonal estimates, in the context of the following results we have $e^{-tL_w^{1/2}} f \rightarrow f$ as $t \rightarrow 0^+$ in L^2 ; see [Auscher and Martell 2007b, Section 4.2]. Further details are left to the interested reader.

We first consider the Dirichlet problem on $\mathbb{R}_+^{n+1} = \mathbb{R}^n \times [0, \infty)$:

$$\begin{cases} \partial_t^2 u - L_w u = 0 & \text{on } \mathbb{R}^n, \\ u|_{\partial\mathbb{R}_+^{n+1}} = f & \text{on } \partial\mathbb{R}_+^{n+1} = \mathbb{R}^n. \end{cases} \tag{12.1}$$

Theorem 12.2. *Given a weight $w \in A_2$, suppose $1 \leq r_w < 1 + \frac{2}{n}$ and $s_w > \frac{n}{2}r_w + 1$. Then for any $f \in L^2(\mathbb{R}^n)$, we have that $u(x, t) = e^{-tL_w^{1/2}} f(x)$ is a solution of (12.1) with convergence to the boundary data as $t \rightarrow 0^+$ in the L^2 -sense. Furthermore, we have*

$$\sup_{t>0} \|u(\cdot, t)\|_{L^2} \leq C \|f\|_{L^2}. \tag{12.3}$$

In particular, this is the case if we assume that $w \in A_1 \cap \text{RH}_{1+n/2}$, or $w \in A_r \cap \text{RH}_{n/2r+1}$ with $1 < r \leq 1 + \frac{2}{n}$, or if we take the power weights

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} < \alpha < 2.$$

Proof. Formally, it is clear that u is a solution to (12.1), and this formalism can be justified by appealing to the theory of maximal accretive operators; see [Kato 1966]. Alternatively, the weighted estimates for the functional calculus in Proposition 4.3 show that both $(\partial^2/\partial t^2)u(\cdot, t)$ and $L_w u(\cdot, t)$ belong to L^2 for each $t > 0$ and that they are equal in the L^2 -sense. To see that inequality (12.3) holds, it suffices to let $\varphi_t(z) = e^{-t\sqrt{z}}$. Then φ_t is a bounded holomorphic function on Σ_μ , and so by Theorem 11.2 we get the desired bound. \square

Remark 12.4. Note that as observed in Remark 11.7, in the previous result we can replace $1 \leq r_w < 1 + \frac{2}{n}$ with the possibly weaker condition $1 \leq r_w < p_+(L_w)/2$. Also, by Proposition 4.3 we also have that for u as in Theorem 12.2 and all $k \geq 1$,

$$\sup_{t>0} \left\| t^k \frac{\partial^k}{\partial t^k} u(\cdot, t) \right\|_{L^2} = \sup_{t>0} \left\| (t^k L_w^{1/2})^k e^{-tL_w^{1/2}} f(\cdot) \right\|_{L^2} \leq C \|f\|_{L^2}. \tag{12.5}$$

For the regularity problem we have the following.

Theorem 12.6. *Given a weight $w \in A_2$, suppose*

$$1 \leq r_w < \frac{q_+(L_w)}{2} \quad \text{and} \quad s_w > \max \left\{ \left(\frac{2}{r_w} \right)', \frac{n}{2}r_w + 1 \right\}.$$

Then for any $f \in H^1(\mathbb{R}^n)$, we have $u(x, t) = e^{-tL_w^{1/2}} f(x)$ is a solution of (12.1) with convergence to the boundary data as $t \rightarrow 0^+$ in the L^2 -sense. Furthermore, we have

$$\sup_{t>0} \|\nabla_{x,t} u(\cdot, t)\|_{L^2} \leq C \|\nabla f\|_{L^2}. \tag{12.7}$$

In particular, (12.7) holds if we assume that $w \in A_1 \cap \text{RH}_{1+n/2}$. Furthermore, it holds if the following is true: given $\Theta \geq 1$ there exists $\varepsilon_0 = \varepsilon_0(\Theta, n, \Lambda/\lambda)$, $0 < \varepsilon_0 \leq \frac{1}{2n}$, such that $w \in A_{1+\varepsilon} \cap \text{RH}_{\max\{(2/(1+\varepsilon))', (n/2)(1+\varepsilon)+1\}}$, $0 \leq \varepsilon < \varepsilon_0$, and $[w]_{A_2} \leq \Theta$.

For power weights, there exists $\varepsilon_1 = \varepsilon_1(n, \Lambda/\lambda)$, $0 < \varepsilon_1 \leq \frac{1}{2}$, such that (12.7) holds if

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{n}{2} < \alpha < \varepsilon_1.$$

Proof. Arguing as before, it suffices to prove that (12.7) holds. For any $t > 0$ we have, by Theorem 11.11,

$$\begin{aligned} \|\nabla_{x,t}u(\cdot, t)\|_{L^2} &\leq \|\nabla L_w^{-1/2} L_w^{1/2} e^{-tL_w^{1/2}} f\|_{L^2} + \|L_w^{1/2} e^{-tL_w^{1/2}} f\|_{L^2} \\ &\lesssim \|L_w^{1/2} e^{-tL_w^{1/2}} f\|_{L^2} = \|e^{-tL_w^{1/2}} L_w^{1/2} f\|_{L^2} \lesssim \|L_w^{1/2} f\|_{L^2} \lesssim \|\nabla f\|_{L^2}. \quad \square \end{aligned}$$

Note that under the hypothesis of Theorem 12.6, and as observed in Remark 12.4, we have that $u(\cdot, t) = e^{-tL_w^{1/2}} f$ satisfies (12.3) and (12.5). Additionally, from the functional calculus estimates on L^2 it follows that

$$\sup_{t>0} \|t\nabla_{x,t}u(\cdot, t)\|_{L^2} \lesssim \|tL_w^{1/2} e^{-tL_w^{1/2}} f\|_{L^2} \lesssim \|f\|_{L^2}. \tag{12.8}$$

Finally, we consider the Neumann problem

$$\begin{cases} \partial_t^2 u - L_w u = 0 & \text{on } \mathbb{R}^n, \\ \partial_t u|_{\partial\mathbb{R}_+^{n+1}} = f & \text{on } \partial\mathbb{R}_+^{n+1} = \mathbb{R}^n. \end{cases} \tag{12.9}$$

Theorem 12.10. *Given a weight $w \in A_2$, suppose $1 \leq r_w < q_+(L_w)/2$ and $s_w > \frac{n}{2}r_w + 1$. Then for any $f \in L^2(\mathbb{R}^n)$, we have $u(x, t) = -L_w^{-1/2} e^{-tL_w^{1/2}} f(x)$ is a solution of (12.9) with convergence of $\partial_t u(\cdot, t) \rightarrow f$ as $t \rightarrow 0^+$ in the L^2 -sense. Furthermore, we have*

$$\sup_{t>0} \|\nabla_{x,t}u(\cdot, t)\|_{L^2} \leq C \|f\|_{L^2}. \tag{12.11}$$

In particular, (12.11) holds if we assume that $w \in A_1 \cap \text{RH}_{1+n/2}$. Furthermore, it holds if the following is true: given $\Theta \geq 1$ there exists $\varepsilon_0 = \varepsilon_0(\Theta, n, \Lambda/\lambda)$, $0 < \varepsilon_0 \leq \frac{1}{2n}$, such that $w \in A_{1+\varepsilon} \cap \text{RH}_{(n/2)(1+\varepsilon)+1}$, $0 \leq \varepsilon < \varepsilon_0$, and $[w]_{A_2} \leq \Theta$.

For power weights, there exists $\varepsilon_1 = \varepsilon_1(n, \Lambda/\lambda)$, $0 < \varepsilon_1 \leq \frac{1}{2}$, such that (12.11) holds if

$$w_\alpha(x) = |x|^\alpha, \quad -\frac{2n}{n+2} < \alpha < \varepsilon_1.$$

Proof. Again, u is clearly a formal solution of (12.9); see [Kato 1966]. The proof that (12.11) holds is similar to the proof of (12.7):

$$\|\nabla_{x,t}u(\cdot, t)\|_{L^2} \leq \|\nabla L_w^{-1/2} e^{-tL_w^{1/2}} f\|_{L^2} + \|e^{-tL_w^{1/2}} f\|_{L^2} \lesssim \|e^{-tL_w^{1/2}} f\|_{L^2} \lesssim \|f\|_{L^2},$$

where we have used Theorem 11.8 (for the Riesz transform) and Theorem 11.2 (for the functional calculus with $\varphi(z) = e^{-t\sqrt{z}}$). □

Remark 12.12. As we noted in Remark 11.3, we can also get unweighted L^p bounds for these operators for values of p close to 2. As a consequence we can also get estimates for L^p boundary value problems for the same values of p . Details are left to the reader.

Acknowledgements

Cruz-Uribe is supported by NSF grant 1362425 and research funds provided by the Dean of Arts & Sciences at the University of Alabama. While substantial portions of this work were done, he was supported by the Stewart–Dorwart faculty development fund at Trinity College. Martell acknowledges financial support from the Spanish Ministry of Economy and Competitiveness, through the “Severo Ochoa Programme for Centres of Excellence in R&D” (SEV-2015-0554). He also acknowledges that the research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ ERC agreement no. 615112 HAPDEGMT. Rios is supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant RT733901. The authors warmly thank P. Auscher and S. Hofmann for several useful discussions and their many suggestions.

References

- [Alberico et al. 2009] T. Alberico, A. Cianchi, and C. Sbordone, “Fractional integrals and A_p -weights: a sharp estimate”, *C. R. Math. Acad. Sci. Paris* **347**:21-22 (2009), 1265–1270. MR Zbl
- [Auscher 2007] P. Auscher, *On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates*, Mem. Amer. Math. Soc. **871**, American Mathematical Society, Providence, RI, 2007. MR Zbl
- [Auscher and Coulhon 2005] P. Auscher and T. Coulhon, “Riesz transform on manifolds and Poincaré inequalities”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **4**:3 (2005), 531–555. MR Zbl
- [Auscher and Martell 2006] P. Auscher and J. M. Martell, “Weighted norm inequalities, off-diagonal estimates and elliptic operators, III: Harmonic analysis of elliptic operators”, *J. Funct. Anal.* **241**:2 (2006), 703–746. MR Zbl
- [Auscher and Martell 2007a] P. Auscher and J. M. Martell, “Weighted norm inequalities, off-diagonal estimates and elliptic operators, I: General operator theory and weights”, *Adv. Math.* **212**:1 (2007), 225–276. MR Zbl
- [Auscher and Martell 2007b] P. Auscher and J. M. Martell, “Weighted norm inequalities, off-diagonal estimates and elliptic operators, II: Off-diagonal estimates on spaces of homogeneous type”, *J. Evol. Equ.* **7**:2 (2007), 265–316. MR Zbl
- [Auscher and Martell 2008] P. Auscher and J. M. Martell, “Weighted norm inequalities, off-diagonal estimates and elliptic operators, IV: Riesz transforms on manifolds and weights”, *Math. Z.* **260**:3 (2008), 527–539. MR Zbl
- [Auscher and Tchamitchian 1998] P. Auscher and P. Tchamitchian, *Square root problem for divergence operators and related topics*, Astérisque **249**, Société Mathématique de France, Paris, 1998. MR Zbl
- [Auscher et al. 2002] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh, and P. Tchamitchian, “The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n ”, *Ann. of Math. (2)* **156**:2 (2002), 633–654. MR Zbl
- [Auscher et al. 2015] P. Auscher, A. Rosén, and D. Rule, “Boundary value problems for degenerate elliptic equations and systems”, *Ann. Sci. Éc. Norm. Supér. (4)* **48**:4 (2015), 951–1000. MR Zbl
- [Bernicot and Zhao 2008] F. Bernicot and J. Zhao, “New abstract Hardy spaces”, *J. Funct. Anal.* **255**:7 (2008), 1761–1796. MR Zbl
- [Björn and Björn 2011] A. Björn and J. Björn, *Nonlinear potential theory on metric spaces*, EMS Tracts in Mathematics **17**, European Mathematical Society, Zürich, 2011. MR Zbl
- [Chen et al. 2016] L. Chen, J. M. Martell, and C. Prisuelos-Arribas, “Conical square functions for degenerate elliptic operators”, preprint, 2016. To appear in *Adv. Cal. Var.* arXiv
- [Cowling et al. 1996] M. Cowling, I. Doust, A. McIntosh, and A. Yagi, “Banach space operators with a bounded H^∞ functional calculus”, *J. Austral. Math. Soc. Ser. A* **60**:1 (1996), 51–89. MR Zbl
- [Cruz-Uribe and Neugebauer 1995] D. Cruz-Uribe and C. J. Neugebauer, “The structure of the reverse Hölder classes”, *Trans. Amer. Math. Soc.* **347**:8 (1995), 2941–2960. MR Zbl

- [Cruz-Uribe and Rios 2008] D. Cruz-Uribe and C. Rios, “Gaussian bounds for degenerate parabolic equations”, *J. Funct. Anal.* **255**:2 (2008), 283–312. Correction in **267**:9 (2014), 3507–3513. MR Zbl
- [Cruz-Uribe and Rios 2012] D. Cruz-Uribe and C. Rios, “The solution of the Kato problem for degenerate elliptic operators with Gaussian bounds”, *Trans. Amer. Math. Soc.* **364**:7 (2012), 3449–3478. MR Zbl
- [Cruz-Uribe and Rios 2015] D. Cruz-Uribe and C. Rios, “The Kato problem for operators with weighted ellipticity”, *Trans. Amer. Math. Soc.* **367**:7 (2015), 4727–4756. MR Zbl
- [Duoandikoetxea 2001] J. Duoandikoetxea, *Fourier analysis*, Graduate Studies in Mathematics **29**, American Mathematical Society, Providence, RI, 2001. MR Zbl
- [Fabes et al. 1982] E. B. Fabes, C. E. Kenig, and R. P. Serapioni, “The local regularity of solutions of degenerate elliptic equations”, *Comm. Partial Differential Equations* **7**:1 (1982), 77–116. MR Zbl
- [García-Cuerva and Rubio de Francia 1985] J. García-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Mathematics Studies **116**, North-Holland, Amsterdam, 1985. MR Zbl
- [Haase 2006] M. Haase, *The functional calculus for sectorial operators*, Operator Theory: Advances and Applications **169**, Birkhäuser, Basel, 2006. MR Zbl
- [Hofmann et al. 2015] S. Hofmann, P. Le, and A. Morris, “Carleson measure estimates and the Dirichlet problem for degenerate elliptic equations”, preprint, 2015.
- [Kato 1966] T. Kato, *Perturbation theory for linear operators*, Die Grundlehren der mathematischen Wissenschaften **132**, Springer, 1966. MR Zbl
- [Le 2015] P. Le, “ L^p bounds of Riesz transform and vertical square functions for degenerate elliptic operators”, preprint, 2015.
- [McIntosh 1986] A. McIntosh, “Operators which have an H_∞ functional calculus”, pp. 210–231 in *Miniconference on operator theory and partial differential equations* (North Ryde, 1986), edited by B. Jefferies et al., Proc. Centre Math. Anal. Austral. Nat. Univ. **14**, Austral. Nat. Univ., Canberra, 1986. MR Zbl
- [Miller 1982] N. Miller, “Weighted Sobolev spaces and pseudodifferential operators with smooth symbols”, *Trans. Amer. Math. Soc.* **269**:1 (1982), 91–109. MR Zbl
- [Pérez 1999] C. Pérez, “Calderón–Zygmund theory related to Poincaré–Sobolev inequalities, fractional integrals and singular integral operators”, in *Function spaces: lectures notes of the spring school on analysis* (Paseky nad Jizerou, Czech Republic, 1999), edited by J. Lukes and L. Pick, Matfyzpress, Prague, 1999.
- [Yang and Zhang 2017] D. Yang and J. Zhang, “Weighted L^p estimates of Kato square roots associated to degenerate elliptic operators”, *Publ. Mat.* **61**:2 (2017), 395–444. MR Zbl

Received 6 Oct 2016. Revised 6 Jun 2017. Accepted 20 Sep 2017.

DAVID CRUZ-URIBE: dcruzuribe@ua.edu

Department of Mathematics, University of Alabama, Tuscaloosa, AL, United States

JOSÉ MARÍA MARTELL: chema.martell@icmat.es

Instituto de Ciencias Matemáticas CSIC-UAM-UC3M-UCM, Consejo Superior de Investigaciones Científicas, Madrid, Spain

and

Department of Mathematics, University of Missouri, Columbia, MO, USA

martellj@missouri.edu

CRISTIAN RIOS: crios@ucalgary.ca

Department of Mathematics and Statistics, University of Calgary, Canada

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 3 2018

The endpoint perturbed Brascamp–Lieb inequalities with examples RUIXIANG ZHANG	555
Square function estimates for discrete Radon transforms MARIUSZ MIREK	583
On the Kato problem and extensions for degenerate elliptic operators DAVID CRUZ-URIBE, JOSÉ MARÍA MARTELL and CRISTIAN RIOS	609
Small data global regularity for half-wave maps JOACHIM KRIEGER and YANNICK SIRE	661
The semigroup generated by the Dirichlet Laplacian of fractional order TSUKASA IWABUCHI	683
Klein’s paradox and the relativistic δ -shell interaction in \mathbb{R}^3 ALBERT MAS and FABIO PIZZICHILLO	705
Dimension-free L^p estimates for vectors of Riesz transforms associated with orthogonal ex- pansions BŁAŻEJ WRÓBEL	745
Reducibility of the quantum harmonic oscillator in d -dimensions with polynomial time-dependent perturbation DARIO BAMBUSI, BENOÎT GRÉBERT, ALBERTO MASPERO and DIDIER ROBERT	775
Eigenfunction scarring and improvements in L^∞ bounds JEFFREY GALKOWSKI and JOHN A. TOTH	801



2157-5045(2018)11:3;1-R