

ANALYSIS & PDE

Volume 11

No. 4

2018

MATTEO BONFORTE, ALESSIO FIGALLI AND JUAN LUIS VÁZQUEZ

**SHARP GLOBAL ESTIMATES FOR LOCAL AND NONLOCAL
POROUS MEDIUM-TYPE EQUATIONS IN BOUNDED DOMAINS**

SHARP GLOBAL ESTIMATES FOR LOCAL AND NONLOCAL POROUS MEDIUM-TYPE EQUATIONS IN BOUNDED DOMAINS

MATTEO BONFORTE, ALESSIO FIGALLI AND JUAN LUIS VÁZQUEZ

We provide a quantitative study of nonnegative solutions to nonlinear diffusion equations of porous medium-type of the form $\partial_t u + \mathcal{L}u^m = 0$, $m > 1$, where the operator \mathcal{L} belongs to a general class of linear operators, and the equation is posed in a bounded domain $\Omega \subset \mathbb{R}^N$. As possible operators we include the three most common definitions of the fractional Laplacian in a bounded domain with zero Dirichlet conditions, and also a number of other nonlocal versions. In particular, \mathcal{L} can be a fractional power of a uniformly elliptic operator with C^1 coefficients. Since the nonlinearity is given by u^m with $m > 1$, the equation is degenerate parabolic.

The basic well-posedness theory for this class of equations was recently developed by Bonforte and Vázquez (2015, 2016). Here we address the regularity theory: decay and positivity, boundary behavior, Harnack inequalities, interior and boundary regularity, and asymptotic behavior. All this is done in a quantitative way, based on sharp a priori estimates. Although our focus is on the fractional models, our results cover also the local case when \mathcal{L} is a uniformly elliptic operator, and provide new estimates even in this setting.

A surprising aspect discovered in this paper is the possible presence of nonmatching powers for the long-time boundary behavior. More precisely, when $\mathcal{L} = (-\Delta)^s$ is a spectral power of the Dirichlet Laplacian inside a smooth domain, we can prove that

- when $2s > 1 - 1/m$, for large times all solutions behave as $\text{dist}^{1/m}$ near the boundary;
- when $2s \leq 1 - 1/m$, different solutions may exhibit different boundary behavior.

This unexpected phenomenon is a completely new feature of the nonlocal nonlinear structure of this model, and it is not present in the semilinear elliptic equation $\mathcal{L}u^m = u$.

1. Introduction	946
2. General class of operators and their kernels	949
3. Reminders about weak dual solutions	954
4. Upper boundary estimates	955
5. Lower bounds	959
6. Summary of the general decay and boundary results	968
7. Asymptotic behavior	970
8. Regularity results	971
9. Numerical evidence	976
10. Complements, extensions and further examples	978
Acknowledgments	980
References	980

MSC2010: 35B45, 35B65, 35K55, 35K65.

Keywords: nonlocal diffusion, nonlinear equations, bounded domains, a priori estimates, positivity, boundary behavior, regularity, Harnack inequalities.

1. Introduction

In this paper we address the question of obtaining a priori estimates, positivity, boundary behavior, Harnack inequalities, and regularity for a suitable class of weak solutions of nonlinear nonlocal diffusion equations of the form

$$\partial_t u + \mathcal{L}F(u) = 0 \quad \text{posed in } Q = (0, \infty) \times \Omega, \quad (1-1)$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with $C^{1,1}$ boundary, $N \geq 2$,¹ and \mathcal{L} is a linear operator representing diffusion of local or nonlocal type, the prototype example being the fractional Laplacian (the class of admissible operators will be precisely described below). Although our arguments hold for a rather general class of nonlinearities $F : \mathbb{R} \rightarrow \mathbb{R}$, for the sake of simplicity we shall focus on the model case $F(u) = u^m$ with $m > 1$.

The use of nonlocal operators in diffusion equations reflects the need to model the presence of long-distance effects not included in evolution driven by the Laplace operator, and this is well documented in the literature. The physical motivation and relevance of the nonlinear diffusion models with nonlocal operators has been mentioned in many references; see for instance [Athanasopoulos and Caffarelli 2010; Bonforte and Vázquez 2014; 2015; de Pablo et al. 2011; 2012; Vázquez 2014b]. Because u usually represents a density, all data and solutions are supposed to be nonnegative. Since the problem is posed in a bounded domain, we need boundary or external conditions that we assume to be of Dirichlet type.

This kind of problem has been extensively studied when $\mathcal{L} = -\Delta$ and $F(u) = u^m$, $m > 1$, in which case the equation becomes the classical porous medium equation [Vázquez 2004; 2007; Dahlberg and Kenig 1988; Daskalopoulos and Kenig 2007]. Here, we are interested in treating nonlocal diffusion operators, in particular fractional Laplacian operators. Note that, since we are working on a bounded domain, the concept of fractional Laplacian operator admits several nonequivalent versions, the best known being the restricted fractional Laplacian (RFL), the spectral fractional Laplacian (SFL), and the censored fractional Laplacian (CFL); see Section 2A for more details. We use these names because they already appeared in some previous works [Bonforte et al. 2015; Bonforte and Vázquez 2016], but we point out that the RFL is usually known as the standard fractional Laplacian, or plainly fractional Laplacian, and the CFL is often called the regional fractional Laplacian.

The case of the SFL operator with $F(u) = u^m$, $m > 1$, was already studied by the first and third authors in [Bonforte and Vázquez 2015; 2016]. In particular, in [Bonforte and Vázquez 2016] the authors presented a rather abstract setting where they were able to treat not only the usual fractional Laplacians but also a large number of variants that will be listed below for the reader's convenience. Besides, rather general increasing nonlinearities F were allowed. The basic questions of existence and uniqueness of suitable solutions for this problem were solved in [Bonforte and Vázquez 2016] in the class of “weak dual solutions”, an extension of the concept of solution introduced in [Bonforte and Vázquez 2015] that has proved to be quite flexible and efficient. A number of a priori estimates (absolute bounds and smoothing effects) were also derived in that generality.

¹Our results work also in dimension $N = 1$ if the fractional exponent (that we shall introduce later) belongs to the range $0 < s < \frac{1}{2}$. The interval $\frac{1}{2} \leq s < 1$ requires some minor modifications that we prefer to avoid in this paper.

Since these basic facts are settled, here we focus our attention on the finer aspects of the theory, mainly sharp boundary estimates and decay estimates. Such upper and lower bounds will be formulated in terms of the first eigenfunction Φ_1 of \mathcal{L} , which under our assumptions will satisfy $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)^\gamma$ for a certain characteristic power $\gamma \in (0, 1]$ that depends on the particular operator we consider. Typical values are $\gamma = s$ (SFL), $\gamma = 1$ (RFL), and $\gamma = s - \frac{1}{2}$ for $s > \frac{1}{2}$ (CFL); see Sections 2A and 10A. As a consequence, we get various kinds of local and global Harnack-type inequalities.

It is worth mentioning that some of the boundary estimates that we obtain for the parabolic case are essentially elliptic in nature. The study of this issue for stationary problems is done in a companion paper [Bonforte et al. 2017b]. This has the advantage that many arguments are clearer, since the parabolic problem is more complicated than the elliptic one. Clarifying such differences is one of the main contributions of our present work.

Thanks to these results, in the last part of the paper we are able to prove both interior and boundary regularity, and to find the large-time asymptotic behavior of solutions.

Let us indicate here some notation of general use. The symbol ∞ will always denote $+\infty$. Given a, b , we use the notation $a \asymp b$ whenever there exist universal constants $c_0, c_1 > 0$ such that $c_0 b \leq a \leq c_1 b$. We also use the symbols $a \vee b = \max\{a, b\}$ and $a \wedge b = \min\{a, b\}$. We will always consider bounded domains Ω with boundary of class C^2 . In the paper we use the short form “solution” to mean “weak dual solution”, unless differently stated.

1A. Presentation of the results on sharp boundary behavior. • A basic principle in the paper is that the sharp boundary estimates depend not only on \mathcal{L} but also on the behavior of the nonlinearity $F(u)$ near $u = 0$, i.e., in our case, on the exponent $m > 1$. The elliptic analysis performed in the companion paper [Bonforte et al. 2017b] combined with some standard arguments will allow us to prove that, in *all* cases, $u(t)$ approaches the separate-variable solution $\mathcal{U}(x, t) = t^{-1/(m-1)}S(x)$ in the sense that

$$\|t^{1/(m-1)}u(t, \cdot) - S\|_{L^\infty(\Omega)} \xrightarrow{t \rightarrow \infty} 0, \quad (1-2)$$

where S is the solution of the elliptic problem (see Theorems 3.2 and 7.1). The behavior of the profile $S(x)$ is shown to be, when $2sm \neq \gamma(m-1)$,

$$S(x) \asymp \Phi_1(x)^{\sigma/m}, \quad \sigma := \min\left\{1, \frac{2sm}{\gamma(m-1)}\right\}. \quad (1-3)$$

Thus, the behavior strongly depends on the new parameter σ , more precisely, on whether this parameter is equal to 1 or less than 1. As we shall see later, σ encodes the interplay between the “elliptic scaling power” $2s/(m-1)$, the “eigenfunction power” γ , and the “nonlinearity power” m . When $2sm = \gamma(m-1)$ we have $\sigma = 1$, but a logarithmic correction appears:

$$S(x) \asymp \Phi_1(x)^{1/m} (1 + |\log \Phi_1(x)|)^{1/(m-1)}. \quad (1-4)$$

• This fact and the results in [Bonforte et al. 2017a] prompted us to look for estimates of the form

$$c_0(t) \frac{\Phi_1^{\sigma/m}(x_0)}{t^{1/(m-1)}} \leq u(t, x_0) \leq c_1 \frac{\Phi_1^{\sigma/m}(x_0)}{t^{1/(m-1)}} \quad \text{for all } t > 0, x_0 \in \Omega, \quad (1-5)$$

where $c_0(t)$ and c_1 are positive and independent of u , eventually with a logarithmic term appearing when $2sm = \gamma(m - 1)$, as in (1-4). We will prove in this paper that the upper bound holds for the three mentioned fractional Laplacian choices, and indeed for the whole class of integrodifferential operators we will introduce below; see [Theorem 4.1](#). Also, separate-variable solutions saturate the upper bound.

The issue of the validity of a lower bound as in (1-5) is instead much more elusive. A first indication for this is the introduction of a function $c_0(t)$ depending on t , instead of a constant. This seems to reflect the fact that the solution may take some time to reach the boundary behavior that is expected to hold uniformly for large times. Indeed, recall that in the classical PME [[Aronson and Peletier 1981](#); [Vázquez 2004](#); [2007](#)], for data supported away from the boundary, some “waiting time” is needed for the support to reach the boundary.

- As proved in [[Bonforte et al. 2017a](#)], the stated lower bound holds for the RFL with $c_0(t) \sim (1 \wedge t)^{m/(m-1)}$. In particular, in this nonlocal setting, infinite speed of propagation holds. Here, we show that this holds also for the CFL and a number of other operators; see [Theorem 5.2](#). Note that for the RFL and the CFL we have $2sm > \gamma(m - 1)$, in particular $\sigma = 1$, which simplifies formula (1-5).

A combination of an upper and a lower bound with matching behavior (with respect to x and t) will be called a *global Harnack principle*, and holds for all $t > 0$ for these operators; see [Theorems 6.1](#) and [6.2](#).

- When \mathcal{L} is the SFL, we shall see that the lower bound may fail. Of course, solutions by separation of variables satisfy the matching estimates in (1-5), eventually with an extra logarithmic term in the limit case, as in (1-4), but it came as a complete surprise to us that for the SFL the situation is not the same for “small” initial data. More precisely:

(i) We can prove that the following bounds always hold for all times:

$$c_0 \left(1 \wedge \frac{t}{t_*} \right)^{m/(m-1)} \frac{\Phi_1(x_0)}{t^{1/(m-1)}} \leq u(t, x_0) \leq c_1 \frac{\Phi_1^{\sigma/m}(x_0)}{t^{1/(m-1)}} \tag{1-6}$$

(when $2sm = \gamma(m - 1)$, a logarithmic correction $(1 + |\log \Phi_1(x)|)^{1/(m-1)}$ appears in the right-hand side); see [Theorem 5.1](#). These are nonmatching estimates.

(ii) For $2sm > \gamma(m - 1)$, the sharp estimate (1-5) holds for any nonnegative nontrivial solution for large times $t \geq t_*$; see [Theorem 5.3](#).

(iii) *Anomalous boundary behavior.* Consider now the SFL with $\sigma < 1$ (resp. $2sm = \gamma(m - 1)$).² In this case we can find initial data for which the upper bound in (1-6) is not sharp. Depending on the initial data, there are several possible rates for the long-time behavior near the boundary. More precisely:

(a) When $u_0 \leq A\Phi_1$, we have $u(t) \leq F(t)\Phi_1^{1/m} \ll \Phi_1^{\sigma/m}$ (resp. $\Phi_1^{1/m} \ll \Phi_1^{1/m}(1 + |\log \Phi_1|)^{1/(m-1)}$) for all times; see [Theorem 5.4](#). In particular

$$\lim_{x \rightarrow \partial\Omega} \frac{u(t, x)}{\Phi_1(x)^{\sigma/m}} = 0 \quad \left(\text{resp. } \lim_{x \rightarrow \partial\Omega} \frac{u(t, x)}{\Phi_1(x)^{1/m}(1 + |\log \Phi_1(x)|)^{1/(m-1)}} = 0 \right) \tag{1-7}$$

for any $t > 0$.

²Since for the SFL $\gamma = 1$, we have $\sigma < 1$ if and only if $0 < s < s_* := (m - 1)/(2m) < \frac{1}{2}$. Note that $s_* \rightarrow 0$ as we tend to the linear case $m = 1$, so this exceptional regime does not appear for linear diffusions, both fractional and standard.

- (b) When $u_0 \leq A\Phi_1^{1-2s/\gamma}$, we have $u(t) \leq F(t)\Phi_1^{1-2s/\gamma}$ for small times; see [Theorem 4.4](#). Notice that when $\sigma < 1$ we have always $1 - 2s/\gamma > \sigma/m$. This sets a limitation on the improvement of the lower bound, which is confirmed by another result: in [Theorem 5.5](#) we show that lower bounds of the form $u(T, x) \geq \kappa\Phi_1^\alpha(x)$ for data $u_0(x) \leq A\Phi_1(x)$ are possible only for $\alpha \geq 1 - 2s/\gamma$.
- (c) On the other hand, for “large” initial data, [Theorem 6.2](#) shows that the desired matching estimates from above and below hold.

After discovering this strange boundary behavior, we looked for numerical confirmation. In [Section 9](#) we will explain the numerical results obtained in [[Cusimano et al. 2017](#)]. Note that, if one looks for universal bounds independent of the initial condition, [Figures 2–3](#) below seem to suggest that the bounds provided by (1-6) are optimal for all times and all operators.

- The current interest in more general types of nonlocal operators led us to a more general analysis where the just-explained alternative has been extended to a wide class of integrodifferential operators, subject only to a list of properties that we call (A1), (A2), (L1), (L2), (K2), (K4); a number of examples are explained in [Section 2](#). These general classes appear also in the study of the elliptic problem [[Bonforte et al. 2017b](#)].

1B. Asymptotic behavior and regularity. Our quantitative lower and upper estimates admit formulations as local or global Harnack inequalities. They are used at the end of the paper to settle two important issues.

Sharp asymptotic behavior. Exploiting the techniques in [[Bonforte et al. 2015](#)], we can prove a sharp asymptotic behavior for our nonnegative and nontrivial solutions when the upper and lower bounds have matching powers. Such sharp results hold true for a quite general class of local and nonlocal operators. A detailed account is given in [Section 7](#).

Regularity. By a variant of the techniques used in [[Bonforte et al. 2017a](#)], we can show interior Hölder regularity. In addition, if the kernel of the operator satisfies some suitable continuity assumptions, we show that solutions are classical in the interior and are Hölder continuous up to the boundary if the upper and lower bounds have matching powers. We refer to [Section 8](#) for details.

2. General class of operators and their kernels

The interest of the theory developed here lies both in the sharpness of the results and in the wide range of applicability. We have just mentioned the most relevant examples appearing in the literature, and more are listed at the end of this section. Actually, our theory applies to a general class of operators with definite assumptions, and this is what we want to explain now.

Let us present the properties that have to be assumed on the class of admissible operators. Some of them already appeared in [[Bonforte and Vázquez 2016](#)]. However, to further develop our theory, more hypotheses need to be introduced. In particular, while the paper above only uses the properties of the Green function, here we shall make some assumptions also on the kernel of \mathcal{L} (whenever it exists). Note that assumptions on the kernel K of \mathcal{L} are needed for the positivity results, because we need to distinguish between the local and nonlocal cases. The study of the kernel K is performed in [Section 10](#).

For convenience of reference, the list of used assumptions is (A1), (A2), (K2), (K4), (L1), (L2). The first three are assumed in all operators \mathcal{L} that we use.

- *Basic assumptions on \mathcal{L} .* The linear operator $\mathcal{L} : \text{dom}(\mathcal{L}) \subseteq L^1(\Omega) \rightarrow L^1(\Omega)$ is assumed to be densely defined and sub-Markovian; more precisely, it satisfies (A1) and (A2) below:

(A1) \mathcal{L} is m -accretive on $L^1(\Omega)$;

(A2) If $0 \leq f \leq 1$ then $0 \leq e^{-t\mathcal{L}} f \leq 1$.

Under these assumptions, in [Bonforte and Vázquez 2016], the first and third authors proved existence, uniqueness, weighted estimates, and smoothing effects.

- *Assumptions on the kernel.* Whenever \mathcal{L} is defined in terms of a kernel $K(x, y)$ via the formula

$$\mathcal{L}f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y)) K(x, y) dy,$$

assumption (L1) states that there exists $\underline{\kappa}_\Omega > 0$ such that

$$\inf_{x, y \in \Omega} K(x, y) \geq \underline{\kappa}_\Omega > 0. \tag{L1}$$

We note that condition holds both for the RFL and the CFL; see Section 2A.

Whenever \mathcal{L} is defined in terms of a kernel $K(x, y)$ and a zero-order term via the formula

$$\mathcal{L}f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y)) K(x, y) dy + B(x)f(x),$$

assumption (L2) states that

$$K(x, y) \geq c_0 \delta^\gamma(x) \delta^\gamma(y), \quad c_0 > 0, \quad \text{and} \quad B(x) \geq 0, \tag{L2}$$

where, from now on, we adopt the notation $\delta(x) := \text{dist}(x, \partial\Omega)$. This condition is satisfied by the SFL in a stronger form; see Section 10 and Lemma 10.1.

- *Assumptions on \mathcal{L}^{-1} .* In order to prove our quantitative estimates, we need to be more specific about the operator \mathcal{L} . Besides satisfying (A1) and (A2), we will assume that it has a left-inverse $\mathcal{L}^{-1} : L^1(\Omega) \rightarrow L^1(\Omega)$ that can be represented by a kernel \mathbb{G} (the letter “G” standing for Green function) as

$$\mathcal{L}^{-1}[f](x) = \int_{\Omega} \mathbb{G}(x, y) f(y) dy,$$

where \mathbb{G} satisfies the following assumption for some $s \in (0, 1]$: there exist constants $\gamma \in (0, 1]$ and $c_{0,\Omega}, c_{1,\Omega} > 0$ such that, for a.e. $x, y \in \Omega$,

$$c_{0,\Omega} \delta^\gamma(x) \delta^\gamma(y) \leq \mathbb{G}(x, y) \leq \frac{c_{1,\Omega}}{|x - y|^{N-2s}} \left(\frac{\delta^\gamma(x)}{|x - y|^\gamma} \wedge 1 \right) \left(\frac{\delta^\gamma(y)}{|x - y|^\gamma} \wedge 1 \right). \tag{K2}$$

(Here and below we use the labels (K2) and (K4) to be consistent with the notation in [Bonforte and Vázquez 2016].) Hypothesis (K2) introduces an exponent γ which is a characteristic of the operator and will play a big role in the results. Notice that defining an inverse operator \mathcal{L}^{-1} implies that we are taking into account the Dirichlet boundary conditions. See more details in Section 2 of [Bonforte and Vázquez 2016].

The lower bound in (K2) is weaker than the known bounds on the Green function for many examples under consideration; indeed, the following stronger estimate holds in many cases:

$$\mathbb{G}(x, y) \asymp \frac{1}{|x - y|^{N-2s}} \left(\frac{\delta^\gamma(x)}{|x - y|^\gamma} \wedge 1 \right) \left(\frac{\delta^\gamma(y)}{|x - y|^\gamma} \wedge 1 \right). \quad (\text{K4})$$

Remarks. (i) The labels (A1), (A2), (K1), (K2), (K4) are consistent with the notation in [Bonforte and Vázquez 2016]. The label (K3) was used to mean hypothesis (K2) written in terms of Φ_1 instead of δ^γ .

(ii) In the classical local case $\mathcal{L} = -\Delta$, the Green function \mathbb{G} satisfies (K4) only when $N \geq 3$, as the formulas slightly change when $N = 1, 2$. In the fractional case $s \in (0, 1)$ the same problem arises when $N = 1$ and $s \in [\frac{1}{2}, 1)$. Hence, treating also these cases would require a slightly different analysis based on different but related assumptions on \mathbb{G} . Since our approach is very general, we expect it to work also in these remaining cases without any major difficulties. However, to simplify the presentation, from now on we assume that

$$\text{either } N \geq 2 \text{ and } s \in (0, 1) \quad \text{or} \quad N = 1 \text{ and } s \in (0, \frac{1}{2}).$$

The role of the first eigenfunction of \mathcal{L} . We showed in [Bonforte et al. 2017b] that, under assumption (K1), the operator \mathcal{L} is compact, has a discrete spectrum, and has a first nonnegative bounded eigenfunction Φ_1 ; assuming also (K2), we have

$$\Phi_1(x) \asymp \delta^\gamma(x) = \text{dist}(x, \partial\Omega)^\gamma \quad \text{for all } x \in \bar{\Omega}. \quad (2-1)$$

Hence, Φ_1 encodes the parameter γ that takes care of describing the boundary behavior. We recall that we are assuming that the boundary of Ω is smooth enough, for instance $C^{1,1}$.

Remark. We note that our assumptions allow us to cover all the examples of operators described in Sections 2A and 10A.

2A. Main examples of operators and properties. When working in the whole \mathbb{R}^N , the fractional Laplacian admits different definitions that can be shown to be all equivalent. On the other hand, when we deal with bounded domains, there are at least three different operators in the literature, which we call the restricted (RFL), the spectral (SFL) and the censored fractional Laplacian (CFL). We will show below that these different operators exhibit quite different behaviors, so the distinction between them has to be taken into account. Let us present the statement and results for the three model cases, and we refer to Section 10A for further examples. Here, we collect the sharp results about the boundary behavior, namely the global Harnack inequalities from Theorems 6.1, 6.2, and 6.3.

The parameters γ and σ . The strong difference between the various operators \mathcal{L} is reflected in the different boundary behavior of their nonnegative solutions. We will often use the exponent γ , which represents the boundary behavior of the first eigenfunction $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)^\gamma$; see [Bonforte et al. 2017b]. Both in the parabolic theory of this paper and the elliptic theory of [Bonforte et al. 2017b] the parameter $\sigma = \min\{1, 2sm/(\gamma(m-1))\}$ introduced in (1-3) plays a big role.

2A1. The RFL. We define the fractional Laplacian operator acting on a bounded domain by using the integral representation on the whole space in terms of a hypersingular kernel; namely

$$(-\Delta_{\mathbb{R}^N})^s g(x) = c_{N,s} \text{P.V.} \int_{\mathbb{R}^N} \frac{g(x) - g(z)}{|x - z|^{N+2s}} dz, \tag{2-2}$$

where $c_{N,s} > 0$ is a normalization constant, and we “restrict” the operator to functions that are zero outside Ω . We denote such operator by $\mathcal{L} = (-\Delta|_{\Omega})^s$, and call it the *restricted fractional Laplacian*³ (RFL). The initial and boundary conditions associated to the fractional diffusion equation (1-1) are $u(t, x) = 0$ in $(0, \infty) \times \mathbb{R}^N \setminus \Omega$ and $u(0, \cdot) = u_0$. As explained in [Bonforte et al. 2015], such boundary conditions can also be understood via the Caffarelli–Silvestre extension [2007]. The sharp expression of the boundary behavior for the RFL was investigated in [Ros-Oton and Serra 2014]. We refer to [Bonforte et al. 2015] for a careful construction of the RFL in the framework of fractional Sobolev spaces, and [Blumenthal and Gettoor 1960] for a probabilistic interpretation.

This operator satisfies the assumptions (A1), (A2), (L1), and also (K2) and (K4) with $\gamma = s < 1$. Let us present our results in this case. Note that we have $\sigma = 1$ for all $0 < s < 1$, and Theorem 6.1 shows the sharp boundary behavior for all times; namely for all $t > 0$ and a.e. $x \in \Omega$ we have

$$\underline{\kappa} \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\text{dist}(x, \partial\Omega)^{s/m}}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\text{dist}(x, \partial\Omega)^{s/m}}{t^{1/(m-1)}}. \tag{2-3}$$

The critical time t_* is given by a weighted L^1 norm; namely

$$t_* := \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)},$$

where $\kappa_* > 0$ is a universal constant. Moreover, solutions are classical in the interior and we prove sharp Hölder continuity up to the boundary. These regularity results were first obtained in [Bonforte et al. 2017a]; we give here different proofs valid in the more general setting of this paper. See Section 8 for further details.

2A2. The SFL. Starting from the classical Dirichlet Laplacian Δ_{Ω} on the domain Ω , the so-called *spectral definition* of the fractional power of Δ_{Ω} may be defined via a formula in terms of the semigroup associated to the Laplacian; namely

$$(-\Delta_{\Omega})^s g(x) = \frac{1}{\Gamma(-s)} \int_0^{\infty} (e^{t\Delta_{\Omega}} g(x) - g(x)) \frac{dt}{t^{1+s}} = \sum_{j=1}^{\infty} \lambda_j^s \hat{g}_j \phi_j(x), \tag{2-4}$$

where (λ_j, ϕ_j) , $j = 1, 2, \dots$, is the normalized spectral sequence of the standard Dirichlet Laplacian on Ω , $\hat{g}_j = \int_{\Omega} g(x) \phi_j(x) dx$, and $\|\phi_j\|_{L^2(\Omega)} = 1$. We denote this operator by $\mathcal{L} = (-\Delta_{\Omega})^s$, and call it the *spectral fractional Laplacian* (SFL) as in [Cabr e and Tan 2010]. The initial and boundary conditions associated to the fractional diffusion equation (1-1) are $u(t, x) = 0$ on $(0, \infty) \times \partial\Omega$ and $u(0, \cdot) = u_0$.

³In the literature this is often called the fractional Laplacian on domains, but this simpler name may be confusing when the spectral fractional Laplacian is also considered; see [Bonforte and V azquez 2015]. As discussed in this paper, there are other natural versions.

Such boundary conditions can also be understood via the Caffarelli–Silvestre extension; see [Bonforte et al. 2015]. Following ideas of [Song and Vondraček 2003], we use the fact that this operator admits a kernel representation,

$$(-\Delta_\Omega)^s g(x) = c_{N,s} \text{P.V.} \int_\Omega [g(x) - g(z)]K(x, z) dz + B(x)g(x), \quad (2-5)$$

where K is a singular and compactly supported kernel, which degenerates at the boundary, and $B \asymp \text{dist}(\cdot, \partial\Omega)^{-2s}$ (see [Song and Vondraček 2003] or Lemma 10.1 for further details). This operator satisfies the assumptions (A1), (A2), (L2), and also (K2) and (K4) with $\gamma = 1$. Therefore, σ can be less than 1, depending on the values of s and m .

As we shall see, in our parabolic setting, the degeneracy of the kernel is responsible for a peculiar change of the boundary behavior of the solutions (with respect to the previous case) for small and large times. Here, the lower bounds change both for short and large times, and they strongly depend on σ and on u_0 : we called this phenomenon *anomalous boundary behavior* in Section 1A. More precisely, Theorem 6.3 shows that for all $t > 0$ and all $x \in \Omega$ we have

$$\underline{\kappa} \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\text{dist}(x, \partial\Omega)}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\text{dist}(x, \partial\Omega)^{\sigma/m}}{t^{1/(m-1)}} \quad (2-6)$$

(when $2sm = \gamma(m-1)$, a logarithmic correction $(1 + |\log \Phi_1(x)|)^{1/(m-1)}$ appears in the right-hand side). Such lower behavior is somehow minimal, in the sense that it holds in all cases. The basic asymptotic result (see (1-2) or Theorem 7.1) suggests that the lower bound in (2-6) could be improved by replacing $\text{dist}(x, \partial\Omega)$ with $\text{dist}(x, \partial\Omega)^{\sigma/m}$, at least for large times. This is shown to be true for $\sigma = 1$ (see Theorem 5.3), but it is false for $\sigma < 1$ (see Theorem 5.4), since there are “small” solutions with nonmatching boundary behavior for all times; see (1-7).

It is interesting that, in this case, one can appreciate the interplay between the “elliptic scaling power” $2s/(m-1)$ related to the invariance of the equation $\mathcal{L}S^m = S$ under the scaling $S(x) \mapsto \lambda^{-2s/(m-1)}S(\lambda x)$, the “eigenfunction power” $\gamma = 1$, and the “nonlinearity power” m , made clear through the parameter σ/m . Also in this case, thanks to the strict positivity in the interior, we can show interior space-time regularity of solutions, as well as sharp boundary Hölder regularity for large times whenever upper and lower bounds match.

2A3. The CFL. In the simplest case, the infinitesimal operator of the censored stochastic processes has the form

$$\mathcal{L}g(x) = \text{P.V.} \int_\Omega \frac{g(x) - g(y)}{|x - y|^{N+2s}} dy, \quad \text{with } \frac{1}{2} < s < 1. \quad (2-7)$$

This operator was introduced in [Bogdan et al. 2003] (see also [Chen et al. 2010] and [Bonforte and Vázquez 2016] for further details and references).

In this case $\gamma = s - \frac{1}{2} < 2s$; hence $\sigma = 1$ for all $\frac{1}{2} < s < 1$, and Theorem 6.1 shows that for all $t > 0$ and $x \in \Omega$ we have

$$\underline{\kappa} \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\text{dist}(x, \partial\Omega)^{(s-1/2)/m}}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\text{dist}(x, \partial\Omega)^{(s-1/2)/m}}{t^{1/(m-1)}}.$$

Again, we have interior space-time regularity of solutions, as well as sharp boundary Hölder regularity for all times.

2A4. Other examples. There a number of examples to which our theory applies, besides the RFL, CFL and SFL, since they satisfy the list of assumptions listed in the previous section. Some are listed in the last Section 10; see more detail in [Bonforte and Vázquez 2016].

3. Reminders about weak dual solutions

We denote by $L^p_{\Phi_1}(\Omega)$ the weighted L^p space $L^p(\Omega, \Phi_1 dx)$, endowed with the norm

$$\|f\|_{L^p_{\Phi_1}(\Omega)} = \left(\int_{\Omega} |f(x)|^p \Phi_1(x) dx \right)^{1/p}.$$

Weak dual solutions: existence and uniqueness. We recall the definition of weak dual solutions used in [Bonforte and Vázquez 2016]. This is expressed in terms of the inverse operator \mathcal{L}^{-1} , and encodes the Dirichlet boundary condition. This is needed to build a theory of bounded nonnegative unique solutions to (1-1) under the assumptions of the previous section. Note that in [Bonforte and Vázquez 2016] we used the setup with the weight $\delta^\gamma = \text{dist}(\cdot, \partial\Omega)^\gamma$, but the same arguments generalize immediately to the weight Φ_1 ; indeed under assumption (K2), these two setups are equivalent.

Definition 3.1. A function u is a *weak dual* solution to the Dirichlet problem for (1-1) in $(0, \infty) \times \Omega$ if:

- $u \in C((0, \infty) : L^1_{\Phi_1}(\Omega))$ and $u^m \in L^1((0, \infty) : L^1_{\Phi_1}(\Omega))$.
- The identity

$$\int_0^\infty \int_{\Omega} \mathcal{L}^{-1}u \frac{\partial \psi}{\partial t} dx dt - \int_0^\infty \int_{\Omega} u^m \psi dx dt = 0 \tag{3-1}$$

holds for every test function ψ such that $\psi/\Phi_1 \in C^1_c((0, \infty) : L^\infty(\Omega))$.

- A weak dual solution to the Cauchy–Dirichlet problem (CDP) is a weak dual solution to the homogeneous Dirichlet problem for (1-1) such that $u \in C([0, \infty) : L^1_{\Phi_1}(\Omega))$ and $u(0, x) = u_0 \in L^1_{\Phi_1}(\Omega)$.

This kind of solution was first introduced in [Bonforte and Vázquez 2015]; see also [Bonforte and Vázquez 2016]. Roughly speaking, we are considering the weak solution to the “dual equation” $\partial_t U = -u^m$, where $U = \mathcal{L}^{-1}u$, posed on the bounded domain Ω with homogeneous Dirichlet conditions. Such a weak solution is obtained by approximation from below as the limit of the unique mild solution provided by the semigroup theory [Bonforte and Vázquez 2016], and it was used in [Vázquez 2014a] with space domain \mathbb{R}^N in the study of Barenblatt solutions. We call those solutions *minimal weak dual solutions*, and it has been proven in Theorems 4.4 and 4.5 of [Bonforte and Vázquez 2016] that such solutions exist and are unique for any nonnegative data $u_0 \in L^1_{\Phi_1}(\Omega)$. The class of weak dual solutions includes the classes of weak, mild and strong solutions, and is included in the class of very weak solutions. In this class of solutions the standard comparison result holds.

Explicit solution. When trying to understand the behavior of positive solutions with general nonnegative data, it is natural to look for solutions obtained by separation of variables. These are given by

$$\mathcal{U}_T(t, x) := (T + t)^{-1/(m-1)} S(x), \quad T \geq 0, \quad (3-2)$$

where S solves the elliptic problem

$$\begin{cases} \mathcal{L}S^m = S & \text{in } (0, +\infty) \times \Omega, \\ S = 0 & \text{on the boundary.} \end{cases} \quad (3-3)$$

The properties of S have been thoroughly studied in the companion paper [Bonforte et al. 2017b], and we summarize them here for the reader's convenience.

Theorem 3.2 (properties of asymptotic profiles). *Assume that \mathcal{L} satisfies (A1), (A2), and (K2). Then there exists a unique positive solution S to the Dirichlet problem (3-3) with $m > 1$. Moreover, let σ be as in (1-3), and assume that*

- either $\sigma = 1$ and $2sm \neq \gamma(m - 1)$;
- or $\sigma < 1$ and (K4) holds.

Then there exist positive constants c_0 and c_1 such that the following sharp absolute bounds hold true for all $x \in \Omega$:

$$c_0 \Phi_1(x)^{\sigma/m} \leq S(x) \leq c_1 \Phi_1(x)^{\sigma/m}. \quad (3-4)$$

When $2sm = \gamma(m - 1)$ then, assuming (K4), for all $x \in \Omega$ we have

$$c_0 \Phi_1(x)^{1/m} (1 + |\log \Phi_1(x)|)^{1/(m-1)} \leq S(x) \leq c_1 \Phi_1(x)^{1/m} (1 + |\log \Phi_1(x)|)^{1/(m-1)}. \quad (3-5)$$

Remark. As observed in the proof of Theorem 7.2, by applying Theorem 6.1 to the separate-variables solution $t^{-1/(m-1)} S(x)$ we deduce that (3-4) is still true when $\sigma < 1$ if, instead of assuming (K4), we suppose that

$$K(x, y) \leq c_1 |x - y|^{-(N+2s)}$$

for a.e. $x, y \in \mathbb{R}^N$ and that $\Phi_1 \in C^\gamma(\Omega)$.

When $T = 0$, the solution \mathcal{U}_0 in (3-2) is commonly named “friendly giant”, because it takes initial data $u_0 \equiv +\infty$ (in the sense of a pointwise limit as $t \rightarrow 0$) but is bounded for all $t > 0$. This term was coined in the study of the standard porous medium equation.

In Sections 4 and 5 we will state and prove our general results concerning upper and lower bounds respectively. These sections are the crux of this paper. The combination of such upper and lower bounds will then be summarized in Section 6. Consequences of these results in terms of asymptotic behavior and regularity estimates will be studied in Sections 7 and 8 respectively.

4. Upper boundary estimates

We present a general upper bound that holds under the sole assumptions (A1), (A2), and (K2), and hence is valid for all our examples.

Theorem 4.1 (absolute boundary estimates). *Let (A1), (A2), and (K2) hold. Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$, and let σ be as in (1-3). Then, there exists a computable constant $k_1 > 0$, depending only on N, s, m , and Ω , such that for all $t \geq 0$ and all $x \in \Omega$*

$$u(t, x) \leq \frac{k_1}{t^{1/(m-1)}} \begin{cases} \Phi_1(x_0)^{\sigma/m} & \text{if } \gamma \neq 2sm/(m-1), \\ \Phi_1(x_0)^{1/m}(1 + |\log \Phi_1(x_0)|)^{1/(m-1)} & \text{if } \gamma = 2sm/(m-1). \end{cases} \tag{4-1}$$

This absolute bound proves a strong regularization which is independent of the initial datum. It improves the absolute bound in [Bonforte and Vázquez 2016] in the sense that it exhibits a precise boundary behavior. The estimate gives the correct behavior for the solutions \mathcal{U}_T in (3-2) obtained by separation of variables; see Theorem 3.2. It turns out that the estimate will be sharp for all nonnegative, nontrivial solutions in the case of the RFL and the CFL. We will also see below that the estimate is not always the correct behavior for the SFL when data are small, as explained in the Introduction (see Section 4A, and Theorem 5.4 in Section 5).

Proof of Theorem 4.1. This subsection is devoted to the proof of Theorem 4.1. The first steps are based on a few basic results of [Bonforte and Vázquez 2016] that will also be used in the rest of the paper.

Step 1: pointwise and absolute upper estimates.

Pointwise estimates. We begin by recalling the basic pointwise estimates which are crucial in the proof of all the upper and lower bounds of this paper.

Proposition 4.2 [Bonforte and Vázquez 2015; 2016]. *It holds that*

$$\int_{\Omega} u(t, x) \mathbb{G}(x, x_0) \, dx \leq \int_{\Omega} u_0(x) \mathbb{G}(x, x_0) \, dx \quad \text{for all } t > 0. \tag{4-2}$$

Moreover, for every $0 < t_0 \leq t_1 \leq t$ and almost every $x_0 \in \Omega$, we have

$$\frac{t_0^{m/(m-1)}}{t_1^{m/(m-1)}}(t_1 - t_0) u^m(t_0, x_0) \leq \int_{\Omega} [u(t_0, x) - u(t_1, x)] \mathbb{G}(x, x_0) \, dx \leq (m - 1) \frac{t^{m/(m-1)}}{t_0^{1/(m-1)}} u^m(t, x_0). \tag{4-3}$$

Absolute upper bounds. Using the estimates above, in Theorem 5.2 of [Bonforte and Vázquez 2016] the authors proved that solutions corresponding to initial data $u_0 \in L^1_{\Phi_1}(\Omega)$ satisfy

$$\|u(t)\|_{L^\infty(\Omega)} \leq \frac{K_1}{t^{1/(m-1)}} \quad \text{for all } t > 0, \tag{4-4}$$

with a constant K_1 independent of u_0 . For this reason, this is called ‘‘absolute bound’’.

Step 2: upper bounds via Green function estimates. The proof of Theorem 4.1 requires the following general statement; see [Bonforte et al. 2017b, Proposition 6.5]:

Lemma 4.3. *Let (A1), (A2), and (K2) hold, and let $v : \Omega \rightarrow \mathbb{R}$ be a nonnegative bounded function. Let σ be as in (1-3), and assume that, for a.e. $x_0 \in \Omega$,*

$$v(x_0)^m \leq \kappa_0 \int_{\Omega} v(x) \mathbb{G}(x, x_0) \, dx. \tag{4-5}$$

Then, there exists a constant $\bar{\kappa}_\infty > 0$, depending only on s, γ, m, N, Ω , such that the following bound holds true for a.e. $x_0 \in \Omega$:

$$\int_\Omega v(x) \mathbb{G}(x, x_0) \, dx \leq \bar{\kappa}_\infty \kappa_0^{1/(m-1)} \begin{cases} \Phi_1(x_0)^\sigma & \text{if } \gamma \neq 2sm/(m-1), \\ \Phi_1(x_0)(1 + |\log \Phi_1(x_0)|^{m/(m-1)}) & \text{if } \gamma = 2sm/(m-1). \end{cases} \quad (4-6)$$

Step 3: end of the proof of [Theorem 4.1](#). We already know that $u(t) \in L^\infty(\Omega)$ for all $t > 0$ by [\(4-4\)](#). Also, choosing $t_1 = 2t_0$ in [\(4-3\)](#) we deduce that, for $t \geq 0$ and a.e. $x_0 \in \Omega$,

$$u^m(t, x_0) \leq \frac{2^{m/(m-1)}}{t} \int_\Omega u(t, x) \mathbb{G}(x, x_0) \, dx. \quad (4-7)$$

The above inequality corresponds exactly to hypothesis [\(4-5\)](#) of [Lemma 4.3](#) with the value $\kappa_0 = 2^{m/(m-1)}t^{-1}$. As a consequence, inequality [\(4-6\)](#) holds, and we conclude that for a.e. $x_0 \in \Omega$ and all $t > 0$

$$\int_\Omega u(t, x) \mathbb{G}(x, x_0) \, dx \leq \frac{\bar{\kappa}_\infty 2^{m/(m-1)^2}}{t^{1/(m-1)}} \begin{cases} \Phi_1(x_0)^\sigma & \text{if } \gamma \neq 2sm/(m-1), \\ \Phi_1(x_0)(1 + |\log \Phi_1(x_0)|^{m/(m-1)}) & \text{if } \gamma = 2sm/(m-1). \end{cases} \quad (4-8)$$

Hence, combining this bound with [\(4-7\)](#), we get

$$u^m(t, x_0) \leq \frac{k_1^m}{t^{m/(m-1)}} \begin{cases} \Phi_1(x_0)^\sigma & \text{if } \gamma \neq 2sm/(m-1), \\ \Phi_1(x_0)(1 + |\log \Phi_1(x_0)|^{m/(m-1)}) & \text{if } \gamma = 2sm/(m-1). \end{cases}$$

This proves the upper bounds [\(4-1\)](#) and concludes the proof. □

4A. Upper bounds for small data and small times. As mentioned in the [Introduction](#), the above upper bounds may not be realistic when $\sigma < 1$. We have the following estimate for small times if the initial data are sufficiently small.

Theorem 4.4. *Let \mathcal{L} satisfy [\(A1\)](#), [\(A2\)](#), and [\(L2\)](#). Suppose also that \mathcal{L} has a first eigenfunction $\Phi_1 \asymp \text{dist}(x, \partial\Omega)^\gamma$, and assume that $\sigma < 1$. Finally, we assume that for all $x, y \in \Omega$*

$$K(x, y) \leq \frac{c_1}{|x - y|^{N+2s}} \left(\frac{\Phi_1(x)}{|x - y|^\gamma} \wedge 1 \right) \left(\frac{\Phi_1(y)}{|x - y|^\gamma} \wedge 1 \right) \quad \text{and} \quad B(x) \leq c_1 \Phi_1(x)^{-2s/\gamma}. \quad (4-9)$$

Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Then, for every initial data $u_0 \leq A\Phi_1^{1-2s/\gamma}$ for some $A > 0$, we have

$$u(t) \leq \frac{\Phi_1^{1-2s/\gamma}}{(A^{1-m} - \tilde{C}t)^{m-1}} \quad \text{on } [0, T_A], \quad \text{where } T_A := \frac{1}{\tilde{C}A^{m-1}},$$

and the constant $\tilde{C} > 0$, that depends only on N, s, m, λ_1, c_1 , and Ω .

Remark. This result applies to the SFL. Notice that when $\sigma < 1$ we have always $1 - 2s/\gamma > \sigma/m$; hence in this situation small data have a smaller behavior at the boundary than the one predicted in [Theorem 4.1](#). This is not true for “big” data, for instance for solutions obtained by separation of variables, as already said.

Proof. In view of our assumption on the initial datum, namely $u_0 \leq A\Phi_1^{1-2s/\gamma}$, by comparison it is enough to prove that the function

$$\bar{u}(t, x) = F(t)\Phi_1(x)^{1-2s/\gamma}, \quad F(t) = \frac{1}{(A^{1-m} - \tilde{C}t)^{m-1}},$$

is a supersolution (i.e., $\partial_t \bar{u} \geq -\mathcal{L}\bar{u}^m$) in $(0, T_A) \times \Omega$ provided we choose \tilde{C} sufficiently large.

To this aim, we use the following elementary inequality, whose proof is left to the interested reader: for any $\eta > 1$ and any $M > 0$ there exists $\tilde{b} = \tilde{b}(M) > 0$ such that letting $\tilde{\eta} := \eta \wedge 2$

$$a^\eta - b^\eta \leq \eta b^{\eta-1}(a - b) + \tilde{b}|a - b|^{\tilde{\eta}} \quad \text{for all } 0 \leq a, b \leq M. \tag{4-10}$$

We apply inequality (4-10) to $a = \Phi_1(y)$ and $b = \Phi_1(x)$, $\eta = m(1 - 2s/\gamma)$, noticing that $\eta > 1$ if and only if $\sigma < 1$, and we obtain (recall that Φ_1 is bounded)

$$\begin{aligned} \bar{u}^m(t, y) - \bar{u}^m(t, x) &= F(t)^m (\Phi_1(y)^{m(1-2s/\gamma)} - \Phi_1(x)^{m(1-2s/\gamma)}) = F(t)^m (\Phi_1(y)^\eta - \Phi_1(x)^\eta) \\ &\leq \eta F(t)^m \Phi_1(x)^{\eta-1} [\Phi_1(y) - \Phi_1(x)] + \tilde{b} F(t)^m |\Phi_1(y) - \Phi_1(x)|^{\tilde{\eta}} \\ &\leq \eta F(t)^m \Phi_1(x)^{\eta-1} [\Phi_1(y) - \Phi_1(x)] + \tilde{b} F(t)^m c_\gamma^{\tilde{\eta}} |x - y|^{\tilde{\eta}\gamma}, \end{aligned}$$

where in the last step we have used that $|\Phi_1(y) - \Phi_1(x)| \leq c_\gamma |x - y|^\gamma$. Since $B \leq c_1 \Phi_1^{-2s/\gamma}$,

$$\int_{\mathbb{R}^N} [\Phi_1(y) - \Phi_1(x)]K(x, y) dy = -\mathcal{L}\Phi_1(x) + B(x)\Phi_1(x) \leq -\lambda_1 \Phi_1(x) + c_1 \Phi_1(x)^{1-2s/\gamma}.$$

Thus, recalling that $\eta, \tilde{\eta} > 1$ and that Φ_1 is bounded, it follows that

$$\begin{aligned} -\mathcal{L}[\bar{u}^m](x) &= \int_{\mathbb{R}^N} [\bar{u}^m(t, y) - \bar{u}^m(t, x)]K(x, y) dy + B(x)\bar{u}^m(t, x) \\ &\leq \eta F(t)^m \Phi_1(x)^{\eta-1} [-\lambda_1 \Phi_1(x) + c_1 \Phi_1(x)^{1-2s/\gamma}] \\ &\quad + B(x)F(t)^m \Phi_1^\eta(x) + \tilde{b} c_\gamma^{\tilde{m}} F(t)^m \int_{\mathbb{R}^N} |x - y|^{\tilde{\eta}\gamma} K(x, y) dy \\ &\leq \tilde{c}F(t)^m \left(\Phi_1(x)^{\eta-2s/\gamma} + \int_{\mathbb{R}^N} |x - y|^{\tilde{\eta}\gamma} K(x, y) dy \right). \end{aligned} \tag{4-11}$$

Next, we claim that, as a consequence of (4-9),

$$\int_{\mathbb{R}^N} |x - y|^{\tilde{\eta}\gamma} K(x, y) dy \leq c_4 \Phi_1(x)^{1-2s/\gamma}. \tag{4-12}$$

Postponing for the moment the proof of the above inequality, we first show how to conclude: combining (4-11) and (4-12) we have

$$-\mathcal{L}\bar{u}^m \leq c_5 F(t)^m \Phi_1(x)^{1-2s/\gamma} = F'(t)\Phi_1(x)^{1-2s/\gamma} = \partial_t \bar{u},$$

where we used that $F'(t) = c_5 F(t)^{\tilde{m}}$ provided $\tilde{C} = c_5(m - 1)$. This proves that \bar{u} is a supersolution in $(0, T) \times \Omega$. Hence the proof is concluded once we prove inequality (4-12); for this, using hypothesis

(4-9) and choosing $r = \Phi_1(x)^{1/\gamma}$ we have

$$\begin{aligned} \int_{\mathbb{R}^N} |x - y|^{\tilde{\eta}\gamma} K(x, y) \, dy &\leq c_1 \int_{B_r(x)} \frac{1}{|x - y|^{N+2s-\tilde{\eta}\gamma}} \, dy + c_1 \Phi_1(x) \int_{\Omega \setminus B_r(x)} \frac{1}{|x - y|^{N+2s+\gamma-\tilde{\eta}\gamma}} \, dy \\ &\leq c_2 r^{\tilde{\eta}\gamma-2s} + c_1 \frac{\Phi_1(x)}{r^{2s}} \int_{\Omega \setminus B_r(x)} \frac{1}{|x - y|^{N+\gamma-\tilde{\eta}\gamma}} \, dy \\ &= c_2 r^{\tilde{\eta}\gamma-2s} + c_3 \frac{\Phi_1(x)}{r^{2s}} \leq c_4 \Phi_1(x)^{1-2s/\gamma}, \end{aligned}$$

where we used that $\tilde{\eta}\gamma - 2s > 0$ and $\tilde{\eta} > 1$. □

Remark. For operators for which the previous assumptions hold with $B \equiv 0$, we can actually prove a better upper bound for “smaller data”:

Corollary 4.5. *Under the assumptions of Theorem 4.4, assume that moreover $B \equiv 0$ and $u_0 \leq A\Phi_1$ for some $A > 0$. Then, we have*

$$u(t) \leq \frac{\Phi_1}{(A^{1-m} - \tilde{C}t)^{m-1}} \quad \text{on } [0, T_A], \quad \text{where } T_A := \frac{1}{\tilde{C}A^{m-1}},$$

and the constant $\tilde{C} > 0$ depends only on N, s, m, λ_1, c_1 , and Ω .

Proof. We have to show that $\bar{u}(t, x) = F(t)\Phi_1(x)$ is a supersolution: we essentially repeat the proof of Theorem 4.4 with $\gamma = m$ (formally replace $1 - 2s/\gamma$ by 1), taking into account that $B \equiv 0$ and $u_0 \leq A\Phi_1$. □

5. Lower bounds

This section is devoted to the proofs of all the lower bounds summarized later in the main theorems, Theorems 6.1, 6.2, and 6.3. The general situation is quite involved to describe, so we will separate several cases and we will indicate for which examples it holds for the sake of clarity.

Infinite speed of propagation: universal lower bounds. First, we are going to quantitatively establish that all nonnegative weak dual solutions of our problems are in fact positive in Ω for all $t > 0$. This result is valid for all nonlocal operators considered in this paper.

Theorem 5.1. *Let \mathcal{L} satisfy (A1), (A2), and (L2). Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Then there exists a constant $\underline{\kappa}_0 > 0$ such that the following inequality holds:*

$$u(t, x) \geq \underline{\kappa}_0 \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\Phi_1(x)}{t^{1/(m-1)}} \quad \text{for all } t > 0 \text{ and a.e. } x \in \Omega. \tag{5-1}$$

Here $t_* = \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, and the constants $\underline{\kappa}_0$ and κ_* depend only on $N, s, \gamma, m, c_0, c_1$, and Ω .

Notice that, for $t \geq t_*$, the dependence on the initial data disappears from the lower bound, as the inequality reads as

$$u(t) \geq \underline{\kappa}_0 \frac{\Phi_1}{t^{1/(m-1)}} \quad \text{for all } t \geq t_*,$$

where $\underline{\kappa}_0$ is an absolute constant. Assumption (L2) on the kernel K of \mathcal{L} holds for all examples mentioned in Section 10A.

Clearly, the power in this lower bound does not match the one of the general upper bounds of Theorem 4.1; hence we cannot expect these bounds to be sharp. However, when $\sigma < 1$, for small times and small data and when $B \equiv 0$, the lower bounds (5-1) match the upper bounds of Corollary 4.5; hence they are sharp. Theorem 5.1 shows that, even in the “worst case scenario”, there is a quantitative lower bound for all positive times, and shows infinite speed of propagation.

Matching lower bounds, I. Actually, in many cases the kernel of the nonlocal operator satisfies a stronger property, namely $\inf_{x,y \in \Omega} K(x, y) \geq \underline{\kappa}_\Omega > 0$ and $B \equiv 0$, in which case we can actually obtain sharp lower bounds for all times. Here we do not consider the potential logarithmic correction that may appear in the “critical case” $2sm = \gamma(m - 1)$; indeed, as far as examples are concerned, the next theorem applies to the RFL and the CFL, for which $2sm > \gamma(m - 1)$.

Theorem 5.2. *Let \mathcal{L} satisfy (A1), (A2), and (L1). Furthermore, suppose that \mathcal{L} has a first eigenfunction $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)^\gamma$. Let σ be as in (1-3) and assume that*

- either $\sigma = 1$;
- or $\sigma < 1$, $K(x, y) \leq c_1|x - y|^{-(N+2s)}$ for a.e. $x, y \in \mathbb{R}^N$, and $\Phi_1 \in C^\gamma(\bar{\Omega})$.

Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Then there exists a constant $\underline{\kappa}_1 > 0$ such that the following inequality holds:

$$u(t, x) \geq \underline{\kappa}_1 \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \quad \text{for all } t > 0 \text{ and a.e. } x \in \Omega, \tag{5-2}$$

where $t_* = \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$. The constants κ_* and $\underline{\kappa}_1$ depend only on $N, s, \gamma, m, \underline{\kappa}_\Omega, c_1, \Omega$, and $\|\Phi_1\|_{C^\gamma(\Omega)}$.

- Remarks.** (i) As in the case of Theorem 5.1, for large times the dependence on the initial data disappears from the lower bound and we have absolute lower bounds.
- (ii) The boundary behavior is sharp when $2sm \neq \gamma(m - 1)$ in view of the upper bound from Theorem 4.1.
- (iii) This theorem applies to the RFL and the CFL, but not to the SFL (or, more generally, spectral powers of elliptic operators); see Sections 2A and 2. In the case of the RFL, this result was obtained in Theorem 1 of [Bonforte et al. 2017a].

We have already seen the example of the separate-variables solutions (3-2) that have a very definite behavior at the boundary $\partial\Omega$. The analysis of general solutions leads to completely different situations for $\sigma = 1$ and $\sigma < 1$.

Matching lower bounds, II: The case $\sigma = 1$. When $\sigma = 1$ we can establish a quantitative lower bound near the boundary that matches the separate-variables behavior for large times (except in the case $2sm = \gamma(m - 1)$ where the result is false, see Theorem 5.4 below). We do not need the assumption of nondegenerate kernel, so the SFL can be considered.

Theorem 5.3. *Let (A1), (A2), and (K2) hold, and let $\sigma = 1$. Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. There exists a constant $\underline{\kappa}_2 > 0$ such that*

$$u(t, x) \geq \underline{\kappa}_2 \frac{\Phi_1(x)^{1/m}}{t^{1/(m-1)}} \quad \text{for all } t \geq t_* \text{ and a.e. } x \in \Omega. \quad (5-3)$$

Here, $t_* = \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, and the constants κ_* and $\underline{\kappa}_2$ depend only on N, s, γ, m , and Ω .

Remarks. (i) At first sight, this theorem may seem weaker than the previous positivity result. However, this result has wider applicability since it holds under the only assumption (K2) on \mathbb{G} . In particular it is valid in the local case $s = 1$, where the finite speed of propagation makes it impossible to have global lower bounds for small times.

(ii) When $\mathcal{L} = -\Delta$ the result has been proven in [Aronson and Peletier 1981; Vázquez 2004] by quite different methods. On the other hand, our method is very general and immediately applies to the case when \mathcal{L} is an elliptic operator with C^1 coefficients; see Section 10A.

(iii) This result fixes a small error in Theorem 7.1 of [Bonforte and Vázquez 2015], where the power σ was not present.

The anomalous lower bounds with small data. As shown in Theorem 5.1, the lower bound $u(t) \gtrsim \Phi_1$ is always valid. We now discuss the possibility of improving this bound.

Let S solve the elliptic problem (3-3). It follows by comparison whenever $u_0 \geq \epsilon_0 S$ with $\epsilon_0 > 0$ then $u(t) \geq S/(T_0 + t)^{1/(m-1)}$, where $T_0 = \epsilon_0^{1-m}$. Since $S \asymp \Phi_1^{\sigma/m}$ under (K4) (up to a possible logarithmic correction in the critical case, see Theorem 3.2), there are initial data for which the lower behavior is dictated by $\Phi_1(x)^{\sigma/m} t^{-1/(m-1)}$. More generally, as we shall see in Theorem 7.1, given any initial datum $u_0 \in L^1_{\Phi_1}(\Omega)$ the function $v(t, x) := t^{1/(m-1)} u(t, x)$ always converges to S in $L^\infty(\Omega)$ as $t \rightarrow \infty$, independently of the value of σ . Hence, one may conjecture that there should exist a waiting time $t_* > 0$ after which the lower behavior is dictated by $\Phi_1(x)^{\sigma/m} t^{-1/(m-1)}$, in analogy with what happens for the classical porous medium equation. As we shall see, this is actually *false* when $\sigma < 1$ or $2sm = \gamma(m-1)$. Since for large times $v(t, x)$ must look like $S(x)$ in uniform norm away from the boundary (by the interior regularity that we will prove later), the contrasting situation for large times could be described as a “dolphin’s head” with the “snout” flatter than the “forehead”. As $t \rightarrow \infty$ the forehead progressively fills the whole domain.

The next result shows that, in general, we cannot hope to prove that $u(t)$ is larger than $\Phi_1^{1/m}$. In particular, when $\sigma < 1$ or $2sm = \gamma(m-1)$, this shows that the behavior $u(t) \asymp S$ cannot hold.

Theorem 5.4. *Let (A1), (A2), and (K2) hold, and $u \geq 0$ be a weak dual solution to the CDP corresponding to a nonnegative initial datum $u_0 \in L^1_{\Phi_1}(\Omega)$. Assume that $u_0(x) \leq C_0 \Phi_1(x)$ a.e. in Ω for some $C_0 > 0$. Then there exists a constant $\hat{\kappa}$, depending only on N, s, γ, m , and Ω , such that*

$$u(t, x)^m \leq C_0 \hat{\kappa} \frac{\Phi_1(x)}{t} \quad \text{for all } t > 0 \text{ and a.e. } x \in \Omega.$$

In particular, if $\sigma < 1$ (resp. $2sm = \gamma(m - 1)$), then

$$\lim_{x \rightarrow \partial\Omega} \frac{u(t, x)}{\Phi_1(x)^{\sigma/m}} = 0 \quad \left(\text{resp. } \lim_{x \rightarrow \partial\Omega} \frac{u(t, x)}{\Phi_1(x)^{1/m}(1 + |\log \Phi_1(x)|)^{1/(m-1)}} = 0 \right) \quad \text{for any } t > 0.$$

The proposition above could make one wonder whether the sharp general lower bound could be given by $\Phi_1^{1/m}$, as in the case $\sigma = 1$. Recall that, under rather minimal assumptions on the kernel K associated to \mathcal{L} , we have a universal lower bound for $u(t)$ in terms of Φ_1 (see [Theorem 5.1](#)). Here we shall see that, under [\(K4\)](#), the bound $u(t) \gtrsim \Phi_1^{1/m}$ is false for $\sigma < 1$.

Theorem 5.5. *Let [\(A1\)](#), [\(A2\)](#), and [\(K4\)](#) hold, and let $u \geq 0$ be a weak dual solution to the CDP corresponding to a nonnegative initial datum $u_0 \leq C_0\Phi_1$ for some $C_0 > 0$. Assume that there exist constants $\underline{\kappa}$, T , $\alpha > 0$ such that*

$$u(T, x) \geq \underline{\kappa}\Phi_1^\alpha(x) \quad \text{for a.e. } x \in \Omega.$$

Then $\alpha \geq 1 - 2s/\gamma$. In particular $\alpha > 1/m$ if $\sigma < 1$.

We devote the rest of this section to the proof of the above results, and to this end we collect in the first two subsections some preliminary lower bounds and results about approximate solutions.

5A. Lower bounds for weighted norms. Here we prove some useful lower bounds for weighted norms, which follow from the L^1 -continuity for ordered solutions in the version proved in [Proposition 8.1](#) of [\[Bonforte and Vázquez 2016\]](#).

Lemma 5.6 (backward-in-time $L^1_{\Phi_1}$ lower bounds). *Let u be a solution to the CDP corresponding to the initial datum $u_0 \in L^1_{\Phi_1}(\Omega)$. For all*

$$0 \leq \tau_0 \leq t \leq \tau_0 + \frac{1}{(2\bar{K})^{1/(2s\vartheta_\gamma)} \|u(\tau_0)\|_{L^1_{\Phi_1}(\Omega)}^{m-1}} \tag{5-4}$$

we have

$$\frac{1}{2} \int_{\Omega} u(\tau_0, x)\Phi_1(x) \, dx \leq \int_{\Omega} u(t, x)\Phi_1(x) \, dx, \tag{5-5}$$

where $\vartheta_\gamma := 1/[2s + (N + \gamma)(m - 1)]$ and $\bar{K} > 0$ is a computable constant.

Proof. We recall the inequality of [Proposition 8.1](#) of [\[Bonforte and Vázquez 2016\]](#), adapted to our case: for all $0 \leq \tau_0 \leq \tau$, t we have

$$\int_{\Omega} u(\tau, x)\Phi_1(x) \, dx \leq \int_{\Omega} u(t, x)\Phi_1(x) \, dx + \bar{K} \|u(\tau_0)\|_{L^1_{\Phi_1}(\Omega)}^{2s(m-1)\vartheta_\gamma+1} |t - \tau|^{2s\vartheta_\gamma}. \tag{5-6}$$

Choosing $\tau = \tau_0$ in the above inequality, we get

$$\left[1 - \bar{K}_9 \|u(\tau_0)\|_{L^1_{\Phi_1}(\Omega)}^{2s(m-1)\vartheta_\gamma} |t - \tau_0|^{2s\vartheta_\gamma} \right] \int_{\Omega} u(\tau_0, x)\Phi_1(x) \, dx \leq \int_{\Omega} u(t, x)\Phi_1(x) \, dx. \tag{5-7}$$

Then [\(5-5\)](#) follows from [\(5-4\)](#). □

We also need a lower bound for $L^p_{\Phi_1}(\Omega)$ norms.

Lemma 5.7. *Let u be a solution to the CDP corresponding to the initial datum $u_0 \in L^1_{\Phi_1}(\Omega)$. Then the following lower bound holds true for any $t \in [0, t_*]$ and $p \geq 1$:*

$$c_2 \left(\int_{\Omega} u_0(x) \Phi_1(x) \, dx \right)^p \leq \int_{\Omega} u^p(t, x) \Phi_1(x) \, dx. \tag{5-8}$$

Here $t_* = c_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, where $c_2, c_* > 0$ are positive constants that depend only on N, s, m, p, Ω .

The proof of this lemma is an easy adaptation of the proof of Lemma 2.2 of [Bonforte et al. 2017a], so we skip it. Notice that c_* has explicit form given in [Bonforte and Vázquez 2015; 2016; Bonforte et al. 2017a], while the form of c_2 is given in the proof of Lemma 2.2 of [Bonforte et al. 2017a].

5B. Approximate solutions. To prove our lower bounds, we will need a special class of approximate solutions u_δ . We will list now the necessary details. In the case when \mathcal{L} is the restricted fractional Laplacian (RFL), see Section 10A, these solutions have been used in Appendix II of [Bonforte et al. 2017a], where complete proofs can be found; the proof there holds also for the operators considered here. The interested reader can easily adapt the proofs in that paper to the current case.

Let us fix $\delta > 0$ and consider the problem

$$\begin{cases} \partial_t v_\delta = -\mathcal{L}[(v_\delta + \delta)^m - \delta^m] & \text{for any } (t, x) \in (0, \infty) \times \Omega, \\ v_\delta(t, x) = 0 & \text{for any } (t, x) \in (0, \infty) \times (\mathbb{R}^N \setminus \Omega), \\ v_\delta(0, x) = u_0(x) & \text{for any } x \in \Omega. \end{cases} \tag{5-9}$$

Next, we define

$$u_\delta := v_\delta + \delta.$$

We summarize here the basic properties of u_δ .

Approximate solutions u_δ exist, are unique, and bounded for all $(t, x) \in (0, \infty) \times \bar{\Omega}$ whenever $0 \leq u_0 \in L^1_{\Phi_1}(\Omega)$. Also, they are uniformly positive: for any $t \geq 0$,

$$u_\delta(t, x) \geq \delta > 0 \quad \text{for a.e. } x \in \Omega. \tag{5-10}$$

This implies that the equation for u_δ is never degenerate in the interior, so solutions are as smooth as the linear parabolic theory with the kernel K allows them to be (in particular, in the case of the fractional Laplacian, they are C^∞ in space and C^1 in time). Also, by a comparison principle, for all $\delta > \delta' > 0$,

$$u_\delta(t, x) \geq u_{\delta'}(t, x) \quad \text{for all } t \geq 0 \text{ and a.e. } x \in \Omega, \tag{5-11}$$

$$u_\delta(t, x) \geq u(t, x) \quad \text{for all } t \geq 0 \text{ and a.e. } x \in \Omega. \tag{5-12}$$

Furthermore, they converge in $L^1_{\Phi_1}(\Omega)$ to u as $\delta \rightarrow 0$:

$$\|u_\delta(t) - u(t)\|_{L^1_{\Phi_1}(\Omega)} \leq \|u_\delta(0) - u_0\|_{L^1_{\Phi_1}(\Omega)} = \delta \|\Phi_1\|_{L^1(\Omega)}. \tag{5-13}$$

As a consequence of (5-11) and (5-13), we deduce that u_δ converges pointwise to u at almost every point: more precisely, for all $t \geq 0$,

$$u(t, x) = \lim_{\delta \rightarrow 0^+} u_\delta(t, x) \quad \text{for a.e. } x \in \Omega. \tag{5-14}$$

5C. Proof of Theorem 5.1. The proof consists in showing that

$$u(t, x) \geq \underline{u}(t, x) := k_0 t \Phi_1(x)$$

for all $t \in [0, t_*]$ and a.e. $x \in \Omega$, where the parameter $k_0 > 0$ will be fixed later. Note that, once the inequality $u \geq \underline{u}$ on $[0, t_*]$ is proved, we conclude as follows: since $t \mapsto t^{1/(m-1)} u(t, x)$ is nondecreasing in $t > 0$ for a.e. $x \in \Omega$ [Bonforte and Vázquez 2016, (2.3)], we have

$$u(t, x) \geq \left(\frac{t_*}{t}\right)^{1/(m-1)} u(t_*, x) \geq k_0 t_* \left(\frac{t_*}{t}\right)^{1/(m-1)} \Phi_1(x) \quad \text{for all } t \geq t_*.$$

Then, the result will follow $\underline{\kappa}_0 = k_0 t_*^{m/(m-1)}$ (note that, as we shall see below, $k_0 t_*^{m/(m-1)}$ can be chosen independently of u_0). Hence, we are left with proving that $u \geq \underline{u}$ on $[0, t_*]$.

Step 1: reduction to an approximate problem. Let us fix $\delta > 0$ and consider the approximate solutions u_δ constructed in Section 5B. We shall prove that $u_\delta \geq \underline{u}$ on $[0, t_*] \times \Omega$, so that the result will follow by the arbitrariness of δ .

Step 2: We claim that $\underline{u}(t, x) < u_\delta(t, x)$ for all $0 \leq t \leq t_*$ and $x \in \Omega$, for a suitable choice of $k_0 > 0$. Assume that the inequality $\underline{u} < u_\delta$ is false in $[0, t_*] \times \bar{\Omega}$, and let (t_c, x_c) be the first contact point between \underline{u} and u_δ . Since $u_\delta = \delta > 0 = \underline{u}$ on the lateral boundary, $(t_c, x_c) \in (0, t_*] \times \Omega$. Now, since $(t_c, x_c) \in (0, t_*] \times \Omega$ is the first contact point, we necessarily have

$$u_\delta(t_c, x_c) = \underline{u}(t_c, x_c) \quad \text{and} \quad u_\delta(t, x) \geq \underline{u}(t, x) \quad \text{for all } t \in [0, t_c], x \in \bar{\Omega}. \tag{5-15}$$

Thus, as a consequence,

$$\partial_t u_\delta(t_c, x_c) \leq \partial_t \underline{u}(t_c, x_c) = k_0 \Phi_1(x_c). \tag{5-16}$$

Next, we observe the following Kato-type inequality holds: for any nonnegative function f ,

$$\mathcal{L}(f^m) \leq m f^{m-1} \mathcal{L} f. \tag{5-17}$$

Indeed, by convexity, $f(x)^m - f(y)^m \leq m[f(x)]^{m-1}(f(x) - f(y))$; therefore

$$\begin{aligned} \mathcal{L}(f^m)(x) &= \int_{\mathbb{R}^N} [f(x)^m - f(y)^m] K(x, y) dy + B(x) f(x)^m \\ &\leq m[f(x)]^{m-1} \int_{\mathbb{R}^N} [f(x) - f(y)] K(x, y) dy + B(x) f(x)^m \\ &= m[f(x)]^{m-1} \left(\int_{\mathbb{R}^N} [f(x) - f(y)] K(x, y) dy + B(x) f(x) \right) - (m-1)B(x) f(x)^m \\ &\leq m[f(x)]^{m-1} \mathcal{L} f(x). \end{aligned}$$

As a consequence of (5-17), since $t_c \leq t_*$ and Φ_1 is bounded,

$$\begin{aligned} \mathcal{L}(\underline{u}^m)(t, x) &\leq m \underline{u}^{m-1} \mathcal{L}(\underline{u}) = m[k_0 t \Phi_1(x)]^{m-1} k_0 t \mathcal{L}(\Phi_1)(x) \\ &= m \lambda_1 [k_0 t \Phi_1(x)]^m \leq \kappa_1 (t_* k_0)^m \Phi_1(x). \end{aligned} \tag{5-18}$$

Then, using (5-16) and (5-18), we establish an upper bound for $-\mathcal{L}(u_\delta^m - \underline{u}^m)(t_c, x_c)$ as follows:

$$-\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) = \partial_t u_\delta(t_c, x_c) + \mathcal{L}(\underline{u}^m)(t_c, x_c) \leq k_0 [1 + \kappa_1 t_*^m k_0^{m-1}] \Phi_1(x_c). \tag{5-19}$$

Next, we want to prove lower bounds for $-\mathcal{L}(u_\delta^m - \psi^m)(t_c, x_c)$, and this is the point where the nonlocality of the operator enters, since we make essential use of hypothesis (L2). We recall that by (5-15) we have $u_\delta^m(t_c, x_c) = \underline{u}^m(t_c, x_c)$, so that assumption (L2) gives

$$\begin{aligned} -\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) &= -\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) + B(x_c)[u_\delta^m(t_c, x_c) - \underline{u}^m(t_c, x_c)] \\ &= -\int_{\mathbb{R}^N} [(u_\delta^m(t_c, x_c) - u_\delta^m(t_c, y)) - (\underline{u}^m(t_c, x_c) - \underline{u}^m(t_c, y))] K(x_c, y) dy \\ &= \int_{\Omega} [u_\delta^m(t_c, y) - \underline{u}^m(t_c, y)] K(x_c, y) dy \\ &\geq c_0 \Phi_1(x_c) \int_{\Omega} [u_\delta^m(t_c, y) - \underline{u}^m(t_c, y)] \Phi_1(y) dy, \end{aligned}$$

from which it follows (since $\underline{u}^m = [k_0 t \Phi_1(x)]^m \leq \kappa_2 (t_* k_0)^m$)

$$\begin{aligned} -\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) &\geq c_0 \Phi_1(x_c) \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) dy - c_0 \Phi_1(x_c) \int_{\Omega} \underline{u}^m(t_c, y) \Phi_1(y) dy. \\ &\geq c_0 \Phi_1(x_c) \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) dy - c_0 \Phi_1(x_c) \kappa_3 (t_* k_0)^m. \end{aligned} \tag{5-20}$$

Combining the upper and lower bounds (5-19) and (5-20) we obtain

$$c_0 \Phi_1(x_c) \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) dy \leq k_0 [1 + (\kappa_1 + \kappa_3) t_*^m k_0^{m-1}] \Phi_1(x_c). \tag{5-21}$$

Hence, recalling (5-8), we get

$$c_2 \left(\int_{\Omega} u_0(x) \Phi_1(x) dx \right)^m \leq \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) dy \leq \frac{k_0}{c_0} [1 + (\kappa_1 + \kappa_3) t_*^m k_0^{m-1}].$$

Since $t_* = \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, this yields

$$c_2 \kappa_*^{m/(m-1)} t_*^{-m/(m-1)} \leq \frac{k_0}{c_0} [1 + (\kappa_1 + \kappa_3) t_*^m k_0^{m-1}],$$

which gives the desired contradiction provided we choose k_0 so that $\underline{k}_0 := k_0 t_*^{m/(m-1)}$ is universally small. □

5D. Proof of Theorem 5.2. The proof proceeds along the lines of the proof of Theorem 5.1, so we will just briefly mention the common parts.

We want to show that

$$\underline{u}(t, x) := \kappa_0 t \Phi_1(x)^{\sigma/m}, \tag{5-22}$$

is a lower barrier for our problem on $[0, t_*] \times \Omega$ provided κ_0 is small enough. More precisely, as in the proof of Theorem 5.1, we aim to prove that $\underline{u} < u_\delta$ on $[0, t_*] \times \Omega$, as the lower bound for $t \geq t_*$ then follows by monotonicity.

Assume by contradiction that the inequality $\underline{u}(t, x) < u_\delta(t, x)$ is false inside $[0, t_*] \times \bar{\Omega}$. Since $\underline{u} < u_\delta$ on the parabolic boundary, letting (t_c, x_c) be the first contact point, we necessarily have $(t_c, x_c) \in (0, t_*] \times \Omega$. The desired contradiction will be obtained by combining the upper and lower bounds (which we prove below) for the quantity $-\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c)$, and then choosing $\kappa_0 > 0$ suitably small. In this direction, it is convenient in what follows to assume that

$$\kappa_0 \leq 1 \wedge t_*^{-m/(m-1)} \quad \text{so that} \quad \kappa_0^{m-1} t_*^m \leq 1. \tag{5-23}$$

Upper bound. We first establish the following upper bound: there exists a constant $\bar{A} > 0$ such that

$$-\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) \leq \partial_t u_\delta(t_c, x_c) + \mathcal{L}\underline{u}^m(t_c, x_c) \leq \bar{A}\kappa_0. \tag{5-24}$$

To prove this, we estimate $\partial_t u_\delta(t_c, x_c)$ and $\mathcal{L}\underline{u}^m(t_c, x_c)$ separately. First we notice that, since (t_δ, x_δ) is the first contact point, we have

$$u_\delta(t_\delta, x_\delta) = \underline{u}(t_\delta, x_\delta) \quad \text{and} \quad u_\delta(t, x) \geq \underline{u}(t, x) \quad \text{for all } t \in [0, t_\delta], x \in \Omega. \tag{5-25}$$

Hence, since $t_\delta \leq t_*$,

$$\partial_t u_\delta(t_\delta, x_\delta) \leq \partial_t \underline{u}(t_\delta, x_\delta) = \kappa_0 \Phi_1(x)^{\sigma/m} \leq \kappa_0 \|\Phi_1\|_{L^\infty(\Omega)}^{\sigma/m} = A_1 \kappa_0, \tag{5-26}$$

where we defined $A_1 := \|\Phi_1\|_{L^\infty(\Omega)}^{\sigma/m}$. Next we estimate $\mathcal{L}\underline{u}^m(t_c, x_c)$, using the Kato-type inequality (5-17); namely $\mathcal{L}[u^m] \leq m u^{m-1} \mathcal{L}u$. This implies

$$\begin{aligned} \mathcal{L}[\underline{u}^m](t, x) &\leq m \underline{u}^{m-1}(t, x) \mathcal{L}\underline{u}(t, x) = m(\kappa_0 t)^m \Phi_1(x)^{\sigma(m-1)/m} \mathcal{L}\Phi_1^\sigma(x) \\ &\leq m(\kappa_0 t_*)^m \|\Phi_1\|_{L^\infty(\Omega)}^{\sigma(m-1)/m} \|\mathcal{L}\Phi_1^\sigma\|_{L^\infty(\Omega)} := A_2 \kappa_0. \end{aligned} \tag{5-27}$$

Since $\kappa_0^{m-1} t_*^m \leq 1$, see (5-23), in order to prove that A_2 is finite it is enough to bound $\|\mathcal{L}\Phi_1^\sigma\|_{L^\infty(\Omega)}$. When $\sigma = 1$ we simply have $\mathcal{L}\Phi_1 = -\lambda_1 \Phi_1$; hence $A_2 \leq m\lambda_1 \|\Phi_1\|_{L^\infty(\Omega)}^{2-1/m}$. When $\sigma < 1$, we use the assumption $\Phi_1 \in C^\gamma(\Omega)$ to estimate

$$|\Phi_1^\sigma(x) - \Phi_1^\sigma(y)| \leq |\Phi_1(x) - \Phi_1(y)|^\sigma \leq C|x - y|^{\gamma\sigma} \quad \text{for all } x, y \in \Omega. \tag{5-28}$$

Hence, since $\gamma\sigma = 2sm/(m-1) > 2s$ and $K(x, y) \leq c_1|x - y|^{-(N+2s)}$, we see that

$$\begin{aligned} |\mathcal{L}\Phi_1^\sigma(x)| &= \left| \int_{\mathbb{R}^N} [\Phi_1^\sigma(x) - \Phi_1^\sigma(y)] K(x, y) dy \right| \\ &\leq \int_{\Omega} |x - y|^{\gamma\sigma} K(x, y) dy + C \|\Phi_1\|_{L^\infty(\Omega)}^\sigma \int_{\mathbb{R}^N \setminus B_1} |y|^{-(N+2s)} dy < \infty; \end{aligned}$$

hence A_2 is again finite. Combining (5-26) and (5-27), we obtain (5-24) with $\bar{A} := A_1 + A_2$.

Lower bound. We want to prove that there exists $\underline{A} > 0$ such that

$$-\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) \geq \frac{\kappa \Omega}{\|\Phi_1\|_{L^\infty(\Omega)}} \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) dy - \underline{A}\kappa_0. \tag{5-29}$$

This follows by (L1) and (5-25):

$$\begin{aligned}
 -\mathcal{L}[u_\delta^m - \underline{u}^m](t_c, x_c) &= - \int_{\mathbb{R}^N} [(u_\delta^m(t_c, x_c) - u_\delta^m(t_c, y)) - (\underline{u}^m(t_c, x_c) - \underline{u}^m(t_c, y))] K(x, y) \, dy \\
 &= \int_{\Omega} [u_\delta^m(t_c, y) - \underline{u}^m(t_c, y)] K(x, y) \, dy \\
 &\geq \underline{\kappa} \int_{\Omega} [u_\delta^m(t_c, y) - \underline{u}^m(t_c, y)] \, dy \\
 &\geq \frac{\underline{\kappa} \Omega}{\|\Phi_1\|_{L^\infty(\Omega)}} \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) \, dy - \underline{A} \kappa_0,
 \end{aligned} \tag{5-30}$$

where in the last step we used that $\underline{u}^m(t_c, y) = [\kappa_0 t_1^{\sigma/m}(y)]^m \leq \kappa_2(\kappa_0 t_*)^m$ and $\kappa_0^{m-1} t_*^m \leq 1$; see (5-23).

End of the proof. The contradiction can be now obtained by joining the upper and lower bounds (5-24) and (5-29). More precisely, we have proved

$$\int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) \, dy \leq \frac{\|\Phi_1\|_{L^\infty(\Omega)}}{\underline{\kappa} \Omega} (\bar{A} + \underline{A}) \kappa_0 := \bar{\kappa} \kappa_0,$$

which combined with the lower bound (5-8) yields

$$c_2 \left(\int_{\Omega} u_0(x) \Phi_1(x) \, dx \right)^m \leq \int_{\Omega} u_\delta^m(t_c, y) \Phi_1(y) \, dy \leq \bar{\kappa} \kappa_0.$$

Setting $\kappa_0 := (1 \wedge c_2/\bar{\kappa}) t_*^{-m/(m-1)}$, we obtain the desired contradiction. □

5E. Proof of Theorem 5.3. We first recall the upper pointwise estimates (4-3): for all $0 \leq t_0 \leq t_1 \leq t$ and a.e. $x_0 \in \Omega$, we have

$$\int_{\Omega} u(t_0, x) \mathbb{G}(x, x_0) \, dx - \int_{\Omega} u(t_1, x) \mathbb{G}(x, x_0) \, dx \leq (m-1) \frac{t^{m/(m-1)}}{t_0^{1/(m-1)}} u^m(t, x_0). \tag{5-31}$$

The proof follows by estimating the two integrals on the left-hand side separately.

We begin by using the upper bounds (4-8) to get

$$\int_{\Omega} u(t_1, x) \mathbb{G}(x, x_0) \, dx \leq \bar{\kappa} \frac{\Phi_1(x_0)}{t_1^{1/(m-1)}} \quad \text{for all } (t_1, x) \in (0, +\infty) \times \Omega. \tag{5-32}$$

Then we note that, as a consequence of (K2) and Lemma 5.6,

$$\int_{\Omega} u(t_0, x) \mathbb{G}(x, x_0) \, dx \geq \underline{\kappa} \Omega \Phi_1(x_0) \int_{\Omega} u(t_0, x) \Phi_1(x) \, dx \geq \frac{1}{2} \underline{\kappa} \Omega \Phi_1(x_0) \int_{\Omega} u_0(x) \Phi_1(x) \, dx \tag{5-33}$$

provided $t_0 \leq \tau_0 / \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{m-1}$. Combining (5-31), (5-32), and (5-33), for all $t \geq t_1 \geq t_0 \geq 0$ we obtain

$$u^m(t, x_0) \geq \frac{t_0^{1/(m-1)}}{m-1} \left(\frac{1}{2} \underline{\kappa} \Omega \|u_0\|_{L^1_{\Phi_1}(\Omega)} - \bar{\kappa} t_1^{-1/(m-1)} \right) \frac{\Phi_1(x_0)}{t^{m/(m-1)}}.$$

Choosing

$$t_0 := \frac{\tau_0}{\|u_0\|_{L^1_{\Phi_1}(\Omega)}^{m-1}} \leq t_1 := t_* = \frac{\kappa_*}{\|u_0\|_{L^1_{\Phi_1}(\Omega)}^{m-1}} \quad \text{with} \quad \kappa_* \geq \tau_0 \vee \left(\frac{\kappa_\Omega}{4\bar{\kappa}}\right)^{m-1}$$

so that $\frac{1}{2}\kappa_\Omega \|u_0\|_{L^1_{\Phi_1}(\Omega)} - \bar{\kappa} t_1^{-1/(m-1)} \geq \frac{1}{4}\kappa_\Omega \|u_0\|_{L^1_{\Phi_1}(\Omega)}$, the result follows. □

5F. Proofs of Theorems 5.4 and 5.5.

Proof of Theorem 5.4. Since $u_0 \leq C_0\Phi_1$ and $\mathcal{L}\Phi = \lambda_1\Phi_1$, we have

$$\int_{\Omega} u_0(x) \mathbb{G}(x, x_0) dx \leq C_0 \int_{\Omega} \Phi_1(x) \mathbb{G}(x, x_0) dx = C_0 \mathcal{L}^{-1}\Phi_1(x_0) = \frac{C_0}{\lambda_1} \Phi_1(x_0).$$

Since $t \mapsto \int_{\Omega} u(t, y) \mathbb{G}(x, y) dy$ is decreasing, see (4-2), it follows that

$$\int_{\Omega} u(t, y) \mathbb{G}(x_0, y) dy \leq \frac{C_0}{\lambda_1} \Phi_1(x_0) \quad \text{for all } t \geq 0. \tag{5-34}$$

Combining this estimate with (4-7) concludes the proof. □

Proof of Theorem 5.5. Given $x_0 \in \Omega$, set $R_0 := \text{dist}(x_0, \partial\Omega)$. Since $\mathbb{G}(x, x_0) \gtrsim |x - x_0|^{-(N-2s)}$ inside $B_{R_0/2}(x_0)$ by (K4), using our assumption on $u(T)$ we get

$$\int_{\Omega} \mathbb{G}(x, x_0) u(T, x) dx \gtrsim \int_{B_{R_0/2}(x_0)} \frac{\Phi_1(x)^\alpha}{|x - x_0|^{N-2s}} \gtrsim \Phi_1(x_0)^\alpha R_0^{2s}.$$

Recalling that $\Phi_1(x_0) \asymp R_0^\gamma$, this yields

$$\Phi_1(x_0)^{\alpha+2s/\gamma} \lesssim \int_{\Omega} \mathbb{G}(x, x_0) u(T, x) dx.$$

Combining the above inequality with (5-34) gives

$$\Phi_1(x_0)^{\alpha+2s/\gamma} \lesssim \Phi_1(x_0) \quad \text{for all } x_0 \in \Omega,$$

which implies

$$\alpha \geq 1 - \frac{2s}{\gamma}.$$

Noticing that $1 - 2s/\gamma > 1/m$ if and only if $\sigma < 1$, this concludes the proof. □

6. Summary of the general decay and boundary results

We now present a summary of the main results, which can be summarized in various forms of upper and lower bounds, which we call the global Harnack principle, GHP for short. As already mentioned, such inequalities are important for regularity issues (see Section 8), and they play a fundamental role in formulating the sharp asymptotic behavior (see Section 7). The proof of such a GHP is obtained by combining upper and lower bounds, stated and proved in Sections 4 and 5 respectively. There are cases when the bounds do not match, for which the complicated panorama described in the Introduction holds. As explained before, as far as examples are concerned, the latter anomalous situation happens only for the SFL.

Theorem 6.1 (global Harnack principle I). *Let \mathcal{L} satisfy (A1), (A2), (K2), and (L1). Furthermore, suppose that \mathcal{L} has a first eigenfunction $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)^\gamma$. Let σ be as in (1-3) and assume that $2sm \neq \gamma(m - 1)$ and*

- either $\sigma = 1$
- or $\sigma < 1$, $K(x, y) \leq c_1|x - y|^{-(N+2s)}$ for a.e. $x, y \in \mathbb{R}^N$, and $\Phi_1 \in C^\gamma(\bar{\Omega})$.

Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Then, there exist constants $\underline{\kappa}, \bar{\kappa} > 0$ such that the following inequality holds:

$$\underline{\kappa} \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \quad \text{for all } t > 0, x \in \Omega. \tag{6-1}$$

The constants $\underline{\kappa}, \bar{\kappa}$ depend only on $N, s, \gamma, m, c_1, \underline{\kappa}_\Omega, \Omega$, and $\|\Phi_1\|_{C^\gamma(\Omega)}$.

Proof. We combine the upper bound (4-1) with the lower bound (5-2). The expression of t_* is explicitly given in Theorem 5.2. □

Degenerate kernels. When the kernel K vanishes on $\partial\Omega$, there are two combinations of upper/lower bounds that provide Harnack inequalities, one for small times and one for large times. As we have already seen, there is a strong difference between the cases $\sigma = 1$ and $\sigma < 1$.

Theorem 6.2 (global Harnack principle II). *Let (A1), (A2), and (K2) hold. Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Assume that*

- either $\sigma = 1$ and $2sm \neq \gamma(m - 1)$;
- or $\sigma < 1$, $u_0 \geq \underline{\kappa}_0 \Phi_1^{\sigma/m}$ for some $\underline{\kappa}_0 > 0$, and (K4) holds.

Then there exist constants $\underline{\kappa}, \bar{\kappa} > 0$ such that the following inequality holds:

$$\underline{\kappa} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \quad \text{for all } t \geq t_*, x \in \Omega.$$

If $2sm = \gamma(m - 1)$, assuming (K4) and that $u_0 \geq \underline{\kappa}_0 \Phi_1(1 + |\log \Phi_1|)^{1/(m-1)}$ for some $\underline{\kappa}_0 > 0$, then for all $t \geq t_$ and all $x \in \Omega$*

$$\underline{\kappa} \frac{\Phi_1(x)^{1/m}}{t^{1/(m-1)}} (1 + |\log \Phi_1(x)|)^{1/(m-1)} \leq u(t, x) \leq \bar{\kappa} \frac{\Phi_1(x)^{1/m}}{t^{1/(m-1)}} (1 + |\log \Phi_1(x)|)^{1/(m-1)}.$$

The constants $\underline{\kappa}, \bar{\kappa}$ depend only on $N, s, \gamma, m, \underline{\kappa}_0, \underline{\kappa}_\Omega$, and Ω .

Proof. In the case $\sigma = 1$, we combine the upper bound (4-1) with the lower bound (5-3). The expression of t_* is explicitly given in Theorem 5.3. When $\sigma < 1$, the upper bound is still given by (4-1), while the lower bound follows by comparison with the solution $S(x)(\underline{\kappa}_0^{1-m} + t)^{-1/(m-1)}$, recalling that $S \asymp \Phi_1^{\sigma/m}$ (see Theorem 3.2). □

Remark. Local Harnack inequalities of elliptic/backward type follow as a consequence of Theorems 6.1 and 6.2, for all times and for large times respectively, see Theorem 8.2.

Note that, for small times, we cannot find matching powers for a global Harnack inequality (except for some special initial data), and such a result is actually false for $s = 1$ (in view of the finite speed of propagation). Hence, in the remaining cases, we have only the following general result.

Theorem 6.3 (nonmatching upper and lower bounds). *Let \mathcal{L} satisfy (A1), (A2), (K2), and (L2). Let $u \geq 0$ be a weak dual solution to the CDP corresponding to $u_0 \in L^1_{\Phi_1}(\Omega)$. Then, there exist constants $\underline{\kappa}, \bar{\kappa} > 0$ such that the following inequality holds when $2sm \neq \gamma(m - 1)$:*

$$\underline{\kappa} \left(1 \wedge \frac{t}{t_*}\right)^{m/(m-1)} \frac{\Phi_1(x)}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{\Phi_1(x)^{\sigma/m}}{t^{1/(m-1)}} \quad \text{for all } t > 0, x \in \Omega. \tag{6-2}$$

When $2sm = \gamma(m - 1)$, a logarithmic correction $(1 + |\log \Phi_1(x)|)^{1/(m-1)}$ appears in the right-hand side.

Proof. We combine the upper bound (4-1) with the lower bound (5-1). The expression of t_* is explicitly given in Theorem 5.1. □

Remark. As already mentioned in the Introduction, in the nonmatching case, which in examples can only happen for spectral-type operators, we have the appearance of an *anomalous behavior of solutions* corresponding to “small data”: it happens for all times when $\sigma < 1$ or $2sm = \gamma(m - 1)$, and it can eventually happen for short times when $\sigma = 1$.

7. Asymptotic behavior

An important application of the global Harnack inequalities of the previous section concerns the sharp asymptotic behavior of solutions. More precisely, we first show that for large times all solutions behave like the separate-variables solution $\mathcal{U}(t, x) = S(x) t^{-1/(m-1)}$ introduced at the end of Section 3. Then, whenever the GHP holds, we can improve this result to an estimate in relative error.

Theorem 7.1 (asymptotic behavior). *Assume that \mathcal{L} satisfies (A1), (A2), and (K2), and let S be as in Theorem 3.2. Let u be any weak dual solution to the CDP. Then, unless $u \equiv 0$,*

$$\|t^{1/(m-1)}u(t, \cdot) - S\|_{L^\infty(\Omega)} \xrightarrow{t \rightarrow \infty} 0. \tag{7-1}$$

Proof. The proof uses rescaling and time monotonicity arguments, and it is a simple adaptation of the proof of Theorem 2.3 of [Bonforte et al. 2015]. In those arguments, the interior $C^\alpha_x(\Omega)$ continuity is needed to improve the $L^1(\Omega)$ convergence to $L^\infty(\Omega)$, but the interior Hölder continuity is guaranteed by Theorem 8.1(i) below. □

We now exploit the GHP to get a stronger result.

Theorem 7.2 (sharp asymptotic behavior). *Under the assumptions of Theorem 7.1, assume that $u \not\equiv 0$. Furthermore, suppose that either the assumptions of Theorem 6.1 or of Theorem 6.2 hold. Set $\mathcal{U}(t, x) := t^{-1/(m-1)}S(x)$. Then there exists $c_0 > 0$ such that, for all $t \geq t_0 := c_0\|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, we have*

$$\left\| \frac{u(t, \cdot)}{\mathcal{U}(t, \cdot)} - 1 \right\|_{L^\infty(\Omega)} \leq \frac{2}{m-1} \frac{t_0}{t_0 + t}. \tag{7-2}$$

We remark that the constant $c_0 > 0$ only depends on $N, s, \gamma, m, \underline{\kappa}_0, \underline{\kappa}_\Omega$, and Ω .

Remark. This asymptotic result is sharp, as it can be checked by considering $u(t, x) = \mathcal{U}(t + 1, x)$. For the classical case, that is $\mathcal{L} = \Delta$, we recover the classical results of [Aronson and Peletier 1981; Vázquez 2004] with a different proof.

Proof. Notice that we are in the position to use Theorem 6.1 or 6.2; namely we have

$$u(t) \asymp t^{-1/(m-1)} S = \mathcal{U}(t, \cdot) \quad \text{for all } t \geq t_*,$$

where the last equivalence follows by Theorem 3.2. Hence, we can rewrite the bounds above saying that there exist $\underline{\kappa}, \bar{\kappa} > 0$ such that

$$\underline{\kappa} \frac{S(x)}{t^{1/(m-1)}} \leq u(t, x) \leq \bar{\kappa} \frac{S(x)}{t^{1/(m-1)}} \quad \text{for all } t \geq t_* \text{ and a.e. } x \in \Omega. \quad (7-3)$$

Since $t_* = \kappa_* \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)}$, the first inequality implies

$$\frac{S}{(t_* + t_0)^{1/(m-1)}} \leq \underline{\kappa} \frac{S}{t_*^{1/(m-1)}} \leq u(t_*)$$

for some $t_0 = c_0 \|u_0\|_{L^1_{\Phi_1}(\Omega)}^{-(m-1)} \geq t_*$. Hence, by the comparison principle,

$$\frac{S}{(t + t_0)^{1/(m-1)}} \leq u(t) \quad \text{for all } t \geq t_*.$$

On the other hand, it follows by (7-3) that $u(t, x) \leq \mathcal{U}_T(t, x) := S(x)(t - T)^{-1/(m-1)}$ for all $t \geq T$ provided T is large enough. If we now start to reduce T , the comparison principle combined with the upper bound (4-1) shows that u can never touch \mathcal{U}_T from below in $(T, \infty) \times \Omega$. Hence we can reduce T until $T = 0$, proving that $u \leq \mathcal{U}_0$; for an alternative proof, see Lemma 5.4 in [Bonforte et al. 2015]. Since $t_0 \geq t_*$, this shows that

$$\frac{S(x)}{(t + t_0)^{1/(m-1)}} \leq u(t, x) \leq \frac{S(x)}{t^{1/(m-1)}} \quad \text{for all } t \geq t_0 \text{ and a.e. } x \in \Omega.$$

Therefore

$$\left| 1 - \frac{u(t, x)}{\mathcal{U}(t, x)} \right| \leq 1 - \left(1 - \frac{t_0}{t_0 + t} \right)^{1/(m-1)} \leq \frac{2}{m-1} \frac{t_0}{t_0 + t} \quad \text{for all } t \geq t_0 \text{ and a.e. } x \in \Omega,$$

as desired. \square

8. Regularity results

In order to obtain the regularity results, we basically require the validity of a global Harnack principle, namely Theorem 6.1, 6.2, or 6.3, depending on the situation under study. For some higher-regularity results, we will eventually need some extra assumptions on the kernels. For simplicity we assume that \mathcal{L} is described by a kernel, without any lower-order term. However, it is clear that the presence of lower-order terms does not play any role in the interior regularity.

Theorem 8.1 (interior regularity). *Assume that*

$$\mathcal{L}f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y))K(x, y) dy + B(x)f(x),$$

with

$$K(x, y) \asymp |x - y|^{-(N+2s)} \quad \text{in } B_{2r}(x_0) \subset \Omega, \quad K(x, y) \lesssim |x - y|^{-(N+2s)} \quad \text{in } \mathbb{R}^N \setminus B_{2r}(x_0).$$

Let u be a nonnegative bounded weak dual solution to the CDP on $(T_0, T_1) \times \Omega$, and assume that there exist $\delta, M > 0$ such that

$$\begin{aligned} 0 < \delta \leq u(t, x) & \quad \text{for a.e. } (t, x) \in (T_0, T_1) \times B_{2r}(x_0), \\ 0 \leq u(t, x) \leq M & \quad \text{for a.e. } (t, x) \in (T_0, T_1) \times \Omega. \end{aligned}$$

(i) Then u is Hölder continuous in the interior. More precisely, there exists $\alpha > 0$ such that, for all $0 < T_0 < T_2 < T_1$,

$$\|u\|_{C_{t,x}^{\alpha/2s,\alpha}((T_2,T_1) \times B_r(x_0))} \leq C. \tag{8-1}$$

(ii) Assume in addition $|K(x, y) - K(x', y)| \leq c|x - x'|^\beta |y|^{-(N+2s)}$ for some $\beta \in (0, 1 \wedge 2s)$ such that $\beta + 2s$ is not an integer. Then u is a classical solution in the interior. More precisely, for all $0 < T_0 < T_2 < T_1$,

$$\|u\|_{C_{t,x}^{1+\beta/2s,2s+\beta}((T_2,T_1) \times B_r(x_0))} \leq C. \tag{8-2}$$

The constants in the above regularity estimates depend on the solution only through the upper and lower bounds on u . These bounds can be made quantitative by means of local Harnack inequalities, of elliptic and forward type, which follow from the global ones.

Theorem 8.2 (local Harnack inequalities of elliptic/backward type). *Under the assumptions of Theorem 6.1, there exists a constant $\widehat{H} > 0$, depending only on $N, s, \gamma, m, c_1, \underline{\kappa}_\Omega, \Omega$, such that for all balls $B_R(x_0)$ such that $B_{2R}(x_0) \subset \Omega$*

$$\sup_{x \in B_R(x_0)} u(t, x) \leq \frac{\widehat{H}}{(1 \wedge t/t_*)^{m/(m-1)}} \inf_{x \in B_R(x_0)} u(t, x) \quad \text{for all } t > 0. \tag{8-3}$$

Moreover, for all $t > 0$ and all $h > 0$ we have

$$\sup_{x \in B_R(x_0)} u(t, x) \leq \widehat{H} \left[\left(1 + \frac{h}{t}\right) \left(1 \wedge \frac{t}{t_*}\right)^{-m} \right]^{1/(m-1)} \inf_{x \in B_R(x_0)} u(t+h, x). \tag{8-4}$$

Proof. Recalling (6-1), the bound (8-3) follows easily from the following Harnack inequality for the first eigenfunction, see for instance [Bonforte et al. 2017b]:

$$\sup_{x \in B_R(x_0)} \Phi_1(x) \leq H_{N,s,\gamma,\Omega} \inf_{x \in B_R(x_0)} \Phi_1(x).$$

Since $u(t, x) \leq (1 + h/t)^{1/(m-1)} u(t+h, x)$, by the time monotonicity of $t \mapsto t^{1/(m-1)} u(t, x)$, (8-4) follows. □

Remark. The same result holds for large times $t \geq t_*$ as a consequence of Theorem 6.2. Already in the local case $s = 1$, these Harnack inequalities are stronger than the known two-sided inequalities valid for solutions to the Dirichlet problem for the classical porous medium equation, see [Aronson and Caffarelli 1983; Daskalopoulos and Kenig 2007; DiBenedetto 1988; 1993; DiBenedetto et al. 2012], which are

of forward type and are often stated in terms of the so-called intrinsic geometry. Note that elliptic and backward Harnack-type inequalities usually occur in the fast diffusion range $m < 1$ [Bonforte et al. 2012; Bonforte and Vázquez 2006; 2010; 2014], or for linear equations in bounded domains [Fabes et al. 1986; Safonov and Yuan 1999].

For sharp boundary regularity we need a GHP with matching powers, like Theorems 6.1 or 6.2, and when $s > \gamma/2$, we can also prove Hölder regularity up to the boundary. We leave to the interested reader to check that the presence of an extra term $B(x)u^m(t, x)$ with $0 \leq B(x) \leq c_1 \text{dist}(x, \partial\Omega)^{-2s}$ (as in the SFL) does not affect the validity of the next result. Indeed, when considering the scaling in (8-6), the lower term scales as $\widehat{B}_r u_r^m$ with $0 \leq \widehat{B}_r \leq c_1$ inside the unit ball B_1 .

Theorem 8.3 (Hölder continuity up to the boundary). *Under assumptions of Theorem 8.1(ii), assume in addition that $2s > \gamma$. Then u is Hölder continuous up to the boundary. More precisely, for all $0 < T_0 < T_2 < T_1$ there exists a constant $C > 0$ such that*

$$\|u\|_{C_{t,x}^{\gamma/m\vartheta, \gamma/m}((T_2, T_1) \times \Omega)} \leq C \quad \text{with } \vartheta := 2s - \gamma\left(1 - \frac{1}{m}\right). \tag{8-5}$$

Remark. Since we have $u(t, x) \asymp \Phi_1(x)^{1/m} \asymp \text{dist}(x, \partial\Omega)^{\gamma/m}$ (note that $2s > \gamma$ implies that $\sigma = 1$ and that $2sm \neq \gamma(m - 1)$), the spacial Hölder exponent is sharp, while the Hölder exponent in time is the natural one by scaling.

8A. Proof of interior regularity. The strategy to prove Theorem 8.1 follows the lines of [Bonforte et al. 2017a] but with some modifications. The basic idea is that, because u is bounded away from zero and infinity, the equation is nondegenerate and we can use parabolic regularity for nonlocal equations to obtain the results. More precisely, interior Hölder regularity will follow by applying $C_{t,x}^{\alpha/2s, \alpha}$ estimates of [Felsinger and Kassmann 2013] for a “localized” linear problem. Once Hölder regularity is established, under a Hölder continuity assumption on the kernel we can use the Schauder estimates proved in [Dong and Zhang 2016] to conclude.

8A1. Localization of the problem. Up to a rescaling, we can assume $r = 2$, $T_0 = 0$, $T_1 = 1$. Also, by a standard covering argument, it is enough to prove the results with $T_2 = \frac{1}{2}$.

Take a cutoff function $\rho \in C_c^\infty(B_4)$ such that $\rho \equiv 1$ on B_3 and $\eta \in C_c^\infty(B_2)$ a cutoff function such that $\eta \equiv 1$ on B_1 , and define $v = \rho u$. By construction $u = v$ on $(0, 1) \times B_3$. Since $\rho \equiv 1$ on B_3 , we can write the equation for v on the small cylinder $(0, 1) \times B_1$ as

$$\partial_t v(t, x) = -\mathcal{L}[v^m](t, x) + g(t, x) = -L_a v(t, x) + f(t, x) + g(t, x),$$

where

$$\begin{aligned} L_a[v](t, x) &:= \int_{\mathbb{R}^N} (v(t, x) - v(t, y))a(t, x, y)K(x, y) dy, \\ a(t, x, y) &:= \frac{v^m(t, x) - v^m(t, y)}{v(t, x) - v(t, y)}\eta(x - y) + [1 - \eta(x - y)] \\ &= m\eta(x - y) \int_0^1 [(1 - \lambda)v(t, x) + \lambda v(t, y)]^{m-1} d\lambda + [1 - \eta(x - y)], \end{aligned}$$

$$f(t, x) := \int_{\mathbb{R}^N \setminus B_1(x)} (v^m(t, x) - v^m(t, y) - v(t, x) + v(t, y))[1 - \eta(x - y)]K(x, y) \, dy,$$

$$g(t, x) := -\mathcal{L}[(1 - \rho^m)u^m](t, x) = \int_{\mathbb{R}^N \setminus B_3} (1 - \rho^m(y))u^m(t, y)K(x, y) \, dy$$

(recall that $(1 - \rho^m)u^m \equiv 0$ on $(0, 1) \times B_3$).

8A2. Hölder continuity in the interior. Set $b := f + g$, with f and g as above. It is easy to check that, since $K(x, y) \lesssim |x - y|^{-(N+2s)}$, we have $b \in L^\infty((0, 1) \times B_1)$. Also, since $0 < \delta \leq u \leq M$ inside $(0, 1) \times B_1$, there exists $\Lambda > 1$ such that $\Lambda^{-1} \leq a(t, x, y) \leq \Lambda$ for a.e. $(t, x, y) \in (0, 1) \times B_1 \times B_1$ with $|x - y| \leq 1$. This guarantees that the linear operator L_a is uniformly elliptic, so we can apply the results in [Felsinger and Kassmann 2013] to ensure that

$$\|v\|_{C_{t,x}^{\alpha/2s,\alpha}((1/2,1) \times B_{1/2})} \leq C(\|b\|_{L^\infty((0,1) \times B_1)} + \|v\|_{L^\infty((0,1) \times \mathbb{R}^N)})$$

for some universal exponent $\alpha > 0$. This proves Theorem 8.1(i).

8A3. Classical solutions in the interior. Now that we know that $u \in C^{\alpha/2s,\alpha}((\frac{1}{2}, 1) \times B_{1/2})$, we repeat the localization argument above with cutoff functions ρ and η supported inside $(\frac{1}{2}, 1) \times B_{1/2}$ to ensure that $v := \rho u$ is Hölder continuous in $(\frac{1}{2}, 1) \times \mathbb{R}^N$. Then, to obtain higher regularity we argue as follows.

Set $\beta_1 := \min\{\alpha, \beta\}$. Thanks to the assumption on K and Theorem 8.1(i), it is easy to check that $K_a(t, x, y) := a(t, x, y)K(x, y)$ satisfies

$$|K_a(t, x, y) - K_a(t', x', y)| \leq C(|x - x'|^{\beta_1} + |t - t'|^{\beta_1/2s})|y|^{-(N+2s)}$$

inside $(\frac{1}{2}, 1) \times B_{1/2}$. Also, $f, g \in C^{\beta_1/2s,\beta_1}((\frac{1}{2}, 1) \times B_{1/2})$. This allows us to apply the Schauder estimates from [Dong and Zhang 2016], see also [Chang-Lara and Kriventsov 2017], to obtain that

$$\|v\|_{C_{t,x}^{1+\beta_1/2s,2s+\beta}((3/4,1) \times B_{1/4})} \leq C(\|b\|_{C_{t,x}^{\beta/2s,\beta}((1/2,1) \times B_{1/2})} + \|v\|_{C_{t,x}^{\beta/2s,\beta}((1/2,1) \times \mathbb{R}^N)}).$$

In particular, $u \in C^{1+\beta_1/2s,2s+\beta_1}((\frac{3}{4}, 1) \times B_{1/8})$. In the case $\beta_1 = \beta$ we stop here. Otherwise we set $\alpha_1 := 2s + \beta$ and we repeat the argument above with $\beta_2 := \min\{\alpha_1, \beta\}$ in place of β_1 . In this way, we obtain that $u \in C^{1+\beta_1/2s,2s+\beta_1}((1 - 2^{-4}, 1) \times B_{2^{-5}})$. Iterating this procedure finitely many times, we finally obtain

$$u \in C^{1+\beta/2s,2s+\beta}((1 - 2^{-k}, 1) \times B_{2^{-k-1}})$$

for some universal k . Finally, a covering argument completes the proof of Theorem 8.1(ii).

8B. Proof of boundary regularity. The proof of Theorem 8.3 follows by scaling and interior estimates. Notice that the assumption $2s > \gamma$ implies $\sigma = 1$; hence $u(t)$ has matching upper and lower bounds.

Given $x_0 \in \Omega$, set $r = \text{dist}(x_0, \partial\Omega)/2$ and define

$$u_r(t, x) := r^{-\gamma/m} u(t_0 + r^\vartheta t, x_0 + rx), \quad \text{with } \vartheta := 2s - \gamma\left(1 - \frac{1}{m}\right). \tag{8-6}$$

Note that, because $2s > \gamma$, we have $\vartheta > 0$. With this definition, we see that u_r satisfies the equation $\partial_t u_r + \mathcal{L}_r u_r^m = 0$ in $\Omega_r := (\Omega - x_0)/r$, where

$$\mathcal{L}_r f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y)) K_r(x, y) \, dy, \quad K_r(x, y) := r^{N+2s} K(x_0 + rx, x_0 + ry).$$

Note that, since $\sigma = 1$, it follows by the GHP that $u(t) \asymp \text{dist}(x, \partial\Omega)^{\gamma/m}$. Hence,

$$0 < \delta \leq u_r(t, x) \leq M \quad \text{for all } t \in [r^{-\vartheta} T_0, r^{-\vartheta} T_1], \, x \in B_1,$$

with constants $\delta, M > 0$ that are independent of r and x_0 . In addition, using again that $u(t) \asymp \text{dist}(x, \partial\Omega)^{\gamma/m}$, we see that

$$u_r(t, x) \leq C(1 + |x|^{\gamma/m}) \quad \text{for all } t \in [r^{-\vartheta} T_0, r^{-\vartheta} T_1] \text{ and } x \in \mathbb{R}^N.$$

Noticing that $u_r^m(t, x) \leq C(1 + |x|^\gamma)$ and that $\gamma < 2s$ by assumption, we see that the tails of u_r will not create any problem. Indeed, for any $x \in B_1$,

$$\int_{\mathbb{R}^N \setminus B_2} u_r^m(t, y) K_r(x, y)^{-(N+2s)} \, dy \leq C \int_{\mathbb{R}^N \setminus B_2} |y|^\gamma |y|^{-(N+2s)} \, dy \leq \bar{C}_0,$$

where \bar{C}_0 is independent of r . This means that we can localize the problem using cutoff functions as done in [Section 8A1](#), and the integrals defining the functions f and g will converge uniformly with respect to x_0 and r . Hence, we can apply [Theorem 8.1\(ii\)](#) to get

$$\|u_r\|_{C^{1+\beta/2s, 2s+\beta}([r^{-\vartheta} T+1/2, r^{-\vartheta} T+1] \times B_{1/2})} \leq C \tag{8-7}$$

for all $T \in [T_0, T_1 - r^{-\vartheta}]$. Since $\gamma/m < 2s + \beta$ (because $\gamma < 2s$), it follows that

$$\|u_r\|_{L^\infty([r^{-\vartheta} T+1/2, r^{-\vartheta} T+1], C^{\gamma/m}(B_{1/2}))} \leq \|u_r\|_{C^{1+\beta/2s, 2s+\beta}([r^{-\vartheta} T+1/2, r^{-\vartheta} T+1] \times B_{1/2})} \leq C.$$

Noticing that

$$\sup_{t \in [r^{-\vartheta} T+1/2, r^{-\vartheta} T+1]} [u_r]_{C^{\gamma/m}(B_{1/2})} = \sup_{t \in [T+r^\vartheta/2, r^{-\vartheta} T+r^{-\vartheta}]} [u]_{C^{\gamma/m}(B_r(x_0))},$$

and that $T \in [T_0, T_1 - r^{-\vartheta}]$ and x_0 are arbitrary, arguing as in [\[Ros-Oton and Serra 2014\]](#) we deduce that, given $T_2 \in (T_0, T_1)$,

$$\sup_{t \in [T_2, T_1]} [u]_{C^{\gamma/m}(\Omega)} \leq C. \tag{8-8}$$

This proves the global Hölder regularity in space. To show the regularity in time, we start again from [\(8-7\)](#) to get

$$\|\partial_t u_r\|_{L^\infty([r^{-\vartheta} T+1/2, r^{-\vartheta} T+1] \times B_{1/2})} \leq C.$$

By scaling, this implies

$$\|\partial_t u\|_{L^\infty([T+r^\vartheta/2, r^{-\vartheta} T+r^{-\vartheta}] \times B_r(x_0))} \leq Cr^{\gamma/m-\vartheta},$$

and by the arbitrariness of T and x_0 we obtain (recall that $r = \text{dist}(x_0, \partial\Omega)/2$)

$$|\partial_t u(t, x)| \leq C \text{dist}(x, \partial\Omega)^{\gamma/m-\vartheta} \quad \text{for all } t \in [T_2, T_1], \, x \in \Omega. \tag{8-9}$$

Note that $\gamma/m - \vartheta = \gamma - 2s < 0$ by our assumption.

Now, given $t_0, t_1 \in [T_2, T_1]$ and $x \in \Omega$, we argue as follows: if $|t_0 - t_1| \leq \text{dist}(x, \partial\Omega)^\vartheta$ then we use (8-9) to get (recall that $\gamma/m - \vartheta < 0$)

$$|u(t_1, x) - u(t_0, x)| \leq C \text{dist}(x, \partial\Omega)^{\gamma/m - \vartheta} |t_0 - t_1| \leq C |t_0 - t_1|^{\gamma/m\vartheta}.$$

On the other hand, if $|t_0 - t_1| \geq \text{dist}(x, \partial\Omega)^\vartheta$, then we use (8-8) and the fact that u vanishes on $\partial\Omega$ to obtain

$$|u(t_1, x) - u(t_0, x)| \leq |u(t_1, x)| + |u(t_0, x)| \leq C \text{dist}(x, \partial\Omega)^{\gamma/m} \leq C |t_0 - t_1|^{\gamma/m\vartheta}.$$

This proves that u is $(\gamma/m\vartheta)$ -Hölder continuous in time, and completes the proof of Theorem 8.3. □

9. Numerical evidence

After discovering the unexpected boundary behavior, we looked for numerical confirmation. This has been given to us by the authors of [Cusimano et al. 2017], who exploited the analytical tools developed in this paper to support our results by means of accurate numerical simulations. We include here some of these simulations, courtesy of the authors. In all the figures we shall consider the spectral fractional Laplacian, so that $\gamma = 1$ (see Section 2A for more details).

We take $\Omega = (-1, 1)$, and we consider as initial datum the compactly supported function

$$u_0(x) = e^{4-1/((x-1/2)(x+1/2))} \chi_{|x| < 1/2}$$

appearing in the left of Figure 1. In all the other figures, the solid line represents either $\Phi_1^{1/m}$ or Φ_1^{1-2s} , while the dotted lines represent $t^{1/(m-1)}u(t)$ for different values of t , where $u(t)$ is the solution starting from u_0 . These choices are motivated by Theorems 5.3 and 5.5. Since the map $t \mapsto t^{1/(m-1)}u(t, x)$ is nondecreasing for all $x \in \Omega$ [Bonforte and Vázquez 2016, (2.3)], the lower dotted line corresponds to an earlier time with respect to the higher one.




Figure 1. On the left, the initial condition u_0 . On the right, the solid line represents $\Phi_1^{1/m}$, and the dotted lines represent $t^{1/(m-1)}u(t)$ at $t = 1$ and $t = 5$. The parameters are $m = 2$ and $s = \frac{1}{2}$; hence $\sigma = 1$. While $u(t)$ appears to behave as $\Phi_1 \propto \text{dist}(\cdot, \partial\Omega)$ for very short times, already at $t = 5$ it exhibits the matching boundary behavior predicted by Theorem 5.3.




Figure 2. In both pictures, the solid line represents $\Phi_1^{1/m}$. On the left, the dotted lines represent $t^{1/(m-1)}u(t)$ at $t = 30$ and $t = 150$, with parameters $m = 4$ and $s = \frac{3}{4}$; hence $\sigma = 1$. In this case $u(t)$ appears to behave as $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)$ for quite some time, and only around $t = 150$ it exhibits the matching boundary behavior predicted by [Theorem 5.3](#). On the right, the dotted lines represent $t^{1/(m-1)}u(t)$ at $t = 150$ and $t = 600$ with parameters $m = 4$ and $s = \frac{1}{5}$; hence $\sigma = \frac{8}{15} < 1$. In this case $u(t)$ seems to exhibit a linear boundary behavior even after long time (this linear boundary behavior is a universal lower bound for all times by [Theorem 5.1](#)). The second picture may lead one to conjecture that, in the case $\sigma < 1$ and $u_0 \lesssim \Phi_1$, the behavior $u(t) \asymp \Phi_1$ holds for all times. However, as shown in [Figure 3](#), there are cases when $u(t) \gg \Phi_1^{1-2s}$ for large times.




Figure 3. In both pictures we use the parameters $m = 2$ and $s = \frac{1}{10}$; hence $\sigma = \frac{2}{5} < 1$, and the solid line represents Φ_1^{1-2s} . On the left, the dotted lines represent $t^{1/(m-1)}u(t)$ at $t = 4$ and $t = 25$, on the right we see $t = 40$ and $t = 150$. Note that $u(t) \asymp \Phi_1$ for short times. Then, after some time, $u(t)$ starts looking more like Φ_1^{1-2s} , and for large times ($t = 150$) it becomes much larger than Φ_1^{1-2s} .

Comparing [Figures 2](#) and [3](#), it seems that when $\sigma < 1$ there is no hope of finding a universal behavior of solutions for large times. In particular, the bound provided by [\(1-6\)](#) seems to be optimal.

10. Complements, extensions and further examples

Elliptic versus parabolic. The exceptional boundary behaviors we have found for some operators and data came as a surprise to us, since the solution to the corresponding “elliptic setting” $\mathcal{L}S^m = S$ satisfies $S \asymp \Phi_1^{\sigma/m}$ (with a logarithmic correction when $2sm \neq \gamma(m - 1)$); hence separate-variable solutions always satisfy (1-5) (see (3-2) and Theorem 3.2).

About the kernel of operators of the spectral type. In this section we study the properties of the kernel of \mathcal{L} . While in some situations \mathcal{L} may not have a kernel (for instance, in the local case), in other situations that may not be so obvious from its definition. In the next lemma it is shown in particular that the SFL, defined by (2-4), admits a representation of the form (2-5). We state hereby the precise result, mentioned in [Abatangelo 2015] and proven in [Song and Vondraček 2003] for the SFL.

Lemma 10.1 (spectral kernels). *Let $s \in (0, 1)$, and let \mathcal{L} be the s -th spectral power of a linear elliptic second-order operator \mathcal{A} , and let $\Phi_1 \asymp \text{dist}(\cdot, \partial\Omega)^\gamma$ be the first positive eigenfunction of \mathcal{A} . Let $H(t, x, y)$ be the heat kernel of \mathcal{A} , and assume that it satisfies the following bounds: there exist constants $c_0, c_1, c_2 > 0$ such that for all $0 < t \leq 1$*

$$c_0 \left(\frac{\Phi_1(x)}{t^{\gamma/2}} \wedge 1 \right) \left(\frac{\Phi_1(y)}{t^{\gamma/2}} \wedge 1 \right) \frac{e^{-c_1|x-y|^2/t}}{t^{N/2}} \leq H(t, x, y) \leq c_0^{-1} \left(\frac{\Phi_1(x)}{t^{\gamma/2}} \wedge 1 \right) \left(\frac{\Phi_1(y)}{t^{\gamma/2}} \wedge 1 \right) \frac{e^{-|x-y|^2/(c_1 t)}}{t^{N/2}} \tag{10-1}$$

and

$$0 \leq H(t, x, y) \leq c_2 \Phi_1(x) \Phi_1(y) \quad \text{for all } t \geq 1. \tag{10-2}$$

Then the operator \mathcal{L} can be expressed in the form

$$\mathcal{L}f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y)) K(x, y) dy + B(x)u(x) \tag{10-3}$$

with a kernel $K(x, y)$ supported in $\bar{\Omega} \times \bar{\Omega}$ satisfying

$$K(x, y) \asymp \frac{1}{|x - y|^{N+2s}} \left(\frac{\Phi_1(x)}{|x - y|^\gamma} \wedge 1 \right) \left(\frac{\Phi_1(y)}{|x - y|^\gamma} \wedge 1 \right) \quad \text{and} \quad B(x) \asymp \Phi_1(x)^{-2s/\gamma}. \tag{10-4}$$

The proof of this lemma follows the ideas of [Song and Vondraček 2003]; indeed assumptions of Lemma 10.1 allow us to adapt the proof of that paper to our case with minor changes.

Method and generality. Our work is part of a current effort aimed at extending the theory of evolution equations of parabolic type to a wide class of nonlocal operators, in particular operators with general kernels that have been studied by various authors; see for instance [del Teso et al. 2017; de Pablo et al. 2016; Serra 2015]. Our approach is different from many others: indeed, even if the equation is nonlinear, we concentrate on the properties of the inverse operator \mathcal{L}^{-1} (more precisely, on its kernel given by the Green function \mathbb{G}), rather than on the operator \mathcal{L} itself. Once this setting is well-established and good linear estimates for the Green function are available, the calculations and estimates are very general. Hence, the method is applicable to a very large class of equations, both for elliptic and parabolic problems, as well

as to more general nonlinearities than $F(u) = u^m$; see also related comments in the works [Bonforte and Vázquez 2015; 2016; Bonforte et al. 2015].

Finite and infinite propagation. In all cases considered in the paper for $s < 1$ we prove that the solution becomes strictly positive inside the domain at all positive times. This is called *infinite speed of propagation*, a property that does not hold in the limit $s = 1$ for any $m > 1$ [Vázquez 2007] (in that case, finite speed of propagation holds and a free boundary appears). Previous results on this infinite speed of propagation can be found in [Bonforte et al. 2017a; de Pablo et al. 2012]. We recall that infinite speed of propagation is typical of the evolution with nonlocal operators representing long-range interactions, but it is not true for the standard porous medium equation; hence a trade-off takes place when both effects are combined. All our models fall on the side of infinite propagation, but we recall that finite propagation holds for a related nonlocal model called “nonlinear porous medium flow with fractional potential pressure”; see [Caffarelli and Vázquez 2011].

The local case. Since $2sm > \gamma(m - 1)$ when $s = 1$ (independently of $m > 1$), our results give a sharp behavior in the local case after a “waiting time”. Although this is well-known for the classical porous medium equation, our results apply also to the case of the uniformly elliptic operator in divergence form with C^1 coefficients, and yield new results in this setting. Actually one can check that, even when the coefficients are merely measurable, many of our results are still true and they provided universal upper and lower estimates. At least to our knowledge, such general results are completely new.

10A. Further examples of operators. Here we briefly exhibit a number of examples to which our theory applies, besides the RFL, CFL and SFL already discussed in Section 2. These include a wide class of local and nonlocal operators. We just sketch the essential points, referring to [Bonforte and Vázquez 2016] for a more detailed exposition.

Censored fractional Laplacian (CFL) and operators with more general kernels. As already mentioned in Section 2A, assumptions (A1), (A2), and (K2) are satisfied with $\gamma = s - \frac{1}{2}$. Moreover, it follows by [Bogdan et al. 2003; Chen et al. 2010] that we can also consider operators of the form:

$$\mathcal{L}f(x) = \text{P.V.} \int_{\Omega} (f(x) - f(y)) \frac{a(x, y)}{|x - y|^{N+2s}} dy, \quad \text{with } \frac{1}{2} < s < 1,$$

where $a(x, y)$ is a symmetric function of class C^1 bounded between two positive constants. The Green function $\mathbb{G}(x, y)$ of \mathcal{L} satisfies the stronger assumption (K4); see Corollary 1.2 of [Chen et al. 2010].

Fractional operators with more general kernels. Consider integral operators of the form

$$\mathcal{L}f(x) = \text{P.V.} \int_{\mathbb{R}^N} (f(x) - f(y)) \frac{a(x, y)}{|x - y|^{N+2s}} dy,$$

where a is a measurable symmetric function, bounded between two positive constants, and satisfying

$$|a(x, y) - a(x, x)| \chi_{|x-y|<1} \leq c|x - y|^\sigma, \quad \text{with } 0 < s < \sigma \leq 1,$$

for some $c > 0$ (actually, one can allow even more general kernels; see [Bonforte and Vázquez 2016; Kim and Kim 2014]). Then, for all $s \in (0, 1]$, the Green function $\mathbb{G}(x, y)$ of \mathcal{L} satisfies (K4) with $\gamma = s$; see Corollary 1.4 of [Kim and Kim 2014].

Spectral powers of uniformly elliptic operators. Consider a linear operator \mathcal{A} in divergence form,

$$\mathcal{A} = - \sum_{i,j=1}^N \partial_i (a_{ij} \partial_j),$$

with uniformly elliptic C^1 coefficients. The uniform ellipticity allows one to build a self-adjoint operator on $L^2(\Omega)$ with discrete spectrum (λ_k, ϕ_k) . Using the spectral theorem, we can construct the spectral power of such operator as

$$\mathcal{L} f(x) := \mathcal{A}^s f(x) := \sum_{k=1}^{\infty} \lambda_k^s \hat{f}_k \phi_k(x), \quad \text{where } \hat{f}_k = \int_{\Omega} f(x) \phi_k(x) dx$$

(we refer to the books [Davies 1990; 1995] for further details), and the Green function satisfies (K2) with $\gamma = 1$; see [Davies 1995, Chapter 4.6]. Then, the first eigenfunction Φ_1 is comparable to $\text{dist}(\cdot, \partial\Omega)$. Also, Lemma 10.1 applies, see for instance [Davies 1995], and allow us to get sharp upper and lower estimates for the kernel K of \mathcal{L} , as in (10-4).

Other examples. As explained in Section 3 of [Bonforte and Vázquez 2016], our theory may also be applied to: (i) sums of two fractional operators; (ii) the sum of the Laplacian and a nonlocal operator kernel; (iii) Schrödinger equations for nonsymmetric diffusions; (iv) gradient perturbation of restricted fractional Laplacians. Finally, it is worth mentioning that our arguments readily extend to operators on manifolds for which the required bounds hold.

Acknowledgments

Bonforte and Vázquez are partially funded by Project MTM2011-24696 and MTM2014-52240-P (Spain). Figalli was supported by NSF Grants DMS-1262411 and DMS-1361122, and by the ERC Grant “Regularity and Stability in Partial Differential Equations (RSPDE)”. Bonforte and Vázquez would like to acknowledge the hospitality of the Mathematics Department of the University of Texas at Austin, where part of this work was done. Vázquez was also invited by BCAM, Bilbao. We thank an anonymous referee for pointing out that Lemma 10.1 was proved in [Song and Vondraček 2003] and mentioned in [Abatangelo 2015].

References

- [Abatangelo 2015] N. Abatangelo, *Large solutions for fractional Laplacian operators*, Ph.D. thesis, Université de Picardie Jules Verne, 2015, available at <https://arxiv.org/abs/1511.00571>.
- [Aronson and Caffarelli 1983] D. G. Aronson and L. A. Caffarelli, “The initial trace of a solution of the porous medium equation”, *Trans. Amer. Math. Soc.* **280**:1 (1983), 351–366. MR Zbl
- [Aronson and Peletier 1981] D. G. Aronson and L. A. Peletier, “Large time behaviour of solutions of the porous medium equation in bounded domains”, *J. Differential Equations* **39**:3 (1981), 378–412. MR Zbl

- [Athanasopoulos and Caffarelli 2010] I. Athanasopoulos and L. A. Caffarelli, “Continuity of the temperature in boundary heat control problems”, *Adv. Math.* **224**:1 (2010), 293–315. [MR](#) [Zbl](#)
- [Blumenthal and Gettoor 1960] R. M. Blumenthal and R. K. Gettoor, “Some theorems on stable processes”, *Trans. Amer. Math. Soc.* **95** (1960), 263–273. [MR](#) [Zbl](#)
- [Bogdan et al. 2003] K. Bogdan, K. Burdzy, and Z.-Q. Chen, “Censored stable processes”, *Probab. Theory Related Fields* **127**:1 (2003), 89–152. [MR](#) [Zbl](#)
- [Bonforte and Vázquez 2006] M. Bonforte and J. L. Vázquez, “Global positivity estimates and Harnack inequalities for the fast diffusion equation”, *J. Funct. Anal.* **240**:2 (2006), 399–428. [MR](#) [Zbl](#)
- [Bonforte and Vázquez 2010] M. Bonforte and J. L. Vázquez, “Positivity, local smoothing, and Harnack inequalities for very fast diffusion equations”, *Adv. Math.* **223**:2 (2010), 529–578. [MR](#) [Zbl](#)
- [Bonforte and Vázquez 2014] M. Bonforte and J. L. Vázquez, “Quantitative local and global a priori estimates for fractional nonlinear diffusion equations”, *Adv. Math.* **250** (2014), 242–284. [MR](#) [Zbl](#)
- [Bonforte and Vázquez 2015] M. Bonforte and J. L. Vázquez, “A priori estimates for fractional nonlinear degenerate diffusion equations on bounded domains”, *Arch. Ration. Mech. Anal.* **218**:1 (2015), 317–362. [MR](#) [Zbl](#)
- [Bonforte and Vázquez 2016] M. Bonforte and J. L. Vázquez, “Fractional nonlinear degenerate diffusion equations on bounded domains, I: Existence, uniqueness and upper bounds”, *Nonlinear Anal.* **131** (2016), 363–398. [MR](#) [Zbl](#)
- [Bonforte et al. 2012] M. Bonforte, G. Grillo, and J. L. Vázquez, “Behaviour near extinction for the fast diffusion equation on bounded domains”, *J. Math. Pures Appl.* (9) **97**:1 (2012), 1–38. [MR](#) [Zbl](#)
- [Bonforte et al. 2015] M. Bonforte, Y. Sire, and J. L. Vázquez, “Existence, uniqueness and asymptotic behaviour for fractional porous medium equations on bounded domains”, *Discrete Contin. Dyn. Syst.* **35**:12 (2015), 5725–5767. [MR](#) [Zbl](#)
- [Bonforte et al. 2017a] M. Bonforte, A. Figalli, and X. Ros-Oton, “Infinite speed of propagation and regularity of solutions to the fractional porous medium equation in general domains”, *Comm. Pure Appl. Math.* **70**:8 (2017), 1472–1508. [MR](#) [Zbl](#)
- [Bonforte et al. 2017b] M. Bonforte, A. Figalli, and J. L. Vázquez, “Sharp boundary behaviour of solutions to semilinear nonlocal elliptic equations”, preprint, 2017. [arXiv](#)
- [Cabré and Tan 2010] X. Cabré and J. Tan, “Positive solutions of nonlinear problems involving the square root of the Laplacian”, *Adv. Math.* **224**:5 (2010), 2052–2093. [MR](#) [Zbl](#)
- [Caffarelli and Silvestre 2007] L. Caffarelli and L. Silvestre, “An extension problem related to the fractional Laplacian”, *Comm. Partial Differential Equations* **32**:7-9 (2007), 1245–1260. [MR](#) [Zbl](#)
- [Caffarelli and Vázquez 2011] L. Caffarelli and J. L. Vázquez, “Nonlinear porous medium flow with fractional potential pressure”, *Arch. Ration. Mech. Anal.* **202**:2 (2011), 537–565. [MR](#) [Zbl](#)
- [Chang-Lara and Kriventsov 2017] H. A. Chang-Lara and D. Kriventsov, “Further time regularity for nonlocal, fully nonlinear parabolic equations”, *Comm. Pure Appl. Math.* **70**:5 (2017), 950–977. [MR](#) [Zbl](#)
- [Chen et al. 2010] Z.-Q. Chen, P. Kim, and R. Song, “Two-sided heat kernel estimates for censored stable-like processes”, *Probab. Theory Related Fields* **146**:3-4 (2010), 361–399. [MR](#) [Zbl](#)
- [Cusimano et al. 2017] N. Cusimano, F. del Teso, L. Gerardo-Giorda, and G. Pagnini, “Discretizations of the spectral fractional Laplacian on general domains with Dirichlet, Neumann, and Robin boundary conditions”, preprint, 2017. [arXiv](#)
- [Dahlberg and Kenig 1988] B. E. J. Dahlberg and C. E. Kenig, “Nonnegative solutions of the initial-Dirichlet problem for generalized porous medium equations in cylinders”, *J. Amer. Math. Soc.* **1**:2 (1988), 401–412. [MR](#) [Zbl](#)
- [Daskalopoulos and Kenig 2007] P. Daskalopoulos and C. E. Kenig, *Degenerate diffusions: initial value problems and local regularity theory*, EMS Tracts in Mathematics **1**, European Mathematical Society, Zürich, 2007. [MR](#) [Zbl](#)
- [Davies 1990] E. B. Davies, *Heat kernels and spectral theory*, Cambridge Tracts in Mathematics **92**, Cambridge University Press, 1990. [MR](#) [Zbl](#)
- [Davies 1995] E. B. Davies, *Spectral theory and differential operators*, Cambridge Studies in Advanced Mathematics **42**, Cambridge University Press, 1995. [MR](#) [Zbl](#)
- [DiBenedetto 1988] E. DiBenedetto, “Intrinsic Harnack type inequalities for solutions of certain degenerate parabolic equations”, *Arch. Rational Mech. Anal.* **100**:2 (1988), 129–147. [MR](#) [Zbl](#)

- [DiBenedetto 1993] E. DiBenedetto, *Degenerate parabolic equations*, Springer, 1993. [MR](#) [Zbl](#)
- [DiBenedetto et al. 2012] E. DiBenedetto, U. Gianazza, and V. Vespi, *Harnack's inequality for degenerate and singular parabolic equations*, Springer, 2012. [MR](#) [Zbl](#)
- [Dong and Zhang 2016] H. Dong and H. Zhang, “On Schauder estimates for a class of nonlocal fully nonlinear parabolic equations”, preprint, 2016. [arXiv](#)
- [Fabes et al. 1986] E. B. Fabes, N. Garofalo, and S. Salsa, “A backward Harnack inequality and Fatou theorem for nonnegative solutions of parabolic equations”, *Illinois J. Math.* **30**:4 (1986), 536–565. [MR](#) [Zbl](#)
- [Felsinger and Kassmann 2013] M. Felsinger and M. Kassmann, “Local regularity for parabolic nonlocal operators”, *Comm. Partial Differential Equations* **38**:9 (2013), 1539–1573. [MR](#) [Zbl](#)
- [Kim and Kim 2014] K.-Y. Kim and P. Kim, “Two-sided estimates for the transition densities of symmetric Markov processes dominated by stable-like processes in $C^{1,\eta}$ open sets”, *Stochastic Process. Appl.* **124**:9 (2014), 3055–3083. [MR](#) [Zbl](#)
- [de Pablo et al. 2011] A. de Pablo, F. Quirós, A. Rodríguez, and J. L. Vázquez, “A fractional porous medium equation”, *Adv. Math.* **226**:2 (2011), 1378–1409. [MR](#) [Zbl](#)
- [de Pablo et al. 2012] A. de Pablo, F. Quirós, A. Rodríguez, and J. L. Vázquez, “A general fractional porous medium equation”, *Comm. Pure Appl. Math.* **65**:9 (2012), 1242–1284. [MR](#) [Zbl](#)
- [de Pablo et al. 2016] A. de Pablo, F. Quirós, and A. Rodríguez, “Nonlocal filtration equations with rough kernels”, *Nonlinear Anal.* **137** (2016), 402–425. [MR](#) [Zbl](#)
- [Ros-Oton and Serra 2014] X. Ros-Oton and J. Serra, “The Dirichlet problem for the fractional Laplacian: regularity up to the boundary”, *J. Math. Pures Appl.* (9) **101**:3 (2014), 275–302. [MR](#) [Zbl](#)
- [Safonov and Yuan 1999] M. V. Safonov and Y. Yuan, “Doubling properties for second order parabolic equations”, *Ann. of Math.* (2) **150**:1 (1999), 313–327. [MR](#) [Zbl](#)
- [Serra 2015] J. Serra, “Regularity for fully nonlinear nonlocal parabolic equations with rough kernels”, *Calc. Var. Partial Differential Equations* **54**:1 (2015), 615–629. [MR](#) [Zbl](#)
- [Song and Vondraček 2003] R. Song and Z. Vondraček, “Potential theory of subordinate killed Brownian motion in a domain”, *Probab. Theory Related Fields* **125**:4 (2003), 578–592. [MR](#) [Zbl](#)
- [del Teso et al. 2017] F. del Teso, J. Endal, and E. R. Jakobsen, “Uniqueness and properties of distributional solutions of nonlocal equations of porous medium type”, *Adv. Math.* **305** (2017), 78–143. [MR](#) [Zbl](#)
- [Vázquez 2004] J. L. Vázquez, “The Dirichlet problem for the porous medium equation in bounded domains: asymptotic behavior”, *Monatsh. Math.* **142**:1-2 (2004), 81–111. [MR](#) [Zbl](#)
- [Vázquez 2007] J. L. Vázquez, *The porous medium equation: mathematical theory*, Clarendon, New York, 2007. [MR](#) [Zbl](#)
- [Vázquez 2014a] J. L. Vázquez, “Barenblatt solutions and asymptotic behaviour for a nonlinear fractional heat equation of porous medium type”, *J. Eur. Math. Soc. (JEMS)* **16**:4 (2014), 769–803. [MR](#) [Zbl](#)
- [Vázquez 2014b] J. L. Vázquez, “Recent progress in the theory of nonlinear diffusion with fractional Laplacian operators”, *Discrete Contin. Dyn. Syst. Ser. S* **7**:4 (2014), 857–885. [MR](#) [Zbl](#)

Received 2 Feb 2017. Revised 31 Jul 2017. Accepted 22 Nov 2017.

MATTEO BONFORTE: matteo.bonforte@uam.es

Departamento de Matemáticas, Universidad Autónoma de Madrid, Campus de Cantoblanco, Madrid, Spain

ALESSIO FIGALLI: alessio.figalli@math.ethz.ch

ETH Zürich, Department of Mathematics, Zürich, Switzerland

JUAN LUIS VÁZQUEZ: juanluis.vazquez@uam.es

Departamento de Matemáticas, Universidad Autónoma de Madrid, Campus de Cantoblanco, Madrid, Spain

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rhm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 4 2018

C^1 regularity of orthotropic p -harmonic functions in the plane	813
PIERRE BOUSQUET and LORENZO BRASCO	
Applications of small-scale quantum ergodicity in nodal sets	855
HAMID HEZARI	
On rank-2 Toda systems with arbitrary singularities: local mass and new estimates	873
CHANG-SHOU LIN, JUN-CHENG WEI, WEN YANG and LEI ZHANG	
Beyond the BKM criterion for the 2D resistive magnetohydrodynamic equations	899
LÉO AGÉLAS	
On a bilinear Strichartz estimate on irrational tori	919
CHENJIE FAN, GIGLIOLA STAFFILANI, HONG WANG and BOBBY WILSON	
Sharp global estimates for local and nonlocal porous medium-type equations in bounded domains	945
MATTEO BONFORTE, ALESSIO FIGALLI and JUAN LUIS VÁZQUEZ	
Blow-up of a critical Sobolev norm for energy-subcritical and energy-supercritical wave equations	983
THOMAS DUYCKAERTS and JIANWEI YANG	
Global weak solutions for generalized SQG in bounded domains	1029
HUY QUANG NGUYEN	
Scale-free unique continuation principle for spectral projectors, eigenvalue-lifting and Wegner estimates for random Schrödinger operators	1049
IVICA NAKIĆ, MATTHIAS TÄUFER, MARTIN TAUTENHAHN and IVAN VESELIĆ	