

ANALYSIS & PDE

Volume 11

No. 7

2018

SUSANNA TERRACINI, GIORGIO TORTONE AND STEFANO VITA

ON s -HARMONIC FUNCTIONS ON CONES



ON s -HARMONIC FUNCTIONS ON CONES

SUSANNA TERRACINI, GIORGIO TORTONE AND STEFANO VITA

We deal with nonnegative functions satisfying

$$\begin{cases} (-\Delta)^s u_s = 0 & \text{in } C, \\ u_s = 0 & \text{in } \mathbb{R}^n \setminus C, \end{cases}$$

where $s \in (0, 1)$ and C is a given cone on \mathbb{R}^n with vertex at zero. We consider the case when s approaches 1, wondering whether solutions of the problem do converge to harmonic functions in the same cone or not. Surprisingly, the answer will depend on the opening of the cone through an auxiliary eigenvalue problem on the upper half-sphere. These conic functions are involved in the study of the nodal regions in the case of optimal partitions and other free boundary problems and play a crucial role in the extension of the Alt–Caffarelli–Friedman monotonicity formula to the case of fractional diffusions.

1. Introduction	1653
2. Homogeneous s -harmonic functions on cones	1659
3. Characteristic exponent $\gamma_s(C)$: properties and asymptotic behavior	1665
4. The limit for $s \nearrow 1$	1681
5. Uniform-in- s estimates in $C^{0,\alpha}$ on annuli	1686
References	1689

1. Introduction

Let $n \geq 2$ and let C be an open cone in \mathbb{R}^n with vertex in 0; for a given $s \in (0, 1)$, we consider the problem of the classification of nontrivial functions which are s -harmonic inside the cone and vanish identically outside, that is,

$$\begin{cases} (-\Delta)^s u_s = 0 & \text{in } C, \\ u_s \geq 0 & \text{in } \mathbb{R}^n, \\ u_s \equiv 0 & \text{in } \mathbb{R}^n \setminus C. \end{cases} \quad (1-1)$$

Here we define (see [Section 2](#) for the details)

$$(-\Delta)^s u(x) = C(n, s) \text{ p.v. } \int_{\mathbb{R}^n} \frac{u(x) - u(\eta)}{|x - \eta|^{n+2s}} d\eta,$$

Work partially supported by the ERC Advanced Grant 2013 no. 339958 *Complex patterns for strongly interacting dynamical systems - COMPAT* and by the INDAM-GNAMPA project *Aspetti non-locali in fenomeni di segregazione*. The authors wish to thank Alessandro Zilio for many fruitful conversations.

MSC2010: primary 35R11; secondary 35B45, 35B08.

Keywords: fractional Laplacian, conic functions, asymptotic behavior, Martin kernel.

where u is a sufficiently smooth function and

$$C(n, s) = \frac{2^{2s} s \Gamma(n/2 + s)}{\pi^{n/2} \Gamma(1 - s)} > 0, \tag{1-2}$$

where

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt.$$

The principal value is taken at $\eta = x$; hence, though u needs not to decay at infinity, it has to keep an algebraic growth with a power strictly smaller than $2s$ in order to make the above expression meaningful. By Theorem 3.2 in [Bañuelos and Bogdan 2004], it is known that there exists a homogeneous, nonnegative and nontrivial solution to (1-1) of the form

$$u_s(x) = |x|^{\gamma_s} u_s\left(\frac{x}{|x|}\right),$$

where $\gamma_s := \gamma_s(C)$ is a definite homogeneity degree (characteristic exponent), which depends on the cone. Moreover, such a solution is continuous in \mathbb{R}^n and unique, up to multiplicative constants. We can normalize it in such a way that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. We consider the case when s approaches 1, wondering whether solutions of the problem do converge to a harmonic function in the same cone and, in that case, which are the suitable spaces for convergence.

Such conic s -harmonic functions appear as limiting blow-up profiles and play a major role in many free boundary problems with fractional diffusions and in the study of the geometry of nodal sets, also in the case of partition problems; see, e.g., [Allen 2012; Barrios et al. 2015; Caffarelli et al. 2017; Dipierro et al. 2017; Garofalo and Ros-Oton 2017]. Moreover, as we shall see later, they are strongly involved with the possible extensions of the Alt–Caffarelli–Friedman monotonicity formula to the case of fractional diffusion. The study of their properties and, ultimately, their classification is therefore a major achievement in this setting. The problem of homogeneous s -harmonic functions on cones has been deeply studied in [Bañuelos and Bogdan 2004; Bogdan and Byczkowski 1999; Bogdan et al. 2015; Michalik 2006]. The present paper mainly focuses on the limiting behavior as $s \nearrow 1$.

Our problem (1-1) can be linked to a specific spectral problem of local nature in the upper half-sphere; indeed let us look at the extension technique popularized by the authors in [Caffarelli and Silvestre 2007], characterizing the fractional Laplacian in \mathbb{R}^n as the Dirichlet-to-Neumann map for a variable v depending on one more space dimension and satisfying

$$\begin{cases} L_s v = \operatorname{div}(y^{1-2s} \nabla v) = 0 & \text{in } \mathbb{R}_+^{n+1}, \\ v(x, 0) = u(x) & \text{on } \mathbb{R}^n. \end{cases} \tag{1-3}$$

Such an extension exists and is unique for a suitable class of functions u , see (2-1), and it is given by the formula

$$v(x, y) = \gamma(n, s) \int_{\mathbb{R}^n} \frac{y^{2s} u(\eta)}{(|x - \eta|^2 + y^2)^{n/2+s}} d\eta, \quad \text{where } \gamma(n, s)^{-1} := \int_{\mathbb{R}^n} \frac{1}{(|\eta|^2 + 1)^{n/2+s}} d\eta.$$

Then, the nonlocal original operator translates into a boundary derivative operator of Neumann type:

$$-\frac{C(n, s)}{\gamma(n, s)} \lim_{y \rightarrow 0} y^{1-2s} \partial_y v(x, y) = (-\Delta)^s u(x).$$

Now, let us consider an open region $\omega \subseteq S^{n-1} = \partial S_+^n$, with $S_+^n = S^n \cap \{y > 0\}$, and define the eigenvalue

$$\lambda_1^s(\omega) = \inf \left\{ \frac{\int_{S_+^n} y^{1-2s} |\nabla_{S^n} u|^2 \, d\sigma}{\int_{S_+^n} y^{1-2s} u^2 \, d\sigma} : u \in H^1(S_+^n; y^{1-2s} \, d\sigma) \setminus \{0\} \text{ and } u \equiv 0 \text{ in } S^{n-1} \setminus \omega \right\}.$$

Next, define the *characteristic exponent* of the cone C_ω spanned by ω (see Definition 2.1) as

$$\gamma_s(C_\omega) = \gamma_s(\lambda_1^s(\omega)), \tag{1-4}$$

where the function $\gamma_s(t)$ is defined by

$$\gamma_s(t) := \sqrt{\left(\frac{1}{2}(n - 2s)\right)^2 + t} - \frac{1}{2}(n - 2s).$$

Remark 1.1. There is a remarkable link between the nonnegative $\lambda_1^s(\omega)$ -eigenfunctions and the $\gamma_s(\lambda_1^s(\omega))$ -homogeneous L_s -harmonic functions: Let consider the spherical coordinates (r, θ) with $r > 0$ and $\theta \in S^n$. Let φ_s be the first nonnegative eigenfunction to $\lambda_1^s(\omega)$ and let v_s be its $\gamma_s(\lambda_1^s(\omega))$ -homogeneous extension to \mathbb{R}_+^{n+1} , i.e.,

$$v_s(r, \theta) = r^{\gamma_s(\lambda_1^s(\omega))} \varphi_s(\theta),$$

which is well-defined as soon as $\gamma_s(\lambda_1^s(\omega)) < 2s$ (as we shall see, this fact is always granted). By [Rüland 2015], the operator L_s can be decomposed as

$$L_s u = \sin^{1-2s}(\theta_n) \frac{1}{r^n} \partial_r (r^{n+1+2s} \partial_r u) + \frac{1}{r^{1+2s}} L_s^{S^n} u,$$

where $y = r \sin(\theta_n)$ and the Laplace–Beltrami-type operator is defined as

$$L_s^{S^n} u = \operatorname{div}_{S^n} (\sin^{1-2s}(\theta_n) \nabla_{S^n} u)$$

with ∇_{S^n} the tangential gradient on S^n . Then, we easily get that v_s is L_s -harmonic in the upper half-space. Moreover, its trace $u_s(x) = v_s(x, 0)$ is s -harmonic in the cone C_ω spanned by ω , vanishing identically outside; in other words u_s is a solution of our problem (1-1).

In a symmetric way, for the standard Laplacian, we consider the problem of γ -homogeneous functions which are harmonic inside the cone spanned by ω and vanish outside:

$$\begin{cases} -\Delta u_1 = 0 & \text{in } C_\omega, \\ u_1 \geq 0 & \text{in } \mathbb{R}^n, \\ u_1 = 0 & \text{in } \mathbb{R}^n \setminus C_\omega. \end{cases} \tag{1-5}$$

It is well known that the associated eigenvalue problem on the sphere is that of the Laplace–Beltrami operator with Dirichlet boundary conditions

$$\lambda_1(\omega) = \inf \left\{ \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 \, d\sigma}{\int_{S^{n-1}} u^2 \, d\sigma} : u \in H^1(S^{n-1}) \setminus \{0\} \text{ and } u = 0 \text{ in } S^{n-1} \setminus \omega \right\},$$

and the *characteristic exponent* of the cone C_ω is

$$\gamma(C_\omega) = \sqrt{\left(\frac{1}{2}(n-2)\right)^2 + \lambda_1(\omega)} - \frac{1}{2}(n-2) = \gamma_{s|s=1}(\lambda_1(\omega)). \tag{1-6}$$

In the classical case, the characteristic exponent enjoys a number of nice properties: It is minimal on spherical caps among sets having a given measure. Moreover, for the spherical caps, the eigenvalues enjoy a fundamental convexity property with respect to the colatitude θ [Alt et al. 1984; Friedland and Hayman 1976]. The convexity plays a major role in the proof of the Alt–Caffarelli–Friedman monotonicity formula, a key tool in the free boundary theory [Caffarelli and Salsa 2005].

Since the standard Laplacian can be viewed as the limiting operator of the family $(-\Delta)^s$ as $s \nearrow 1$, some questions naturally arise:

Problem 1.2. Is it true that:

- (a) $\lim_{s \rightarrow 1} \gamma_s(C) = \gamma(C)$?
- (b) $\lim_{s \rightarrow 1} u_s = u_1$ uniformly on compact sets, or better, in Hölder local norms?
- (c) for spherical caps of opening θ there is any convexity of the map $\theta \mapsto \lambda_1^s(\theta)$, at least, for s near 1?

We therefore addressed the problem of the asymptotic behavior of the solutions of problem (1-1) for $s \nearrow 1$, obtaining a rather unexpected result: our analysis shows high sensitivity to the opening solid angle ω of the cone C_ω , as evaluated by the value of $\gamma(C)$. In the case of wide cones, when $\gamma(C) < 2$ (that is, $\theta \in (\pi/4, \pi)$ for spherical caps of colatitude θ), our solutions do converge to the harmonic homogeneous function of the cone; in the case of narrow cones, when $\gamma(C) \geq 2$ (that is, $\theta \in (0, \pi/4]$ for spherical caps), the limit of the homogeneity degree will always be 2 and the limiting profile will be something different, though related, of course, through a correction term. Similar transition phenomena have been detected in other contexts for some types of free boundary problems on cones [Allen and Lara 2015; Shahgholian 2004]. As a consequence of our main result, we will see a lack of convexity of the eigenvalue as a function of the colatitude. Our main result is the following theorem.

Theorem 1.3. *Let C be an open cone with vertex at the origin. There exist the following finite limits:*

$$\bar{\gamma}(C) := \lim_{s \rightarrow 1^-} \gamma_s(C) = \min\{\gamma(C), 2\}, \tag{1-7}$$

$$\mu(C) := \lim_{s \rightarrow 1^-} \frac{C(n, s)}{2s - \gamma_s(C)} = \begin{cases} 0 & \text{if } \gamma(C) \leq 2, \\ \mu_0(C) & \text{if } \gamma(C) \geq 2, \end{cases} \tag{1-8}$$

where $C(n, s)$ is defined in (1-2) and

$$\mu_0(C) := \inf \left\{ \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 - 2nu^2 \, d\sigma}{\left(\int_{S^{n-1}} |u| \, d\sigma\right)^2} : u \in H^1(S^{n-1}) \setminus \{0\} \text{ and } u = 0 \text{ in } S^{n-1} \setminus C \right\}.$$

Let us consider the family (u_s) of nonnegative solutions to (1-1) such that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. Then, as $s \nearrow 1$, up to a subsequence, we have:

- (1) $u_s \rightarrow \bar{u}$ in $L^2_{\text{loc}}(\mathbb{R}^n)$ for some $\bar{u} \in H^1_{\text{loc}}(\mathbb{R}^n) \cap L^\infty(S^{n-1})$.




Figure 1. Characteristic exponents of spherical caps of aperture 2θ for $s < 1$ and $s = 1$.

- (2) The convergence is uniform on compact subsets of C and \bar{u} is nontrivial with $\|\bar{u}\|_{L^\infty(S^{n-1})} = 1$ and is $\bar{\gamma}(C)$ -homogeneous.
- (3) The limit \bar{u} solves

$$\begin{cases} -\Delta \bar{u} = \mu(C) \int_{S^{n-1}} \bar{u} \, d\sigma & \text{in } C, \\ \bar{u} = 0 & \text{in } \mathbb{R}^n \setminus C. \end{cases} \tag{1-9}$$

Remark 1.4. Uniqueness of the limit \bar{u} and therefore existence of the limit of u_s as $s \nearrow 1$ hold in the case of connected cones and, in any case, whenever $\gamma(C) > 2$. We will see in Remark 4.2 that under symmetry assumptions on the cone C , the limit function \bar{u} is unique and hence it does not depend on the choice of the subsequence.

A nontrivial improvement of the main theorem concerns uniform bounds in Hölder spaces holding uniformly for $s \rightarrow 1$.

Theorem 1.5. Assume the cone is $C^{1,1}$. Let $\alpha \in (0, 1)$, $s_0 \in (\max\{\frac{1}{2}, \alpha\}, 1)$ and A be an annulus centered at zero. Then the family of solutions u_s to (1-1) is uniformly bounded in $C^{0,\alpha}(A)$ for any $s \in [s_0, 1)$.

On the fractional Alt–Caffarelli–Friedman monotonicity formula. In the case of reaction-diffusion systems with strong competition between a number of densities which spread in space, one can observe a segregation phenomenon: as the interspecific competition rate grows, the populations tend to separate their supports in nodal sets, separated by a free boundary. For the case of standard diffusion, both the asymptotic analysis and the properties of the segregated limiting profiles are fairly well understood, we refer to [Caffarelli and Lin 2008; Conti et al. 2005; Dancer et al. 2012; Noris et al. 2010; Tavares and Terracini 2012]. Instead, when the diffusion is nonlocal and modeled by the fractional Laplacian, the only known results are contained in [Terracini et al. 2014; 2016; Terracini and Vita 2017; Wang and Wei 2016]. As shown in [Terracini et al. 2014; 2016], estimates in Hölder spaces can be obtained by the use of fractional versions of the Alt–Caffarelli–Friedman (ACF) and Almgren monotonicity formulas. For the statement, proof and applications of the original ACF monotonicity formula we refer to the book

[Caffarelli and Salsa 2005] on free boundary problems. Let us state here the fractional version of the spectral problem beyond the ACF formula used in [Terracini et al. 2014; 2016]: consider the set of 2-partitions of S^{n-1} as

$$\mathcal{P}^2 := \{(\omega_1, \omega_2) : \omega_i \subseteq S^{n-1} \text{ open, } \omega_1 \cap \omega_2 = \emptyset, \bar{\omega}_1 \cup \bar{\omega}_2 = S^{n-1}\}$$

and define the optimal partition value as

$$v_s^{\text{ACF}} := \frac{1}{2} \inf_{(\omega_1, \omega_2) \in \mathcal{P}^2} \sum_{i=1}^2 \gamma_s(\lambda_1^s(\omega_i)). \tag{1-10}$$

It is easy to see, by a Schwarz symmetrization argument, that v_s^{ACF} is achieved by a pair of complementary spherical caps $(\omega_\theta, \omega_{\pi-\theta}) \in \mathcal{P}^2$ with aperture 2θ and $\theta \in (0, \pi)$ (for a detailed proof of this kind of symmetrization we refer to [Terracini and Vita 2017]); that is,

$$v_s^{\text{ACF}} = \min_{\theta \in [0, \pi]} \Gamma^s(\theta) = \min_{\theta \in [0, \pi]} \frac{\gamma_s(\theta) + \gamma_s(\pi - \theta)}{2}.$$

This gives a further motivation to our study of (1-1) for spherical caps. A classical result in [Friedland and Hayman 1976] yields $v^{\text{ACF}} = 1$ (in the case $s = 1$), and the minimal value is achieved for two half-spheres; this equality is the core of the proof of the classical Alt–Caffarelli–Friedman monotonicity formula.

It was proved in [Terracini et al. 2014] that v_s^{ACF} is linked to the threshold for uniform bounds in Hölder norms for competition-diffusion systems, as the interspecific competition rate diverges to infinity, as well as the exponent of the optimal Hölder regularity for their limiting profiles. It was also conjectured that $v_s^{\text{ACF}} = s$ for every $s \in (0, 1)$. Unfortunately, the exact value of v_s^{ACF} is still unknown, and we only know that $0 < v_s^{\text{ACF}} \leq s$; see [Terracini et al. 2014; 2016]. Actually one can easily give a better lower bound given by $v_s^{\text{ACF}} \geq \max\{\frac{1}{2}s, s - \frac{1}{4}\}$ when $n = 2$ and $v_s^{\text{ACF}} \geq \frac{1}{2}s$ otherwise, which however is not satisfactory. As already remarked in [Allen 2012], this lack of information implies also the lack of an exact Alt–Caffarelli–Friedman monotonicity formula for the case of fractional Laplacians. Our contribution to this open problem is a byproduct of the main result, Theorem 1.3, and is depicted in Figure 2.

Corollary 1.6. *In any space dimension we have*

$$\lim_{s \rightarrow 1} v_s^{\text{ACF}} = 1.$$

The paper is organized as follows. In Section 2 we introduce our setting and we state the relevant known properties of homogeneous s -harmonic functions on cones. After this, we will obtain local $C^{0,\alpha}$ -estimates in compact subsets of C and local H^s -estimates in compact subsets of \mathbb{R}^n for solutions u_s of (1-1). We will see that an important quantity which appears in these estimates and plays a fundamental role is

$$\frac{C(n, s)}{2s - \gamma_s(C)},$$

where $C(n, s) > 0$ is the normalization constant given in (1-2). It will be therefore very important to bound this quantity uniformly in s . In Section 3 we analyze the asymptotic behavior of $\gamma_s(C)$ as s converges to 1, in order to understand the quantities $\bar{\gamma}(C)$ and $\mu(C)$. To do this, we will establish a distributional




Figure 2. Possible values of $\Gamma^s(\theta) = \Gamma^s(\omega_\theta, \omega_{\pi-\theta})$ for $s < 1$ and $s = 1$ and $n = 2$.

semigroup property for the fractional Laplacian for functions which grow at infinity. In [Section 4](#) we prove [Theorem 1.3](#) and [Corollary 1.6](#). Eventually, in [Section 5](#), we prove [Theorem 1.5](#).

2. Homogeneous s -harmonic functions on cones

In this section, we focus our attention on the local properties of homogeneous s -harmonic functions on *regular cones*. Since in the next section we will study the behavior of the characteristic exponent as s approaches 1, in this section we recall some known results related to the boundary behavior of the solution of (1-1) restricted to the unitary sphere S^{n-1} and some estimates of the Hölder and H^s seminorms.

Definition 2.1. Let $\omega \subset S^{n-1}$ be an open set, which may be disconnected. We define the *unbounded cone* with vertex in 0, spanned by ω , to be the open set

$$C_\omega = \{rx : r > 0, x \in \omega\}.$$

Moreover we say that $C = C_\omega$ is narrow if $\gamma(C) \geq 2$ and wide if $\gamma(C) < 2$. We call C_ω a *regular cone* if ω is connected and of class $C^{1,1}$. Let $\theta \in (0, \pi)$ and $\omega_\theta \subset S^{n-1}$ be an open spherical cap of colatitude θ . Then we denote by $C_\theta = C_{\omega_\theta}$ the *right circular cone* of aperture 2θ .

Hence, let C be a fixed unbounded open cone in \mathbb{R}^n with vertex in 0 and consider

$$\begin{cases} (-\Delta)^s u_s = 0 & \text{in } C, \\ u_s \equiv 0 & \text{in } \mathbb{R}^n \setminus C \end{cases}$$

with the condition $\|u_s\|_{L^\infty(S^{n-1})} = 1$. By Theorem 3.2 in [\[Bañuelos and Bogdan 2004\]](#) there exists, up to a multiplicative constant, a unique nonnegative function u_s smooth in C and $\gamma_s(C)$ -homogeneous, i.e.,

$$u_s(x) = |x|^{\gamma_s(C)} u_s\left(\frac{x}{|x|}\right),$$

where $\gamma_s(C) \in (0, 2s)$. As is well known, see for example [Bogdan and Byczkowski 1999; Silvestre 2005], the fractional Laplacian $(-\Delta)^s$ is a nonlocal operator well-defined in the class of integrability $\mathcal{L}_s^1 := \mathcal{L}^1(dx/(1 + |x|)^{n+2s})$, namely the normed space of all Borel functions u satisfying

$$\|u\|_{\mathcal{L}_s^1} := \int_{\mathbb{R}^n} \frac{|u(x)|}{(1 + |x|)^{n+2s}} dx < \infty. \tag{2-1}$$

Hence, for every $u \in \mathcal{L}_s^1$, $\varepsilon > 0$ and $x \in \mathbb{R}^n$ we define

$$(-\Delta)_\varepsilon^s u(x) = C(n, s) \int_{\mathbb{R}^n \setminus B_\varepsilon(x)} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy,$$

where

$$C(n, s) = \frac{2^{2s}s\Gamma(n/2 + s)}{\pi^{n/2}\Gamma(1 - s)} \in (0, 4\Gamma(n/2 + 1)].$$

and we can consider the fractional Laplacian as the limit

$$(-\Delta)^s u(x) = \lim_{\varepsilon \downarrow 0} (-\Delta)_\varepsilon^s u(x) = C(n, s) \text{ p.v. } \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+2s}} dy.$$

We remark that $u \in \mathcal{L}_s^1$ is such that $u \in \mathcal{L}_{s+\delta}^1$ for any $\delta > 0$, which will be an important tool in this section of the paper, in order to compute high-order fractional Laplacians. Another definition of the fractional Laplacian, which can be constructed by a double change of variables as in [Di Nezza et al. 2012], is

$$(-\Delta)^s u(x) = \frac{C(n, s)}{2} \int_{\mathbb{R}^n} \frac{2u(x) - u(x + y) - u(x - y)}{|y|^{n+2s}} dy,$$

which emphasizes that given $u \in C^2(D) \cap \mathcal{L}_s^1$, we obtain that $x \mapsto (-\Delta)^s u(x)$ is a continuous and bounded function on D for some bounded $D \subset \mathbb{R}^n$.

By [Michalik 2006, Lemma 3.3], if we consider a regular unbounded cone C symmetric with respect to a fixed axis, there exist two positive constants $c_1 = c_1(n, s, C)$ and $c_2 = c_2(n, s, C)$ such that

$$c_1|x|^{\gamma_s-s} \text{dist}(x, \partial C)^s \leq u_s(x) \leq c_2|x|^{\gamma_s-s} \text{dist}(x, \partial C)^s \tag{2-2}$$

for every $x \in C$. We remark that this result can be easily generalized to regular unbounded cones C_ω with $\omega \subset S^{n-1}$ a finite union of connected $C^{1,1}$ domains ω_i such that $\bar{\omega}_i \cup \bar{\omega}_j = \emptyset$ for $i \neq j$, since the reasoning in [Michalik 2006] relies on a boundary Harnack principle and on sharp estimates for the Green function for bounded $C^{1,1}$ domains which are not necessarily connected; for more details see [Chen and Song 1998].

Throughout the paper we will call the coefficient of homogeneity γ_s the “characteristic exponent”, since it is strictly related to an eigenvalue partition problem.

As we already mentioned, our solutions are smooth in the interior of the cone and locally $C^{0,s}$ near the boundary $\partial C \setminus \{0\}$, see for example [Michalik 2006], but we need some quantitative estimates in order to better understand the dependence of the Hölder seminorm on the parameter $s \in (0, 1)$.

Before showing the main result of Hölder regularity, we need the following estimates about the fractional Laplacian of smooth compactly supported functions; this result can be found in [Bogdan and

Byczkowski 1999, Lemma 3.5; Dávila et al. 2015, Lemma 5.1], but here we compute the formula with a deep attention on the dependence of the constant with respect to $s \in (0, 1)$.

Proposition 2.2. *Let $s \in (0, 1)$ and $\varphi \in C_c^2(\mathbb{R}^n)$. Then*

$$|(-\Delta)^s \varphi(x)| \leq \frac{c}{(1 + |x|)^{n+2s}} \quad \text{for all } x \in \mathbb{R}^n, \tag{2-3}$$

where the constant $c > 0$ depends only on n and the choice of φ .

Proof. Let $K \subset \mathbb{R}^n$ be the compact support of φ and $k = \max_{x \in K} |\varphi(x)|$. There exists $R > 1$ such that $K \subset B_{R/2}(0)$.

Let $|x| > R$. Then

$$\begin{aligned} |(-\Delta)^s \varphi(x)| &= \left| C(n, s) \int_{\mathbb{R}^n} \frac{\varphi(x) - \varphi(y)}{|x - y|^{n+2s}} dy \right| = \left| C(n, s) \int_K \frac{\varphi(y)}{|x - y|^{n+2s}} dy \right| \\ &\leq \frac{C(n, s)k}{|x|^{n+2s}} \int_K \frac{1}{(1 - |y/x|)^{n+2s}} dy \leq \frac{C(n, s)k2^{n+2s}|K|}{|x|^{n+2s}} \\ &\leq \frac{C(n, s)k2^{2(n+2s)}|K|}{(1 + |x|)^{n+2s}} \leq \frac{c}{(1 + |x|)^{n+2s}}, \end{aligned}$$

where $c > 0$ depends only on n and the choice of φ .

Let now $|x| \leq R$. We use the fact that any derivative of φ of first or second order is uniformly continuous in the compact set K and the fact that in $B_R(0)$ the function $(1 + |x|)^{n+2s}$ has maximum given by $(1 + R)^{n+2s}$. Hence there exist $0 < \delta < 1$ and a constant $M > 0$, both depending only on n and the choice of φ , such that

$$|\varphi(x + z) + \varphi(x - z) - 2\varphi(x)| \leq M|z|^2 \quad \text{for all } z \in B_\delta(0).$$

Hence

$$\begin{aligned} |(-\Delta)^s \varphi(x)| &= \left| C(n, s) \int_{\mathbb{R}^n \setminus B_\delta(x)} \frac{\varphi(x) - \varphi(y)}{|x - y|^{n+2s}} dy + C(n, s) \int_{B_\delta(x)} \frac{\varphi(x) - \varphi(y)}{|x - y|^{n+2s}} dy \right| \\ &\leq 2kC(n, s) \int_{\mathbb{R}^n \setminus B_\delta(x)} \frac{1}{|x - y|^{n+2s}} dy + \frac{C(n, s)}{2} \int_{B_\delta(0)} \frac{|\varphi(x + z) + \varphi(x - z) - 2\varphi(x)|}{|z|^{n+2s}} dz \\ &\leq 2kC(n, s)\omega_{n-1} \int_\delta^\infty r^{-1-2s} dr + \frac{C(n, s)\omega_{n-1}M}{2} \int_0^\delta r^{1-2s} dr \\ &= \frac{kC(n, s)\omega_{n-1}}{s\delta^{2s}} + \frac{C(n, s)\omega_{n-1}M\delta^{2-2s}}{4(1 - s)} \\ &\leq \frac{c}{\delta^2} + c = c \frac{(1 + |x|)^{n+2s}}{(1 + |x|)^{n+2s}} \leq \frac{c(1 + R)^{n+2}}{(1 + |x|)^{n+2s}} = \frac{c}{(1 + |x|)^{n+2s}}, \end{aligned}$$

where $c > 0$ depends only on n and the choice of φ . □

By the previous calculations we have also the following result.

Remark 2.3. Let $s \in (0, 1)$ and $\varphi \in C_c^2(\mathbb{R}^n)$. Then there exists a constant $c = c(n, \varphi) > 0$ and a radius $R = R(\varphi) > 0$ such that

$$|(-\Delta)^s \varphi(x)| \leq c \frac{C(n, s)}{(1 + |x|)^{n+2s}} \quad \text{for all } x \in \mathbb{R}^n \setminus B_R(0). \tag{2-4}$$

The following result provides interior estimates for the Hölder norm of our solutions.

Proposition 2.4. Let C be a cone, $K \subset C$ be a compact set and $s_0 \in (0, 1)$. Then there exist a constant $c > 0$ and $\bar{\alpha} \in (0, 1)$, both dependent only on s_0, K, n, C , such that

$$\|u_s\|_{C^{0,\alpha}(K)} \leq c \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)}\right)$$

for any $\alpha \in (0, \bar{\alpha}]$ and any $s \in [s_0, 1)$.

By a standard covering argument, there exists a finite number of balls such that $K \subset \bigcup_{j=1}^k B_r(x_j)$ for a given radius $r > 0$ such that $\bigcup_{j=1}^k \overline{B_{2r}(x_j)} \subset C$. Thus, it is enough to prove:

Proposition 2.5. Let $\overline{B_{2r}(\bar{x})} \subset C$ be a closed ball and $s_0 \in (0, 1)$. Then there exist a constant $c > 0$ and $\bar{\alpha} \in (0, 1)$, both dependent only on s_0, r, \bar{x}, n, C , such that

$$\|u_s\|_{C^{0,\alpha}(\overline{B_r(\bar{x})})} \leq c \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)}\right)$$

for any $\alpha \in (0, \bar{\alpha}]$ and any $s \in [s_0, 1)$.

In order to achieve the desired result, we need to estimate locally the value of the fractional Laplacian of u_s in a ball compactly contained in the cone C .

Lemma 2.6. Let $\eta \in C_c^\infty(B_{2r}(\bar{x}))$ be a cut-off function such that $0 \leq \eta \leq 1$ with $\eta \equiv 1$ in $B_r(\bar{x})$. Under the same assumptions as [Proposition 2.5](#),

$$\|(-\Delta)^s(u_s \eta)\|_{L^\infty(B_{2r}(\bar{x}))} \leq C_0 \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)}\right)$$

for any $s \in [s_0, 1)$, where $C_0 > 0$ depends on s_0, n, \bar{x}, r, C , and the choice of the function η .

Proof. Let $R > 1$ be such that $\overline{B_{2r}(\bar{x})} \subset B_{R/2}(0)$. Hence, let us fix a point $x \in B_{2r}(\bar{x})$. We can express the fractional Laplacian of $u_s \eta$ in the following way:

$$\begin{aligned} (-\Delta)^s(u_s \eta)(x) &= \eta(x)(-\Delta)^s u_s(x) + C(n, s) \int_{\mathbb{R}^n} u_s(y) \frac{\eta(x) - \eta(y)}{|x - y|^{n+2s}} dy \\ &= C(n, s) \int_{B_R(0)} u_s(y) \frac{\eta(x) - \eta(y)}{|x - y|^{n+2s}} dy + C(n, s) \int_{\mathbb{R}^n \setminus B_R(0)} u_s(y) \frac{\eta(x) - \eta(y)}{|x - y|^{n+2s}} dy. \end{aligned}$$

We recall that $u_s(x) = |x|^{\gamma_s(C)} u_s(x/|x|)$ and that for any $s \in (0, 1)$ the functions u_s are normalized such that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. Moreover we remark that $\eta(x) - \eta(y) = \eta(x) \geq 0$ in $B_{2r}(\bar{x}) \times (\mathbb{R}^n \setminus B_R(0))$. Hence,

using Proposition 2.2 and the fact that $\gamma_s(C) < 2s$, we obtain

$$\begin{aligned} |(-\Delta)^s(u_s \eta)(x)| &\leq C(n, s) \left| \int_{B_R(0)} u_s(y) \frac{\eta(x) - \eta(y)}{|x - y|^{n+2s}} dy \right| + C(n, s) \left| \int_{\mathbb{R}^n \setminus B_R(0)} u_s(y) \frac{\eta(x) - \eta(y)}{|x - y|^{n+2s}} dy \right| \\ &\leq R^{\gamma_s(C)} |(-\Delta)^s \eta(x)| + C(n, s) 2^{n+2s} \int_{\mathbb{R}^n \setminus B_R(0)} \frac{1}{|y|^{n+2s-\gamma_s(C)}} dy \\ &\leq \frac{cR^2}{(1 + |x|)^{n+2s}} + C(n, s) 2^{n+2} \omega_{n-1} \int_R^\infty r^{-1-2s+\gamma_s(C)} dr \\ &\leq \frac{cR^2}{(1 + |x|)^{n+2s}} + \frac{cC(n, s)}{R^{2s-\gamma_s(C)}(2s - \gamma_s(C))} \\ &\leq C_0 \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)} \right). \end{aligned} \quad \square$$

Proof of Proposition 2.5. Let as before $\eta \in C_c^\infty(B_{2r}(\bar{x}))$ be a cut-off function such that $0 \leq \eta \leq 1$ with $\eta \equiv 1$ in $B_r(\bar{x})$. First, we remark that there exists a constant $c_0 > 0$ such that for any $s \in (0, 1)$

$$\|u_s \eta\|_{L^\infty(\mathbb{R}^n)} \leq c_0, \tag{2-5}$$

where c_0 depends only on n, \bar{x}, r . In fact, let $R > 0$ be such that $\overline{B_{2r}(\bar{x})} \subset B_R(0)$. Then, for any $x \in \mathbb{R}^n$, we have $0 \leq u_s \eta(x) \leq R^{\gamma_s(C)} \leq R^2$. Using the bound (2-5) and the previous lemma, we can apply [Caffarelli and Silvestre 2009, Theorem 12.1] obtaining the existence of $\bar{\alpha} \in (0, 1)$ and $C > 0$, both depending only on n, s_0 and the choice of $B_r(\bar{x})$ such that

$$\begin{aligned} \|u_s \eta\|_{C^{0,\alpha}(\overline{B_r(\bar{x})})} &\leq C(\|u_s \eta\|_{L^\infty(\mathbb{R}^n)} + \|(-\Delta)^s(u_s \eta)\|_{L^\infty(B_{2r}(\bar{x}))}) \\ &\leq C \left(c_0 + C_0 \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)} \right) \right) \end{aligned}$$

for any $s \in [s_0, 1)$ and any $\alpha \in (0, \bar{\alpha}]$. Since $\eta \equiv 1$ in $B_r(\bar{x})$ we obtain the result. □

Similarly, now we need to construct some estimate related to the H^s seminorm of the solution u_s . Since the functions do not belong to $H^s(\mathbb{R}^n)$, we need to truncate the solution with some cut-off function in order to avoid the problems related to the growth at infinity. In such a way, we can use

$$[v]_{H^s(\mathbb{R}^n)}^2 = \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 = \int_{\mathbb{R}^n} v(-\Delta)^s v dx, \tag{2-6}$$

which holds for every $v \in H^s(\mathbb{R}^n)$. So, let $\eta \in C_c^\infty(B_2)$ be a radial cut-off function such that $\eta \equiv 1$ in B_1 and $0 \leq \eta \leq 1$ in B_2 , and consider $\eta_R(x) = \eta((x - x_0)/R)$, the rescaled cut-off function defined in $B_{2R}(x_0)$ for some $R > 0$ and $x_0 \in \mathbb{R}^n$.

Proposition 2.7. *Let $s_0 \in (0, 1)$ and $\eta_R \in C_c^\infty(B_{2R}(x_0))$ be previously defined. Then*

$$[u_s \eta_R]_{H^s(\mathbb{R}^n)}^2 \leq c \left(1 + \frac{C(n, s)}{2s - \gamma_s(C)} \right)$$

for any $s \in [s_0, 1)$, where $c > 0$ is a constant that depends on x_0, R, C, s_0 and η .

Proof. Let $\eta \in C_c^\infty(B_2)$ be a radial cut-off function such that $\eta \equiv 1$ in B_1 and $0 \leq \eta \leq 1$ in B_2 , and consider the collection of $(\eta_R)_R$ with $R > 0$ defined by $\eta_R(x) = \eta((x - x_0)/R)$ with some $x_0 \in \mathbb{R}^n$. By (2-6), for every $R > 0$ we obtain

$$[u_s \eta_R]_{H^s(\mathbb{R}^n)}^2 = \|(-\Delta)^{s/2}(u_s \eta_R)\|_{L^2(\mathbb{R}^n)}^2 = \int_{\mathbb{R}^n} u_s \eta_R (-\Delta)^s (u_s \eta_R) \, dx.$$

By the definition of the fractional Laplacian we have

$$\begin{aligned} \int_{\mathbb{R}^n} u_s \eta_R (-\Delta)^s (u_s \eta_R) \, dx &= C(n, s) \int_{\mathbb{R}^n \times \mathbb{R}^n} u_s(x) \eta_R(x) \frac{u_s(x) \eta_R(x) - u_s(y) \eta_R(y)}{|x - y|^{n+2s}} \, dy \, dx \\ &= \int_{\mathbb{R}^n} \eta_R^2 u_s (-\Delta)^s u_s \, dx + C(n, s) \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} u_s(x) u_s(y) \eta_R(x) \, dy \, dx \\ &= \frac{C(n, s)}{2} \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx, \end{aligned}$$

where the last equation is obtained by the symmetrization of the previous integral with respect to the variable $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n$. Before splitting the domain of integration into different subsets, it is easy to see that

$$\begin{aligned} \eta_R(x) - \eta_R(y) &\equiv 0 \quad \text{in } B_R(x_0) \times B_R(x_0) \cup (\mathbb{R}^n \setminus B_{2R}(x_0)) \times (\mathbb{R}^n \setminus B_{2R}(x_0)), \\ |\eta_R(x) - \eta_R(y)| &\equiv 1 \quad \text{in } B_R(x_0) \times (\mathbb{R}^n \setminus B_{2R}(x_0)) \cup (\mathbb{R}^n \setminus B_{2R}(x_0)) \times B_R(x_0), \end{aligned}$$

where all the previous balls are centered at the point x_0 . Hence, given the sets

$$\begin{aligned} \Omega_1 &= B_{3R}(x_0) \times B_{3R}(x_0), \\ \Omega_2 &= B_{2R}(x_0) \times (\mathbb{R}^n \setminus B_{3R}(x_0)) \cup (\mathbb{R}^n \setminus B_{3R}(x_0)) \times B_{2R}(x_0), \end{aligned}$$

we have

$$\begin{aligned} \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx \\ \leq \int_{\Omega_1} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx + \int_{\Omega_2} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx. \end{aligned}$$

In particular

$$\begin{aligned} \int_{\Omega_1} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx &\leq \sup_{B_{3R}(x_0)} u_s^2 \int_{B_{3R}(x_0) \times B_{3R}(x_0)} \frac{\|\nabla \eta_R\|_{L^\infty(\mathbb{R}^n)}^2}{|x - y|^{n+2s-2}} \, dy \, dx \\ &\leq \|\nabla \eta_R\|_{L^\infty}^2 \sup_{B_{3R}(x_0)} u_s^2 \int_{B_{3R}(0)} \, dx \int_{B_{6R}(x)} \frac{1}{|x - y|^{n+2s-2}} \, dy \\ &\leq \frac{\|\nabla \eta\|_{L^\infty}^2}{R^2} \sup_{B_{3R}(x_0)} u_s^2 |B_{3R}| |S^{n-1}| \frac{(6R)^{2-2s}}{2(1-s)} \\ &\leq C \|\nabla \eta\|_{L^\infty}^2 \frac{R^{n-2s}}{2(1-s)} \max\{|x_0|^{2\gamma_s}, (3R)^{2\gamma_s}\} \|u_s\|_{L^\infty(S^{n-1})}, \end{aligned}$$

where in the second inequality we use the changes of variables $x - x_0$ and $y - x_0$ and the fact that $B_{3R}(0) \times B_{3R}(0) \subset B_{3R}(0) \times B_{6R}(x)$ for every $x \in B_{3R}(0)$. Similarly we have

$$\begin{aligned} \int_{\Omega_2} \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{n+2s}} u_s(x) u_s(y) \, dy \, dx &\leq 2 \int_{B_{2R}(x_0)} u_s(x) \left(\int_{\mathbb{R}^n \setminus B_{3R}(x_0)} \frac{u_s(y)}{|x - y|^{n+2s}} \, dy \right) \, dx \\ &\leq 2 \int_{B_{2R}(0)} u_s(x + x_0) \left(\int_{\mathbb{R}^n \setminus B_{3R}(0)} \frac{u_s(y + x_0)}{|y|^{n+2s} (1 - |x|/|y|)^{n+2s}} \, dy \right) \, dx \\ &\leq 2 \cdot 3^{n+2s} \int_{B_{2R}(0)} u_s(x + x_0) \left(\int_{\mathbb{R}^n \setminus B_{3R}(0)} \frac{C(|y| + |x_0|)^{\gamma_s}}{|y|^{n+2s}} \, dy \right) \, dx \\ &\leq C \sup_{B_{2R}(x_0)} u_s |B_{2R}| |S^{n-1}| 2^{\gamma_s} G(x_0, R) \end{aligned}$$

with

$$\begin{aligned} G(x_0, R) &= \begin{cases} |x_0|^{\gamma_s} / (2s - \gamma_s) (3R)^{-2s} & \text{if } |x_0| \geq 3R, \\ (3R)^{\gamma_s - 2s} / (2s - \gamma_s) & \text{if } |x_0| \leq 3R \end{cases} \\ &\leq \frac{(3R)^{-2s}}{2s - \gamma_s} \max\{|x_0|, 3R\}^{\gamma_s}. \end{aligned}$$

Finally, we obtain the desired bound for the seminorm $[u_s \eta_R]_{H^s(\mathbb{R}^n)}^2$ summing the two terms and recalling that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. □

3. Characteristic exponent $\gamma_s(C)$: properties and asymptotic behavior

In this section we start the analysis of the asymptotic behavior of the homogeneity degree $\gamma_s(C)$ as s converges to 1. We get two main results: first we get a monotonicity result for the map $s \mapsto \gamma_s(C)$ for a fixed regular cone C , which ensures the existence of the limit and, using a comparison result, a bound on the possible value of the limit exponent. Secondly we study the asymptotic behavior of the quotient $C(n, s)/(2s - \gamma_s(C))$.

In order to prove the first result and compare different orders of s -harmonic functions for different power of $(-\Delta)^s$, we need to introduce some results which give a natural extension of the classic semigroup property of the fractional Laplacian for functions defined on cones which grow at infinity.

Distributional semigroup property. It is well known that if we deal with smooth functions with compact support, or more generally with functions in the Schwartz space $\mathcal{S}(\mathbb{R}^n)$, a semigroup property holds for the fractional Laplacian; i.e., $(-\Delta)^{s_1} \circ (-\Delta)^{s_2} = (-\Delta)^{s_1+s_2}$, where $s_1, s_2 \in (0, 1)$ with $s_1 + s_2 < 1$. Since we have to deal with functions in \mathcal{L}_s^1 that grow at infinity, we have to construct a distributional counterpart of the semigroup property in order to compute high-order fractional Laplacians for solutions of the problem given in (1-1).

First of all, we remark that a solution u_s to (1-1) for a fixed cone C belongs to \mathcal{L}_s^1 since $0 \leq u_s(x) \leq |x|^{\gamma_s(C)}$ in \mathbb{R}^n with $\gamma_s(C) \in (0, 2s)$. Moreover, by the homogeneity one can rewrite the norm (2-1) as

$$\|u_s\|_{\mathcal{L}_s^1} = \int_{\mathbb{R}^n} \frac{u_s(x)}{(1+|x|)^{n+2s}} \, dx = \int_{S^{n-1}} u_s \, d\sigma \int_0^\infty \frac{\rho^{n-1+\gamma_s(C)}}{(1+\rho)^{n+2s}} \, d\rho = \frac{\Gamma(n+\gamma_s(C))\Gamma(2s-\gamma_s(C))}{\Gamma(n+2s)} \int_{S^{n-1}} u_s \, d\sigma.$$

In the recent paper [Dipierro et al. 2016] the authors introduced a new notion of fractional Laplacian applying to a wider class of functions which grow more than linearly at infinity. This is achieved by defining an equivalence class of functions modulo polynomials of a fixed order. However, it can hardly be applied to the solutions of (1-1) as they annihilate on a set of nonempty interior.

As shown in [Bogdan and Byczkowski 1999, Definition 3.6], if we consider a smooth function with compact support $\varphi \in C_c^\infty(\mathbb{R}^n)$ (or $\varphi \in C_c^2(\mathbb{R}^n)$), we can define the distribution k^{2s} by the formula

$$(-\Delta)^s \varphi(0) = (k^{2s}, \varphi).$$

By this definition, it follows that $(-\Delta)^s \varphi(x) = k^{2s} * \varphi(x)$.

Definition 3.1 [Bogdan and Byczkowski 1999, Definition 3.7]. For $u \in \mathcal{L}_s^1$ we define the *distributional fractional Laplacian* $(-\tilde{\Delta})^s u$ by the formula

$$((-\tilde{\Delta})^s u, \varphi) = (u, (-\Delta)^s \varphi) \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n).$$

In particular, since given an open subset $D \subset \mathbb{R}^n$ and $u \in C^2(D) \cap \mathcal{L}_s^1$, the fractional Laplacian exists as a continuous function of $x \in D$ and $(-\tilde{\Delta})^s u = (-\Delta)^s u$ as a distribution in D [Bogdan and Byczkowski 1999, Lemma 3.8], throughout the paper we will always use $(-\Delta)^s$ both for the classical and the distributional fractional Laplacian. The following is a useful tool for computing the distributional fractional Laplacian.

Lemma 3.2 [Bogdan and Byczkowski 1999, Lemma 3.3]. Assume that

$$\iint_{|y-x|>\varepsilon} \frac{|f(x)g(y)|}{|y-x|^{n+2s}} dx dy < \infty \quad \text{and} \quad \int_{\mathbb{R}^n} |f(x)g(x)| dx < \infty. \tag{3-1}$$

Then $((-\Delta)_\varepsilon^s f, g) = (f, (-\Delta)_\varepsilon^s g)$. Moreover if $f \in \mathcal{L}_s^1$ and $g \in C_c(\mathbb{R}^n)$, the assumptions (3-1) are satisfied for every $\varepsilon > 0$.

Before proving the semigroup property, we prove the following lemma which ensures the existence of the δ -Laplacian of the s -Laplacian for $0 < \delta < 1$.

Lemma 3.3. Let u_s be a solution of (1-1) with C a regular cone. Then we have $(-\Delta)^s u_s \in \mathcal{L}_\delta^1$ for any $\delta > 0$, i.e.,

$$\int_{\mathbb{R}^n} \frac{|(-\Delta)^s u_s(x)|}{(1+|x|)^{n+2\delta}} dx < \infty.$$

Proof. Since the function u_s is s -harmonic in C , namely $(-\Delta)^s u_s(x) = 0$ for all $x \in C$, we can restrict the domain of integration to $\mathbb{R}^n \setminus C$.

By homogeneity and the results in [Bogdan and Byczkowski 1999], we have that the function $(-\Delta)^s u_s$ is $(\gamma_s - 2s)$ -homogeneous and in particular $x \mapsto (-\Delta)^s u_s(x)$ is a continuous negative function, for every $x \in D \Subset \mathbb{R}^n \setminus C$. In order to compute the previous integral, we focus our attention on the restriction of the fractional Laplacian to the sphere S^{n-1} ; in particular, we prove that there exist $\bar{\varepsilon} > 0$ and $C > 0$ such that

$$|(-\Delta)^s u_s(x)| \leq \frac{C}{\text{dist}(x, \partial C)^s} \quad \text{for all } x \in N_{\bar{\varepsilon}}(\partial C) \cap S^{n-1}, \tag{3-2}$$

where $N_\varepsilon(\partial C) = \{x \in \mathbb{R}^n \setminus C : \text{dist}(x, \partial C) \leq \varepsilon\}$ is the tubular neighborhood of ∂C .

Hence, fixing $R > 0$ small enough, consider initially $\varepsilon < R$ and $x \in S^{n-1} \cap N_\varepsilon(\partial C)$; since $u_s(y) \leq |y|^{\gamma_s}$ in \mathbb{R}^n and by (2-2) there exists a constant $C > 0$ such that for every $y \in C$ we have

$$u_s(y) \leq C|y|^{\gamma_s-s} \text{dist}(y, \partial C)^s,$$

it follows, defining $\delta(x) := \text{dist}(x, \partial C) > 0$, that

$$\begin{aligned} |(-\Delta)^s u_s(x)| &= C(n, s) \int_{C \cap B_R(x)} \frac{u_s(y)}{|x-y|^{n+2s}} dy + C(n, s) \int_{C \setminus B_R(x)} \frac{u_s(y)}{|x-y|^{n+2s}} dy \\ &\leq C(n, s) \int_{C \cap B_R(x)} \frac{C|y|^{\gamma_s-s} \text{dist}(y, \partial C)^s}{|x-y|^{n+2s}} dy + C(n, s) \int_{C \setminus B_R(x)} \frac{|y|^{\gamma_s}}{|x-y|^{n+2s}} dy. \end{aligned}$$

Since $C \cap B_R(x) \subset B_R(x) \setminus B_{\delta(x)}(x)$, we have

$$\begin{aligned} |(-\Delta)^s u_s(x)| &\leq C \int_{R \geq |x-y| \geq \delta(x)} \frac{|y|^{\gamma_s-s}}{|x-y|^{n+s}} dy + \int_{|x-y| \geq R} \frac{(|x-y|+1)^{\gamma_s}}{|x-y|^{n+2s}} dy \\ &\leq C \int_{R \geq |x-y| \geq \delta(x)} \frac{1}{|x-y|^{n+s}} dy + \omega_{n-1} \int_R^\infty \frac{(t+1)^{\gamma_s}}{t^{1+2s}} dt \\ &\leq C \int_{\delta(x)}^R \frac{1}{r^{1+s}} dr + M \\ &\leq C \frac{1}{\text{dist}(x, \partial C)^s} + M. \end{aligned}$$

Moreover, again since $s \in (0, 1)$, considering a smaller neighborhood $N_\varepsilon(\partial C)$, we obtain that there exists a constant $\bar{\varepsilon} > 0$ small enough and $C > 0$ such that

$$|(-\Delta)^s u_s(x)| \leq \frac{C}{\text{dist}(x, \partial C)^s} \quad \text{for every } x \in N_{\bar{\varepsilon}}(\partial C) \cap S^{n-1}.$$

Now, fixing $\delta > 0$ and considering $\bar{\varepsilon} > 0$ of (3-2), we have

$$\begin{aligned} \int_{\mathbb{R}^n \setminus C} \frac{|(-\Delta)^s u_s(x)|}{(1+|x|)^{n+2\delta}} dx &= \int_{\mathbb{R}^n \setminus C} \frac{|x|^{\gamma_s-2s} |(-\Delta)^s u_s(x/|x|)|}{(1+|x|)^{n+2\delta}} dx \\ &= \int_0^\infty \int_{S^{n-1} \cap (\mathbb{R}^n \setminus C)} \frac{r^{\gamma_s-2s} |(-\Delta)^s u_s(z)|}{(1+r)^{n+2\delta}} r^{n-1} d\sigma(z) dr \\ &= \int_0^\infty \frac{r^{n-1+\gamma_s-2s}}{(1+r)^{n+2\delta}} dr \int_{S^{n-1} \cap (\mathbb{R}^n \setminus C)} |(-\Delta)^s u_s(z)| d\sigma. \end{aligned}$$

Since $\gamma_s \in (0, 2s)$ and $s \in (0, 1)$, it follows that

$$\begin{aligned} \int_{\mathbb{R}^n \setminus C} \frac{|(-\Delta)^s u_s(x)|}{(1+|x|)^{n+2\delta}} dx &\leq C \int_{S^{n-1} \cap N_{\bar{\varepsilon}}(\partial C)} |(-\Delta)^s u_s(z)| d\sigma + C \int_{((\mathbb{R}^n \setminus C) \setminus N_{\bar{\varepsilon}}(\partial C)) \cap S^{n-1}} |(-\Delta)^s u_s(z)| d\sigma \\ &\leq C \int_{S^{n-1} \cap N_{\bar{\varepsilon}}(\partial C)} \frac{1}{\text{dist}(z, \partial C)^s} d\sigma + M < \infty \end{aligned}$$

where in the second inequality we used that $z \mapsto (-\Delta)^s u_s(z)$ is continuous in every $A \Subset S^{n-1} \cap (\mathbb{R}^n \setminus C)$ and in the last one that $\text{dist}(x, \partial C)^{-s} \in L^1(S^{n-1} \cap N_{\bar{\varepsilon}}(\partial C), d\sigma)$. □

Proposition 3.4 (distributional semigroup property). *Let u_s be a solution of (1-1) with C a regular cone and consider $\delta \in (0, 1 - s)$. Then*

$$(-\Delta)^{s+\delta}u_s = (-\Delta)^\delta [(-\Delta)^s u_s] \quad \text{in } \mathcal{D}'(C)$$

or equivalently

$$((-\Delta)^{s+\delta}u_s, \varphi) = ((-\Delta)^\delta [(-\Delta)^s u_s], \varphi) \quad \text{for all } \varphi \in C_c^\infty(C).$$

Proof. Since $|u_s(x)| \leq |x|^{\gamma_s}$, with $\gamma_s \in (0, 2s)$, it is easy to see that $u_s \in \mathcal{L}_s^1 \cap C^2(C)$. Moreover, as we have already remarked, if $u_s \in \mathcal{L}_s^1$ then $u_s \in \mathcal{L}_{s+\delta}^1$ for every $\delta > 0$. In particular, $(-\Delta)^{s+\delta}u_s$ does exist and it is a continuous function of $x \in C$ for every $\delta \in (0, 1 - s)$. By the definition of the distributional fractional Laplacian, we obtain

$$((-\Delta)^{s+\delta}u_s, \varphi) = (u_s, (-\Delta)^{s+\delta}\varphi),$$

and since for $\varphi \in C_c^\infty(C) \subset \mathcal{S}(\mathbb{R}^n)$ in the Schwarz space, the classic semigroup property holds, we obtain

$$((-\Delta)^{s+\delta}u_s, \varphi) = (u_s, (-\Delta)^s [(-\Delta)^\delta \varphi]).$$

On the other hand, since by Lemma 3.3 we have $(-\Delta)^s u_s \in \mathcal{L}_\delta^1$, it follows that

$$((-\Delta)_\varepsilon^\delta [(-\Delta)^s u_s], \varphi) = ((-\Delta)^s u_s, (-\Delta)_\varepsilon^\delta \varphi) \tag{3-3}$$

for every $\varepsilon > 0$. Since $(-\Delta)^s u_s \in \mathcal{L}_\delta^1$ and $\varphi \in C_c^\infty(\mathbb{R}^n)$, the δ -Laplacian of $(-\Delta)^s u_s$ does exist in a distributional sense and hence the left-hand side in (3-3) does converge to $((-\Delta)^\delta [(-\Delta)^s u_s], \varphi)$ as $\varepsilon \rightarrow 0$. Moreover the right-hand side in (3-3) does converge to $((-\Delta)^s u_s, (-\Delta)^\delta \varphi)$ by the dominated convergence theorem, using Proposition 2.2 and Lemma 3.3, which give

$$\int_{\mathbb{R}^n} (-\Delta)^s u_s(x) (-\Delta)_\varepsilon^\delta \varphi(x) \, dx \leq \int_{\mathbb{R}^n} \frac{|(-\Delta)^s u_s(x)|}{(1 + |x|)^{n+2\delta}} \, dx < \infty.$$

By the previous remarks,

$$((-\Delta)^\delta [(-\Delta)^s u_s], \varphi) = ((-\Delta)^s u_s, (-\Delta)^\delta \varphi).$$

In order to conclude the proof of the distributional semigroup property, we need to show that

$$(u_s, (-\Delta)^s [(-\Delta)^\delta \varphi]) = ((-\Delta)^s u_s, (-\Delta)^\delta \varphi), \tag{3-4}$$

which is not a trivial equality, since $(-\Delta)^\delta \varphi \in C^\infty(\mathbb{R}^n)$ is no more compactly supported.

Let $\eta \in C_c^\infty(B_2(0))$ be a radial cut-off function such that $\eta \equiv 1$ in $B_1(0)$ and $0 \leq \eta \leq 1$ in $B_2(0)$, and define $\eta_R(x) = \eta(x/R)$ for $R > 0$. Obviously, since $u_s \eta_R \in C_c(\mathbb{R}^n)$ and $(-\Delta)^\delta \varphi \in \mathcal{L}_s^1$, by Lemma 3.2 we have

$$(u_s \eta_R, (-\Delta)_\varepsilon^s [(-\Delta)^\delta \varphi]) = ((-\Delta)_\varepsilon^s (u_s \eta_R), (-\Delta)^\delta \varphi) \tag{3-5}$$

for every $\varepsilon, R > 0$. First, for $R > 0$ fixed, we want to pass to the limit for $\varepsilon \rightarrow 0$. For the left-hand side in (3-5), we get the convergence to $(u_s \eta_R, (-\Delta)^s [(-\Delta)^\delta \varphi])$ since we can apply the dominated convergence theorem. In fact

$$\int_{\mathbb{R}^n} u_s \eta_R (-\Delta)_\varepsilon^s [(-\Delta)^\delta \varphi] \leq c \int_K (-\Delta)^{s+\delta} \varphi < \infty,$$

where K denotes the support of $u_s \eta_R$. For the right-hand side in (3-5) we observe that, for any $x \in \mathbb{R}^n$,

$$(-\Delta)_\varepsilon^s(u_s \eta_R)(x) = \eta_R(x)(-\Delta)_\varepsilon^s u_s(x) + u_s(x)(-\Delta)_\varepsilon^s \eta_R(x) - I_\varepsilon(u_s, \eta_R)(x),$$

where

$$I_\varepsilon(u_s, \eta_R)(x) = C(n, s) \int_{\mathbb{R}^n \setminus B_\varepsilon(x)} \frac{(u_s(x) - u_s(y))(\eta_R(x) - \eta_R(y))}{|x - y|^{n+2s}} dy.$$

Obviously the first term $((-\Delta)_\varepsilon^s u_s, \eta_R(-\Delta)^\delta \varphi)$ converges to $((-\Delta)^s u_s, \eta_R(-\Delta)^\delta \varphi)$ by the definition of the distributional s -Laplacian, since $u_s \in \mathcal{L}_s^1$ and $\eta_R(-\Delta)^\delta \varphi \in C_c^\infty(\mathbb{R}^n)$. The second term $(u_s(-\Delta)_\varepsilon^s \eta_R, (-\Delta)^\delta \varphi)$ converges to $(u_s(-\Delta)^s \eta_R, (-\Delta)^\delta \varphi)$ by dominated convergence, since

$$\int_{\mathbb{R}^n} u_s(-\Delta)_\varepsilon^s \eta_R(-\Delta)^\delta \varphi dx \leq c \int_{\mathbb{R}^n} \frac{u_s(x)}{(1 + |x|)^{n+2s}} dx.$$

Finally, the last term $(I_\varepsilon(u_s, \eta_R), (-\Delta)^\delta \varphi)$ converges to $(I(u_s, \eta_R), (-\Delta)^\delta \varphi)$ by dominated convergence, since

$$\int_{\mathbb{R}^n} I_\varepsilon(u_s, \eta_R)(-\Delta)^\delta \varphi dx \leq C \int_{\mathbb{R}^n} |(-\Delta)^\delta \varphi| dx,$$

which is integrable by Proposition 2.2. Finally, passing to the limit for $\varepsilon \rightarrow 0$, from (3-5) we get

$$(u_s \eta_R, (-\Delta)^s [(-\Delta)^\delta \varphi]) = ((-\Delta)^s (u_s \eta_R), (-\Delta)^\delta \varphi) \tag{3-6}$$

for every $R > 0$.

Now we want to prove (3-4), concluding this proof, by passing to the limit in (3-6) for $R \rightarrow \infty$. Since we know, by dominated convergence, that the left-hand side of (3-6) converges to $(u_s, (-\Delta)^s (-\Delta)^\delta \varphi)$ for $R \rightarrow \infty$, we focus our attention on the other side. At this point, we need to prove that for any $\varphi \in C_c^\infty(C)$,

$$\int_{\mathbb{R}^n} (-\Delta)^s (u_s \eta_R)(-\Delta)^\delta \varphi \rightarrow \int_{\mathbb{R}^n} (-\Delta)^s u_s (-\Delta)^\delta \varphi \tag{3-7}$$

as $R \rightarrow \infty$. First of all, we remark that $(-\Delta)^s (u_s \eta_R) \rightarrow (-\Delta)^s u_s$ in $L^1_{\text{loc}}(\mathbb{R}^n)$. In fact, let $K \subset \mathbb{R}^n$ be a compact set. There exists $\bar{r} > 0$ such that $K \subset B_{\bar{r}}$. Then, considering any radius $R > \bar{r}$, we have $\eta_R(x) = 1$ for any $x \in K$. Hence, for any $R > \bar{r}$, using the fact that $u_s(x) = |x|^{\gamma_s} u_s(x/|x|)$, we obtain

$$\begin{aligned} & \int_K |(-\Delta)^s (u_s \eta_R)(x) - (-\Delta)^s u_s(x)| dx \\ &= \int_K dx \left| C(n, s) \text{ p.v.} \int_{\mathbb{R}^n} \frac{u_s(x)\eta_R(x) - u_s(y)\eta_R(y) + u_s(y) - u_s(x)}{|x - y|^{n+2s}} dy \right| \\ &= C(n, s) \int_K dx \left(\text{p.v.} \int_{C \setminus B_R} \frac{u_s(y)[1 - \eta_R(y)]}{|x - y|^{n+2s}} dy \right) \\ &\leq C(n, s) \int_K dx \left(\text{p.v.} \int_{C \setminus B_R} \frac{|y|^{\gamma_s}}{(|y| - \bar{r})^{n+2s}} dy \right) \\ &\leq C(n, s) \int_K dx \left(\text{p.v.} \int_{C \setminus B_R} \frac{|y|^{\gamma_s}}{|y|^{n+2s} (1 - \bar{r}/R)^{n+2s}} dy \right) \\ &= C \left(\frac{R}{R - \bar{r}} \right)^{n+2s} \lim_{\rho \rightarrow \infty} \int_R^\rho \frac{1}{r^{2s - \gamma_s + 1}} dr = C \left(\frac{R}{R - \bar{r}} \right)^{n+2s} \frac{1}{R^{2s - \gamma_s}} \rightarrow 0 \end{aligned}$$

as $R \rightarrow \infty$. Hence we obtain also pointwise convergence almost everywhere. Moreover, we can give the following expression:

$$(-\Delta)^s(u_s \eta_R)(x) = \eta_R(x)(-\Delta)^s u_s(x) + C(n, s) \text{ p.v. } \int_{\mathbb{R}^n} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy. \tag{3-8}$$

We remark that $\eta_R(x)(-\Delta)^s u_s(x) \rightarrow (-\Delta)^s u_s(x)$ and

$$\int_{\mathbb{R}^n} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \rightarrow 0$$

pointwisely. Moreover, we can dominate the first term in the following way:

$$\eta_R(x)(-\Delta)^s u_s(x) \leq (-\Delta)^s u_s(x),$$

and

$$\int_{\mathbb{R}^n} (-\Delta)^s u_s(x)(-\Delta)^\delta \varphi(x) dx < \infty$$

since $(-\Delta)^s u_s \in \mathcal{L}_\delta^1$ and using [Proposition 2.2](#) over $\varphi \in C_c^\infty(C)$. In order to prove (3-7), we want to apply the dominated convergence theorem, and hence we need the following condition for any $R > 0$:

$$I := \left| \int_{\mathbb{R}^n} (-\Delta)^\delta \varphi(x) \left(\text{p.v. } \int_{\mathbb{R}^n} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right) dx \right| \leq c.$$

Therefore, we will obtain a stronger condition, that is, the existence of a value $k > 0$ such that for any $R > 1$

$$I \leq \frac{c}{R^k}.$$

We split the region of integration $\mathbb{R}^n \times \mathbb{R}^n$ into five different parts

$$\begin{aligned} \Omega_1 &:= (\mathbb{R}^n \setminus B_{2R}) \times \mathbb{R}^n, & \Omega_2 &:= B_{2R} \times B_{2R}, & \Omega_3 &:= (B_{2R} \setminus B_R) \times (B_{3R} \setminus B_{2R}), \\ \Omega_4 &:= (B_{2R} \setminus B_R) \times (\mathbb{R}^n \setminus B_{3R}), & \Omega_5 &:= B_R \times (\mathbb{R}^n \setminus B_{2R}). \end{aligned}$$

First of all, we remark that

$$(-\Delta)^s \eta_R(x) = R^{-2s} (-\Delta)^s \eta(x/R)$$

and also that $\|(-\Delta)^s \eta\|_{L^\infty(\mathbb{R}^n)} < \infty$. For the first term, using the fact that $\eta_R(x) - \eta_R(y) = 0$ if $(x, y) \in (\mathbb{R}^n \setminus B_{2R}) \times (\mathbb{R}^n \setminus B_{2R})$,

$$\begin{aligned} I_1 &:= \int_{\mathbb{R}^n \setminus B_{2R}} |(-\Delta)^\delta \varphi(x)| \left| \int_{\mathbb{R}^n} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx \\ &\leq \int_{\mathbb{R}^n \setminus B_{2R}} |(-\Delta)^\delta \varphi(x)| \left| \int_{B_{2R}} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx \\ &\leq \int_{\mathbb{R}^n \setminus B_{2R}} |(-\Delta)^\delta \varphi(x)| (\sup_{B_{2R}} u_s) |(-\Delta)^s \eta_R(x)| dx \\ &\leq \frac{c}{R^{2s-\gamma_s}} \int_{\mathbb{R}^n} \frac{1}{(1 + |x|)^{n+2\delta}} dx \leq \frac{c}{R^{2s-\gamma_s}}. \end{aligned}$$

For the second term, using the fact that $\eta_R(x) - \eta_R(y) \geq 0$ if $(x, y) \in B_{2R} \times (\mathbb{R}^n \setminus B_{2R})$, we obtain as before

$$\begin{aligned} I_2 &:= \int_{B_{2R}} |(-\Delta)^\delta \varphi(x)| \left| \int_{B_{2R}} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx \\ &\leq \int_{B_{2R}} |(-\Delta)^\delta \varphi(x)| (\sup_{B_{2R}} u_s) |(-\Delta)^s \eta_R(x)| dx \\ &\leq \frac{c}{R^{2s-\gamma_s}} \int_{\mathbb{R}^n} \frac{1}{(1 + |x|)^{n+2\delta}} dx \leq \frac{c}{R^{2s-\gamma_s}}. \end{aligned}$$

For the third part

$$I_3 := \int_{B_{2R} \setminus B_R} |(-\Delta)^\delta \varphi(x)| \left| \int_{B_{3R} \setminus B_{2R}} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx,$$

we consider the change of variables

$$\xi = \frac{x}{R} \in B_2 \setminus B_1, \quad \zeta = \frac{y}{R} \in B_3 \setminus B_2.$$

Hence, using the γ_s -homogeneity of u_s and the definition of our cut-off functions, we obtain

$$I_3 \leq \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} |(-\Delta)^\delta \varphi(R\xi)| u_s(\zeta) \frac{\eta(\xi) - \eta(\zeta)}{|\xi - \zeta|^{n+2s}} d\xi d\zeta.$$

We use the fact that $u_s \in C^{0,s}(B_3 \setminus B_1)$, (see (2-2) proved in [Michalik 2006]) and the cut-off function $\eta \in \text{Lip}(B_3 \setminus B_1)$; that is, there exists a constant $c > 0$ such that

$$\begin{aligned} |u_s(\xi) - u_s(\zeta)| &\leq c|\xi - \zeta|^s, \\ |\eta(\xi) - \eta(\zeta)| &\leq c|\xi - \zeta| \end{aligned} \tag{3-9}$$

for every $\xi, \zeta \in B_3 \setminus B_1$. Hence,

$$\begin{aligned} I_3 &\leq \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} |(-\Delta)^\delta \varphi(R\xi)| \frac{|u_s(\zeta) - u_s(\xi)| |\eta(\xi) - \eta(\zeta)|}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &\quad + \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} |(-\Delta)^\delta \varphi(R\xi)| u_s(\xi) \frac{|\eta(\xi) - \eta(\zeta)|}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &= J_1 + J_2. \end{aligned}$$

By (3-9), we obtain

$$\begin{aligned} J_1 &\leq c \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} |(-\Delta)^\delta \varphi(R\xi)| \frac{|\xi - \zeta|^{s+1}}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &\leq c \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} \frac{1}{(1 + R|\xi|)^{n+2\delta}} \frac{1}{|\xi - \zeta|^{n+s-1}} d\xi d\zeta \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} \frac{1}{|\xi - \zeta|^{n+s-1}} d\xi d\zeta \leq \frac{c}{R^{2s+2\delta-\gamma_s}}. \end{aligned}$$

Moreover, using the additional changes of variable

$$(\xi, \zeta) \mapsto (\xi, \xi + h), \quad (\xi, \zeta) \mapsto (\xi, \xi - h),$$

we obtain

$$\begin{aligned} J_2 &\leq \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} |(-\Delta)^\delta \varphi(R\xi)| u_s(\xi) \frac{\eta(\xi) - \eta(\zeta)}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &\leq \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} \frac{1}{(1 + R|\xi|)^{n+2\delta}} u_s(\xi) \frac{\eta(\xi) - \eta(\zeta)}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (B_3 \setminus B_2)} \frac{\eta(\xi) - \eta(\zeta)}{|\xi - \zeta|^{n+2s}} d\xi d\zeta \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \iint_{(B_2 \setminus B_1) \times B_2} \frac{2\eta(\xi) - \eta(\xi + h) - \eta(\xi - h)}{|h|^{n+2s}} d\xi dh \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \left(c + \iint_{(B_2 \setminus B_1) \times B_\varepsilon} \frac{\langle \nabla^2 \eta(\xi) h, h \rangle}{|h|^{n+2s}} d\xi dh \right) \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \left(c + \iint_{(B_2 \setminus B_1) \times B_\varepsilon} \frac{1}{|h|^{n+2s-2}} d\xi dh \right) \leq \frac{c}{R^{2s+2\delta-\gamma_s}}. \end{aligned}$$

For the fourth part

$$I_4 := \int_{B_{2R} \setminus B_R} |(-\Delta)^\delta \varphi(x)| \left| \int_{\mathbb{R}^n \setminus B_{3R}} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx,$$

we consider, as before, the change of variables

$$\xi = \frac{x}{R} \in B_2 \setminus B_1, \quad \zeta = \frac{y}{R} \in \mathbb{R}^n \setminus B_3.$$

Hence,

$$\begin{aligned} I_4 &\leq c \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (\mathbb{R}^n \setminus B_3)} |(-\Delta)^\delta \varphi(R\xi)| \frac{|\zeta|^{\gamma_s}}{|\zeta - \xi|^{n+2s}} d\xi d\zeta \\ &\leq c \frac{R^{2n}}{R^{n+2s-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (\mathbb{R}^n \setminus B_3)} \frac{1}{(1 + R|\xi|)^{n+2\delta}} \frac{|\zeta|^{\gamma_s}}{|\zeta - 2\xi/|\xi||^{n+2s}} d\xi d\zeta \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (\mathbb{R}^n \setminus B_3)} \frac{|\zeta|^{\gamma_s}}{|\zeta|^{n+2s} (1 - 2/|\zeta|)^{n+2s}} d\xi d\zeta \\ &\leq \frac{c}{R^{2s+2\delta-\gamma_s}} \iint_{(B_2 \setminus B_1) \times (\mathbb{R}^n \setminus B_3)} \frac{1}{|\zeta|^{n+2s-\gamma_s}} d\xi d\zeta \leq \frac{c}{R^{2s+2\delta-\gamma_s}}. \end{aligned}$$

Finally we consider the last term

$$I_5 := \int_{B_R} |(-\Delta)^\delta \varphi(x)| \left| \int_{\mathbb{R}^n \setminus B_{2R}} u_s(y) \frac{\eta_R(x) - \eta_R(y)}{|x - y|^{n+2s}} dy \right| dx.$$

Hence we obtain

$$\begin{aligned}
 I_5 &\leq c \int_{B_R} |(-\Delta)^\delta \varphi(x)| \left(\int_{\mathbb{R}^n \setminus B_{2R}} \frac{|y|^{\gamma_s}}{|y-x|^{n+2s}} dy \right) dx \\
 &\leq c \int_{B_R} |(-\Delta)^\delta \varphi(x)| \left(\int_{\mathbb{R}^n \setminus B_{2R}} \frac{|y|^{\gamma_s}}{|y-Ry/|y||^{n+2s}} dy \right) dx \\
 &\leq c \int_{B_R} |(-\Delta)^\delta \varphi(x)| \left(\int_{\mathbb{R}^n \setminus B_{2R}} \frac{|y|^{\gamma_s}}{|y|^{n+2s}(1-R/|y|)^{n+2s}} dy \right) dx \\
 &\leq c \int_{B_R} |(-\Delta)^\delta \varphi(x)| \left(\int_{\mathbb{R}^n \setminus B_{2R}} \frac{1}{|y|^{n+2s-\gamma_s}} dy \right) dx \\
 &\leq c \left(\int_{\mathbb{R}^n} \frac{1}{(1+|x|)^{n+2\delta}} dx \right) \left(\int_{2R}^\infty \frac{1}{r^{1+2s-\gamma_s}} dr \right) \\
 &= c \left(\int_{\mathbb{R}^n} \frac{1}{(1+|x|)^{n+2\delta}} dx \right) \left(\lim_{\rho \rightarrow \infty} \int_{2R}^\rho \frac{1}{r^{1+2s-\gamma_s}} dr \right) \leq \frac{c}{R^{2s-\gamma_s}}.
 \end{aligned}$$

Since $I \leq \sum_{i=1}^5 I_i$, we obtain the desired result. □

At this point, fixing $s \in (0, 1)$, by the distributional semigroup property we can compute easily the high-order fractional Laplacian $(-\Delta)^{s+\delta}$ viewing it as the δ -Laplacian of the s -Laplacian.

Corollary 3.5. *Let C be a regular cone. For every $\delta \in (0, 1 - s)$, the solution u_s of (1-1) is $(s + \delta)$ -superharmonic in C in the sense of distribution; i.e.,*

$$((-\Delta)^{s+\delta} u_s, \varphi) \geq 0$$

for every test function $\varphi \in C_c^\infty(C)$ nonnegative in C .

Moreover, u_s is also superharmonic in C in the sense of distribution; i.e.,

$$(-\Delta u_s, \varphi) \geq 0$$

for every test function $\varphi \in C_c^\infty(C)$ nonnegative in C .

Proof. As said before, the facts that $u_s \in \mathcal{L}_{s+\delta}^1$ and $u_s \in C^2(A)$ for every $A \Subset C$ ensure the existence of the $(-\Delta)^{s+\delta} u_s$ and the continuity of the map $x \mapsto (-\Delta)^{s+\delta} u_s(x)$ for every $x \in A \Subset C$. Hence at this point, the only part we need to prove is the positivity of the $(s + \delta)$ -Laplacian in the sense of the distribution, which is a direct consequence of the previous result. Indeed, since u_s is a solution of the problem (1-1), by Proposition 3.4 we know that for every $\varphi \in C_c^\infty(C)$ we have

$$\begin{aligned}
 ((-\Delta)^{s+\delta} u_s, \varphi) &= ((-\Delta)^\delta [(-\Delta)^s u_s], \varphi) \\
 &= \int_C \varphi(x) \text{p.v.} \int_{\mathbb{R}^n} \frac{(-\Delta)^s u_s(x) - (-\Delta)^s u_s(y)}{|x-y|^{n+2\delta}} dy dx,
 \end{aligned}$$

where $(-\Delta)^\delta [(-\Delta)^s u_s]$ is well-defined since $(-\Delta)^s u_s \equiv 0 \in C^2(A)$ for every $A \Subset C$ and, by Lemma 3.3, $(-\Delta)^s u_s \in \mathcal{L}_\delta^1$ for every $\delta \in (0, 1 - s)$.

Consider now the nonnegative test function $\varphi \geq 0$ in C . Since $(-\Delta)^s u_s(x) = 0$ for every $x \in C$, we have for every $x \in \mathbb{R}^n \setminus \bar{C}$

$$(-\Delta)^s u_s(x) = - \int_C \frac{u_s(y)}{|x - y|^{n+2s}} dy \leq 0.$$

Similarly,

$$((-\Delta)^\delta [(-\Delta)^s u_s], \varphi) = \int_C \varphi(x) \int_{\mathbb{R}^n} \frac{-(-\Delta)^s u_s(y)}{|x - y|^{n+2\delta}} dy dx \geq 0,$$

since the support of φ is compact in the cone C , and so there exists $\varepsilon > 0$ such that $|x - y| > \varepsilon$ in the above integral. We have obtained that for any $\delta \in (0, 1 - s)$ and any nonnegative $\varphi \in C_c^\infty(C)$

$$((-\Delta)^{s+\delta} u_s, \varphi) \geq 0.$$

Then, passing to the limit for $\delta \rightarrow 1 - s$, the function u_s is superharmonic in the distributional sense

$$0 \leq \lim_{\delta \rightarrow 1-s} ((-\Delta)^{s+\delta} u_s, \varphi) = \lim_{\delta \rightarrow 1-s} (u_s, (-\Delta)^{s+\delta} \varphi) = (u_s, -\Delta \varphi) = (-\Delta u_s, \varphi). \quad \square$$

Monotonicity of $s \mapsto \gamma_s(C)$. The following proposition is a consequence of [Corollary 3.5](#) and it follows essentially the proof of Lemma 2 in [\[Bogdan et al. 2015\]](#).

Proposition 3.6. *For any fixed regular cone C with vertex in 0 , the map $s \mapsto \gamma_s(C)$ is monotone nondecreasing in $(0, 1)$.*

Proof. Fixing the cone C , let us denote by γ_s and $\gamma_{s+\delta}$ respectively the homogeneities of u_s and $u_{s+\delta}$. Let us suppose by way of contradiction that $\gamma_s > \gamma_{s+\delta}$ for a $\delta \in (0, 1 - s)$, and let us consider the function

$$h(x) = u_{s+\delta}(x) - u_s(x) \quad \text{in } \mathbb{R}^n,$$

where u_s is the homogeneous solution of (1-1) and $u_{s+\delta}$ is the unique, up to multiplicative constants, nonnegative nontrivial homogeneous and continuous-in- \mathbb{R}^n solution for

$$\begin{cases} (-\Delta)^{s+\delta} u = 0 & \text{in } C, \\ u = 0 & \text{in } \mathbb{R}^n \setminus C \end{cases}$$

of the form

$$u_{s+\delta}(x) = |x|^{\gamma_{s+\delta}} u_{s+\delta} \left(\frac{x}{|x|} \right).$$

The function h is continuous in \mathbb{R}^n and $h(x) = 0$ in $\mathbb{R}^n \setminus C$. We want to prove that $h(x) \leq 0$ in $\mathbb{R}^n \setminus (C \cap B_1)$. Since $h = 0$ outside the cone, we can consider only what happens in $C \setminus B_1$. As we already quoted, we have

$$c_1(s) |x|^{\gamma_s - s} \text{dist}(x, \partial C)^s \leq u_s(x) \leq c_2(s) |x|^{\gamma_s - s} \text{dist}(x, \partial C)^s \tag{3-10}$$

for any $x \in \bar{C} \setminus \{0\}$, and there exist two constants $c_1(s + \delta), c_2(s + \delta) > 0$ such that

$$c_1(s + \delta) |x|^{\gamma_{s+\delta} - (s+\delta)} \text{dist}(x, \partial C)^{s+\delta} \leq u_{s+\delta}(x) \leq c_2(s + \delta) |x|^{\gamma_{s+\delta} - (s+\delta)} \text{dist}(x, \partial C)^{s+\delta}.$$

We can choose u_s and $u_{s+\delta}$ so that $c := c_1(s) = c_2(s + \delta)$ since they are defined up to a multiplicative constant. Then, for any $x \in C \setminus B_1$, since $|x|^{\gamma_{s+\delta}} \leq |x|^{\gamma_s}$, we have

$$h(x) \leq c|x|^{\gamma_s} \operatorname{dist}(x, \partial C)^s \left[\frac{\operatorname{dist}(x, \partial C)^\delta}{|x|^\delta} - 1 \right] \leq 0. \tag{3-11}$$

In fact, if we take x such that $\operatorname{dist}(x, \partial C) \leq 1$, then (3-11) follows by

$$\frac{\operatorname{dist}(x, \partial C)^\delta}{|x|^\delta} - 1 \leq \operatorname{dist}(x, \partial C)^\delta - 1 \leq 0.$$

Instead, if we consider x so that $\operatorname{dist}(x, \partial C) > 1$, then $\operatorname{dist}(x, \partial C)^\delta < |x|^\delta$ and hence (3-11) follows.

Now we want to show that there exists a point $x_0 \in C \cap B_1$ such that $h(x_0) > 0$. Let us take a point $\bar{x} \in S^{n-1} \cap C$ and let $\alpha := u_{s+\delta}(\bar{x}) > 0$ and $\beta := u_s(\bar{x}) > 0$. Hence, there exists a small $r > 0$ so that $\alpha r^{\gamma_{s+\delta}} > \beta r^{\gamma_s}$, and so, taking x_0 with $|x_0| = r$ and so that $x_0/|x_0| = \bar{x}$, we obtain $h(x_0) > 0$.

If we consider the restriction of h to $\overline{C \cap B_1}$, which is continuous on a compact set, by the previous arguments and the Weierstrass theorem, there exists a maximum point $x_1 \in C \cap B_1$ for the function h which is global in \mathbb{R}^n and is strict at least in a set of positive measure. Hence,

$$(-\Delta)^{s+\delta} h(x_1) = C(n, s) \operatorname{p.v.} \int_{\mathbb{R}^n} \frac{h(x_1) - h(y)}{|x_1 - y|^{n+2(s+\delta)}} dy > 0,$$

and since $(-\Delta)^{s+\delta} h$ is a continuous function in the open cone, there exists an open set $U(x_1)$ with $\overline{U(x_1)} \subset C$ such that

$$(-\Delta)^{s+\delta} h(x) > 0 \quad \text{for all } x \in U(x_1).$$

But thanks to Corollary 3.5 we obtain a contradiction since for any nonnegative $\varphi \in C_c^\infty(U(x_1))$

$$((-\Delta)^{s+\delta} h, \varphi) = ((-\Delta)^{s+\delta} u_{s+\delta}, \varphi) - ((-\Delta)^{s+\delta} u_s, \varphi) = -((-\Delta)^{s+\delta} u_s, \varphi) \leq 0. \quad \square$$

With the same arguments as the previous proof we can show also the following useful upper bound.

Proposition 3.7. *For any fixed regular cone C with vertex in 0 and any $s \in (0, 1)$, we have $\gamma_s(C) \leq \gamma(C)$.*

Proof. Seeking a contradiction, we suppose that there exists $s \in (0, 1)$ such that $\gamma_s > \gamma$. Hence we define the function

$$h(x) = u(x) - u_s(x) \quad \text{in } \mathbb{R}^n,$$

where u_s and u are respectively solutions to (1-1) and

$$\begin{cases} -\Delta u = 0 & \text{in } C, \\ u = 0 & \text{in } \mathbb{R}^n \setminus C. \end{cases} \tag{3-12}$$

We recall that these solutions are unique, up to multiplicative constants, nonnegative, nontrivial, homogeneous and continuous in \mathbb{R}^n and of the form

$$u(x) = |x|^\gamma u\left(\frac{x}{|x|}\right), \quad u_s(x) = |x|^{\gamma_s} u_s\left(\frac{x}{|x|}\right)$$

for some $\gamma_s \in (0, 2s)$ and $\gamma \in (0, \infty)$. The function h is continuous in \mathbb{R}^n and $h(x) = 0$ in $\mathbb{R}^n \setminus C$. We want to prove that $h(x) \leq 0$ in $\mathbb{R}^n \setminus (C \cap B_1)$. Since $h = 0$ outside the cone, we can consider only what happens in $C \setminus B_1$. So, there exist two constants $c_1(s), c_2(s) > 0$ such that, for any $x \in \bar{C} \setminus \{0\}$, (3-10) holds. Moreover there exist two constants $c_1, c_2 > 0$ such that

$$c_1|x|^{\gamma-1} \operatorname{dist}(x, \partial C) \leq u(x) \leq c_2|x|^{\gamma-1} \operatorname{dist}(x, \partial C).$$

We can choose u_s and u so that $c := c_1(s) = c_2$ since they are defined up to a multiplicative constant. Then, for any $x \in C \setminus B_1$, since $|x|^\gamma \leq |x|^{\gamma_s}$, we have

$$h(x) \leq c|x|^{\gamma_s} \operatorname{dist}(x, \partial C)^s \left[\frac{\operatorname{dist}(x, \partial C)^{1-s}}{|x|^{1-s}} - 1 \right] \leq 0$$

by the same arguments as the previous proof.

Now we want to show that there exists a point $x_0 \in C \cap B_1$ such that $h(x_0) > 0$. Let us take a point $\bar{x} \in S^{n-1} \cap C$ and let $\alpha := u(\bar{x}) > 0$ and $\beta := u_s(\bar{x}) > 0$. Hence, there exists a small $r > 0$ such that $\alpha r^\gamma > \beta r^{\gamma_s}$, and so, taking x_0 with $|x_0| = r$ and so that $x_0/|x_0| = \bar{x}$, we obtain $h(x_0) > 0$.

If we consider the restriction of h to $\bar{C} \cap \bar{B}_1$, which is continuous on a compact set, by the previous arguments and the Weierstrass theorem, there exists at least a maximum point in $C \cap B_1$ for the function h which is global in \mathbb{R}^n . Moreover, since h cannot be constant on $C \cap B_1$ and it is of class C^2 inside the cone, there exists a global maximum $y \in C \cap B_1$ such that, up to a rotation, $\partial_{x_i}^2 h(y) \leq 0$ for any $i = 1, \dots, n$ and $\partial_{x_j x_j}^2 h(y) < 0$ for at least a coordinate direction. Hence

$$\Delta h(y) = \sum_{i=1}^n \partial_{x_i x_i}^2 h(y) < 0.$$

By the continuity of Δh in the open cone, there exists an open set $U(y)$ with $\overline{U(y)} \subset C$ such that

$$\Delta h(x) < 0 \quad \text{for all } x \in U(y).$$

Since, by Corollary 3.5 for any nonnegative $\varphi \in C_c^\infty(U(y))$

$$(-\Delta u_s, \varphi) \geq 0,$$

we have

$$(\Delta h, \varphi) = (\Delta u, \varphi) - (\Delta u_s, \varphi) = (-\Delta u_s, \varphi) \geq 0,$$

and this is a contradiction. □

Asymptotic behavior of $C(n, s)/(2s - \gamma_s(C))$. Let us define for any regular cone C the limit

$$\mu(C) = \lim_{s \rightarrow 1^-} \frac{C(n, s)}{2s - \gamma_s(C)} \in [0, \infty].$$

Obviously, thanks to the monotonicity of $s \mapsto \gamma_s(C)$ in $(0, 1)$, this limit does exist, but we want to show that $\mu(C)$ cannot be infinite. At this point, this situation can happen since $2s - \gamma_s(C)$ can converge to zero and we do not have enough information about this convergence. The study of this limit depends

on the cone C itself and so we will consider separately the cases of wide cones and narrow cones, which are respectively when $\gamma(C) < 2$ and when $\gamma(C) \geq 2$. In this section, we prove this result just for regular cones, while in Section 4 we will extend the existence of a finite limit $\mu(C)$ to any unbounded cone, without the monotonicity result of Proposition 3.6.

Wide cones: $\gamma(C) < 2$. We remark that, fixing a wide cone $C \subset \mathbb{R}^n$, there exist $\varepsilon > 0$ and $s_0 \in (0, 1)$, both depending on C , such that for any $s \in [s_0, 1)$

$$2s - \gamma_s(C) \geq \varepsilon > 0.$$

In fact we know that $s \mapsto \gamma_s(C)$ is monotone nondecreasing in $(0, 1)$ and $0 < \gamma_s(C) \leq \gamma(C) < 2$. Hence, defining $\bar{\gamma}(C) = \lim_{s \rightarrow 1} \gamma_s(C) \in (0, 2)$ we can choose

$$s_0 := \frac{1}{4}(\bar{\gamma}(C) - 2) + 1 \in \left(\frac{1}{2}, 1\right) \quad \text{and} \quad \varepsilon := \frac{1}{2}(2 - \bar{\gamma}(C)) > 0,$$

obtaining

$$2s - \gamma_s(C) \geq 2s_0 - \bar{\gamma}(C) = \varepsilon > 0.$$

As a consequence we have $\mu(C) = 0$ for any wide cone.

Narrow cones: $\gamma(C) \geq 2$. Before addressing the asymptotic analysis for any regular cone, we focus our attention on the spherical-caps ones with “small” aperture. Hence, let us fix $\theta_0 \in (0, \pi/4)$ and for any $\theta \in (0, \theta_0]$, let

$$\lambda_1(\theta) := \lambda_1(\omega_\theta) = \min_{\substack{u \in H_0^1(S^{n-1} \cap C_\theta) \\ u \neq 0}} \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 \, d\sigma}{\int_{S^{n-1}} u^2 \, d\sigma}.$$

We have that $\lambda_1(\theta) > 2n$, and hence the following problem is well defined:

$$\mu_0(\theta) := \min_{\substack{u \in H_0^1(S^{n-1} \cap C_\theta) \\ u \neq 0}} \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 - 2nu^2 \, d\sigma}{\left(\int_{S^{n-1}} |u| \, d\sigma\right)^2}. \tag{3-13}$$

This number $\mu_0(\theta)$ is strictly positive and achieved by a nonnegative $\varphi \in H_0^1(S^{n-1} \cap C_\theta) \setminus \{0\}$ which is strictly positive on $S^{n-1} \cap C_\theta$ and is obviously a solution to

$$\begin{cases} -\Delta_{S^{n-1}} \varphi = 2n\varphi + \mu_0(\theta) \int_{S^{n-1}} \varphi \, d\sigma & \text{in } S^{n-1} \cap C_\theta, \\ \varphi = 0 & \text{in } S^{n-1} \setminus C_\theta, \end{cases} \tag{3-14}$$

where $-\Delta_{S^{n-1}}$ is the Laplace–Beltrami operator on the unitary sphere S^{n-1} .

Let now v be the 0-homogeneous extension of φ to the whole of \mathbb{R}^n and $r(x) := |x|$. Such a function will be a solution to

$$\begin{cases} -\Delta v = 2nv/r^2 + \mu_0(\theta)/r^2 \int_{S^{n-1}} v \, d\sigma & \text{in } C_\theta, \\ v = 0 & \text{in } \mathbb{R}^n \setminus C_\theta. \end{cases} \tag{3-15}$$

Since the spherical cap $C_\theta \cap S^{n-1}$ is an analytic submanifold of S^{n-1} and the data $(\partial C_\theta \cap S^{n-1}, 0, \partial_\nu \varphi)$ are not characteristic, by the classic theorem of Cauchy and Kovalevskaya we can extend the solution φ

of (3-14) to a function $\tilde{\varphi}$, which is defined in an enlarged cone and satisfies

$$\begin{cases} -\Delta_{S^{n-1}} \tilde{\varphi} = 2n\tilde{\varphi} + \mu_0(\theta) \int_{S^{n-1}} \varphi \, d\sigma & \text{in } S^{n-1} \cap C_{\theta+\varepsilon}, \\ \tilde{\varphi} = \varphi & \text{in } S^{n-1} \cap C_\theta \end{cases}$$

for some $\varepsilon > 0$. As in (3-15), we can define \tilde{v} as the 0-homogeneous extension of $\tilde{\varphi}$. Finally, we introduce the function

$$v_s(x) := r(x)^{\gamma_s^*(\theta)} v(x),$$

where the choice of the homogeneity exponent $\gamma_s^*(\theta) \in (0, 2s)$ is suggested by the following important result.

Theorem 3.8. *Let $\theta \in (0, \theta_0]$. Then there exists $s_0 = s_0(\theta) \in (0, 1)$ such that*

$$(-\Delta)^s v_s(x) \leq 0 \quad \text{in } C_\theta$$

for any $s \in [s_0, 1)$.

Proof. By the $\gamma_s^*(\theta)$ -homogeneity of v_s , it is sufficient to prove that

$$(-\Delta)^s v_s \leq 0 \quad \text{on } C_\theta \cap S^{n-1},$$

since $x \mapsto (-\Delta)^s v_s$ is $(\gamma_s^*(\theta) - 2s)$ -homogeneous. In order to ease the notation, through the following computations we will simply use γ instead of $\gamma_s^*(\theta)$ and $o(1)$ for the terms which converge to zero as s goes to 1. Hence, for $x \in S^{n-1} \cap C_\theta$, we have

$$(-\Delta)^s v_s(x) = |x|^\gamma (-\Delta)^s v(x) + v(x) (-\Delta)^s r^\gamma(x) - C(n, s) \int_{\mathbb{R}^n} \frac{(r^\gamma(x) - r^\gamma(y))(v(x) - v(y))}{|x - y|^{n+2s}} \, dy.$$

First for $R > 0$,

$$\begin{aligned} (-\Delta)^s r^\gamma(x) &= C(n, s) \int_{B_R(x)} \frac{|x|^\gamma - |y|^\gamma}{|x - y|^{n+2s}} \, dy + C(n, s) \int_{\mathbb{R}^n \setminus B_R(x)} \frac{|x|^\gamma - |y|^\gamma}{|x - y|^{n+2s}} \, dy \\ &= \frac{C(n, s)}{2} \int_{B_R(0)} \frac{2|x|^\gamma - |x+z|^\gamma - |x-z|^\gamma}{|z|^{n+2s}} \, dz + C(n, s) \int_{\mathbb{R}^n \setminus B_R(x)} \frac{1 - |y|^\gamma}{|x - y|^{n+2s}} \, dy \\ &= -\frac{C(n, s)}{2} \int_0^R \frac{\rho^2 \rho^{n-1}}{\rho^{n+2s}} \, d\rho \int_{S^{n-1}} \langle \nabla^2 |x|^\gamma z, z \rangle \, d\sigma + o(1) \\ &\quad + C(n, s) |S^{n-1}| \int_R^\infty \frac{1}{\rho^{1+2s}} \, d\rho - C(n, s) \int_{\mathbb{R}^n \setminus B_R(x)} \frac{|y|^\gamma}{|x - y|^{n+2s}} \, dy \\ &= -\frac{C(n, s)}{2} \frac{R^{2-2s}}{2-2s} \int_{S^{n-1}} \langle \nabla^2 |x|^\gamma z, z \rangle \, d\sigma \\ &\quad - C(n, s) \int_R^\infty \frac{\rho^{n-1+\gamma}}{\rho^{n+2s}} \int_{S^{n-1}} \left| \frac{x}{\rho} - \vartheta \right|^\gamma \, d\sigma(\vartheta) \, d\rho + o(1). \end{aligned}$$

Since for every symmetric matrix A we have

$$\int_{S^{n-1}} \langle Az, z \rangle \, d\sigma = \frac{\text{tr } A}{n} \omega_{n-1},$$

where ω_{n-1} is the Lebesgue measure of the $(n - 1)$ -sphere S^{n-1} , we can simplify the first term since $\text{tr } \nabla^2|x|^\gamma = \Delta(|x|^\gamma)$ and checking that

$$\left| \frac{x}{\rho} - \vartheta \right|^\gamma = 1 + \gamma \rho^{-1} \langle \vartheta, x \rangle + o(\rho^{-1})$$

as $\rho \rightarrow \infty$ it follows that

$$\begin{aligned} (-\Delta)^s r^\gamma(x) &= -\frac{C(n, s)}{2} \frac{R^{2-2s}}{2-2s} \frac{\Delta(|x|^\gamma)\omega_{n-1}}{n} - C(n, s)\omega_{n-1} \int_R^\infty \frac{\rho^{n-1+\gamma}}{\rho^{n+2s}} d\rho + o(1) \\ &= -\frac{C(n, s)\omega_{n-1}}{4n(1-s)} \gamma(n-2+\gamma)|x|^{\gamma-2} R^{2-2s} - \frac{C(n, s)}{2s-\gamma} \omega_{n-1} R^{\gamma-2s} + o(1) \\ &= -\frac{C(n, s)\omega_{n-1}}{4n(1-s)} \gamma(n-2+\gamma)R^{2-2s} - \frac{C(n, s)}{2s-\gamma} \omega_{n-1} R^{\gamma-2s} + o(1) \\ &= -\frac{C(n, s)\omega_{n-1}}{4n(1-s)} \gamma(n-2+\gamma) - \frac{C(n, s)}{2s-\gamma} \omega_{n-1} + o(1), \end{aligned}$$

where in the last equality we choose $\gamma = \gamma_s^*(\theta)$ such that $\gamma_s^*(\theta) - 2s \rightarrow 0$ as s goes to 1.

Similarly, if \tilde{v} is the 0-homogeneous extension of v in an enlarged cone, which is such that $v \geq \tilde{v}$ and $v = \tilde{v}$ on $C_\theta \cap S^{n-1}$, it follows that

$$\begin{aligned} (-\Delta)^s v(x) &= \frac{C(n, s)}{2} \int_{|z|<1} \frac{2v(x)-v(x+z)-v(x-z)}{|z|^{n+2s}} dz + C(n, s) \int_{|x-y|>1} \frac{v(x)-v(y)}{|x-y|^{n+2s}} dy \\ &\leq \frac{C(n, s)}{2} \int_{|z|<1} \frac{2\tilde{v}(x)-\tilde{v}(x+z)-\tilde{v}(x-z)}{|z|^{n+2s}} dz + C(n, s) \int_1^\infty \frac{\rho^{n-1}}{\rho^{n+2s}} \int_{S^{n-1}} v(x)-v(y) d\sigma d\rho \\ &= -\frac{C(n, s)}{2} \int_0^1 \frac{\rho^{n-1}\rho^2}{\rho^{n+2s}} \int_{S^{n-1}} \langle \nabla^2 \tilde{v}(x)z, z \rangle d\sigma d\rho + o(1) \\ &= \frac{C(n, s)\omega_{n-1}}{4n(1-s)} (-\Delta)\tilde{v}(x) + o(1), \end{aligned}$$

where we can use that \tilde{v} solves

$$-\Delta \tilde{v} = 2n\tilde{v} + \mu_0 \int_{S^{n-1}} v d\sigma$$

in the enlarged cap $S^{n-1} \cap C_{\theta+\varepsilon}$. Finally,

$$\begin{aligned} C(n, s) \int_{\mathbb{R}^n} \frac{(|x|^\gamma - |y|^\gamma)(v(x) - v(y))}{|x - y|^{n+2s}} dy \\ = C(n, s) \left[\int_{|y|<1} \frac{(1 - |y|^\gamma)(v(x) - v(y))}{|x - y|^{n+2s}} dy + \int_{|y|>1} \frac{(1 - |y|^\gamma)(v(x) - v(y))}{|x - y|^{n+2s}} dy \right], \end{aligned}$$

where the first term is $o(1)$ since

$$\begin{aligned} \int_0^1 (1-\rho^\gamma)\rho^{n-1} \int_{S^{n-1}} \frac{v(x)-v(y)}{|x-\rho y|^{n+2s}} d\sigma d\rho &= \int_0^1 (1-\rho^\gamma)\rho^{n-1} \int_{S^{n-1}} (v(x)-v(y))(1+o(\rho)) d\sigma d\rho \\ &\quad + \int_0^R (1-\rho^\gamma)\rho^{n-1} \int_{S^{n-1}} (v(x)-v(y))(n+2s)\rho \langle x, y \rangle d\sigma d\rho. \end{aligned}$$

Hence, we obtain

$$\begin{aligned}
 C(n, s) & \int_{\mathbb{R}^n} \frac{(|x|^\gamma - |y|^\gamma)(v(x) - v(y))}{|x - y|^{n+2s}} dy \\
 & = C(n, s) \int_{|y|>1} \frac{(1 - |y|^\gamma)(v(x) - v(y))}{|x - y|^{n+2s}} dy + o(1) \\
 & = o(1) - C(n, s) \int_{|y|>1} \frac{|y|^\gamma(v(x) - v(y))}{|x - y|^{n+2s}} dy + o(1) \\
 & = o(1) - C(n, s) \int_1^\infty \rho^\gamma \rho^{n-1} \int_{S^{n-1}} \frac{v(x) - v(y)}{|x - \rho y|^{n+2s}} d\sigma d\rho \\
 & = o(1) - C(n, s) \int_1^\infty \rho^{-1+\gamma-2s} \int_{S^{n-1}} (v(x) - v(y))(1 + o(\rho^{-1})) d\sigma d\rho \\
 & \quad - C(n, s) \int_1^\infty \rho^{-1+\gamma-2s} \int_{S^{n-1}} (v(x) - v(y))(n + 2s)\langle y, x \rangle \rho^{-1} d\sigma d\rho \\
 & = o(1) - \frac{C(n, s)\omega_{n-1}}{2s - \gamma} v(x) + \frac{C(n, s)}{2s - \gamma} \int_{S^{n-1}} v(y) d\sigma.
 \end{aligned}$$

Hence, recalling that $\gamma = \gamma_s^*(\theta)$, for $x \in S^{n-1} \cap C_\theta$ we have

$$\begin{aligned}
 (-\Delta)^s v_s(x) & \leq \left(\mu_0(\theta) \frac{C(n, s)\omega_{n-1}}{4n(1-s)} - \frac{C(n, s)}{2s - \gamma_s^*(\theta)} \right) \int_{S^{n-1}} v_s d\sigma + \frac{C(n, s)\omega_{n-1}}{4n(1-s)} (n + \gamma_s^*(\theta))(2 - \gamma_s^*(\theta)) v_s \\
 & \leq \left(\mu_0(\theta) - \frac{C(n, s)}{2s - \gamma_s^*(\theta)} \right) \int_{S^{n-1}} v_s d\sigma + o(1),
 \end{aligned}$$

where $o(1)$ is uniform with respect to $\gamma_s^*(\theta)$ as $s \rightarrow 1$. In order to obtain a negative right-hand side, it is sufficient to choose $\gamma_s^*(\theta) < 2s$ in such a way to make the denominator $2s - \gamma_s^*(\theta)$ small enough and the quotient $C(n, s)/(2s - \gamma_s^*(\theta))$ still bounded. \square

The previous result suggests the following choice of the homogeneity exponent:

$$\gamma_s^*(\theta) := 2s - s \frac{C(n, s)}{\mu_0(\theta)}.$$

We can finally prove the main result of this section.

Corollary 3.9. *For any regular cone C , we have $\mu(C) < \infty$.*

Proof. We will show that $\mu(\theta) < \infty$ for any $\theta \in (0, \theta_0]$. Then, fixing an unbounded regular cone C , there exists a spherical cone C_θ such that $\theta \in (0, \theta_0]$ and $C_\theta \subset C$. Since by inclusion $\gamma_s(C) < \gamma_s(\theta)$, we obtain

$$\mu(C) \leq \mu(\theta) < \infty.$$

We want to show that fixing $\theta \in (0, \theta_0]$, we have $\gamma_s(\theta) \leq \gamma_s^*(\theta)$ for any $s \in [s_0(\theta), 1)$, where the choice of $s_0(\theta) \in (0, 1)$ is given in [Theorem 3.8](#). The proof of this fact is based on considerations done in [Proposition 3.6](#). By way of contradiction, suppose $\gamma_s(\theta) > \gamma_s^*(\theta)$. Let

$$h(x) = v_s(x) - u_s(x).$$

The function h is continuous in \mathbb{R}^n and $h(x) = 0$ in $\mathbb{R}^n \setminus C_\theta$. We want to prove that $h(x) \leq 0$ in $\mathbb{R}^n \setminus (C_\theta \cap B_1)$. Since $h = 0$ outside the cone, we can consider only what happens in $C_\theta \setminus B_1$. By (3-10), there exist two constants $c_1(s), c_2(s) > 0$ such that, for any $x \in \bar{C}_\theta \setminus \{0\}$,

$$c_1(s)|x|^{\gamma_s - s} \text{dist}(x, \partial C_\theta)^s \leq u_s(x) \leq c_2(s)|x|^{\gamma_s - s} \text{dist}(x, \partial C_\theta)^s,$$

and there exist two constants $c_1, c_2 > 0$ such that

$$c_1|x|^{\gamma_s^* - 1} \text{dist}(x, \partial C_\theta) \leq v_s(x) \leq c_2|x|^{\gamma_s^* - 1} \text{dist}(x, \partial C_\theta).$$

We can choose v_s so that $c := c_1(s) = c_2$ since it is defined up to a multiplicative constant. Then, for any $x \in C_\theta \setminus B_1$, since $|x|^{\gamma_s^*} \leq |x|^{\gamma_s}$, we have

$$h(x) \leq c|x|^{\gamma_s} \text{dist}(x, \partial C_\theta)^s \left[\frac{\text{dist}(x, \partial C_\theta)^{1-s}}{|x|^{1-s}} - 1 \right] \leq 0.$$

Now we want to show that there exists a point $x_0 \in C_\theta \cap B_1$ such that $h(x_0) > 0$. Let us consider for example the point $\bar{x} \in S^{n-1} \cap C_\theta$ determined by the angle $\vartheta = \theta/2$, and let $\alpha := v_s(\bar{x}) > 0$ and $\beta := u_s(\bar{x}) > 0$. Hence, there exists a small $r > 0$ such that $\alpha r^{\gamma_s^*} > \beta r^{\gamma_s}$, and so, taking x_0 with angle $\vartheta = \theta/2$ and $|x_0| = r$, we obtain $h(x_0) > 0$.

If we consider the restriction of h to $\overline{C_\theta \cap B_1}$, which is continuous on a compact set, by the previous arguments and the Weierstrass theorem, there exists a maximum point $x_1 \in C_\theta \cap B_1$ for the function h which is global in \mathbb{R}^n and is strict at least in a set of positive measure. Hence,

$$(-\Delta)^s h(x_1) = C(n, s) \text{ p.v. } \int_{\mathbb{R}^n} \frac{h(x_1) - h(y)}{|x_1 - y|^{n+2s}} dy > 0,$$

and since $(-\Delta)^s h$ is a continuous function in the open cone, there exists an open set $U(x_1)$ with $\overline{U(x_1)} \subset C_\theta$ such that

$$(-\Delta)^s h(x) > 0 \quad \text{for all } x \in U(x_1).$$

But thanks to Theorem 3.8 we obtain a contradiction since for any nonnegative $\varphi \in C_c^\infty(U(x_1))$

$$((-\Delta)^s h, \varphi) = ((-\Delta)^s v_s, \varphi) - ((-\Delta)^s u_s, \varphi) = ((-\Delta)^s v_s, \varphi) \leq 0,$$

where the last inequality holds for any $s \in [s_0(\theta), 1)$. Hence, for any $\theta \in (0, \theta_0]$

$$\mu(\theta) = \lim_{s \rightarrow 1^-} \frac{C(n, s)}{2s - \gamma_s(\theta)} \leq \lim_{s \rightarrow 1^-} \frac{C(n, s)}{2s - \gamma_s^*(\theta)} = \mu_0(\theta) < \infty. \quad \square$$

4. The limit for $s \nearrow 1$

In this section we prove the main result, Theorem 1.3, emphasizing the difference between wide and narrow cones. Then we improve the asymptotic analysis, proving uniqueness of the limit under assumptions on the geometry and the regularity of C .

Let $C \subset \mathbb{R}^n$ be an open cone and consider the minimization problem

$$\lambda_1(C) = \inf \left\{ \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 \, d\sigma}{\int_{S^{n-1}} u^2 \, d\sigma} : u \in H^1(S^{n-1}) \setminus \{0\} \text{ and } u = 0 \text{ in } S^{n-1} \setminus C \right\}, \tag{4-1}$$

which is strictly related to the homogeneity of the solution of (3-12) by

$$\lambda_1(C) = \gamma(C)(\gamma(C) + n - 2).$$

Moreover, if $\gamma(C) > 2$, equivalently if $\lambda_1(C) > 2n$, the problem

$$\mu_0(C) := \inf \left\{ \frac{\int_{S^{n-1}} |\nabla_{S^{n-1}} u|^2 - 2nu^2 \, d\sigma}{\left(\int_{S^{n-1}} |u| \, d\sigma\right)^2} : u \in H^1(S^{n-1}) \setminus \{0\} \text{ and } u = 0 \text{ in } S^{n-1} \setminus C \right\} \tag{4-2}$$

is well-defined and the number $\mu_0(C)$ is strictly positive.

By a standard argument due to the variational characterization of the previous quantities, we already know the existence of a nonnegative eigenfunction $\varphi \in H_0^1(S^{n-1} \cap C) \setminus \{0\}$ associated to the minimization problem (4-1) and a nonnegative function $\psi \in H_0^1(S^{n-1} \cap C) \setminus \{0\}$ that achieves the minimum (4-2), since the numerator in (4-2) is a coercive quadratic form equivalent to the one in (4-1).

Since the cone C may be disconnected, it is well known that φ is not necessarily unique. Instead, the function ψ is unique up to a multiplicative constant, since it solves

$$\begin{cases} -\Delta_{S^{n-1}} \psi = 2n\psi + \mu_0(C) \int_{S^{n-1}} \psi \, d\sigma & \text{in } S^{n-1} \cap C, \\ \psi = 0 & \text{in } S^{n-1} \setminus C. \end{cases} \tag{4-3}$$

In fact, due to the integral term in the equation, the solution ψ must be strictly positive in every connected component of C and localizing the equation in a generic component we can easily get uniqueness by the maximum principle.

The next result highlights the functional space in which the limit of the s -harmonic functions on cones for $s \rightarrow 1$ will be defined.

Proposition 4.1 [Bourgain et al. 2001, Corollary 7]. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. For $1 < p < \infty$, let $f_s \in W^{s,p}(\Omega)$, and assume that*

$$[f_s]_{W^{s,p}(\Omega)} \leq C_0.$$

Then, up to a subsequence, (f_s) converges in $L^p(\Omega)$ as $s \rightarrow 1$ (and, in fact, in $W^{1,p}(\Omega)$ for all $t < 1$) to some $f \in W^{1,p}(\Omega)$.

We use a different notation than that in [Bourgain et al. 2001] since in our paper the normalization constant $C(n, s)$ is incorporated in the seminorm $[\cdot]_{H^s}$ in order to obtain a continuity of the norm $\|\cdot\|_{H^s}$ for $s \in (0, 1]$.

Proof. Let C be an open cone and C_R be a regular cone with section on S^{n-1} of class $C^{1,1}$ such that $C_R \subset C$ and $\partial C_R \cap \partial C = \{0\}$.

By monotonicity of the homogeneity degree $\gamma_s(\cdot)$ with respect to the inclusion, we directly obtain $\gamma_s(C) < \gamma_s(C_R)$ and consequently, up to considering a subsequence, we obtain the existence of the

finite limits

$$\bar{\gamma}(C) = \lim_{s \rightarrow 1} \gamma_s(C), \quad \mu(C) = \lim_{s \rightarrow 1} \frac{C(n, s)}{2s - \gamma_s(C)}. \tag{4-4}$$

Since $\gamma_s(C) < 2s$, we know $\bar{\gamma}(C) \leq 2$ and similarly $\mu(C) \in [0, \infty)$.

Let $K \subset \mathbb{R}^n$ be a compact set and consider $x_0 \in K$ and $R > 0$ such that $K \subset B_R(x_0)$. Given $\eta \in C_c^\infty(B_2)$, a radial cut-off function such that $\eta \equiv 1$ in B_1 and $0 \leq \eta \leq 1$ in B_2 , consider the rescaled function $\eta_K(x) = \eta((x - x_0)/R)$ which satisfies $\eta_K \equiv 1$ on K .

By Proposition 2.7, we have

$$[u_s \eta_K]_{H^s(B_{2R}(x_0))}^2 \leq [u_s \eta_K]_{H^s(\mathbb{R}^n)}^2 \leq M(n, K) \left[\frac{C(n, s)}{2(1-s)} + \frac{C(n, s)}{2s - \gamma_s} \right],$$

and similarly

$$\begin{aligned} \|u_s \eta_K\|_{H^s(B_{2R}(x_0))}^2 &\leq \|u_s \eta_K\|_{L^2(\mathbb{R}^n)}^2 + [u_s \eta_K]_{H^s(\mathbb{R}^n)}^2 \\ &\leq M(n, K) \left[\frac{C(n, s)}{2(1-s)} + \frac{C(n, s)}{2s - \gamma_s} + 1 \right] \leq M(n, K) \left[\frac{2n}{\omega_{n-1}} + c\mu(C) + 1 \right]. \end{aligned}$$

By applying Proposition 4.1 with $\Omega = B_{2R}(x_0)$, we obtain that, up to a subsequence, $u_s \eta_K \rightarrow \bar{u} \eta_K$ in $L^2(B_{2R}(x_0))$ and

$$\|\bar{u} \eta_K\|_{H^1(B_{2R}(x_0))}^2 \leq M(n, K)$$

up to relabeling the constant $M(n, K)$.

By construction, since $\eta_K \equiv 1$ on K and $\eta_K \in [0, 1]$, we obtain that $u_s \rightarrow \bar{u}$ in $L^2(K)$ and similarly

$$\|\bar{u}\|_{H^1(K)} \leq \|\bar{u} \eta_K\|_{H^1(K)} \leq \|\bar{u} \eta_K\|_{H^1(B_{2R}(x_0))} < \infty,$$

which gives us the local integrability in $H^1(\mathbb{R}^n)$.

By Proposition 2.4 and Corollary 3.9 we obtain, up to passing to a subsequence, a bound in $C_{\text{loc}}^{0,\alpha}(C)$ for (u_s) that is uniform in s . Then, since we obtain uniform convergence on compact subsets of C , the limit must be necessary nontrivial with $\|\bar{u}\|_{L^\infty(S^{n-1})} = 1$, nonnegative and $\bar{\gamma}(C)$ -homogeneous.

Let $\varphi \in C_c^\infty(C)$ be a positive smooth function compactly supported such that $\text{supp } \varphi \subset B_\rho$ for some $\rho > 0$. By the definition of the distributional fractional Laplacian

$$0 = \int_{\mathbb{R}^n} \varphi(-\Delta)^s u_s \, dx = \int_{\mathbb{R}^n} u_s(-\Delta)^s \varphi \, dx = \int_{\mathbb{R}^n \setminus B_\rho} u_s(-\Delta)^s \varphi \, dx + \int_{B_\rho} u_s(-\Delta)^s \varphi \, dx.$$

Since

$$\frac{1}{|x - y|^{n+2s}} = \frac{1}{|x|^{n+2s}} \left(1 - (n + 2s) \frac{y}{|x|} \int_0^1 \frac{x/|x| - ty/|x|}{|x/|x| - ty/|x||^{n+2s+2}} \, dt \right),$$

by the definition of the fractional Laplacian for regular functions, it follows that

$$\begin{aligned} &\int_{\mathbb{R}^n \setminus B_\rho} u_s(-\Delta)^s \varphi \, dx \\ &= C(n, s) \int_{\mathbb{R}^n \setminus B_\rho} u_s(x) \int_{\text{supp } \varphi} \frac{-\varphi(y)}{|y - x|^{n+2s}} \, dy \, dx \\ &= C(n, s) \int_{\mathbb{R}^n \setminus B_\rho} \frac{u_s(x)}{|x|^{n+2s}} \int_{\text{supp } \varphi} -\varphi(y) \, dy \, dx + C(n, s)(n + 2s) \int_{\mathbb{R}^n \setminus B_\rho} \frac{u_s(x)}{|x|^{n+2s+1}} \psi(x) \, dx \end{aligned}$$

for some $\psi \in L^\infty$. Moreover, since u_s is $\gamma_s(C)$ -homogeneous with $\gamma_s(C) < 2s$, we have

$$C(n, s) \int_{\mathbb{R}^n \setminus B_\rho} \frac{u_s(x)}{|x|^{n+2s}} dx = \frac{C(n, s)}{2s - \gamma_s(C)} \rho^{\gamma_s(C)-2s} \int_{S^{n-1}} u_s(\theta) d\sigma$$

and similarly

$$C(n, s) \left| \int_{\mathbb{R}^n \setminus B_\rho} \frac{u_s(x)}{|x|^{n+2s+1}} \psi(x) dx \right| \leq \frac{C(n, s) \|\psi\|_{L^\infty}}{2s - \gamma_s(C) + 1} \rho^{\gamma_s(C)-2s-1} \int_{S^{n-1}} u_s(\theta) d\sigma = o(1).$$

Hence, for each $s \in (0, 1)$

$$\begin{aligned} \int_{B_\rho} u_s(-\Delta)^s \varphi dx &= \int_{\mathbb{R}^n \setminus B_\rho} u_s(-\Delta)^s \varphi dx \\ &= C(n, s) \int_{\mathbb{R}^n \setminus B_\rho} u_s(x) \int_{\text{supp } \varphi} \frac{\varphi(y)}{|x-y|^{n+2s}} dy dx \\ &= \frac{C(n, s)}{2s - \gamma_s(C)} \int_{\text{supp } \varphi} \varphi(x) dx \int_{S^{n-1}} u_s d\sigma + o(1), \end{aligned}$$

and passing through the limit, up to a subsequence, we obtain

$$\int_{B_\rho} \bar{u}(-\Delta)\varphi dx = \mu(C) \int_{S^{n-1}} \bar{u} d\sigma \int_{\text{supp } \varphi} \varphi(x) dx = \int_{B_\rho} \left(\mu(C) \int_{S^{n-1}} \bar{u} d\sigma \right) \varphi(x) dx,$$

which implies, integrating by parts, that

$$-\Delta \bar{u} = \mu(C) \int_{S^{n-1}} \bar{u} d\sigma \quad \text{in } \mathcal{D}'(C).$$

Since the function \bar{u} is $\bar{\gamma}(C)$ -homogeneous, we get

$$-\Delta_{S^{n-1}} \bar{u} = \bar{\lambda} \bar{u} + \mu(C) \int_{S^{n-1}} \bar{u} d\sigma \quad \text{on } S^{n-1} \cap C, \tag{4-5}$$

where $\bar{\lambda} = \bar{\gamma}(C)(\bar{\gamma}(C) + n - 2)$ is the eigenvalue associated to the critical exponent $\bar{\gamma}(C) \leq 2$.

Consider now a nonnegative $\varphi \in H_0^1(S^{n-1} \cap C) \setminus \{0\}$, strictly positive on $S^{n-1} \cap C$ which achieves (4-1). Then

$$-\Delta_{S^{n-1}} \varphi = \lambda_1(C) \varphi \quad \text{in } H^{-1}(S^{n-1} \cap C). \tag{4-6}$$

By testing this equation with \bar{u} and integrating by parts, we obtain

$$(\lambda_1(C) - \bar{\lambda}) \int_{S^{n-1}} \bar{u} \varphi d\sigma = \mu(C) \int_{S^{n-1}} \bar{u} d\sigma \int_{S^{n-1}} \varphi d\sigma \geq 0, \tag{4-7}$$

which implies that in general $\gamma(C) \geq \bar{\gamma}(C)$ and $\gamma(C) = \bar{\gamma}(C)$ if and only if $\mu(C) = 0$.

Wide cones: $\gamma(C) < 2$. By the previous remark we have $\bar{\gamma}(C) < 2$ and by the definition of $\mu(C)$, it follows that $\mu(C) = 0$. Since φ is the trace on S^{n-1} of a homogeneous harmonic function on C , we obtain that $\bar{\gamma}(C) = \gamma(C)$ and \bar{u} is a homogeneous nonnegative harmonic function on C such that $\|\bar{u}\|_{L^\infty(S^{n-1})} = 1$.

Narrow cones: $\gamma(C) \geq 2$. If $\bar{\gamma}(C) < 2$ we have $\mu(C) = 0$ and consequently $\lambda_1(C) = \bar{\lambda}$, which is a contradiction since $\gamma(C) \geq 2 > \bar{\gamma}(C)$. Hence, if C is a narrow cone we get $\bar{\gamma}(C) = 2$. Since $\gamma(C) = 2$ is trivial and it follows directly from the previous computations, consider now $\mu_0(C)$ as the minimum defined in (4-2), which is well-defined and strictly positive since we are focusing on the remaining case $\gamma(C) > 2$. We already remarked that it is achieved by a nonnegative $\psi \in H_0^1(S^{n-1} \cap C) \setminus \{0\}$ which is strictly positive on $S^{n-1} \cap C$ and a solution of

$$-\Delta_{S^{n-1}} \psi = 2n\psi + \mu_0(C) \int_{S^{n-1}} \psi \, d\sigma \quad \text{in } H^{-1}(S^{n-1} \cap C).$$

As we already did in the previous cases, by testing this equation with \bar{u} we obtain $\mu(C) = \mu_0(C)$.

By uniqueness of the limits $\bar{\gamma}(C)$ and $\mu(C)$, the result in (4-4) holds for $s \rightarrow 1$ and not just up to a subsequence. □

Remark 4.2. The possible obstruction to the existence of the limit of u_s as s converges to 1 lies in the possible lack of uniqueness of nonnegative solutions to (1-9) such that $\|\bar{u}\|_{L^\infty(S^{n-1})} = 1$. This is the reason why we need to extract subsequences in the asymptotic analysis of Theorem 1.3. More precisely, uniqueness of (4-1) implies uniqueness of the limit \bar{u} in the case $\gamma(C) \leq 2$ and uniqueness of (4-2) in the case $\gamma(C) > 2$. When C is connected (4-1) is attained by a unique normalized nonnegative solution via a standard argument based upon the maximum principle. On the other hand, as we already remarked, when $\gamma(C) > 2$, problem (4-2) always admits a unique solution. Ultimately, the main obstacle in this analysis is the disconnection of the cone C when $\gamma(C) \leq 2$: in this case we cannot always ensure the uniqueness of the solution of the limit problem and even the positivity of the limit function \bar{u} on every connected component of C .

The following example shows uniqueness of the limit function \bar{u} due to the nonlocal nature of the fractional Laplacian under a symmetry assumption on the cone C .

Proposition 4.3. *Let $C = C_1 \cup \dots \cup C_m$ be a union of disconnected cones such that C_1 is connected and there are orthogonal maps $\Phi_2, \dots, \Phi_m \in O(n)$ (e.g., reflections about hyperplanes) such that $C_i = \Phi_i(C_1)$ and $\Phi_i(C) = (C)$ for $i = 2, \dots, m$. Let (u_s) be the family of nonnegative solutions to (1-1) such that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. Then there exists the limit of u_s as $s \nearrow 1$ in $L_{loc}^2(\mathbb{R}^n)$ and uniformly on compact subsets of C .*

Proof. We remark that, for any element of the orthogonal group $\Phi: \mathbb{R}^n \rightarrow \mathbb{R}^n$,

$$(-\Delta)^s(u \circ \Phi)(x) = C(n, s) \text{ p.v. } \int_{\mathbb{R}^n} \frac{u(\Phi(x)) - u(y)}{|\Phi(x) - y|^{n+2s}} \, dy = (-\Delta)^s u(\Phi(x)).$$

By the uniqueness result [Bañuelos and Bogdan 2004, Theorem 3.2] of s -harmonic functions on cones, we infer that $u_s \equiv u_s \circ \Phi_i$ for every $i = 2, \dots, m$. Therefore, we have convergence to \bar{u} , which satisfies $\|\bar{u}\|_{L^\infty(S^{n-1})} = 1$ and is a solution of

$$\begin{cases} -\Delta \bar{u} = \mu(C) \int_{S^{n-1}} \bar{u} \, d\sigma & \text{in } C, \\ \bar{u} \geq 0 & \text{in } C, \\ \bar{u} = 0 & \text{in } \mathbb{R}^n \setminus C \end{cases} \tag{4-8}$$

such that $\bar{u} \equiv \bar{u} \circ \Phi_i$ for every $i = 2, \dots, m$. Finally, connectedness of C_1 yields uniqueness of such a solution also for narrow cones. □




Figure 3. Values of the limit $\bar{\Gamma}(\theta) = \lim_{s \rightarrow 1} \Gamma^s(\theta)$ and $\Gamma(\theta)$ for $n = 2$.

Proof of Corollary 1.6. This corollary is an easy application of our main result, [Theorem 1.3](#), since it is a consequence of Dini’s theorem for a monotone sequence of continuous functions which converges pointwisely to a continuous function on a compact set. In fact, fixing $s \in (0, 1)$, the function $\theta \mapsto \gamma_s(\theta)$ is continuous in $[0, \pi)$ with $\gamma_s(0) = 2s$ and $\gamma_s(\pi) = 0$. Moreover this function is also monotone decreasing in $[0, \pi]$ and since there exists the limit

$$\lim_{\theta \rightarrow \pi^-} \gamma_s(\theta) = \begin{cases} \frac{1}{2}(2s - 1) & \text{if } n = 2 \text{ and } s > \frac{1}{2}, \\ \gamma_s(\pi) = 0 & \text{otherwise,} \end{cases}$$

we can extend $\theta \mapsto \gamma_s(\theta)$ to a continuous function in $[0, \pi]$; see [\[Michalik 2006\]](#). Nevertheless, the limit $\bar{\gamma}(\theta) = \lim_{s \rightarrow 1} \gamma_s(\theta) = \min\{\gamma(\theta), 2\}$ is continuous on $[0, \pi]$ with

$$\bar{\gamma}(\pi) = \begin{cases} \frac{1}{2} & \text{if } n = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Eventually, for any fixed $\theta \in [0, \pi]$, the function $s \mapsto \gamma_s(\theta)$ is monotone nondecreasing in $(0, 1)$. By Dini’s theorem the convergence is uniform on $[0, \pi]$. This fact obviously implies the uniform convergence

$$\Gamma^s(\theta) = \frac{\gamma_s(\theta) + \gamma_s(\pi - \theta)}{2} \rightarrow \bar{\Gamma}(\theta) = \frac{\bar{\gamma}(\theta) + \bar{\gamma}(\pi - \theta)}{2}$$

in $[0, \pi]$, and hence (see [Figure 3](#))

$$v_s^{\text{ACF}} = \min_{\theta \in [0, \pi]} \Gamma^s(\theta) \rightarrow \min_{\theta \in [0, \pi]} \bar{\Gamma}(\theta) = v^{\text{ACF}}. \quad \square$$

5. Uniform-in- s estimates in $C^{0,\alpha}$ on annuli

We have already remarked in [Section 2](#) that, if you take a cone $C = C_\omega$ with $\omega \subset S^{n-1}$ a finite union of connected $C^{1,1}$ domains ω_i such that $\bar{\omega}_i \cup \bar{\omega}_j = \emptyset$ for $i \neq j$, by [\[Michalik 2006, Lemma 3.3\]](#) we have (2-2).

Hence solutions u_s to (1-1) are $C^{0,s}(S^{n-1})$ and for any fixed $\alpha \in (0, 1)$, any solution u_s with $s \in (\alpha, 1)$ is $C^{0,\alpha}(S^{n-1})$; that is, there exists $L_s > 0$ such that

$$\sup_{x,y \in S^{n-1}} \frac{|u_s(x) - u_s(y)|}{|x - y|^\alpha} = L_s.$$

Let us consider an annulus $A = A_{r_1,r_2} = B_{r_2} \setminus \bar{B}_{r_1}$ with $0 < r_1 < r_2 < \infty$. We have the following result.

Lemma 5.1. *Let $\alpha \in (0, 1)$, $s_0 \in (\max\{\frac{1}{2}, \alpha\}, 1)$ and A be an annulus centered at zero. Then there exists a constant $c > 0$ such that any solution u_s to (1-1) with $s \in [s_0, 1)$ satisfies*

$$\sup_{x,y \in A} \frac{|u_s(x) - u_s(y)|}{|x - y|^\alpha} \leq cL_s.$$

Proof. First of all we remark that

$$\sup_{x,y \in S_r^{n-1}} \frac{|u_s(x) - u_s(y)|}{|x - y|^\alpha} \leq cL_s \tag{5-1}$$

for any $r \in (r_1, r_2)$. In fact, by the γ_s -homogeneity of our solutions, we have

$$\sup_{x,y \in S_r^{n-1}} \frac{|u_s(x) - u_s(y)|}{|x - y|^\alpha} = L_s r^{\gamma_s - \alpha},$$

and since $(2s_0 - 1)/2 \leq \gamma_s(C) < 2$ for any $s \in [s_0, 1)$ by the inclusion $C \subset \mathbb{R}^n \setminus \{\text{half-line from } 0\}$, we obtain (5-1).

Now we can show what happens considering $x, y \in A$ which are not on the same sphere. We can suppose without loss of generality that $x \in S_R^{n-1}$, $y \in S_r^{n-1}$ with $r_1 < r < R < r_2$. Hence let us take the point z obtained by the intersection between S_r^{n-1} and the half-line connecting 0 and x (z may be y itself). Hence

$$\begin{aligned} |u_s(x) - u_s(y)| &\leq |u_s(x) - u_s(z)| + |u_s(z) - u_s(y)| \\ &\leq u_s\left(\frac{x}{|x|}\right) \left||x|^{\gamma_s} - |z|^{\gamma_s}\right| + cL_s |z - y|^\alpha \\ &\leq cL_s |x - y|^\alpha. \end{aligned}$$

In fact we remark that $\|u_s\|_{L^\infty(S^{n-1})} = 1$. Moreover, since $\beta = \widehat{xzy} \in (\pi/2, \pi]$, obviously $|z - y|^\alpha \leq |x - y|^\alpha$. Thus by the α -Hölder continuity of $t \mapsto t^{\gamma_s}$ in (r_1, r_2) and the bounds $(2s_0 - 1)/2 \leq \gamma_s(C) < 2$, one can find a universal constant $c > 0$ such that

$$\left||x|^{\gamma_s} - |z|^{\gamma_s}\right| \leq c \left||x| - |z|\right|^\alpha \leq c|x - z|^\alpha \leq c|x - y|^\alpha,$$

where the last inequality holds since z is the point on S_r^{n-1} which minimizes the distance $\text{dist}(x, S_r^{n-1})$. \square

Proof of Theorem 1.5. Seeking a contradiction,

$$\max_{x,y \in S^{n-1}} \frac{|u_{s_k}(x) - u_{s_k}(y)|}{|x - y|^\alpha} = L_{s_k} = L_k \rightarrow \infty \quad \text{as } s_k \rightarrow 1. \tag{5-2}$$

We can consider the sequence of points $x_k, y_k \in S^{n-1}$ which realizes L_k at any step. It is easy to see that this pair belongs to $\bar{C} \cap S^{n-1}$. Moreover we can always think of x_k as the one closer to the boundary $\partial C \cap S^{n-1}$. Therefore, to have (5-2), we have $r_k = |x_k - y_k| \rightarrow 0$. Hence, without loss of generality, we can assume that x_k, y_k belong definitively to the same connected component of C and

$$\frac{|u_{s_k}(y_k) - u_{s_k}(x_k)|}{r_k^\alpha} = L_k, \quad \frac{y_k - x_k}{r_k} \rightarrow e_1.$$

Let us define

$$u^k(x) = \frac{u_{s_k}(x_k + r_k x) - u_{s_k}(x_k)}{r_k^\alpha L_k}, \quad x \in \Omega_k = \frac{C - x_k}{r_k}.$$

We remark that $u^k(0) = 0$ and $u^k((y_k - x_k)/r_k) = 1$.

Moreover, we can have two different situations.

Case 1: If

$$\frac{r_k}{\text{dist}(x_k, \partial C)} \rightarrow 0,$$

then the limit of Ω_k is \mathbb{R}^n .

Case 2: If

$$\frac{r_k}{\text{dist}(x_k, \partial C)} \rightarrow l \in (0, \infty],$$

then the limit of Ω_k is a half-space $\mathbb{R}^n \cap \{x_1 > 0\}$.

In any case let us define Ω_∞ to be this limit set. Let us consider the annulus $A^* := B_{3/2} \setminus \bar{B}_{1/2}$. By Lemma 5.1 and the definition of u^k , we obtain, for any k ,

$$\sup_{x,y \in A_k^*} \frac{|u^k(x) - u^k(y)|}{|x - y|^\alpha} \leq c, \tag{5-3}$$

where $A_k^* := (A^* - x_k)/r_k \rightarrow \mathbb{R}^n$ and the constant $c > 0$ depends only on α and A^* . Let us consider a compact subset K of Ω_∞ . Since for k large enough $K \subset A_k^*$, functions u^k are $C^{0,\alpha}(K)$ uniformly in k . This is due also to the fact that they are uniformly in $L^\infty(K)$, since $|u^k(x) - u^k(0)| \leq c|x|^\alpha$ on K . Hence $u^k \rightarrow \bar{u}$ uniformly on compact subsets of Ω_∞ . Moreover \bar{u} is globally α -Hölder continuous and it is not constant, since $\bar{u}(e_1) - \bar{u}(0) = 1$. To conclude, we will show that \bar{u} is harmonic in the limit domain Ω_∞ ; that is, for any $\varphi \in C_c^\infty(\Omega_\infty)$

$$\int_{\Omega_\infty} \varphi(-\Delta)\bar{u} \, dx = 0,$$

and this fact will be a contradiction to the global Hölder continuity. In fact we can apply Corollary 2.3 in [Noris et al. 2010], if $\Omega_\infty = \mathbb{R}^n$ directly on the function \bar{u} and if $\Omega_\infty = \mathbb{R}^n \cap \{x_1 > 0\}$; since $\bar{u} = 0$ in $\partial\Omega_\infty$, we can use the same result over its odd reflection. Hence we want to prove

$$\int_{\Omega_\infty} \varphi(-\Delta)\bar{u} \, dx = \int_{\Omega_\infty} \bar{u}(-\Delta)\varphi \, dx = \lim_{k \rightarrow \infty} \int_{B_R} u^k(-\Delta)^{s_k} \varphi \, dx = 0,$$

where B_R contains the support of φ and the second equality holds by the uniform convergences $u^k \rightarrow \bar{u}$ and $(-\Delta)^{s_k} \varphi \rightarrow (-\Delta)\varphi$ on compact subsets of Ω_∞ , since φ is a smooth function compactly supported.

Moreover, since u^k is s_k -harmonic on Ω_k , and for k large enough the support of φ is contained in this domain, we have

$$\int_{\mathbb{R}^n} u^k (-\Delta)^{s_k} \varphi \, dx = \int_{\mathbb{R}^n} \varphi (-\Delta)^{s_k} u^k \, dx = 0.$$

In order to conclude we want

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}^n \setminus B_R} u^k (-\Delta)^{s_k} \varphi \, dx = 0.$$

Hence, defining $\eta = x_k + r_k x$ and using Remark 2.3, we obtain

$$\left| \int_{\mathbb{R}^n \setminus B_R} u^k (-\Delta)^{s_k} \varphi \, dx \right| \leq \frac{C(n, s_k)}{L_k} r_k^{2s_k - \alpha} \int_{|\eta - x_k| > Rr_k} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta.$$

For k large enough, we notice that we can choose $\varepsilon > 0$ such that the set $\{\eta \in \mathbb{R}^n : Rr_k < |\eta - x_k| < \varepsilon\}$ is contained in A^* . So, we can split the integral obtaining

$$\int_{|\eta - x_k| > Rr_k} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta \leq \int_{Rr_k < |\eta - x_k| < \varepsilon} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta + \int_{|\eta - x_k| > \varepsilon} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta,$$

where we have

$$\begin{aligned} \frac{C(n, s_k) r_k^{2s_k - \alpha}}{L_k} \int_{Rr_k < |\eta - x_k| < \varepsilon} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta &\leq C(n, s_k) r_k^{2s_k - \alpha} c \omega_{n-1} \int_{Rr_k}^{\varepsilon} t^{-1+\alpha-2s_k} \, dt \\ &= \frac{C(n, s_k) c \omega_{n-1}}{2s_k - \alpha} \left(R^{\alpha-2s_k} - \frac{r_k^{2s_k - \alpha}}{\varepsilon^{2s_k - \alpha}} \right) \end{aligned}$$

and similarly

$$\begin{aligned} \frac{C(n, s_k) r_k^{2s_k - \alpha}}{L_k} \int_{|\eta - x_k| > \varepsilon} \frac{|u_{s_k}(\eta) - u_{s_k}(x_k)|}{|\eta - x_k|^{n+2s_k}} \, d\eta &\leq \frac{C(n, s_k) r_k^{2s_k - \alpha} c \omega_{n-1}}{L_k} \int_{\varepsilon}^{\infty} \frac{(1+t)^{\gamma_{s_k}}}{t^{1+2s_k}} \, dt \\ &= \frac{C(n, s_k) r_k^{2s_k - \alpha} c \omega_{n-1}}{L_k} \left(1 + \frac{\varepsilon^{\gamma_{s_k} - 2s_k}}{2s_k - \gamma_{s_k}} \right). \end{aligned}$$

Finally, recalling that $r_k \rightarrow 0$, $C(n, s_k) \rightarrow 0$, $L_k \rightarrow \infty$ and $2s_k - \alpha > 0$ taking $s_0 > \frac{1}{2}$, we obtain

$$\left| \int_{\mathbb{R}^n \setminus B_R} u^k (-\Delta)^{s_k} \varphi \, dx \right| \leq \left(C(n, s_k) + \frac{C(n, s_k) r_n^{2s_k - \alpha}}{2s_k - \gamma_{s_k} L_k} \right) M,$$

which converges to zero as we claimed, since

$$\frac{C(n, s_k)}{2s_k - \gamma_{s_k}(C)} \rightarrow \mu(C) \in [0, \infty)$$

in any regular cone $C \subset \mathbb{R}^n$. □

References

- [Allen 2012] M. Allen, “Separation of a lower dimensional free boundary in a two-phase problem”, *Math. Res. Lett.* **19**:5 (2012), 1055–1074. [MR](#) [Zbl](#)
- [Allen and Lara 2015] M. Allen and H. C. Lara, “Free boundaries on two-dimensional cones”, *J. Geom. Anal.* **25**:3 (2015), 1547–1575. [MR](#) [Zbl](#)

- [Alt et al. 1984] H. W. Alt, L. A. Caffarelli, and A. Friedman, “Variational problems with two phases and their free boundaries”, *Trans. Amer. Math. Soc.* **282**:2 (1984), 431–461. [MR](#) [Zbl](#)
- [Bañuelos and Bogdan 2004] R. Bañuelos and K. Bogdan, “Symmetric stable processes in cones”, *Potential Anal.* **21**:3 (2004), 263–288. [MR](#) [Zbl](#)
- [Barrios et al. 2015] B. Barrios, A. Figalli, and X. Ros-Oton, “Global regularity for the free boundary in the obstacle problem for the fractional Laplacian”, preprint, 2015. To appear in *Amer. J. Math.* [arXiv](#)
- [Bogdan and Byczkowski 1999] K. Bogdan and T. Byczkowski, “Potential theory for the α -stable Schrödinger operator on bounded Lipschitz domains”, *Studia Math.* **133**:1 (1999), 53–92. [MR](#) [Zbl](#)
- [Bogdan et al. 2015] K. Bogdan, B. Siudeja, and A. Stós, “Martin kernel for fractional Laplacian in narrow cones”, *Potential Anal.* **42**:4 (2015), 839–859. [MR](#) [Zbl](#)
- [Bourgain et al. 2001] J. Bourgain, H. Brezis, and P. Mironescu, “Another look at Sobolev spaces”, pp. 439–455 in *Optimal control and partial differential equations*, edited by J. L. Menaldi et al., IOS, Amsterdam, 2001. [MR](#) [Zbl](#)
- [Caffarelli and Lin 2008] L. A. Caffarelli and F.-H. Lin, “Singularly perturbed elliptic systems and multi-valued harmonic functions with free boundaries”, *J. Amer. Math. Soc.* **21**:3 (2008), 847–862. [MR](#) [Zbl](#)
- [Caffarelli and Salsa 2005] L. Caffarelli and S. Salsa, *A geometric approach to free boundary problems*, Graduate Studies in Mathematics **68**, American Mathematical Society, Providence, RI, 2005. [MR](#) [Zbl](#)
- [Caffarelli and Silvestre 2007] L. Caffarelli and L. Silvestre, “An extension problem related to the fractional Laplacian”, *Comm. Partial Differential Equations* **32**:7-9 (2007), 1245–1260. [MR](#) [Zbl](#)
- [Caffarelli and Silvestre 2009] L. Caffarelli and L. Silvestre, “Regularity theory for fully nonlinear integro-differential equations”, *Comm. Pure Appl. Math.* **62**:5 (2009), 597–638. [MR](#) [Zbl](#)
- [Caffarelli et al. 2017] L. Caffarelli, D. De Silva, and O. Savin, “The two membranes problem for different operators”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **34**:4 (2017), 899–932. [MR](#) [Zbl](#)
- [Chen and Song 1998] Z.-Q. Chen and R. Song, “Estimates on Green functions and Poisson kernels for symmetric stable processes”, *Math. Ann.* **312**:3 (1998), 465–501. [MR](#) [Zbl](#)
- [Conti et al. 2005] M. Conti, S. Terracini, and G. Verzini, “Asymptotic estimates for the spatial segregation of competitive systems”, *Adv. Math.* **195**:2 (2005), 524–560. [MR](#) [Zbl](#)
- [Dancer et al. 2012] E. N. Dancer, K. Wang, and Z. Zhang, “The limit equation for the Gross–Pitaevskii equations and S. Terracini’s conjecture”, *J. Funct. Anal.* **262**:3 (2012), 1087–1131. [MR](#) [Zbl](#)
- [Dávila et al. 2015] J. Dávila, M. del Pino, S. Dipierro, and E. Valdinoci, “Concentration phenomena for the nonlocal Schrödinger equation with Dirichlet datum”, *Anal. PDE* **8**:5 (2015), 1165–1235. [MR](#) [Zbl](#)
- [Di Nezza et al. 2012] E. Di Nezza, G. Palatucci, and E. Valdinoci, “Hitchhiker’s guide to the fractional Sobolev spaces”, *Bull. Sci. Math.* **136**:5 (2012), 521–573. [MR](#) [Zbl](#)
- [Dipierro et al. 2016] S. Dipierro, O. Savin, and E. Valdinoci, “Definition of fractional Laplacian for functions with polynomial growth”, preprint, 2016. [arXiv](#)
- [Dipierro et al. 2017] S. Dipierro, O. Savin, and E. Valdinoci, “All functions are locally s -harmonic up to a small error”, *J. Eur. Math. Soc. (JEMS)* **19**:4 (2017), 957–966. [MR](#) [Zbl](#)
- [Friedland and Hayman 1976] S. Friedland and W. K. Hayman, “Eigenvalue inequalities for the Dirichlet problem on spheres and the growth of subharmonic functions”, *Comment. Math. Helv.* **51**:2 (1976), 133–161. [MR](#) [Zbl](#)
- [Garofalo and Ros-Oton 2017] N. Garofalo and X. Ros-Oton, “Structure and regularity of the singular set in the obstacle problem for the fractional Laplacian”, preprint, 2017. [arXiv](#)
- [Michalik 2006] K. Michalik, “Sharp estimates of the Green function, the Poisson kernel and the Martin kernel of cones for symmetric stable processes”, *Hiroshima Math. J.* **36**:1 (2006), 1–21. [MR](#) [Zbl](#)
- [Noris et al. 2010] B. Noris, H. Tavares, S. Terracini, and G. Verzini, “Uniform Hölder bounds for nonlinear Schrödinger systems with strong competition”, *Comm. Pure Appl. Math.* **63**:3 (2010), 267–302. [MR](#) [Zbl](#)
- [Rüland 2015] A. Rüland, “Unique continuation for fractional Schrödinger equations with rough potentials”, *Comm. Partial Differential Equations* **40**:1 (2015), 77–114. [MR](#) [Zbl](#)

- [Shahgholian 2004] H. Shahgholian, “When does the free boundary enter into corner points of the fixed boundary?”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **310** (2004), 213–225. [MR](#) [Zbl](#)
- [Silvestre 2005] L. E. Silvestre, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Ph.D. thesis, The University of Texas at Austin, 2005, available at <https://search.proquest.com/docview/305384256>. [MR](#)
- [Tavares and Terracini 2012] H. Tavares and S. Terracini, “Regularity of the nodal set of segregated critical configurations under a weak reflection law”, *Calc. Var. Partial Differential Equations* **45**:3-4 (2012), 273–317. [MR](#) [Zbl](#)
- [Terracini and Vita 2017] S. Terracini and S. Vita, “On the asymptotic growth of positive solutions to a nonlocal elliptic blow-up system involving strong competition”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* (online publication September 2017).
- [Terracini et al. 2014] S. Terracini, G. Verzini, and A. Zilio, “Uniform Hölder regularity with small exponent in competition-fractional diffusion systems”, *Discrete Contin. Dyn. Syst.* **34**:6 (2014), 2669–2691. [MR](#) [Zbl](#)
- [Terracini et al. 2016] S. Terracini, G. Verzini, and A. Zilio, “Uniform Hölder bounds for strongly competing systems involving the square root of the laplacian”, *J. Eur. Math. Soc. (JEMS)* **18**:12 (2016), 2865–2924. [MR](#) [Zbl](#)
- [Wang and Wei 2016] K. Wang and J. Wei, “On the uniqueness of solutions of a nonlocal elliptic system”, *Math. Ann.* **365**:1-2 (2016), 105–153. [MR](#) [Zbl](#)

Received 18 May 2017. Revised 18 Sep 2017. Accepted 18 Feb 2018.

SUSANNA TERRACINI: susanna.terracini@unito.it

Dipartimento di Matematica “Giuseppe Peano”, Università di Torino, Torino, Italy

GIORGIO TORTONE: giorgio.tortone@unito.it

Dipartimento di Matematica “Giuseppe Peano”, Università di Torino, Torino, Italy

STEFANO VITA: stefano.vita@unito.it

Dipartimento di Matematica “Giuseppe Peano”, Università di Torino, Torino, Italy

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Clément Mouhot	Cambridge University, UK c.mouhot@dpmms.cam.ac.uk
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Alessio Figalli	ETH Zurich, Switzerland alessio.figalli@math.ethz.ch	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zvorski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2018 is US \$275/year for the electronic version, and \$480/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 11 No. 7 2018

On the global stability of a beta-plane equation FABIO PUSATERI and KLAUS WIDMAYER	1587
Airy-type evolution equations on star graphs DELIO MUGNOLO, DIEGO NOJA and CHRISTIAN SEIFERT	1625
On s -harmonic functions on cones SUSANNA TERRACINI, GIORGIO TORTONE and STEFANO VITA	1653
Weighted little bmo and two-weight inequalities for Journé commutators IRINA HOLMES, STEFANIE PETERMICHL and BRETT D. WICK	1693
Estimates for eigenvalues of Aharonov–Bohm operators with varying poles and non-half-integer circulation LAURA ABATANGELO, VERONICA FELLI, BENEDETTA NORIS and MANON NYS	1741
Complex rotation numbers: bubbles and their intersections NATALIYA GONCHARUK	1787
Quantitative stability of the free boundary in the obstacle problem SYLVIA SERFATY and JOAQUIM SERRA	1803