

ANALYSIS & PDE

Volume 16

No. 2

2023

ROBERT XIN DONG, SONG-YING LI AND JOHN N. TREUER

SHARP POINTWISE AND UNIFORM ESTIMATES FOR $\bar{\delta}$



SHARP POINTWISE AND UNIFORM ESTIMATES FOR $\bar{\partial}$

ROBERT XIN DONG, SONG-YING LI AND JOHN N. TREUER

We use weighted L^2 -methods to obtain sharp pointwise estimates for the canonical solution to the equation $\bar{\partial}u = f$ on smoothly bounded strictly convex domains and the Cartan classical domains when f is bounded in the Bergman metric g . We provide examples to show our pointwise estimates are sharp. In particular, we show that on the Cartan classical domains Ω of rank 2 the maximum blow-up order is greater than $-\log \delta_\Omega(z)$, which was obtained for the unit ball case by Berndtsson. For example, for Ω of type IV(n) with $n \geq 3$, the maximum blow-up order is $\delta(z)^{1-n/2}$ because of the contribution of the Bergman kernel. Additionally, we obtain uniform estimates for the canonical solutions on the polydiscs, strictly pseudoconvex domains and the Cartan classical domains under stronger conditions on f .

1. Introduction

The existence and regularity of solutions to the Cauchy–Riemann equation $\bar{\partial}u = f$ on a bounded pseudoconvex domain Ω in \mathbb{C}^n is a fundamental topic in several complex variables. Since the kernel of $\bar{\partial}$ is the set of holomorphic functions, a solution to the Cauchy–Riemann equation is not unique if it exists. However, let $A^2(\Omega) := L^2(\Omega) \cap \ker(\bar{\partial})$ denote the Bergman space over Ω . Then the solution to $\bar{\partial}u = f$ with $u \perp A^2(\Omega)$ is unique, and it is called the canonical solution or L^2 -minimal solution because it has minimal L^2 -norm among all solutions. Hörmander [1965] showed that if Ω is bounded and pseudoconvex and $f \in L^2_{(0,1)}(\Omega)$ is $\bar{\partial}$ -closed, then there exists a solution u that satisfies the estimate $\|u\|_{L^2} \leq C\|f\|_{L^2}$ for some constant C depending only on the diameter of Ω . In view of Hörmander’s result, a natural question arises: does there exist a constant C depending only on Ω such that for any $\bar{\partial}$ -closed $f \in L^\infty_{(0,1)}(\Omega)$ there exists a solution to $\bar{\partial}u = f$ with $\|u\|_\infty \leq C\|f\|_\infty$? If the answer is yes, we say the $\bar{\partial}$ -equation can be solved with uniform estimates on Ω . An important method for solving the $\bar{\partial}$ -equation is the integral representation for solutions. In this method, one constructs a differential form $B(z, w)$ on $\Omega \times \Omega$ which is an $(n, n-1)$ form in w such that solutions to $\bar{\partial}u = f$ can be written as

$$u(z) = \int_{\Omega} B(z, w) \wedge f(w). \quad (1-1)$$

The method of integral representation of solutions was initiated by Cauchy, Leray, Fantappi , etc. On a smoothly bounded strictly pseudoconvex domain Ω in \mathbb{C}^n , Henkin [1970] and Grauert and Lieb [1970] constructed integral kernels $B(z, w)$ such that u given by (1-1) satisfies $\|u\|_\infty \leq C\|f\|_\infty$. Kerzman [1971] improved the estimate by showing that $\|u\|_{C^\alpha(\bar{\Omega})} \leq C_\alpha\|f\|_\infty$ for any $0 < \alpha < \frac{1}{2}$. Henkin and Romanov [1971] obtained the sharp estimate $\|u\|_{C^{1/2}(\bar{\Omega})} \leq C\|f\|_\infty$. For more results on strictly pseudoconvex

MSC2020: primary 32A25; secondary 32A36, 32M15, 32W05.

Keywords: L^2 minimal solution, canonical solution, $\bar{\partial}$ equation, Cartan classical domain, Bergman kernel, Bergman metric.

domains, the reader may consult the papers [Krantz 1976; Range and Siu 1972; 1973] and the books [Chen and Shaw 2001; Fornæss and Stensønes 1987; Range 1986].

When the class of domains under consideration is changed from strictly pseudoconvex to weakly pseudoconvex, it is no longer possible to conclude in generality the existence of uniform estimates for $\bar{\partial}$. Berndtsson [1993], Fornæss [1986] and Sibony [1980] constructed examples of weakly pseudoconvex domains in \mathbb{C}^2 and \mathbb{C}^3 where uniform estimates for $\bar{\partial}$ fail. More strikingly, Fornæss and Sibony [1993] constructed a smoothly bounded pseudoconvex domain $\Omega \subset \mathbb{C}^2$ such that $\partial\Omega$ is strictly pseudoconvex except at one point, but any solution to $\bar{\partial}u = f$ for some given $\bar{\partial}$ -closed $f \in L^\infty_{(0,1)}(\Omega)$ does not belong to $L^p(\Omega)$ for any $2 < p \leq \infty$. Range [1978] gave uniform estimates on bounded convex domains in \mathbb{C}^2 with real analytic boundaries, and in [Range 1990] gave Hölder estimates on pseudoconvex domains of finite type in \mathbb{C}^2 . See [Laurent-Thiébaud and Leiterer 1993; Michel and Shaw 1999] for related results. Of particular interest is the unit polydisc $\mathbb{D}^n := \mathbb{D}(0, 1)^n \subset \mathbb{C}^n$, which is pseudoconvex with nonsmooth boundary. When $n = 2$, Henkin [1971] showed that there exists a constant C such that $\|u\|_\infty \leq C\|f\|_\infty$ for any $f \in C^1_{(0,1)}(\bar{\mathbb{D}}^2)$. Landucci [1975] obtained the same uniform estimate for the canonical solution on \mathbb{D}^2 . Chen and McNeal [2020] and Fassina and Pan [2019] generalized Henkin’s result to higher dimensions when additional regularity assumptions on f are imposed. It remains open whether uniform estimates hold on \mathbb{D}^n with $n \geq 2$ when f is only assumed to be bounded. See [Dong et al. 2020; Fornæss et al. 2011; Grundmeier et al. 2022] for related results.

A class of pseudoconvex domains in \mathbb{C}^n including \mathbb{D}^n and the unit ball \mathbb{B}^n are the so-called bounded symmetric domains, which up to biholomorphism are Cartesian product(s) of the Cartan classical domains of types I to IV and two domains of exceptional types. In [Henkin and Leiterer 1983, p. 200], the authors asked whether there exists a uniform estimate for the $\bar{\partial}$ -equation on the Cartan classical domains of rank at least 2. Additionally, [Sergeev 1994] conjectured that the $\bar{\partial}$ -equation cannot be solved with uniform estimates on the Cartan classical domains of type IV of dimension $n \geq 3$.

Let $g = (g_{j\bar{k}})_{j,k=1}^n$ be the Bergman metric on a domain Ω . For a $(0, 1)$ -form $f = \sum_{j=1}^n f_j d\bar{z}_j$, one defines

$$\|f\|_{g,\infty}^2 := \text{ess sup} \left\{ \sum_{j,k=1}^n g^{j\bar{k}}(z) f_k(z) \overline{f_j(z)} : z \in \Omega \right\},$$

where $(g^{j\bar{k}})^\tau = (g_{j\bar{k}})^{-1}$; see (3-1) for details. Berndtsson used weighted L^2 estimates of Donnelly–Fefferman type to prove the following pointwise and uniform estimates.

Theorem 1.1 [Berndtsson 1997, 2001]. *There is a numerical constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f on \mathbb{B}^n , the canonical solution to $\bar{\partial}u = f$ satisfies*

$$|u(z)| \leq C\|f\|_{g,\infty} \log \frac{2}{1-|z|}, \tag{1-2}$$

and for any $\epsilon > 0$,

$$\|u\|_\infty \leq \frac{C}{\epsilon} \|(1-|z|^2)^{-\epsilon} f\|_{g,\infty}. \tag{1-3}$$

The estimate (1-2) is sharp. If $f(z) := \sum_{k=1}^n z_k (|z|^2 - 1)^{-1} d\bar{z}_k$ then f is $\bar{\partial}$ -closed, $\|f\|_{g,\infty} = 1$ and the canonical solution to $\bar{\partial}u = f$ is $u = \log(1 - |z|^2) - C_n$, which shows the sharpness of (1-2).

Berndtsson [2001] also pointed out that his proof should generalize to other domains when enough information about the Bergman kernel is known. This result [Berndtsson 1997, 2001] was improved in [Schuster and Varolin 2014] via the “double twisting” method.

Motivated by Berndtsson’s results (1-2) and (1-3) and the problems raised by Henkin and Leiterer [1983] and Sergeev [1994], we study sharp pointwise estimates for $\bar{\partial}u = f$ for any $\bar{\partial}$ -closed $(0, 1)$ -form f with $\|f\|_{g,\infty} < \infty$ and uniform estimates under stronger conditions on f . We generalize Berndtsson’s results from \mathbb{B}^n to smoothly bounded strictly pseudoconvex domains and the Cartan classical domains. Our main theorem, [Theorem 1.2](#) (see also [Theorem 3.3](#)), for pointwise estimates is stated as follows.

Theorem 1.2. *Let Ω be a smoothly bounded strictly convex domain, a Cartan classical domain or the polydisc, whose Bergman kernel and metric are denoted by K and g , respectively. Then there is a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f with $\|f\|_{g,\infty} < \infty$, the canonical solution to $\bar{\partial}u = f$ satisfies*

$$|u(z)| \leq C \|f\|_{g,\infty} \int_{\Omega} |K(z, w)| dv_w, \quad z \in \Omega. \quad (1-4)$$

Remarks. (i) When Ω is a smoothly bounded strictly pseudoconvex domain, by Fefferman’s asymptotic expansion for the Bergman kernel,

$$\int_{\Omega} |K(z, w)| dv_w \approx C \log \frac{1}{\delta_{\Omega}(z)} \approx \log K(z, z), \quad z \rightarrow \partial\Omega.$$

In this case, the estimate (1-4) is sharp. Take for example $\Omega = \mathbb{B}^n$ and $u(z) = \log K(z, z) - c$, where c is chosen so that $P[u] = 0$.

(ii) We will show in [Section 3](#), [Theorem 3.4](#), that if Ω is a smoothly bounded strictly pseudoconvex domain, then (1-4) holds for a solution u which may not be canonical.

(iii) When Ω is the unit polydisc \mathbb{D}^n , one has

$$\int_{\Omega} |K(z, w)| dv_w \approx \prod_{j=1}^n \log \frac{2}{1-|z_j|}, \quad z \rightarrow \partial\Omega.$$

(iv) When Ω is a Cartan classical domain of rank greater than or equal to 2, the blow-up order of $\int_{\Omega} |K(z, w)| dv_w$ depends on the direction in which z approaches $\partial\Omega$ and it may be larger than $-\log \delta_{\Omega}(z)$. For example, if $z = tI_2 \in \Omega := \text{II}(2)$, then $\int_{\Omega} |K(z, w)| dv_w \approx \delta_{\Omega}(z)^{-1/2}$ as $t \rightarrow 1^-$. Moreover, if $z = te_1 \oplus te_2 \in \text{IV}(n)$ with $e_j \in \mathcal{U}$ and $n \geq 3$, then $\int_{\Omega} |K(z, w)| dv_w \approx \delta_{\Omega}(z)^{1-(n/2)}$ as $t \rightarrow 1^-$. Here \mathcal{U} denotes the characteristic boundary of Ω .

(v) In [Section 6B](#), we show the estimate (1-4) is sharp on the Cartan classical domains.

Our main theorem for uniform estimates is stated as follows, as a combination of [Theorems 4.1](#) and [4.2](#).

Theorem 1.3. *Let Ω be either the unit polydisc or a smoothly bounded strictly convex domain, whose Bergman kernel and metric are denoted by K and g , respectively. Then for any $p \in (1, \infty)$, there is a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f , the canonical solution to $\bar{\partial}u = f$ satisfies*

$$\|u\|_{\infty} \leq C \left\| f(\cdot) \left(\int_{\Omega} |K(\cdot, w)| dv_w \right)^p \right\|_{g,\infty}.$$

For Cartan classical domains, we give a uniform estimate under condition (5-2) in Theorem 5.4.

This paper is organized as follows: In Section 2, we recall and prove some properties of the Bergman kernel and metric which will be used later. In Section 3, we use L^2 -methods to establish pointwise estimates on strictly pseudoconvex domains and the Cartan classical domains. In Sections 4 and 5, we obtain uniform estimates on the polydiscs, strictly pseudoconvex domains and the Cartan classical domains under various conditions on f . In Section 6, we verify the sharpness of our pointwise estimates on the Cartan classical domains; in particular, on $IV(n)$ with $n \geq 3$ we show the estimate has maximum blow-up order $\delta^{1-(n/2)}(z)$.

2. Bergman kernel and metric

The Bergman space $A^2(\Omega)$ on a domain $\Omega \subset \mathbb{C}^n$ is the closed holomorphic subspace of $L^2(\Omega)$. The Bergman projection is the orthogonal projection $P_\Omega : L^2(\Omega) \rightarrow A^2(\Omega)$ given by

$$P_\Omega[f](z) = \int_\Omega K(z, w)f(w) dv(w),$$

where $K(z, w)$ is the Bergman kernel on Ω and dv is the Lebesgue \mathbb{R}^{2n} measure. We will write $K(z)$ to denote the on-diagonal Bergman kernel $K(z, z)$. When Ω is bounded, the complex Hessian of $\log K(z)$ induces the Bergman metric $B_\Omega(z; X)$ defined by

$$B_\Omega(z; X) := \left(\sum_{j,k=1}^n g_{j\bar{k}} X_j \bar{X}_k \right)^{1/2}, \quad g_{j\bar{k}}(z) := \frac{\partial^2}{\partial z_j \partial \bar{z}_k} \log K(z), \quad \text{for } z \in \Omega, \quad X \in \mathbb{C}^n.$$

The Bergman distance between $z, w \in \Omega$ is

$$\beta_\Omega(z, w) := \inf \left\{ \int_0^1 B_\Omega(\gamma(t); \gamma'(t)) dt \right\},$$

where the infimum is taken over all piecewise C^1 -curves $\gamma : [0, 1] \rightarrow \Omega$ such that $\gamma(0) = z$ and $\gamma(1) = w$. Throughout the paper,

$$B_a(\epsilon) := \{z \in \Omega : \beta_\Omega(z, a) \leq \epsilon\} \tag{2-1}$$

will denote the hyperbolic ball in the Bergman metric centered at $a \in \Omega$ of radius ϵ . Additionally, $K(z, w)$, P_Ω and g will always denote the Bergman kernel, the Bergman projection on Ω and the Bergman metric, respectively.

Consider a convex domain Ω that contains no complex lines and $a \in \Omega$. Choose any $a^1 \in \partial\Omega$ such that $\tau_1(a) := |a - a^1| = \text{dist}(a, \partial\Omega)$ and define $V_1 = a + \text{span}(a^1 - a)^\perp$. Let $\Omega_1 = \Omega \cap V_1$ and choose any $a^2 \in \partial\Omega_1$ such that $\tau_2(a) := \|a - a^2\| = \text{dist}(a, \partial\Omega_1)$. Let $V_2 = a + \text{span}(a^1 - a, a^2 - a)^\perp$ and $\Omega_2 = \Omega \cap V_2$. Repeat this process to obtain a^1, \dots, a^n and $w_k = (a^k - a)/\|a^k - a\|$, $1 \leq k \leq n$. Define

$$D(a; w, r) = \{z \in \mathbb{C}^n : |\langle z - a, w_k \rangle| < r_k, 1 \leq k \leq n\} \tag{2-2}$$

and

$$D(a, r) = \{z \in \mathbb{C}^n : |z_k - a_k| < r_k, 1 \leq k \leq n\}.$$

By [Nikolov and Pflug 2003, Theorem 2], for convex domains that contain no complex lines, the Kobayashi metric and the Bergman metric are comparable. It follows by [Nikolov and Trybuła 2015, Corollary 2] that if Ω is a convex domain with no complex lines, then for every $\epsilon > 0$ there exists constants C_1 and C_2 such that for any a ,

$$D(a; w, C_1\tau(a)) \subset B_a(\epsilon) \subset D(a; w, C_2\tau(a)). \quad (2-3)$$

By [Nikolov and Pflug 2003, Theorem 1],

$$\frac{1}{4^n} \leq K(a) \prod_{j=1}^n \pi \tau_j^2(a) \leq \frac{(2n)!}{2^n},$$

which implies that

$$\left(\frac{C_1}{2}\right)^{2n} \leq K(a)v(B_a(\epsilon)) \leq (2n)! \left(\frac{C_2}{2}\right)^n.$$

For any open subset A of Ω , we define

$$\|\bar{\partial}u\|_{g,\infty,A} = \|\bar{\partial}u(z)|_g\|_{L^\infty(A)}.$$

In the proofs of this paper, C will denote a numerical constant which may be different at each appearance. The Cauchy–Pompeiu formula gives the following useful proposition.

Proposition 2.1. *Let Ω be a bounded convex domain. For any $\epsilon > 0$ sufficiently small, there exists a constant C such that for any complex-valued C^1 function u on Ω*

$$|u(a)| \leq C \oint_{B_a(\epsilon)} |u(z)| dv_z + C \|\bar{\partial}u\|_{g,\infty,B_a(\epsilon)}.$$

Proof. After a complex rotation, without loss of generality, we may assume the standard basis for \mathbb{C}^n is $(w_k)_{k=1}^n$, using the notation of (2-2). Let $r_k(a) = C_1\tau_k(a)$, where C_1 is the same constant as in (2-3). Define the metric

$$M_A(z; X) = \left(\sum_{k=1}^n \frac{|X_k|^2}{\tau_k(z)^2} \right)^{1/2}, \quad X \in \mathbb{C}^n.$$

It was proved in [McNeal 2001] (see also [McNeal 1994; Nikolov and Pflug 2003]) that

$$M_A(z; X) \approx B_\Omega(z; X), \quad X \in \mathbb{C}^n,$$

where \approx is independent of z and X . So we can choose holomorphic coordinates such that

$$\frac{1}{C} D \left[\frac{1}{\tau_1^2}, \dots, \frac{1}{\tau_n^2} \right] \leq [g_{i\bar{j}}] \leq CD \left[\frac{1}{\tau_1^2}, \dots, \frac{1}{\tau_n^2} \right],$$

where $D[a_1, \dots, a_n]$ is a diagonal matrix with diagonal entries a_1, \dots, a_n . Therefore

$$\frac{1}{C} D[\tau_1^2, \dots, \tau_n^2] \leq [g_{i\bar{j}}]^{-1} \leq CD[\tau_1^2, \dots, \tau_n^2].$$

Additionally, by (2-3) and the definition of the hyperbolic ball (2-1),

$$\tau_k(a) \leq C\tau_k(z), \quad z \in D(a; w, C_1\tau(a)),$$

and the constant C is independent of a . Therefore, for $a = (a_1, \dots, a_n) \in \Omega$,

$$\begin{aligned} r_1(a) \|\bar{\partial}_1 u(\cdot, a_2, \dots, a_n)\|_{L^\infty(D(a_1, r_1))} &\leq C \sup_{z \in D(a; w, r)} \sum_{k=1}^n r_k(z) \left| \frac{\partial u}{\partial \bar{w}_k}(z) \right| \\ &\leq C \sup_{z \in D(a; w, r)} \left(\sum_{k=1}^n \tau_k^2(z) \left| \frac{\partial u}{\partial \bar{w}_k}(z) \right|^2 \right)^{1/2} \\ &\leq C \|\bar{\partial} u\|_{g, \infty, D(a; w, r)} \leq C \|\bar{\partial} u\|_{g, \infty, B_a(\epsilon)}. \end{aligned}$$

By Stokes' theorem, for $0 < s_k < r_k$,

$$u(a) = \frac{1}{2\pi i} \int_{|w_1 - a_1| = s_1} \frac{u(w_1, a_2, \dots, a_n)}{w_1 - a_1} dw_1 + \frac{1}{2\pi i} \int_{|w_1 - a_1| < s_1} \frac{\partial u}{\partial \bar{w}_1} \frac{1}{w_1 - a_1} dw_1 \wedge d\bar{w}_1.$$

By polar coordinates and (2-3),

$$\begin{aligned} |u(a)| &\leq \frac{1}{\pi r_1^2} \int_{|z_1 - a_1| < r_1} |u(w_1, a_2, \dots, a_n)| dv_{z_1} + \frac{2r_1}{3} \|\bar{\partial}_1 u(\cdot, a_2, \dots, a_n)\|_{L^\infty(D(a_1, r_1))} \\ &\leq \frac{1}{\pi r_1^2} \int_{|w_1 - a_1| < r_1} |u(w_1, a_2, \dots, a_n)| dv_{w_1} + C \|\bar{\partial} u\|_{g, \infty, B_a(\epsilon)}. \end{aligned}$$

Using the same estimate on the disc $|w_k - a_k| < s_k$ for $2 \leq k \leq n$,

$$\begin{aligned} |u(a)| &\leq \oint_{D(a, C_1 \tau(a))} |u(w_1, \dots, w_n)| dv_w + C \|\bar{\partial} u\|_{g, \infty, B_a(\epsilon)} \\ &\leq C^{2n} \oint_{B_a(\epsilon)} |u(w)| dv_w + C \|\bar{\partial} u\|_{g, \infty, B_a(\epsilon)}. \end{aligned} \quad \square$$

We remark that Proposition 2.1 also holds for smoothly bounded strictly pseudoconvex domains.

For positive real-valued functions f and g on Ω , we say $f \approx g$ for $z \in B_a(\epsilon)$ if for every $\epsilon > 0$ sufficiently small, there exists a constant $C = C(\epsilon, \Omega)$ such that

$$C^{-1} \leq f(z)g(z)^{-1} \leq C, \quad z \in B_a(\epsilon)$$

for all $a \in \Omega$. A similar definition holds for $f \approx g$ for $z \in \Omega$.

A domain Ω is homogeneous if it has a transitive (holomorphic) automorphism group. For convex homogeneous domains, the following results are known; see [Ishi and Yamaji 2011].

Proposition 2.2. *Let Ω be a bounded homogeneous convex domain. Then*

$$|K(z, a)| \approx K(a) \approx \frac{1}{v(B_a(\epsilon))}, \quad z \in B_a(\epsilon),$$

and for any $\epsilon > 0$, there is a C_ϵ such that for any $a \in \Omega$

$$\max_{w \in B_a(\epsilon)} \left| \frac{K(z, w)}{K(z, a)} \right| \leq C_\epsilon, \quad z \in \Omega.$$

Let Ω be a smoothly bounded strictly pseudoconvex domain in \mathbb{C}^n and let $-r \in C^\infty(\bar{\Omega})$ be a strictly plurisubharmonic defining function for Ω . Define

$$X(z, w) = r(w) + \sum_{j=1}^n \frac{\partial r(w)}{\partial z_j} (z_j - w_j) + \frac{1}{2} \sum_{i,j=1}^n \frac{\partial^2 r(w)}{\partial z_i \partial z_j} (z_i - w_i)(z_j - w_j).$$

It was proved by Fefferman [1974] that there is a $\delta > 0$ such that

$$K(z, w) = \frac{F(z, w)}{X(z, w)^{n+1}} + G(z, w) \log X(z, w)$$

for all $(z, w) \in R_\delta(\Omega) = \{(z, w) \in \Omega \times \Omega : r(z) + r(w) + |z - w|^2 < \delta\}$, where $F, G \in C^\infty(\bar{\Omega} \times \bar{\Omega})$ and $F(z, z) > 0$ on $\partial\Omega$.

When Ω is a smoothly bounded strictly convex domain, the definition of $X(z, w)$ can be simplified. In fact, one can take

$$X(z, a) = h_a(z) = r(a) + \sum_{j=1}^n \frac{\partial r}{\partial z_j}(a)(z_j - a_j).$$

We can take $-r(z)$ to be strictly convex. Then by Taylor's theorem, one can easily see that

$$\operatorname{Re} h_a(z) \approx \operatorname{Re} X(z, w).$$

Moreover, for any $a, z \in \Omega$, we will write $\tilde{z} = (x_j)_{j=1}^{2n}$ and $\tilde{a} = (a_j)_{j=1}^{2n}$ if $z = (x_{2j-1} + ix_{2j})_{j=1}^n$ and $a = (a_{2j-1} + ia_{2j})_{j=1}^n$. If we apply Taylor's theorem on the line segment between \tilde{a} and \tilde{z} , then there is a $\theta \in (0, 1)$ such that

$$\operatorname{Re} h_a(z) = r(z) - \sum_{i,j=1}^{2n} \frac{\partial^2 r(\tilde{a} + \theta(\tilde{z} - \tilde{a}))}{\partial \tilde{z}_i \partial \tilde{z}_j} (\tilde{z}_i - \tilde{a}_i)(\tilde{z}_j - \tilde{a}_j) \approx r(z) + |z - a|^2.$$

Therefore, for any $a \in \Omega$,

$$|h_a(z)| \approx \frac{r(z) + r(a)}{2} + |z - a|^2 + \left| \operatorname{Im} \sum_{j=1}^n \frac{\partial r(a)}{\partial z_j} (z_j - a_j) \right|, \quad z \in \Omega.$$

In particular, this implies $h_a(z) \neq 0$. Therefore, by Fefferman's asymptotic expansion [1974] on strictly pseudoconvex domains mentioned above, we know the following.

Lemma 2.3 [Fefferman 1974]. *Let Ω be a smoothly bounded strictly convex domain. Then,*

$$|h_a(z)|^{-n-1} \approx K(a) \approx \frac{1}{v(B_a(\epsilon))}, \quad z \in B_a(\epsilon),$$

and there is a constant C such that

$$\int_{\Omega} |h_a(z)|^{-n-1} dv_z \approx \int_{\Omega} |K(z, a)| dv_z \approx \log \frac{C}{\delta_{\Omega}(a)}, \quad a \in \Omega,$$

where $\delta_{\Omega}(\cdot)$ is the Euclidean distance function to $\partial\Omega$. Moreover, for any $\epsilon > 0$, there is a constant C_ϵ such that for any $a \in \Omega$

$$\max_{w \in B_a(\epsilon)} |K(z, w) h_a(z)^{n+1}| \leq C_\epsilon, \quad z \in \Omega.$$

Note. We provide some insight into the integration of $|K(z, w)|$ as a Forelli–Rudin-type integral. Roughly, one can view $\partial\Omega$ as a space of homogeneous type with Borel measure dX and quasidistance $|X(z, w)|$. Write

$$|K(z, w)| \approx (\delta(z) + t + |X(\pi(z), \pi(w))|)^{-n-1},$$

where $\pi(z)$ and $\pi(w)$ are the projections of z and w on $\partial\Omega$ along the outer normal direction and $z, w \in R_\delta$. It follows that

$$\int_{\partial\Omega} (\delta(z) + t + |X|)^{-n-1} dX \approx (\delta + t)^{-1}.$$

Consequently,

$$\int_{\Omega} |K(z, w)| dv(w) \approx \int_0^C (\delta(z) + t)^{-1} dt \approx \log \frac{1}{\delta(z)}.$$

For more information, one can consult the paper of Beatrous and the second author [Beatrous and Li 1993] and the papers of Krantz and the second author [Krantz and Li 2001a; 2001b].

Lemma 2.4. *Let Ω be either a smoothly bounded strictly pseudoconvex domain or a Cartan classical domain. Let $\phi(z) := \gamma \log K(z)$ with $\gamma > 0$. Then, for γ sufficiently small,*

$$\int_{\Omega} e^{\phi(z)} dv_z < \infty \quad \text{and} \quad \|\bar{\partial}\phi\|_{i\bar{\partial}\bar{\partial}\phi}^2 \leq \frac{1}{2}.$$

Proof. When Ω is a smoothly bounded strictly pseudoconvex domain, from Fefferman’s asymptotic expansion for the Bergman kernel, one has

$$\phi(z) \approx \gamma \log \frac{1}{\delta(z)} \quad \text{and} \quad \int_{\Omega} e^{\phi(z)} dv(z) \approx \int_{\Omega} \left(\frac{1}{\delta(z)}\right)^\gamma dv \approx \int_0^1 t^{-\gamma} dt < \infty.$$

Notice that

$$\|\bar{\partial}\phi\|_{i\bar{\partial}\bar{\partial}\phi}^2 = \gamma \|\bar{\partial} \log K\|_g^2,$$

where g is the Bergman metric. From Fefferman’s asymptotic expansion formula (see also [Donnelly 1994]), one gets the boundedness of $\|\bar{\partial} \log K\|_g^2$. Choose $\gamma > 0$ small enough so that $\|\bar{\partial}\phi\|_g^2 < \frac{1}{2}$. For the Cartan classical domains, the first inequality follows from explicit formulas of the Bergman kernel [Hua 1963], and we compute the precise value of $\|\bar{\partial}\phi\|_{i\bar{\partial}\bar{\partial}\phi}$ in Section 6A. □

3. Pointwise estimates

An upper semicontinuous function ϕ defined on a domain $\Omega \subset \mathbb{C}^n$ with values in $\mathbb{R} \cup \{-\infty\}$ is called plurisubharmonic if its restriction to every complex line is subharmonic. Let $L^2(\Omega, \phi)$ denote the set of measurable functions h satisfying $\int_{\Omega} |h(z)|^2 e^{-\phi(z)} dv_z < \infty$. A C^2 function ϕ is called strongly plurisubharmonic if $i\bar{\partial}\bar{\partial}\phi$ is strictly positive definite. Now, let Ω be a bounded pseudoconvex domain and ϕ be strongly plurisubharmonic on Ω . Then, for any $(0, 1)$ -form $f = \sum_{k=1}^n f_k(z) d\bar{z}_k$, define the norm of f induced by $i\bar{\partial}\bar{\partial}\phi$ as (see also [Błocki 2005])

$$|f|_{i\bar{\partial}\bar{\partial}\phi}^2(z) := \sum_{j,k=1}^n \phi^{j\bar{k}}(z) \overline{f_j(z)} f_k(z), \tag{3-1}$$

where $(\phi^{j\bar{k}})^\tau$ equals the inverse of the complex Hessian matrix $H(\phi)$. Demailly's reformulation [1982; 2012] of Hörmander's theorem [1965] says that for any $\bar{\partial}$ -closed $(0, 1)$ -form f , the $L^2(\Omega, \phi)$ minimal solution to $\bar{\partial}u = f$ satisfies

$$\int_{\Omega} |u|^2 e^{-\phi} dv \leq \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\phi} dv. \quad (3-2)$$

From this we see that when the $(0, 1)$ -form f is bounded in the Bergman metric g , the canonical solution u to $\bar{\partial}u = f$ exists and the right-hand side of the estimate (3-2) is finite because it is dominated by a constant times a positive power of the Euclidean volume.

Donnelly and Fefferman [1983] (see also [Berndtsson 1993; 1996; 1997; McNeal 1996; Siu 1996]) modified Hörmander's theorem further as follows.

Theorem 3.1 (Donnelly–Fefferman-type estimate). *Let Ω be a bounded pseudoconvex domain in \mathbb{C}^n , and let ψ and ϕ be plurisubharmonic functions on Ω such that $i\bar{\partial}\bar{\partial}\phi > 0$ and $|\partial\phi|_{i\bar{\partial}\bar{\partial}\phi}^2 \leq \frac{1}{2}$. Then the $L^2(\Omega, \psi + \frac{1}{2}\phi)$ minimal solution u_0 to $\bar{\partial}u = f$ satisfies*

$$\int_{\Omega} |u_0|^2 e^{-\psi} dv \leq 4 \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi} dv. \quad (3-3)$$

Next we prove the following lemma, using the estimates (3-2) and (3-3).

Lemma 3.2. *Let Ω be a bounded pseudoconvex domain and f be a $\bar{\partial}$ -closed $(0, 1)$ -form on Ω . Let ψ and ϕ be plurisubharmonic on Ω and u_0 and u_1 be the L^2 -minimal solutions to $\bar{\partial}u = f$ in $L^2(\Omega, \psi + \frac{1}{2}\phi)$ and $L^2(\Omega, \phi)$, respectively. Suppose B is a compact subset of Ω and $h \in L^\infty(\Omega)$ with support in B .*

(i) *If $i\bar{\partial}\bar{\partial}\phi > 0$ and $\|\partial\phi\|_{i\bar{\partial}\bar{\partial}\phi}^2 \leq \frac{1}{2}$ on Ω , then*

$$\int_B |u_0| dv \leq 2 \left(\int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi} dv \right)^{1/2} \left(\int_B e^{\psi} dv \right)^{1/2} \quad (3-4)$$

and

$$\left| \int_{\Omega} u_0 \overline{P(h)} dv \right| \leq 2v(B) \|h\|_{\infty} \left(\int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi} dv \right)^{1/2} \left(\int_{\Omega} \max_{w \in B} |K(z, w)|^2 e^{\psi(z)} dv_z \right)^{1/2}. \quad (3-5)$$

(ii) *Additionally,*

$$\int_B |u_1| dv \leq 2 \left(\int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\phi} dv \right)^{1/2} \left(\int_B e^{\phi} dv \right)^{1/2}$$

and

$$\left| \int_{\Omega} u_1 \overline{P(h)} dv \right| \leq 2v(B) \|h\|_{\infty} \left(\int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\phi} dv \right)^{1/2} \left(\int_{\Omega} \max_{w \in B} |K(z, w)|^2 e^{\phi(z)} dv_z \right)^{1/2}.$$

Proof. Let χ_B denote the characteristic function on B , and let $\beta := \chi_B(u_0(z)/|u_0(z)|)$. By (3-3),

$$\left(\int_B |u_0| dv \right)^2 = \left| \int_{\Omega} u_0 \bar{\beta} dv \right|^2 \leq \int_{\Omega} |u_0|^2 e^{-\psi} dv \int_B |\beta|^2 e^{\psi} dv \leq 4 \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi} dv \int_B e^{\psi} dv,$$

which proves (3-4). Notice that

$$\begin{aligned} \left| \int_{\Omega} u_0 \overline{P(h)} dv \right|^2 &\leq \int_{\Omega} |u_0|^2 e^{-\psi} dv \int_{\Omega} |P(h)|^2 e^{\psi} dv \\ &\leq 4 \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi} dv \cdot v^2(B) \|h\|_{\infty}^2 \int_{\Omega} \max_{w \in B} |K(z, w)|^2 e^{\psi(z)} dv_z, \end{aligned}$$

which proves (3-5). Part (ii) can be proved similarly using Hörmander’s estimate (3-2) in place of Donnelly–Fefferman’s estimate (3-3). \square

Theorem 3.3 (key estimate). *Let Ω be a Cartan classical domain or a smoothly bounded strictly convex domain. Then there is a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f on Ω with $\|f\|_{g, \infty} < \infty$, the canonical solution to $\bar{\partial}u = f$ satisfies*

$$|u(z)| \leq C \|f\|_{g, \infty} \int_{\Omega} |K(z, w)| dv_w, \quad z \in \Omega.$$

Proof. From the discussion after (3-2) we see that the canonical solution to $\bar{\partial}u = f$ exists. Suppose first that Ω is a Cartan classical domain. For an arbitrary $a \in \Omega$ and any sufficiently small $\epsilon > 0$, let $\beta := \chi_{B_a(\epsilon)}(u(z)/|u(z)|)$, where $\chi_{B_a(\epsilon)}$ is the characteristic function of the hyperbolic ball $B_a(\epsilon)$. Let $\phi := \gamma \log K(z)$ be a plurisubharmonic function on Ω for some chosen γ that satisfies the condition in Lemma 2.4. Define $\psi(z) := \psi_a(z) := -\log|K(z, a)|$. Then ψ_a is pluriharmonic and bounded on Ω . Also define the function

$$\phi_0 := \psi_a + \frac{1}{2}\phi,$$

and let u_0 be the $L^2(\Omega, \phi_0)$ minimal solution to the equation $\bar{\partial}v = f$. Then by Theorem 3.1,

$$\int_{\Omega} |u_0|^2 e^{-\psi} dv \leq 4\gamma^{-1} \int_{\Omega} |f|_g^2 e^{-\psi} dv \leq 4\gamma^{-1} \|f\|_{g, \infty}^2 \int_{\Omega} e^{-\psi} dv < \infty,$$

which implies that $u_0 \in L^2(\Omega)$. So $u - u_0 \in A^2(\Omega)$ and

$$\int_{B_a(\epsilon)} |u| dv = \int_{\Omega} u \bar{\beta} dv = \int_{\Omega} u(\overline{\beta - P(\beta)}) dv = \int_{\Omega} u_0(\overline{\beta - P(\beta)}) dv = \int_{\Omega} u_0 \bar{\beta} dv - \int_{\Omega} u_0 \overline{P(\beta)} dv.$$

By Lemma 2.4 and (3-4) in Lemma 3.2,

$$\begin{aligned} \left| \int_{\Omega} u_0 \bar{\beta} dv \right|^2 &\leq 4 \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi_a} dv \int_{B_a(\epsilon)} e^{\psi_a} dv \\ &\leq C \|f\|_{g, \infty}^2 \int_{\Omega} |K(z, a)| dv_z \int_{B_a(\epsilon)} |K(z, a)|^{-1} dv_z \\ &\leq C_{\epsilon} \|f\|_{g, \infty}^2 \int_{\Omega} |K(z, a)| dv_z \cdot v(B_a(\epsilon)) K(a)^{-1} \\ &\leq C_{\epsilon} \|f\|_{g, \infty}^2 v^2(B_a(\epsilon)) \int_{\Omega} |K(z, a)| dv_z, \end{aligned}$$

where the last two inequalities hold due to [Proposition 2.2](#) and C_ϵ is a constant depending on ϵ . On the other hand, by (3-5) in [Lemma 3.2](#) and [Proposition 2.2](#) again,

$$\begin{aligned} \left| \int_{\Omega} u_0 \overline{P(\beta)} dv \right|^2 &\leq C v^2(B_a(\epsilon)) \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\psi_a} dv \int_{\Omega} \max_{w \in \overline{B_a(\epsilon)}} |K(z, w)|^2 e^{\psi_a(z)} dv_z \\ &\leq C_\epsilon v^2(B_a(\epsilon)) \int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2(z) |K(z, a)| dv_z \int_{\Omega} |K(z, a)|^{2-1} dv_z \\ &\leq C_\epsilon \|f\|_{g, \infty}^2 v^2(B_a(\epsilon)) \left(\int_{\Omega} |K(z, a)| dv_z \right)^2. \end{aligned}$$

Combining the above estimates, one easily sees that

$$\frac{1}{v(B_a(\epsilon))} \int_{B_a(\epsilon)} |u| dv \leq C_\epsilon \|f\|_{g, \infty} \int_{\Omega} |K(z, a)| dv_z.$$

Fix $\epsilon > 0$. By [Proposition 2.1](#), there exists a constant C depending only on Ω such that

$$|u(a)| \leq C \|f\|_{g, \infty} \int_{\Omega} |K(z, a)| dv_z.$$

If Ω is instead a smoothly bounded strictly convex domain, then let $\psi_a(z) = (n + 1) \log|h_a(z)|$, repeat the argument for the Cartan classical domains and use [Lemma 2.3](#). \square

In [Section 6](#), [Proposition 6.1](#), we show that the estimate in [Theorem 3.3](#) is sharp for the Cartan classical domains. When Ω is the unit ball \mathbb{B}^n , the *key estimate* reduces to Berndtsson’s result (1-2). Now we generalize (1-2) and (1-4) to smoothly bounded strictly pseudoconvex domains.

Theorem 3.4. *Let Ω be a smoothly bounded strictly pseudoconvex domain. Then there is a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f on Ω with $\|f\|_{g, \infty} < \infty$, there is a solution u to $\bar{\partial}u = f$ such that*

$$|u(z)| \leq C \|f\|_{g, \infty} \log(1 + K(z)), \quad z \in \Omega.$$

Proof. Let $r(z)$ be a strongly plurisubharmonic defining function for Ω such that $r(z) \in C^\infty(\bar{\Omega})$ and $r > 0$ in Ω . Consider the polynomial

$$X(z, w) := r(w) + \sum_{j=1}^n \left(\frac{\partial r}{\partial w_j} \Big|_w (z_j - w_j) \right) + \frac{1}{2} \sum_{j,k=1}^n \left(\frac{\partial^2 r}{\partial w_j \partial w_k} \Big|_w (z_j - w_j)(z_k - w_k) \right).$$

Define the region $R_\epsilon = \{(z, w) : z, w \in \Omega, r(z) + r(w) + |z - w|^2 < \epsilon\}$. For $(z, w) \in R_\epsilon$, Fefferman [1974] showed the Bergman kernel on Ω can be expressed as

$$K(z, w) = \frac{F(z, w)}{X(z, w)^{n+1}} + G(z, w) \log X(z, w), \tag{3-6}$$

where $G, F \in C^\infty(\bar{\Omega} \times \bar{\Omega})$, $F(z, z) > 0$ on $(\bar{\Omega} \times \bar{\Omega}) \cap R_\epsilon$ and “log” denotes the principal branch of the logarithm defined on $\mathbb{C} \setminus (-\infty, 0]$. The asymptotic expansion (3-6) implies

$$\int_{\Omega} |K(z, w)| dv_w \leq C(1 + \log K(z)), \quad z \in \Omega. \tag{3-7}$$

Since the boundary $\partial\Omega$ is compact, for any $\delta > 0$ there are finitely many $b^j \in \partial\Omega$, $j = 1, \dots, m$, such that $\partial\Omega \subset \cup_{j=1}^m \mathbb{B}(b^j, \delta)$. Choose smoothly bounded strictly pseudoconvex domains Ω^j , $j = 1, \dots, m$, such that

$$\mathbb{B}(b^j, 3\delta) \cap \Omega \subset \Omega^j \subset \Omega \cap \mathbb{B}(b^j, 4\delta),$$

where δ is chosen small enough such that for each j , after a polynomial change of variables, each Ω^j is a strictly convex domain. Let $\{\Omega^j\}_{j=m+1}^{m+k}$ be a finite open cover of $\Omega \setminus \cup_{j=1}^m (\mathbb{B}(b^j, \delta) \cap \Omega)$ consisting of balls contained in Ω . In the argument of [Theorem 3.3](#) by letting $\phi_0 = \gamma \log K_\Omega(z)$ (instead of $\gamma \log K_{\Omega^j}(z)$) and using [\(3-7\)](#), we can solve the equation $\bar{\partial}u^j = f$ on Ω^j with minimal solution u^j satisfying

$$|u^j(z)| \leq C \|f\|_{g,\infty} \log(1 + K(z)). \tag{3-8}$$

Let $\{\eta_j\}_{j=1}^{m+k}$ be a partition of unity of $\bar{\Omega}$ subordinate to the cover $\{B(b^j, \delta)\}_{j=1}^m \cup \{\Omega^j\}_{j=m+1}^{m+k}$, and let $v(z) := \sum_{j=1}^{m+k} \eta_j(z)u^j(z)$. Then $\bar{\partial}v = f + h$, where $h := \sum_{j=1}^{m+k} u^j \bar{\partial}\eta_j$ is a $\bar{\partial}$ -closed $(0, 1)$ -form on Ω . By the integral formula in [\[Grauert and Lieb 1970; Henkin 1970\]](#), there is a bounded solution v_0 to the equation $\bar{\partial}v_0 = h$. Let $u = v - v_0$. Then $\bar{\partial}u = f$ and by [\(3-8\)](#),

$$|u(z)| \leq \sum_{j=1}^k \eta_j(z) C \|f\|_{g,\infty} \log(1 + K(z)) + C \leq C \|f\|_{g,\infty} \log(1 + K(z)). \quad \square$$

Remark. For a smoothly bounded strictly pseudoconvex domain Ω , if the canonical solution is u_0 , then for $h \in L^\infty(\Omega)$ with $\|h\|_\infty \leq 1$,

$$|P[h(\cdot) \log K(\cdot)](z)| \leq C(1 + \log K(z))^2.$$

In fact, letting $\omega_t = \{z \in \Omega : \delta(z) > t\}$, by Fefferman’s expansion theorem on the Bergman kernel [\[Fefferman 1974\]](#), we know that

$$\begin{aligned} |P[h(\cdot) \log K(\cdot)](z)| &\leq \int_\Omega |h(w)| \log K(w) |K(z, w)| dv(w) \leq \|h\|_\infty \int_\Omega \log K(w) |K(z, w)| dv(w) \\ &\approx \|h\|_\infty \int_0^c \int_{\partial\Omega_t} \log K(w) |K(z, w)| d\sigma(w) dt \\ &\leq C \|h\|_\infty \int_0^c (-\log t) \frac{1}{\delta(z)+t} dt \\ &\leq C \|h\|_\infty \left(\frac{1}{\delta(z)} \int_0^{\delta(z)} (-\log t) dt + \int_{\delta(z)}^c -\log t \frac{1}{\delta(z)+t} dt \right) \\ &\leq C \|h\|_\infty \left(\log \frac{C}{\delta(z)} + \int_{\delta(z)}^c -\log(\delta(z)+t) \frac{1}{\delta(z)+t} dt \right) \leq C \|h\|_\infty \left(\log \frac{C}{\delta(z)} \right)^2. \end{aligned}$$

Combining this with [Theorem 3.4](#) one gets

$$|u_0(z)| \leq C(1 + \log K(z))^2.$$

4. Uniform estimates

In this section, we obtain uniform estimates for the equation $\bar{\partial}u = f$ on the unit polydisc \mathbb{D}^n and strictly pseudoconvex domains by imposing conditions on f stronger than $\|f\|_{g,\infty} < \infty$.

Theorem 4.1. *For any $p \in (1, \infty)$, there is a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f on \mathbb{D}^n , the canonical solution u to $\bar{\partial}u = f$ satisfies*

$$\|u\|_\infty \leq C \left\| f \prod_{j=1}^n \left(\log \left(\frac{2}{1-|z_j|^2} \right) \right)^p \right\|_{g,\infty}.$$

Proof. Let $A_0 = 2p + \log v(\mathbb{D}^n)$. Choose $0 < \gamma < \frac{1}{2}$ such that $\phi(z) := \gamma \log K(z)$ satisfies [Lemma 2.4](#). Choose $\alpha > 1$ such that $\alpha - \gamma = 1$. Let $K_j(z) := \pi^{-1}(1 - |z_j|^2)^{-2}$, and let

$$\phi_0(z) := \phi(z) - \sum_{j=1}^n p \log(A_0 + \gamma \log K_j(z)) - \alpha \log |K(z, a)|.$$

Since on \mathbb{D}^n , $\log |K(z, a)|$ is pluriharmonic and

$$A_0 + \gamma \log K_j = 2p + n \log \pi - \gamma \log \pi - 2\gamma \log(1 - |z_j|^2) \geq 2p,$$

we know that

$$i\bar{\partial}\bar{\partial}\phi = \sum_{j=1}^n \left[i\bar{\partial}\bar{\partial}(\gamma \log K_j) \left(1 - \frac{p}{A_0 + \gamma \log K_j} \right) + \pi \frac{\bar{\partial}(\gamma \log K_j) \wedge \bar{\partial}(\gamma \log K_j)}{(A_0 + \gamma \log K_j)^2} \right] \geq \frac{1}{2} \sum_{j=1}^n i\bar{\partial}\bar{\partial}(\gamma \log K_j).$$

Thus $|f|_{i\bar{\partial}\bar{\partial}\phi_0}^2 \leq 2|f|_{i\bar{\partial}\bar{\partial}\phi}^2 = 2|f|_g^2/\gamma$. Let

$$A_p(f) = \left\| f \left(\prod_{j=1}^n (A_0 - \gamma \log \pi - 2\gamma \log(1 - |z_j|^2)) \right)^p \right\|_{g,\infty}.$$

As in [Theorem 3.3](#), let u_0 be the $L^2(\mathbb{D}^n, \phi_0)$ minimal solution to $\bar{\partial}u_0 = f$ and $\beta := e^{i\theta(z)}\chi_{B_a(\epsilon)}$, where $u(z) = |u(z)|e^{i\theta(z)}$. Then

$$\begin{aligned} \int_{B_a(\epsilon)} |u| dv &\leq C v(B_a(\epsilon)) \left(\int_{D(0,1)^n} |f|_g^2 e^{-\phi_0} dv \right)^{1/2} \left(\int_{D(0,1)^n} |K(z, a)|^2 e^{\phi_0(z)} dv_z \right)^{1/2} \\ &\leq C A_p(f) v(B_a(\epsilon)) \left(\int_{D(0,1)^n} \frac{e^{-\phi_0} dv_z}{\prod_{j=1}^n (A_0 + \gamma \log K_j(z))^{2p}} \int_{D(0,1)^n} |K(z, a)|^2 e^{\phi_0} dv_z \right)^{1/2} \\ &= C A_p(f) v(B_a(\epsilon)) \prod_{j=1}^n \left(\int_{D(0,1)} \frac{|K_j(z, a)|^\alpha K_j(z)^{-\gamma} dv_z}{(A_0 + \gamma \log K_j(z))^p} \int_{D(0,1)} \frac{|K_j(z, a)|^{2-\alpha} K_j(z)^\gamma dv_z}{(A_0 + \gamma \log K_j(z))^p} \right)^{1/2} \\ &\leq C A_p(f) v(B_a(\epsilon)). \end{aligned}$$

Fix $\epsilon > 0$ sufficiently small. By [Proposition 2.1](#),

$$\begin{aligned} |u(a)| &\leq CA_p(f) + C\|f\|_{g,\infty} \\ &\leq C\left\|f\prod_{j=1}^n(2(n+p)-\log(1-|z_j|^2))^p\right\|_{g,\infty}. \end{aligned}$$

Notice that

$$2(n+p)-\log(1-|z_j|^2)\leq 5(n+p)\log\frac{2}{1-|z_j|^2},$$

which completes the proof of the theorem. □

In fact, the finiteness of the right-hand side of the estimate in [Theorem 4.1](#) is a stronger condition than f being L^∞ on \mathbb{D}^n . Recently, in [\[Dong et al. 2020\]](#), the first author, Pan and Zhang obtained uniform estimates for the canonical solution to $\bar{\partial}u = f$ when f is continuous up to the boundary of \mathbb{D}^n , and more generally the Cartesian product of smoothly bounded planar domains.

However, the situation for strictly pseudoconvex domains is quite different. The finiteness of the right-hand side of the estimate in either [Theorem 4.2](#) or [Theorem 4.3](#) is a much weaker condition than f being L^∞ on each smooth domain considered. In fact, we allowed f to blow-up on $\partial\Omega$ to order less than $\frac{1}{2}$.

Theorem 4.2. *Let Ω be a smoothly bounded strictly convex domain. For any $p \in (1, \infty)$ and sufficiently small $\gamma > 0$, there exists a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f , the canonical solution u to $\bar{\partial}u = f$ satisfies*

$$\|u\|_\infty \leq C\|(1 + \log v(\Omega) + \gamma \log K(z))^p f\|_{g,\infty}.$$

Proof. Choose $0 < \gamma < 1/(n+2)$ such that $\phi(z) := \gamma \log K(z)$ satisfies [Lemma 2.4](#), and let $\alpha = \gamma + 1$. Let

$$A_0 := 2p + \gamma \log v(\Omega),$$

and let

$$\phi_0(z) = \phi(z) - (n+1)\alpha \log|h_a(z)| - p \log(A_0 + \gamma \log K(z)).$$

Notice that

$$i\partial\bar{\partial}\phi_0 \geq \left(1 - \frac{p}{A_0 + \phi}\right)i\partial\bar{\partial}\phi \geq \frac{i\partial\bar{\partial}\phi}{2}$$

and

$$\gamma K(z) > \gamma v(\Omega)^{-1}.$$

Therefore

$$|f|_{i\partial\bar{\partial}\phi_0}^2 \leq \frac{2}{\gamma}|f|_g^2.$$

Define

$$A_p(f) := \|(A_0 + \gamma \log K(z))^p f\|_{g,\infty}.$$

Using arguments similar to those in Theorems 3.3 and 4.1 and $\alpha - \gamma = 1$,

$$\begin{aligned} & \int_{B_a(\epsilon)} |u| dv \\ & \leq C v(B_a(\epsilon)) \left(\int_{\Omega} |f|_{i\bar{\partial}\bar{\partial}\phi}^2 e^{-\phi_0} dv \right)^{1/2} \left(\int_{\Omega} |K(z, a)|^2 e^{\phi_0(z)} dv_z \right)^{1/2} \\ & \leq C A_p(f) v(B_a(\epsilon)) \left(\int_{\Omega} \frac{e^{-\phi_0}}{(A_0 + \gamma \log K(z))^{2p}} dv_z \int_{\Omega} |K(z, a)|^2 e^{\phi_0} dv_z \right)^{1/2} \\ & = C A_p(f) v(B_a(\epsilon)) \left(\int_{\Omega} \frac{K(z)^{-\gamma} |h_a(z)|^{\alpha(n+1)}}{(A_0 + \gamma \log K(z))^p} dv_z \int_{\Omega} \frac{|K(z, a)|^2 K(z)^{\gamma} |h_a(z)|^{-\alpha(n+1)}}{(A_0 + \gamma \log K(z))^p} dv_z \right)^{1/2} \\ & \leq C A_p(f) v(B_a(\epsilon)) \left(\int_{\Omega} \frac{|K(z, a)| dv_z}{(A_0 + \gamma \log K(z))^p} \int_{\Omega} \frac{|K(z, a)|^{2-\alpha} K(z)^{\gamma}}{(A_0 + \gamma \log K(z))^p} dv_z \right)^{1/2} \leq C A_p(f) v(B_a(\epsilon)), \end{aligned}$$

where the last inequality follows from Fefferman’s asymptotic expansion. In fact, since $n/(n+1) < 2 - \alpha$, if $\Omega_t = \{z : r(z) > t\}$ where $r(z)$ is a defining function satisfying the definition of $h_a(z)$, then

$$\begin{aligned} \int_{\Omega} \frac{|K(z, a)|^{2-\alpha} K(z)^{\gamma}}{(A_0 + \gamma \log K(z))^p} dv_z & \leq C \left(1 + \int_0^{\epsilon} \int_{\partial\Omega_t} \frac{|K(z, a)|^{2-\alpha} K(z)^{\gamma}}{(A_0 + \gamma \log K(z))^p} d\sigma_t(z) dt \right) \\ & \leq C \left(1 + \int_0^{\epsilon} \frac{t^{-(2-\alpha)(n+1)+n} t^{-\gamma(n+1)}}{(A_0 - \gamma(n+1) \log t)^p} dt \right) \\ & \leq C \left(1 + \int_0^{\epsilon} \frac{1}{t(\log t)^p} dt \right) \leq C \left(1 - \frac{\log \epsilon}{p-1} \right). \end{aligned}$$

By Proposition 2.1, for a fixed $\epsilon > 0$ sufficiently small, $|u(a)| \leq C A_p(f) + \|f\|_{g,\infty} \leq C A_p(f)$. □

Using an argument similar to the proof of Theorem 3.4 we get the following generalization of Theorem 4.2 to smoothly bounded strictly pseudoconvex domains.

Theorem 4.3. *Let Ω be a smoothly bounded strictly pseudoconvex domain. Then, for any $p \in (1, \infty)$, there exists a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f , there is a solution u to $\bar{\partial}u = f$ that satisfies*

$$\|u\|_{\infty} \leq C \|(\log K(\cdot))^p f(\cdot)\|_{g,\infty}.$$

Remark. Let $f \in L^{\infty}_{(0,1)}(\Omega)$ be a $\bar{\partial}$ -closed form on a smoothly bounded strictly pseudoconvex domain Ω . Henkin and Romanov’s theorem [1971] states that there exists a solution $u \in C^{1/2}(\Omega)$. Theorem 3.4 implies that one can find a bounded solution when $(\log(1/\delta(z)))^p \delta(z) |f(z)|^2$ is bounded. Moreover, [Lieb and Range 1986, Theorem 2 (i)] shows that uniform estimates hold for the canonical solutions to the $\bar{\partial}$ -equations on Ω .

5. Additional estimates for Cartan classical domains

A domain Ω is symmetric if, for all $a \in \Omega$, there is an involutive automorphism G such that a is isolated in the set of fixed points of G . All bounded symmetric domains are convex and homogeneous. E. Cartan

classical domain	rank r	multiplicity \mathbf{a}	genus p	dimension N	index k
I(m, n), $m \leq n$	m	2	$m+n$	mn	1
II(n)	n	1	$n+1$	$\frac{1}{2}n(n+1)$	1
III($2n+\epsilon$), $\epsilon = 0$ or 1	n	4	$2(2n+\epsilon-1)$	$n(2n+2\epsilon-1)$	$\frac{1}{2}$
IV(n)	2	$n-2$	n	n	1

Table 1. Characteristics of classical domains.

proved that all bounded symmetric domains in \mathbb{C}^N up to biholomorphism are the Cartesian product(s) of the following four types of Cartan classical domains and two domains of exceptional types.

Definition 5.1. A Cartan classical domain is a domain of one of the following types:

- (i) $I(m, n) := \{z \in M_{(m,n)}(\mathbb{C}) : I_m - zz^* > 0\}$, $m \leq n$.
- (ii) $II(n) := \{z \in I(n, n) : z^\tau = z\}$.
- (iii) $III(n) := \{z \in I(n, n) : z^\tau = -z\}$.
- (iv) $IV(n) := \{z \in \mathbb{C}^n : 1 - 2|z|^2 + |s(z)|^2 > 0 \text{ and } |s(z)| < 1\}$, where $s(z) := \sum_{j=1}^n z_j^2$ and $n > 2$.

Here $z^* := \bar{z}^\tau$ is the conjugate transpose of z .

Let Ω be a Cartan classical domain. Denote the rank, characteristic multiplicity, genus, complex dimension and kernel index of Ω by r, \mathbf{a}, p, N and k , respectively. Their values are given in [Table 1](#).

Hua [[1963](#)] obtained explicit formulas for the Bergman kernels on the Cartan classical domains. For a domain Ω of type I, II or III,

$$K(z, w) = C_\Omega [\det(I - zw^*)]^{-pk},$$

and for a domain of type IV,

$$K(z, w) = C_n \left[1 - 2 \sum_{j=1}^n z_j \bar{w}_j + s(z) \overline{s(w)} \right]^{-n}.$$

For any $z \in \Omega$,

$$\delta_\Omega(z) \leq K(z)^{-1/(rpk)}.$$

Let $\lambda = pk$. By [[Faraut and Korányi 1990](#), Theorem 3.8], one can write the Bergman kernel on a Cartan classical domain Ω as

$$K(z, w) = h(z, w)^{-\lambda} = \sum_{\mathbf{m} \geq 0} (\lambda)_{\mathbf{m}} K_{\mathbf{m}}(z, w),$$

where

$$\mathbf{m} = (m_1, \dots, m_r) \text{ and } \mathbf{m} \geq 0 \iff m_1 \geq m_2 \geq \dots \geq m_r \geq 0,$$

and

$$(\lambda)_{\mathbf{m}} = \frac{\Gamma_\Omega(\lambda + \mathbf{m})}{\Gamma_\Omega(\lambda)}, \quad \Gamma_\Omega(s) = c_\Omega \prod_{j=1}^r \Gamma\left(s_j - (j-1)\frac{\mathbf{a}}{2}\right), \quad \lambda = (\lambda, \dots, \lambda).$$

Here, K_m is the Bergman kernel for homogeneous polynomials in \mathbb{C}^r of degree $|m| = m_1 + \cdots + m_r$. For each Cartan domain Ω , there is a subgroup $\mathcal{K}(\Omega)$ of the unitary group such that for each $z \in \Omega$ there is $k \in \mathcal{K}(\Omega)$ such that $z = k\tilde{z}$ where $\tilde{z} \in \mathbb{C}^r \times \prod_{j=r+1}^N \{0\}$ and $K_m(z, z) =: K_m(\tilde{z}, \tilde{z})$.

The following Forelli–Rudin-type integral was studied in [Faraut and Korányi 1990]:

$$J_{\beta,c}(z) := \int_{\Omega} K(w)^{\beta} |K(w, z)|^{1+c-\beta} dv_w.$$

By the proof of Theorem 4.1 in [Faraut and Korányi 1990], one has

$$J_{\beta,c}(z) = \sum_{m \geq 0} \frac{|(\mu)_m|^2}{((1-k\beta)p)_m} K_m(z, z), \quad \mu = \frac{1}{2}kp(1+c-\beta).$$

Using Stirling’s formula, one can show (see [Faraut and Korányi 1990, (4.3)] or [Engliš and Zhang 2004, (2.9)]) that as m varies,

$$\frac{|\left(\frac{1}{2}pk(1-\beta)\right)_m|^2}{((1-k\beta)p)_m} \approx \frac{\left(\frac{1}{2}kp\right)_m^2}{(p)_m},$$

which implies that

$$J_{\beta,0}(z) \approx J_{0,0}(z) = \int_{\Omega} |K(z, w)| dv(w), \quad \beta < \frac{1}{pk}. \quad (5-1)$$

Further computations were carried out by Faraut and Korányi [1990].

Theorem 5.2 [Faraut and Korányi 1990]. *For any $\beta < 1/(pk)$,*

- (i) $J_{\beta,c}(z)$ is bounded for all $z \in \Omega$ if and only if $c < -(r-1)\mathbf{a}/(2p)$,
- (ii) $J_{\beta,c}(z) \approx K(z)^c$ if $c > (r-1)\mathbf{a}/(2p)$.

When $|c| \leq (r-1)\mathbf{a}/(2p)$, it is difficult to compute $J_{\beta,c}(z)$; see [Korányi 1991; Yan 1992]. Theorem 1 of [Engliš and Zhang 2004], whose parameters are chosen as $\frac{1}{2}p(1+c-\beta)$, $\frac{1}{2}p(1+c-\beta)$ and $p(1-\beta)$, is stated as follows.

Theorem 5.3. *Let Ω be a Cartan classical domain of rank 2 with characteristic boundary \mathcal{U} . Then for any $z = te_1 + Te_2$ with $0 \leq t \leq T < 1$ and $e_1, e_2 \in \mathcal{U}$ the following statements hold:*

- (i) *If $2pc = \mathbf{a}$, then $J_{\beta,c}(z) \approx (1-t)^{-\mathbf{a}/2}(1-T)^{-\mathbf{a}/2}[1 - \log(1-t)]$.*
- (ii) *If $0 < 2pc < \mathbf{a}$, then $J_{\beta,c} \approx (1-t)^{-\mathbf{a}/2}(1-T)^{-pc}$.*
- (iii) *If $c = 0$, then $J_{\beta,c}(z) \approx (1-t)^{-\mathbf{a}/2}[1 + \log[(1-t)/(1-T)]]$.*
- (iv) *If $-\mathbf{a} < 2pc < 0$, then $J_{\beta,c}(z) \approx (1-t)^{-pc-\mathbf{a}/2}$.*
- (v) *If $2pc = -\mathbf{a}$, then $J_{\beta,c}(z) \approx 1 - \log(1-t)$.*

As a consequence, when Ω is a Cartan classical domain of rank 2 and $z = te_1 + te_2$ with $0 \leq t < 1$ and $e_i \in \mathcal{U}$, one has

$$\int_{\Omega} |K(z, w)| dv_w \approx (1-t)^{-\mathbf{a}/2} \approx \delta_{\Omega}(z)^{-\mathbf{a}/2}.$$

On the Cartan classical domains, we impose a stronger assumption on f to get bounded solutions to $\bar{\partial}u = f$. The following result provides a partial answer to the problems raised by Henkin and Leiterer [1983] and Sergeev [1994].

Theorem 5.4. *Let Ω be a Cartan classical domain and $\alpha > 1 + (r - 1)\mathbf{a}/(2p)$. Then there exists a constant C such that for any $\bar{\partial}$ -closed $(0, 1)$ -form f , the canonical solution u to $\bar{\partial}u = f$ satisfies*

$$\|u\|_\infty \leq C \left\| \int_\Omega |f|_g^2(z) |K(z, \cdot)|^\alpha dv_z \right\|_\infty^{1/2} + C \|f\|_{g,\infty}. \tag{5-2}$$

Proof. As in the proof of Theorem 3.3, for any $a \in \Omega$, let $\beta := \chi_{B_a(\epsilon)} u(z)/|u(z)|$, $\phi := \gamma \log K(z)$ and $\psi_a(z) := -\alpha \log |K(z, a)|$. Then

$$\int_{B_a(\epsilon)} |u| dv \leq \int_\Omega |u_0 \bar{\beta}| dv + \int_\Omega |u_0 \overline{P(\beta)}| dv.$$

By Lemma 2.4 and (3-4),

$$\begin{aligned} \left| \int_\Omega u_0 \bar{\beta} dv \right| &\leq C \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2} \left(\int_{B_a(\epsilon)} |K(z, a)|^{-\alpha} dv_z \right)^{1/2} \\ &\leq C \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2} (v(B_a(\epsilon)) K(a)^{-\alpha})^{1/2} \\ &\leq C v(B_a(\epsilon))^{(1+\alpha)/2} \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2}. \end{aligned}$$

On the other hand, by (3-5),

$$\begin{aligned} \left| \int_\Omega u_0 \overline{P(\beta)} dv \right| &\leq C \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2} v(B_a(\epsilon)) \left(\int_\Omega \max_{w \in B_a(\epsilon)} |K(z, w)|^2 e^{\psi_a} dv_z \right)^{1/2} \\ &\leq C \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2} v(B_a(\epsilon)) \left(\int_\Omega |K(z, a)|^{2-\alpha} dv_z \right)^{1/2}. \end{aligned}$$

If $\alpha > 1 + (r - 1)\mathbf{a}/(2p) \geq 1$, then $|K(z, a)|^{2-\alpha}$ is integrable on Ω by Theorem 5.2. Therefore, for any $a \in \Omega$,

$$\frac{1}{v(B_a(\epsilon))} \int_{B_a(\epsilon)} |u| dv \leq C \left(\int_\Omega |f|_g^2(z) |K(z, a)|^\alpha dv_z \right)^{1/2}.$$

Coupling this estimate with Proposition 2.1, we have proved u is bounded. □

6. Sharpness of the pointwise estimates

For the Cartan classical domains, we show that the logarithm of the Bergman kernel has a bounded gradient with respect to the Bergman metric and also verify that Theorem 3.3 is sharp.

6A. Solutions with logarithmic growth.

Example 1. Let Ω be a Cartan classical domain and $u(z) = \log K(z)$. Then $P[u](z)$ is a constant function on Ω and there exists a constant c such that $|\bar{\partial}u|_g^2 = c \operatorname{Tr}(zz^*)$.

Proof. Notice that for all $z \in \Omega$,

$$\begin{aligned} P[u](z) &= \int_{\Omega} u(w) K(z, w) dv_w \\ &= \int_{\Omega} \frac{1}{2\pi} \int_0^{2\pi} u(e^{i\theta} w) K(z, e^{i\theta} w) d\theta dv_w \\ &= \int_{\Omega} \frac{1}{2\pi} \int_0^{2\pi} u(w) K(z, e^{i\theta} w) d\theta dv_w \\ &= \int_{\Omega} u(w) K(z, 0) dv_w = \frac{1}{v(\Omega)} \int_{\Omega} u(w) dv_w, \end{aligned}$$

where the third equality follows by the transformation rule of the Bergman kernel, and the fourth equality follows by the mean value property of (anti)holomorphic functions.

Now we show the second part of the example. For $z \in M_{(m,n)}(\mathbb{C})$, define $V(z) := I_m - zz^*$ and let V_{uv} denote the (u, v) entry of V . Then by [Hua 1963; Lu 1997] (see also [Chen and Li 2019, Proposition 2.1]), for domains of type I, II and III,

$$g^{j\alpha, \bar{k}\beta}(z) = \begin{cases} V_{jk}(\delta_{\alpha\beta} - \sum_{l=1}^m z_{l\alpha} \bar{z}_{l\beta}), & z \in \text{I}(m, n), \\ V_{jk} \frac{V_{\alpha\beta}}{(2 - \delta_{j\alpha})(2 - \delta_{k\beta})}, & z \in \text{II}(n), \\ \frac{1}{4} V_{jk} V_{\alpha\beta} (1 - \delta_{j\alpha})(1 - \delta_{k\beta}), & z \in \text{III}(n). \end{cases}$$

For matrices $E_{j\alpha} := (\delta_{ju} \delta_{\alpha v})_{u,v}$, $A := (a_{uv})_{u,v} \in M_{(n,m)}(\mathbb{C})$,

$$E_{j\alpha} A = (\delta_{ju} a_{\alpha v})_{u,v} \quad \text{and} \quad \frac{\partial V}{\partial z_{j\alpha}} = -E_{j\alpha} z^*.$$

Then for $z \in \text{I}(m, n)$,

$$\begin{aligned} \frac{\partial \log \det V(z)}{\partial z_{j\alpha}} &= \operatorname{Tr} \left(V^{-1}(z) \frac{\partial V(z)}{\partial z_{j\alpha}} \right) = -\operatorname{Tr} (V^{-1}(z) E_{j\alpha} z^*) = -\operatorname{Tr} (E_{j\alpha} z^* V^{-1}(z)) \\ &= -\sum_u \delta_{ju} [z^* V^{-1}]_{\alpha u} = -[z^* V^{-1}]_{\alpha j}. \end{aligned}$$

Since $u(z) = \log(\det(V(z)))^{-(m+n)} - \log v(\text{I}(m, n))$ is real-valued,

$$\begin{aligned} |\bar{\partial}u|_g^2(z) &= \sum_{j,\beta,k,\alpha} g^{j\alpha, \bar{k}\beta} \frac{\partial u}{\partial z_{j\alpha}} \frac{\bar{\partial}u}{\partial z_{k\beta}} = (m+n)^2 \sum_{j,\beta,k,\alpha} V_{jk} [I - z^\tau \bar{z}]_{\alpha\beta} [z^* V^{-1}(z)]_{\alpha j} [\bar{z}^* \bar{V}^{-1}]_{\beta k} \\ &= (m+n)^2 \sum_{\alpha,k} [z^*]_{\alpha k} [(I - z^\tau \bar{z}) z^\tau \bar{V}^{-1}]_{\alpha k} \\ &= (m+n)^2 \sum_{\alpha,k} [z^*]_{\alpha k} [z^\tau]_{\alpha k} = (m+n)^2 \operatorname{Tr}(zz^*). \end{aligned}$$

For $z \in \text{II}(n)$, using the symmetry of z , we know

$$\frac{\partial V(z)}{\partial z_{j\alpha}} = -(1 - \frac{1}{2}\delta_{j\alpha})(E_{j\alpha} + E_{\alpha j})z^* \quad \text{and} \quad z^*V^{-1}(z) = (z^*V^{-1}(z))^{\tau}.$$

Hence,

$$\begin{aligned} \frac{\partial \log \det V(z)}{\partial z_{j\alpha}} &= \text{Tr}\left(V^{-1}(z) \frac{\partial V}{\partial z_{j\alpha}}(z)\right) = -(1 - \frac{1}{2}\delta_{j\alpha}) \text{Tr}(E_{j\alpha}z^*V^{-1}(z) + z^*V^{-1}(z)E_{\alpha j}) \\ &= -(2 - \delta_{j\alpha}) \text{Tr}(E_{j\alpha}z^*V^{-1}(z)) = -(2 - \delta_{j\alpha})[z^*V^{-1}(z)]_{j\alpha}. \end{aligned}$$

Since $u(z) = \log(\det(V(z)))^{-(n+1)} - \log v(\text{II}(n))$ and for z symmetric $\overline{z^*V(z)^{-1}} = V(z)^{-1}z$,

$$\begin{aligned} |\bar{\partial}u|_g^2(z) &= \sum_{j,\beta,k,\alpha} g^{j\alpha,\bar{k}\beta} \frac{\partial u}{\partial z_{j\alpha}} \frac{\bar{\partial}u}{\partial \bar{z}_{k\beta}} = (n+1)^2 \sum_{\alpha,\beta,k,j} V_{jk}V_{\alpha\beta}[z^*V^{-1}(z)]_{j\alpha}[\overline{z^*V^{-1}(z)}]_{k\beta} \\ &= (n+1)^2 \sum_{j,\beta} [z^*]_{j\beta}[V(z)V^{-1}(z)z]_{j\beta} = (n+1)^2 \text{Tr}(zz^*). \end{aligned}$$

The proof for skew-symmetric $z \in \text{III}(n)$ is similar to the preceding proofs.

For a Cartan classical domain $\text{IV}(n)$, let $s(z) := \sum z_j^2$ and $r(z) := 1 - 2|z|^2 + |s(z)|^2$ for $z \in \mathbb{C}^n$. By [Hua 1963], the Bergman kernel $K(z, z)$ equals $cr(z)^{-n}$. Also,

$$g^{j,\bar{k}}(z) = r(z)(\delta_{jk} - 2z_j\bar{z}_k) + 2(\bar{z}_j - \overline{s(z)}z_j)(z_k - s(z)\bar{z}_k).$$

Notice that

$$(\log(r(z)^{-n}))_{z_j}(\log(r(z)^{-n}))_{\bar{z}_k} = \frac{4n^2}{r(z)^2}[z_j s(\bar{z}) - \bar{z}_j][\bar{z}_k s(z) - z_k]$$

and

$$\begin{aligned} |\bar{\partial}u|_g^2(z) &= 4n^2 \sum_{j,k=1}^n [r(z)(\delta_{jk} - 2z_j\bar{z}_k) + 2(\bar{z}_j - \overline{s(z)}z_j)(z_k - s(z)\bar{z}_k)] \frac{(\bar{z}_j - \overline{s(z)}z_j)(z_k - s(z)\bar{z}_k)}{r(z)^2} \\ &= \sum_{j,k=1}^n \frac{4n^2}{r(z)} (\delta_{jk} - 2z_j\bar{z}_k)[\bar{z}_j - \overline{s(z)}z_j][z_k - s(z)\bar{z}_k] + \sum_{j,k=1}^n \frac{8n^2(\bar{z}_j - \overline{s(z)}z_j)^2(z_k - s(z)\bar{z}_k)^2}{r(z)^2} \\ &=: F(z) + G(z). \end{aligned}$$

Thus,

$$\begin{aligned} F(z) \frac{r}{4n^2} &= \sum_{j=1}^n |z_j|^2 - s\bar{z}_j^2 - z_j^2\bar{s} + |s|^2|z_j|^2 - 2 \sum_{j,k=1}^n |z_j|^2|z_k|^2 - s|z_j|^2\bar{z}_k^2 - z_j^2|z_k|^2\bar{s} + |s|^2z_j^2\bar{z}_k^2 \\ &= |z|^2 - 2|s|^2 + |s|^2|z|^2 - 2(|z|^4 - s|z|^2\bar{s} - s|z|^2\bar{s} + |s|^2s\bar{s}) \\ &= -2|z|^4 + 5|s|^2|z|^2 - 2|s|^2 + |z|^2 - 2|s|^4 \end{aligned}$$

and

$$G(z) = \frac{8n^2}{r^2} \left| \sum_{j=1}^n (z_j - s\bar{z}_j)^2 \right|^2 = \frac{8n^2}{r^2} |s - 2s|z|^2 + s^2\bar{s}|^2 = 8n^2|s|^2.$$

Therefore

$$\begin{aligned} |\bar{\partial}u|_g^2 &= \frac{4n^2}{r}[-2|z|^4 + 5|s|^2|z|^2 - 2|s|^2 + |z|^2 - 2|s|^4] + \frac{4n^2}{r}2|s(z)|^2r(z) \\ &= \frac{4n^2}{r}[-2|z|^4 + |z|^2|s|^2 + |z|^2] \\ &= 4n^2|z|^2 = 4n^2 \operatorname{Tr}(zz^*). \end{aligned} \quad \square$$

Example 2 shows that the canonical solution to the equation $\bar{\partial}u = f := \bar{\partial} \log K(z)$ (here $\|f\|_{g,\infty} < \infty$) given by $\log K(z) - C_\Omega$ is unbounded with logarithmic growth near the boundary of the polydisc.

Example 2. Consider $f(z) := -\sum_{j=1}^n z_j(1 - |z_j|^2)^{-1} d\bar{z}_j$ defined on \mathbb{D}^n . Then f is $\bar{\partial}$ -closed, $\|f\|_{g,\infty} \leq \frac{1}{2}$ and the canonical solution to $\bar{\partial}u = f$ on \mathbb{D}^n is

$$u(z) := \sum_{j=1}^n \log(1 - |z_j|^2) - n \int_0^1 \log(1 - r) dr. \tag{6-1}$$

Proof. We compute directly that u given by (6-1) satisfies $\bar{\partial}u = f$, and that

$$|f(z)|_g^2 = \frac{1}{2} \sum_{j=1}^n \frac{(1 - |z_j|^2)^2}{(1 - |z_j|^2)^2} |z_j|^2 = \frac{|z|^2}{2}.$$

To verify that u is canonical, notice that

$$\begin{aligned} P_{\mathbb{D}^n} \left[\sum_{j=1}^n \log(1 - |w_j|^2) \right] (z) &= \frac{1}{\pi^n} \int_{\mathbb{D}^n} \prod_{j=1}^n \frac{1}{(1 - \langle z_j, w_j \rangle)^2} \sum_{k=1}^n \log(1 - |w_k|^2) dv_{w_1} \cdots dv_{w_n} \\ &= \sum_{k=1}^n \frac{1}{\pi} \int_{\mathbb{D}^n} \frac{\log(1 - |w_k|^2)}{(1 - \langle z_k, w_k \rangle)^2} dv_{w_k} \\ &= \sum_{k=1}^n 2 \int_0^1 \log(1 - r_k^2) r_k dr_k = n \int_0^1 \log(1 - r) dr. \end{aligned} \quad \square$$

6B. A sharp example. The maximum blow-up order for a solution to $\bar{\partial}u = f$ with $\|f\|_{g,\infty} < \infty$ is $\int_\Omega |K(\cdot, w)| dv_w$. Here we provide an example to show that [Theorem 3.3](#) is sharp on the Cartan classical domains.

Proposition 6.1. *Let Ω be a Cartan classical domain. Then there is a constant c such that for each $z \in \Omega$, there is a $\bar{\partial}$ -closed $(0, 1)$ -form f_z on Ω with $\|f_z\|_{g,\infty} = 1$ and the canonical solution to $\bar{\partial}u = f_z$ satisfies*

$$|u(z)| \geq c \int_\Omega |K(z, w)| dv_w.$$

Proof. For any point $z \in \Omega$, consider the functions $U_z(\cdot) := K(\cdot)^{-1}K(\cdot, z)$ and

$$f_z(\cdot) := \bar{\partial}U_z(\cdot) = K(\cdot, z)\bar{\partial}(K(\cdot)^{-1}).$$

Then, by [Example 1](#),

$$\begin{aligned} \|f_z\|_{g,\infty} &= \|K(\cdot, z)K(\cdot)^{-2}\bar{\partial}(K(\cdot))\|_{g,\infty} = \|K(\cdot, z)K(\cdot)^{-1}\bar{\partial}(\log K(\cdot))\|_{g,\infty} \\ &\leq \|K(\cdot, z)K(\cdot)^{-1}\|_\infty \|\bar{\partial}(\log K(\cdot))\|_{g,\infty} \leq C. \end{aligned}$$

The Bergman projection of U_z is

$$P[U_z](\cdot) = \int_\Omega U_z(w)K(\cdot, w) dv_w = \int_\Omega K(w)^{-1}K(w, z)K(\cdot, w) dv_w.$$

In particular, by [\(5-1\)](#) with $\beta = -1$,

$$P[U_z](z) = \int_\Omega K(w)^{-1}K(w, z)K(z, w) dv_w = \int_\Omega K(w)^{-1}|K(w, z)|^2 dv_w \approx \int_\Omega |K(z, w)| dv_w.$$

The canonical solution to $\bar{\partial}u = f$ is $u_z := U_z - P[U_z]$ and

$$|u_z(z)| = |1 - J_{-1,0}(z)| \geq c \int_\Omega |K(z, w)| dv_w - 1$$

for a uniform constant $c > 0$, independent of z . □

6C. Blow-up order greater than log. With the previous example and [Theorem 5.3](#) we will provide the maximum blow-up order when Ω is a Cartan classical domain of rank 2. By [Theorem 5.3](#), for $z = te_1 + te_2$ where $e_1, e_2 \in \mathcal{U}$,

$$\int_\Omega |K(z, w)| dv_w \approx (1-t)^{-a/2} \approx \delta_\Omega(z)^{-a/2} \quad \text{as } t \rightarrow 1^-.$$

When Ω is IV(n) with $n \geq 3$,

$$\int_\Omega |K(z, w)| dv_w \approx \delta_\Omega(z)^{-n/2+1}.$$

When Ω is III(4) or III(5),

$$\int_\Omega |K(z, w)| dv_w \approx \delta_\Omega(z)^{-2}.$$

When Ω is I(2, n) with $n \geq 2$,

$$\int_\Omega |K(z, w)| dv_w \approx \delta_\Omega(z)^{-1}.$$

When Ω is II(2),

$$\int_\Omega |K(z, w)| dv_w \approx \delta_\Omega(z)^{-1/2}.$$

Acknowledgements

Dong sincerely thanks Professors Bo-Yong Chen and Jinhao Zhang for their suggestions and warm encouragement throughout the years.

We greatly appreciate the referees who carefully read our original version and raised many valuable questions, which were very helpful when revising our paper.

References

- [Beatrous and Li 1993] F. Beatrous and S.-Y. Li, “On the boundedness and compactness of operators of Hankel type”, *J. Funct. Anal.* **111**:2 (1993), 350–379. [MR](#) [Zbl](#)
- [Berndtsson 1993] B. Berndtsson, “A smooth pseudoconvex domain in \mathbb{C}^2 for which L^∞ -estimates for $\bar{\partial}$ do not hold”, *Ark. Mat.* **31**:2 (1993), 209–218. [MR](#) [Zbl](#)
- [Berndtsson 1996] B. Berndtsson, “The extension theorem of Ohsawa–Takegoshi and the theorem of Donnelly–Fefferman”, *Ann. Inst. Fourier (Grenoble)* **46**:4 (1996), 1083–1094. [MR](#) [Zbl](#)
- [Berndtsson 1997] B. Berndtsson, “Uniform estimates with weights for the $\bar{\partial}$ -equation”, *J. Geom. Anal.* **7**:2 (1997), 195–215. [MR](#) [Zbl](#)
- [Berndtsson 2001] B. Berndtsson, “Weighted estimates for the $\bar{\partial}$ -equation”, pp. 43–57 in *Complex analysis and geometry* (Columbus, OH, 1999), edited by J. D. McNeal, Ohio State Univ. Math. Res. Inst. Publ. **9**, de Gruyter, Berlin, 2001. [MR](#) [Zbl](#)
- [Błocki 2005] Z. Błocki, “The Bergman metric and the pluricomplex Green function”, *Trans. Amer. Math. Soc.* **357**:7 (2005), 2613–2625. [MR](#) [Zbl](#)
- [Chen and Li 2019] R.-Y. Chen and S.-Y. Li, “Graham type theorem on classical bounded symmetric domains”, *Calc. Var. Partial Differential Equations* **58**:2 (2019), art. id. 81. [MR](#) [Zbl](#)
- [Chen and McNeal 2020] L. Chen and J. D. McNeal, “Product domains, multi-Cauchy transforms, and the $\bar{\partial}$ equation”, *Adv. Math.* **360** (2020), art. id. 106930. [MR](#) [Zbl](#)
- [Chen and Shaw 2001] S.-C. Chen and M.-C. Shaw, *Partial differential equations in several complex variables*, AMS/IP Stud. Adv. Math. **19**, Amer. Math. Soc., Providence, RI, 2001. [MR](#) [Zbl](#)
- [Demailly 1982] J.-P. Demailly, “Estimations L^2 pour l’opérateur $\bar{\partial}$ d’un fibré vectoriel holomorphe semi-positif au-dessus d’une variété kählérienne complète”, *Ann. Sci. École Norm. Sup. (4)* **15**:3 (1982), 457–511. [MR](#) [Zbl](#)
- [Demailly 2012] J.-P. Demailly, *Analytic methods in algebraic geometry*, Surv. Modern Monogr. **1**, International, Somerville, MA, 2012. [MR](#) [Zbl](#)
- [Dong et al. 2020] R. X. Dong, Y. Pan, and Y. Zhang, “Uniform estimates for the canonical solution to the $\bar{\partial}$ -equation on product domains”, preprint, 2020. [arXiv 2006.14484](#)
- [Donnelly 1994] H. Donnelly, “ L_2 cohomology of pseudoconvex domains with complete Kähler metric”, *Michigan Math. J.* **41**:3 (1994), 433–442. [MR](#) [Zbl](#)
- [Donnelly and Fefferman 1983] H. Donnelly and C. Fefferman, “ L^2 -cohomology and index theorem for the Bergman metric”, *Ann. of Math. (2)* **118**:3 (1983), 593–618. [MR](#) [Zbl](#)
- [Engliš and Zhang 2004] M. Engliš and G. Zhang, “On the Faraut–Korányi hypergeometric functions in rank two”, *Ann. Inst. Fourier (Grenoble)* **54**:6 (2004), 1855–1875. [MR](#) [Zbl](#)
- [Faraut and Korányi 1990] J. Faraut and A. Korányi, “Function spaces and reproducing kernels on bounded symmetric domains”, *J. Funct. Anal.* **88**:1 (1990), 64–89. [MR](#) [Zbl](#)
- [Fassina and Pan 2019] M. Fassina and Y. Pan, “Supnorm estimates for $\bar{\partial}$ on product domains in \mathbb{C}^n ”, preprint, 2019. To appear in *Acta Math. Sin. (Engl. Ser.)*. [arXiv 1903.10475](#)
- [Fefferman 1974] C. Fefferman, “The Bergman kernel and biholomorphic mappings of pseudoconvex domains”, *Invent. Math.* **26** (1974), 1–65. [MR](#) [Zbl](#)
- [Fornæss 1986] J. E. Fornæss, “Sup-norm estimates for $\bar{\partial}$ in \mathbb{C}^2 ”, *Ann. of Math. (2)* **123**:2 (1986), 335–345. [MR](#) [Zbl](#)
- [Fornæss and Sibony 1993] J. E. Fornæss and N. Sibony, “Smooth pseudoconvex domains in \mathbb{C}^2 for which the corona theorem and L^p estimates for $\bar{\partial}$ fail”, pp. 209–222 in *Complex analysis and geometry*, edited by V. Ancona and A. Silva, Plenum, New York, 1993. [MR](#) [Zbl](#)
- [Fornæss and Stensønes 1987] J. E. Fornæss and B. Stensønes, *Lectures on counterexamples in several complex variables*, Math. Notes **33**, Princeton Univ. Press, 1987. [MR](#) [Zbl](#)
- [Fornæss et al. 2011] J. E. Fornæss, L. Lee, and Y. Zhang, “On supnorm estimates for $\bar{\partial}$ on infinite type convex domains in \mathbb{C}^{2n} ”, *J. Geom. Anal.* **21**:3 (2011), 495–512. [MR](#) [Zbl](#)

- [Grauert and Lieb 1970] H. Grauert and I. Lieb, “Das Ramirezsche Integral und die Lösung der Gleichung $\bar{\partial} f = \alpha$ im Bereich der beschränkten Formen”, *Rice Univ. Stud.* **56:2** (1970), 29–50. [MR](#) [Zbl](#)
- [Grundmeier et al. 2022] D. Grundmeier, L. Simon, and B. Stensønes, “Sup-norm estimates for $\bar{\partial}$ ”, *Pure Appl. Math. Q.* **18:2** (2022), 531–571. [MR](#) [Zbl](#)
- [Henkin 1970] G. M. Henkin, “Integral representation of functions in strictly pseudoconvex domains and applications to the $\bar{\partial}$ -problem”, *Mat. Sb. (N.S.)* **82(124):2(6)** (1970), 300–308. In Russian; translated in *Math. USSR-Sb.* **11:2** (1970), 273–281. [MR](#) [Zbl](#)
- [Henkin 1971] G. M. Henkin, “A uniform estimate for the solution of the $\bar{\partial}$ -problem in a Weil region”, *Uspekhi Mat. Nauk* **26:3(159)** (1971), 211–212. In Russian. [MR](#)
- [Henkin and Leiterer 1983] G. M. Henkin and J. Leiterer, *Theory of functions on complex manifolds*, Monogr. Math. **79**, Birkhäuser, Basel, 1983. [MR](#) [Zbl](#)
- [Hörmander 1965] L. Hörmander, “ L^2 estimates and existence theorems for the $\bar{\partial}$ operator”, *Acta Math.* **113** (1965), 89–152. [MR](#) [Zbl](#)
- [Hua 1963] L. K. Hua, *Harmonic analysis of functions of several complex variables in the classical domains*, Transl. Math. Monogr. **6**, Amer. Math. Soc., Providence, RI, 1963. [MR](#) [Zbl](#)
- [Ishi and Yamaji 2011] H. Ishi and S. Yamaji, “Some estimates of the Bergman kernel of minimal bounded homogeneous domains”, *J. Lie Theory* **21:4** (2011), 755–769. [MR](#) [Zbl](#)
- [Kerzman 1971] N. Kerzman, “Hölder and L^p estimates for solutions of $\bar{\partial} u = f$ in strongly pseudoconvex domains”, *Comm. Pure Appl. Math.* **24** (1971), 301–379. [MR](#) [Zbl](#)
- [Korányi 1991] A. Korányi, “Hua-type integrals, hypergeometric functions and symmetric polynomials”, pp. 169–180 in *International Symposium in Memory of Hua Loo Keng, II: Analysis* (Beijing, 1988), edited by S. Gong et al., Springer, 1991. [MR](#) [Zbl](#)
- [Krantz 1976] S. G. Krantz, “Optimal Lipschitz and L^p regularity for the equation $\bar{\partial} u = f$ on strongly pseudo-convex domains”, *Math. Ann.* **219:3** (1976), 233–260. [MR](#) [Zbl](#)
- [Krantz and Li 2001a] S. G. Krantz and S.-Y. Li, “Boundedness and compactness of integral operators on spaces of homogeneous type and applications, I”, *J. Math. Anal. Appl.* **258:2** (2001), 629–641. [MR](#) [Zbl](#)
- [Krantz and Li 2001b] S. G. Krantz and S.-Y. Li, “Boundedness and compactness of integral operators on spaces of homogeneous type and applications, II”, *J. Math. Anal. Appl.* **258:2** (2001), 642–657. [MR](#) [Zbl](#)
- [Landucci 1975] M. Landucci, “On the projection of $L^2(D)$ into $H(D)$ ”, *Duke Math. J.* **42** (1975), 231–237. [MR](#) [Zbl](#)
- [Laurent-Thiébaud and Leiterer 1993] C. Laurent-Thiébaud and J. Leiterer, “Uniform estimates for the Cauchy–Riemann equation on q -convex wedges”, *Ann. Inst. Fourier (Grenoble)* **43:2** (1993), 383–436. [MR](#) [Zbl](#)
- [Lieb and Range 1986] I. Lieb and R. M. Range, “Integral representations and estimates in the theory of the $\bar{\partial}$ -Neumann problem”, *Ann. of Math. (2)* **123:2** (1986), 265–301. [MR](#) [Zbl](#)
- [Lu 1997] Q. Lu, *New results of classical manifolds and classical domains*, Shanghai Sci. Tech., 1997. In Chinese.
- [McNeal 1994] J. D. McNeal, “Estimates on the Bergman kernels of convex domains”, *Adv. Math.* **109:1** (1994), 108–139. [MR](#) [Zbl](#)
- [McNeal 1996] J. D. McNeal, “On large values of L^2 holomorphic functions”, *Math. Res. Lett.* **3:2** (1996), 247–259. [MR](#) [Zbl](#)
- [McNeal 2001] J. D. McNeal, “Invariant metric estimates for $\bar{\partial}$ on some pseudoconvex domains”, *Ark. Mat.* **39:1** (2001), 121–136. [MR](#) [Zbl](#)
- [Michel and Shaw 1999] J. Michel and M.-C. Shaw, “The $\bar{\partial}$ problem on domains with piecewise smooth boundaries with applications”, *Trans. Amer. Math. Soc.* **351:11** (1999), 4365–4380. [MR](#) [Zbl](#)
- [Nikolov and Pflug 2003] N. Nikolov and P. Pflug, “Estimates for the Bergman kernel and metric of convex domains in \mathbb{C}^n ”, *Ann. Polon. Math.* **81:1** (2003), 73–78. [MR](#) [Zbl](#)
- [Nikolov and Trybuła 2015] N. Nikolov and M. Trybuła, “The Kobayashi balls of $(\mathbb{C}-)$ convex domains”, *Monatsh. Math.* **177:4** (2015), 627–635. [MR](#) [Zbl](#)
- [Range 1978] R. M. Range, “On Hölder estimates for $\bar{\partial} u = f$ on weakly pseudoconvex domains”, pp. 247–267 in *Several complex variables* (Cortona, Italy, 1976/1977), edited by J. J. Kohn and E. Vesentini, Scuola Norm. Sup. Pisa, 1978. [MR](#) [Zbl](#)

- [Range 1986] R. M. Range, *Holomorphic functions and integral representations in several complex variables*, Grad. Texts Math. **108**, Springer, 1986. [MR](#) [Zbl](#)
- [Range 1990] R. M. Range, “Integral kernels and Hölder estimates for $\bar{\partial}$ on pseudoconvex domains of finite type in \mathbb{C}^2 ”, *Math. Ann.* **288**:1 (1990), 63–74. [MR](#) [Zbl](#)
- [Range and Siu 1972] R. M. Range and Y.-T. Siu, “Uniform estimates for the $\bar{\partial}$ -equation on intersections of strictly pseudoconvex domains”, *Bull. Amer. Math. Soc.* **78** (1972), 721–723. [MR](#) [Zbl](#)
- [Range and Siu 1973] R. M. Range and Y.-T. Siu, “Uniform estimates for the $\bar{\partial}$ -equation on domains with piecewise smooth strictly pseudoconvex boundaries”, *Math. Ann.* **206** (1973), 325–354. [MR](#) [Zbl](#)
- [Romanov and Henkin 1971] A. V. Romanov and G. M. Henkin, “Exact Hölder estimates for the solutions of the $\bar{\partial}$ -equation”, *Izv. Akad. Nauk SSSR Ser. Mat.* **35**:5 (1971), 1171–1183. In Russian; translated in *Math. USSR-Izv.* **5**:5 (1971), 1180–1192. [MR](#) [Zbl](#)
- [Schuster and Varolin 2014] A. P. Schuster and D. Varolin, “New estimates for the minimal L^2 solution of $\bar{\partial}$ and applications to geometric function theory in weighted Bergman spaces”, *J. Reine Angew. Math.* **691** (2014), 173–201. [MR](#) [Zbl](#)
- [Sergeev 1994] A. G. Sergeev, “ $\bar{\partial}$ in the future tube”, pp. 356–359 in *Linear and complex analysis: problem book 3, II*, edited by V. P. Havin and N. K. Nikolski, Lecture Notes in Math. **1574**, Springer, 1994.
- [Sibony 1980] N. Sibony, “Un exemple de domaine pseudoconvexe régulier où l’équation $\bar{\partial}u = f$ n’admet pas de solution bornée pour f bornée”, *Invent. Math.* **62**:2 (1980), 235–242. [MR](#) [Zbl](#)
- [Siu 1996] Y.-T. Siu, “The Fujita conjecture and the extension theorem of Ohsawa–Takegoshi”, pp. 577–592 in *Geometric complex analysis* (Hayama, Japan, 1995), edited by J. Noguchi et al., World Sci., River Edge, NJ, 1996. [MR](#) [Zbl](#)
- [Yan 1992] Z. M. Yan, “A class of generalized hypergeometric functions in several variables”, *Canad. J. Math.* **44**:6 (1992), 1317–1338. [MR](#) [Zbl](#)

Received 26 May 2020. Revised 29 Mar 2021. Accepted 10 Jun 2021.

ROBERT XIN DONG: dong@uconn.edu

Department of Mathematics, University of Connecticut, Storrs, CT, United States

SONG-YING LI: sli@math.uci.edu

Department of Mathematics, University of California, Irvine, CA, United States

JOHN N. TREUER: jtreuer@uci.edu

Department of Mathematics, University of California, Irvine, CA, United States

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard Université Paris Sud XI, France
patrick.gerard@universite-paris-saclay.fr

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Isabelle Gallagher	Université Paris-Diderot, IMJ-PRG, France gallagher@math.ens.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	András Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 2 2023

Riesz transform and vertical oscillation in the Heisenberg group KATRIN FÄSSLER and TUOMAS ORPONEN	309
A Wess–Zumino–Witten type equation in the space of Kähler potentials in terms of Hermitian–Yang–Mills metrics KUANG-RU WU	341
The strong topology of ω -plurisubharmonic functions ANTONIO TRUSIANI	367
Sharp pointwise and uniform estimates for $\bar{\partial}$ ROBERT XIN DONG, SONG-YING LI and JOHN N. TREUER	407
Some applications of group-theoretic Rips constructions to the classification of von Neumann algebras IONUȚ CHIFAN, SAYAN DAS and KRISHNENDU KHAN	433
Long time existence of Yamabe flow on singular spaces with positive Yamabe constant JØRGEN OLSEN LYE and BORIS VERTMAN	477
Disentanglement, multilinear duality and factorisation for nonpositive operators ANTHONY CARBERY, TIMO S. HÄNNINEN and STEFÁN INGI VALDIMARSSON	511
The Green function with pole at infinity applied to the study of the elliptic measure JOSEPH FENEUIL	545
Talagrand’s influence inequality revisited DARIO CORDERO-ERAUSQUIN and ALEXANDROS ESKENAZIS	571