

ANALYSIS & PDE

Volume 16

No. 3

2023

ILAN HIRSHBERG AND N. CHRISTOPHER PHILLIPS

**A SIMPLE NUCLEAR C^* -ALGEBRA WITH AN INTERNAL
ASYMMETRY**

A SIMPLE NUCLEAR C^* -ALGEBRA WITH AN INTERNAL ASYMMETRY

ILAN HIRSHBERG AND N. CHRISTOPHER PHILLIPS

We construct an example of a simple approximately homogeneous C^* -algebra such that its Elliott invariant admits an automorphism which is not induced by an automorphism of the algebra.

Classification theory for simple nuclear C^* -algebras reached a milestone recently. The results of [Elliott et al. 2015; Tikuisis et al. 2017], building on decades of work by many authors, show that simple separable unital C^* -algebras with finite nuclear dimension satisfying the universal coefficient theorem are classified via the Elliott invariant, $\text{Ell}(\cdot)$, which consists of the ordered K_0 -group along with the class of the identity, the K_1 -group, the trace simplex, and the pairing between the trace simplex and the K_0 -group. Earlier counterexamples due to Toms [2008] and Rørdam [2003], related to ideas of Villadsen [1998], show that one cannot expect to be able to extend this classification theorem beyond the case of finite nuclear dimension, at least not without either extending the invariant or restricting to another class of C^* -algebras. An important facet of the classification theorems is a form of rigidity. Starting with two C^* -algebras A and B and an isomorphism $\Phi : \text{Ell}(A) \rightarrow \text{Ell}(B)$, one not only shows that A and B are isomorphic, but rather that there exists an isomorphism from A to B which induces the given isomorphism Φ on the level of the Elliott invariant.

The goal of this paper is to illustrate how this existence property may fail in the infinite nuclear dimension setting, even when restricting to a class consisting of a single C^* -algebra. Namely, we construct an example of a simple unital nuclear separable AH algebra C , along with an automorphism of $\text{Ell}(C)$, which is not induced by any automorphism of C . This can be viewed as a companion of sorts to [Toms 2008, Theorem 1.2], where it was shown that when such automorphisms exist, they need not be unique in the sense described. The mechanism of the example is that if there were such an automorphism φ , there would be projections $p, q \in \mathbb{C}$ such that $\varphi(p) = q$ but such that the corners pCp and qCq have different radii of comparison [Toms 2006] (the definition is recalled at the beginning of Section 1). This further shows that simple unital AH algebras can be quite inhomogeneous. In particular, extending the Elliott invariant by adding something as simple as the radius of comparison will not help for the classification of AH algebras which are not Jiang–Su stable.

This material is based upon work supported by the US National Science Foundation under grant DMS-1501144, by the US-Israel Binational Science Foundation and by the Israel Science Foundation grant no. 476/16.

MSC2020: 46L35, 46L40, 46L80.

Keywords: C^* -algebras, Elliott invariant.

We now give an overview of our construction. We start with the counterexample from [Toms 2008, Theorem 1.1]. We consider two direct systems, described diagrammatically as follows:

$$\begin{array}{c}
 C(X_0) \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C(X_1) \otimes M_{r(1)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C(X_2) \otimes M_{r(2)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \cdots \\
 \\
 C([0, 1]) \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C([0, 1]) \otimes M_{r(1)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C([0, 1]) \otimes M_{r(2)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \cdots
 \end{array} \tag{0-1}$$

The ordinary arrows indicate a large (and rapidly increasing) number of embeddings which are carefully chosen, and the dotted arrows indicate a small number of point evaluation maps, thrown in so as to ensure that the resulting direct limit is simple. The spaces in the upper diagram are contractible CW complexes whose dimension increases rapidly compared to the sizes of the matrix algebras. (Toms uses cubes; in our construction we found it easier to use cones over products of spheres, but the underlying idea is similar.) The direct system is constructed so as to have positive radius of comparison. We use [Thomsen 1994] to choose the lower diagram so as to mimic the upper diagram, and produce the same Elliott invariant. As the resulting algebra on the bottom is AI, it has strict comparison, and therefore is not isomorphic to the one on the top. (In [Toms 2008] it isn't important for the two diagrams to match up nicely in terms of the ranks of the matrices involved. However, we will show that it can be done, as it is important for us.)

Our construction involves moving the point evaluations across, so as to merge the two systems:

$$\begin{array}{c}
 C(X_0) \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C(X_1) \otimes M_{r(1)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C(X_2) \otimes M_{r(2)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \cdots \\
 \begin{array}{ccc} \swarrow & \searrow & \swarrow \\ & & \searrow \\ \swarrow & \searrow & \swarrow \\ & & \searrow \\ \swarrow & \searrow & \swarrow \\ & & \searrow \end{array} \\
 C([0, 1]) \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C([0, 1]) \otimes M_{r(1)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} C([0, 1]) \otimes M_{r(2)} \begin{array}{c} \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \\ \text{====} \end{array} \cdots
 \end{array} \tag{0-2}$$

With care, one can arrange for the flip between the two levels of the diagram to make sense as an automorphism of the Elliott invariant. The resulting C^* -algebra has positive radius of comparison and behaves roughly as badly as Toms' example. Nevertheless, we can distinguish a part of it which roughly corresponds to the rapid dimension growth diagram on the top from a part which roughly corresponds to the AI part on the bottom. Namely, if at the first level $C(X_0) \oplus C([0, 1])$ we denote by q the function which is 1 on X_0 and 0 on $[0, 1]$, and we define $q^\perp = 1 - q$, then the K_0 -classes of q and q^\perp will be switched by the automorphism of the Elliott invariant we construct. However, we can tell apart the corners qCq and $q^\perp Cq^\perp$ by considering their radii of comparison.

Section 1 develops the choices needed to get different radii of comparison in different corners of the algebra we construct. Section 2 contains the work needed to assemble the ingredients of the construction into a simple C^* -algebra whose Elliott invariant admits an appropriate automorphism. The main theorem is in Section 3.

1. Upper and lower bounds on the radius of comparison

We recall the required standard definitions and notation related to the Cuntz semigroup. See Section 2 of [Rørdam 1992] for details. For a unital C^* -algebra A , we denote its tracial state space by $T(A)$. We take $M_\infty(A) = \bigcup_{n=1}^\infty M_n(A)$, using the usual embeddings $M_n(A) \hookrightarrow M_{n+1}(A)$. For $\tau \in T(A)$, we

define $d_\tau : M_\infty(A)_+ \rightarrow [0, \infty)$ by $d_\tau(a) = \lim_{n \rightarrow \infty} \tau(a^{1/n})$. If $a, b \in M_\infty(A)_+$, then $a \precsim b$ (a is Cuntz subequivalent to b) if there is a sequence $(v_n)_{n=1}^\infty$ in $M_\infty(A)$ such that $\lim_{n \rightarrow \infty} v_n b v_n^* = a$.

Following [Toms 2006, Definition 6.1], for $\rho \in [0, \infty)$, we say that A has ρ -comparison if whenever $a, b \in M_\infty(A)_+$ satisfy $d_\tau(a) + \rho < d_\tau(b)$ for all $\tau \in T(A)$, then $a \precsim b$. The radius of comparison of A , denoted by $\text{rc}(A)$, is

$$\text{rc}(A) = \inf(\{\rho \in [0, \infty) \mid A \text{ has } \rho\text{-comparison}\}).$$

We take $\text{rc}(A) = \infty$ if there is no ρ such that A has ρ -comparison. Since AH algebras are nuclear, all quasitraces on them are traces by [Haagerup 2014, Theorem 5.11]. Thus, we ignore quasitraces. Also, by [Phillips 2014, Proposition 6.12], the radius of comparison remains unchanged if we replace $M_\infty(A)$ by $K \otimes A$ throughout. Thus, we may work only in $M_\infty(A)$.

Our construction uses a specific setup, with a number of parameters of various kinds which must be chosen to satisfy specific conditions. Construction 1.1 lists for reference many of the objects used in it, and some of the conditions they must satisfy. It abstracts the diagram (0-2). Construction 1.6 specifies the choices of spaces and maps needed for the results on Cuntz comparison, and Construction 2.17, together with the additional maps in parts (11), (12), and (13) of Construction 1.1, is used to arrange the existence of a suitable automorphism of the tracial state space of the algebra we construct. Because of the necessity of passing to a subsystem at one stage in this process, we must start the proof of the main theorem with a version of just the top row in the diagram (0-1); this is Construction 3.3. Many of the lemmas use only a few of the objects and their properties, so that the reader can refer back to just the relevant parts of the constructions. In particular, many details are used only in this section or only in Section 2. Some of the details are used for just one lemma each.

Construction 1.1. For much of this paper, we will consider algebras constructed in the following way and using the following notation:

(1) $(d(n))_{n=0,1,2,\dots}$ and $(k(n))_{n=0,1,2,\dots}$ are sequences in $\mathbb{Z}_{\geq 0}$, with $d(0) = 1$ and $k(0) = 0$. Moreover, for $n \in \mathbb{Z}_{\geq 0}$,

$$l(n) = d(n) + k(n), \quad r(n) = \prod_{j=0}^n l(j), \quad \text{and} \quad s(n) = \prod_{j=0}^n d(j).$$

Further define $t(n)$ inductively as follows. Set $t(0) = 0$, and

$$t(n + 1) = d(n + 1)t(n) + k(n + 1)[r(n) - t(n)].$$

(See Lemma 1.14 for the significance of $t(n)$.)

(2) We will assume that $k(n) < d(n)$ for all $n \in \mathbb{Z}_{\geq 0}$.

(3) We define

$$\kappa = \inf_{n \in \mathbb{Z}_{> 0}} \frac{s(n)}{r(n)}.$$

For estimates involving the radius of comparison, we will assume $\kappa > \frac{1}{2}$.

(4) The numbers $\omega, \omega' \in (0, \infty]$ are defined by

$$\omega = \frac{k(1)}{k(1) + d(1)} \quad \text{and} \quad \omega' = \sum_{n=2}^{\infty} \frac{k(n)}{k(n) + d(n)}.$$

We will require $\omega' < \omega < \frac{1}{2}$. In particular,

$$\sum_{n=1}^{\infty} \frac{k(n)}{k(n) + d(n)} < \infty.$$

(5) We will also eventually require that κ as in (3) and ω as in (4) are related by $2\kappa - 1 > 2\omega$. This can easily be arranged with a suitable choice of $d(1)$ and $k(1)$.

(6) $(X_n)_{n=0,1,2,\dots}$ and $(Y_n)_{n=0,1,2,\dots}$ are sequences of compact metric spaces. (They will be further specified in Construction 1.6.)

(7) For $n \in \mathbb{Z}_{\geq 0}$, the algebra C_n is

$$C_n = M_{r(n)} \otimes (C(X_n) \oplus C(Y_n)).$$

We further make the identifications

$$\begin{aligned} C(X_{n+1}, M_{r(n+1)}) &= M_{l(n+1)} \otimes C(X_{n+1}, M_{r(n)}), \\ C(Y_{n+1}, M_{r(n+1)}) &= M_{l(n+1)} \otimes C(Y_{n+1}, M_{r(n)}), \\ C(X_n) \oplus C(Y_n) &= C(X_n \amalg Y_n), \\ C(X_n, M_{r(n)}) \oplus C(Y_n, M_{r(n)}) &= C(X_n \amalg Y_n, M_{r(n)}). \end{aligned}$$

(8) For $n \in \mathbb{Z}_{> 0}$, we are given a unital homomorphism

$$\gamma_n : C(X_n) \oplus C(Y_n) \rightarrow M_{l(n+1)}(C(X_{n+1}) \oplus C(Y_{n+1})),$$

and the homomorphism

$$\Gamma_{n+1, n} : C_n \rightarrow C_{n+1}$$

is given by $\Gamma_{n+1, n} = \text{id}_{M_{r(n)}} \otimes \gamma_n$. Moreover, for $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$,

$$\Gamma_{n, m} = \Gamma_{n, n-1} \circ \Gamma_{n-1, n-2} \circ \cdots \circ \Gamma_{m+1, m} : C_m \rightarrow C_n.$$

In particular, $\Gamma_{n, n} = \text{id}_{C_n}$.

(9) We require that the maps

$$\gamma_n : C(X_n \amalg Y_n) \rightarrow M_{l(n+1)}(C(X_{n+1} \amalg Y_{n+1}))$$

in (8) be diagonal; that is, that there exist continuous functions

$$S_{n,1}, S_{n,2}, \dots, S_{n, l(n+1)} : X_{n+1} \amalg Y_{n+1} \rightarrow X_n \amalg Y_n$$

such that for all $f \in C(X_n \amalg Y_n)$, we have

$$\gamma_n(f) = \text{diag}(f \circ S_{n,1}, f \circ S_{n,2}, \dots, f \circ S_{n, l(n+1)}).$$

(These maps will be specified further in Construction 1.6.)

(10) We set $C = \varinjlim_n C_n$, taken with respect to the maps $\Gamma_{n,m}$. The maps associated with the direct limit will be called $\Gamma_{\infty,m} : C_m \rightarrow C$ for $m \in \mathbb{Z}_{\geq 0}$.

As we need to work with two diagrams which are similar in most positions, as in diagrams (0-1) and (0-2), we sometimes use additional objects and conditions in the construction, as follows:

(11) For $n \in \mathbb{Z}_{>0}$, we may be given an additional unital homomorphism

$$\gamma_n^{(0)} : C(X_n) \oplus C(Y_n) \rightarrow M_{l(n+1)}(C(X_{n+1}) \oplus C(Y_{n+1})).$$

Then the maps $\Gamma_{n+1,n}^{(0)} : C_n \rightarrow C_{n+1}$, $\Gamma_{n,m}^{(0)} : C_m \rightarrow C_n$ are defined analogously to (8), the algebra $C^{(0)}$ is given as $C^{(0)} = \varinjlim_n C_n$, taken with respect to the maps $\Gamma_{n,m}^{(0)}$, and the maps $\Gamma_{\infty,m}^{(0)} : C_m \rightarrow C^{(0)}$ are defined analogously to (10).

(12) In (11), analogously to (9), we may require that there be

$$S_{n,1}^{(0)}, S_{n,2}^{(0)}, \dots, S_{n,l(n+1)}^{(0)} : X_{n+1} \amalg Y_{n+1} \rightarrow X_n \amalg Y_n$$

such that for all $f \in C(X_n \amalg Y_n)$ we have

$$\gamma_n^{(0)}(f) = \text{diag}(f \circ S_{n,1}^{(0)}, f \circ S_{n,2}^{(0)}, \dots, f \circ S_{n,l(n+1)}^{(0)}).$$

(These maps will be specified further in Construction 1.6.)

(13) Assuming diagonal maps as in (9), we may require that they agree in the coordinates $1, 2, \dots, d(n+1)$; that is, for $n \in \mathbb{Z}_{>0}$ and $k = 1, 2, \dots, d(n+1)$, we have $S_{n,k}^{(0)} = S_{n,k}$.

Lemma 1.2. *In Construction 1.1(1), the sequence $(s(n)/r(n))_{n=1,2,\dots}$ is strictly decreasing.*

Proof. The proof is straightforward. □

Lemma 1.3. *In Construction 1.1(1), and assuming Construction 1.1(2), we have*

$$0 = \frac{t(0)}{r(0)} < \frac{t(1)}{r(1)} < \frac{t(2)}{r(2)} < \dots < \frac{1}{2}.$$

Proof. We have $t(0) = 0$ by definition. We prove by induction on $n \in \mathbb{Z}_{>0}$ that

$$\frac{t(n-1)}{r(n-1)} < \frac{t(n)}{r(n)} < \frac{1}{2}. \tag{1-1}$$

This will finish the proof. For $n = 1$, we have

$$\frac{t(1)}{r(1)} = \frac{k(1)}{k(1) + d(1)},$$

which is in $(0, \frac{1}{2})$ by Construction 1.1(2). Now assume (1-1); we prove this relation with $n + 1$ in place of n . We have $r(n) - t(n) > t(n)$, so

$$\frac{t(n+1)}{r(n+1)} = \frac{d(n+1)t(n) + k(n+1)[r(n) - t(n)]}{[d(n+1) + k(n+1)]r(n)} > \frac{d(n+1)t(n) + k(n+1)t(n)}{[d(n+1) + k(n+1)]r(n)} = \frac{t(n)}{r(n)}. \tag{1-2}$$

Also, with

$$\alpha = \frac{d(n+1)}{d(n+1) + k(n+1)} \quad \text{and} \quad \beta = \frac{t(n)}{r(n)},$$

starting with the first step in (1-2), and at the end using $\alpha > \frac{1}{2}$ (by Construction 1.1(2)) and $\beta < \frac{1}{2}$ (by the induction hypothesis), we have

$$\frac{t(n+1)}{r(n+1)} = \alpha\beta + (1-\alpha)(1-\beta) = \frac{1}{2}[1 - (2\alpha - 1)(1 - 2\beta)] < \frac{1}{2}.$$

This completes the induction, and the proof. \square

Lemma 1.4. *With the notation of Constructions 1.1(1) and 1.1(4), and assuming the conditions in Constructions 1.1(2) and 1.1(4), for all $n \in \mathbb{Z}_{>0}$ we have*

$$\omega \leq \frac{t(n)}{r(n)} \leq \omega + \omega' < 2\omega.$$

Proof. The third inequality is immediate from Construction 1.1(4).

By Lemma 1.3, the sequence $(t(n)/r(n))_{n=1,2,\dots}$ is strictly increasing. Also,

$$\frac{t(1)}{r(1)} = \frac{k(1)}{k(1) + d(1)} = \omega. \tag{1-3}$$

The first inequality in the statement now follows.

Next, we claim that

$$\frac{t(n)}{r(n)} \leq \sum_{j=1}^n \frac{k(j)}{k(j) + d(j)}$$

for all $n \in \mathbb{Z}_{>0}$. The case $n = 1$ is (1-3). Assume this inequality is known for n . Then

$$\begin{aligned} \frac{t(n+1)}{r(n+1)} &= \left(\frac{d(n+1)}{k(n+1) + d(n+1)} \right) \left(\frac{t(n)}{r(n)} \right) + \left(\frac{k(n+1)}{k(n+1) + d(n+1)} \right) \left(\frac{r(n) - t(n)}{r(n)} \right) \\ &\leq \frac{t(n)}{r(n)} + \frac{k(n+1)}{k(n+1) + d(n+1)} \leq \sum_{j=1}^{n+1} \frac{k(j)}{k(j) + d(j)}, \end{aligned}$$

as desired.

The second inequality in the statement now follows. \square

Notation 1.5. For a topological space X , we define

$$\text{cone}(X) = (X \times [0, 1]) / (X \times \{0\}).$$

Then $\text{cone}(X)$ is contractible, and $\text{cone}(\cdot)$ is a covariant functor: if $T : X \rightarrow Y$ is a continuous map, then it induces a continuous map $\text{cone}(T) : \text{cone}(X) \rightarrow \text{cone}(Y)$. We identify X with the image of $X \times \{1\}$ in $\text{cone}(X)$.

Construction 1.6. We give further details on the spaces X_n and Y_n in Construction 1.1(6).

(14) The space X_n is chosen as follows. First set $Z_0 = S^2$. With $(d(n))_{n=0,1,2,\dots}$ and $(s(n))_{n=0,1,2,\dots}$ as in Construction 1.1(1), define inductively

$$Z_n = Z_{n-1}^{d(n)} = (S^2)^{s(n)}.$$

Then set $X_n = \text{cone}(Z_n)$. (In particular, X_n is contractible, and $Z_n \subset X_n$ as in Notation 1.5.) Further, for $n \in \mathbb{Z}_{\geq 0}$ and $j = 1, 2, \dots, d(n+1)$, we let $P_j^{(n)} : Z_{n+1} \rightarrow Z_n$ be the j -th coordinate projection, and we set $Q_j^{(n)} = \text{cone}(P_j^{(n)}) : X_{n+1} \rightarrow X_n$.

(15) $Y_n = [0, 1]$ for all $n \in \mathbb{Z}_{>0}$. (In particular, Y_n is contractible.)

(16) We assume we are given points $x_m \in X_m$ for $m \in \mathbb{Z}_{\geq 0}$ such that, using the notation in (14), for all $n \in \mathbb{Z}_{\geq 0}$, the set

$$\left\{ (Q_{v_1}^{(n)} \circ Q_{v_2}^{(n+1)} \circ \dots \circ Q_{v_{m-n}}^{(m-1)})(x_m) \mid m = n+1, n+2, \dots \text{ and } v_j = 1, 2, \dots, d(n+j) \text{ for } j = 1, 2, \dots, m-n \right\}$$

is dense in X_n .

(17) We assume we are given a sequence $(y_k)_{k=0,1,2,\dots}$ in $[0, 1]$ such that for all $n \in \mathbb{Z}_{\geq 0}$ the set $\{y_k \mid k \geq n\}$ is dense in $[0, 1]$.

(18) The maps

$$\gamma_n : C(X_n \amalg Y_n) \rightarrow M_{l(n+1)}(C(X_{n+1} \amalg Y_{n+1}))$$

will be as in Construction 1.1(9), with the maps $S_{n,j} : X_{n+1} \amalg Y_{n+1} \rightarrow X_n \amalg Y_n$ appearing there defined as follows:

(a) With $Q_j^{(n)}$ as in (14), we set $S_{n,j}(x) = Q_j^{(n)}(x)$ for $x \in X_{n+1}$ and $j = 1, 2, \dots, d(n+1)$.

(b) $S_{n,j}(x) = y_n$ for

$$x \in X_{n+1} \quad \text{and} \quad j = d(n+1) + 1, d(n+1) + 2, \dots, l(n+1).$$

(c) There are continuous functions

$$R_{n,1}, R_{n,2}, \dots, R_{n,d(n+1)} : Y_{n+1} \rightarrow Y_n$$

(which will be taken from Proposition 2.14 below) such that $S_{n,j}(y) = R_{n,j}(y)$ for $y \in Y_{n+1}$ and $j = 1, 2, \dots, d(n+1)$.

(d) $S_{n,j}(y) = x_n$ for

$$y \in Y_{n+1} \quad \text{and} \quad j = d(n+1) + 1, d(n+1) + 2, \dots, l(n+1).$$

(19) The maps

$$\gamma_n^{(0)} : C(X_n \amalg Y_n) \rightarrow M_{l(n+1)}(C(X_{n+1} \amalg Y_{n+1}))$$

will be as in Construction 1.1(12), with the maps $S_{n,j}^{(0)} : X_{n+1} \amalg Y_{n+1} \rightarrow X_n \amalg Y_n$ appearing there given by $S_{n,j}^{(0)} = S_{n,j}$ for $j = 1, 2, \dots, d(n+1)$ and to be specified later for $j = d(n+1) + 1, d(n+1) + 2, \dots, l(n+1)$.

With the choices in Construction 1.6(18), the map

$$\gamma_n : C(X_n) \oplus C(Y_n) \rightarrow C(X_{n+1}, M_{l(n+1)}) \oplus C(Y_{n+1}, M_{l(n+1)})$$

in Construction 1.1(8), as further specified in Construction 1.1(9), is given as follows. With $\mathbb{C}^{d(n)}$ viewed as embedded in $M_{d(n)}$ as the diagonal matrices, there is a homomorphism

$$\delta_n : C(Y_n) \rightarrow C(Y_{n+1}, \mathbb{C}^{d(n+1)}) \subset C(Y_{n+1}, M_{d(n+1)})$$

such that

$$\gamma_n(f, g) = (\text{diag}(f \circ Q_1^{(n)}, f \circ Q_2^{(n)}, \dots, f \circ Q_{d(n+1)}^{(n)}, \underbrace{g(y_n), g(y_n), \dots, g(y_n)}_{k(n+1) \text{ times}}, \text{diag}(\delta_n(g), \underbrace{f(x_n), f(x_n), \dots, f(x_n)}_{k(n+1) \text{ times}})). \quad (1-4)$$

For the purposes of this section, we need no further information on the maps δ_n , except that they send constant functions to constant functions.

Lemma 1.7. *Assume the notation and choices in parts (1), (7), (8), and (10) of Construction 1.1, and in Construction 1.6 (except part (19)) and the parts of Construction 1.1 referred to there. Then the algebra C is simple.*

Proof. Using Construction 1.6(16), this is easily deduced from [Dădărlat et al. 1992, Proposition 2.1]. \square

Notation 1.8. Let $p \in C(S^2, M_2)$ denote the Bott projection, and let L be the tautological line bundle over $S^2 \cong \mathbb{C}P^1$. (Thus, the range of p is the section space of L .) Recalling that $X_0 = \text{cone}(S^2)$, parametrized as in Notation 1.5, define $b \in C(X_0, M_2)$ by $b(\lambda) = \lambda \cdot p$ for $\lambda \in [0, 1]$. Assuming the notation and choices in parts (1), (6), (7), (8), and (10) of Construction 1.1 and in Construction 1.6, for $n \in \mathbb{Z}_{\geq 0}$ set $b_n = (\text{id}_{M_2} \otimes \Gamma_{n,0})(b, 0) \in M_2(C_n)$.

We require the following simple lemma concerning characteristic classes. It gives us a way of estimating the radius of comparison, which is similar to the one used in [Villadsen 1998, Lemma 1], but more suitable for the types of estimates we need here.

Lemma 1.9. *The Cartesian product $L^{\times k}$ does not embed in a trivial bundle over $(S^2)^k$ of rank less than $2k$.*

Proof. We refer the reader to [Milnor and Stasheff 1974, Section 14] for an account of Chern classes. The Chern character $c(L)$ is of the form $1 + \varepsilon$, where ε is a generator of $H^2(S^2, \mathbb{Z})$, and the product operation satisfies $\varepsilon^2 = 0$. Let $P_1, P_2, \dots, P_k : (S^2)^k \rightarrow S^2$ be the coordinate projections. For $j = 1, 2, \dots, k$, set $\varepsilon_j = P_j^*(\varepsilon)$. The elements $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_k \in H^2((S^2)^k, \mathbb{Z})$, along with $1 \in H^0((S^2)^k, \mathbb{Z})$ (the standard generator) generate the cohomology ring of $(S^2)^k$ and satisfy $\varepsilon_j^2 = 0$ for $j = 1, 2, \dots, k$. By naturality of the Chern character [Milnor and Stasheff 1974, Lemma 14.2] and the product theorem [Milnor and Stasheff 1974, (14.7) on page 164], we have $c(L^{\times k}) = \prod_{j=1}^k (1 + \varepsilon_j)$. Now, suppose $L^{\times k}$ embeds as a subbundle of a trivial bundle E . Let F be the complementary bundle, so that $L^{\times k} \oplus F = E$. By the product theorem, $c(L^{\times k})c(F) = c(L^{\times k} \oplus F) = c(E) = 1$. Thus, $c(F) = c(L^{\times k})^{-1} = \prod_{j=1}^k (1 - \varepsilon_j)$. Since $c(F)$ has a nonzero term in the top cohomology group $H^{2k}((S^2)^k)$, it follows that $\text{rank}(F)$ is at least k . Thus, $\text{rank}(E) = \text{rank}(L^{\times k}) + \text{rank}(F) \geq 2k$, as required. \square

Lemma 1.10. *Adopt the assumptions and notation of Notation 1.8. Let $n \in \mathbb{Z}_{>0}$. Then $b_n|_{Z_n}$ is the orthogonal sum of a projection p_n whose range is isomorphic to the section space of the Cartesian product bundle $L^{\times s(n)}$ and a constant function of rank at most $r(n) - s(n) - t(n)$.*

We don't expect $b_n|_{Z_n}$ to be a projection, since some of the point evaluations occurring in the maps of the direct system will be at points $x \in \text{cone}(Z_m) \setminus Z_m$ for values of $m < n$, and $b_m(x)$ is not a projection for such x .

We don't need the estimate on the rank of the second part of the description of $b_n|_{Z_n}$; it is included to make the construction more explicit. If there are no evaluations at the "cone points"

$$(Z_m \times \{0\}) / (Z_m \times \{0\}) \in (Z_m \times [0, 1]) / (Z_m \times \{0\})$$

(following the parametrization in Notation 1.5), then this rank will be exactly $r(n) - s(n) - t(n)$.

Proof of Lemma 1.10. For $n \in \mathbb{Z}_{\geq 0}$ write $b_n = (c_n, g_n)$, with

$$c_n \in M_2(C(X_n, M_{r(n)})) \quad \text{and} \quad g_n \in M_2(C(Y_n, M_{r(n)})).$$

Further, for $j = 1, 2, \dots, s(n)$ let $T_j^{(n)} : (S^2)^{s(n)} \rightarrow S^2$ be the j -th coordinate projection. We claim that c_n is an orthogonal sum $c_{n,0} + c_{n,1}$, in which $c_{n,0}$ is the direct sum of the functions $b \circ \text{cone}(T_j^{(n)})$ for $j = 1, 2, \dots, s(n)$ and $c_{n,1}$ is a constant function of rank at most $r(n) - s(n) - t(n)$, and moreover that g_n is a constant function of rank at most $t(n)$. The statement of the lemma follows from this claim.

The proof of the claim is by induction on n . The claim is true for $n = 0$, by the definition of b and since $s(0) = 1$, $t(0) = 0$, and $r(0) - s(0) - t(0) = 0$.

Now assume that the claim is known for n , recall that $\Gamma_{n+1,n} = \text{id}_{M_{r(n)}} \otimes \gamma_n$ (see Construction 1.1(8)), and examine the summands in the description (1-4) of the map γ_n (after Construction 1.6). With this convention, first take (f, g) in (1-4) to be $(c_{n,0}, 0)$. The first coordinate $\Gamma_{n+1,n}(c_{n,0}, 0)_1$ is of the form required for $c_{n+1,0}$, while $\Gamma_{n+1,n}(c_{n,0}, 0)_2$ is a constant function of rank $k(n+1)s(n)$ unless $c_n(x_n) = 0$, in which case it is zero. In the same manner, we see that:

- $\Gamma_{n+1,n}(c_{n,1}, 0)_1$ is constant of rank at most $d(n+1)[r(n) - s(n) - t(n)]$.
- $\Gamma_{n+1,n}(c_{n,1}, 0)_2$ is constant of rank at most $k(n+1)[r(n) - s(n) - t(n)]$.
- $\Gamma_{n+1,n}(0, g_n)_1$ is constant of rank at most $k(n+1)t(n)$.
- $\Gamma_{n+1,n}(0, g_n)_2$ is constant of rank at most $d(n+1)t(n)$.

Putting these together, we get in the first coordinate of $\Gamma_{n+1,n}(b_n)$ the direct sum of $c_{n+1,0}$ as described and a constant function of rank at most

$$d(n+1)[r(n) - s(n) - t(n)] + k(n+1)t(n).$$

A computation shows that this expression is equal to $r(n+1) - s(n+1) - t(n+1)$. In the second coordinate we get a constant function of rank at most

$$k(n+1)s(n) + k(n+1)[r(n) - s(n) - t(n)] + d(n+1)t(n) = t(n+1).$$

This completes the induction, and the proof. □

Corollary 1.11. *Adopt the assumptions and notation of Notation 1.8. Let $n \in \mathbb{Z}_{\geq 0}$. Let $e = (e_1, e_2)$ be an element in $M_\infty(C_n) \cong M_\infty(C(X_n) \oplus C(Y_n))$ such that e_1 is a projection which is equivalent to a constant projection. If there exists $x \in M_\infty(C_n)$ such that $\|xex^* - b_n\| < \frac{1}{2}$ then $\text{rank}(e_1) \geq 2s(n)$.*

Proof. Recall from Construction 1.6(14) and Notation 1.5 that

$$Z_n = (S^2)^{s(n)} \quad \text{and} \quad Z_n \subset \text{cone}(Z_n) = X_n \subset X_n \amalg Y_n.$$

Also recall the line bundle L and the projection p from Notation 1.8.

It follows from Lemma 1.10 that there is a projection $q \in M_{2r(n)}(C(Z_n))$ whose range is isomorphic to the section space of the $s(n)$ -dimensional vector bundle $L^{\times s(n)}$ and such that $q(b_n|_{Z_n})q = q$. Now $\|xex^* - b_n\| < \frac{1}{2}$ implies $\|q(xex^*|_{Z_n})q - q\| < \frac{1}{2}$. Since $e|_{Z_n}$ and $q|_{Z_n}$ are projections, it follows that $q|_{Z_n}$ is Murray–von Neumann equivalent to a subprojection of $e|_{Z_n} = e_1|_{Z_n}$. Therefore $\text{rank}(e_1|_{Z_n}) \geq 2s(n)$ by Lemma 1.9. So $\text{rank}(e_1) \geq 2s(n)$. □

Although not strictly needed for the sequel, we record the following.

Corollary 1.12. *Assume the notation and choices in parts (1), (3) (including $\kappa > \frac{1}{2}$), (7), (8), and (10) of Construction 1.1, and in Construction 1.6 (except part (19)) and the parts of Construction 1.1 referred to there. Then the algebra C satisfies $\text{rc}(C) \geq 2\kappa - 1 > 0$.*

Proof. Suppose $\rho < 2\kappa - 1$. We show that C does not have ρ -comparison. Choose $n \in \mathbb{Z}_{>0}$ such that $1/r(n) < 2\kappa - 1 - \rho$. Choose $M \in \mathbb{Z}_{\geq 0}$ such that $\rho + 1 < M/r(n) < 2\kappa$. Let $e \in M_\infty(C_n)$ be a trivial projection of rank M . By slight abuse of notation, we use $\Gamma_{m,n}$ to denote the amplified map from $M_\infty(C_n)$ to $M_\infty(C_m)$ as well. For $m > n$, the rank of $\Gamma_{m,n}(e)$ is $Mr(m)/r(n)$, and the choice of M guarantees that this rank is strictly less than $2s(m)$. Now, for any trace τ on C_m (and thus for any trace on C), and justifying the last step afterwards, we have

$$d_\tau(\Gamma_{m,n}(e)) = \tau(\Gamma_{m,n}(e)) = \frac{1}{r(m)} \cdot M \cdot \frac{r(m)}{r(n)} \geq 1 + \rho > d_\tau(b_m) + \rho.$$

To explain the last step, recall b_m from Notation 1.8, and use Lemma 1.10 to see that the ranks of its components $(b_m)_1 \in M_2(C(X_m, M_{r(m)}))$ and $(b_m)_2 \in M_2(C(Y_m, M_{r(m)}))$ are both less than $r(m)$, while the identity element has rank $r(m)$.

On the other hand, if $\Gamma_{\infty,0}(b) \precsim \Gamma_{\infty,n}(e)$ then, in particular, there exists some $m > n$ and $x \in M_\infty(C_m)$ such that $\|x\Gamma_{m,n}(e)x^* - b_m\| < \frac{1}{2}$, which contradicts Corollary 1.11. □

Notation 1.13. We assume the notation and choices in parts (1), (6), (7), (8), and (10) of Construction 1.1. In particular, $C_0 = C(X_0) \oplus C(Y_0)$. Define $q_0 = (1, 0) \in C(X_0) \oplus C(Y_0)$ and $q_0^\perp = 1 - q_0$. For $n \in \mathbb{Z}_{>0}$ define $q_n = \Gamma_{n,0}(q_0) \in C_n$ and $q_n^\perp = 1 - q_n$, and finally, define $q = \Gamma_{\infty,0}(q_0) \in C$ and $q^\perp = 1 - q$.

Lemma 1.14. *Make the assumptions in Notation 1.13. Further assume the notation and choices in Construction 1.6 (except part (19)). Then the projection*

$$1 - q_n \in M_{l(n)}(C(X_n)) \oplus M_{l(n)}(C(Y_n))$$

has the form (e, f) for a constant projection $e \in M_{l(n)}(C(X_n)) = C(X_n, M_{l(n)})$ of rank $t(n)$ and a constant projection $f \in M_{l(n)}(C(Y_n)) = C(Y_n, M_{l(n)})$ of rank $r(n) - t(n)$.

From Construction 1.6, we don't actually need to know anything about the spaces X_n and Y_n , we don't need to know anything about the points x_n and y_n except which spaces they are in, and we don't need to know anything about the maps $Q_j^{(n)}$ and $R_{n,j}$ except their domains and codomains.

Proof of Lemma 1.14. The proof is an easy induction argument, using the fact that the image of a constant function under a diagonal map is again a constant function. \square

Lemma 1.15. *Assume the notation and choices in parts (1)–(10) of Construction 1.1, Construction 1.6 (except part (19)), and Notation 1.13, including $k(n) < d(n)$ for all $n \in \mathbb{Z}_{\geq 0}$, $\kappa > \frac{1}{2}$, $\omega > \omega'$, and $2\kappa - 1 > 2\omega$. Then*

$$\text{rc}(q^\perp C q^\perp) \geq \frac{2\kappa - 1}{2\omega}.$$

Proof. We proceed as in the proof of Corollary 1.12, although the rank computations are somewhat more involved. The difference is in the definition of d_τ . In this corner, d_τ is normalized so that $d_\tau(q^\perp) = 1$ for all $\tau \in \text{T}(C)$. To avoid redefining the notation, we will use τ to denote a tracial state on C , and therefore our dimension functions will be of the form $a \mapsto d_\tau(a)/\tau(q^\perp)$, noting that $\tau(q^\perp) = d_\tau(q^\perp)$ since q^\perp is a projection.

It suffices to show that for all $\rho \in (1, (2\kappa - 1)/(2\omega)) \cap \mathbb{Q}$, we have $\text{rc}(q^\perp C q^\perp) \geq \rho$.

Fix $\delta \in (0, \omega)$ such that

$$\rho < (1 - \delta) \left(\frac{2\kappa - 1}{2\omega} \right). \tag{1-5}$$

Set

$$\varepsilon = \frac{\delta}{2\rho(1 - \delta)} > 0. \tag{1-6}$$

Since the sequence $(s(n)/r(n))_{n=0,1,2,\dots}$ is nonincreasing and converges to a nonzero limit κ , there exists $n_0 \in \mathbb{Z}_{\geq 0}$ such that, for all n and m with $m \geq n \geq n_0$, we have

$$0 \leq 1 - \frac{r(n)}{s(n)} \cdot \frac{s(m)}{r(m)} < \varepsilon.$$

This implies that

$$\frac{r(m)}{r(n)} - \frac{s(m)}{s(n)} < \varepsilon \cdot \frac{r(m)}{r(n)}. \tag{1-7}$$

Using (1-5) and $\delta < \omega$ at the first step, we get

$$1 - \omega + 2\rho\omega < 1 - \delta + 2(1 - \delta) \left(\frac{2\kappa - 1}{2\omega} \right) \omega = 2\kappa(1 - \delta).$$

Now write $\rho = \alpha/\beta$ with $\alpha, \beta \in \mathbb{Z}_{>0}$. Choose $n \geq n_0$ such that

$$\frac{\beta}{r(n)} < 2\kappa(1 - \delta) - (1 - \omega + 2\rho\omega).$$

Then there exists $N_1 \in \mathbb{Z}_{>0}$ such that $\rho N_1 \in \mathbb{Z}_{>0}$ and

$$2\kappa(1 - \delta) > \frac{N_1}{r(n)} > 1 - \omega + 2\rho\omega. \tag{1-8}$$

Set

$$N_2 = \rho N_1. \quad (1-9)$$

Using $\rho > 1$ at the last step, we have

$$\frac{N_2}{r(n)} = \frac{\rho N_1}{r(n)} > \rho(1 - \omega + 2\rho\omega) > \rho(1 - \omega) + 2\omega.$$

Now suppose $e \in M_\infty(C_n) = M_\infty(C(X_n) \oplus C(Y_n))$ is an ordered pair whose first component is a trivial projection on X_n of rank N_1 and whose second component is a (trivial) projection on Y_n of rank N_2 . Let $m > n$, and let f be the first component of $\Gamma_{m,n}(e)$; we estimate $\text{rank}(f)$. (The second component is a trivial projection over Y_m whose rank we don't care about.) Now f is the direct sum of $r(m)/r(n)$ trivial projections, coming from $C(X_n, M_{r(n)})$ and $C(Y_n, M_{r(n)})$. At least $s(m)/s(n)$ of these summands come from $C(X_n, M_{r(n)})$. So at most $r(m)/r(n) - s(m)/s(n)$ of these summands come from $C(Y_n, M_{r(n)})$. The summands coming from $C(X_n, M_{r(n)})$ have rank N_1 and the summands coming from $C(Y_n, M_{r(n)})$ have rank N_2 . Since $N_2 > N_1$, we get

$$\text{rank}(f) \leq \left(\frac{r(m)}{r(n)} - \frac{s(m)}{s(n)} \right) N_2 + \frac{s(m)}{s(n)} \cdot N_1 = \frac{r(m)}{r(n)} \cdot N_1 + \left(\frac{r(m)}{r(n)} - \frac{s(m)}{s(n)} \right) (N_2 - N_1).$$

Combining this with (1-7) at the first step, and using (1-9) at the second step, (1-6) at the third step, (1-8) at the fifth step, and Construction 1.1(3) at the sixth step, we get

$$\begin{aligned} \text{rank}(f) &< \frac{r(m)}{r(n)} \cdot (N_1 + \varepsilon N_2) = \frac{r(m)}{r(n)} \cdot (1 + \varepsilon\rho) \cdot N_1 \\ &= \frac{r(m)}{r(n)} \cdot \frac{2 - \delta}{2(1 - \delta)} \cdot N_1 < \frac{r(m)}{r(n)} \cdot \frac{N_1}{1 - \delta} < 2\kappa r(m) \leq 2s(m). \end{aligned}$$

So Corollary 1.11 implies that there is no $x \in M_\infty(C_m)$ for which $\|x\Gamma_{n,m}(e)x^* - b_m\| < \frac{1}{2}$. Since $m > n$ is arbitrary,

$$\Gamma_{\infty,n}(e) \not\prec b. \quad (1-10)$$

Now let τ be a trace on C , and restrict it to $C_n \cong M_{r(n)}(C(X_n) \oplus C(Y_n))$. Denote by tr the normalized trace on $M_{r(n)}$. There is a probability measure μ on $X_n \amalg Y_n$ such that $\tau(a) = \int_{X_n \amalg Y_n} \text{tr}(a) d\mu$ for all $a \in C_n$. Define $\lambda = \mu(X_n)$, so $1 - \lambda = \mu(Y_n)$. Then, using (1-9) at the second step,

$$\tau(e) = \frac{\lambda N_1 + (1 - \lambda)N_2}{r(n)} = \frac{[\lambda + \rho(1 - \lambda)]N_1}{r(n)}.$$

Using Lemma 1.14 to calculate the ranks of the components of q_n^\perp , we get

$$\tau(q_n^\perp) = \frac{\lambda t(n) + (1 - \lambda)[r(n) - t(n)]}{r(n)}, \quad (1-11)$$

$$\tau(q_n) = 1 - \tau(q_n^\perp) = \frac{\lambda[r(n) - t(n)] + (1 - \lambda)t(n)}{r(n)}. \quad (1-12)$$

It follows from Lemmas 1.10 and 1.14 that $d_\tau(b_n) \leq \tau(q_n)$. Using this at the first step, and (1-11) and (1-12) at the second step, we get

$$\frac{d_\tau(b_n)}{\tau(q_n^\perp)} \leq \frac{\tau(q_n)}{\tau(q_n^\perp)} = \frac{\lambda[r(n) - t(n)] + (1 - \lambda)t(n)}{\lambda t(n) + (1 - \lambda)[r(n) - t(n)]}.$$

So

$$\frac{\tau(e) - d_\tau(b_n)}{\tau(q_n^\perp)} \geq \frac{(\lambda + \rho(1 - \lambda))N_1 - (\lambda[r(n) - t(n)] + (1 - \lambda)t(n))}{\lambda t(n) + (1 - \lambda)[r(n) - t(n)]}.$$

The last expression is a fractional linear function in λ and is defined for all values of λ in the interval $[0, 1]$. Any such function is monotone on $[0, 1]$. In the following calculations, we recall from Lemma 1.4 that $\omega \leq t(n)/r(n) < 2\omega$. If we set $\lambda = 1$ and use (1-8), the value we obtain is

$$\frac{N_1/r(n) - (1 - t(n)/r(n))}{t(n)/r(n)} > \frac{(1 - \omega + 2\rho\omega) - (1 - \omega)}{2\omega} = \rho.$$

If we set $\lambda = 0$, we get, using (1-8) at the first step and $\rho > 1$ at the last step,

$$\frac{\rho N_1/r(n) - t(n)/r(n)}{1 - t(n)/r(n)} > \frac{\rho(1 - \omega + 2\rho\omega) - 2\omega}{1 - \omega} = \rho + \frac{2\rho^2\omega - 2\omega}{1 - \omega} > \rho.$$

Therefore

$$\frac{d_\tau(\Gamma_{\infty,n}(e))}{d_\tau(q^\perp)} > \frac{d_\tau(b)}{d_\tau(q^\perp)} + \rho$$

for all traces τ on C , so $\text{rc}(q^\perp C q^\perp) > \rho$, as required. □

We now turn to the issue of finding upper bounds on the radius of comparison. For this, we appeal to results from [Niu 2014]. Niu [2014, Definition 3.6] introduced a notion of mean dimension for a diagonal AH-system. Suppose we are given a direct system of homogeneous algebras of the form

$$A_n = C(K_{n,1}) \otimes M_{j_{n,1}} \oplus C(K_{n,2}) \otimes M_{j_{n,2}} \oplus \cdots \oplus C(K_{n,m(n)}) \otimes M_{j_{n,m(n)}},$$

in which each of the spaces involved is a connected finite CW complex, and the connecting maps are unital diagonal maps. Let γ denote the mean dimension of this system, in the sense of Niu. It follows trivially from [Niu 2014, Definition 3.6] that

$$\gamma \leq \lim_{n \rightarrow \infty} \max \left(\left\{ \frac{\dim(K_{n,l})}{j_{n,l}} \mid l = 1, 2, \dots, m(n) \right\} \right).$$

Theorem 6.2 of [Niu 2014] states that if A is the direct limit of a system as above, and A is simple, then $\text{rc}(A) \leq \gamma/2$. Since the system we are considering here is of this type, Niu's theorem applies. With that at hand, we can derive an upper bound for the radius of comparison of the complementary corner.

Lemma 1.16. *Under the same assumptions as in Lemma 1.15, we have*

$$\text{rc}(qCq) \leq \frac{1}{1 - 2\omega}.$$

Proof. The algebra C is simple by Lemma 1.7, so qCq is also simple. This fact and Lemma 1.14 allow us to apply the discussion above, getting

$$\text{rc}(qCq) \leq \frac{1}{2} \lim_{n \rightarrow \infty} \max \left(\frac{\dim(X_n)}{\text{rank}(q_n|_{X_n})}, \frac{\dim(Y_n)}{\text{rank}(q_n|_{Y_n})} \right).$$

As $\dim(Y_n) = 1$ for all n , the second term converges to 0. As for the first term, by Construction 1.6(14), we have $\dim(X_n) = 2s(n) + 1$. Also, $\text{rank}(q_n|_{X_n}) = r(n) - t(n)$ by Lemma 1.14. Thus, by Construction 1.1(1) and Lemma 1.4, and using $d(n) \rightarrow \infty$ (which follows from Construction 1.1(4)) at the last step,

$$\lim_{n \rightarrow \infty} \frac{\dim(X_n)}{\text{rank}(q_n|_{X_n})} = \lim_{n \rightarrow \infty} \frac{2s(n) + 1}{r(n) - t(n)} \leq \lim_{n \rightarrow \infty} \frac{2r(n) + 1}{r(n) - t(n)} \leq \frac{2}{1 - 2\omega}.$$

This gives us the required estimate. \square

Lemma 1.17. *Let the assumptions and notation be as in Notation 1.13, Construction 1.6(14), and Construction 1.6(15). If $e \in C$ is a projection which has the same K_0 -class as q then e is unitarily equivalent to q . The same holds with q^\perp in place of q .*

Proof. This can be seen directly from the construction. For each $n \in \mathbb{Z}_{\geq 0}$, since X_n and Y_n are contractible (Constructions 1.6(14) and (15)), if $e \in M_\infty(C_n)$ is a projection which has the same K_0 -class as q , then e is actually unitarily equivalent to q_n . The same holds for q_n^\perp . It follows that this is the case in C as well. \square

We point out that this lemma can also be deduced using cancellation. By [Elliott et al. 2009, Theorem 4.1], simple unital AH algebras which arise from AH systems with diagonal maps have stable rank 1. Rieffel has shown that C^* -algebras with stable rank 1 have cancellation; see [Blackadar 1998, Theorem 6.5.1].

2. The tracial state space

For a compact Hausdorff space X , we will need all of $C(X, \mathbb{R})$ (the space of real-valued continuous functions on X), the tracial state space of $C(X)$ (and of $C(X, M_n)$), and the space of affine functions on the tracial state space. This last space is an order unit space, and much of our work will be done there.

For later reference, we recall some of the definitions, and then describe how to move between these spaces. We begin with the definition of an order unit space from the discussion before Proposition II.1.3 of [Alfsen 1971]. We suppress the order unit in our notation, since (except in several abstract results) our order unit spaces will always be sets of affine continuous functions on compact convex sets with order unit the constant function 1.

Definition 2.1. An *order unit space* V is a partially ordered real Banach space (see page 1 of [Goodearl 1986] for the axioms of a partially ordered real vector space) which is *Archimedean* (if $v \in V$ and $\{\lambda v \mid \lambda \in (0, \infty)\}$ has an upper bound, then $v \leq 0$), with a distinguished element $e \in V$ which is an *order unit* (that is, for every $v \in V$ there is $\lambda \in (0, \infty)$ such that $-\lambda e \leq v \leq \lambda e$), and such that the norm on V satisfies

$$\|v\| = \inf(\{\lambda \in (0, \infty) \mid -\lambda e \leq v \leq \lambda e\})$$

for all $v \in V$.

The morphisms of order unit spaces are the positive linear maps which preserve the order units.

The morphisms of compact convex sets (compact convex subsets of locally convex topological vector spaces) are just the continuous affine maps.

Definition 2.2. If K is a compact convex set, we denote by $\text{Aff}(K)$ the order unit space of continuous affine functions $f : K \rightarrow \mathbb{R}$, with the supremum norm and with order unit the constant function 1.

If K and L are compact convex sets and $\lambda : K \rightarrow L$ is continuous and affine, we let $\lambda^* : \text{Aff}(L) \rightarrow \text{Aff}(K)$ be the positive linear order unit preserving map given by $\lambda^*(f) = f \circ \lambda$ for $f \in \text{Aff}(L)$.

This definition makes $K \mapsto \text{Aff}(K)$ a functor.

Definition 2.3. If V is an order unit space with order unit e , we denote by $S(V)$ (or $S(V, e)$ if e is not understood) its state space (the order unit space morphisms to $(\mathbb{R}, 1)$), which is a compact convex set with the weak* topology.

If W is another order unit space and $\varphi : V \rightarrow W$ is positive, linear, and order unit preserving, we let $S(\varphi) : S(W) \rightarrow S(V)$ be the continuous affine map given by $S(\varphi)(\omega) = \omega \circ \varphi$ for $\omega \in S(W)$.

This definition makes $V \mapsto S(V)$ a functor.

Theorem 2.4 [Goodearl 1986, Theorem 7.1]. *There is a natural isomorphism $S(\text{Aff}(K)) \cong K$ for compact convex sets K , given by sending $x \in K$ to the evaluation map $\text{ev}_x : \text{Aff}(K) \rightarrow \mathbb{R}$ defined by $\text{ev}_x(f) = f(x)$ for $f \in \text{Aff}(K)$.*

Definition 2.5. For a unital C^* -algebra A , we denote its tracial state space by $T(A)$. If A and B are unital C^* -algebras and $\varphi : A \rightarrow B$ is a unital homomorphism, we let $T(\varphi) : T(B) \rightarrow T(A)$ be the continuous affine map given by $T(\varphi)(\tau) = \tau \circ \varphi$ for $\tau \in T(B)$. We let $\hat{\varphi} : \text{Aff}(T(A)) \rightarrow \text{Aff}(T(B))$ be the positive order unit preserving map given by $\hat{\varphi}(f) = f \circ T(\varphi)$ for $f \in \text{Aff}(T(A))$. (Thus, $\hat{\varphi} = T(\varphi)^*$.)

Lemma 2.6. *Let X be a compact Hausdorff space. Then $C(X, \mathbb{R})$, with the supremum norm and distinguished element the constant function 1, is a complete order unit space. Restriction of tracial states on $C(X)$ is an affine homeomorphism from $T(C(X))$ to $S(C(X, \mathbb{R}))$. The map from X to $S(C(X, \mathbb{R}))$ which sends $x \in X$ to the point evaluation $\text{ev}_x : C(X, \mathbb{R}) \rightarrow \mathbb{R}$ is a homeomorphism onto its image, and the map $R_X : \text{Aff}(S(C(X, \mathbb{R}))) \rightarrow C(X, \mathbb{R})$, given by $R_X(f)(x) = f(\text{ev}_x)$ for $f \in \text{Aff}(S(C(X, \mathbb{R})))$ and $x \in X$, is an isomorphism of order unit spaces.*

If Y is another compact Hausdorff space, then the function which sends a positive linear order unit preserving map $Q : C(X, \mathbb{R}) \rightarrow C(Y, \mathbb{R})$ to $S(Q) : S(C(Y, \mathbb{R})) \rightarrow S(C(X, \mathbb{R}))$, as in Definition 2.3, is a bijection to the continuous affine maps from $S(C(Y, \mathbb{R}))$ to $S(C(X, \mathbb{R}))$. Its inverse is the map E given as follows. For a continuous affine map $\lambda : S(C(Y, \mathbb{R})) \rightarrow S(C(X, \mathbb{R}))$, using the notation of Definition 2.2, define $E(\lambda) : C(X, \mathbb{R}) \rightarrow C(Y, \mathbb{R})$ by $E(\lambda) = R_Y \circ \lambda^ \circ R_X^{-1}$.*

A positive linear order unit preserving map from $C(X, \mathbb{R})$ to $C(Y, \mathbb{R})$ is called a *Markov operator*.

Proof of Lemma 2.6. It is immediate that $C(X, \mathbb{R})$ is a complete order unit space. The identification of $S(C(X, \mathbb{R}))$ is also immediate. The fact that R_X is bijective follows from [Goodearl 1986, Corollary 11.20] using the identification of X with the extreme points of $S(C(X, \mathbb{R}))$.

For the second paragraph, it is immediate that S sends positive linear order unit preserving maps to continuous affine maps, and that E does the reverse. For the rest, we must show that $S \circ E$ and $E \circ S$ are the identity maps on the appropriate sets.

We first claim that for $g \in \text{Aff}(S(C(X, \mathbb{R})))$ and $\rho \in S(C(X, \mathbb{R}))$ we have

$$g(\rho) = \rho(R_X(g)). \quad (2-1)$$

This formula is true by definition when $\rho = \text{ev}_x$ for some $x \in X$. Since, for fixed g , both sides of (2-1) are continuous affine functions of ρ , and since $S(C(X, \mathbb{R}))$ is the closed convex hull of $\{\text{ev}_x \mid x \in X\}$, the claim follows.

We next claim that if $\lambda : S(C(Y, \mathbb{R})) \rightarrow S(C(X, \mathbb{R}))$ is continuous and affine, $\omega \in S(C(Y, \mathbb{R}))$, and $g \in \text{Aff}(S(C(X, \mathbb{R})))$, then

$$(\omega \circ R_Y)(g \circ \lambda) = (\lambda(\omega) \circ R_X)(g). \quad (2-2)$$

To prove this claim, for the same reasons as in the proof of the first claim, it suffices to prove this when there is $y \in Y$ such that $\omega = \text{ev}_y$. In this case, using the definition of R_Y at the second step, and the previous claim with $\rho = \lambda(\text{ev}_y)$ at the third step,

$$(\text{ev}_y \circ R_Y)(g \circ \lambda) = R_Y(g \circ \lambda)(y) = (g \circ \lambda)(\text{ev}_y) = (\lambda(\text{ev}_y) \circ R_X)(g),$$

as desired.

Now let $\lambda : S(C(Y, \mathbb{R})) \rightarrow S(C(X, \mathbb{R}))$ be continuous and affine; we prove that $S(E(\lambda)) = \lambda$. Let $\omega \in S(C(X, \mathbb{R}))$ and let $f \in C(Y, \mathbb{R})$. Working through the definitions gives

$$S(E(\lambda))(\omega)(f) = (\omega \circ R_Y)(R_X^{-1}(f) \circ \lambda).$$

By (2-2) with $g = R_X^{-1}(f)$, the right-hand side is $\lambda(\omega)(f)$, as desired.

Finally, let $Q : C(X, \mathbb{R}) \rightarrow C(Y, \mathbb{R})$ be a positive linear order unit preserving map; we show that $E(S(Q)) = Q$. Let $f \in C(X, \mathbb{R})$ and let $y \in Y$. Working through the definitions gives

$$E(S(Q))(f)(y) = R_X^{-1}(f)(\text{ev}_y \circ Q).$$

Applying (2-1) with $g = R_X^{-1}(f)$ and $\rho = \text{ev}_y \circ Q$, we see that the right-hand side is $(\text{ev}_y \circ Q)(f) = Q(f)(y)$. This proves that $E(S(Q)) = Q$, and the proof is complete. \square

Direct limits of direct systems of order unit spaces are constructed at the beginning of Section 3 of [Thomsen 1994], including Lemma 3.1 there.

Proposition 2.7. *Let $((D_n)_{n=0,1,2,\dots}, (\varphi_{n,m})_{0 \leq m \leq n})$ be a direct system of unital C^* -algebras and unital homomorphisms. Set $D = \varinjlim_n D_n$. Then there are a natural homeomorphism*

$$T(D) \rightarrow \varprojlim_n T(D_n)$$

and a natural isomorphism

$$\text{Aff}(T(D)) \rightarrow \varinjlim_n \text{Aff}(T(D_n))$$

of order unit spaces.

Proof. The first part is Lemma 3.3 of [Thomsen 1994].

The second part is Lemma 3.2 of [Thomsen 1994], combined with the fact (Theorem 2.4) that the state space of $\text{Aff}(K)$ is naturally identified with K . \square

Definition 2.8. Let V and W be order unit spaces, with order units $e \in V$ and $f \in W$. We define the *direct sum* $V \oplus W$ to be the vector space direct sum $V \oplus W$ as a real vector space, with the order $(v_1, w_1) \leq (v_2, w_2)$ for $v_1, v_2 \in V$ and $w_1, w_2 \in W$ if and only if $v_1 \leq v_2$ and $w_1 \leq w_2$, with the order unit (e, f) , and the norm $\|(v, w)\| = \max(\|v\|, \|w\|)$.

Lemma 2.9. *Let V and W be order unit spaces. Then $V \oplus W$ as in Definition 2.8 is an order unit space, which is complete if V and W are.*

Proof. The proof is straightforward. \square

Lemma 2.10. *Let A and B be unital C^* -algebras. Then, taking the direct sum on the right to be as in Definition 2.8, there is an isomorphism*

$$\text{Aff}(T(A \oplus B)) \cong \text{Aff}(T(A)) \oplus \text{Aff}(T(B)),$$

given as follows. Identify $T(A)$ with a subset of $T(A \oplus B)$ by, for $\tau \in T(A)$, defining $i(\tau)(a, b) = \tau(a)$ for all $a \in A$ and $b \in B$, and similarly identify $T(B)$ with a subset of $T(A \oplus B)$. Then the map $\text{Aff}(T(A \oplus B)) \rightarrow \text{Aff}(T(A)) \oplus \text{Aff}(T(B))$ is $f \mapsto (f|_{T(A)}, f|_{T(B)})$.

Proof. It is clear that if $f \in \text{Aff}(T(A \oplus B))$, then $f|_{T(A)} \in \text{Aff}(T(A))$ and $f|_{T(B)} \in \text{Aff}(T(B))$, and moreover that the map of the lemma is linear, positive, and preserves the order units. One easily checks that every tracial state on $A \oplus B$ is a convex combination of tracial states on A and B , from which it follows that if $f|_{T(A)} = 0$ and $f|_{T(B)} = 0$ then $f = 0$.

It remains to prove that the map of the lemma is surjective. Let $g \in \text{Aff}(T(A))$ and $h \in \text{Aff}(T(B))$. Define $f : T(A \oplus B) \rightarrow \mathbb{R}$ by, for $\tau \in T(A \oplus B)$,

$$f(\tau) = \tau(1, 0)g(\tau(1, 0)^{-1}\tau|_A) + \tau(0, 1)g(\tau(0, 1)^{-1}\tau|_B)$$

(taking the first summand to be zero if $\tau(1, 0) = 0$ and the second summand to be zero if $\tau(0, 1) = 0$). Straightforward but somewhat tedious calculations show that f is weak* continuous and affine, and clearly $f|_{T(A)} = g$ and $f|_{T(B)} = h$. \square

The following result generalizes Lemma 3.4 of [Thomsen 1994]. It still isn't the most general Elliott approximate intertwining result for order unit spaces, because we assume that the underlying order unit spaces of the two direct systems are the same. The main effect of this assumption is to simplify the notation.

Proposition 2.11. *Let $(V_m)_{m=0,1,2,\dots}$ be a sequence of separable complete order unit spaces, and let*

$$((V_m)_{m=0,1,2,\dots}, (\varphi_{n,m})_{0 \leq m \leq n}) \quad \text{and} \quad ((V_m)_{m=0,1,2,\dots}, (\varphi'_{n,m})_{0 \leq m \leq n})$$

be two direct systems of order unit spaces, using the same spaces, and with maps $\varphi_{n,m}, \varphi'_{n,m} : V_m \rightarrow V_n$ which are linear, positive, and preserve the order units. Let V and V' be the direct limits

$$V = \varinjlim ((V_m)_{m=0,1,2,\dots}, (\varphi_{n,m})_{0 \leq m \leq n}) \quad \text{and} \quad V' = \varinjlim ((V_m)_{m=0,1,2,\dots}, (\varphi'_{n,m})_{0 \leq m \leq n}),$$

with corresponding maps

$$\varphi_{\infty,n} : V_n \rightarrow V \quad \text{and} \quad \varphi'_{\infty,n} : V_n \rightarrow V'$$

for $n \in \mathbb{Z}_{\geq 0}$. For $n \in \mathbb{Z}_{\geq 0}$ further let

$$v_0^{(n)}, v_1^{(n)}, \dots \in V_n$$

be a dense sequence in the closed unit ball of V_n , and define $F_n \subset V_n$ to be the finite set

$$F_n = \bigcup_{m=0}^n [\{\varphi_{n,m}(v_k^{(m)}) : 0 \leq k \leq n\} \cup \{\varphi'_{n,m}(v_k^{(m)}) : 0 \leq k \leq n\}].$$

Suppose that there are $\delta_0, \delta_1, \dots \in (0, \infty)$ such that

$$\sum_{n=0}^{\infty} \delta_n < \infty \tag{2-3}$$

and for all $n \in \mathbb{Z}_{\geq 0}$ and all $v \in F_n$ we have

$$\|\varphi_{n+1,n}(v) - \varphi'_{n+1,n}(v)\| < \delta_n.$$

Then there is a unique isomorphism $\rho : V \rightarrow V'$ such that for all $m \in \mathbb{Z}_{\geq 0}$ and all $v \in V_m$ we have

$$\rho(\varphi_{\infty,m}(v)) = \lim_{n \rightarrow \infty} (\varphi'_{\infty,n} \circ \varphi_{n,m})(v).$$

Its inverse is determined by

$$\rho^{-1}(\varphi'_{\infty,m}(v)) = \lim_{n \rightarrow \infty} (\varphi_{\infty,n} \circ \varphi'_{n,m})(v)$$

for $m \in \mathbb{Z}_{\geq 0}$ and $v \in V_m$.

Proof. We first claim that for $m \in \mathbb{Z}_{\geq 0}$ and $v \in F_m$, the sequence $((\varphi'_{\infty,n} \circ \varphi_{n,m})(v))_{n \geq m}$ is a Cauchy sequence in V' . For $n \geq m$, we estimate, using $\|\varphi'_{\infty,n+1}\| \leq 1$, $\|v\| \leq 1$, and $\varphi_{n,m}(v) \in F_n$ at the last step:

$$\begin{aligned} \|(\varphi'_{\infty,n+1} \circ \varphi_{n+1,m})(v) - (\varphi'_{\infty,n} \circ \varphi_{n,m})(v)\| &= \|(\varphi'_{\infty,n+1} \circ \varphi_{n+1,n} \circ \varphi_{n,m})(v) - (\varphi'_{\infty,n+1} \circ \varphi'_{n+1,n} \circ \varphi_{n,m})(v)\| \\ &\leq \|\varphi'_{\infty,n+1}\| \|\varphi_{n+1,n}(\varphi_{n,m}(v)) - \varphi'_{n+1,n}(\varphi_{n,m}(v))\| \leq \delta_n. \end{aligned}$$

The claim now follows from (2-3).

Next, we claim that for $m \in \mathbb{Z}_{\geq 0}$ and $k \in \mathbb{Z}_{>0}$, the sequence $((\varphi'_{\infty,n} \circ \varphi_{n,m})(v_k^{(m)}))_{n \geq m}$ is a Cauchy sequence in V' . Indeed, taking $m_0 = \max(m, k)$, this follows from the previous claim and the fact that $\varphi_{m_0,m}(v_k^{(m)}) \in F_{m_0}$.

Now we claim that for $m \in \mathbb{Z}_{\geq 0}$ and $v \in V_m$, the sequence $((\varphi'_{\infty,n} \circ \varphi_{n,m})(v))_{n \geq m}$ is a Cauchy sequence in V' . Without loss of generality $\|v\| \leq 1$. This claim follows from a standard $\varepsilon/3$ argument: to show that

$$\|(\varphi'_{\infty,n_1} \circ \varphi_{n_1,m})(v) - (\varphi'_{\infty,n_2} \circ \varphi_{n_2,m})(v)\| < \varepsilon$$

for all sufficiently large n_1 and n_2 , choose $k \in \mathbb{Z}_{>0}$ such that $\|v - v_k^{(m)}\| < \varepsilon/3$, and use the previous claim.

Since V' is complete, it follows that $\lim_{n \rightarrow \infty} (\varphi'_{\infty,n} \circ \varphi_{n,m})(v)$ exists for all $m \in \mathbb{Z}_{\geq 0}$ and $v \in V_m$. Since $\|\varphi'_{\infty,n} \circ \varphi_{n,m}\| \leq 1$ whenever $m, n \in \mathbb{Z}_{\geq 0}$ satisfy $m \leq n$, it follows that for $m \in \mathbb{Z}_{>0}$ there is a unique bounded linear map $\rho_m : V_m \rightarrow V'$ such that $\|\rho_m\| \leq 1$ and $\rho_m(v) = \lim_{n \rightarrow \infty} (\varphi'_{\infty,n} \circ \varphi_{n,m})(v)$ for all $v \in V_m$.

It is clear from the construction that $\rho_n \circ \varphi_{n,m} = \rho_m$ whenever $m, n \in \mathbb{Z}_{\geq 0}$ satisfy $m \leq n$. By the universal property of the direct limit, there is a unique bounded linear map $\rho : V \rightarrow V'$ such that $\rho \circ \varphi_{\infty,m} = \rho_m$ for all $m \in \mathbb{Z}_{\geq 0}$. It is clearly contractive, order preserving, order unit preserving, and uniquely determined as in the statement of the proposition.

The same argument shows that there is a unique contractive linear map $\lambda : V' \rightarrow V$ determined in the analogous way. For all $m \in \mathbb{Z}_{\geq 0}$, we have

$$\lambda \circ \rho \circ \varphi_{\infty,m} = \lambda \circ \varphi'_{\infty,m} = \varphi_{\infty,m},$$

so the universal property of the direct limit implies $\lambda \circ \rho = \text{id}_V$. Similarly $\rho \circ \lambda = \text{id}_{V'}$. □

Proposition 2.12. *The isomorphism of Proposition 2.11 has the following naturality property. Let the notation be as there, and suppose that, in addition, we are given separable complete order unit spaces W_n for $n \in \mathbb{Z}_{\geq 0}$, direct systems*

$$((W_m)_{m=0,1,2,\dots}, (\psi_{n,m})_{0 \leq m \leq n}) \quad \text{and} \quad ((W_m)_{m=0,1,2,\dots}, (\psi'_{n,m})_{0 \leq m \leq n})$$

using the same spaces, with positive linear order unit preserving maps, with direct limits W and W' , and with corresponding maps

$$\psi_{\infty,n} : W_n \rightarrow W \quad \text{and} \quad \psi'_{\infty,n} : W_n \rightarrow W'$$

for $n \in \mathbb{Z}_{\geq 0}$. Also suppose that for $n \in \mathbb{Z}_{> 0}$ there is a sequence

$$w_0^{(n)}, w_1^{(n)}, \dots \in W_n$$

which is dense in the closed unit ball of W_n , and that there is a sequence $(\varepsilon_n)_{n=0,1,2,\dots}$ in $(0, \infty)$ such that $\sum_{n=0}^{\infty} \varepsilon_n < \infty$ and, with

$$G_n = \bigcup_{m=0}^n [\{\psi_{n,m}(w_k^{(m)}) \mid 0 \leq k \leq n\} \cup \{\psi'_{n,m}(w_k^{(m)}) \mid 0 \leq k \leq n\}],$$

for all $n \in \mathbb{Z}_{\geq 0}$ and all $w \in G_n$ we have

$$\|\psi_{n+1,n}(w) - \psi'_{n+1,n}(w)\| < \varepsilon_n.$$

Let $\sigma : W \rightarrow W'$ be the isomorphism of Proposition 2.11. Suppose further that we have positive linear order unit preserving maps $\mu_n, \mu'_n : V_n \rightarrow W_n$ for $n \in \mathbb{Z}_{\geq 0}$ such that

$$\mu_n \circ \varphi_{n,m} = \psi_{n,m} \circ \mu_m \quad \text{and} \quad \mu'_n \circ \varphi'_{n,m} = \psi'_{n,m} \circ \mu'_m$$

for all $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$. Let $\mu : V \rightarrow W$ and $\mu' : V' \rightarrow W'$ be the induced maps of the direct limits. Then $\mu' \circ \rho = \sigma \circ \mu$.

Proof. By construction, $\rho : V \rightarrow V'$ and $\sigma : W \rightarrow W'$ are determined by

$$\rho(\varphi_{\infty,m}(v)) = \lim_{n \rightarrow \infty} (\varphi'_{\infty,n} \circ \varphi_{n,m})(v) \tag{2-4}$$

for $m \in \mathbb{Z}_{\geq 0}$ and $v \in V_m$, and

$$\sigma(\psi_{\infty,m}(w)) = \lim_{n \rightarrow \infty} (\psi'_{\infty,n} \circ \psi_{n,m})(w) \tag{2-5}$$

for $m \in \mathbb{Z}_{> 0}$ and $w \in W_m$. Using (2-4) at the first step and (2-5) at the last step, for $m \in \mathbb{Z}_{\geq 0}$ and $v \in V_m$ we therefore have

$$\begin{aligned} (\mu' \circ \rho)(\varphi_{\infty,m}(v)) &= \mu' \left(\lim_{n \rightarrow \infty} (\varphi'_{\infty,n} \circ \varphi_{n,m})(v) \right) = \lim_{n \rightarrow \infty} (\mu' \circ \varphi'_{\infty,n} \circ \varphi_{n,m})(v) \\ &= \lim_{n \rightarrow \infty} (\psi'_{\infty,n} \circ \psi_{n,m} \circ \mu_m)(v) = (\sigma \circ \mu)(\varphi_{\infty,m}(v)). \end{aligned}$$

Since $\bigcup_{m=0}^{\infty} \varphi_{\infty,m}(V_m)$ is dense in V , the result follows. □

Proposition 2.14 below can essentially be extracted from the proof of Lemma 3.7 of [Thomsen 1994]. We give here a precise formulation which is needed for our purposes. The difference between our formulation and that of [Thomsen 1994] is that we need more control over the matrix sizes in the construction. In the argument, the following result substitutes for Lemma 3.6 there.

Lemma 2.13 (based on [Thomsen 1994]). *Let X and Y be compact Hausdorff spaces, with X path connected. Let $\lambda : T(C(Y)) \rightarrow T(C(X))$ be affine and continuous. Let $E(\lambda) : C(X, \mathbb{R}) \rightarrow C(Y, \mathbb{R})$ be as in Lemma 2.6. Then for every $\varepsilon > 0$ and every finite set $F \subset C(X, \mathbb{R})$ there exists $N_0 \in \mathbb{Z}_{> 0}$ such that for every $N \in \mathbb{Z}_{> 0}$ with $N \geq N_0$ there are continuous functions $g_1, g_2, \dots, g_N : Y \rightarrow X$ such that for every $f \in F$ we have*

$$\left\| E(\lambda)(f) - \frac{1}{N} \sum_{j=1}^N f \circ g_j \right\|_{\infty} < \varepsilon.$$

Proof. It suffices to prove the result under the additional assumption that $\|f\| \leq 1$ for all $f \in F$.

Let $\varepsilon > 0$. Since $E(\lambda)$ is a Markov operator, Theorem 2.1 of [Thomsen 1994] provides $n \in \mathbb{Z}_{> 0}$, unital homomorphisms $\psi_1, \psi_2, \dots, \psi_n : C(X) \rightarrow C(Y)$, and $\alpha_1, \alpha_2, \dots, \alpha_n \in [0, 1]$ with $\sum_{l=1}^n \alpha_l = 1$ such that

$$\left\| E(\lambda)(f) - \sum_{l=1}^n \alpha_l \psi_l(f) \right\|_{\infty} < \frac{\varepsilon}{2}$$

for all $f \in F$. Note that if $\beta_1, \beta_2, \dots, \beta_n \in [0, 1]$ satisfy $\sum_{l=1}^n |\alpha_l - \beta_l| < \varepsilon/2$ then

$$\left\| E(\lambda)(f) - \sum_{l=1}^n \beta_l \psi_l(f) \right\|_{\infty} < \varepsilon$$

for all $f \in F$. Choose $N_0 \in \mathbb{Z}_{> 0}$ such that $N_0 > 4n/\varepsilon$. Let $N \in \mathbb{Z}_{> 0}$ satisfy $N \geq N_0$. For $l = 1, 2, \dots, n-1$ choose $\beta_l \in (\alpha_l - 1/N, \alpha_l] \cap (1/N)\mathbb{Z}$, and set $\beta_n = 1 - \sum_{l=1}^{n-1} \beta_l$. Then

$$\beta_1, \beta_2, \dots, \beta_n \in \frac{1}{N}\mathbb{Z}_{\geq 0}, \quad \sum_{l=1}^n \beta_l = 1, \quad \text{and} \quad \sum_{l=1}^n |\alpha_l - \beta_l| < \frac{\varepsilon}{2}.$$

Set $m_l = N\beta_l$ for $l = 1, 2, \dots, n$. Then for all $f \in F$ we have

$$\left\| E(\lambda)(f) - \frac{1}{N} \sum_{l=1}^n m_l \psi_l(f) \right\|_{\infty} < \varepsilon.$$

Now for $l = 1, 2, \dots, n$ let $h_l : Y \rightarrow X$ be the continuous function such that $\psi_l(f) = f \circ h_l$ for all $f \in C(X)$, and for $j = 1, 2, \dots, N$ define $g_j = h_l$ when

$$\sum_{k=1}^{l-1} m_k < j \leq \sum_{k=1}^l m_k.$$

Then

$$\frac{1}{N} \sum_{l=1}^n m_l \psi_l(f) = \frac{1}{N} \sum_{j=1}^N f \circ g_j$$

for all $f \in C(X)$. □

Proposition 2.14. *Let K be a metrizable Choquet simplex, and let $(l(n))_{n=0,1,2,\dots}$ be a sequence of integers such that $l(n) \geq 2$ for all $n > 0$. For $n \in \mathbb{Z}_{\geq 0}$ set $r(n) = \prod_{j=1}^n l(j)$. Then there exist $n_0 < n_1 < n_2 < \dots \in \mathbb{Z}_{\geq 0}$, with $n_0 = 0$ and $n_1 = 1$, and a direct system*

$$C([0, 1]) \otimes M_{r(n_0)} \xrightarrow{\alpha_{1,0}} C([0, 1]) \otimes M_{r(n_1)} \xrightarrow{\alpha_{2,1}} C([0, 1]) \otimes M_{r(n_2)} \xrightarrow{\alpha_{3,2}} \dots$$

with injective maps which are diagonal (in the sense analogous to Construction 1.1 (9)) and such that the direct limit A satisfies $T(A) \cong K$.

It is easy to arrange that the algebra A in this proposition be simple: by Proposition 2.11, replacement of a small enough fraction of the maps $g_{k,l}$ in the proof with suitable point evaluations does not change the tracial state space. However, doing so at this stage does not help with later work.

The conditions $n_0 = 0$ and $n_1 = 1$ are needed because we will later need to pass to a corresponding subsystem of a system as in Construction 1.1 (more accurately, Construction 3.3 below), and we want to avoid later complexity of the argument by preserving the value of ω .

Proof of Proposition 2.14. We mostly follow the proof of Lemma 3.7 of [Thomsen 1994], using Lemma 2.13 in place of Lemma 3.6 of [Thomsen 1994], and slightly changing the order of the steps to accommodate the difference between our conclusion and that of Theorem 3.9 of [Thomsen 1994]. For convenience, we will use Proposition 2.11 in place of Lemma 3.4 of [Thomsen 1994].

For convenience of notation, and following [Thomsen 1994], set $P = T(C([0, 1]))$. Lemma 3.8 of [Thomsen 1994] provides an inverse system $((P_k)_{k=0,1,\dots}, (\lambda_{j,k})_{0 \leq j \leq k})$ with continuous affine maps $\lambda_{j,k} : P_k \rightarrow P_j$ such that $P_k = P$ for all $k \in \mathbb{Z}_{\geq 0}$ and

$$\varprojlim ((P_k)_{k=0,1,\dots}, (\lambda_{j,k})_{0 \leq j \leq k}) \cong K. \tag{2-6}$$

Choose $f_0, f_1, \dots \in C([0, 1], \mathbb{R})$ such that $\{f_0, f_1, \dots\}$ is dense in $C([0, 1], \mathbb{R})$.

We now construct numbers $n_k \in \mathbb{Z}_{>0}$ for $k \in \mathbb{Z}_{\geq 0}$, finite subsets $F_k \subset C([0, 1], \mathbb{R})$ for $k \in \mathbb{Z}_{\geq 0}$, positive unital linear maps $\psi_{k+1,k} : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ for $k \in \mathbb{Z}_{>0}$, and continuous functions

$$g_{k,1}, g_{k,2}, \dots, g_{k,r(n_{k+1})/r(n_k)} : [0, 1] \rightarrow [0, 1]$$

such that the following conditions are satisfied:

- (1) $F_0 = \{f_0\}$ and for $k \in \mathbb{Z}_{\geq 0}$,

$$F_{k+1} = F_k \cup \{f_{k+1}\} \cup E(\lambda_{k,k+1})(F_k \cup \{f_{k+1}\}) \cup \psi_{k+1,k}(F_k \cup \{f_{k+1}\}).$$

- (2) $n_0 = 0, n_1 = 1,$ and $n_2 = 2,$ and for $k \in \mathbb{Z}_{>0}$ with $k \geq 2$ we have $n_{k+1} > n_k$ and $r(n_{k+1})/r(n_k) > 2^k.$
- (3) For $k \in \mathbb{Z}_{\geq 0}$ and $f \in C([0, 1], \mathbb{R}),$

$$\psi_{k+1,k}(f) = \frac{r(n_k)}{r(n_{k+1})} \sum_{l=1}^{r(n_{k+1})/r(n_k)} f \circ g_{k,l}.$$

- (4) $\|E(\lambda_{k,k+1})(f) - \psi_{k+1,k}(f)\| < 2^{-k}$ for $k \geq 2$ and $f \in F_k.$

We carry out the construction by induction on $k.$ Define $F_0 = \{f_0\}, n_0 = 0,$ and $n_1 = 1.$ Take $g_{0,l} : [0, 1] \rightarrow [0, 1]$ to be the identity map for $l = 1, 2, \dots, r(1).$ Then define $\psi_{1,0}$ by (3) and define F_1 by (1). Now suppose $k \geq 1$ and we have F_k and $n_k;$ we construct

$$F_{k+1}, \quad n_{k+1}, \quad g_{k,1}, g_{k,2}, \dots, g_{k,r(n_{k+1})/r(n_k)}, \quad \text{and} \quad \psi_{k+1,k}.$$

Apply Lemma 2.13 with $\lambda = \lambda_{k,k+1},$ with $\varepsilon = 2^{-k},$ and with $F = F_k,$ obtaining $N_0 \in \mathbb{Z}_{>0}.$ Choose $n_{k+1} > n_k$ and so large that

$$\frac{r(n_{k+1})}{r(n_k)} > \max(N_0, 2^k).$$

This gives (2). Apply the conclusion of Lemma 2.13 with $N = r(n_{k+1})/r(n_k),$ calling the resulting functions $g_{k,1}, g_{k,2}, \dots, g_{k,r(n_{k+1})/r(n_k)}.$ Then define $\psi_{k+1,k}$ by (3). This gives (4). Finally, define F_{k+1} by (1). This completes the induction.

For $j, k \in \mathbb{Z}_{\geq 0}$ with $j \leq k,$ define $\psi_{k,j} : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ by

$$\psi_{k,j} = \psi_{k,k-1} \circ \psi_{k-1,k-2} \circ \dots \circ \psi_{j+1,j}.$$

An induction argument shows that for $j, k \in \mathbb{Z}_{\geq 0}$ with $j \leq k,$ we have

$$E(\lambda_{j,k})(f_j) \in F_k \quad \text{and} \quad \psi_{k,j}(f_j) \in F_k.$$

This condition, together with Proposition 2.11, allows us to conclude that, as order unit spaces, we have

$$\varinjlim((C([0, 1], \mathbb{R}))_{k=0,1,\dots}, (E(\lambda_{j,k}))_{0 \leq j \leq k}) \cong \varinjlim((C([0, 1], \mathbb{R}))_{k=0,1,\dots}, (\psi_{k,j})_{0 \leq j \leq k}). \tag{2-7}$$

For $k \in \mathbb{Z}_{\geq 0}$ define

$$\alpha_{k+1,k} : C([0, 1], M_{r(n_k)}) \rightarrow C([0, 1], M_{r(n_{k+1})}) = M_{r(n_{k+1})/r(n_k)}(C([0, 1], M_{r(n_k)}))$$

by

$$\alpha_{k+1,k}(f) = \text{diag}(f \circ g_{k,1}, f \circ g_{k,2}, \dots, f \circ g_{k,r(n_{k+1})/r(n_k)})$$

for $f \in C([0, 1], M_{r(n_k)}).$ Let A be the resulting direct limit C^* -algebra.

It is easy to check, and is stated as Lemma 3.5 of [Thomsen 1994], that $\widehat{\alpha_{k+1,k}} = \psi_{k+1,k}.$ Letting V and W be the order unit spaces

$$V = \varinjlim((C([0, 1], \mathbb{R}))_{k=0,1,\dots}, (E(\lambda_{j,k}))_{0 \leq j \leq k}),$$

$$W = \varinjlim((C([0, 1], \mathbb{R}))_{k=0,1,\dots}, (\widehat{\alpha_{k,j}})_{0 \leq j \leq k}),$$

(2-7) now says $V \cong W.$ Lemma 3.2 of [Thomsen 1994] and (2-6) imply that $V \cong \text{Aff}(K).$ Proposition 2.7 implies that $\text{Aff}(T(A)) \cong W.$ So $\text{Aff}(T(A)) \cong \text{Aff}(K),$ whence $T(A) \cong K$ by Theorem 2.4. □

Proposition 2.15. *Let $(D_n)_{n=0,1,2,\dots}$ and $(C_n)_{n=0,1,2,\dots}$ be sequences of unital C^* -algebras. Let*

$$\begin{aligned} &((D_n)_{n=0,1,2,\dots}, (\varphi_{n,m})_{0 \leq m \leq n}), \quad ((D_n)_{n=0,1,2,\dots}, (\varphi'_{n,m})_{0 \leq m \leq n}), \\ &((C_n)_{n=0,1,2,\dots}, (\psi_{n,m})_{0 \leq m \leq n}), \quad ((C_n)_{n=0,1,2,\dots}, (\psi'_{n,m})_{0 \leq m \leq n}) \end{aligned}$$

be direct systems with unital homomorphisms, and call the direct limits (in order) $D, D', C,$ and C' . Suppose further that we have unital homomorphisms $\mu_n, \mu'_n : D_n \rightarrow C_n$ for $n \in \mathbb{Z}_{\geq 0}$ such that

$$\mu_n \circ \varphi_{n,m} = \psi_{n,m} \circ \mu_m \quad \text{and} \quad \mu'_n \circ \varphi'_{n,m} = \psi'_{n,m} \circ \mu'_m$$

for all $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$. Let $\mu : D \rightarrow C$ and $\mu' : D' \rightarrow C'$ be the induced maps of the direct limits. Assume that for all $m \in \mathbb{Z}_{\geq 0}$ we have

$$\sum_{n=m}^{\infty} \|\widehat{\varphi_{n,m}} - \widehat{\varphi'_{n,m}}\| < \infty \quad \text{and} \quad \sum_{n=m}^{\infty} \|\widehat{\psi_{n,m}} - \widehat{\psi'_{n,m}}\| < \infty.$$

Then there exist isomorphisms

$$\rho : \text{Aff}(T(D)) \rightarrow \text{Aff}(T(D')) \quad \text{and} \quad \sigma : \text{Aff}(T(C)) \rightarrow \text{Aff}(T(C'))$$

such that $\widehat{\mu'} \circ \rho = \sigma \circ \widehat{\mu}$. Moreover, if $C_n = D_n$ for all $n \in \mathbb{Z}_{\geq 0}$ and $\psi_{n,m} = \varphi_{n,m}$ and $\psi'_{n,m} = \varphi_{n,m}$ for all m and n , then we can take $\sigma = \rho$.

Proof. We can apply Propositions 2.11 and 2.12 using arbitrary countable dense subsets of the closed unit balls of $\text{Aff}(T(D_n))$ and $\text{Aff}(T(C_n))$ for $n \in \mathbb{Z}_{>0}$. Under the hypotheses of the last statement, the uniqueness statement in Proposition 2.11 implies that $\sigma = \rho$. □

Lemma 2.16. *Adopt the notation of Construction 1.1, including (11) (a second set of maps), and (9) and (13) (diagonal maps, agreeing in the coordinates $1, 2, \dots, d(n+1)$). Then*

$$\|\widehat{\Gamma_{n+1,n}^{(0)}} - \widehat{\Gamma_{n+1,n}}\| \leq \frac{2k(n+1)}{d(n+1) + k(n+1)}$$

for all $n \in \mathbb{Z}_{\geq 0}$.

Proof. For a compact metrizable space Z , let $M(Z)$ be the real Banach space consisting of all signed Borel measures on Z . (That is, $M(Z)$ is the dual space of $C(Z, \mathbb{R})$.) Identify Z with the set of point masses in $M(Z)$. For $n \in \mathbb{Z}_{\geq 0}$, we can identify $T(C_n)$ with the weak* compact convex subset of $M(X_n \sqcup Y_n)$ consisting of probability measures. Thus $X_n \sqcup Y_n \subset T(C_n)$. For every function $f \in \text{Aff}(T(C_n))$, the function $\iota_n(f)(z) = f(z) \cdot 1_{M_r(n)}$ for $z \in X_n \sqcup Y_n$ is in $C(X_n \sqcup Y_n, M_r(n)) = C_n$; and $\tau(\iota_n(f)) = f(\tau)$ for all $\tau \in X_n \sqcup Y_n \subset T(C_n)$, hence also all $\tau \in T(C_n)$ by linearity and continuity.

For $f \in \text{Aff}(T(C_n))$ and $\tau \in T(C_{n+1})$, we can apply the formula in Construction 1.1(9) to $\iota_n(f)$ and apply τ to everything, to get

$$\widehat{\Gamma_{n+1,n}^{(0)}}(f)(\tau) = \frac{1}{l(n+1)} \sum_{k=1}^{l(n+1)} \tau(f \circ S_{n,1}^{(0)}) \quad \text{and} \quad \widehat{\Gamma_{n+1,n}}(f)(\tau) = \frac{1}{l(n+1)} \sum_{k=1}^{l(n+1)} \tau(f \circ S_{n,1}).$$

Using (13), we get

$$\begin{aligned} |\widehat{\Gamma_{n+1,n}^{(0)}}(f)(\tau) - \widehat{\Gamma_{n+1,n}}(f)(\tau)| &= \frac{1}{l(n+1)} \left| \sum_{k=d(n+1)+1}^{l(n+1)} [\tau(f \circ S_{n,1}^{(0)}) - \tau(f \circ S_{n,1})] \right| \\ &\leq \frac{l(n+1) - d(n+1)}{l(n+1)} (2\|f\|_\infty). \end{aligned}$$

The conclusion follows. □

We add additional parts to Constructions 1.1 and 1.6.

Construction 2.17. Adopt the assumptions and notation of all parts of Construction 1.1 (except (13)), and in addition make the following assumptions and definitions:

(20) For all $m \in \mathbb{Z}_{\geq 0}$, the maps $S_{m,j}^{(0)}, S_{m,j} : X_{m+1} \sqcup Y_{m+1} \rightarrow X_m \sqcup Y_m$ satisfy

$$S_{m,j}^{(0)}(X_{m+1}) \subset X_m \quad \text{and} \quad S_{m,j}^{(0)}(Y_{m+1}) \subset Y_m$$

for $j = 1, 2, \dots, l(m)$,

$$S_{m,j}(X_{m+1}) \subset X_m \quad \text{and} \quad S_{m,j}(Y_{m+1}) \subset Y_m$$

for $j = 1, 2, \dots, d(m)$, and

$$S_{m,j}(X_{m+1}) \subset Y_m \quad \text{and} \quad S_{m,j}(Y_{m+1}) \subset X_m$$

for $j = d(m) + 1, d(m) + 2, \dots, l(m)$.

(21) For $m \in \mathbb{Z}_{\geq 0}$, define $D_m = M_{r(m)} \oplus M_{r(m)}$. Define $\varphi_{m+1,m}^{(0)}, \varphi_{m+1,m} : D_m \rightarrow D_{m+1}$ by, for $a, b \in M_{r(m)}$,

$$\begin{aligned} \varphi_{m+1,m}^{(0)}(a, b) &= (\text{diag}(a, a, \dots, a), \text{diag}(b, b, \dots, b)), \\ \varphi_{m+1,m}(a, b) &= (\text{diag}(a, a, \dots, a, b, b, \dots, b), \text{diag}(b, b, \dots, b, a, a, \dots, a)), \end{aligned}$$

in which a occurs $d(m)$ times in the first entry in the second line on the right and $k(m)$ times in the second entry, while b occurs $k(m)$ times in the first entry and $d(m)$ times in the second entry. For $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$, define

$$\varphi_{n,m} = \varphi_{n,n-1} \circ \varphi_{n-1,n-2} \circ \dots \circ \varphi_{m+1,m} : D_m \rightarrow D_n,$$

and define $\varphi_{n,m}^{(0)} : D_m \rightarrow D_n$ similarly. Define AF algebras by

$$D = \varinjlim_m (D_m, \varphi_{m+1,m}) \quad \text{and} \quad D^{(0)} = \varinjlim_m (D_m, \varphi_{m+1,m}^{(0)}),$$

and for $m \in \mathbb{Z}_{>0}$ let $\varphi_{\infty,m} : D_m \rightarrow D$ and $\varphi_{\infty,m}^{(0)} : D_m \rightarrow D^{(0)}$ be the maps associated to these direct limits.

(22) For $m \in \mathbb{Z}_{\geq 0}$, define $\mu_m : D_m \rightarrow C_m$ as follows. For $a, b \in M_{r(m)}$ let $f \in C(X_m, M_{r(m)})$ and $g \in C(Y_m, M_{r(m)})$ be the constant functions with values a and b . Then set $\mu_m(a, b) = (f, g)$. Further, following Lemma 2.18(2) below, let $\mu : D \rightarrow C$ and $\mu^{(0)} : D^{(0)} \rightarrow C^{(0)}$ be the direct limits of the maps μ_m .

(23) For $m \in \mathbb{Z}_{\geq 0}$, define $\theta_m : D_m \rightarrow D_m$ by $\theta_m(a, b) = (b, a)$ for $a, b \in M_{r(m)}$. Further, following Lemma 2.18(3) below, let $\theta \in \text{Aut}(D)$ and $\theta^{(0)} \in \text{Aut}(D^{(0)})$ be the direct limits of the maps θ_m .

Lemma 2.18. *Under the assumptions of Constructions 1.1 (except (13)), 1.6, and 2.17, the following hold:*

(1) *The direct system $((C_n^{(0)})_{n=0,1,2,\dots}, (\Gamma_{n,m}^{(0)})_{0 \leq m \leq n})$ is the direct sum of two direct systems*

$$\begin{aligned} & ((C(X_n, M_{r(n)}))_{n=0,1,2,\dots}, (\Gamma_{n,m}^{(0)}|_{C(X_m, M_{r(m)})})_{0 \leq m \leq n}), \\ & ((C(Y_n, M_{r(n)}))_{n=0,1,2,\dots}, (\Gamma_{n,m}^{(0)}|_{C(Y_m, M_{r(m)})})_{0 \leq m \leq n}), \end{aligned}$$

and $C^{(0)}$ is isomorphic to the direct sum of the direct limits A and B of these systems.

(2) *For all $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$,*

$$\Gamma_{n,m}^{(0)} \circ \mu_m = \mu_n \circ \varphi_{n,m}^{(0)} \quad \text{and} \quad \Gamma_{n,m} \circ \mu_m = \mu_n \circ \varphi_{n,m}.$$

Moreover, the maps μ_m induce unital homomorphisms $\mu^{(0)} : D^{(0)} \rightarrow C^{(0)}$ and $\mu : D \rightarrow C$, and for all $m \in \mathbb{Z}_{\geq 0}$,

$$\Gamma_{\infty,m}^{(0)} \circ \mu_m = \mu^{(0)} \circ \varphi_{\infty,m}^{(0)} \quad \text{and} \quad \Gamma_{\infty,m} \circ \mu_m = \mu \circ \varphi_{\infty,m}.$$

(3) *For all $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$,*

$$\varphi_{n,m}^{(0)} \circ \theta_m = \theta_n \circ \varphi_{n,m}^{(0)} \quad \text{and} \quad \varphi_{n,m} \circ \theta_m = \theta_n \circ \varphi_{n,m}.$$

The maps θ_m induce automorphisms $\theta : D \rightarrow D$ and $\theta^{(0)} : D^{(0)} \rightarrow D^{(0)}$ such that

$$\varphi_{\infty,m} \circ \theta_m = \theta \circ \varphi_{\infty,m} \quad \text{and} \quad \varphi_{\infty,m}^{(0)} \circ \theta_m = \theta^{(0)} \circ \varphi_{\infty,m}^{(0)}$$

for all $m \in \mathbb{Z}_{\geq 0}$.

(4) *For all $m \in \mathbb{Z}_{\geq 0}$, $(\mu_m)_* : K_*(D_m) \rightarrow K_*(C_m)$ is an isomorphism, and*

$$\mu_* : K_*(D) \rightarrow K_*(C) \quad \text{and} \quad (\mu^{(0)})_* : K_*(D^{(0)}) \rightarrow K_*(C^{(0)})$$

are isomorphisms.

Proof. The fact that all the maps in (4) are isomorphisms on K -theory comes from the assumption that the spaces X_m and Y_m are contractible ((14) and (15) in Construction 1.6). Everything else is essentially immediate from the constructions. □

3. The main theorem

We now have the ingredients to deduce the main theorem of this paper, Theorem 3.2.

To state the theorem, we first need to define automorphisms of Elliott invariants, so we need a category in which they lie. For convenience, we restrict to unital C^* -algebras, and we give a very basic list of conditions.

Definition 3.1. An *abstract unital Elliott invariant* is a tuple $G = (G_0, (G_0)_+, g, G_1, K, \rho)$ in which $(G_0, (G_0)_+, g)$ is a preordered abelian group with distinguished positive element g which is an order unit, G_1 is an abelian group, K is a Choquet simplex (possibly empty), and $\rho : G_0 \rightarrow \text{Aff}(K)$ is an order preserving group homomorphism such that $\rho(g)$ is the constant function 1. (If $K = \emptyset$, we take $\text{Aff}(K) = \{0\}$, and we take ρ to be the constant function with value 0.)

If

$$G^{(0)} = (G_0^{(0)}, (G_0^{(0)})_+, g^{(0)}, G_1^{(0)}, K^{(0)}, \rho^{(0)}) \quad \text{and} \quad G^{(1)} = (G_0^{(1)}, (G_0^{(1)})_+, g^{(1)}, G_1^{(1)}, K^{(1)}, \rho^{(1)})$$

are abstract unital Elliott invariants, then a *morphism* from $G^{(0)}$ to $G^{(1)}$ is a triple $F = (F_0, F_1, S)$ in which $F_0 : G_0^{(0)} \rightarrow G_0^{(1)}$ is a group homomorphism satisfying

$$F_0((G_0^{(0)})_+) \subset (G_0^{(1)})_+ \quad \text{and} \quad F_0(g^{(0)}) = g^{(1)},$$

$F_1 : G_1^{(0)} \rightarrow G_1^{(1)}$ is a group homomorphism, and $S : K^{(1)} \rightarrow K^{(0)}$ is a continuous affine map satisfying

$$\rho^{(1)}(F_0(\eta)) = \rho^{(0)}(\eta) \circ S \tag{3-1}$$

for all $\eta \in G_0^{(0)}$.

If

$$F^{(0)} : G^{(0)} \rightarrow G^{(1)} \quad \text{and} \quad F^{(1)} = (F_0^{(1)}, F_1^{(1)}, S^{(1)}) : G^{(1)} \rightarrow G^{(2)}$$

are morphisms of abstract unital Elliott invariants, then define

$$F^{(1)} \circ F^{(0)} = (F_0^{(1)} \circ F_0^{(0)}, F_1^{(1)} \circ F_1^{(0)}, S^{(0)} \circ S^{(1)}).$$

(Note: $S^{(0)} \circ S^{(1)}$, not $S^{(1)} \circ S^{(0)}$.)

The *Elliott invariant of a unital C^* -algebra A* is

$$\text{Ell}(A) = (K_0(A), K_0(A)_+, [1], K_1(A), T(A), \rho_A),$$

in which $\rho_A : K_0(A) \rightarrow \text{Aff}(T(A))$ is given by $\rho_A(\eta)(\tau) = \tau_*(\eta)$ for $\eta \in K_0(A)$ and $\tau \in T(A)$.

If A and B are unital C^* -algebras and $\varphi : A \rightarrow B$ is a unital homomorphism, then we define $\varphi_* : \text{Ell}(A) \rightarrow \text{Ell}(B)$ to consist of the maps φ_* from $K_0(A)$ to $K_0(B)$ and from $K_1(A)$ to $K_1(B)$, together with the map $T(\varphi)$ of Definition 2.5. We write it as $(\varphi_{*,0}, \varphi_{*,1}, T(\varphi))$.

Definition 3.1 is enough to make the abstract unital Elliott invariants into a category such that $\text{Ell}(\cdot)$ is a functor from unital C^* -algebras and unital homomorphisms to abstract unital Elliott invariants.

Theorem 3.2. *There exists a simple unital separable AH algebra C with stable rank 1 and with the following property. There exists an automorphism F of $\text{Ell}(C)$ such that there is no automorphism α of C satisfying $\alpha_* = F$. Moreover, the automorphism F in this example can be chosen so that $F \circ F$ is the identity morphism of $\text{Ell}(C)$.*

We outline the proof. We make a first pass through Constructions 1.1 and 1.6, without the spaces Y_n , and without specifying the point evaluation maps. This is Construction 3.3 below. We get a direct system; call its direct limit \tilde{C} . Apply Proposition 2.14 using the sequence of matrix sizes in this system and $K = T(\tilde{C})$. Doing so requires passing to a subsequence of the sequence of matrix sizes. Replace the original system with the corresponding subsystem; Lemma 3.5 below justifies this. Then make a second pass through Constructions 1.1 and 1.6, taking the spaces X_n and the maps between them from this subsystem and the spaces Y_n and the maps between them from the system gotten from Proposition 2.14, as needed substituting appropriate point evaluations for the diagonal entries of the formulas for the maps. This

requires sufficiently few changes that, by our work in Section 2, the tracial state space remains the same. Therefore the algebra obtained from these constructions has an order two automorphism of its tracial state space which corresponds to exchanging the two rows in the diagram (0-2). The constructions have been designed so that there is also a corresponding automorphism of the K-theory. Our work in Section 1 rules out the possibility of a corresponding automorphism of the algebra, because such an automorphism would necessarily send a particular corner of the algebra to another one with a different radius of comparison.

We start with the following construction, which is “half” of Construction 1.1, and gives just the top row of the diagram (0-1).

Construction 3.3. We will consider direct systems and their associated direct limits constructed as follows.

(1) The sequences $(d(n))_{n=0,1,2,\dots}$ and $(k(n))_{n=0,1,2,\dots}$ in $\mathbb{Z}_{\geq 0}$ are as in Construction 1.1(1) and satisfy the condition of Construction 1.1(2). We further define $(l(n))_{n=0,1,2,\dots}$, $(r(n))_{n=0,1,2,\dots}$, $(s(n))_{n=0,1,2,\dots}$, and $(t(n))_{n=0,1,2,\dots}$ as in Construction 1.1(1).

(2) Following Constructions 1.1(3) and (4), we define

$$\kappa = \inf_{n \in \mathbb{Z}_{>0}} \frac{s(n)}{r(n)}, \quad \omega = \frac{k(1)}{k(1) + d(1)}, \quad \text{and} \quad \omega' = \sum_{n=2}^{\infty} \frac{k(n)}{k(n) + d(n)}.$$

(These will not be used directly in connection with this direct system.)

(3) As in Construction 1.6(14), we define compact metric spaces by $X_n = \text{cone}((S^2)^{s(n)})$ for $n \in \mathbb{Z}_{\geq 0}$, and we define maps $Q_j^{(n)} : X_{n+1} \rightarrow X_n$ for $n \in \mathbb{Z}_{\geq 0}$ and $j = 1, 2, \dots, d(n+1)$ to be the cones over the projection maps

$$(S^2)^{s(n+1)} = ((S^2)^{s(n)})^{d(n+1)} \rightarrow (S^2)^{s(n)}.$$

(4) We are given maps $\delta_n : C(X_n) \rightarrow C(X_{n+1}, M_{l(n+1)})$ (as in Construction 1.1(8), but with only one summand) which are diagonal; that is, there are continuous maps

$$T_{n,1}, T_{n,2}, \dots, T_{n,l(n+1)} : X_{n+1} \rightarrow X_n$$

such that

$$\delta_n(f) = \text{diag}(f \circ T_{n,1}, f \circ T_{n,2}, \dots, f \circ T_{n,l(n+1)})$$

for $f \in C(X_n)$. (Compare with Construction 1.1(9).) Moreover, $T_{n,j} = Q_j^{(n)}$ for $j = 1, 2, \dots, d(n+1)$. The maps $T_{n,j}$ are unspecified for $j = d(n+1) + 1, d(n+1) + 2, \dots, l(n+1)$.

(5) Set $A_n = M_{r(n)} \otimes C(X_n)$ (like in Construction 1.1(7) but with only one summand). Following Construction 1.1(8), set

$$\Delta_{n+1,n} = \text{id}_{M_{r(n)}} \otimes \delta_n : A_n \rightarrow A_{n+1},$$

and for $m, n \in \mathbb{Z}_{\geq 0}$ with $m \leq n$, take

$$\Delta_{n,m} = \Delta_{n,n-1} \circ \Delta_{n-1,n-2} \circ \dots \circ \Delta_{m+1,m} : A_m \rightarrow A_n.$$

(6) Define $A = \varinjlim_n A_n$, taken with respect to the maps $\Delta_{n,m}$. For $n \in \mathbb{Z}_{\geq 0}$, let $\Delta_{\infty,n} : A_n \rightarrow A$ be the map associated with the direct limit.

To avoid confusing notation, we isolate the following computation as a lemma.

Lemma 3.4. *Let $n \in \mathbb{Z}_{>0}$ and let $\kappa_1, \kappa_2, \dots, \kappa_n, \delta_1, \delta_2, \dots, \delta_n \in (0, \infty)$. Then*

$$\sum_{j=1}^n \frac{\kappa_j}{\delta_j + \kappa_j} \geq \frac{\prod_{j=1}^n (\delta_j + \kappa_j) - \prod_{j=1}^n \delta_j}{\prod_{j=1}^n (\delta_j + \kappa_j)}.$$

Proof. For $j = 1, 2, \dots, n$ define

$$\lambda_j = \frac{\kappa_j}{\delta_j + \kappa_j}.$$

Then $\lambda_j \in (0, 1)$. Some calculation shows that the conclusion of the lemma becomes

$$\sum_{j=1}^n \lambda_j \geq 1 - \prod_{j=1}^n (1 - \lambda_j). \tag{3-2}$$

We prove (3-2) by induction on n . For $n = 1$ it is trivial. Suppose (3-2) is known for some value of n . Given $\lambda_1, \lambda_2, \dots, \lambda_{n+1} \in (0, 1)$, set $\mu = 1 - (1 - \lambda_n)(1 - \lambda_{n+1})$. Then

$$\mu \in (0, 1) \quad \text{and} \quad \mu = \lambda_n + \lambda_{n+1} - \lambda_n \lambda_{n+1} \leq \lambda_n + \lambda_{n+1}.$$

Applying the induction hypothesis on $\lambda_1, \lambda_2, \dots, \lambda_{n-1}, \mu$ at the second step, we then have

$$\sum_{j=1}^{n+1} \lambda_j \geq \sum_{j=1}^{n-1} \lambda_j + \mu \geq 1 - \left[\prod_{j=1}^{n-1} (1 - \lambda_j) \right] (1 - \mu) = 1 - \prod_{j=1}^{n+1} (1 - \lambda_j).$$

This completes the induction, and the proof of the lemma. □

Lemma 3.5. *Let a direct system as in Construction 3.3 be given, but using sequences $(\tilde{d}(n))_{n=0,1,2,\dots}$ and $(\tilde{k}(n))_{n=0,1,2,\dots}$ in place of $(d(n))_{n=0,1,2,\dots}$ and $(k(n))_{n=0,1,2,\dots}$. Denote the additional sequences analogous to those in Construction 3.3(1) by \tilde{l}, \tilde{r} , and \tilde{s} . Denote the numbers analogous to those in Construction 3.3(2) by $\tilde{\kappa}, \tilde{\omega}$, and $\tilde{\omega}'$. Denote the spaces used in the system by \tilde{X}_n . Let $v : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$ be a strictly increasing function such that $v(0) = 0$ and $v(1) = 1$. Then the direct system $(C(\tilde{X}_{v(m)}), M_{\tilde{r}(v(m))})_{m=0,1,2,\dots}$ is isomorphic to a system as in Construction 3.3, with the choices $d(0) = 1, k(0) = 0,$*

$$d(m) = \tilde{d}(v(m-1) + 1) \tilde{d}(v(m-1) + 2) \cdots \tilde{d}(v(m)), \tag{3-3}$$

$$k(m) = \tilde{l}(v(m-1) + 1) \tilde{l}(v(m-1) + 2) \cdots \tilde{l}(v(m)) - d(m) \tag{3-4}$$

for $m \in \mathbb{Z}_{>0}$. Moreover, following the notation of Construction 3.3,

$$l(m) = \tilde{l}(v(m-1) + 1) \tilde{l}(v(m-1) + 2) \cdots \tilde{l}(v(m)), \tag{3-5}$$

$$r(m) = \tilde{r}(v(m)), \quad \text{and} \quad s(m) = \tilde{s}(v(m))$$

for $m \in \mathbb{Z}_{\geq 0}$, and

$$\kappa = \tilde{\kappa}, \quad \omega = \tilde{\omega}, \quad \text{and} \quad \omega' \leq \tilde{\omega}'.$$

Proof. Given the definitions of d and k , the proofs of the formulas for l, r , and s are easy.

Using Lemma 1.2 at the first and fourth steps, we now get

$$\tilde{\kappa} = \lim_{n \rightarrow \infty} \frac{\tilde{s}(n)}{\tilde{r}(n)} = \lim_{m \rightarrow \infty} \frac{\tilde{r}(v(m))}{\tilde{s}(v(m))} = \lim_{m \rightarrow \infty} \frac{s(m)}{r(m)} = \kappa.$$

We have $\omega = \tilde{\omega}$ because $v(1) = 1$.

Using Lemma 3.4 at the second step and (3-3), (3-4) and (3-5) at the third step, we have

$$\begin{aligned} \tilde{\omega}' &= \sum_{m=2}^{\infty} \sum_{j=v(m-1)+1}^{v(m)} \frac{\tilde{k}(j)}{\tilde{k}(j) + \tilde{d}(j)} \\ &\geq \sum_{m=2}^{\infty} \frac{\prod_{j=v(m-1)+1}^{v(m)} [\tilde{d}(j) + \tilde{k}(j)] - \prod_{j=v(m-1)+1}^{v(m)} \tilde{d}(j)}{\prod_{j=v(m-1)+1}^{v(m)} [\tilde{d}(j) + \tilde{k}(j)]} = \sum_{m=2}^{\infty} \frac{k(m)}{k(m) + d(m)} = \omega'. \end{aligned}$$

Define $X_m = \tilde{X}_{v(m)}$ for $m \in \mathbb{Z}_{\geq 0}$. Clearly $X_m = \text{cone}((S^2)^{s(m)})$, as required. Denote the maps in the system of the hypotheses by

$$\tilde{\delta}_n : C(\tilde{X}_n) \rightarrow C(\tilde{X}_{n+1}, M_{\tilde{l}(n+1)}) \quad \text{and} \quad \tilde{\Delta}_{n,m} : \tilde{C}_m \rightarrow \tilde{C}_n,$$

with $\tilde{\delta}_n$ being built using maps

$$\tilde{T}_{n,1}, \tilde{T}_{n,2}, \dots, \tilde{T}_{n,l(n+1)} : \tilde{X}_{n+1} \rightarrow \tilde{X}_n,$$

as in Construction 3.3(4). For $p = v(m), v(m) + 1, \dots, v(m + 1) - 1$, set

$$j(p) = \frac{\tilde{r}(p)}{\tilde{r}(v(m))} = \tilde{l}(v(m) + 1)\tilde{l}(v(m) + 2) \cdots \tilde{l}(p).$$

Then define

$$\delta_m^{(0)} : C(\tilde{X}_{v(m)}) \rightarrow C(\tilde{X}_{v(m+1)}, M_{l(n+1)})$$

by

$$\delta_m^{(0)} = \text{id}_{M_{j(v(m+1)-1)}} \otimes \tilde{\delta}_{v(m+1)-1} \circ \text{id}_{M_{j(v(m+1)-2)}} \otimes \tilde{\delta}_{v(m+1)-2} \circ \cdots \circ \tilde{\delta}_{v(m)}.$$

(In the last term we omit $\text{id}_{M_{j(v(m))}}$ since $j(v(m)) = 1$.) With this definition, one checks that $\text{id}_{M_{\tilde{r}(v(m))}} \otimes \tilde{\delta}_m = \tilde{\Delta}_{v(m+1), v(m)}$, so that the direct system gotten using the maps $\delta_m^{(0)}$ in Construction 3.3 is a subsystem of the system given in the hypotheses.

We claim that $\delta_m^{(0)}$ is unitarily equivalent to a map $\delta_m : C(X_m) \rightarrow C(X_{m+1}, M_{l(n+1)})$ as in Construction 3.3. This will imply isomorphism of the direct systems, and complete the proof of the lemma. First, $\delta_m^{(0)}$ is given as in Construction 3.3(4) using some maps from $\tilde{X}_{v(m+1)}$ to $\tilde{X}_{v(m)}$, namely all possible compositions

$$\tilde{T}_{v(m), i_{v(m)}} \circ \tilde{T}_{v(m)+1, i_{v(m)+1}} \circ \cdots \circ \tilde{T}_{v(m+1)-1, i_{v(m+1)-1}},$$

with $i_p = 1, 2, \dots, \tilde{l}(p+1)$ for $p = v(m), v(m) + 1, \dots, v(m + 1) - 1$. Moreover, since the composition of projection maps is a projection map, restricting to $i_p = 1, 2, \dots, \tilde{d}(p+1)$ for all p gives exactly all the maps $Q_j^{(m)} : X_{m+1} \rightarrow X_m$ for $j = 1, 2, \dots, d(n+1)$. Therefore $\delta_m^{(0)}$ is unitarily equivalent to a map as in Construction 3.3 by a permutation matrix. □

Proof of Theorem 3.2. Choose $N \in \mathbb{Z}_{>0}$ such that

$$N > 5 \quad \text{and} \quad \exp\left(-\frac{1}{N-1}\right) > \frac{3}{4}. \tag{3-6}$$

(For example, $N = 6$ will work.) In Construction 1.1(1) we make preliminary choices of the numbers $d(n)$ etc., calling them $\tilde{d}(n)$ etc. Take $\tilde{d}(0) = 1$ and $\tilde{k}(0) = 0$, and take $\tilde{d}(n) = N^n$ and $\tilde{k}(n) = 1$ for $n \in \mathbb{Z}_{>0}$. Then

$$\tilde{l}(n) = N^n + 1, \quad \tilde{r}(n) = \prod_{j=1}^n (N^j + 1), \quad \text{and} \quad \tilde{s}(n) = \prod_{j=1}^n N^j$$

for $n \in \mathbb{Z}_{>0}$. We obtain numbers as in Construction 3.3(2) (equivalently, Constructions 1.1(3) and (4)), which we call $\tilde{\kappa}$, $\tilde{\omega}$, and $\tilde{\omega}'$. Further, adopt the definitions and notation of Construction 3.3, except that we use \tilde{X}_n instead of X_n and similarly throughout. That is, in Construction 3.3(3) we call the spaces \tilde{X}_n instead of X_n , the projection maps $\tilde{Q}_j^{(n)}$, in Construction 3.3(4) we call the maps of algebras $\tilde{\delta}_n$ and the maps of spaces $\tilde{T}_{n,j} : \tilde{X}_{n+1} \rightarrow \tilde{X}_n$, in Construction 3.3(5) we call the algebras \tilde{A}_n and the maps $\tilde{\Delta}_{n,m}$, and in Construction 3.3(6) we call the direct limit \tilde{A} and the maps to it $\tilde{\Delta}_{\infty,n}$. As in Construction 3.3(4), we take $\tilde{T}_{n,j} = \tilde{Q}_j^{(n)}$ for $j = 1, 2, \dots, \tilde{d}(n+1)$. For $n \in \mathbb{Z}_{\geq 0}$ choose an arbitrary point $\tilde{x}_n \in \tilde{X}_n$, and for $j = \tilde{d}(n+1) + 1$ let $\tilde{T}_{n,j}$ be the constant function on \tilde{X}_{n+1} with value \tilde{x}_n . (Note that $\tilde{d}(n+1) + 1 = \tilde{l}(n+1)$.)

We claim that the conditions in Constructions 1.1(3), 1.1(4), and 1.1(5) are satisfied, and moreover that

$$\frac{1}{1 - 2\tilde{\omega}} < \frac{2\tilde{\kappa} - 1}{2\tilde{\omega}}.$$

For $n \in \mathbb{Z}_{>0}$ we have, using $\log(m+1) - \log(m) < 1/m$ at the third step,

$$\frac{\tilde{s}(n)}{\tilde{r}(n)} = \prod_{j=1}^n \frac{N^j}{N^j + 1} = \exp\left(\sum_{j=1}^n -[\log(N^j + 1) - \log(N^j)]\right) \geq \exp\left(-\sum_{j=1}^n \frac{1}{N^j}\right) > \exp\left(-\frac{1}{N-1}\right).$$

So $\tilde{\kappa} \geq \exp(-1/(N-1)) > \frac{3}{4}$ by (3-6). Furthermore,

$$\tilde{\omega} = \frac{1}{N+1} < \frac{1}{4} \quad \text{and} \quad \tilde{\omega}' = \sum_{j=2}^{\infty} \frac{1}{N^j + 1} < \sum_{j=2}^{\infty} \frac{1}{N^j} = \frac{1}{N(N-1)},$$

so the conditions $\tilde{\omega}' < \tilde{\omega} < \frac{1}{2}$ in Construction 1.1(4) and $2\tilde{\kappa} - 1 > 2\tilde{\omega}$ in Construction 1.1(5) are satisfied. Moreover,

$$\frac{1}{1 - 2\tilde{\omega}} = \frac{N+1}{N-1} < \frac{N+1}{4} = \frac{1}{4\tilde{\omega}} = \frac{2\left(\frac{3}{4}\right) - 1}{2\tilde{\omega}} < \frac{2\tilde{\kappa} - 1}{2\tilde{\omega}}.$$

The claim is proved.

Apply Proposition 2.14 with $K = T(\tilde{A})$ and with $\tilde{l}(n)$ and $\tilde{r}(n)$ in place of $l(n)$ and $r(n)$, getting a strictly increasing sequence, which we call $(\nu(n))_{n=0,1,2,\dots}$, with $\nu(j) = j$ for $j = 0, 1$, an AI algebra B_0 (called A in Proposition 2.14) which is the direct limit of a unital system

$$C([0, 1]) \otimes M_{r(\nu(0))} \xrightarrow{\alpha_{1,0}} C([0, 1]) \otimes M_{r(\nu(1))} \xrightarrow{\alpha_{2,1}} C([0, 1]) \otimes M_{r(\nu(2))} \xrightarrow{\alpha_{3,2}} \dots,$$

with injective diagonal maps $\alpha_{n+1, n}$ given by

$$f \mapsto \text{diag}(f \circ R_{n,1}, f \circ R_{n,2}, \dots, f \circ R_{n, r(v_{n+1})/r(v_n)})$$

for continuous functions

$$R_{n,1}, R_{n,2}, \dots, R_{n, r(v(n+1))/r(v(n))} : [0, 1] \rightarrow [0, 1],$$

and an isomorphism $T(B_0) \rightarrow T(\tilde{A})$.

Apply Lemma 3.5 with this choice of ν . Define the sequences $(d(n))_{n=0,1,2,\dots}$ and $(k(n))_{n=0,1,2,\dots}$ as in Lemma 3.5, and then make all the definitions in Constructions 1.1 and 1.6. (Some are also given in the statement of Lemma 3.5.) Then, as in the proof of Lemma 3.5, $X_n = \tilde{X}_{\nu(n)}$. We make the following choices for the unspecified objects in these constructions. We choose points $x_n \in X_n$ and $y_n \in [0, 1]$ for $n \in \mathbb{Z}_{\geq 0}$ such that the conditions in Constructions 1.6(16) and (17) are satisfied. (It is easy to see that this can be done.) Use these points in parts (b) and (d) of Construction 1.6(18). Take the maps

$$R_{n,1}, R_{n,2}, \dots, R_{n, d(n+1)} : Y_{n+1} \rightarrow Y_n$$

in part (c) of Construction 1.6(18) to be those from the application of Proposition 2.14 above. For $j = 1, 2, \dots, l(n+1)$, let $S_{n,j}^{(0)}|_{X_{n+1}} : X_{n+1} \rightarrow X_n$ be the maps in the system obtained from Lemma 3.5, and take $S_{n,j}^{(0)}|_{Y_{n+1}} = R_{n,j}$. The requirement $S_{n,j}^{(0)} = S_{n,j}$ for $j = 1, 2, \dots, d(n+1)$ in Construction 1.6(19) is then satisfied, so that the condition in Construction 1.1(13) is also satisfied. Moreover, with these choices, the conditions in Construction 2.17(20) are satisfied.

By Lemma 3.5, the numbers $\kappa, \omega,$ and ω' from Constructions 1.1(3) and (4) satisfy

$$\kappa = \tilde{\kappa}, \quad \omega = \tilde{\omega}, \quad \text{and} \quad \omega' \leq \tilde{\omega}'.$$

Therefore $\kappa > \frac{1}{2}, \omega' < \omega < \frac{1}{2}$, and $2\kappa - 1 > 2\omega$, as required in Constructions 1.1(3), (4), and (5); moreover

$$\frac{1}{1 - 2\omega} < \frac{2\kappa - 1}{2\omega}. \tag{3-7}$$

The algebra C is simple by Lemma 1.7.

The algebras A and B of Lemma 2.18(1) are now $A = \tilde{A}$ and $B = B_0$, so $C^{(0)}$, as in Construction 1.1(11), is isomorphic to $\tilde{A} \oplus B_0$. The isomorphism $T(B_0) \rightarrow T(\tilde{A})$ gives an isomorphism $\zeta_0^{(0)} : \text{Aff}(T(A)) \rightarrow \text{Aff}(T(B))$. This provides an automorphism of $\text{Aff}(T(A)) \oplus \text{Aff}(T(B))$, given by

$$(f, g) \mapsto ((\zeta_0^{(0)})^{-1}(g), \zeta_0^{(0)}(f)).$$

Let $\zeta^{(0)}$ be the corresponding automorphism of $\text{Aff}(T(A \oplus B)) = \text{Aff}(T(C^{(0)}))$ gotten using Lemma 2.10. Clearly $\zeta^{(0)} \circ \zeta^{(0)}$ is the identity map on $\text{Aff}(T(C^{(0)}))$.

Adopt the notation of Construction 2.17: C and $C^{(0)}$ are as already described, D and $D^{(0)}$ are the AF algebras from Construction 2.17(21), $\mu : D \rightarrow C$ and $\mu^{(0)} : D^{(0)} \rightarrow C^{(0)}$ are the maps of Construction 2.17(22) (which are isomorphisms on K-theory by Lemma 2.18(4)), and $\theta \in \text{Aut}(D)$ and $\theta^{(0)} \in \text{Aut}(D^{(0)})$ are as in Construction 2.17(23).

Define $E = \varinjlim_n M_{r(m)}$, with respect to the maps $a \mapsto \text{diag}(a, a, \dots, a)$, with a repeated $l(n)$ times. The direct system defining $D^{(0)}$ is the direct sum of two copies of the direct system just defined, so

$$D^{(0)} \cong E \oplus E \quad \text{and} \quad \text{Aff}(T(D^{(0)})) \cong \text{Aff}(T(E \oplus E)).$$

Since E is a UHF algebra, we have $\text{Aff}(T(E)) \cong \mathbb{R}$ with the usual order and order unit 1. Using $\text{id}_{\widehat{\text{Aff}(T(E))}}$ in place of $\zeta_0^{(0)}$ above, we get an automorphism of $\text{Aff}(T(D^{(0)}))$. But this automorphism is just $\widehat{\theta}^{(0)}$.

We claim that $\zeta^{(0)} \circ \widehat{\mu}^{(0)} = \widehat{\mu}^{(0)} \circ \widehat{\theta}^{(0)}$. To prove the claim, we work with

$$\text{Aff}(T(E)) \oplus \text{Aff}(T(E)) \quad \text{and} \quad \text{Aff}(T(A)) \oplus \text{Aff}(T(B))$$

in place of $\text{Aff}(T(D^{(0)}))$ and $\text{Aff}(T(C^{(0)}))$, but keep the same names for the maps.

Since $\mu^{(0)} : E \oplus E \rightarrow A \oplus B$ is the direct sum of unital maps from the first summand to A and the second summand to B , the map $\widehat{\mu}^{(0)}$ is similarly a direct sum of maps $\text{Aff}(T(E)) \rightarrow \text{Aff}(T(A))$ and $\text{Aff}(T(E)) \rightarrow \text{Aff}(T(B))$. Let e and f be the order units of $\text{Aff}(T(A))$ and $\text{Aff}(T(B))$. The unique positive order unit preserving maps $\text{Aff}(T(E)) \rightarrow \text{Aff}(T(A))$ and $\text{Aff}(T(E)) \rightarrow \text{Aff}(T(B))$ are $\alpha \mapsto \alpha e$ and $\beta \mapsto \beta f$ for $\alpha, \beta \in \mathbb{R}$. Therefore $\widehat{\mu}^{(0)}(\alpha, \beta) = (\alpha e, \beta f)$. Since $\zeta_0^{(0)}$ is order unit preserving, we have $\zeta_0^{(0)}(e) = f$, so

$$\zeta^{(0)}(\alpha e, \beta f) = (\beta e, \alpha f) = \widehat{\mu}^{(0)}(\beta, \alpha) = (\widehat{\mu}^{(0)} \circ \widehat{\theta}^{(0)})(\alpha, \beta).$$

The claim follows.

Using conditions (4) and (13) in Construction 1.1, Lemma 2.16, and Proposition 2.15, we get isomorphisms

$$\rho : \text{Aff}(T(D^{(0)})) \rightarrow \text{Aff}(T(D)) \quad \text{and} \quad \sigma : \text{Aff}(T(C^{(0)})) \rightarrow \text{Aff}(T(C))$$

such that $\widehat{\mu} \circ \rho = \sigma \circ \widehat{\mu}^{(0)}$. Define

$$\eta = \rho \circ \widehat{\theta}^{(0)} \circ \rho^{-1} \in \text{Aut}(\text{Aff}(T(D))) \quad \text{and} \quad \zeta = \sigma \circ \zeta^{(0)} \circ \sigma^{-1} \in \text{Aut}(\text{Aff}(T(C))).$$

A calculation now shows that the claim above implies

$$\zeta \circ \widehat{\mu} = \widehat{\mu} \circ \eta. \tag{3-8}$$

We also have $\zeta \circ \zeta = \text{id}_{\text{Aff}(T(C))}$.

We want to apply Proposition 2.15 with D_n and $\varphi_{n,m}$ as in Construction 2.17(21), and $\varphi_{n,m}^{(0)}$ as there in place of $\varphi'_{n,m}$, so that D and $D^{(0)}$ are as already given, with $C_n = D_n$ for all $n \in \mathbb{Z}_{\geq 0}$ and $\psi_{n,m} = \varphi_{n,m}$ and $\psi'_{n,m} = \varphi'_{n,m}$ for all m and n , and with $\theta_n, \theta_n^{(0)}, \theta$, and $\theta^{(0)}$ from Construction 2.17(23) in place of μ_n, μ'_n, μ , and μ' . As before, this application is justified by conditions (4) and (13) in Construction 1.1, and Lemma 2.16. The outcome is an isomorphism $\rho' : \text{Aff}(T(D^{(0)})) \rightarrow \text{Aff}(T(D))$ such that

$$\widehat{\theta} = \rho' \circ \widehat{\theta}^{(0)} \circ (\rho')^{-1}. \tag{3-9}$$

We claim that $\eta = \widehat{\theta}$. The “right” way to do this is presumably to show that $\rho' = \rho$ above, but the following argument is easier to write. We have

$$\text{Aff}(T(D)) \cong \text{Aff}(T(D^{(0)})) \cong \mathbb{R}^2,$$

with order $(\alpha, \beta) \geq 0$ if and only if $\alpha \geq 0$ and $\beta \geq 0$ and order unit $(1, 1)$. Since the state space $S(\mathbb{R}^2)$ of \mathbb{R}^2 with this order unit space structure is an interval, and automorphisms of order unit spaces preserve the extreme points of the state space, there is only one possible action of a nontrivial automorphism of \mathbb{R}^2 on $S(\mathbb{R}^2)$. Theorem 2.4 implies that $\mathbb{R}^2 \cong \text{Aff}(S(\mathbb{R}^2))$, so there is only one nontrivial automorphism of \mathbb{R}^2 . Since $\widehat{\theta}^{(0)}$ is nontrivial, so is $\hat{\theta}$ by (3-9), and so is η by its definition. The claim follows.

The claim and (3-8) imply

$$\zeta \circ \hat{\mu} = \hat{\mu} \circ \hat{\theta}. \tag{3-10}$$

Passing to state spaces and applying Theorem 2.4, we get an affine homeomorphism $H : T(C) \rightarrow T(C)$ such that $\zeta(f) = f \circ H$ for all $f \in \text{Aff}(T(C))$, and moreover $H \circ H = \text{id}_{T(C)}$. By Lemma 2.18(4), the expression $\mu_* \circ \theta_* \circ (\mu_*)^{-1}$ is a well-defined automorphism of $K_*(C)$, of order 2. We claim that $F = (\mu_* \circ \theta_* \circ (\mu_*)^{-1}, H)$ is an order two automorphism of $\text{Ell}(C)$. We use the notation of Definition 3.1 for the Elliott invariant of a C^* -algebra; in particular, ρ_C and ρ_D are not related to the maps ρ and ρ' above. The only part needing work is the compatibility condition (3-1) in Definition 3.1, which amounts to showing that

$$\rho_C \circ \mu_* \circ \theta_* \circ (\mu_*)^{-1} = \zeta \circ \rho_C.$$

To see this, we calculate, using at the second and last steps the notation of Definition 2.5 and the fact that the morphisms of Elliott invariants defined by μ and θ satisfy (3-1) in Definition 3.1, and using (3-10) at the third step,

$$\begin{aligned} \zeta \circ \rho_C &= \zeta \circ \rho_C \circ \mu_* \circ (\mu_*)^{-1} = \zeta \circ \hat{\mu} \circ \rho_D \circ (\mu_*)^{-1} \\ &= \hat{\mu} \circ \hat{\theta} \circ \rho_D \circ (\mu_*)^{-1} = \rho_C \circ \mu_* \circ \theta_* \circ (\mu_*)^{-1}, \end{aligned}$$

as desired.

Thus, we have constructed an automorphism F of $\text{Ell}(C)$ of order 2. It remains to show that F is not induced by any automorphism of C .

Using (3-10) on the last components, one easily sees that $F \circ \mu_* = \mu_* \circ \theta_*$. Let q and q^\perp be as in Notation 1.13. In the construction of D as in Construction 2.17(21), set $e = \varphi_{\infty,0}((1, 0))$ and $e^\perp = 1 - e = \varphi_{\infty,0}((0, 1))$. Then $\theta(e) = e^\perp$, $\mu(e) = q$, and $\mu(e^\perp) = q^\perp$. Therefore $F([q]) = [q^\perp]$.

Suppose now that there exists an automorphism α such that $\alpha_* = F$. Then $[\alpha(q)] = [q^\perp]$. By Lemma 1.17, $\alpha(q)$ is unitarily equivalent to q^\perp . Let u be a unitary such that $u\alpha(q)u^* = q^\perp$. Thus, since $\alpha(qAq) = \alpha(q)A\alpha(q) = u^*q^\perp Aq^\perp u$, it follows that the qAq and $q^\perp Aq^\perp$ have the same radius of comparison. By (3-7), this contradicts Lemmas 1.15 and 1.16. \square

Remark 3.6. One can easily check that, with C as in the proof of Theorem 3.2, there is a unique automorphism of $\text{Ell}(C)$ whose component automorphism of the tracial state space is as in the proof. Therefore the conclusion can be slightly strengthened: there is an automorphism of $T(C)$ which is compatible with an automorphism of $\text{Ell}(C)$ but which is not induced by any automorphism of C .

Question 3.7. *Does there exist a compact metric space X and a minimal homeomorphism $h : X \rightarrow X$ such that the crossed product $C^*(\mathbb{Z}, X, h)$ has the same features as the example we construct here?*

Our construction provides an example of an automorphism of order 2 of the Elliott invariant which is not induced by any automorphism of the C^* -algebra. The question of whether there exists an example of such an automorphism of the invariant which is induced by an automorphism of the algebra but not by one of order 2 is an older question by Blackadar, which we record below. For Kirchberg algebras in the UCT class, it is known that any order two automorphism of the Elliott invariant is induced by an order two automorphism of the C^* -algebra [Benson et al. 2003]; also see [Katsura 2008] for a generalization to actions of many other finite groups. However, very little seems to be known in the stably finite case, even for classifiable C^* -algebras (and in fact even for AF algebras).

Question 3.8 (Blackadar). *Does there exist a simple separable stably finite unital nuclear C^* -algebra C and an automorphism F of $\text{Ell}(C)$ such that:*

- (1) $F \circ F$ is the identity morphism of $\text{Ell}(C)$.
- (2) There is an automorphism α of C such that $\alpha_* = F$.
- (3) There is no α as in (2) which in addition satisfies $\alpha \circ \alpha = \text{id}_C$.

Can such an algebra be chosen to be AH and have stable rank 1?

Our method of proof suggests that, instead of being just a number, the radius of comparison should be taken to be a function from $V(A)$ to $[0, \infty]$. If one uses the generalization to nonunital algebras in [Blackadar et al. 2012, Section 3.3], one could presumably even get a function from $\text{Cu}(A)$ to $[0, \infty]$.

Acknowledgement

The second author is grateful to M. Ali Asadi-Vasfi for a careful reading of Section 1, and in particular finding a number of misprints.

References

- [Alfsen 1971] E. M. Alfsen, *Compact convex sets and boundary integrals*, Ergebnisse der Mathematik (2) **57**, Springer, 1971. MR Zbl
- [Benson et al. 2003] D. J. Benson, A. Kumjian, and N. C. Phillips, “Symmetries of Kirchberg algebras”, *Canad. Math. Bull.* **46**:4 (2003), 509–528. MR Zbl
- [Blackadar 1998] B. Blackadar, *K-theory for operator algebras*, 2nd ed., Math. Sci. Res. Inst. Publ. **5**, Cambridge University Press, 1998. MR Zbl
- [Blackadar et al. 2012] B. Blackadar, L. Robert, A. P. Tikuisis, A. S. Toms, and W. Winter, “An algebraic approach to the radius of comparison”, *Trans. Amer. Math. Soc.* **364**:7 (2012), 3657–3674. MR Zbl
- [Dădărlat et al. 1992] M. Dădărlat, G. Nagy, A. Némethi, and C. Pasnicu, “Reduction of topological stable rank in inductive limits of C^* -algebras”, *Pacific J. Math.* **153**:2 (1992), 267–276. MR Zbl
- [Elliott et al. 2009] G. A. Elliott, T. M. Ho, and A. S. Toms, “A class of simple C^* -algebras with stable rank one”, *J. Funct. Anal.* **256**:2 (2009), 307–322. MR Zbl
- [Elliott et al. 2015] G. A. Elliott, G. Gong, H. Lin, and Z. Niu, “On the classification of simple amenable C^* -algebras with finite decomposition rank, II”, preprint, 2015. arXiv 1507.03437v2
- [Goodearl 1986] K. R. Goodearl, *Partially ordered abelian groups with interpolation*, Math. Surv. Monogr. **20**, Amer. Math. Soc., Providence, RI, 1986. MR Zbl

- [Haagerup 2014] U. Haagerup, “Quasitraces on exact C^* -algebras are traces”, *C. R. Math. Acad. Sci. Soc. R. Can.* **36**:2-3 (2014), 67–92. MR Zbl
- [Katsura 2008] T. Katsura, “A construction of actions on Kirchberg algebras which induce given actions on their K -groups”, *J. Reine Angew. Math.* **617** (2008), 27–65. MR Zbl
- [Milnor and Stasheff 1974] J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Ann. of Math. Stud. **76**, Princeton Univ. Press, 1974. MR Zbl
- [Niu 2014] Z. Niu, “Mean dimension and AH-algebras with diagonal maps”, *J. Funct. Anal.* **266**:8 (2014), 4938–4994. MR Zbl
- [Phillips 2014] N. C. Phillips, “Large subalgebras”, preprint, 2014. arXiv 1408.5546v1
- [Rørdam 1992] M. Rørdam, “On the structure of simple C^* -algebras tensored with a UHF-algebra, II”, *J. Funct. Anal.* **107**:2 (1992), 255–269. MR Zbl
- [Rørdam 2003] M. Rørdam, “A simple C^* -algebra with a finite and an infinite projection”, *Acta Math.* **191**:1 (2003), 109–142. MR Zbl
- [Thomsen 1994] K. Thomsen, “Inductive limits of interval algebras: the tracial state space”, *Amer. J. Math.* **116**:3 (1994), 605–620. MR Zbl
- [Tikuisis et al. 2017] A. Tikuisis, S. White, and W. Winter, “Quasidiagonality of nuclear C^* -algebras”, *Ann. of Math. (2)* **185**:1 (2017), 229–284. MR Zbl
- [Toms 2006] A. S. Toms, “Flat dimension growth for C^* -algebras”, *J. Funct. Anal.* **238**:2 (2006), 678–708. MR Zbl
- [Toms 2008] A. S. Toms, “On the classification problem for nuclear C^* -algebras”, *Ann. of Math. (2)* **167**:3 (2008), 1029–1044. MR Zbl
- [Villadsen 1998] J. Villadsen, “Simple C^* -algebras with perforation”, *J. Funct. Anal.* **154**:1 (1998), 110–116. MR Zbl

Received 1 Aug 2020. Revised 16 Jul 2021. Accepted 23 Aug 2021.

ILAN HIRSHBERG: ilan@math.bgu.ac.il

Department of Mathematics, Ben Gurion University of the Negev, Be'er Sheva, Israel

N. CHRISTOPHER PHILLIPS: Department of Mathematics, University of Oregon, Eugene, OR, United States

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard Université Paris Sud XI, France
patrick.gerard@universite-paris-saclay.fr

Clément Mouhot Cambridge University, UK
c.mouhot@dpms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Isabelle Gallagher	Université Paris-Diderot, IMJ-PRG, France gallagher@math.ens.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	András Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 3 2023

Global regularity for the nonlinear wave equation with slightly supercritical power MARIA COLOMBO and SILJA HAFFTER	613
Subelliptic wave equations are never observable CYRIL LETROUIT	643
Quantitative Alexandrov theorem and asymptotic behavior of the volume preserving mean curvature flow VESA JULIN and JOONAS NIINIKOSKI	679
A simple nuclear C^* -algebra with an internal asymmetry ILAN HIRSHBERG and N. CHRISTOPHER PHILLIPS	711
Partial regularity of Leray–Hopf weak solutions to the incompressible Navier–Stokes equations with hyperdissipation WOJCIECH S. OZAŃSKI	747
The Peskin problem with viscosity contrast EDUARDO GARCÍA-JUÁREZ, YOICHIRO MORI and ROBERT M. STRAIN	785
Solution of the qc Yamabe equation on a 3-Sasakian manifold and the quaternionic Heisenberg group STEFAN IVANOV, IVAN MINCHEV and DIMITER VASSILEV	839
Garland’s method with Banach coefficients IZHAR OPPENHEIM	861