

ANALYSIS & PDE

Volume 16

No. 5

2023

ROWAN KILLIP, MARIA NTEKOUME AND MONICA VIŞAN

**ON THE WELL-POSEDNESS PROBLEM FOR THE
DERIVATIVE NONLINEAR SCHRÖDINGER EQUATION**

ON THE WELL-POSEDNESS PROBLEM FOR THE DERIVATIVE NONLINEAR SCHRÖDINGER EQUATION

ROWAN KILLIP, MARIA NTEKOUME AND MONICA VIŞAN

We consider the derivative nonlinear Schrödinger equation in one space dimension, posed both on the line and on the circle. This model is known to be completely integrable and L^2 -critical with respect to scaling. We first discuss whether ensembles of orbits with L^2 -equicontinuous initial data remain equicontinuous under evolution. We prove that this is true under the restriction $M(q) = \int |q|^2 < 4\pi$. We conjecture that this restriction is unnecessary. Further, we prove that the problem is globally well posed for initial data in $H^{1/6}$ under the same restriction on M . Moreover, we show that this restriction would be removed by a successful resolution of our equicontinuity conjecture.

1. Introduction

The derivative nonlinear Schrödinger equation

$$iq_t + q'' + i(|q|^2 q)' = 0 \quad (\text{DNLS})$$

describes the evolution of a complex-valued field q defined either on the line \mathbb{R} or the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. This equation was introduced as an effective model in magnetohydrodynamics; see [Ichikawa and Watanabe 1977; Mio et al. 1976; Mjølhus 1976]. It was soon shown to be completely integrable [Kaup and Newell 1978] and has received enduring attention since that time.

As we shall document more fully below, well-posedness questions for (DNLS), particularly *global* well-posedness, have been particularly stubborn. Local well-posedness is already very challenging: the nonlinearity contains a full derivative, like KdV or mKdV, while the linear part gives only Schrödinger-like smoothing.

The task of converting local into global well-posedness is typically a matter of exploiting conservation laws. As a completely integrable system, (DNLS) has an infinite family of conserved quantities. The first three are as follows:

$$M(q) = \int |q(x)|^2 dx, \quad (1-1)$$

$$H(q) = -\frac{1}{2} \int i(q\bar{q}' - \bar{q}q') + |q|^4 dx, \quad (1-2)$$

$$H_2(q) = \int |q'|^2 + \frac{3}{4}i|q|^2(q\bar{q}' - \bar{q}q') + \frac{1}{2}|q|^6 dx. \quad (1-3)$$

Killip was supported by NSF grant DMS-1856755 and Vişan by grant DMS-1763074.
MSC2020: 35Q55.

Keywords: derivative nonlinear Schrödinger equation, well-posedness.

The striking fact about (DNLS) is that, with the exception of $M(q)$, none of the Hamiltonians in the hierarchy are coercive. Indeed, algebraic solitons have $M = 4\pi$ but all other Hamiltonians are identically zero. Applying the scaling symmetry

$$q(t, x) \mapsto \sqrt{\lambda} q(\lambda^2 t, \lambda x) \quad (1-4)$$

to an algebraic soliton yields a one-parameter family of solutions with identical values for *all* the conserved quantities. However, this family is unbounded in H^s for every $s > 0$.

The quantity $H(q)$ serves as the Hamiltonian for (DNLS) with respect to the Poisson structure

$$\{F, G\} = \int \frac{\delta F}{\delta q} \left(\frac{\delta G}{\delta \bar{q}} \right)' + \frac{\delta F}{\delta \bar{q}} \left(\frac{\delta G}{\delta q} \right)' dx, \quad (1-5)$$

while $M(q)$ generates translations, albeit at speed 2. Although the momentum is given by $\frac{1}{2}M(q)$, our definition of M leads to a more seamless connection to the existing literature.

Given that $M(q)$ is invariant under both (DNLS) and the scaling (1-4), it is natural to ask whether or not (DNLS) is well posed in L^2 . This is not known. Indeed, the existing local well-posedness theory requires H^s initial data with $s \geq \frac{1}{2}$. (We will make some further progress on this question in this paper.) It is important to recognize that because $M(q)$ is scaling critical, the mere fact that it forms a coercive conservation law would not suffice to render local well-posedness in L^2 automatically global. One must fear the solution concentrates at one (or more) points in space, a scenario known as type-II blowup. We do not believe this happens.

Conjecture 1.1. *For any $Q \subseteq \mathcal{S}$ that is L^2 -bounded and equicontinuous, the totality of states reached by (DNLS) orbits originating from Q , that is*

$$Q_* = \{e^{tJ\nabla H} q : q \in Q \text{ and } t \in \mathbb{R}\}, \quad (1-6)$$

is also L^2 -equicontinuous.

Here \mathcal{S} denotes Schwartz class in the line case and C^∞ on the torus. In the line case, recent works (discussed below) guarantee that all such initial data lead to global Schwartz solutions. The analogous claim is unknown on the torus, though we believe it to be true. Nevertheless, one can still ask if equicontinuity holds for as long as the orbits do exist. By the arguments presented in this paper, solutions cannot break down without losing equicontinuity. Therefore, a positive resolution of the conjecture for such partial solutions would already guarantee that they are global and so settle the conjecture in its entirety; see Corollary 4.2.

We phrased the conjecture in terms of \mathcal{S} initial data because it is a class that is dense in all relevant spaces. It also serves to emphasize that the central question to be addressed is not inherently tied to low regularity.

Equicontinuity in L^2 is most easily understood via Fourier transformation: it means that $|\hat{q}|^2$ forms a tight family of measures. Notice that, in view of the uncertainty principle, concentration on the physical side must be accompanied by a loss of tightness on the Fourier side.

In setting this conjecture, we have in mind four principal reasons: (1) It is challenging, yet recent developments give us hope for a successful resolution. (2) It encapsulates a single essential obstacle, namely, understanding conservation laws for (DNLS). (3) A proof of this conjecture would have significant consequences for the well-posedness problem. Indeed, such equicontinuity results form an essential part of a recent program developed in [Bringmann et al. 2021; Harrop-Griffiths et al. 2020; Killip and Viřan 2019] that has proved successful in obtaining optimal well-posedness results for completely integrable PDE. (4) We are able to verify that it is true in the regime $M(q) < 4\pi$; see Theorem 1.3 below.

Given the nature of completely integrable systems, it is natural to imagine that an equicontinuity conjecture of the same form holds for all other PDE in the (DNLS) hierarchy. Indeed, we truly believe that this is so and will shortly formulate just such a conjecture. However, the particular claim that we believe will be of greatest use in understanding the hierarchy is best expressed through the perturbation determinant. Let us turn our attention now to presenting this object, beginning with the requisite background.

The Lax pair introduced by Kaup and Newell [1978] for (DNLS) employs

$$L_{\text{KN}} = \begin{bmatrix} -i\lambda^2 - \partial & \lambda q \\ -\lambda \bar{q} & i\lambda^2 - \partial \end{bmatrix}.$$

For what follows, it will be convenient to make some cosmetic changes to this choice. Specifically, we set $\lambda = e^{i\pi/4} \sqrt{\kappa}$ with $\kappa \geq 1$ and replace $e^{i\pi/4} q \mapsto q$. This yields

$$L(\kappa) := \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \kappa - \partial & \sqrt{\kappa} q \\ i\sqrt{\kappa} \bar{q} & \kappa + \partial \end{bmatrix} \quad \text{and, for } q \equiv 0, \quad L_0(\kappa) := \begin{bmatrix} \kappa - \partial & 0 \\ 0 & -(\kappa + \partial) \end{bmatrix}.$$

These modifications maintain the crucial property that for smooth functions,

$$q(t) \text{ solves (DNLS)} \iff \frac{d}{dt} L(t; \kappa) = [P(t; \kappa), L(t; \kappa)],$$

where

$$P(\kappa) = \begin{bmatrix} 2i\kappa^2 - \kappa|q|^2 & 2i\kappa^{3/2}q - \kappa^{1/2}|q|^2q + i\kappa^{1/2}q' \\ 2\kappa^{3/2}\bar{q} + i\kappa^{1/2}|q|^2\bar{q} - \kappa^{1/2}\bar{q}' & -2i\kappa^2 + \kappa|q|^2 \end{bmatrix}.$$

This guarantees that the Lax operators L at different times are conjugate, at least formally. This in turn suggests that the perturbation determinant $\det[L_0^{-1}(\kappa)L(\kappa)]$ should be well defined and conserved by the flow.

To make this precise, it is convenient for us to mimic the analysis of the AKNS-ZS system employed in [Killip et al. 2018]: Let us first define $(\kappa \pm \partial)^{-1/2}$ as the Fourier multipliers $(\kappa \pm i\xi)^{-1/2}$, where the complex square root is determined by $\sqrt{\kappa} > 0$ and continuity. We then define

$$\Lambda(q) := (\kappa - \partial)^{-1/2} q (\kappa + \partial)^{-1/2} \quad \text{and} \quad \Gamma(q) := (\kappa + \partial)^{-1/2} \bar{q} (\kappa - \partial)^{-1/2}, \tag{1-7}$$

which are Hilbert–Schmidt operators for $q \in L^2$; see Lemma 2.1. Thus

$$a(\kappa; q) = \det[1 - i\kappa \Lambda \Gamma] \tag{1-8}$$

is well defined for $q \in L^2$ (and extends holomorphically to all $\text{Re } \kappa > 0$); moreover, for $q \in S$ it agrees with the formal notion of the perturbation determinant.

While $a(\kappa)$ does encode all the Hamiltonians of the (DNLS) hierarchy, this is more easily seen through its logarithm,

$$\alpha(\kappa; q) := -\log[a(\kappa; q)] = \sum_{\ell \geq 1} \frac{1}{\ell} \operatorname{tr}\{(i\kappa \Lambda \Gamma)^\ell\}, \quad (1-9)$$

which serves as a generating function for these conservation laws. Due to the possibility of $a(\kappa)$ vanishing, $\alpha(\kappa)$ may not be defined for all $\kappa \geq 1$. Nevertheless, the series in (1-9) does converge for fixed $q \in L^2$ and κ sufficiently large; see Proposition 2.6.

We have not yet addressed the conservation of $a(\kappa; q)$ under the (DNLS) flow. In the line case, this could be effected by demonstrating that $a(\kappa; q)$ coincides with the reciprocal of the transmission coefficient and then appealing to the inverse scattering theory. However, two direct proofs have appeared recently in the literature: Klaus and Schippa [2022] argued by differentiating the series (following a model introduced in [Killip et al. 2018]), while Tang and Xu [2021] developed a microscopic representation of this conservation law (in the style of [Harrop-Griffiths et al. 2020]). While these papers impose a small $M(q)$ requirement, this is solely to guarantee the convergence of the series (1-9). This issue is remedied by our Proposition 2.6.

To state the grand version of Conjecture 1.1, covering a wide range of commuting flows, let us first introduce a replacement for the set Q_* defined in (1-6). Given $q \in \mathcal{S}$, we first define

$$C_q = \{\tilde{q} \in \mathcal{S} : a(\kappa; \tilde{q}) = a(\kappa; q) \text{ for all } \kappa > 0\} \quad (1-10)$$

and write C_q^0 for the connected component (in the L^2 topology) of C_q containing q . Finally, given a set $Q \subseteq \mathcal{S}$, we define

$$Q_{**} = \bigcup_{q \in Q} C_q^0. \quad (1-11)$$

Conjecture 1.2. *If $Q \subseteq \mathcal{S}$ is L^2 -bounded and equicontinuous, then so too is the set Q_{**} defined in (1-11).*

We have several motivations in choosing connected components when defining Q_{**} . This formulation of the conjecture retains a vestige of the behavior of orbits, while emphasizing that this is a question about conservation laws and is ultimately independent of the well-posedness of any flow. Note also that while the zero solution and the family of algebraic solitons all share $a(\kappa) \equiv 1$, they are not in the same connected component under the (DNLS) hierarchy.

Our most compelling evidence in favor of these two conjectures is that both hold in the regime where $M(q) < 4\pi$.

Theorem 1.3. *Let $Q \subseteq \mathcal{S}$ be an L^2 -equicontinuous set satisfying*

$$\sup\{\|q\|_{L^2}^2 : q \in Q\} < 4\pi. \quad (1-12)$$

*Then the set Q_{**} defined in (1-11) is L^2 -bounded and equicontinuous.*

The significance of 4π is this: It is the value of M at which the polynomial conservation laws lose their efficacy. It is also the value of M for the algebraic soliton, which is maximal among all solitary

wave solutions. Unlike mass-critical NLS, (DNLS) admits solitons of arbitrarily small L^2 -norm, and consequently, there is no notion of a scattering threshold.

The proof of Theorem 1.3, which will be given in Section 3, is both short and simple. Indeed, the hypothesis (1-12) even allows us to forgo the restriction to connected components.

It has been observed before that $\text{tr}(i\kappa\Lambda\Gamma)$ may be used to understand how the L^2 -norm of q is distributed across frequencies (compare Lemma 2.2). The key observation that allows us to reach all the way to 4π (as opposed to mere smallness, compare [Klaus and Schippa 2022; Tang and Xu 2021]) is the manner in which we handle the remainder, specifically, the observation that the remainder may be summed in κ for any $q \in L^2$; see (3-8).

While the 4π restriction is crucial to our proof of Theorem 1.3, it does not play any role in our subsequent analysis of the consequences of such equicontinuity. For this reason, we introduce a general threshold M_* .

Definition 1.4. Let M_* denote the maximal constant such that for any L^2 -equicontinuous set $Q \subseteq \mathcal{S}$ satisfying

$$\sup\{\|q\|_{L^2}^2 : q \in Q\} < M_*, \tag{1-13}$$

the set defined in (1-11) is L^2 -equicontinuous.

Evidently, Theorem 1.3 shows that $M_* \geq 4\pi$ and we conjecture that $M_* = \infty$. Our primary contribution to the well-posedness problem is low-regularity well-posedness below the M_* threshold.

Theorem 1.5. Fix $\frac{1}{6} \leq s < \frac{1}{2}$. The (DNLS) evolution is globally well posed, both on the line and on the circle, in the space

$$B_{M_*}^s = \{q \in H^s : \|q\|_{L^2}^2 < M_*\} \tag{1-14}$$

endowed with the H^s topology.

A natural prerequisite for proving this theorem is a priori H^s bounds. In Section 4, we show how such bounds follow from L^2 -equicontinuity; see Theorem 4.3.

To prove Theorem 1.5 we employ the method of commuting flows introduced in [Killip and Viřan 2019]. In that paper, the method was used to prove well-posedness of the Korteweg–de Vries equation. It has also been adapted and extended to treat the well-posedness problem for other completely integrable PDE [Bringmann et al. 2021; Harrop-Griffiths et al. 2020], to prove symplectic non-squeezing [Ntekoume 2022], and to construct dynamics for KdV in thermal equilibrium [Killip et al. 2020].

In contrast to those papers, we do not employ a change of unknown; this simplifies some of the analysis. On the other hand, new difficulties attend the construction of regularized flows: Because they are rooted in $\alpha(\kappa; q)$, the regularized Hamiltonians $H_\kappa(q)$ cannot be defined throughout $B_{M_*}^0$ for any single value of κ . Instead, we need to use an exhaustion by equicontinuous subsets. Ultimately, these problems originate in the L^2 -criticality of the problem. Nevertheless, we will be able to prove that the regularized flows admit a satisfactory notion of well-posedness all the way down to L^2 ! The $s \geq \frac{1}{6}$ restriction arises later when we show that the regularized flows converge to the full (DNLS) evolution.

At this moment we do not know whether $s = \frac{1}{6}$ is sharp in either geometry or indeed, whether the threshold regularity will differ between the line and the circle. Moreover, we do not know of any results (in either geometry) that would preclude well-posedness all the way down to the scaling critical space L^2 . On the other hand, the self-similar solutions constructed in [Fujiwara et al. 2020] (see also [Kitaev 1985]) show that smooth solutions can break-down in a dramatic way if one permits mere weak- L^2 decay at spatial infinity.

The restriction $s < \frac{1}{2}$ in Theorem 1.5 does not represent a meaningful breakdown of our methods. However, treating larger values would require additional arguments. This seems unwarranted given that a great deal is already known about H^s -solutions for $s \geq \frac{1}{2}$, as we shall now discuss.

Local well-posedness in H^s for $s > \frac{3}{2}$ was proved by Tsutsumi and Fukuda [1980; 1981]. This was extended to $s \geq \frac{1}{2}$ by Takaoka [1999] for (DNLS) posed on the line and by Herr [2006] for the periodic problem. The endpoint $s = \frac{1}{2}$ is significant: for lesser s , the data-to-solution map can no longer be uniformly continuous on bounded sets; see [Biagioni and Linares 2001; Takaoka 1999].

Global well-posedness in $H^1(\mathbb{R})$ for initial data satisfying $M(q) < 2\pi$ was obtained by Hayashi and Ozawa [1992]. This result was extended first to $s > \frac{2}{3}$ and then to $s > \frac{1}{2}$ by Colliander, Keel, Staffilani, Takaoka, and Tao [Colliander et al. 2001; 2002], under the same L^2 restriction. See [Miao et al. 2011] for a refinement of these arguments to handle the endpoint case $s = \frac{1}{2}$, as well as [Takaoka 2001] for earlier efforts in this direction.

Hayashi and Ozawa [1992] also proved that solutions with initial data in \mathcal{S} remain in \mathcal{S} for as long as they remain bounded in H^1 .

Wu [2015] proved global well-posedness in $H^1(\mathbb{R})$ for initial data satisfying $M(q) < 4\pi$; see also his earlier work [Wu 2013] which first overcame the 2π barrier. An alternate variational proof was given in [Fukaya et al. 2017], which also constructed global solutions for highly modulated initial data of arbitrary L^2 size. The result in [Wu 2015] was extended to the periodic setting in [Mosincat and Oh 2015]. Finally, the argument in [Colliander et al. 2002] was further advanced in [Guo and Wu 2017; Mosincat 2017] to treat the endpoint case $s = \frac{1}{2}$ and $M(q) < 4\pi$; see also [Win 2010] for earlier work in the periodic setting.

We note that the results of this paper provide an alternate proof of the main results in [Mosincat and Oh 2015; Wu 2015]; see Corollary 4.2. In particular, Proposition 4.1 shows that H^1 bounds follow from Theorem 1.3.

The well-posedness of (DNLS) has also been investigated in Fourier–Lebesgue spaces; [Deng et al. 2021; Grünrock 2005; Grünrock and Herr 2008]. This allowed the authors to obtain a uniformly continuous data-to-solution map in spaces that are closer to the critical scaling; recall that this property breaks down in H^s spaces when $s < \frac{1}{2}$. An almost sure global well-posedness result for randomized initial data was proved in [Nahmod et al. 2012].

As a completely integrable PDE, (DNLS) is also amenable to inverse scattering techniques. Building on the pioneering work of Liu [2017], global well-posedness and asymptotic analysis of soliton-free solutions in $H^{2,2}(\mathbb{R}) = \{f \in H^2(\mathbb{R}) : x^2 f \in L^2(\mathbb{R})\}$ were addressed in [Liu et al. 2016; 2018].

Global well-posedness for *all* $H^{2,2}(\mathbb{R})$ initial data was proved by Jenkins, Liu, Perry, and Sulem in [Jenkins et al. 2020b]. This work builds on the authors' prior successes in [Jenkins et al. 2018b]. These

authors also proved a soliton resolution result [Jenkins et al. 2018a] for generic data in $H^{2,2}(\mathbb{R})$. See also their excellent review article [Jenkins et al. 2020a].

The inverse scattering approach was also applied by Pelinovsky and Shimabukuro [2018] to prove global well-posedness in $H^{1,1}(\mathbb{R}) \cap H^2(\mathbb{R})$ for soliton-free solutions and then in joint work with Saalmann [Pelinovsky et al. 2017] for data giving rise to finitely many solitons; see also [Saalmann 2017].

Recently there has been a surge of activity on the well-posedness problem for (DNLS). We first note the paper [Klaus and Schippa 2022], which showed a priori H^s bounds, $0 < s < \frac{1}{2}$, for solutions with $M(q)$ small. The smallness assumption allows them to guarantee that the series (1-9) converges rapidly for κ large, and so the series can be conflated with its first term. The paper [Tang and Xu 2021] presents a microscopic representation of the conservation of $\alpha(\kappa; q)$. In [Bahouri and Perelman 2022], the authors achieve the major breakthrough of proving that for every initial datum in $H^{1/2}(\mathbb{R})$, the orbit remains bounded in the same space (irrespective of the size of $M(q)$). For the periodic (DNLS), the paper [Isom et al. 2020] shows that for $s \geq 1$ and $M(q)$ small, the $H^s(\mathbb{T})$ -norm of solutions grows at most polynomially in time.

While these exciting results appeared too recently to affect what we do in this paper, their novelty and insightfulness give us every hope that the conjectures presented herein may soon be resolved.

2. Preliminaries

Our conventions for the Fourier transform are

$$\hat{f}(\xi) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-i\xi x} f(x) dx, \quad \text{so} \quad f(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{i\xi x} \hat{f}(\xi) d\xi$$

for functions on the line, and

$$\hat{f}(\xi) = \int_0^1 e^{-i\xi x} f(x) dx, \quad \text{so} \quad f(x) = \sum_{\xi \in 2\pi\mathbb{Z}} \hat{f}(\xi) e^{i\xi x}$$

for functions on the torus \mathbb{T} . These definitions of the Fourier transform are unitary on L^2 and yield the Plancherel identities

$$\|f\|_{L^2(\mathbb{R})} = \|\hat{f}\|_{L^2(\mathbb{R})} \quad \text{and} \quad \|f\|_{L^2(\mathbb{T})} = \sum_{\xi \in 2\pi\mathbb{Z}} |\hat{f}(\xi)|^2,$$

as well as the following convolution identity on \mathbb{R} :

$$\widehat{fg} = \frac{1}{\sqrt{2\pi}} \hat{f} * \hat{g}.$$

We use the standard Littlewood–Paley decomposition of a function,

$$q = \sum_{N \in 2^{\mathbb{N}}} q_N,$$

based on a smooth partition of unity on the Fourier side. Here q_1 denotes the projection onto frequencies $|\xi| \leq 1$; for $N \geq 2$, frequencies $|\xi| \sim N$ are contained in q_N .

The fact that the operators Λ and Γ defined in (1-7) are Hilbert–Schmidt was noticed already in [Killip et al. 2018, Lemma 4.1]:

Lemma 2.1. *For $q \in L^2$ and $\kappa > 0$, we have*

$$\|\Lambda\|_{\mathfrak{J}_2(\mathbb{R})}^2 = \|\Gamma\|_{\mathfrak{J}_2(\mathbb{R})}^2 \approx \int_{\mathbb{R}} \log\left(4 + \frac{\xi^2}{\kappa^2}\right) \frac{|\hat{q}(\xi)|^2}{\sqrt{4\kappa^2 + \xi^2}} d\xi \lesssim \kappa^{-1} \|q\|_{L^2}^2, \tag{2-1}$$

$$\|\Lambda\|_{\mathfrak{J}_2(\mathbb{T})}^2 = \|\Gamma\|_{\mathfrak{J}_2(\mathbb{T})}^2 \approx \sum_{\xi \in 2\pi\mathbb{Z}} \log\left(4 + \frac{\xi^2}{\kappa^2}\right) \frac{|\hat{q}(\xi)|^2}{\sqrt{4\kappa^2 + \xi^2}} \lesssim \kappa^{-1} \|q\|_{L^2}^2. \tag{2-2}$$

Proof. The estimate (2-1) follows from the computation

$$\|\Lambda\|_{\mathfrak{J}_2(\mathbb{R})}^2 = \frac{1}{2\pi} \int_{\mathbb{R}} |\hat{q}(\xi)|^2 \int_{\mathbb{R}} \frac{1}{\sqrt{\kappa^2 + \eta^2} \sqrt{\kappa^2 + (\eta + \xi)^2}} d\eta d\xi \approx \int_{\mathbb{R}} \log\left(4 + \frac{\xi^2}{\kappa^2}\right) \frac{|\hat{q}(\xi)|^2}{\sqrt{4\kappa^2 + \xi^2}} d\xi.$$

To compute the above integral in η , one treats separately the regions $|\eta| \leq 2|\xi|$ and $|\eta| > 2|\xi|$; the logarithm term arises only when considering the first region.

On the torus, similar arguments yield

$$\|\Lambda\|_{\mathfrak{J}_2(\mathbb{T})}^2 = \sum_{\xi \in 2\pi\mathbb{Z}} |\hat{q}(\xi)|^2 \sum_{\eta \in 2\pi\mathbb{Z}} \frac{1}{\sqrt{\kappa^2 + \eta^2} \sqrt{\kappa^2 + (\eta + \xi)^2}} \approx \sum_{\xi \in 2\pi\mathbb{Z}} \log\left(4 + \frac{\xi^2}{\kappa^2}\right) \frac{|\hat{q}(\xi)|^2}{\sqrt{4\kappa^2 + \xi^2}},$$

which settles (2-2). □

These Hilbert–Schmidt bounds ensure that $i\kappa\Lambda\Gamma$ is trace class and thus that the determinant in (1-8) is well defined. The trace of this operator will also be important and is easily evaluated.

Lemma 2.2. *Let $q \in L^2$ and $\kappa > 0$. Then*

$$\text{tr}(i\kappa\Lambda\Gamma) = \int \frac{i\kappa|\hat{q}(\xi)|^2}{2\kappa - i\xi} d\xi \quad \text{on } \mathbb{R}, \tag{2-3}$$

$$\text{tr}(i\kappa\Lambda\Gamma) = \frac{1 + e^{-\kappa}}{1 - e^{-\kappa}} \sum_{\xi \in 2\pi\mathbb{Z}} \frac{i\kappa|\hat{q}(\xi)|^2}{2\kappa - i\xi} \quad \text{on } \mathbb{T}. \tag{2-4}$$

Proof. To prove (2-3), we simply compute the trace on the Fourier side:

$$\text{tr}(i\kappa\Lambda\Gamma) = \frac{i\kappa}{2\pi} \iint \frac{|\hat{q}(\xi)|^2}{(\eta - i\kappa)(\eta + \xi + i\kappa)} d\eta d\xi = \int \frac{i\kappa|\hat{q}(\xi)|^2}{2\kappa - i\xi} d\xi.$$

In the circle setting, we use the partial fraction decomposition of the cotangent:

$$\sum_{\eta \in 2\pi\mathbb{Z}} \left(\frac{1}{\kappa + i\eta} + \frac{1}{\kappa - i\eta} \right) = \coth\left(\frac{\kappa}{2}\right) = \frac{1 + e^{-\kappa}}{1 - e^{-\kappa}}. \tag{2-5}$$

In this way, we find

$$\text{tr}(i\kappa\Lambda\Gamma) = i\kappa \sum_{\xi \in 2\pi\mathbb{Z}} |\hat{q}(\xi)|^2 \frac{1}{\xi + 2i\kappa} \sum_{\eta \in 2\pi\mathbb{Z}} \left(\frac{1}{\eta - i\kappa} - \frac{1}{\eta + \xi + i\kappa} \right) = \sum_{\xi \in 2\pi\mathbb{Z}} \frac{i\kappa|\hat{q}(\xi)|^2}{2\kappa - i\xi} \frac{1 + e^{-\kappa}}{1 - e^{-\kappa}}.$$

Notice that the sum over η simplifies to (2-5) because $\xi \in 2\pi\mathbb{Z}$. □

In Section 5, it will be convenient to express the next term in the series (1-9) as a paraproduct. This is the role of the next lemma.

Lemma 2.3. *Let $q \in L^2$ and $\kappa > 0$. Then*

$$\text{tr}([\Lambda\Gamma]^2) = \int_{\mathbb{R}} \left(\frac{1}{2\kappa + \partial} \bar{q}\right)^2 (4\kappa - \partial) \left(\frac{1}{2\kappa - \partial} q\right)^2 dx \quad \text{on } \mathbb{R}, \tag{2-6}$$

$$\text{tr}([\Lambda\Gamma]^2) = \frac{1 + e^{-\kappa}}{1 - e^{-\kappa}} \int_{\mathbb{T}} \left(\frac{1}{2\kappa + \partial} \bar{q}\right)^2 (4\kappa - \partial) \left(\frac{1}{2\kappa - \partial} q\right)^2 dx \quad \text{on } \mathbb{T}. \tag{2-7}$$

Proof. The method is exactly that of the previous lemma, only the details change. In the line case, we have a more complicated (but still elementary) contour integral. In the circle case, one must verify that

$$\begin{aligned} \sum_{\xi \in 2\pi\mathbb{Z}} \frac{1}{(\kappa + i\xi)(\kappa - i[\xi + \eta_1])(\kappa + i[\xi + \eta_1 + \eta_2])(\kappa - i[\xi + \eta_1 + \eta_2 + \eta_3])} \\ = \frac{1 + e^{-\kappa}}{1 - e^{-\kappa}} \cdot \frac{4\kappa - i(\eta_1 + \eta_3)}{(2\kappa - i\eta_1)(2\kappa + i\eta_2)(2\kappa - i\eta_3)(2\kappa + i\eta_4)}. \end{aligned}$$

This follows from (2-5) via a careful partial fraction decomposition. □

Our next lemma records operator estimates for frequency localized potentials.

Lemma 2.4 (operator estimates). *Fix $q \in L^2$, $N \in 2^{\mathbb{N}}$, and $\kappa \geq 1$, and write $\Lambda_N = \Lambda(q_N)$ and $\Gamma_N = \Gamma(q_N)$. Then*

$$\|\Lambda_N\|_{\mathfrak{S}_2} = \|\Gamma_N\|_{\mathfrak{S}_2} \approx \sqrt{\frac{1}{\kappa + N} \log\left(4 + \frac{N^2}{\kappa^2}\right)} \|q_N\|_{L^2}, \tag{2-8}$$

$$\|\Lambda_N\|_{\text{op}} = \|\Gamma_N\|_{\text{op}} \lesssim \min\left\{\frac{\sqrt{N}}{\kappa}, \sqrt{\frac{1}{\kappa + N} \log\left(4 + \frac{N^2}{\kappa^2}\right)}\right\} \|q_N\|_{L^2}, \tag{2-9}$$

$$\sum_{N \leq N_0} \|\Lambda_N\|_{\text{op}} \lesssim \kappa^{-1} \min\{\sqrt{N_0}, \sqrt{\kappa}\} \|q\|_{L^2}. \tag{2-10}$$

Proof. The claim (2-8) follows immediately from Lemma 2.1.

Using the Bernstein inequality, we estimate

$$\|\Lambda_N\|_{\text{op}} \leq \|(\kappa - \partial)^{-1/2}\|_{\text{op}} \|q_N\|_{\text{op}} \|(\kappa + \partial)^{-1/2}\|_{\text{op}} \leq \frac{1}{\kappa} \|q_N\|_{L^\infty} \lesssim \frac{\sqrt{N}}{\kappa} \|q_N\|_{L^2}.$$

Combining this with (2-8) yields (2-9).

The case $N_0 \leq \kappa$ of (2-10) is clear. If $N_0 > \kappa$, an application of (2-9) yields

$$\sum_{N \leq N_0} \|\Lambda_N\|_{\text{op}} \lesssim \sum_{N \leq \kappa} \frac{\sqrt{N}}{\kappa} \|q\|_{L^2} + \sum_{\kappa < N \leq N_0} \sqrt{\frac{1}{N} \log\left(4 + \frac{N^2}{\kappa^2}\right)} \|q\|_{L^2} \lesssim \frac{\sqrt{\kappa}}{\kappa} \|q\|_{L^2},$$

as desired. □

Lemma 2.5. *For all $\kappa \geq 1$, we have*

$$\|(\kappa + \partial)^{-1} f(\kappa - \partial)^{-1}\|_{\mathfrak{J}_2} \lesssim \kappa^{-1/2} \|f\|_{H^{-1}}, \tag{2-11}$$

$$\|q(\kappa + \partial)^{-3/4}\|_{\mathfrak{J}_2} \lesssim \kappa^{-1/4} \|q\|_{L^2}, \tag{2-12}$$

$$\|(\kappa - \partial)^{-1/4} q(\kappa + \partial)^{-1/4}\|_{\text{op}} \lesssim \|q\|_{L^2}. \tag{2-13}$$

Proof. We first turn to (2-11). We will only consider here the line setting; in the periodic case, one can apply a similar argument to the one in the proof of Lemma 2.1. A straightforward computation yields

$$\|(\kappa + \partial)^{-1} f(\kappa - \partial)^{-1}\|_{\mathfrak{J}_2}^2 = \frac{1}{2\pi} \iint \frac{|\hat{f}(\xi)|^2}{[\kappa^2 + (\xi + \eta)^2](\kappa^2 + \eta^2)} d\eta d\xi.$$

Considering separately the regions $|\eta| \leq 2|\xi|$ and $|\eta| > 2|\xi|$ when integrating in η , we find

$$\|(\kappa + \partial)^{-1} f(\kappa - \partial)^{-1}\|_{\mathfrak{J}_2}^2 \lesssim \int \frac{|\hat{f}(\xi)|^2}{\kappa(\kappa^2 + \xi^2)} d\xi \lesssim \kappa^{-1} \|f\|_{H^{-1}}^2.$$

By direct computation (compare [Simon 2005, Theorem 4.1]), we have

$$\|q(\kappa + \partial)^{-3/4}\|_{\mathfrak{J}_2} \lesssim \|q\|_{L^2} \|(\kappa + i\xi)^{-3/4}\|_{L^2_\xi} \lesssim \kappa^{-1/4} \|q\|_{L^2},$$

which settles (2-12).

Similarly, by Cwikel’s theorem (see [Cwikel 1977] or [Simon 2005, Theorem 4.2]), we find that

$$\begin{aligned} \|(\kappa - \partial)^{-1/4} q(\kappa + \partial)^{-1/4}\|_{\text{op}} &\leq \|(\kappa - \partial)^{-1/4} \sqrt{|q|}\|_{\text{op}} \left\| \frac{q}{\sqrt{|q|}} (\kappa + \partial)^{-1/4} \right\|_{\text{op}} \\ &\lesssim \|(\kappa \pm i\xi)^{-1/4}\|_{L^4_{\text{weak}}}^2 \|\sqrt{|q|}\|_{L^4}^2 \lesssim \|q\|_{L^2}. \end{aligned} \quad \square$$

Proposition 2.6. *Let Q be a bounded and equicontinuous subset of L^2 . Then*

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q} \sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} = 0. \tag{2-14}$$

Moreover, there exists $\kappa_0 \geq 1$ such that the series (1-9) converges uniformly for $\kappa \geq \kappa_0$ and $q \in Q$.

Proof. Fix $\varepsilon > 0$ and let $\eta > 0$ be a small parameter to be chosen later. Using (2-10) and Lemma 2.1, we get

$$\sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} \lesssim \sqrt{\kappa} \|\Lambda(q_{>\eta\kappa})\|_{\text{op}} + \sqrt{\kappa} \sum_{N \leq \eta\kappa} \|\Lambda_N(q)\|_{\text{op}} \lesssim \|q_{>\eta\kappa}\|_{L^2} + \sqrt{\eta} \|q\|_{L^2}.$$

Choosing η small enough depending on the L^2 bound of Q , and then κ sufficiently large depending on η and the equicontinuity property of Q , we may ensure that

$$\sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} < \varepsilon \quad \text{for all } q \in Q,$$

which yields (2-14).

To continue, we choose κ_0 sufficiently large such that for any $\kappa \geq \kappa_0$ we have $\sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} \leq \frac{1}{2}$ uniformly for $q \in Q$. Lemma 2.1 then yields

$$\|(i\kappa \Lambda \Gamma)^{\ell+1}\|_{\mathfrak{J}_1} \leq \kappa^{\ell+1} \|\Lambda\|_{\mathfrak{J}_2}^2 \|\Lambda\|_{\text{op}}^{2\ell} \lesssim 2^{-\ell} \|q\|_{L^2}^2, \tag{2-15}$$

uniformly for $\kappa \geq \kappa_0$ and $q \in Q$, which ensures convergence of the series (1-9). □

As discussed in the introduction, this convergence result allows the arguments of [Klaus and Schippa 2022; Tang and Xu 2021] to be extended beyond the regime of small L^2 -norm and so show that $\alpha(\kappa; q)$ is conserved under the (DNLS) flow, for κ sufficiently large. This conservation is inherited by $a(\kappa; q)$ for all $\text{Re } \kappa > 0$ because this is a holomorphic function in this region.

3. Equicontinuity in L^2

The goal of this section is to prove Theorem 1.3. We begin with a convenient notion of the momentum at high frequencies in each geometry:

$$\beta_{\mathbb{R}}^{[2]}(\kappa; q) := \int_{\mathbb{R}} \frac{\xi^2 |\hat{q}(\xi)|^2}{4\kappa^2 + \xi^2} d\xi \quad \text{and} \quad \beta_{\mathbb{T}}^{[2]}(\kappa; q) := \sum_{\xi \in 2\pi\mathbb{Z}} \frac{\xi^2 |\hat{q}(\xi)|^2}{4\kappa^2 + \xi^2}. \tag{3-1}$$

The curious notation is explained by the fact that these expressions coincide with the quadratic (in q) parts of the quantities in (4-3). For our immediate purposes, however, the following relation with the formulas of Lemma 2.2 is more important:

$$\begin{aligned} \text{Im tr}(i\kappa \Lambda \Gamma) &= \frac{1}{2} [M(q) - \beta_{\mathbb{R}}^{[2]}(\kappa; q)] && \text{on } \mathbb{R}, \\ \text{Im tr}(i\kappa \Lambda \Gamma) &= \frac{1}{2} \frac{1+e^{-\kappa}}{1-e^{-\kappa}} [M(q) - \beta_{\mathbb{T}}^{[2]}(\kappa; q)] && \text{on } \mathbb{T}. \end{aligned} \tag{3-2}$$

Given an infinite subset $\mathcal{K} \subseteq 2^{\mathbb{N}}$, we then define a norm via

$$\|q\|_{\mathcal{K}}^2 := \|q\|_{L^2}^2 + \sum_{\kappa \in \mathcal{K}} \beta^{[2]}(\kappa; q). \tag{3-3}$$

This in turn leads to a very convenient formulation of equicontinuity.

Lemma 3.1. *A set $Q \subseteq L^2$ is bounded and equicontinuous if and only if there exists an infinite set $\mathcal{K} \subseteq 2^{\mathbb{N}}$ such that $\sup_{q \in Q} \|q\|_{\mathcal{K}} < \infty$.*

Proof. This is immediately evident from the observation that

$$\|q\|_{\mathcal{K}}^2 \approx \|q\|_{L^2}^2 + \sum_{\kappa \in \mathcal{K}} \|q_{>\kappa}\|_{L^2}^2 \approx \|q\|_{L^2}^2 + \sum_{N \in 2^{\mathbb{N}}} \#\{\kappa \in \mathcal{K} : \kappa < N\} \|q_N\|_{L^2}^2. \tag{3-4}$$

Before beginning the proof of Theorem 1.3, we need two further preliminaries. The first will allow us to pass from the determinant to the exponentiated trace, and the second to take logarithms.

Lemma 3.2. *Let $A \in \mathfrak{J}_1$. Then*

$$|\det(1 + A) - \exp\{\text{tr}(A)\}| \leq \frac{1}{2} \|A\|_{\mathfrak{J}_2}^2 \exp\{\|A\|_{\mathfrak{J}_1}\}. \tag{3-4}$$

Proof. Let λ_i enumerate the nonzero eigenvalues of A repeated according to algebraic multiplicity. By relating eigenvalues and singular values, Weyl proved that

$$\sum |\lambda_i| \leq \|A\|_{\mathfrak{J}_1} \quad \text{and} \quad \sum |\lambda_i|^2 \leq \|A\|_{\mathfrak{J}_2}^2.$$

Now let us compare

$$\det(1 + A) = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{\substack{i_1, \dots, i_n \\ \text{distinct}}} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_n}, \quad \text{and} \quad \exp\{\text{tr}(A)\} = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1, \dots, i_n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_n}.$$

Evidently, the difference contains only sums over n -tuples (i_1, \dots, i_n) that contain at least one pair of identical indices. Thus,

$$\text{LHS of (3-4)} \leq \sum_{n=2}^{\infty} \frac{1}{n!} \binom{n}{2} \left[\sum_j |\lambda_j|^2 \right] \left[\sum_i |\lambda_i| \right]^{n-2} \leq \frac{1}{2} \|A\|_{\mathfrak{J}_2}^2 \sum_{n=2}^{\infty} \frac{1}{(n-2)!} \|A\|_{\mathfrak{J}_1}^{n-2},$$

and so (3-4) follows. □

Lemma 3.3. *Given $C > 0$ and $0 < \varepsilon < \pi$, let*

$$\mathcal{R} = \{z : |\text{Re } z| \leq C \text{ and } 0 < \text{Im } z < 2\pi - \varepsilon\}. \tag{3-5}$$

Then

$$|\text{Im}(z - w)| \leq \frac{\pi e^C}{\sin(\varepsilon/2)} |e^w - e^z| \quad \text{uniformly for } z, w \in \mathcal{R}. \tag{3-6}$$

Proof. This reduces to elementary trigonometry once one realizes that the worst-case scenario is $\text{Re } z = \text{Re } w = -C$. □

We are now ready for the climax of the section.

Proof of Theorem 1.3. Let us begin right away with the key computation. Given any $q \in L^2$, we may apply (2-8), (2-10), and (in the final step) Cauchy–Schwarz to deduce that

$$\begin{aligned} \sum_{\kappa \in 2^{\mathbb{N}}} \|i\kappa \Lambda(q) \Gamma(q)\|_{\mathfrak{J}_2}^2 &\lesssim \sum_{\kappa \in 2^{\mathbb{N}}} \kappa^2 \sum_{N_1 \sim N_2 \geq N_3, N_4} \|\Lambda_{N_1}(q)\|_{\mathfrak{J}_2} \|\Lambda_{N_2}(q)\|_{\mathfrak{J}_2} \|\Lambda_{N_3}(q)\|_{\text{op}} \|\Lambda_{N_4}(q)\|_{\text{op}} \\ &\lesssim M(q) \sum_{\kappa \in 2^{\mathbb{N}}} \sum_{N_1 \sim N_2} \frac{1}{N_2 + \kappa} \log\left(4 + \frac{N_2^2}{\kappa^2}\right) \|q_{N_1}\|_{L^2} \|q_{N_2}\|_{L^2} \min\{N_2, \kappa\} \\ &\lesssim M(q) \sum_{N_1 \sim N_2} \|q_{N_1}\|_{L^2} \|q_{N_2}\|_{L^2} \left(\sum_{\kappa \leq N_2} \frac{\kappa}{N_2} \log\left(4 + \frac{N_2^2}{\kappa^2}\right) + \sum_{\kappa > N_2} \frac{N_2}{\kappa} \right) \\ &\lesssim M(q)^2. \end{aligned} \tag{3-7}$$

Combining this with Lemmas 2.1 and 3.2, we find

$$\sum_{\kappa \in 2^{\mathbb{N}}} |a(\kappa; q) - \exp\{-\text{tr}[i\kappa \Lambda(q) \Gamma(q)]\}| \leq CM(q)^2 e^{CM(q)} \tag{3-8}$$

for some absolute C .

As we did not explicitly require that $M(\tilde{q}) = M(q)$ for $\tilde{q} \in C_q^0$, let us pause to see that this follows from the equality $a(\kappa; \tilde{q}) \equiv a(\kappa; q)$. From (3-4) and Lemma 2.2 we see that for $\kappa \rightarrow \infty$,

$$\begin{aligned} 0 &= |a(\kappa; \tilde{q}) - a(\kappa; q)| = \left| \exp\{-\operatorname{tr}[i\kappa \Lambda(\tilde{q})\Gamma(\tilde{q})]\} - \exp\{-\operatorname{tr}[i\kappa \Lambda(q)\Gamma(q)]\} \right| + o(1) \\ &= \left| \exp\left\{-\frac{i}{2}M(\tilde{q})\right\} - \exp\left\{-\frac{i}{2}M(q)\right\} \right| + o(1). \end{aligned}$$

Thus $M(\tilde{q})$ is preserved modulo $4\pi\mathbb{Z}$. As \tilde{q} belongs to the same connected component as q , we must have that $M(\tilde{q}) = M(q)$. For later use, we note the consequence

$$\sup_{q \in Q_{**}} M(q) = \sup_{q \in Q} M(q). \tag{3-9}$$

While this argument did not require the hypothesis (1-12), we will need it to unwrap this phase ambiguity when we address equicontinuity. This is our next topic.

Given an equicontinuous set Q satisfying (1-12), choose $\varepsilon > 0$ and an infinite subset $\mathcal{K} \subseteq 2^{\mathbb{N}}$ such that

$$\sup_{q \in Q, \kappa \in \mathcal{K}} \frac{1+e^{-\kappa}}{1-e^{-\kappa}} M(q) \leq 4\pi - 2\varepsilon \quad \text{and} \quad \sup_{q \in Q} \|q\|_{\mathcal{K}} < \infty.$$

Proceeding very much as we did above, we see that

$$\sum_{\kappa \in \mathcal{K}} \left| \exp\{-\operatorname{tr}[i\kappa \Lambda(\tilde{q})\Gamma(\tilde{q})]\} - \exp\{-\operatorname{tr}[i\kappa \Lambda(q)\Gamma(q)]\} \right| \leq 2CM(q)^2 e^{CM(q)}$$

for any $\tilde{q} \in C_q^0$. Combining this with (3-2) and Lemma 3.3, we deduce that

$$\sum_{\kappa \in \mathcal{K}} |\beta^{[2]}(\kappa; \tilde{q}) - \beta^{[2]}(\kappa; q)| \lesssim_{\varepsilon} 1.$$

This in turn guarantees that

$$\sup\{\|\tilde{q}\|_{\mathcal{K}}^2 : \tilde{q} \in Q_{**}\} \leq \sup\{\|q\|_{\mathcal{K}}^2 : q \in Q\} + O_{\varepsilon}(1) < \infty,$$

from which equicontinuity follows via Lemma 3.1. □

4. Conservation laws and equicontinuity

The primary goal of this section is to prove H^s bounds for (DNLS) solutions, for $0 < s < \frac{1}{2}$, as a prerequisite for proving Theorem 1.5. In addition, we will prove equicontinuity in these spaces, which is also needed to prove that theorem.

Before turning to that subject, we pause to show how L^2 -equicontinuity can be used to restore coercivity to the traditional polynomial conservation laws. As a representative example, we show how $H_2(q)$ can be used to control the H^1 -norm.

Proposition 4.1. *Let $Q \subseteq H^1$ be L^2 -bounded and equicontinuous. Then*

$$\|q\|_{H^1}^2 \lesssim H_2(q) + M(q)^3, \tag{4-1}$$

uniformly for all $q \in Q$.

Proof. Splitting into low and high frequency parts and estimating using the Bernstein and Gagliardo–Nirenberg inequalities, respectively, we obtain

$$\|q\|_{L^6}^6 \lesssim \|q_{\leq N}\|_{L^6}^6 + \|q_{>N}\|_{L^6}^6 \lesssim N^2 \|q\|_{L^2}^6 + \|q_{>N}\|_{L^2}^4 \|q'\|_{L^2}^2.$$

This allows us to control the quartic term in H_2 , and hence the H^1 -norm, as follows:

$$\begin{aligned} \|q'\|_{L^2}^2 &\leq H_2(q) + \frac{3}{2} \int |q(x)|^3 |q'(x)| dx \\ &\leq H_2(q) + \varepsilon \|q'\|_{L^2}^2 + \frac{9}{16\varepsilon} \|q\|_{L^6}^6 \\ &\leq H_2(q) + \left(\varepsilon + C \frac{9}{16\varepsilon} \|q_{>N}\|_{L^2}^4\right) \|q'\|_{L^2}^2 + C \frac{9}{16\varepsilon} N^2 M(q)^3, \end{aligned}$$

for any $\varepsilon > 0$. The claim (4-1) now follows by choosing ε small and then N large, exploiting the equicontinuity of Q . □

Proposition 4.1 allows us to extend local H^1 solutions globally in time, provided we remain below the M_* bound introduced in Definition 1.4.

Corollary 4.2. *The (DNLS) evolution is globally well posed, both on the line and on the circle, in the space*

$$B_{M_*}^1 = \{q \in H^1 : \|q\|_{L^2}^2 < M_*\} \tag{4-2}$$

endowed with the H^1 topology. Moreover, initial data in \mathcal{S} leads to solutions that belong to \mathcal{S} at all times.

Proof. In the line case, this result can be deduced from [Bahouri and Perelman 2022]; indeed, the restriction $M(q) < M_*$ is not needed in this case. Below we give an alternate argument that works also in the periodic setting.

As discussed in the introduction, local well-posedness in H^1 was proved already in [Takaoka 1999; Herr 2006]. Thus, given initial data $q(0) \in B_{M_*}^1 \cap \mathcal{S}$, there is a corresponding maximal lifespan solution $q \in C_t([0, T); H^1)$ to (DNLS). Moreover, [Hayashi and Ozawa 1992] shows that $q(t) \in \mathcal{S}$ for all $t \in [0, T)$. Combining [Klaus and Schippa 2022; Tang and Xu 2021] with Proposition 2.6 yields that $a(\kappa; q(t)) = a(\kappa, q(0))$ for all $t \in [0, T)$ and $\kappa > 0$. By the definition of M_* and Proposition 4.1, the solution q satisfies a priori H^1 bounds on $[0, T)$, which in turn guarantees that $T = \infty$.

Finally, global well-posedness in $B_{M_*}^1$ follows from local well-posedness and the density of \mathcal{S} in H^1 . □

Let us now turn to low-regularity questions. Bounded sets in H^s , with $s > 0$, are automatically bounded and equicontinuous in L^2 . As we shall work only below the M_* threshold in this section, such L^2 -equicontinuity is retained globally in time. Our goal is to propagate H^s bounds. The key to doing this is a certain renormalization of $\alpha(\kappa; q)$ that we introduce now:

$$\begin{aligned} \beta_{\mathbb{R}}(\kappa; q) &:= \|q\|_{L^2}^2 - 2 \operatorname{Im} \alpha(\kappa; q) && \text{on } \mathbb{R}, \\ \beta_{\mathbb{T}}(\kappa; q) &:= \|q\|_{L^2}^2 - \frac{1-e^{-\kappa}}{1+e^{-\kappa}} 2 \operatorname{Im} \alpha(\kappa; q) && \text{on } \mathbb{T}. \end{aligned} \tag{4-3}$$

Proposition 2.6 guarantees that these quantities are well defined for κ sufficiently large across our whole family of orbits.

The quadratic (in q) parts of these expressions were presented already in (3-1). As we saw there, these provide a sense of the L^2 -norm of the high-frequency part of q . To address higher regularity, for $0 < s < \frac{1}{2}$ we consider the quantity

$$\beta_s(\kappa; q) := \int_{\kappa}^{\infty} \beta(x; q) x^{2s} \frac{dx}{x}.$$

The quadratic term in this expression is given by

$$\beta_s^{[2]}(\kappa; q) = \int_{\kappa}^{\infty} \beta^{[2]}(x; q) x^{2s} \frac{dx}{x} = \int_{\kappa}^{\infty} \left\langle \frac{-\partial^2}{4x^2 - \partial^2} q, q \right\rangle x^{2s} \frac{dx}{x} \approx_s \left\langle \frac{-\partial^2}{(\kappa^2 - \partial^2)^{1-s}} q, q \right\rangle.$$

From this we see that for any $0 < \eta < 1$,

$$\|q_{>\kappa}\|_{H^s}^2 \lesssim \beta_s^{[2]}(\kappa; q) \lesssim \eta^{2(1-s)} \|q\|_{H^s}^2 + \|q_{>\eta\kappa}\|_{H^s}^2, \tag{4-4}$$

and so $\beta_s^{[2]}(\kappa; q)$ captures the H^s -norm of the high-frequency part of q . Indeed, a bounded set $Q \subseteq H^s$ is equicontinuous in H^s if and only if $\beta_s^{[2]}(\kappa; q) \rightarrow 0$ uniformly on Q as $\kappa \rightarrow \infty$.

Theorem 4.3. Fix $0 < s < \frac{1}{2}$ and let $Q \subseteq S$ be H^s -bounded and satisfy (1-13). Then, recalling the notation Q_{**} from (1-11), we have

$$\sup_{q \in Q_{**}} \|q\|_{H^s} \lesssim C \left(\sup_{q \in Q} \|q\|_{L^2}, \sup_{q \in Q} \|q\|_{H^s} \right). \tag{4-5}$$

Moreover, if Q is H^s -equicontinuous, then so is Q_{**} .

Proof. As Q is H^s -bounded, it is automatically L^2 -bounded and equicontinuous. By (3-9), Q_{**} inherits L^2 -boundedness from Q . As Q satisfies (1-13), we deduce that Q_{**} is also L^2 -equicontinuous. By Proposition 2.6, we may choose $\kappa_0 \geq 1$ such that

$$\sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} \leq \frac{1}{2} \quad \text{uniformly for } q \in Q_{**} \text{ and } \kappa \geq \kappa_0. \tag{4-6}$$

As shown there, this ensures that $\alpha(\kappa; q)$ and so also $\beta(\kappa; q)$ are well defined for all $q \in Q_{**}$ and $\kappa \geq \kappa_0$.

Arguing as in (2-15), we also see that (4-6) implies

$$\begin{aligned} |\beta_s(\kappa; q) - \beta_s^{[2]}(\kappa; q)| &\lesssim \int_{\kappa}^{\infty} x^{2s+2} \|\Lambda(q)\Gamma(q)\|_{\mathfrak{J}_2}^2 \frac{dx}{x} \\ &\lesssim \int_{\kappa}^{\infty} x^{2s+2} \sum_{N_1 \sim N_2 \geq N_3 \geq N_4} \|\Lambda_{N_1}\|_{\mathfrak{J}_2} \|\Lambda_{N_2}\|_{\mathfrak{J}_2} \|\Lambda_{N_3}\|_{\text{op}} \|\Lambda_{N_4}\|_{\text{op}} \frac{dx}{x} \end{aligned} \tag{4-7}$$

uniformly for $q \in Q_{**}$ and $\kappa \geq \kappa_0$. To continue from here, we decompose the full sum into the subregions S_j defined by

$$\begin{aligned} S_1 &= \{N_2 \leq \kappa\}, & S_2 &= \{\kappa < N_2 \leq x \text{ and } N_3 \leq \eta\kappa\}, & S_4 &= \{N_2 > x \text{ and } N_3 \leq \eta\kappa\}, \\ S_3 &= \{\kappa < N_2 \leq x \text{ and } N_3 > \eta\kappa\}, & S_5 &= \{N_2 > x \text{ and } N_3 > \eta\kappa\}, \end{aligned}$$

where $\eta \in (0, 1)$ is a small parameter to be chosen later. We will estimate separately each of the contributions

$$I_j(\kappa; q) := \int_{\kappa}^{\infty} \chi^{2s+2} \sum_{S_j} \|\Lambda_{N_1}\|_{\mathcal{J}_2} \|\Lambda_{N_2}\|_{\mathcal{J}_2} \|\Lambda_{N_3}\|_{\text{op}} \|\Lambda_{N_4}\|_{\text{op}} \frac{d\chi}{\chi}.$$

Applying (2-8) and (2-10) from Lemma 2.4, we have

$$\begin{aligned} I_1 &\lesssim \int_{\kappa}^{\infty} \chi^{2s+2} \sum_{N_1 \sim N_2 \leq \kappa} \frac{N_2}{\chi^3} \|q_{N_1}\|_{L^2} \|q_{N_2}\|_{L^2} \|q\|_{L^2}^2 \frac{d\chi}{\chi} \\ &\lesssim \kappa^{2s-1} \|q\|_{L^2}^2 \sum_{N_1 \sim N_2 \leq \kappa} N_2 \|q_{N_1}\|_{L^2} \|q_{N_2}\|_{L^2} \lesssim \kappa^{2s} \|q\|_{L^2}^4. \end{aligned}$$

Proceeding analogously and using (4-4), we find

$$\begin{aligned} I_2 &\lesssim \sum_{N_1 \sim N_2 > \kappa} \int_{N_2}^{\infty} \eta \kappa \chi^{2s-1} N_2^{-2s} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2}^2 \frac{d\chi}{\chi} \\ &\lesssim \sum_{N_1 \sim N_2 > \kappa} \eta \kappa N_2^{-1} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2}^2 \lesssim \eta \|q\|_{L^2}^2 \beta_s^{[2]}(\kappa; q), \\ I_3 &\lesssim \sum_{N_1 \sim N_2 > \kappa} \int_{N_2}^{\infty} \chi^{2s-1} N_2^{1-2s} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \frac{d\chi}{\chi} \\ &\lesssim \sum_{N_1 \sim N_2 > \kappa} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \lesssim \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \beta_s^{[2]}(\kappa; q), \\ I_4 &\lesssim \sum_{N_1 \sim N_2 > \kappa} \int_{\kappa}^{N_2} \eta \kappa \chi^{2s} \log\left(4 + \frac{N_2^2}{\chi^2}\right) N_2^{-1-2s} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2}^2 \frac{d\chi}{\chi} \\ &\lesssim \sum_{N_1 \sim N_2 > \kappa} \eta \kappa N_2^{-1} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2}^2 \lesssim \eta \|q\|_{L^2}^2 \beta_s^{[2]}(\kappa; q), \end{aligned}$$

and finally,

$$\begin{aligned} I_5 &\lesssim \sum_{N_1 \sim N_2 > \kappa} \int_{\kappa}^{N_2} \chi^{2s} \log\left(4 + \frac{N_2^2}{\chi^2}\right) N_2^{-2s} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \frac{d\chi}{\chi} \\ &\lesssim \sum_{N_1 \sim N_2 > \kappa} \|q_{N_1}\|_{H^s} \|q_{N_2}\|_{H^s} \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \lesssim \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2} \beta_s^{[2]}(\kappa; q). \end{aligned}$$

Collecting all our estimates, we conclude that

$$|\beta_s(\kappa; q) - \beta_s^{[2]}(\kappa; q)| \lesssim \kappa^{2s} \|q\|_{L^2}^4 + (\eta \|q\|_{L^2}^2 + \|q\|_{L^2} \|q_{>\eta\kappa}\|_{L^2}) \beta_s^{[2]}(\kappa; q)$$

uniformly on Q_{**} . As Q_{**} is L^2 -bounded and equicontinuous, we may choose η small and then $\kappa_1 \geq \kappa_0$ large to deduce that

$$\sup_{q \in Q_{**}} \beta_s^{[2]}(\kappa; q) \lesssim \sup_{q \in Q} \beta_s^{[2]}(\kappa; q) + \kappa^{2s} \sup_{q \in Q} \|q\|_{L^2}^4, \quad \text{for all } \kappa \geq \kappa_1. \quad (4-8)$$

The claim (4-5) now follows from (4-4) by choosing $\kappa = \kappa_1$.

It remains to prove that H^s -equicontinuity for Q is inherited by Q_{**} . This requires a different estimate for I_1 . Using (2-8) and (2-9), we obtain

$$I_1 \lesssim \sum_{N_4 \leq \dots \leq N_1 \leq \kappa} \sqrt{N_3 N_4} \|q_{N_1}\|_{L^2} \|q_{N_2}\|_{L^2} \|q_{N_3}\|_{L^2} \|q_{N_4}\|_{L^2} \int_{\kappa}^{\infty} \chi^{2s-1} \frac{d\chi}{\chi} \lesssim \kappa^{2s-4\sigma} \|q\|_{H^s}^4,$$

where $\sigma = \min\{s, \frac{1}{4}\}$. Now that we know (4-5), we may employ it here to deduce the following analogue of (4-8):

$$\sup_{q \in Q_{**}} \beta_s^{[2]}(\kappa; q) \lesssim \sup_{q \in Q} \beta_s^{[2]}(\kappa; q) + \kappa^{2s-4\sigma} C \left(\sup_{q \in Q} \|q\|_{L^2}^2, \sup_{q \in Q} \|q\|_{H^s}^2 \right)^4 \tag{4-9}$$

uniformly for $\kappa \geq \kappa_1$. As $4\sigma > 2s$, equicontinuity follows by sending $\kappa \rightarrow \infty$. □

5. Global well-posedness in H^s for $s \geq \frac{1}{6}$

In order to treat the line and circle simultaneously, it is convenient to introduce

$$A(\kappa; q) = \alpha(\kappa; q) \quad \text{on } \mathbb{R} \quad \text{and} \quad A(\kappa; q) = \frac{1-e^{-\kappa}}{1+e^{-\kappa}} \alpha(\kappa; q) \quad \text{on } \mathbb{T}. \tag{5-1}$$

This leads to parallel leading asymptotic expansions:

$$A(\kappa; q) = \frac{i}{2} M(q) + \frac{1}{4\kappa} H(q) + O\left(\frac{1}{\kappa^2}\right),$$

as follows from Lemmas 2.2 and 2.3. This expansion is important; it guides our choice of regularized Hamiltonian flows. We choose

$$H_{\kappa}(q) := 4\kappa \operatorname{Re} A(\kappa; q),$$

since, formally at least, $H(q) = H_{\kappa}(q) + O(\kappa^{-1})$, which suggests that the flow generated by $H_{\kappa}(q)$ approximates the (DNLS) flow as the parameter κ diverges to infinity.

The flow generated by $H_{\kappa}(q)$ with respect to the Poisson structure (1-5) is

$$\frac{d}{dt} q = \left(\frac{\delta H_{\kappa}}{\delta \bar{q}} \right)' = 2\kappa \left(\frac{\delta A(\kappa; q)}{\delta \bar{q}} + \frac{\overline{\delta A(\kappa; q)}}{\delta q} \right)', \quad \text{since } \frac{\delta \bar{A}}{\delta \bar{q}} = \frac{\delta A}{\delta q}. \tag{H_{\kappa}}$$

Our first task in this section is to prove that the H_{κ} flow is well posed on L^2 -equicontinuous sets of Schwartz initial data satisfying (1-13), provided κ is chosen sufficiently large depending on the equicontinuous family; see Proposition 5.3. Moreover, we will show that the corresponding solutions belong to \mathcal{S} for all times.

In Lemma 5.2, the H_{κ} flow will be shown to conserve $M(q)$ and $\alpha(\kappa; q)$; thus, it satisfies both the H^s -bounds and the H^s -equicontinuity guaranteed by Theorem 4.3. Together with Proposition 5.3, this immediately yields well-posedness of the H_{κ} flow on H^s for all $0 \leq s < \frac{1}{2}$ under the restriction (1-13); see Corollary 5.4.

To prove that the (DNLS) flow is well posed in H^s for $\frac{1}{6} \leq s < \frac{1}{2}$, it then suffices to prove that this is well approximated by (H_{κ}) flows as $\kappa \rightarrow \infty$. An important ingredient in our argument is the commutativity of the H_{κ} and (DNLS) flows, at least on \mathcal{S} . This follows from Lemma 5.2 and the well-posedness of these

flows on \mathcal{S} by mimicking the arguments in [Arnold 1989, §39]. In view of this commutativity, proving convergence of the (H_κ) flows to the (DNLS) flow amounts to showing that the flow generated by the difference of the Hamiltonians $H(q) - H_\kappa(q)$ converges to the identity as $\kappa \rightarrow \infty$. This final stage of the proof will be carried out in Theorem 5.5.

In order to make sense of (H_κ) , we must prove that $\alpha(\kappa; q)$ is in fact differentiable. To solve (H_κ) locally in time, we further need to show that this functional derivative is itself a Lipschitz function of q . These goals require us to define $\alpha(\kappa; q)$ on open sets in L^2 , rather than merely equicontinuous sets. The next result addresses these issues.

Here and below we write Q_ε to denote the ε neighborhood of Q in the L^2 -metric.

Lemma 5.1. *Let Q be a bounded and equicontinuous subset of L^2 . Then there exist $\varepsilon > 0$ and $\kappa_0 \geq 1$ such that for all $\kappa \geq \kappa_0$, we have that $\alpha(\kappa; q)$ is a real-analytic function of $q \in Q_\varepsilon$. Moreover, we have the bounds*

$$\left\| \frac{\delta\alpha(\kappa; q)}{\delta q} \right\|_{H^1} + \left\| \frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} \right\|_{H^1} \lesssim \kappa \|q\|_{L^2}, \tag{5-2}$$

$$\left\| \frac{\delta\alpha(\kappa; q)}{\delta q} - \frac{\delta\alpha(\kappa; \tilde{q})}{\delta q} \right\|_{H^1} + \left\| \frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} - \frac{\delta\alpha(\kappa; \tilde{q})}{\delta \bar{q}} \right\|_{H^1} \lesssim \kappa \|q - \tilde{q}\|_{L^2}, \tag{5-3}$$

where the implicit constants depend only on Q . Additionally, for every $\kappa \geq \kappa_0$ and $q \in Q_\varepsilon$, there exists $\gamma(\kappa; q) \in H^1$ such that

$$\left(\frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} \right)' = 2\kappa \frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} - i\kappa q[\gamma(\kappa; q) + 1], \tag{5-4}$$

$$\left(\frac{\delta\alpha(\kappa; q)}{\delta q} \right)' = -2\kappa \frac{\delta\alpha(\kappa; q)}{\delta q} + i\kappa \bar{q}[\gamma(\kappa; q) + 1], \tag{5-5}$$

$$\gamma(\kappa; q)' = 2\bar{q} \frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} - 2q \frac{\delta\alpha(\kappa; q)}{\delta q}. \tag{5-6}$$

Lastly, for each integer $m \geq 0$ we have

$$\left\| \left(\frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} \right)' \right\|_{H^m} \lesssim_m \kappa \|q\|_{H^m}, \tag{5-7}$$

$$\left\| \langle x \rangle^{2m} \left(\frac{\delta\alpha(\kappa; q)}{\delta \bar{q}} \right)' \right\|_{L^2} \lesssim_m \kappa \|\langle x \rangle^{2m} q\|_{L^2}, \tag{5-8}$$

uniformly for $q \in Q_\varepsilon$ and $\kappa \geq \kappa_0$.

Proof. Proposition 2.6 shows that given $\delta \in (0, 1]$, there exists $\kappa_0 \geq 1$ such that

$$\sup_{q \in Q} \sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} \leq \frac{\delta}{4} \quad \text{uniformly for } \kappa \geq \kappa_0.$$

As $\Lambda(q)$ is linear in q , Lemma 2.1 allows us to deduce

$$\sup_{q \in Q_\varepsilon} \sqrt{\kappa} \|\Lambda(q)\|_{\text{op}} \leq \frac{\delta}{2} \quad \text{uniformly for } \kappa \geq \kappa_0, \tag{5-9}$$

provided ε is chosen sufficiently small (depending on δ).

Now we must explain how to choose δ . In view of (2-15), $\delta \leq 1$ guarantees that the series (1-9) converges on Q_ε . We place an additional requirement to aid in the proofs of (5-2) and (5-3). From Lemma 2.5 we find that

$$\|(\kappa + \partial)^{-1/4} q(\kappa - \partial)^{-1/4}\|_{\text{op}} \cdot \|(\kappa - \partial)^{-3/4} q(\kappa + \partial)^{-3/4}\|_{\text{op}} \lesssim \|q\|_{L^2} \cdot \kappa^{-1/2} \|\Lambda(q)\|_{\text{op}}.$$

Thus, we may choose δ even smaller if necessary to ensure also that

$$\kappa \|(\kappa + \partial)^{-1/4} q(\kappa - \partial)^{-1/4}\|_{\text{op}} \cdot \|(\kappa - \partial)^{-3/4} q(\kappa + \partial)^{-3/4}\|_{\text{op}} \leq \frac{1}{2} \tag{5-10}$$

uniformly for $q \in Q_\varepsilon$ and $\kappa \geq \kappa_0$.

Turning now to (5-2), we argue by duality. For $f \in H^{-1}$, we have

$$\left\langle f, \frac{\delta \alpha(\kappa; q)}{\delta \bar{q}} \right\rangle = \sum_{\ell \geq 0} (i\kappa)^{\ell+1} \text{tr}\{[(\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{q}]^\ell (\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{f}\}.$$

The $\ell = 0$ term is readily computed exactly via Lemma 2.2. For example,

$$i\kappa \text{tr}\{(\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{f}\} = i\kappa \left\langle \frac{1}{2\kappa + \partial} f, q \right\rangle \text{ in the line case.}$$

In either geometry, this is easily seen to satisfy the desired bound.

For $\ell \geq 1$, we employ (5-10) and Lemma 2.5 to estimate

$$\begin{aligned} & |(i\kappa)^{\ell+1} \text{tr}\{[(\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{q}]^\ell (\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{f}\}| \\ & \lesssim \kappa^{\ell+1} \|(\kappa + \partial)^{-1} \bar{f}(\kappa - \partial)^{-1}\|_{\mathfrak{J}_2} \|q(\kappa + \partial)^{-3/4}\|_{\mathfrak{J}_2}^2 \|(\kappa + \partial)^{-1/4} q(\kappa - \partial)^{-1/4}\|_{\text{op}}^\ell \\ & \hspace{20em} \times \|(\kappa - \partial)^{-3/4} q(\kappa + \partial)^{-3/4}\|_{\text{op}}^{\ell-1} \\ & \lesssim 2^{-\ell} \kappa \|f\|_{H^{-1}} \|q\|_{L^2}^3, \end{aligned}$$

with an implicit constant independent of ℓ . This proves that the estimate (5-2) holds for the \bar{q} derivative; the bound on the q derivative follows in a parallel fashion.

The proof of (5-3) proceeds analogously, noting that one can always exhibit the difference $q - \tilde{q}$ in place of a q .

We define $\gamma(\kappa; q)$ via the associated linear functional

$$\begin{aligned} \langle f, \gamma(\kappa; q) \rangle = & \sum_{\ell \geq 1} (i\kappa)^\ell \text{tr}\{[(\kappa - \partial)^{-1} q(\kappa + \partial)^{-1} \bar{q}]^\ell (\kappa - \partial)^{-1} \bar{f}\} \\ & + \sum_{\ell \geq 1} (i\kappa)^\ell \text{tr}\{[(\kappa + \partial)^{-1} \bar{q}(\kappa - \partial)^{-1} q]^\ell (\kappa + \partial)^{-1} \bar{f}\}, \tag{5-11} \end{aligned}$$

and will prove $\gamma \in H^1$ by showing that this functional is bounded for $f \in H^{-1}$.

Regarding the $\ell = 1$ terms, Lemma 2.5 and direct computation show that

$$|\kappa \text{tr}\{(\kappa \mp \partial)^{-1} q(\kappa \pm \partial)^{-1} \bar{q}(\kappa \mp \partial)^{-1} \bar{f}\}| \lesssim \sqrt{\kappa} \|q(\kappa \pm \partial)^{-1} \bar{q}\|_{\mathfrak{J}_2} \|f\|_{H^{-1}} \lesssim \sqrt{\kappa} \|q\|_{L^2}^2 \|f\|_{H^{-1}}.$$

For $\ell \geq 2$, we employ Lemma 2.5 and (5-10) as follows:

$$\begin{aligned} & |\kappa^\ell \operatorname{tr}\{[(\kappa \mp \partial)^{-1} q(\kappa \pm \partial)^{-1} \bar{q}]^\ell (\kappa \mp \partial)^{-1} f\}| \\ & \lesssim \kappa^{\ell-1/2} \|f\|_{H^{-1}} \|q(\kappa + \partial)^{-3/4}\|_{\mathfrak{S}_2}^2 \|(\kappa - \partial)^{-1/2}\|_{\text{op}} \|(\kappa + \partial)^{-1/4} q(\kappa - \partial)^{-1/4}\|_{\text{op}}^\ell \\ & \hspace{15em} \times \|(\kappa - \partial)^{-3/4} q(\kappa + \partial)^{-3/4}\|_{\text{op}}^{\ell-2} \\ & \lesssim 2^{-\ell} \sqrt{\kappa} \|q\|_{L^2}^4 \|f\|_{H^{-1}}, \end{aligned}$$

where the implicit constant is independent of ℓ . Thus $\gamma \in H^1$ and

$$\|\gamma(\kappa; q)\|_{H^1} \lesssim \sqrt{\kappa} \|q\|_{L^2}^2.$$

The proofs of (5-4) and (5-5) follow parallel arguments. In the former case, we pair $\delta\alpha(\kappa; q)/\delta\bar{q}$ with f' , which we then rewrite as a trace. The result then follows by noting the operator identity $f' = -(\kappa - \partial)f - f(\kappa + \partial) + 2\kappa f$ and simplifying.

The proof of (5-6) follows the same style: one pairs $\gamma(\kappa; q)$ with f' and employs the operator identity $f' = [\kappa + \partial, f] = -[\kappa - \partial, f]$.

The proof of (5-7) mimics closely that of (5-2), once one understands how to move the derivatives from the test function f to copies of q . Introducing the notation $f_h(x) = f(x - h)$, we observe that by the translation invariance of the trace,

$$\begin{aligned} \left\langle f^{(m)}, \left(\frac{\delta\alpha(\kappa; q)}{\delta\bar{q}} \right)' \right\rangle &= -\frac{\partial^m}{\partial h^m} \Big|_{h=0} \left\langle f'_h, \frac{\delta\alpha(\kappa; q)}{\delta\bar{q}} \right\rangle = -\frac{\partial^m}{\partial h^m} \Big|_{h=0} \left\langle f', \frac{\delta}{\delta\bar{q}} \alpha(\kappa; q_{-h}) \right\rangle \\ &= -\frac{\partial^m}{\partial h^m} \Big|_{h=0} \sum_{\ell \geq 0} (i\kappa)^{\ell+1} \operatorname{tr}\{[(\kappa - \partial)^{-1} q_{-h}(\kappa + \partial)^{-1} \bar{q}_{-h}]^\ell (\kappa - \partial)^{-1} q_{-h}(\kappa + \partial)^{-1} \bar{f}'\}. \end{aligned}$$

Next, we apply the estimates used to prove (5-2) together with the elementary inequality

$$\|q^{(n)}\|_{L^2} \lesssim \|q\|_{L^2}^{1-n/m} \|q\|_{H^m}^{n/m} \quad \text{for all } 0 < n \leq m.$$

This yields the estimate (5-7). Note that summability in ℓ is guaranteed by (5-10), just as before.

Lastly, we turn to (5-8). The argument is very similar; the key ingredient is to move the polynomial weight $\langle x \rangle^{2m}$ from the test function f to a copy of q . This is achieved via the identity

$$q(\kappa + \partial)^{-1} P \bar{f} = \sum_{n \geq 0} (-1)^n [P^{(n)} q] (\kappa + \partial)^{-n-1} \bar{f},$$

valid for any polynomial $P(x)$, which follows easily by induction using

$$[(\kappa + \partial)^{-1}, P(x)] = -(\kappa + \partial)^{-1} P'(x) (\kappa + \partial)^{-1}. \quad \square$$

Lemma 5.2. *Let $Q \subseteq \mathcal{S}$ be L^2 -bounded and equicontinuous, and let ε and κ_0 be as in Lemma 5.1. Then for all $\kappa, \varkappa \geq \kappa_0$,*

$$\{H, \alpha(\kappa)\} = 0, \quad \{M, \alpha(\kappa)\} = 0, \quad \text{and} \quad \{\alpha(\varkappa), \alpha(\kappa)\} = 0$$

on Q_ε . Consequently, $A(\kappa)$, $A(\varkappa)$, M , H , and H_κ all Poisson commute on Q_ε .

Proof. As discussed in the introduction, the commutativity of $\alpha(\kappa)$ with the Hamiltonian H was proved in [Klaus and Schippa 2022; Tang and Xu 2021] whenever the series defining $\alpha(\kappa)$ can be guaranteed to converge. Such convergence is guaranteed by Lemma 5.1.

Recalling (1-5) and employing (5-4), (5-5), and (5-6), we find

$$\{M, \alpha(\kappa)\} = -2\kappa \int \gamma' dx = 0.$$

Notice that (5-6) guarantees $\gamma' \in L^1$.

If $\kappa = \varkappa$ the third equality is clear. When $\kappa \neq \varkappa$, we may proceed to compute the Poisson bracket by applying (5-4) and (5-5) directly to the derivatives of $\alpha(\kappa)$ or by employing integration by parts and then the corresponding formulae for the partial derivatives of $\alpha(\varkappa)$. Comparing the two approaches yields

$$\varkappa\{\alpha(\kappa), \alpha(\varkappa)\} = \kappa\{\alpha(\kappa), \alpha(\varkappa)\}, \quad \text{and so } \{\alpha(\varkappa), \alpha(\kappa)\} = 0. \quad \square$$

Proposition 5.3. *For each L^2 -equicontinuous set $Q \subseteq S$ satisfying (1-13), there exists $\kappa_0 \geq 1$ sufficiently large such that for all $\kappa \geq \kappa_0$, the (H_κ) flow is globally well posed for initial data in Q . Moreover, the solutions remain in S for all time. Lastly, the set*

$$Q_* := \{e^{tJ\nabla H_\kappa} q : q \in Q, t \in \mathbb{R}, \text{ and } \kappa \geq \kappa_0\}$$

is bounded and equicontinuous in L^2 .

Proof. Recall the set Q_{**} introduced in (1-11). By (3-9), the hypothesis (1-13), and the definition of M_* , this set is bounded and equicontinuous in L^2 . We fix $\varepsilon > 0$ and $\kappa_0 \geq 1$ as the values obtained by applying Lemma 5.1 to the set Q_{**} .

Next we construct a local solution for initial data $q(0) \in Q_{**}$. For $\kappa \geq \kappa_0$, Lemma 5.1 ensures that one can run the usual contraction mapping argument for the integral equation

$$q(t) = q(0) + \int_0^t 2\kappa \left(\frac{\delta A(\kappa; q(s))}{\delta \bar{q}} + \overline{\frac{\delta A(\kappa; q(s))}{\delta q}} \right)' ds$$

to find a unique solution $q \in C([0, T]; L^2)$, provided T is chosen sufficiently small. In fact, T is chosen so small that $q(t)$ and indeed all Picard iterates remain in the ε -neighborhood of Q_{**} .

Combining the estimates (5-7) and (5-8) with the Gronwall inequality shows that $q(t) \in S$ for all $t \in [0, T]$. This in turn allows us to apply Lemma 5.2 to conclude that $\alpha(\varkappa; q(t))$ and hence $a(\varkappa; q(t))$ are conserved. Taken together, these observations guarantee that $q([0, T]) \subseteq Q_{**}$ and so the local solutions may be concatenated to yield a global solution lying wholly within Q_{**} . Finally, as Q_* is a subset of Q_{**} , it is L^2 -bounded and equicontinuous. □

Combining Proposition 5.3 with Theorem 4.3 immediately yields well-posedness of the (H_κ) flow in the following sense:

Corollary 5.4. *Fix $0 < s < \frac{1}{2}$ and let $Q \subseteq S$ be H^s -bounded and satisfy (1-13). Then there exists $\kappa_0 \geq 1$ such that for all $\kappa \geq \kappa_0$ the (H_κ) flow is globally well posed for initial data in Q . Moreover,*

$$Q_* := \{e^{tJ\nabla H_\kappa} q : q \in Q, t \in \mathbb{R}, \text{ and } \kappa \geq \kappa_0\} \subseteq S \quad \text{is } H^s\text{-bounded.}$$

If Q is H^s -equicontinuous, then so is Q_ .*

In order to complete the proof of Theorem 1.5, we must prove that H^s -Cauchy sequences of initial data $q_n(0) \in \mathcal{S}$ satisfying (1-13) lead to Cauchy sequences of solutions to (DNLS). As mentioned above, this will be accomplished by showing that the flow

$$\frac{d}{dt}q = \left[iq' - |q|^2q - 2\kappa \left(\frac{\delta A(\kappa; q)}{\delta \bar{q}} + \frac{\overline{\delta A(\kappa; q)}}{\delta q} \right) \right]', \tag{H_\kappa^{\text{diff}}}$$

generated by $H(q) - H_\kappa(q)$, converges to the identity as $\kappa \rightarrow \infty$. Due to commutativity of the flows, \mathcal{S} -valued solutions to (H_κ^{diff}) can be built via

$$e^{tJ\nabla H_\kappa^{\text{diff}}}q = e^{tJ\nabla H}e^{-tJ\nabla H_\kappa}q$$

using Corollaries 4.2 and 5.4. In view of Lemma 5.2, these solutions conserve M and $\alpha(x)$.

The proof of our final theorem makes a fitting end for this paper by highlighting the power of equicontinuity. It is also here that we will finally see the origin of the restriction $s \geq \frac{1}{6}$. It is needed to make sense of the nonlinearity in (H_κ^{diff}) pointwise in time.

Theorem 5.5. *Fix $\frac{1}{6} \leq s < \frac{1}{2}$ and $T > 0$. Given a sequence $q_n(0) \in \mathcal{S}$ of initial data that converges in H^s and satisfies (1-13), let $q_n(t)$ denote the corresponding solutions to (DNLS). Then $q_n(t)$ converges in H^s , uniformly for $|t| \leq T$.*

Proof. By hypothesis, the set $Q = \{q_n(0) : n \in \mathbb{N}\}$ is bounded and equicontinuous in the H^s -metric. Let $\kappa_0 \geq 1$ be as given by Corollary 5.4. Then for $\kappa \geq \kappa_0$, the (H_κ) flow is well posed for initial data in Q , and the set

$$Q_* := \{e^{tJ\nabla H_\kappa}q_n(0) : n \in \mathbb{N}, t \in \mathbb{R}, \text{ and } \kappa \geq \kappa_0\} \subseteq \mathcal{S}$$

is bounded and equicontinuous in H^s .

The commutativity of the (H_κ) and the (DNLS) flows allows us to rewrite our sequence of solutions as

$$q_n(t) = e^{tJ\nabla H_\kappa^{\text{diff}}}e^{tJ\nabla H_\kappa}q_n(0).$$

Moreover, by Theorem 4.3, the set

$$\{e^{tJ\nabla H_\kappa^{\text{diff}}}q : q \in Q_*, t \in \mathbb{R}, \text{ and } \kappa \geq \kappa_0\} \subseteq Q_{**}$$

is bounded and equicontinuous in H^s .

We will show that $q_n(t)$ forms a Cauchy sequence in H^s , uniformly for $|t| \leq T$. By the definition of Q_* , we estimate

$$\sup_{|t| \leq T} \|q_n(t) - q_m(t)\|_{H^s} \leq 2 \sup_{q \in Q_*} \sup_{|t| \leq T} \|e^{tJ\nabla H_\kappa^{\text{diff}}}q - q\|_{H^s} + \sup_{|t| \leq T} \|e^{tJ\nabla H_\kappa}q_n(0) - e^{tJ\nabla H_\kappa}q_m(0)\|_{H^s}$$

for all $\kappa \geq \kappa_0$. For any such fixed κ , the well-posedness of the (H_κ) flow ensures that the last term of the right-hand side converges to 0 as $n, m \rightarrow \infty$. Thus, it suffices to prove that the difference flow converges to the identity uniformly on Q_* :

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q_*} \sup_{|t| \leq T} \|e^{tJ\nabla H_\kappa^{\text{diff}}}q - q\|_{H^s} = 0.$$

In fact, as Q_{**} is H^s -equicontinuous, it suffices to show that

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q_{**}} \sup_{|t| \leq T} \|e^{tJ\nabla H_{\kappa}^{\text{diff}}} q - q\|_{H^{-4}} = 0. \quad (5-12)$$

By the fundamental theorem of calculus and $(H_{\kappa}^{\text{diff}})$, proving (5-12) reduces to showing that

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q_{**}} \|F\|_{H^{-3}} = 0, \quad \text{where } F := iq' - |q|^2 q - 2\kappa \left(\frac{\delta A(\kappa; q)}{\delta \bar{q}} + \overline{\frac{\delta A(\kappa; q)}{\delta q}} \right). \quad (5-13)$$

A straightforward computation shows that $F^{[1]}$, the linear (in q) term in F , is given by $-i\partial^3/(4\kappa^2 - \partial^2)q$. This clearly converges to zero in H^{-3} as $\kappa \rightarrow \infty$, uniformly on Q_{**} , or indeed, on any L^2 -bounded set.

We turn now to the contribution of $F^{[3]}$, the term in F that is cubic in q . Employing Lemma 2.3, we find the cubic terms

$$\begin{aligned} \left(\frac{\delta A(\kappa; q)}{\delta \bar{q}} \right)^{[3]} &= -\frac{\kappa^2}{2\kappa - \partial} \left\{ \left(\frac{1}{2\kappa + \partial} \bar{q} \right) (4\kappa - \partial) \left(\frac{1}{2\kappa - \partial} q \right)^2 \right\} = -\frac{2\kappa^2}{2\kappa - \partial} \left\{ q \left(\frac{1}{2\kappa - \partial} q \right) \left(\frac{1}{2\kappa + \partial} \bar{q} \right) \right\}, \\ \left(\frac{\delta A(\kappa; q)}{\delta q} \right)^{[3]} &= -\frac{\kappa^2}{2\kappa + \partial} \left\{ \left(\frac{1}{2\kappa - \partial} \bar{q} \right) (4\kappa + \partial) \left(\frac{1}{2\kappa + \partial} q \right)^2 \right\} = -\frac{2\kappa^2}{2\kappa + \partial} \left\{ q \left(\frac{1}{2\kappa + \partial} q \right) \left(\frac{1}{2\kappa - \partial} \bar{q} \right) \right\}. \end{aligned}$$

This allows us to compute the full cubic term as follows:

$$\begin{aligned} F^{[3]} &= 2\kappa^2 \frac{\partial}{2\kappa - \partial} \left[q \left(\frac{1}{2\kappa - \partial} q \right) \left(\frac{1}{2\kappa + \partial} \bar{q} \right) \right] - 2\kappa^2 \frac{\partial}{2\kappa + \partial} \left[q \left(\frac{1}{2\kappa + \partial} q \right) \left(\frac{1}{2\kappa - \partial} \bar{q} \right) \right] \\ &\quad + 2\kappa^2 q \left(\frac{1}{2\kappa - \partial} q \right) \left(\frac{1}{2\kappa + \partial} \bar{q} \right) + 2\kappa^2 q \left(\frac{1}{2\kappa + \partial} q \right) \left(\frac{1}{2\kappa - \partial} \bar{q} \right) - q^2 \bar{q} \\ &= \frac{2\partial}{2\kappa - \partial} \left[q \left(\frac{\kappa}{2\kappa - \partial} q \right) \left(\frac{\kappa}{2\kappa + \partial} \bar{q} \right) \right] - \frac{2\partial}{2\kappa + \partial} \left[q \left(\frac{\kappa}{2\kappa + \partial} q \right) \left(\frac{\kappa}{2\kappa - \partial} \bar{q} \right) \right] + q^2 \left(\frac{\partial^2}{4\kappa^2 - \partial^2} \bar{q} \right) \\ &\quad + q \bar{q} \left(\frac{\partial^2}{4\kappa^2 - \partial^2} q \right) - \frac{1}{2} q \left(\frac{\partial}{2\kappa - \partial} q \right) \left(\frac{\partial}{2\kappa + \partial} \bar{q} \right) - \frac{1}{2} q \left(\frac{\partial}{2\kappa + \partial} q \right) \left(\frac{\partial}{2\kappa - \partial} \bar{q} \right). \end{aligned}$$

To estimate its contribution, we pair with $f \in H^3$ and apply Hölder's inequality. Boundedness is easily deduced from

$$\|f\|_{L^\infty} + \kappa \left\| \frac{\partial}{2\kappa \pm \partial} f \right\|_{L^\infty} \lesssim \|f\|_{H^3}, \quad (5-14)$$

$$\|q\|_{L^3} + \left\| \frac{\partial}{2\kappa \pm \partial} q \right\|_{L^3} + \left\| \frac{\partial^2}{4\kappa^2 - \partial^2} q \right\|_{L^3} \lesssim \|q\|_{H^s}. \quad (5-15)$$

Evidently (5-15) requires $s \geq \frac{1}{6}$. The gain of a power of κ in (5-14) guarantees that the contribution of the first two terms in $F^{[3]}$ decays to zero as $\kappa \rightarrow \infty$. For the remaining terms, we use H^s -equicontinuity to obtain decay: as $s \geq \frac{1}{6}$, we have that

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q_{**}} \left\| \frac{\partial}{2\kappa \pm \partial} q \right\|_{L^3} \lesssim \lim_{\kappa \rightarrow \infty} \sup_{q \in Q_{**}} \left\| \frac{\partial}{2\kappa \pm \partial} q \right\|_{H^s} = 0.$$

Finally, we turn our attention to the remaining terms (quintic and higher) in the series expansion of F . By Lemma 2.1, (1-9), and the embedding $H^3 \hookrightarrow L^\infty$,

$$\left| \int f F^{[\geq 5]} dx \right| \lesssim \sum_{\ell \geq 2} \kappa^{\ell+2} \|\Lambda(q)\|_{\dot{H}^2}^2 \|\Lambda(q)\|_{\text{op}}^{2\ell-1} \|\Lambda(f)\|_{\text{op}} \lesssim \|q\|_{L^2}^2 \|f\|_{H^3} \sum_{\ell \geq 2} \kappa^\ell \|\Lambda(q)\|_{\text{op}}^{2\ell-1}.$$

The convergence we require does not follow from Proposition 2.6; we would lose by a factor of $\sqrt{\kappa}$. However arguing in the same fashion, we find

$$\begin{aligned} \|\Lambda(q)\|_{\text{op}} &\lesssim \|\Lambda(q_{\leq \eta\kappa})\|_{\text{op}} + \|\Lambda(q_{> \eta\kappa})\|_{\text{op}} \lesssim \kappa^{-1} \|q_{\leq \eta\kappa}\|_{L^\infty} + \kappa^{-1/2} \|q_{> \eta\kappa}\|_{L^2} \\ &\lesssim \kappa^{-1/2-s} (\eta^{1/2-s} \|q\|_{H^s} + \eta^{-s} \|q_{> \eta\kappa}\|_{H^s}), \end{aligned}$$

for any $\eta > 0$. When $s > \frac{1}{6}$, we may simply take $\eta = 1$ to deduce that

$$\lim_{\kappa \rightarrow \infty} \sup_{q \in Q_{**}} \|F^{[5]}\|_{H^{-3}} = 0.$$

For the endpoint case $s = \frac{1}{6}$, this follows from the H^s -equicontinuity of Q_{**} by choosing η small and then κ large. This completes the proof of the theorem. \square

References

- [Arnold 1989] V. I. Arnold, *Mathematical methods of classical mechanics*, 2nd ed., Grad. Texts in Math. **60**, Springer, 1989. MR Zbl
- [Bahouri and Perelman 2022] H. Bahouri and G. Perelman, “Global well-posedness for the derivative nonlinear Schrödinger equation”, *Invent. Math.* **229**:2 (2022), 639–688. MR Zbl
- [Biagioni and Linares 2001] H. A. Biagioni and F. Linares, “Ill-posedness for the derivative Schrödinger and generalized Benjamin–Ono equations”, *Trans. Amer. Math. Soc.* **353**:9 (2001), 3649–3659. MR Zbl
- [Bringmann et al. 2021] B. Bringmann, R. Killip, and M. Vişan, “Global well-posedness for the fifth-order KdV equation in $H^{-1}(\mathbb{R})$ ”, *Ann. PDE* **7**:2 (2021), art. id. 21. MR Zbl
- [Colliander et al. 2001] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, “Global well-posedness for Schrödinger equations with derivative”, *SIAM J. Math. Anal.* **33**:3 (2001), 649–669. MR Zbl
- [Colliander et al. 2002] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, “A refined global well-posedness result for Schrödinger equations with derivative”, *SIAM J. Math. Anal.* **34**:1 (2002), 64–86. MR Zbl
- [Cwikel 1977] M. Cwikel, “Weak type estimates for singular values and the number of bound states of Schrödinger operators”, *Ann. of Math. (2)* **106**:1 (1977), 93–100. MR Zbl
- [Deng et al. 2021] Y. Deng, A. R. Nahmod, and H. Yue, “Optimal local well-posedness for the periodic derivative nonlinear Schrödinger equation”, *Comm. Math. Phys.* **384**:2 (2021), 1061–1107. MR Zbl
- [Fujiwara et al. 2020] K. Fujiwara, V. Georgiev, and T. Ozawa, “Self-similar solutions to the derivative nonlinear Schrödinger equation”, *J. Differential Equations* **268**:12 (2020), 7940–7961. MR Zbl
- [Fukaya et al. 2017] N. Fukaya, M. Hayashi, and T. Inui, “A sufficient condition for global existence of solutions to a generalized derivative nonlinear Schrödinger equation”, *Anal. PDE* **10**:5 (2017), 1149–1167. MR Zbl
- [Grünrock 2005] A. Grünrock, “Bi- and trilinear Schrödinger estimates in one space dimension with applications to cubic NLS and DNLS”, *Int. Math. Res. Not.* **2005**:41 (2005), 2525–2558. MR Zbl
- [Grünrock and Herr 2008] A. Grünrock and S. Herr, “Low regularity local well-posedness of the derivative nonlinear Schrödinger equation with periodic initial data”, *SIAM J. Math. Anal.* **39**:6 (2008), 1890–1920. MR Zbl

- [Guo and Wu 2017] Z. Guo and Y. Wu, “Global well-posedness for the derivative nonlinear Schrödinger equation in $H^{1/2}(\mathbb{R})$ ”, *Discrete Contin. Dyn. Syst.* **37**:1 (2017), 257–264. MR Zbl
- [Harrop-Griffiths et al. 2020] B. Harrop-Griffiths, R. Killip, and M. Vişan, “Sharp well-posedness for the cubic NLS and mKdV in $H^s(\mathbb{R})$ ”, preprint, 2020. arXiv 2003.05011
- [Hayashi and Ozawa 1992] N. Hayashi and T. Ozawa, “On the derivative nonlinear Schrödinger equation”, *Phys. D* **55**:1-2 (1992), 14–36. MR Zbl
- [Herr 2006] S. Herr, “On the Cauchy problem for the derivative nonlinear Schrödinger equation with periodic boundary condition”, *Int. Math. Res. Not.* **2006** (2006), art. id. 96763. MR Zbl
- [Ichikawa and Watanabe 1977] Y. H. Ichikawa and S. Watanabe, “Solitons, envelope solitons in collisionless plasmas”, *J. Phys. Colloques* **38**:C6 (1977), 15–26.
- [Isom et al. 2020] B. Isom, D. Mantzavinos, and A. Stefanov, “Growth bound and nonlinear smoothing for the periodic derivative nonlinear Schrödinger equation”, preprint, 2020. arXiv 2012.09933
- [Jenkins et al. 2018a] R. Jenkins, J. Liu, P. Perry, and C. Sulem, “Soliton resolution for the derivative nonlinear Schrödinger equation”, *Comm. Math. Phys.* **363**:3 (2018), 1003–1049. MR Zbl
- [Jenkins et al. 2018b] R. Jenkins, J. Liu, P. A. Perry, and C. Sulem, “Global well-posedness for the derivative non-linear Schrödinger equation”, *Comm. Partial Differential Equations* **43**:8 (2018), 1151–1195. MR Zbl
- [Jenkins et al. 2020a] R. Jenkins, J. Liu, P. Perry, and C. Sulem, “The derivative nonlinear Schrödinger equation: global well-posedness and soliton resolution”, *Quart. Appl. Math.* **78**:1 (2020), 33–73. MR Zbl
- [Jenkins et al. 2020b] R. Jenkins, J. Liu, P. Perry, and C. Sulem, “Global existence for the derivative nonlinear Schrödinger equation with arbitrary spectral singularities”, *Anal. PDE* **13**:5 (2020), 1539–1578. MR Zbl
- [Kaup and Newell 1978] D. J. Kaup and A. C. Newell, “An exact solution for a derivative nonlinear Schrödinger equation”, *J. Math. Phys.* **19**:4 (1978), 798–801. MR Zbl
- [Killip and Vişan 2019] R. Killip and M. Vişan, “KdV is well-posed in H^{-1} ”, *Ann. of Math. (2)* **190**:1 (2019), 249–305. MR Zbl
- [Killip et al. 2018] R. Killip, M. Vişan, and X. Zhang, “Low regularity conservation laws for integrable PDE”, *Geom. Funct. Anal.* **28**:4 (2018), 1062–1090. MR Zbl
- [Killip et al. 2020] R. Killip, J. Murphy, and M. Vişan, “Invariance of white noise for KdV on the line”, *Invent. Math.* **222**:1 (2020), 203–282. MR Zbl
- [Kitaev 1985] A. V. Kitaev, “Self-similar solutions of a modified nonlinear Schrödinger equation”, *Teoret. Mat. Fiz.* **64**:3 (1985), 347–369. In Russian; translated in *Theoret. and Math. Phys.* **64**:3 (1986), 878–894. MR Zbl
- [Klaus and Schippa 2022] F. Klaus and R. Schippa, “A priori estimates for the derivative nonlinear Schrödinger equation”, *Funkcial. Ekvac.* **65**:3 (2022), 329–346. MR Zbl
- [Liu 2017] J. Liu, *Global well-posedness for the derivative nonlinear Schrödinger equation through inverse scattering*, Ph.D. thesis, University of Kentucky, 2017, available at https://uknowledge.uky.edu/math_etds/50/.
- [Liu et al. 2016] J. Liu, P. A. Perry, and C. Sulem, “Global existence for the derivative nonlinear Schrödinger equation by the method of inverse scattering”, *Comm. Partial Differential Equations* **41**:11 (2016), 1692–1760. MR Zbl
- [Liu et al. 2018] J. Liu, P. A. Perry, and C. Sulem, “Long-time behavior of solutions to the derivative nonlinear Schrödinger equation for soliton-free initial data”, *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **35**:1 (2018), 217–265. MR Zbl
- [Miao et al. 2011] C. Miao, Y. Wu, and G. Xu, “Global well-posedness for Schrödinger equation with derivative in $H^{1/2}(\mathbb{R})$ ”, *J. Differential Equations* **251**:8 (2011), 2164–2195. MR Zbl
- [Mio et al. 1976] K. Mio, T. Ogino, K. Minami, and S. Takeda, “Modulational instability and envelope-solitons for nonlinear Alfvén waves propagating along the magnetic field in plasmas”, *J. Phys. Soc. Japan* **41**:2 (1976), 667–673.
- [Mjølhus 1976] E. Mjølhus, “On the modulational instability of hydromagnetic waves parallel to the magnetic field”, *J. Plasma Phys.* **16**:3 (1976), 321–334.
- [Mosincat 2017] R. Mosincat, “Global well-posedness of the derivative nonlinear Schrödinger equation with periodic boundary condition in $H^{1/2}$ ”, *J. Differential Equations* **263**:8 (2017), 4658–4722. MR Zbl

- [Mosincat and Oh 2015] R. Mosincat and T. Oh, “A remark on global well-posedness of the derivative nonlinear Schrödinger equation on the circle”, *C. R. Math. Acad. Sci. Paris* **353**:9 (2015), 837–841. MR Zbl
- [Nahmod et al. 2012] A. R. Nahmod, T. Oh, L. Rey-Bellet, and G. Staffilani, “Invariant weighted Wiener measures and almost sure global well-posedness for the periodic derivative NLS”, *J. Eur. Math. Soc.* **14**:4 (2012), 1275–1330. MR Zbl
- [Ntekoume 2022] M. Ntekoume, “Symplectic nonsqueezing for the KdV flow on the line”, *Pure Appl. Anal.* **4**:3 (2022), 401–448. MR Zbl
- [Pelinovsky and Shimabukuro 2018] D. E. Pelinovsky and Y. Shimabukuro, “Existence of global solutions to the derivative NLS equation with the inverse scattering transform method”, *Int. Math. Res. Not.* **2018**:18 (2018), 5663–5728. MR Zbl
- [Pelinovsky et al. 2017] D. E. Pelinovsky, A. Saalman, and Y. Shimabukuro, “The derivative NLS equation: global existence with solitons”, *Dyn. Partial Differ. Equ.* **14**:3 (2017), 271–294. MR Zbl
- [Saalman 2017] A. Saalman, “Global existence for the derivative NLS equation in the presence of solitons”, preprint, 2017. Zbl arXiv 1704.00071
- [Simon 2005] B. Simon, *Trace ideals and their applications*, 2nd ed., Math. Surv. Monogr. **120**, Amer. Math. Soc., Providence, RI, 2005. MR Zbl
- [Takaoka 1999] H. Takaoka, “Well-posedness for the one-dimensional nonlinear Schrödinger equation with the derivative nonlinearity”, *Adv. Differential Equations* **4**:4 (1999), 561–580. MR Zbl
- [Takaoka 2001] H. Takaoka, “Global well-posedness for Schrödinger equations with derivative in a nonlinear term and data in low-order Sobolev spaces”, *Electron. J. Differential Equations* **2001** (2001), art. id. 42. MR Zbl
- [Tang and Xu 2021] X. Tang and G. Xu, “Microscopic conservation laws for the derivative nonlinear Schrödinger equation”, *Lett. Math. Phys.* **111**:6 (2021), art. id. 138. MR Zbl
- [Tsutsumi and Fukuda 1980] M. Tsutsumi and I. Fukuda, “On solutions of the derivative nonlinear Schrödinger equation: existence and uniqueness theorem”, *Funkcial. Ekvac.* **23**:3 (1980), 259–277. MR Zbl
- [Tsutsumi and Fukuda 1981] M. Tsutsumi and I. Fukuda, “On solutions of the derivative nonlinear Schrödinger equation, II”, *Funkcial. Ekvac.* **24**:1 (1981), 85–94. MR Zbl
- [Win 2010] Y. Y. S. Win, “Global well-posedness of the derivative nonlinear Schrödinger equations on T ”, *Funkcial. Ekvac.* **53**:1 (2010), 51–88. MR Zbl
- [Wu 2013] Y. Wu, “Global well-posedness for the nonlinear Schrödinger equation with derivative in energy space”, *Anal. PDE* **6**:8 (2013), 1989–2002. MR Zbl
- [Wu 2015] Y. Wu, “Global well-posedness on the derivative nonlinear Schrödinger equation”, *Anal. PDE* **8**:5 (2015), 1101–1112. MR Zbl

Received 29 Jan 2021. Revised 28 Jun 2021. Accepted 6 Oct 2021.

ROWAN KILLIP: killip@math.ucla.edu

Department of Mathematics, University of California, Los Angeles, Los Angeles, CA, United States

MARIA NTEKOUME: maria.ntekoume@rice.edu

Department of Mathematics, Rice University, Houston, TX, United States

MONICA VIŞAN: visan@math.ucla.edu

Math Sciences, University of California, Los Angeles, Los Angeles, CA, United States

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard Université Paris Sud XI, France
patrick.gerard@universite-paris-saclay.fr

Clément Mouhot Cambridge University, UK
c.mouhot@dpms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	András Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 5 2023

Nonuniform stability of damped contraction semigroups	1089
RALPH CHILL, LASSI PAUNONEN, DAVID SEIFERT, REINHARD STAHN and YURI TOMILOV	
Long time solutions for quasilinear Hamiltonian perturbations of Schrödinger and Klein–Gordon equations on tori	1133
ROBERTO FEOLA, BENOÎT GRÉBERT and FELICE IANDOLI	
An extension problem, trace Hardy and Hardy’s inequalities for the Ornstein–Uhlenbeck operator	1205
PRITAM GANGULY, RAMESH MANNA and SUNDARAM THANGAVELU	
On the well-posedness problem for the derivative nonlinear Schrödinger equation	1245
ROWAN KILLIP, MARIA NTEKOUME and MONICA VIŞAN	
Exponential integrability in Gauss space	1271
PAATA IVANISVILI and RYAN RUSSELL	