

ANALYSIS & PDE

Volume 16

No. 6

2023

EVGENIOS T.A. KAKARIADIS, ELIAS G. KATSOULIS,
MARCELO LACA AND XIN LI

**COUNIVERSALITY AND CONTROLLED MAPS
ON PRODUCT SYSTEMS OVER RIGHT LCM SEMIGROUPS**

COUNIVERSALITY AND CONTROLLED MAPS ON PRODUCT SYSTEMS OVER RIGHT LCM SEMIGROUPS

EVGENIOS T.A. KAKARIADIS, ELIAS G. KATSOUNIS, MARCELO LACA AND XIN LI

We study the structure of C^* -algebras associated with compactly aligned product systems over group embeddable right LCM semigroups. Towards this end we employ controlled maps and a controlled elimination method that associates the original cores to those of the controlling pair, and we combine these with applications of the C^* -envelope theory for cosystems of nonselfadjoint operator algebras recently produced. We derive several applications of these methods that generalize results on single C^* -correspondences.

First we show that if the controlling group is exact then the couniversal C^* -algebra of the product system coincides with the quotient of the Fock C^* -algebra by the ideal of strong covariance relations. We show that if the controlling group is amenable then the product system is amenable. In particular if the controlling group is abelian then the couniversal C^* -algebra is the C^* -envelope of the tensor algebra.

Secondly we give necessary and sufficient conditions for the Fock C^* -algebra to be nuclear and exact. When the controlling group is amenable we completely characterize nuclearity and exactness of any equivariant injective Nica-covariant representation of the product system.

Thirdly we consider controlled maps that enjoy a saturation property. In this case we induce a compactly aligned product system over the controlling pair that shares the same Fock representation, and preserves injectivity. By using couniversality, we show that they share the same reduced covariance algebras. If in addition the controlling pair is a total order then the fixed point algebra of the controlling group induces a super product system that has the same reduced covariance algebra and is moreover reversible.

1. Introduction

Framework. In the present project we study further the effect of nonselfadjoint operator algebras and boundary theory of group coactions on the theory of C^* -algebras recently initiated by the authors and Dor-On in [Dor-On et al. 2022]. We work in the class of algebras of a compactly aligned product system X over a right LCM semigroup P in a group G with coefficients in a C^* -algebra A (for brevity we will say that such a pair (G, P) is a *weak right LCM inclusion*). Continuous product systems of Hilbert spaces were coined by Arveson [1989] for \mathbb{R}^+ , and their discrete counterparts were studied by Dinh [1991]. Motivated by Pimsner's seminal work [1997], Fowler [1999] studied product systems of correspondences over quasilattices. Since then discrete product systems have been studied by many authors (far too many to list here) and constitute an active area of research in their own right. Recently there has been a growing interest in passing from quasilattices to right LCM semigroups. Kwaśniewski and Larsen [2019]

MSC2020: 46L05, 46L08.

Keywords: product systems, Nica–Pimsner algebras, C^* -envelope.

studied the Toeplitz–Nica–Pimsner C^* -algebra $\mathcal{NT}(X)$ for right LCM semigroups proving Toeplitz–Cuntz–Krieger-type uniqueness theorems. Here we turn our focus to equivariant quotients with an eye towards Cuntz-type covariant realizations.

One of the main questions in this direction has been to identify the appropriate quotient of $\mathcal{NT}(X)$ so that faithful representations of A lift to faithful representations of the quotient. This cannot be expected to hold unconditionally. The next best hope is thus to locate the quotient of $\mathcal{NT}(X)$ so that faithful representations of A lift to faithful representations of its fixed point algebra. Sehnem [2019] has provided a full answer by introducing the strongly covariant representations. This generalizes the study of Cuntz–Nica–Pimsner relations, initiated by Sims and Yeend [2010], and later continued by Carlsen, Larsen, Sims and Vittadello [Carlsen et al. 2011]. A second aim of [Carlsen et al. 2011] was to use these relations and provide a couniversal object by passing to an appropriate reduced quotient. This was achieved under extra conditions on the product system (such as injectivity or directness of the quasilattice).

Couniversality and boundary representations arise naturally in the context of nonselfadjoint operator algebras and their C^* -envelope in the sense of Arveson. With Dor-On, in [Dor-On et al. 2022] we introduced a coaction variant of the C^* -envelope and used it to fully answer the problem of Carlsen, Larsen, Sims and Vittadello [Carlsen et al. 2011] without any assumptions on the product system X . Even more, the results of [Dor-On et al. 2022] apply to weak right LCM inclusions (G, P) rather than just quasilattices; more specifically, the C^* -envelope $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$ of the Fock tensor algebra $\mathcal{T}_\lambda(X)^+$ with its normal coaction is couniversal for equivariant injective Nica-covariant representations of X . Seeing Sehnem’s covariance algebra $A \times_X P$ as the universal C^* -algebra of an induced Fell bundle we further showed that $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta})$ coincides with the reduced C^* -algebra of this Fell bundle, here denoted by $A \times_{X, \lambda} P$.

The algebraic structure of $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$ was studied in [Dor-On et al. 2022]. Pivotal in this endeavor was the remark that the strong covariance relations of Sehnem are actually filtered through the Fock representation. Following Sehnem [2019], we will denote by $A \times_X P$ the universal C^* -algebra with respect to the strongly covariant representations of X . We further consider the induced quotient $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ of $\mathcal{T}_\lambda(X)$ by the strong covariance relations. In [Dor-On et al. 2022] it is shown that the canonical map

$$q_{\text{sc}}(\mathcal{T}_\lambda(X)) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}) \simeq A \times_{X, \lambda} P \quad (1-1)$$

is faithful if and only if the normal coaction of $\mathcal{T}_\lambda(X)$ descends to a normal coaction on $q_{\text{sc}}(\mathcal{T}_\lambda(X))$, e.g., when G is exact.

The motivation for the present work is two-fold. On one hand we wish to explore further general settings that entail normality of the coaction of $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ and thus identify the algebraic structure of the couniversal object. Our main theorem here is that this happens when (G, P) is controlled by another weak right LCM inclusion $(\mathcal{G}, \mathcal{P})$ with \mathcal{G} exact. When \mathcal{G} is abelian we can further induce dual actions on the C^* -algebras. This has the remarkable consequence that the canonical $*$ -epimorphism

$$C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+) \quad (1-2)$$

is faithful. On the other hand we wish to use the couniversal property in such a context and apply it in the identification of C^* -algebras. The quotient by the strong covariance relations is used as a model in several constructions, and this line of reasoning allows to show functoriality without checking a long list of C^* -properties. This is quite pleasing in particular because reduced C^* -algebras do not enjoy a priori universal properties. In fact we follow the reverse route of using the identification of reduced objects and then lift them to $*$ -isomorphisms of the universal ones.

Main results. Controlled maps $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ between quasilattice ordered groups were introduced by Laca and Raeburn [1996] with the purpose of extending the range of application of the faithfulness and uniqueness theorems for Toeplitz algebras of quasilattice ordered groups. The key idea is that (G, P) is amenable in the sense of Nica [1992], provided that \mathcal{G} is an amenable group. A similar notion of controlled maps was formulated simultaneously and independently by Crisp [1999] to prove that some Artin monoids inject in their groups. The combination of these two sets of ideas led to the amenability and nonamenability results for Artin monoids by Crisp and the third author [Crisp and Laca 2002]. Similar results can be derived for the Fock algebra $\mathcal{T}_\lambda(X)$ of a product system over P , as it has a \mathcal{P} -core that can be expressed as a direct sum of matrix algebras (see for example the proof of Theorem 6.4). As a consequence one obtains for example that compactly aligned product systems over the free semigroup \mathbb{F}_+^n are amenable, although the group \mathbb{F}^n is not, the reason being that the pair $(\mathbb{F}^n, \mathbb{F}_+^n)$ is controlled by its abelianization or by its length map on $(\mathbb{Z}, \mathbb{Z}_+)$.

However this type of argument is no longer valid for equivariant quotients as these relations live in the diagonal of the P -core (and thus in the \mathcal{P} -core). An elimination method was recently developed in [Kakariadis 2020] when $(\mathcal{G}, \mathcal{P}) = (\mathbb{Z}^n, \mathbb{Z}_+^n)$ with the purpose of studying nuclearity and exactness properties. By building further on these techniques, in the subsection on page 1458 we give a controlled elimination method for passing from the \mathcal{P} -cores to the P -cores of injective Nica-covariant representations. Essentially the method asserts that any relation in a \mathcal{P} -core must live at the diagonal and thus in a P -core. We then use this to lift all properties from the realm of the P -fixed point algebras to the \mathcal{P} -fixed point algebras. For example this applies to the fixed point algebra property of Sehnen’s algebra [2019]; see Corollary 5.8. In particular exactness of \mathcal{G} impacts on the maps appearing in (1-1).

Theorem A (Theorem 6.1). *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Consider the canonical $*$ -epimorphisms*

$$q_{sc}(\mathcal{T}_\lambda(X)) \rightarrow A \times_{X, \lambda} P \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \rightarrow C_{env}^*(\mathcal{T}_\lambda(X)^+). \tag{1-3}$$

If \mathcal{G} is exact then the left map is faithful. If in addition \mathcal{G} is abelian then the right map is also faithful.

Theorem A implies that the coaction on $q_{sc}(\mathcal{T}_\lambda(X))$ is normal when \mathcal{G} is exact. As pointed out in [Dor-On et al. 2022], this implies that the reduced Hao–Ng problem over discrete group actions has a positive answer (Remark 6.3). A similar method applies whenever the C^* -envelope functor is stable under crossed products, e.g., for dynamics over abelian locally compact groups or when the tensor algebra is hyperrigid [Katsoulis 2020; Katsoulis and Ramsey 2019], and we leave this to the interested

reader. A further consequence of Theorem A is that amenability of \mathcal{G} implies amenability of the product system and thus universality of the reduced constructions (Theorem 6.4). The case of abelian \mathcal{G} directly generalizes the results of [Dor-On and Katsoulis 2020]. There is further potential for Takai duality results even when (G, P) does not admit a dual. A further consequence of Theorem A provides a generalization of the extension theorem of [Katsoulis and Ramsey 2019], which recognizes a Fock tensor algebra by the presence of a coaction (Corollary 6.7).

Another application of the controlled elimination method concerns nuclearity and exactness results. It has been observed by Katsura [2004] that nuclearity of a Cuntz–Pimsner algebra is equivalent to the coefficient algebra being nuclearly embedded in the fixed point algebra. Kakariadis [2020] produced similar results for \mathbb{Z}_+^n . In Theorem 6.11 we first give an equivalent characterization for nuclearity of $\mathcal{T}_\lambda(X)$ for right LCM semigroups. Although our original goal was to exploit $A \times_X P$, we tackle any equivariant quotient of $\mathcal{NT}(X)$ that is injective on A .

Theorem B (Theorem 6.12, Theorem 6.13). *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable and let X be a compactly aligned product system over P with coefficients in A . Let (π, t) be an equivariant injective Nica-covariant representation of X . Then:*

- (i) *A is exact if and only if $C^*(\pi, t)$ is exact.*
- (ii) *$A \hookrightarrow C^*(\pi, t)$ is nuclear if and only if $C^*(\pi, t)$ is nuclear.*

We emphasize that the controlled elimination process occurs at the level of representations. One might be intrigued to introduce a product system Y over \mathcal{P} that would share the same algebras with X over P . However it is not clear that such a procedure gives a compactly aligned product system. For this reason we introduce the notion of saturation for controlled maps, which preserves inclusions of ideals in the semigroups. Under this condition we do get a super product system on the *same* coefficient algebra that does the job.

Theorem C (Theorem 7.7). *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a saturated controlled map between weak right LCM inclusions. Let X be an (injective) compactly aligned product system over P with coefficients in A and let*

$$Y_h := \sum_{p \in \vartheta^{-1}(h)}^\oplus X_p \quad \text{for } h \in \mathcal{P}.$$

Then the collection $Y = \{Y_h\}_{h \in \mathcal{P}}$ is an (injective) compactly aligned product system over \mathcal{P} with coefficients in A such that $\mathcal{T}_\lambda(X)^+ \simeq \mathcal{T}_\lambda(Y)^+$ with

$$\mathcal{T}_\lambda(X) \simeq \mathcal{T}_\lambda(Y) \quad \text{and} \quad A \times_{X, \lambda} P \simeq A \times_{Y, \lambda} \mathcal{P},$$

by $$ -isomorphisms that preserve the inclusions $X_p \mapsto Y_{\vartheta(p)}$ for all $p \in P$. These $*$ -isomorphisms further lift to $*$ -isomorphisms*

$$\mathcal{NT}(X) \simeq \mathcal{NT}(Y) \quad \text{and} \quad A \times_X P \simeq A \times_Y \mathcal{P}$$

that preserve the inclusions $X_p \hookrightarrow Y_{\vartheta(p)}$ for all $p \in P$.

Our method here is to show that the $*$ -isomorphism $\mathcal{T}_\lambda(X) \simeq \mathcal{T}_\lambda(Y)$ is canonical on the tensor algebras and then apply the C^* -envelope machinery to induce the $*$ -isomorphism $A \times_{X,\lambda} P \simeq A \times_{Y,\lambda} \mathcal{P}$. The saturation property can be induced by free products of abelian total orders, and is preserved by semidirect products. As a notable application of this method we deduce that Sehnen’s covariance algebra of a product system over \mathbb{F}_n^+ is nothing more than the Cuntz–Pimsner algebra of a single C^* -correspondence, in a similar way that the Nica–Cuntz–Pimsner algebra of \mathbb{F}_n^+ coincides with \mathcal{O}_n (Corollary 7.8).

We then take a closer look at total orders. To further motivate these results, recall that the Cuntz algebra \mathcal{O}_n may be viewed as the Cuntz–Pimsner algebra of a Hilbert bimodule over the n^∞ -hyperfinite C^* -algebra. In spite of the coefficient algebra of the latter being much larger, Hilbert bimodules are better behaved than other types of C^* -correspondences and they allow for a rich theory, including versions of Takai duality. Here we will show that the situation with \mathcal{O}_n generalizes to product systems that are controlled by exact total orders. Towards this end we consider *reversible* product systems for which the image of every fiber in $A \times_{X,\lambda} P$ is a Hilbert bimodule. We then show that reversible product systems produce all possible covariance algebras for weak right LCM inclusions that are controlled by total orders in a saturated way. The construction relies on using the fixed point algebra and generalizes results of Pimsner [1997], Abadie, Eilers and Exel [Abadie et al. 1998], Schweizer [2001], Kakariadis and Katsoulis [2012], and Meyer and Sehnen [2019]. However our proof uses the C^* -envelope machinery and thus avoids categorical arguments.

Theorem D (Theorem 7.15). *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a saturated controlled map between weak right LCM inclusions and suppose that $(\mathcal{G}, \mathcal{P})$ is a total order. Let X be an (injective) product system over P with coefficients in A . Then there exists an (injective) reversible product system Z over \mathcal{P} with coefficients in a C^* -algebra B such that*

$$A \subseteq B \quad \text{and} \quad X_p \subseteq Z_{\vartheta(p)} \quad \text{for all } p \in P \tag{1-4}$$

that satisfies

$$A \times_X P \simeq B \times_Z \mathcal{P} \quad \text{and} \quad A \times_{X,\lambda} P \simeq B \times_{Z,\lambda} \mathcal{P}, \tag{1-5}$$

by $*$ -isomorphisms that preserve the inclusions $X_p \hookrightarrow Z_{\vartheta(p)}$ for all $p \in P$.

Semigroup C^* -algebras have been an important source of inspiration for this study. Our results have a direct application to C^* -algebras of right LCM semigroups where $X_p = \mathbb{C}$ for every $p \in P$. In this case the Nica–Toeplitz C^* -algebra is denoted by $C_s^*(P)$ for the Nica-covariant representations of P and Theorem A (and in particular Theorem 6.4) is a direct generalization of [Crisp and Laca 2007, Theorem 4.7]. Faithfulness of the maps of Theorem A has been further investigated in [Kakariadis et al. 2022] for (not-necessarily right LCM) semigroups that embed in exact groups. Theorem B asserts that every quotient of $C_s^*(P)$ is nuclear and aligns with [Li 2013, Corollary 8.3] for quasilattices. Under the saturation property, Theorem C asserts that the operator algebras of P coincide with those of a product system Y over \mathcal{P} with $Y_h = \mathbb{C}^{|\vartheta^{-1}(h)|}$ for $h \in \mathcal{P}$. This follows a recurring idea of obtaining realizations of the same C^* -algebra in different classes. It has been shown in [Li 2017] that $\mathbb{C} \times_{\mathbb{C},\lambda} P$ can be realized as the partial crossed product of the smallest \mathcal{G} -invariant subspace of the fixed point algebra of $C_s^*(P)$ by G . Theorem D provides a similar (augmented) realization when ϑ is saturated and $(\mathcal{G}, \mathcal{P})$ is a total order.

Let us close with a remark on controlled maps. It has been known that controlled maps cannot handle HNN extensions of quasilattices as the height map does not have a trivial kernel on the semigroup. In order to resolve this, recently an Huef, Nucinkis, Sehnm and Yang [an Huef et al. 2021] introduced a more general definition of controlled maps for weak quasilattices that allows infinite descending chains and thus produces direct limits of matrix algebras. The controlled elimination arguments we provide here should be compatible with this general definition, as they refer to ideals of representations, which are compatible with direct limits.

Structure of the paper. In Section 2 we review the boundary theory and the theory of the cosystems from [Dor-On et al. 2022]. In Sections 3 and 4 we review the main elements of the product systems theory, and we see how they are enriched under the presence of a controlled map. We have included more details from the aforementioned paper in order to set the ground for the next sections, and also prove additional results that are not covered there. In Section 5 we present the controlled elimination method. Section 6 contains the applications to Sehnm’s covariance algebra, the structure of the couniversal C^* -algebra, amenable product systems, nuclearity and exactness, and the reduced Hao–Ng problem. In Section 7 we give the product system reparametrizations under the saturation property with applications to reversible product systems.

2. Operator algebras and their coactions

Operator algebras. The reader may refer to [Blecher and Le Merdy 2004; Paulsen 2002] for the general theory of nonselfadjoint operator algebras and dilations of their representations.

Let \mathcal{A} be an operator algebra, which in this paper means a subalgebra of $\mathcal{B}(H)$ for a Hilbert space H . We say that (C, ι) is a C^* -cover of \mathcal{A} if $\iota : \mathcal{A} \rightarrow C$ is a completely isometric representation with $C = C^*(\iota(\mathcal{A}))$. The C^* -envelope $C_{\text{env}}^*(\mathcal{A})$ of \mathcal{A} is a C^* -cover $(C_{\text{env}}^*(\mathcal{A}), \iota)$ with the following couniversal property: if (C', ι') is a C^* -cover of \mathcal{A} then there exists a (necessarily unique) $*$ -epimorphism $\Phi : C' \rightarrow C_{\text{env}}^*(\mathcal{A})$ such that $\Phi(\iota'(a)) = \iota(a)$ for all $a \in \mathcal{A}$. Arveson [1969] defined the C^* -envelope in and computed it for a variety of operator algebras, predicting its existence in general. Ten years later Hamana [1979] confirmed Arveson’s prediction by proving the existence of injective envelopes for the unital case. The C^* -envelope is the C^* -algebra generated in the injective envelope of \mathcal{A} once this is endowed with the Choi–Effros C^* -structure.

Dritschel and McCullough [2005] provided an alternative proof based on maximal dilations for the unital case. A *dilation* of a representation $\phi : \mathcal{A} \rightarrow \mathcal{B}(H)$ is a representation $\phi' : \mathcal{A} \rightarrow \mathcal{B}(H')$ such that $H \subseteq H'$ and $\phi(a) = P_H \phi'(a)|_H$ for all $a \in \mathcal{A}$. A completely contractive map $\phi : \mathcal{A} \rightarrow \mathcal{B}(H)$ is called *maximal* if every dilation $\phi' : \mathcal{A} \rightarrow \mathcal{B}(H')$ is trivial, i.e., $P_H \phi'(a) = \phi(a) = \phi'(a)|_H$ for all $a \in \mathcal{A}$. It follows that the C^* -envelope is the C^* -algebra generated by a maximal completely isometric representation.

It does not hold in general that if $\pi : C_{\text{env}}^*(\mathcal{A}) \rightarrow \mathcal{B}(H)$ is a $*$ -representation then it is the unique contractive completely positive extension of $\pi|_{\mathcal{A}}$. The algebra \mathcal{A} is called *hyperrigid* if this is the case for any representation π of $C_{\text{env}}^*(\mathcal{A})$. An operator algebra \mathcal{A} is said to be *Dirichlet* if

$$C_{\text{env}}^*(\mathcal{A}) = \overline{\mathcal{A} + \mathcal{A}^*}.$$

Equivalently, \mathcal{A} is Dirichlet if there exists a C^* -cover (C, ι) of \mathcal{A} such that $C = \overline{\iota(\mathcal{A}) + \iota(\mathcal{A})^*}$, in which case $C = C_{\text{env}}^*(\mathcal{A})$. It follows that Dirichlet algebras are automatically hyperrigid.

Coactions on operator algebras. If \mathcal{X} and \mathcal{Y} are subspaces of some $\mathcal{B}(H)$ then we write

$$[\mathcal{X}\mathcal{Y}] := \overline{\text{span}\{xy \mid x \in \mathcal{X}, y \in \mathcal{Y}\}}.$$

All groups and semigroups we consider are discrete and unital. Further, we denote the spatial tensor product by \otimes .

For a discrete group G we write u_g for the unitary generator associated with $g \in G$ in the full group C^* -algebra $C^*(G)$. We write λ_g for the generators of the left regular representation $C_\lambda^*(G)$. We write $\lambda : C^*(G) \rightarrow C_\lambda^*(G)$ for the canonical $*$ -epimorphism. Recall that $C^*(G)$ admits a faithful $*$ -homomorphism

$$\Delta : C^*(G) \rightarrow C^*(G) \otimes C^*(G) \quad \text{such that } \Delta(u_g) = u_g \otimes u_g,$$

given by the universal property of $C^*(G)$, and with left inverse given by $\text{id} \otimes \chi$ for the character χ of $C^*(G)$. We will require some preliminaries from [Dor-On et al. 2022] on coactions on operator algebras.

Definition 2.1 [Dor-On et al. 2022, Definition 3.1]. Let \mathcal{A} be an operator algebra. A *coaction of G on \mathcal{A}* is a completely isometric representation $\delta : \mathcal{A} \rightarrow \mathcal{A} \otimes C^*(G)$ such that the linear span of the induced subspaces

$$\mathcal{A}_g := \{a \in \mathcal{A} \mid \delta(a) = a \otimes u_g\}$$

is norm-dense in \mathcal{A} , in which case δ satisfies the coaction identity

$$(\delta \otimes \text{id}_{C^*(G)})\delta = (\text{id}_{\mathcal{A}} \otimes \Delta)\delta.$$

If, in addition, the map $(\text{id} \otimes \lambda)\delta$ is injective, then the coaction δ is called *normal*.

If \mathcal{A} is an operator algebra and $\delta : \mathcal{A} \rightarrow \mathcal{A} \otimes C^*(G)$ is a coaction on \mathcal{A} , then we will refer to the triple (\mathcal{A}, G, δ) as a *cosystem*. A map $\phi : \mathcal{A} \rightarrow \mathcal{A}'$ between two cosystems (\mathcal{A}, G, δ) and $(\mathcal{A}', G, \delta')$ is said to be *G -equivariant*, or simply *equivariant*, if $\delta'\phi = (\phi \otimes \text{id})\delta$.

If (\mathcal{A}, G, δ) is a cosystem then $\mathcal{A}_r \cdot \mathcal{A}_s \subseteq \mathcal{A}_{r,s}$ for all $r, s \in G$, since δ is a homomorphism.

Remark 2.2 [Dor-On et al. 2022]. Suppose that (\mathcal{A}, G, δ) is a cosystem and that δ extends to a $*$ -homomorphism $\delta : C^*(\mathcal{A}) \rightarrow C^*(\mathcal{A}) \otimes C^*(G)$ that satisfies the coaction identity

$$(\delta \otimes \text{id})\delta(c) = (\text{id} \otimes \Delta)\delta(c) \quad \text{for all } c \in C^*(\mathcal{A}).$$

Then δ is automatically nondegenerate on $C^*(\mathcal{A})$ in the sense that

$$[\delta(C^*(\mathcal{A}))C^*(\mathcal{A}) \otimes C^*(G)] = C^*(\mathcal{A}) \otimes C^*(G).$$

Moreover, Definition 2.1 covers that of full coactions of Quigg [1996] when \mathcal{A} is a C^* -algebra. In this case δ is a faithful $*$ -homomorphism and we have that

$$(\mathcal{A}_g)^* = \{a^* \in \mathcal{A} \mid \delta(a^*) = a^* \otimes u_{g^{-1}}\} = \mathcal{A}_{g^{-1}}.$$

Due to the Fell absorption principle, the existence of a “reduced” coaction implies that of a normal coaction.

Proposition 2.3 [Dor-On et al. 2022, Proposition 3.4]. *Let \mathcal{A} be an operator algebra. Suppose there is a group G that induces a grading on \mathcal{A} , i.e., there are subspaces $\{\mathcal{A}_g\}_{g \in G}$ such that $\sum_{g \in G} \mathcal{A}_g$ is norm-dense in \mathcal{A} , and a completely isometric homomorphism*

$$\delta_\lambda : \mathcal{A} \rightarrow \mathcal{A} \otimes C_\lambda^*(G)$$

such that

$$\delta_\lambda(a_g) = a_g \otimes \lambda_g \text{ for all } a_g \in \mathcal{A}_g, \text{ for all } g \in G.$$

Then \mathcal{A} admits a normal coaction δ of G such that $\delta_\lambda = (\text{id} \otimes \lambda)\delta$.

Example 2.4. The reduced group C^* -algebra $C_\lambda^*(G)$ admits a faithful $*$ -homomorphism

$$\Delta_\lambda : C_\lambda^*(G) \rightarrow C_\lambda^*(G) \otimes C_\lambda^*(G) \text{ such that } \Delta_\lambda(\lambda_g) = \lambda_g \otimes \lambda_g.$$

Thus $C_\lambda^*(G)$ admits a normal coaction δ of G such that $\Delta_\lambda = (\text{id} \otimes \lambda)\delta$.

Definition 2.5 [Dor-On et al. 2022, Definition 3.6]. Let (\mathcal{A}, G, δ) be a cosystem. A triple (C, ι, δ_C) is called a C^* -cover for (\mathcal{A}, G, δ) if (C, ι) is a C^* -cover of \mathcal{A} and $\delta_C : C \rightarrow C \otimes C^*(G)$ is a coaction on C such that the diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\iota} & C \\ \downarrow \delta & & \downarrow \delta_C \\ \mathcal{A} \otimes C^*(G) & \xrightarrow{\iota \otimes \text{id}} & C \otimes C^*(G) \end{array}$$

commutes. When the coaction is understood we will say that C is a C^* -cover for \mathcal{A} over G .

Definition 2.6 [Dor-On et al. 2022, Definition 3.7]. Let (\mathcal{A}, G, δ) be a cosystem. The C^* -envelope of (\mathcal{A}, G, δ) is a C^* -cover $(C_{\text{env}}^*(\mathcal{A}, G, \delta), \iota, \delta_{\text{env}})$ such that: for every C^* -cover (C', ι', δ') of (\mathcal{A}, G, δ) there exists a $*$ -epimorphism $\Phi : C' \rightarrow C_{\text{env}}^*(\mathcal{A}, G, \delta)$ that fixes \mathcal{A} and intertwines the coactions, i.e., the diagram

$$\begin{array}{ccc} \iota'(\mathcal{A}) & \xrightarrow{\delta'} & C' \otimes C^*(G) \\ \downarrow \Phi & & \downarrow \Phi \otimes \text{id} \\ \iota(\mathcal{A}) & \xrightarrow{\delta_{\text{env}}} & C_{\text{env}}^*(\mathcal{A}, G, \delta) \otimes C^*(G) \end{array}$$

is commutative on \mathcal{A} , and thus is commutative on C' .

The existence of the C^* -envelope of a cosystem was proved in [Dor-On et al. 2022] by a direct computation that uses the C^* -envelope of the ambient operator algebra. In order to state the result explicitly we need to make some preliminary remarks and establish the notation. Suppose (\mathcal{A}, G, δ)

is a cosystem, let $i : \mathcal{A} \rightarrow C_{\text{env}}^*(\mathcal{A})$ be the C^* -envelope of \mathcal{A} , and recall that the spatial tensor product of completely isometric maps is completely isometric. Then the representation of \mathcal{A} obtained via the composition

$$\mathcal{A} \xrightarrow{\delta} \mathcal{A} \otimes C^*(G) \xrightarrow{i \otimes \text{id}} C_{\text{env}}^*(\mathcal{A}) \otimes C^*(G)$$

is completely isometric, and the C^* -algebra

$$C^*((i \otimes \text{id})\delta(\mathcal{A})) := C^*(i(a_g) \otimes u_g \mid g \in G)$$

becomes a C^* -cover of \mathcal{A} . This C^* -cover is special because it admits a coaction $\text{id} \otimes \Delta$, such that the triple

$$(C^*(i(a_g) \otimes u_g \mid g \in G), (i \otimes \text{id})\delta, \text{id} \otimes \Delta)$$

is a C^* -cover for (\mathcal{A}, G, δ) . The following theorem summarizes fundamental results about existence and representations of C^* -envelopes for cosystems.

Theorem 2.7 [Dor-On et al. 2022, Theorem 3.8, Corollary 3.9 and Corollary 3.10]. *Let (\mathcal{A}, G, δ) be a cosystem and let $i : \mathcal{A} \rightarrow C_{\text{env}}^*(\mathcal{A})$ be the inclusion map. Then*

$$(C_{\text{env}}^*(\mathcal{A}, G, \delta), \iota, \delta_{\text{env}}) \simeq (C^*(i(a_g) \otimes u_g \mid g \in G), (i \otimes \text{id})\delta, \text{id} \otimes \Delta).$$

If in addition δ is normal on \mathcal{A} , then δ_{env} is normal on $C_{\text{env}}^(\mathcal{A}, G, \delta)$.*

Moreover, if $\Phi : C_{\text{env}}^(\mathcal{A}, G, \delta) \rightarrow B$ is a $*$ -homomorphism that is completely isometric on \mathcal{A} , then it is faithful on the fixed point algebra of $C_{\text{env}}^*(\mathcal{A}, G, \delta)$.*

Remark 2.8. A coaction of an abelian group G is equivalent to point-norm continuous actions $\{\beta_\gamma\}_{\gamma \in \hat{G}}$ of the dual group \hat{G} . Since every β_γ is a completely isometric automorphism it extends to the C^* -envelope. Hence the C^* -envelope of a cosystem coincides with the usual C^* -envelope of the ambient operator algebra when G is abelian. Equivalently, every coaction of an abelian group on an operator algebra lifts to a coaction on its C^* -envelope. As pointed out in [Dor-On et al. 2022], it is unknown if this is the case for general amenable groups.

Group homomorphisms implement coactions. Note that the following proposition for $\mathcal{G} = \{e_G\}$ says nothing more than that every C^* -cover of a cosystem is a C^* -cover of the ambient operator algebra.

Proposition 2.9. *Let $(\mathcal{A}, G, \delta_G)$ be a (normal) cosystem and let $\vartheta : G \rightarrow \mathcal{G}$ be a group homomorphism. Then \mathcal{G} induces a (normal) coaction $\delta_{\mathcal{G}}$ on \mathcal{A} . Thus every C^* -cover of \mathcal{A} over G is also a C^* -cover of \mathcal{A} over \mathcal{G} .*

Proof. By the universal property of $C^*(G)$ we have a $*$ -homomorphism

$$\tilde{\vartheta} : C^*(G) \rightarrow C^*(\mathcal{G}), \quad u_g \mapsto u_{\vartheta(g)}.$$

We then have the canonical completely contractive homomorphism

$$\delta_{\mathcal{G}} : \mathcal{A} \xrightarrow{\delta_G} \mathcal{A} \otimes C^*(G) \xrightarrow{\text{id} \otimes \tilde{\vartheta}} \mathcal{A} \otimes C^*(\mathcal{G}),$$

which has $\text{id} \otimes \chi$ as a completely contractive left inverse. By definition we have that

$$\mathcal{A}_h := \{a \in \mathcal{A} \mid \delta_G(a) = a \otimes u_h\} \supseteq \{a \in \mathcal{A}_g \mid \vartheta(g) = h\}$$

and thus

$$\mathcal{A} = \overline{\sum_{g \in G} \mathcal{A}_g} \subseteq \overline{\sum_{h \in G} \mathcal{A}_h} \subseteq \mathcal{A}.$$

Hence $(\text{id} \otimes \tilde{\vartheta})\delta_G$ defines a coaction of \mathcal{G} on \mathcal{A} .

Next suppose that δ_G is normal and let $\delta_{G,\lambda} = (\text{id} \otimes \lambda)\delta_G$. Let $\delta_{\mathcal{G}}$ be the coaction induced by δ_G . By Fell’s absorption principle we have that the map $\lambda_g \mapsto \lambda_g \otimes \lambda_{\vartheta(g)}$ gives a faithful $*$ -homomorphism of $C^*_\lambda(G)$ and thus we get the induced completely isometric representation

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\delta_{G,\lambda}} & \overline{\text{alg}\{a_g \otimes \lambda_g \mid g \in G\}} \\ \delta_{G,\lambda} \downarrow & & \downarrow \\ \overline{\text{alg}\{a_g \otimes \lambda_{\vartheta(g)} \mid g \in G\}} & \xleftarrow{\delta_{G,\lambda}^{-1} \otimes \text{id}} & \overline{\text{alg}\{a_g \otimes \lambda_g \otimes \lambda_{\vartheta(g)} \mid g \in G\}} \end{array}$$

which induces a faithful $*$ -homomorphism $\delta_{\mathcal{G},\lambda}$. It follows that $\delta_{G,\lambda} = (\text{id} \otimes \lambda)\delta_G$ and thus $\delta_{\mathcal{G}}$ is a normal coaction of \mathcal{G} on \mathcal{A} . □

Let us close this section with some remarks on topological gradings from [Exel 1997; 2017]. Recall that a *topological grading* $\{\mathcal{B}_g\}_{g \in G}$ of a C^* -algebra \mathcal{B} consists of linearly independent subspaces that span a dense subspace of \mathcal{B} and are compatible with the group G , i.e., $\mathcal{B}_g^* = \mathcal{B}_{g^{-1}}$ and $\mathcal{B}_g \cdot \mathcal{B}_h \subseteq \mathcal{B}_{gh}$. By [Exel 1997, Theorem 3.3] the linear independence condition can be substituted by the existence of a conditional expectation on \mathcal{B}_e . The maximal C^* -algebra $C^*(\mathcal{B})$ of \mathcal{B} is defined as universal with respect to the representations of \mathcal{B} . The reduced C^* -algebra $C^*_\lambda(\mathcal{B})$ of \mathcal{B} is defined by the left regular representation of \mathcal{B} on $\ell^2(\mathcal{B})$.

Definition 2.10. Let $\mathcal{B} = \{\mathcal{B}_g\}_{g \in G}$ be a topological grading over a group G in a C^* -algebra $C^*(\mathcal{B})$ that it generates, with completely contractive Fourier maps $E_g : C^*(\mathcal{B}) \rightarrow \mathcal{B}_g$, i.e.,

$$E_g(b) = \delta_{g,h}b \quad \text{for all } b \in \mathcal{B}_h \text{ and } g, h \in G.$$

An ideal $\mathcal{I} \triangleleft C^*(\mathcal{B})$ is called *induced* if $\mathcal{I} = \langle \mathcal{I} \cap \mathcal{B}_e \rangle$. An ideal $\mathcal{I} \triangleleft C^*(\mathcal{B})$ is called *Fourier* if $E_g(f) \subseteq \mathcal{I}$ for every $f \in \mathcal{I}$.

Remark 2.11. It follows that an ideal $\mathcal{I} \triangleleft C^*(\mathcal{B})$ is Fourier if and only if $E_e(f^*f) \in \mathcal{I}$ for all $f \in \mathcal{I}$. Every induced ideal is a Fourier ideal. The converse holds if G is exact and E_e is a faithful conditional expectation. These can be found at [Exel 2017, Proposition 23.9].

A topological grading defines a *Fell bundle* and once a representation of a Fell bundle is established the two notions are the same. In a loose sense a Fell bundle \mathcal{B} over a discrete group G is a collection of Banach spaces $\{\mathcal{B}_g\}_{g \in G}$, often called *the fibers of \mathcal{B}* , that satisfy canonical algebraic properties and the C^* -norm properties; see [Exel 2017, Definition 16.1]. So we will alternate freely between Fell bundles

and topologically graded C^* -algebras. Spectral subspaces of coactions on C^* -algebras are an important source of topological gradings.

Definition 2.12. Let δ be a coaction of G on a C^* -algebra C and let $\mathcal{I} \triangleleft C$ be an ideal of C . We say that the quotient map is G -equivariant, or that the quotient C/\mathcal{I} is G -equivariant if δ descends to a coaction of G on C/\mathcal{I} .

Remark 2.13. If $\delta : C \rightarrow C \otimes C^*(G)$ is a coaction and $\mathcal{I} \triangleleft C$ is an induced ideal then δ descends to a faithful coaction of G on C/\mathcal{I} ; see for example [Carlsen et al. 2011, Proposition A.1]. The same holds for the normal actions when G is exact; see for example [Carlsen et al. 2011, Proposition A.5].

3. Operator algebras of product systems

C^* -correspondences. A C^* -correspondence X over A is a right Hilbert module over A with a left action given by a $*$ -homomorphism $\varphi_X : A \rightarrow \mathcal{L}X$. We write $\mathcal{L}X$ and $\mathcal{K}X$ for the adjointable operators and the compact operators of X , respectively. For two C^* -correspondences X, Y over the same A we write $X \otimes_A Y$ for the balanced tensor product over A . We say that X is unitarily equivalent to Y (and write $X \simeq Y$) if there is a surjective adjointable operator $U \in \mathcal{L}(X, Y)$ such that $\langle U\xi, U\eta \rangle = \langle \xi, \eta \rangle$ and $U(a\xi b) = aU(\xi)b$ for all $\xi, \eta \in X$ and $a, b \in A$. A C^* -correspondence is called *injective* if the left action is injective.

A *representation* (π, t) of a C^* -correspondence is a left module map that preserves the inner product. Then (π, t) is automatically a bimodule map. Moreover there exists a $*$ -homomorphism ψ on $\mathcal{K}X$ such that $\psi(\theta_{\xi, \eta}) = t(\xi)t(\eta)^*$ for all $\theta_{\xi, \eta} \in \mathcal{K}X$. When π is injective, then both t and ψ are isometric. A representation (π, t) is called *covariant* if it satisfies $\pi(a) = \psi(\varphi_X(a))$ for all a in Katsura’s ideal $J_X := \ker \varphi_X^\perp \cap \varphi_X^{-1}(\mathcal{K}X)$.

Toeplitz algebras. Let P be a unital subsemigroup of a group G . We will write P^* for the set of elements in P that are invertible in P . A *product system* X over P is a family $\{X_p \mid p \in P\}$ of C^* -correspondences over the same C^* -algebra A such that:

- (i) $X_e = A$.
- (ii) There are multiplication rules $X_p \otimes_A X_q \simeq_{u_{p,q}} X_{pq}$ for every $p, q \in P \setminus \{e\}$.
- (iii) There are multiplication rules $A \otimes_A X_p \simeq_{u_{e,p}} [A \cdot X_p]$ and $X_p \otimes_A A \simeq_{u_{p,e}} [X_p \cdot A] = X_p$ for all $p \in P$.
- (iv) The multiplication rules are associative in the sense that

$$u_{pq,r}(u_{p,q} \otimes \text{id}_{X_r}) = u_{p,q,r}(\text{id}_{X_p} \otimes u_{q,r}) \quad \text{for all } p, q, r \in P.$$

We say that X is *injective* if every X_p is injective. If $x \in P^*$ then the multiplication rules impose that

$$X_x \otimes_A X_{x^{-1}} \simeq A \simeq X_{x^{-1}} \otimes_A X_x.$$

In particular every such X_x is nondegenerate since

$$A \otimes_A X_x \simeq X_x \otimes_A X_{x^{-1}} \otimes_A X_x \simeq X_x \otimes_A A = X_x.$$

Throughout this work we will be assuming that all left actions are nondegenerate. We do this in order to be able to use freely the results from [Dor-On et al. 2022; Sehnem 2019]. Nevertheless it is possible that this assumption can be removed.

Henceforth we will suppress the use of symbols for the multiplication rules. Thus we write $\xi_p \xi_q$ for the image of $\xi_p \otimes \xi_q$ under $u_{p,q}$, and so

$$\varphi_{pq}(a)(\xi_p \xi_q) = (\varphi_p(a)\xi_p)\xi_q \quad \text{for all } a \in A \quad \text{and} \quad \xi_p \in X_p, \quad \xi_q \in X_q.$$

The product system structure gives maps

$$i_p^{pq} : \mathcal{L}X_p \rightarrow \mathcal{L}X_{pq} \quad \text{such that} \quad i_p^{pq}(S)(\xi_p \xi_q) = (S\xi_p)\xi_q.$$

If $x \in P^*$ then $i_r^{rx} : \mathcal{L}X_r \rightarrow \mathcal{L}X_{rx}$ is a $*$ -isomorphism with inverse $i_{rx}^{rx^{-1}} : \mathcal{L}X_{rx} \rightarrow \mathcal{L}X_r$.

Definition 3.1. Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . A *Toeplitz representation* (π, t) of X consists of a family of representations (π, t_p) of X_p over A such that

$$t_p(\xi_p)t_q(\xi_q) = t_{pq}(\xi_p \xi_q) \quad \text{for all } \xi_p \in X_p, \quad \xi_q \in X_q.$$

The *Toeplitz algebra* $\mathcal{T}(X)$ of X is the universal C^* -algebra generated by A and X with respect to the representations of X . The *Toeplitz tensor algebra* $\mathcal{T}(X)^+$ of X is the subalgebra of $\mathcal{T}(X)$ generated by A and X .

If (π, t) is a Toeplitz representation then we write ψ_p for the induced representation on $\mathcal{K}X_p$. We obtain a bimodule triple $(\psi_r, \psi_{r,s}, \psi_s)$ on the bimodule $(\mathcal{K}X_r, \mathcal{K}(X_s, X_r), \mathcal{K}X_s)$ so that $\psi_{r,s}(\theta_{\xi_r, \xi_s}) = t_r(\xi_r)t_s(\xi_s)^*$. We will often interpret π as t_e or ψ_e to simplify our notation henceforth.

Proposition 3.2 [Dor-On et al. 2022, Proposition 2.4]. *Let X be a product system over P with coefficients in A . Let (π, t) be a Toeplitz representation of X . If $x \in P^*$ then*

$$t_x(X_x)^* = t_{x^{-1}}(X_{x^{-1}}).$$

If $w \in P$ and $x \in P^*$ then

$$i_w^{wx}(k_w) \in \mathcal{K}X_{wx} \quad \text{and} \quad \psi_{wx}(i_w^{wx}(k_w)) = \psi_w(k_w) \quad \text{for all } k_w \in \mathcal{K}X_w.$$

Suppose that $\mathcal{T}(X)$ is faithfully represented by $(\tilde{\pi}, \tilde{t})$. By the universal property of $\mathcal{T}(X)$ there is a canonical $*$ -homomorphism

$$\tilde{\delta} : \mathcal{T}(X) \rightarrow \mathcal{T}(X) \otimes C^*(G), \quad \tilde{t}(\xi_p) \mapsto \tilde{t}(\xi_p) \otimes u_p.$$

Sehnem [2019, Lemma 2.2] has shown that $\tilde{\delta}$ is a nondegenerate and faithful coaction of $\mathcal{T}(X)$ when X is nondegenerate, with each spectral space $\mathcal{T}(X)_g$, with $g \in G$, given by the products

$$\tilde{t}_{p_1}(\xi_{p_1})\tilde{t}_{p_2}(\xi_{p_2})^* \cdots \tilde{t}_{p_n}(\xi_{p_n})^* \quad \text{for } p_1 p_2^{-1} \cdots p_n^{-1} = g.$$

We will do a little bit more for semigroup homomorphisms.

Definition 3.3. Let P and \mathcal{P} be unital subsemigroups of the groups G and \mathcal{G} , respectively. If $\vartheta : G \rightarrow \mathcal{G}$ is a group homomorphism such that $\vartheta(P) \subseteq \mathcal{P}$, we write $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ and say that ϑ is a *semigroup preserving homomorphism*.

Proposition 3.4. Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a semigroup preserving homomorphism and suppose that $(\tilde{\pi}, \tilde{\iota})$ is a faithful representation of $\mathcal{T}(X)$. Then there is a coaction of \mathcal{G} on $\mathcal{T}(X)$ such that

$$\tilde{\delta} : \mathcal{T}(X) \rightarrow \mathcal{T}(X) \otimes C^*(\mathcal{G}), \quad \tilde{\iota}(\xi_p) \mapsto \tilde{\iota}(\xi_p) \otimes u_{\vartheta(p)}.$$

Moreover, each spectral space $\mathcal{T}(X)_h$ with $h \in \mathcal{G}$ is given by the products of the form

$$\tilde{\iota}_{p_1}(\xi_{p_1})\tilde{\iota}_{p_2}(\xi_{p_2})^* \cdots \tilde{\iota}_{p_n}(\xi_{p_n})^* \quad \text{for } \vartheta(p_1)\vartheta(p_2)^{-1} \cdots \vartheta(p_n)^{-1} = h,$$

where we impose that $\tilde{\iota}_{p_i}(\xi_{p_i}) = I$ when $p_i = e_P$ and $h \neq e_{\mathcal{P}}$.

Proof. The universal property induces a $*$ -homomorphism $\tilde{\delta} : \mathcal{T}(X) \rightarrow \mathcal{T}(X) \otimes C^*(G)$. Moreover $\tilde{\delta}$ is injective with left inverse given by $\text{id} \otimes \chi$. By construction the fibers $[\mathcal{T}(X)]_g$ contain the generators of $\mathcal{T}(X)$. By Remark 2.2 and the definition of $\mathcal{T}(X)^+$, this gives the coaction of G . Proposition 2.9 provides the coaction of \mathcal{G} . \square

Remark 3.5. The Fock space representation of Fowler [2002] ensures that A , and thus X , embeds isometrically in $\mathcal{T}(X)$. In short, let $\mathcal{F}(X) = \sum_{q \in P} X_q$ and for $a \in A$ and $\xi_p \in X_p$ define $(\bar{\pi}, \bar{\iota}_p)$ by

$$\bar{\pi}(a)\xi_q = \varphi_q(a)\xi_q \quad \text{and} \quad \bar{\iota}_p(\xi_p)\xi_q = \xi_p\xi_q \quad \text{for all } \xi_q \in X_q.$$

Then every $(\bar{\pi}, \bar{\iota}_p)$ defines a representation of X_p and hence it induces a representation of $\mathcal{T}(X)$. By taking the compression at the (e, e) -entry we see that $\bar{\pi}$, and thus $\bar{\iota}_p$, is injective.

Definition 3.6. Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . The *Fock algebra* $\mathcal{T}_\lambda(X)$ is the C^* -algebra generated by the Fock representation $(\bar{\pi}, \bar{\iota})$. The *Fock tensor algebra* $\mathcal{T}_\lambda(X)^+$ of X is the subalgebra of $\mathcal{T}_\lambda(X)$ generated by A and X .

It is shown in [Dor-On et al. 2022, Proposition 4.1] that the Fock algebra admits an analogous normal coaction. Proposition 2.9 yields the next proposition.

Proposition 3.7. Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A , and let $\mathcal{T}_\lambda(X) = C^*(\bar{\pi}, \bar{\iota})$ be its associated Fock algebra. If $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ is a semigroup preserving homomorphism then there is a normal coaction of \mathcal{G} on $\mathcal{T}_\lambda(X)$ such that

$$\bar{\delta}_{\mathcal{G}} : \mathcal{T}_\lambda(X) \rightarrow \mathcal{T}_\lambda(X) \otimes C^*(\mathcal{G}), \quad \bar{\iota}(\xi_p) \mapsto \bar{\iota}(\xi_p) \otimes u_{\vartheta(p)}.$$

Moreover for each $h \in \mathcal{G}$ the spectral space $\mathcal{T}_\lambda(X)_h$ is the closed linear span of the products of the form

$$\bar{\iota}_{p_1}(\xi_{p_1})\bar{\iota}_{p_2}(\xi_{p_2})^* \cdots \bar{\iota}_{p_{n-1}}(\xi_{p_{n-1}})\bar{\iota}_{p_n}(\xi_{p_n})^* \quad \text{for } \vartheta(p_1)\vartheta(p_2)^{-1} \cdots \vartheta(p_n)^{-1} = h,$$

where we impose that $\bar{\iota}_{p_i}(\xi_{p_i}) = I$ when $p_i = e_P$ and $h \neq e_{\mathcal{P}}$.

In turn the coaction of \mathcal{G} induces a faithful conditional expectation of the following form.

Proposition 3.8. *Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a semigroup preserving homomorphism. Then $\mathcal{T}_\lambda(X)$ admits a faithful conditional expectation $\bar{E}_\mathcal{P}$ such that*

$$\bar{E}_\mathcal{P}(\bar{\psi}_{r,s}(k_{r,s})) = \delta_{\vartheta(r), \vartheta(s)} \bar{\psi}_{r,s}(k_{r,s}) \quad \text{for all } k_{r,s} \in \mathcal{K}(X_s, X_r).$$

Proof. Let $\bar{\delta}_\mathcal{G} : \mathcal{T}_\lambda(X) \rightarrow \mathcal{T}_\lambda(X) \otimes C^*(\mathcal{G})$ be the normal coaction and let $\omega_{e,e}$ be the faithful conditional expectation on $C^*_\lambda(\mathcal{G})$. Then $\mathcal{T}_\lambda(X)$ admits the faithful conditional expectation

$$\bar{E}_\mathcal{P} := (\text{id} \otimes \omega_{e,e})(\text{id} \otimes \lambda) \bar{\delta}_\mathcal{G}.$$

On the other hand for $h \in \mathcal{P}$ let $Y_h := \sum_{\vartheta(p)=h}^\oplus X_h$ and consider the projections $Q_h : \mathcal{F}(X) \rightarrow Y_h$. We will show that

$$\bar{E}_\mathcal{P}(\cdot) = \sum_{h \in \mathcal{P}} Q_h \cdot Q_h.$$

It suffices to check on the spanning elements of the form

$$f := \bar{t}_{p_1}(\xi_{p_1}) \bar{t}_{p_2}(\xi_{p_2})^* \cdots \bar{t}_{p_{n-1}}(\xi_{p_{n-1}}) \bar{t}_{p_n}(\xi_{p_n})^*,$$

where we impose that $\bar{t}_{p_i}(\xi_{p_i}) = I$ when $p_i = e_P$. For $p \in P$ we directly compute

$$\bar{E}_\mathcal{P}(f) = \begin{cases} f \xi_p & \text{if } \vartheta(p_1^{-1} p_2 \cdots p_{n-1} p_n) = e_\mathcal{G}, \\ 0 & \text{otherwise.} \end{cases}$$

If $f \xi_p \neq 0$ then it is in some X_r with $r = p_1^{-1} p_2 \cdots p_{n-1} p_n p$ which gives $\vartheta(r) = \vartheta(p)$. On the other hand we have that

$$\left(\sum_{h \in \mathcal{P}} Q_h f Q_h \right) \xi_p = \begin{cases} f \xi_p & \text{if } \vartheta(p_1^{-1} p_2 \cdots p_{n-1} p_n p) = \vartheta(p), \\ 0 & \text{otherwise.} \end{cases}$$

We have that $\vartheta(p_1^{-1} p_2 \cdots p_{n-1} p_n p) = \vartheta(p)$ if and only if $\vartheta(p_1^{-1} p_2 \cdots p_{n-1} p_n) = e_\mathcal{G}$ and so

$$\bar{E}_\mathcal{P}(f) = \sum_{h \in \mathcal{P}} Q_h f Q_h.$$

For the second part let $r, s \in P$ and $\xi_p \in Y_h$ so that $\vartheta(p) = h$. Then we directly compute

$$\begin{aligned} \bar{E}_\mathcal{P}(\bar{\psi}_{r,s}(k_{r,s})) \xi_p &= Q_h \bar{\psi}_{r,s}(k_{r,s}) \xi_p = \begin{cases} \bar{\psi}_{r,s}(k_{r,s}) \xi_p & \text{if } p = ss', \vartheta(p) = \vartheta(rs'), \\ 0 & \text{otherwise,} \end{cases} \\ &= \delta_{\vartheta(r), \vartheta(s)} \bar{\psi}_{r,s}(k_{r,s}) \xi_p, \end{aligned}$$

where we used that ϑ is a group homomorphism and so $\vartheta(s)\vartheta(s') = \vartheta(p) = \vartheta(r)\vartheta(s')$. As $p \in P$ is arbitrary the proof is complete. □

Covariance algebras and Cuntz–Nica–Pimsner algebras. Let us review Sehnm’s strong covariance relations [2019]. We will be using a description presented in [Dor-On et al. 2022]. Let P be a unital subsemigroup of a group G . For a finite set $F \subseteq G$ let

$$K_F := \bigcap_{g \in F} gP.$$

For $r \in P$ and $g \in F$ define the ideal of A given by

$$I_{r^{-1}K_{\{r,g\}}} := \begin{cases} \bigcap_{t \in K_{\{r,g\}}} \ker \varphi_{r^{-1}t} & \text{if } K_{\{r,g\}} \neq \emptyset \text{ and } r \notin K_{\{r,g\}}, \\ A & \text{otherwise.} \end{cases}$$

Then let

$$I_{r^{-1}(r \vee F)} := \bigcap_{g \in F} I_{r^{-1}K_{\{r,g\}}},$$

and define the C^* -correspondences

$$X_F := \bigoplus_{r \in P} X_r I_{r^{-1}(r \vee F)} \quad \text{and} \quad X_F^+ := \bigoplus_{g \in G} X_g F.$$

For every $p \in P$ define the representation $(\pi_F, t_{F,p})$ to X_F^+ given by

$$t_{F,p}(\xi_p)(\eta_r) = u_{p,r}(\xi_p \otimes \eta_r) \in X_{pr} I_{(pr)^{-1}(pr \vee pF)}, \quad \text{for all } \eta_r \in X_r I_{r^{-1}(r \vee F)}.$$

It is well defined as $I_{r^{-1}(r \vee F)} = I_{(pr)^{-1}(pr \vee pF)}$ for all $r \in P$, and $I_{r^{-1}(r \vee F)} = I_{(s^{-1}r)^{-1}(s^{-1}r \vee s^{-1}F)}$ for all $r \in sP$. This provides a representation (π_F, t_F) of X on $\mathcal{L}(X_F^+)$ that integrates to a representation

$$\Phi_F : \mathcal{T}(X) \rightarrow \mathcal{L}(X_F^+).$$

Now consider the projections $Q_{g,F} : X_F^+ \rightarrow X_g F$ and define

$$\|f\|_F := \|Q_{e,F} \Phi_F(f) Q_{e,F}\| \quad \text{for all } f \in [\mathcal{T}(X)]_e.$$

In particular we have that

$$t_{F,p}(\xi_p) Q_{g,F} = Q_{pg,F} t_{F,p}(\xi_p) \quad \text{and} \quad t_{F,p}(\xi_p)^* Q_{g,F} = Q_{p^{-1}g,F} t_{F,p}(\xi_p)^*,$$

and so $Q_{e,F}$ is reducing for the fixed point algebra $[\mathcal{T}(X)]_e$ under Φ_F .

Definition 3.9 [Sehnm 2019, Definition 3.2]. A Toeplitz representation is called *strongly covariant* if it vanishes on the ideal $\mathcal{I}_e \triangleleft [\mathcal{T}(X)]_e$ given by

$$\mathcal{I}_e := \{f \in [\mathcal{T}(X)]_e \mid \lim_F \|f\|_F = 0\},$$

where the limit is taken with respect to the partial order induced by inclusion on finite sets of P . The universal C^* -algebra with respect to the strongly covariant representations of X is denoted by $A \times_X P$.

That is, $A \times_X P$ is the quotient $\mathcal{T}(X)/\mathcal{I}_\infty$ for the ideal $\mathcal{I}_\infty \triangleleft \mathcal{T}(X)$ of *strong covariance relations* generated by \mathcal{I}_e . One of the important points of Sehnm’s theory is that the inclusion $A \hookrightarrow A \times_X P$ is faithful. As a quotient by an induced ideal of $\mathcal{T}(X)$, the C^* -algebra $A \times_X P$ inherits the coaction of G . The following is the main theorem of [Sehnm 2019].

Theorem 3.10 [Sehnm 2019, Theorem 3.10]. *Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . Then a $*$ -homomorphism of $A \times_X P$ is faithful on A if and only if it is faithful on the fixed point algebra $[A \times_X P]_e$.*

Due to the grading $A \times_X P$ is the maximal C^* -algebra of a Fell bundle over G . We consider two reduced versions.

Definition 3.11. Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . We write $A \times_{X,\lambda} P$ for the reduced C^* -algebra of the Fell bundle in $A \times_X P$. If $q : \mathcal{T}(X) \rightarrow \mathcal{T}_\lambda(X)$ is the canonical $*$ -epimorphism, then we write $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ for the quotient of $\mathcal{T}_\lambda(X)$ by the ideal $q(\mathcal{I}_\infty)$.

Remark 3.12. The notation SCX is used in [Dor-On et al. 2022] to denote the G -Fell bundle inside $A \times_X^G P$. Therefore we have two ways of writing the related C^* -algebras in the sense that

$$A \times_X P = C^*(SCX) \quad \text{and} \quad A \times_{X,\lambda} P = C_\lambda^*(SCX).$$

Sehnm [2019, Lemma 3.9] shows that the strong covariance relations do not depend on the group embedding in the following sense. Suppose that P admits two group embeddings $i_G : P \rightarrow G$ and $i_H : P \rightarrow H$ and write $C_{\max}^*(SC_G X) = C^*(\pi^G, t^G)$ and $C_{\max}^*(SC_H X) = C^*(\pi^H, t^H)$. Then there exists a $*$ -isomorphism

$$C_{\max}^*(SC_G X) \rightarrow C_{\max}^*(SC_H X), \quad t_{i_G(p)}^G(\xi_{i_G(p)}) \mapsto t_{i_H(p)}^H(\xi_{i_H(p)}).$$

The $*$ -isomorphism between $C_{\max}^*(SC_G X)$ and $C_{\max}^*(SC_H X)$ descends to a $*$ -isomorphism that fixes X at the reduced level, as well, and thus $A \times_{X,\lambda} P$ does not depend on the group embedding either. Indeed suppose that G is the enveloping group of P and thus there exists a group homomorphism $\gamma : G \rightarrow H$ that is injective on P . We then have that there is a $*$ -homomorphism between the maximal C^* -algebras induced by the G -Fell bundle and the H -Fell bundle on Sehnm’s covariance algebra. Sehnm’s result [2019, Lemma 3.9] is that this $*$ -homomorphism is faithful. By Fell bundle theory we then get a canonical $*$ -epimorphism

$$C_\lambda^*(SC_G X) \rightarrow C_\lambda^*(SC_H X)$$

that fixes X . Hence by construction it intertwines the normal faithful conditional expectations. Their fixed point algebras are $*$ -isomorphic to the fixed point algebras in the maximal C^* -algebras and these are $*$ -isomorphic by [Sehnm 2019, Lemma 3.9]. Thus the $*$ -epimorphism on the reduced models is faithful.

We see that the representations Φ_F used to define the strong covariance relations are subrepresentations of $\bar{\delta}_{G,\lambda} : \mathcal{T}_\lambda(X) \rightarrow \mathcal{T}_\lambda(X) \otimes C_\lambda^*(G)$ for $\bar{\delta}_{G,\lambda} = (\text{id} \otimes \lambda)\bar{\delta}_G$, where $\bar{\delta}_G$ is the normal coaction on the Fock representation. Indeed we can identify

$$X_F^+ = \bigoplus_{\substack{g \in G \\ r \in P}} X_r I_{r^{-1}(r \vee g F)}$$

with a submodule of $\mathcal{F}X \otimes \ell^2(G)$ through the isometry given by

$$X_r I_{r^{-1}(r \vee g F)} \ni \eta_r \mapsto \eta_r \otimes \delta_g \in X_r \otimes \ell^2(G).$$

Recall here that $\mathcal{F}X \otimes \ell^2(G)$ is the exterior tensor product of two modules (seeing $\ell^2(G)$ as a module over \mathbb{C}), and there is a faithful $*$ -homomorphism

$$\mathcal{T}_\lambda(X) \otimes C_\lambda^*(G) \subseteq \mathcal{L}(\mathcal{F}X) \otimes \mathcal{B}(\ell^2(G)) \hookrightarrow \mathcal{L}(\mathcal{F}X \otimes \ell^2(G)).$$

We then see that

$$t_{F,p}(\xi_p) = (\bar{t}_p(\xi_p) \otimes \lambda_p)|_{X_F^+} = \bar{\delta}_{G,\lambda}(\bar{t}_p(\xi_p))|_{X_F^+} \quad \text{for all } p \in P,$$

and likewise for their adjoints. Thus X_F^+ is reducing under $\bar{\delta}_{G,\lambda}(\mathcal{T}_\lambda(X))$. Recall also that X_F is reducing for $[\mathcal{T}(X)]_e$ as the range of the projection $Q_{e,F}$ and so we obtain the representation

$$\bigoplus_{\text{fin } F \subseteq G} \Phi_F(\cdot)|_{X_F} : [\mathcal{T}(X)]_e \rightarrow [\mathcal{T}_\lambda(X)]_e \rightarrow \prod_{\text{fin } F \subseteq G} \mathcal{L}(X_F).$$

In particular, by definition we have for an $f \in \mathcal{T}(X)$ that

$$f \in \mathcal{I}_e \iff \bigoplus_{\text{fin } F \subseteq G} \Phi_F(f)|_{X_F} \in c_0(\mathcal{L}(X_F) \mid \text{fin } F \subseteq G).$$

By definition we then get that the diagram

$$\begin{array}{ccccc} [\mathcal{T}(X)]_e & \longrightarrow & [\mathcal{T}_\lambda(X)]_e & \longrightarrow & \prod_{\text{fin } F \subseteq G} \mathcal{L}(X_F) \\ \downarrow & & \downarrow q_{\text{sc}} & & \downarrow \\ [A \times_X P]_e & \longrightarrow & [q_{\text{sc}}(\mathcal{T}_\lambda(X))]_e & \longrightarrow & (\prod_{\text{fin } F \subseteq G} \mathcal{L}(X_F)) / (c_0(\mathcal{L}(X_F) \mid \text{fin } F \subseteq G)) \end{array}$$

is commutative. Consequently the e -graded $*$ -algebraic relations in $\mathcal{T}_\lambda(X)$ and $A \times_X P$ induce relations in $q_{\text{sc}}(\mathcal{T}_\lambda(X))$. In particular, since by [Sehnem 2019, Proposition 3.5] A is represented faithfully in the bottom right corner of the above diagram, we obtain the following corollary.

Corollary 3.13 [Dor-On et al. 2022, Corollary 5.5]. *Let P be a unital subsemigroup of a group G and X be a product system over P with coefficients in A . Then we have $A \hookrightarrow q_{\text{sc}}(\mathcal{T}_\lambda(X))$. Moreover, a $*$ -homomorphism of $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ is faithful on A if and only if it is faithful on $[q_{\text{sc}}(\mathcal{T}_\lambda(X))]_e$. Likewise for the reduced C^* -algebra $A \times_{X,\lambda} P$.*

4. Compactly aligned product systems over weak right LCM inclusions

Weak right LCM inclusions. A semigroup P is said to be a *right LCM semigroup* if it is left-cancellative and satisfies *Clifford’s condition* [Lawson 2012; Norling 2014]:

$$\text{for every } p, q \in P \text{ with } pP \cap qP \neq \emptyset \text{ there exists a } w \in P \text{ such that } pP \cap qP = wP.$$

In other words, if $p, q \in P$ have a right common multiple then they have a right least common multiple. As we always see a semigroup P inside a group G , it follows that P is by default cancellative, and we will refer to (G, P) simply as a *weak right LCM inclusion*. We use the adjective “weak” here to emphasize that we do not assume that the least common multiple property holds for *all* elements in G .

It is clear that w is a right LCM for p, q if and only if wx is a right LCM of p, q for every $x \in P^*$. A *weak quasilattice* (G, P) is a weak right LCM inclusion with $P \cap P^{-1} = \{e\}$, i.e., when least common multiples are unique (whenever they exist).

Definition 4.1. Let (G, P) be a right weak LCM inclusion. A finite set F is said to be \vee -closed if for any $p, q \in F$ with $pP \cap qP \neq \emptyset$ there exists a unique $w \in F$ such that $pP \cap qP = wP$.

Equivalently, a finite $F \subseteq P$ is \vee -closed if and only if the familiar relation

$$p \leq q \iff q^{-1}p \in P$$

defines a partial order on F . In particular, if F is \vee -closed, then $pP \neq qP$ for any $p, q \in F$ with $p \neq q$. Furthermore, any \vee -closed set admits maximal and minimal elements. Our terminology here regarding \vee -closed sets extends the familiar one from the case where (G, P) is a weak quasilattice order. There is an alternative way for describing \vee -closed sets in the context of weak right LCM inclusions. Given a finite subset $F \subseteq P$ we write

$$\mathcal{I}(F) := \{pP \mid p \in F\}$$

for the set of principal ideals defined by F . It then follows that F is \vee -closed if and only if $\mathcal{I}(F)$ is closed under intersections and the partial order defined on $\mathcal{I}(F)$ by set theoretic inclusion lifts to a partial order on F .

Let $F \subseteq P$ be a finite set so that $\mathcal{I}(F)$ is closed under intersections. From such a set F we can produce a \vee -closed subset F^\vee such that $\mathcal{I}(F) = \mathcal{I}(F^\vee)$ by choosing a minimal set of distinct representatives for the principal ideals. This process does not produce a unique F^\vee in general.

Nica-covariant representations. Following Fowler’s work [2002], Brownlowe, Larsen and Stammeier [Brownlowe et al. 2018] and Kwaśniewski and Larsen [2019] considered product systems of right LCM semigroups.

Definition 4.2. A product system X over a weak right LCM semigroup P with coefficients in A is called *compactly aligned* if for $p, q \in P$ with $pP \cap qP = wP$ we have that

$$i_p^w(S)i_q^w(T) \in \mathcal{K}X_w \quad \text{whenever } S \in \mathcal{K}X_p, \quad T \in \mathcal{K}X_q.$$

A note is in order for clarifying that this is independent of the choice of w . Recall that if w' is a right LCM of p, q then $w' = wx$ for some $x \in P^*$. Since $\mathcal{L}X_w \simeq \mathcal{L}X_{wx}$ we have that $i_p^w(S)i_q^w(T) \in \mathcal{K}X_w$ if and only if $i_p^{wx}(S)i_q^{wx}(T) = i_w^{wx}(i_p^w(S)i_q^w(T)) \in \mathcal{K}X_{wx}$ for all $x \in P^*$.

Definition 4.3. Let X be a compactly aligned product system over a right LCM semigroup P with coefficients in A . A *Nica-covariant representation* (π, t) is a Toeplitz representation of A that in addition satisfies the *Nica-covariance* condition: for $S \in \mathcal{K}X_p$ and $T \in \mathcal{K}X_q$ we have that

$$\psi_p(S)\psi_q(T) = \begin{cases} \psi_w(i_p^w(S)i_q^w(T)) & \text{if } pP \cap qP = wP, \\ 0 & \text{otherwise.} \end{cases}$$

The *Toeplitz–Nica–Pimsner algebra* $\mathcal{NT}(X)$ of X is the universal C^* -algebra generated by A and X with respect to the representations of X . The *Toeplitz–Nica–Pimsner tensor algebra* $\mathcal{NT}(X)^+$ of X is the subalgebra of $\mathcal{T}(X)$ generated by A and X .

Remark 4.4. As noted in [Dor-On et al. 2022], the definition of Nica-covariance requires that the right-hand side is independent of the choice of the least common multiple, i.e., if $pP \cap qP = wP$ and $x \in P^*$ then

$$\psi_w(i_p^w(S)i_q^w(T)) = \psi_{wx}(i_p^{wx}(S)i_q^{wx}(T)) \quad \text{for all } S \in \mathcal{KX}_p, \quad T \in \mathcal{KX}_q.$$

This is verified in [Dor-On et al. 2022, Proposition 2.4] (see Proposition 3.2 herein) and completes the definition of Nica-covariance in [Kwaśniewski and Larsen 2019].

Remark 4.5. By definition $\mathcal{NT}(X)$ is a quotient of $\mathcal{T}(X)$ by an ideal generated by a subspace of $[\mathcal{T}(X)]_e$. Even though $\mathcal{NT}(X) = \mathcal{T}(X)$ when $P = \mathbb{Z}_+$, this is not the case even when $P = \mathbb{Z}_+^n$. Dor-On and Katsoulis provide a counterexample to this effect in [Dor-On and Katsoulis 2020, Example 5.2]. The same example further shows that $\mathcal{T}(X)^+$ is not completely isometric to $\mathcal{NT}(X)^+$.

Under the assumption of compact alignment, one can check that the Fock representation is automatically Nica-covariant. Thus $\mathcal{NT}(X)$ is nontrivial. As $\mathcal{NT}(X)$ is a quotient of $\mathcal{T}(X)$ by an induced ideal, by [Carlsen et al. 2011, Proposition A.1] the nondegenerate and faithful coaction of $\mathcal{T}(X)$ descends canonically to one on $\mathcal{NT}(X)$. Alternatively one may use the arguments of the proof of Proposition 3.4 for the Toeplitz–Nica–Pimsner tensor algebra to deduce the following.

Proposition 4.6. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Suppose that $(\hat{\pi}, \hat{t})$ is a faithful representation of $\mathcal{NT}(X)$. Then the canonical $*$ -homomorphism*

$$\hat{\delta} : \mathcal{NT}(X) \rightarrow \mathcal{NT}(X) \otimes C^*(G), \quad \hat{t}(\xi_p) \mapsto \hat{t}(\xi_p) \otimes u_p$$

defines a coaction of G on $\mathcal{NT}(X)$.

We have refrained from describing the spectral spaces for the coaction on $\mathcal{NT}(X)$ because of the following additional property of Nica-covariant representations. Let (π, t) be a Nica-covariant representation of X . We compute

$$t_p(X_p)^* t_p(X_p) \cdot t_p(\xi_p)^* t_q(\xi_q) \cdot t_q(X_q)^* t_q(X_q) \subseteq [t_p(X_p)^* \psi_p(\mathcal{KX}_p) \psi_q(\mathcal{KX}_q) t_q(X_q)].$$

Next take a limit by contractive approximate identities in $[t_p(X_p)^* t_p(X_p)]$ and in $[t_q(X_q)^* t_q(X_q)]$, and derive that

$$t_p(\xi_p)^* t_q(\xi_q) \in [t_{p'}(X_{p'})^* t_{q'}(X_{q'})^*] \quad \text{for } wP = pP \cap qP, \quad p' = p^{-1}w, \quad q' = q^{-1}w,$$

and

$$t_p(\xi_p)^* t_q(\xi_q) = 0 \quad \text{for } pP \cap qP = \emptyset.$$

Hence the C^* -algebra $C^*(\pi, t)$ generated by $\pi(A)$ and $t_p(X_p)$ admits a *Wick ordering* in the sense that

$$C^*(\pi, t) = \overline{\text{span}}\{t_p(\xi_p)t_q(\xi_q)^* \mid \xi_p \in X_p, \xi_q \in X_q \text{ and } p, q \in P\}.$$

In particular if $\mathcal{NT}(X) = C^*(\hat{\pi}, \hat{t})$ then the spectral spaces that only matter are of the form

$$\mathcal{NT}(X)_{pq^{-1}} = \overline{\text{span}}\{\hat{t}_p(\xi_p)\hat{t}_q(\xi_q)^* \mid \xi_p \in X_p, \xi_q \in X_q\},$$

that is, only for $g \in G$ of the form $g = pq^{-1}$ for some $p, q \in P$.

The following proposition gives a direct criterion to check compact alignment.

Proposition 4.7. *Let (G, P) be a weak right LCM inclusion and let $X = \{X_p\}_{p \in P}$ be a product system over $X_e = A$. Let (π, t) be an injective representation of X . Then X is compactly aligned, if and only if for all $p, q \in P$ we have that*

$$t_p(X_p)^*t_q(X_q) \subseteq [t_{p^{-1}w}(X_{p^{-1}w})t_{q^{-1}w}(X_{q^{-1}w})^*] \quad \text{for } wP = pP \cap qP,$$

if and only if for all $p, q \in P$ we have that

$$t_p(X_p)t_p(X_p)^*t_q(X_q)t_q(X_q)^* \subseteq [t_w(X_w)t_w(X_w)^*] = \psi_w(\mathcal{K}X_w) \quad \text{for } wp = pP \cap qP,$$

with the understanding that the left-hand sides are the zero space when p and q have no right common multiple.

Proof. The first equivalence follows in the same way as [Katsoulis 2020, Proposition 3.2] and it is omitted. By using that $X_p X_p^* X_p$ is dense in X_p for every $p \in P$, we get the second equivalence. \square

Let us now pass to the analysis of the cores of a Nica-covariant representation (π, t) of X . For a finite $F \subseteq P$ that is \vee -closed we write

$$B_F := \text{span}\{\psi_p(k_p) \mid k_p \in \mathcal{K}X_p, p \in F\}.$$

Since F is \vee -closed, Nica-covariance implies that B_F is a $*$ -subalgebra of $C^*(\pi, t)$. In [Dor-On et al. 2022, Proposition 2.10] we show that every B_F is actually a C^* -subalgebra. Moreover for such an F we write

$$B_{F \cdot P} := \overline{\text{span}}\{\psi_q(k_q) \mid k_q \in \mathcal{K}X_q, q \in F \cdot P\}.$$

Likewise this is also a (closed) $*$ -subalgebra. Finally we write

$$B_{P \setminus \{e\}} := \overline{\text{span}}\{\psi_p(k_p) \mid k_p \in \mathcal{K}X_p, e \neq p \in P\} \quad \text{and} \quad B_P := \pi(A) + B_{P \setminus \{e\}}.$$

We see that $B_{P \setminus \{e\}}$ is an ideal in B_P and thus the sum $\pi(A) + B_{P \setminus \{e\}}$ is indeed closed. We refer to these sets as the *cores* of the representation (π, t) . In [Dor-On et al. 2022, Proposition 2.11] we showed that we can exhaust the cores by using finite \vee -closed sets, in the sense that

$$B_P = \overline{\bigcup\{B_F \mid F \subseteq P \text{ finite and } \vee\text{-closed}\}}.$$

We denote by \bar{B}_F the cores of $\mathcal{T}_\lambda(X) = C^*(\bar{\pi}, \bar{t})$. Recall that $\mathcal{T}_\lambda(X)$ admits the faithful conditional expectation

$$\bar{E}_P : \mathcal{T}_\lambda(X) \rightarrow \bar{B}_P, \quad \bar{t}_p(\xi_p)\bar{t}_q(\xi_q)^* = \delta_{p,q}\bar{t}_p(\xi_p)\bar{t}_q(\xi_q)$$

given by the sum of compressions to the (r, r) -entries in $\mathcal{L}(\mathcal{F}X)$ (see Proposition 3.8).

The Toeplitz–Nica–Pimsner algebra is modeled after the Fock algebra in this context. A compactly aligned product system X over P with coefficients in A is called *amenable* if the Fock representation is faithful on $\mathcal{NT}(X)$. Let us give some equivalent conditions for this to happen.

Theorem 4.8. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . The following are equivalent:*

- (i) *The coaction of G on $\mathcal{NT}(X)$ is normal.*
- (ii) *The conditional expectation on $\mathcal{NT}(X)$ is faithful.*
- (iii) *The Fock representation is faithful on $\mathcal{NT}(X)$.*
- (iv) *The representation*

$$\mathcal{NT}(X) \rightarrow C^*(\pi, t) \otimes C_\lambda^*(P), \quad \tilde{t}_p(\xi_p) \mapsto t_p(\xi_p) \otimes V_p$$

is faithful for any injective Nica-covariant pair (π, t) .

Proof. By the universal property there exists a canonical $*$ -representation

$$\mathcal{NT}(X) \rightarrow \mathcal{NT}(X) \otimes C_\lambda^*(G)$$

that intertwines the conditional expectations. Thus items (i) and (ii) are equivalent. For the same reason items (ii) and (iii) are equivalent.

Assuming item (iii) we have to show that the representation

$$\mathcal{NT}(X) \rightarrow C^*(\pi, t) \otimes C_\lambda^*(P)$$

is faithful on the fixed point algebra. It suffices to show injectivity on the F -boxes for arbitrary \vee -closed $F \subseteq P$. To this end suppose that

$$\sum_{p \in F} \psi_p(k_p) = 0$$

for some $k_p \in \mathcal{K}X_p$ and let p_0 be minimal so that $k_{p_0} \neq 0$. Injectivity of π then implies that $\psi_{p_0}(k_{p_0}) \neq 0$ as well. However, if $Q_{p_0} : \ell^2(P) \rightarrow \mathbb{C}e_{p_0}$ is the canonical projection, minimality of p_0 yields

$$\psi_{p_0}(k_{p_0}) = I \otimes Q_{p_0} \left(\sum_{p \in F} \psi_p(k_p) \right) I \otimes Q_{p_0} = 0,$$

which is a contradiction. This shows that item (iii) implies item (iv).

Since the $*$ -representation

$$\mathcal{NT}(X) \rightarrow C^*(\pi, t) \otimes C_\lambda^*(P)$$

intertwines the conditional expectations, we finally have that item (iv) implies item (i). □

On the other hand strongly covariant representations are Nica-covariant (which is expected as Nica-covariance is an e -graded relation in $[\mathcal{T}_\lambda(X)]_e$). It is proven by Sehnem [2019, Proposition 4.2] for quasilattices, but the same proof passes to right LCM semigroups as well [Dor-On et al. 2022, Proposition 5.4]. Hence $A \times_X P$ is a quotient of $\mathcal{NT}(X)$.

Proposition 4.9 [Dor-On et al. 2022, Proposition 5.4; Sehne 2019, Proposition 4.2]. *Let X be a compactly aligned product system over a right LCM semigroup P with coefficients in A . Let*

$$\psi_{F,p} : \mathcal{K}X_p \rightarrow \mathcal{L}(X_F^+)$$

be the induced representations from $(\pi_F, t_{F,p})$. A representation (π, t) of X is strongly covariant if and only if it is Nica-covariant and it satisfies

$$\sum_{p \in F} \psi_{F,p}(k_p)|_{X_F} = 0 \implies \sum_{p \in F} \psi_p(k_p) = 0$$

for any finite $F \subseteq P$ and $k_p \in \mathcal{K}X_p$.

Carlsen, Larsen, Sims and Vittadello [Carlsen et al. 2011] explored the idea of finding the couniversal C^* -algebra with respect to injective equivariant Nica-covariant representations of X . By using the C^* -envelope machinery we can prove that this object always exists, thus completing the couniversal aspect of their program in the more general context of right weak LCM inclusions.

Definition 4.10. Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . We say that a representation (π, t) of X is *couniversal for $\mathcal{NT}(X)$* if

- (i) π is faithful,
- (ii) $C^*(\pi, t)$ is an equivariant quotient of $\mathcal{NT}(X)$,
- (iii) (π, t) factors through any other equivariant quotient of $\mathcal{NT}(X)$ that is injective on A .

Of course the C^* -algebras of couniversal representations are automatically $*$ -isomorphic by an equivariant homomorphism. In [Dor-On et al. 2022] we proved that the equivariant representation

$$\mathcal{NT}(X) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$$

that is given by the diagram

$$\begin{array}{ccc} \mathcal{NT}(X) & \xrightarrow{\dots\dots\dots} & C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \\ & \searrow & \nearrow \\ & \mathcal{T}_\lambda(X) & \end{array}$$

is couniversal. Let us review the main arguments and see what more we can obtain.

Proposition 4.11 [Dor-On et al. 2022, Proposition 4.4]. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Let $\Phi : \mathcal{T}_\lambda(X) \rightarrow B$ be a $*$ -representation such that $\Phi|_{\bar{\pi}(A)}$ is faithful. Then there exists a faithful $*$ -homomorphism*

$$\mathcal{T}_\lambda(X) \rightarrow B \otimes C_\lambda^*(P), \quad \bar{t}_p(\xi_p) \mapsto \Phi \bar{t}_p(\xi_p) \otimes V_p.$$

As a consequence the injective equivariant representations on product systems generate C^* -covers for the cosystem $(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$.

Proposition 4.12 [Dor-On et al. 2022, Proposition 4.5]. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Let $\Phi : \mathcal{T}_\lambda(X) \rightarrow B$ be an equivariant *-epimorphism such that $\Phi|_{\overline{\pi(A)}}$ is faithful. Then B is a C^* -cover for the cosystem $(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$.*

Another consequence of Proposition 4.11 provides a generalization of the extension theorem of [Katsoulis and Ramsey 2019]. It essentially allows us to recognize a Fock tensor algebra by the presence of a coaction.

Theorem 4.13 (extension theorem). *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Let $\Phi : \mathcal{T}_\lambda(X) \rightarrow B$ be a representation of X and set*

$$\mathcal{A} := \overline{\text{alg}}\{\Phi\bar{\pi}(A), \Phi\bar{t}_p(X_p) \mid p \in P\}.$$

Then the following are equivalent:

- (i) $\Phi|_{\mathcal{T}_\lambda(X)^+}$ is completely isometric.
- (ii) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C^*(G), \quad \Phi\bar{t}_p(\xi_p) \mapsto \Phi\bar{t}_p(\xi_p) \otimes u_p.$$


- (iii) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C_\lambda^*(G), \quad \Phi\bar{t}_p(\xi_p) \mapsto \Phi\bar{t}_p(\xi_p) \otimes \lambda_p.$$

- (iv) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C_\lambda^*(P), \quad \Phi\bar{t}_p(\xi_p) \mapsto \Phi\bar{t}_p(\xi_p) \otimes V_p.$$

Proof. Below we have a diagram of completely contractive representations induced by Propositions 4.11 and 4.12, which are completely positive maps fixing the nonselfadjoint part. If any of the items holds then it makes the representation of $\mathcal{T}_\lambda(X)^+$ to \mathcal{A} completely isometric and the proof is complete.



□

We now come to the last part of [Dor-On et al. 2022] that connects reduced C^* -algebras with the C^* -envelope. By Corollary 3.13 and Proposition 4.12 we get a canonical $*$ -epimorphism

$$q_{\text{sc}}(\overline{\mathcal{T}}_\lambda(X)) \rightarrow A \times_{X,\lambda} P \simeq C_{\text{env}}^*(\overline{\mathcal{T}}_\lambda(X)^+, G, \bar{\delta}_G).$$

We remind the reader of the notation used here and in [Dor-On et al. 2022] as explained in Remark 3.12. The same remark asserts that the C^* -envelope of the cosystem is independent of the group embedding in this setting.

Theorem 4.14 [Dor-On et al. 2022, Theorem 4.9, Theorem 5.3 and Corollary 5.6]. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Then the equivariant $*$ -epimorphism*

$$\mathcal{NT}(X) \rightarrow C_{\text{env}}^*(\overline{\mathcal{T}}_\lambda(X)^+, G, \bar{\delta}_G)$$

is couniversal. Moreover we have an equivariant $$ -isomorphism*

$$C_{\text{env}}^*(\overline{\mathcal{T}}_\lambda(X)^+, G, \bar{\delta}_G) \simeq A \times_{X,\lambda} P.$$

The equivariant $$ -epimorphism*

$$q_{\text{sc}}(\overline{\mathcal{T}}_\lambda(X)) \rightarrow A \times_{X,\lambda} P \simeq C_{\text{env}}^*(\overline{\mathcal{T}}_\lambda(X)^+, G, \bar{\delta}_G)$$

is faithful if and only if the coaction of G on $q_{\text{sc}}(\overline{\mathcal{T}}_\lambda(X))$ is normal.

5. Controlled maps

Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a semigroup preserving homomorphism between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . By Proposition 2.9 the Toeplitz algebra admits a \mathcal{G} -grading that contains the G -grading, and the same is true for the fixed point algebras. Of course this may be useless; for example the ϑ -fixed point algebra for the map $\vartheta : G \rightarrow \{e\}$ is the entire C^* -algebra. Nevertheless more can be obtained for weak right LCM inclusions as long as we impose axioms that control the map. The following extends the controlled maps on quasilattice ordered groups from [Laca and Raeburn 1996], see also [Fowler 2002] and [Crisp and Laca 2007], to the context of weak right LCM inclusions.

Definition 5.1. A controlled map $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ between weak right LCM inclusions is a semigroup preserving homomorphism such that:

- (A1) If $pP \cap qP \neq \emptyset$, then $\vartheta(p)\mathcal{P} \cap \vartheta(q)\mathcal{P} = \vartheta(pP \cap qP)\mathcal{P}$.
- (A2) If $pP \cap qP \neq \emptyset$ and $\vartheta(p) = \vartheta(q)$, then $p = q$.

It is worth pointing out that in the case where $P = G$ then there is only one right ideal (generated by the identity). Therefore a controlled map in this case is simply an injective group homomorphism due to (A2).

Remark 5.2. It is clear that having $\vartheta(p)\mathcal{P} \cap \vartheta(q)\mathcal{P} = \vartheta(w)\mathcal{P}$ whenever $pP \cap qP = wP$ is equivalent to (A1). Moreover, because of (A2) we have that $\vartheta^{-1}(e_G) \cap P = \{e_G\}$. Indeed as ϑ is a group homomorphism we have that $\vartheta(e_G) = e_G$. Now if $\vartheta(p) = e_G$ for some $p \in P$, then since $pP \cap e_G P = pP \neq \emptyset$ we get by (A1) that $p = e_G$. This extra generality is crucial when we wish to consider the generalized length function given by abelianization on the free monoid \mathbb{F}_n^+ [Laca and Raeburn 1996], and, more generally, the Artin monoids of rectangular type [Crisp and Laca 2002].

Remark 5.3. Similar types of maps appear in [Crisp 1999] and [Brownlowe et al. 2018, Section 3]. However the maps therein satisfy the stronger requirement that $\vartheta(p)\mathcal{P} \cap \vartheta(q)\mathcal{P} = \vartheta(pP \cap qP)\mathcal{P}$ for all $p, q \in P$. This means that $p, q \in P$ have a right LCM if and only if $\vartheta(p), \vartheta(q) \in \mathcal{P}$ also do. In our Definition 5.1, the condition (A1) allows the possibility that $\vartheta(p), \vartheta(q)$ have a right LCM in \mathcal{P} even when $pP \cap qP = \emptyset$.

We will investigate the impact of the existence of a controlled map on Nica-covariant representations. Henceforth fix a controlled map $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ between two weak right LCM inclusions. Suppose that (π, t) is a Nica-covariant representation of a compactly aligned product system X over P with coefficients in A . If $p, q \in P$ with $\vartheta(p) = \vartheta(q)$ then by (A2) either $p = q$ or $pP \cap qP = \emptyset$; thus Nica-covariance yields the orthogonality

$$t_p(\xi_p)^* t_q(\xi_q) = \delta_{p,q} \pi(\langle \xi_p, \xi_q \rangle).$$

Hence the C^* -algebra

$$B_{\vartheta^{-1}(h)} := \overline{\text{span}}\{\psi_{p,q}(k_{p,q}) \mid k_{p,q} \in \mathcal{K}(X_q, X_p), \vartheta(p) = h = \vartheta(q)\}$$

is a matrix C^* -algebra. For a \vee -closed $\mathcal{F} \subseteq \mathcal{P}$ we define

$$B_{\vartheta^{-1}(\mathcal{F})} := \overline{\text{span}}\{B_{\vartheta^{-1}(h)} \mid h \in \mathcal{F}\}.$$

By conditions (A1) and (A2) of Definition 5.1 we get that $\vartheta^{-1}(\mathcal{F})$ is also \vee -closed (and thus the above space is a C^* -algebra). Therefore every $B_{\vartheta^{-1}(\mathcal{F})}$ is the inductive limit of the matrix C^* -subalgebras

$$\text{span}\{\psi_{p,q}(k_{p,q}) \mid k_{p,q} \in \mathcal{K}(X_q, X_p), p, q \in F, \vartheta(p) = \vartheta(q)\} \quad \text{for finite } \vee\text{-closed } F \subseteq \vartheta^{-1}(\mathcal{F}).$$

Taking the closure of the union we obtain the ϑ -fixed point algebra

$$B_{\vartheta^{-1}(\mathcal{P})} := \overline{\text{span}}\{\psi_{p,q}(k_{p,q}) \mid k_{p,q} \in \mathcal{K}(X_q, X_p), \vartheta(p) = \vartheta(q)\}.$$

It follows that

$$B_P = \overline{\text{span}}\{\psi_p(\mathcal{K}X_p) \mid p \in P\} \subseteq B_{\vartheta^{-1}(\mathcal{P})}.$$

It is clear that the faithful conditional expectation $\bar{E}_{\mathcal{P}}$ on $C^*(\bar{\pi}, \bar{t}) = \mathcal{T}_{\lambda}(X)$ described in Proposition 3.8 is onto $\bar{B}_{\vartheta^{-1}(\mathcal{P})}$. We already commented on the effect of semigroup preserving homomorphisms of the form $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ on $\mathcal{T}(X)$ and $\mathcal{T}_{\lambda}(X)$. We give some basic facts about the effect of controlled maps on $\mathcal{T}_{\lambda}(X)$.

Proposition 5.4. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Let $\Phi : \mathcal{T}_\lambda(X) \rightarrow B$ be a $*$ -representation such that $\Phi|_{\overline{\pi(A)}}$ is faithful. Then there exists a faithful $*$ -homomorphism*

$$\mathcal{T}_\lambda(X) \rightarrow B \otimes C_\lambda^*(\mathcal{P}), \quad \bar{t}_p(\xi_p) \mapsto \Phi \bar{t}_p(\xi_p) \otimes V_{\vartheta(p)}.$$

Proof. The proof follows the same lines as Proposition 4.11 with the observation that $\bar{B}_{\vartheta^{-1}(h)}$ for $h \in \mathcal{G}$ is a matrix algebra. □

As an immediate consequence we have the following corollary which extends Theorem 4.13 to the controlled setting.

Corollary 5.5. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Let $\Phi : \mathcal{T}_\lambda(X) \rightarrow B$ be a $*$ -representation and set*

$$\mathcal{A} := \overline{\text{alg}}\{\Phi \bar{\pi}(A), \Phi \bar{t}_p(X_p) \mid p \in P\}.$$

Then the following are equivalent:

- (i) $\Phi|_{\mathcal{T}_\lambda(X)^+}$ is completely isometric.
- (ii) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C^*(\mathcal{G}), \quad \Phi \bar{t}_p(\xi_p) \mapsto \Phi \bar{t}_p(\xi_p) \otimes u_{\vartheta(p)}.$$

- (iii) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C_\lambda^*(\mathcal{G}), \quad \Phi \bar{t}_p(\xi_p) \mapsto \Phi \bar{t}_p(\xi_p) \otimes \lambda_{\vartheta(p)}.$$

- (iv) There exists a completely contractive map

$$\mathcal{A} \rightarrow B \otimes C_\lambda^*(\mathcal{P}), \quad \Phi \bar{t}_p(\xi_p) \mapsto \Phi \bar{t}_p(\xi_p) \otimes V_{\vartheta(p)}.$$

Proof. The proof follows as in Theorem 4.13, modulo Proposition 3.7 and Proposition 5.4. □

Controlled elimination. We will require the following lemma for solving polynomial equations in the ϑ -fixed point algebra.

Lemma 5.6. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Let (π, t) be an injective Nica-covariant representation of X .*

- (i) Let p, q be distinct in $\vartheta^{-1}(h)$. For $r, s \in \vartheta^{-1}(h)$ with $(r, s) \neq (p, q)$, we get

$$t_p(X_p)^* \psi_{r,s}(k_{r,s}) t_q(X_q) = (0) \quad \text{for all } k_{r,s} \in \mathcal{K}(X_s, X_r).$$

- (ii) Let $\mathcal{F} \subseteq \mathcal{P}$ be \vee -closed and $F \subseteq \vartheta^{-1}(\mathcal{F})$ be finite and \vee -closed. Let $(r, s) \in F \times F$ with $\vartheta(r) = \vartheta(s)$ and $k_{r,s} \in \mathcal{K}(X_s, X_r)$ such that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s)}} \psi_{r,s}(k_{r,s}) = 0,$$

and suppose that $h \neq e_G$ is minimal in \mathcal{F} so that $k_{p,q} \neq 0$ for distinct $p, q \in \vartheta^{-1}(h)$. Then there exists a \vee -closed $\mathcal{F}' \subseteq \mathcal{P}$ and a finite \vee -closed $F' \subseteq \vartheta^{-1}(\mathcal{F}')$ with $e_G \notin \mathcal{F}'$ and $|\mathcal{F}'| \leq |\mathcal{F}| - 1$ such that

$$t_p(X_p)^* \psi_{p,q}(k_{p,q}) t_q(X_q) \subseteq B_{F'}.$$

Proof. (i) First we note that condition (A2) of Definition 5.1 yields $pP \cap qP = \emptyset$. By Nica-covariance we have that $t_p(X_p)^* \psi_{r,s}(k_{r,s}) t_q(X_q) = (0)$, unless

$$\exists w, z, v \in P \quad \text{such that} \quad pP \cap rP = wP, \quad qP \cap sP = zP \quad \text{and} \quad r^{-1}wP \cap s^{-1}zP = vP. \quad (5-1)$$

If $(r, s) \neq (p, q)$ and $\vartheta(r) = h = \vartheta(s)$, then condition (A2) of Definition 5.1 implies that $pP \cap rP = \emptyset$ or $qP \cap sP = \emptyset$, in which case

$$t_p(X_p)^* \psi_{r,s}(k_{r,s}) t_q(X_q) = (0).$$

(ii) Minimality of h in \mathcal{F} forces minimality of p, q in F . If (5-1) holds, then Nica-covariance yields

$$t_p(\xi_p)^* \psi_{r,s}(k_{r,s}) t_q(\xi_q) \in \psi_{p^{-1}rv, q^{-1}sv}(\mathcal{K}(X_{q^{-1}sv}, X_{p^{-1}rv})),$$

otherwise the product is zero. If $r = s$ and v exists then there are $p', q', x, x', y, y' \in P$ such that

$$pp' = rx, \quad qq' = ry \quad \text{and} \quad xx' = yy'.$$

But then

$$pp'x' = rxx' = ryy' = qq'y,$$

giving the contradiction that $pP \cap qP \neq \emptyset$. Hence in this case the product is zero. We will show that the product is zero also when $\vartheta(p^{-1}rv) = e_G = \vartheta(q^{-1}sv)$ for $r \neq s$ unless $(r, s) = (p, q)$. If $\vartheta(p^{-1}rv) = e_G$ then condition (A2) of Definition 5.1 yields $p \in rP$. Likewise $q \in sP$. Minimality of p, q in F forces either $(p, q) = (r, s)$ or $k_{r,s} = 0$. Set

$$\mathcal{F}' := \{h^{-1}g \mid g \in \mathcal{F}, g > h\} \quad \text{and} \quad F' := \{u^{-1}v \mid u, v \in F, \vartheta(u) = h, u > v\} \subseteq \vartheta^{-1}(\mathcal{F}').$$

We see that \mathcal{F}' is \vee -closed with $|\mathcal{F}'| \leq |\mathcal{F}| - 1$ and so F' is \vee -closed with

$$|F'| \leq |F \setminus \{p, q\}| = |F| - 2.$$

Moreover we see that $p^{-1}rv, q^{-1}sv \in F'$ whenever v exists. Hence for every $\xi_p \in X_p$ and $\xi_q \in X_q$ there are suitable $k'_{r',s'}$ with nontrivial $r', s' \in F'$ so that

$$0 = \sum_{r,s} t_p(\xi_p)^* \psi_{r,s}(k_{r,s}) t_q(\xi_q) = t_p(\xi_p)^* \psi_{p,q}(k_{p,q}) t_q(\xi_q) + \sum_{r',s'} \psi_{r',s'}(k'_{r',s'}), \quad \square$$

In the next proposition we show that we can eliminate elements of the form $\psi_{r,s}(k_{r,s})$ for $r \neq s$ with $\vartheta(r) = \vartheta(s)$, from a polynomial equation in the ϑ -fixed point algebra. Such arguments for the left-regular representation appear in [Dinh 1991, Proposition 2.10] and [Laca and Raeburn 1996, Lemma 4.1] for semigroups over quasilattices, i.e., when $X_p = \mathbb{C}$ for every $p \in P$ and (G, P) is a quasilattice. Here we need to move in three directions: (a) beyond one-dimensional fibers, (b) beyond quasilattices, and (c) beyond just the left-regular representation. A step towards this direction is done in [Kakariadis 2020] for quasilattices that are controlled by $(\mathbb{Z}^n, \mathbb{Z}_+^n)$, and here we expand further on this approach.

Proposition 5.7. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Let (π, t) and (π', t') be injective Nica-covariant representations such that there exists a canonical $*$ -epimorphism*

$$\Phi : C^*(\pi', t') \rightarrow C^*(\pi, t), \quad \text{with } \Phi(\pi'(a)) = \pi(a), \quad \Phi(t'_p(\xi_p)) = t_p(\xi_p).$$

Then Φ is injective on B'_p if and only if it is injective on $B'_{\vartheta^{-1}(\mathcal{P})}$.

Proof. As $B'_p \subseteq B'_{\vartheta^{-1}(\mathcal{P})}$ we need to show just one direction. To this end suppose that Φ is injective on the C^* -subalgebras of the form

$$B'_F = \overline{\text{span}}\{\psi'_p(\mathcal{K}X_p) \mid p \in F\}$$

for every finite \vee -closed $F \subseteq P$. We will show that Φ is injective on every

$$B'_{\vartheta^{-1}(\mathcal{F})} = \overline{\text{span}}\{\psi'_{r,s}(k_{r,s}) \mid \vartheta(r) = \vartheta(s) \in \mathcal{F}\}$$

for all \vee -closed $\mathcal{F} \subseteq \mathcal{P}$. Our strategy is to show the implication

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) \in \mathcal{F}}} \psi'_{r,s}(k_{r,s}) \in \ker \Phi \quad \implies \quad k_{r,s} = 0 \quad \text{whenever } r \neq s$$

for every finite \vee -closed $F \subseteq \vartheta^{-1}(\mathcal{F})$. Then injectivity of Φ in the smaller cores yields

$$\sum_{r \in F} \psi_r(k_r) = \sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) \in \mathcal{F}}} \psi_{r,s}(k_{r,s}) = 0 \quad \implies \quad \sum_{r \in F} \psi'_r(k_r) = 0,$$

and so

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) \in \mathcal{F}}} \psi'_{r,s}(k_{r,s}) = \sum_{r \in F} \psi'_r(k_r) = 0.$$

Since F is arbitrary this proves injectivity of Φ on $B'_{\vartheta^{-1}(\mathcal{F})}$. We proceed by induction on the size of \mathcal{F} .

Case 1. Assume that $\mathcal{F} = \{h\}$ and let F be a finite \vee -closed subset of $\vartheta^{-1}(\mathcal{F})$. Suppose that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) = h}} \psi_{r,s}(k_{r,s}) = 0,$$

and fix $p, q \in \vartheta^{-1}(h)$. Then condition (A2) of Definition 5.1 implies that

$$\psi_p(\mathcal{K}X_p)\psi_{p,q}(k_{p,q})\psi_q(\mathcal{K}X_q) = \psi_p(\mathcal{K}X_p) \left(\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) = h}} \psi_{r,s}(k_{r,s}) \right) \psi_q(\mathcal{K}X_q) = (0).$$

Using an approximate identity on both sides gives that $\psi_{p,q}(k_{p,q}) = 0$, and the injectivity of ψ implies that $k_{p,q} = 0$. As (p, q) was arbitrary we have that $k_{r,s} = 0$ for all $r, s \in \vartheta^{-1}(h)$ and so

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) = h}} \psi'_{r,s}(k_{r,s}) = 0.$$

Hence Φ is injective on $B'_{\vartheta^{-1}(\mathcal{F})}$ whenever $|\mathcal{F}| = 1$.

Case 2. Assume that $\mathcal{F} = \{e_G, h\}$ and let F be a finite \vee -closed subset of $\vartheta^{-1}(\mathcal{F})$. Suppose that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s)}} \psi_{r,s}(k_{r,s}) = 0.$$

By condition (A1) of Definition 5.1 we have that if $p \neq q$ with $\vartheta(p) = \vartheta(q) \in \mathcal{F}$ then $p, q \in \vartheta^{-1}(h)$. As before and by using item (i) of Lemma 5.6 on p, q we get that

$$\psi_p(\mathcal{K}X_p)\psi_{p,q}(k_{p,q})\psi_q(\mathcal{K}X_q) = \psi_p(\mathcal{K}X_p) \left(\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s)}} \psi_{r,s}(k_{r,s}) \right) \psi_q(\mathcal{K}X_q) = (0).$$

Using an approximate identity eventually gives that $k_{p,q} = 0$ whenever $p \neq q$. Hence $k_{r,s} = 0$ whenever $r \neq s$ in F and injectivity of Φ on B'_F gives that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s)}} \psi'_{r,s}(k_{r,s}) = 0.$$

Hence Φ is injective on $B'_{\vartheta^{-1}(\mathcal{F})}$ whenever $\mathcal{F} = \{e, h\}$.

Case 3. Assume that $\mathcal{F} = \{h_1, h_2\}$ and let F be a finite \vee -closed subset of $\vartheta^{-1}(\mathcal{F})$. Suppose that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) \in \{h_1, h_2\}}} \psi'_{r,s}(k_{r,s}) \in \ker \Phi.$$

Without loss of generality assume that it is written with the understanding that for every $\psi'_{r,s}(k_{r,s})$ we have that either $\psi'_{r,s}(k_{r,s}) = 0$ or that

$$\psi'_{r,s}(k_{r,s}) \notin B'_{\vartheta^{-1}(\vartheta(r)\mathcal{P})}.$$

Choose $h \in \mathcal{F}$ to be minimal such that $\psi'_{p,q}(k_{p,q}) \neq 0$ for distinct $p, q \in \vartheta^{-1}(h)$. Hence $k_{p,q} \neq 0$ and so

$$0 \neq \psi'_{p,q}(k_{p,q}) \notin B'_{\vartheta^{-1}(h\mathcal{P})}.$$

By using Lemma 5.6 item (ii) we have that

$$t_p(X_p)^* \psi_{p,q}(k_{p,q}) t_q(X_q) \subseteq B_{\vartheta^{-1}(\mathcal{F}')} \quad \text{for } |\mathcal{F}'| \leq 1,$$

with $e_G \notin \mathcal{F}'$. By using injectivity of Case 2 we then derive that

$$\psi'_p(\mathcal{K}X_p)\psi'_{p,q}(k_{p,q})\psi'_q(\mathcal{K}X_q) \subseteq B'_{\vartheta^{-1}(h\mathcal{P})}.$$

By using approximate identities on both sides we get the contradiction

$$\psi'_{p,q}(k_{p,q}) \in B'_{\vartheta^{-1}(h\mathcal{P})}.$$

Hence Φ is injective on $B'_{\vartheta^{-1}(\mathcal{F})}$ whenever $|\mathcal{F}| \leq 2$.

Case 4. Let $\mathcal{F} \subseteq \mathcal{P}$ be \vee -closed with $|\mathcal{F}| = n + 1$ and assume that Φ is injective on $B'_{\vartheta^{-1}(\mathcal{F}')}$ for all $\mathcal{F}' \subseteq \mathcal{P}$ with $|\mathcal{F}'| \leq n$. We will show that Φ is injective on $B'_{\vartheta^{-1}(\mathcal{F})}$. To this end let F be a finite \vee -closed subset of $\vartheta^{-1}(\mathcal{F})$ and suppose that

$$\sum_{\substack{r,s \in F \\ \vartheta(r) = \vartheta(s) \in \mathcal{F}}} \psi'_{r,s}(k_{r,s}) \in \ker \Phi,$$

with the understanding that for every $\psi'_{r,s}(k_{r,s})$ we have that either $\psi'_{r,s}(k_{r,s}) = 0$ or that

$$\psi'_{r,s}(k_{r,s}) \notin B'_{\vartheta^{-1}(\vartheta(r)\mathcal{P})}.$$

Choose $h \in \mathcal{F}$ to be minimal such that $\psi'_{p,q}(k_{p,q}) \neq 0$ for distinct $p, q \in \vartheta^{-1}(h)$. Hence $k_{p,q} \neq 0$ and so

$$0 \neq \psi'_{p,q}(k_{p,q}) \notin B'_{\vartheta^{-1}(h\mathcal{P})}.$$

By using Lemma 5.6 item (ii) we then have that

$$t_p(X_p)^* \psi'_{p,q}(k_{p,q}) t_q(X_q) \subseteq B_{\vartheta^{-1}(\mathcal{F}')} \quad \text{for } |\mathcal{F}'| \leq |\mathcal{F}| - 1 = n.$$

Using the induction hypothesis we then derive that

$$\psi'_p(\mathcal{K}X_p) \psi'_{p,q}(k_{p,q}) \psi'_q(\mathcal{K}X_q) \subseteq B_{\vartheta^{-1}(\mathcal{F}')} \subseteq B'_{\vartheta^{-1}(h\mathcal{P})}.$$

By using approximate identities on both sides we have the contradiction

$$\psi'_{p,q}(k_{p,q}) \in B'_{\vartheta^{-1}(h\mathcal{P})}.$$

This concludes the proof of the proposition. □

Combining with [Sehnm 2019, Theorem 3.10] we get the following corollary.

Corollary 5.8. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Then the following are equivalent for a strongly covariant representation (π, t) of $A \times_X P$:*

- (i) *The $*$ -representation π is faithful on A .*
- (ii) *The induced $*$ -representation is faithful on the fixed point algebra B_P of $A \times_X P$.*
- (iii) *The induced $*$ -representation is faithful on the ϑ -fixed point algebra $B_{\vartheta^{-1}(\mathcal{P})}$ of $A \times_X P$.*

In particular this holds for the $$ -representations of $q_{sc}(\mathcal{T}_\lambda(X))$ and $A \times_{X,\lambda} P$.*

A second application of the controlled elimination allows us to pass in between the C^* -envelopes induced by G and \mathcal{G} .

Proposition 5.9. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions, and let X be a compactly aligned product system over P with coefficients in A . Let $\bar{\delta}_G$ be the induced coaction of \mathcal{G} on $\mathcal{T}_\lambda(X)$ and $\mathcal{T}_\lambda(X)^+$. Then $C_{env}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$ inherits a normal coaction of \mathcal{G} and there exists a \mathcal{G} -equivariant $*$ -isomorphism*

$$C_{env}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_G)$$

that fixes $\mathcal{T}_\lambda(X)^+$.

Proof. By Theorem 2.7 and Propositions 2.9 and 3.7, we get that $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$ admits a normal coaction of \mathcal{G} and therefore there exists a \mathcal{G} -equivariant $*$ -epimorphism

$$\Phi : C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_\mathcal{G})$$

that fixes $\mathcal{T}_\lambda(X)^+$. By construction Φ is \mathcal{G} -equivariant, and so it intertwines the faithful conditional expectations induced by \mathcal{G} . On the other hand, by Theorem 2.7 the map Φ is faithful on the G -fixed point algebra of $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$. By Proposition 5.7 the map Φ is faithful on the \mathcal{G} -fixed point algebra of $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$. Consequently Φ is injective. \square

6. Applications

Couniversality of Sehnm’s covariance algebra. We will consider weak right LCM inclusions that are controlled by exact groups. In this case we get normality of the coaction of G on $q_{\text{sc}}(\mathcal{T}_\lambda(X))$, and thus the latter coincides with $A \times_{X,\lambda} P$ and, by [Dor-On et al. 2022, Theorem 5.3], with $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$. This provides another algebraic description of $C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G)$ by the strong covariance relations in the Fock space representation.

Theorem 6.1. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . Consider the canonical $*$ -epimorphisms*

$$q_{\text{sc}}(\mathcal{T}_\lambda(X)) \rightarrow A \times_{X,\lambda} P \simeq C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+).$$

If \mathcal{G} is exact then the left map is faithful. If in addition \mathcal{G} is abelian then the right map is also faithful.

Proof. First we show that the ideal of the strong covariance relations is \mathcal{G} -induced. Let \mathcal{I}_λ be the image of the strong covariance relations in $\mathcal{T}_\lambda(X)$ so that $\mathcal{T}_\lambda(X)/\mathcal{I}_\lambda = q_{\text{sc}}(\mathcal{T}_\lambda(X))$. Let us denote by \bar{B}_F the cores of the Fock representation $(\bar{\pi}, \bar{t})$ and let $q_{\mathcal{I}_\lambda} : \mathcal{T}_\lambda(X) \rightarrow q_{\text{sc}}(\mathcal{T}_\lambda(X))$ be the canonical $*$ -epimorphism. Proposition 5.7 implies that

$$\mathcal{I}_\lambda \cap \bar{B}_{\vartheta^{-1}(\mathcal{P})} = \bigcup_{\substack{\text{finite } \vee\text{-closed} \\ \mathcal{F} \subseteq \mathcal{P}}} \ker q_{\mathcal{I}_\lambda} \cap \bar{B}_{\vartheta^{-1}(\mathcal{F})} = \bigcup_{\substack{\text{finite } \vee\text{-closed} \\ F \subseteq P}} \ker q_{\mathcal{I}_\lambda} \cap \bar{B}_F = \mathcal{I}_\lambda \cap \bar{B}_P.$$

Therefore we get that

$$\mathcal{I}_\lambda = \langle \mathcal{I}_\lambda \cap \bar{B}_P \rangle = \langle \mathcal{I}_\lambda \cap \bar{B}_{\vartheta^{-1}(\mathcal{P})} \rangle,$$

showing that \mathcal{I}_λ is indeed \mathcal{G} -induced.

Consequently, by exactness of \mathcal{G} we derive that the normal coaction of \mathcal{G} on $\mathcal{T}_\lambda(X)$ descends to a normal coaction on the quotient $q_{\text{sc}}(\mathcal{T}_\lambda(X))$. Thus by Proposition 4.12 we have that $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ is a C^* -cover for $(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_\mathcal{G})$. Therefore there exists a \mathcal{G} -equivariant $*$ -epimorphism

$$\Phi : q_{\text{sc}}(\mathcal{T}_\lambda(X)) \rightarrow C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_\mathcal{G})$$

that fixes $\mathcal{T}_\lambda(X)^+$. The $*$ -epimorphism Φ intertwines the coactions (and thus the faithful conditional expectations implemented by normality and exactness of \mathcal{G}), and it is faithful on the G -fixed point algebra

$[q_{sc}(\mathcal{T}_\lambda(X))]_e$ by Corollary 3.13. Hence we derive that Φ is faithful by Corollary 5.8. By Proposition 5.9 we conclude that

$$q_{sc}(\mathcal{T}_\lambda(X)) \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_{\mathcal{G}}) \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G).$$

Now if in addition \mathcal{G} is abelian then $C_{env}^*(\mathcal{T}_\lambda(X)^+)$ inherits the coaction of \mathcal{G} by the dual gauge action $\hat{\mathcal{G}}$. Due to couniversality we thus derive

$$C_{env}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_{\mathcal{G}}) \simeq C_{env}^*(\mathcal{T}_\lambda(X)^+). \quad \square$$

Combining with Proposition 2.9 and Proposition 4.12 we get the following corollary.

Corollary 6.2. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions such that \mathcal{G} is exact, and let X be a compactly aligned product system over P with coefficients in A . Then $q_{sc}(\mathcal{T}_\lambda(X))$ is couniversal with respect to both G -equivariant and \mathcal{G} -equivariant quotients of $\mathcal{T}_\lambda(X)$ that are faithful on A .*

Remark 6.3. As an immediate consequence of Theorem 6.1 we get that the coaction of G on $q_{sc}(\mathcal{T}_\lambda(X))$ is normal. Therefore one can use the results of [Dor-On et al. 2022] to derive that the reduced Hao–Ng problem for discrete group actions on $A \times_{X,\lambda} P$ has a positive answer when (G, P) is controlled by $(\mathcal{G}, \mathcal{P})$ with \mathcal{G} exact. A similar method applies whenever the C^* -envelope functor is stable under crossed products, e.g., for dynamics over abelian locally compact groups or when the tensor algebra is hyperrigid [Katsoulis 2020; Katsoulis and Ramsey 2019], and we leave this to the interested reader.

Next we consider *amenably controlled* weak right LCM inclusions, i.e., the range of the controlled map is inside an amenable group. In this case the reduced C^* -algebras become universal with respect to classes of representations. First we consider $\mathcal{NT}(X)$. (A variant of) the following has been obtained by Fowler [2002] for nondegenerate product systems over quasilattices. Here we extend it to the weak right LCM inclusions framework with a different approach that does not require nondegeneracy of X .

Theorem 6.4. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable and let X be a compactly aligned product system over P with coefficients in A . Then the Fock representation is faithful on $\mathcal{NT}(X)$.*

Conversely, suppose that (π, t) is an injective \mathcal{G} -equivariant Nica-covariant representation of X and for every \vee -closed $\mathcal{F} \subseteq \mathcal{P}$ we have linear independence in the ϑ -cores in the sense that

$$B_{\vartheta^{-1}(\mathcal{F})} = \sum_{h \in \mathcal{F}}^{\oplus} B_{\vartheta^{-1}(h)}.$$

Then (π, t) integrates to a faithful representation of $\mathcal{NT}(X)$.

In particular, a Nica-covariant pair (π, t) defines a faithful representation of $\mathcal{NT}(X)$ if and only if the associated representation is \mathcal{G} -equivariant and satisfies the condition

$$\sum_{p \in F} \psi_p(k_p) = 0 \iff k_p = 0 \quad \text{for all } p \in F,$$

for every \vee -closed $\mathcal{F} \subseteq \mathcal{P}$ and every finite \vee -closed $F \subseteq \vartheta^{-1}(\mathcal{F})$.

Proof. Let $(\hat{\pi}, \hat{\iota})$ be a faithful representation of $\mathcal{NT}(X)$ and consider the canonical $*$ -epimorphism

$$\Phi : \mathcal{NT}(X) = C^*(\hat{\pi}, \hat{\iota}) \rightarrow \mathcal{T}_\lambda(X) = C^*(\bar{\pi}, \bar{\iota}).$$

Let \bar{E}_G be the faithful conditional expectation induced by Proposition 3.8 on $\mathcal{T}_\lambda(X)$. Let \hat{E}_G be the faithful conditional expectation on $\mathcal{NT}(X)$ induced by the amenable \mathcal{G} . Since $\Phi \hat{E}_G = \bar{E}_G \Phi$, it suffices to show injectivity of Φ on \hat{B}_F for every \vee -closed $F \subseteq P$. To this end fix a finite \vee -closed $F \subseteq \vartheta^{-1}(\mathcal{F})$ and suppose that

$$f := \sum \{ \hat{\psi}_{r_1, r_2}(k_{r_1, r_2}) \mid k_{r_1, r_2} \in \mathcal{K}(X_{r_2}, X_{r_1}), r_1, r_2 \in F, \vartheta(r_1) = \vartheta(r_2) \} \in \ker \Phi.$$

Let h be minimal in \mathcal{F} such that $k_{q_1, q_2} \neq 0$ with $\vartheta(q_1) = \vartheta(q_2) = h$. Using condition (A2) of Definition 5.1 and the Fock space representation we have that

$$k_{q_1, q_2} = Q_{q_1} \Phi(f) Q_{q_2} = 0$$

for the projections $Q_p : \mathcal{FX} \rightarrow X_p$, which gives the required contradiction. Thus the Fock representation is injective and also we have linear independence of the cores. The converse follows with a similar proof.

For the last part it is clear that the condition with $F = \{e_G\}$ and $\mathcal{F} = \{e_G\}$ implies that π is injective. Moreover the condition shows that the canonical $*$ -epimorphism Φ is injective on the C^* -subalgebras

$$\hat{B}_F = \text{span}\{\hat{\psi}_r(k_r) \mid r \in F\}$$

for every finite \vee -closed $F \subseteq P$, and so Φ is injective on \hat{B}_P . Thus by Proposition 5.7 the map Φ is injective on $\hat{B}_{\vartheta^{-1}(\mathcal{P})}$ and hence on $\mathcal{NT}(X)$. □

Next we consider the universal covariance algebra $A \times_X P$.

Theorem 6.5. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable and let X be a compactly aligned product system over P with coefficients in A . Then a strongly covariant representation of X integrates to a faithful representation of $A \times_X P$, if and only if it is injective and G -equivariant, if and only if it is injective and \mathcal{G} -equivariant.*

Proof. By Theorem 6.4 we have that $A \times_X P$ coincides with $q_{\text{sc}}(\mathcal{T}_\lambda(X))$ and $A \times_{X, \lambda} P$. Thus the result follows from Corollary 6.2. □

Remark 6.6. When (G, P) is amenably controlled then we have a wider selection for a coaction that implements the extension theorem. The diagram in the proof of Corollary 6.7 on page 1466 depicts those. We denote restrictions of $*$ -homomorphisms by solid arrows, and we have used Proposition 4.11 for the upper and lower completely isometric maps. Recall that if \mathcal{G} is amenable then $C^*(\mathcal{G}) \simeq C^*_\lambda(\mathcal{G})$ is nuclear, and by [Li 2013] $C^*_\lambda(\mathcal{P})$ is also nuclear.

Corollary 6.7. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable. Suppose that $A, X_p \subseteq B(\mathcal{H})$ for $p \in P$ define a compactly aligned product system $X = \{X_p\}_{p \in P}$ and set*

$$\mathcal{A} := \overline{\text{alg}}\{A, X_p \mid p \in P\}.$$

Then the following are equivalent:

(i) *There is a completely isometric isomorphism*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+, \quad \xi_p \mapsto \bar{t}(\xi_p).$$

(ii) *There is a completely contractive map*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+ \otimes C^*(G), \quad \xi_p \mapsto \bar{t}(\xi_p) \otimes u_p.$$

(iii) *There is a completely contractive map*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+ \otimes C_\lambda^*(G), \quad \xi_p \mapsto \bar{t}(\xi_p) \otimes \lambda_p.$$

(iv) *There is a completely contractive map*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+ \otimes C_\lambda^*(P), \quad \xi_p \mapsto \bar{t}(\xi_p) \otimes V_p.$$


(v) *There is a completely contractive map*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+ \otimes C^*(\mathcal{G}), \quad \xi_p \mapsto \bar{t}(\xi_p) \otimes u_{\vartheta(p)}.$$

(vi) *There is a completely contractive map*

$$\mathcal{A} \rightarrow \mathcal{T}_\lambda(X)^+ \otimes C_\lambda^*(\mathcal{P}), \quad \xi_p \mapsto \bar{t}(\xi_p) \otimes V_{\vartheta(p)}.$$

Proof. The proof follows by the system of completely positive maps fixing the nonselfadjoint part below, where the solid arrows denote the maps that arise from restrictions of $*$ -homomorphisms from the appropriate C^* -algebras to the required subalgebras.



□

Exactness and nuclearity. We will require some results about nuclearity which we record here for convenience.

Lemma 6.8 [Katsura 2004, Proposition B.8]. *Let (π, t) be a representation of a C^* -correspondence X over A such that $\pi(A) \subseteq B$ and $t(X) \subseteq Y$ for a second C^* -correspondence Y over B . If $\pi : A \rightarrow B$ is nuclear then the induced map $\psi : \mathcal{K}X \rightarrow \mathcal{K}Y$ is nuclear.*

Lemma 6.9 [Kakariadis 2020, Proposition 3.1]. *Let A, A' be C^* -algebras and consider the ideals $I \triangleleft A$ and $I' \triangleleft A'$. Suppose we have the commutative diagram of short exact sequences*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \longrightarrow & A & \longrightarrow & A/I & \longrightarrow & 0 \\ & & \downarrow \varphi_0 & & \downarrow \varphi & & \downarrow \tilde{\varphi} & & \\ 0 & \longrightarrow & I' & \longrightarrow & A' & \longrightarrow & A'/I' & \longrightarrow & 0 \end{array}$$

where $\varphi : A \rightarrow A'$ is an injective $*$ -homomorphism that satisfies $\varphi(I) \subseteq I'$, $\tilde{\varphi} : A/I \rightarrow A'/I'$ is the induced map and $\varphi_0 := \varphi|_I$. If $\varphi : A \rightarrow A'$ is nuclear, then φ_0 and $\tilde{\varphi}$ are both nuclear.

Lemma 6.10 [Kakariadis 2020, Proposition 3.3]. *Let A, A' be C^* -algebras and consider the ideals $I \triangleleft A$ and $I' \triangleleft A'$. Suppose we have the commutative diagram of short exact sequences*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \longrightarrow & A & \longrightarrow & A/I & \longrightarrow & 0 \\ & & \downarrow \varphi_0 & & \downarrow \varphi & & \downarrow \tilde{\varphi} & & \\ 0 & \longrightarrow & I' & \longrightarrow & A' & \longrightarrow & A'/I' & \longrightarrow & 0 \end{array}$$

where $\varphi : A \rightarrow A'$ is an injective $*$ -homomorphism that satisfies $\varphi(I) \subseteq I'$, $\tilde{\varphi} : A/I \rightarrow A'/I'$ is the induced map and $\varphi_0 := \varphi|_I$. Suppose further that there exists a contractive approximate identity (e_i) of I' such that $\varphi(a)e_i \in \varphi_0(I)$ for all $a \in A$. If φ_0 and $\tilde{\varphi}$ are nuclear, then so is φ .

First we provide a nuclearity and exactness result for $\mathcal{T}_\lambda(X)$.

Theorem 6.11. *Let (G, P) be a weak right LCM inclusion and X be a compactly aligned product system over P with coefficients in A . Let $\bar{E}_P : \mathcal{T}_\lambda(X) \rightarrow \bar{B}_P$ be the faithful conditional expectation that arises by compressing to the diagonal. Then the following are equivalent:*

- (i) *A is nuclear (resp. exact) and $\bar{E}_P \otimes_{\max} \text{id}_D$ is a faithful conditional expectation on $\mathcal{T}_\lambda(X) \otimes_{\max} D$ for all C^* -algebras D .*
- (ii) *$\mathcal{T}_\lambda(X)$ is nuclear (resp. exact).*

Proof. We will show nuclearity; exactness follows in the same way. Notice that for any C^* -algebra D we have the commutative diagram

$$\begin{array}{ccc} C^*(\bar{\pi}, \bar{t}) \otimes_{\max} D & \longrightarrow & C^*(\bar{\pi}, \bar{t}) \otimes D \\ \downarrow \bar{E}_P \otimes_{\max} \text{id} & & \downarrow \bar{E}_P \otimes \text{id} \\ \bar{B}_P \otimes_{\max} D & \longrightarrow & \bar{B}_P \otimes D \end{array}$$

and we recall that $\bar{E}_P \otimes \text{id}$ is faithful on $C^*(\bar{\pi}, \bar{t}) \otimes D$.

Suppose first that $C^*(\bar{\pi}, \bar{t})$ is nuclear. Then trivially $\bar{E}_P \otimes_{\max} \text{id}$ is faithful on $C^*(\bar{\pi}, \bar{t}) \otimes_{\max} D$. Since A is the corner of $C^*(\bar{\pi}, \bar{t})$ at the (e, e) -place we have that A is nuclear, as the compression of a nuclear C^* -algebra.

For the converse, the diagram above implies that it suffices to show that \bar{B}_P is nuclear. Equivalently it suffices to show that \bar{B}_F is nuclear for every finite \vee -closed $F \subseteq P$. To this end let $F = \{p_1, \dots, p_n\}$. We choose the enumeration so that it covers the partial order in F in the sense that if $p_m > p_{m'}$ then $m < m'$. We will use induction on n .

For the first step we have that $\bar{\psi}_{p_1}(\mathcal{K}X_{p_1})$ is nuclear as A is nuclear by [Katsura 2004, Proposition B.7]. For the inductive step suppose that \bar{B}_{F_k} is nuclear for $F_k = \{p_1, \dots, p_k\}$ (which is \vee -closed by the choice of the enumeration). We will show that so is $\bar{B}_{F_{k+1}}$ for $F_{k+1} = \{p_1, \dots, p_k, p_{k+1}\}$. The enumeration shows that p_{k+1} is minimal in F_{k+1} and hence

$$\bar{B}_{F_{k+1}} = \bar{B}_{F_k} \oplus \bar{\psi}_{p_{k+1}}(\mathcal{K}X_{p_{k+1}}).$$

Indeed let $k_{p_i} \in \mathcal{K}X_{p_i}$ such that

$$\sum_{i=1}^{k+1} \bar{\psi}_{p_i}(k_{p_i}) = 0.$$

Due to minimality of p_{k+1} in F_{k+1} we have that

$$k_{p_{k+1}} = Q_{p_{k+1}} \left(\sum_{i=1}^{k+1} \bar{\psi}_{p_i}(k_{p_i}) \right) Q_{p_{k+1}} = 0,$$

for the projection $Q_{p_{k+1}} : \mathcal{F}X \rightarrow X_{p_{k+1}}$. Minimality of p_{k+1} also gives that \bar{B}_{F_k} is an ideal in $\bar{B}_{F_{k+1}}$, and we thus derive the following short exact sequence

$$0 \longrightarrow \bar{B}_{F_k} \longrightarrow \bar{B}_{F_{k+1}} \longrightarrow \bar{\psi}_{p_{k+1}}(\mathcal{K}X_{p_{k+1}}) \longrightarrow 0.$$

Since \bar{B}_{F_k} is nuclear by the inductive hypothesis and $\bar{\psi}_{p_{k+1}}(\mathcal{K}X_{p_{k+1}})$ is nuclear by the base case we have that $\bar{B}_{F_{k+1}}$ is nuclear. Inducing on k gives that $\bar{B}_F = \bar{B}_{F_n}$ is nuclear. □

In the amenable controlled case, and by combining with Theorem 6.4, we can deduce nuclearity and exactness of $\mathcal{NT}(X)$ from nuclearity and exactness of A , and conversely. The exactness equivalence passes to $A \times_X P$, however this fails for nuclearity even for $P = \mathbb{Z}_+$ due to a counterexample of Ozawa in [Katsura 2004]. In [Kakariadis 2020] it is shown that $A \times_X P$ is nuclear if and only if the embedding $A \hookrightarrow A \times_X P$ is nuclear when (G, P) is a quasilattice controlled by $(\mathbb{Z}^n, \mathbb{Z}_+^n)$ that satisfies a minimality condition. In fact this holds for any quotient in between the Toeplitz–Nica–Pimsner and the covariance algebra. Here we generalize to controlled maps by amenable weak right LCM inclusions. Recall that in the amenable controlled case the reduced C^* -algebras are universal.

Theorem 6.12. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable and let X be a compactly aligned product system over P with coefficients in A . Let (π, t) be an equivariant injective Nica-covariant representation of X . Then A is exact if and only if $C^*(\pi, t)$ is exact.*

Proof. We are going to introduce new product systems from X . Therefore in order to make a distinction we will write \bar{E}_p^X for the faithful conditional expectation on the Fock C^* -algebra $\mathcal{T}_\lambda(X)$ of X .

If $C^*(\pi, t)$ is exact then so is A , since exactness passes to C^* -subalgebras. For the converse, by Theorem 6.4 we have that X is amenable and thus $C^*(\pi, t)$ is a quotient of $\mathcal{T}_\lambda(X)$. Hence it suffices to show that $\mathcal{T}_\lambda(X)$ is exact. In view of Theorem 6.11 it suffices to show that $\bar{E}_p^X \otimes_{\max} \text{id}_D$ is faithful on $\mathcal{T}_\lambda(X) \otimes_{\max} D$ for all C^* -algebras D .

Towards this end let the product system $Y = \{Y_p\}_{p \in P}$ be defined by

$$Y_p := \overline{\bar{t}_p(X_p) \odot D}^{\otimes_{\max}} \subseteq \mathcal{T}_\lambda(X) \otimes_{\max} D.$$

That Y is a product system follows as X is a product system. Since X is compactly aligned we have that

$$Y_p Y_p^* Y_q Y_q^* \subseteq \overline{\bar{\psi}_p(\mathcal{K}X_p) \bar{\psi}_q(\mathcal{K}X_q) \odot D}^{\otimes_{\max}} = \overline{\bar{\psi}_w(\mathcal{K}X_w) \odot D}^{\otimes_{\max}} = [Y_w Y_w^*]$$

for $wP = pP \cap qP$, with the understanding that $Y_p Y_p^* Y_q Y_q^* = (0)$ when p and q have no common right common multiple. Thus by Proposition 4.7 we get that Y is a compactly aligned product system over P with coefficients in A .

Again by Theorem 6.4 we have that Y is amenable. Our goal is to show that the identity representation on Y is faithful on $\mathcal{NT}(Y) \simeq \mathcal{T}_\lambda(Y)$, and thus we have that

$$\mathcal{NT}(Y) \simeq \mathcal{T}_\lambda(Y) \simeq \mathcal{T}_\lambda(X) \otimes_{\max} D.$$

We then derive that the faithful conditional expectation \bar{E}_p^Y on $\mathcal{T}_\lambda(Y)$ coincides with $\bar{E}_p^X \otimes_{\max} \text{id}_D$ and the proof will be completed. We will invoke Theorem 6.4.

First we see that the identity representation is \mathcal{G} -equivariant. Indeed we have that $(\bar{\pi}, \bar{t})$ admits a coaction $\bar{\delta}_\mathcal{G}$ of \mathcal{G} and thus we have an equivariant $*$ -homomorphism

$$\bar{\delta}_\mathcal{G} \otimes_{\max} \text{id}_D : \mathcal{T}_\lambda(X) \otimes_{\max} D \rightarrow (\mathcal{T}_\lambda(X) \otimes C^*(\mathcal{G})) \otimes_{\max} D.$$

By amenability of \mathcal{G} and associativity of the maximal tensor product we get that

$$\begin{aligned} (\mathcal{T}_\lambda(X) \otimes C^*(\mathcal{G})) \otimes_{\max} D &\simeq \mathcal{T}_\lambda(X) \otimes_{\max} C^*(\mathcal{G}) \otimes_{\max} D \\ &\simeq (\mathcal{T}_\lambda(X) \otimes_{\max} D) \otimes_{\max} C^*(\mathcal{G}) \simeq (\mathcal{T}_\lambda(X) \otimes_{\max} D) \otimes C^*(\mathcal{G}), \end{aligned}$$

and thus we deduce that $\bar{\delta}_\mathcal{G} \otimes_{\max} \text{id}_D$ is a coaction of \mathcal{G} on $\mathcal{T}_\lambda(X) \otimes_{\max} D$. By construction $\bar{\delta}_\mathcal{G} \otimes_{\max} \text{id}_D$ satisfies the coaction identity with aligned fibers in the sense that

$$[\mathcal{T}_\lambda(X) \otimes_{\max} D]_\mathcal{G} = \overline{[\mathcal{T}_\lambda(X)]_\mathcal{G} \odot D}^{\otimes_{\max}}.$$

Secondly let $F \subseteq P$ be a \vee -closed finite set and let $k'_p \in \mathcal{K}Y_p$ such that $\sum_{p \in F} \text{id}(k'_p) = 0$. For every state $\phi \in \mathcal{S}(D)$ we have the completely contractive map

$$\text{id} \otimes_{\max} \phi : [Y_p Y_p^*] \otimes_{\max} D \rightarrow \bar{\psi}_p(\mathcal{K}X_p), \quad \bar{\psi}_p(k_p) \otimes d \mapsto \phi(d) \bar{\psi}_p(k_p).$$

Therefore we derive

$$\sum_{p \in F} (\text{id} \otimes_{\max} \phi)(k'_p) = (\text{id} \otimes_{\max} \phi) \left(\sum_{p \in F} k'_p \right) = 0.$$

Note here that this is a relation in $\mathcal{T}_\lambda(X)$ with every $(\text{id} \otimes_{\max} \phi)(k'_p) \in \bar{\psi}_p(\mathcal{K}X_p)$. Thus if p_0 is a minimal element in F such that $(\text{id} \otimes_{\max} \phi)(k'_{p_0}) \neq 0$, then we get

$$P_{p_0}(\text{id} \otimes_{\max} \phi)(k'_{p_0})P_{p_0} = P_{p_0}(\text{id} \otimes_{\max} \phi)\left(\sum_{p \in F} k'_p\right)P_{p_0} = 0,$$

where $P_{p_0} : \mathcal{F}X \rightarrow X_{p_0}$ is the canonical projection. However the compression to X_{p_0} is a faithful $*$ -representation on $\bar{\psi}_{p_0}(\mathcal{K}X_{p_0})$, and thus we get the contradiction that $(\text{id} \otimes_{\max} \phi)(k'_{p_0}) = 0$. Continuing inductively we deduce that $(\text{id} \otimes_{\max} \phi)(k'_p) = 0$ for all $p \in F$ (one by one for fixed ϕ). As this holds for all ϕ and the family $\{\text{id} \otimes_{\max} \phi\}_{\phi \in \mathcal{S}(D)}$ separates $[Y_p Y_p^*] \otimes_{\max} D$ we get that $k'_p = 0$ for all $p \in F$. Hence the assumptions of Theorem 6.4 hold for Y and the proof is complete. \square

Theorem 6.13. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions with \mathcal{G} amenable and let X be a compactly aligned product system over P with coefficients in A . Let (π, t) be an equivariant injective Nica-covariant representation of X . Then $A \hookrightarrow C^*(\pi, t)$ is nuclear if and only if $C^*(\pi, t)$ is nuclear.*

Proof. It is clear that if $C^*(\pi, t)$ is nuclear then $A \hookrightarrow C^*(\pi, t)$ is nuclear. Let us prove the converse. By Theorem 6.4 we have that $\mathcal{T}_\lambda(X) \simeq \mathcal{NT}(X)$ and so (π, t) promotes to a $*$ -representation of $\mathcal{T}_\lambda(X)$. Due to amenability, $C^*(\mathcal{G}) = C^*_\lambda(\mathcal{G})$ is nuclear (and so the minimal and the maximal tensor product coincide). Let $\delta : C^*(\pi, t) \rightarrow C^*(\pi, t) \otimes C^*_\lambda(\mathcal{G})$ be the coaction of \mathcal{G} and let $E = (\text{id} \otimes E_\mathcal{G})\delta$ be the faithful conditional expectation induced on $C^*(\pi, t)$ by the faithful conditional expectation $E_\mathcal{G}$ of $C^*_\lambda(\mathcal{G})$. Let D be any C^* -algebra. Associativity of \otimes_{\max} and nuclearity of $C^*_\lambda(\mathcal{G})$ yields

$$D \otimes_{\max} C^*(\pi, t) \otimes_{\max} C^*_\lambda(\mathcal{G}) \simeq (D \otimes_{\max} C^*(\pi, t)) \otimes C^*_\lambda(\mathcal{G}),$$

and so $\text{id}_D \otimes_{\max} \text{id} \otimes_{\max} E_\mathcal{G} = (\text{id}_D \otimes_{\max} \text{id}) \otimes E_\mathcal{G}$ is faithful on $D \otimes_{\max} C^*(\pi, t) \otimes_{\max} C^*_\lambda(\mathcal{G})$. Hence

$$\text{id}_D \otimes_{\max} E := (\text{id}_D \otimes_{\max} \text{id} \otimes_{\max} E_\mathcal{G})(\text{id}_D \otimes_{\max} \delta)$$

is a faithful conditional expectation of $D \otimes_{\max} C^*(\pi, t)$ on $D \otimes_{\max} B_\mathcal{P}$. Therefore we have the commutative diagram

$$\begin{array}{ccc} C^*(\pi, t) \otimes_{\max} D & \longrightarrow & C^*(\pi, t) \otimes D \\ \downarrow & & \downarrow \\ B_\mathcal{P} \otimes_{\max} D & \longrightarrow & B_\mathcal{P} \otimes D \end{array}$$

where the vertical arrows are faithful conditional expectations. Hence it suffices to show that if $\pi : A \rightarrow B_\mathcal{P}$ is nuclear then the fixed point algebra $B_\mathcal{P}$ is nuclear. As the latter is an inductive limit, it suffices to show that nuclearity of π in $B_\mathcal{P}$ induces nuclearity of the embedding $B_{\vartheta^{-1}(F)} \hookrightarrow B_{\vartheta^{-1}(\mathcal{P})}$ for every finite \vee -closed $\mathcal{F} \subseteq \mathcal{P}$. We will actually show nuclearity of the embedding

$$B_{\vartheta^{-1}(\mathcal{F})} \hookrightarrow B_{\vartheta^{-1}(\mathcal{F} \cdot \mathcal{P})} \subseteq B_{\vartheta^{-1}(\mathcal{P})},$$

where we write

$$\vartheta^{-1}(\mathcal{F} \cdot \mathcal{P}) = \{pP \mid \vartheta(p) \in \mathcal{F}\}.$$

First we remark that $B_{\mathcal{F}}$ contains a contractive approximate identity for $B_{\mathcal{F}\cdot P}$. Indeed let (e_i) be a contractive approximate identity for $B_{\mathcal{F}}$ so that $\lim_i e_i \psi_p(k_p) = \psi_p(k_p)$ for every $p \in \vartheta^{-1}(\mathcal{F})$. Consequently $\lim_i e_i t_p(\xi_p) = t_p(\xi_p)$ for every $p \in \vartheta^{-1}(\mathcal{F})$ and thus

$$\lim_i e_i t_p(\xi_p) t_r(\xi_r) t_s(\xi_s)^* = t_p(\xi_p) t_r(\xi_r) t_s(\xi_s)^* \quad \text{for all } r, s \in P.$$

Thus $\lim_i e_i \psi_{p,q}(k_{p,q}) = \psi_{p,q}(k_{p,q})$ for every $p, q \in \vartheta^{-1}(\mathcal{F} \cdot P)$.

Now fix a finite \vee -closed \mathcal{F} . By using maximal elements we can write \mathcal{F} in levels, i.e.,

$$\mathcal{F} = \{h_{11}, \dots, h_{1n_1}, h_{21}, \dots, h_{2n_2}, \dots, h_{m1}, \dots, h_{mn_m}\},$$

such that every

$$\mathcal{F}_i := \{h_{i1}, \dots, h_{in_i}\}, \quad \text{with } i \in \{1, \dots, m\},$$

consists of the maximal elements of $\mathcal{F} \setminus \cup_{j=1}^{i-1} \mathcal{F}_j$ and \mathcal{F}_1 consists of the maximal elements of \mathcal{F} .

We now proceed by induction. For the base case let $h \in \mathcal{P}$ and consider the space

$$Y_h := \sum_{p \in \vartheta^{-1}(h)} t_p(X_p).$$

By using condition (A2) of Definition 5.1 we can equip Y_h with the A -valued bilinear map defined by

$$\langle y_h, y'_h \rangle := y_h^* y'_h \in \pi(A) \quad \text{for all } y_h, y'_h \in Y_h.$$

Then each Y_h becomes a C^* -correspondence over A , since π is faithful. The embedding $Y_h \hookrightarrow [Y_h B_{\vartheta^{-1}(\mathcal{P})}]$ and nuclearity of $\pi(A) \hookrightarrow B_{\vartheta^{-1}(\mathcal{P})}$ imply nuclearity of the embedding

$$B_{\vartheta^{-1}(h)} = [Y_h Y_h^*] \hookrightarrow [Y_h B_{\vartheta^{-1}(\mathcal{P})} Y_h^*] = B_{\vartheta^{-1}(h\mathcal{P})} \quad \text{for all } h \in \mathcal{P},$$

by [Katsura 2004, Proposition B.8]. Maximality of the h_{1j} in \mathcal{F} yields that the $h_{1j}\mathcal{P}$ are minimal in $\{h\mathcal{P} \mid h \in \mathcal{F}\}$ with respect to inclusions. As \mathcal{F} is \vee -closed we have that $h_{1j}\mathcal{P} \cap h_{1j'}\mathcal{P} = \emptyset$ for $j \neq j'$. Hence the C^* -algebras $B_{\vartheta^{-1}(h_{1j})}$ are orthogonal and thus the embedding

$$B_{\mathcal{F}_1} = \sum_{j=1}^{n_1} \oplus B_{\vartheta^{-1}(h_{1j})} \hookrightarrow \sum_{j=1}^{n_1} B_{\vartheta^{-1}(h_{1j}\mathcal{P})} \subseteq B_{\vartheta^{-1}(\mathcal{F}_1 \cdot \mathcal{P})}$$

is nuclear. For the inductive hypothesis suppose we have shown that the embedding $B_{\vartheta^{-1}(\mathcal{F}')} \hookrightarrow B_{\vartheta^{-1}(\mathcal{F}' \cdot \mathcal{P})}$ is nuclear for

$$\mathcal{F}' = \{h_{11}, \dots, h_{1n_1}, \dots, h_{i1}, \dots, h_{ij}\}$$

for some $j \in \{1, \dots, n_i\}$. If $j < n_i$ then set $h := h_{i(j+1)}$; if $j = n_i$ then set $h = h_{(i+1)1}$. We will show that the embedding

$$B_{\vartheta^{-1}(\mathcal{F}'')} \hookrightarrow B_{\vartheta^{-1}(\mathcal{F}'' \cdot \mathcal{P})} \quad \text{for } \mathcal{F}'' := \mathcal{F}' \cup \{h\}$$

is nuclear. By construction $B_{\vartheta^{-1}(\mathcal{F}'')}$ is an ideal in $B_{\vartheta^{-1}(\mathcal{F}'')}$ and $B_{\vartheta^{-1}(\mathcal{F}'')} = B_{\vartheta^{-1}(h)} + B_{\vartheta^{-1}(\mathcal{F}')}$; thus

$$B_{\vartheta^{-1}(\mathcal{F}'')}/B_{\vartheta^{-1}(\mathcal{F}')} \simeq B_{\vartheta^{-1}(h)}/(B_{\vartheta^{-1}(h)} \cap B_{\vartheta^{-1}(\mathcal{F}')}).$$

Likewise $B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})}$ is an ideal of $B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})}$. From the base case we have nuclearity of the map

$$B_{\vartheta^{-1}(h)} \hookrightarrow B_{\vartheta^{-1}(h\mathcal{P})} \subseteq B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})}.$$

By applying Lemma 6.9 on the commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & B_{\vartheta^{-1}(h)}/B_{\vartheta^{-1}(\mathcal{F}')} & \longrightarrow & B_{\vartheta^{-1}(h)} & \longrightarrow & B_{\vartheta^{-1}(h)}/(B_{\vartheta^{-1}(h)} \cap B_{\vartheta^{-1}(\mathcal{F}')} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})}/B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} & \longrightarrow & 0 \end{array}$$

we get that the right vertical arrow is nuclear, i.e., the map

$$B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})}/B_{\vartheta^{-1}(\mathcal{F}')} \simeq B_{\vartheta^{-1}(h)}/(B_{\vartheta^{-1}(h)} \cap B_{\vartheta^{-1}(\mathcal{F}')} \rightarrow B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})}/B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})}$$

is nuclear. Let $(e_i) \subseteq B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})}$ be a contractive approximate identity for $B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})}$, and note that

$$B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} \cdot e_i \subseteq B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} \cdot B_{\vartheta^{-1}(\mathcal{F}')} = B_{\vartheta^{-1}(\mathcal{F}')}$$

Using the inductive hypothesis and Lemma 6.10 on the commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}')} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}'')} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}'')}/B_{\vartheta^{-1}(\mathcal{F}')} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})} & \longrightarrow & B_{\vartheta^{-1}(\mathcal{F}''\cdot\mathcal{P})}/B_{\vartheta^{-1}(\mathcal{F}'\cdot\mathcal{P})} & \longrightarrow & 0 \end{array}$$

we derive that the middle vertical arrow is nuclear, as required. This concludes the inductive step. Now by using induction we derive that $B_{\vartheta^{-1}(\mathcal{F})} \hookrightarrow B_{\vartheta^{-1}(\mathcal{F}\cdot\mathcal{P})}$ is nuclear, and the proof is complete. \square

7. Saturated controlled maps

A product system reparametrization. Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map between weak right LCM inclusions and let X be a compactly aligned product system over P with coefficients in A . We can then define the C^* -correspondence

$$Y_h := \sum_{p \in \vartheta^{-1}(h)}^{\oplus} X_p \quad \text{for all } h \in \mathcal{P}.$$

One is tempted to consider the family $Y = \{Y_h\}_{h \in \mathcal{P}}$ and associate its C^* -algebras with those of X . However it is not clear that Y is in general a product system (let alone compactly aligned). Nevertheless this happens for controlled maps that satisfy one extra condition.

Definition 7.1. Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a controlled map of weak right LCM inclusions. We say that ϑ is *saturated* if for any $h \in \mathcal{P}$ and $t \in \vartheta^{-1}(h\mathcal{P})$ there exists an $s \in P$ with $\vartheta(s)\mathcal{P} = h\mathcal{P}$ and $t \in sP$.

Remark 7.2. In particular, saturated maps satisfy the following property:

(A3) If $z \in \mathcal{P}^*$ then there exists an $x \in P^*$ such that $\vartheta(x) = z$.

Indeed, we apply the saturation property for $z \in \mathcal{P}^*$ and $t = e_G \in \vartheta^{-1}(z\mathcal{P})$ to obtain an $x \in P$ with $e_G \in xP$. Hence we get that $P = xP$ giving that $x \in P^*$.

The following provides a good supply of saturated controlled maps. Recall that a pair (G, P) is a *total order* if $G = P^{-1} \cup P$ and $P^{-1} \cap P = \{e_G\}$. It is clear that total orders, being lattices, form weak right LCM inclusions.

Proposition 7.3. *Let (G, P) be an abelian total order. For $n \in \mathbb{N} \cup \{\infty\}$ consider the free product $(\ast_{i=1}^n G, \ast_{i=1}^n P)$ of n copies of (G, P) . Then the map*

$$\vartheta : (\ast_{i=1}^n G, \ast_{i=1}^n P) \rightarrow (G, P), \quad (g_1)_{i_1} (g_2)_{i_2} \cdots (g_k)_{i_k} \mapsto g_1 + g_2 + \cdots + g_k$$

is a saturated controlled map.

Proof. For condition (A1) of Definition 5.1, if $\bar{p}, \bar{q} \in \ast_{i=1}^n P$ with

$$\bar{p}(\ast_{i=1}^n P) \cap \bar{q}(\ast_{i=1}^n P) \neq \emptyset,$$

then the freeness construction implies that either $\bar{p} \leq \bar{q}$ or $\bar{q} \leq \bar{p}$.

For condition (A2) of Definition 5.1 suppose that \bar{p}, \bar{q} have a right LCM and they satisfy $\vartheta(\bar{p}) = \vartheta(\bar{q})$. Without loss of generality assume that $\bar{r} = \bar{p}^{-1} \bar{q} \in \ast_{i=1}^n P$. Then $\vartheta(\bar{r}) = 0$. If $\bar{r} = (r_1)_{i_1} \cdots (r_k)_{i_k}$ then $r_1 + \cdots + r_k = 0$ giving that $r_k \in -P \cap P = \{0\}$. Inductively we get that $r_1 = \cdots = r_k = 0$ and so $\bar{p} = \bar{q}$.

Next we verify that ϑ is saturated. To this end let

$$\bar{p} = (p_1)_{i_1} (p_2)_{i_2} \cdots (p_k)_{i_k},$$

and let $h \in P$ with

$$h \leq \vartheta(\bar{p}) = p_1 + p_2 + \cdots + p_k.$$

Let $\ell \in \{1, \dots, k\}$ be the smallest index such that $h \leq p_1 + p_2 + \cdots + p_\ell$. Set

$$h' = \begin{cases} h & \text{if } \ell = 1, \\ h - (p_1 + p_2 + \cdots + p_{\ell-1}) & \text{otherwise,} \end{cases}$$

and notice that $h' \in P$ with $h' \leq p_\ell$. Let

$$\bar{q} = \begin{cases} (h')_{i_1} & \text{if } \ell = 1, \\ (p_1)_{i_1} (p_2)_{i_2} \cdots (p_{\ell-1})_{i_{\ell-1}} (h')_{i_\ell} & \text{otherwise.} \end{cases}$$

Then $\bar{q} \leq \bar{p}$ and $\vartheta(\bar{q}) = h$, as desired. \square

Example 7.4. A second example comes from types of semidirect products. Let (G, P) and (H, S) be quasilattice ordered groups and consider an action $\alpha : H \rightarrow \text{Aut}(G)$ such that $\alpha|_S : S \rightarrow \text{Aut}(P)$ restricts to automorphisms of P . Then we can form the semidirect products $G \rtimes_\alpha H$ and $P \rtimes_\alpha S$ with respect to the relations $\alpha_h(g)h = hg$. The condition on α makes $P \cdot S$ a subsemigroup of the semidirect product, and in [Kakariadis 2020] it is shown that the pair $(G \rtimes_\alpha H, P \rtimes_\alpha S)$ is quasilattice ordered. Now suppose that (G, P) admits an abelian controlled map ϑ_1 in $(\mathcal{G}_1, \mathcal{P}_1)$ and (H, S) admits an abelian controlled map ϑ_2 in $(\mathcal{G}_2, \mathcal{P}_2)$. In order for the semidirect product to inherit the obvious controlled map on $(\mathcal{G}_1 \oplus \mathcal{G}_2, \mathcal{P}_1 \oplus \mathcal{P}_2)$, it is necessary that α is ϑ_1 -invariant in the sense that $\vartheta_1 \alpha_h = \vartheta_1$ for all $h \in H$. We can then define the homomorphism

$$\vartheta : (G \rtimes_\alpha H, P \rtimes_\alpha S) \rightarrow (\mathcal{G}_1 \oplus \mathcal{G}_2, \mathcal{P}_1 \oplus \mathcal{P}_2) \quad \text{such that } \vartheta(gh) = (\vartheta_1(g), \vartheta_2(h)).$$

We claim that if ϑ_1 and ϑ_2 are saturated, then so is ϑ . Suppose that $\vartheta(gh) = (\vartheta_1(g), \vartheta_2(h)) \geq (m, \ell)$. Then there are $s_1, r_1 \in G$ and $s_2, r_2 \in H$ such that

$$g = s_1 r_1, \quad \vartheta_1(s_1) = m \quad \text{and} \quad h = s_2 r_2, \quad \vartheta_2(s_2) = \ell.$$

It follows that $\vartheta(s_1 s_2) = (m, \ell)$ and $gh = s_1 s_2 \alpha_{s_2}^{-1}(r_1) r_2$.

The following examples show that surjectivity is not enough to render a controlled map saturated.

Example 7.5. Take the free quasilattice on two symbols a, b and take ϑ to be its abelianization map. Then for ab and $(0, 1) \in \mathbb{Z}^2$ we have that $\vartheta(ab) = (1, 1) \geq (0, 1)$. However $\{b\} = \vartheta^{-1}((0, 1))$ and $ab \not\leq b$. (Although, Proposition 7.3 induces a saturated map on free quasilattices.)

Example 7.6. Consider the Baumslag–Solitar group $B(3, 3) = \langle a, b \mid a^3 b = b a^3 \rangle$. Recall that every element $x \in B(3, 3)$ admits a unique normal form

$$x = a^{p_1} b^{\varepsilon_1} a^{p_2} \dots a^{p_k} b^{\varepsilon_k} a^{p_{k+1}} \quad \text{with } p_1, \dots, p_k \in \{0, 1, 2\}, \quad p_{k+1} \in \mathbb{Z}, \quad k \in \mathbb{Z}_+.$$

Let $B_+(3, 3)$ be its subsemigroup generated by a, b . If x is in its normal form as above, it follows that

$$x = a^{p_1} b^{\varepsilon_1} a^{p_2} \dots a^{p_k} b^{\varepsilon_k} a^{p_{k+1}} \in B_+(3, 3) \iff \varepsilon_1, \dots, \varepsilon_k = 1, \quad p_{k+1} \geq 0.$$

By [Spielberg 2012, Theorem 2.11] we have that the pair $(B(3, 3), B_+(3, 3))$ is a quasilattice ordered group. In [Kakariadis 2020] it is shown that the abelianization gives a surjective controlled map

$$\vartheta : (B(3, 3), B_+(3, 3)) \rightarrow (\mathbb{Z}^2, \mathbb{Z}_+^2), \quad a^{p_1} b a^{p_2} b \dots a^{p_k} b a^{p_{k+1}} \mapsto (p_1 + \dots + p_{k+1}, k).$$

However this map is not saturated. Take $t = a^2 b$ and $h = (1, 1)$ so that

$$\vartheta(t) = (2, 1) \in (1, 1) + \mathbb{Z}_+^2.$$

We have that $\vartheta^{-1}(1, 1) = \{ab, ba\}$ and thus these are the only choices for a possible s with $\vartheta(s) = (1, 1)$ and $s \leq t$. However we see that

$$(ab)^{-1} t = b^{-1} a b \notin B_+(3, 3) \quad \text{and} \quad (ba)^{-1} t = a^{-1} b^{-1} a^2 b \notin B_+(3, 3).$$

Theorem 7.7. Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a saturated controlled map between weak right LCM inclusions. Let X be an (injective) compactly aligned product system over P with coefficients in A and let

$$Y_h := \sum_{p \in \vartheta^{-1}(h)}^\oplus X_p \quad \text{for } h \in \mathcal{P}.$$

Then the collection $Y = \{Y_h\}_{h \in \mathcal{P}}$ is an (injective) compactly aligned product system over \mathcal{P} with coefficients in A such that $\mathcal{T}_\lambda(X)^+ \simeq \mathcal{T}_\lambda(Y)^+$ with

$$\mathcal{T}_\lambda(X) \simeq \mathcal{T}_\lambda(Y) \quad \text{and} \quad A \times_{X, \lambda} P \simeq A \times_{Y, \lambda} \mathcal{P},$$

by $*$ -homomorphisms that preserve the inclusions $X_p \mapsto Y_{\vartheta(p)}$ for all $p \in P$. These $*$ -isomorphisms further lift to $*$ -isomorphisms

$$\mathcal{NT}(X) \simeq \mathcal{NT}(Y) \quad \text{and} \quad A \times_X P \simeq A \times_Y \mathcal{P}$$

that preserve the inclusions $X_p \hookrightarrow Y_{\vartheta(p)}$ for all $p \in P$.

Proof. Let A act on both the left and right of each Y_h with $h \in \mathcal{P}$ via the usual multiplication of operators. By using condition (A2) of Definition 5.1 we can equip Y_h with the A -valued bilinear map defined by

$$\langle y_h, y'_h \rangle := y_h^* y'_h \in A \subseteq \mathcal{T}_\lambda(X) \quad \text{for all } y_h, y'_h \in Y_h.$$

Then each Y_h becomes a C^* -correspondence over A . Since $\ker \varphi_{Y_h} = \bigcap_{p \in \vartheta^{-1}(h)} \ker \varphi_{X_p}$, we have that every Y_h is injective when every X_p is so.

We now show that $Y := \{Y_h\}_{h \in \mathcal{P}}$ is a product system. Since $[Y_h Y_g] \simeq Y_h \otimes_A Y_g$ we have to show that

$$[Y_h Y_g] = Y_{hg} \quad \text{for all } h, g \in \mathcal{P}.$$

As $(\bar{\pi}, \bar{t})$ is a Toeplitz representation we have that $Y_h Y_g \subseteq Y_{hg}$ for all $h, g \in \mathcal{P}$. For the reverse inclusion, let $p \in P$ with $\vartheta(p) = hg$, and we will show that $\bar{t}_p(X_p) \in [Y_h Y_g]$. Since ϑ is saturated there are $q, q' \in P$ such that

$$p = qq' \quad \text{and} \quad \vartheta(q)\mathcal{P} = h\mathcal{P}.$$

We can write $\vartheta(q) = hz$ for some $z \in \mathcal{P}^*$ and let $w \in P^*$ with $\vartheta(w) = z$ by condition (A3) of the saturation property. Since $\vartheta(q)\vartheta(q') = \vartheta(p) = hg$ it follows that $\vartheta(q') = z^{-1}g$. We thus conclude that

$$\vartheta(qw^{-1}) = h \quad \text{and} \quad \vartheta(wq') = g.$$

Recall that X_w satisfies $[\bar{t}_{w^{-1}}(X_{w^{-1}})\bar{t}_w(X_w)] = \bar{\pi}(A)$. By taking elementary vectors we get the required

$$\begin{aligned} \bar{t}_p(X_p) &= [\bar{t}_q(X_q)\bar{t}_{q'}(X_{q'})] = [\bar{t}_q(X_q)\bar{\pi}(A)\bar{t}_{q'}(X_{q'})] \\ &= [\bar{t}_q(X_q)\bar{t}_{w^{-1}}(X_{w^{-1}})\bar{t}_w(X_w)\bar{t}_{q'}(X_{q'})] \\ &\subseteq [\bar{t}_{qw^{-1}}(X_{qw^{-1}})\bar{t}_{wq'}(X_{wq'})] \subseteq [Y_{\vartheta(q)\vartheta(w^{-1})}Y_{\vartheta(w)\vartheta(q')}] = [Y_h Y_g]. \end{aligned}$$

Next we show that Y is compactly aligned. Let $h, h' \in \mathcal{P}$ and take $p \in \vartheta^{-1}(h)$ and $q \in \vartheta^{-1}(h')$. If $h \vee h' = \infty$ then $p \vee q = \infty$ as well for all $p \in \vartheta^{-1}(h)$ and $q \in \vartheta^{-1}(h')$, and so

$$Y_h^* Y_{h'} = \sum_{\substack{p \in \vartheta^{-1}(h) \\ q \in \vartheta^{-1}(h')}} \bar{t}_p(X_p)^* \bar{t}_q(X_q) = (0).$$

On the other hand, if $h \vee h' < \infty$ and $p \vee q < \infty$ for $p \in \vartheta^{-1}(h)$ and $q \in \vartheta^{-1}(h')$, then

$$\bar{t}_p(X_p)^* \bar{t}_q(X_q) \subseteq [\bar{t}_{p^{-1}w}(X_{p^{-1}w})\bar{t}_{q^{-1}w}(X_{q^{-1}w})^*].$$

Since $w = px = qy$ we have that $\vartheta(p^{-1}w) = h^{-1}\vartheta(w) = (h')^{-1}\vartheta(w) = \vartheta(q^{-1}w)$ and also $\vartheta(w) = h \vee h'$. Hence

$$Y_h^* Y_{h'} = \sum_{\substack{p \in \vartheta^{-1}(h) \\ q \in \vartheta^{-1}(h') \\ p \vee q < \infty}} \bar{t}_p(X_p)^* \bar{t}_q(X_q) \subseteq [Y_{h^{-1}(h \vee h')} Y_{h'(h \vee h')}].$$

Thus Proposition 4.7 gives that Y is compactly aligned.

By definition we have that $\mathcal{F}X \simeq \mathcal{F}Y$ (by grouping together summands with the same ϑ -image), and therefore we have that $\mathcal{T}_\lambda(X) \simeq \mathcal{T}_\lambda(Y)$ and that $\mathcal{T}_\lambda(X)^+ \simeq \mathcal{T}_\lambda(Y)^+$. Notice that these identifications are \mathcal{G} -compatible. By applying Proposition 5.9 and Theorem 4.14 we then get

$$A \times_{X,\lambda} P \simeq C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, G, \bar{\delta}_G) \simeq C_{\text{env}}^*(\mathcal{T}_\lambda(X)^+, \mathcal{G}, \bar{\delta}_\mathcal{G}) \simeq C_{\text{env}}^*(\mathcal{T}_\lambda(Y)^+, \mathcal{G}, \bar{\delta}_\mathcal{G}) \simeq A \times_{Y,\lambda} \mathcal{P}.$$

The second part of the proof is treated likewise. First note that any representation of X lifts to a representation of Y in a unique way, as every fiber of Y is spanned independently by the corresponding fibers of X . Applying similar arguments as above for a representation (π, t) in the place of the Fock representation we see that this correspondence preserves Nica-covariant representations. Hence we get that $\mathcal{NT}(X) \simeq \mathcal{NT}(Y)$.

Finally the $*$ -isomorphisms $A \times_{X,\lambda} P \simeq A \times_{Y,\lambda} \mathcal{P}$ gives an injective map

$$[A \times_X P]_{pq^{-1}} \simeq [A \times_{X,\lambda} P]_{pq^{-1}} \hookrightarrow [A \times_{Y,\lambda} \mathcal{P}]_{\vartheta(pq^{-1})} \simeq [A \times_Y \mathcal{P}]_{\vartheta(pq^{-1})}.$$

Therefore we get a commutative diagram

$$\begin{array}{ccc} \mathcal{NT}(X) & \xrightarrow{\Phi} & \mathcal{NT}(Y) \\ \downarrow q_X & & \downarrow q_Y \\ A \times_X P & \xrightarrow{\Psi} & A \times_Y \mathcal{P} \end{array}$$

where the upper horizontal arrow is a $*$ -isomorphism. Since the ideals of strong covariance relations are induced, it suffices to show that

$$\ker \Phi q_Y \cap [\mathcal{NT}(X)]_{e_G} \subseteq \ker q_X.$$

Equivalently it suffices to show that Ψ is faithful on the \mathcal{G} -fixed point algebra defined on $A \times_X P$. However this follows by Corollary 5.8 as Ψ is by definition faithful on A . □

Theorem 7.7 gives a very clear picture for the covariance algebras of a product system over a free product order of the form $(\ast_{i=1}^n G, \ast_{i=1}^n P)$ for an abelian total order (G, P) . It is well known that the Cuntz C^* -algebra \mathcal{O}_n for $n \in \mathbb{N}$ can be viewed as either the Nica–Cuntz–Pimsner C^* -algebra of the trivial product system over the free semigroup on n generators or as the Cuntz–Pimsner C^* -algebra of the C^* -correspondence $(\mathbb{C}^n, \mathbb{C})$. Our next result generalizes this fact to arbitrary product systems over the free semigroup.

Corollary 7.8. *Let X be a compactly aligned product system over the free semigroup $\mathbb{F}_n^+ = \langle i_1, \dots, i_n \rangle$. Then $A \times_X \mathbb{F}_n^+ \simeq \mathcal{O}_Y$ for the C^* -correspondence $Y = \sum_{j=1, \dots, n}^\oplus X_{i_j}$.*

Reversible product systems and total orders. An application of the theorem of Burns and Hale [1972] asserts that G admits a total order if and only if for every nontrivial finitely generated subgroup H of G there exists a totally ordered L and a nontrivial homomorphism $H \rightarrow L$. If $L = \mathbb{Z}$ then the group is called *left indicable*. There are plenty of abelian total orders. Examples include \mathbb{R}^2 with the lexicographical order and \mathbb{Z}^2 with the semigroup given by the half-plane defined by any line through the origin with irrational slope. Conrad’s theorem [1959] asserts that if (G, P) is a total order and G is Archimedean

then G embeds in \mathbb{R} so that P embeds in \mathbb{R}^+ . Here we say that G is *Archimedean* if whenever $e_G < x < y$, there exists an $n \in \mathbb{N}$ such that $y < x^n$. We refer the reader to [Clay and Rolfsen 2016] for an exposition of these results.

There are not many ways for a total order to be controlled by an *abelian* total order.

Proposition 7.9. *Let (G, P) be a total order and let $\vartheta_{\text{ab}} : (G, P) \rightarrow (G_{\text{ab}}, P_{\text{ab}})$ be the abelianization map. Then the following are equivalent:*

- (i) *There is a controlled map $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ where $(\mathcal{G}, \mathcal{P})$ is an abelian total order.*
- (ii) $\vartheta_{\text{ab}}^{-1}(0) \cap P = \{e_G\}$.
- (iii) *The abelianization map is a controlled map and $(G_{\text{ab}}, P_{\text{ab}})$ is a total order.*

If any (and thus all) of the above hold then the abelianization is a saturated controlled map.

Proof. If item (i) holds then ϑ factors through the abelianization. Since $\vartheta^{-1}(e_G) \cap P = \{e_G\}$ is a controlled map, we have that $\vartheta_{\text{ab}}^{-1}(0) \cap P = \{e_G\}$ as well.

Assume that item (ii) holds, and we will show that $(G_{\text{ab}}, P_{\text{ab}})$ is a total order. First we clearly have that

$$-P_{\text{ab}} \cup P_{\text{ab}} = \vartheta_{\text{ab}}(P^{-1} \cup P) = G_{\text{ab}}.$$

Next suppose that $-P_{\text{ab}} \cap P_{\text{ab}} \neq \{0\}$ so that there are $h, g \in P_{\text{ab}}$ with $h + g = 0$. As the abelianization map is surjective there are $p, q \in P$ with $pq = e_G$ with $\vartheta(p) = h$ and $\vartheta(q) = g$. As (G, P) is a total order we derive that $p = q = e_G$ and thus $h = g = 0$. Next we show that ϑ_{ab} satisfies conditions (A1) and (A2) of Definition 5.1. Let $p, q \in P$. Then either $p \leq q$ or $q \leq p$ and condition (A1) follows. For condition (A2) suppose without loss of generality that $p \leq q$ with $\vartheta_{\text{ab}}(p) = \vartheta_{\text{ab}}(q)$. Then $q = ps$ for $s \in P \cap \vartheta_{\text{ab}}^{-1}(0)$. Then $s = e_G$ and so $p = q$.

If item (iii) holds then clearly item (i) holds, concluding the equivalences between all items.

For the saturation property let a $t \in P$ and an $h \in P_{\text{ab}}$ such that $\vartheta_{\text{ab}}(t) = h + h'$. Take an $s \in \vartheta_{\text{ab}}^{-1}(h)$ since the abelianization map is surjective. Then either $s \leq t$ or $s > t$. But if $s > t$ then $h = \vartheta_{\text{ab}}(s) > \vartheta_{\text{ab}}(t)$ which is a contradiction. Thus we must have that $s \leq t$. □

Remark 7.10. There are exact total orders for which the abelianization map is not controlled. An example is given by the Klein bottle group

$$\mathbb{K} := \langle x, y \mid x^{-1}yx = y^{-1} \rangle = \langle x, y \mid x = yxy \rangle$$

with the total order induced by the semigroup \mathbb{K}^+ generated by x, y in \mathbb{K} . It is not hard to see that \mathbb{K}^+ induces a total order on \mathbb{K} , being left indicable (or since \mathbb{K} is the extension $\mathbb{Z} \rtimes \mathbb{Z}$). Alternatively one can see that every element in \mathbb{K} is written (uniquely) in the form $x^m y^n$ for $m, n \in \mathbb{Z}$ and we take cases: if $m, n \geq 0$ then $x^m y^n \in \mathbb{K}$; if $m \geq 1$ and $n \leq 0$ then we have that $x^m y^n = x^{m-1} y^{-n} x \in \mathbb{K}^+$; if $m = 0$ and $n \leq 0$ then $x^m y^n = y^n \in (\mathbb{K}^+)^{-1}$. By symmetry these cover all cases. We see that $\vartheta_{\text{ab}}(yxy) = \vartheta_{\text{ab}}(x)$ and so $e_{\mathbb{K}} \neq y^2 \in \mathbb{K}^+ \cap \vartheta_{\text{ab}}^{-1}(0)$. In fact we have that $\mathbb{K}_{\text{ab}} = \mathbb{Z} \times \mathbb{Z}_2$ and $\mathbb{K}_{\text{ab}}^+ = \mathbb{Z}^+ \times \mathbb{Z}_2$ and thus it does not define a total order as $-\mathbb{K}_{\text{ab}}^+ \cap \mathbb{K}_{\text{ab}}^+ = \mathbb{Z}_2$.

Definition 7.11. Let (G, P) be a total order and let X be a product system over P with coefficients in A . We say that X is a *reversible* product system if every X_p is a Hilbert bimodule in $A \times_{X,\lambda} P$, i.e., if $A \times_{X,\lambda} P = C^*(\pi, t)$ then $t_p(X_p)t_p(X_p)^* \subseteq A$ for all $p \in P$.

It follows that reversible product systems consist of Hilbert bimodules. The converse holds also for injective product systems, as in this case every strongly covariant representation is Katsura-covariant fiberwise.

Proposition 7.12. *Let (G, P) be a total order and let X be a product system over P with coefficients in A . Suppose that every X_p is injective. If (π, t) is a strongly covariant representation of X then (π, t_p) is a covariant representation of X_p , in the sense of Katsura, for every $p \in P$.*

Therefore an injective product system X is reversible if and only if every X_p is a Hilbert bimodule.

Proof. Fix $p \in P$ and $a \in A$ such that $\varphi_p(a) = k_p \in KX_p$. In view of strong covariance of Proposition 4.9 and Katsura covariance we have to show that

$$[\pi_F(a) + \psi_{p,F}(k_p)]_{X_F} = 0 \quad \text{for } F = \{e, p\},$$

where

$$X_F = \bigoplus_{r \in P} X_r I_{r^{-1}(r \vee F)}.$$

Let $r \in P$ with $r = ps$ for some $s \in P$. Then for every $\xi_r = \xi_p \xi_s \in X_r$ and $b \in I_{r^{-1}(r \vee F)}$ we have that

$$\pi_F(a)\xi_r b = (\varphi_p(a)\xi_p)\xi_s b = (k_p \xi_p)\xi_s b = \psi_{F,p}(k_p)\xi_r b.$$

Now suppose that $r < p$. Then by construction $\psi_{F,p}(k_p)\xi_r b = 0$ and we have to show that $\pi_F(a)\xi_r b = 0$ as well. To this end it suffices to show that

$$I_{r^{-1}(r \vee F)} := I_{r^{-1}K_{\{r,e\}}} \bigcap I_{r^{-1}K_{\{r,p\}}} = (0).$$

Since $r < p$ we have that $r \notin K_{\{r,p\}} \subseteq pP$ while $p \in K_{\{r,p\}}$. Therefore $r^{-1}p \neq e_G$ and so

$$I_{r^{-1}K_{\{r,p\}}} = \bigcap_{t \in K_{\{r,p\}}} \ker \varphi_{r^{-1}t} \subseteq \ker \varphi_{r^{-1}p} = (0),$$

and the proof is complete. □

In the case of $(G, P) = (\mathbb{Z}, \mathbb{Z}_+)$, the following result was established in [Kakariadis 2013].

Proposition 7.13. *Let (G, P) be a total order and let X be a product system over P with coefficients in A . Then X is a reversible product system if and only if the tensor algebra $\mathcal{T}_\lambda(X)^+$ is Dirichlet in $A \times_{X,\lambda} P$.*

Proof. Let (π, t) be a faithful representation of $A \times_{X,\lambda} P$. Suppose first that X is a reversible product system so that $t_p(X_p)t_p(X_p)^* \subseteq \pi(A)$ for all $p \in P$. We will show that

$$A \times_{X,\lambda} P = \overline{\text{span}}\{t_s(X_s) + t_r(X_r)^* \mid s, r \in P\}.$$

Let $s, r \in P$. If $rs^{-1} \in P$ then we have that

$$t_s(X_s)t_r(X_r)^* \subseteq [t_s(X_s)t_s(X_s)^*t_{rs^{-1}}(X_{rs^{-1}})^*] \subseteq [\pi(A)t_{rs^{-1}}(X_{rs^{-1}})^*] = t_{rs^{-1}}(X_{rs^{-1}})^*.$$

If $sr^{-1} \in P$ then we have that

$$t_s(X_s)t_r(X_r)^* \subseteq [t_{sr^{-1}}(X_{sr^{-1}})t_r(X_r)t_r(X_r)^*] \subseteq [t_{sr^{-1}}(X_{sr^{-1}})\pi(A)] = t_{sr^{-1}}(X_{sr^{-1}}).$$

Hence

$$A \times_{X,\lambda} P = \overline{\text{span}}\{t_s(X_s)t_r(X_r)^* \mid s, r \in P\} \subseteq \overline{\text{span}}\{t_s(X_s) + t_r(X_r)^* \mid s, r \in P\} \subseteq A \times_{X,\lambda} P,$$

and so $\mathcal{T}_\lambda(X)^+$ is Dirichlet in $A \times_{X,\lambda} P$.

Conversely, assume that $\mathcal{T}_\lambda(X)^+$ is Dirichlet in $A \times_{X,\lambda} P$ and let E be the conditional expectation induced by the coaction of G on $A \times_{X,\lambda} P$. Then $E(\mathcal{T}_\lambda(X)^+) = \pi(A)$ and

$$E(A \times_{X,\lambda} P) = E(\overline{(\mathcal{T}_\lambda(X)^+ + (\mathcal{T}_\lambda(X)^+)^*)}) = \pi(A).$$

Thus for each $p \in P$ we have that $t_p(X_p)t_p(X_p)^* \subseteq E(A \times_{X,\lambda} P) = \pi(A)$ as desired. □

The next corollary squares with the fact that Popescu’s noncommutative disc algebra is not Dirichlet. Recall that for abelian coactions the C^* -envelope of a cosystem coincides with the usual C^* -envelope of the ambient operator algebra.

Corollary 7.14. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a saturated controlled map between weak right LCM inclusions and suppose that $(\mathcal{G}, \mathcal{P})$ is an abelian total order. Let X be an injective product system over P with coefficients in A . Then $\mathcal{T}_\lambda(X)^+$ is Dirichlet if and only if every strongly covariant representation (π, t) of X satisfies $t_p(X_p)t_q(X_q)^* \subseteq A$ whenever $\vartheta(p) = \vartheta(q)$.*

Proof. By Theorem 6.1, and since the controlling pair is abelian, the C^* -envelope of $\mathcal{T}_\lambda(X)^+$ is $A \times_X P$. For the injective X , let Y be the injective product system over \mathcal{P} with coefficients in A constructed in Theorem 7.7. By construction we see that Y_h with $h \in \mathcal{P}$ is a Hilbert bimodule if and only if $t_p(X_p)t_q(X_q)^* \subseteq A$ for all $p, q \in \vartheta^{-1}(h)$. By applying Remark 2.8, Theorems 6.1 and 7.7, and Propositions 7.12 and 7.13 we have that the Fock tensor algebra $\mathcal{T}_\lambda(X)^+$ is Dirichlet in $A \times_X P$, if and only if $\mathcal{T}_\lambda(Y)^+$ is Dirichlet in $A \times_Y \mathcal{P}$, if and only if every Y_h with $h \in \mathcal{P}$ is a Hilbert bimodule, if and only if $t_p(X_p)t_q(X_q)^* \subseteq \pi(A)$ whenever $\vartheta(p) = \vartheta(q) = h$ for all $h \in \mathcal{P}$. □

The next theorem shows that, for weak right LCM inclusions that are controlled by total orders in a saturated way, reversible product systems produce all possible covariance algebras.

Theorem 7.15. *Let $\vartheta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ be a saturated controlled map between weak right LCM inclusions and suppose that $(\mathcal{G}, \mathcal{P})$ is a total order. Let X be an (injective) product system over P with coefficients in A . Then there exists an (injective) reversible product system Z over \mathcal{P} with coefficients in a C^* -algebra B such that*

$$A \subseteq B \quad \text{and} \quad X_p \subseteq Z_{\vartheta(p)} \quad \text{for all } p \in P \tag{7-1}$$

that satisfies

$$A \times_X P \simeq B \times_Z \mathcal{P} \quad \text{and} \quad A \times_{X,\lambda} P \simeq B \times_{Z,\lambda} \mathcal{P}, \tag{7-2}$$

by $*$ -homomorphisms that preserve the inclusions $X_p \hookrightarrow Z_{\vartheta(p)}$ for all $p \in P$.

Proof. By Theorem 7.7 we can assume that $(G, P) = (\mathcal{G}, \mathcal{P})$. Let (π, t) be a faithful representation of $A \times_{X,\lambda} P$, and let

$$B := B_P = C^*({t_s(X_s)t_s(X_s)^* \mid s \in P}) \quad \text{and} \quad Z_p := [t_p(X_p)B] \quad \text{for all } p \in P \setminus \{e\}.$$

The trivial C^* -correspondence structure on $A \times_{X,\lambda} P$ descends to a C^* -correspondence structure on each Z_p over B . Notice here that since (G, P) is totally ordered we automatically have that the product system $Z = \{Z_p\}_{p \in P}$ is compactly aligned. Also $C^*(B, Z) = C^*(\pi, t)$ admits a coaction of G from $A \times_{X,\lambda} P$. Hence by Theorem 4.13 we have that

$$\overline{\text{alg}\{B, Z_p \mid p \in P\}} \simeq \overline{\mathcal{T}_\lambda(Z)}^+.$$

By construction

$$A \times_{X,\lambda} P = \overline{\mathcal{T}_\lambda(Z)^+ + (\mathcal{T}_\lambda(Z)^+)^*},$$

thus the cosystem of $\mathcal{T}_\lambda(Z)^+$ over G is Dirichlet in a C^* -cover. This gives at the same time that this C^* -cover $A \times_{X,\lambda} P$ is the C^* -envelope of the cosystem $\mathcal{T}_\lambda(Z)^+$ over G , and that Z is reversible by Proposition 7.13. Theorem 4.14 then concludes that

$$B \times_{Z,\lambda} P \simeq C_{\text{env}}^*(\mathcal{T}_\lambda(Z)^+, G, \bar{\delta}_G) \simeq A \times_{X,\lambda} P.$$

For the case of the universal C^* -algebras we proceed as in Theorem 7.7. That is, first we notice that the $*$ -isomorphism between the reduced C^* -algebras implies an embedding of the Fell bundles

$$[A \times_X P]_{pq^{-1}} \simeq [A \times_{X,\lambda} P]_{pq^{-1}} \hookrightarrow [B \times_{Z,\lambda} P]_{pq^{-1}} \simeq [B \times_Z P]_{pq^{-1}}$$

which lifts to a $*$ -epimorphism $\Psi : A \times_X P \rightarrow B \times_Z P$. Since $X \subseteq Z$ we also have a $*$ -epimorphism at the level of the Nica–Toeplitz–Pimsner algebras and thus the diagram

$$\begin{array}{ccc} \mathcal{NT}(X) & \xrightarrow{\Phi} & \mathcal{NT}(Z) \\ \downarrow q_X & & \downarrow q_Z \\ A \times_X P & \xrightarrow{\Psi} & B \times_Z P \end{array}$$

is commutative, and fixes X . Since the ideals of strong covariance relations are induced, it suffices to show that

$$\ker \Phi q_Z \cap [\mathcal{NT}(X)]_e \subseteq \ker q_X.$$

Equivalently it suffices to show that Ψ is faithful on the G -fixed point algebra defined on $A \times_X P$, which by definition is B . However this follows by the property of $A \times_X P$ as $\Psi|_A$ is by construction faithful.

It is left to show that injectivity of X implies injectivity of Z . By Theorem 7.7 we can still assume that $(G, P) = (\mathcal{G}, \mathcal{P})$. To this end let $p \in P$ and $f \in \ker \varphi_p^Z$. We need to show that $f = 0$.

As B_{pP} is an ideal in B we have that $B = B_{\{s < p\}} + B_{pP}$, and let $f_1 \in B_{\{s < p\}}$ and $f_2 \in B_{pP}$ be such that $f = f_1 + f_2$. Let (e_i) be a contractive approximate identity of $\psi_p(\mathcal{K}X_p)$ so that

$$0 = f e_i = f_1 e_i + f_2 e_i.$$

However (e_i) is also a contractive approximate identity for B_{pP} and so

$$\lim_i f_1 e_i = - \lim_\lambda f_2 e_i = f_2.$$

By Nica-covariance $f_1 e_i \in B_p$ for all i , and so we have that $f_2 \in B_p$. Thus we can assume without loss of generality that $f \in B_{\{s \leq p\}}$. As $B_{\{s \leq p\}}$ is the inductive limit of B_F for $F = \{p_1 < p_2 < \dots < p_n = p\}$ we may assume that

$$f = \sum_{i=1}^n \psi_{p_i}(k_{p_i}) \quad \text{with } k_{p_i} \in \mathcal{K}X_{p_i} \text{ and } p_1 < p_2 < \dots < p_n = p.$$

Recall the representation (π_F, t_F) on

$$X_F = \bigoplus_{r \in P} X_r I_{r^{-1}(r \vee F)}^X,$$

and we will show that

$$\sum_{i=1}^n \psi_{F, p_i}(k_{p_i})|_{X_F} = 0.$$

As (π, t) is strongly covariant this will give that $f = 0$ by Proposition 4.9. For $r \geq p$ we have that $f \in \ker \varphi_p^Z \subseteq \ker \varphi_r^Z$, and for every $\eta_r \in X_r I_{r^{-1}(r \vee F)}^X$ we have that $t_r(\eta_r) \in t_r(X_r) \subseteq Z_r$. Hence

$$t_r \left(\sum_{i=1}^n i_{p_i}^r(k_{p_i})(\eta_r) \right) = \sum_{i=1}^n \psi_{p_i}(k_{p_i}) t_r(\eta_r) = f t_r(\eta_r) = 0.$$

As t is isometric we obtain

$$\sum_{i=1}^n \psi_{F, p_i}(k_{p_i})|_{X_r I_{r^{-1}(r \vee F)}^X} = \sum_{i=1}^n i_{p_i}^r(k_{p_i}) = 0, \quad \text{for } r \geq p. \tag{7-3}$$

On the other hand for $r < p$ we have that $r^{-1}p \neq e_G$, and so

$$I_{r^{-1}(r \vee F)}^X \subseteq I_{r^{-1}K_{\{r, p\}}}^X \subseteq \ker \varphi_{r^{-1}p}^X = (0).$$

Hence trivially

$$\sum_{i=1}^n \psi_{F, p_i}(k_{p_i})|_{X_r I_{r^{-1}(r \vee F)}^X} = 0, \quad \text{for } r < p. \tag{7-4}$$

By (7-3) and (7-4) we have that $\sum_{i=1}^n \psi_{p_i, F}(k_{p_i})|_{X_F} = 0$, and the proof is complete. \square

Acknowledgements

Kakariadis was partially supported by EPSRC grant EP/T02576X/1 and LMS grant 41908. Katsoulis was partially supported by NSF grant 2054781. Laca was partially supported by NSERC Discovery Grant #RGPIN-2017-04052. Li has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement no. 817597).

References

- [Abadie et al. 1998] B. Abadie, S. Eilers, and R. Exel, “Morita equivalence for crossed products by Hilbert C^* -bimodules”, *Trans. Amer. Math. Soc.* **350**:8 (1998), 3043–3054. MR Zbl
- [Arveson 1969] W. B. Arveson, “Subalgebras of C^* -algebras”, *Acta Math.* **123** (1969), 141–224. MR Zbl
- [Arveson 1989] W. Arveson, *Continuous analogues of Fock space*, Mem. Amer. Math. Soc. **409**, Amer. Math. Soc., Providence, RI, 1989. MR Zbl
- [Blecher and Le Merdy 2004] D. P. Blecher and C. Le Merdy, *Operator algebras and their modules: an operator space approach*, Lond. Math. Soc. Monogr. (N.S.) **30**, Oxford Univ. Press, 2004. MR Zbl
- [Brownlowe et al. 2018] N. Brownlowe, N. S. Larsen, and N. Stammeier, “ C^* -algebras of algebraic dynamical systems and right LCM semigroups”, *Indiana Univ. Math. J.* **67**:6 (2018), 2453–2486. MR Zbl
- [Burns and Hale 1972] R. G. Burns and V. W. D. Hale, “A note on group rings of certain torsion-free groups”, *Canad. Math. Bull.* **15** (1972), 441–445. MR Zbl
- [Carlsen et al. 2011] T. M. Carlsen, N. S. Larsen, A. Sims, and S. T. Vittadello, “Co-universal algebras associated to product systems, and gauge-invariant uniqueness theorems”, *Proc. Lond. Math. Soc.* (3) **103**:4 (2011), 563–600. MR Zbl
- [Clay and Rolfsen 2016] A. Clay and D. Rolfsen, *Ordered groups and topology*, Grad. Studies in Math. **176**, Amer. Math. Soc., Providence, RI, 2016. MR Zbl
- [Conrad 1959] P. Conrad, “Right-ordered groups”, *Michigan Math. J.* **6** (1959), 267–275. MR Zbl
- [Crisp 1999] J. Crisp, “Injective maps between Artin groups”, pp. 119–137 in *Geometric group theory down under* (Canberra, 1996), edited by J. Cossey et al., de Gruyter, Berlin, 1999. MR Zbl
- [Crisp and Laca 2002] J. Crisp and M. Laca, “On the Toeplitz algebras of right-angled and finite-type Artin groups”, *J. Aust. Math. Soc.* **72**:2 (2002), 223–245. MR Zbl
- [Crisp and Laca 2007] J. Crisp and M. Laca, “Boundary quotients and ideals of Toeplitz C^* -algebras of Artin groups”, *J. Funct. Anal.* **242**:1 (2007), 127–156. MR Zbl
- [Dinh 1991] H. T. Dinh, “Discrete product systems and their C^* -algebras”, *J. Funct. Anal.* **102**:1 (1991), 1–34. MR Zbl
- [Dor-On and Katsoulis 2020] A. Dor-On and E. Katsoulis, “Tensor algebras of product systems and their C^* -envelopes”, *J. Funct. Anal.* **278**:7 (2020), art. id. 108416. MR Zbl
- [Dor-On et al. 2022] A. Dor-On, E. T. A. Kakariadis, E. Katsoulis, M. Laca, and X. Li, “ C^* -envelopes for operator algebras with a coaction and co-universal C^* -algebras for product systems”, *Adv. Math.* **400** (2022), art. id. 108286. MR Zbl
- [Dritschel and McCullough 2005] M. A. Dritschel and S. A. McCullough, “Boundary representations for families of representations of operator algebras and spaces”, *J. Operator Theory* **53**:1 (2005), 159–167. MR Zbl
- [Exel 1997] R. Exel, “Amenability for Fell bundles”, *J. Reine Angew. Math.* **492** (1997), 41–73. MR Zbl
- [Exel 2017] R. Exel, *Partial dynamical systems, Fell bundles and applications*, Math. Surv. Monogr. **224**, Amer. Math. Soc., Providence, RI, 2017. MR Zbl
- [Fowler 1999] N. J. Fowler, “Compactly-aligned discrete product systems, and generalization of \mathcal{O}_∞ ”, *Int. J. Math.* **10**:6 (1999), 721–738. MR Zbl
- [Fowler 2002] N. J. Fowler, “Discrete product systems of Hilbert bimodules”, *Pacific J. Math.* **204**:2 (2002), 335–375. MR Zbl
- [Hamana 1979] M. Hamana, “Injective envelopes of operator systems”, *Publ. Res. Inst. Math. Sci.* **15**:3 (1979), 773–785. MR Zbl
- [an Huef et al. 2021] A. an Huef, B. Nucinkis, C. F. Sehnm, and D. Yang, “Nuclearity of semigroup C^* -algebras”, *J. Funct. Anal.* **280**:2 (2021), art. id. 108793. MR Zbl
- [Kakariadis 2013] E. T. A. Kakariadis, “The Dirichlet property for tensor algebras”, *Bull. Lond. Math. Soc.* **45**:6 (2013), 1119–1130. MR Zbl
- [Kakariadis 2020] E. T. A. Kakariadis, “Finite-dimensional approximations for Nica–Pimsner algebras”, *Ergodic Theory Dynam. Systems* **40**:12 (2020), 3375–3402. MR Zbl
- [Kakariadis and Katsoulis 2012] E. T. A. Kakariadis and E. G. Katsoulis, “Contributions to the theory of C^* -correspondences with applications to multivariable dynamics”, *Trans. Amer. Math. Soc.* **364**:12 (2012), 6605–6630. MR Zbl

- [Kakariadis et al. 2022] E. T. A. Kakariadis, E. G. Katsoulis, M. Laca, and X. Li, “Boundary quotient C^* -algebras of semigroups”, *J. Lond. Math. Soc. (2)* **105**:4 (2022), 2136–2166. MR
- [Katsoulis 2020] E. Katsoulis, “Product systems of C^* -correspondences and Takai duality”, *Israel J. Math.* **240**:1 (2020), 223–251. MR Zbl
- [Katsoulis and Ramsey 2019] E. G. Katsoulis and C. Ramsey, *Crossed products of operator algebras*, Mem. Amer. Math. Soc. **1240**, Amer. Math. Soc., Providence, RI, 2019. MR Zbl
- [Katsura 2004] T. Katsura, “On C^* -algebras associated with C^* -correspondences”, *J. Funct. Anal.* **217**:2 (2004), 366–401. MR Zbl
- [Kwaśniewski and Larsen 2019] B. K. Kwaśniewski and N. S. Larsen, “Nica–Toeplitz algebras associated with product systems over right LCM semigroups”, *J. Math. Anal. Appl.* **470**:1 (2019), 532–570. MR Zbl
- [Laca and Raeburn 1996] M. Laca and I. Raeburn, “Semigroup crossed products and the Toeplitz algebras of nonabelian groups”, *J. Funct. Anal.* **139**:2 (1996), 415–440. MR Zbl
- [Lawson 2012] M. V. Lawson, “Non-commutative Stone duality: inverse semigroups, topological groupoids and C^* -algebras”, *Int. J. Algebra Comput.* **22**:6 (2012), art. id. 1250058. MR Zbl
- [Li 2013] X. Li, “Nuclearity of semigroup C^* -algebras and the connection to amenability”, *Adv. Math.* **244** (2013), 626–662. MR Zbl
- [Li 2017] X. Li, “Partial transformation groupoids attached to graphs and semigroups”, *Int. Math. Res. Not.* **2017**:17 (2017), 5233–5259. MR Zbl
- [Meyer and Sehnem 2019] R. Meyer and C. F. Sehnem, “A bicategorical interpretation for relative Cuntz–Pimsner algebras”, *Math. Scand.* **125**:1 (2019), 84–112. MR Zbl
- [Nica 1992] A. Nica, “ C^* -algebras generated by isometries and Wiener–Hopf operators”, *J. Operator Theory* **27**:1 (1992), 17–52. MR Zbl
- [Norling 2014] M. D. Norling, “Inverse semigroup C^* -algebras associated with left cancellative semigroups”, *Proc. Edinb. Math. Soc. (2)* **57**:2 (2014), 533–564. MR Zbl
- [Paulsen 2002] V. Paulsen, *Completely bounded maps and operator algebras*, Cambridge Stud. Adv. Math. **78**, Cambridge Univ. Press, 2002. MR Zbl
- [Pimsner 1997] M. V. Pimsner, “A class of C^* -algebras generalizing both Cuntz–Krieger algebras and crossed products by \mathbb{Z} ”, pp. 189–212 in *Free probability theory* (Waterloo, ON, 1995), edited by D.-V. Voiculescu, Fields Inst. Commun. **12**, Amer. Math. Soc., Providence, RI, 1997. MR Zbl
- [Quigg 1996] J. C. Quigg, “Discrete C^* -coactions and C^* -algebraic bundles”, *J. Austral. Math. Soc. Ser. A* **60**:2 (1996), 204–221. MR Zbl
- [Schweizer 2001] J. Schweizer, “Dilations of C^* -correspondences and the simplicity of Cuntz–Pimsner algebras”, *J. Funct. Anal.* **180**:2 (2001), 404–425. MR Zbl
- [Sehnem 2019] C. F. Sehnem, “On C^* -algebras associated to product systems”, *J. Funct. Anal.* **277**:2 (2019), 558–593. MR Zbl
- [Sims and Yeend 2010] A. Sims and T. Yeend, “ C^* -algebras associated to product systems of Hilbert bimodules”, *J. Operator Theory* **64**:2 (2010), 349–376. MR Zbl
- [Spielberg 2012] J. Spielberg, “ C^* -algebras for categories of paths associated to the Baumslag–Solitar groups”, *J. Lond. Math. Soc. (2)* **86**:3 (2012), 728–754. MR Zbl

Received 20 Aug 2021. Accepted 20 Dec 2021.

EVGENIOS T.A. KAKARIADIS: evgenios.kakariadis@ncl.ac.uk
School of Mathematics, Statistics and Physics, Newcastle University, Newcastle upon Tyne, United Kingdom

ELIAS G. KATSOULIS: katsoulise@ecu.edu
Department of Mathematics, East Carolina University, Greenville, NC, United States

MARCELO LACA: laca@uvic.ca
Department of Mathematics and Statistics, University of Victoria, Victoria, BC, Canada

XIN LI: xin.li@glasgow.ac.uk
School of Mathematics and Statistics, University of Glasgow, Glasgow, United Kingdom

Analysis & PDE

msp.org/apde

EDITORS-IN-CHIEF

Patrick Gérard Université Paris Sud XI, France
patrick.gerard@universite-paris-saclay.fr

Clément Mouhot Cambridge University, UK
c.mouhot@dpms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Błocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	András Vasy	Stanford University, USA andras@math.stanford.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2023 is US \$405/year for the electronic version, and \$630/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 16 No. 6 2023

The Cauchy problem for the infinitesimal model in the regime of small variance FLORIAN PATOUT	1289
The “good” Boussinesq equation: long-time asymptotics CHRISTOPHE CHARLIER, JONATAN LENELLS and DENG-SHAN WANG	1351
Quantitative Obata’s theorem FABIO CAVALLETTI, ANDREA MONDINO and DANIELE SEMOLA	1389
Couniversality and controlled maps on product systems over right LCM semigroups EVGENIOS T.A. KAKARIADIS, ELIAS G. KATSOULIS, MARCELO LACA and XIN LI	1433