

ANALYSIS & PDE

Volume 17

No. 1

2024

TOKE MEIER CARLSEN, ADAM DOR-ON AND SØREN EILERS

**SHIFT EQUIVALENCES
THROUGH THE LENS OF CUNTZ-KRIEGER ALGEBRAS**

SHIFT EQUIVALENCES THROUGH THE LENS OF CUNTZ–KRIEGER ALGEBRAS

TOKE MEIER CARLSEN, ADAM DOR-ON AND SØREN EILERS

Motivated by Williams’ problem of measuring novel differences between shift equivalence (SE) and strong shift equivalence (SSE), we introduce three equivalence relations that provide new ways to obstruct SSE while merely assuming SE.

Our shift equivalence relations arise from studying graph C^* -algebras, where a variety of intermediary equivalence relations naturally arise. As a consequence we realize a goal sought after by Muhly, Pask and Tomforde, measure a delicate difference between SSE and SE in terms of Pimsner dilations for C^* -correspondences of adjacency matrices, and use this distinction to refute a proof from a previous paper.

1. Introduction

Initially recognized in the 40’s as the right object to model quantum phenomena, C^* -algebras are applied today in a variety of areas including theoretical physics, topology, differential geometry and dynamical systems. Such applications drive the impetus for obtaining structural and classification results for C^* -algebras, especially in relation with Elliott’s classification programme [14; 45; 49; 50]. One fantastic application of C^* -algebras in dynamics, using tools from classification of C^* -algebras, is the classification of Cantor minimal systems up to orbit equivalence by their dimension groups [17; 18]. Similar to this, our work here deals with subtle invariants arising from C^* -algebras associated to subshifts of finite type (SFTs), with the aim of distinguishing SFTs up to conjugacy.

In a seminal 1973 paper [47], Williams recast conjugacy and eventual conjugacy for SFTs purely in terms of equivalence relations between adjacency matrices of the directed graphs. These are called strong shift equivalence (SSE) and shift equivalence (SE) respectively. Williams expected SSE and SE to be the same [47, Proposition 7.2], but after around 25 years the last hope for a positive answer to Williams’ problem, even under the most restrictive conditions, was extinguished by Kim and Roush [28]. Although these counterexamples are concrete, aperiodic and irreducible 7×7 matrices, showing that they are not strong shift equivalent requires an invariant which is very difficult to compute. Thus, finding new

Toke M. Carlsen was supported by the Research Council of the Faroe Islands. Adam Dor-On was supported by NSF grant DMS-1900916 and the European Union’s Horizon 2020 Marie Skłodowska-Curie grant 839412-IRIOA. Søren Eilers was supported by the DFF-Research Project 2 “Automorphisms and invariants of operator algebras”, no. 7014-00145B. Finally, this project was partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy-EXC 2044-390685587, Mathematics Münster: Dynamics-Geometry-Structure; the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Project-ID 427320536-SFB 1442, and ERC Advanced Grant 834267-AMAREC.

MSC2020: primary 37A55, 46L55; secondary 37B10, 37A35, 46L08, 46L35.

Keywords: shift equivalence, Williams’ problem, Cuntz–Krieger algebras, Cuntz–Pimsner algebras, compatible shift equivalence, Pimsner dilations.

obstructions to strong shift equivalence when two matrices are only assumed to be shift equivalent is an important endeavor, even just for 2×2 matrices (see [31, Example 7.3.13]).

Definition 1.1 [47]. Let A and B be matrices indexed by sets V and W respectively, with (possibly infinite) cardinal entries. We say that A and B are:

- (1) *Shift equivalent* with lag $m \in \mathbb{N} \setminus \{0\}$ if there are a $V \times W$ matrix R and a $W \times V$ matrix S with cardinal entries such that

$$\begin{aligned} A^m &= RS, & B^m &= SR, \\ SA &= BS, & AR &= RB. \end{aligned}$$

- (2) *Elementary shift related* if they are shift equivalent with lag 1.

- (3) *Strong shift equivalent* if they are equivalent in the transitive closure of elementary shift relation.

In tandem with early attacks on Williams' problem, Cuntz and Krieger [8] created a bridgehead between symbolic dynamics and operator algebras, where several natural properties of subshifts of finite type are expressed through associated C^* -algebras. In fact, by [8, Proposition 2.17] we know that strong shift equivalence of A and B implies that the Cuntz–Krieger C^* -algebras \mathcal{O}_A and \mathcal{O}_B are stably isomorphic in a way preserving their gauge actions γ^A and γ^B and their diagonal subalgebras \mathcal{D}_A and \mathcal{D}_B . On the other hand, by a theorem of Krieger [29] we know that the dimension group triples of SFTs are isomorphic if and only if the associated matrices are SE. Since these dimension group triples coincide with K -theoretical data of crossed products of Cuntz–Krieger C^* -algebras by their gauge action, Krieger's theorem implies as a corollary (see Section 7) that if two Cuntz–Krieger algebras \mathcal{O}_A and \mathcal{O}_B are stably isomorphic in a way preserving their gauge actions γ^A and γ^B , then their defining adjacency matrices A and B are shift equivalent. Through the lens of Cuntz–Krieger algebras, this provides several natural equivalence relations between SSE and SE, and it then becomes important to orient them and determine whether they coincide with SSE, SE or perhaps a completely new equivalence relation strictly between SSE and SE. Such distinctions may pave the way towards more concrete and computable invariants that distinguish SFTs up to conjugacy.

In this paper we introduce, study and orient three equivalence relations that provide new ways of measuring the difference between SSE and SE. Before we discuss these, let us first mention the state of the art.

A partial converse of the corollary to Krieger's theorem was obtained by Bratteli and Kishimoto [4], using deep machinery from C^* -algebra K -theory classification, involving the essential concept of Rokhlin towers. More precisely, using their work it can be shown that if A and B are two *aperiodic and irreducible* adjacency matrices then

$$A \text{ is SE to } B \iff (\mathcal{O}_A \otimes \mathbb{K}, \gamma^A \otimes \text{id}) \simeq (\mathcal{O}_B \otimes \mathbb{K}, \gamma^B \otimes \text{id}). \quad (1-1)$$

This *converse to Krieger's corollary* for essential matrices remains a subtle and important classification problem in this line of research, and is one of the key motivating problems for our work here.

After a major undertaking pioneered by Matsumoto [33; 34], it is now known that several key concepts in symbolic dynamics may be fully understood in terms of operator algebraic descriptions. Indeed, both

SSE as well as flow equivalence (FE) of subshifts of finite type may be given an inherently operator-algebraic characterization. Suppose A and B are finite adjacency matrices defining two-sided SFTs (X_A, σ_A) and (X_B, σ_B) respectively (see Section 2). When A and B are irreducible, Matsumoto and Matui [35] established that

$$(X_A, \sigma_A) \text{ is FE to } (X_B, \sigma_B) \iff (\mathcal{O}_A \otimes \mathbb{K}, \mathcal{D}_A \otimes c_0) \simeq (\mathcal{O}_B \otimes \mathbb{K}, \mathcal{D}_B \otimes c_0),$$

where \mathbb{K} is the C^* -algebra of compact operators on $\ell^2(\mathbb{N})$ and c_0 is its subalgebra of diagonal operators. This was later extended to cover all two-sided SFTs by the first-named and third-named authors with Ortega and Restorff [6].

For conjugacy, the first-named author and Rout [5] proved that (X_A, σ_A) is conjugate to (X_B, σ_B) precisely when the associated Cuntz–Krieger algebras \mathcal{O}_A and \mathcal{O}_B are stably isomorphic in a way preserving both the gauge actions and the diagonals (this time with no additional restrictions on A and B). More precisely, and combining with Williams’ characterization of conjugacy, we have

$$A \text{ is SSE to } B \iff (\mathcal{O}_A \otimes \mathbb{K}, \gamma^A \otimes \text{id}, \mathcal{D}_A \otimes c_0) \simeq (\mathcal{O}_B \otimes \mathbb{K}, \gamma^B \otimes \text{id}, \mathcal{D}_B \otimes c_0). \quad (1-2)$$

The equivalences in (1-1) and (1-2) provide a fresh perspective for symbolic dynamics via C^* -algebras, and reinterpret Williams’ problem via the counterexamples provided by Kim and Roush [27; 28]. More precisely, the examples of Kim and Roush show that two adjacency matrices can be shift equivalent and with flow equivalent two-sided SFTs without being strong shift equivalent. Hence, in terms of C^* -algebras, this shows it is not always possible to trade in two isomorphisms — one respecting the diagonal, and one respecting the gauge action — for one which respects both.

Another research agenda motivating this work is the *graded isomorphism problem* of Hazrat from the theory of Leavitt path algebras, where one studies graded isomorphisms of pure algebras by means of graded K -theory. In work of Hazrat [19], it was shown that the relevant graded K -theoretical data is in fact the same as Krieger’s dimension group triple, and Hazrat conjectured that this invariant is complete when the class of the unit is added to the dimension triple as part of the invariant. Thus, the converse to Krieger’s corollary is a topological analogue of Hazrat’s conjecture. Although substantial advances have been made [2; 3], Hazrat’s conjecture remains elusive.

For a $V \times W$ matrix $F = [F_{ij}]$ with cardinal entries, we define

$$E_F := \{(v, w, \alpha) \mid 0 \leq \alpha < F_{vw}, v \in V, w \in W\},$$

so that $r(v, w, \alpha) = w$ and $s(v, w, \alpha) = v$, α is an ordinal, and F_{vw} is interpreted as the least ordinal with cardinality F_{vw} . When $V = W$, this makes $G_F := (V, E_F)$ into a directed graph in its own right. For two matrices C over $V \times W$ and D over $W \times X$ with cardinal entries, we define the *fibred product*

$$E_C \times E_D := \{cd \mid c \in E_C, d \in E_D, r(c) = s(d)\}.$$

Note here that we write cd to mean the pair (c, d) for which $r(c) = s(d)$, which should be thought of as concatenation of edges, even if there are no actual graphs. When $V = W$, for any $n \in \mathbb{N}$ we denote by E_C^n the n -fold product of E_C with itself, so that E_C^n is naturally identified with E_{C^n} .

One might also consider $E_C \times E_D$ as a pullback of W , and indicate the role of W in the notation. We refrain from doing so for the sake of readability.

The following definition was dubbed “specified equivalence” by Nasu [38] in his study of shift equivalences between textile systems.

Definition 1.2. Let A and B be matrices with cardinal entries over $V \times W$. A *path isomorphism* is a bijection $\phi : E_A \rightarrow E_B$ such that $s(\phi(e)) = s(e)$ and $r(\phi(e)) = r(e)$ for every $e \in E_A$.

Using path isomorphism, we define our first equivalence relation of *compatible shift equivalence* (CSE) (see Definition 4.1). The mere existence of path isomorphisms $\phi_R, \phi_S, \psi_A, \psi_B$ in Definition 4.1 implies shift equivalence, so that CSE is a direct strengthening of shift equivalence. In Theorem 7.3 we show that CSE coincides with SSE, and this allows for a direct comparison between SSE and SE via CSE. Motivated by Williams’ problem, another related notion called *adapted shift equivalence* was studied by Parry [39] and was also shown to be equivalent to SSE. More precisely, instead of a requirement on compatibility of path isomorphisms between shift equivalent matrices, adapted shift equivalence is a shift equivalence of lag $m \in \mathbb{N} \setminus \{0\}$ between the adjacency matrices of the m -line graphs of A and B .

It is known that shift equivalence is decidable by Kim and Roush [25; 26], but the problem of decidability of strong shift equivalence remains a fundamental open problem in symbolic dynamics. In fact, this was the original motivation in Williams’ paper [47]. The identification between path isomorphisms in the definition of CSE illustrates in what way the algorithm of Kim and Roush would need to improve were SSE turn out to be decidable.

Our second equivalence relation is called *representable shift equivalence* (RSE) (see Definition 5.1). It arises naturally when one attempts to represent shift equivalence as bounded operators on Hilbert space.

Surprisingly, by merely representing the relations of shift equivalence as bounded operator on Hilbert space, we get SSE (see Theorem 7.3). Considering the counterexamples of Kim and Roush, we see that if A and B are shift equivalent but not strong shift equivalent, then it follows that one of the four relations of shift equivalence must fail when representing everything on the same Hilbert space (see Section 5 for more details).

In what follows, we will say that a matrix is *essential* if it has no zero rows and no zero columns. In the work of Pimsner [40], Pimsner dilations were introduced and were subsequently studied by several authors [13; 15; 22; 37]. Pimsner dilations offer a “reversible” perspective for Cuntz–Pimsner C^* -algebras, showing that they are always generated by an imprimitivity bimodule that contains the original C^* -correspondence. In [37, Remark 5.5] an equivalent formulation of strong shift equivalence in terms of Pimsner dilations was sought. This leads us to our third equivalence relation, which we call *strong Morita shift equivalence* (SMSE) (see Definition 6.1). Let A and B be finite essential matrices. Denote by $X(A)_\infty$ and $X(B)_\infty$ the Pimsner dilations of their graph C^* -correspondences (see Section 3) and by $C^*(G_A)$ and $C^*(G_B)$ their Cuntz–Krieger graph C^* -algebras. Our main theorem provides an equivalent formulation sought by Muhly, Pask and Tomforde, and orients CSE and RSE in one fell swoop (see Theorem 7.3).

Theorem 1.3. *Suppose A and B are two finite essential matrices with entries in \mathbb{N} . Then the following are equivalent:*

- (1) A and B are strong shift equivalent.
- (2) A and B are compatible shift equivalent.
- (3) A and B are representable shift equivalent.
- (4) $X(A)_\infty$ and $X(B)_\infty$ are strong Morita shift equivalent.
- (5) $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic in a way that respects the diagonals.

For some time now, several experts have been perplexed about certain Pimsner dilation techniques (see for instance [37, Remark 5.5], the incorrect proof of [22, Theorem 5.8], and the subsequent corrigendum of [22]), and many are still wondering whether they can be used to show that shift equivalence of A and B implies strong Morita equivalence of the Pimsner dilations $X(A)_\infty$ and $X(B)_\infty$ (in the sense of Abadie, Eilers and Exel [1, Section 4] or Muhly and Solel [36]). The importance of this question is further elevated because of Theorem 3.17 and the discussion succeeding it, where we show that a positive answer to it is *equivalent* to the *converse to Krieger’s corollary*. Combining our main theorem with the celebrated counterexamples of Kim and Roush [28], as well as the work of Bratteli and Kishimoto [4], we obtain the following cutoff result. This result addresses the ambiguities mentioned above, and refutes the proof of [22, Theorem 5.8] (see Theorem 7.4).

Theorem 1.4. *There exist finite aperiodic 7×7 irreducible matrices A and B with entries in \mathbb{N} such that $X(A)_\infty$ and $X(B)_\infty$ are strong Morita equivalent, but not strong Morita shift equivalent.*

Since the validity of [22, Theorem 5.8] is in question, so is the validity of [22, Corollary 5.11]. This latter result states that shift equivalence of A and B implies the (not necessarily equivariant) stable isomorphisms of $C^*(G_A)$ and $C^*(G_B)$. However, thanks to Ara, Hazrat and Li [3] this result can still be recovered. Indeed, in work of the third author with Restorff, Ruiz and Sørensen [12] it was shown that filtered, ordered K-theory classifies unital graph C*-algebras up to stable isomorphism. By showing that filtered ordered K-theory is an invariant of shift equivalence, Ara, Hazrat and Li [3] ipso facto show that shift equivalence implies the stable isomorphisms of unital graph C*-algebras.

This paper contains eight sections, including this introductory section. In Section 2 we discuss some of the basic theory of directed graphs, subshifts of finite type, groupoid C*-algebra description of graph C*-algebras and Cuntz–Krieger C*-algebras. In Section 3 we discuss some of the theory of C*-correspondences, Cuntz–Pimsner algebras, Pimsner dilations and equivariant isomorphisms. We provide there a characterization of the existence of an equivariant stable isomorphism between Cuntz–Pimsner algebras in terms of Pimsner dilations. In Section 4 we introduce CSE, show it is an equivalence relation, and that it is implied by SSE. In Section 5 we introduce RSE, show that it is implied by CSE, and upgrade representations to be faithful on associated graph C*-algebras. In Section 6 we explain how to concretely construct inductive limits related to Pimsner dilations, introduce SMSE, show that RSE implies SMSE and show that SMSE implies the existence of a stable equivariant diagonal-preserving isomorphism between graph C*-algebras. Finally in Section 7 we discuss dimension triples for graph C*-algebras, orient different equivalence relations on adjacency matrices and prove the main results stated above.

2. Preliminaries

In this section we explain some of the basic theory of directed graphs, subshifts of finite type, groupoid C^* -algebra descriptions of graph C^* -algebras, Cuntz–Krieger algebras and dimension triples. We recommend [31] for the basics of symbolic dynamics.

A directed graph $G = (V, E, r, s)$ is comprised of a vertex set V and an edge set E together with range and source maps $r, s : E \rightarrow V$.

We say that a directed graph $G = (V, E)$ has *finite out-degrees* if $s^{-1}(v)$ is finite for every $v \in V$. We say that G is *finite* if both V and E are finite. When G has finite out-degrees and has no sources (i.e., r is surjective) and no sinks (i.e., s is surjective), we may define the *two-sided edge shift* to be the pair (X_E, σ_E) where X_E is the set of bi-infinite paths

$$X_E = \{(e_n)_{n \in \mathbb{Z}} \mid s(e_{i+1}) = r(e_i)\} \subseteq \prod_{i \in \mathbb{Z}} E,$$

with the product topology, and $\sigma_E : X_E \rightarrow X_E$ is the left shift homeomorphism given by $\sigma_E((e_n)_{n \in \mathbb{Z}}) = (e_{n+1})_{n \in \mathbb{Z}}$. The *one-sided edge shift* (X_E^+, σ_E^+) is defined as above, by replacing every occurrence of \mathbb{Z} with \mathbb{N} .

Given a directed graph $G = (V, E)$, we may always form its $V \times V$ adjacency matrix with cardinal entries given for $v, w \in V$ by

$$A_E(v, w) = |\{e \in E \mid s(e) = v, r(e) = w\}|.$$

Conversely, we have seen that given a matrix A indexed by V with cardinal entries, one may form a directed graph $G_A = (V, E_A, r, s)$, where E_A is the set of triples (v, w, α) such that $v, w \in V$, $0 \leq \alpha < A_{vw}$ is an ordinal, A_{vw} is interpreted as the least ordinal with cardinality A_{vw} , while $r(v, w, \alpha) = w$ and $s(v, w, \alpha) = v$. It is clear that G_{A_E} and G are isomorphic directed graphs and that $A_{E_A} = A$.

The following shows that under countability/finiteness assumptions, shift equivalence with arbitrary cardinals becomes the standard notion we know from the literature.

Proposition 2.1. *Let A and B be matrices with cardinal entries, indexed by sets V and W . Suppose that V and W are countable/finite, and suppose that A and B are over $\mathbb{N} \cup \{\aleph_0\}$ / over \mathbb{N} , respectively. If A and B are shift equivalent, then the matrices R and S realizing shift equivalence can be chosen to be with entries in $\mathbb{N} \cup \{\aleph_0\}$ / in \mathbb{N} , respectively.*

Proof. Suppose now that V and W are countable/finite, and that A and B are with entries in $\mathbb{N} \cup \{\aleph_0\}$ / in \mathbb{N} respectively. If R and S implement shift equivalence of lag m between A and B , denote by R' and S' the matrices obtained from R and S by replacing all noncountable/nonfinite entries with zeros (respectively). As the matrices A^m and B^m are with entries in $\mathbb{N} \cup \{\aleph_0\}$ / in \mathbb{N} (respectively), we still have that $R'S' = A^m$ and $S'R' = B^m$, as well as $S'A = BS'$ and $AR' = R'B$. Hence, A and B are shift equivalent via R' and S' , and we are done. \square

In this paper we will conduct our study through the lens of graph C^* -algebras, which include the class of Cuntz–Krieger C^* -algebras. We recommend [16; 41] for more on graph C^* -algebras. We will sometimes assume that our graphs have finite out-degree, which is often called “row-finiteness” in the literature.

Definition 2.2. Let $G = (V, E, r, s)$ be a directed graph. A family of operators $(S_v, S_e)_{v \in V, e \in E}$ on a Hilbert space \mathcal{H} is called a *Cuntz–Krieger family* if

- (1) $(S_v)_{v \in V}$ is a family of pairwise orthogonal projections,
- (2) $S_e^* S_e = S_{r(e)}$ for all $e \in E$, and
- (3) $\sum_{e \in s^{-1}(v)} S_e S_e^* = S_v$ for all $v \in V$ with $0 < |s^{-1}(v)| < \infty$.

The graph C^* -algebra $C^*(G)$ of G is the universal C^* -algebra generated by a Cuntz–Krieger families.

Universality of $C^*(G)$ gives rise to a point-norm continuous *gauge action* of the unit circle $\gamma : \mathbb{T} \rightarrow \text{Aut}(C^*(G))$ given by

$$\gamma_z(S_v) = S_v \quad \text{and} \quad \gamma_z(S_e) = z \cdot S_e \quad \text{for } z \in \mathbb{T}, v \in V, e \in E.$$

With this gauge action, $C^*(G)$ becomes a \mathbb{Z} -graded C^* -algebra whose graded components are $C^*(G)_n := \{T \in C^*(G) \mid \gamma_z(T) = z^n \cdot T\}$.

We will sometime assume that our graphs have finite out-degree, which is often called “row-finiteness” in the literature. We shall need the groupoid C^* -algebra description of $C^*(G)$, as specified in [5]. Indeed, if $G = (V, E)$ is a directed graph with finite out-degree, no sources and no sinks, we may construct the locally compact, Hausdorff étale groupoid

$$\mathcal{G}_E := \{(x, m - n, y) \in X_E^+ \times \mathbb{Z} \times X_E^+ \mid x, y \in X_E, m, n \in \mathbb{N}, \sigma_E^m(x) = \sigma_E^n(y)\},$$

with product $(x, k, y)(w, \ell, z) = (x, k + \ell, z)$ if $y = w$ (otherwise undefined), and inverse $(x, k, y)^{-1} = (y, -k, x)$. The topology on \mathcal{G}_E is generated by subsets of the form

$$Z(U, m, n, V) = \{(x, m - n, y) \in \mathcal{G}_E \mid x \in U, y \in V\},$$

where $m, n \in \mathbb{N}$ and $U, V \subseteq X_E^+$ are clopen such that $\sigma_E^m|_U$ is injective, $\sigma_E^n|_V$ is injective, and $\sigma_E^m(U) = \sigma_E^n(V)$. The map $x \mapsto (x, 0, x)$ then provides a homeomorphism from X_E^+ into the unit space \mathcal{G}_E^0 of \mathcal{G}_E . It is well known that $C^*(G) \cong C^*(\mathcal{G}_E)$ is the groupoid C^* -algebra of \mathcal{G}_E , and that $\mathcal{D}_E := C_0(\mathcal{G}_E^{(0)}) \cong C_0(X_E^+)$ is the subalgebra of continuous functions on units of \mathcal{G}_E . This subalgebra is often called the *diagonal subalgebra* of $C^*(G)$, and is given by

$$\mathcal{D}_E = \overline{\text{span}}\{S_\mu S_\mu^* \mid \mu \in E^n, n \in \mathbb{N}\}.$$

In what follows, recall that \mathbb{K} denotes compact operators on $\ell^2(\mathbb{N})$, which contains a natural copy of diagonal compact operators $c_0 \subseteq \mathbb{K}$. We will consider $\gamma \otimes \text{id}_{\mathbb{K}}$ as the standard gauge action on the stabilization $C^*(G) \otimes \mathbb{K}$. It was shown in [5, Theorem 5.1] that for any two finite graphs without sources and sinks $G = (V, E)$ and $G' = (V', E')$ we have that (X_E, σ_E) and $(X_{E'}, \sigma_{E'})$ are conjugate if and only if there is an equivariant isomorphism $\varphi : C^*(G) \otimes \mathbb{K} \rightarrow C^*(G') \otimes \mathbb{K}$ such that $\varphi(\mathcal{D}_E \otimes c_0) = \mathcal{D}_{E'} \otimes c_0$.

Suppose that $A = (A_{vw})_{v,w=0}^{n-1}$ is an essential $n \times n$ matrix with nonnegative integer entries. The *Cuntz–Krieger algebra* of A is the universal C^* -algebra \mathcal{O}_A generated by a family

$$\{S_{(v,w,m)} \mid v, w, m \in \mathbb{N}, 0 \leq v, w < n, 0 \leq m < A_{vw}\}$$

of partial isometries satisfying

- (1) $S_{(v,w,m)}^* S_{(v,w,m)} = \sum_{u=0}^{n-1} \sum_{\ell=0}^{A_{wu}-1} S_{(w,u,\ell)} S_{(w,u,\ell)}^*$,
- (2) $\sum_{v,w=0}^{n-1} \sum_{m=0}^{A_{vw}-1} S_{(v,w,m)} S_{(v,w,m)}^* = 1$.

For each family as above, we get projections $S_w = S_{(v,w,m)}^* S_{(v,w,m)}$ (independent of v and m), so that the family $(S_v, S_{(v',w',m)})$ becomes a Cuntz–Krieger family for the directed graph G_A (see [8, Remark 2.16] and [41, Remark 2.8]). Hence, there is a $*$ -isomorphism between \mathcal{O}_A and $C^*(G_A)$ which maps the generators $S_{(v,w,m)}$ of \mathcal{O}_A to edge generators for $C^*(G_A)$. This isomorphism between \mathcal{O}_A and $C^*(G_A)$ induces the usual gauge action on \mathcal{O}_A from [8] (which is also built from the universality of \mathcal{O}_A , see [20, Remark 2.2(2)]) and sends the natural diagonal subalgebra of \mathcal{O}_A to the diagonal subalgebra \mathcal{D}_E inside $C^*(G_A)$. Thus, there is no loss of generality arising from considering graph C^* -algebras instead of Cuntz–Krieger C^* -algebras.

Notation 2.3. Whenever $X, Y \subseteq \mathcal{L}(E)$ are norm-closed subspaces, we denote by XY or by $X \cdot Y$ the closed linear span of products $x \cdot y$ with $x \in X$ and $y \in Y$.

3. Shift equivalence and Cuntz–Pimsner algebras

In this section we discuss C^* -correspondences of adjacency matrices, Cuntz–Pimsner C^* -algebras and Pimsner dilations. The main result of this section is a characterization of equivariant stable isomorphism of Cuntz–Pimsner algebras in terms of Pimsner dilations.

We will need some of the theory of C^* -correspondences. We mention some of the basic definitions, but will assume some familiarity with the theory of Hilbert C^* -modules as in [30; 32].

Definition 3.1. A C^* -correspondence from \mathcal{A} to \mathcal{B} (or \mathcal{B} - \mathcal{A} C^* -correspondence) is a right Hilbert \mathcal{A} -module E with a $*$ -representation $\phi_E : \mathcal{B} \rightarrow \mathcal{L}(E)$, where $\mathcal{L}(E)$ is the C^* -algebra of adjointable operators on E . We denote by $\mathcal{K}(E)$ the ideal of $\mathcal{L}(E)$ which is closed linear span of rank-1 operators $\theta_{\xi,\eta}$ given by $\theta_{\xi,\eta}(\zeta) = \xi \langle \eta, \zeta \rangle$.

We will say that two \mathcal{B} - \mathcal{A} correspondences E and F are *unitarily isomorphic* (and denote this by $E \cong F$) if there is an isometric surjection $U : E \rightarrow F$ such that for every $a \in \mathcal{A}$, $b \in \mathcal{B}$ and $\xi \in E$ we have

$$U(\phi_E(b)\xi a) = \phi_F(b)U(\xi)a.$$

We will assume throughout this note that C^* -correspondences E are nondegenerate (sometimes called essential) in the sense that $\phi_E(\mathcal{B})E = E$. We say that a \mathcal{B} - \mathcal{A} C^* -correspondence E is *regular* if its left action ϕ_E is injective and $\phi_E(\mathcal{B}) \subseteq \mathcal{K}(E)$. When the context is clear, we write $b\xi$ to mean $\phi_E(b)\xi$ for $b \in \mathcal{B}$ and $\xi \in E$. Finally, we say that a \mathcal{B} - \mathcal{A} C^* -correspondence E is *full* if \mathcal{A} is equal to the ideal $\langle E, E \rangle$ defined as the closed linear span of $\langle \xi, \eta \rangle$ for $\xi, \eta \in E$.

The most important examples of C^* -correspondences in our study are the ones coming from adjacency matrices. Let V and W be sets, and let C be a $V \times W$ matrix so that C_{vw} is some cardinal. We define

$$E_C := \{(v, w, \alpha) \mid 0 \leq \alpha < C_{vw}, v \in V, w \in W\}.$$

When C is a $V \times W$ matrix with cardinal entries, we may construct a $c_0(V)$ - $c_0(W)$ correspondence $X(C)$ by taking the Hausdorff completion of all finitely supported functions on E_C with respect to the inner product

$$\langle \xi, \eta \rangle(w) = \sum_{(v,w,\alpha) \in E_C} \overline{\xi(v, w, \alpha)} \eta(v, w, \alpha),$$

where ξ and η are finitely supported on E_C . The left actions of $c_0(V)$ and the right action of $c_0(W)$ on $X(C)$ are then given by

$$(f \cdot \xi \cdot g)(v, w, \alpha) = f(v)\xi(v, w, \alpha)g(w) \quad \text{for } f \in c_0(V) \text{ and } g \in c_0(W).$$

When $V = W$, it is clear that $X(C)$ coincides with the graph C^* -correspondence $X(G_C)$ of the directed $G_C = (V, E_C)$, as is explained in the discussion preceding [10, Theorem 6.2], with range and source interchanged in the definitions of inner product and bimodule actions.

We will need to know that in the above way we obtain all $c_0(V)$ - $c_0(W)$ correspondences. This type of result was first shown by Kaliszewski, Patani and Quigg [23] when C is a square matrix indexed by a countable set and with countable entries.

Proposition 3.2. *Let V and W be sets, and E a $c_0(V)$ - $c_0(W)$ correspondence. Then there exists a unique $V \times W$ matrix C with cardinal entries such that E is unitarily isomorphic to $X(C)$.*

Proof. Let $v \in V$ and $w \in W$, and let $p_v \in c_0(V)$ and $p_w \in c_0(W)$ be the characteristic functions of $\{v\}$ and $\{w\}$ respectively. Since $p_v E p_w$ is a Hilbert space, we let C_{vw} be its dimension, and let $\{e_{(v,w,\alpha)}\}_{0 \leq \alpha < C_{vw}}$ be an orthonormal basis for it, indexed by ordinals $0 \leq \alpha < C_{vw}$. Then clearly C is a $V \times W$ matrix with cardinal values. We define a map $U : X(C) \rightarrow E$ on finitely supported functions by setting $U(\xi) = \sum_{(v,w,\alpha) \in E_C} \xi(v, w, \alpha)e_{(v,w,\alpha)}$ for finitely supported $\xi \in X(C)$. Now, for a finitely supported function $\xi \in X(C)$ we have for fixed $w \in W$ that

$$|U(\xi)|^2(w) = \sum_{(v,w,\alpha) \in E_C} |\xi(v, w, \alpha)|^2 = |\xi|^2(w).$$

Hence, U extends to an isometry on $X(C)$, and since the linear span of

$$\{e_{(v,w,\alpha)} \mid v \in V, w \in W, 0 \leq \alpha < C_{vw}\}$$

is dense in E , we get that U is a unitary isomorphism.

As for uniqueness, suppose F is another $c_0(W)$ - $c_0(V)$ -correspondence which is unitarily isomorphic to E via a unitary U . Hence, if C^E and C^F are the matrices associated to E and F via the first paragraph, we must have $C_{vw}^F = C_{vw}^E$, and we are done. \square

For a \mathcal{B} - \mathcal{A} correspondence E and a \mathcal{C} - \mathcal{B} correspondence F , we can form the *interior tensor product* \mathcal{C} - \mathcal{A} correspondence $F \otimes_{\mathcal{B}} E$ of E and F as follows. Let $F \otimes_{\text{alg}} E$ denote the quotient of the algebraic tensor product by the subspace generated by elements of the form

$$\eta b \otimes \xi - \eta \otimes b \xi \quad \text{for } \xi \in E, \eta \in F, b \in \mathcal{B}.$$

Define an \mathcal{A} -valued semi-inner product and left \mathcal{C} -action by setting

$$\begin{aligned} \langle \eta_1 \otimes \xi_1, \eta_2 \otimes \xi_2 \rangle &= \langle \xi_1, \langle \eta_1, \eta_2 \rangle \cdot \xi_2 \rangle \quad \text{for } \xi_1, \xi_2 \in E, \eta_1, \eta_2 \in F, \\ c \cdot (\eta \otimes \xi) &= (c \cdot \eta) \otimes \xi \quad \text{for } \eta \in E, \xi \in F, c \in \mathcal{C}. \end{aligned}$$

We denote by $F \otimes_{\mathcal{B}} E$ the separated completion of $F \otimes_{\text{alg}} E$ with respect to the \mathcal{A} -valued semi-inner product defined above. One then verifies that $F \otimes_{\mathcal{B}} E$ is a \mathcal{C} - \mathcal{A} correspondence (see for example [30, Proposition 4.5]). We will often abuse notation and write $F \otimes E$ for $F \otimes_{\mathcal{B}} E$ when the context is clear. If E is an \mathcal{A} - \mathcal{A} correspondence (a C^* -correspondence over \mathcal{A}), then we denote by $E^{\otimes n}$ the n -fold interior tensor product of E with itself.

Proposition 3.3. *Let I, J, K be sets, and let C be an $I \times J$ matrix and D be a $J \times K$ matrix, both with cardinal entries. Then there is a unitary isomorphism $U : X(C) \otimes X(D) \rightarrow X(CD)$.*

Proof. For each $i \in I, j \in J$ and $k \in K$, let $C_{ij} = |X_{ij}|, D_{jk} = |Y_{jk}|$ for some sets X_{ij} and Y_{jk} . Then $(CD)_{ik}$ is the cardinality of the disjoint union of Cartesian products $\bigsqcup_{j \in J} X_{ij} \times Y_{jk}$.

Now, clearly both $X(C) \otimes X(D)$ and $X(CD)$ are $c_0(I)$ - $c_0(K)$ correspondences, so that by the uniqueness part of Proposition 3.2 it will suffice to show for every $i \in I$ and $k \in K$ that the dimension of the Hilbert space $p_i X(C) \otimes X(D) p_k$ is equal to $(CD)_{ik}$.

So let $\{e_x\}_{x \in X_{ij}}$ be an orthonormal basis for $p_i X(C) p_j$ for $i \in I$ and $j \in J$ and let $\{e_y\}_{y \in Y_{j'k}}$ be an orthonormal basis for $p_{j'} X(D) p_k$ for $j' \in J$ and $k \in K$. Then, for $x \in X_{ij}$ and $y \in Y_{j'k}$ we have $e_x \otimes e_y \neq 0$ if and only if $j = j'$. Hence, an orthonormal basis for $p_i X(C) \otimes X(D) p_k$ is given by $\{e_x \otimes e_y \mid x \in X_{ij}, y \in Y_{jk}, j \in J\}$. The cardinality of this basis is clearly equal to $(CD)_{ik}$, so the proof is concluded. \square

The following definition of shift equivalence of C^* -correspondences first appeared in [22].

Definition 3.4. Let E and F be C^* -correspondences over \mathcal{A} and \mathcal{B} respectively. We say that E and F are:

- (1) *Shift equivalent* with lag m if there are $m \in \mathbb{N} \setminus \{0\}$, an \mathcal{A} - \mathcal{B} correspondence R and a \mathcal{B} - \mathcal{A} correspondence S , together with unitary isomorphisms

$$\begin{aligned} E^{\otimes m} &\cong R \otimes S, & F^{\otimes m} &\cong S \otimes R, \\ S \otimes E &\cong F \otimes S, & E \otimes R &\cong R \otimes F. \end{aligned}$$

- (2) *Elementary shift related* if they are shift equivalent with lag 1.
- (3) *Strong shift equivalent* if they are equivalent in the transitive closure of the elementary shift relation.

The following shows that shift equivalence between two C^* -correspondences generalizes shift equivalence of adjacency matrices.

Proposition 3.5. *Let A and B be matrices with cardinal entries, indexed by sets V and W . Then, A and B are shift equivalent if and only if $X(A)$ and $X(B)$ are shift equivalent.*

Proof. By Proposition 3.3 we see that if A and B are shift equivalent with lag m via matrices C and D , then the C^* -correspondences $X(A)$ and $X(B)$ are shift equivalent with lag m via the C^* -correspondences $X(C)$ and $X(D)$.

Conversely, suppose R and S are $c_0(V)$ - $c_0(W)$ and $c_0(W)$ - $c_0(V)$ correspondences implementing shift equivalence of lag m for $X(A)$ and $X(B)$. By Proposition 3.2 there are a $V \times W$ matrix C and $W \times V$ matrix D with unitary isomorphisms $R \cong X(C)$ and $S \cong X(D)$. Hence, the uniqueness portion in Proposition 3.2 guarantees that C and D implement a shift equivalence of A and B with lag m . \square

Every path isomorphism $\phi : E_A \rightarrow E_B$ induces a unitary isomorphism $\Phi : X(A) \rightarrow X(B)$ by setting $\Phi(\xi)(v, w, \alpha) = \xi(\phi^{-1}(v, w, \alpha))$. However, the converse is in general false; it is easy to construct a unitary isomorphism $U : X(A) \rightarrow X(B)$ for which there is no path isomorphism $\phi : E_A \rightarrow E_B$ such that U is the induced unitary isomorphism from ϕ . The point of Proposition 3.5 is that we only need to find *some* path isomorphisms for each one of the four relations appearing in the definition of shift equivalence, and not necessarily path isomorphisms which induce *the same* four isomorphisms we started with at the level of C^* -correspondences.

Remark 3.6. From considerations similar to the above we see that strong shift equivalence of matrices implies the strong shift equivalence of their associated C^* -correspondences. The converse, however, is unknown.

Next we discuss Cuntz–Pimsner algebras and Pimsner dilations. More material on Cuntz–Pimsner algebras, with a special emphasis on C^* -correspondences of graphs, can be found in [41, Chapter 8]. We note immediately that what we call a *rigged* representation here is often referred to as an *isometric* representation in the literature (see [36]).

Definition 3.7. Let E be a C^* -correspondence from \mathcal{A} to \mathcal{B} , and \mathcal{C} some C^* -algebra. A *rigged representation* of E is a triple $(\pi_{\mathcal{A}}, \pi_{\mathcal{B}}, t)$ such that $\pi_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{C}$ and $\pi_{\mathcal{B}} : \mathcal{B} \rightarrow \mathcal{C}$ are $*$ -homomorphisms and $t : E \rightarrow \mathcal{C}$ is a linear map such that

- (1) $\pi_{\mathcal{B}}(b)t(\xi)\pi_{\mathcal{A}}(a) = t(b \cdot \xi \cdot a)$ for $a \in \mathcal{A}$, $b \in \mathcal{B}$ and $\xi \in E$,
- (2) $t(\xi)^*t(\eta) = \pi_{\mathcal{A}}(\langle \xi, \eta \rangle)$.

We say that $(\pi_{\mathcal{A}}, \pi_{\mathcal{B}}, t)$ is *injective* if both $\pi_{\mathcal{A}}$ and $\pi_{\mathcal{B}}$ are injective $*$ -homomorphisms. We denote by $C^*(\pi_{\mathcal{A}}, \pi_{\mathcal{B}}, t)$ the C^* -algebra generated by the images of $\pi_{\mathcal{A}}$, $\pi_{\mathcal{B}}$ and t .

We will mostly be concerned with the situation where $\mathcal{A} = \mathcal{B}$ and $\pi_{\mathcal{A}} = \pi_{\mathcal{B}}$, in which case we will define $\pi := \pi_{\mathcal{A}} = \pi_{\mathcal{B}}$, and refer to the representation as a pair (π, t) , and the generated C^* -algebra by $C^*(\pi, t)$.

Now let E be a C^* -correspondence over \mathcal{A} . The Toeplitz–Pimsner algebra $\mathcal{T}(E)$ is then the universal C^* -algebra generated by a rigged representation of E . Universality of $\mathcal{T}(E)$ implies that it comes equipped with a point-norm continuous gauge action $\gamma : \mathbb{T} \rightarrow \text{Aut}(\mathcal{T}(E))$ given by

$$\gamma_z(\pi(a)) = \pi(a) \quad \text{and} \quad \gamma_z(t(\xi)) = z \cdot t(\xi) \quad \text{for } z \in \mathbb{T}, \xi \in E, a \in \mathcal{A}.$$

Toeplitz–Pimsner algebras have a canonical quotient, also known as the Cuntz–Pimsner algebra, which was originally defined by Pimsner in [40] and refined by Katsura in [24].

Definition 3.8. For a C^* -correspondence E over \mathcal{A} , we define *Katsura’s ideal* J_E in \mathcal{A} by

$$J_E := \{a \in \mathcal{A} \mid \phi_E(a) \in \mathcal{K}(E) \text{ and } ab = 0 \text{ for all } b \in \ker \phi_E\}.$$

For a rigged representation (π, t) of a C^* -correspondence E over \mathcal{A} , it is known there is a well-defined $*$ -homomorphism $\psi_t : \mathcal{K}(E) \rightarrow C^*(\pi, t)$ given by $\psi_t(\theta_{\xi, \eta}) = t(\xi)t(\eta)^*$ for $\xi, \eta \in E$ (see for instance [21, Lemma 2.2]).

Definition 3.9. A rigged representation (π, t) is said to be *covariant* if $\pi(a) = \psi_t(\phi_E(a))$ for all $a \in J_E$.

The Cuntz–Pimsner algebra $\mathcal{O}(E)$ is then the universal C^* -algebra generated by a *covariant* representation of E . Suppose now that (π, t) and $(\bar{\pi}, \bar{t})$ are universal rigged and covariant representations respectively, so that $\mathcal{T}(E) = C^*(\pi, t)$ and $\mathcal{O}(E) = C^*(\bar{\pi}, \bar{t})$. We denote by \mathcal{J}_E the kernel of the natural quotient map from $\mathcal{T}(E)$ onto $\mathcal{O}(E)$, which is the ideal generated by elements of the form $\pi(a) - \psi_t(\phi_E(a))$ for $a \in J_E$. Since the ideal \mathcal{J}_E is gauge-invariant, we see there is an induced point-norm continuous circle action $\gamma : \mathbb{T} \rightarrow \text{Aut}(\mathcal{O}(E))$ given by

$$\gamma_z(\pi(a)) = \pi(a) \quad \text{and} \quad \gamma_z(t(\xi)) = z \cdot t(\xi) \quad \text{for } z \in \mathbb{T}, \xi \in E, a \in \mathcal{A}.$$

It follows that $\mathcal{O}(E)$ then becomes a \mathbb{Z} -graded C^* -algebra, with its n -th graded component given by $\mathcal{O}(E)_n := \{c \in \mathcal{O}(E) \mid \gamma_z(c) = z^n \cdot c\}$. By [42, Theorem 3] we see that there is a bijective correspondence between topologically \mathbb{Z} -graded C^* -algebras, with graded $*$ -homomorphisms and C^* -algebras equipped with a circle action, together with equivariant $*$ -homomorphisms. In particular, an isomorphism $\varphi : \mathcal{O}(E) \rightarrow \mathcal{O}(F)$ between two Cuntz–Pimsner algebras is equivariant if and only if it is graded.

Example 3.10. When $G = (V, E)$ is a directed graph we know that the graph C^* -algebra $C^*(G)$ coincides with the Cuntz–Pimsner algebra of the correspondence $X(A_E)$ by an isomorphism that intertwines the gauge action of $C^*(G)$ and the gauge action of $\mathcal{O}(X(A_E))$. See [41, Section 8] for more details.

Definition 3.11. Let E, F be C^* -correspondences over \mathcal{A} and \mathcal{B} , respectively. Denote by γ^E and γ^F the gauge actions on $\mathcal{O}(E)$ and $\mathcal{O}(F)$ respectively. We say that $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are *equivariantly stably isomorphic* if there is a $*$ -isomorphism $\varphi : \mathcal{O}(E) \otimes \mathbb{K} \rightarrow \mathcal{O}(F) \otimes \mathbb{K}$ such that $\varphi \circ (\gamma_z^E \otimes \text{id}) = (\gamma_z^F \otimes \text{id}) \circ \varphi$ for every $z \in \mathbb{T}$.

Equivariant stable isomorphisms arise naturally in the classification of groupoid C^* -algebras, but always in a way which respects diagonal subalgebras. For instance in [7], equivariant stable isomorphisms which respect diagonal subalgebras are characterized in terms of isomorphisms of graded groupoids. We will get back to such isomorphisms in Section 6.

The following definition is similar to Definition 3.1, and we will provide a precise distinction between the two in the remark that follows.

Definition 3.12. Suppose E and F are C^* -correspondences over C^* -algebras \mathcal{A} and \mathcal{B} respectively. We say that E and F are *unitarily isomorphic* (denoted by $E \cong F$) if there exist a surjective, isometric map $U : E \rightarrow F$ and a $*$ -isomorphism $\rho : \mathcal{A} \rightarrow \mathcal{B}$ such that $U(b \cdot \xi \cdot a) = \rho(b) \cdot U(\xi) \cdot \rho(a)$ for all $a, b \in \mathcal{A}$, $\xi \in E$.

Remark 3.13. Suppose now that E and F are C^* -correspondences over C^* -algebras \mathcal{A} and \mathcal{B} respectively, and that $U : E \rightarrow F$ is a unitary isomorphism implemented by a $*$ -isomorphism $\rho : \mathcal{A} \rightarrow \mathcal{B}$. By the discussion in [9, Subsection 2.1] we may “twist” the C^* -correspondence F to a C^* -correspondence F_ρ

over \mathcal{A} so that $U : E \rightarrow F_\rho$ becomes a unitary isomorphism as in Definition 3.1. More precisely, the new operations on F_ρ are given by

$$\begin{aligned} \langle \xi, \eta \rangle_\rho &:= \rho^{-1}(\langle \xi, \eta \rangle_\mathcal{B}) \quad \text{for } \xi, \eta \in F, \\ a \cdot \xi &= \rho(a) \cdot \xi \quad \text{and} \quad \xi \cdot a := \xi \cdot \rho(a) \quad \text{for all } \xi \in F \text{ and } a \in \mathcal{A}, \end{aligned}$$

and the identity map $\text{id}_F^\rho : F \rightarrow F_\rho$ becomes a unitary isomorphism as in Definition 3.12. Then, the isometric surjection $\text{id}_F^\rho \circ U$ is a unitary isomorphism as in Definition 3.1. Hence, we can go back and forth between the two definitions of unitary isomorphism.

We warn the reader that unitary isomorphism as in Definition 3.12 is not the same as having an isometric surjection U implemented via two potentially *different* $*$ -isomorphisms $\rho_1 : \mathcal{A} \rightarrow \mathcal{B}$ and $\rho_2 : \mathcal{A} \rightarrow \mathcal{B}$ in the sense that $U(b \cdot \xi \cdot a) = \rho_1(a)U(\xi)\rho_2(b)$ for $a, b \in \mathcal{A}$. We also note that whenever one of our C^* -correspondences has possibly different left and right coefficient C^* -algebras, we only consider one notion of unitary isomorphism, which is the one in Definition 3.1.

In what follows, we say that M is an *imprimitivity* \mathcal{A} - \mathcal{B} correspondence (from \mathcal{B} to \mathcal{A}) if it is full and its left action ϕ_M is a $*$ -isomorphism onto $\mathcal{K}(E)$. The following was introduced by Muhly and Solel [36] in their study of tensor algebras of C^* -correspondences.

Definition 3.14. Let E and F be C^* -correspondences over C^* -algebras \mathcal{A} and \mathcal{B} respectively. We say that E and F are *strongly Morita equivalent* if there are an imprimitivity \mathcal{A} - \mathcal{B} bimodule M and an isometric surjective linear map $U : E \otimes M \rightarrow M \otimes F$ such that for every $\xi \in E \otimes M$, $a \in \mathcal{A}$ and $b \in \mathcal{B}$ we have $U(a\xi b) = aU(\xi)b$.

When M is an imprimitivity \mathcal{A} - \mathcal{B} -bimodule, there is an “inverse” imprimitivity \mathcal{B} - \mathcal{A} -bimodule M^* satisfying $M \otimes M^* \cong \mathcal{A}$ and $M^* \otimes M \cong \mathcal{B}$. If moreover $\mathcal{A} = \mathcal{B}$, it follows from [1, Theorem 2.9] that the \mathbb{Z} -graded components of $\mathcal{O}(M)$ are given for $n \in \mathbb{N}$ by

- $\mathcal{O}(M)_n \cong M^{\otimes n}$ for $n > 0$,
- $\mathcal{O}(M)_0 \cong \mathcal{A}$, and
- $\mathcal{O}(M)_n \cong (M^{\otimes n})^*$ for $n < 0$.

Proposition 3.15. *Let E and F be C^* -correspondences over C^* -algebras \mathcal{A} and \mathcal{B} respectively. If E and F are unitarily equivalent, then $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly isomorphic. If moreover E and F are imprimitivity bimodules, then E and F are unitarily equivalent if and only if $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly isomorphic.*

Proof. Let $(\bar{\pi}, \bar{t})$ be a universal covariant representation for $\mathcal{O}(E)$. Assume that E and F are unitarily equivalent. By the implication (1) \Rightarrow (4) of [10, Corollary 3.5] we get that $\mathcal{T}(E)$ and $\mathcal{T}(F)$ are graded isomorphic, and hence equivariantly isomorphic. Then, [10, Theorem 3.1] gives rise to an induced isomorphism between $\mathcal{O}(E)$ and $\mathcal{O}(F)$, which is actually equivariant since the ideals \mathcal{J}_E and \mathcal{J}_F are gauge-invariant.

Conversely, if E and F are imprimitivity bimodules, and $\varphi : \mathcal{O}(E) \rightarrow \mathcal{O}(F)$ is an equivariant isomorphism, then φ is \mathbb{Z} -graded, and we get that the restriction $U := \varphi|_{\mathcal{O}(E)_1} : \mathcal{O}(E)_1 \rightarrow \mathcal{O}(F)_1$ is an isometric

$\mathcal{O}(E)_0$ - $\mathcal{O}(F)_0$ bimodule isomorphism, implemented by the $*$ -isomorphism $\rho := \varphi|_{\mathcal{O}(E)_0} : \mathcal{O}(E)_0 \rightarrow \mathcal{O}(F)_0$. From the identifications in the discussion preceding the theorem, we get that E and F are unitarily isomorphic. \square

Given a C^* -correspondence E over \mathcal{A} , we may form the external minimal tensor product $E \otimes \mathbb{K}$, which is a C^* -correspondence over $\mathcal{A} \otimes \mathbb{K}$ as defined in [30, p. 34]. A consequence of [10, Proposition 2.10] is that $\mathcal{O}(E \otimes \mathbb{K})$ is canonically isomorphic to $\mathcal{O}(E) \otimes \mathbb{K}$ via a map induced by the representation $(\bar{\pi} \otimes \text{id}, \bar{\tau} \otimes \text{id})$.

Corollary 3.16. *Let E and F be C^* -correspondences over C^* -algebras \mathcal{A} and \mathcal{B} respectively. If $E \otimes \mathbb{K}$ and $F \otimes \mathbb{K}$ are unitarily equivalent, then $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly stably isomorphic. If moreover E and F are **imprimitivity bimodules**, then $E \otimes \mathbb{K}$ and $F \otimes \mathbb{K}$ are unitarily equivalent if and only if $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly stably isomorphic.*

Proof. Since $E \otimes \mathbb{K}$ and $F \otimes \mathbb{K}$ are unitarily equivalent, by Proposition 3.15 we get that $\mathcal{O}(E \otimes \mathbb{K}) \cong \mathcal{O}(F \otimes \mathbb{K})$ equivariantly.

Suppose now that $(\bar{\pi}, \bar{\tau})$ is a universal covariant representation of E . By [10, Proposition 2.10], we get that the representation $(\bar{\pi} \otimes \text{id}, \bar{\tau} \otimes \text{id})$ induces an isomorphism $\bar{\rho} : \mathcal{O}(E \otimes \mathbb{K}) \rightarrow \mathcal{O}(E) \otimes \mathbb{K}$ satisfying $\bar{\rho}(\xi \otimes K) = \bar{\tau}(\xi) \otimes K$ for every $\xi \in E$ and $K \in \mathbb{K}$. Hence $\bar{\rho}$ must be equivariant. It is similarly shown that $\mathcal{O}(F \otimes \mathbb{K}) \cong \mathcal{O}(F) \otimes \mathbb{K}$ equivariantly. Hence, we get equivariantly that

$$\mathcal{O}(E) \otimes \mathbb{K} \cong \mathcal{O}(E \otimes \mathbb{K}) \cong \mathcal{O}(F \otimes \mathbb{K}) \cong \mathcal{O}(F) \otimes \mathbb{K}.$$

With this, we obtain the first part of our result.

Conversely, if moreover E and F are imprimitivity bimodules, then so are $E \otimes \mathbb{K}$ and $F \otimes \mathbb{K}$. Thus, we are done by Proposition 3.15 and the above identifications. \square

Let E be a C^* -correspondence over a C^* -algebra \mathcal{A} , and $(\bar{\pi}, \bar{\tau})$ a universal covariant representation of E . We denote by

$$\mathcal{A}_\infty = \mathcal{O}(E)_0 = \overline{\text{span}}\{\bar{\pi}(\mathcal{A}), \psi_{\bar{\tau}_m}(\mathcal{K}(E^{\otimes m})) \mid m \geq 1\}$$

the fixed point algebra of $\mathcal{O}(E)$ under the gauge action γ . It is well known that \mathcal{A}_∞ is the direct limit of C^* -subalgebras \mathcal{A}_n given by

$$\mathcal{A}_n = \overline{\text{span}}\{\bar{\pi}(\mathcal{A}), \psi_{\bar{\tau}_m}(\mathcal{K}(E^{\otimes m})) \mid 1 \leq m \leq n\}.$$

When E has an injective left action, we get that $\mathcal{A}_n = \psi_{\bar{\tau}_n}(\mathcal{K}(E^{\otimes n}))$ for each $n \in \mathbb{N}$, and we define $E_n := \bar{\tau}(E)\mathcal{A}_n$ inside $\mathcal{O}(E)$. Then, E_n becomes a C^* -correspondence from \mathcal{A}_n to \mathcal{A}_{n+1} , where the left action of \mathcal{A}_{n+1} is defined by left multiplication in $\mathcal{O}(E)$. By taking the direct limit $E_\infty := \bar{\tau}(E) \cdot \mathcal{A}_\infty = \overline{\bigcup_{n=1}^\infty E_n}$, we obtain a C^* -correspondence over \mathcal{A}_∞ called the *Pimsner dilation* of E .

It was shown by Pimsner in [40, Theorem 2.5(2)] that the identification $E \otimes \mathcal{A}_\infty \cong E_\infty$ gives rise to an equivariant $*$ -injection $\mathcal{T}(E) \rightarrow \mathcal{T}(E_\infty)$, which then induces an equivariant isomorphism $\mathcal{O}(E) \cong \mathcal{O}(E_\infty)$ between the quotients. When E is also *regular* and *full*, the C^* -correspondence E_∞ becomes an imprimitivity bimodule.

Theorem 3.17. *Let E and F be regular and full C^* -correspondences over σ -unital C^* -algebras \mathcal{A} and \mathcal{B} respectively. Then, $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly stably isomorphic if and only if E_∞ and F_∞ are strong Morita equivalent.*

Proof. Since E and F are regular and full, we get that E_∞ and F_∞ are imprimitivity bimodules. Hence, by Corollary 3.16 we get that $\mathcal{O}(E)$ and $\mathcal{O}(F)$ are equivariantly stably isomorphic if and only if $E_\infty \otimes \mathbb{K}$ and $F_\infty \otimes \mathbb{K}$ are unitarily isomorphic.

Since \mathcal{A} and \mathcal{B} are σ -unital C^* -algebras, so are \mathcal{A}_∞ and \mathcal{B}_∞ . Hence, every strong Morita equivalence M for E_∞ and F_∞ must be a σ -TRO in the sense of [13, p. 6], so that a combination of [13, Theorem 5.2] and [13, Proposition 3.1] shows that $E_\infty \otimes \mathbb{K}$ and $F_\infty \otimes \mathbb{K}$ are unitarily equivalent if and only if E_∞ and F_∞ are strongly Morita equivalent. \square

It is important to say a few words about the assumptions in Theorem 3.17 and what they mean for C^* -correspondences of adjacency matrices with entries in \mathbb{N} . Suppose A is a square adjacency matrix with entries in \mathbb{N} , and indexed by a set V . First note that V is countable if and only if $c_0(V)$ is σ -unital if and only if $C^*(G_A)_0$ is σ -unital. Moreover, by [41, Proposition 8.8] the following hold:

- (1) A has finitely supported rows if and only if G_A has finite out-degrees, if and only if the left action of $X(A)$ has image in $\mathcal{K}(X(A))$.
- (2) A has no zero rows if and only if G_A has no sinks, if and only if $X(A)$ has an injective left action.
- (3) A has no zero columns if and only if G_A has no sources, if and only if $X(A)$ is full.

So we see that in order to apply Theorem 3.17 to $X(A)$, we must verify that A is over a countable set V , has finitely supported rows and is essential. In this case, Theorem 3.17 shows that the existence of an equivariant stable isomorphism between graph C^* -algebras $C^*(G_A)$ and $C^*(G_B)$ coincides with the existence of a strong Morita equivalence of the Pimsner dilations $X(A)_\infty$ and $X(B)_\infty$.

4. Compatible shift equivalence

In this section we introduce and study *compatible shift equivalence*, which is formulated in terms of adjacency matrices and path isomorphisms. We show that it is indeed an equivalence relation, and that strong shift equivalence implies compatible shift equivalence. In what follows, for a matrix A we write id_A to mean id_{E_A} .

Definition 4.1. Let A and B be matrices indexed by V and W respectively, with entries in \mathbb{N} . Suppose there are a lag $m \in \mathbb{N} \setminus \{0\}$ and matrices R over $V \times W$ and S over $W \times V$ with entries in \mathbb{N} together with path isomorphisms

$$\begin{aligned} \phi_R : E_A \times E_R &\rightarrow E_R \times E_B, & \phi_S : E_B \times E_S &\rightarrow E_S \times E_A, \\ \psi_A : E_R \times E_S &\rightarrow E_A^m, & \psi_B : E_S \times E_R &\rightarrow E_B^m. \end{aligned}$$

We say that R and S are *compatible* if

$$\phi_R^{(m)} = (\text{id}_R \times \psi_B)(\psi_A^{-1} \times \text{id}_R), \quad \phi_S^{(m)} = (\text{id}_S \times \psi_A)(\psi_B^{-1} \times \text{id}_S), \tag{4-1}$$

where

$$\begin{aligned}\phi_R^{(m)} &:= (\phi_R \times \text{id}_{B^{m-1}})(\text{id}_A \times \phi_R \times \text{id}_{B^{m-2}}) \cdots (\text{id}_{A^{m-1}} \times \phi_R), \\ \phi_S^{(m)} &:= (\phi_S \times \text{id}_{A^{m-1}})(\text{id}_B \times \phi_S \times \text{id}_{A^{m-2}}) \cdots (\text{id}_{B^{m-1}} \times \phi_S).\end{aligned}$$

Finally, we say that A and B are *compatibly shift equivalent* if they are shift equivalent via a compatible pair of matrices R and S .

The following lemma shows that certain relations between the maps $\phi_R, \phi_S, \psi_A, \psi_B$ are automatic when R and S are compatible.

Lemma 4.2. *Suppose A and B are essential matrices over V and W respectively, with entries in \mathbb{N} . Suppose A and B are compatibly shift equivalent with lag m , matrices R and S , and path isomorphisms $\phi_R, \phi_S, \psi_A, \psi_B$. Then*

$$\begin{aligned}(\psi_A \times \text{id}_A)(\text{id}_R \times \phi_S) &= (\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S), \\ (\psi_B \times \text{id}_B)(\text{id}_S \times \phi_R) &= (\text{id}_B \times \psi_B)(\phi_S^{-1} \times \text{id}_R).\end{aligned}$$

Proof. First note that by compatible shift equivalence we have that

$$\begin{aligned}(\psi_A \times \text{id}_R \times \text{id}_S) &= (\phi_R^{(m)} \times \text{id}_S)^{-1}(\text{id}_R \times \psi_B \times \text{id}_S) \\ &= (\phi_R^{(m)} \times \text{id}_S)^{-1}(\text{id}_R \times \phi_S^{(m)})^{-1}(\text{id}_R \times \text{id}_S \times \psi_A).\end{aligned}$$

However, since

$$(\psi_A \times \text{id}_R \times \text{id}_S)(\text{id}_R \times \text{id}_S \times \psi_A^{-1}) = (\text{id}_{A^m} \times \psi_A^{-1})(\psi_A \times \text{id}_{A^m}),$$

we actually get that

$$(\psi_A \times \text{id}_{A^m}) = (\text{id}_{A^m} \times \psi_A)(\phi_R^{(m)} \times \text{id}_S)^{-1}(\text{id}_R \times \phi_S^{(m)})^{-1}. \quad (4-2)$$

Now let $(r, s, a) \in E_R \times E_S \times E_A$. Define $r_0 = r$, $s_0 = s$ and $a_0 = a$. Since A is essential, we can find $a_1, \dots, a_m, a_{m+1}, \dots, a_{2m-1} \in E_A$ so that $a_0 a_1 \cdots a_m a_{m+1} \cdots a_{2m-1} \in E_A^{m+1}$. We may then define inductively $r_k \in E_R$, $s_k \in E_S$ and $a'_k \in E_A$ for $0 \leq k \leq 2m-1$ so that

$$(\phi_R^{-1} \times \text{id}_S)(\text{id}_R \times \phi_S^{-1})(r_k, s_k, a_k) = (a'_k, r_{k+1}, s_{k+1}).$$

From (4-2) we get that $\psi_A(r_0, s_0) = a'_0 \cdots a'_{m-1}$, that $\psi_A(r_1, s_1) = a'_1 \cdots a'_m$ and that $a_0 \cdots a_{m-1} = \psi_A(r_m, s_m) = a'_m \cdots a'_{2m-1}$. In particular, we see that $a = a_0 = a'_m$.

To prove the first equality in the statement, we compute

$$\begin{aligned}(\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S)(\text{id}_R \times \phi_S^{-1})(r, s, a) &= (\text{id}_A \times \psi)(a'_0, r_1, s_1) \\ &= a'_0 a'_1 \cdots a'_m = \psi_A(r, s) a'_m = \psi_A(r, s) a = (\psi_A \times \text{id})(r, s, a).\end{aligned}$$

This shows that $(\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S)(\text{id}_R \times \phi_S^{-1}) = (\psi_A \times \text{id})$, which is then equivalent to the first equality $(\psi_A \times \text{id}_A)(\text{id}_R \circ \phi_S) = (\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S)$. A symmetric argument works to show the second equality as well. \square

Remark 4.3. Notice from the proof that we also get that ψ_A and ψ_B are uniquely determined by ϕ_R and ϕ_S . Indeed, from (4-2) we get that

$$(\psi_A \times \psi_A^{-1}) = (\phi_R^{(m)} \times \text{id}_S)^{-1} (\text{id}_R \times \phi_S^{(m)})^{-1},$$

so that compressing to the first part yields back ψ_A . A similar argument then works for ψ_B as well.

Proposition 4.4. *Compatible shift equivalence is an equivalence relation on the collection of essential matrices with entries in \mathbb{N} .*

Proof. That compatible shift equivalence is reflexive and symmetric is clear. Thus, we need to show transitivity.

Let A, B and C be essential matrices over \mathbb{N} . Suppose that A and B are compatibly shift equivalent with lag m , matrices R, S and path isomorphisms $\psi_A, \psi_B, \phi_R, \phi_S$, while B and C are compatibly shift equivalent with lag m' , matrices R', S' and path isomorphisms $\psi'_B, \psi'_C, \phi_{R'}, \phi_{S'}$. We claim that A and C are compatibly shift equivalent with lag $m + m'$, matrices $RR', S'S$ and path isomorphisms $\psi'_A, \psi_C, \phi_{RR'}, \phi_{S'S}$ given by

$$\begin{aligned}\psi'_A &:= (\text{id}_{A^{m'}} \times \psi_A)((\phi_R^{(m')})^{-1} \times \text{id}_S)(\text{id}_R \times \psi'_B \times \text{id}_S), \\ \psi_C &:= (\text{id}_{C^m} \times \psi'_C)((\phi_{S'}^{(m)})^{-1} \times \text{id}_{R'})(\text{id}_{S'} \times \psi_B \times \text{id}_{R'}), \\ \phi_{RR'} &:= (\text{id}_R \times \phi_{R'})(\phi_R \times \text{id}_{R'}), \\ \phi_{S'S} &:= (\text{id}_S \times \phi_{S'})(\phi_S \times \text{id}_{S'}).\end{aligned}$$

It is easy to see that $RR'S'S = A^{m+m'}$, $S'SRR' = C^{m+m'}$, $RR'C = ARR'$ and $CS'S = S'SA$. Moreover, it is clear that $\psi'_A, \psi_C, \phi_{RR'}, \phi_{S'S}$ are path isomorphisms. Thus, in order to show that the above data yields a compatible shift equivalence, we need only show

$$\begin{aligned}\phi_{RR'}^{(m+m')}(\psi'_A \times \text{id}_{RR'}) &= \text{id}_{RR'} \times \psi_C, \\ \phi_{S'S}^{(m+m')}(\psi_C \times \text{id}_{S'S}) &= \text{id}_{S'S} \times \psi'_A.\end{aligned}$$

We will show the first of these equalities, and the second will follow from a symmetric argument.

First, let $r_1 r'_1 s' s' r_2 r'_2 \in E_{RR'} \times E_{S'S} \times E_{RR'}$, and define

$$\mu_A r_3 := (\phi_R^{(m')})^{-1} (r_1 \psi'_B (r'_1 s' s')) \quad \text{and} \quad r'_3 \mu_C := \phi_{R'}^{(m)} (\psi_B (s r_2) r'_2). \quad (4-3)$$

Then we get that

$$\begin{aligned}\psi'_A (r_1 r'_1 s' s) &= (\text{id}_{A^m} \times \psi_A)((\phi_R^{(m)})^{-1} \times \text{id}_S)(r_1 \psi'_B (r'_1 s' s)) \\ &= (\text{id}_{A^m} \times \psi_A)(\mu_A r_3) = \mu_A \psi_A (r_3 s),\end{aligned}$$

and from Lemma 4.2 we also get that

$$\begin{aligned}\psi_C (s' s' r_2 r'_2) &= (\text{id}_{C^m} \times \psi'_C)((\phi_{S'}^{(m)})^{-1} \times \text{id}_{R'})(s' \psi_B (s r_2) r'_2) \\ &= (\psi'_C \times \text{id}_{C^m})(\text{id}_{S'} \times \phi_{R'}^{(m)})(s' \psi_B (s r_2) r'_2) = \psi'_C (s' r'_3) \mu_C.\end{aligned}$$

Thus, together we obtain

$$\psi'_A (r_1 r'_1 s' s) = \mu_A \psi_A (r_3 s) \quad \text{and} \quad \psi_C (s' s' r_2 r'_2) = \psi'_C (s' r'_3) \mu_C. \quad (4-4)$$

Next, from compatibility we also get

$$\phi_R^{(m)}(\psi_A(r_3s)r_2) = r_3\psi_B(sr_2) \quad \text{and} \quad \phi_{R'}^{(m)}(\psi'_B(r'_1s')r'_3) = r'_1\psi'_C(s'r'_3). \quad (4-5)$$

Combining (4-3), (4-4), (4-5), we compute

$$\begin{aligned} \phi_{RR'}^{(m+m')}(\psi'_A \times \text{id}_{RR'})(r_1r'_1s'sr_2r'_2) &= \phi_{RR'}^{(m+m')}(\mu_A\psi_A(r_3s)r_2r'_2) \\ &= (\text{id}_R \times \phi_{R'}^{(m+m')})(\phi_R^{(m+m')} \times \text{id}_{R'})(\mu_A\psi_A(r_3s)r_2r'_2) \\ &= (\text{id}_R \times \phi_{R'}^{(m+m')})(\phi_R^{(m')} \times \text{id}_{B^m} \times \text{id}_{R'})(\mu_A r_3 \psi_B(sr_2)r'_2) \\ &= (\text{id}_R \times \phi_{R'}^{(m+m')})(r_1\psi'_B(r'_1s')\psi_B(sr_2)r'_2) \\ &= (\text{id}_R \times \phi_{R'}^{(m')} \times \text{id}_{C^m})(r_1\psi'_B(r'_1s')r'_3\mu_C) \\ &= r_1r'_1\psi'_C(s'r'_3)\mu_C \\ &= r_1r'_1\psi_C(s'sr_2r'_2) \\ &= (\text{id}_{RR'} \times \psi_C)(r_1r'_1s'sr_2r'_2). \end{aligned}$$

Thus, we have shown $\phi_{RR'}^{(m+m')}(\psi'_A \times \text{id}_{RR'}) = \text{id}_{RR'} \times \psi_C$ as desired. \square

Corollary 4.5. *Let A and B be essential matrices over V and W respectively, with entries in \mathbb{N} . If A and B are strong shift equivalent, then they are compatibly shift equivalent.*

Proof. By Proposition 4.4 we know that compatible shift equivalence is an equivalence relation. Hence, it will suffice to show that if A and B are elementary shift related via R and S , then R and S are compatible.

Since A and B are elementary shift related via R and S , we have that $A = RS$ and $B = SR$. So choose some path isomorphisms $\psi_A : E_R \times E_S \rightarrow E_A$ and $\psi_B : E_S \times E_S \rightarrow E_B$ and define

$$\phi_R : E_A \times E_R \rightarrow E_R \times E_A \quad \text{and} \quad \phi_S : E_B \times E_S \rightarrow E_S \times E_A$$

by setting $\phi_R := (\text{id}_R \times \psi_B)(\psi_A^{-1} \times \text{id}_R)$ and $\phi_S := (\text{id}_S \times \psi_A)(\psi_B^{-1} \times \text{id}_S)$. Since the lag is $m = 1$, compatibility follows by definition of ϕ_R and ϕ_S . \square

Remark 4.6. When A and B are essential matrices with entries in \mathbb{N} over V and W respectively, it can be shown directly that compatible shift equivalence implies conjugacy of (X_{E_A}, σ_{E_A}) and (X_{E_B}, σ_{E_B}) with a formula for the homeomorphism $h : X_{E_A} \rightarrow X_{E_B}$ which implements the conjugacy. Thus, by Williams' theorem, it follows that A and B are strong shift equivalent. We skip the argument here, because it will follow from Theorem 7.3 that the converse of Corollary 4.5 holds.

5. Representable shift equivalence

Our goal in this section is to determine when a shift equivalence between two matrices can be represented as operators on Hilbert space. This leads to the notion of representable shift equivalence. We show that compatible shift equivalence implies representable shift equivalence, and that a representation of shift equivalence can be chosen so that the graph C^* -algebras act *faithfully* on the Hilbert space.

Let A and B be matrices over \mathbb{N} indexed by V and W respectively, and suppose that R and S are matrices over \mathbb{N} indexed by $V \times W$ and $W \times V$ respectively. To ease some of our notation, we define two matrices indexed by $V \sqcup W$,

$$C = \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \quad \text{and} \quad D = \begin{bmatrix} 0 & R \\ S & 0 \end{bmatrix},$$

so that A and B are shift equivalent with lag m via R and S if and only if

$$CD = DC \quad \text{and} \quad D^2 = C^m.$$

From shift equivalence, there exist path isomorphisms

$$\begin{aligned} \phi_R : E_A \times E_R &\rightarrow E_R \times E_B, & \phi_S : E_B \times E_S &\rightarrow E_S \times E_A, \\ \psi_A : E_R \times E_S &\rightarrow E_A^m, & \psi_B : E_S \times E_R &\rightarrow E_B^m. \end{aligned}$$

We may define path isomorphisms $\phi : E_C \times E_D \rightarrow E_D \times E_C$ and $\psi : E_D^2 \rightarrow E_C^m$ given by

$$\psi = \begin{bmatrix} \psi_A & 0 \\ 0 & \psi_B \end{bmatrix} \quad \text{and} \quad \phi = \begin{bmatrix} 0 & \phi_R \\ \phi_S & 0 \end{bmatrix}.$$

Conversely, all path isomorphisms $\phi : E_C \times E_D \rightarrow E_D \times E_C$ and $\psi : E_D^2 \rightarrow E_C^m$ must be of the above form for some path isomorphisms $\phi_R, \phi_S, \psi_A, \psi_B$ as above. Hence, we see that A and B are compatibly shift equivalent with lag m via R and S if and only if there exist path isomorphisms $\phi : E_C \times E_D \rightarrow E_D \times E_C$ and $\psi : E_D^2 \rightarrow E_C^m$ such that

$$\phi^{(m)} = (\text{id}_D \times \psi)(\psi^{-1} \times \text{id}_D), \tag{5-1}$$

where $\phi^{(m)} : E_C^m \times E_D \rightarrow E_D \times E_C^m$ is the path isomorphism given by

$$\phi^{(m)} := (\phi \times \text{id}_{C^{m-1}})(\text{id}_C \times \phi \times \text{id}_{C^{m-2}}) \cdots (\text{id}_{C^{m-1}} \times \phi).$$

The following is the natural way to represent shift equivalence as bounded operators on Hilbert space, via some choice of path isomorphisms as above.

Definition 5.1. Let A and B be square matrices indexed by V and W respectively, with entries in \mathbb{N} . Let $m \in \mathbb{N} \setminus \{0\}$ and suppose there are matrices R over $V \times W$ and S over $W \times V$ with entries in \mathbb{N} for which there exist path isomorphisms

$$\phi : E_C \times E_D \rightarrow E_D \times E_C \quad \text{and} \quad \psi : E_D^2 \rightarrow E_C^m.$$

We say that R and S are *representable* via ϕ and ψ if there are Cuntz–Krieger families (P_v, S_c) for G_C and (P_v, T_d) for G_D on the same Hilbert space \mathcal{H} with $P_v \neq 0$ for all $v \in V \sqcup W$ such that

$$T_{d_1 d_2} = S_{c_1 \cdots c_m}, \quad \text{when } \psi(d_1 d_2) = c_1 \cdots c_m \text{ for } d_1 d_2 \in E_D^2, \tag{5-2}$$

$$S_c T_d = T_{d'} S_{c'}, \quad \text{when } \phi(cd) = d' c' \text{ for } cd \in E_C \times E_D. \tag{5-3}$$

We say that A and B are *representable shift equivalent* if there are R and S and ϕ and ψ as above so that R and S are representable via ϕ and ψ .

Remark 5.2. When A and B both have no zero columns we may define R and S to be representable via ϕ and ψ by specifying a priori separate Cuntz–Krieger families (S_v, S_c) for G_C and (T_w, T_d) for G_D . It then follows from (5-2) that $S_v = T_v$ for every $v \in V \sqcup W$. Indeed, since no vertex is a source, for every $v \in V$ there exists $c_1 \cdots c_m \in E_C^m$ with $r(c_m) = v$, so we may take $\psi^{-1}(c_1 \cdots c_m) = d_1 d_2$ with $d_1 d_2 \in E_D^2$ and $r(d_2) = v$. So that

$$S_v = S_{c_1 \cdots c_m}^* S_{c_1 \cdots c_m} = T_{d_1 d_2}^* T_{d_1 d_2} = T_v.$$

Proposition 5.3. *Let A and B be essential matrices indexed by V and W respectively, with entries in \mathbb{N} and finitely supported rows. Suppose there are a lag $m \in \mathbb{N} \setminus \{0\}$ and matrices R over $V \times W$ and S over $W \times V$ with entries in \mathbb{N} , and path isomorphisms*

$$\begin{aligned} \phi_R : E_A \times E_R &\rightarrow E_R \times E_B, & \phi_S : E_B \times E_S &\rightarrow E_S \times E_A, \\ \psi_A : E_R \times E_S &\rightarrow E_A^m, & \psi_B : E_S \times E_R &\rightarrow E_B^m \end{aligned}$$

such that

$$(\psi_A \times \text{id}_A)(\text{id}_R \times \phi_S) = (\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S) \quad (5-4)$$

and

$$\phi_S^{(m)} = (\text{id}_S \times \psi_A)(\psi_B^{-1} \times \text{id}_S). \quad (5-5)$$

Then R and S are representable via ϕ and ψ .

Proof. Let $X_{E_D}^+$ be the one-sided subshift for $G_D = (V \sqcup W, E_D)$, and let $\{e_x\}_{x \in X_{E_D}^+}$ be an orthonormal basis for $\ell^2(X_{E_D}^+)$. We denote by $VX_{E_D}^+$ and $WX_{E_D}^+$ the clopen subsets of those $x \in X_{E_D}^+$ such that $s(x) \in V$ and $s(x) \in W$ respectively. We then have the homeomorphisms $\psi_A^\infty : VX_{E_D}^+ \rightarrow X_{E_A}^+$ and $\psi_B^\infty : WX_{E_D}^+ \rightarrow X_{E_B}^+$ given by

$$\begin{aligned} \psi_A^\infty(r_0 s_0 r_1 s_1 \cdots) &= \psi_A(r_0 s_0) \psi_A(r_1 s_1) \cdots, \\ \psi_B^\infty(s_0 r_0 s_1 r_1 \cdots) &= \psi_B(s_0 r_0) \psi_B(s_1 r_1) \cdots \end{aligned}$$

for $r_i \in E_R$ and $s_i \in E_S$.

We define P_v for $v \in V \sqcup W$ by

$$P_v(e_x) = \begin{cases} e_x & \text{if } v = s(x), \\ 0 & \text{otherwise.} \end{cases}$$

For $a \in E_A$ and $x \in X_{E_D}^+$, define S_a via

$$S_a(e_x) = \begin{cases} e_y & \text{if } r(a) = s(x), \\ 0 & \text{otherwise,} \end{cases}$$

where $y := (\psi_A^\infty)^{-1}(a \psi_A^\infty(r_0 s_0 r_1 s_1 \cdots))$ when we write $x = r_0 s_0 \cdots$ for elements $r_i \in E_R$ and $s_i \in E_S$ in case that $r(a) = s(x)$.

For $b \in E_B$ and $x \in X_{E_D}^+$, define S_b via

$$S_b(e_x) = \begin{cases} e_y & \text{if } r(b) = s(x), \\ 0 & \text{otherwise,} \end{cases}$$

where $y := (\text{id}_S \times \psi_A^\infty)^{-1}(\phi_S(b s_0) \psi_A^\infty(r_0 s_1 r_1 s_2 \cdots))$ when we write $x = s_0 r_0 \cdots$ for elements $s_i \in E_S$, $r_i \in E_R$ in the case that $r(b) = s(x)$.

Finally, we define for $d \in E_D$ the operator T_d via

$$T_d(e_x) = \begin{cases} e_{dx} & \text{if } r(d) = s(x), \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to verify that (P_v, T_d) is a Cuntz–Krieger family for G_D . So we first verify that (P_v, S_c) is a Cuntz–Krieger family for G_C . Since concatenation for S_a is done simply through the homeomorphism ψ_A^∞ , it is easy to show that, for $a \in E_A$ or $v \in V$, we have $S_a^* S_a = P_{r(a)}$ and $\sum_{s(e)=v} S_e S_e^* = P_v$. Now, to show the same for $b \in E_B$, for $y = s'_0 r'_0 s'_1 r'_1 \cdots$, when $s'_i \in E_S$ and $r'_i \in E_R$, we write $(\text{id}_B \times \text{id}_S \times \psi_A^\infty)^{-1}(\phi_S^{-1} \times \text{id})(s'_0 \psi_A^\infty(r'_0 s'_1 r'_1 \cdots)) = b' s_0 r_0 \cdots$ for $b' \in E_B$, $s_i \in E_S$, $r_i \in E_R$, so that

$$S_b^*(e_y) = \begin{cases} e_{s_0 r_0 \cdots} & \text{if } r(b) = s(x), b = b', \\ 0 & \text{otherwise.} \end{cases}$$

From this formula it follows that $S_b^* S_b = P_{r(b)}$, and that for $w \in W$ we have $\sum_{s(e)=w} S_e S_e^* = P_w$. Thus, we see that (P_v, S_c) is a Cuntz–Krieger family for G_C . Since clearly $P_v \neq 0$ for every $v \in V \sqcup W$, we are left with verifying (5-2) and (5-3).

It is clear from the definition of S_a for $a \in E_A$ that for $r \in E_R$ and $s \in E_S$ we have $T_{rs} = S_{a_1 \cdots a_m}$ when $\psi_A(rs) = a_1 \cdots a_m$. We next show that $S_b T_s = T_{s'} S_a$ when $\phi_S(bs) = s'a$ for $b \in E_B$ and $s \in E_S$. Indeed, let $x \in X_{E_D}^+$ with $s(x) \in V$, and write $x = r_0 s_0 \cdots$ so that $S_b T_s(e_x) = e_z$ and $T_{s'} S_a(e_x) = e_{z'}$, where

$$\begin{aligned} z &= (\text{id}_S \times \psi_A^\infty)^{-1} \phi_S(bs) \psi_A^\infty(r_0 s_0 \cdots), \\ z' &= s'(\psi_A^\infty)^{-1}(a \psi_A^\infty(r_0 s_0 \cdots)). \end{aligned}$$

Since $\phi(bs) = s'a$, it follows that $z = z'$, so that $S_b T_s = T_{s'} S_a$.

Next, we show that $T_{sr} = S_{b_1 \cdots b_m}$ when $\psi_B(sr) = b_1 \cdots b_m$. Indeed, let $x \in X_{E_D}^+$ with $s(x) \in W$, and write $x = s_0 r_0 \cdots$. Then we have that $S_{b_1 \cdots b_m}(e_x) = e_z$, where

$$z = (\text{id}_S \times \psi_A^\infty)^{-1}(\phi_S^{(m)}(b_1 \cdots b_m s_0) \psi_A^\infty(r_0 s_1 \cdots)).$$

From (5-5) it follows that

$$z = (\psi_B)^{-1}(b_1 \cdots b_m) s_0 r_0 s_1 \cdots.$$

Thus, we see that if $\psi_B(sr) = b_1 \cdots b_m$, then $T_{sr} = S_{b_1 \cdots b_m}$.

Finally, we show that when $\phi_R(ar) = r'b$ we have $S_a T_r = T_{r'} S_b$. Indeed, let $x \in X_{E_D}^+$ with $s(x) \in W$, and write $x = s_0 r_0 \cdots$ so that $S_a T_r(e_x) = e_z$ and $T_{r'} S_b(e_x) = e_{z'}$, where

$$\begin{aligned} z &= (\psi_A^\infty)^{-1}(a \psi_A^\infty(r s_0 r_0 \cdots)), \\ z' &= r'(\text{id}_S \times \psi_A^\infty)^{-1}(\phi_S(bs_0) \psi_A^\infty(r_0 s_1 r_1 \cdots)). \end{aligned}$$

But by (5-4) and the fact that $\phi_R(ar) = r'b$ we get

$$\begin{aligned} z' &= (\psi_A^\infty)^{-1}(\text{id}_A \times \psi_A)(\phi_R^{-1} \times \text{id}_S)(r' b s_0) \psi_A^\infty(r_0 s_1 r_1 \cdots) \\ &= (\psi_A^\infty)^{-1}(\text{id}_A \times \psi_A)(a r s_0 \cdot \psi_A^\infty(r_0 s_1 r_1 \cdots)) = (\psi_A^\infty)^{-1}(a \psi_A^\infty(r s_0 r_0 \cdots)) = z. \end{aligned}$$

Hence, we get that $z' = z$, so that $S_a T_r = T_{r'} S_b$. Thus, we have shown that R and S are representable via ϕ and ψ . \square

Remark 5.4. We note that for two essential matrices A and B with entries in \mathbb{N} to be representable shift equivalent, we need only know the validity of the two asymmetric equations (5-4) and (5-5), as opposed to the symmetric equations in (4-1).

Using Szymański’s uniqueness theorem [44, Theorem 1.2], we can upgrade a representation of a shift equivalence to be injective in the following sense.

Corollary 5.5. *Let A and B be essential matrices indexed by V and W respectively, with entries in \mathbb{N} and finitely supported rows. Suppose that A and B are compatibly shift equivalent via R and S and path isomorphisms ϕ and ψ . Then R and S are representable via ϕ and ψ . In fact, there are Cuntz–Krieger families (P_v, S_c) for G_C and (P_v, T_d) for G_D satisfying (5-2) and (5-3) so that both of the canonical surjections $q_C : C^*(G_C) \rightarrow C^*(P_v, S_c)$ and $q_D : C^*(G_D) \rightarrow C^*(P_v, T_d)$ are injective.*

Proof. From Lemma 4.2 we know (5-4) holds. Since (5-5) holds by definition, by Proposition 5.3 there are Cuntz–Krieger families $S := (P_v, S_c)$ for G_C and $T := (P_v, T_d)$ for G_D satisfying (5-2) and (5-3).

Let $z \in \mathbb{T}$ be some unimodular scalar. Then we may define two operator families $S^{(z)} := (P_v, z \cdot S_c)$ and $T^{(z)} := (P_v, T_d^{(z)})$, where

$$T_d^{(z)} := \begin{cases} T_d & \text{if } d \in E_R, \\ z^m \cdot T_d & \text{if } d \in E_S. \end{cases}$$

Then clearly $S^{(z)}$ and $T^{(z)}$ are Cuntz–Krieger families satisfying (5-2) and (5-3).

Let $(z_n)_{n \in \mathbb{N}}$ be a countable dense subset of \mathbb{T} . We take $S' := \bigoplus_{n=1}^\infty S^{(z_n)}$ and $T' := \bigoplus_{n=1}^\infty T^{(z_n)}$, which are still Cuntz–Krieger families satisfying (5-2) and (5-3). By Szymański’s uniqueness theorem [44, Theorem 1.2], it suffices to show that for every cycle $c_1 \cdots c_\ell$ without exits in G_C , and every cycle $d_1 \cdots d_{2\ell}$ without exits in G_D (which is necessarily of even length since G_D is bipartite), the spectrum of the operators $S_{c_1 \cdots c_\ell}$ and $T_{d_1 \cdots d_{2\ell}}$ contains the entire unit circle.

Since $S_{c_1 \cdots c_\ell}$ and $T_{d_1 \cdots d_{2\ell}}$ are unitaries on the ranges of $P_{s(c_1)}$ and $P_{s(d_1)}$ respectively, each of their spectra must contain some element in the unit circle. But since for every $n \in \mathbb{N}$ we have that $z_n^\ell \cdot S_{c_1 \cdots c_\ell}$ is a direct summand of $S'_{c_1 \cdots c_\ell}$, and $z_n^{m \cdot \ell} \cdot T_{d_1 \cdots d_{2\ell}}$ is a direct summand of $T'_{d_1 \cdots d_{2\ell}}$, we see that the spectra of $S'_{c_1 \cdots c_\ell}$ and $T'_{d_1 \cdots d_{2\ell}}$ both contain a dense subset of \mathbb{T} , and hence \mathbb{T} itself. Thus, by Szymański’s uniqueness theorem we get that the canonical surjections $C^*(G_C) \rightarrow C^*(P'_v, S'_c)$ and $C^*(G_D) \rightarrow C^*(P'_v, T'_d)$ are injective. □

Remark 5.6. Suppose we have two essential matrices A and B with entries in \mathbb{N} indexed by V and W respectively, and that R, S are matrices that comprise a representable shift equivalence of lag m via path isomorphisms ϕ and ψ . It can be shown directly that A and B are compatible shift equivalent with lag m via R and S , together with the 1-1 and 2-2 corners of ψ and the 1-2 and 2-1 corners of ϕ . We skip the proof since representable shift equivalence implies compatible shift equivalence by Theorem 7.3.

6. Strong Morita shift equivalence

In this section we introduce and study strong Morita shift equivalence. This equivalence relation is expressed in terms of a specific strong Morita equivalence between Pimsner dilations and is implied by

representable shift equivalence. Strong Morita shift equivalence turns out to imply the existence of a stable equivariant isomorphism of graph C*-algebras that also preserves the diagonal subalgebras.

Suppose A and B are essential matrices over \mathbb{N} indexed by V and W respectively and have finitely supported rows. Suppose there are matrices R and S with entries in \mathbb{N} , and let C and D be as described in the previous section. Suppose further that we (only) have a path isomorphism $\psi : E_D^2 \rightarrow E_C^m$, which is then the direct sum of path isomorphisms

$$\psi_A : E_R \times E_S \rightarrow E_A^m \quad \text{and} \quad \psi_B : E_S \times E_R \rightarrow E_B^m.$$

Now let (S_v, S_c) be a CK family that generates the graph C*-algebra $C^*(G_C) = C^*(G_A) \oplus C^*(G_B)$ of G_C , and let (T_v, T_d) be a CK family that generates the graph C*-algebra $C^*(G_D)$. We define

$$\begin{aligned} \mathcal{A}_n^C &:= \overline{\text{span}}\{S_\lambda S_{\lambda'}^* \mid \lambda, \lambda' \in E_C^n, r(\lambda) \in V\}, & \mathcal{B}_n^C &:= \overline{\text{span}}\{S_\lambda S_{\lambda'}^* \mid \lambda, \lambda' \in E_C^n, r(\lambda) \in W\}, \\ \mathcal{A}_n^D &:= \overline{\text{span}}\{T_\mu T_{\mu'}^* \mid \mu, \mu' \in E_D^n, r(\mu) \in V\}, & \mathcal{B}_n^D &:= \overline{\text{span}}\{T_\mu T_{\mu'}^* \mid \mu, \mu' \in E_D^n, r(\mu) \in W\}, \end{aligned}$$

with direct limits $\mathcal{A}_\infty^C, \mathcal{B}_\infty^C, \mathcal{A}_\infty^D$ and \mathcal{B}_∞^D . Then it follows that $\mathcal{A}_\infty^C \oplus \mathcal{B}_\infty^C$ is canonically isomorphic to the fixed-point algebra $C^*(G_C)_0$ of $C^*(G_C)$ with its canonical gauge action, and that $\mathcal{A}_\infty^D \oplus \mathcal{B}_\infty^D$ is canonically isomorphic to the fixed-point algebra $C^*(G_D)_0$ of $C^*(G_D)$ with its canonical gauge action.

Thinking of $X(A)$ and $X(B)$ as block diagonal C*-subcorrespondences of $X(C)$ in the natural way, we get that the Pimsner dilations $X(A)_\infty$ and $X(B)_\infty$ are naturally identified with the direct limits of C*-correspondences

$$X(A)_n := \overline{\text{span}}\{S_\lambda S_{\lambda'}^* \mid \lambda \in E_C^{n+1}, \lambda' \in E_C^n, s(\lambda) \in V\},$$

which is a C*-correspondence from \mathcal{A}_n^C to \mathcal{A}_{n+1}^C , and

$$X(B)_n := \overline{\text{span}}\{S_\lambda S_{\lambda'}^* \mid \lambda \in E_C^{n+1}, \lambda' \in E_C^n, s(\lambda) \in W\},$$

which is a C*-correspondence from \mathcal{B}_n^C to \mathcal{B}_{n+1}^C .

Similarly, thinking of $X(R)$ and $X(S)$ as the off-diagonal C*-subcorrespondences of $X(D)$ in the natural way, we get the C*-correspondences (which are only denoted as) $X(R)_\infty$ and $X(S)_\infty$ as the direct limits of C*-correspondences

$$X(R)_n := \overline{\text{span}}\{T_\mu T_{\mu'}^* \mid \mu \in E_D^{n+1}, \mu' \in E_D^n, s(\mu) \in V\},$$

which is a C*-correspondence from \mathcal{B}_n^D to \mathcal{A}_{n+1}^D , and

$$X(S)_n := \overline{\text{span}}\{T_\mu T_{\mu'}^* \mid \mu \in E_D^{n+1}, \mu' \in E_D^n, s(\mu) \in W\},$$

which is a C*-correspondence from \mathcal{A}_n^D to \mathcal{B}_{n+1}^D . Note, however, that $X(R)_\infty$ and $X(S)_\infty$ are not Pimsner dilations in the sense described in Section 3, because they are C*-correspondences over possibly different left and right coefficient C*-algebras.

Recall that $X(D)_\infty$ is the C*-correspondence over $C^*(G_D)_0 = \mathcal{A}_\infty^D \oplus \mathcal{B}_\infty^D$ given by $X(D)_\infty = X(D) \cdot (\mathcal{A}_\infty^D \oplus \mathcal{B}_\infty^D)$, which is identified with the direct limit

$$\overline{\text{span}}\{T_\mu T_{\mu'}^* \mid \mu \in E_D^{n+1}, \mu' \in E_D^n, n \in \mathbb{N}\}.$$

The $\mathcal{B}_\infty^D\text{-}\mathcal{A}_\infty^D$ correspondence $X(R)_\infty$ and the $\mathcal{A}_\infty^D\text{-}\mathcal{B}_\infty^D$ correspondence $X(S)_\infty$ coincide with the 1-2 corner and 2-1 corner (respectively) of the Pimsner dilation $X(D)_\infty$ and satisfy the equalities

$$X(R)_\infty = X(R) \cdot \mathcal{B}_\infty^D, \quad X(S)_\infty = X(S) \cdot \mathcal{A}_\infty^D.$$

Now, using the map $\psi : E_D^2 \rightarrow E_C^m$, for each $k \in \mathbb{N}$ and $\mu_1, \dots, \mu_k \in E_D^{2k}$ such that $\mu_1 \cdots \mu_k \in E_D^{2k}$ we may define $\psi_k : E_D^{2k} \rightarrow E_C^{mk}$ by setting $\psi_k(\mu_1 \cdots \mu_k) = \psi(\mu_1) \cdots \psi(\mu_k)$. This then gives rise to $*$ -isomorphisms $\Psi_k^A : \mathcal{A}_{2k}^D \rightarrow \mathcal{A}_{mk}^C$ and $\Psi_k^B : \mathcal{B}_{2k}^D \rightarrow \mathcal{B}_{mk}^C$ by setting $\Psi_k^A(T_\mu T_{\mu'}^*) = S_{\psi_k(\mu)} S_{\psi_k(\mu')}^*$ and $\Psi_k^B(T_\mu T_{\mu'}^*) = S_{\psi_k(\mu)} S_{\psi_k(\mu')}^*$. Since these maps are compatible with direct limits, we obtain two $*$ -isomorphisms $\Psi_\infty^A : \mathcal{A}_\infty^D \rightarrow \mathcal{A}_\infty^C$ and $\Psi_\infty^B : \mathcal{B}_\infty^D \rightarrow \mathcal{B}_\infty^C$ that we will use as identifications between the coefficient C^* -algebras. For instance, this allows us to turn $X(R)_\infty$ into a $\mathcal{B}_\infty^C\text{-}\mathcal{A}_\infty^C$ -bimodule, where the left and right actions are implemented via Ψ_∞^A and Ψ_∞^B respectively, and the inner product via Ψ_∞^B .

Definition 6.1. Let A and B be essential matrices over \mathbb{N} indexed by V and W respectively, with finitely supported rows. We say that $X(A)_\infty$ and $X(B)_\infty$ are *strong Morita shift equivalent* if there are a lag $m \in \mathbb{N} \setminus \{0\}$ and matrices R over $V \times W$ and S over $W \times V$ over \mathbb{N} together with path isomorphisms

$$\psi_A : E_R \times E_S \rightarrow E_A^m, \quad \psi_B : E_S \times E_R \rightarrow E_B^m$$

such that $X(R)_\infty$ is a strong Morita equivalence between $X(A)_\infty$ and $X(B)_\infty$, up to the identifications Ψ_∞^A and Ψ_∞^B . More precisely, when $X(R)_\infty$ is considered as a $\mathcal{B}_\infty^C\text{-}\mathcal{A}_\infty^C$ -bimodule via Ψ_∞^A and Ψ_∞^B , there exists a unitary bimodule isomorphism $U : X(A)_\infty \otimes X(R)_\infty \rightarrow X(R)_\infty \otimes X(B)_\infty$ such that for every $a \in \mathcal{A}_\infty^C$, $b \in \mathcal{B}_\infty^C$ and $\xi \in X(A)_\infty \otimes X(R)_\infty$ we have

$$U(a \cdot \xi \cdot b) = a \cdot U(\xi) \cdot b.$$

Henceforth, we will no longer belabor the point of distinguishing between \mathcal{A}_∞^C and \mathcal{A}_∞^D and between \mathcal{B}_∞^C and \mathcal{B}_∞^D . However, we emphasize that this identification is important in the above definition, and depends on the choice of the maps ψ_A and ψ_B . We have already seen that

$$\begin{aligned} X(A)_\infty &= X(A) \cdot \mathcal{A}_\infty, & X(B)_\infty &= X(B) \cdot \mathcal{B}_\infty, \\ X(R)_\infty &= X(R) \cdot \mathcal{B}_\infty, & X(S)_\infty &= X(S) \cdot \mathcal{A}_\infty, \end{aligned}$$

so that by the above discussion and identifications using Ψ_∞^A and Ψ_∞^B , as $\mathcal{B}_\infty\text{-}\mathcal{A}_\infty$ correspondences we may canonically identify

$$\begin{aligned} X(A)_\infty \otimes_{\mathcal{A}_\infty} X(R)_\infty &\cong X(A) \otimes_{\mathcal{A}} X(R) \cdot \mathcal{B}_\infty, \\ X(R)_\infty \otimes_{\mathcal{B}_\infty} X(B)_\infty &\cong X(R) \otimes_{\mathcal{B}} X(B) \cdot \mathcal{B}_\infty. \end{aligned}$$

Proposition 6.2. *Let A and B be essential matrices over \mathbb{N} indexed by V and W respectively, with finitely supported rows. Suppose A and B are representable shift equivalent with lag $m \in \mathbb{N}$ via R and S , together with path isomorphisms ϕ and ψ . Then $X(A)_\infty$ and $X(B)_\infty$ are strong Morita shift equivalent with lag m via R , S and ψ .*

Proof. By Corollary 5.5 we have a Cuntz–Krieger family (P_v, S_c) for G_C generating $C^*(G_C)$ and a Cuntz–Krieger family (P_v, T_d) for G_D generating $C^*(G_D)$ on the same Hilbert space \mathcal{H} , satisfying (5-2) and (5-3). Thus, we are in the context of the discussion above.

By the identifications preceding the theorem, we may define maps

$$\begin{aligned} U_{AR} &: X(A)_\infty \otimes_{\mathcal{A}_\infty} X(R)_\infty \rightarrow X(A) \cdot X(R) \cdot \mathcal{B}_\infty, \\ U_{RB} &: X(R)_\infty \otimes_{\mathcal{B}_\infty} X(B)_\infty \rightarrow X(R) \cdot X(B) \cdot \mathcal{B}_\infty \end{aligned}$$

(where the notation $X(A) \cdot X(R) \cdot \mathcal{B}_\infty$ and $X(R) \cdot X(B) \cdot \mathcal{B}_\infty$ are understood as the closed linear span of products), by setting

$$U_{AR}(S_a \otimes T_r \cdot w) = S_a T_r \cdot w \quad \text{and} \quad U_{RB}(T_r \otimes S_b \cdot w) = T_r S_b \cdot w$$

for $a \in E_A$, $r \in E_R$, $b \in E_B$ and $w \in \mathcal{B}_\infty$. It is straightforward to show that U_{AR} and U_{RB} are well-defined unitary right \mathcal{B}_∞ -module maps. Thus, we are left with showing that U_{AR} and U_{RB} are left \mathcal{A}_∞ -module maps.

We first show that U_{AR} is a left \mathcal{A}_∞ -module map. We let $S_\lambda S_{\lambda'}^* \in \mathcal{A}_\infty$ for $\lambda, \lambda' \in E_A^{mk}$, and note that it will suffice to show that for $a \in E_A$ and $r \in E_R$ we have

$$U_{AR}(S_\lambda S_{\lambda'}^*(S_a \otimes T_r)) = S_\lambda S_{\lambda'}^* S_a T_r.$$

To prove this, let $e \in E_A$ be some edge so that $s(e) = r(\lambda) = r(\lambda')$, and suppose that $\lambda = a_1 v$ and $\lambda' = a'_1 v'$ for $a_1, a'_1 \in E_A$. Write

$$v e = \psi(r_1 s_1) \cdots \psi(r_k s_k) \quad \text{and} \quad v' e = \psi(r'_1 s'_1) \cdots \psi(r'_k s'_k)$$

for $r_i, r'_i \in E_R$ and $s_i, s'_i \in E_S$. Then, we have

$$\begin{aligned} S_\lambda S_{\lambda'}^* S_a \otimes T_r &= \delta_{a, a'_1} \cdot S_{a_1} S_v S_{v'}^* \otimes T_r = \delta_{a, a'_1} \cdot S_{a_1} \sum_{e \in s^{-1}(r(\lambda))} S_{v e} S_{v' e}^* \otimes T_r \\ &= \delta_{a, a'_1} \cdot S_{a_1} \otimes \sum_{e \in s^{-1}(r(\lambda))} T_{r_1 s_1 \cdots r_k s_k} T_{r'_1 s'_1 \cdots r'_k s'_k}^* T_r \\ &= \delta_{a, a'_1} \cdot S_{a_1} \otimes T_{r_1} \sum_{e \in s^{-1}(r(\lambda))} \delta_{r'_1, r} \cdot T_{s_1 \cdots r_k s_k} T_{s'_1 \cdots r'_k s'_k}^* \\ &= \delta_{a, a'_1} \cdot S_{a_1} \otimes T_{r_1} \sum_{e \in s^{-1}(r(\lambda))} \delta_{r'_1, r} \cdot \sum_{f \in s^{-1}(r(\lambda e))} S_{\psi_k^{-1}(s_1 \cdots r_k s_k f)} S_{\psi_k^{-1}(s'_1 \cdots r'_k s'_k f)}, \end{aligned}$$

where in the above calculations we used the identifications via Ψ_∞^A and Ψ_∞^B . Essentially the same calculation, using (5-2) instead, will show that

$$S_\lambda S_{\lambda'}^* S_a T_r = \delta_{a, a'_1} \cdot S_{a_1} T_{r_1} \sum_{e \in s^{-1}(r(\lambda))} \delta_{r_1, r} \cdot \sum_{f \in s^{-1}(r(\lambda e))} S_{\psi_k^{-1}(s_1 \cdots r_k s_k f)} S_{\psi_k^{-1}(s'_1 \cdots r'_k s'_k f)}.$$

Thus, we see that U_{AR} is a left \mathcal{A}_∞ -module map.

Next, we show that U_{RB} is a left \mathcal{A}_∞ -module map. We let $S_\lambda S_{\lambda'}^* \in \mathcal{A}_\infty$ with $\lambda, \lambda' \in E_A^{mk}$, and note that it will suffice to show that for $r \in E_R$ and $b \in E_B$ we have

$$U_{RB}(S_\lambda S_{\lambda'}^* T_r \otimes S_b) = S_\lambda S_{\lambda'}^* T_r S_b.$$

Write $\lambda = \psi(r_1 s_1) \cdots \psi(r_k s_k)$ and $\lambda' = \psi(r'_1 s'_1) \cdots \psi(r'_k s'_k)$, and then further write $\psi(s_1 r_2) = b_1 \sigma$ and $\psi(s'_1 r'_2) = b'_1 \sigma'$ for $b_1, b'_1 \in E_B$. Then, we have

$$\begin{aligned} S_\lambda S_{\lambda'}^* T_r \otimes S_b &= T_{r_1 s_1 \cdots r_k s_k} T_{r'_1 s'_1 \cdots r'_k s'_k}^* T_r \otimes S_b \\ &= \delta_{r'_1, r} T_{r_1} \sum_{r \in r^{-1}(\lambda)} T_{s_1 r_2 \cdots s_k r} T_{s'_1 r'_2 \cdots s'_k r}^* \otimes S_b \\ &= \delta_{r'_1, r} T_{r_1} \otimes \sum_{r \in r^{-1}(\lambda)} S_{\psi(s_1 r_2) \cdots \psi(s_k r)} S_{\psi(s'_1 r'_2) \cdots \psi(s'_k r)}^* S_b \\ &= \delta_{r'_1, r} \delta_{b'_1, b} \cdot T_{r_1} \otimes S_{b_1} \sum_{r \in r^{-1}(\lambda)} S_{\sigma \psi(s_2 r_3) \cdots \psi(s_k r)} S_{\sigma' \psi(s'_2 r'_3) \cdots \psi(s'_k r)}^*, \end{aligned}$$

where in the above calculation we used the identifications via Ψ_∞^A and Ψ_∞^B . Essentially the same calculation, using (5-2) instead, will show that

$$S_\lambda S_{\lambda'}^* T_r S_b = \delta_{r'_1, r} \delta_{b'_1, b} \cdot T_{r_1} S_{b_1} \sum_{r \in r^{-1}(\lambda)} S_{\sigma \psi(s_2 r_3) \cdots \psi(s_k r)} S_{\sigma' \psi(s'_2 r'_3) \cdots \psi(s'_k r)}^*.$$

Thus, we see that U_{BR} is a left \mathcal{A}_∞ -module map.

Thus, to conclude the proof we need only show the equality

$$X(A) \cdot X(R) \cdot \mathcal{B}_\infty = X(R) \cdot X(B) \cdot \mathcal{B}_\infty.$$

However, it is clear from (5-3) that $X(A) \cdot X(R) = X(R) \cdot X(B)$, so we are done. Thus, the map $U_{RB}^{-1} \circ U_{AR}$ is a unitary isomorphism, showing that $X(R)_\infty$ is a strong Morita equivalence between $X(A)_\infty$ and $X(B)_\infty$. \square

Suppose A and B are matrices indexed by V and W respectively, and suppose there are matrices R and S , so that C and D are as described in the previous section. Suppose we have a path isomorphism $\psi : E_D^2 \rightarrow E_C^m$.

Assume we have a unitary $U : X(A)_\infty \otimes X(R)_\infty \rightarrow X(R)_\infty \otimes X(B)_\infty$ such that for every $\xi \in X(A)_\infty \otimes X(R)_\infty$ and every $a \in \mathcal{A}_\infty$ and $b \in \mathcal{B}_\infty$ we have $U(b\xi a) = bU(\xi)a$. We refer the reader to [1] and [36, Section 2] for the basic theory of Morita equivalence, linking algebras and crossed products by Hilbert bimodules that we shall need in what follows. We denote by \mathcal{L} the linking algebra of $X(R)_\infty$ given by

$$\mathcal{L} := \begin{bmatrix} \mathcal{A}_\infty & X(R)_\infty \\ X(R)_\infty^* & \mathcal{B}_\infty \end{bmatrix},$$

and by W the \mathcal{L} -imprimitivity bimodule

$$W := \begin{bmatrix} X(A)_\infty & X(R)_\infty \otimes_{\mathcal{B}_\infty} X(B)_\infty \\ X(R)_\infty^* \otimes_{\mathcal{A}_\infty} X(A)_\infty & X(B)_\infty \end{bmatrix}.$$

It is shown in the proof of [1, Theorem 4.2] that there are two complementary and full projections $p_A, p_B \in \mathcal{M}(\mathcal{O}(W))$ (corresponding to the 2×2 block form of W) together with two $*$ -isomorphisms $\varphi_E : \mathcal{O}(X(A)_\infty) \rightarrow p_A \mathcal{O}(W) p_A$ and $\varphi_F : \mathcal{O}(X(B)_\infty) \rightarrow p_B \mathcal{O}(W) p_B$ given as follows. For $\xi \in X(A)_\infty$

and $\eta \in X(B)_\infty$ we let

$$\tilde{\xi} := \begin{bmatrix} \xi & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad \tilde{\eta} := \begin{bmatrix} 0 & 0 \\ 0 & \eta \end{bmatrix},$$

so that the maps φ_E and φ_F are given by

$$\varphi_E(S_\xi) = S_{\tilde{\xi}} \quad \text{and} \quad \varphi_F(S_\eta) = S_{\tilde{\eta}}.$$

In particular, we see that φ_E and φ_F are gauge equivariant.

Since W is an imprimitivity bimodule, again by [1, Theorem 2.9] we get that $\mathcal{O}(W)_0 \cong \mathcal{L}$. In particular, we get that up to the same identification, the direct sum of diagonal subalgebras $\mathcal{D}_{E_A} \oplus \mathcal{D}_{E_B}$, according to $p_A, p_B \in \mathcal{M}(\mathcal{O}(W))$, coincides with

$$\mathcal{D}_W := \overline{\text{span}}\{S_\zeta S_{\zeta'}^* \mid \zeta, \zeta' \in W^{\otimes n}, n \in \mathbb{N}\}.$$

Hence, we get that $\varphi_E(\mathcal{D}_{E_A}) = p_A \mathcal{D}_W p_A$ and $\varphi_F(\mathcal{D}_{E_B}) = p_B \mathcal{D}_W p_B$. Let

$$N(\mathcal{D}_W) = \{n \in \mathcal{O}(W) \mid n^* \mathcal{D}_W n, n \mathcal{D}_W n^* \subseteq \mathcal{D}_W\}$$

be the set of normalizers of \mathcal{D}_W , and denote by

$$N_\star(\mathcal{D}_W) = \{n \in N(\mathcal{D}_W) \mid \text{there exists } k \in \mathbb{Z} \text{ such that } \gamma_z(n) = z^k n\}$$

the *homogeneous* normalizers of \mathcal{D}_W .

Proposition 6.3. *Let A and B be essential matrices over \mathbb{N} , indexed by V and W respectively, with finitely supported rows. Suppose that $X(A)_\infty$ and $X(B)_\infty$ are strong Morita shift equivalent. Then there is an equivariant \ast -isomorphism $\varphi : C^\ast(G_A) \otimes \mathbb{K} \rightarrow C^\ast(G_B) \otimes \mathbb{K}$, with $\varphi(\mathcal{D}_{E_A} \otimes c_0) = \mathcal{D}_{E_B} \otimes c_0$.*

Proof. We are in the situation where we can apply the implication (7) \Rightarrow (8) in [7, Corollary 11.3]. By the description of $C^\ast(G_A) \cong C^\ast(\mathcal{G}_{E_A})$ and $C^\ast(G_B) \cong C^\ast(\mathcal{G}_{E_B})$ as groupoid C^\ast -algebras, we get that item (8) in [7, Corollary 11.3] is equivalent to $C^\ast(G_A)$ and $C^\ast(G_B)$ being stably equivariant diagonal-isomorphic. Indeed, this is because \mathbb{Z} -coactions correspond to topological \mathbb{Z} -gradings by [42, Remark 6], which in turn correspond to \mathbb{T} -actions by [42, Theorem 3]. This means that the second equality in item (8) of [7, Corollary 11.3] is equivalent to equivariance of the isomorphism.

Thus, to prove our result it will suffice to prove item (7) in [7, Corollary 11.3]. Everything is set up in the discussion preceding the proposition, except for one thing. We are left with showing that $p_A \mathcal{O}(W) p_B$ is the closed linear span of $p_A N_\star(\mathcal{D}_W) p_B$. However, we know that $\mathcal{O}(W)$ is the closed linear span of its graded subspaces $\mathcal{O}(W)_n$ for $n \in \mathbb{Z}$. Hence, $p_A \mathcal{O}(W) p_B$ is the closed linear span of $p_A \mathcal{O}(W)_n p_B$. By [1, Theorem 2.9] we have for all $n \in \mathbb{Z}$ that

- (1) $\mathcal{O}(W)_n \cong W^{\otimes n}$ for $n > 0$,
- (2) $\mathcal{O}(W)_0 \cong \mathcal{L}$, and
- (3) $\mathcal{O}(W)_n \cong (W^{\otimes n})^\ast$ for $n < 0$.

Thus, we are left with showing that each $p_A \mathcal{O}(W)_n p_B$ is the closed linear span of its normalizers (which are automatically n -homogeneous).

For $n = 0$ we get that $p_A \mathcal{O}(W)_0 p_B \cong X(R)_\infty$, for $n > 0$ we get that $p_A \mathcal{O}(W)_n p_B \cong X(A)_\infty^n \otimes X(R)_\infty$, and for $n < 0$ we get that $p_A \mathcal{O}(W)_n p_B \cong (X(A)_\infty^*)^n \otimes X(R)_\infty$.

Thus, $p_A \mathcal{O}(W)_n p_B$ is the closed linear span of elements of the form $S_\lambda S_{\lambda'}^* S_\alpha T_r$ (for $n \geq 0$) or $S_\lambda S_{\lambda'}^* S_\alpha^* T_r$ (for $n < 0$) for $\lambda, \lambda' \in E_A^\ell$, $\alpha \in E_A^{|n|}$ and $r \in E_R$ for some $\ell \in \mathbb{N}$. In order to show that these elements are normalizers for \mathcal{D}_W , and since $S_\lambda S_{\lambda'}^* S_\alpha, S_\lambda S_{\lambda'}^* S_\alpha^* \in N(\mathcal{D}_{E_A})$, it will suffice to show that $T_r \mathcal{D}_{E_B} T_r^* \subseteq \mathcal{D}_{E_A}$ and that $T_r^* \mathcal{D}_{E_A} T_r \subseteq \mathcal{D}_{E_B}$ for $r \in E_R$.

We know already from the inclusions $\mathcal{D}_{E_A} \subseteq \mathcal{A}_\infty$ and $\mathcal{D}_{E_B} \subseteq \mathcal{B}_\infty$ that \mathcal{D}_{E_A} is the closed linear span of elements of the form

$$S_{\psi_k(r_1 s_1 \dots r_k s_k)} S_{\psi_k(r_1 s_1 \dots r_k s_k)}^*$$

for paths $r_1 s_1 \dots r_k s_k \in E_D^{2k}$ with $r_i \in E_R$ and $s_i \in E_S$, and that \mathcal{D}_{E_B} is the closed linear span of elements of the form

$$S_{\psi_k(s_1 r_1 \dots s_k r_k)} S_{\psi_k(s_1 r_1 \dots s_k r_k)}^*$$

for paths $s_1 r_1 \dots s_k r_k \in E_D^{2k}$, with $r_i \in E_R$ and $s_i \in E_S$. So for such paths we compute

$$T_r S_{\psi_k(s_1 r_1 \dots s_k r_k)} S_{\psi_k(s_1 r_1 \dots s_k r_k)}^* T_r^* = \sum_{s \in E_S} T_r T_{s_1 r_1 \dots s_k r_k} T_{s_1 r_1 \dots s_k r_k}^* T_r^* = \sum_{s \in E_S} S_{\psi_{k+1}(r s_1 r_1 \dots s_k r_k s)} S_{\psi_{k+1}(r s_1 r_1 \dots s_k r_k s)}^*$$

is in \mathcal{D}_{E_B} . On the other hand,

$$T_r^* S_{\psi_k(r_1 s_1 \dots r_k s_k)} S_{\psi_k(r_1 s_1 \dots r_k s_k)}^* T_r = \delta_{r, r_1} \cdot \sum_{r \in E_R} S_{\psi_k(s_1 \dots r_k s_k r)} S_{\psi_k(s_1 \dots r_k s_k r)}^*$$

is in \mathcal{D}_{E_A} . Thus, $T_r \mathcal{D}_{E_B} T_r^* \subseteq \mathcal{D}_{E_A}$ and $T_r^* \mathcal{D}_{E_A} T_r \subseteq \mathcal{D}_{E_B}$ as required. \square

7. Shift equivalences through the lens

In this section we orient various equivalence relations between strong shift equivalence and shift equivalence. We will assume some familiarity with crossed product C^* -algebras and K-theory of C^* -algebras. We recommend [48] for the basic theory of crossed product C^* -algebras, [43] for the basic K-theory for C^* -algebras, and especially [41, Chapter 7] for K-theory and crossed products of graph algebras by their gauge actions.

Suppose now that $G = (V, E)$ is a directed graph, and let γ be the gauge unit circle action on $C^*(G)$. We denote by $\hat{\gamma}$ the dual \mathbb{Z} action on $C^*(G) \rtimes_\gamma \mathbb{T}$. By [48, Lemma 2.75] we have that $[C^*(G) \rtimes \mathbb{T}] \otimes \mathbb{K}$ and $[C^*(G) \otimes \mathbb{K}] \rtimes \mathbb{T}$ are $*$ -isomorphic via the map $f \otimes K \mapsto f \cdot K$ defined for $f \in C(\mathbb{T}; C^*(G))$ and $K \in \mathbb{K}$, and that this map intertwines the action $\hat{\gamma}^G \otimes \text{id}$ with the dual action $\gamma^G \otimes \text{id}$. Using this equivariant identification together with [48, Corollary 2.48] we obtain the following standard fact which we leave for the reader to verify.

Proposition 7.1. *Let G and G' be directed graphs, and suppose there exists a $*$ -isomorphism $\varphi : C^*(G) \otimes \mathbb{K} \rightarrow C^*(G') \otimes \mathbb{K}$ such that $\varphi \circ (\gamma^G \otimes \text{id})_z = (\gamma^{G'} \otimes \text{id})_z \circ \varphi$ for all $z \in \mathbb{T}$. Then there exists a $*$ -isomorphism $\varphi \rtimes \text{id} : [C^*(G) \rtimes_{\gamma, G} \mathbb{T}] \otimes \mathbb{K} \rightarrow [C^*(G') \rtimes_{\gamma, G'} \mathbb{T}] \otimes \mathbb{K}$ such that $[\varphi \rtimes \text{id}] \circ [\hat{\gamma}^G \otimes \text{id}] = [\hat{\gamma}^{G'} \otimes \text{id}] \circ [\varphi \rtimes \text{id}]$.*

Let A be a finite essential matrix with entries in \mathbb{N} . We denote by (D_A, D_A^+) the inductive limit of the inductive system of ordered abelian groups acting on columns

$$(\mathbb{Z}^V, \mathbb{Z}_+^V) \xrightarrow{A^T} (\mathbb{Z}^V, \mathbb{Z}_+^V) \xrightarrow{A^T} (\mathbb{Z}^V, \mathbb{Z}_+^V) \xrightarrow{A^T} \dots$$

and let $d_A : D_A \rightarrow D_A$ be the homomorphism induced by the following diagram:

$$\begin{array}{ccccccc} \mathbb{Z}^V & \xrightarrow{A^T} & \mathbb{Z}^V & \xrightarrow{A^T} & \mathbb{Z}^V & \xrightarrow{A^T} & \dots D_A \\ \downarrow A^T & & \downarrow A^T & & \downarrow A^T & & \downarrow d_A \\ \mathbb{Z}^V & \xrightarrow{A^T} & \mathbb{Z}^V & \xrightarrow{A^T} & \mathbb{Z}^V & \xrightarrow{A^T} & \dots D_A \end{array}$$

The triple (D_A, D_A^+, d_A) is called the *dimension group triple* of A .

Let H_i be an abelian group, H_i^+ a submonoid of H_i , and α_i an automorphism of H_i for $i = 1, 2$. We say two triples (H_i, H_i^+, α_i) are isomorphic if there is a group isomorphism $\phi : H_1 \rightarrow H_2$ such that $\phi(H_1^+) = H_2^+$ and $\phi \circ \alpha_1 = \alpha_2 \circ \phi$.

From work of Wagoner [46, Corollary 2.9] we have that (D_A, D_A^+, d_A) is isomorphic to a dimension group triple constructed from the edge shift (X_{E_A}, σ_{E_A}) associated to A . On the other hand, the dimension group triple (D_A, D_A^+, d_A) also coincides with a K-theory triple arising from the crossed product C*-algebra $C^*(G_A) \rtimes_{\gamma^A} \mathbb{T}$, where γ^A is the gauge unit circle action. More precisely, noting that arrow directions in [41] are reversed to ours, it follows from [41, Corollary 7.14] together with the discussion preceding [41, Lemma 7.15] that the triple (D_A, D_A^+, d_A) is isomorphic to the triple

$$(K_0(C^*(G_A) \rtimes_{\gamma^A} \mathbb{T}), K_0(C^*(G_A) \rtimes_{\gamma^A} \mathbb{T})^+, K_0(\hat{\gamma}_1^A)^{-1}),$$

where $\hat{\gamma}^A$ is the dual action of \mathbb{Z} on $C^*(G_A) \rtimes_{\gamma^A} \mathbb{T}$. The implication (3) \Rightarrow (4) below is what we referred to as *Krieger’s corollary* in the Introduction.

Corollary 7.2. *Suppose now that A and B are two finite essential matrices with entries in \mathbb{N} . The former conditions imply the latter:*

- (1) A and B are strong shift equivalent.
- (2) $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic in a way that respects the diagonals.
- (3) $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic.
- (4) A and B are shift equivalent.

Proof. By Williams’ theorem [47, Theorem 7.5.8] (see also [31]), strong shift equivalence of A and B coincides with conjugacy of two-sided edge shifts (X_{E_A}, σ_{E_A}) and (X_{E_B}, σ_{E_B}) (see [31, Theorem 7.2.7]), so we get that (1) implies (2) by the implication (III) \Rightarrow (II) of [5, Theorem 5.1] (see also [8, Proposition 2.17]).

Clearly (2) \Rightarrow (3), so we are left with showing (3) \Rightarrow (4). To do this, we use Proposition 7.1 to get that $[C^*(G) \rtimes_{\gamma^A} \mathbb{T}] \otimes \mathbb{K}$ and $[C^*(G') \rtimes_{\gamma^B} \mathbb{T}] \otimes \mathbb{K}$ are equivariantly isomorphic with actions $\hat{\gamma}^A \otimes \text{id}$ and

$\hat{\gamma}^B \otimes \text{id}$ respectively. Hence, by applying K-theory we get an isomorphism of the triples

$$\begin{aligned} & (K_0(C^*(G) \rtimes_{\gamma^A} \mathbb{T}), K_0(C^*(G) \rtimes_{\gamma^A} \mathbb{T})^+, K_0(\hat{\gamma}_1^A)), \\ & (K_0(C^*(G') \rtimes_{\gamma^B} \mathbb{T}), K_0(C^*(G') \rtimes_{\gamma^B} \mathbb{T})^+, K_0(\hat{\gamma}_1^B)). \end{aligned}$$

By Krieger's theorem [11, Theorem 6.4] (see also [29]), and up to the identification with dimension triples, we get that the triples above are isomorphic if and only if A and B are shift equivalent. Hence, (3) \Rightarrow (4). \square

From the perspective of C^* -algebras, we get that compatible, representable, and strong Morita shift equivalences are between strong shift equivalence and shift equivalence. The work done in the previous sections allows us to orient them, and show that each one provides a new way of obstructing SSE when merely assuming SE. The equivalence between (1) and (4) in the theorem below realizes a goal sought after by Muhly, Pask and Tomforde in [37, Remark 5.5], and provides a characterization of strong shift equivalence of matrices in terms of strong Morita shift equivalence.

Theorem 7.3. *Suppose A and B are two finite essential matrices with entries in \mathbb{N} . Then the following are equivalent:*

- (1) A and B are strong shift equivalent.
- (2) A and B are compatibly shift equivalent.
- (3) A and B are representable shift equivalent.
- (4) $X(A)_\infty$ and $X(B)_\infty$ are strong Morita shift equivalent.
- (5) $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic in a way that respects the diagonals.

Proof. It follows from Corollary 4.5 that (1) \Rightarrow (2), from Corollary 5.5 that (2) \Rightarrow (3), and from Proposition 6.2 that (3) \Rightarrow (4). The implication (4) \Rightarrow (5) is provided by Proposition 6.3. Finally, the implication (5) \Rightarrow (1) is granted to us by combining the implication (II) \Rightarrow (III) of [5, Theorem 5.1] and the fact that conjugacy of two-sided subshifts coincides with strong shift equivalence (Williams' theorem [47]). \square

We note here that if the construction in [22, Section 5] is applied to $E := X(A)$, $F := X(B)$ and $X := X(D)$ (as in the notation of [22]), we obtain again the correspondences E_∞ , F_∞ and X_∞ as in [22, Section 5]. This follows from uniqueness of Pimsner dilations [22, Theorem 3.9]. In particular, the correspondence $X(R)_\infty$ coincides with " R_∞ " in [22, Section 5] as the 1-2 corner of $X_\infty = X(D)_\infty$.

Hence, when we specify to $E := X(A)$ and $F := X(B)$, in the proof of [22, Theorem 5.8] it is erroneously claimed that $X(R)_\infty$ is the imprimitivity bimodule that implements a strong Morita equivalence between $X(A)_\infty$ and $X(B)_\infty$, through the identifications coming from ψ_A and ψ_B . However, by our result this would show that shift equivalence implies strong shift equivalence. Hence, the strategy of proof in [22, Theorem 5.8] cannot be made to work, as the following cutoff result demonstrates.

Theorem 7.4. *There exist finite aperiodic irreducible matrices A and B with entries in \mathbb{N} that such that $X(A)_\infty$ and $X(B)_\infty$ are strong Morita equivalent, but not strong Morita shift equivalent.*

Proof. Let A and B be the aperiodic and irreducible counterexamples of Kim and Roush from [28], so that they are shift equivalent but not strong shift equivalent. By the result of Bratteli and Kishimoto [4] we know that $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic, so that by Theorem 3.17 we get that $X(A)_\infty$ and $X(B)_\infty$ are strong Morita equivalent.

On the other hand, since A and B are not strong shift equivalent, by Theorem 7.3 we get that $X(A)_\infty$ and $X(B)_\infty$ cannot be strong Morita shift equivalent. \square

Remark 7.5. Suppose A and B are aperiodic and irreducible matrices over \mathbb{N} , and not 1×1 . Then the following conditions are equivalent:

- (1) A and B are shift equivalent.
- (2) $X(A)$ and $X(B)$ are shift equivalent.
- (3) $X(A)_\infty$ and $X(B)_\infty$ are strong Morita equivalent.
- (4) $C^*(G_A)$ and $C^*(G_B)$ are equivariantly stably isomorphic.
- (5) The triples (D_A, D_A^+, d_A) and (D_B, D_B^+, d_B) are isomorphic.

Indeed, Krieger’s theorem shows that (1) \Leftrightarrow (5), Proposition 3.5 shows that (1) \Leftrightarrow (2), Theorem 3.17 shows that (3) \Leftrightarrow (4), and Corollary 7.2 shows that (1) \Rightarrow (4). Finally, from Bratteli and Kishimoto [4, Corollary 4.3] we get (5) \Rightarrow (4), which finishes the proof.

Note that the above proof avoids the implication (2) \Rightarrow (3). Since now the validity of [22, Theorem 5.8] is in question, it is unknown whether one can prove (2) \Rightarrow (3) directly, without classification techniques as in [4].

Acknowledgments

The authors are grateful to Mike Boyle for bringing their attention to the work of Parry [39], as well as to Kevin Brix and Efren Ruiz for some remarks on earlier versions of this paper. The authors are also thankful for suggestions and remarks made by anonymous referees, ultimately leading to a more streamlined and readable version of the paper.

References

- [1] B. Abadie, S. Eilers, and R. Exel, “Morita equivalence for crossed products by Hilbert C^* -bimodules”, *Trans. Amer. Math. Soc.* **350**:8 (1998), 3043–3054. MR Zbl
- [2] P. Ara and E. Pardo, “Towards a K -theoretic characterization of graded isomorphisms between Leavitt path algebras”, *J. K-Theory* **14**:2 (2014), 203–245. MR Zbl
- [3] P. Ara, R. Hazrat, and H. Li, “Graded K -theory, filtered K -theory and the classification of graph algebras”, *Ann. K-Theory* **7**:4 (2022), 731–795. MR Zbl
- [4] O. Bratteli and A. Kishimoto, “Trace scaling automorphisms of certain stable AF algebras, II”, *Q. J. Math.* **51**:2 (2000), 131–154. MR Zbl
- [5] T. M. Carlsen and J. Rout, “Diagonal-preserving gauge-invariant isomorphisms of graph C^* -algebras”, *J. Funct. Anal.* **273**:9 (2017), 2981–2993. MR Zbl
- [6] T. M. Carlsen, S. Eilers, E. Ortega, and G. Restorff, “Flow equivalence and orbit equivalence for shifts of finite type and isomorphism of their groupoids”, *J. Math. Anal. Appl.* **469**:2 (2019), 1088–1110. MR Zbl

- [7] T. M. Carlsen, E. Ruiz, A. Sims, and M. Tomforde, “Reconstruction of groupoids and C^* -rigidity of dynamical systems”, *Adv. Math.* **390** (2021), art. id. 107923. MR Zbl
- [8] J. Cuntz and W. Krieger, “A class of C^* -algebras and topological Markov chains”, *Invent. Math.* **56**:3 (1980), 251–268. MR Zbl
- [9] A. Dor-On, “Isomorphisms of tensor algebras arising from weighted partial systems”, *Trans. Amer. Math. Soc.* **370**:5 (2018), 3507–3549. MR Zbl
- [10] A. Dor-On, S. Eilers, and S. Geffen, “Classification of irreversible and reversible Pimsner operator algebras”, *Compos. Math.* **156**:12 (2020), 2510–2535. MR Zbl
- [11] E. G. Effros, *Dimensions and C^* -algebras*, CBMS Reg. Conf. Ser. Math. **46**, Conf. Board Math. Sci., Washington, DC, 1981. MR Zbl
- [12] S. Eilers, G. Restorff, E. Ruiz, and A. P. W. Sørensen, “The complete classification of unital graph C^* -algebras: geometric and strong”, *Duke Math. J.* **170**:11 (2021), 2421–2517. MR Zbl
- [13] G. K. Eleftherakis, E. T. A. Kakariadis, and E. G. Katsoulis, “Morita equivalence of C^* -correspondences passes to the related operator algebras”, *Israel J. Math.* **222**:2 (2017), 949–972. MR Zbl
- [14] G. A. Elliott, “On the classification of C^* -algebras of real rank zero”, *J. Reine Angew. Math.* **443** (1993), 179–219. MR Zbl
- [15] M. Eryüzü, “Passing C^* -correspondence relations to the Cuntz–Pimsner algebras”, *Münster J. Math.* **15**:2 (2022), 441–471. MR Zbl
- [16] N. J. Fowler, M. Laca, and I. Raeburn, “The C^* -algebras of infinite graphs”, *Proc. Amer. Math. Soc.* **128**:8 (2000), 2319–2327. MR Zbl
- [17] T. Giordano, I. F. Putnam, and C. F. Skau, “Topological orbit equivalence and C^* -crossed products”, *J. Reine Angew. Math.* **469** (1995), 51–111. MR Zbl
- [18] T. Giordano, H. Matui, I. F. Putnam, and C. F. Skau, “Orbit equivalence for Cantor minimal \mathbb{Z}^d -systems”, *Invent. Math.* **179**:1 (2010), 119–158. MR Zbl
- [19] R. Hazrat, “The graded Grothendieck group and the classification of Leavitt path algebras”, *Math. Ann.* **355**:1 (2013), 273–325. MR Zbl
- [20] A. an Huef and I. Raeburn, “The ideal structure of Cuntz–Krieger algebras”, *Ergodic Theory Dynam. Systems* **17**:3 (1997), 611–624. MR Zbl
- [21] T. Kajiwara, C. Pinzari, and Y. Watatani, “Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules”, *J. Funct. Anal.* **159**:2 (1998), 295–322. MR Zbl
- [22] E. T. A. Kakariadis and E. G. Katsoulis, “ C^* -algebras and equivalences for C^* -correspondences”, *J. Funct. Anal.* **266**:2 (2014), 956–988. Correction in **283**:6 (2022), art. id. 109564. MR Zbl
- [23] S. Kaliszewski, N. Patani, and J. Quigg, “Characterizing graph C^* -correspondences”, *Houston J. Math.* **38**:3 (2012), 751–759. MR Zbl
- [24] T. Katsura, “On C^* -algebras associated with C^* -correspondences”, *J. Funct. Anal.* **217**:2 (2004), 366–401. MR Zbl
- [25] K. H. Kim and F. W. Roush, “Some results on decidability of shift equivalence”, *J. Combin. Inform. System Sci.* **4**:2 (1979), 123–146. MR Zbl
- [26] K. H. Kim and F. W. Roush, “Decidability of shift equivalence”, pp. 374–424 in *Dynamical systems* (College Park, MD, 1986–1987), edited by J. C. Alexander, Lecture Notes in Math. **1342**, Springer, 1988. MR Zbl
- [27] K. H. Kim and F. W. Roush, “Williams’s conjecture is false for reducible subshifts”, *J. Amer. Math. Soc.* **5**:1 (1992), 213–215. MR Zbl
- [28] K. H. Kim and F. W. Roush, “The Williams conjecture is false for irreducible subshifts”, *Ann. of Math. (2)* **149**:2 (1999), 545–558. MR Zbl
- [29] W. Krieger, “On dimension functions and topological Markov chains”, *Invent. Math.* **56**:3 (1980), 239–250. MR Zbl
- [30] E. C. Lance, *Hilbert C^* -modules: a toolkit for operator algebraists*, Lond. Math. Soc. Lect. Note Ser. **210**, Cambridge Univ. Press, 1995. MR Zbl

- [31] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge Univ. Press, 1995. MR Zbl
- [32] V. M. Manuilov and E. V. Troitsky, *Hilbert C^* -modules*, Transl. Math. Monogr. **226**, Amer. Math. Soc., Providence, RI, 2005. MR Zbl
- [33] K. Matsumoto, “Orbit equivalence of topological Markov shifts and Cuntz–Krieger algebras”, *Pacific J. Math.* **246**:1 (2010), 199–225. MR Zbl
- [34] K. Matsumoto, “Uniformly continuous orbit equivalence of Markov shifts and gauge actions on Cuntz–Krieger algebras”, *Proc. Amer. Math. Soc.* **145**:3 (2017), 1131–1140. MR Zbl
- [35] K. Matsumoto and H. Matui, “Continuous orbit equivalence of topological Markov shifts and Cuntz–Krieger algebras”, *Kyoto J. Math.* **54**:4 (2014), 863–877. MR Zbl
- [36] P. S. Muhly and B. Solel, “On the Morita equivalence of tensor algebras”, *Proc. Lond. Math. Soc.* (3) **81**:1 (2000), 113–168. MR Zbl
- [37] P. S. Muhly, D. Pask, and M. Tomforde, “Strong shift equivalence of C^* -correspondences”, *Israel J. Math.* **167** (2008), 315–346. MR Zbl
- [38] M. Nasu, *Textile systems for endomorphisms and automorphisms of the shift*, Mem. Amer. Math. Soc. **546**, Amer. Math. Soc., Providence, RI, 1995. MR Zbl
- [39] W. Parry, “The classification of topological Markov chains: adapted shift equivalence”, *Israel J. Math.* **38**:4 (1981), 335–344. MR Zbl
- [40] M. V. Pimsner, “A class of C^* -algebras generalizing both Cuntz–Krieger algebras and crossed products by \mathbb{Z} ”, pp. 189–212 in *Free probability theory* (Waterloo, ON, 1995), edited by D.-V. Voiculescu, Fields Inst. Commun. **12**, Amer. Math. Soc., Providence, RI, 1997. MR Zbl
- [41] I. Raeburn, *Graph algebras*, CBMS Reg. Conf. Ser. Math. **103**, Amer. Math. Soc., Providence, RI, 2005. MR Zbl
- [42] I. Raeburn, “On graded C^* -algebras”, *Bull. Aust. Math. Soc.* **97**:1 (2018), 127–132. MR Zbl
- [43] M. Rørdam, F. Larsen, and N. Laustsen, *An introduction to K -theory for C^* -algebras*, Lond. Math. Soc. Stud. Texts **49**, Cambridge Univ. Press, 2000. MR Zbl
- [44] W. Szymański, “General Cuntz–Krieger uniqueness theorem”, *Int. J. Math.* **13**:5 (2002), 549–555. MR Zbl
- [45] A. Tikuisis, S. White, and W. Winter, “Quasidiagonality of nuclear C^* -algebras”, *Ann. of Math.* (2) **185**:1 (2017), 229–284. MR Zbl
- [46] J. B. Wagoner, “Topological Markov chains, C^* -algebras, and K_2 ”, *Adv. Math.* **71**:2 (1988), 133–185. MR Zbl
- [47] R. F. Williams, “Classification of subshifts of finite type”, *Ann. of Math.* (2) **98**:1 (1973), 120–153. Correction in **99**:2 (1974), 380–381. MR Zbl
- [48] D. P. Williams, *Crossed products of C^* -algebras*, Math. Surv. Monogr. **134**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [49] W. Winter, “Localizing the Elliott conjecture at strongly self-absorbing C^* -algebras”, *J. Reine Angew. Math.* **692** (2014), 193–231. MR Zbl
- [50] W. Winter and J. Zacharias, “The nuclear dimension of C^* -algebras”, *Adv. Math.* **224**:2 (2010), 461–498. MR Zbl

Received 10 Mar 2022. Revised 8 Aug 2022. Accepted 15 Sep 2022.

TOKE MEIER CARLSEN: toke.carlsen@gmail.com

Department of Mathematics, University of the Faroe Islands, Tórshavn, Faroe Islands

ADAM DOR-ON: adoron.math@gmail.com

Mathematisches Institut, WWU Münster, Münster, Germany

SØREN EILERS: eilers@math.ku.dk

Department of Mathematical Sciences, University of Copenhagen, Copenhagen, Denmark

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 1 2024

The prescribed curvature problem for entire hypersurfaces in Minkowski space CHANGYU REN, ZHIZHANG WANG and LING XIAO	1
Anisotropic micropolar fluids subject to a uniform microtorque: the stable case ANTOINE REMOND-TIEDREZ and IAN TICE	41
Strong ill-posedness for SQG in critical Sobolev spaces IN-JEE JEONG and JUNHA KIM	133
Large-scale regularity for the stationary Navier–Stokes equations over non-Lipschitz boundaries MITSUO HIGAKI, CHRISTOPHE PRANGE and JINPING ZHUGE	171
On a family of fully nonlinear integrodifferential operators: from fractional Laplacian to non-local Monge–Ampère LUIS A. CAFFARELLI and MARÍA SORIA-CARRO	243
Propagation of singularities for gravity-capillary water waves HUI ZHU	281
Shift equivalences through the lens of Cuntz–Krieger algebras TOKE MEIER CARLSEN, ADAM DOR-ON and SØREN EILERS	345