

ANALYSIS & PDE

Volume 17

No. 2

2024

KARLHEINZ GRÖCHENIG AND ANDREAS KLOTZ

**NECESSARY DENSITY CONDITIONS FOR SAMPLING AND
INTERPOLATION IN SPECTRAL SUBSPACES OF ELLIPTIC
DIFFERENTIAL OPERATORS**



NECESSARY DENSITY CONDITIONS FOR SAMPLING AND INTERPOLATION IN SPECTRAL SUBSPACES OF ELLIPTIC DIFFERENTIAL OPERATORS

KARLHEINZ GRÖCHENIG AND ANDREAS KLOTZ

We prove necessary density conditions for sampling in spectral subspaces of a second-order uniformly elliptic differential operator on \mathbb{R}^d with slowly oscillating symbol. For constant-coefficient operators, these are precisely Landau’s necessary density conditions for bandlimited functions, but for more general elliptic differential operators it has been unknown whether such a critical density even exists. Our results prove the existence of a suitable critical sampling density and compute it in terms of the geometry defined by the elliptic operator. In dimension $d = 1$, functions in a spectral subspace can be interpreted as functions with variable bandwidth, and we obtain a new critical density for variable bandwidth. The methods are a combination of the spectral theory and the regularity theory of elliptic partial differential operators, some elements of limit operators, certain compactifications of \mathbb{R}^d , and the theory of reproducing kernel Hilbert spaces.

1. Introduction

The classical Paley–Wiener space is the subspace $\text{PW}_\Omega = \{f \in L^2(\mathbb{R}) : \text{supp } \hat{f} \subseteq [-\Omega, \Omega]\}$ of $L^2(\mathbb{R})$. Using Fourier inversion, one sees that the point evaluation $f \mapsto f(x)$ is bounded on PW_Ω . The fundamental questions about PW_Ω are originally motivated by problems in signal processing and information theory: when is $f \in \text{PW}_\Omega$ completely and stably determined by its samples $\{f(s) : s \in S\}$ on a set $S \subseteq \mathbb{R}$? On which sets $S \subseteq \mathbb{R}$ can every sequence $(a_s)_{s \in S} \in \ell^2(S)$ be interpolated by a function f in PW_Ω , so that $f(s) = a_s$ for all $s \in S$? These questions were answered by Beurling [1989] and Landau [1967].

Theorem A. (i) *Assume that S is uniformly separated and*

$$A \|f\|_2^2 \leq \sum_{s \in S} |f(s)|^2 \leq B \|f\|_2^2 \quad \text{for all } f \in \text{PW}_\Omega. \quad (1-1)$$

Then

$$D^-(S) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}} \frac{\#(S \cap [x - r, x + r])}{2r} \geq \frac{\Omega}{\pi}. \quad (1-2)$$

(ii) *If for all $a \in \ell^2(S)$ there exists $f \in \text{PW}_\Omega$ such that $f(s) = a_s$, $s \in S$, then*

$$D^+(S) = \limsup_{r \rightarrow \infty} \sup_{x \in \mathbb{R}} \frac{\#(S \cap [x - r, x + r])}{2r} \leq \frac{\Omega}{\pi}. \quad (1-3)$$

This research was funded in whole or in part by the Austrian Science Fund (FWF) 10.55776/P31887. For open access purposes, the authors have applied a CC BY public copyright license to any author-accepted manuscript version arising from this submission. MSC2020: primary 35J99, 46E22, 47B32, 54D35, 94A20; secondary 42C40.

Keywords: spectral subspace, Paley–Wiener space, bandwidth, Beurling density, sampling, interpolation, elliptic operator, regularity theory, slow oscillation, Higson compactification.

In the established terminology, a set that satisfies a sampling inequality of the form (1-1) is called a sampling set for the underlying space PW_Ω , or a set of stable sampling. A set on which arbitrary ℓ^2 -data can be interpolated is called a set of interpolation. The expressions $D^-(S)$ and $D^+(S)$ are called the lower and the upper Beurling density.

The number Ω/π in (1-2) and (1-3) is an important invariant of the space PW_Ω and has an interpretation in information theory. Since, roughly speaking, the densities $D^\pm(S)$ measure the average number of samples in S per unit length, the *necessary density conditions* of Theorem A say that at least Ω/π samples per unit length are required to recover a function in PW_Ω from $f|_S$, whereas at most Ω/π values per unit length are permitted to solve the interpolation problem in PW_Ω . Thus the density Ω/π represents a critical value below which (stable) sampling is impossible, and above which interpolation is impossible. Indeed, these questions about sampling and interpolation were at the origin of Shannon’s information theory [1948], and the uniform sampling theorem with $S = \alpha\mathbb{Z}$ is still considered the basis of analog-digital conversion in modern signal processing. The ratio $D^\pm(S)/\Omega$ is a measure for the redundancy, thus for the performance quality, of the sampling set S . The theory of Beurling, Kahane, and Landau provides a rigorous mathematical formulation for the existence of a critical density for arbitrary sets S (in place of $\alpha\mathbb{Z}$). Although we will not touch this question here, we mention that the conditions of Theorem A yield almost a characterization of sets of sampling and of interpolation: *in dimension $d = 1$, if S is uniformly separated and $D^-(S) > 1$, then S is a sampling set for PW_Ω , and if $D^+(S) < 1$, then S is a set of interpolation for PW_Ω .* See [Kahane 1962; Beurling 1989; Seip 2004] for an exposition of the sampling theory in the classical Paley–Wiener space.

The connection with partial differential operators comes from the observation that PW_Ω is a spectral subspace of the differential operator $H = -d^2/dx^2$. Using the Fourier transform \mathcal{F} , this differential operator is unitarily equivalent to the multiplication operator $\mathcal{F}(-d^2f/dx^2)(\xi) = \xi^2 \hat{f}(\xi)$. In this representation of $-d^2/dx^2$ the spectral projection on the interval $[0, \Omega]$ is given by $\chi_{[0,\Omega]}(H)f = \mathcal{F}^{-1}(\chi_{[0,\Omega]}(\xi^2)\hat{f})$. This implies

$$PW_\Omega = \chi_{[0,\Omega^2]}(H)L^2(\mathbb{R}).$$

This observation is the starting point for many generalizations of Paley–Wiener spaces and sampling theorems. In this work we study the question of necessary density conditions for sampling and interpolation in the spectral subspaces of a self-adjoint uniformly elliptic differential operator

$$H_a = - \sum_{j,k=1}^d \partial_j a_{jk}(x) \partial_k$$

acting on $L^2(\mathbb{R}^d)$ with a smooth positive definite (matrix) symbol $a = (a_{jk}(x))_{j,k=1,\dots,d}$. The Paley–Wiener space associated to H_a is the spectral subspace

$$PW_\Omega(H_a) = \chi_{[0,\Omega]}(H_a)L^2(\mathbb{R}^d),$$

where, as usual, $\chi_{[0,\Omega]}(H_a)$ is the orthogonal projection corresponding to the spectrum $[0, \Omega]$.

If the symbol $a(x) = a$ is constant, then H_a is similar to the Laplace operator, and the corresponding spectral subspace can be described with Fourier techniques. For this case necessary density conditions for sampling and interpolation are already contained in [Landau 1967]. Optimal sufficient conditions

for sampling in \mathbb{R}^d in terms of a covering density were obtained in [Beurling 1966]. However, if H_a is a uniformly elliptic differential operator with variable coefficients, then the standard techniques break down, and it was an open question whether a critical density exists for sampling and interpolation in the spectral subspaces of H_a , and how to compute this critical density.

We will answer this question for a class uniformly elliptic operators. We say a smooth symbol with all derivatives bounded, $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$, is slowly oscillating if $\lim_{|x| \rightarrow \infty} |\partial_k a(x)| = 0$ for $k = 1, \dots, d$.

Theorem B. *If a is slowly oscillating, then there exists a critical density for sampling and interpolation for $\text{PW}_\Omega(H_a)$.*

Adapting the measure to the geometry associated to the differential operator H_a , the critical density can be determined explicitly. This is our main result.

Theorem C. *Assume $H_a = -\sum_{j,k=1}^d \partial_j a_{jk} \partial_k$ is a self-adjoint uniformly elliptic operator with slowly oscillating symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. Let $d\nu(x) = (\det a(x))^{-1/2} dx$ be the associated measure.*

(i) *If $S \subseteq \mathbb{R}^d$ is a set of stable sampling for $\text{PW}_\Omega(H_a)$ then*

$$D_v^-(S) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}^d} \frac{\#(S \cap B_r(x))}{\nu(B_r(x))} \geq \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}.$$

(ii) *If $S \subseteq \mathbb{R}^d$ is a set of interpolation for $\text{PW}_\Omega(H_a)$, then*

$$D_v^+(S) = \limsup_{r \rightarrow \infty} \sup_{x \in \mathbb{R}^d} \frac{\#(S \cap B_r(x))}{\nu(B_r(x))} \leq \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}.$$

Except for the modified definition of the density, the formulation of the theorem is identical to Landau’s theorem [1967]. By contrast, the method of proof is vastly different as Fourier methods are not available for the proof of Theorem C. In addition we draw the new insight that the appropriate notion of density must be linked to the geometry defined by a . For compact manifolds the link between density and geometry was already observed in [Ortega-Cerdà and Pridhni 2012].

For the special case of a symbol that is asymptotically constant at infinity we can use the standard Beurling densities in \mathbb{R}^d from (1-2) and (1-3) (with intervals replaced by Euclidean balls $B_r(x)$) and obtain the following consequence.

Corollary D. *Assume that $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ is asymptotically constant, i.e., $\lim_{x \rightarrow \infty} a(x) = b$. Let $\Sigma_\Omega^b = \{\xi \in \mathbb{R}^d : b\xi \cdot \xi \leq \Omega\}$.*

(i) *If $S \subseteq \mathbb{R}^d$ is a set of sampling for the Paley–Wiener space $\text{PW}_\Omega(H_a)$, then*

$$D^-(S) \geq \frac{|\Sigma_\Omega^b|}{(2\pi)^d} = (\det b)^{-1/2} \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}.$$

(ii) *If $S \subseteq \mathbb{R}^d$ is a set of interpolation for the Paley–Wiener space $\text{PW}_\Omega(H_a)$, then*

$$D^+(S) \leq \frac{|\Sigma_\Omega^b|}{(2\pi)^d}.$$

We note that the same critical density holds for the Paley–Wiener space of the constant-coefficient differential operator H_b . Since H_a may be considered a perturbation of H_b and since the Beurling density $D^\pm(S)$ is an asymptotic quantity, it is to be expected that the necessary density for $\text{PW}_\Omega(H_a)$ coincides with the necessary density for $\text{PW}_\Omega(H_b)$.

Let us put these statements into context.

Sampling in spectral subspaces. Several researchers have created an extensive qualitative theory of sampling in spectral subspaces of a general unbounded, positive, self-adjoint operator H on a Hilbert space \mathcal{H} . In this case the abstract Paley–Wiener space is defined as $\text{PW}_{[0,\Omega]}(H) = \chi_{[0,\Omega]}(H)\mathcal{H}$. Usually $\mathcal{H} = L^2(X, \mu)$ and $\text{PW}_{[0,\Omega]}(H)$ is a reproducing kernel Hilbert space. In this situation many authors have proved the existence of sampling sets [Coulhon et al. 2012; Feichtinger et al. 2016; Filbir and Mhaskar 2011; Feichtinger and Pesenson 2004; Pesenson 2000; 2001; Pesenson and Zayed 2009]. In particular the set-up of [Coulhon et al. 2012; Pesenson 1999; Pesenson and Zayed 2009] covers the case of H being a self-adjoint uniformly elliptic differential operator on $L^2(\mathbb{R}^d)$. The construction of sampling sets in these abstract Paley–Wiener spaces requires some smoothness properties of functions in $\text{PW}_\Omega(H)$ and a Bernstein-type inequality (see (2-1) below). The result then is that a “sufficiently dense” subset in X is a sampling set and a “sufficiently sparse” subset of X is a set of interpolation. What remained unknown is the existence of a critical density against which one could compare the quality of the construction. Theorems B, C, and Corollary D address this gap for uniformly elliptic differential operators. Once a critical sampling density is established, one may aim for sampling sets near the critical density. The question of optimal sampling sets in spectral subspaces is wide open; in fact, it has become meaningful only after the critical density is known explicitly. This problem is already difficult for multivariate bandlimited functions $\text{PW}_K = \{f \in L^2(\mathbb{R}^d) : \text{supp } \hat{f} \subseteq K\}$ for compact spectrum $K \subseteq \mathbb{R}^d$ and was solved only recently in [Matei and Meyer 2010; Olevskii and Ulanovskii 2008]. A possible general approach is via the construction of Fekete sets and weak limits, as was carried out in [Gröchenig et al. 2019] for Fock spaces with a general weight.

Insight for partial differential operators. Although the spectral subspaces of a partial differential operator are natural objects, they seem to have received little attention. To the best of our knowledge, nothing is known about the nature of the corresponding reproducing kernel and the behavior of functions in the spectral subspaces PW_Ω . Our investigation reveals several properties of the reproducing kernel, such as the behavior of its diagonal and some form of off-diagonal decay. These are key properties for the proofs of Theorems B and C, and we hope that these also hold some interest for partial differential operators.

Variable bandwidth. Our original motivation comes from a new concept of variable bandwidth. In [Gröchenig and Klotz 2017] we argued that the spectral subspaces of the Sturm–Liouville operator $-(d/dx)a(d/dx)$ on $L^2(\mathbb{R})$ for some function $a > 0$ can be taken as spaces with variable bandwidth. We proved that $a(x)^{-1/2}$ is a measure for the bandwidth near x (the largest active frequency at position x). The function a thus parametrizes the local bandwidth. For $a = \text{const.}$, the spectral subspace is just the classical Paley–Wiener space PW_Ω . For the special case of an eventually constant parametrizing function a , i.e., a is constant outside an interval $[-R, R]$, we computed the critical density for sampling in

$\text{PW}_\Omega(-(d/dx)a(d/dx))$. The proof required intricate details of the scattering theory of one-dimensional Schrödinger operators. [Theorem C](#), formulated for dimension $d = 1$, yields a significant extension of the density theorem for the sampling of functions of variable bandwidth.

Corollary E. *Assume that $a \in C_b^\infty(\mathbb{R})$ is bounded, $a > 0$, and $\lim_{x \rightarrow \pm\infty} a'(x) = 0$. Let $\text{PW}_\Omega(H_a)$ be the Paley–Wiener space associated to H_a .*

If S is a sampling set for $\text{PW}_\Omega(H_a)$, then

$$D_v^-(S) \geq \frac{\Omega^{1/2}}{\pi}.$$

Similarly, if S is a set of interpolation for $\text{PW}_\Omega(H_a)$, then

$$D_v^+(S) \leq \frac{\Omega^{1/2}}{\pi}.$$

Methods. The proofs of [Theorems B and C](#) combine ideas and techniques from several areas of analysis.

Critical density in reproducing kernel Hilbert spaces. Originally, density theorems in the style of Landau — and there are dozens in analysis — were proved from scratch. In our approach we apply the results on sampling and interpolation in general reproducing kernel Hilbert spaces from [\[Führ et al. 2017\]](#). The main insight was that it suffices to verify some geometric conditions on the measure space, such as a doubling condition of the underlying measure, and of the reproducing kernel, such as some form of off-diagonal decay. Once these conditions are satisfied, one obtains the existence of a critical density and can calculate it in terms of the averaged trace of the reproducing kernel. Since the geometric conditions are trivially satisfied for \mathbb{R}^d , our main technical difficulty is to understand the reproducing kernel of the spectral subspaces of a self-adjoint uniformly elliptic differential operator.

Regularity theory and heat kernel estimates. To study this reproducing kernel, we use the fundamental results of the regularity theory of elliptic differential operators. With these tools we investigate the smoothness of the reproducing kernel and compare various Sobolev norms on $\text{PW}_\Omega(H_a)$. See [Lemma 2.1](#) and [Proposition 2.2](#). For an important technical detail ([Proposition 2.2](#)) we will need Gaussian estimates for the heat kernel, which we expect to play a key role in extensions of our theory.

Limit operators and slowly varying symbols. To connect asymptotic properties of the symbol a of a partial differential operator H_a to the spectral theory of H_a , we use the notion of limit operators. Although we do not use any elaborate results from this theory (see [\[Georgescu 2011; Rabinovich et al. 2004a; Špakula and Willett 2017\]](#)), limit operators are central to our arguments.

Higson compactification of \mathbb{R}^d . An important structure underlying the proof of [Theorem C](#) is a compactification of \mathbb{R}^d , the so-called Higson compactification. This is the compactification arising as the maximal ideal space of the C^* -algebra of slowly oscillating functions on \mathbb{R}^d . By Gelfand theory every slowly oscillating function can be identified with a continuous function on the Higson compactification $h\mathbb{R}^d$; see, e.g., [\[Rabinovich et al. 2004a; Roe 2003; Shteinberg 2000\]](#). On a technical level we will show that for slowly oscillating symbols the mapping $x \rightarrow T_{-x}k_x$ of centered reproducing kernels can be extended continuously to the compactification $h\mathbb{R}^d$ ([Proposition 6.3](#)).

The underlying philosophy is summarized in the following diagram; we write $T_x f(z) = f(z - x)$ for the translation operator and k_x for the reproducing kernel of $\text{PW}_\Omega(H_a)$:

$$\{T_{-x}a : x \in \mathbb{R}^d\} \text{ compact} \implies \{T_{-x}H_aT_x : x \in \mathbb{R}^d\} \text{ compact} \implies \{T_{-x}k_x : x \in \mathbb{R}^d\} \text{ compact}.$$

Thus, if $T_{x_n}a \rightarrow b$ in a suitable topology, then $T_{-x_n}H_aT_{x_n} \rightarrow H_b$ and the sequence of centered reproducing kernels $T_{-x_n}k_{x_n}$ converges to the reproducing kernel of $\text{PW}_\Omega(H_b)$. In the considered examples the limit operator H_b is simpler than the original operator H_a , and this facilitates information about the reproducing kernel of $\text{PW}_\Omega(H_a)$.

The paper is organized as follows. [Section 2](#) prepares the background material on regularity theory, symbol classes for partial differential operators, and reproducing kernel Hilbert spaces. We prove the basic properties of the Paley–Wiener space $\text{PW}_\Omega(H_a)$. [Section 3](#) gives the precise formulation of the general density theorem for $\text{PW}_\Omega(H_a)$. Its proof is given in [Sections 4](#) and [5](#). In [Section 6](#) we calculate the critical density for sampling in $\text{PW}_\Omega(H_a)$ for the class of slowly varying symbols ([Theorem C](#) and [Corollary D](#)). We conclude with an outlook and collect additional material in [Appendices A](#) and [B](#).

2. Preliminaries

2A. Notation. For a function f on \mathbb{R}^d and $x, z \in \mathbb{R}^d$ we define the translation operator $T_x f(z) = f(z - x)$. The open Euclidean ball of radius r at x is $B_r(x)$, and $B_r = B_r(0)$.

We use standard multi-index notation; thus the differential operator D^α is $\partial^{|\alpha|}/(\partial x_1^{\alpha_1} \cdots \partial x_d^{\alpha_d})$ and the multivariate binomial symbol is $\binom{\alpha}{\gamma} = \prod_{j=1}^d \binom{\alpha_j}{\gamma_j}$ for multi-indices $\alpha, \gamma \in \mathbb{N}_0^d$.

We will denote the space of uniformly continuous and bounded functions on \mathbb{R}^d with values in a Banach space X by $C_b^u(\mathbb{R}^d, X)$. The indices c, ∞ , and 0 refer to the subspaces of compactly supported, smooth, and vanishing-at-infinity functions in $C(\mathbb{R}^d)$. Thus $C_b^\infty(\mathbb{R}^d, X)$ consists of all smooth X -valued functions with bounded derivatives of all orders. The space $C^\infty(\mathbb{R}^d, X)$ has the Fréchet space topology induced by the seminorms $|f|_{R,\alpha} = \sup_{x \in B_R(0)} \|D^\alpha f(x)\|_X$. If $X = \mathbb{C}$, we write $C_b^\infty(\mathbb{R}^d)$, etc.

The Fourier transform of $f \in L^1(\mathbb{R}^d)$ is

$$\mathcal{F} f(\omega) = \hat{f}(\omega) = (2\pi)^{-d/2} \int_{\mathbb{R}^d} f(x)e^{-ix \cdot \omega} dx,$$

and \mathcal{F} extends to a unitary operator on $L^2(\mathbb{R}^d)$ as usual. For every $s \geq 0$ the Sobolev space W_2^s is defined by

$$W_2^s = \left\{ f \in L^2(\mathbb{R}^d) : \|f\|_{W_2^s} = \left[(2\pi)^{-d/2} \int_{\mathbb{R}^d} |\hat{f}(\omega)|^2 (1 + |\omega|^2)^s d\omega \right]^{1/2} < \infty \right\}.$$

If $s \in \mathbb{N}$, then $\|f\|_{W_2^s} \asymp \sum_{|\alpha| \leq s} \|D^\alpha f\|_2$. By the Sobolev embedding theorem, $W_2^s \hookrightarrow C_0(\mathbb{R}^d)$ for $s > d/2$.

Recall that a reproducing kernel Hilbert space \mathcal{H} is a Hilbert space of functions defined on a set X such that $f(x) = \langle f, k_x \rangle_{\mathcal{H}}$ for all $f \in \mathcal{H}$ and $x \in X$. We write $k(x, y) = \overline{k_x(y)}$ for the reproducing kernel of \mathcal{H} . See, e.g., [\[Aronszajn 1950\]](#). In particular, the Sobolev space W_2^s is a reproducing kernel Hilbert space with reproducing kernel $T_x \kappa$, $x \in \mathbb{R}^d$, where $\hat{\kappa}(\omega) = \hat{\kappa}_s(\omega) = (1 + |\omega|^2)^{-s}$, by direct computation or by [\[Wendland 2005\]](#).

2B. The generalized Paley–Wiener space and its basic properties. Pesenson’s idea [1998; 2001] (see also [Pesenson and Zayed 2009]) was to define an abstract Paley–Wiener space as a spectral subspace associated to an arbitrary positive, self-adjoint operator $H \geq 0$ with domain $\mathcal{D}(H)$ on a Hilbert space \mathcal{H} and a spectral interval $[0, \Omega]$. Let $\chi_{[0, \Omega]}(H)$ be the spectral projection of H . Then the generalized Paley–Wiener space is defined as

$$\text{PW}_\Omega(H) = \chi_{[0, \Omega]}(H)\mathcal{H}.$$

Equivalently, for a positive, self-adjoint operator, one can define the Paley–Wiener space $\text{PW}_\Omega(H)$ by a Bernstein-type inequality: $f \in \text{PW}_\Omega(H)$, if and only if $f \in \mathcal{D}(H^k)$ for all $k \in \mathbb{N}$, and

$$\|H^k f\|_2 \leq \Omega^k \|f\|_2 \quad \text{for all } k \in \mathbb{N}. \tag{2-1}$$

This is an easy consequence of the spectral theorem; see [Gröchenig and Klotz 2010; Pesenson 2001; Pesenson and Zayed 2009].

If $H = -d^2/dx^2$ on $L^2(\mathbb{R})$, then

$$\text{PW}_\Omega(H) = \{f \in L^2(\mathbb{R}) : \text{supp } \hat{f} \subseteq [-\sqrt{\Omega}, \sqrt{\Omega}]\}$$

is precisely the classical Paley–Wiener space, or in engineering language the space of band-limited functions with bandwidth $2\sqrt{\Omega}$.

Convention. In this work we consider positive, formally self-adjoint differential expressions $H = H_a$ of the form

$$H_a f = - \sum_{j,k=1}^d \partial_j a_{jk} \partial_k f, \quad f \in W_2^2. \tag{2-2}$$

Here the *matrix symbol* $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ is positive definite; i.e., $a_{jk} = \bar{a}_{kj} \in C_b^\infty(\mathbb{R}^d)$ and there exists $\theta > 0$ such that $a(x)\xi \cdot \xi \geq \theta|\xi|^2$ for all $\xi, x \in \mathbb{R}^d$. Then H_a is a positive, uniformly elliptic self-adjoint operator on \mathbb{R}^d with domain $\mathcal{D}(H_a) = W_2^2$. In particular $C_c^\infty(\mathbb{R}^d)$ is a core for H_a ; i.e., H_a is the operator closure of $H_a|_{C_c^\infty(\mathbb{R}^d)}$. The regularity theory of elliptic differential operators asserts that for every $k \in \mathbb{N}_0$ there is a $c_k \in \mathbb{R}$ such that

$$H_a^k + c_k : W_2^{2k} \rightarrow L^2(\mathbb{R}^d)$$

is a Hilbert space isomorphism. See [Zimmer 1990, Theorem 6.3.12] or the standard references [Agmon 1965; Shubin 1992]. For further use we record the fact that a uniformly elliptic operator is one-to-one on its domain and thus

$$0 \text{ is not an eigenvalue of } H_a. \tag{2-3}$$

To see this, we use the ellipticity and $f \in W_2^2$. Then the identity

$$\langle H_a f, f \rangle = \int \sum_{j,k} a_{jk} \partial_k f(x) \overline{\partial_j f(x)} dx = 0$$

implies that $\partial_j f \equiv 0$; thus $f = 0$.

Remark. We regard the mapping $a \mapsto H_a$ as a mapping from functions to operators (a symbolic calculus) and refer to a as the (matrix) symbol of the operator. This terminology differs slightly from

the usage in PDE, where the (principal) symbol of the differential operator $\sum_{|\alpha| \leq m} a_\alpha D^\alpha$ is the function $p(x, \xi) = \sum_{|\alpha|=m} a_\alpha \xi^\alpha$ on \mathbb{R}^{2d} . For the second-order differential operator H_a in (2-2) the principal symbol is $p(x, \xi) = a(x)\xi \cdot \xi$. Since H_a is self-adjoint, the coefficients a_α are all real for $|\alpha| = 2$.

First we verify that $\text{PW}_\Omega(H_a)$ embeds in every Sobolev space.

Lemma 2.1. *The Paley–Wiener space $\text{PW}_\Omega(H_a)$ is continuously embedded in all Sobolev spaces W_2^s , $s \geq 0$, and in $C_0^\infty(\mathbb{R}^d)$. As a consequence, on $\text{PW}_\Omega(H_a)$, the L^2 -norm and the Sobolev norms are equivalent.*

Proof. Let $f \in \text{PW}_\Omega(H_a)$ and $k \in \mathbb{N}$. By elliptic regularity and Bernstein’s inequality (2-1), $\|f\|_{W_2^{2k}} \asymp \|(H^k + c_k)f\|_2 \leq (\Omega^k + |c_k|)\|f\|_2$. Consequently, $f \in \bigcap_{k \in \mathbb{N}} W_2^{2k} = \bigcap_{s \geq 0} W_2^s \subseteq C_0^\infty(\mathbb{R}^d)$ via the Sobolev embedding. □

Embeddings of Paley–Wiener spaces different from Lemma 2.1 can be found in [Feichtinger and Pesenson 2004].

Next we show that $\text{PW}_\Omega(H_a)$ is a reproducing kernel Hilbert space in $L^2(\mathbb{R}^d)$.

Proposition 2.2. *There exists a reproducing kernel $k_x \in \text{PW}_\Omega(H_a)$ such that $\chi_{[0, \Omega]}(H_a)f(x) = \langle f, k_x \rangle$ for all $f \in L^2(\mathbb{R}^d)$ and all $x \in \mathbb{R}^d$. In addition, there are positive constants c, C such that*

$$0 < c \leq \|k_x\|_2 \leq C \quad \text{for all } x \in \mathbb{R}^d. \tag{2-4}$$

Proof. Let $f \in \text{PW}_\Omega(H_a)$ and $s > d/2$. By Lemma 2.1, $f \in W_2^s$ and $\|f\|_2 \asymp \|f\|_{W_2^s}$. Since W_2^s is a reproducing kernel Hilbert space, we obtain

$$|f(x)| = |\langle f, T_x \kappa \rangle_{W_2^s}| \leq \|T_x \kappa\|_{W_2^s} \|f\|_{W_2^s} \leq C \|f\|_2.$$

Thus $\text{PW}_\Omega(H_a)$ is a reproducing kernel Hilbert space with kernel $k_x \in \text{PW}_\Omega(H_a)$.

For the lower bound in (2-4) we do not have a proof based exclusively on regularity theory. Instead we refer to [Coulhon et al. 2012, Lemma 3.19], where the lower bound for the reproducing kernel was derived by means of heat kernel estimates. As some details and notation differ, we reproduce the proof in Appendix B. □

Proposition 2.3. *The mapping $x \mapsto k_x$ is continuous from \mathbb{R}^d to W_2^s , $s \geq 0$.*

Proof. Since $k_x \in \text{PW}_\Omega(H_a)$ and $\|k_x\|_2$ is bounded by (2-4), Lemma 2.1 and the Sobolev embedding theorem imply that $C_1 = \sup_{x, y \in \mathbb{R}^d} |\nabla k_x(y)|$ is finite; therefore

$$\begin{aligned} \|k_x - k_y\|_{W_2^s}^2 &\leq C \|k_x - k_y\|_2^2 = C(k_x(x) - k_x(y) - k_y(x) + k_y(y)) \\ &\leq 2C \sup_{z, w} |\nabla k_z(w)| |x - y| \leq C' |x - y|. \end{aligned}$$

Consequently $x \rightarrow k_x$ is continuous. □

2C. Sampling and interpolation in $\text{PW}_\Omega(H_a)$ and the Beurling densities. Let μ be a Borel measure on \mathbb{R}^d that is equivalent to Lebesgue measure in the sense that $d\mu = h dx$ for a measurable function h with $0 < c \leq h(x) \leq C$ for all $x \in \mathbb{R}^d$.

The lower Beurling density of S with respect to μ is defined as

$$D_\mu^-(S) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}^d} \frac{\#(S \cap B_r(x))}{\mu(B_r(x))}, \tag{2-5}$$

and the upper Beurling density of S is

$$D_\mu^+(S) = \limsup_{r \rightarrow \infty} \sup_{x \in \mathbb{R}^d} \frac{\#(S \cap B_r(x))}{\mu(B_r(x))}.$$

If $d\mu = dx$ we omit the subscript and write $D^\pm(S)$.

For sampling in reproducing kernel Hilbert spaces the relevant measure is $d\mu(x) = k(x, x) dx$. We call the Beurling density with respect to this measure the dimension-free density and write $D_0^\pm(S)$ for $D_\mu^\pm(S)$.

We say that the reproducing kernel k of a reproducing kernel Hilbert space $\mathcal{H} \subseteq L^2(\mathbb{R}^d, dx)$ satisfies the *weak localization* property (WL) if for every $\varepsilon > 0$ there is a constant $r = r(\varepsilon)$ such that

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_r(x)} |k(x, y)|^2 dy < \varepsilon^2. \tag{WL}$$

The discrete analog of the weak localization is the so-called *homogeneous approximation property* (HAP) of the reproducing kernel: Assume that S is such that $\{k_s : s \in S\}$ is a *Bessel sequence* for \mathcal{H} ; i.e., S satisfies the upper sampling inequality $\sum_{s \in S} |f(s)|^2 \leq C \|f\|_2^2$ for all $f \in \mathcal{H}$. Then for every $\varepsilon > 0$ there is a constant $r = r(\varepsilon)$ such that

$$\sup_{x \in \mathbb{R}^d} \sum_{s \in S \setminus B_r(x)} |k(x, s)|^2 < \varepsilon^2. \tag{HAP}$$

Under the assumptions of weak localization (WL) and (2-4), an upper sampling inequality implies that for some (and hence all) $\rho > 0$

$$\max_{x \in \mathbb{R}^d} \#(S \cap B_\rho(x)) < \infty.$$

We call such a set S *relatively separated*. See also [Führ et al. 2017, Lemma 3.7].

The two localization properties (WL) and (HAP) are the key properties of the reproducing kernel required for an abstract density theorem to hold. For reproducing kernel Hilbert spaces embedded in $L^2(\mathbb{R}^d)$ this can be stated as follows [Führ et al. 2017, Corollary 4.1].

Theorem 2.4. *Let $\mathcal{H} \subseteq L^2(\mathbb{R}^d, dx)$ be a reproducing kernel Hilbert space with kernel k . Assume that k satisfies the boundedness property (2-4) on the diagonal, the weak localization (WL) and the homogeneous approximation property (HAP).*

- (i) *If S is a sampling set for \mathcal{H} , then $D_0^-(S) \geq 1$.*
- (ii) *If S is an interpolating set for \mathcal{H} , then $D_0^+(S) \leq 1$.*

This result holds under a set of natural assumptions on metric measure spaces and conditions on the reproducing kernel. We will not dwell on the geometric conditions, e.g., doubling measure, as these are clearly satisfied for \mathbb{R}^d with μ equivalent to Lebesgue measure. We want to verify Theorem 2.4 for $\mathcal{H} = \text{PW}_\Omega(H_a)$ for a suitable class of symbols a . The boundedness of the diagonal of the kernel was

already established in Proposition 2.2, (2-4). To prove Theorems B and C we therefore need to verify the properties (WL) and (HAP) for the reproducing kernel Hilbert space $\text{PW}_\Omega(H_a)$.

Observe that (WL) is equivalent to the condition

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus B_r(0)} |T_{-x}k_x(y)|^2 dy < \varepsilon^2 \tag{2-6}$$

for the centered reproducing kernels. We will show the stronger statement that the set $\{T_{-x}k_x : x \in \mathbb{R}^d\}$ is relatively compact in $L^2(\mathbb{R}^d)$. The Riesz–Kolmogorov compactness theorem then implies (2-6) and thus (WL).

The proof of (HAP) requires some additional local regularity of k_x . We will use prominently elliptic regularity theory to show that $\{T_{-x}k_x : x \in \mathbb{R}^d\}$ is relatively compact in all Sobolev spaces W_2^s . For the proof of (HAP) it is fundamental that the point evaluation on $\text{PW}_\Omega(H_a)$ can be expressed two-fold as

$$f(x) = \langle f, k_x \rangle_{L^2} = \langle f, T_x \kappa \rangle_{W_2^s} \quad \text{for all } f \in \text{PW}_\Omega(H_a). \tag{2-7}$$

2D. Classes of symbols, limit operators. First we define the relevant symbol classes. Let

$$\tau_x(H_a) = T_{-x}H_aT_x = H_{T_{-x}a} \tag{2-8}$$

be the conjugation of H_a by the translation T_x . If $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$, observe that $\tau_x(H_a)$ is again a self-adjoint, uniformly elliptic operator with domain W_2^2 and core $C_c^\infty(\mathbb{R}^d)$. In this section we describe symbol classes that ensure that $\{\tau_x(H_a)f : x \in \mathbb{R}^d\}$ is relatively compact in $L^2(\mathbb{R}^d)$ for all $f \in C_c^\infty(\mathbb{R}^d)$. Equivalently, every sequence $\tau_{x_k}(H_a)f$ has a norm-convergent subsequence. If (x_k) is bounded, this follows from the continuity of $x \mapsto T_x f$. To treat unbounded sequences we need some terminology.

Since in Section 6 we will deal with a nonmetrizable compactification of \mathbb{R}^d , we formulate most results for nets $(x_\lambda)_{\lambda \in \Lambda}$ instead of sequences. (Here Λ is a directed set with a partial order \succeq and we write $\lim_\lambda x_\lambda$ for the limit of a net when it exists.)

Definition 2.5. Assume $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. If the net $(x_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^d$ diverges to infinity and there is an operator $H \in \mathcal{B}(W_2^2, L^2(\mathbb{R}^d))$ such that $\lim_\lambda \tau_{x_\lambda}(H_a)f = Hf$ for all $f \in C_c^\infty(\mathbb{R}^d)$, then H is called a *limit operator* of H_a .

Remark 2.6. (i) Existence and uniqueness of the limit operator follow from the Banach–Steinhaus theorem.

(ii) We do not even scratch the surface of the method of limit operators: see, amongst many others, [Rabinovich et al. 2004a; 2004b; Špakula and Willett 2017], and in the C^* -algebra setting [Davies and Georgescu 2013; Georgescu 2011; 2018].

(iii) Limit operators are related to compactifications of \mathbb{R}^d . An example can be found in Section 6B.

2D1. Compact orbits. Identity (2-8) suggests that compactness properties of $\{\tau_x(H_a) : x \in \mathbb{R}^d\}$ are related to compactness properties of $\{T_{-x}a : x \in \mathbb{R}^d\}$, so we investigate these first.

Lemma 2.7. (i) If $f \in C_b^\infty(\mathbb{R}^d)$, then $\{T_x f : x \in \mathbb{R}^d\}$ is relatively compact in the Fréchet space $C^\infty(\mathbb{R}^d)$ with respect to its topology of uniform convergence of all derivatives on compact sets.

(ii) In particular, if $\lim_{\lambda} T_{x_{\lambda}} f = g$ pointwise, then $\lim_{\lambda} T_{x_{\lambda}} f = g$ in $C^{\infty}(\mathbb{R}^d)$. The limit function g is again in $C_b^{\infty}(\mathbb{R}^d)$.

Proof. (i) The space $C^{\infty}(\mathbb{R}^d)$ has the Heine–Borel property [Rudin 1973, 1.46], so it suffices to verify that $\{T_x f : x \in \mathbb{R}^d\}$ is bounded in $C^{\infty}(\mathbb{R}^d)$, which means that

$$\|D^{\alpha} T_x f\|_{L^{\infty}(B_r(0))} < C_{\alpha,r} \quad \text{for all } x \in \mathbb{R}^d \text{ and all } r > 0, \alpha \in \mathbb{N}_0^d.$$

But this is trivial for $f \in C_b^{\infty}(\mathbb{R}^d)$, since all derivatives are globally bounded.

(ii) We apply the following observation: A net converges to a limit g if and only if every subnet has a subnet that converges to g . By (i) every subnet of $(T_{x_{\lambda}} f)_{k \in \mathbb{N}}$ has a subnet $(T_{z_{\lambda}} f)_{k \in \mathbb{N}}$ that converges in $C^{\infty}(\mathbb{R}^d)$ (to the limit function g). We conclude that $(T_{x_{\lambda}} f)_{k \in \mathbb{N}}$ converges to g in $C^{\infty}(\mathbb{R}^d)$. As all functions and their derivatives of all orders are bounded and continuous, this is true for the limit as well. \square

Proposition 2.8. *Let $a \in C_b^{\infty}(\mathbb{R}^d, \mathbb{C}^{d \times d})$, $k, m \in \mathbb{N}_0$, and assume $\lim_{\lambda} T_{-x_{\lambda}} a = b$ pointwise. Then, for every $f \in W_2^{2m+2k}$*

$$\lim_{\lambda} \|(\tau_{x_{\lambda}}(H_a^k) - H_b^k) f\|_{W_2^{2m}} = 0.$$

Proof. We treat the case $k = 1$ first and assume for the moment that $f \in C_c^{\infty}(\mathbb{R}^d)$. Set $a^{(\lambda)} = T_{-x_{\lambda}} a$. We can express H_a in the form $H_a = \sum_{|\beta| \leq 2} a_{\beta} D^{\beta}$, with coefficients $a_{\beta} \in C_b^{\infty}(\mathbb{R}^d)$, and estimate, for every multindex α with $|\alpha| \leq 2m$,

$$|D^{\alpha}(H_{a^{(\lambda)}} - H_b) f| = \left| D^{\alpha} \sum_{|\beta| \leq 2} (a_{\beta}^{(\lambda)} - b_{\beta}) D^{\beta} f \right| = \left| \sum_{|\beta| \leq 2} \sum_{|\gamma| \leq |\alpha|} \binom{\alpha}{\gamma} D^{\gamma} (a_{\beta}^{(\lambda)} - b_{\beta}) D^{\alpha - \gamma + \beta} f \right|.$$

By Lemma 2.7 we have $\lim_{\lambda} D^{\gamma} a^{(\lambda)} = D^{\gamma} b$ uniformly on compact sets, so the convergence is actually uniform on $\text{supp } f$, and thus

$$\lim_{\lambda} \|D^{\alpha}(H_{a^{(\lambda)}} - H_b) f\|_{\infty} = 0.$$

Consequently

$$\begin{aligned} \|(H_{a^{(\lambda)}} - H_b) f\|_{W_2^{2m}} &\leq C \max_{|\alpha| \leq 2m} \|D^{\alpha}(H_{a^{(\lambda)}} - H_b) f\|_2 \\ &\leq C |\text{supp } f|^{1/2} \max_{|\alpha| \leq 2m} \|D^{\alpha}(H_{a^{(\lambda)}} - H_b) f\|_{\infty} \rightarrow 0. \end{aligned}$$

As $C_c^{\infty}(\mathbb{R}^d)$ is dense in W_2^{2m+2} , and the operators $H_{a^{(\lambda)}}$ are uniformly bounded from W_2^{2m+2} to W_2^{2m} , a standard density argument (see, e.g., [Teschl 2009, Lemma 1.14]) implies $\|(H_{a^{(\lambda)}} - H_b) f\|_{W_2^{2m}} \rightarrow 0$ for all $f \in W_2^{2m+2}$.

For $k > 1$ observe that

$$H_a^k f - H_b^k f = H_a^{k-1}(H_a f - H_b f) + (H_a^{k-1} - H_b^{k-1})H_b f.$$

As $\lim_{\lambda} \|(H_{a^{(\lambda)}} - H_b) f\|_{W_2^{2m}} = 0$ for $f \in W_2^{2m+2}$, the result follows by induction on k . \square

Remark. The statement of the proposition and its proof are valid under the following more general conditions: $a_{\lambda}, b \in C_b^{\infty}(\mathbb{R}^d)$, $a_{\lambda} \xrightarrow{C^{\infty}} b$, and $(H_{a_{\lambda}})$ is uniformly bounded from W_2^2 to $L^2(\mathbb{R}^d)$.

Though not needed in the sequel, we state an interesting corollary that shows how compactness properties of the orbit $\{T_x a : x \in \mathbb{R}^d\}$ are transferred to compactness properties of $\{\tau_x(H_a) : x \in \mathbb{R}^d\}$.

Corollary 2.9. *If $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and $f \in C_c^\infty(\mathbb{R}^d)$ the set $\{\tau_x(H_a)f : x \in \mathbb{R}^d\}$ is relatively compact in every Sobolev space W_2^s , $s > 0$.*

Proof. The set $\{T_x a : x \in \mathbb{R}^d\}$ is relatively compact in $C^\infty(\mathbb{R}^d)$ by Lemma 2.7, and Proposition 2.8 says that the mapping $a \mapsto H_a f$ is continuous from $\overline{\{T_x a : x \in \mathbb{R}^d\}}^{C^\infty(\mathbb{R}^d)}$ to W_2^s . \square

2D2. Slowly oscillating symbols. In the next step we single out a subclass of operators for which the spectral theory is sufficiently simple. In our approach it is essential that the limit operators do not have the endpoint 0 and Ω of the spectrum as eigenvalues. The limits of translates of *slowly oscillating* symbols are constant, if they exist (Lemma 2.13 below), so the limit operators are similar to the Laplacian. This will be used in Section 6 to compute the critical density.

Definition 2.10. An X -valued function $f \in C_b^u(\mathbb{R}^d, X)$ is slowly oscillating¹ if for all compact subsets $M \subset \mathbb{R}^d$

$$\lim_{|x| \rightarrow \infty} \sup_{m \in M} \|f(x) - f(x + m)\|_X = 0.$$

In fact, it suffices to use the closed unit ball \bar{B}_1 instead of an arbitrary compact set M .

We denote the space of all slowly oscillating functions on \mathbb{R}^d by $C_h(\mathbb{R}^d, X)$ and define $C_h^\infty(\mathbb{R}^d, X) = C_h(\mathbb{R}^d, X) \cap C_b^\infty(\mathbb{R}^d, X)$.

The space $C_h(\mathbb{R}^d)$ with the $\|\cdot\|_\infty$ -norm and pointwise multiplication is a commutative C^* -subalgebra of $C_b^u(\mathbb{R}^d)$.

We will need the following characterization of $C_h^\infty(\mathbb{R}^d, X)$. The statement is folklore, but we do not know a formal reference. For completeness we sketch the simple proof.

Lemma 2.11. *A function f is in $C_h^\infty(\mathbb{R}^d, X)$ if and only if $f \in C_b^\infty(\mathbb{R}^d, X)$ and $\lim_{|x| \rightarrow \infty} \partial_k f(x) = 0$ for all $1 \leq k \leq d$.*

Proof. Assume that $f \in C_b^\infty(\mathbb{R}^d, X)$ and $\lim_{|x| \rightarrow \infty} \partial_k f(x) = 0$ for all $1 \leq k \leq d$ and choose $M = [-h, h]^d$. Writing $m \in M$ as $m = \sum_{k=1}^d h_k e_k$ with $|h_k| \leq h$, the difference in Definition 2.10 is

$$f\left(x + \sum_{k=1}^d h_k e_k\right) - f(x) = \sum_{k=0}^{d-1} \int_{x + \sum_{l \leq k} h_l e_l}^{x + \sum_{l \leq k+1} h_l e_l} \partial_{k+1} f.$$

This implies that $\sup_{m \in M} \|f(x + m) - f(x)\|_X \rightarrow 0$ for $|x| \rightarrow \infty$.

Conversely, assume that $f \in C_h^\infty(\mathbb{R}^d, X)$. Fix $\varepsilon > 0$ and let $\eta \in C_c^\infty(\mathbb{R}^d)$ with $\text{supp } \eta \subset B_1$, $\eta \geq 0$, $\int_{\mathbb{R}^d} \eta(x) dx = 1$. Then $\eta_\tau(x) = \tau^{-d} \eta(\tau^{-1}x)$, $\tau > 0$, is an approximate unit. This implies that for $f \in C_b^u(\mathbb{R}^d, X)$ bounded and uniformly continuous

$$\lim_{\tau \rightarrow 0^+} \sup_{x \in \mathbb{R}^d} \|f(x) - f * \eta_\tau(x)\|_X = 0. \tag{2-9}$$

To estimate the partial derivative of $f \in C_h^\infty(\mathbb{R}^d, X)$ we introduce the approximate unit:

$$\|\partial_k f(x)\|_X \leq \|\partial_k f(x) - \eta_\tau * \partial_k f(x)\|_X + \|\eta_\tau * \partial_k f(x)\|_X = \text{I}_\tau + \text{II}_\tau.$$

¹In the literature f is also called “of vanishing oscillation at infinity” or a Higson function.

Since all derivatives of $f \in C_h^\infty(\mathbb{R}^d, X)$ are bounded and uniformly continuous, by (2-9) there exists τ_0 such that $I_\tau < \varepsilon/2$ for $t < \tau_0$. Fix $\tau < \tau_0$ and observe that $\eta_\tau * \partial_k f = \partial_k \eta_\tau * f$ and that $\int_{\mathbb{R}^d} \partial_k \eta_\tau(y) dy = 0$. So

$$\begin{aligned} \Pi_\tau &= \|\eta_\tau * \partial_k f(x)\|_X = \left\| \int_{\mathbb{R}^d} (f(x) - f(y)) \partial_k \eta_\tau(x - y) dy \right\|_X \\ &\leq \sup_{y \in B_\tau(x)} \|f(x) - f(y)\|_X \int_{\mathbb{R}^d} |\partial_k \eta_\tau(x - y)| dy \\ &\leq C_\tau \sup_{y \in B_\tau(0)} \|f(x) - f(x + y)\|_X. \end{aligned}$$

As $f \in C_h^\infty(\mathbb{R}^d, X)$, there is $R > 0$ such that $\Pi_\tau \leq \varepsilon/(2C_\tau)$ for $|x| > R$, and thus $\|\eta_\tau * \partial_k f(x)\|_X < \varepsilon$ for $|x| > R$. □

Example 2.12. A typical example of a genuinely slowly oscillating function is $a(x) = \sin |x|^{1/2}(1 - \varphi(x))$ for some $\varphi \in C_c^\infty(\mathbb{R}^d)$ with $\varphi(x) = 1$ near 0. (The cut-off of the singularity at 0 serves to make all derivatives of a bounded, but, of course, it is immaterial for the asymptotic behavior.)

Our interest in $C_h(\mathbb{R}^d, X)$ stems from the following fact (see [Rabinovich et al. 2004a, Proposition 2.4.1]):

Lemma 2.13. Assume that $f \in C_h(\mathbb{R}^d, X)$ and $(x_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^d$ diverges to infinity, $|x_\lambda| \rightarrow \infty$. If $\lim_\lambda T_{-x_\lambda} f(x) = g(x)$ exists for all $x \in \mathbb{R}^d$, then g is constant.

Proof. Let $x, x' \in \mathbb{R}^d$. Definition 2.10 with $M = \{x, x'\}$ shows that for all $\varepsilon > 0$ there exists an index $\lambda_\varepsilon = \lambda_\varepsilon(x, x')$ such that $\|f(x + x_\lambda) - f(x_\lambda)\|_X < \varepsilon/2$ and $\|f(x' + x_\lambda) - f(x_\lambda)\|_X < \varepsilon/2$ for all $\lambda \geq \lambda_\varepsilon$. So $\|f(x + x_\lambda) - f(x' + x_\lambda)\|_X < \varepsilon$ for all $\lambda \geq \lambda_\varepsilon$. If $g = \lim_\lambda T_{-x_\lambda} f$ exists, it follows that $\|g(x) - g(x')\|_X \leq \varepsilon$. As $\varepsilon > 0$ was arbitrary, g must be constant. □

3. Statement of the density theorem

We state our main theorems. A first version describes a general setup for symbols in the class $C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ under additional assumptions on the spectra of the limit operators. We then formulate a corollary for slowly oscillating symbols, where the assumptions on the limit operators are automatically satisfied. We discuss possible applications of the general version in Section 7.

Theorem 3.1. Assume that $H_a = -\sum_{j,k=1}^d \partial_j a_{jk} \partial_k$ is uniformly elliptic with symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. Let $\text{PW}_\Omega(H_a)$ be the Paley–Wiener space as defined in Section 2B. Assume that Ω is not an eigenvalue of any limit operator H_b .

- If S is a set of stable sampling for $\text{PW}_\Omega(H_a)$, then

$$D_0^-(S) \geq 1.$$

- If S is a set of interpolation for $\text{PW}_\Omega(H_a)$, then

$$D_0^+(S) \leq 1.$$

The following consequence is a more explicit version of [Theorem B](#) of the [Introduction](#), where we have used the equivalence of [Lemma 2.11](#) to avoid the formal definition of $C_h^\infty(\mathbb{R}^d)$.

Corollary 3.2. *Assume that $H_a = -\sum_{j,k=1}^d \partial_j a_{jk} \partial_k$ is uniformly elliptic with symbol $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$.*

- *If S is a set of stable sampling for $\text{PW}_\Omega(H_a)$, then $D_0^-(S) \geq 1$.*
- *If S is a set of interpolation for $\text{PW}_\Omega(H_a)$, then $D_0^+(S) \leq 1$.*

Proof of Corollary 3.2. If $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$, then by [Lemma 2.7](#) every net $(x_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^d$ that diverges to infinity has a subnet $(x_\mu)_{\mu \in M}$ such that $\lim_\mu T_{-x_\mu} a = b$ in the topology of $C^\infty(\mathbb{R}^d)$ for a symbol b . This symbol b is constant by [Lemma 2.13](#) and positive definite; so H_b is similar to the Laplacian and has no point spectrum. □

4. Proof of weak localization of the kernel

To prove [Theorem 3.1](#) we invoke [Theorem 2.4](#) and verify its main hypotheses (WL) and (HAP) on the reproducing kernel.

Let $Q_h = [-h/2, h/2]^d$ be the cube of side-length h , and let $\varphi_x^h(y) = h^{-d} \chi_{Q_h}((y-x)/h) = T_x \varphi_0^h(y)$ be the usual approximate unit.

Lemma 4.1. *We have $\lim_{h \rightarrow 0} \|\chi_{[0,\Omega]}(H_a)\varphi_x^h - k_x\|_2 = 0$ uniformly in $x \in \mathbb{R}^d$.*

Proof. Let $f \in \text{PW}_\Omega(H_a)$. Then

$$\begin{aligned} |\langle f, \chi_{[0,\Omega]}(H_a)\varphi_x^h - k_x \rangle| &= |\langle f, \varphi_x^h \rangle - f(x)| = h^{-d} \left| \int_{Q_h(x)} (f(y) - f(x)) dy \right| \\ &\leq h^{-d} \int_{Q_h(x)} |f(y) - f(x)| dy \leq \sup_{z \in \mathbb{R}^d} |\nabla f(z)| h^{-d} \int_{Q_h(x)} |y - x| dy \\ &\leq C \|\nabla f\|_\infty h. \end{aligned}$$

Since $f \in W_2^s(\mathbb{R}^d)$ for all $s \geq 0$, we apply first the Sobolev embedding (with $s > d/2 + 1$) and then [Lemma 2.1](#) and obtain

$$\|\nabla f\|_\infty \leq C_1 \|f\|_{W_2^s} \leq C \|f\|_2,$$

since $f \in \text{PW}_\Omega(H_a)$. Consequently,

$$|\langle f, \chi_{[0,\Omega]}(H_a)\varphi_x^h - k_x \rangle| \leq Ch \|f\|_2.$$

Taking the supremum over $f \in \text{PW}_\Omega(H_a)$, we obtain

$$\|\chi_{[0,\Omega]}(H_a)\varphi_x^h - k_x\|_2 = \sup_{f \in \text{PW}_\Omega(H_a), \|f\|_2=1} \langle f, \chi_{[0,\Omega]}(H_a)\varphi_x^h - k_x \rangle \leq Ch.$$

As this estimate is independent of x , we have shown that $\chi_{[0,\Omega]}(H_a)\varphi_x^h \rightarrow k_x$ in $L^2(\mathbb{R}^d)$ uniformly in x . □

The following result relates the reproducing kernel of a limit operator of H_a to the original kernel. It expresses a form of continuous dependence of the reproducing kernel of the matrix symbol of H_a . We will denote the point spectrum of an operator H by $\sigma_p(H)$.

Theorem 4.2. *Let H_a with symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$, and let $(x_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^d$ be an unbounded net such that $\lim_\lambda T_{-x_\lambda} a = b$ pointwise. Assume that $\Omega \notin \sigma_p(H_b)$. Let \tilde{k} be the reproducing kernel of $\text{PW}_\Omega(H_b)$. Then*

$$\lim_\lambda T_{-x_\lambda} k_{x_\lambda} = \tilde{k}_0,$$

with convergence in W_2^s for every $s \geq 0$.

Before the proof we remind the reader of the following standard facts of spectral theory; see, e.g., [Teschl 2009, Chapter 6.6]. Although in the literature these results are formulated for sequences of operators, the statements and proofs are equally valid for nets.²

Let $H_\lambda, \lambda \in \Lambda$, and H_b be self-adjoint operators with a common core \mathcal{D} . If $H_\lambda f \rightarrow H_b f$ for all $f \in \mathcal{D}$, then, for every $F \in C_b(\mathbb{R})$,

$$F(H_\lambda) f \rightarrow F(H_b) f \quad \text{for all } f \in L^2(\mathbb{R}^d). \tag{4-1}$$

Furthermore, if $\chi_{\{\alpha\}}(H_b) = \chi_{\{\beta\}}(H_b) = 0$, i.e., $\alpha, \beta \notin \sigma_p(H_b)$, then

$$\chi_{[\alpha, \beta]}(H_\lambda) f \rightarrow \chi_{[\alpha, \beta]}(H_b) f \quad \text{for all } f \in L^2(\mathbb{R}^d). \tag{4-2}$$

Proof of Theorem 4.2. We split the difference $T_{-x_\lambda} k_{x_\lambda} - \tilde{k}_0$ into three terms and then estimate their W_2^s -norms separately:

$$\begin{aligned} & \|T_{-x_\lambda} k_{x_\lambda} - \tilde{k}_0\|_{W_2^s} \\ & \leq \|T_{-x_\lambda} k_{x_\lambda} - T_{-x_\lambda} \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h\|_{W_2^s} + \|T_{-x_\lambda} \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h - \chi_{[0, \Omega]}(H_b) \varphi_0^h\|_{W_2^s} + \|\chi_{[0, \Omega]}(H_b) \varphi_0^h - \tilde{k}_0\|_{W_2^s} \\ & = (I) + (II) + (III). \end{aligned}$$

Choose $\varepsilon > 0$.

Step 1: Expression (I) can be estimated by

$$\|T_{-x_\lambda} k_{x_\lambda} - T_{-x_\lambda} \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h\|_{W_2^s} = \|k_{x_\lambda} - \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h\|_{W_2^s} \leq C_s \|k_{x_\lambda} - \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h\|_2.$$

The first equality holds by the translation invariance of the Sobolev norm; the second inequality is a consequence of Lemma 2.1. By Lemma 4.1 there exists $h_\varepsilon > 0$ such that, for every $0 < h < h_\varepsilon$,

$$\|k_x - \chi_{[0, \Omega]}(H_a) \varphi_x^h\|_2 < \frac{\varepsilon}{3C_s}$$

for all $x \in \mathbb{R}^d$. So for $h < h_\varepsilon$, we obtain (I) $< \varepsilon/3$. Similarly, we achieve (III) $< \varepsilon/3$ for every $h < h'_\varepsilon$.

Step 2: To bound the decisive term (II), we bring in limit operators and elliptic regularity theory. Set $a_\lambda = T_{-x_\lambda} a$. First note that

$$T_{-x_\lambda} \chi_{[0, \Omega]}(H_a) \varphi_{x_\lambda}^h = T_{-x_\lambda} \chi_{[0, \Omega]}(H_a) T_{x_\lambda} \varphi_0^h = \chi_{[0, \Omega]}(\tau_{x_\lambda} H_a) \varphi_0^h = \chi_{[0, \Omega]}(H_{a_\lambda}) \varphi_0^h.$$

We have to verify that

$$\lim_\lambda \|\chi_{[0, \Omega]}(H_{a_\lambda}) \varphi_0^h - \chi_{[0, \Omega]}(H_b) \varphi_0^h\|_{W_2^s} = 0. \tag{4-3}$$

²The cited results use the strong operator topology. As this topology is metrizable on bounded sets, the convergence of nets is equivalent to the convergence of sequences.

For L^2 -convergence ($s = 0$) we argue as follows. By Lemma 2.7 the translates $T_{-x_\lambda} a$ converge to the matrix b uniformly on compact sets. Proposition 2.8 implies that $H_{T_{-x_\lambda} a} f \rightarrow H_b f$ for $f \in W_2^s$, $s \geq 0$. To apply (4-2), we note that $C_c^\infty(\mathbb{R}^d)$ is a common core for all H_{a_λ} and for H_b and that $0 \notin \sigma_p(H_b)$ by (2-3) and $\Omega \notin \sigma_p(H_b)$ by assumption. Therefore (4-3) follows from (4-2).

For the convergence of (4-3) in general Sobolev spaces W_2^s it suffices to treat the case $s = 2k$ for every integer k . Recall that by the results on elliptic regularity in Section 2B the operator $(H_a^k + c_k)$ defines an isomorphism from $W_2^{2k}(\mathbb{R}^d)$ to $L^2(\mathbb{R}^d)$, and since $\tau_{x_\lambda}(H_a^k + c_k) = H_{a_\lambda}^k + c_k$ we obtain

$$\|H_{a_\lambda}^k + c_k\|_{W_2^{2k} \rightarrow L^2} = \|H_a^k + c_k\|_{W_2^{2k} \rightarrow L^2} < \infty.$$

The Sobolev norm can be estimated by the L^2 -norm

$$\|f\|_{W_2^{2k}} = \|T_x f\|_{W_2^{2k}} \leq C_s \|(H_a^k + c_k)T_x f\|_2 = C_s \|T_{-x}(H_a^k + c_k)T_x f\|_2$$

independently of $x \in \mathbb{R}^d$. Thus (II) can be estimated by the L^2 -norm, namely

$$\begin{aligned} \|\chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h - \chi_{[0,\Omega]}(H_b)\varphi_0^h\|_{W_2^s} &\leq C_s \|(H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h - (H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h\|_2 \\ &\leq C_s \|(H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h - (H_b^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h\|_2 \\ &\quad + C_s \|(H_b^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h - (H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h\|_2 \\ &= A_\lambda + B_\lambda. \end{aligned}$$

By Proposition 2.8 we have $(H_{a_\lambda}^k + c_k)f \rightarrow (H_b^k + c_k)f$ in L^2 -norm for all $f \in W_2^{2k}$. In particular, this holds for $f = \chi_{[0,\Omega]}(H_b)\varphi_0^h$; thus $\lim_\lambda B_\lambda = 0$.

For the first term we use spectral theory again. Define $F \in C_c(\mathbb{R})$ such that its restriction to $[0, \Omega]$ satisfies

$$F(t) = t^k + c_k \quad \text{for } t \in [0, \Omega].$$

Then $F(t)\chi_{[0,\Omega]}(t) = (t^k + c_k)\chi_{[0,\Omega]}(t)$, and $\lim_\lambda F(\tau_{x_\lambda}(H_a))f = F(H_b)f$ for all $f \in L^2(\mathbb{R}^d)$ by (4-1). Since the product of bounded operators is continuous in the strong operator topology, it follows that

$$\begin{aligned} \lim_\lambda (H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h &= \lim_\lambda F(H_{a_\lambda})(\lim_\lambda \chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h) \\ &= F(H_b)\chi_{[0,\Omega]}(H_b)\varphi_0^h = (H_b^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h, \end{aligned}$$

and so $\lim_\lambda A_\lambda = 0$.

We can finish the proof as follows. We have already chosen $h < \min\{h_\varepsilon, h'_\varepsilon\}$ so that the terms (I) and (III) are $< \varepsilon/3$ for all $\lambda \in \Lambda$. For this fixed $h > 0$ we can find an index λ_0 such that

$$(II) \leq C_s \|(H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_{a_\lambda})\varphi_0^h - (H_{a_\lambda}^k + c_k)\chi_{[0,\Omega]}(H_b)\varphi_0^h\|_2 < \frac{\varepsilon}{3}$$

for all $\lambda \geq \lambda_0$. Altogether we obtain $\|T_{-x_\lambda} k_{x_\lambda} - \tilde{k}_0\|_2 \leq (I) + (II) + (III) < \varepsilon$. □

Theorem 4.3. Assume that H_a is uniformly elliptic with symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and that no limit operator has the eigenvalue Ω . Then the set $\{T_{-x} k_x : x \in \mathbb{R}^d\}$ is relatively compact in W_2^s for every $s \geq 0$.

Proof. This follows directly from [Theorem 4.2](#). Let $(x_n)_{n \in \mathbb{N}} \subseteq \mathbb{R}^d$ be an arbitrary sequence. By [Lemma 2.7](#) the sequence $T_{-x_n} a$ has a C^∞ -convergent subsequence $T_{-x_{n_l}} a$. If $(x_{n_l})_{l \in \mathbb{N}}$ is bounded, we can assume without loss of generality that $x_{n_l} \rightarrow x \in \mathbb{R}^d$, and $T_{-x_{n_l}} k_{x_{n_l}} \rightarrow T_{-x} k_x$ in W_2^s by the continuity of the translations and [Proposition 2.3](#). If $(x_{n_l})_{l \in \mathbb{N}}$ is unbounded, we can assume $|x_{n_l}| \rightarrow \infty$. This case is settled by [Theorem 4.2](#) and yields the convergence of $T_{-x_{n_l}} k_{x_{n_l}}$. \square

A combination of the above arguments yields the weak localization [\(WL\)](#).

Theorem 4.4. *Assume that H_a is uniformly elliptic with symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and that no limit operator has the eigenvalue Ω . Let k be the reproducing kernel of $\text{PW}_\Omega(H_a)$. Then k satisfies the weak localization property [\(WL\)](#), i.e.,*

$$\lim_{R \rightarrow \infty} \int_{|y-x| > R} |k(x, y)|^2 dy = 0.$$

Proof. By [Theorem 4.3](#) (for $s = 0$) the set $\{T_{-x} k_x : x \in \mathbb{R}^d\}$ is relatively compact in $L^2(\mathbb{R}^d)$. The Riesz–Kolmogorov theorem implies that for all $\varepsilon > 0$ there is $R > 0$ such that for all $x \in \mathbb{R}^d$

$$\int_{\mathbb{R}^d \setminus B_R(0)} |T_{-x} k_x(y)|^2 dy < \varepsilon^2.$$

By a change of variable this expression reads as

$$\int_{|y-x| > R} |k(x, y)|^2 dy < \varepsilon^2,$$

and this is [\(WL\)](#). \square

5. Proof of the homogeneous approximation property (HAP)

Next we prove the homogeneous approximation property. Recall that $T_x \kappa$ is the reproducing kernel for W_2^s with $\hat{\kappa}(\omega) = (1 + |\omega|^2)^{-s}$.

Lemma 5.1. *If S is a relatively separated set in \mathbb{R}^d , then $\{T_x \kappa : x \in S\}$ is a Bessel sequence for W_2^s , $s > d/2$.*

Proof. By standard facts of frame theory (see, e.g., [\[Heil 2011, Theorem 7.6\]](#)) the Bessel property is equivalent to the boundedness of the Gramian $G = (\langle T_x \kappa, T_y \kappa \rangle_{W_2^s})_{x, y \in S}$ on $\ell^2(S)$. To deduce the boundedness of G we first show that G possesses exponential off-diagonal decay and then apply Schur’s test. The off-diagonal decay follows from a (well-known) calculation. Let \mathcal{J}_r denote the Bessel function of the first kind and \mathcal{K}_r the modified Bessel function of the second kind. Then by [\[Wendland 2005, Theorem 6.13\]](#) or [\[Grafakos 2004, Appendix B\]](#)

$$\begin{aligned} \langle T_x \kappa, T_y \kappa \rangle_{W_2^s} &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} \widehat{T_x \kappa}(\omega) \overline{\widehat{T_y \kappa}(\omega)} (1 + |\omega|^2)^s d\omega \\ &= (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{-i(x-y)\omega} (1 + |\omega|^2)^{-s} d\omega \\ &= C|x - y|^{-(d-2)/2} \int_0^\infty (1 + r^2)^{-s} \mathcal{J}_{(d-2)/2}(r|x - y|) r^{d/2} dr \\ &= C'|x - y|^{s-d/2} \mathcal{K}_{s-d/2}(|x - y|). \end{aligned}$$

Using the asymptotic decay $\mathcal{K}_r(x) \sim \sqrt{\pi/(2x)}e^{-x}$ for $x \rightarrow \infty$, see, e.g., [DLMF 2020, equation 10.25.3], the off-diagonal decay of G is

$$|\langle T_x \kappa, T_y \kappa \rangle_{W_2^s}| \leq C'' |x - y|^{s-d/2-1/2} e^{-|x-y|} \quad (|x - y| \rightarrow \infty). \tag{5-1}$$

The off-diagonal decay of the Gramian implies the boundedness of the Gramian as follows. By (5-1) there exists $N_0 \in \mathbb{N}$ such that $|G_{xy}| \leq C e^{-c|x-y|}$ if $|x - y| > N_0$. Obviously, $|G_{xy}| \leq \|\kappa\|_{W_2^s}^2$ is bounded for all x, y .

For $x \in S$ and $k \in \mathbb{N}_0$ set $A_k(x) = \{y \in S : k < |y - x| \leq k + 1\}$. Since $S \subset \mathbb{R}^d$ is relatively separated, there exists $r > 0$ such that

$$\max \#(S \cap B_r(x)) < \infty.$$

A covering argument (of a large ball $B_R(z)$ by balls $B_r(x)$) implies that $\#(S \cap B_R(z)) = \mathcal{O}(R^d)$ for arbitrary $R > 0$. Consequently we also obtain $\#A_k(x) \leq Ck^d$ independent of x . Then

$$\begin{aligned} \sum_{y \in S} |G_{xy}| &= \sum_{k=0}^{\infty} \sum_{y \in A_k(x)} |G_{xy}| = \sum_{k=0}^{N_0} \sum_{y \in A_k(x)} |G_{xy}| + \sum_{k > N_0} \sum_{y \in A_k(x)} |G_{xy}| \\ &\leq C_0 \#(B_{N_0+1}(x) \cap S) + C \sum_{k > N_0} e^{-ck} \#A_k(x) \\ &\leq C_1(N_0 + 1)^d + C_2 \sum_{k > N_0} e^{-ck} k^d. \end{aligned}$$

This expression is bounded independently of x . Now Schur’s test implies that the Gramian is bounded on $\ell^2(S)$. □

Theorem 5.2 (HAP). *Assume that H_a is uniformly elliptic with symbol $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and that $\Omega \notin \sigma_p(H_b)$ for every limit operator H_b . Let $\{k_x : x \in S\}$ be a Bessel sequence in $\text{PW}_\Omega(H_a)$. Then for every $\varepsilon > 0$ there exists an $R > 0$ such that for all $y \in \mathbb{R}^d$*

$$\sum_{x \in S \setminus B_R(y)} |k(y, x)|^2 < \varepsilon^2.$$

Proof. If $\{k_x : x \in S\}$ is a Bessel sequence of reproducing kernels, then S is relatively separated in \mathbb{R}^d (see [Führ et al. 2017, Lemma 3.7]). Lemma 5.1 implies that $\{T_x \kappa : x \in S\}$ is also a Bessel sequence in W_2^s for $s > d/2$.

Choose $\varepsilon > 0$. Since $\{T_{-x} k_x : x \in \mathbb{R}^d\}$ is relatively compact in W_2^s for $s \geq 0$ by Theorem 4.3, the Riesz–Kolmogorov theorem for translation-invariant Banach spaces [Feichtinger 1984] asserts that there exists a $R = R_\varepsilon > 0$ and a function $\psi \in C_c^\infty(\mathbb{R}^d)$ satisfying $\psi|_{B_{R/2}(0)} = 1$, $\text{supp } \psi \subseteq B_R(0)$ such that

$$\|T_{-x} k_x (1 - \psi)\|_{W_2^s} \leq \varepsilon \quad \text{for all } x \in \mathbb{R}^d.$$

We now use the fundamental observation (2-7) that the point evaluation in $\text{PW}_\Omega(H_a)$ can be expressed in two ways. For $f = k_x$ we have

$$k(y, x) = \langle k_y, k_x \rangle_{L^2} = \langle k_y, T_x \kappa \rangle_{W_2^s}. \tag{5-2}$$

Since $\{T_x\kappa : x \in S\}$ is a Bessel sequence in W_2^s with bound B , the set $\{T_{x-y}\kappa : x \in S\}$ is a Bessel sequence with the same bound. Observe that for $|u| > R$ we obtain

$$\langle \psi T_{-y}k_y, T_u\kappa \rangle_{W_2^s} = T_{-y}k_y(u)\psi(u) = 0.$$

This implies

$$\begin{aligned} \sum_{x \in S \setminus B_R(y)} |\langle k_y, T_x\kappa \rangle_{W_2^s}|^2 &= \sum_{x \in S \setminus B_R(y)} |\langle T_{-y}k_y, T_{x-y}\kappa \rangle_{W_2^s}|^2 \\ &= \sum_{x \in S \setminus B_R(y)} |\langle (1 - \psi)T_{-y}k_y, T_{x-y}\kappa \rangle_{W_2^s}|^2 \\ &\leq B \|(1 - \psi)T_{-y}k_y\|_{W_2^s}^2 \leq B\varepsilon^2, \end{aligned}$$

and this is the homogeneous approximation property, (HAP). □

Proof of Theorem 3.1. After the verification of the properties (WL) and (HAP) of the reproducing kernel, the version for the dimension-free density of Theorem 2.4 is applicable and yields Theorem 3.1. The statement asserts the existence of a critical density for sampling and interpolation with the dimension-free Beurling density $D_0^\pm(S)$. □

6. Geometric Beurling densities

In this section we derive results for the geometric densities (2-5). According to Theorem 2.4 this step requires the computation of the averaged trace $|\mu(B_r(x))|^{-1} \int_{B_r(x)} k(x, x) d\mu(x)$ of the reproducing kernel. This version of the density theorems is of interest because it relates the critical density in $\text{PW}_\Omega(H_a)$ to the geometry defined by the differential operator H_a . The explicit computation of the averaged trace becomes possible by introducing a suitable compactification of \mathbb{R}^d and then extending the centered kernels $T_{-x}k_x$ to this compactification.

6A. The basic computation: constant coefficients. For reference we mention the case when $H_b = -\sum_{j,k} \partial_j b_{jk} \partial_k = -\sum_{j,k} b_{jk} \partial_j \partial_k$ is a differential operator with constant coefficients b_{jk} . Define

$$\Sigma_\Omega^b = \{\xi \in \mathbb{R}^d : b\xi \cdot \xi \leq \Omega\} = b^{-1/2} \overline{B_{\Omega^{1/2}}(0)},$$

with volume

$$|\Sigma_\Omega^b| = \det(b^{-1/2}) |B_{\Omega^{1/2}}(0)| = \det(b^{-1/2}) \Omega^{d/2} |B_1|. \tag{6-1}$$

Since $\widehat{H_b f}(\xi) = \sum_{j,k} b_{jk} \xi_j \xi_k \widehat{f}(\xi) = (b\xi \cdot \xi) \widehat{f}(\xi)$, the spectral subspace is

$$\text{PW}_\Omega(H_b) = \chi_{[0,\Omega]}(H_b) L^2(\mathbb{R}^d) = \{f \in L^2(\mathbb{R}^d) : \text{supp } \widehat{f} \subseteq \Sigma_\Omega^b\}.$$

The kernel of $\text{PW}_\Omega(H_b)$ is

$$\tilde{k}(x, y) = (2\pi)^{-d/2} (\mathcal{F}^{-1} \chi_{\Sigma_\Omega^b})(x - y), \tag{6-2}$$

whence

$$\tilde{k}(x, x) = (2\pi)^{-d/2} (\mathcal{F}^{-1} \chi_{\Sigma_\Omega^b})(0) = \frac{|\Sigma_\Omega^b|}{(2\pi)^d} = \frac{|B_1|}{(2\pi)^d} \det(b^{-1/2}) \Omega^{d/2}.$$

By Landau's theorem [1967] a sampling set S for $\text{PW}_\Omega(H_b)$ has Beurling density at least $D^-(S) \geq |\Sigma_\Omega^b| / (2\pi)^d$.

6B. The Higson compactification. We recall how a compactification arises in Gelfand theory. Let $C_\gamma(\mathbb{R}^d)$ be a unital C^* algebra of functions on \mathbb{R}^d satisfying the embeddings $C_0(\mathbb{R}^d) \subset C_\gamma(\mathbb{R}^d) \subset C_b(\mathbb{R}^d)$. The maximal ideal space M_γ of $C_\gamma(\mathbb{R}^d)$ is the space of all multiplicative homomorphisms $\varphi : C_\gamma(\mathbb{R}^d) \rightarrow \mathbb{C}$. Equipped with the weak-star topology M_γ is a compact Hausdorff space. The point evaluations $\delta_x(f) = f(x)$ constitute an embedding γ of \mathbb{R}^d into M_γ via $\gamma(x) = \delta_x$, and $\gamma(\mathbb{R}^d)$ is dense in M_γ and homeomorphic to \mathbb{R}^d . Thus, the pair (γ, M_γ) is a compactification of \mathbb{R}^d , which we will call $\gamma\mathbb{R}^d$. The corona of $\gamma\mathbb{R}^d$ is $\partial_\gamma\mathbb{R}^d = \gamma\mathbb{R}^d \setminus \gamma(\mathbb{R}^d)$. By abuse of notation we will identify a point $x \in \mathbb{R}^d$ with its point evaluation δ_x . Then $C_\gamma(\mathbb{R}^d)$ is isometrically isomorphic to $C(\gamma\mathbb{R}^d)$. We denote the image of $f \in C_\gamma(\mathbb{R}^d)$ in $C(\gamma\mathbb{R}^d)$ by \tilde{f} . See, e.g., [Engelking 1977] for the basics of compactifications, and [Gamelin 1969] for compactifications of function algebras.

As noted in Section 2D2 the space $C_h(\mathbb{R}^d)$ of slowly oscillating functions with supremum norm is a commutative unital C^* -algebra. Thus there is a compactification $h\mathbb{R}^d$ of \mathbb{R}^d , the Higson compactification, such that $C_h(\mathbb{R}^d)$ is isometrically isomorphic to $C(h\mathbb{R}^d)$. It is known that $h\mathbb{R}^d$ is nonmetrizable, and even more, points of the corona $h\mathbb{R}^d \setminus \mathbb{R}^d$ can only be reached by nets; see, e.g., [Rabinovich et al. 2004a, 2.4.10]. Therefore we need to work with nets instead of sequences.

The relevance of the Higson compactification and the algebra of slowly oscillating functions in our context is given by the fact that translations act trivially on the corona $\partial_h\mathbb{R}^d$.

Lemma 6.1. *If $x_\lambda \rightarrow \eta \in \partial_h\mathbb{R}^d$, then $x + x_\lambda \rightarrow \eta$ for all $x \in \mathbb{R}^d$.*

Proof. By definition, $x_\lambda \rightarrow \eta \in \partial_h\mathbb{R}^d$ if $f(x_\lambda) \rightarrow \tilde{f}(\eta) = \eta(f)$ for every $f \in C_h(\mathbb{R}^d)$. From the definition of $C_h(\mathbb{R}^d)$ we obtain

$$\lim_\lambda |f(x_\lambda) - f(x + x_\lambda)| = 0$$

for every $x \in \mathbb{R}^d$, so $f(x_\lambda + x) \rightarrow \tilde{f}(\eta)$ for every $f \in C_h(\mathbb{R}^d)$ as well. □

One can show that $h\mathbb{R}^d$ is the maximal compactification of \mathbb{R}^d with this property: every $C_\gamma(\mathbb{R}^d)$ as above with translations acting trivially on $\partial_\gamma\mathbb{R}^d$ is a subalgebra of $C_h(\mathbb{R}^d)$.

We need the following fact [Roe 2003].

Proposition 6.2. *Let $C_\gamma(\mathbb{R}^d)$ be a C^* -algebra of functions on \mathbb{R}^d as above with corresponding compactification $\gamma\mathbb{R}^d$ of \mathbb{R}^d . If $f \in C_\gamma(\mathbb{R}^d)$ satisfies*

$$\tilde{f}|_{\partial_\gamma\mathbb{R}^d} \equiv 0$$

then $f \in C_0(\mathbb{R}^d)$.

Proof. Let $(x_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^d$ be an unbounded net, $\lim_\lambda |x_\lambda| = \infty$. As $\gamma\mathbb{R}^d$ is compact, every subnet of (x_λ) has a convergent subnet $(x_\mu)_{\mu \in M}$, and $\lim_\mu x_\mu = \eta \in \partial_\gamma\mathbb{R}^d$ by the assumption of unboundedness. So $\lim_\mu f(x_\mu) = \tilde{f}(\eta) = 0$ for a subnet of a given subnet, and therefore $\lim_\lambda f(x_\lambda) = 0$. This means $f \in C_0(\mathbb{R}^d)$. □

We next study uniformly elliptic operators H_a with a symbol in $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. By definition a has a continuous extension to $h\mathbb{R}^d$. Thus for $\eta \in \partial_h\mathbb{R}^d$ the symbol $\bar{a}(\eta) = \lim_{x \rightarrow \eta} a(x)$ is well-defined, and by Lemma 2.13 $H_{\bar{a}(\eta)}$ is a differential operator with constant coefficients. Let k^η denote the reproducing kernel of $\text{PW}_\Omega(H_{\bar{a}(\eta)})$. We show that the mapping $x \mapsto T_{-x}k_x$ has a continuous extension to $h\mathbb{R}^d$.

Proposition 6.3. *Let the symbol $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ be the symbol of the operator H_a and let k_x be the reproducing kernel of $\text{PW}_\Omega(H_a)$. Set $K(x) = T_{-x}k_x \in L^2(\mathbb{R}^d)$. Then K extends to a continuous function from $h\mathbb{R}^d$ to $L^2(\mathbb{R}^d)$ by setting $K(\eta) = k_0^\eta$ for $\eta \in h\mathbb{R}^d$. In particular, the diagonal $k(x, x) = \|k_x\|_2^2$ is a slowly oscillating function.*

Proof. By Proposition 2.3 the centered reproducing kernel K is continuous from \mathbb{R}^d to $L^2(\mathbb{R}^d)$. To show that $K \in C_h(\mathbb{R}^d, L^2)$, we need to extend K to the Higson corona $\partial_h\mathbb{R}^d$.

This is accomplished by means of Theorem 4.2. Let $(x_\lambda) \subseteq \mathbb{R}^d$ be an unbounded net such that $x_\lambda \rightarrow \eta \in \partial_h\mathbb{R}^d$. Since $a \in C_h(\mathbb{R}^d, \mathbb{C}^{d \times d})$, there is a continuous function $\bar{a} \in C(h\mathbb{R}^d, \mathbb{C}^{d \times d})$ such that $\lim_\lambda a(x_\lambda) = \bar{a}(\eta)$. Furthermore, for $x \in \mathbb{R}^d$ arbitrary, $x + x_\lambda \rightarrow \eta$ by Lemma 6.1, and this fact implies the pointwise convergence

$$\lim_\lambda T_{-x_\lambda} a(x) = \bar{a}(\eta).$$

Clearly, the spectrum of the (constant-coefficient) operator $H_{\bar{a}(\eta)}$ is continuous and does not contain any eigenvalues. The assumptions of Theorem 4.2 are thus satisfied.

To formulate its conclusion, denote the reproducing kernel of $\text{PW}_\Omega(H_{\bar{a}(\eta)})$ by k^η . Then by Theorem 4.2

$$\lim_\lambda T_{-x_\lambda} k_{x_\lambda} = k_0^\eta$$

in the L^2 -norm, and this holds for every net (x_λ) with $x_\lambda \rightarrow \eta$. Thus we must take the limiting function to be

$$K(\eta) = k_0^\eta = (2\pi)^{-d/2} \mathcal{F}^{-1}(\chi_{\Sigma_\Omega^{\bar{a}(\eta)}}),$$

with the explicit formula for the kernel given by (6-2).

It remains to be shown that the limiting kernel K is continuous on $\partial_h\mathbb{R}^d$. Let $\eta_\lambda \rightarrow \eta \in \partial_h\mathbb{R}^d$. Then with the definition of Σ_Ω^b and (6-1) we obtain

$$\begin{aligned} \|k_0^{\eta_\lambda} - k_0^\eta\|_2^2 &= (2\pi)^{-d} \|\chi_{\Sigma_\Omega^{\bar{a}(\eta_\lambda)}} - \chi_{\Sigma_\Omega^{\bar{a}(\eta)}}\|_2^2 \\ &= (2\pi)^{-d} (|\Sigma_\Omega^{\bar{a}(\eta_\lambda)}| + |\Sigma_\Omega^{\bar{a}(\eta)}| - 2|\Sigma_\Omega^{\bar{a}(\eta_\lambda)} \cap \Sigma_\Omega^{\bar{a}(\eta)}|). \end{aligned}$$

As $a \in C_h(\mathbb{R}^d, \mathbb{C}^{d \times d})$, \bar{a} is continuous on $\partial_h\mathbb{R}^d$, and this expression tends to 0, whence K is also continuous on the corona $\partial_h\mathbb{R}^d$. □

6C. Geometric densities for slowly oscillating symbols. In order to obtain values for the critical densities $D_\mu^\pm(S)$ we need the averaged traces

$$\text{tr}_\mu^-(k) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}^d} \frac{1}{\mu(B_r(x))} \int_{B_r(x)} k(z, z) dz$$

and $\text{tr}_\mu^+(k)$. For the comparison of averaged traces we will need the following well-known fact, whose proof is supplied in Appendix A for completeness.

For $f \in C_b(\mathbb{R}^d)$ set

$$\text{tr}^-(f) = \liminf_{r \rightarrow \infty} \inf_{y \in \mathbb{R}^d} \frac{1}{|B_r(y)|} \int_{B_r(y)} f(x) dx,$$

and define $\text{tr}^+(f)$ similarly with sup instead of inf.

Lemma 6.4. *Assume that $f, g \in C_b(\mathbb{R}^d)$ and $\lim_{x \rightarrow \infty} |f(x) - g(x)| = 0$. Then*

$$\text{tr}^-(f) = \text{tr}^-(g) \quad \text{and} \quad \text{tr}^+(f) = \text{tr}^+(g).$$

Proposition 6.5. *If the symbol a is in $C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$, then the trace of the reproducing kernel satisfies*

$$\lim_{r \rightarrow \infty} \sup_{y \in \mathbb{R}^d} \left| \frac{1}{|B_r(y)|} \int_{B_r(y)} \left(k(x, x) - \frac{|B_1|}{(2\pi)^d} \frac{\Omega^{d/2}}{\det a(x)^{1/2}} \right) dx \right| = 0. \tag{6-3}$$

Equivalently, using the Borel measure $\nu(B) = \int_B \det a(x)^{-1/2} dx$,

$$\lim_{r \rightarrow \infty} \sup_{y \in \mathbb{R}^d} \left| \frac{1}{\nu(B_r(y))} \int_{B_r(y)} k(x, x) dx - \frac{|B_1|}{(2\pi)^d} \Omega^{d/2} \right| = 0. \tag{6-4}$$

Consequently, the averaged trace is

$$\text{tr}_\nu^+(k) = \text{tr}_\nu^-(k) = \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}. \tag{6-5}$$

Proof. We apply Lemma 6.4 to the functions $f(x) = k(x, x)$ and $g(x) = (|B_1| \Omega^{d/2} / (2\pi)^d) \det a(x)^{-1/2}$. Then $k(x, x)$ is bounded by Proposition 2.2 and continuous by Proposition 2.3. Likewise $\det a(x)^{-1/2}$ is bounded and continuous by elliptic regularity. By assumption on a and Proposition 6.3 both functions are in $C_h(\mathbb{R}^d)$ and thus possess the limits $\bar{a}(\eta)$ and $\|K(\eta)\|_2^2$; in particular for a this means that

$$\lim_{x_\lambda \rightarrow \eta} \det a(x_\lambda)^{-1/2} = \det \bar{a}(\eta)^{-1/2}.$$

Using the notation of Section 6A and Proposition 6.3, we obtain

$$\lim_{x_\lambda \rightarrow \eta} \|k_{x_\lambda}\|_2^2 = \lim_{x_\lambda \rightarrow \eta} \|T_{-x_\lambda} k_{x_\lambda}\|_2^2 = \|k_0^\eta\|_2^2 = |\Sigma_\Omega^{\bar{a}(\eta)}| = \frac{|B_1| \Omega^{d/2}}{(2\pi)^d} \det \bar{a}(\eta)^{-1/2}.$$

We conclude that both f and g have the same limit function, and therefore $f - g \in C_0(\mathbb{R}^d)$ by means of Proposition 6.2. Lemma 6.4 now yields (6-3). Equation (6-4) follows after multiplying with $|B_r(y)|/\nu(B_r(y))$ and taking limits. Finally, (6-5) is a direct consequence of (6-4). \square

Equation (6-4) allows us to state our main result on geometric Beurling densities for operators with slowly oscillating symbols in a simple form. In order to do so we need an elementary result on the relation between density and a change of measure.

Lemma 6.6. *Let $d\mu = h dx$ for a positive, continuous function h on \mathbb{R}^d , bounded above and below, $0 < c \leq h(z) \leq C$ for all $z \in \mathbb{R}^d$. Then the dimension-free density condition*

$$D_0^-(S) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}^d} \frac{\#(S \cap B_r(x))}{\int_{B_r(x)} k(y, y) dy} \geq 1$$

holds, if and only if

$$D_\mu^-(S) \geq \text{tr}_\mu^-(k).$$

Similarly

$$D_0^+(S) \leq 1 \quad \text{if and only if} \quad D_\mu^+(S) \leq \text{tr}_\mu^+(k).$$

Proof. The inequality $D_0^-(S) \geq 1$ means that for all $\varepsilon > 0$ there is an $r_\varepsilon > 0$ such that for all $r > r_\varepsilon$

$$\#(S \cap B_r(x)) \geq (1 - \varepsilon) \int_{B_r(x)} k(y, y) dy,$$

or equivalently,

$$\frac{\#(S \cap B_r(x))}{\mu(B_r(x))} \geq (1 - \varepsilon) \frac{1}{\mu(B_r(x))} \int_{B_r(x)} k(y, y) dy.$$

Written in terms of the Beurling density, this is

$$D_\mu^-(S) \geq \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}^d} \frac{\int_{B_r(x)} k(y, y) dy}{\mu(B_r(x))} = \text{tr}_\mu^-(k).$$

The converse is obtained by reading the argument backwards. □

As a direct consequence we obtain the main result on geometric Beurling densities for uniformly elliptic operators with slowly oscillating symbols. This is [Theorem C](#) of the [Introduction](#).

Theorem 6.7. *Assume that $H_a = -\sum_{j,k=1}^d \partial_j a_{jk} \partial_k$ is uniformly elliptic with symbol $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. Let $\text{PW}_\Omega(H_a) = \chi_{[0,\Omega]}(H_a)L^2(\mathbb{R}^d)$ be the corresponding Paley–Wiener space and set $dv(x) = \det(a(x))^{-1/2} dx$.*

- If $S \subseteq \mathbb{R}^d$ is a set of stable sampling for $\text{PW}_\Omega(H_a)$ then

$$D_v^-(S) \geq \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}.$$

- If $S \subseteq \mathbb{R}^d$ is a set of interpolation for $\text{PW}_\Omega(H_a)$, then

$$D_v^+(S) \leq \frac{|B_1|}{(2\pi)^d} \Omega^{d/2}.$$

Proof. We only verify the first assertion. By [Corollary 3.2](#), if S is a set of stable sampling, then $D_0^-(S) \geq 1$. By [Lemma 6.6](#) this is equivalent to

$$D_v^-(S) \geq \text{tr}_v^-(k).$$

The averaged trace $\text{tr}_v^-(k)$ was computed in [\(6-5\)](#) to be $(|B_1|/(2\pi)^d)\Omega^{d/2}$. □

Example 6.8. We consider some special cases of [Theorem 6.7](#).

(i) *Asymptotically constant symbols.* Assume that $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and $\lim_{x \rightarrow \infty} a(x) = b$. Then it is straightforward to verify that $a \in C_h^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ and $D_v^\pm(S) = (\det b)^{1/2} D^\pm(S)$. Thus we may use the original Beurling density, and [Theorem 6.7](#) implies that a sampling set $S \subseteq \mathbb{R}^d$ for $\text{PW}_\Omega(H_a)$ must have density

$$D^-(S) = (\det b)^{-1/2} D_v^-(S) \geq (\det b)^{-1/2} \frac{|B_1|}{(2\pi)^d} \Omega^{d/2} = \frac{|\Sigma_\Omega^b|}{(2\pi)^d},$$

and a set of interpolation S in $\text{PW}_\Omega(H_a)$ must satisfy $D^+(S) \leq |\Sigma_\Omega^b|/(2\pi)^d$. This is [Corollary D](#) of the [Introduction](#). As was to be expected, this coincides with the critical density for the classical Paley–Wiener space $\text{PW}_\Omega(H_b)$; see [\[Landau 1967\]](#).

(ii) *Symbols with radial limits.* Let us consider the class of symbols that possess radial limits at ∞ . We say that $a \in C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$ is spherically continuous, if it possesses radial limits in the following sense. There exists a continuous matrix function $b \in C(S^{d-1}, \mathbb{C}^{d \times d})$ such that

$$\lim_{r \rightarrow \infty} \sup_{\eta \in S^{d-1}} \|a(r\eta) - b(\eta)\|_X = 0.$$

A 3ε -argument shows that these symbols are slowly oscillating. Consequently [Theorem 6.7](#) holds for spherically continuous symbols in $C_b^\infty(\mathbb{R}^d, \mathbb{C}^{d \times d})$. Spherically continuous symbols are related to another compactification, the spherical compactification with corona S^{d-1} . In contrast to the Higson compactification, it is metrizable, but it is much smaller. See [[Cordes 1979](#)] for its use in partial differential equations.

6D. Variable bandwidth in dimension $d = 1$. Let H_a be the differential operator

$$H_a f = -\frac{d}{dx} \left(a \frac{d}{dx} f \right)$$

on $L^2(\mathbb{R})$. This is a Sturm–Liouville operator on \mathbb{R} , and the ellipticity assumption amounts to the conditions $\inf_{x \in \mathbb{R}} a(x) > 0$ and $a \in C_b^\infty(\mathbb{R}^d)$. In [[Gröchenig and Klotz 2017](#)] we argued that the spectral subspaces of H_a can be interpreted as spaces of locally variable bandwidth. Intuitively, the quantity $a(x)^{-1/2}$ is a measure for the bandwidth in a neighborhood of x . We apply [Theorem 6.7](#) to H_a . The relevant measure is $dv(x) = a^{-1/2}(x) dx$, and $v(I) = \int_I a(x)^{-1/2} dx$ for $I \subseteq \mathbb{R}$. Then we have the following necessary density condition for functions of variable bandwidth ([Corollary E](#) of the [Introduction](#)).

Corollary 6.9. *Assume that $a \in C_b^\infty(\mathbb{R})$ and $\lim_{x \rightarrow \pm\infty} a'(x) = 0$. Let $\text{PW}_\Omega(H_a)$ be the Paley–Wiener space associated to H_a .*

(i) *If S is a sampling set for $\text{PW}_\Omega(H_a)$, then*

$$D_v^-(S) = \liminf_{r \rightarrow \infty} \inf_{x \in \mathbb{R}} \frac{\#(S \cap [x - r, x + r])}{v([x - r, x + r])} \geq \frac{\Omega^{1/2}}{\pi}. \tag{6-6}$$

(ii) *If S is a set of interpolation for $\text{PW}_\Omega(H_a)$, then $D_v^+(S) \leq \Omega^{1/2}/\pi$.*

Arguing as in [Lemma 6.6](#), equation (6-6) says that for $\varepsilon > 0$ and r large enough we have

$$\#(S \cap [x - r, x + r]) \geq \left(\frac{\Omega^{1/2}}{\pi} - \varepsilon \right) \int_{x-r}^{x+r} a(y)^{-1/2} dy.$$

Thus the number of samples in an interval $[x - r, x + r]$ is determined by $a(x)^{-1/2}$, which is in line with our interpretation of $a^{-1/2}$ as the local bandwidth.

[Corollary 6.9](#) is precisely the formulation of the necessary density conditions in [[Gröchenig and Klotz 2017](#)]. However, the main result of that work was proved under the restrictive assumption that a is constant outside an interval $[-R, R]$. The proof there dwelt heavily on the scattering theory of one-dimensional Schrödinger operators. The method of this paper yields a significantly more general result with a completely different method of proof. [Corollary 6.9](#) was our dream that motivated this work.

Finally we remark that the density conditions of [Theorem 6.7](#) suggest that the Paley–Wiener spaces $PW_\Omega(H_a)$ associated to a uniformly elliptic differential operator may be taken as an appropriate generalization of variable bandwidth to higher dimensions.

7. Outlook

We have proved necessary density conditions for sampling and interpolation in spectral subspaces of uniformly elliptic partial differential operators with slowly oscillating coefficients. These spectral subspaces may be taken as a suitable generalization of the notion of variable bandwidth to higher dimensions. The emphasis has been on a new method that combines elements from limit operators, regularity theory and heat kernel estimates, and the use of compactifications.

Clearly one can envision manifold extensions of our results and methods. [Theorem 3.1](#) is stated for a significantly larger class of operators and symbols. For instance, it could be applied to higher-order partial differential operators or to Schrödinger operators and to symbols with less smoothness or to almost periodic symbols. However, the spectral theory of such operators is more involved and one needs to find conditions that prevent their limit operators from having a point spectrum at the ends of the spectral interval. As these questions belong to spectral theory rather than sampling theory, we plan to pursue them in a separate publication.

In a different direction one may consider the graph Laplacian on an infinite graph or even a metric measure space endowed with a kernel that satisfies Gaussian estimates [[Coulhon et al. 2012](#)]. While many steps of our proofs remain in place, this set-up opens numerous new questions.

Finally several hidden connections beg to be explored. The identity (6-3) resembles the famous Weyl formula for the asymptotic density of eigenvalues in a spectral interval [[Hörmander 1968](#)]. This observation invites the comparison of the Beurling density with the density of states in spectral theory. We plan to investigate some of these issues in future work.

Appendix A: Averaged traces

For completeness we provide the proof of [Lemma 6.4](#). Recall that

$$\text{tr}^-(f) = \liminf_{r \rightarrow \infty} \inf_{y \in \mathbb{R}^d} \frac{1}{|B_r(y)|} \int_{B_r(y)} f(x) dx.$$

If $f, g \in C_b(\mathbb{R}^d)$ and $\lim_{|x| \rightarrow \infty} |f(x) - g(x)| = 0$, then

$$\text{tr}^-(f) = \text{tr}^-(g) \quad \text{and} \quad \text{tr}^+(f) = \text{tr}^+(g).$$

Proof. Set $h = f - g$, then $\lim_{|x| \rightarrow \infty} h(x) = 0$. We split the relevant averages as

$$\frac{1}{|B_r(y)|} \int_{B_r(y)} |h(x)| dx = \frac{1}{|B_r(y)|} \left[\int_{B_r(y) \cap B_{R'}^c(0)} + \int_{B_r(y) \cap B_{R'}(0)} \right] |h(x)| dx = (I) + (II).$$

Given $\varepsilon > 0$, there exists an $R_\varepsilon > 0$ such that $\sup_{|x| \geq R_\varepsilon} |h(x)| < \varepsilon/2$. So,

$$(I) < \frac{|B_r(y) \cap B_{R_\varepsilon}^c(0)|}{|B_r(y)|} \frac{\varepsilon}{2} < \frac{\varepsilon}{2},$$

independent of y . For the second term observe that

$$(II) < \|h\|_\infty \frac{|B_r(y) \cap B_{R_\varepsilon}(0)|}{|B_r(y)|} \leq \|h\|_\infty \frac{|B_{R_\varepsilon}|}{|B_r(y)|} < \frac{\varepsilon}{2}$$

for $r > \left(\frac{2}{\varepsilon} \|h\|_\infty\right)^{1/d} R_\varepsilon$. Consequently,

$$\lim_{r \rightarrow \infty} \sup_{y \in \mathbb{R}^d} \frac{1}{|B_r(y)|} \int_{B_r(y)} |f - g| = 0.$$

It follows that

$$\text{tr}^-(f) \leq \liminf_{r \rightarrow \infty} \inf_{y \in \mathbb{R}^d} \frac{1}{|B_r(y)|} \int_{B_r(y)} g + \limsup_{r \rightarrow \infty} \sup_{y \in \mathbb{R}^d} \frac{1}{|B_r(y)|} \int_{B_r(y)} |f - g| = \text{tr}^-(g).$$

Interchanging f and g yields equality. The equality $\text{tr}^+(f) = \text{tr}^+(g)$ is proved in the same way. \square

Appendix B: The lower bound for the reproducing kernel

We verify the lower bound for $\|k_x\|_2$ in the proof of [Proposition 2.2](#). This fact is proved in [\[Coulhon et al. 2012, Lemma 3.19\(a\)\]](#). For completeness we reproduce that proof with some necessary modifications and adjustments. The idea is to relate the reproducing kernel to the heat kernel of e^{-tH_a} via functional calculus.

We write k^Ω for the reproducing kernel of $\text{PW}_\Omega(H_a)$. For a bounded, nonnegative Borel function $F \geq 0$ with support in $[0, \Omega]$ we define $k_x^F = F(H_a)k_x^\Omega$ and the corresponding integral kernel $k^F(x, y) := F(H_a)(x, y) := k_x^F(y)$. The last expression is well-defined, as $k_x^F \in \text{PW}_\Omega(H_a)$. The kernel $k^F(x, y)$ is symmetric, because $F(H_a)$ is self-adjoint:

$$k_x^F(y) = \langle F(H_a)k_x^\Omega, k_y^\Omega \rangle = \langle k_x^\Omega, F(H_a)k_y^\Omega \rangle = \overline{\langle F(H_a)k_y^\Omega, k_x^\Omega \rangle} = \overline{k_y^F(x)}.$$

Consequently, $F(H_a)$ is an integral operator. For $f \in L^2(\mathbb{R}^d)$

$$F(H_a)f(x) = \langle F(H_a)f, k_x^\Omega \rangle = \langle f, F(H_a)k_x^\Omega \rangle = \langle f, k_x^F \rangle = \int_{\mathbb{R}^d} k^F(y, x) f(y) dy.$$

If $0 \leq G \leq F$, then

$$0 \leq k^G(x, x) \leq k^F(x, x) \tag{B-1}$$

for all $x \in \mathbb{R}^d$. For the proof observe that $F - G \geq 0$ implies that

$$k^F(x, x) = \langle F(H_a)k_x^\Omega, k_x^\Omega \rangle \geq \langle G(H_a)k_x^\Omega, k_x^\Omega \rangle = k^G(x, x).$$

The heat operator e^{-tH_a} is bounded and has a kernel $p_t(x, y)$ that satisfies *on diagonal estimates*. There are positive constants c, C such that for all $x \in \mathbb{R}^d$ and $t > 0$

$$ct^{-d/2} \leq p_t(x, x) \leq Ct^{-d/2}.$$

This is well known; see, e.g., [Ouhabaz 2006].

Claim. We have $0 < c < k^\Omega(x, x) < C$ for all $x \in \mathbb{R}^d$.

Proof. As $\chi_\Omega(u) \leq e \cdot e^{-u/\Omega} \chi_{[0, \infty)}(u)$ and $\chi_{[0, \infty)}(H_a) = \text{Id}$, we obtain

$$k^\Omega(x, x) = \chi_{[0, \Omega]}(H_a)(x, x) \leq ee^{-\Omega^{-1}H_a}(x, x) \leq C\Omega^{d/2}, \quad (\text{B-2})$$

which gives an explicit upper bound for $\|k_x\|_2^2$ in Proposition 2.2.

For the proof of the lower bound, we use a dyadic decomposition:

$$\begin{aligned} \chi_{[0, T]}(u)e^{-tu} &\leq \chi_{[0, \infty)}(u)e^{-tu} = \chi_{[0, \Omega]}(u)e^{-tu} + \sum_{k \geq 0} \chi_{[2^k \Omega, 2^{k+1} \Omega]}(u)e^{-tu} \\ &\leq \chi_{[0, \Omega]}(u) + \sum_{k \geq 0} \chi_{[0, 2^{k+1} \Omega]}(u)e^{-t2^k \Omega}, \quad t > 0. \end{aligned}$$

One can verify that this inequality remains true as an operator inequality

$$\chi_{[0, T]}(H_a)e^{-tH_a} \leq \chi_{[0, \Omega]}(H_a) + \sum_{k \geq 0} \chi_{[0, 2^{k+1} \Omega]}(H_a)e^{-t2^k \Omega},$$

with strong convergence of the sum, and every term is an integral operator. By (B-1) and (B-2) the operator inequality can be transferred to a corresponding inequality of the diagonals of the integral kernel as follows:

$$\begin{aligned} (\chi_{[0, T]}(H_a)e^{-tH_a})(x, x) &\leq \chi_{[0, \Omega]}(H_a)(x, x) + \sum_{k \geq 0} \chi_{[0, 2^{k+1} \Omega]}(H_a)(x, x)e^{-t2^k \Omega} \\ &\leq \chi_{[0, \Omega]}(H_a)(x, x) + C\Omega^{d/2} \sum_{k \geq 0} 2^{(k+1)d/2} e^{-t2^k \Omega}. \end{aligned}$$

In [Coulhon et al. 2012, equation (3.46)] it is shown that $p_t(x, y) = \lim_{T \rightarrow \infty} (\chi_{[0, T]}(H_a)e^{-tH_a})(x, y)$, consequently

$$ct^{-d/2} \leq p_t(x, x) \leq \chi_{[0, \Omega]}(H_a)(x, x) + C\Omega^{d/2} \sum_{k \geq 0} 2^{(k+1)d/2} e^{-t2^k \Omega}.$$

We choose $t = 2^r / \Omega$ for $r \in \mathbb{N}$ to be specified later. Then

$$\begin{aligned} c\Omega^{d/2}2^{-rd/2} &\leq \chi_{[0, \Omega]}(H_a)(x, x) + C\Omega^{d/2} \sum_{k \geq 0} e^{-2^k 2^r} 2^{(k+1)d/2} \\ &= \chi_{[0, \Omega]}(H_a)(x, x) + C2^{d/2}\Omega^{d/2}2^{-rd/2} \sum_{k \geq 0} e^{-2^{k+r}} 2^{(k+r)d/2} \\ &\leq \chi_{[0, \Omega]}(H_a)(x, x) + C2^{d/2}\Omega^{d/2}2^{-rd/2} \sum_{k \geq r} e^{-2^k} 2^{kd/2}. \end{aligned}$$

Hence,

$$\Omega^{d/2} 2^{-rd/2} \left(c - C' 2^{d/2} \sum_{k \geq r} e^{-2^k} 2^{kd/2} \right) \leq \chi_{[0, \Omega]}(H_a)(x, x) = k^\Omega(x, x).$$

For $r \in \mathbb{N}$ sufficiently large, this implies the lower bound for $k^\Omega(x, x)$. \square

References

- [Agmon 1965] S. Agmon, *Lectures on elliptic boundary value problems*, Van Nostrand Math. Stud. **2**, Van Nostrand, Princeton, NJ, 1965. [MR](#) [Zbl](#)
- [Aronszajn 1950] N. Aronszajn, “Theory of reproducing kernels”, *Trans. Amer. Math. Soc.* **68** (1950), 337–404. [MR](#) [Zbl](#)
- [Beurling 1966] A. Beurling, “Local harmonic analysis with some applications to differential operators”, pp. 109–125 in *Some recent advances in the basic sciences, I* (New York, 1962–1964), edited by A. Gelbart, Yeshiva Univ., New York, 1966. [MR](#)
- [Beurling 1989] A. Beurling, *The collected works of Arne Beurling, II: Harmonic analysis*, Birkhäuser, Boston, 1989. [MR](#) [Zbl](#)
- [Cordes 1979] H. O. Cordes, *Elliptic pseudo-differential operators: an abstract theory*, Lecture Notes in Math. **756**, Springer, 1979. [MR](#) [Zbl](#)
- [Coulhon et al. 2012] T. Coulhon, G. Kerkycharian, and P. Petrushev, “Heat kernel generated frames in the setting of Dirichlet spaces”, *J. Fourier Anal. Appl.* **18**:5 (2012), 995–1066. [MR](#) [Zbl](#)
- [Davies and Georgescu 2013] E. B. Davies and V. Georgescu, “ C^* -algebras associated with some second order differential operators”, *J. Operator Theory* **70**:2 (2013), 437–450. [MR](#) [Zbl](#)
- [DLMF 2020] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain (editors), “NIST digital library of mathematical functions”, electronic reference, Nat. Inst. Standards Tech., 2020, available at <http://dlmf.nist.gov>. Release 1.0.28.
- [Engelking 1977] R. Engelking, *General topology*, Monografie Mat. **60**, PWN, Warsaw, 1977. [MR](#) [Zbl](#)
- [Feichtinger 1984] H. G. Feichtinger, “Compactness in translation invariant Banach spaces of distributions and compact multipliers”, *J. Math. Anal. Appl.* **102**:2 (1984), 289–327. [MR](#) [Zbl](#)
- [Feichtinger and Pesenson 2004] H. Feichtinger and I. Pesenson, “Recovery of band-limited functions on manifolds by an iterative algorithm”, pp. 137–152 in *Wavelets, frames and operator theory* (College Park, MD, 2003), edited by C. Heil et al., Contemp. Math. **345**, Amer. Math. Soc., Providence, RI, 2004. [MR](#) [Zbl](#)
- [Feichtinger et al. 2016] H. G. Feichtinger, H. Führ, and I. Z. Pesenson, “Geometric space-frequency analysis on manifolds”, *J. Fourier Anal. Appl.* **22**:6 (2016), 1294–1355. [MR](#) [Zbl](#)
- [Filbir and Mhaskar 2011] F. Filbir and H. N. Mhaskar, “Marcinkiewicz–Zygmund measures on manifolds”, *J. Complexity* **27**:6 (2011), 568–596. [MR](#) [Zbl](#)
- [Führ et al. 2017] H. Führ, K. Gröchenig, A. Haimi, A. Klotz, and J. L. Romero, “Density of sampling and interpolation in reproducing kernel Hilbert spaces”, *J. Lond. Math. Soc.* (2) **96**:3 (2017), 663–686. [MR](#) [Zbl](#)
- [Gamelin 1969] T. W. Gamelin, *Uniform algebras*, Prentice-Hall, Englewood Cliffs, NJ, 1969. [MR](#) [Zbl](#)
- [Georgescu 2011] V. Georgescu, “On the structure of the essential spectrum of elliptic operators on metric spaces”, *J. Funct. Anal.* **260**:6 (2011), 1734–1765. [MR](#) [Zbl](#)
- [Georgescu 2018] V. Georgescu, “On the essential spectrum of elliptic differential operators”, *J. Math. Anal. Appl.* **468**:2 (2018), 839–864. [MR](#) [Zbl](#)
- [Grafakos 2004] L. Grafakos, *Classical and modern Fourier analysis*, Pearson, Upper Saddle River, NJ, 2004. [MR](#) [Zbl](#)
- [Gröchenig and Klotz 2010] K. Gröchenig and A. Klotz, “Noncommutative approximation: inverse-closed subalgebras and off-diagonal decay of matrices”, *Constr. Approx.* **32**:3 (2010), 429–466. [MR](#) [Zbl](#)
- [Gröchenig and Klotz 2017] K. Gröchenig and A. Klotz, “What is variable bandwidth?”, *Comm. Pure Appl. Math.* **70**:11 (2017), 2039–2083. [MR](#) [Zbl](#)
- [Gröchenig et al. 2019] K. Gröchenig, A. Haimi, J. Ortega-Cerdà, and J. L. Romero, “Strict density inequalities for sampling and interpolation in weighted spaces of holomorphic functions”, *J. Funct. Anal.* **277**:12 (2019), art. id. 108282. [MR](#) [Zbl](#)

- [Heil 2011] C. Heil, *A basis theory primer*, expanded ed., Birkhäuser, New York, 2011. [MR](#) [Zbl](#)
- [Hörmander 1968] L. Hörmander, “The spectral function of an elliptic operator”, *Acta Math.* **121** (1968), 193–218. [MR](#) [Zbl](#)
- [Kahane 1962] J.-P. Kahane, “Pseudo-périodicité et séries de Fourier lacunaires”, *Ann. Sci. École Norm. Sup.* (3) **79** (1962), 93–150. [MR](#) [Zbl](#)
- [Landau 1967] H. J. Landau, “Necessary density conditions for sampling and interpolation of certain entire functions”, *Acta Math.* **117** (1967), 37–52. [MR](#)
- [Matei and Meyer 2010] B. Matei and Y. Meyer, “Simple quasicrystals are sets of stable sampling”, *Complex Var. Elliptic Equ.* **55**:8-10 (2010), 947–964. [MR](#) [Zbl](#)
- [Olevskiĭ and Ulanovskii 2008] A. Olevskiĭ and A. Ulanovskii, “Universal sampling and interpolation of band-limited signals”, *Geom. Funct. Anal.* **18**:3 (2008), 1029–1052. [MR](#) [Zbl](#)
- [Ortega-Cerdà and Pridhnani 2012] J. Ortega-Cerdà and B. Pridhnani, “Beurling–Landau’s density on compact manifolds”, *J. Funct. Anal.* **263**:7 (2012), 2102–2140. [MR](#) [Zbl](#)
- [Ouhabaz 2006] E. M. Ouhabaz, “Sharp Gaussian bounds and L^p -growth of semigroups associated with elliptic and Schrödinger operators”, *Proc. Amer. Math. Soc.* **134**:12 (2006), 3567–3575. [MR](#) [Zbl](#)
- [Pesenson 1998] I. Pesenson, “Sampling of Paley–Wiener functions on stratified groups”, *J. Fourier Anal. Appl.* **4**:3 (1998), 271–281. [MR](#) [Zbl](#)
- [Pesenson 1999] I. Pesenson, “A reconstruction formula for band limited functions in $L_2(\mathbb{R}^d)$ ”, *Proc. Amer. Math. Soc.* **127**:12 (1999), 3593–3600. [MR](#) [Zbl](#)
- [Pesenson 2000] I. Pesenson, “A sampling theorem on homogeneous manifolds”, *Trans. Amer. Math. Soc.* **352**:9 (2000), 4257–4269. [MR](#) [Zbl](#)
- [Pesenson 2001] I. Pesenson, “Sampling of band-limited vectors”, *J. Fourier Anal. Appl.* **7**:1 (2001), 93–100. [MR](#) [Zbl](#)
- [Pesenson and Zayed 2009] I. Pesenson and A. I. Zayed, “Paley–Wiener subspace of vectors in a Hilbert space with applications to integral transforms”, *J. Math. Anal. Appl.* **353**:2 (2009), 566–582. [MR](#) [Zbl](#)
- [Rabinovich et al. 2004a] V. Rabinovich, S. Roch, and B. Silbermann, *Limit operators and their applications in operator theory*, Oper. Theory Adv. Appl. **150**, Birkhäuser, Basel, 2004. [MR](#) [Zbl](#)
- [Rabinovich et al. 2004b] V. S. Rabinovich, S. Roch, and J. Roe, “Fredholm indices of band-dominated operators”, *Integral Equations Operator Theory* **49**:2 (2004), 221–238. [MR](#) [Zbl](#)
- [Roe 2003] J. Roe, *Lectures on coarse geometry*, Univ. Lect. Ser. **31**, Amer. Math. Soc., Providence, RI, 2003. [MR](#) [Zbl](#)
- [Rudin 1973] W. Rudin, *Functional analysis*, McGraw-Hill, New York, 1973. [MR](#) [Zbl](#)
- [Seip 2004] K. Seip, *Interpolation and sampling in spaces of analytic functions*, Univ. Lect. Ser. **33**, Amer. Math. Soc., Providence, RI, 2004. [MR](#) [Zbl](#)
- [Shannon 1948] C. E. Shannon, “A mathematical theory of communication”, *Bell System Tech. J.* **27** (1948), 379–423, 623–656. [MR](#) [Zbl](#)
- [Shteinberg 2000] B. Y. Shteinberg, “Compactification of a locally compact group and the Noethericity of convolution operators with coefficients on quotient groups”, pp. 209–223 in *Proceedings of the St. Petersburg Mathematical Society, VI*, edited by N. N. Uraltseva, Amer. Math. Soc. Transl. Ser. 2 **199**, Amer. Math. Soc., Providence, RI, 2000. [MR](#) [Zbl](#)
- [Shubin 1992] M. A. Shubin, “Spectral theory of elliptic operators on noncompact manifolds”, pp. 35–108 in *Méthodes semi-classiques, I* (Nantes, France, 1991), Astérisque **207**, Soc. Math. France, Paris, 1992. [MR](#) [Zbl](#)
- [Špakula and Willett 2017] J. Špakula and R. Willett, “A metric approach to limit operators”, *Trans. Amer. Math. Soc.* **369**:1 (2017), 263–308. [MR](#) [Zbl](#)
- [Teschl 2009] G. Teschl, *Mathematical methods in quantum mechanics: with applications to Schrödinger operators*, Grad. Stud. Math. **99**, Amer. Math. Soc., Providence, RI, 2009. [MR](#) [Zbl](#)
- [Wendland 2005] H. Wendland, *Scattered data approximation*, Cambridge Monogr. Appl. Computat. Math. **17**, Cambridge Univ. Press, 2005. [MR](#) [Zbl](#)
- [Zimmer 1990] R. J. Zimmer, *Essential results of functional analysis*, Univ. Chicago Press, 1990. [MR](#) [Zbl](#)

Received 20 Sep 2021. Revised 7 Jun 2022. Accepted 11 Jul 2022.

KARLHEINZ GRÖCHENIG: karlheinz.groechenig@univie.ac.at

Faculty of Mathematics, University of Vienna, Vienna, Austria

ANDREAS KLOTZ: andreas.klotz@univie.ac.at

Faculty of Mathematics, University of Vienna, Vienna, Austria

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 2 2024

On a spatially inhomogeneous nonlinear Fokker–Planck equation: Cauchy problem and diffusion asymptotics	379
FRANCESCA ANCESCHI and YUZHE ZHU	
Strichartz inequalities with white noise potential on compact surfaces	421
ANTOINE MOUZARD and IMMANUEL ZACHHUBER	
Curvewise characterizations of minimal upper gradients and the construction of a Sobolev differential	455
SYLVESTER ERIKSSON-BIQUE and ELEFTERIOS SOULTANIS	
Smooth extensions for inertial manifolds of semilinear parabolic equations	499
ANNA KOSTIANKO and SERGEY ZELIK	
Semiclassical eigenvalue estimates under magnetic steps	535
WAFAA ASSAAD, BERNARD HELFFER and AYMAN KACHMAR	
Necessary density conditions for sampling and interpolation in spectral subspaces of elliptic differential operators	587
KARLHEINZ GRÖCHENIG and ANDREAS KLOTZ	
On blowup for the supercritical quadratic wave equation	617
ELEK CSOBO, IRFAN GLOGIĆ and BIRGIT SCHÖRKHUBER	
Arnold’s variational principle and its application to the stability of planar vortices	681
THIERRY GALLAY and VLADIMÍR ŠVERÁK	
Explicit formula of radiation fields of free waves with applications on channel of energy	723
LIANG LI, RUIPENG SHEN and LIJUAN WEI	
On L^∞ estimates for Monge–Ampère and Hessian equations on nef classes	749
BIN GUO, DUONG H. PHONG, FREID TONG and CHUWEN WANG	