

ANALYSIS & PDE

Volume 17

No. 3

2024

HAÏM BREZIS, ANDREAS SEEGER,
JEAN VAN SCHAFTINGEN AND PO-LAM YUNG

FAMILIES OF FUNCTIONALS REPRESENTING SOBOLEV NORMS



FAMILIES OF FUNCTIONALS REPRESENTING SOBOLEV NORMS

HAÏM BREZIS, ANDREAS SEEGER, JEAN VAN SCHAFTINGEN AND PO-LAM YUNG

We obtain new characterizations of the Sobolev spaces $\dot{W}^{1,p}(\mathbb{R}^N)$ and the bounded variation space $\dot{BV}(\mathbb{R}^N)$. The characterizations are in terms of the functionals $v_\gamma(E_{\lambda,\gamma/p}[u])$, where

$$E_{\lambda,\gamma/p}[u] = \left\{ (x, y) \in \mathbb{R}^N \times \mathbb{R}^N : x \neq y, \frac{|u(x) - u(y)|}{|x - y|^{1+\gamma/p}} > \lambda \right\}$$

and the measure v_γ is given by $dv_\gamma(x, y) = |x - y|^{\gamma-N} dx dy$. We provide characterizations which involve the $L^{p,\infty}$ -quasinorms $\sup_{\lambda>0} \lambda v_\gamma(E_{\lambda,\gamma/p}[u])^{1/p}$ and also exact formulas via corresponding limit functionals, with the limit for $\lambda \rightarrow \infty$ when $\gamma > 0$ and the limit for $\lambda \rightarrow 0^+$ when $\gamma < 0$. The results unify and substantially extend previous work by Nguyen and by Brezis, Van Schaftingen and Yung. For $p > 1$ the characterizations hold for all $\gamma \neq 0$. For $p = 1$ the upper bounds for the $L^{1,\infty}$ quasinorms fail in the range $\gamma \in [-1, 0)$; moreover, in this case the limit functionals represent the L^1 norm of the gradient for C_c^∞ -functions but not for generic $\dot{W}^{1,1}$ -functions. For this situation we provide new counterexamples which are built on self-similar sets of dimension $\gamma + 1$. For $\gamma = 0$ the characterizations of Sobolev spaces fail; however, we obtain a new formula for the Lipschitz norm via the expressions $v_0(E_{\lambda,0}[u])$.

1. Introduction

We are concerned with various ways in which we can recover the Sobolev seminorm $\|\nabla u\|_{L^p(\mathbb{R}^N)}$ via positive nonconvex functionals involving differences $u(x) - u(y)$.

We begin by mentioning two relevant results already in the literature. A theorem of H.-M. Nguyen [2006] (see also [Brezis and Nguyen 2018; 2020]) states that, for $1 < p < \infty$ and u in the inhomogeneous Sobolev space $W^{1,p}(\mathbb{R}^N)$,

$$\lim_{\lambda \searrow 0} \lambda^p \iint_{|u(x)-u(y)|>\lambda} |x - y|^{-p-N} dx dy = \frac{\kappa(p, N)}{p} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p, \quad (1-1)$$

with

$$\kappa(p, N) := \int_{\mathbb{S}^{N-1}} |e \cdot \omega|^p d\omega = \frac{2\Gamma((p+1)/2)\pi^{(N-1)/2}}{\Gamma((N+p)/2)}, \quad (1-2)$$

and e is any unit vector in \mathbb{R}^N . As shown in [Brezis and Nguyen 2018], (1-1) still holds for all $u \in C_c^1(\mathbb{R}^N)$ when $p = 1$ but fails for general $u \in W^{1,1}(\mathbb{R}^N)$. The limit formula (1-1) may be compared to a theorem of [Brezis et al. 2021b], which states that, for all $u \in C_c^\infty(\mathbb{R}^N)$ and $1 \leq p < \infty$, one has

$$\lim_{\lambda \rightarrow \infty} \lambda^p \mathcal{L}^{2N}(\{(x, y) \in \mathbb{R}^N \times \mathbb{R}^N : |u(x) - u(y)| > \lambda|x - y|^{1+N/p}\}) = \frac{\kappa(p, N)}{N} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p, \quad (1-3)$$

MSC2020: primary 26D10, 26A33; secondary 35A23, 42B25, 42B35, 46E30, 46E35.

Keywords: Sobolev norms, nonconvex functionals, nonlocal functionals, Marcinkiewicz spaces, Cantor sets and functions.

where \mathcal{L}^{2N} denotes the Lebesgue measure on $\mathbb{R}^N \times \mathbb{R}^N$. Our first result, namely [Theorem 1.1](#) below, provides an extension of [\(1-1\)](#) and [\(1-3\)](#) that unifies the two statements. Before we state the theorem, we introduce some notation that will be used throughout the paper.

First, for Lebesgue measurable subsets E of $\mathbb{R}^{2N} = \mathbb{R}^N \times \mathbb{R}^N$ and $\gamma \in \mathbb{R}$, we define

$$v_\gamma(E) := \iint_{\substack{(x,y) \in E \\ x \neq y}} |x - y|^{\gamma - N} \, dx \, dy. \tag{1-4}$$

In particular, when $\gamma = N$, v_N is just the Lebesgue measure on \mathbb{R}^{2N} . If u is a measurable function on \mathbb{R}^N and $b \in \mathbb{R}$, we define, for $(x, y) \in \mathbb{R}^N \times \mathbb{R}^N$ with $x \neq y$, a difference quotient

$$Q_b u(x, y) := \frac{u(x) - u(y)}{|x - y|^{1+b}}; \tag{1-5}$$

moreover, we define, for $\lambda > 0$, the superlevel set of $Q_b u$ at height λ by

$$E_{\lambda,b}[u] := \{(x, y) \in \mathbb{R}^N \times \mathbb{R}^N : x \neq y, |Q_b u(x, y)| > \lambda\}. \tag{1-6}$$

We will denote by $\dot{W}^{1,p}(\mathbb{R}^N)$, $p \geq 1$, the homogeneous Sobolev space, i.e., the space of $L^1_{\text{loc}}(\mathbb{R}^N)$ functions for which the distributional gradient ∇u belongs to $L^p(\mathbb{R}^N)$, with the seminorm $\|u\|_{\dot{W}^{1,p}} := \|\nabla u\|_{L^p(\mathbb{R}^N)}$. The inhomogeneous Sobolev space $W^{1,p}$ is the subspace of $\dot{W}^{1,p}$ -functions u for which $u \in L^p$, and we set $\|u\|_{W^{1,p}} := \|u\|_{L^p} + \|\nabla u\|_{L^p}$. For $p = 1$ we will also consider the space $\dot{\text{BV}}(\mathbb{R}^N)$ of functions of bounded variations, i.e., locally integrable functions u for which the gradient $\nabla u \in \mathcal{M}$ belongs to the space \mathcal{M} of \mathbb{R}^N -valued bounded Borel measures and we put $\|u\|_{\dot{\text{BV}}} := \|\nabla u\|_{\mathcal{M}}$; furthermore, let $\text{BV} := \dot{\text{BV}} \cap L^1$. In the dual formulation, with C^1_c denoting the space of C^1 functions with compact support,

$$\|u\|_{\dot{\text{BV}}} := \sup \left\{ \left| \int_{\mathbb{R}^N} u \operatorname{div}(\phi) \right| : \phi \in C^1_c(\mathbb{R}^N, \mathbb{R}^N), \|\phi\|_\infty \leq 1 \right\}.$$

For general background material on Sobolev spaces, see [\[Brezis 2011; Stein 1970\]](#).

Theorem 1.1. *Suppose $N \geq 1$, $1 \leq p < \infty$, $\gamma \in \mathbb{R} \setminus \{0\}$.*

(a) *If $\gamma > 0$, then, for all $u \in \dot{W}^{1,p}(\mathbb{R}^N)$,*

$$\lim_{\lambda \rightarrow +\infty} \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u]) = \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{1-7}$$

(b) *If either $\gamma < 0$, $p > 1$ or $\gamma < -1$, $p = 1$ then, for all $u \in \dot{W}^{1,p}(\mathbb{R}^N)$,*

$$\lim_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u]) = \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{1-8}$$

(c) *If $p = 1$ and $-1 \leq \gamma < 0$ then [\(1-8\)](#) remains true for all $u \in C^1_c(\mathbb{R}^N)$ but fails for generic $u \in \dot{W}^{1,1}(\mathbb{R}^N)$. However, we still have, for all $u \in \dot{W}^{1,1}(\mathbb{R}^N)$,*

$$\liminf_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda,\gamma}[u]) \geq \frac{\kappa(1, N)}{|\gamma|} \|\nabla u\|_{L^1(\mathbb{R}^N)}. \tag{1-9}$$

Formula (1-1) is the special case of (1-8) with $\gamma = -p$, and formula (1-3) is the special case of (1-7) with $\gamma = N$. Note that our result concerns functions in the homogeneous Sobolev space $\dot{W}^{1,p}$; we do not require u to be in L^p .

Remarks. (i) The reader will note the resemblance of (1-8) and (1-7) and may wonder why in (1-8), for $\gamma < 0$, one is concerned with the limit as $\lambda \searrow 0$ and in (1-7), for $\gamma > 0$, one takes the limit as $\lambda \rightarrow \infty$. In the proofs of these formulas one relates limits involving $\lambda v_\gamma(E_{\lambda,\gamma/p}[u])^{1/p}$ to (the absolute value of) limits of directional difference quotients $\delta^{-1}(u(x + \delta\theta) - u(x))$ with increment $\delta = \lambda^{-p/\gamma}$, and in order to recover the directional derivative $\langle \theta, \nabla u(x) \rangle$ we need to let $\delta \rightarrow 0$, which suggests that we need to take $\lambda \rightarrow \infty$ or $\lambda \searrow 0$ depending on the sign of γ . For the calculations see the proofs of Lemmas 3.2 and 3.3 below.

(ii) The failure of (1-8) for $p = 1$, $\gamma \in [-1, 0)$ and $u \in \dot{W}^{1,1}(\mathbb{R}^N)$ is generic in the sense of Baire category. It may happen that $\lim_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \infty$. This phenomenon was originally revealed when $\gamma = -1$ by A. Ponce and is presented in [Nguyen 2006]; see also [Brezis and Nguyen 2018, Pathology 1]. For stronger statements and more information, see Theorem 1.8. For $\gamma \in (-1, 0)$ we provide new examples based on self-similarity considerations. For discussion of failure in the case $\gamma = 0$, see Theorem 1.5 below. The special case of (1-9) for $\gamma = -1$ was already established in [Brezis and Nguyen 2018, Proposition 1].

When $p = 1$ we can also consider what happens if one allows functions in $\dot{B}\mathcal{V}(\mathbb{R}^N)$ in (1-7) and (1-8). For $\gamma = N$ in particular Poliakovsky [2022] asked whether the limit formulas remain valid in this generality (with $\|\nabla u\|_{L^1}$ replaced by $\|\nabla u\|_{\mathcal{M}}$). We provide a negative answer:

Proposition 1.2. (i) *The analogues of the limiting formulas (1-7) for $\gamma > 0$, $p = 1$ and (1-8) for $\gamma < 0$, $p = 1$, with $\|\nabla u\|_{\mathcal{M}}$ on the right-hand side, fail for suitable $u \in \dot{B}\mathcal{V}$.*

(ii) *Specifically, let $\Omega \subset \mathbb{R}^N$ be a bounded convex domain with smooth boundary and let u be the characteristic function of Ω . The limits $\lim_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda,\gamma}[u])$ for $\gamma > 0$ and $\lim_{\lambda \rightarrow 0+} \lambda v_\gamma(E_{\lambda,\gamma}[u])$ for $\gamma < -1$ exist, but they are not equal to $|\gamma|^{-1} \kappa(1, N) \|\nabla u\|_{\mathcal{M}}$.*

For a more detailed discussion we refer to Section 3F. See also Section 7B for a discussion about some related open problems.

Motivated by [Brezis et al. 2021b], we will also be interested in what happens to the larger quantity obtained by replacing the limits on the left-hand sides of (1-7) and (1-8) by $\sup_{\lambda > 0}$. This will be formulated in terms of the Marcinkiewicz space $L^{p,\infty}(\mathbb{R}^{2N}, v_\gamma)$ (a.k.a. weak-type L^p) defined by the condition

$$[F]_{L^{p,\infty}(\mathbb{R}^{2N}, v_\gamma)}^p := \sup_{\lambda > 0} \lambda^p v_\gamma(\{(x, y) \in \mathbb{R}^N \times \mathbb{R}^N : |F(x, y)| > \lambda\}) < \infty. \tag{1-10}$$

As an immediate consequence of Theorem 1.1 we have, for $N \geq 1$, $1 \leq p < \infty$, $\gamma \neq 0$ and all $u \in C_c^\infty(\mathbb{R}^N)$,

$$[Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, v_\gamma)}^p \geq C(N, p, \gamma) \|\nabla u\|_{L^p(\mathbb{R}^N)}^p, \tag{1-11}$$

where $C(N, p, \gamma)$ is a positive constant depending only on N, p and γ . Moreover, the same conclusion holds for all $u \in \dot{W}^{1,p}(\mathbb{R}^N)$ when $p > 1$, with any $\gamma \neq 0$, and when $p = 1$, with any $\gamma \notin [-1, 0]$. We shall show that the conditions in the last statement can in fact be relaxed; see the inequalities (1-14) and

(1-16) below. In addition we have the important upper bounds for $Q_{\gamma/p}u$, extending the case $\gamma = N$ already dealt with in [Brezis et al. 2021b] for $u \in C_c^\infty(\mathbb{R}^N)$. The result in [Brezis et al. 2021b] states that, for every $N \geq 1$, there exists a constant $C(N)$ such that

$$[Q_{N/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_N)}^p \leq C(N) \|\nabla u\|_{L^p(\mathbb{R}^N)}^p \tag{1-12}$$

for all $u \in C_c^\infty(\mathbb{R}^N)$ and all $1 \leq p < \infty$. In light of Theorem 1.1, it is natural to ask whether one can replace the limits on the left-hand sides of (1-7) and (1-8) by $\sup_{\lambda>0}$ and still obtain a quantity that is comparable to $\|\nabla u\|_{L^p(\mathbb{R}^N)}^p$. As suggested by Theorem 1.1 the answer to our question is sensitive to the values of γ and p .

Theorem 1.3. *Suppose that $N \geq 1$, $1 < p < \infty$ and $\gamma \in \mathbb{R}$. Then the following hold:*

(i) *The inequality*

$$[Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)} \leq C(N, p, \gamma) \|\nabla u\|_{L^p(\mathbb{R}^N)} \tag{1-13}$$

holds for all $u \in C_c^\infty(\mathbb{R}^N)$ if and only if $\gamma \neq 0$. In this case (1-13) extends to all $u \in \dot{W}^{1,p}(\mathbb{R}^N)$.

(ii) *Suppose that $u \in L^1_{\text{loc}}(\mathbb{R}^N)$ and $Q_{\gamma/p}u \in L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)$. Then $u \in \dot{W}^{1,p}(\mathbb{R}^N)$ and we have the inequality*

$$\|\nabla u\|_{L^p(\mathbb{R}^N)} \leq C_{N,p,\gamma} [Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}. \tag{1-14}$$

There is a new phenomenon for $p = 1$, namely the upper bounds for $Q_\gamma u$ only hold for the more restrictive range $\gamma \in (-\infty, -1) \cup (0, \infty)$. Here it is also natural to replace $\dot{W}^{1,1}$ with $\mathring{B}\mathring{V}$.

Theorem 1.4. *Suppose that $N \geq 1$ and $\gamma \in \mathbb{R}$. Then the following hold:*

(i) *The inequality*

$$[Q_\gamma u]_{L^{1,\infty}(\mathbb{R}^{2N}, \nu_\gamma)} \leq C(N, \gamma) \|\nabla u\|_{L^1(\mathbb{R}^N)} \tag{1-15}$$

holds for all $u \in C_c^\infty(\mathbb{R}^N)$ if and only if $\gamma \notin [-1, 0]$. In this case (1-15) extends to all $u \in \dot{W}^{1,1}(\mathbb{R}^N)$, and, if $\|\nabla u\|_{L^1(\mathbb{R}^N)}$ is replaced by $\|\nabla u\|_{\mathcal{M}}$, to all $u \in \mathring{B}\mathring{V}(\mathbb{R}^N)$.

(ii) *Suppose that $u \in L^1_{\text{loc}}(\mathbb{R}^N)$ and $Q_\gamma u \in L^{1,\infty}(\mathbb{R}^{2N}, \nu_\gamma)$. Then $u \in \mathring{B}\mathring{V}(\mathbb{R}^N)$ and we have the inequality*

$$\|\nabla u\|_{\mathcal{M}} \leq C_{N,\gamma} [Q_\gamma u]_{L^{1,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}. \tag{1-16}$$

We note that the quantitative bounds (1-13) and (1-15) in Theorems 1.3 and 1.4 are crucial tools for establishing the limiting relations for all $\dot{W}^{1,p}$ functions in Theorem 1.1. Note that there is no restriction on γ in (1-14) and (1-16). The constants in the inequalities will be quantified further later in the paper. In particular, $C(N, p, \gamma)$ in (1-13) remains bounded as $p \searrow 1$ only in the range $\gamma \in (0, \infty) \cup (-\infty, -1)$ (see Theorem 2.2 and Proposition 6.1).

Historical comments. Some special cases of the above quantitative estimates have been known. Estimate (1-13) for $\gamma = -p$ and $1 < p < \infty$ was discovered independently by H.-M. Nguyen [2006], and by A. Ponce and J. Van Schaftingen (unpublished communication to H. Brezis and H.-M. Nguyen), both relying on the Hardy–Littlewood maximal inequality. A. Poliakovsky [2022] recently proved generalizations of results in [Brezis et al. 2021b] to Sobolev spaces on domains; moreover, he obtained Theorems 1.3 and 1.4 in the special case $\gamma = N$ under the additional assumption that $u \in L^p$. Other far-reaching generalizations to one-parameter families of operators were obtained by Ó. Domínguez and M. Milman [2022].

The case $\gamma = 0$. We shall now return to the necessity of the assumption $\gamma \notin [-1, 0]$ in parts of Theorems 1.1, 1.3 and 1.4. When $\gamma = 0$, the bounds for $[Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}$ fail in a striking way. We begin by formulating a result illustrating this failure, which also gives a characterization of the seminorm in the Lipschitz space $\dot{W}^{1,\infty}$.

Theorem 1.5. *Suppose $N \geq 1$, u is locally integrable on \mathbb{R}^N and $\nabla u \in L^1_{\text{loc}}(\mathbb{R}^N)$. Then*

$$\|\nabla u\|_{L^\infty(\mathbb{R}^N)} = \inf\{\lambda > 0 : \nu_0(E_{\lambda,0}[u]) < \infty\}. \tag{1-17}$$

Indeed in Proposition 5.1 we shall prove the stronger statement that $\nu_0(E_{\lambda,0}[u]) = 0$ for $\lambda > \|\nabla u\|_\infty$, and $\nu_0(E_{\lambda,0}[u]) = \infty$ for $\lambda < \|\nabla u\|_\infty$. As an immediate consequence of Theorem 1.5 we get:

Corollary 1.6. *Let u be locally integrable on \mathbb{R}^N . If $\nabla u \in L^1_{\text{loc}}(\mathbb{R}^N)$ and if $\nu_0(E_{\lambda,0}[u])$ is finite for all $\lambda > 0$, then u is almost everywhere equal to a constant function.*

In view of other known results [Brezis 2002; Brezis et al. 2021a] on how to recognize constant functions, a natural question arises whether the hypothesis on the local integrability of ∇u in the corollary could be relaxed; one can ask whether the constancy conclusion holds for all locally integrable functions satisfying $\nu_0(E_{\lambda,0}[u]) < \infty$ for all $\lambda > 0$. However, the following example shows that such an extension fails (for details, see Lemma 5.2).

Example 1.7. Let $\Omega \subset \mathbb{R}^N$ be a bounded Lipschitz domain and let u be the characteristic function of Ω . Then $u \in \text{BV}(\mathbb{R}^N) \setminus \dot{W}^{1,1}(\mathbb{R}^N)$ and $\sup_{\lambda > 0} \lambda \nu_0(E_{\lambda,0}[u]) < \infty$.

More on counterexamples. We now make more explicit the exclusion of the parameters $\gamma \in [-1, 0]$ in part (c) of Theorem 1.1 and in (1-15). We shall show in Section 6B that for $\gamma \in (-1, 0)$ these negative results can be related to self-similar Cantor subsets of \mathbb{R} , of dimension $1 + \gamma$.

Theorem 1.8. *Suppose $N \geq 1$. Then the following hold:*

(i) *Let $-1 \leq \gamma < 0$. There exists a C^∞ function $u \in \dot{W}^{1,1}(\mathbb{R}^N)$, rapidly decreasing as $|x| \rightarrow \infty$ and such that*

$$\lim_{\lambda \searrow 0} \lambda \nu_\gamma(E_{\lambda,\gamma}[u]) = \infty. \tag{1-18}$$

(ii) *Let $-1 \leq \gamma < 0$. There exists a compactly supported $u \in W^{1,1}(\mathbb{R}^N)$ for which (1-18) holds. The set*

$$\{u \in W^{1,1}(\mathbb{R}^N) : \limsup_{\lambda \searrow 0} \lambda \nu_\gamma(E_{\lambda,\gamma}[u]) < \infty\}$$

is meager in $W^{1,1}(\mathbb{R}^N)$, i.e., of first category in the sense of Baire.

(iii) *Let $-1 \leq \gamma < 0$, $N \geq 2$ or $-1 < \gamma < 0$, $N = 1$. There exists a compactly supported $u \in W^{1,1}(\mathbb{R}^N)$ such that $\nu_\gamma(E_{\lambda,\gamma}[u]) = \infty$ for all $\lambda > 0$; moreover, the set*

$$\{u \in W^{1,1}(\mathbb{R}^N) : \nu_\gamma(E_{\lambda,\gamma}[u]) < \infty \text{ for some } \lambda \in (0, \infty)\}$$

is meager in $W^{1,1}(\mathbb{R}^N)$.

The case $N = 1 = -\gamma$ plays a special role and is excluded in the strongest statement (iii) since for all compactly supported $u \in \dot{W}^{1,1}(\mathbb{R})$ one has $v_{-1}(E_{\lambda,-1}[u]) < \infty$ for all $\lambda > 0$ (see Lemma 6.5 below). The proofs of existence of counterexamples are constructive and the Baire category statements will be obtained as rather straightforward consequences of the constructions.

Outline of the paper. In Section 2 we provide the upper bounds for $[Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, v_\gamma)}$, i.e., the proof of inequalities (1-13) and (1-15) in Theorems 1.3 and 1.4. We first derive these for a dense subclass, relying on covering lemmas, and then extend in Sections 2C and 2D to general $\dot{W}^{1,p}$ and $\dot{B}V$ -functions. In Section 3 we derive the limit formulas of Theorem 1.1; specifically in Section 3B we prove the sharp lower bounds involving a $\liminf \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u])$ for general functions in $\dot{W}^{1,p}$ and in Section 3C we obtain the sharp upper bounds for $\limsup \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u])$, under the assumption that $u \in C^1$ is compactly supported. Then in Section 3D we extend these limits to general $\dot{W}^{1,p}$ functions. In Section 3F we show that the limit formulas for $\dot{W}^{1,1}$ do not extend to general $\dot{B}V$ functions and prove Proposition 1.2. In Section 4 we prove the reverse inequalities (1-14) and (1-16) in Theorems 1.3 and 1.4. In Section 5 we prove Theorem 1.5 on a characterization of the Lipschitz norm and also discuss Example 1.7. In Section 6 we provide various constructions of counterexamples and in particular prove Theorem 1.8. We discuss some further perspectives and open problems in Section 7.

2. Bounding $[Q_{\gamma/p}u]_{L^{p,\infty}(\mathbb{R}^{2N}, v_\gamma)}$ by the Sobolev norm

In this section we prove inequalities (1-13) and (1-15) in Theorems 1.3 and 1.4.

2A. The bound (1-13) via the Hardy–Littlewood maximal operator. Following [Brezis et al. 2021b], one can prove the result of Theorem 1.3 for $p > 1$ by an elementary argument involving the Hardy–Littlewood maximal function $M|\nabla u|$ of $|\nabla u|$; however, the behavior of the constants as $p \searrow 1$ will only be sharp in the range $-1 \leq \gamma < 0$.

Proposition 2.1. *Let $N \geq 1$ and $1 < p < \infty$. There exists a constant C_N such that, for all $\gamma \neq 0$ and all $u \in \dot{W}^{1,p}(\mathbb{R}^N)$,*

$$\sup_{\lambda > 0} \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u]) \leq \frac{C_N}{|\gamma|} \left(\frac{p}{p-1} \right)^p \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{2-1}$$

Proof. We assume first that $u \in C^1$ and that ∇u is compactly supported. As in [Brezis et al. 2021b, Remark 2.3], one uses the Lusin–Lipschitz inequality

$$\frac{|u(x) - u(y)|}{|x - y|} \leq C[M(|\nabla u|)(x) + M(|\nabla u|)(y)] \tag{2-2}$$

and observes that (2-2) implies

$$E_{\lambda,\gamma/p}[u] \subseteq \{|x - y|^{\gamma/p} < 2C\lambda^{-1}M(|\nabla u|)(x)\} \cup \{|x - y|^{\gamma/p} < 2C\lambda^{-1}M(|\nabla u|)(y)\}.$$

As a consequence

$$v_\gamma(E_{\lambda,\gamma/p}[u]) \leq 2 \int_x \int_{|h|^\gamma < 2C[\lambda^{-1}M(|\nabla u|)(x)]^p} |h|^{\gamma-N} dh dx.$$

Direct computation of the inner integral (distinguishing the cases $\gamma > 0$ and $\gamma < 0$) yields

$$v_\gamma(E_{\lambda,\gamma/p}[u]) \lesssim_N C^p |\gamma|^{-1} \lambda^{-p} \int_{\mathbb{R}^N} [M(|\nabla u|)(x)]^p dx.$$

Inequality (2-1) follows then from the standard maximal inequality $\|Mf\|_p^p \leq [C(N)p']^p \|f\|_p^p$ for $p > 1$; see [Stein 1970] (here $p' = p/(p - 1)$). The extension to general $\dot{W}^{1,p}$ functions will be taken up in Section 2C. □

2B. The case $\gamma \in \mathbb{R} \setminus [-1, 0]$. We shall prove the following more precise versions of the estimates (1-13) and (1-15) when $\gamma \notin [-1, 0]$, with constants that stay bounded as $p \searrow 1$; indeed we cover all $p \in [1, \infty)$. We denote by σ_{N-1} the surface area of the sphere \mathbb{S}^{N-1} . In the proof of the following theorem we will first establish the estimates for functions $u \in C^1(\mathbb{R}^N)$ whose gradient is compactly supported. The extension to $\dot{W}^{1,p}$ and \mathbf{BV} will be taken up in Sections 2C and 2D.

Theorem 2.2. *There exists an absolute constant $C > 0$ such that, for every $N \geq 1$, every $1 \leq p < \infty$, and every $u \in \dot{W}^{1,p}(\mathbb{R}^N)$:*

(i) *If $\gamma > 0$, then*

$$\sup_{\lambda>0} \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u]) \leq C \sigma_{N-1} \frac{5^\gamma}{\gamma} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{2-3}$$

(ii) *If $\gamma < -1$, then*

$$\sup_{\lambda>0} \lambda^p v_\gamma(E_{\lambda,\gamma/p}[u]) \leq \frac{C \sigma_{N-1}}{|\gamma|} \left(1 + \frac{1}{|\gamma + 1|}\right) \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{2-4}$$

When $p = 1$ the above assertions hold for all $u \in \mathbf{BV}(\mathbb{R}^N)$ provided that $\|\nabla u\|_{L^1(\mathbb{R}^N)}$ is replaced by $\|\nabla u\|_{\mathcal{M}}$.

The proof of Theorem 2.2 relies on the following proposition, in which $[x, y] \subset \mathbb{R}^N$ denotes the closed line segment connecting two points $x, y \in \mathbb{R}^N$.

Proposition 2.3. *Let*

$$E(f, \gamma) := \left\{ (x, y) \in \mathbb{R}^N \times \mathbb{R}^N : x \neq y, \int_{[x,y]} |f| ds > |x - y|^{\gamma+1} \right\} \tag{2-5}$$

for $f \in C_c(\mathbb{R}^N)$. There exists an absolute constant $C > 0$ such that, for every $N \geq 1$ and every $f \in C_c(\mathbb{R}^N)$:

(i) *If $\gamma > 0$, then*

$$\iint_{E(f,\gamma)} |x - y|^{\gamma-N} dx dy \leq C \sigma_{N-1} \frac{5^\gamma}{\gamma} \|f\|_{L^1(\mathbb{R}^N)}. \tag{2-6}$$

(ii) *If $\gamma < -1$, then*

$$\iint_{E(f,\gamma)} |x - y|^{\gamma-N} dx dy \leq \frac{C \sigma_{N-1}}{|\gamma|} \left(1 + \frac{1}{|\gamma + 1|}\right) \|f\|_{L^1(\mathbb{R}^N)}. \tag{2-7}$$

Indeed, to deduce Theorem 2.2 from Proposition 2.3 one argues as in the proof of (1-12) in [Brezis et al. 2021b]; for $u \in C^1(\mathbb{R}^N)$ and $1 \leq p < \infty$, one has

$$|u(x) - u(y)|^p \leq \left[\int_{[x,y]} |\nabla u| ds \right]^p \leq \int_{[x,y]} |\nabla u|^p ds |x - y|^{p-1}$$

for all $x, y \in \mathbb{R}^N$, which implies

$$E_{\lambda, \gamma/p}[u] \subseteq E(\lambda^{-p}|\nabla u|^p, \gamma).$$

Hence for $u \in C^1(\mathbb{R}^N)$ whose gradient is compactly supported, one establishes [Theorem 2.2](#) by applying [Proposition 2.3](#) with $f := \lambda^{-p}|\nabla u|^p$. The extension to $u \in \dot{W}^{1,p}$ will be taken up in [Section 2C](#).

Proof of Proposition 2.3. As in the proof of [[Brezis et al. 2021b](#), Proposition 2.2], using the method of rotation, we only need to prove [Proposition 2.3](#) for $N = 1$. Indeed,

$$\iint_{E(f, \gamma)} |x - y|^{\gamma - N} \, dx \, dy = \frac{1}{2} \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \iint_{E(f_{\omega, x'}, \gamma)} |r - s|^{\gamma - 1} \, dr \, ds \, dx' \, d\omega,$$

where for every $\omega \in \mathbb{S}^{N-1}$ and every $x' \in \omega^\perp$, $f_{\omega, x'}$ is a function of one real variable defined by

$$f_{\omega, x'}(t) := f(x' + t\omega).$$

The innermost double integral can be estimated by the case $N = 1$ of [Proposition 2.3](#), and

$$\int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \int_{\mathbb{R}} |f_{\omega, x'}(t)| \, dt \, dx' \, d\omega = \sigma_{N-1} \|f\|_{L^1(\mathbb{R}^N)}.$$

Thus from now on, we assume $N = 1$ and $f \in C_c(\mathbb{R})$.

If $\gamma > 0$, the desired estimate (2-6) is the content of [[Brezis et al. 2021b](#), Proposition 2.1]. On the other hand, suppose now $\gamma < -1$. Without loss of generality, assume $f \geq 0$ on \mathbb{R} . In addition, we may assume that f is not identically zero, for otherwise there is nothing to prove.

Let

$$E_+(f, \gamma) := \{(x, y) \in E(f, \gamma) : y < x\}.$$

Then by symmetry,

$$\iint_{E(f, \gamma)} |x - y|^{\gamma - 1} \, dx \, dy = 2 \iint_{E_+(f, \gamma)} |x - y|^{\gamma - 1} \, dx \, dy,$$

and it suffices to estimate the latter integral.

In what follows we will need to always keep in mind that in view of our assumption $\gamma < -1$ we have $-(\gamma + 1) = |\gamma| - 1 > 0$. We will now use a simple stopping-time argument based on the fact that for all $c \in \mathbb{R}$ the continuous function

$$x \mapsto (x - c)^{-(\gamma+1)} \int_c^x f(s) \, ds, \quad x \geq c,$$

increases from 0 to ∞ on $[c, \infty)$.

Assume that $\text{supp } f \subseteq [a, b]$. We construct a finite sequence of intervals I_1, \dots, I_K , that are disjoint up to endpoints, that cover $\text{supp } f = [a, b]$, and that satisfy

$$|I_i|^{-(\gamma+1)} \int_{I_i} f = \frac{1}{2} \quad \text{for } 1 \leq i \leq K. \tag{2-8}$$

Indeed, we may take $a_1 := a$, and $a_2 > a_1$ to be the unique number for which

$$(a_2 - a_1)^{-(\gamma+1)} \int_{a_1}^{a_2} f = \frac{1}{2},$$

and set $I_1 := [a_1, a_2]$. If $a_2 < b$, we may now repeat, and take $I_2 := [a_2, a_3]$, where $a_3 > a_2$ is the unique number for which $(a_3 - a_2)^{-(\gamma+1)} \int_{a_2}^{a_3} f = \frac{1}{2}$. Note that the a_i 's chosen as such satisfy

$$(a_{i+1} - a_i)^{-(\gamma+1)} \geq \frac{1}{2} \|f\|_{L^1(\mathbb{R})}^{-1},$$

so that $a_{i+1} - a_i \geq (2\|f\|_{L^1(\mathbb{R})})^{1/(\gamma+1)}$. This shows that in finitely many steps, we would reach $a_{K+1} \geq b$ for some $K \geq 1$, with $a_K < b$ if $1 \leq K$. Then we have our sequence of disjoint (up to endpoints) intervals I_1, \dots, I_K that cover $[a, b]$ and satisfy (2-8). We also write $I_0 := (-\infty, a_1]$ and $I_{K+1} := [a_{K+1}, +\infty)$.

We now claim that $I_i \times I_i \cap E_+(f, \gamma) = \emptyset$ for every $0 \leq i \leq K + 1$. This being trivially the case when $i \in \{0, K + 1\}$, we consider the case $i \in \{1, \dots, K\}$: any $x, y \in I_i$ satisfy

$$|x - y|^{-(\gamma+1)} \left| \int_y^x f \right| \leq |I_i|^{-(\gamma+1)} \int_{I_i} f = \frac{1}{2} < 1.$$

It follows thus that

$$E_+(f, \gamma) = \bigcup_{i=1}^{K+1} E_+(f, \gamma) \cap ((a_i, +\infty) \times (-\infty, a_i)). \tag{2-9}$$

Furthermore, for $i \in \{2, \dots, K\}$, if $y < a_i < x$ and $x - y < \min\{|I_i|, |I_{i-1}|\}$, then

$$\begin{aligned} |x - y|^{-(\gamma+1)} \left| \int_y^x f \right| &< \min\{|I_i|, |I_{i-1}|\}^{-(\gamma+1)} \left(\int_{I_{i-1}} f + \int_{I_i} f \right) \\ &\leq |I_{i-1}|^{-(\gamma+1)} \int_{I_{i-1}} f + |I_i|^{-(\gamma+1)} \int_{I_i} f \leq \frac{1}{2} + \frac{1}{2} = 1 \end{aligned}$$

(again we used $\gamma < -1$ so that $-(\gamma+1) > 0$ here), from which it follows that $(x, y) \notin E_+(f, \gamma)$. Combining this with a similar argument for $i \in \{1, K + 1\}$, we get that if $(x, y) \in E_+(f, \gamma) \cap (a_i, +\infty) \times (-\infty, a_i)$, then $|x - y| \geq \min\{|I_i|, |I_{i-1}|\}$, and thus

$$\begin{aligned} \int_{E_+(f, \gamma) \cap (a_i, +\infty) \times (-\infty, a_i)} |x - y|^{\gamma-1} dx dy &\leq \int_{a_i}^{\infty} \int_{-\infty}^{\min\{a_i, x - \min\{|I_i|, |I_{i-1}|\}\}} |x - y|^{\gamma-1} dy dx \\ &= \frac{1}{|\gamma|} \int_{a_i}^{\infty} (\max\{x - a_i, \min\{|I_i|, |I_{i-1}|\}\})^\gamma dx \\ &= \frac{1}{|\gamma|} \left(1 + \frac{1}{|\gamma + 1|} \right) \min\{|I_i|, |I_{i-1}|\}^{\gamma+1} \\ &\leq \frac{2}{|\gamma|} \left(1 + \frac{1}{|\gamma + 1|} \right) \int_{I_{i-1} \cup I_i} f. \end{aligned}$$

(The computation of these integrals uses our assumption $\gamma + 1 < 0$.) Summing the estimates, we get in view of (2-9)

$$\int_{E_+(f, \gamma)} |x - y|^{\gamma-1} dx dy \leq \frac{4}{|\gamma|} \left(1 + \frac{1}{|\gamma + 1|} \right) \int_{\mathbb{R}} f.$$

We have thus completed the proof of (2-7) under the assumption $\gamma < -1$ and $N = 1$. □

2C. Proof of Proposition 2.1 and Theorem 2.2 for general $\dot{W}^{1,p}$ functions. We use a limiting argument, together with the following fact: if $u \in \dot{W}^{1,p}(\mathbb{R}^N)$, $N \geq 1$, and $1 \leq p < \infty$, then there exists a Lebesgue measurable set $X \subset \mathbb{R}^{2N}$, with $\mathcal{L}^{2N}(X) = 0$, so that, for every $(x, h) \in \mathbb{R}^{2N} \setminus X$, we have

$$u(x+h) - u(x) = \int_0^1 \langle h, \nabla u(x+th) \rangle dt. \tag{2-10}$$

Indeed, both sides are measurable functions of $(x, h) \in \mathbb{R}^{2N}$, and if X is the set of all (x, h) where the two sides are not equal, then X is a measurable subset of \mathbb{R}^{2N} , and the assertion will follow from Fubini’s theorem if, for every fixed $h \in \mathbb{R}^N$, we have $\mathcal{L}^N(\{x \in \mathbb{R}^N : (x, h) \in X\}) = 0$, i.e., (2-10) holds for \mathcal{L}^N almost every x . This follows since for every $\phi \in C_c^\infty(\mathbb{R}^N)$, one has

$$\begin{aligned} \int_{\mathbb{R}^N} [u(x+h) - u(x)]\phi(x) dx &= \int_{\mathbb{R}^N} u(x)[\phi(x-h) - \phi(x)] dx = - \int_{\mathbb{R}^N} u(x) \int_0^1 \langle h, \nabla \phi(x-th) \rangle dt dx \\ &= \int_{\mathbb{R}^N} \int_0^1 \langle h, \nabla u(x) \rangle \phi(x-th) dt dx = \int_{\mathbb{R}^N} \int_0^1 \langle h, \nabla u(x+th) \rangle dt \phi(x) dx. \end{aligned}$$

Now given $u \in \dot{W}^{1,p}(\mathbb{R}^N)$, there exists a sequence $u_n \in C^\infty(\mathbb{R}^N)$ such that ∇u_n are compactly supported, and

$$\|\nabla(u_n - u)\|_{L^p(\mathbb{R}^N)} \rightarrow 0. \tag{2-11}$$

Indeed if $N > 1$ and $p \geq 1$, or if $N = 1$ and $p > 1$, then this follows from the density of $C_c^\infty(\mathbb{R}^N)$ in $\dot{W}^{1,p}(\mathbb{R}^N)$ as asserted in [Hajlasz and Kalamajska 1995] (in this case one may choose $u_n \in C_c^\infty(\mathbb{R}^N)$). The density of $C_c^\infty(\mathbb{R}^N)$ in $\dot{W}^{1,p}$ fails when $N = p = 1$ (again see [Hajlasz and Kalamajska 1995]); the issue is that if ∇u is supported in a convex set in \mathbb{R}^N , $N \geq 2$, then u is constant in the complement of the set, but this fails for $N = 1$ since the complement of a bounded interval has two connected components. On the other hand, in the anomalous case $N = 1$ and $p = 1$, one can choose an approximation of the identity to get a sequence v_n of C_c^∞ functions on \mathbb{R} such that $\|v_n - u'\|_{L^1(\mathbb{R})} \rightarrow 0$. One can then take $u_n(x) := \int_0^x v_n(t) dt$, and (2-11) follows with $u'_n = v_n$ being compactly supported (even though u_n may not be compactly supported).

Let, for $R > 1$,

$$K_R = \{(x, y) \in \mathbb{R}^{2N} : |x| \leq R, |y| \leq R \text{ and } R^{-1} \leq |x - y|\}.$$

By monotone convergence it suffices to prove

$$v_\gamma(E_{\lambda,\gamma/p}[u] \cap K_R) \leq C \frac{\|\nabla u\|_{L^p(\mathbb{R}^N)}^p}{\lambda^p}. \tag{2-12}$$

with C independent of R .

Under the assumptions of Proposition 2.1 and Theorem 2.2 on p and γ , since $u_n \in C_c^\infty(\mathbb{R}^N)$, we already know

$$v_\gamma(E_{\lambda,\gamma/p}[u_n]) \leq C \frac{\|\nabla u_n\|_{L^p(\mathbb{R}^N)}^p}{\lambda^p}.$$

Moreover, the sequence $Q_{\gamma/p}u_n$ converges to $Q_{\gamma/p}u$ in $L^p(K_R)$ as $n \rightarrow \infty$. Indeed, using (2-10) we may write

$$Q_{\gamma/p}u(x, y) = \frac{1}{|x - y|^{\gamma/p}} \int_0^1 \left\langle \frac{x - y}{|x - y|}, \nabla u((1 - t)y + tx) \right\rangle dt$$

for \mathcal{L}^{2N} a.e. $(x, y) \in \mathbb{R}^{2N}$, and similarly for u_n in place of u , which allows us to estimate

$$\begin{aligned} & \left(\iint_{K_R} |Q_{\gamma/p}u_n(x, y) - Q_{\gamma/p}u(x, y)|^p dx dy \right)^{1/p} \\ & \leq R^{\gamma/p} \int_0^1 \left(\int_{|x| \leq R} \int_{|y| \leq R} |\nabla(u_n - u)((1 - s)x + sy)|^p dx dy \right)^{1/p} ds \\ & \leq 2^{N/p} (2R)^{N/p} R^{\gamma/p} \|\nabla(u_n - u_{n+1})\|_p \rightarrow 0. \end{aligned}$$

By passing to a subsequence if necessary, we may assume that $Q_{\gamma/p}u_n$ converges \mathcal{L}^{2N} -a.e. to $Q_{\gamma/p}u$ on K_R as $n \rightarrow \infty$. Thus

$$K_R \cap E_{\lambda, \gamma/p}[u] \subseteq K_R \cap \left(\bigcup_{n \in \mathbb{N}} \bigcap_{\ell \geq n} E_{\lambda, \gamma/p}[u_\ell] \right),$$

which implies

$$\begin{aligned} v_\gamma(K_R \cap E_{\lambda, \gamma/p}[u]) & \leq \lim_{n \rightarrow \infty} v_\gamma \left(K_R \cap \bigcap_{\ell \geq n} E_{\lambda, \gamma/p}[u_\ell] \right) \leq \liminf_{n \rightarrow \infty} v_\gamma(K_R \cap E_{\lambda, \gamma/p}[u_n]) \\ & \leq C \liminf_{n \rightarrow \infty} \frac{\|\nabla u_n\|_{L^p(\mathbb{R}^N)}^p}{\lambda^p} \leq C \frac{\|\nabla u\|_{L^p(\mathbb{R}^N)}^p}{\lambda^p}. \end{aligned}$$

2D. Proof of Theorem 2.2 for BV-functions. We choose a sequence $\rho_n \in C_c^\infty(\mathbb{R}^N)$, with $\rho_n = 2^{2n} \rho(2^n \cdot)$ and $\int_{\mathbb{R}^N} \rho dx = 1$, and set $u_n := u * \rho_n$. Then $u_n \in \dot{W}^{1,1}(\mathbb{R}^N)$ and $u_n \rightarrow u$ almost everywhere. This means if $G_L := \{(x, h) \in \mathbb{R}^N \times \mathbb{R}^N : |x| \leq L, L^{-1} \leq |h| \leq L\}$ then

$$\lim_{n \rightarrow \infty} v_\gamma(E_{\lambda, \gamma}[u_n] \cap G_L) = v_\gamma(E_{\lambda, \gamma}[u] \cap G_L),$$

by dominated convergence. Also

$$\|\nabla u_n\|_{L^1(\mathbb{R}^N)} = \sup_{\substack{\vec{\phi} \in C_c^\infty \\ \|\vec{\phi}\|_\infty \leq 1}} \left| \int u_n(x) \operatorname{div} \vec{\phi}(x) dx \right| = \sup_{\substack{\vec{\phi} \in C_c^\infty \\ \|\vec{\phi}\|_\infty \leq 1}} \left| \int u(x) \operatorname{div}(\rho_n * \vec{\phi})(x) dx \right| \leq \|\nabla u\|_{\mathcal{M}};$$

here we used $\|\rho_n * \vec{\phi}\|_\infty \leq \|\vec{\phi}\|_\infty$ for the last inequality. Combining these two limiting identities with Theorem 2.2 we get the desired inequalities with $E_{\lambda, \gamma}[u]$ replaced by $E_{\lambda, \gamma}[u] \cap G_L$. By monotone convergence we may finish the proof letting $L \rightarrow \infty$. \square

3. Proof of Theorem 1.1

We extend and refine arguments from [Brezis and Nguyen 2018; Brezis et al. 2021b], which are partially inspired by techniques developed in [Bourgain et al. 2001].

3A. A Lebesgue differentiation lemma. Our argument uses the following standard variant of the Lebesgue differentiation theorem. For lack of a proper reference, a proof is provided for the convenience of the reader.

Lemma 3.1. *Let $u \in \dot{W}^{1,1}(\mathbb{R}^N)$ and let $\{\delta_n\}$ be a sequence of positive numbers with $\lim_{n \rightarrow \infty} \delta_n = 0$. Then*

$$\lim_{n \rightarrow \infty} \frac{u(x + \delta_n h) - u(x)}{\delta_n} = \langle h, \nabla u(x) \rangle$$

for almost every $(x, h) \in \mathbb{R}^N \times \mathbb{R}^N$.

Proof. If $u \in C^1$ with compact support the limit relation clearly holds for all (x, h) . We shall below consider for each $\theta \in \mathbb{S}^{N-1}$ the maximal function

$$\mathfrak{M}_\theta F(x) = \sup_{t>0} \frac{1}{t} \int_0^t |F(x + r\theta)| \, dr,$$

which is well-defined for all θ , a measurable function on $\mathbb{R}^N \times \mathbb{S}^{N-1}$, and satisfies a weak-type (1, 1) inequality

$$\mathcal{L}^N(\{x \in \mathbb{R}^N : \mathfrak{M}_\theta F(x) > a\}) \leq 5a^{-1} \|F\|_1.$$

Let $u \in \dot{W}^{1,1}(\mathbb{R}^N)$ and $\mathcal{A}_M = \{h \in \mathbb{R}^N : 2^{-M} \leq |h| \leq 2^M\}$. It suffices to prove the limit relation for almost every $(x, h) \in \mathbb{R}^N \times \mathcal{A}_M$. From (2-10) we get that, for every $n \geq 1$,

$$\frac{u(x + \delta_n h) - u(x)}{\delta_n} = \frac{1}{\delta_n |h|} \int_0^{\delta_n |h|} \left\langle h, \nabla u \left(x + r \frac{h}{|h|} \right) \right\rangle dr$$

for \mathcal{L}^{2N} almost every $(x, h) \in \mathbb{R}^N \times \mathcal{A}_M$; as a result, there exist representatives of $u, \nabla u$ and a null set $\mathcal{N} \in \mathbb{R}^N \times \mathcal{A}_M$ such that the identity holds for all $(x, h) \in \mathcal{N}^c$ and all $n \geq 1$. It suffices to show that, for every $\alpha > 0, \varepsilon > 0$,

$$\mathcal{L}^{2N} \left(\left\{ (x, h) \in \mathbb{R}^N \times \mathcal{A}_M : \limsup_{n \rightarrow \infty} \left| \frac{1}{\delta_n |h|} \int_0^{\delta_n |h|} \langle h, \nabla u(x + rh) \rangle \, dr - \langle h, \nabla u(x) \rangle \right| > \alpha \right\} \right) \leq \varepsilon. \tag{3-1}$$

Let $v \in C_c^1$ so that $\|\nabla(v - u)\|_1 \leq \alpha\varepsilon / (12\mathcal{L}^N(\mathcal{A}_M))$. Let $g = u - v$. Since the asserted limiting relation holds for v , we see that the expression on the left-hand side of (3-1) is dominated by

$$\begin{aligned} \mathcal{L}^{2N} \left(\left\{ (x, h) \in \mathbb{R}^N \times \mathcal{A}_M : |\nabla g(x)| + \sup_{n>0} \frac{1}{\delta_n |h|} \int_0^{\delta_n |h|} \left| \nabla g \left(x + r \frac{h}{|h|} \right) \right| \, dr > \alpha \right\} \right) \\ \leq 2\mathcal{L}^N(\mathcal{A}_M) \alpha^{-1} \|\nabla g\|_1 + \int_{\mathcal{A}_M} \mathcal{L}^N \left(\left\{ x : \mathfrak{M}_{h/|h|} |\nabla g|(x) > \frac{\alpha}{2} \right\} \right) \, dh \\ \leq 12\mathcal{L}^N(\mathcal{A}_M) \alpha^{-1} \|\nabla g\|_1 \leq \varepsilon \end{aligned}$$

since $\|\nabla g\|_1 \leq \alpha\varepsilon / (12\mathcal{L}^N(\mathcal{A}_M))$. □

3B. The lower bounds for $\liminf \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u])$. We use Lemma 3.1 to establish lower bounds, relying on an idea in [Brezis and Nguyen 2018], where the case $\gamma = -1$ was considered.

Lemma 3.2. *Let $1 \leq p < \infty$ and $u \in \dot{W}^{1,p}(\mathbb{R}^N)$. Then:*

(i) *For $\gamma > 0$,*

$$\liminf_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \geq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p.$$

(ii) *For $\gamma < 0$,*

$$\liminf_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \geq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p.$$

Proof. We write, for $\lambda > 0$ and $\delta > 0$,

$$\begin{aligned} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) &= \lambda^p \iint_{|u(x+h)-u(x)|/|h|^{1+\gamma/p} > \lambda} |h|^{\gamma-N} \, dh \, dx \\ &= \lambda^p \delta^\gamma \iint_{|(u(x+\delta h)-u(x))/(\delta|h)||^p > \lambda^p \delta^\gamma |h|^\gamma} |h|^{\gamma-N} \, dh \, dx; \end{aligned}$$

here we have changed variables replacing h by δh . Hence

$$\lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) = \iint \mathbb{1}_{(|h|^\gamma, \infty)} \left(\left| \frac{u(x + \delta h) - u(x)}{\delta|h|} \right|^p \right) |h|^{\gamma-N} \, dh \, dx, \quad \text{with } \delta = \lambda^{-p/\gamma}. \tag{3-2}$$

We now take a sequence $\{\lambda_n\}$ of positive numbers, set $\delta_n = \lambda_n^{-p/\gamma}$ and note that

$$\lim_{n \rightarrow \infty} \delta_n = 0 \quad \text{if} \quad \begin{cases} \lim_{n \rightarrow \infty} \lambda_n = \infty \text{ and } \gamma > 0, \\ \lim_{n \rightarrow \infty} \lambda_n = 0 \text{ and } \gamma < 0. \end{cases} \tag{3-3}$$

Also observe that

$$\liminf_{n \rightarrow \infty} \mathbb{1}_{(|h|^\gamma, \infty)}(s_n) \geq \mathbb{1}_{(|h|^{-\gamma}, \infty)}(t) \quad \text{if} \quad \lim_{n \rightarrow \infty} s_n = t.$$

Now assume that $\lambda_n \rightarrow \infty$ if $\gamma > 0$ and $\lambda_n \rightarrow 0^+$ if $\gamma < 0$ and stay with $\delta_n = \lambda_n^{-p/\gamma}$, a sequence which converges to 0 in both cases. Use Fatou’s lemma in (3-2) and combine it with Lemma 3.1 to get

$$\begin{aligned} \liminf_{n \rightarrow \infty} \lambda_n^p v_\gamma(E_{\lambda_n, \gamma/p}[u]) &\geq \iint \liminf_{n \rightarrow \infty} \mathbb{1}_{(|h|^\gamma, \infty)} \left(\left| \frac{u(x + \delta_n h) - u(x)}{\delta_n |h|} \right|^p \right) |h|^{\gamma-N} \, dh \, dx \\ &\geq \iint \mathbb{1}_{(|h|^\gamma, \infty)} \left(\lim_{n \rightarrow \infty} \left| \frac{u(x + \delta_n h) - u(x)}{\delta_n |h|} \right|^p \right) |h|^{\gamma-N} \, dh \, dx \\ &= \iint_{|h|^\gamma < |\langle h/|h|, \nabla u(x) \rangle|^p} |h|^{\gamma-N} \, dh \, dx =: J_\gamma. \end{aligned}$$

We use polar coordinates $h = r\theta$ and write the last expression as

$$\begin{aligned} J_\gamma &= \iint_{\mathbb{R}^N \times \mathbb{S}^{N-1}} \int_{r^\gamma < |\langle \theta, \nabla u(x) \rangle|^p} r^{\gamma-1} \, dr \, d\theta \, dx \\ &= \frac{1}{|\gamma|} \iint_{\mathbb{R}^N \times \mathbb{S}^{N-1}} |\langle \theta, \nabla u(x) \rangle|^p \, d\theta \, dx = \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p, \end{aligned}$$

with the calculation valid in both cases $\gamma > 0$ and $\gamma < 0$. □

3C. Upper bounds for $\limsup \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u])$, for C_c^1 functions. We assume that $u \in C^1$ is compactly supported and obtain the sharp upper bounds for $\limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u])$ when $\gamma > 0$ and $\limsup_{\lambda \rightarrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u])$ when $\gamma < 0$.

Lemma 3.3. *Suppose $u \in C_c^1(\mathbb{R}^N)$ and $1 \leq p < \infty$. Then the following hold:*

(i) *If $\gamma > 0$ then*

$$\limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p.$$

(ii) *If $\gamma < 0$ then*

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p.$$

(iii) *The statement in part (i) continues to hold for $u \in C^1(\mathbb{R}^N)$ whose gradient is compactly supported.*

Remark 3.4. The subtlety in part (iii) above is only relevant in dimension $N = 1$, since if $N \geq 2$, then any function in $C^1(\mathbb{R}^N)$ with a compactly supported gradient is constant outside a compact set.

Proof of Lemma 3.3. We distinguish the cases $\gamma > 0$ and $\gamma < 0$.

The case $\gamma > 0$. We assume that ∇u is compactly supported. To prove part (iii) (and thus part (i)) assume

$$\lambda \geq L := \left\| \left(\sum_{i=1}^N |\partial_i u|^2 \right)^{1/2} \right\|_{L^\infty(\mathbb{R}^N)}. \tag{3-4}$$

Then

$$(x, y) \in E_{\lambda, \gamma/p}[u] \implies \lambda |x - y|^{\gamma/p} \leq L \implies |x - y| \leq 1. \tag{3-5}$$

Furthermore, if $(x, y) \in E_{\lambda, \gamma/p}[u]$, then writing $y = x + r\omega$ with $r > 0$ and $\omega \in \mathbb{S}^{N-1}$, we have

$$\lambda r^{\gamma/p} \leq |\nabla u(x) \cdot \omega| + \rho(r), \quad \text{with } \rho(r) := \sup_{x \in \mathbb{R}^N} \sup_{|h| \leq r} |\nabla u(x+h) - \nabla u(x)|; \tag{3-6}$$

since ∇u is uniformly continuous on \mathbb{R}^N , we have $\rho(r) \searrow 0$ as $r \searrow 0$. This, together with the first implication of (3-5), shows

$$\lambda r^{\gamma/p} \leq |\nabla u(x) \cdot \omega| + \rho\left(\left(\frac{L}{\lambda}\right)^{p/\gamma}\right). \tag{3-7}$$

Let B be a ball centered at the origin containing $\text{supp}(\nabla u)$, and let \tilde{B} be the expanded ball with radius $1 + \text{rad}(B)$. Then for $x \notin \tilde{B}$, we have $Q_{\gamma/p} u(x, y) = 0$ for every y with $|x - y| \leq 1$, and (3-5) shows $(x, y) \notin E_{\lambda, \gamma/p}[u]$ for every y with $|x - y| > 1$, so $E_{\lambda, \gamma/p}[u] \subseteq \tilde{B} \times \mathbb{R}^N$. Define, for $x \in \tilde{B}$, $\omega \in \mathbb{S}^{N-1}$, and $\lambda > 0$

$$\bar{R}(x, \omega, \lambda) := \left(\lambda^{-1} \left(|\nabla u(x) \cdot \omega| + \rho\left(\left(\frac{L}{\lambda}\right)^{p/\gamma}\right) \right) \right)^{p/\gamma}.$$

Then by (3-7),

$$\begin{aligned} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) &\leq \lambda^p \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} \int_0^{\bar{R}(x, \omega, \lambda)} r^{\gamma-1} dr d\omega dx \\ &= \gamma^{-1} \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} \left(|\nabla u(x) \cdot \omega| + \rho\left(\left(\frac{L}{\lambda}\right)^{p/\gamma}\right) \right)^p d\omega dx. \end{aligned}$$

Letting $\lambda \rightarrow \infty$ we get

$$\limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \gamma^{-1} \kappa(p, N) \int_{\tilde{B}} |\nabla u(x)|^p dx$$

and hence the assertion.

The case $\gamma < 0$. We first note that if $(x, y) \in E_{\lambda, \gamma/p}[u]$, then writing $y = x + r\omega$, we have again (3-6).

Now let $\varepsilon > 0$, and let $\delta(\varepsilon) > 0$ be such that $\rho(r) \leq \varepsilon$ for $0 < r \leq \delta(\varepsilon)$. Let

$$r_\lambda(x, \omega, \varepsilon) = \min \left\{ \delta(\varepsilon), \left(\frac{\lambda}{|\nabla u(x) \cdot \omega| + \varepsilon} \right)^{-p/\gamma} \right\}.$$

Note that $r_\lambda(x, \omega, \varepsilon) > 0$ for $\lambda > 0$. Also if $(x, x + r\omega) \in E_{\lambda, \gamma/p}[u]$ then $r \geq r_\lambda(x, \omega, \varepsilon)$; indeed, either $r_\lambda(x, \omega, \varepsilon) \geq \delta(\varepsilon)$ already, or else $r_\lambda(x, \omega, \varepsilon) < \delta(\varepsilon)$, in which case (3-6) shows

$$r_\lambda(x, \omega, \varepsilon) \geq \left(\frac{\lambda}{|\nabla u(x) \cdot \omega| + \varepsilon} \right)^{-p/\gamma}.$$

Finally let B be any ball in \mathbb{R}^N containing the support of u , and let \tilde{B} be the double ball. Then

$$\begin{aligned} \limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u] \cap (\tilde{B} \times \mathbb{R}^N)) &\leq \limsup_{\lambda \searrow 0} \lambda^p \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} \int_{r_\lambda(x, \omega, \varepsilon)}^\infty r^{\gamma-1} dr d\omega dx \\ &= \limsup_{\lambda \searrow 0} \lambda^p \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} \frac{1}{|\gamma|} [r_\lambda(x, \omega, \varepsilon)]^\gamma d\omega dx \\ &= \limsup_{\lambda \searrow 0} \frac{1}{|\gamma|} \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} \max\{\lambda^p \delta(\varepsilon)^\gamma, (|\nabla u(x) \cdot \omega| + \varepsilon)^p\} d\omega dx \\ &= \frac{1}{|\gamma|} \int_{\tilde{B}} \int_{\mathbb{S}^{N-1}} (|\nabla u(x) \cdot \omega| + \varepsilon)^p d\omega dx. \end{aligned}$$

Since $\varepsilon > 0$ was arbitrary we obtain

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u] \cap (\tilde{B} \times \mathbb{R}^N)) \leq \frac{1}{|\gamma|} \kappa(p, N) \|\nabla u\|_{L^p(\mathbb{R}^N)}^p. \tag{3-8}$$

Since $u = 0$ in $\mathbb{R}^N \setminus B$, if $(x, y) \in E_{\lambda, \gamma/p}[u] \cap ((\mathbb{R}^N \setminus \tilde{B}) \times \mathbb{R}^N)$ then $y \in B$. Therefore

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u] \cap ((\mathbb{R}^N \setminus \tilde{B}) \times \mathbb{R}^N)) \leq \limsup_{\lambda \searrow 0} \lambda^p \int_B \int_{\mathbb{R}^N \setminus \tilde{B}} |x - y|^{\gamma-N} dx dy = 0.$$

This finishes the proof of part (ii). □

In dimension $N = 1$, when $\gamma < -1$, one can also weaken the hypothesis $u \in C_c^1(\mathbb{R})$ in Lemma 3.3 to $u \in C^1(\mathbb{R})$ and u' is compactly supported:

Lemma 3.5. *Suppose $u \in C^1(\mathbb{R})$, u' is compactly supported, and $1 \leq p < \infty$. If $\gamma < -1$ then*

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \frac{\kappa(p, N)}{|\gamma|} \|u'\|_{L^p(\mathbb{R})}^p.$$

Proof. Let $\text{supp}(u') \subset B := (-\beta, \beta)$. By (3-8) we have

$$\limsup_{\lambda \searrow 0} v_\gamma(E_{\lambda, \gamma/p}[u] \cap (-2\beta, 2\beta) \times \mathbb{R}) \leq \frac{1}{|\gamma|} \kappa(p, 1) \|u'\|_{L^p(\mathbb{R})}^p.$$

Moreover, since u is constant on (β, ∞) and constant on $(-\infty, -\beta)$, if $(x, y) \in E_{\lambda, \gamma/p}[u]$ and $x < -2\beta$ then $y > -\beta$, and if $(x, y) \in E_{\lambda, \gamma/p}[u]$ and $x > 2\beta$ then $y < \beta$. Since $\gamma < -1$,

$$v_\gamma(E_{\lambda, \gamma/p}[u] \cap (\mathbb{R} \setminus (-2\beta, 2\beta)) \times \mathbb{R}) \leq \int_{2\beta}^\infty \int_{-\infty}^\beta (x - y)^{\gamma-1} dy dx + \int_{-\infty}^{-2\beta} \int_{-\beta}^\infty (y - x)^{\gamma-1} dy dx < \infty.$$

We conclude

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u] \cap (\mathbb{R} \setminus (-2\beta, 2\beta)) \times \mathbb{R}) = 0. \quad \square$$

3D. Upper bounds for $\limsup \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u])$, for general $\dot{W}^{1,p}$ functions. Let $N \geq 1$, $1 \leq p < \infty$ and $u \in \dot{W}^{1,p}(\mathbb{R}^N)$. In light of Lemma 3.2, to prove the limiting relations (1-7) and (1-8) in Theorem 1.1, we need only show that

$$\limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p \tag{3-9}$$

if $\gamma > 0$ and

$$\limsup_{\lambda \searrow 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) \leq \frac{\kappa(p, N)}{|\gamma|} \|\nabla u\|_{L^p(\mathbb{R}^N)}^p \tag{3-10}$$

if $\gamma < 0$ and $p > 1$, or $\gamma < -1$ and $p = 1$. Lemma 3.3(i)–(ii) asserts that these desired inequalities hold for functions in $C_c^1(\mathbb{R}^N)$. When $N \geq 2$ or $p > 1$, a general $\dot{W}^{1,p}(\mathbb{R}^N)$ function can be approximated in $\dot{W}^{1,p}(\mathbb{R}^N)$ by functions in $C_c^1(\mathbb{R}^N)$; by [Hajlasz and Kałamajska 1995], there exists a sequence $\{u_n\}$ in $C_c^\infty(\mathbb{R}^N)$ such that $\lim_{n \rightarrow \infty} \|\nabla(u_n - u)\|_{L^p(\mathbb{R}^N)} = 0$. If further $\gamma > 0$, or $\gamma < 0$ and $p > 1$, or $\gamma < -1$ and $p = 1$, then by parts (i) of Theorems 1.3 and 1.4 (proved in Section 2), we have

$$\sup_{\lambda > 0} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u_n - u]) \leq C_{N,p,\gamma}^p \|\nabla(u_n - u)\|_{L^p(\mathbb{R}^N)}^p. \tag{3-11}$$

It follows that, for every n and every $\delta \in (0, 1)$,

$$\begin{aligned} \limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]) &\leq \limsup_{\lambda \rightarrow \infty} \lambda^p v_\gamma(E_{(1-\delta)\lambda, \gamma/p}[u_n]) + \sup_{\lambda > 0} \lambda^p v_\gamma(E_{\delta\lambda, \gamma/p}[u_n - u]) \\ &\leq \frac{\kappa(p, N)}{|\gamma|(1-\delta)} \|\nabla u_n\|_{L^p(\mathbb{R}^N)}^p + \frac{C_{N,p,\gamma}^p \|\nabla(u_n - u)\|_{L^p(\mathbb{R}^N)}^p}{\delta^p} \end{aligned} \tag{3-12}$$

if $\gamma > 0$, and a similar inequality holds with $\limsup_{\lambda \rightarrow \infty}$ replaced by $\limsup_{\lambda \searrow 0}$ if $\gamma < 0$, $p > 1$ or $\gamma < -1$, $p = 1$. Letting first $n \rightarrow \infty$ and then $\delta \rightarrow 0$, we get the desired conclusions (3-9) and (3-10) under the corresponding conditions on γ and p .

It remains to tackle the case $N = p = 1$, in which case we only need to prove (3-9) when $\gamma > 0$ and (3-10) when $\gamma < -1$. Using (2-11), we approximate u by finding a sequence $\{u_n\}$ in $C^\infty(\mathbb{R})$ so that u'_n are compactly supported for each n , and $\lim_{n \rightarrow \infty} \|u'_n - u'\|_{L^1(\mathbb{R})} = 0$. Since the desired inequalities hold for u_n in place of u by Lemma 3.3(iii) and Lemma 3.5, and since part (i) of Theorem 1.4 applies to give

(3-11) when $\gamma > 0$ or $\gamma < -1$, our earlier argument in (3-12) can be repeated to yield (3-9) when $\gamma > 0$ and (3-10) when $\gamma < -1$. This completes our proof of parts (a) and (b) of Theorem 1.1.

3E. Conclusion of the proof of Theorem 1.1. In Section 3D we proved parts (a) and (b) of Theorem 1.1. The lower bound for the \liminf in part (c) has been established in Lemma 3.2(ii), and the limiting equality for $u \in C_c^1(\mathbb{R}^N)$ when $p = 1$ and $-1 \leq \gamma < 0$ follows by combining that with the upper bound for the \limsup in part (ii) of Lemma 3.3. The proof of the negative result in part (c) of the theorem (generic failure for $p = 1, -1 \leq \gamma < 0$) will be given in Proposition 6.6 below. \square

3F. On limit formulas for $\dot{B}\dot{V}(\mathbb{R})$ -functions: the proof of Proposition 1.2. When $p = 1$, Poliakovsky [2022] asked whether (1-7) still holds for $u \in \dot{B}\dot{V}(\mathbb{R}^N)$ instead of $\dot{W}^{1,1}(\mathbb{R}^N)$ if $\gamma = N$. More generally, one may wonder whether it is possible that, for all $u \in \dot{B}\dot{V}(\mathbb{R}^N)$, one has

$$\lim_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \frac{\kappa(1, N)}{|\gamma|} \|\nabla u\|_{\mathcal{M}} \quad \text{when } \gamma > 0, \tag{3-13}$$

$$\lim_{\lambda \rightarrow 0^+} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \frac{\kappa(1, N)}{|\gamma|} \|\nabla u\|_{\mathcal{M}} \quad \text{when } \gamma < 0. \tag{3-14}$$

We show that this is *not* the case.

First, when $-1 \leq \gamma < 0$, Theorem 1.8(i) (proved in Proposition 6.3 below) shows that even if $u \in \dot{W}^{1,1}(\mathbb{R}^N)$, it may happen that $\lim_{\lambda \rightarrow 0^+} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \infty$. So (3-14) cannot hold for all $u \in \dot{B}\dot{V}(\mathbb{R}^N)$ for such γ .

The following lemma provides examples of failure of (3-13) and (3-14) when $\gamma \in \mathbb{R} \setminus [-1, 0]$, since $|\gamma + 1| \neq |\gamma|$ unless $\gamma = -\frac{1}{2}$:

Lemma 3.6. *Suppose $N \geq 1$ and $u = \mathbb{1}_\Omega$, where Ω is any bounded convex domain in \mathbb{R}^N with smooth boundary. Then $u \in \dot{B}\dot{V}(\mathbb{R}^N)$ and*

$$\lim_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \frac{\kappa(1, N)}{|\gamma + 1|} \|\nabla u\|_{\mathcal{M}} \quad \text{for all } \gamma > -1,$$

while

$$\lim_{\lambda \rightarrow 0^+} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \frac{\kappa(1, N)}{|\gamma + 1|} \|\nabla u\|_{\mathcal{M}} \quad \text{for all } \gamma < -1.$$

Proof. First consider the case $N = 1$. If $u = \mathbb{1}_{[0,\infty)}$ (so that $\|u'\|_{\mathcal{M}(\mathbb{R})} = 1$), then, for every $\gamma \in \mathbb{R} \setminus \{-1\}$ and $\lambda > 0$, one has

$$v_\gamma(E_{\lambda,\gamma}[u]) = 2v_\gamma(\{(x, y) \in \mathbb{R} : x \geq 0, y < 0, |x - y|^{-(\gamma+1)} \geq \lambda\}) = \frac{2}{|\gamma + 1|} \frac{1}{\lambda}, \tag{3-15}$$

which follows from a change of variables $s = x - y, t = x + y$: when $\gamma > -1$, one has

$$v_\gamma(E_{\lambda,\gamma}[u]) = \int_0^{\lambda^{-1/(\gamma+1)}} \int_{-s}^s dt s^{\gamma-1} ds = 2 \int_0^{\lambda^{-1/(\gamma+1)}} s^\gamma ds = \frac{2}{\gamma + 1} \frac{1}{\lambda},$$

while when $\gamma < -1$, one has

$$v_\gamma(E_{\lambda,\gamma}[u]) = \int_{\lambda^{-1/(\gamma+1)}}^\infty \int_{-s}^s dt s^{\gamma-1} ds = 2 \int_{\lambda^{-1/(\gamma+1)}}^\infty s^\gamma ds = \frac{2}{|\gamma + 1|} \frac{1}{\lambda}.$$

A similar calculation shows that if $u = \mathbb{1}_I$ is a characteristic function of a bounded open interval (so that $\|u'\|_{\mathcal{M}(\mathbb{R})} = 2$), then

$$\lim_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda, \gamma}[u]) = \frac{2}{|\gamma + 1|} \|u'\|_{\mathcal{M}(\mathbb{R})} \quad \text{for all } \gamma > -1, \tag{3-16}$$

while

$$\lim_{\lambda \rightarrow 0^+} \lambda v_\gamma(E_{\lambda, \gamma}[u]) = \frac{2}{|\gamma + 1|} \|u'\|_{\mathcal{M}(\mathbb{R})} \quad \text{for all } \gamma < -1; \tag{3-17}$$

we also have

$$\sup_{\lambda > 0} \lambda v_\gamma(E_{\lambda, \gamma}[u]) \leq \frac{2}{|\gamma + 1|} \|u'\|_{\mathcal{M}(\mathbb{R})} \quad \text{for all } \gamma \in \mathbb{R} \setminus \{-1\}. \tag{3-18}$$

Now consider the case $N \geq 2$. Let Ω be a bounded convex domain in \mathbb{R}^N with smooth boundary and $u = \mathbb{1}_\Omega$. Then $u \in \mathbf{BV}(\mathbb{R}^N)$ with $\|\nabla u\|_{\mathcal{M}} = \mathcal{L}^{N-1}(\partial\Omega)$. The method of rotation shows

$$\lambda v_\gamma(E_{\lambda, \gamma}[u]) = \frac{1}{2} \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \lambda v_\gamma(E_{\lambda, \gamma}[u_{\omega, x'}]) \, dx' \, d\omega,$$

where $u_{\omega, x'}(t) := u(x' + t\omega)$ for $\omega \in \mathbb{S}^{N-1}$ and $x' \in \omega^\perp$. Note that $\|u'_{\omega, x'}\|_{\mathcal{M}(\mathbb{R})} \leq 2$ for all $\omega \in \mathbb{S}^{N-1}$ and all $x' \in \omega^\perp$, since Ω is convex and every line only meets $\partial\Omega$ at at most two points. Thus (3-16), (3-18) and the dominated convergence theorem allow one to show that

$$\lim_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda, \gamma}[u]) = \frac{1}{|\gamma + 1|} \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \|u'_{\omega, x'}\|_{\mathcal{M}(\mathbb{R})} \, dx' \, d\omega \quad \text{for all } \gamma > -1,$$

and using (3-17) in place of (3-16) we obtain the same conclusion with $\lim_{\lambda \rightarrow \infty}$ replaced by $\lim_{\lambda \rightarrow 0^+}$ if $\gamma < -1$. It remains to observe that

$$\int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \|u'_{\omega, x'}\|_{\mathcal{M}(\mathbb{R})} \, dx' \, d\omega = \kappa(1, N) \|\nabla u\|_{\mathcal{M}}. \tag{3-19}$$

This holds by Fubini's theorem if $u = \mathbb{1}_\Omega$ is replaced by $u_\varepsilon := u * \rho_\varepsilon$, where ρ_ε is a suitable family of mollifiers, because the left-hand side is then just

$$\int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \int_{\mathbb{R}} \left| \frac{d}{dt} u_\varepsilon(x' + t\omega) \right| \, dt \, dx' \, d\omega = \int_{\mathbb{S}^{N-1}} \int_{\mathbb{R}^N} |\omega \cdot \nabla u_\varepsilon(x)| \, dx \, d\omega,$$

which equals $\kappa(1, N) \|\nabla u_\varepsilon\|_{L^1(\mathbb{R}^N)}$. One then just needs to let $\varepsilon \rightarrow 0$ to obtain (3-19): in fact, a standard argument shows that

$$\lim_{\varepsilon \rightarrow 0^+} \|\nabla u_\varepsilon\|_{L^1(\mathbb{R}^N)} = \|\nabla u\|_{\mathcal{M}(\mathbb{R}^N)}.$$

so it remains to prove that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \int_{\mathbb{R}} \left| \frac{d}{dt} u_\varepsilon(x' + t\omega) \right| \, dt \, dx' \, d\omega = \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \|u'_{\omega, x'}\|_{\mathcal{M}(\mathbb{R})} \, dx' \, d\omega. \tag{3-20}$$

But for every $\omega \in \mathbb{S}^{N-1}$, and almost every $x' \in \omega^\perp$ (as long as $t \mapsto x' + t\omega$ parametrizes a line $L_{\omega, x'}$ that is either disjoint from Ω , or intersects $\partial\Omega$ transversely at two different points), we have

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\mathbb{R}} \left| \frac{d}{dt} u_\varepsilon(x' + t\omega) \right| \, dt = \|u'_{\omega, x'}\|_{\mathcal{M}(\mathbb{R})}. \tag{3-21}$$

The validity of (3-21) is clear if $L_{\omega,x'}$ does not intersect Ω , while if $L_{\omega,x'}$ intersects $\partial\Omega$ transversely at two different points, then we can choose a coordinate system so that $\omega = (0, \dots, 0, 1)$, and assume that for some open neighborhood U of x' in ω^\perp , the intersection of $U \times L_{\omega,x'}$ with Ω takes the form

$$\{(y', y_N) : y' \in U, \phi_1(y') < y_N < \phi_2(y')\}$$

for some smooth functions ϕ_1 and ϕ_2 of $y' \in U$. Then, for $\varepsilon > 0$ sufficiently small,

$$\begin{aligned} \int_{\mathbb{R}} \left| \frac{d}{dt} u_\varepsilon(x' + t\omega) \right| dt &= \int_{\mathbb{R}} \left| \int_{\mathbb{R}^N} \mathbb{1}_\Omega(y) \partial_N \rho_\varepsilon(x' - y', t - y_N) dy \right| dt \\ &= \int_{\mathbb{R}} \left| - \int_{\mathbb{R}^N} \mathbb{1}_{\phi_1(y') < y_N < \phi_2(y')} \frac{\partial}{\partial y_N} [\rho_\varepsilon(x' - y', t - y_N)] dy \right| dt \\ &= \int_{\mathbb{R}} \left| \int_{\mathbb{R}^{N-1}} \rho_\varepsilon(x' - y', t - \phi_1(y')) - \rho_\varepsilon(x' - y', t - \phi_2(y')) dy' \right| dt \\ &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}^{N-1}} \rho_\varepsilon(x' - y', t - \phi_1(y')) dy' + \int_{\mathbb{R}^{N-1}} \rho_\varepsilon(x' - y', t - \phi_2(y')) dy' \right) dt \\ &= 2 \int_{\mathbb{R}^{N-1}} \int_{\mathbb{R}} \rho_\varepsilon(x' - y', t) dt dy' \\ &= 2 = \|u'_{\omega,x'}\|_{\mathcal{M}(\mathbb{R})}. \end{aligned}$$

This proves (3-21), and then the dominated convergence theorem allows one to conclude the proof of (3-20). □

Remark. The identity (3-19) for $u = \mathbb{1}_\Omega$ can be derived from Crofton’s formula for rather general (not necessarily convex) domains Ω . See [Federer 1969, Chapter 3.2.26], which showed that when $\partial\Omega$ is rectifiable, then its $(N-1)$ -dimensional Hausdorff measure $\mathcal{H}^{N-1}(\partial\Omega)$ is equal to $\mathcal{I}_1^{N-1}(\partial\Omega)$, where $\mathcal{I}_1^{N-1}(\partial\Omega)$ is given by [Federer 1969, Chapter 2.10.15] as

$$\frac{1}{\beta_1(N, N-1)} \int_{p \in \mathbf{O}^*(N, N-1)} \int_{y \in \mathbb{R}^{N-1}} N(p|\partial\Omega, y) dy dp;$$

here $\mathbf{O}^*(N, N-1)$ is the space of all orthogonal projections p from \mathbb{R}^N onto \mathbb{R}^{N-1} , dp is the right- $\mathbf{O}(N)$ -invariant measure on $\mathbf{O}^*(N, N-1)$ normalized so that $\int_{\mathbf{O}^*(N, N-1)} dp = 1$, $N(p|\partial\Omega, y)$ is the number of points $x \in \partial\Omega$ so that $px = y$, and

$$\beta_1(N, N-1) = \frac{\Gamma(N/2)}{\Gamma((N+1)/2)\Gamma(1/2)}$$

according to [Federer 1969, Chapter 3.2.13]. It follows that, for $u = \mathbb{1}_\Omega$,

$$\begin{aligned} \int_{\mathbb{S}^{N-1}} \int_{\omega^\perp} \|u'_{\omega,x'}\|_{\mathcal{M}(\mathbb{R})} dx' d\omega &= \mathcal{H}^{N-1}(\mathbb{S}^{N-1}) \int_{p \in \mathbf{O}^*(N, N-1)} \int_{y \in \mathbb{R}^{N-1}} N(p|\partial\Omega, y) dy dp \\ &= \frac{2\pi^{N/2}}{\Gamma(N/2)} \beta_1(N, N-1) \mathcal{H}^{N-1}(\partial\Omega) \\ &= \frac{2\pi^{(N-1)/2}}{\Gamma((N+1)/2)} \|\nabla u\|_{\mathcal{M}} = \kappa(1, N) \|\nabla u\|_{\mathcal{M}}, \end{aligned}$$

as asserted in (3-19).

4. From weak-type bounds on quotients to $\dot{W}^{1,p}$ and $\dot{B}\dot{V}$

In this section we complete the proofs of Theorems 1.3 and 1.4 proving part (ii) of these theorems. We use as a key tool the BBM formula discovered in [Bourgain et al. 2001] (see also [Dávila 2002] for additional information for the BV case), in a way that is reminiscent of the proof of [Nguyen 2006, Theorem 2], and we apply duality for Lorentz spaces to control the double integral arising in the BBM formula. The BBM formula stated in [Bourgain et al. 2001] is quite flexible, involving a bounded smooth domain Ω and a sequence of nonnegative radial mollifiers $\rho_n(|x|)$, with $\int_0^\infty \rho_n(r)r^{N-1} dr = 1$ and $\lim_{n \rightarrow \infty} \int_\delta^\infty \rho_n(r)r^{N-1} dr = 0$ for every $\delta > 0$; we will apply it in the case when $\Omega = B_R$, the ball of radius R centered at 0, and $\rho_n(r) = s_n p(2R)^{-s_n p} r^{-N+s_n p} \mathbb{1}_{[0,2R]}(r)$, where $\{s_n\}$ is a sequence of positive numbers tending to 0. As a result, we conclude that if $R > 0$, $1 \leq p < \infty$, $u \in L^p(B_R)$ and

$$\liminf_{s \rightarrow 0^+} s \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy < \infty,$$

then for $p = 1$ we have $u \in \dot{B}\dot{V}(B_R)$ with $\|\nabla u\|_{\mathcal{M}(B_R)}$ being bounded by $\kappa(1, N)$ times the above liminf, and, for $1 < p < \infty$ we have $u \in \dot{W}^{1,p}(B_R)$ and $\|\nabla u\|_{L^p(B_R)}$ being bounded by $\kappa(p, N)/p$ times the above liminf. The assumption $u \in L^p(B_R)$ can easily be relaxed to $u \in L^1(B_R)$, via an observation of Stein as explained in [Brezis 2002, proof of Theorem 2]: if $u \in L^1(B_R)$ and the above liminf is finite for some $1 < p < \infty$, then, for any $\delta > 0$ and any $\varepsilon \in (0, \delta)$, we may consider $u_\varepsilon := u * \phi_\varepsilon(x)$, where $\phi_\varepsilon(x) := \varepsilon^{-N} \phi(\varepsilon^{-1}x)$ and $\phi \in C_c^\infty(B_1)$ is nonnegative and has integral 1. Then u_ε is C^∞ on the closure of the ball $B_{R-\delta}$, so the above formulation of BBM applies, and $\|\nabla u_\varepsilon\|_{L^p(B_{R-\delta})}$ is uniformly bounded independent of $\varepsilon \in (0, \delta)$; indeed Jensen’s inequality implies

$$\iint_{B_{R-\delta} \times B_{R-\delta}} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^p}{|x - y|^{N+p-sp}} dx dy \leq \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy$$

for every ε . This shows that a subsequence of $\{\nabla u_\varepsilon\}$ converges weakly in $L^p(B_{R-\delta})$ to the distributional gradient ∇u on $B_{R-\delta}$, and a desired bound on $\|\nabla u\|_{L^p(B_{R-\delta})}$ follows for every $\delta > 0$.

Suppose now $N \geq 1$, $1 \leq p < \infty$, $\gamma \in \mathbb{R}$, $u \in L^1_{loc}(\mathbb{R}^N)$ and $Q_{\gamma/p}u \in L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)$. Let

$$A := \sup_{R>0} \liminf_{s \rightarrow 0^+} s \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy. \tag{4-1}$$

Suppose A is finite. If $p = 1$, then the BBM formula above implies $u \in \dot{B}\dot{V}(B_R)$ for every $R > 0$, with $\|\nabla u\|_{\mathcal{M}(B_R)} \leq \kappa(1, N)A$ independent of R ; as a result, $u \in \dot{B}\dot{V}(\mathbb{R}^N)$, with $\|\nabla u\|_{\mathcal{M}(\mathbb{R}^N)} \leq \kappa(1, N)A$. Similarly, if $1 < p < \infty$, the above BBM formula (applicable for $u \in L^1_{loc}(\mathbb{R}^N)$) implies $u \in \dot{W}^{1,p}(\mathbb{R}^N)$, with $\|\nabla u\|_{L^p(\mathbb{R}^N)} \leq (\kappa(1, N)A/p)^{1/p}$.

It remains to prove that $A < \infty$. By considering truncations of u we may assume additionally that $u \in L^\infty(\mathbb{R}^N)$; the reduction is based on the pointwise bound

$$Q_{\gamma/p}u_n(x, y) \leq Q_{\gamma/p}u(x, y), \quad \text{where } u_n(x) = \begin{cases} u(x) & \text{if } |u(x)| < n, \\ nu(x)/|u(x)| & \text{if } |u(x)| \geq n. \end{cases}$$

Using the definition of weak derivative we see by a limiting argument that the conclusion $\sup_n \|\nabla u_n\|_p \leq C$ implies $\|\nabla u\|_p \leq C$ if $p > 1$ and $\sup_n \|\nabla u_n\|_{\mathcal{M}} \leq C$ implies $\|\nabla u\|_{\mathcal{M}} \leq C$.

In order to establish our estimate for bounded functions we will use Lorentz duality in the following form: if F, G are measurable functions on \mathbb{R}^{2N} , then, for any $1 < q < \infty$, we have

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} F(x, y)G(x, y) \, dv_\gamma \leq q'[F]_{L^{q,\infty}(\mathbb{R}^{2N}, v_\gamma)}[G]_{L^{q',1}(\mathbb{R}^{2N}, v_\gamma)}, \tag{4-2}$$

where $1/q + 1/q' = 1$,

$$\begin{aligned} [F]_{L^{q,\infty}(\mathbb{R}^{2N}, v_\gamma)} &:= \sup_{\lambda > 0} \lambda v_\gamma(\{|F| > \lambda\})^{1/q} = \sup_{t > 0} t^{1/q} F^*(t), \\ [G]_{L^{q',1}(\mathbb{R}^{2N}, v_\gamma)} &:= \int_0^\infty v_\gamma(\{|G| > \lambda\})^{1/q'} \, d\lambda = \frac{1}{q'} \int_0^\infty t^{1/q'} G^*(t) \frac{dt}{t}; \end{aligned}$$

here $F^*(t) := \inf\{s > 0 : v_\gamma(\{|F| > \lambda\}) \leq s\}$ is the nonincreasing rearrangement of F , and similarly for $G^*(t)$; see [Hunt 1966; Stein and Weiss 1971]. Indeed, (4-2) follows by noticing that

$$\iint_{\mathbb{R}^N \times \mathbb{R}^N} F(x, y)G(x, y) \, dv_\gamma \leq \int_0^\infty F^*(t)G^*(t) \, dt = \int_0^\infty [t^{1/q} F^*(t)][t^{1/q'} G^*(t)] \frac{dt}{t},$$

which is clearly $\leq q'[F]_{L^{q,\infty}(\mathbb{R}^{2N}, v_\gamma)}[G]_{L^{q',1}(\mathbb{R}^{2N}, v_\gamma)}$.

First we consider the case $\gamma > 0$. For sufficiently small $s > 0$, define

$$\theta := \frac{s}{1 + \gamma/p}$$

so that $\theta \in (0, 1)$ and $p - sp = p(1 - \theta)(1 + \gamma/p) - \gamma$. Then, for every $R > 0$,

$$\begin{aligned} \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} \, dx \, dy &= \iint_{\mathbb{R}^N \times \mathbb{R}^N} (\mathcal{Q}_{\gamma/p} u(x, y))^{p(1-\theta)} (|u(x) - u(y)| \mathbb{1}_{B_R \times B_R}(x, y))^{p\theta} \, dv_\gamma \\ &\leq \frac{1}{\theta} [(\mathcal{Q}_{\gamma/p} u)^{p(1-\theta)}]_{L^{1/(1-\theta), \infty}(\mathbb{R}^{2N}, v_\gamma)} [|u(x) - u(y)|^{p\theta}]_{L^{1/\theta, 1}(B_R \times B_R, v_\gamma)} \end{aligned}$$

by (4-2). But

$$[(\mathcal{Q}_{\gamma/p} u)^{p(1-\theta)}]_{L^{1/(1-\theta), \infty}(\mathbb{R}^{2N}, v_\gamma)} = [\mathcal{Q}_{\gamma/p} u]_{L^{p, \infty}(\mathbb{R}^{2N}, v_\gamma)}^{p(1-\theta)}$$

and

$$\begin{aligned} [|u(x) - u(y)|^{p\theta}]_{L^{1/\theta, 1}(B_R \times B_R, v_\gamma)} &\leq (2\|u\|_{L^\infty(\mathbb{R}^N)})^{p\theta} [\mathbb{1}_{B_R \times B_R}]_{L^{1/\theta, 1}(\mathbb{R}^N \times \mathbb{R}^N, v_\gamma)} \\ &= (2\|u\|_{L^\infty(\mathbb{R}^N)})^{p\theta} v_\gamma(B_R \times B_R)^\theta, \end{aligned}$$

from which it follows that

$$s \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} \, dx \, dy \leq \frac{s}{\theta} [\mathcal{Q}_{\gamma/p} u]_{L^{p, \infty}(\mathbb{R}^{2N}, v_\gamma)}^{p(1-\theta)} (2\|u\|_{L^\infty(\mathbb{R}^N)})^{p\theta} v_\gamma(B_R \times B_R)^\theta.$$

Furthermore, since $\gamma > 0$, we have

$$v_\gamma(B_R \times B_R) \leq |B_R| \int_{B_{2R}} \frac{1}{|h|^{N-\gamma}} \, dh < \infty.$$

Recall $\theta = s/(1 + \gamma/p)$. Thus as $s \rightarrow 0^+$, we have

$$\limsup_{s \rightarrow 0^+} s \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy \leq \left(1 + \frac{\gamma}{p}\right) [\mathcal{Q}_{\gamma/p} u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}^p < \infty.$$

Since this upper bound holds uniformly over all $R > 0$, this concludes the argument for the case $\gamma > 0$.

Next we turn to the case $\gamma \leq 0$. We then observe that, for $0 < s < 1$ and every $R > 0$,

$$\begin{aligned} \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy &= \iint_{\mathbb{R}^N \times \mathbb{R}^N} (\mathcal{Q}_{\gamma/p} u(x, y))^{p(1-s/2)} (|u(x) - u(y)| |x - y|^{1-\gamma/p} \mathbb{1}_{B_R \times B_R})^{ps/2} d\nu_\gamma \\ &\leq \frac{2}{s} [(\mathcal{Q}_{\gamma/p} u)^{p(1-s/2)}]_{L^{1/(1-s/2),\infty}(\mathbb{R}^{2N}, \nu_\gamma)} [(|u(x) - u(y)| |x - y|^{1-\gamma/p})^{ps/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)}. \end{aligned}$$

Again

$$[(\mathcal{Q}_{\gamma/p} u)^{p(1-s/2)}]_{L^{1/(1-s/2),\infty}(\mathbb{R}^{2N}, \nu_\gamma)} = [\mathcal{Q}_{\gamma/p} u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}^{p(1-s/2)}$$

and

$$\begin{aligned} [(|u(x) - u(y)| |x - y|^{1-\gamma/p})^{ps/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)} &\leq (2\|u\|_{L^\infty(\mathbb{R}^N)})^{ps/2} [|x - y|^{(p-\gamma)s/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)}. \end{aligned} \tag{4-3}$$

We will show that

$$\limsup_{s \rightarrow 0^+} [|x - y|^{(p-\gamma)s/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)} \leq 1 - \frac{\gamma}{p} \tag{4-4}$$

when $\gamma \leq 0$. We then see that

$$\limsup_{s \rightarrow 0^+} s \iint_{B_R \times B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+p-sp}} dx dy \leq 2 \left(1 - \frac{\gamma}{p}\right) [\mathcal{Q}_{\gamma/p} u]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}^p,$$

which concludes the argument in this case since this bound is uniform in $R > 0$.

It remains to prove (4-4) when $\gamma \leq 0$. Note that in this case $p - \gamma > 0$, so $|x - y|^{(p-\gamma)s/2} \leq (2R)^{(p-\gamma)s/2}$ on $B_R \times B_R$. Thus

$$[|x - y|^{(p-\gamma)s/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)} = \int_0^{(2R)^{(p-\gamma)s/2}} \nu_\gamma \{ (x, y) \in B_R \times B_R : |x - y|^{(p-\gamma)s/2} > \lambda \}^{s/2} d\lambda.$$

If $\gamma < 0$, then

$$\nu_\gamma \{ (x, y) \in B_R \times B_R : |x - y|^{(p-\gamma)s/2} > \lambda \} \leq |B_R| \int_{|h| > \lambda^{2/(s(p-\gamma))}} \frac{1}{|h|^{N-\gamma}} dh \leq \sigma_{N-1} |B_R| \frac{1}{|\gamma|} \lambda^{2\gamma/(s(p-\gamma))},$$

where σ_{N-1} is the surface area of \mathbb{S}^{N-1} . Hence in this case,

$$\begin{aligned} [|x - y|^{(p-\gamma)s/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)} &\leq \left(\sigma_{N-1} |B_R| \frac{1}{|\gamma|} \right)^{s/2} \int_0^{(2R)^{(p-\gamma)s/2}} \lambda^{\gamma/(p-\gamma)} d\lambda \\ &= \left(1 - \frac{\gamma}{p}\right) \left(\sigma_{N-1} |B_R| \frac{1}{|\gamma|} \right)^{s/2} (2R)^{ps/2}. \end{aligned}$$

(Here we used $\gamma/(p - \gamma) = -1/(1 - \gamma/p) \in (-1, 0)$ whenever $\gamma < 0$.) This proves (4-4) when $\gamma < 0$. Next, suppose $\gamma = 0$. Then

$$\begin{aligned} [|x - y|^{(p-\gamma)s/2}]_{L^{2/s,1}(B_R \times B_R, \nu_\gamma)} &= \int_0^{(2R)^{ps/2}} \nu_0\{(x, y) \in B_R \times B_R : |x - y|^{ps/2} > \lambda\}^{s/2} d\lambda \\ &\leq \int_0^{(2R)^{ps/2}} \left(|B_R| \int_{\lambda^{2/(sp)} \leq |h| \leq 2R} \frac{1}{|h|^N} dh \right)^{s/2} d\lambda \\ &= \int_0^{(2R)^{ps/2}} \left(|B_R| \omega_{N-1} \frac{2}{ps} \log \left(\frac{(2R)^{ps/2}}{\lambda} \right) \right)^{s/2} d\lambda \\ &= (2R)^{ps/2} \int_0^1 \left(|B_R| \omega_{N-1} \frac{2}{ps} \log \left(\frac{1}{\lambda} \right) \right)^{s/2} d\lambda, \end{aligned}$$

which shows (4-4) remains valid when $\gamma = 0$ by the dominated convergence theorem. □

5. Finiteness of $\nu_0(E_{\lambda,0}[u])$ and the Lipschitz norm

In this section we prove Theorem 1.5, which we put in the following more precise form.

Proposition 5.1. *Let u be locally integrable on \mathbb{R}^N and $\nabla u \in L^1_{loc}(\mathbb{R}^N)$. Then*

$$\nu_0(E_{\lambda,0}[u]) = \begin{cases} 0 & \text{if } \lambda > \|\nabla u\|_\infty, \\ \infty & \text{if } \lambda < \|\nabla u\|_\infty. \end{cases}$$

Proof. First assume $\nabla u \in L^\infty$ and $\lambda > \|\nabla u\|_\infty$. Then for every $h \in \mathbb{R}^N$ we have $|u(x+h) - u(x)|/|h| \leq \lambda$ for almost every $x \in \mathbb{R}^N$. This immediately implies $\nu_0(E_{\lambda,0}[u]) = 0$.

For the more substantial part assume $\lambda < \|\nabla u\|_\infty$, where $\|\nabla u\|_\infty$ may be finite or infinite. We need to show that $\nu_0(E_{\lambda,0}[u]) = \infty$. We pick λ_1, λ_2 such that

$$\lambda < \lambda_1 < \lambda_2 < \|\nabla u\|_\infty.$$

Let $B_R = \{x \in \mathbb{R}^N : |x| < R\}$ and assume that $R > 1$ is so large that $\|\nabla u\|_{L^\infty(B_R)} > \lambda_2$. Let $\chi \in C_c^\infty$ such that $\chi(x) = 1$ in a neighborhood of \bar{B}_{2R} and set $u_\circ = \chi u$. Then $\nabla u_\circ = \nabla u$ as integrable functions on B_{2R} . There is a measurable set $F_0 \subset B_R$ of positive measure such that $|\nabla u(x)| > \lambda_2$ for all $x \in F_0$.

Fix $0 < \varepsilon \ll 1 - \lambda_1/\lambda_2$. We now consider the set \mathfrak{S}_ε of all spherical balls $S \subset \mathbb{S}^{N-1}$ with positive radius and the property that $\langle \theta_1, \theta_2 \rangle > 1 - \varepsilon$ for all $\theta_1, \theta_2 \in S$. By pigeonholing there exists a spherical ball $S \in \mathfrak{S}_\varepsilon$ and a Lebesgue measurable subset $F \subset F_0$ such that $\mathcal{L}^N(F) > 0$ and $\nabla u(x)/|\nabla u(x)| \in S$ for all $x \in F$. For the remainder of the argument we fix this spherical ball S ; we denote by $\sigma(S)$ its spherical measure.

We first note that, for $|h| \leq 1$ and for almost every $|x| \leq R$,

$$\frac{u(x+h) - u(x)}{|h|} = \frac{u_\circ(x+h) - u_\circ(x)}{|h|} = \left\langle \frac{h}{|h|}, \int_0^1 \nabla u_\circ(x+sh) ds \right\rangle. \tag{5-1}$$

Secondly since the translation operator is continuous in the strong operator topology of L^1 we see that there exists $\delta_0 < 1$ such that

$$\|\nabla u_\circ(\cdot + w) - \nabla u_\circ\|_{L^1(\mathbb{R}^N)} < \frac{\mathcal{L}^N(F)(\lambda_1 - \lambda)}{10} \quad \text{for } |w| \leq \delta_0. \tag{5-2}$$

In what follows we let $\delta \ll \delta_0$ and set

$$S(\delta, \delta_0) = \left\{ h \in \mathbb{R}^N : \delta \leq |h| \leq \delta_0, \frac{h}{|h|} \in S \right\}.$$

Let

$$\mathcal{E}_0 = \left\{ (x, h) : x \in F, h \in S(\delta, \delta_0), \frac{|u(x+h) - u(x)|}{|h|} > \lambda \right\}$$

so that $(x, h) \in \mathcal{E}_0$ implies $(x, x+h) \in E_{\lambda,0}[u]$. We then have by (5-1)

$$\begin{aligned} v_0(E_{\lambda,0}[u]) &\geq v_0(\mathcal{E}_0) = v_0\left(\left\{ (x, h) : x \in F, h \in S(\delta, \delta_0), \left| \left\langle \frac{h}{|h|}, \int_0^1 \nabla u_\circ(x+sh) \, ds \right\rangle \right| > \lambda \right\}\right) \\ &\geq v_0(\mathcal{E}_1) - v_0(\mathcal{E}_2), \end{aligned} \tag{5-3}$$

where

$$\begin{aligned} \mathcal{E}_1 &= \left\{ (x, h) : x \in F, h \in S(\delta, \delta_0), \left| \left\langle \frac{h}{|h|}, \nabla u_\circ(x) \right\rangle \right| > \lambda_1 \right\}, \\ \mathcal{E}_2 &= \left\{ (x, h) : x \in F, h \in S(\delta, \delta_0), \int_0^1 |\nabla u_\circ(x+sh) - \nabla u_\circ(x)| \, ds > \lambda_1 - \lambda \right\}. \end{aligned}$$

Indeed, if $(x, h) \notin \mathcal{E}_0 \cup \mathcal{E}_2$ then

$$\left| \left\langle \frac{h}{|h|}, \nabla u_\circ(x) \right\rangle \right| \leq \left| \left\langle \frac{h}{|h|}, \int_0^1 \nabla u_\circ(x+sh) \, ds \right\rangle \right| + \int_0^1 |\nabla u_\circ(x+sh) - \nabla u_\circ(x)| \, ds,$$

which is then $\leq \lambda_1$, so $(x, h) \notin \mathcal{E}_1$, establishing $\mathcal{E}_1 \subset \mathcal{E}_0 \cup \mathcal{E}_2$ and thus (5-3).

The set \mathcal{E}_1 does not change if we replace u_\circ by u in its definition. Since

$$\left\langle \frac{h}{|h|}, \nabla u(x) \right\rangle \geq (1 - \varepsilon)|\nabla u(x)| > (1 - \varepsilon)\lambda_2 > \lambda_1 \quad \text{for } x \in F, \frac{h}{|h|} \in S,$$

we get

$$v_0(\mathcal{E}_1) \geq \int_F dx \int_{S(\delta, \delta_0)} \frac{dh}{|h|^N} = \mathcal{L}^N(F)\sigma(S) \log\left(\frac{\delta_0}{\delta}\right).$$

Moreover, using (5-2) and Chebyshev's inequality we see that

$$\begin{aligned} v_0(\mathcal{E}_2) &\leq \int_{S(\delta, \delta_0)} \frac{\int_0^1 \|\nabla u_\circ(\cdot + sh) - \nabla u_\circ\|_{L^1(\mathbb{R}^N)} \, ds}{\lambda_1 - \lambda} \frac{dh}{|h|^N} \\ &\leq \int_{S(\delta, \delta_0)} \frac{\mathcal{L}^N(F)(\lambda_1 - \lambda)/10}{\lambda_1 - \lambda} \frac{dh}{|h|^N} = \frac{\mathcal{L}^N(F)}{10} \sigma(S) \log\left(\frac{\delta_0}{\delta}\right), \end{aligned}$$

and hence putting pieces together we obtain for $\delta < \delta_0$

$$v_0(E_{\lambda,0}[u]) \geq v_0(\mathcal{E}_1) - v_0(\mathcal{E}_2) > \frac{\mathcal{L}^N(F)}{2} \sigma(S) \log\left(\frac{\delta_0}{\delta}\right).$$

Here $\delta < \delta_0$ was arbitrary and by letting $\delta \rightarrow 0$ we conclude that $v_0(E_{\lambda,0}[u]) = \infty$. □

We now give a more precise version of [Example 1.7](#).

Lemma 5.2. *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain with Lipschitz boundary and let $u = \mathbb{1}_\Omega$. Then $u \in \text{BV}(\mathbb{R}^N) \setminus \dot{W}^{1,1}(\mathbb{R}^N)$, with*

$$v_0(E_{\lambda,0}[u]) \leq C_\Omega \times \begin{cases} \log(2/\lambda) & \text{if } \lambda \leq 1, \\ \lambda^{-1} & \text{if } \lambda > 1; \end{cases}$$

in particular we have $\sup_{\lambda>0} \lambda v_0(E_{\lambda,0}[u]) < \infty$.

Proof. Let

$$E(r, \lambda) = \{(x, y) \in E_{\lambda,0}[u] : r \leq |x - y| \leq 2r\}.$$

We begin with the observation that $r\lambda \leq 2$ if $v_0(E(r, \lambda)) > 0$. Furthermore, if $(x, y) \in E(r, \lambda)$ for some $y \in \mathbb{R}^N$, then x belongs to the $2r$ -neighborhood of $\partial\Omega$. The Lebesgue measure of such a neighborhood is $O(r)$ if $r \leq r_0$, where r_0 is some positive constant depending on Ω (because the boundary of a bounded Lipschitz domain can be covered by finitely many Lipschitz graphs, and the $2r$ -neighborhood of such graphs can be approximated by a union of $O(r)$ neighborhoods of suitable hyperplanes). Hence for $r \leq r_0$ we have $v_0(E(r, \lambda)) \leq Cr$ if $r \leq 2/\lambda$ and $v_0(E(r, \lambda)) = 0$ if $r > 2/\lambda$. As a result, if $2/\lambda \leq r_0$ we get

$$v_0(E_{\lambda,0}[u]) \leq \sum_{j \in \mathbb{Z}: 2^j \leq 2/\lambda} v_0(E(2^j, \lambda)) \lesssim \lambda^{-1}$$

and if $2/\lambda > r_0$ we get

$$v_0(E_{\lambda,0}[u]) \leq \sum_{j \in \mathbb{Z}: 2^j \leq r_0} v_0(E(2^j, \lambda)) + 2 \int_\Omega \int_{r_0 \leq |x-y| \leq 2/\lambda} \frac{dy}{|x-y|^N} dx \lesssim 1 + \log(\lambda^{-1}). \quad \square$$

6. When the upper bound (1-15) fails

In this section we make various constructions demonstrating the failure of (1-15) in the range $-1 \leq \gamma < 0$, and give the proof of [Theorem 1.8](#). We first establish:

Proposition 6.1. *Suppose $N \geq 1$ and $-1 \leq \gamma < 0$.*

(i) *For every $m > 0$, there exists $u \in C_c^\infty(\mathbb{R}^N)$ such that*

$$v_\gamma(E_{1,\gamma}[u]) > m \|\nabla u\|_{L^1(\mathbb{R}^N)}. \tag{6-1}$$

(ii) *There exists $C = C(N, \gamma) > 0$ and $p_0 = p_0(N, \gamma) > 1$ such that, for all $1 < p < p_0$,*

$$\sup_{\substack{u \in C_c^\infty(\mathbb{R}^N) \\ \|\nabla u\|_{L^p} \leq 1}} v_\gamma(E_{1,\gamma/p}[u]) \geq C \frac{p}{p-1}. \tag{6-2}$$

6A. Proof of Proposition 6.1: the case $\gamma = -1$. Here we may choose, for $m > 1$,

$$v_m = 2 \eta_m * \mathbb{1}_{B_1} \in C_c^\infty(\mathbb{R}^N), \tag{6-3}$$

where $\eta_m(x) := 2^{mN} \eta(2^m x)$ for some nonnegative, radially decreasing $\eta \in C_c^\infty(B_1)$, with $\int_{\mathbb{R}^N} \eta = 1$. Then when $1 \leq p < \infty$ and $m \leq p' = p/(p - 1)$ (which is no restriction on m if $p = 1$), we have $\|\nabla v_m\|_p \lesssim 2^{m/p'} \lesssim 1$, while $E_{1,-1/p}[v_m] \supseteq \{|x| \leq 1 - 2^{-m}, 1 + 2^{-m} \leq |y| \leq 2\}$ (because for (x, y) in the latter set, $|v_m(x) - v_m(y)| = 2$ and $|x - y|^{1-1/p} \leq 2^{1-1/p}$, which means $|Q_{-1/p} v_m(x, y)| \geq 2/2^{1-1/p} = 2^{1/p} > 1$). Hence

$$\begin{aligned} v_{-1}(E_{1,-1/p}[v_m]) &\geq \int_{|x| \leq 1 - 2^{-m}} \int_{1 + 2^{-m} \leq |y| \leq 2} |x - y|^{-1-N} \, dx \, dy \\ &\geq c_N \int_{|x| \leq 1 - 2^{-m}} (1 + 2^{-m} - |x|)^{-1} - (2 - |x|)^{-1} \, dx \geq c'_N m. \end{aligned}$$

This proves both (i) and (ii) of Proposition 6.1 in the case $\gamma = -1$. □

6B. The case $-1 < \gamma < 0$: examples of Cantor–Lebesgue-type on the real line. We now discuss some examples related to self-similar Cantor sets of dimension $\beta = 1 + \gamma$. Recall the definition of v_γ, Q_γ in (1-4), (1-5) and observe the behavior under dilations:

$$v_\gamma(tE) = t^{1+\gamma} v_\gamma(E). \tag{6-4}$$

We have:

Lemma 6.2. *Let $-1 < \gamma < 0$. There exist constants $c_\gamma > 0, C_\gamma > 0$, and a sequence of functions $g_m \in C^\infty(\mathbb{R})$, with $g_m(x) = 0$ for $x \leq 0$ and $g_m(x) = 1$ for $x \geq 1$, such that, for all $1 \leq p < \infty$,*

$$\|g'_m\|_p \leq c_\gamma 2^{m|\gamma|/(1+\gamma)(1-1/p)} \tag{6-5}$$

and if

$$m - 1 \leq \frac{\gamma + 1}{|\gamma|} \frac{p}{p - 1},$$

then

$$v_\gamma\left(\left\{(x, y) \in [0, 1]^2 : |Q_{\gamma/p} g_m(x, y)| > \frac{1}{4}\right\}\right) \geq \frac{m}{C_\gamma}. \tag{6-6}$$

Proof. For $-1 < \gamma < 0$ let

$$\rho = 2^{-1/(1+\gamma)} \tag{6-7}$$

so that $0 < \rho < \frac{1}{2}$. We construct g_m such that its derivative is supported on the m -th step of the construction of symmetric Cantor sets of dimension $\beta = 1 + \gamma = \log 2 / \log(1/\rho)$, with an equal variation on each of its 2^m components [Mattila 2015, Chapter 8.1].

Let $g_0 \in C^\infty(\mathbb{R})$ be such that $0 \leq g_0 \leq 1, g_0(x) = 0$ for $x \leq \rho$ and $g_0(x) = 1$ for $x \geq 1 - \rho$. Set, for $m \in \mathbb{N}$,

$$g_{m+1}(x) := \frac{1}{2} g_m\left(\frac{x}{\rho}\right) + \frac{1}{2} g_m\left(1 - \frac{1-x}{\rho}\right).$$

Since $\rho < \frac{1}{2}$, we have, for $p \in [1, \infty)$, $\|g'_{m+1}\|_{L^p(\mathbb{R})}^p = 2 \times (2\rho)^{-p} \rho \|g'_m\|_{L^p(\mathbb{R})}^p$, and thus

$$\|g'_m\|_{L^p(\mathbb{R})} = (2\rho)^{(1/p-1)m} \|g'_0\|_{L^p(\mathbb{R})} = 2^{(1-1/p)m|\gamma|/(\gamma+1)} \|g'_0\|_{L^p(\mathbb{R})}.$$

Fix now $1 \leq p < \infty$, and for $m \in \mathbb{N}$, $\lambda > 0$ define

$$A_{m,\lambda} := v_\gamma(\{(x, y) \in [0, 1]^2 : |\mathcal{Q}_{\gamma/p} g_m(x, y)| > \lambda\}).$$

Our goal is to estimate $A_{m,1/4}$, which we do by deriving a recursive estimate for $A_{m,\lambda}$. We have the decomposition

$$\begin{aligned} A_{m+1,\lambda} &\geq v_\gamma(\{(x, y) \in [0, \rho]^2 : |\mathcal{Q}_{\gamma/p} g_{m+1}(x, y)| > \lambda\}) \\ &\quad + v_\gamma(\{(x, y) \in [1 - \rho, 1]^2 : |\mathcal{Q}_{\gamma/p} g_{m+1}(x, y)| > \lambda\}) \\ &\quad + v_\gamma(\{(x, y) \in [0, \rho] \times [1 - \rho, 1] : |\mathcal{Q}_{\gamma/p} g_{m+1}(x, y)| > \lambda\}). \end{aligned} \tag{6-8}$$

Using the definition of g_{m+1} , (6-7) and (6-4), we compute the first term in the right-hand side of (6-8) as

$$\begin{aligned} v_\gamma(\{(x, y) \in [0, \rho]^2 : |\mathcal{Q}_\gamma g_{m+1}(x, y)| > \lambda\}) &= v_\gamma(\{(\rho w, \rho z) : (w, z) \in [0, 1]^2, |\mathcal{Q}_\gamma g_m(w, z)| > 2\rho^{1+\gamma/p}\lambda\}) \\ &= \rho^{\gamma+1} v_\gamma(\{(w, z) \in [0, 1]^2 : |\mathcal{Q}_\gamma g_m(w, z)| > 2^{|\gamma|/(p'(\gamma+1))}\lambda\}) = \frac{1}{2} A_{m,s\lambda}, \end{aligned} \tag{6-9}$$

where $s := 2\rho^{1+\gamma/p} = 2^{|\gamma|/(p'(\gamma+1))}$, and similarly the second term as

$$v_\gamma(\{(x, y) \in [1 - \rho, 1]^2 : |\mathcal{Q}_\gamma g_{m+1}(x, y)| > \frac{1}{2}\lambda\}) = \frac{1}{2} A_{m,s\lambda}. \tag{6-10}$$

Thus

$$A_{m+1,\lambda} \geq A_{m,s\lambda} + v_\gamma(\{(x, y) \in [0, \rho] \times [1 - \rho, 1] : |\mathcal{Q}_{\gamma/p} g_{m+1}(x, y)| > \lambda\}),$$

which iterates to give

$$A_{m,1/4} \geq A_{0,s^m/4} + \sum_{j=1}^m v_\gamma(\{(x, y) \in [0, \rho] \times [1 - \rho, 1] : |\mathcal{Q}_{\gamma/p} g_j(x, y)| > \frac{1}{4}s^{m-j}\lambda\}).$$

We drop the first term, and note that as long as

$$m - 1 \leq \frac{\gamma + 1}{|\gamma|} \frac{p}{p - 1},$$

we have $\frac{1}{4}s^{m-j} \leq \frac{1}{2}$ for all $j = 1, \dots, m$. Moreover, for every $x \in [0, \rho^2] \times [1 - \rho^2, 1]$ and every $j \geq 1$, we have $g_j(x) \leq \frac{1}{4}$ and $g_j(y) \geq \frac{3}{4}$, so $|\mathcal{Q}_{\gamma/p} g_j(x, y)| > \frac{1}{2}$. Thus we obtain the desired conclusion

$$A_{m,1/4} \geq m v_\gamma([0, \rho^2] \times [1 - \rho^2, 1]) = \frac{m}{C_\gamma}. \quad \square$$

6C. Conclusion of the proof of Proposition 6.1. We continue with the case $-1 < \gamma < 0$. Let $\eta_1 \in C_c^\infty(\mathbb{R})$ supported in $(-1, 2)$ such that $\eta_1(s) = 1$ on $(-\frac{1}{2}, \frac{3}{2})$ and $0 \leq \eta_1(s) \leq 1$ for all $s \in \mathbb{R}$.

We split $x = (x_1, x')$ with $x' \in \mathbb{R}^{N-1}$, where the variable x' should simply be dropped in the case $N = 1$. Set $\eta(x) = \prod_{i=1}^N \eta_1(x_i)$ and define

$$u_m(x_1, x') = 16g_m(x_1)\eta(x), \tag{6-11}$$

where g_m is as in Lemma 6.2. Then $u_m \in C_c^\infty(\mathbb{R}^N)$, and if $1 \leq p < \infty$ and

$$m - 1 \leq \frac{\gamma + 1}{|\gamma|} \frac{p}{p - 1},$$

we have $\|\nabla u_m\|_p \lesssim 1$. Both parts of [Proposition 6.1](#) will follow, if we can prove that under the same hypotheses on p and m , we have

$$v_\gamma(E_{1,\gamma/p}[u_m]) \geq c(N, \gamma)m - C(N, \gamma)^p. \tag{6-12}$$

We aim to reduce to the one-dimensional situation in [Lemma 6.2](#) and split $Q_{\gamma/p}u_m(x, y)$ as

$$Q_{\gamma/p}u_m(x, y) = 16\eta(x) \frac{g_m(x_1) - g_m(y_1)}{|x - y|^{1+\gamma/p}} + 16g_m(y_1) \frac{\eta(x) - \eta(y)}{|x - y|^{1+\gamma/p}} = I_m(x, y) + II_m(x, y),$$

so that

$$\begin{aligned} v_\gamma(E_{1,\gamma/p}[u_m]) &\geq \iint_{\substack{x_1, y_1 \in [0, 1] \\ |I_m(x, y) + II_m(x, y)| > 1}} |x - y|^{\gamma-N} dx dy \\ &\geq \iint_{\substack{x \in [0, 1]^N, y_1 \in [0, 1] \\ |x_1 - y_1| \geq |x' - y'| \\ |I_m(x, y)| > 2}} |x - y|^{\gamma-N} dx dy - \iint_{|II_m(x, y)| > 1} |x - y|^{\gamma-N} dx dy. \end{aligned} \tag{6-13}$$

Clearly if B_2 is the ball in \mathbb{R}^N of radius 2 centered at the origin then

$$|II_m(x, y)| \leq c_N |x - y|^{-\gamma/p} (\mathbb{1}_{B_2}(x) + \mathbb{1}_{B_2}(y)),$$

and it follows immediately (since $-\gamma > 0$) that

$$\iint_{|II_m(x, y)| > 1} |x - y|^{\gamma-N} dx dy \leq |\gamma|^{-1} C(N)^p.$$

For the first term in (6-13), we prove a lower bound and estimate by integrating in y'

$$\begin{aligned} \iint_{\substack{x \in [0, 1]^N, y_1 \in [0, 1] \\ |x_1 - y_1| \geq |x' - y'| \\ |I_m(x, y)| > 2}} |x - y|^{\gamma-N} dx dy &\geq \iint_{\substack{x \in [0, 1]^N, y_1 \in [0, 1] \\ |x_1 - y_1| \geq |x' - y'| \\ |16g_m(x_1) - 16g_m(y_1)|/|x_1 - y_1|^{1+\gamma/p} > 4}} |x - y|^{\gamma-N} dx dy \\ &\geq c_N \iint_{\substack{x_1, y_1 \in [0, 1] \\ |Q_{\gamma/p}g_m(x_1, y_1)| > 1/4}} |x_1 - y_1|^{\gamma-1} dx_1 dy_1, \end{aligned}$$

but by [Lemma 6.2](#) the last expression is bounded below for large m by $c_N m / C_\gamma$ under our hypothesis on m . This concludes the proof of (6-12). \square

For later purposes, note the inequality (6-13) (with $p = 1$) and the argument that follows proved also that for all sufficiently large $m > m(N, \gamma)$, we have

$$v_\gamma(E_{1,\gamma}[u_m] \cap ([0, 1] \times \mathbb{R}^{N-1})^2) \geq c(N, \gamma)m. \tag{6-14}$$

6D. Examples related to Theorems 1.1 and 1.8. We now consider the limit (1-8) in the range $-1 \leq \gamma < 0$ and provide counterexamples for cases where u is no longer required to be a C_c^∞ function. The following proposition covers part (i) of [Theorem 1.8](#).

Proposition 6.3. *Let $-1 \leq \gamma < 0$. Let $s \mapsto \omega(s)$ be any decreasing function on $[0, \infty)$, with $\omega(0) \leq 1$ and $\omega(s) > 0$ for all $s \geq 0$. Then there exists a C^∞ function $u \in \dot{W}^{1,1}(\mathbb{R}^N)$ such that*

$$|u(x)| \leq C\omega(|x|) \quad \text{for all } x \in \mathbb{R}^N \tag{6-15}$$

and

$$\lim_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \infty. \tag{6-16}$$

Proof. We consider the case $-1 < \gamma < 0$. Let $u_m \in C_c^\infty(\mathbb{R}^N)$ be as in (6-11) and define

$$f_m(x) = u_m(x_1 - 2, x') \tag{6-17}$$

so that $f_m(x) = 0$ if $x_1 \notin [1, 4]$. Let, for $n \in \mathbb{N}$,

$$R_n = 2^{2n}, \quad \lambda_n = R_n^{-(N+\gamma)} \omega(R_{n+1}), \quad m(n) \geq 4 \frac{\lambda_n}{\lambda_{n+1}} \omega(R_{n+1})^{-1} n^3. \tag{6-18}$$

We also assume $m(n) > m(N, \gamma)$ so that by (6-14) in Section 6C,

$$\nu_\gamma(\{(x, y) : x_1, y_1 \in [2, 3], |Q_\gamma f_{m(n)}(x, y)| > 1\}) \geq c(N, \gamma)m(n) \tag{6-19}$$

for all $n \in \mathbb{N}$. Finally let

$$u(x) = \sum_{n=2}^\infty \frac{\omega(R_{n+1})}{R_n^{N-1} n^2} f_{m(n)}\left(\frac{x}{R_n}\right). \tag{6-20}$$

Since $\|f_m\|_{\dot{W}^{1,1}} \leq C$, and ω is bounded, it is easy to see that the sum converges in $\dot{W}^{1,1}(\mathbb{R}^N)$, and that u is in $\dot{W}^{1,1}(\mathbb{R}^N)$. Also, the supports of $f_{m(n)}(R_n^{-1} \cdot)$, namely $[R_n, 4R_n] \times [-4R_n, 4R_n]^{N-1}$, are disjoint as n varies, so clearly $u \in C^\infty(\mathbb{R}^N)$. Since $\|f_m\|_{L^\infty} \leq C$, we have

$$|u(x)| \leq \omega(R_{n+1}) R_n^{-(N-1)} n^{-2} \quad \text{for } |x| \geq R_n,$$

so $|u(x)| \leq C'|x|^{-N+1} \omega(|x|)$ for $|x| \geq 2$. In particular $|u(x)| \leq C\omega(|x|)$.

For $\lambda \in ((n+1)^{-2} \lambda_{n+1}, n^{-2} \lambda_n]$ we estimate

$$\lambda \nu_\gamma(E_{\lambda, \gamma}[u]) \geq (n+1)^{-2} \lambda_{n+1} \nu_\gamma(E_{n^{-2} \lambda_n, \gamma}[u]) \geq \frac{\lambda_{n+1}}{4 \lambda_n} n^{-2} \lambda_n \nu_\gamma(\mathcal{E}_n),$$

where $\mathcal{E}_n := E_{n^{-2} \lambda_n, \gamma}[u] \cap ([2R_n, 3R_n] \times \mathbb{R}^{N-1})^2$. Moreover, for $(x, y) \in ([2R_n, 3R_n] \times \mathbb{R}^{N-1})^2$, we have

$$u(x) - u(y) = R_n^{1-N} n^{-2} \omega(R_{n+1}) (f_{m(n)}(R_n^{-1} x) - f_{m(n)}(R_n^{-1} y)),$$

so

$$|Q_\gamma u(x, y)| > n^{-2} \lambda_n \iff \frac{|f_{m(n)}(R_n^{-1} x) - f_{m(n)}(R_n^{-1} y)|}{|R_n^{-1} x - R_n^{-1} y|^{1+\gamma}} > \frac{R_n^{N+\gamma}}{\omega(R_{n+1})} \lambda_n = 1,$$

where the last equality follows from (6-18). Hence rescaling using (6-4) yields

$$\begin{aligned} n^{-2} \lambda_n \nu_\gamma(\mathcal{E}_n) &= n^{-2} \lambda_n R_n^{\gamma+N} \nu_\gamma(\{(x, y) : x_1, y_1 \in [2, 3], |Q_\gamma f_{m(n)}(x, y)| > 1\}) \\ &\geq c(N, \gamma) m(n) \omega(R_{n+1}) n^{-2}, \end{aligned} \tag{6-21}$$

with $c(N, \gamma) > 0$, by (6-19). Thus we have shown

$$\inf_{\lambda \in ((n+1)^{-2} \lambda_{n+1}, n^{-2} \lambda_n]} \lambda \nu_\gamma(E_{\lambda, \gamma}[u]) \geq c(N, \gamma) \frac{\lambda_{n+1}}{4 \lambda_n} \omega(R_{n+1}) m(n) n^{-2} \geq c(N, \gamma) n,$$

where for the last inequality we have used our assumption (6-18) on $m(n)$. The assertion follows for $-1 < \gamma < 0$.

Finally consider the case $\gamma = -1$. We now choose v_m as in (6-3) and

$$R_n = 2^{2n}, \quad \lambda_n = R_n^{-(N-1)}\omega(R_{n+1}), \quad m(n) \geq 4 \frac{\lambda_n}{\lambda_{n+1}} \frac{n^3}{\omega(R_{n+1})}. \tag{6-22}$$

In analogy to (6-20) we now use

$$u(x) = \sum_{n=2}^{\infty} \frac{\omega(R_{n+1})}{R_n^{N-1}n^2} v_{m(n)}\left(\frac{x}{R_n}\right). \tag{6-23}$$

Since ω is bounded it is immediate that $u \in \dot{W}^{1,1}(\mathbb{R}^N)$ and also that $|u(x)| \lesssim \omega(|x|)$. We need to check that $\lambda v_{-1}(E_{\lambda,-1}[u]) \rightarrow \infty$ as $\lambda \rightarrow 0^+$. If $|x| \leq R_n(1 - 2^{m(n)})$ and $|y| \geq R_n(1 + 2^{m(n)})$, then

$$u(x) - u(y) \geq \frac{\omega(R_{n+1})}{R_n^{N-1}n^2} v_{m(n)}\left(\frac{x}{R_n}\right) = 2 \frac{\omega(R_{n+1})}{R_n^{N-1}n^2} = 2n^{-2}\lambda_n > n^{-2}\lambda_n,$$

so $(x, y) \in E_{n^{-2}\lambda_n,-1}[u]$. Hence we get

$$\begin{aligned} n^{-2}\lambda_n v_{-1}(E_{n^{-2}\lambda_n,-1}[u]) &\geq n^{-2}\lambda_n \iint_{\substack{|x| \leq R_n(1-2^{m(n)}) \\ |y| \geq R_n(1+2^{m(n)})}} |x - y|^{-1-N} dx dy \\ &\geq n^{-2}\lambda_n R_n^{N-1} \iint_{\substack{|x| \leq 1-2^{m(n)} \\ |y| \geq 1+2^{m(n)}}} |x - y|^{-1-N} dx dy \\ &\geq c_N m(n) \omega(R_{n+1}) n^{-2} \end{aligned}$$

(using (6-22) in the last inequality). This, together with our assumption on $m(n)$, implies that

$$\inf_{\lambda \in ((n+1)^{-2}\lambda_{n+1}, n^{-2}\lambda_n]} \lambda v_{-1}(E_{\lambda,-1}[u]) \geq c_N n \rightarrow \infty$$

when $n \rightarrow \infty$, as desired. □

The next proposition is relevant for part (ii) of [Theorem 1.8](#).

Proposition 6.4. *Suppose $-1 \leq \gamma < 0$. Then there exists a compactly supported $u \in W^{1,1}(\mathbb{R}^N)$ such that u is C^∞ for $x \neq 0$,*

$$|u(x)| \leq \frac{C}{|x|^{N-1}[\log(2 + |x|^{-1})]^2} \tag{6-24}$$

and

$$\lim_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda,\gamma}[u]) = \infty. \tag{6-25}$$

If in addition $N \geq 2$ or $-1 < \gamma < 0$ there exists u with the above properties and

$$v_\gamma(E_{\lambda,\gamma}[u]) = \infty \quad \text{for all } \lambda > 0. \tag{6-26}$$

Proof. Consider first the case $-1 < \gamma < 0$. We choose for $n \in \mathbb{N}$

$$R_n = 2^{-2n}, \quad m(n) \geq 2^{2n}, \tag{6-27}$$

and with these choices of R_n and $m(n)$ and f_m as in (6-17) and (6-11) we define again

$$u(x) = \sum_{n=2}^{\infty} \frac{1}{n^2 R_n^{N-1}} f_{m(n)}\left(\frac{x}{R_n}\right).$$

The sum converges in $W^{1,1}$ to a function supported in $[-4, 4]^N$. We have $|u(x)| \leq C2^{2n(N-1)}n^{-2}$ for $0 < x_1 \leq 2^{-2n}$; moreover, $|x'| \lesssim |x_1|$ on the support of u . This implies $|u(x)| \leq C'[|x|^{1-N} \log(1/|x|)]^{-2}$ for small x . Also, because of the choices of R_n , we see that u is smooth away from 0.

Fix $\lambda > 0$. Since $\lim_{n \rightarrow \infty} R_n^{N+\gamma} n^2 = 0$, we may choose n_0 such that

$$\lambda R_n^{N+\gamma} n^2 \leq 1 \quad \text{for all } n \geq n_0. \tag{6-28}$$

Now $v_\gamma(E_{\lambda,\gamma}[u]) \geq v_\gamma(E_{\lambda,\gamma}[u] \cap ([2R_n, 3R_n] \times \mathbb{R}^{N-1})^2)$, and again $f_{m(n)}(R_n^{-1} \cdot)$ is supported in $\mathcal{R}(n) = [R_n, 4R_n] \times [-4R_n, 4R_n]^{N-1}$. Hence by the same rescaling argument as in (6-21), we obtain

$$v_\gamma(E_{\lambda,\gamma}[u]) \geq R_n^{N+\gamma} v_\gamma(\{(x, y) : x_1, y_1 \in [2, 3], |Q_\gamma f_{m(n)}(x, y)| > \lambda R_n^{N+\gamma} n^2\}).$$

If $n \geq n_0$ then this gives

$$\begin{aligned} v_\gamma(E_{\lambda,\gamma}[u]) &\geq R_n^{N+\gamma} v_\gamma(\{(x, y) : x_1, y_1 \in [2, 3], |Q_\gamma f_{m(n)}(x, y)| > 1\}) \\ &\geq c(N, \gamma)m(n)R_n^{N+\gamma} \end{aligned}$$

by (6-19). Since $\lim_{n \rightarrow \infty} m(n)R_n^{N+\gamma} = \infty$, by (6-27) we conclude $v_\gamma(E_{\lambda,\gamma}[u]) = \infty$.

For the case $\gamma = -1$ and $N \geq 2$, define u as in (6-23) but with the choice of the parameters $R_n, m(n)$ as in (6-27) to obtain a compactly supported $u \in W^{1,1}$ satisfying (6-24). We now fix $\lambda > 0$ and note that when $N \geq 2$ we have $\lambda R_n^{N-1} n^2 \rightarrow 0$ as $n \rightarrow \infty$. The above calculation gives $v_{-1}(E_{\lambda,-1}[u]) \geq c(N)m(n)R_n^{N-1}$ provided that $\lambda R_n^{N-1} n^2 \leq 1$ and thus the conclusion $v_{-1}(E_{\lambda,-1}[u]) = \infty$.

Finally, clearly (6-25) follows from (6-26), and the latter was proved if $-1 < \gamma < 0$ or $N \geq 2$. It remains to consider the case $N = 1, \gamma = -1$. We define u as in the previous paragraph. The above calculation shows that $v_{-1}(E_{\lambda,-1}[u]) \geq cm(n)$ provided that $\lambda < 1/n^2$ which establishes (6-25) in this last case. \square

The case $N = 1, \gamma = -1$ plays a special role. The following lemma shows that the conclusion (6-26) in Proposition 6.4 fails in this case.

Lemma 6.5. *Let $u \in \dot{W}^{1,1}(\mathbb{R})$ be compactly supported. Then $v_{-1}(E_{\lambda,-1}[u]) < \infty$ for all $\lambda > 0$.*

Proof. Let $u \in \dot{W}^{1,1}(\mathbb{R})$ be compactly supported in $[-R, R]$. Then given any $\lambda \in (0, 1)$, there exists $\delta(\lambda) > 0$ such that $\int_I |u'| \leq \lambda/2$ for every interval $I \subset \mathbb{R}$ with length $\leq \delta(\lambda)$. As a result, u is uniformly continuous on \mathbb{R} , with $\sup_{x \in \mathbb{R}} |u(x+h) - u(x)| \leq \lambda/2$ for $|h| \leq \delta(\lambda)$. Thus

$$\begin{aligned} v_{-1}(E_{\lambda,-1}[u]) &= 2 \int_{-\infty}^{\infty} \int_{\substack{h>0 \\ |u(x+h)-u(x)|>\lambda}} \frac{dh}{h^2} dx \\ &\leq \int_{-2R}^{2R} \int_{\delta(\lambda)}^{\infty} \frac{dh}{h^2} dx + \int_{\mathbb{R} \setminus [-2R, 2R]} \int_{|x|-R}^{|x|+R} \frac{dh}{h^2} dx \\ &\leq 4R(\delta(\lambda))^{-1} + 4. \end{aligned} \quad \square$$

6E. Generic failure in $W^{1,1}$ for the case $-1 \leq \gamma < 0$.

Proposition 6.6. *Let $-1 \leq \gamma < 0$, $N \geq 2$ or $-1 < \gamma < 0$, $N \geq 1$. Let*

$$\mathcal{V} = \{f \in W^{1,1}(\mathbb{R}^N) : v_\gamma(E_{\lambda,\gamma}[f]) < \infty \text{ for some } \lambda > 0.\} \tag{6-29}$$

Then \mathcal{V} is of first category in $W^{1,1}(\mathbb{R}^N)$, in the sense of Baire.

Let

$$U_k = \{(x, y) \in \mathbb{R}^{2N} : 2^{k-1} \leq |x - y| \leq 2^k\},$$

$$\Omega_\ell = \bigcup_{k=1-\ell}^\ell U_k. \tag{6-30}$$

For the proof of Proposition 6.6 we use an elementary estimate for the intersections $E_{\lambda,\gamma}[u] \cap \Omega_\ell$.

Lemma 6.7. *For all $\gamma \in \mathbb{R}$, $u \in W^{1,1}(\mathbb{R}^N)$, $\ell > 0$ and Ω_ℓ as in (6-30),*

$$\sup_{\lambda > 0} \lambda v_\gamma(E_{\lambda,\gamma}[u] \cap \Omega_\ell) \leq C(N, \gamma) \ell \|\nabla u\|_1.$$

Proof. For $u \in C^1$ we use the Lusin–Lipschitz inequality (2-2) to see that

$$\begin{aligned} \lambda \iint_{E_{\lambda,\gamma}[u] \cap U_k} |x - y|^{\gamma - N} dx dy &\leq C(\gamma) \lambda 2^{k\gamma} \mathcal{L}^N \{x \in \mathbb{R}^N : M(|\nabla u|)(x) > c 2^{k\gamma} \lambda\} \\ &\leq C(N, \gamma) \|\nabla u\|_1 \end{aligned}$$

by the Hardy–Littlewood maximal inequality. Now sum in $1 - \ell \leq k \leq \ell$. The extension to general $u \in W^{1,1}$ is obtained as in the limiting argument of Section 2C. □

Proof of Proposition 6.6. Let, for $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,

$$\mathcal{V}(m, j) = \{u \in W^{1,1}(\mathbb{R}^N) : v_\gamma(E_{\lambda,\gamma}[u]) \leq m \text{ for all } \lambda > 2^j\}.$$

Since $\lambda \mapsto v_\gamma(E_{\lambda,\gamma}[u])$ is decreasing, we see that \mathcal{V} is contained in $\bigcup_{m \geq 1} \bigcup_{j \in \mathbb{Z}} \mathcal{V}(m, j)$. To show that \mathcal{V} is of first category in $W^{1,1}(\mathbb{R}^N)$, we need to show that for every $m \in \mathbb{N}$, $j \in \mathbb{Z}$, the set $\mathcal{V}(m, j)$ is nowhere dense.

We first show that $\mathcal{V}(m, j)$ is closed in $W^{1,1}(\mathbb{R}^N)$. Let $u_n \in \mathcal{V}(m, j)$ and $u \in W^{1,1}(\mathbb{R}^N)$ such that $\lim_{n \rightarrow \infty} \|u - u_n\|_{W^{1,1}(\mathbb{R}^N)} = 0$. It suffices to show that given $\varepsilon > 0$, we have $v_\gamma(E_{\lambda,\gamma}[u]) \leq m + \varepsilon$ for all $\lambda > 2^j$. By the monotone convergence theorem, we have

$$\lim_{\ell \rightarrow \infty} v_\gamma(E_{\lambda,\gamma}[u] \cap \Omega_\ell) = v_\gamma(E_{\lambda,\gamma}[u]),$$

and it suffices to verify that

$$v_\gamma(E_{\lambda,\gamma}[u] \cap \Omega_\ell) \leq m + \varepsilon \quad \text{for } \lambda > 2^j, \tag{6-31}$$

for all $\ell \in \mathbb{N}$. Now let $\delta > 0$ such that $(1 - \delta)\lambda > 2^j$. Then

$$v_\gamma(E_{\lambda,\gamma}[u] \cap \Omega_\ell) \leq v_\gamma(E_{(1-\delta)\lambda,\gamma}[u_n] \cap \Omega_\ell) + v_\gamma(E_{\delta\lambda,\gamma}[u - u_n] \cap \Omega_\ell)$$

and using that $u_n \in \mathcal{V}(m, j)$ together with $(1 - \delta)\lambda > 2^j$, and [Lemma 6.7](#), we see that for $\lambda > 2^j$

$$v_\gamma(E_{\lambda,\gamma}[u] \cap \Omega_\ell) \leq m + C(N, \gamma)\ell \frac{1 + \delta}{\delta 2^j} \|\nabla(u_n - u)\|_1.$$

Since $\delta > 0$ was arbitrary and since $\|\nabla(u_n - u)\|_{L^1(\mathbb{R}^N)} \rightarrow 0$ by assumption, we obtain [\(6-31\)](#).

To show that the closed set $\mathcal{V}(m, j)$ is nowhere dense when $-1 \leq \gamma < 0$, we need to verify that for every $u \in \mathcal{V}(m, j)$ and $\varepsilon_1 > 0$ there exists $f \in W^{1,1}(\mathbb{R}^N)$ such that $\|f - u\|_{W^{1,1}(\mathbb{R}^N)} < \varepsilon_1$ and $f \notin \mathcal{V}(m, j)$. To see this we use [Proposition 6.4](#), according to which there exists a compactly supported $W^{1,1}$ function f_0 for which $v_\gamma(E_{\lambda,\gamma}[f_0]) = \infty$ for all $\lambda > 0$. It is then clear that

$$f = u + \frac{\varepsilon_1}{2} \frac{f_0}{\|f_0\|_{W^{1,1}}}$$

satisfies $\|f - u\|_{W^{1,1}} \leq \varepsilon_1/2$ and also,

$$v_\gamma(E_{\lambda,\gamma}[f]) \geq v_\gamma\left(E_{2\lambda,\gamma}\left[\frac{\varepsilon_1}{2} \frac{f_0}{\|f_0\|_{W^{1,1}}}\right]\right) - v_\gamma(E_{\lambda,\gamma}[u]) = \infty$$

for every $\lambda > 2^j$, for all $j \in \mathbb{Z}$. The proposition is proved. □

To include a result of generic failure of the limiting relation in the case $N = 1, \gamma = -1$ we give

Proposition 6.8. *Let $-1 \leq \gamma < 0$. Let*

$$\mathcal{W} = \left\{ f \in W^{1,1}(\mathbb{R}) : \limsup_{R \rightarrow 0} \sup_{\lambda > R} R v_\gamma(E_{\lambda,\gamma}[f]) < \infty \right\}.$$

Then \mathcal{W} is of first category in $W^{1,1}$, in the sense of Baire.

Proof. Clearly $\mathcal{W} \subset \mathcal{V}$, where \mathcal{V} is defined in [\(6-29\)](#). We define

$$\mathcal{W}(m, j) = \left\{ u \in W^{1,1}(\mathbb{R}) : \sup_{0 < R \leq 2^{-j}} \sup_{\lambda > R} R v_\gamma(E_{\lambda,\gamma}[u]) \leq m \right\}$$

and note that

$$\mathcal{W} \subset \bigcup_{j \geq 1} \bigcup_{m \geq 1} \mathcal{W}(m, j). \tag{6-32}$$

The arguments in the proof of [Proposition 6.6](#) that were used to show that the sets $\mathcal{V}(m, j)$ are closed in $W^{1,1}(\mathbb{R}^N)$ also show that the sets $\mathcal{W}(m, j)$ are closed in $W^{1,1}(\mathbb{R})$.

Let $u \in \mathcal{W}(m, j)$, and let $\varepsilon_1 > 0$. By [Proposition 6.4](#) there is $f_0 \in W^{1,1}(\mathbb{R})$ such that

$$\lim_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda,\gamma}[f_0]) = \infty.$$

We may normalize so that $\|f_0\|_{W^{1,1}(\mathbb{R})} = 1$. Pick $R \in (0, 2^{-j}]$ so that $\lambda v_\gamma(E_{\lambda,\gamma}[f_0]) > 16m/\varepsilon_1$ for $\lambda \leq 8R/\varepsilon_1$. Let $f = u + (\varepsilon_1/2)f_0$ so that $\|f - u\|_{W^{1,1}(\mathbb{R})} \leq \varepsilon_1/2$. Moreover if $\lambda = 2R$, then $\lambda > R$ and

$$\begin{aligned} R v_\gamma(E_{\lambda,\gamma}[f]) &\geq R v_\gamma\left(E_{2\lambda,\gamma}\left[\frac{\varepsilon_1}{2} f_0\right]\right) - R v_\gamma(E_{\lambda,\gamma}[u]) \\ &= \frac{\varepsilon_1}{8} \frac{8R}{\varepsilon_1} v_\gamma(E_{8R/\varepsilon_1,\gamma}[f_0]) - R v_\gamma(E_{\lambda,\gamma}[u]) > \frac{\varepsilon_1}{8} \frac{16m}{\varepsilon_1} - m = m, \end{aligned}$$

and we see that $f \notin \mathcal{W}(m, j)$. Thus we have shown that $\mathcal{W}(m, j)$ is nowhere dense in $W^{1,1}(\mathbb{R})$. By (6-32) the proof is concluded. \square

7. Perspectives and open problems

7A. Subspaces of $\dot{W}^{1,1}$ and $\dot{B}\dot{V}$ and related spaces. The failure of the upper bounds for $[Q_\gamma u]_{L^{1,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}$ for $\gamma \in [-1, 0)$ raises a number of interesting questions. Consider the space $\dot{B}\dot{V}(\gamma)$ consisting of all $B\dot{V}$ functions satisfying

$$\|u\|_{\dot{B}\dot{V}(\gamma)} := \|\nabla u\|_{\mathcal{M}} + \sup_{\lambda > 0} \lambda \nu_\gamma(E_{\lambda,\gamma}[u]) < \infty \tag{7-1}$$

and the corresponding subspace $\dot{W}^{1,1}(\gamma)$ of $\dot{W}^{1,1}$.

Embeddings. We proved in this paper that for $\gamma \notin [-1, 0]$ we have $\dot{B}\dot{V}(\gamma) = \dot{B}\dot{V}$ and $\dot{W}^{1,1}(\gamma) = \dot{W}^{1,1}$. It is natural to ask how in the range $-1 \leq \gamma < 0$ the proper subspaces $\dot{B}\dot{V}(\gamma)$ and $\dot{W}^{1,1}(\gamma)$ relate to other families of function spaces, in particular to the Hardy–Sobolev space $\dot{F}_{1,2}^1$, another subspace of $\dot{W}^{1,1}$.

Triangle inequalities. The spaces $\dot{W}^{1,1}(\gamma)$ and $\dot{B}\dot{V}(\gamma)$ are defined via $L^{1,\infty}$ -quasinorms, and the space $L^{1,\infty}$ is not normable (unlike $L^{p,\infty}$ for $1 < p < \infty$, which is normable [Hunt 1966]). However Theorem 1.4 tells us that $\dot{W}^{1,1}(\gamma)$ and $\dot{B}\dot{V}(\gamma)$ are normable for $\gamma \notin [-1, 0]$. Are these spaces normable in the range $\gamma \in [-1, 0)$?

Related quasinorms. Consider for $0 < s \leq 1$

$$\|u\|_{(p,s,\gamma)} = \left[\frac{u(x) - u(y)}{|x - y|^{\gamma/p+s}} \right]_{L^{p,\infty}(\mathbb{R}^{2N}, \nu_\gamma)}.$$

It is an obvious consequence of Theorem 1.3 that for $s = 1$ and fixed $p > 1$, these expressions define equivalent (semi/quasi)-norms on C_c^∞ as γ varies over $\mathbb{R} \setminus \{0\}$. It would be interesting to find a more direct proof of this observation which does not involve the relation with $\dot{W}^{1,p}$. We note that the equivalence for varying γ breaks down for $0 < s < 1$. This result, and more about the spaces for which $\|u\|_{(p,s,\gamma)} < \infty$ with $0 < s < 1$, such as their connection to Besov spaces and interpolation, can be found in [Domínguez et al. 2023].

7B. Other limit functionals. Our results, combined with the various developments presented in [Brezis and Nguyen 2018; 2020; Nguyen 2007; 2011], suggest several possible directions of research.

Can one prove a generalization of (1-14), (1-16) where the supremum is replaced by the $\liminf_{\lambda \rightarrow \infty}$ when $\gamma > 0$ and by a $\liminf_{\lambda \rightarrow 0^+}$ when $\gamma < 0$? More precisely, for $1 < p < \infty$, is there a positive constant $C(N, \gamma, p)$ such that, for all $u \in L^1_{\text{loc}}(\mathbb{R}^N)$,

$$\|\nabla u\|_{L^p}^p \leq C(N, \gamma, p) \liminf_{\lambda \rightarrow \infty} \lambda^p \nu_\gamma(E_{\lambda,\gamma/p}[u]) \quad \text{if } \gamma > 0, \tag{7-2a}$$

$$\|\nabla u\|_{L^p}^p \leq C(N, \gamma, p) \liminf_{\lambda \searrow 0} \lambda^p \nu_\gamma(E_{\lambda,\gamma/p}[u]) \quad \text{if } \gamma < 0, \tag{7-2b}$$

in the sense that $\|\nabla u\|_{L^p} = \infty$ if $u \in L^1_{\text{loc}} \setminus \dot{W}^{1,p}$?

For $p = 1$ we can also ask: is there a positive constant $C(N, \gamma)$ such that, for all $u \in L^1_{\text{loc}}(\mathbb{R}^N)$,

$$\|\nabla u\|_{\mathcal{M}} \leq C(N, \gamma) \liminf_{\lambda \rightarrow \infty} \lambda v_\gamma(E_{\lambda, \gamma}[u]) \quad \text{if } \gamma > 0, \tag{7-3a}$$

$$\|\nabla u\|_{\mathcal{M}} \leq C(N, \gamma) \liminf_{\lambda \searrow 0} \lambda v_\gamma(E_{\lambda, \gamma}[u]) \quad \text{if } \gamma < 0, \tag{7-3b}$$

in the sense that $\|\nabla u\|_{\mathcal{M}} = \infty$ if $u \in L^1_{\text{loc}} \setminus \mathring{\text{BV}}$?

Theorem 1.1 gives (7-2a) and (7-2b) if we additionally assume $u \in \dot{W}^{1,p}(\mathbb{R}^N)$. It also gives (7-3a) and (7-3b) if we additionally assume that $u \in \dot{W}^{1,1}(\mathbb{R}^N)$. It would already be interesting to establish (7-3a), (7-3b) for all $\mathring{\text{BV}}$ functions.

When $\gamma = -1, p = 1$, (7-3b) holds for all $u \in L^1_{\text{loc}}(\mathbb{R}^N)$ as established in [Nguyen 2008, Theorem 2] and [Brezis and Nguyen 2018, Section 3.4]. For $\gamma = -p, 1 < p < \infty$, inequality (7-2b) was proved in [Bourgain and Nguyen 2006]. For $\gamma = N$, Poliakovsky [2022] proved weaker versions of (7-2a) and (7-3a) where the \liminf is replaced by a \limsup .

7C. Γ -convergence. This is a far-reaching generalization of the questions raised in Section 7B. For fixed $p \geq 1$ and $\gamma \in \mathbb{R} \setminus \{0\}$ consider the functionals

$$\Phi_\lambda[u] := \lambda^p v_\gamma(E_{\lambda, \gamma/p}[u]), \quad \lambda \in (0, \infty),$$

defined for all $u \in L^1_{\text{loc}}(\mathbb{R}^N)$. It would be very interesting to study the Γ -limit of Φ_λ in $L^1_{\text{loc}}(\mathbb{R}^N)$, in the sense of De Giorgi, as $\lambda \rightarrow \infty$ when $\gamma > 0$ and as $\lambda \searrow 0$ when $\gamma < 0$. More specifically, if $p > 1$, define on $L^1_{\text{loc}}(\mathbb{R}^N)$

$$\Phi_{*,c}[u] = \begin{cases} c \|\nabla u\|_{L^p}^p & \text{if } u \in \dot{W}^{1,p}(\mathbb{R}^N), \\ \infty & \text{otherwise,} \end{cases}$$

and for $p = 1$ define

$$\Phi_{*,c}[u] = \begin{cases} c \|\nabla u\|_{\mathcal{M}} & \text{if } u \in \mathring{\text{BV}}(\mathbb{R}^N), \\ \infty & \text{otherwise.} \end{cases}$$

A challenging question is whether there exists a constant $c = c(p, \gamma, N) > 0$ such that $\Phi_\lambda \rightarrow \Phi_{*,c}$ in the sense of Γ -convergence, meaning

- (1) whenever $u_\lambda \rightarrow u$ in L^1_{loc} then $\liminf \Phi_\lambda[u_\lambda] \geq \Phi_{*,c}[u]$, and
- (2) for each $u \in L^1_{\text{loc}}(\mathbb{R}^N)$ there exist (v_λ) with $v_\lambda \in L^1_{\text{loc}}(\mathbb{R}^N)$, $v_\lambda \rightarrow u$ in L^1_{loc} and $\limsup \Phi_\lambda[v_\lambda] \leq \Phi_{*,c}[u]$.

This question is especially meaningful in the case $p = 1$ where the pointwise limit behaves somewhat pathologically. Indeed, recall that for $p = 1, -1 \leq \gamma < 0$ there is no universal upper bound for $\Phi_\lambda[u]$ in terms of $\|\nabla u\|_{L^1}$. Also when $p = 1$ and $\gamma \in \mathbb{R} \setminus [-1, 0]$ the examples in Section 3F show that the pointwise limit in $\dot{W}^{1,1}$ and on $\mathring{\text{BV}} \setminus \dot{W}^{1,1}$ may differ (by a multiplicative constant). A remarkable result of Nguyen [2007; 2011] states that $\Phi_\lambda \rightarrow \Phi_{*,c}$ as $\lambda \rightarrow 0$, in the sense of Γ -convergence, when $p \geq 1$, and $\gamma = -p$ for some appropriate constant $c = c(p, N)$; see also [Brezis and Nguyen 2020] (note, however, that $\dot{W}^{1,p}$ and $\mathring{\text{BV}}$ are replaced in these papers by $W^{1,p}$ and BV).

7D. More general families of functionals. Consider a monotone nondecreasing function $\varphi : [0, \infty) \rightarrow [0, \infty)$ and set (inspired by [Brezis and Nguyen 2018; 2020])

$$\Psi_\lambda[u] := \lambda^p \iint_{\mathbb{R}^N \times \mathbb{R}^N} \varphi\left(\frac{|u(x) - u(y)|}{\lambda|x - y|^{1+\gamma/p}}\right) |x - y|^{\gamma-N} dx dy.$$

The family Φ_λ in Section 7C corresponds to $\varphi = \mathbb{1}_{(1, \infty)}$. It is an interesting generalization of the above problems to study the limit of Ψ_λ as $\lambda \searrow 0$ when $\gamma < 0$ and the limit of Ψ_λ as $\lambda \rightarrow \infty$ when $\gamma > 0$, both in the sense of pointwise convergence or in the sense of Γ -convergence. A formal computation suggests that our Theorem 1.1 should go over modulo a factor $\int_0^\infty \varphi(s)/s^{p+1} ds$; see [Brezis and Nguyen 2020]. We refer to [Brezis and Nguyen 2018] for a further discussion of applications.

Acknowledgements

Seeger and Yung would like to thank the Hausdorff Research Institute of Mathematics and the organizers of the trimester program “Harmonic analysis and analytic number theory” for a pleasant working environment in the summer of 2021. The research was supported in part by NSF grants DMS-1764295, DMS-2054220 (Seeger) and by a Future Fellowship FT200100399 from the Australian Research Council (Yung). We thank the referee for a careful reading and numerous valuable suggestions.

References

- [Bourgain and Nguyen 2006] J. Bourgain and H.-M. Nguyen, “A new characterization of Sobolev spaces”, *C. R. Math. Acad. Sci. Paris* **343**:2 (2006), 75–80. [MR](#) [Zbl](#)
- [Bourgain et al. 2001] J. Bourgain, H. Brezis, and P. Mironescu, “Another look at Sobolev spaces”, pp. 439–455 in *Optimal control and partial differential equations*, edited by J. L. Menaldi et al., IOS, Amsterdam, 2001. [MR](#) [Zbl](#)
- [Brezis 2002] H. Brezis, “How to recognize constant functions: a connection with Sobolev spaces”, *Uspekhi Mat. Nauk* **57**:4(346) (2002), 59–74. In Russian; translated in *Russian Math. Surv.* **57**:4 (2002), 693–708. [MR](#) [Zbl](#)
- [Brezis 2011] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, 2011. [MR](#) [Zbl](#)
- [Brezis and Nguyen 2018] H. Brezis and H.-M. Nguyen, “Non-local functionals related to the total variation and connections with image processing”, *Ann. PDE* **4**:1 (2018), art. id. 9. [MR](#) [Zbl](#)
- [Brezis and Nguyen 2020] H. Brezis and H.-M. Nguyen, “Non-local, non-convex functionals converging to Sobolev norms”, *Nonlinear Anal.* **191** (2020), art. id. 111626. [MR](#) [Zbl](#)
- [Brezis et al. 2021a] H. Brezis, J. Van Schaftingen, and P.-L. Yung, “Going to Lorentz when fractional Sobolev, Gagliardo and Nirenberg estimates fail”, *Calc. Var. Partial Differential Equations* **60**:4 (2021), art. id. 129. [MR](#) [Zbl](#)
- [Brezis et al. 2021b] H. Brezis, J. Van Schaftingen, and P.-L. Yung, “A surprising formula for Sobolev norms”, *Proc. Natl. Acad. Sci. USA* **118**:8 (2021), art. id. e2025254118. [MR](#)
- [Dávila 2002] J. Dávila, “On an open question about functions of bounded variation”, *Calc. Var. Partial Differential Equations* **15**:4 (2002), 519–527. [MR](#) [Zbl](#)
- [Domínguez and Milman 2022] Ó. Domínguez and M. Milman, “New Brezis–Van Schaftingen–Yung–Sobolev type inequalities connected with maximal inequalities and one parameter families of operators”, *Adv. Math.* **411**:A (2022), art. id. 108774. [MR](#) [Zbl](#)
- [Domínguez et al. 2023] Ó. Domínguez, A. Seeger, B. Street, J. Van Schaftingen, and P.-L. Yung, “Spaces of Besov–Sobolev type and a problem on nonlinear approximation”, *J. Funct. Anal.* **284**:4 (2023), art. id. 109775. [MR](#) [Zbl](#)
- [Federer 1969] H. Federer, *Geometric measure theory*, Grundlehren der Math. Wissenschaften **153**, Springer, 1969. [MR](#) [Zbl](#)

- [Hajłasz and Kałamajska 1995] P. Hajłasz and A. Kałamajska, “Polynomial asymptotics and approximation of Sobolev functions”, *Studia Math.* **113**:1 (1995), 55–64. [MR](#) [Zbl](#)
- [Hunt 1966] R. A. Hunt, “On $L(p, q)$ spaces”, *Enseign. Math. (2)* **12** (1966), 249–276. [MR](#) [Zbl](#)
- [Mattila 2015] P. Mattila, *Fourier analysis and Hausdorff dimension*, Cambridge Stud. Adv. Math. **150**, Cambridge Univ. Press, 2015. [MR](#) [Zbl](#)
- [Nguyen 2006] H.-M. Nguyen, “Some new characterizations of Sobolev spaces”, *J. Funct. Anal.* **237**:2 (2006), 689–720. [MR](#) [Zbl](#)
- [Nguyen 2007] H.-M. Nguyen, “ Γ -convergence and Sobolev norms”, *C. R. Math. Acad. Sci. Paris* **345**:12 (2007), 679–684. [MR](#) [Zbl](#)
- [Nguyen 2008] H.-M. Nguyen, “Further characterizations of Sobolev spaces”, *J. Eur. Math. Soc.* **10**:1 (2008), 191–229. [MR](#) [Zbl](#)
- [Nguyen 2011] H.-M. Nguyen, “ Γ -convergence, Sobolev norms, and BV functions”, *Duke Math. J.* **157**:3 (2011), 495–533. [MR](#) [Zbl](#)
- [Poliakovskiy 2022] A. Poliakovskiy, “Some remarks on a formula for Sobolev norms due to Brezis, Van Schaftingen and Yung”, *J. Funct. Anal.* **282**:3 (2022), art. id. 109312. [MR](#) [Zbl](#)
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Math. Series **30**, Princeton Univ. Press, 1970. [MR](#) [Zbl](#)
- [Stein and Weiss 1971] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Math. Series **32**, Princeton Univ. Press, 1971. [MR](#) [Zbl](#)

Received 24 Sep 2021. Revised 5 Jul 2022. Accepted 11 Aug 2022.

HAÏM BREZIS: brezis@math.rutgers.edu

Department of Mathematics, Rutgers University, Piscataway, NJ, United States

and

Departments of Mathematics and Computer Science, Technion, Israel Institute of Technology, Haifa, Israel

and

Laboratoire Jacques-Louis Lions, Sorbonne Universités, UPMC Université Paris-6, Paris, France

ANDREAS SEEGER: seeger@math.wisc.edu

Department of Mathematics, University of Wisconsin, Madison, WI, United States

JEAN VAN SCHAFTINGEN: jean.vanschafingen@uclouvain.be

Institut de Recherche en Mathématique et Physique, Université catholique de Louvain, Louvain-la-Neuve, Belgium

PO-LAM YUNG: polam.yung@anu.edu.au

Mathematical Sciences Institute, Australian National University, Canberra, Australia

Analysis & PDE

msp.org/apde

EDITOR-IN-CHIEF

Clément Mouhot Cambridge University, UK
c.mouhot@dpmms.cam.ac.uk

BOARD OF EDITORS

Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu
Zbigniew Blocki	Uniwersytet Jagielloński, Poland zbigniew.blocki@uj.edu.pl	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
David Gérard-Varet	Université de Paris, France david.gerard-varet@imj-prg.fr	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Colin Guillarmou	Université Paris-Saclay, France colin.guillarmou@universite-paris-saclay.fr	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Peter Hintz	ETH Zurich, Switzerland peter.hintz@math.ethz.ch	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Vadim Kaloshin	Institute of Science and Technology, Austria vadim.kaloshin@gmail.com	András Vasy	Stanford University, USA andras@math.stanford.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Anna L. Mazzucato	Penn State University, USA alm24@psu.edu	Jim Wright	University of Edinburgh, UK j.r.wright@ed.ac.uk
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France merle@ihes.fr		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2024 is US \$440/year for the electronic version, and \$690/year (+\$65, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 17 No. 3 2024

Schauder estimates for equations with cone metrics, II	757
BIN GUO and JIAN SONG	
The stability of simple plane-symmetric shock formation for three-dimensional compressible Euler flow with vorticity and entropy	831
JONATHAN LUK and JARED SPECK	
Families of functionals representing Sobolev norms	943
HAÏM BREZIS, ANDREAS SEEGER, JEAN VAN SCHAFTINGEN and PO-LAM YUNG	
Schwarz–Pick lemma for harmonic maps which are conformal at a point	981
FRANC FORSTNERIČ and DAVID KALAJ	
An improved regularity criterion and absence of splash-like singularities for g-SQG patches	1005
JUNEKEY JEON and ANDREJ ZLATOŠ	
Spectral gap for obstacle scattering in dimension 2	1019
LUCAS VACOSSIN	