

involve

a journal of mathematics

Hamiltonian labelings of graphs

Willem Renzema and Ping Zhang

 mathematical sciences publishers

2009

Vol. 2, No. 1

Hamiltonian labelings of graphs

Willem Renzema and Ping Zhang

(Communicated by Ron Gould)

For a connected graph G of order n , the detour distance $D(u, v)$ between two vertices u and v in G is the length of a longest $u - v$ path in G . A Hamiltonian labeling of G is a function $c : V(G) \rightarrow \mathbb{N}$ such that $|c(u) - c(v)| + D(u, v) \geq n$ for every two distinct vertices u and v of G . The value $\text{hn}(c)$ of a Hamiltonian labeling c of G is the maximum label (functional value) assigned to a vertex of G by c ; while the Hamiltonian labeling number $\text{hn}(G)$ of G is the minimum value of Hamiltonian labelings of G . Hamiltonian labeling numbers of some well-known classes of graphs are determined. Sharp upper and lower bounds are established for the Hamiltonian labeling number of a connected graph. The corona $\text{cor}(F)$ of a graph F is the graph obtained from F by adding exactly one pendant edge at each vertex of F . For each integer $k \geq 3$, let \mathcal{H}_k be the set of connected graphs G for which there exists a Hamiltonian graph H of order k such that $H \subset G \subseteq \text{cor}(H)$. It is shown that $2k - 1 \leq \text{hn}(G) \leq k(2k - 1)$ for each $G \in \mathcal{H}_k$ and that both bounds are sharp.

1. Introduction

The *distance* $d(u, v)$ between two vertices u and v in a connected graph G is the length of a shortest path between these two vertices. The *eccentricity* $e(v)$ of a vertex v in G is the maximum distance from v to a vertex of G . The *radius* $\text{rad}(G)$ of G is the minimum eccentricity among the vertices of G , while the *diameter* $\text{diam}(G)$ of G is the maximum eccentricity among the vertices of G . A vertex v with $e(v) = \text{rad}(G)$ is called a *central vertex* of G . If $d(u, v) = \text{diam}(G)$, then u and v are *antipodal vertices* of G .

For a connected graph G with diameter d , an *antipodal coloring* of a connected graph G is defined in [Chartrand et al. 2002a] as an assignment $c : V(G) \rightarrow \mathbb{N}$ of colors to the vertices of G such that

$$|c(u) - c(v)| + d(u, v) \geq d,$$

MSC2000: primary 05C12, 05C45; secondary 05C78, 05C15.

Keywords: Hamiltonian labeling, detour distance.

Willem Renzema's research was supported in part by Lee Honors College Dean's Summer Research Award Program at Western Michigan University.

for every two distinct vertices u and v of G . In the case of paths of order $n \geq 2$, this gives

$$|c(u) - c(v)| + d(u, v) \geq n - 1.$$

Antipodal colorings of paths gave rise to the more general Hamiltonian colorings of graphs defined in terms of another distance parameter.

The *detour distance* $D(u, v)$ between two vertices u and v in a connected graph G is the length of a longest path between these two vertices. A $u - v$ path of length $D(u, v)$ is a $u - v$ *detour*. Thus if G is a connected graph of order n , then

$$d(u, v) \leq D(u, v) \leq n - 1,$$

for every two vertices u and v in G , and

$$D(u, v) = n - 1,$$

if and only if G contains a Hamiltonian $u - v$ path. Furthermore $d(u, v) = D(u, v)$ for every two vertices u and v in G if and only if G is a tree. As with standard distance, the detour distance is a metric on the vertex set of a connected graph.

A *Hamiltonian coloring* of a connected graph G of order n is a coloring

$$c : V(G) \rightarrow \mathbb{N}$$

of G such that

$$|c(u) - c(v)| + D(u, v) \geq n - 1,$$

for every two distinct vertices u and v of G . Consequently, if u and v are distinct vertices such that $|c(u) - c(v)| = k$ for some Hamiltonian coloring c of G , then there is a $u - v$ path in G missing at most k vertices of G . The *value* $hc(c)$ of a Hamiltonian coloring c of G is the maximum color assigned to a vertex of G . The *Hamiltonian chromatic number* of G is the minimum value of Hamiltonian colorings of G . Hamiltonian colorings of graphs have been studied in [Chartrand et al. 2002b; 2005a; 2005b; Nebeský 2003; 2006].

For a connected graph G with diameter d , a *radio labeling* of G is defined in [Chartrand et al. 2001] as an assignment $c : V(G) \rightarrow \mathbb{N}$ of labels to the vertices of G such that

$$|c(u) - c(v)| + d(u, v) \geq d + 1,$$

for every two distinct vertices u and v of G . Thus for a radio labeling of a graph, colors assigned to adjacent vertices of G must differ by at least d , colors assigned to two vertices at distance 2 must differ by at least $d - 1$, and so on, up to two vertices at distance d (that is, antipodal vertices), whose colors are only required to differ. The *value* $rn(c)$ of a radio labeling c of G is the maximum color assigned

to a vertex of G . The *radio number* of G is the minimum value of a radio labeling of G . In the case of paths of order $n \geq 2$, this gives

$$|c(u) - c(v)| + d(u, v) \geq n.$$

In a similar manner, radio labelings of paths and detour distance in graphs give rise to a related labeling, which we introduce in this work.

A *Hamiltonian labeling* of a connected graph G of order n is an assignment $c : V(G) \rightarrow \mathbb{N}$ of labels to the vertices of G such that

$$|c(u) - c(v)| + D(u, v) \geq n,$$

for every two distinct vertices u and v of G . Therefore, in a Hamiltonian labeling of G , every two vertices are assigned distinct labels and two vertices u and v can be assigned consecutive labels in G only if G contains a Hamiltonian $u - v$ path. We can assume that every Hamiltonian labeling of a graph uses the integer 1 as one of its labels. The *value* $\text{hn}(c)$ of a Hamiltonian labeling c of G is the maximum label assigned to a vertex of G by c , that is, $\text{hn}(c) = \max\{c(v) : v \in V(G)\}$. The *Hamiltonian labeling number* $\text{hn}(G)$ of G is the minimum value of Hamiltonian labelings of G , that is, $\text{hn}(G) = \min\{\text{hn}(c)\}$, where the minimum is taken over all Hamiltonian labelings c of G . A Hamiltonian labeling c of G with value $\text{hn}(c) = \text{hn}(G)$ is called a *minimum Hamiltonian labeling* of G . Therefore,

$$\text{hn}(G) \geq n. \tag{1}$$

for every connected graph G of order n .

To illustrate these concepts, we consider the Petersen graph P . It is known that $\chi(P) = \text{hc}(P) = 3$. In fact, it is observed in [Chartrand et al. 2005a] that every proper coloring of P is also a Hamiltonian coloring. On the other hand, since the order of P is 10, it follows that $\text{hn}(P) \geq 10$. Observe that $D(u, v) = 8$ if $uv \in E(G)$ and $D(u, v) = 9$ if $uv \notin E(G)$. Thus if c is a Hamiltonian labeling of P , then $|c(u) - c(v)| \geq 2$ if $uv \in E(G)$ and $|c(u) - c(v)| \geq 1$ if $uv \notin E(G)$. Therefore, the labeling shown in Figure 1 is a Hamiltonian labeling and so $\text{hn}(P) = 10$.

2. Bounds for Hamiltonian labeling numbers of graphs

It is convenient to introduce some notation. For a Hamiltonian labeling c of a graph G , an ordering u_1, u_2, \dots, u_n of the vertices of G is called the *c-ordering* of G if

$$1 = c(u_1) < c(u_2) < \dots < c(u_n) = \text{hn}(c).$$

We refer to [Chartrand and Zhang 2008] for graph theory notation and terminology not described in this paper. In order to establish a relationship between the




Figure 1. A Hamiltonian labeling of the Petersen graph.

Hamiltonian chromatic number and Hamiltonian labeling number of a connected graph, we first present a lemma.

Lemma 2.1. *Every connected graph of order $n \geq 3$ with Hamiltonian labeling number n is 2-connected.*

Proof. Assume, to the contrary, that there exists a connected graph G of order $n \geq 3$ with $\text{hn}(G) = n$ such that G is not 2-connected. Then G contains a cut-vertex v . Let c be a minimum Hamiltonian labeling of G and let v_1, v_2, \dots, v_n be the c -ordering of the vertices of G , where then $1 = c(v_1) < c(v_2) < \dots < c(v_n) = n$. Thus $c(v_i) = i$ for $1 \leq i \leq n$. Let $u \in V(G)$ such that u and v are consecutive in the c -ordering. Thus $\{u, v\} = \{v_j, v_{j+1}\}$ for some integer j with $1 \leq j \leq n - 1$. Hence $D(v_j, v_{j+1}) \leq n - 2$. However then,

$$|c(v_j) - c(v_{j+1})| + D(v_j, v_{j+1}) \leq n - 1,$$

which contradicts the fact that c is a Hamiltonian labeling of G . \square

The corollary below now follows immediately.

Corollary 2.2. *No connected graph of order $n \geq 3$ with Hamiltonian labeling number n contains a bridge.*

While $\text{hc}(K_1) = \text{hn}(K_1) = 1$ and $\text{hc}(K_2) = 1$ and $\text{hn}(K_2) = 2$, $\text{hc}(G)$ and $\text{hn}(G)$ must differ by at least 2 for every connected graph G of order 3 or more. In fact, the following result provides upper and lower bounds for the Hamiltonian labeling number of a connected graph in terms of its order and Hamiltonian chromatic number.

Theorem 2.3. *For every connected graph G of order $n \geq 3$,*

$$\text{hc}(G) + 2 \leq \text{hn}(G) \leq \text{hc}(G) + (n - 1).$$

Proof. We first show that $\text{hn}(G) \geq \text{hc}(G) + 2$. Let c be a minimum Hamiltonian labeling of G and let v_1, v_2, \dots, v_n be the c -ordering of the vertices of G , where then $1 = c(v_1) < c(v_2) < \dots < c(v_n) = \text{hn}(c)$. Define a coloring c^* of G by

$$c^*(v_i) = \begin{cases} 1 & \text{if } i = 1, \\ c(v_i) - 1 & \text{if } 2 \leq i \leq n - 1, \\ c(v_i) - 2 & \text{if } i = n. \end{cases}$$

We show that c^* is a Hamiltonian coloring of G . Let $v_i, v_j \in V(G)$, where

$$1 \leq i < j \leq n.$$

We consider two cases.

Case 1. $i = 1$. Suppose first that $2 \leq j \leq n - 2$. Then

$$|c^*(v_j) - c^*(v_1)| + D(v_j, v_1) = c(v_j) - c(v_1) - 1 + D(v_j, v_1) \geq n - 1.$$

Next suppose that $j = n$. Then

$$\begin{aligned} |c^*(v_n) - c^*(v_1)| + D(v_n, v_1) &= c(v_n) - c(v_1) - 2 + D(v_n, v_1) \\ &= c(v_n) - 3 + D(v_n, v_1). \end{aligned}$$

If $c(v_n) \geq n + 1$, then $c(v_n) - 3 + D(v_n, v_1) \geq n - 1$. If $c(v_n) = n$, then $v_1 v_n$ is not a bridge by Corollary 2.2 and so $D(v_n, v_1) \geq 2$. Thus $c(v_n) - 3 + D(v_n, v_1) \geq n - 1$.

Case 2. $i \geq 2$. In this case,

$$|c^*(v_j) - c^*(v_i)| + D(v_j, v_i) = \begin{cases} c(v_j) - c(v_i) + D(v_j, v_i), & \text{if } j \leq n - 1, \\ c(v_j) - c(v_i) - 1 + D(v_j, v_i), & \text{if } j = n, \end{cases} \quad (2)$$

which is greater than or equal to $c(v_j) - c(v_i) - 1 + D(v_j, v_i) \geq n - 1$. Thus c^* is a Hamiltonian coloring of G , as claimed. Therefore,

$$\text{hc}(G) \leq \text{hc}(c^*) = \text{hn}(c) - 2 = \text{hn}(G) - 2,$$

and so $\text{hn}(G) \geq \text{hc}(G) + 2$.

Next, we show that $\text{hn}(G) \leq \text{hc}(G) + (n - 1)$. Let c' be a Hamiltonian coloring of G such that $\text{hc}(c') = \text{hc}(G)$. We may assume that $V(G) = \{v_1, v_2, \dots, v_n\}$ such that

$$1 = c'(v_1) \leq c'(v_2) \leq \dots \leq c'(v_n) = \text{hc}(c').$$

Define a labeling c'' of G by $c''(v_i) = c'(v_i) + (i - 1)$ for $1 \leq i \leq n$. Let v_j and v_k be two distinct vertices of G . Then

$$\begin{aligned} |c''(v_j) - c''(v_k)| + D(v_j, v_k) &= |c'(v_j) - c'(v_k)| + |j - k| + D(v_j, v_k) \\ &\geq (n - 1) + |j - k| \geq n, \end{aligned}$$

and so c'' is a Hamiltonian labeling of G . Since $\text{hn}(c'') = \text{hc}(c) + (n - 1)$, it follows that $\text{hn}(G) \leq \text{hc}(G) + (n - 1)$. \square

While the upper and lower bounds in Theorem 2.3 are sharp (as we will see later), both inequalities in Theorem 2.3 can be strict. For example, consider the Petersen graph P of order $n = 10$ and $\text{hn}(P) = 10$. Thus

$$5 = \text{hc}(P) + 2 < \text{hn}(P) < \text{hc}(P) + (n - 1) = 12.$$

In fact, more can be said. The following result was established in [Chartrand et al. 2005a].

Theorem 2.4 [Chartrand et al. 2005a]. *If G is a Hamiltonian graph of order $n \geq 3$, then $\text{hc}(G) \leq n - 2$. Furthermore, for each pair k, n of integers with $1 \leq k \leq n - 2$, there is a Hamiltonian graph of order n with Hamiltonian chromatic number k .*

On the other hand, every Hamiltonian graph of order n has Hamiltonian labeling number n , as we show next.

Proposition 2.5. *If G is a Hamiltonian graph of order $n \geq 3$, then $\text{hn}(G) = n$.*

Proof. Let $C : v_1, v_2, \dots, v_{n+1} = v_1$ be a Hamiltonian cycle of G . Define the labeling c of G by $c(v_i) = i$ for $1 \leq i \leq n$. Let i, j be two integers with $1 \leq i < j \leq n$. If $j - i \leq n/2$, then $D(v_i, v_j) \geq n - (j - i)$; while if $j - i > n/2$, then $D(v_i, v_j) \geq j - i$. In either case, $|c(v_i) - c(v_j)| + D(v_i, v_j) \geq n$. Thus c is a Hamiltonian labeling and so $\text{hn}(G) = n$ by Equation (1). \square

The converse of Proposition 2.5 is not true. For example, it is well known that the Petersen graph P is a nonHamiltonian graph of order 10 but $\text{hn}(P) = 10$. Whether there exists a connected graph G of order $n \geq 3$ with $\text{hn}(G) = n$ that is neither a Hamiltonian graph nor the Petersen graph is not known. The following realization result is a consequence of Theorem 2.4 and Proposition 2.5.

Corollary 2.6. *For each pair k, n of integers with $2 \leq k \leq n - 1$, there exists a Hamiltonian graph G of order n such that $\text{hn}(G) = \text{hc}(G) + k$.*

In the remainder of this section, we consider the complete bipartite graphs $K_{r,s}$ of order $n = r + s \geq 3$, where $1 \leq r \leq s$. The Hamiltonian chromatic number of a complete bipartite graph has been determined in [Chartrand et al. 2005a]. For positive integers r and s with $r \leq s$ and $r + s \geq 3$,

$$\text{hc}(K_{r,s}) = \begin{cases} r & \text{if } r = s, \\ (s - 1)^2 + 1 & \text{if } 1 = r < s, \\ (s - 1)^2 - (r - 1)^2 & \text{if } 2 \leq r < s. \end{cases} \quad (3)$$

If $r \geq 2$, then $K_{r,r}$ is Hamiltonian and so $\text{hn}(K_{r,r}) = n = 2r$ by Proposition 2.5. Thus, we may assume that $r < s$, beginning with $r = 1$.

Theorem 2.7. *For each integer $n \geq 3$,*

$$\text{hn}(K_{1,n-1}) = n + (n - 2)^2.$$

Proof. Let $G = K_{1,n-1}$ with vertex set $\{v, v_1, v_2, \dots, v_{n-1}\}$, where v is the central vertex of G . By Equation (3) and Theorem 2.3, it suffices to show that

$$\text{hn}(G) \geq n + (n - 2)^2.$$

Let c be a minimum Hamiltonian labeling of G . Since no two vertices of G can be labeled the same, we may assume that

$$c(v_1) < c(v_2) < \dots < c(v_{n-1}).$$

We consider three cases.

Case 1. $c(v) = 1$. Since $D(v_1, v) = 1$ and $D(v_i, v_{i+1}) = 2$ for $1 \leq i \leq n - 2$, it follows that $c(v_1) \geq n$ and

$$c(v_{i+1}) \geq c(v_i) + (n - 2) \geq c(v_1) + i(n - 2) \geq n + i(n - 2)$$

for all $1 \leq i \leq n - 2$. This implies that

$$c(v_{n-1}) \geq n + (n - 2)(n - 2) = n + (n - 2)^2.$$

Therefore, $\text{hn}(G) = \text{hn}(c) \geq n + (n - 2)^2$.

Case 2. $c(v) = \text{hn}(c)$. Then $1 = c(v_1) < c(v_2) < \dots < c(v_{n-1}) < c(v)$. For each i with $2 \leq i \leq n - 1$, it follows that

$$c(v_i) \geq c(v_1) + (i - 1)(n - 2) = 1 + (i - 1)(n - 2).$$

In particular, $c(v_{n-1}) \geq 1 + (n - 2)^2$. Thus

$$c(v) \geq c(v_{n-1}) + n - 1 = n + (n - 2)^2.$$

Therefore, $\text{hn}(G) = \text{hn}(c) \geq n + (n - 2)^2$.

Case 3. $c(v_j) < c(v) < c(v_{j+1})$ for some j with $1 \leq j \leq n - 2$. Thus

$$c(v_j) \geq 1 + (j - 1)(n - 2),$$

$$c(v) \geq c(v_j) + n - 1 \geq n + (j - 1)(n - 2),$$

$$c(v_{j+1}) \geq c(v) + n - 1 \geq 2n - 1 + (j - 1)(n - 2).$$

This implies that

$$\begin{aligned} c(v_{n-1}) &\geq (n - j - 2)(n - 2) + c(v_{j+1}) \\ &\geq (n - j - 2)(n - 2) + (2n - 1) + (j - 1)(n - 2) \\ &= 2n - 1 + (n - 3)(n - 2) = n + 1 + (n - 2)^2 > n + (n - 2)^2. \end{aligned}$$

In each case, we have $\text{hn}(G) \geq n + (n - 2)^2$. □

We now consider $K_{r,s}$, where $2 \leq r < s$, with partite sets V_1 and V_2 such that $|V_1| = r$ and $|V_2| = s$. Then

$$D(u, v) = \begin{cases} 2r - 2 = n - s + r - 2 & \text{if } u, v \in V_1, \\ 2r - 1 = n - s + r - 1 & \text{if } uv \in E(K_{r,s}), \\ 2r = n - s + r & \text{if } u, v \in V_2. \end{cases}$$

Consequently, if c is a Hamiltonian labeling of $K_{r,s}$ ($r < s$), then

$$|c(u) - c(v)| \geq \begin{cases} s - r + 2 & \text{if } u, v \in V_1, \\ s - r + 1 & \text{if } uv \in E(K_{r,s}), \\ s - r & \text{if } u, v \in V_2. \end{cases}$$

Theorem 2.8. *For integers r and s with $2 \leq r < s$,*

$$\text{hn}(K_{r,s}) = (s - 1)^2 - (r - 1)^2 + s + r - 1.$$

Proof. By Equation (3) and Theorem 2.3, it suffices to show that

$$\text{hn}(K_{r,s}) \geq (s - 1)^2 - (r - 1)^2 + s + r - 1.$$

Let $V_1 = \{u_1, u_2, \dots, u_r\}$ and $V_2 = \{v_1, v_2, \dots, v_s\}$ be the partite sets of $K_{r,s}$, and let c be a Hamiltonian labeling of $K_{r,s}$ and let w_1, w_2, \dots, w_{r+s} be the c -ordering of the vertices of $K_{r,s}$. We define a V_1 -block of $K_{r,s}$ to be a set

$$A = \{w_\alpha, w_{\alpha+1}, \dots, w_\beta\},$$

where $1 \leq \alpha \leq \beta \leq r + s$, such that $A \subseteq V_1$, $w_{\alpha-1} \in V_2$ if $\alpha > 1$, and $w_{\beta+1} \in V_2$ if $\beta < r + s$. A V_2 -block of $K_{r,s}$ is defined similarly. Let

$$A_1, A_2, \dots, A_p \quad (p \geq 1)$$

be the distinct V_1 -blocks of $K_{r,s}$ such that if

$$w' \in A_i, \quad w'' \in A_j,$$

where $1 \leq i < j \leq p$, then $c(w') < c(w'')$. If $p \geq 2$, then $K_{r,s}$ contains V_2 -blocks B_1, B_2, \dots, B_{p-1} such that for each integer i ($1 \leq i \leq p - 1$) and for $w' \in A_i$, $w \in B_i$, $w'' \in A_{i+1}$, it follows that

$$c(w') < c(w) < c(w'').$$

The graph $K_{r,s}$ may contain up to two additional V_2 -blocks, namely B_0 and B_p such that if $y \in B_0$ and $y' \in A_1$, then $c(y) < c(y')$; while if $z \in A_p$ and $z' \in B_p$, then $c(z) < c(z')$. If $p = 1$, then at least one of B_0 and B_1 must exist. Hence $K_{r,s}$ contains p V_1 -blocks and $p - 1 + t$ V_2 -blocks, where $t \in \{0, 1, 2\}$. Consequently, there are exactly

- (a) $r - p$ distinct pairs $\{w_i, w_{i+1}\}$ of vertices, both of which belong to V_1 ;

- (b) $2p - 2 + t$ distinct pairs $\{w_i, w_{i+1}\}$ of vertices, exactly one of which belongs to V_1 ;
- (c) $s - (p - 1 + t)$ distinct pairs $\{w_i, w_{i+1}\}$ of vertices, both of which belong to V_2 .

Since (1) the colors of every two vertices w_i and w_{i+1} , both of which belong to V_1 , must differ by at least $s - r + 2$, (2) the colors of every two vertices w_i and w_{i+1} , exactly one of which belongs to V_1 , must differ by at least $s - r + 1$, and (3) the colors of every two vertices w_i and w_{i+1} , both of which belong to V_2 , must differ by at least $s - r$, it follows that

$$\begin{aligned} c(w_{r+s}) &\geq 1 + (r-p)(s-r+2) + (2p-2+t)(s-r+1) + (s-(p-1+t))(s-r) \\ &= (s-1)^2 - (r-1)^2 + s + r - 1 + t. \end{aligned} \tag{4}$$

Since $hn(K_{r,s}) \leq (s-1)^2 - (r-1)^2 + s + r - 1$ and $t \geq 0$, it follows that $t = 0$ and that $hn(K_{r,s}) = (s-1)^2 - (r-1)^2 + s + r - 1$. □

Combining Proposition 2.5 and Theorems 2.7 and 2.8, we obtain the following.

Corollary 2.9. *For integers r and s with $1 \leq r \leq s$,*

$$hn(K_{r,s}) = \begin{cases} r + s & \text{if } r = s, \\ (s-1)^2 + s + 1 & \text{if } r = 1 \text{ and } s \geq 2, \\ (s-1)^2 - (r-1)^2 + r + s - 1 & \text{if } 2 \leq r < s. \end{cases}$$

3. Hamiltonian labeling numbers of subgraphs of coronas of Hamiltonian graphs

A common question in graph theory concerns how the value of a parameter is affected by making a small change in the graph. If G is a Hamiltonian graph and u and v are two nonadjacent vertices of G , then $G + uv$ is also Hamiltonian and so $hn(G) = hn(G + uv)$. On the other hand, if we add a pendant edge to a Hamiltonian graph G producing a nonHamiltonian graph H , then the Hamiltonian labeling number of H can be significantly larger than that of G , as we show in this section. We begin with those graphs obtained from a cycle or a complete graph by adding a single pendant edge.

Theorem 3.1. *If G is the graph of order $n \geq 5$ obtained from C_{n-1} by adding a pendant edge, then $hn(G) = 2n - 2$.*

Proof. Let $C : v_1, v_2, \dots, v_{n-1}, v_1$ and let $v_{n-1}v_n$ be the pendant edge of G . We first show that $hn(G) \leq 2n - 2$. Define a labeling c_0 of G by

$$c_0(v_i) = \begin{cases} 2i & \text{if } 1 \leq i \leq n-1, \\ 1 & \text{if } i = n. \end{cases}$$

We show that c_0 is a Hamiltonian labeling. First let

$$v_i, v_j \in V(C),$$

where $1 \leq i < j \leq n-1$. If $j-i \geq \frac{n-1}{2}$, then $D(v_i, v_j) = j-i$ and so

$$\begin{aligned} |c_0(v_i) - c_0(v_j)| + D(v_i, v_j) &= |2i - 2j| + (j-i) = 3(j-i) \\ &\geq 3\left(\frac{n-1}{2}\right) = \frac{3n}{2} - \frac{3}{2} \geq n, \end{aligned}$$

since $n \geq 3$. If $j-i \leq \frac{n-1}{2}$, then $D(v_i, v_j) = (n-1) - (j-i)$ and so

$$\begin{aligned} |c_0(v_i) - c_0(v_j)| + D(v_i, v_j) &= 2(j-i) + [(n-1) - (j-i)] \\ &= n-1 + (j-i) \geq n. \end{aligned}$$

Next, we consider each pair v_i, v_n where $1 \leq i \leq n-1$. Since $D(v_i, v_n) \geq n-i$ and $|c_0(v_i) - c_0(v_n)| \geq 2i-1$, it follows that

$$|c_0(v_i) - c_0(v_n)| + D(v_i, v_n) \geq n+i-1 \geq n.$$

Therefore, c_0 is a Hamiltonian labeling, as claimed.

Next, we show that $\text{hn}(G) \geq 2n-2$. Let c be a minimum Hamiltonian labeling of G . First, we make some observations.

- (a) For each pair i, j with $1 \leq i \neq j \leq n-1$, $D(v_i, v_j) \leq n-2$ and so $|c(v_i) - c(v_j)| \geq 2$.
- (b) For each i with $i \in \{1, n-2\}$, $D(v_n, v_i) = n-1$ and so $|c(v_n) - c(v_i)| \geq 1$.
- (c) For each i with $1 \leq i \leq n-1$ and $i \notin \{1, n-2\}$, $D(v_n, v_i) \leq n-2$ and so $|c(v_n) - c(v_i)| \geq 2$.

Let u_1, u_2, \dots, u_n be the c -ordering of the vertices of G and let

$$X = \{c(u_{i+1}) - c(u_i) : 1 \leq i \leq n-1\}.$$

By observations (a)–(c), at most two terms in X are 1. If at most one term in X is 1, then $\text{hn}(c) = c(u_n) \geq 1 + 1 + 2(n-2) = 2n-2$. If at least one term in X is 3 or more, then $\text{hn}(c) = c(u_n) \geq 1 + 1 + 1 + 3 + 2(n-4) = 2n-2$. Thus we may assume that exactly two terms in X are 1 and the remaining terms in X are 2. Then $v_n = u_i$ for some i with $2 \leq i \leq n-1$ and $\{v_1, v_{n-2}\} = \{u_{i-1}, u_{i+1}\}$, where $c(u_i) - c(u_{i-1}) = c(u_{i+1}) - c(u_i) = 1$. This implies that $v_{n-1} = u_j$ for some j with $1 \leq j \leq n$ and $j \neq i$. If $2 \leq j \leq n-1$, then $\{u_{j-1}, u_{j+1}\} \neq \{v_1, v_{n-2}\}$; if $j = 1$, then $u_2 \notin \{v_1, v_{n-2}\}$, for otherwise

$$\begin{aligned} |c(v_{n-1}) - c(v_n)| + D(v_{n-1}, v_n) &\leq |c(v_{n-1}) - c(u_2)| + |c(u_2) - c(v_n)| + 1 \\ &\leq 2 + 1 + 1 = 4 < n, \end{aligned}$$

which is impossible; if $j = n$, then $u_{n-1} \notin \{v_1, v_{n-2}\}$, for otherwise

$$\begin{aligned} |c(v_{n-1}) - c(v_n)| + D(v_{n-1}, v_n) &\leq |c(v_{n-1}) - c(u_{n-1})| + |c(u_{n-1}) - c(v_n)| + 1 \\ &\leq 2 + 1 + 1 = 4 < n, \end{aligned}$$

again, which is impossible. Therefore, for each j with $1 \leq j \leq n$, there exists $k \in \{j-1, j+1\}$ such that $u_k \notin \{v_1, v_{n-2}\}$. Assume, without loss of generality, that $u_{j-1} \notin \{v_1, v_{n-2}\}$. Since $D(u_{j-1}, u_j) \leq n-3$, it follows that $c(u_j) - c(u_{j-1}) \geq 3$, which is impossible since each term in X is at most 2. Thus, $\text{hn}(G) \geq 2n-2$. \square

Theorem 3.2. *If G is the graph of order $n \geq 4$ obtained from K_{n-1} by adding a pendant edge, then $\text{hn}(G) = 2n-3$.*

Proof. Let $V(K_{n-1}) = \{v_1, v_2, \dots, v_{n-1}\}$ and let G be obtained from K_{n-1} by adding the pendant edge $v_{n-1}v_n$. We first show that $\text{hn}(G) \leq 2n-3$. Define a labeling c_0 of G by

$$c_0(v) = \begin{cases} 2i-1 & \text{if } v = v_i \text{ for } 1 \leq i \leq n-1, \\ 2 & \text{if } v = v_n. \end{cases}$$

For each pair i, j of integers with $1 \leq i \neq j \leq n-1$,

$$D(v_i, v_j) = n-2 \quad \text{and} \quad |c_0(v_i) - c_0(v_j)| \geq 2.$$

For each i with $1 \leq i \leq n-2$,

$$D(v_n, v_i) = n-1 \quad \text{and} \quad |c_0(v_n) - c_0(v_i)| \geq 1.$$

Furthermore, $D(v_n, v_{n-1}) = 1$ and

$$|c_0(v_n) - c_0(v_{n-1})| \geq (2n-3) - 2 = 2n-5 \geq n-1$$

for $n \geq 4$. In each case,

$$D(v_i, v_j) + |c_0(v_i) - c_0(v_j)| \geq n,$$

for all i, j with $1 \leq i \neq j \leq n$. Therefore, c_0 is a Hamiltonian labeling and so $\text{hn}(G) \leq \text{hn}(c_0) = c_0(v_{n-1}) = 2n-3$.

Next, we show that $\text{hn}(G) \geq 2n-3$. Let c be a minimum Hamiltonian labeling of G . Suppose that the vertices of K_{n-1} in G can be ordered as u_1, u_2, \dots, u_{n-1} such that $c(u_1) < c(u_2) < \dots < c(u_{n-1})$. Since

$$D(u_i, u_j) = n-2$$

for $1 \leq i < j \leq n-1$, it follows that

$$|c(u_i) - c(u_j)| = c(u_j) - c(u_i) \geq 2.$$

This implies that

$$\text{hn}(c) \geq c(u_{n-1}) \geq 1 + 2(n-2) = 2n-3.$$

Therefore, $\text{hn}(G) \geq 2n-3$. \square

Let G be a connected graph containing an edge e that is not a bridge. Then $G-e$ is connected. For every two distinct vertices u and v in $G-e$, the length of a longest $u-v$ path in $G-e$ does not exceed the length of a longest $u-v$ path in G . Thus every Hamiltonian labeling of $G-e$ is a Hamiltonian labeling of G . This observation yields the following useful lemma.

Lemma 3.3. *If F is a connected subgraph of a connected graph G , then*

$$\text{hn}(G) \leq \text{hn}(F).$$

The following is a consequence of Theorems 3.1 and 3.2 and Lemma 3.3.

Corollary 3.4. *Let H be a Hamiltonian graph of order $n-1 \geq 3$. If G is a graph obtained from H by adding a pendant edge, then*

$$2n-3 \leq \text{hn}(G) \leq 2n-2.$$

Proof. Let C be a Hamiltonian cycle in H . If $H = C_{n-1}$, then $\text{hn}(G) = 2n-2$ by Theorem 3.1; while if $H = K_{n-1}$, then $\text{hn}(G) = 2n-3$ by Theorem 3.2. Thus, we may assume that $H \neq C_{n-1}$ and $H \neq K_{n-1}$. Let F be the graph obtained from K_{n-1} by adding a pendant edge and F' be the graph obtained from C_{n-1} by adding a pendant edge. Then G can be obtained from F by deleting nonbridge edges and F' can be obtained from G by deleting nonbridge edges. It then follows by Lemma 3.3 that $\text{hn}(F) \leq \text{hn}(G) \leq \text{hn}(F')$ and so $2n-3 \leq \text{hn}(G) \leq 2n-2$. \square

In fact, there exists a Hamiltonian graph H of order $n-1$ such that adding a pendant edge at a vertex x of H produces a graph G with $\text{hn}(G) = 2n-3$ but adding a pendant edge at a different vertex y of H produces a graph F with $\text{hn}(F) = 2n-2$. For example, let H be the Hamiltonian graph obtained from the cycle $C : v_1, v_2, \dots, v_{n-1}, v_1$ of order $n-1 \geq 4$ by adding the edge v_1v_{n-2} . If G is formed from H by adding a pendant edge at v_{n-1} , then $\text{hn}(G) = 2n-3$; while if F is formed from H by adding the pendant edge v_1 , then $\text{hn}(F) = 2n-2$.

In order to study graphs obtained from a Hamiltonian graph by adding pendant edges, we first establish some additional definitions and notation. For a graph F , the *corona* $\text{cor}(F)$ of F is that graph obtained from F by adding exactly one pendant edge at each vertex of F . For a connected graph G , the *core* $C(G)$ of G is obtained from G by successively deleting vertices of degree 1 until none remain. Thus, if G is a tree, then its core is K_1 ; while if G is not a tree, then the core of G is the induced subgraph F of maximum order with $\delta(F) \geq 2$. For each integer $k \geq 3$, let \mathcal{H}_k be the set of nonHamiltonian graphs that can be obtained from a

Hamiltonian graph of order k by adding pendant edges to this graph in such a way that at most one pendant edge is added to each vertex of the graph. Thus if $G \in \mathcal{H}_k$, then there is a Hamiltonian graph H of order k such that G is a connected subgraph of $\text{cor}(H)$ whose core is H . We now establish lower and upper bounds for the Hamiltonian labeling number of a graph in \mathcal{H}_k in terms of the integer k and the order of the graph, beginning with a lower bound.

Theorem 3.5. *Let $G \in \mathcal{H}_k$ be a graph of order n and $k + 1 \leq n \leq 2k$. Then*

$$\text{hn}(G) \geq (n - 1)(n - k) + (2k - n).$$

Proof. Suppose that H is a Hamiltonian graph of order $k \geq 3$ and that $H \cong C(G)$. If $H \not\cong K_k$, then G can be obtained from some graph $F \in \mathcal{H}_k$ by deleting nonbridge edges from F , where $C(F) \cong K_k$, and $V(G - H) = V(F - K_k)$. That is, G and F possess the same end-vertices. It then follows by Lemma 3.3 that

$$\text{hn}(F) \leq \text{hn}(G).$$

Therefore, it suffices to show that

$$\text{hn}(F) \geq (n - 1)(n - k) + (2k - n).$$

Let $V(F) = U \cup W$, where $U = V(K_k)$ and $W = V(F) - U$. First we make some observations:

- (a) If $x, y \in U$, then $D(x, y) = k - 1$.
- (b) If $x, y \in W$, then $D(x, y) = k + 1$.
- (c) If $x \in U$ and $y \in W$, then $D(x, y) = 1$ if $xy \in E(F)$ and $D(x, y) = k$ otherwise.

Let c be a minimum Hamiltonian labeling of F and let v_1, v_2, \dots, v_n be the c -ordering of the vertices of F . We define the four subsets $S_u, S_w, S_{u,w}$, and $S_{w,u}$ of $V(F)$ as follows:

$$\begin{aligned} S_u &= \{v_i : v_{i-1}, v_i \in U \text{ for } 2 \leq i \leq n\}, \\ S_w &= \{v_i : v_{i-1}, v_i \in W \text{ for } 2 \leq i \leq n\}, \\ S_{u,w} &= \{v_i : v_{i-1} \in U \text{ and } v_i \in W \text{ for } 2 \leq i \leq n\}, \\ S_{w,u} &= \{v_i : v_{i-1} \in W \text{ and } v_i \in U \text{ for } 2 \leq i \leq n\}. \end{aligned}$$

Let $|S_u| = n_u, |S_w| = n_w, |S_{u,w}| = n_{u,w}, |S_{w,u}| = n_{w,u}$. Since

$$S_u \cup S_w \cup S_{u,w} \cup S_{w,u} = V(F) - \{v_1\},$$

it follows that

$$n_u + n_w + n_{u,w} + n_{w,u} = n - 1. \tag{5}$$

For each integer i with $2 \leq i \leq n$,

- (A) if $v_i \in S_u$, then $c(v_i) - c(v_{i-1}) \geq n - k + 1$ by (a);
 (B) if $v_i \in S_w$, then $c(v_i) - c(v_{i-1}) \geq n - k - 1$ by (b);
 (C) if $v_i \in S_u \cup S_w$, then either $c(v_i) - c(v_{i-1}) \geq n - 1$ or $c(v_i) - c(v_{i-1}) \geq n - k$ by (iii), and so $c(v_i) - c(v_{i-1}) \geq n - k$ in this case.

It then follows by (A)–(C) and (5) that

$$\begin{aligned} \text{hn}(c) &= c(v_n) \geq 1 + n_u(n - k + 1) + n_w(n - k - 1) + (n_{u,w} + n_{w,u})(n - k) \\ &= 1 + (n_u + n_w + n_{u,w} + n_{w,u})(n - k) + (n_u - n_w) \\ &= 1 + (n - 1)(n - k) + (n_u - n_w). \end{aligned}$$

We claim that $n_u - n_w \geq 2k - n - 1$. Since

$$S_u \cup S_{u,w} = \{v_i : v_{i-1} \in U \text{ for } 2 \leq i \leq n\},$$

it follows that

$$|S_u \cup S_{u,w}| = \begin{cases} |U| - 1 & \text{if } v_n \in U, \\ |U| & \text{otherwise;} \end{cases}$$

and so

$$n_u + n_{u,w} = k \text{ or } n_u + n_{u,w} = k - 1. \quad (6)$$

Since

$$\begin{aligned} S_w \cup S_{u,w} &= \{v_i : v_i \in W \text{ for } 2 \leq i \leq n\} \\ &= \begin{cases} W - \{v_1\} & \text{if } v_1 \in W, \\ W & \text{otherwise,} \end{cases} \end{aligned}$$

it follows that

$$n_w + n_{u,w} = n - k \text{ or } n_w + n_{u,w} = n - k - 1. \quad (7)$$

By Equations (6) and (7), we obtain

$$n_u - n_w = (n_u + n_{u,w}) - (n_w + n_{u,w}) \geq (k - 1) - (n - k) = 2k - n - 1,$$

as claimed. Therefore,

$$\text{hn}(G) = \text{hn}(c) \geq 1 + (n - 1)(n - k) + (n_u - n_w) \geq (n - 1)(n - k) + (2k - n).$$

This completes the proof. \square

Theorem 3.6. *Let $G \in \mathcal{H}_k$ be a graph of order n and $k + 2 \leq n \leq 2k$. Then*

$$\text{hn}(G) \leq 1 + n + (n - k - 1)^2 + (k - 2)(n - k + 1).$$

Proof. Suppose that H is a Hamiltonian graph of order $k \geq 3$ and that $H \cong C(G)$. If $H \not\cong C_k$, then C_k can be obtained from H by deleting edges. Thus there exists $F \in \mathcal{H}_k$ such that $C(F) \cong C_k$ and F can be obtained from G by deleting edges that are not bridges. It then follows by Lemma 3.3 that

$$\text{hn}(G) \leq \text{hn}(F).$$

Therefore, we may assume that $H \cong C_k : x_1, x_2, \dots, x_k, x_1$. Now let

$$X = \{x_1, x_2, \dots, x_k\} \quad \text{and} \quad Y = V(G) - X = \{y_1, y_2, \dots, y_{n-k}\}$$

such that y_i is adjacent to x_{j_i} , for $1 \leq i \leq n-k$, and $1 = j_1 < j_2 < \dots < j_{n-k} \leq k$. For each i with $1 \leq i \leq n-k$, let

$$g_i = j_{i+1} - j_i - 1, \tag{8}$$

where $j_{n-k+1} = j_1$; that is, g_i is the number of vertices of degree 2 between x_{j_i} and $x_{j_{i+1}}$ on C_k . Thus if $x_{j_i}y_i \in E(G)$, then $x_{j_i+g_i+1}y_{i+1} \in E(G)$, for $1 \leq i \leq n-k$, and

$$\sum_{i=1}^{n-k} g_i = 2k - n.$$

Now define the labeling c of G by

$$c(v) = \begin{cases} 1 & \text{if } v = x_k, \\ 1 + n - k & \text{if } v = y_1, \\ c(y_{i-1}) + (n - k - 1) + g_{i-1} & \text{if } v = y_i \text{ and } 2 \leq i \leq n - k, \\ c(y_{n-k}) + n - k + g_{n-k} & \text{if } v = x_1, \\ c(x_{j-1}) + (n - k + 1) & \text{if } v = x_j \text{ and } 2 \leq j \leq k - 1. \end{cases} \tag{9}$$

Thus the c -ordering of the vertices of G is

$$x_k, y_1, y_2, \dots, y_{n-k}, x_1, x_2, \dots, x_{k-1},$$

and by Equation (9)

$$\begin{aligned} c(x_k) &= 1, \\ c(y_i) &= 1 + n - k + (i - 1)(n - k - 1) + \sum_{\ell=1}^{i-1} g_\ell \text{ for } 1 \leq i \leq n - k, \\ c(x_1) &= 1 + n + (n - k - 1)^2, \\ c(x_j) &= 1 + n + (n - k - 1)^2 + (j - 1)(n - k + 1) \text{ for } 2 \leq j \leq k - 1. \end{aligned} \tag{10}$$

Therefore, the value of c is

$$\text{hn}(c) = c(x_{k-1}) = 1 + n + (n - k - 1)^2 + (k - 2)(n - k + 1).$$

Thus it remains to show that c is a Hamiltonian labeling of G . First, we make some observations. Let $u, v \in V(G)$, where $u \neq v$.

(α) If $u = x_i$ and $v = x_j$ where $1 \leq i \neq j \leq k$, then $D(u, v) = \max\{|i - j|, k - |i - j|\}$.

(β) If $u = y_i$ and $v = y_j$ where $1 \leq i < j \leq n - k$, then

$$D(u, v) = 2 + \max \left\{ j - i + \sum_{\ell=i}^{j-1} g_\ell, k - \left(j - i + \sum_{\ell=i}^{j-1} g_\ell \right) \right\}.$$

(γ) If $u = x_i, v \in Y$, and $vx_j \in E(G)$ where $1 \leq i, j \leq k$ (possibly $i = j$), then $D(u, v) = 1$ if $i = j$ and $D(u, v) = 1 + \max\{|i - j|, k - |i - j|\}$ if $i \neq j$.

We show that

$$D(u, v) + |c(u) - c(v)| \geq n, \quad (11)$$

for every pair u, v of distinct vertices of G . We consider three cases.

Case 1. $u, v \in X$. Let $u = x_i$ and $v = x_j$, where $1 \leq i, j \leq k$. We may assume, without loss of generality, that $i < j$. If $j = k$, then

$$\begin{aligned} |c(x_i) - c(x_j)| &= c(x_i) - c(x_k) \\ &= [1 + n + (n - k - 1)^2 + (i - 1)(n - k + 1)] - 1 \geq n, \end{aligned}$$

and so condition (11) is satisfied. Thus we may assume that $j \neq k$.

If $j - i = 1$, then $D(x_i, x_j) = k - 1$ and $|c(x_i) - c(x_j)| = n - k + 1$. Thus (11) holds in this case. If $j - i \geq \frac{k}{2}$, then

$$\begin{aligned} D(x_i, x_j) + |c(x_i) - c(x_j)| &= c(x_j) - c(x_i) + D(x_i, x_j) \\ &= (j - i)(n - k + 1) + (j - i) = (j - i)(n - k + 2) \\ &\geq \frac{k}{2}(n - k + 2) = k \left(\frac{n - k}{2} + 1 \right) \geq 2k \geq n. \end{aligned}$$

If $2 \leq j - i \leq \frac{k}{2}$, then

$$\begin{aligned} D(x_i, x_j) + |c(x_i) - c(x_j)| &= c(x_j) - c(x_i) + D(x_i, x_j) \\ &= (j - i)(n - k + 1) + (k - (j - i)) \\ &= (j - i)(n - k) + k \\ &\geq 2(n - k) + k = 2n - k \geq n. \end{aligned}$$

Case 2. $u, v \in Y$. Let $u = y_i$ and $v = y_j$, where $1 \leq i, j \leq n - k$. We may assume, without loss of generality, that $i < j$. Then

$$|c(y_i) - c(y_j)| = c(y_j) - c(y_i) = (j - i)(n - k - 1) + \sum_{\ell=i}^{j-1} g_\ell.$$

If $j - i + \sum_{\ell=i}^{j-1} g_\ell \geq \frac{k}{2}$, then

$$D(y_i, y_j) = 2 + j - i + \sum_{\ell=i}^{j-1} g_\ell$$

by (β) , and so

$$\begin{aligned} D(y_i, y_j) + |c(y_i) - c(y_j)| &= (j - i)(n - k - 1) + \left(\sum_{\ell=i}^{j-1} g_\ell \right) + 2 + j - i + \left(\sum_{\ell=i}^{j-1} g_\ell \right) \\ &\geq (j - i)(n - k - 1) + 2 + (j - i) + [k - 2(j - i)] \\ &= (j - i)(n - k - 2) + k + 2 \geq n. \end{aligned}$$

If $1 \leq j - i + \sum_{\ell=i}^{j-1} g_\ell \leq \frac{k}{2}$, then

$$D(y_i, y_j) = 2 + k - \left(j - i + \sum_{\ell=i}^{j-1} g_\ell \right)$$

by (β) , and so

$$\begin{aligned} D(y_i, y_j) + |c(y_i) - c(y_j)| &= (j - i)(n - k - 1) + \left(\sum_{\ell=i}^{j-1} g_\ell \right) + 2 + k - \left(j - i + \sum_{\ell=i}^{j-1} g_\ell \right) \\ &= (j - i)(n - k - 2) + k + 2 \\ &\geq n - k - 2 + k + 2 = n. \end{aligned}$$

Case 3. One of u and v is in X and the other is in Y , say $u \in X$ and $v \in Y$. Let $u = x_i$ and $v = y_j$, where $1 \leq i \leq k$ and $1 \leq j \leq n - k$. We consider two subcases, according to whether $x_i y_j \in E(G)$ or $x_i y_j \notin E(G)$.

Subcase 3.1. $x_i y_j \in E(G)$. We proceed by induction to show that

$$c(x_i) - c(y_j) \geq n - 1$$

when $x_i y_j \in E(G)$. For $i = j = 1$,

$$\begin{aligned} |c(x_1) - c(y_1)| = c(x_1) - c(y_1) &= [1 + n + (n - k - 1)^2] - (1 + n - k) \\ &= (n - k - 1)^2 + k \geq n - 1 \text{ for } n \geq k + 2. \end{aligned}$$

Assume that $c(x_i) - c(y_j) \geq n - 1$. Since $x_{i+1+g_j} y_{j+1} \in E(G)$ by (8), we show that $c(x_{i+1+g_j}) - c(y_{j+1}) \geq n - 1$. Observe that

$$c(x_{i+1+g_j}) = c(x_i) + (g_j + 1)(n - k + 1) \quad \text{and} \quad c(y_{j+1}) = c(y_j) + (n - k - 1) + g_j.$$

It then follows by the induction hypothesis that

$$\begin{aligned} c(x_{i+1+g_j}) - c(y_{j+1}) &\geq n - 1 + (g_j + 1)(n - k + 1) - (n - k - 1) - g_j \\ &= n + 1 + g_j(n - k) \geq n - 1. \end{aligned}$$

Therefore if $x_i y_j \in E(G)$, then $|c(x_i) - c(y_j)| + D(x_i, y_j) \geq n - 1 + 1 = n$. Thus condition (11) is satisfied.

Subcase 3.2. $x_i y_j \notin E(G)$. Then $i \neq j$. By (8), if $y_j x_m \in E(G)$, then

$$\sum_{\ell=1}^{j-1} g_\ell = m - j,$$

and

$$\begin{aligned} D(x_i, y_j) + |c(x_i) - c(y_j)| &= c(x_i) - c(y_j) + D(x_i, x_m) + 1 \quad (12) \\ &= (n - k - 1)^2 + (i - j)(n - k) + i + j - (m - j) + k - 1 + D(x_i, x_m) \\ &= [(n - k - 1)^2 + (i - j)(n - k) + k + 2j - 1] + (i - m) + D(x_i, x_m). \end{aligned}$$

Now observe, if $i > j$, then $(i - j)(n - k) + k \geq n$; whereas if $1 \leq i < j \leq n - k$, then

$$\begin{aligned} (n - k - 1)^2 + (i - j)(n - k) + k + 2j - 1 \\ &= [(n - k)^2 - 2(n - k)] + i(n - k) - j(n - k - 2) + k \\ &\geq [(n - k)^2 - 2(n - k)] + i(n - k) - [(n - k)^2 - 2(n - k)] + k \geq n. \end{aligned}$$

Therefore, by Equation (12)

$$D(x_i, y_j) + |c(x_i) - c(y_j)| \geq n + (i - m) + D(x_i, x_m). \quad (13)$$

We then have three possible situations. If $i > m$, then $i - m > 0$ and so by condition (13), (11) is satisfied. If $m > i$ and $m - i \geq k/2$, then $D(x_i, x_m) = m - i$ and so by (13)

$$D(x_i, y_j) + |c(x_i) - c(y_j)| \geq n + (i - m) + (m - i) = n.$$

Finally, if $m > i$ and $m - i \leq k/2$, then $D(x_i, x_m) = k - (m - i)$ and so from (13)

$$\begin{aligned} D(x_i, y_j) + |c(x_i) - c(y_j)| &\geq n + (i - m) + [k - (m - i)] \\ &= n + k - 2(m - i) \geq n + k - k = n. \end{aligned}$$

For each situation, condition (11) is satisfied. Therefore c is a Hamiltonian labeling of G . \square

We now present two corollaries of Theorems 3.5 and 3.6.

Corollary 3.7. *If G is a graph of order n that is the corona of a Hamiltonian graph, then*

$$\text{hn}(G) = \binom{n}{2}.$$

Proof. Suppose that H is a Hamiltonian graph of order $k \geq 3$ and that $G = \text{cor}(H)$. Then the order of G is $n = 2k$. We show that

$$\text{hn}(G) = \binom{n}{2} = k(2k - 1).$$

If $H \neq C_k$ and $H \neq K_k$, then G can be obtained from $\text{cor}(K_k)$ by deleting nonbridge edges and $\text{cor}(C_k)$ can be obtained from G by deleting edges that are not bridges. It then follows by Lemma 3.3 that

$$\text{hn}(\text{cor}(K_k)) \leq \text{hn}(G) \leq \text{hn}(\text{cor}(C_k)).$$

Therefore, it suffices to show that

$$k(2k - 1) \leq \text{hn}(\text{cor}(K_k)) \text{ and } \text{hn}(\text{cor}(C_k)) \leq k(2k - 1).$$

From Theorems 3.5 and 3.6, we find that

$$\text{hn}(\text{cor}(K_k)) \geq (2k - 1)(2k - k) + (2k - 2k) = k(2k - 1)$$

and

$$\begin{aligned} \text{hn}(\text{cor}(C_k)) &\leq 1 + 2k + (2k - k - 1)^2 + (k - 2)(2k - k + 1) \\ &= 1 + 2k + k^2 - 2k + 1 + k^2 - k - 2 = k(2k - 1). \end{aligned}$$

Therefore, $\text{hn}(G) = k(2k - 1)$. □

Corollary 3.8. *For each graph $G \in \mathcal{H}_k$,*

$$2k - 1 \leq \text{hn}(G) \leq k(2k - 1).$$

Proof. Let

$$f(x) = (x - 1)(x - k) + (2k - x),$$

for $k + 1 \leq x \leq 2k$ and let

$$g(x) = 1 + x + (x - k - 1)^2 + (k - 2)(x - k + 1),$$

for $k + 2 \leq x \leq 2k$. Let $G \in \mathcal{H}_k$ be a graph of order n where $k + 1 \leq n \leq 2k$. Then by Corollary 3.4 and Theorems 3.5 and 3.6,

$$f(n) \leq \text{hn}(G) \leq g(n).$$

Since each $f(x)$ and $g(x)$ is an increasing function in its domain, it follows that $f(x) \geq f(k+1) = 2k-1$ and $g(x) \leq g(2k) = k(2k-1)$, implying the desired result. \square

Both lower and upper bound in Corollary 3.8 are sharp. For example, if $G' \in \mathcal{H}_k$ is a graph of order $k+1$ whose core is K_k , then $\text{hn}(G') = 2n-3 = 2k-1$ by Theorem 3.2; while if $G'' \in \mathcal{H}_k$ is a graph of order $2k$ whose core is K_k , then

$$\text{hn}(G'') = \binom{n}{2} = k(2k-1)$$

by Corollary 3.7.

Acknowledgements

We are grateful to the referee whose valuable suggestions resulted in an improved paper.

References

- [Chartrand and Zhang 2008] G. Chartrand and P. Zhang, *Chromatic graph theory*, Chapman & Hall/CRC, Boca Raton, FL, 2008.
- [Chartrand et al. 2001] G. Chartrand, D. Erwin, P. Zhang, and F. Harary, “Radio labelings of graphs”, *Bull. Inst. Combin. Appl.* **33** (2001), 77–85. MR 2003d:05185 Zbl 0989.05102
- [Chartrand et al. 2002a] G. Chartrand, D. Erwin, and P. Zhang, “Radio antipodal colorings of graphs”, *Math. Bohem.* **127**:1 (2002), 57–69. MR 2003b:05065 Zbl 0995.05056
- [Chartrand et al. 2002b] G. Chartrand, L. Nebeský, and P. Zhang, “Bounds for the Hamiltonian chromatic number of a graph”, pp. 113–125 in *Proceedings of the Thirty-third Southeastern International Conference on Combinatorics, Graph Theory and Computing (Boca Raton, FL, 2002)*, vol. 157, 2002. MR 2004e:05062 Zbl 1029.05059
- [Chartrand et al. 2005a] G. Chartrand, L. Nebeský, and P. Zhang, “Hamiltonian colorings of graphs”, *Discrete Appl. Math.* **146**:3 (2005), 257–272. MR 2005i:05061 Zbl 1056.05054
- [Chartrand et al. 2005b] G. Chartrand, L. Nebesky, and P. Zhang, “On Hamiltonian colorings of graphs”, *Discrete Math.* **290**:2-3 (2005), 133–143. MR 2006a:05058 Zbl 1059.05046
- [Nebeský 2003] L. Nebeský, “Hamiltonian colorings of graphs with long cycles”, *Math. Bohem.* **128**:3 (2003), 263–275. MR 2004h:05051 Zbl 1050.05055
- [Nebeský 2006] L. Nebeský, “The Hamiltonian chromatic number of a connected graph without large Hamiltonian-connected subgraphs”, *Czech. Math. J.* **56**:2 (2006), 317–338. MR 2007k:05125

Received: 2008-08-21 Revised: Accepted: 2008-11-15

Wille.a.renzema@wmich.edu *Department of Mathematics, Western Michigan University,
Kalamazoo, MI 49008, United States*

Ping.zhang@wmich.edu *Department of Mathematics, Western Michigan University,
Kalamazoo, MI 49008, United States*