

# involve

a journal of mathematics

Binary frames with prescribed dot products  
and frame operator

Veronika Furst and Eric P. Smith



# Binary frames with prescribed dot products and frame operator

Veronika Furst and Eric P. Smith

(Communicated by David Royal Larson)

This paper extends three results from classical finite frame theory over real or complex numbers to binary frames for the vector space  $\mathbb{Z}_2^d$ . Without the notion of inner products or order, we provide an analog of the “fundamental inequality” of tight frames. In addition, we prove the binary analog of the characterization of dual frames with given inner products and of general frames with prescribed norms and frame operator.

## 1. Introduction

Due to applications in signal and image processing, data compression, sampling theory, and other problems in engineering and computer science, frames in finite-dimensional spaces have received much attention from pure and applied mathematicians alike, over the past thirty years; see, for example, Chapter 1 of [Casazza and Kutyniok 2013]. The redundant representation of vectors inherent to frame theory is central to the idea of efficient data storage and transmission that is robust to noise and erasures.

Frames for  $\mathbb{C}^d$  and  $\mathbb{R}^d$  have been extensively studied; see [Christensen 2003] for a standard introduction to frame theory, [Kovačević and Chebira 2007] for applications, and [Han et al. 2007] for an exposition at the undergraduate level. Noting the similarity between frames and error-correcting codes, Bodmann, Le, Reza, Tobin, and Tomforde [Bodmann et al. 2009] introduced the concept of *binary frames*, that is, finite frames for the vector space  $\mathbb{Z}_2^d$ . Binary Parseval frames robust to erasures were characterized in [Bodmann et al. 2014], and their Gramian matrices were studied in [Baker et al. 2018]. A more generalized approach to binary frames was taken in [Hotovy et al. 2015].

We begin with a brief introduction to classical frame theory terminology. Let  $\mathbb{H}^d$  denote the Hilbert space  $\mathbb{C}^d$  or  $\mathbb{R}^d$ .

---

MSC2010: 15A03, 15A23, 15B33, 42C15.

Keywords: frames, binary vector spaces, frame operators, Gramian matrices.

**Definition 1.1.** A (finite) *frame* for a Hilbert space  $\mathbb{H}^d$  is a collection  $\{x_j\}_{j=1}^K$  of vectors in  $\mathbb{H}^d$  for which there exist finite constants  $A, B > 0$  such that for every  $y \in \mathbb{H}^d$ ,

$$A\|y\|^2 \leq \sum_{j=1}^K |\langle y, x_j \rangle|^2 \leq B\|y\|^2.$$

The constants  $A$  and  $B$  are known as *frame bounds*. An  $A$ -*tight frame* is one for which  $A = B$ , and a *Parseval frame* is one for which  $A = B = 1$ .

The vectors  $x_j$  in the above definition need not be orthogonal or even linearly independent. An orthonormal basis is most closely resembled by a Parseval frame, for which we have the (not necessarily unique) reconstruction formula:

**Proposition 1.2.** A collection of vectors  $\{x_j\}_{j=1}^K$  in a finite-dimensional Hilbert space  $\mathbb{H}^d$  is a Parseval frame for  $\mathbb{H}^d$  if and only if

$$y = \sum_{j=1}^K \langle y, x_j \rangle x_j$$

for each  $y \in \mathbb{H}^d$ .

**Definition 1.3.** Let  $\{x_j\}_{j=1}^K$  be a frame for the finite-dimensional Hilbert space  $\mathbb{H}^d$ . The corresponding *frame operator*  $S : \mathbb{H}^d \rightarrow \mathbb{H}^d$  is defined by

$$S(x) = \sum_{j=1}^K \langle x, x_j \rangle x_j.$$

It can be seen as the composition  $S = \Theta\Theta^*$  of the *synthesis operator*  $\Theta : \mathbb{C}^K \rightarrow \mathbb{H}^d$  and its adjoint, the *analysis operator*  $\Theta^* : \mathbb{H}^d \rightarrow \mathbb{C}^K$ , given by the formulas

$$\Theta \left( \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_K \end{pmatrix} \right) = \sum_{j=1}^K c_j x_j \quad \text{and} \quad \Theta^*(x) = \begin{pmatrix} \langle x, x_1 \rangle \\ \langle x, x_2 \rangle \\ \vdots \\ \langle x, x_K \rangle \end{pmatrix}.$$

The frame operator is a bounded, invertible, self-adjoint operator satisfying  $AI_d \leq S \leq BI_d$ . Here and in what follows, we use  $I_d$  to denote the  $d \times d$  identity matrix and  $0_d$  to denote the  $d \times d$  zero matrix. A frame is Parseval if and only if its frame operator is the identity operator.

From both a pure and an applied point-of-view, construction of frames with desired properties has been a central question [Bownik and Jasper 2015]. In particular, much attention has been paid to tight frames with prescribed norms and general frames with both prescribed norms and frame operator. In the case of tight frames, the answer, the so-called ‘‘fundamental frame inequality’’, was provided by Casazza, Fickus, Kovačević, Leon, and Tremain:

**Theorem 1.4** [Casazza et al. 2006, Corollary 4.11]. *Given real numbers  $a_1 \geq a_2 \geq \dots \geq a_K > 0$ ,  $K \geq d$ , there exists a  $\lambda$ -tight frame  $\{x_j\}_{j=1}^K$  for a  $d$ -dimensional Hilbert space  $\mathbb{H}^d$  with prescribed norms  $\|x_j\|^2 = a_j$  for  $1 \leq j \leq K$  if and only if*

$$\lambda = \frac{1}{d} \sum_{j=1}^K a_j \geq a_1.$$

Casazza and Leon generalized this result to frames with prescribed frame operators (the classical case is when  $S = \lambda I_d$ ):

**Theorem 1.5** [Casazza and Leon 2010, Theorem 2.1]. *Let  $S$  be a positive self-adjoint operator on a  $d$ -dimensional Hilbert space  $\mathbb{H}^d$  with eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d > 0$ . Given real numbers  $a_1 \geq a_2 \geq \dots \geq a_K > 0$ ,  $K \geq d$ , there is a frame  $\{x_j\}_{j=1}^K$  for  $\mathbb{H}^d$  with frame operator  $S$  and  $\|x_j\|^2 = a_j$  for all  $1 \leq j \leq K$  if and only if*

$$\sum_{j=1}^K a_j = \sum_{j=1}^d \lambda_j \quad \text{and} \quad \sum_{j=1}^k a_j \leq \sum_{j=1}^k \lambda_j$$

for every  $1 \leq k \leq d$ .

This can be seen as a consequence of the classical Schur–Horn theorem [Bownik and Jasper 2015]. Cahill, Fickus, Mixon, Poteet, and Strawn [Cahill et al. 2013] introduced a so-called *eigenstep* method for constructing all frames with a given frame operator and set of norms; see also [Fickus et al. 2013], and [Bownik and Jasper 2015] for a survey of the topic.

A different approach was taken by Christensen, Powell, and Xiao [Christensen et al. 2012], extending Theorem 1.4 to the setting of dual frame pairs. Given a frame  $\{x_j\}_{j=1}^K$ , a sequence  $\{y_j\}_{j=1}^K$  is called a *dual frame* if, for every  $y \in \mathbb{H}^d$ ,

$$y = \sum_{j=1}^K \langle y, x_j \rangle y_j = \sum_{j=1}^K \langle y, y_j \rangle x_j.$$

**Theorem 1.6** [Christensen et al. 2012, Theorem 3.1]. *Given a sequence of numbers  $\{a_j\}_{j=1}^K \subset \mathbb{H}$  with  $K > d$ , the following are equivalent:*

- (1) *There exist dual frames  $\{x_j\}_{j=1}^K$  and  $\{y_j\}_{j=1}^K$  for  $\mathbb{H}^d$  such that  $\langle x_j, y_j \rangle = a_j$  for all  $1 \leq j \leq K$ .*
- (2) *There exists a tight frame  $\{x_j\}_{j=1}^K$  and dual frame  $\{y_j\}_{j=1}^K$  for  $\mathbb{H}^d$  such that  $\langle x_j, y_j \rangle = a_j$  for all  $1 \leq j \leq K$ .*
- (3)  $\sum_{j=1}^K a_j = d$ .

The goal of this paper is to extend the theory of frames with prescribed norms (or inner products) from the classical Hilbert spaces of  $\mathbb{C}^d$  and  $\mathbb{R}^d$  to the binary space  $\mathbb{Z}_2^d$ . We provide analogs of Theorems 1.4, 1.5, and 1.6 for binary frames.

The challenge, of course, is the lack of an inner product, positive elements, and guaranteed eigenvalues. [Section 2](#) contains background material on binary frames. In [Section 3](#), we explore dual binary frames and prove the binary versions of [Theorems 1.6](#) and [1.4](#). In [Section 4](#), we construct binary frames with prescribed “norms” and frame operator, as an analog to [Theorem 1.5](#). We conclude in [Section 5](#) with examples and a catalog.

## 2. Binary frames

Bodmann, Le, Reza, Tobin, and Tomforde [[Bodmann et al. 2009](#)] introduced a theory of frames over the  $d$ -dimensional binary space  $\mathbb{Z}_2^d$ , where  $\mathbb{Z}_2^d$  is the direct product  $\mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$  having  $d \geq 1$  copies of  $\mathbb{Z}_2$ . The main trouble in defining frames in a binary space stems from the lack of an ordering on  $\mathbb{Z}_2$ . Without an order, there can be no inner product defined for binary space. In spite of this, [[Bodmann et al. 2009](#)] establishes the dot product as the analog of the inner product on  $\mathbb{R}^d$  and  $\mathbb{C}^d$ .

**Definition 2.1** [[Bodmann et al. 2009](#)]. The dot product on  $\mathbb{Z}_2^d$  is defined as the map  $(\cdot, \cdot) : \mathbb{Z}_2^d \times \mathbb{Z}_2^d \rightarrow \mathbb{Z}_2$  given by

$$(a, b) = \sum_{n=1}^d a[n]b[n].$$

Due to the degenerate nature of the dot product (note that  $(a, a) = 0$  need not imply  $a = 0$ ), it fails to help define a frame in the manner of [Definition 1.1](#). However, when working over finite-dimensional spaces in the classical case, a frame is merely a spanning sequence of vectors. This motivates the definition of a frame in binary space.

**Definition 2.2** [[Bodmann et al. 2009](#)]. A frame is a sequence of vectors  $\mathcal{F} = \{f_j\}_{j=1}^K$  in  $\mathbb{Z}_2^d$  such that  $\text{Span}(\mathcal{F}) = \mathbb{Z}_2^d$ .

The synthesis, analysis, and frame operators of  $\mathcal{F}$  are defined similarly to [Definition 1.3](#) and are denoted by  $\Theta_{\mathcal{F}}$ ,  $\Theta_{\mathcal{F}}^*$ , and  $S_{\mathcal{F}}$ , respectively.

**Definition 2.3** [[Bodmann et al. 2009](#)]. The synthesis operator of a frame  $\mathcal{F} = \{f_j\}_{j=1}^K$  is the  $d \times K$  matrix whose  $i$ -th column is the  $i$ -th vector in  $\mathcal{F}$ . The analysis operator  $\Theta_{\mathcal{F}}^*$  is the transpose of the synthesis operator. Explicitly,

$$\Theta_{\mathcal{F}} = \begin{bmatrix} | & & | \\ f_1 & \cdots & f_K \\ | & & | \end{bmatrix} \quad \text{and} \quad \Theta_{\mathcal{F}}^* = \begin{bmatrix} -f_1^* - \\ \vdots \\ -f_K^* - \end{bmatrix}.$$

The frame operator is  $S_{\mathcal{F}} = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$ .

It is demonstrated in [[Bodmann et al. 2009](#)] that the spanning property of  $\mathcal{F}$  is necessary and sufficient for  $\mathcal{F}$  to have a reconstruction identity with a dual family  $\mathcal{G}$ .

This fact is summed up in the following theorem and is shown by choosing a basis consisting of  $d$  vectors in  $\mathcal{F}$  (without loss of generality, assumed to be  $f_1, \dots, f_d$ ) and applying the Riesz representation theorem to the linear functionals  $\gamma_i$  defined by  $\gamma_i(f_j) = \delta_{ij}$ .

**Theorem 2.4** [Bodmann et al. 2009, Theorem 2.4]. *The family  $\mathcal{F} = \{f_j\}_{j=1}^K$  in  $\mathbb{Z}_2^d$  is a frame if and only if there exist vectors  $\mathcal{G} = \{g_j\}_{j=1}^K$  such that, for all  $y \in \mathbb{Z}_2^d$ ,*

$$y = \sum_{j=1}^K (y, g_j) f_j. \quad (1)$$

In the proof,  $g_i$  is defined as the unique vector satisfying  $\gamma_i(y) = (y, g_i)$  for every  $y$  for  $1 \leq i \leq d$ , and  $g_i = 0$  for  $d < i \leq K$ . Equation (1) can be rewritten as

$$\Theta_{\mathcal{F}} \Theta_{\mathcal{G}}^* = I_d,$$

which is equivalent to  $\Theta_{\mathcal{G}} \Theta_{\mathcal{F}}^* = I_d$ . Consequently,  $\mathcal{G}$  is a dual frame to  $\mathcal{F}$ . We will refer to the dual frame  $\mathcal{G}$  as a *natural dual* to  $\mathcal{F}$ . Note that this definition is unrelated to the usual definition of the canonical dual in  $\mathbb{C}^d$  or  $\mathbb{R}^d$  as  $\{S^{-1}(f_j)\}$ , where  $S = \Theta \Theta^*$  is the frame operator from Definition 1.3. Although  $S_{\mathcal{F}}$  is no longer necessarily invertible, we still have

$$S_{\mathcal{F}}(g_i) = \sum_{j=1}^K (g_i, f_j) f_j = \sum_{j=1}^K \delta_{ij} f_j = f_i$$

for  $i \leq d$ .

Propositions 2.5 and 2.6 make clear that the natural dual frame is unique, up to permutation, if and only if  $K = d$ .

**Proposition 2.5.** *Let  $\mathcal{F} = \{f_j\}_{j=1}^K$  be a frame for  $\mathbb{Z}_2^d$  with a natural dual frame  $\mathcal{G}$ . Then  $\mathcal{H}$  is a dual frame of  $\mathcal{F}$  if and only if  $\Theta_{\mathcal{H}}^* = \Theta_{\mathcal{G}}^* + C$  for some  $K \times d$  matrix  $C$  satisfying  $\Theta_{\mathcal{F}} C = 0_d$ .*

*Proof.* Given the existence of a matrix  $C$  with  $\Theta_{\mathcal{H}}^* = \Theta_{\mathcal{G}}^* + C$  and  $\Theta_{\mathcal{F}} C = 0_d$ , it is immediate that  $\Theta_{\mathcal{F}} \Theta_{\mathcal{H}}^* = \Theta_{\mathcal{F}} \Theta_{\mathcal{G}}^* = I_d$ . Conversely, if  $\mathcal{H}$  is a dual frame of  $\mathcal{F}$ , then letting  $C = \Theta_{\mathcal{H}}^* - \Theta_{\mathcal{G}}^*$  gives  $\Theta_{\mathcal{F}} C = \Theta_{\mathcal{F}} \Theta_{\mathcal{H}}^* - \Theta_{\mathcal{F}} \Theta_{\mathcal{G}}^* = I_d - I_d = 0_d$ .  $\square$

The following result is well known in  $\mathbb{R}^d$  and  $\mathbb{C}^d$ ; see [Han et al. 2007, Proposition 6.3]. Since the proof in that text uses the invertibility of the frame operator, we provide a modified proof for  $\mathbb{Z}_2^d$  here.

**Proposition 2.6.** *Let  $\mathcal{F} = \{f_j\}_{j=1}^K$  be a frame for  $\mathbb{Z}_2^d$ . Then  $\mathcal{F}$  has a unique dual frame if and only if  $\mathcal{F}$  is a basis.*

*Proof.* Since a frame is a spanning set,  $\mathcal{F}$  is a basis if and only if the vectors  $f_j$  are linearly independent and  $K = d$ . This is equivalent to the only  $K \times d$  matrix  $C$  satisfying  $\Theta_{\mathcal{F}} C = 0_d$  being the zero matrix. By Proposition 2.5, this happens if and only if the (unique choice of) natural dual  $\mathcal{G}$  is the only dual frame of  $\mathcal{F}$ .  $\square$

The diagonal of the *Gramian* matrix  $\Theta_{\mathcal{F}}^* \Theta_{\mathcal{F}}$  is the vector whose  $i$ -th entry is  $(f_i, f_i)$ ; when  $\mathcal{F}$  and  $\mathcal{H}$  are a dual frame pair, the diagonal of the *cross-Gramian* matrix  $\Theta_{\mathcal{F}}^* \Theta_{\mathcal{H}}$  is the vector whose  $i$ -th element is  $(h_i, f_i)$ .

**Definition 2.7** [Bodmann et al. 2009]. A *Parseval frame* for  $\mathbb{Z}_2^d$  is a sequence of vectors  $\mathcal{F} = \{f_j\}_{j=1}^K \subset \mathbb{Z}_2^d$  such that

$$y = \sum_{j=1}^K (y, f_j) f_j$$

for all  $y \in \mathbb{Z}_2^d$ .

Note that a binary Parseval frame must be a binary frame. In matrix notation,  $\mathcal{F}$  is a Parseval frame for  $\mathbb{Z}_2^d$  if and only if  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^* = I_d$ . If a collection of vectors  $\{x_j\} \subset \mathbb{Z}_2^d$  satisfies  $(x_i, x_j) = 0$  for all  $i \neq j$  and  $(x_i, x_i) = 1$  for all  $i$ , we say, through a slight abuse of terminology, that  $\{x_j\}$  is an orthonormal set. An easy, matrix-theoretical consequence of the definitions of frame and Parseval frame is the following proposition:

**Proposition 2.8.** Let  $\mathcal{F} = \{f_j\}_{j=1}^K$  be a sequence of vectors in  $\mathbb{Z}_2^d$ .

- (1) The rows of  $\Theta_{\mathcal{F}}$  are linearly independent if and only if  $\mathcal{F}$  is a frame.
- (2) The rows of  $\Theta_{\mathcal{F}}$  are orthonormal if and only if  $\mathcal{F}$  is a Parseval frame.

In the rest of this paper, unless otherwise noted, all vectors are elements of the binary vector spaces  $\mathbb{Z}_2^d$  or  $\mathbb{Z}_2^K$ . All operations are performed modulo 2; for example,  $\text{Tr}(A)$  represents the usual trace of a matrix  $A$ , but  $\text{Tr}(AB) \equiv \text{Tr}(BA) \pmod{2}$  for two binary matrices  $A$  and  $B$ . All *frames* refer to *binary frames*. Throughout, we denote the standard orthonormal basis in  $\mathbb{Z}_2^d$  by  $\{\varepsilon_1, \varepsilon_2, \dots, \varepsilon_d\}$ .

### 3. Dual and Parseval binary frames

If  $\mathcal{F} = \{f_j\}_{j=1}^K$  is a frame for  $\mathbb{Z}_2^d$  and  $K = d$ , then the vectors  $\{f_j\}$  must be linearly independent and hence a basis with a unique (natural) dual  $\mathcal{G}$ . In this case,  $(f_j, g_j) = \gamma_j(f_j) = 1$  for all  $1 \leq j \leq K$ . In this section, we are largely concerned with the question of which sequences  $\alpha \in \mathbb{Z}_2^K$  satisfy  $\alpha[j] = (f_j, h_j)$  for a dual frame pair  $(\mathcal{F}, \mathcal{H})$ ; so we assume  $K > d$ . We use  $\|\alpha\|_0$  to denote the number of nonzero entries in a vector  $\alpha$ , the parity of which will be fundamental in this paper.

**Lemma 3.1.** Let  $\mathcal{F} = \{f_j\}_{j=1}^K$  be a frame for  $\mathbb{Z}_2^d$  and let  $\pi$  be a permutation of the set  $\{1, 2, \dots, K\}$ . Denote by  $\mathcal{F}_{\pi}$  the frame  $\{f_{\pi(j)}\}_{j=1}^K$ . Then a frame  $\mathcal{H}$  is a dual frame of  $\mathcal{F}$  if and only if  $\mathcal{H}_{\pi}$  is a dual frame of  $\mathcal{F}_{\pi}$ . Furthermore, if  $\alpha$  is a sequence in  $\mathbb{Z}_2^K$ , then the dual frame pair  $(\mathcal{F}, \mathcal{H})$  satisfies  $(f_j, h_j) = \alpha[j]$  for every  $j$  if and only if the dual frame pair  $(\mathcal{F}_{\pi}, \mathcal{H}_{\pi})$  satisfies  $(f_{\pi(j)}, h_{\pi(j)}) = \alpha[\pi(j)]$  for every  $j$ .

*Proof.* Suppose  $\Theta_{\mathcal{F}} \Theta_{\mathcal{H}}^* = I_d$  and  $\text{diag}(\Theta_{\mathcal{H}}^* \Theta_{\mathcal{F}}) = \alpha$ . Then

$$\begin{aligned} \Theta_{\mathcal{F}_{\pi}} \Theta_{\mathcal{H}_{\pi}}^* &= \begin{bmatrix} | & & | \\ f_{\pi(1)} & \cdots & f_{\pi(K)} \\ | & & | \end{bmatrix} \begin{bmatrix} -h_{\pi(1)}^* - \\ \vdots \\ -h_{\pi(K)}^* - \end{bmatrix} \\ &= \begin{bmatrix} | & & | \\ f_{\pi(1)} & \cdots & f_{\pi(K)} \\ | & & | \end{bmatrix} P^* P \begin{bmatrix} -h_{\pi(1)}^* - \\ \vdots \\ -h_{\pi(K)}^* - \end{bmatrix} \\ &= \begin{bmatrix} | & & | \\ f_1 & \cdots & f_K \\ | & & | \end{bmatrix} \begin{bmatrix} -h_1^* - \\ \vdots \\ -h_K^* - \end{bmatrix} \\ &= \Theta_{\mathcal{F}} \Theta_{\mathcal{H}}^* = I_d, \end{aligned}$$

where

$$P = \begin{bmatrix} -\varepsilon_{\pi^{-1}(1)}^* - \\ \vdots \\ -\varepsilon_{\pi^{-1}(K)}^* - \end{bmatrix},$$

and here we use  $\varepsilon_i$  to indicate the  $i$ -th standard basis vector in  $\mathbb{Z}_2^K$ . Thus  $\mathcal{F}_{\pi}$  and  $\mathcal{H}_{\pi}$  are dual frames and  $(f_j, h_j) = \alpha[j]$  for each  $j$  implies  $(f_{\pi(j)}, h_{\pi(j)}) = \alpha[\pi(j)]$ . For the converse statements, let  $\sigma = \pi^{-1}$ .  $\square$

The next theorem and corollary are the analog of [Theorem 1.6](#) in binary space.

**Theorem 3.2.** *Given  $\alpha \in \mathbb{Z}_2^K$ , there exists a dual frame pair  $(\mathcal{F}, \mathcal{H})$  for  $\mathbb{Z}_2^d$  with  $(f_i, h_i) = \alpha[i]$  for every  $i$  if and only if  $\|\alpha\|_0 \equiv d \pmod{2}$ .*

*Proof.* Suppose  $(\mathcal{F}, \mathcal{H})$  is a dual frame pair for  $\mathbb{Z}_2^d$  such that  $(f_i, h_i) = \alpha[i]$  for every  $i$ . Then

$$\|\alpha\|_0 \equiv \text{Tr}(\Theta_{\mathcal{H}}^* \Theta_{\mathcal{F}}) \equiv \text{Tr}(\Theta_{\mathcal{F}} \Theta_{\mathcal{H}}^*) \equiv \text{Tr}(I_d) \equiv d \pmod{2}.$$

Conversely, suppose  $\|\alpha\|_0 \equiv d \pmod{2}$ . We consider three cases.

Case 1:  $\|\alpha\|_0 = d$ . Let  $f_j = \varepsilon_j$  for  $1 \leq j \leq d$  and let  $f_j$  be arbitrary for  $d < j \leq K$ . A natural dual frame is then given by  $g_j = \varepsilon_j$  for  $1 \leq j \leq d$  and  $g_j = 0$  for  $d < j \leq K$ . Define  $\beta \in \mathbb{Z}_2^K$  by  $\beta[j] = 1$  for  $1 \leq j \leq d$  and  $\beta[j] = 0$  for  $d < j \leq K$ , and let  $\pi$  be a permutation of  $\{1, 2, \dots, K\}$  such that  $\beta[j] = \alpha[\pi(j)]$ . It follows that

$$\text{diag}(\Theta_G^* \Theta_{\mathcal{F}}) = \beta.$$

By [Lemma 3.1](#),  $(\mathcal{F}_{\pi^{-1}}, \mathcal{G}_{\pi^{-1}})$  is the desired dual frame pair with  $(f_{\pi^{-1}(j)}, g_{\pi^{-1}(j)}) = \beta[\pi^{-1}(j)] = \alpha[j]$  for each  $j$ .

Case 2:  $\|\alpha\|_0 = d + 2t$  for some positive integer  $t \leq \frac{1}{2}(K - d)$ . Consider the frame  $\mathcal{F}$  defined in Case 1 above, but set  $f_j = \varepsilon_1$  for all  $d + 1 \leq j \leq d + 2t$ . A



natural dual frame  $\mathcal{G}$  of  $\mathcal{F}$  is the same as that defined in Case 1 above. Let  $C$  be the  $K \times d$  matrix whose top  $d \times d$  block is  $0_d$  and rows  $d + 1$  through  $d + 2t$  are equal to  $\varepsilon_1^*$ . The remaining rows of  $C$  are zeros. Due to the introduction of an even number of  $\varepsilon_1^*$ 's, we see that  $\Theta_{\mathcal{F}} C = 0_d$ , and hence  $\mathcal{H}$  given by the rows of  $\Theta_{\mathcal{H}}^* = \Theta_{\mathcal{G}}^* + C$  is a dual frame of  $\mathcal{F}$ , by [Proposition 2.5](#). Since  $\text{diag}(\Theta_{\mathcal{H}}^* \Theta_{\mathcal{F}})$  is the vector composed of  $d + 2t$  ones followed by  $K - (d + 2t)$  zeros, [Lemma 3.1](#) again implies the existence of the desired dual frame pair.

**Case 3:**  $\|\alpha\|_0 = d - 2t$  for some positive integer  $t \leq \frac{1}{2}d$ . Consider again the frame  $\mathcal{F}$  defined in Case 1, except set  $f_{d+1} = \varepsilon_d + \varepsilon_{d-1} + \cdots + \varepsilon_{d-2t+2} + \varepsilon_{d-2t+1}$ . We again take the same natural dual frame  $\mathcal{G}$  as in Case 1 above. Let  $C$  be the  $K \times d$  matrix whose top  $d - 2t$  rows are zeros, rows  $d - 2t + 1$  through  $d + 1$  are  $f_{d+1}^*$ , and the remaining rows are zeros. Then  $\Theta_{\mathcal{F}} C = 0_d$ , and hence  $\mathcal{H}$  defined by  $\Theta_{\mathcal{H}}^* = \Theta_{\mathcal{G}}^* + C$  is a dual frame of  $\mathcal{F}$ . Due to the presence of an even number of ones in  $f_{d+1}$ , we have  $(f_{d+1}, f_{d+1}) = 0$ , while  $(f_j, h_j) = 0$  for  $d - 2t + 1 \leq j \leq d$ . Consequently,  $\text{diag}(\Theta_{\mathcal{H}}^* \Theta_{\mathcal{F}})$  consist of ones in its first  $d - 2t$  entries followed by  $K - (d - 2t)$  zeros. [Lemma 3.1](#) again completes the proof.  $\square$

**Corollary 3.3.** *Given  $\alpha \in \mathbb{Z}_2^K$ , there exists a Parseval frame  $\mathcal{F}$  and a corresponding dual frame  $\mathcal{H}$  for  $\mathbb{Z}_2^d$  with  $(f_i, h_i) = \alpha[i]$  for every  $i$  if and only if  $\|\alpha\|_0 \equiv d \pmod{2}$ .*

*Proof.* The necessity of the condition on  $\|\alpha\|_0$  follows immediately by [Theorem 3.2](#). The sufficiency depends on slight modifications of the frame  $\mathcal{F}$  constructed in the proof of [Theorem 3.2](#). In Case 1, instead of letting  $f_j$  for  $d < j \leq K$  be arbitrary, set each of those vectors to be the zero vector,  $\vec{0}$ , in  $\mathbb{Z}_2^d$ . [Proposition 2.8](#) implies  $\mathcal{F}$  is a Parseval frame. Similarly, the frame built in Case 2 is a Parseval frame if we set  $f_j = \vec{0}$  for  $2t + 1 \leq j \leq K$ . The frame built in Case 3 is not a Parseval frame; however, consider instead the frame  $\mathcal{F}'$  defined as  $f'_j = f_j$  for  $1 \leq j \leq d + 1$ ,  $f'_{d+2} = f_{d+1}$ , and  $f'_j = \vec{0}$  for  $d + 3 \leq j \leq K$ . By [Proposition 2.8](#),  $\mathcal{F}'$  is a Parseval frame. Note that each column of the matrix  $C$  constructed in Case 3 is still a (possibly trivial) dependence relation among the columns of  $\Theta_{\mathcal{F}'}$ , which implies  $\Theta_{\mathcal{F}'} C = 0_d$ . Since the natural dual  $\mathcal{G}$  of  $\mathcal{F}$  constructed in Case 3 is still a natural dual of  $\mathcal{F}'$ , the frame  $\mathcal{H}$  with analysis operator  $\Theta_{\mathcal{H}}^* = \Theta_{\mathcal{G}}^* + C$  is a dual frame of  $\mathcal{F}'$ ; moreover,  $(f'_j, h_j) = (f_j, h_j)$  for all  $j$  since  $h_{d+2} = 0$ .  $\square$

**Remark 3.4.** The Parseval frames built in Cases 1 and 2 of the above corollary, in fact, satisfy  $(f_j, f_j) = \alpha[j]$  for each  $j$  after a suitable permutation, as allowed by [Lemma 3.1](#). However, this is not true in Case 3. By constructing a Parseval frame that satisfies  $(f_j, f_j) = \alpha[j]$  for each  $j$  in the case when  $\|\alpha\|_0 = d - 2t$  for some positive integer  $t \leq \frac{1}{2}d$ , we will prove the binary analog of [Theorem 1.4](#).

**Theorem 3.5.** *Given nonzero  $\alpha \in \mathbb{Z}_2^K$ , there exists a Parseval frame  $\mathcal{F}$  for  $\mathbb{Z}_2^d$  with  $(f_i, f_i) = \alpha[i]$  for every  $i$  if and only if  $\|\alpha\|_0 \equiv d \pmod{2}$ .*

*Proof.* Since a Parseval frame is self-dual, the necessity of the condition on  $\|\alpha\|_0$  follows immediately from [Theorem 3.2](#). For sufficiency, [Remark 3.4](#) implies that we need only construct Parseval frames satisfying  $(f_i, f_i) = \alpha[i]$  for every  $i$  for  $\|\alpha\|_0 < d$ . Since  $\|\alpha\|_0 \equiv d \pmod{2}$  and  $\|\alpha\|_0 \neq 0$ , we must have  $d \geq 3$ .

For  $d = 3$  and  $\|\alpha\|_0 = 1$ , we build the Parseval frame

$$\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\},$$

and note that we may permute the vectors as needed. Moreover, we may insert any number of copies of  $\vec{0}$  to satisfy any  $K > 4$ . By augmenting each vector with a last entry of 0 and inserting the vector  $\varepsilon_4 \in \mathbb{Z}_2^4$ , we construct the Parseval frame

$$\left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

for  $\mathbb{Z}_2^4$  that, after suitable permutation and inclusion of copies of  $\vec{0}$ , satisfies any  $\|\alpha\|_0 = 2$ .

Given any odd dimension  $d$ , suppose we have constructed, without zero vectors, the Parseval frames  $\mathcal{F}^1, \mathcal{F}^3, \dots, \mathcal{F}^{d-4}$  for  $\mathbb{Z}_2^{d-2}$  corresponding to  $\|\alpha\|_0 = 1, 3, \dots, d-4$ . For each odd  $n$ , create the collection  $\tilde{\mathcal{F}}^{n+2}$  by augmenting each vector of  $\mathcal{F}^n$  with two zero entries and unioning the augmented vectors with  $\varepsilon_{d-1}, \varepsilon_d \in \mathbb{Z}_2^d$ . Then  $\tilde{\mathcal{F}}^3, \tilde{\mathcal{F}}^5, \dots, \tilde{\mathcal{F}}^{d-2}$  are Parseval frames for  $\mathbb{Z}_2^d$  corresponding to  $\|\alpha\|_0 = 3, 5, \dots, d-2$ . Let  $\tilde{\mathcal{F}}^1 = \{\vec{1} + \varepsilon_1, \vec{1} + \varepsilon_2, \dots, \vec{1} + \varepsilon_d, \vec{1}\}$ , where  $\vec{1}$  represents the vector with  $d$  ones in  $\mathbb{Z}_2^d$ . Then  $\tilde{\mathcal{F}}^1$  is a Parseval frame for  $\mathbb{Z}_2^d$  corresponding to  $\|\alpha\|_0 = 1$ .

If the dimension  $d$  is even, we create the Parseval frames  $\tilde{\mathcal{F}}^{n+1}$  from  $\tilde{\mathcal{F}}^n$  for each  $n = 1, 3, \dots, d-3$ , corresponding to  $\|\alpha\|_0 = 2, 4, \dots, d-2$ : augment each vector of  $\tilde{\mathcal{F}}^n$  with a last entry of 0 and insert the vector  $\varepsilon_d \in \mathbb{Z}_2^d$ .

In both cases, permutation of the vectors and possible inclusion of copies of  $\vec{0}$  finishes the proof.  $\square$

#### 4. Binary frames with prescribed frame operator

In the previous section, we gave a necessary and sufficient condition on  $\alpha \in \mathbb{Z}_2^K$  for the existence of a Parseval frame  $\mathcal{F}$  for  $\mathbb{Z}_2^d$  with  $(f_j, f_j) = \alpha[j]$  for every  $j$ . In classical frame theory over  $\mathbb{R}$  or  $\mathbb{C}$ , the characterization has been broadened to frames with a given frame operator and specified values for  $\|f_j\|$  (the case of a Parseval frame is when  $S = I$ ), as in [Theorem 1.5](#). In the classical case, the frame operator is a symmetric, invertible, positive definite matrix. For a binary frame  $\mathcal{F}$ ,  $S_{\mathcal{F}} = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$

is not necessarily invertible; for example, the zero matrix is the frame operator of any frame in which every vector occurs twice. Consequently, we must first characterize those binary symmetric matrices that are frame operators of binary frames.

Throughout this section, we rely heavily on the idea of vector parity in  $\mathbb{Z}_2^d$ .

**Definition 4.1.** Describe a vector  $v \in \mathbb{Z}_2^d$  as *even* if  $(v, v) = 0$ . Equivalently, a vector is even if  $\|v\|_0 \equiv 0 \pmod{2}$ . If a vector is not even, then it is *odd*.

**Lemma 4.2.** (1) *The sum of two even vectors is an even vector.*

(2) *The sum of two odd vectors is an even vector.*

(3) *The sum of an odd vector and an even vector is an odd vector.*

*Proof.* This follows from the above definition and the observation that if  $u, v \in \mathbb{Z}_2^d$ , then

$$(u + v, u + v) = (u, u) + (u, v) + (v, u) + (v, v) = (u, u) + (v, v). \quad \square$$

As a consequence of this lemma, we note that a collection of only even vectors cannot span  $\mathbb{Z}_2^d$ .

Given a  $d \times d$  symmetric matrix  $S$ , we call  $A$  a *factor* of  $S$  if  $S = AA^*$ . We say  $A$  is a *minimal factor* if it has the minimum number of columns over all factors of  $S$ . Minimal factorization of symmetric binary matrices also arises in the computation of the covering radius of Reed–Muller codes [Cohen et al. 1997].

**Theorem 4.3** [Lempel 1975, Theorem 1]. *Every binary symmetric matrix  $S$  can be factorized as  $S = AA^*$  for some binary matrix  $A$ . The number of columns of a minimal factor of  $S$  is  $\text{rank}(S)$  if  $\text{diag}(S) \neq \vec{0}$  and  $\text{rank}(S) + 1$  if  $\text{diag}(S) = \vec{0}$ .*

**Proposition 4.4.** *If  $S = AA^*$  for some  $d \times m$  matrix  $A$ , where  $m = \text{rank}(S)$  or  $m = \text{rank}(S) + 1$ , then  $\text{rank}(A) = \text{rank}(S)$ .*

*Proof.* Frobenius' rank inequality and the fact that, for properly sized matrices  $C$  and  $D$ ,  $\text{rank}(CD) \leq \min\{\text{rank}(C), \text{rank}(D)\}$  [Horn and Johnson 1985] imply

$$\text{rank}(A) + \text{rank}(A^*) \leq m + \text{rank}(S) \leq m + \text{rank}(A).$$

If  $m = \text{rank}(S)$ , the above inequalities simplify to

$$\text{rank}(A) \leq \text{rank}(S) \leq \text{rank}(A),$$

and hence  $\text{rank}(A) = \text{rank}(S)$ . If  $m = \text{rank}(S) + 1$ , we instead have

$$2 \text{rank}(A) \leq 2 \text{rank}(S) + 1 \leq 2 \text{rank}(A) + 1.$$

Since  $2 \text{rank}(A) = 2 \text{rank}(S) + 1$  is impossible, we must have  $\text{rank}(A) = \text{rank}(S)$ .  $\square$

We can use factors of a matrix to construct frames with a given frame operator. Minimal factors correspond to *minimal frames*, that is, frames with the fewest number of elements. In the previous section, we disregarded frames  $\{f_j\}_{j=1}^K$  that

were bases since they corresponded to unique duals  $\{g_j\}_{j=1}^K$  with predetermined values for the dot products  $(f_j, g_j)$ . In this section, however, we are concerned with  $(f_j, f_j)$ , so we do not rule the case  $K = d$  out of consideration.

**Theorem 4.5.** *Let  $S$  be a  $d \times d$  symmetric matrix with  $\text{rank}(S) = d$ . There exists a (minimal)  $d$ -element frame  $\mathcal{F}$  (i.e., basis) such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  if and only if  $\text{diag}(S) \neq \vec{0}$ . If  $\text{diag}(S) = \vec{0}$ , then there exists a (minimal)  $(d+1)$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$ .*

*Proof.* Suppose there is a  $d$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$ . If  $\text{diag}(S) = \vec{0}$ , then  $\Theta_{\mathcal{F}}$  is a  $d \times d$  square matrix with all even rows, which cannot span  $\mathbb{Z}_2^d$ . Hence, the columns of  $\Theta_{\mathcal{F}}$  cannot span  $\mathbb{Z}_2^d$ , contradicting  $\mathcal{F}$  being a frame. Conversely, suppose  $\text{diag}(S) \neq \vec{0}$ . By [Theorem 4.3](#), there exists a  $d \times k$  matrix  $A$  such that  $S = AA^*$  and  $k = \text{rank}(S)$ . By [Proposition 4.4](#),  $\text{rank}(A) = \text{rank}(S) = k = d$ . Thus,  $A$  is a  $d \times d$  matrix whose columns span  $\mathbb{Z}_2^d$ . Defining  $\mathcal{F}$  by  $\Theta_{\mathcal{F}} = A$  constructs the  $d$ -element frame.

Now suppose  $\text{diag}(S) = \vec{0}$ . By [Theorem 4.3](#), there exists a minimal  $d \times k$  factor  $A$  such that  $S = AA^*$  and  $k = \text{rank}(S) + 1$ . By [Proposition 4.4](#),  $\text{rank}(A) = \text{rank}(S) = d$ . Therefore, the columns of  $A$  span  $\mathbb{Z}_2^d$ , and we can construct a  $(d+1)$ -element frame by taking  $\Theta_{\mathcal{F}} = A$ .  $\square$

**Remark 4.6.** The columns of  $\Theta_{\mathcal{F}}$  can be augmented by copies of the zero vector without affecting  $S$ , so nonminimal frames can always be constructed from minimal frames by including any number of copies of the zero vector.

Next we construct minimal frames whose prescribed frame operators are not of full rank. Let  $r(A)$  and  $c(A)$  denote the number of rows and columns of a matrix  $A$ , respectively.

**Lemma 4.7** [[Lempel 1975](#), Section 4]. *Let  $S$  be a  $d \times d$  symmetric matrix of  $\text{rank}(S) < d$ . There exists a permutation matrix  $P$  and a nonsingular matrix  $T$  such that*

$$S = P^* \begin{bmatrix} L & M \\ M^* & K \end{bmatrix} P = P^* T \begin{bmatrix} L & 0 \\ 0 & 0 \end{bmatrix} T^* P,$$

where  $L$  is a symmetric matrix with  $r(L) = \text{rank}(L) = \text{rank}(S)$ .

**Corollary 4.8.** *A  $d \times d$  symmetric matrix  $S$  with  $\text{diag}(S) = \vec{0}$  must have even rank.*

*Proof.* It is known that if  $\text{rank}(S) = d$ , then  $d$  must be even; see, for example, [[Cohen et al. 1997](#), Section 9.3]. Suppose  $\text{rank}(S) < d$ . The  $\text{rank}(S) \times \text{rank}(S)$  symmetric matrix  $L$  constructed in [Lemma 4.7](#) has  $\text{rank}(L) = \text{rank}(S)$ . Since the diagonal elements of  $S$  are the diagonal elements of  $L$  and  $K$ ,  $\text{diag}(L) = \vec{0}$ . So  $\text{rank}(S)$  must be even.  $\square$

**Theorem 4.9.** *Let  $S$  be a  $d \times d$  symmetric matrix with  $\text{rank}(S) < d$ . There exists a  $k$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  if and only if  $k \geq 2d - \text{rank}(S)$ .*

*Proof.* Necessity follows from Frobenius’s rank inequality

$$\text{rank}(\Theta_{\mathcal{F}}) + \text{rank}(\Theta_{\mathcal{F}}^*) \leq k + \text{rank}(S)$$

since frames are spanning sets.

Conversely, let  $k$  be an integer such that  $k \geq 2d - \text{rank}(S)$ . Let  $L, P, T$  be the matrices guaranteed by Lemma 4.7, and let  $V = P^*T$ . Suppose  $\text{diag}(L) \neq \vec{0}$ . By Theorem 4.3 there exists a factor  $H$  of  $L$  such that

$$\begin{bmatrix} L & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} H \\ 0 \end{bmatrix} \begin{bmatrix} H^* & 0 \end{bmatrix}$$

and  $r(H) = c(H) = \text{rank}(L)$ . Consider the augmented matrix

$$A = \begin{bmatrix} H & 0 \\ 0 & B \end{bmatrix},$$

where the columns of  $B$  are the standard basis vectors of  $\mathbb{Z}_2^{d-\text{rank}(S)}$ , each repeated twice. Then  $r(A) = d$ ,  $c(A) = 2d - \text{rank}(S)$ , and  $AA^* = \begin{bmatrix} L & 0 \\ 0 & 0 \end{bmatrix}$ . By construction,  $\text{rank}(A) = d$ . Since  $V$  is nonsingular,  $VA$  is a  $d \times (2d - \text{rank}(S))$  matrix of rank  $d$  such that  $S = VAA^*V^*$ . If  $k = 2d - \text{rank}(S)$ , let a minimal frame  $\mathcal{F}$  be the columns of  $VA$ ; if  $k > 2d - \text{rank}(S)$ , augment the columns of  $VA$  with the necessary number of zero vectors.

Now suppose  $\text{diag}(L) = \vec{0}$ . By Corollary 4.8,  $\text{rank}(L) = \text{rank}(S)$  must be even. As above (by Theorem 4.3) we can factorize  $L$  with a matrix  $H$ , but now  $r(H) = \text{rank}(L)$  and  $c(H) = \text{rank}(L) + 1$ . In this case, we build the augmented matrix

$$\tilde{A} = \left[ \begin{array}{c|c} H & 0 \\ \hline r_1 & r_2 \\ \hline 0 & \tilde{B} \end{array} \right],$$

where  $r_1 = [1 \ 1 \ 1 \ \cdots \ 1]$  is a vector of length  $c(H)$ ,  $r_2 = [1 \ 0 \ 0 \ \cdots \ 0]$  has length  $2(d - \text{rank}(S)) - 1$ , and the columns of  $\tilde{B}$  are the zero vector followed by the standard basis vectors of  $\mathbb{Z}_2^{d-(\text{rank}(S)+1)}$ , each repeated twice. Since the  $(i, j)$  entry of the product  $\tilde{A}\tilde{A}^*$  can be viewed as the dot product of the  $i$ -th and  $j$ -th rows of  $\tilde{A}$ , it is easy to see that  $\tilde{A}\tilde{A}^* = \begin{bmatrix} L & 0 \\ 0 & 0 \end{bmatrix}$ . Indeed, since  $\text{diag}(L) = \vec{0}$ , each row of  $H$  is an even vector. Since  $c(H)$  is odd, the vector  $[r_1 \ | \ r_2]$  is even, as is each row of  $\tilde{B}$ . By construction,  $\text{rank}(\tilde{A}) = d$ ,  $r(\tilde{A}) = r(H) + 1 + d - \text{rank}(S) - 1 = d$ , and  $c(\tilde{A}) = c(H) + 1 + 2(d - \text{rank}(S)) - 1 = 2d - \text{rank}(S)$ . As above let  $\mathcal{F}$  consist of the columns of  $V\tilde{A}$  if  $k = 2d - \text{rank}(S)$  or these columns together with  $k - 2d + \text{rank}(S)$  copies of the zero vector if  $k > 2d - \text{rank}(S)$ . □

Theorems 4.5 and 4.9 provide minimal (and nonminimal) frames with frame operator  $S$ , subject only to restrictions based on  $\text{rank}(S)$ . In what follows, sometimes

we will make additional assumptions on  $S$ , which allow the construction of frames with frame operator  $S$  in different, and sometimes more intuitive, ways.

**Definition 4.10.** A  $d \times d$  symmetric matrix  $S$  is said to be *parity indicative* if, for every  $1 \leq i \leq d$ , the diagonal entry  $S_{ii}$  is equal to 1 if and only if the  $i$ -th row of  $S$  is odd.

**Lemma 4.11.** *Let  $S$  be a  $d \times d$  symmetric matrix, and suppose  $S = AA^*$  for some  $d \times m$  matrix  $A$ . If every column of  $A$  is odd, then  $S$  is parity indicative. Conversely, if  $S$  is parity indicative and the columns of  $A$  are linearly independent, then every column of  $A$  must be odd.*

*Proof.* Assume  $S = AA^* = \sum_{i=1}^m a_i a_i^*$ , where each column  $a_i$  of  $A$  is odd. The  $i$ -th row of  $S$  equals the sum of those  $a_j^*$  satisfying  $a_j[i] = 1$ . Suppose the  $i$ -th row of  $S$  is odd. Then this sum must be composed of an odd number of nonzero terms by Lemma 4.2. That is, there is an odd number of indices  $j$  having  $a_j[i] = 1$ . Consequently, the  $i$ -th row of  $A$  is an odd vector, and  $S_{ii} = \sum_{j=1}^m a_j[i] a_j[i] = 1$ . On the other hand, if the  $i$ -th row of  $S$  is even, then Lemma 4.2 implies that an even number of  $a_j$  have  $a_j[i] = 1$ , resulting in  $S_{ii} = 0$ .

To show the converse, assume that the columns of  $A$  are linearly independent. Suppose some of the columns of  $A$  are even; denote the odd columns by  $\{o_j\}$  and the even columns by  $\{e_j\}$ . If for every index  $l$ , we have  $e_j[l] = 1$  for an even number of the vectors  $\{e_j\}$ , then  $\sum e_j = \vec{0}$ , contradicting the linear independence of the columns of  $A$ . So, assume that there exists an index  $i$  such that the number of the vectors  $\{e_j\}$  that satisfy  $e_j[i] = 1$  is odd. If an odd number of the vectors  $\{o_j\}$  are such that  $o_j[i] = 1$ , then the  $i$ -th row of  $A$  is even, so  $S_{ii} = 0$ ; on the other hand, the  $i$ -th row of  $S$  equals the sum of an odd number of rows  $e_j^*$  plus an odd number of rows  $o_j^*$ , which is odd, by Lemma 4.2. If there are an even number of the vectors  $\{o_j\}$  with  $o_j[i] = 1$ , then the  $i$ -th row of  $A$  is odd, so  $S_{ii} = 1$ ; on the other hand, the  $i$ -th row of  $S$  equals the sum of an odd number of  $e_j^*$  plus an even number of  $o_j^*$ , which is even, by Lemma 4.2. Therefore,  $S$  is not parity indicative.  $\square$

**Lemma 4.12.** *Let  $S$  be a  $d \times d$  symmetric matrix, and suppose  $S = AA^*$  for some  $d \times m$  matrix  $A$ . If  $S$  is parity indicative,  $\text{diag}(S) = \vec{0}$ , and  $c(A) = \text{rank}(A) + 1$ , then either every column of  $A$  is even or every column of  $A$  is odd.*

*Proof.* Denote the odd columns and even columns of  $A$  by  $\{o_j\}$  and  $\{e_j\}$ , respectively, and assume both sets are nonempty. Since each row of  $S$  is even, for every  $i$ , an even number of the vectors  $\{o_j\}$  must have  $o_j[i] = 1$ , by Lemma 4.2. It follows that  $\sum o_j = \vec{0}$ ; that is,  $Ax = \vec{0}$  where  $x[j] = 1$  if  $j$  is the index of an odd column and  $x[j] = 0$  if  $j$  is the index of an even column. But since every row of  $A$  is even,  $A\vec{1} = \vec{0}$ . Hence  $Ay = \vec{0}$  for  $y = \vec{1} + x$ . Since the nonzero linearly independent vectors  $x$  and  $y$  are both contained in the null space of  $A$ , the rank-nullity theorem implies

$$\text{rank}(A) \leq c(A) - 2 = \text{rank}(A) + 1 - 2 = \text{rank}(A) - 1,$$

a contradiction. Therefore, either  $\{o_j\}$  or  $\{e_j\}$  must be empty. □

One additional useful fact is required before we state our main result.

**Lemma 4.13.** (1) *Suppose  $e, o_1, o_2, o_3 \in \mathbb{Z}_2^d$  are four vectors such that  $e$  is even and  $o_1, o_2, o_3$  are odd. Then there exists three even vectors  $f_1, f_2, f_3$  and an odd vector  $p$  such that*

$$ee^* + o_1o_1^* + o_2o_2^* + o_3o_3^* = f_1f_1^* + f_2f_2^* + f_3f_3^* + pp^*,$$

$$\text{and Span}\{e, o_1, o_2, o_3\} = \text{Span}\{f_1, f_2, f_3, p\}.$$

(2) *Suppose  $e_1, e_2, e_3, o \in \mathbb{Z}_2^d$  are four vectors such that  $e_1, e_2, e_3$  are even and  $o$  is odd. Then there exists an even vector  $f$  and odd vectors  $p_1, p_2, p_3$  such that*

$$e_1e_1^* + e_2e_2^* + e_3e_3^* + oo^* = ff^* + p_1p_1^* + p_2p_2^* + p_3p_3^*,$$

$$\text{and Span}\{e_1, e_2, e_3, o\} = \text{Span}\{f, p_1, p_2, p_3\}.$$

*Proof.* For part (1), let  $f_1 = e + o_1 + o_2$ ,  $f_2 = e + o_1 + o_3$ ,  $f_3 = e + o_2 + o_3$ , and  $p = o_1 + o_2 + o_3$ . By Lemma 4.2,  $f_1, f_2$  and  $f_3$  are even and  $p$  is odd. For part (2), let  $f = e_1 + e_2 + e_3$ , and  $p_1 = e_1 + e_2 + o$ ,  $p_2 = e_1 + e_3 + o$ ,  $p_3 = e_2 + e_3 + o$ . Lemma 4.2 implies  $f$  is even and  $p_1, p_2, p_3$  are odd. Easy computations show that the given equalities are satisfied. □

We are now ready for the binary analog of Theorem 1.5: necessary and sufficient conditions on pairs of matrices  $S$  and vectors  $\alpha$  such that  $S$  is the frame operator of a frame with vector “norms” determined by  $\alpha$ . The necessary condition is easy.

**Theorem 4.14.** *Let  $\mathcal{F} = \{f_i\}_{i=1}^K$  be a frame such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$ . Let  $\alpha$  be the vector in  $\mathbb{Z}_2^K$  defined by  $\alpha[i] = (f_i, f_i)$  for each  $i$ . Then  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$ .*

*Proof.*  $\|\alpha\|_0 \equiv \text{Tr}(\Theta_{\mathcal{F}}^* \Theta_{\mathcal{F}}) \equiv \text{Tr}(\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*) \equiv \text{Tr}(S) \pmod{2}$ . □

Sufficiency breaks down into three possible scenarios. If  $S$  is parity indicative, then a minimal frame  $\mathcal{F}$  with frame operator  $S$  must consist of only odd vectors or can attain any nonzero vector  $\alpha$  with  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$  in the sense that  $(f_i, f_i) = \alpha[i]$  for each  $i$ ; if  $S$  is not parity indicative, a minimal frame must contain at least one even vector. This is shown in Theorems 4.15 and 4.17. Nonminimal frames can be constructed to correspond to any nonzero  $\alpha$  with  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$  if  $S$  is parity indicative or to any such  $\alpha$  with at least one zero entry if  $S$  is not parity indicative (Corollary 4.16 and Theorem 4.17).

The frame elements can be permuted in any way without affecting the frame operator. Indeed, if  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^* = S$  and  $\Theta_{\tilde{\mathcal{F}}} = \Theta_{\mathcal{F}} P^*$  for some permutation matrix  $P$ , then

$$\Theta_{\tilde{\mathcal{F}}} \Theta_{\tilde{\mathcal{F}}}^* = \Theta_{\mathcal{F}} P^* P \Theta_{\mathcal{F}}^* = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^* = S.$$

Therefore, in what follows, we need only construct frames with the correct number of odd elements, corresponding to  $\|\alpha\|_0$ , in order to attain the dot products prescribed by  $\alpha$ .

**Theorem 4.15.** *Let  $S$  be a  $d \times d$  parity indicative symmetric matrix. Let  $K = 2d - \text{rank}(S)$ , and let  $\alpha \in \mathbb{Z}_2^K$  be a nonzero vector with  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$ .*

(1) *Suppose  $\text{diag}(S) \neq \vec{0}$ .*

(a) *If  $\text{rank}(S) = d$ , there exists a (minimal)  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$  only if  $\|\alpha\|_0 = K$ .*

(b) *If  $\text{rank}(S) < d$ , there exists a (minimal)  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$ .*

(2) *Suppose  $\text{diag}(S) = \vec{0}$ . Then  $\text{rank}(S) < d$ .*

(a) *If  $\text{rank}(S) = d - 1$ , there exists a (minimal)  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$  only if  $\|\alpha\|_0 = K$ .*

(b) *If  $\text{rank}(S) < d - 1$ , there exists a (minimal)  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$ .*

*Proof.* In case (1a), [Theorem 4.5](#) implies the existence of a  $K$ -element frame  $\mathcal{F}$ , where  $K = d$ , whose frame operator is  $S$ . The  $d$  columns of  $\Theta_{\mathcal{F}}$  must be linearly independent, so every  $f_i$  must be odd, by [Lemma 4.11](#). (As a corollary of [Theorem 4.14](#), we note that  $d \equiv \text{Tr}(S) \pmod{2}$  for any  $d \times d$ , full rank, parity indicative symmetric matrix  $S$  with  $\text{diag}(S) \neq \vec{0}$ .)

For case (1b), instead of using the result of [Theorem 4.9](#), we rely directly on [Theorem 4.3](#) to construct a  $d \times \text{rank}(S)$  matrix  $A$  with  $\text{rank}(A) = \text{rank}(S)$  such that  $AA^* = S$ . Since the columns of  $A$  are linearly independent, [Lemma 4.11](#) implies that they are all odd. As in the proof of [Theorem 4.9](#), we consider an augmented matrix

$$\Theta_{\mathcal{F}} = [A \ B]$$

but with more care taken in the choice of  $B$ . By letting the columns of  $B$  be  $d - \text{rank}(S)$  of the standard basis vectors not in the column space of  $A$ , each repeated twice, we construct a frame  $\mathcal{F}$  for  $\|\alpha\|_0 = 2d - \text{rank}(S)$ . Replacing any identical pair of columns of  $B$ , say  $\{\varepsilon_l, \varepsilon_l\}$ , with  $\{\varepsilon_l + \varepsilon_n, \varepsilon_l + \varepsilon_n\}$  for any other basis vector  $\varepsilon_n \neq \varepsilon_l$ , the columns of  $\Theta_{\mathcal{F}}$  still span,  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  is still equal to  $S$ , but now  $\mathcal{F}$  contains two fewer odd vectors. In this way, we are able to construct a frame  $\mathcal{F}$  with dot products  $(f_i, f_i)$  satisfying any  $\|\alpha\|_0 = \text{rank}(S) + 2m$  for  $0 \leq m \leq d - \text{rank}(S)$ . (Note that, by the proof of [Theorem 4.14](#),  $\text{rank}(S) \equiv \text{Tr}(S) \pmod{2}$ .) By [Lemma 4.13](#), any four vectors consisting of three odds and one even can be substituted by four vectors consisting of three evens and one odd, having the same span and no effect on  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$ . Each substitution allows us to increase the number of even vectors by



two, until only two odd vectors remain in  $\mathcal{F}$  if  $\text{rank}(S)$  is even or one odd vector remains if  $\text{rank}(S)$  is odd. Therefore, we can build a frame with  $2d - \text{rank}(S)$  elements, corresponding to any nonzero  $\alpha$  with  $\|\alpha\|_0 \equiv \text{rank}(S) \equiv \text{Tr}(S) \pmod{2}$ .

Now let  $S$  be parity indicative with  $\text{diag}(S) = \vec{0}$ . Since every row of  $S$  is even,  $\text{rank}(S) < d$ . In case (2a), [Theorem 4.9](#) implies the existence of a  $(d+1)$ -element frame  $\mathcal{F}$  whose frame operator is  $S$ . Since  $\mathcal{F}$  is a spanning set, it must contain an odd vector. By [Lemma 4.12](#), every vector in  $\mathcal{F}$  must be odd. (Note that, by [Corollary 4.8](#), case (2a) can only occur if  $\text{rank}(S)$  is even, and hence  $d$  is odd.)

Lastly, we assume in case (2b) that  $\text{rank}(S) \leq d - 2$ . By [Theorem 4.3](#) and [Proposition 4.4](#), there exists a  $d \times (\text{rank}(S) + 1)$  matrix  $A$  with  $\text{rank}(A) = \text{rank}(S)$  such that  $AA^* = S$ . By [Lemma 4.12](#), either every column of  $A$  is even or every column is odd. Since  $S_{ii} = 0$  for every  $i$ , every row of  $A$  is even, and hence the sum of the columns of  $A$  is  $\vec{0}$ . Moreover, since  $\text{diag}(S) = \vec{0}$ , we know that  $\text{rank}(S)$  must be even, by [Corollary 4.8](#). If every column of  $A$  were odd, then the sum of all  $\text{rank}(S) + 1$  columns would have to be odd, by [Lemma 4.2](#), yielding a contradiction. So every column of  $A$  must be even.

Augment  $A$  with a column of zeros and call the resulting matrix  $B$ . Then each column of  $B$  is even, each row of  $B$  is even, and  $B$  has an even number of columns. Consider a row  $b_n^*$  of  $B$  such that  $b_n^* \in \text{Span}\{b_j^* : b_j^* \text{ is a row of } B \text{ and } j \neq n\}$ . Replace  $b_n^*$  by its complement (that is, add  $\vec{1}^*$  for  $\vec{1} \in \mathbb{Z}_2^{\text{rank}(S)+2}$  to  $b_n^*$ ), and call the resulting matrix  $C$ . Then  $CC^* = S$ ,  $\text{rank}(C) = \text{rank}(S) + 1$ , and  $C$  is composed of  $\text{rank}(S) + 2$  odd columns. As in case (1b), we now augment  $C$  with  $d - (\text{rank}(S) + 1)$  of the standard basis vectors not in the column space of  $C$ , each repeated twice, to construct  $\Theta_{\mathcal{F}}$ . In doing so, we construct a frame  $\mathcal{F}$  consisting of  $\text{rank}(S) + 2 + 2(d - (\text{rank}(S) + 1)) = 2d - \text{rank}(S)$  vectors, with frame operator  $S$ , such that every element of  $\mathcal{F}$  is odd. By [Theorem 4.14](#),  $\|\alpha\|_0$  must be even. As in case (1b), we can replace pairs of odd elements of  $\mathcal{F}$  by even vectors until only two odd vectors remain.  $\square$

**Corollary 4.16.** *Let  $S$  be a  $d \times d$  parity indicative symmetric matrix. Let  $K > 2d - \text{rank}(S)$ . Let  $\alpha \in \mathbb{Z}_2^K$  be a nonzero vector such that  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$ . Then there exists a  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$ .*

*Proof.* Since  $S$  is parity indicative, a  $K$ -element frame with  $K > 2d - \text{rank}(S)$  is necessarily nonminimal and can be constructed by augmenting the minimal frames of the previous theorem. Consider first the minimal frame  $\mathcal{F}$  guaranteed by case (1a) of [Theorem 4.15](#). Adding the zero vector to  $\mathcal{F}$  allows us to apply [Lemma 4.13](#) and create frames satisfying  $(f_i, f_i) = \alpha[i]$  for any  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$  with  $0 < \|\alpha\|_0 < d$ . Similarly, for case (2a), including the zero vector allows the construction of a frame corresponding to any  $\|\alpha\|_0 = 2, 4, 6, \dots, d + 1$ . In either

case, the addition of two identical copies of odd vectors or two identical copies of even vectors provides frames for any  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$  when  $\|\alpha\|_0 \geq d + 2$  or  $\|\alpha\|_0 \geq d + 3$ , corresponding to cases (1a) and (2a), respectively. Similarly in cases (1b) and (2b), the addition of two identical copies of an odd vector or two identical copies of an even vector yield frames for  $2d - \text{rank}(S) < \|\alpha\|_0$ .  $\square$

**Theorem 4.17.** *Let  $S$  be a  $d \times d$  symmetric matrix that is not parity indicative. Let  $K \geq 2d - \text{rank}(S)$  or if  $\text{rank}(S) = d$  and  $\text{diag}(S) = \vec{0}$ , let  $K \geq d + 1$ . Let  $\alpha \in \mathbb{Z}_2^K$  be a nonzero vector such that  $\|\alpha\|_0 \equiv \text{Tr}(S) \pmod{2}$ . Then there exists a  $K$ -element frame  $\mathcal{F}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $(f_i, f_i) = \alpha[i]$  for every  $i$  only if  $\|\alpha\|_0 \neq K$ .*

*Proof.* We use [Theorem 4.5](#) and [Remark 4.6](#) or [Theorem 4.9](#) to construct a  $K$ -element frame  $\mathcal{F}$  such that  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^* = S$ . By [Lemma 4.11](#),  $\mathcal{F}$  must contain an even vector. Of course,  $\mathcal{F}$  must also contain an odd vector, in order to span. Let  $m \equiv \text{Tr}(S) \pmod{2}$  represent the number of odd elements of  $\mathcal{F}$  and  $K - m$  be the number of even elements. By [Lemma 4.13](#),  $\mathcal{F}$  may be replaced by a frame with two more or two fewer odd vectors. Through repeated applications, we can construct a frame  $\mathcal{F}$  corresponding to any  $\|\alpha\|_0 = 1, 3, 5, \dots, K - 1$  if  $m$  is odd and  $K$  is even, any  $\|\alpha\|_0 = 1, 3, 5, \dots, K - 2$  if  $m$  is odd and  $K$  is odd, any  $\|\alpha\|_0 = 2, 4, 6, \dots, K - 1$  if  $m (\geq 2)$  is even and  $K (> m)$  is odd, or any  $\|\alpha\|_0 = 2, 4, 6, \dots, K - 2$  if  $m (\geq 2)$  is even and  $K (> m)$  is even.  $\square$

## 5. Examples and data

**Examples.** In this subsection we consider two symmetric matrices  $S$  and build frames with various  $\alpha$ 's to illustrate the main result of [Section 4](#). The algorithm for factorizing a matrix as  $S = AA^*$  and for reducing  $A$  into a minimal factor can be found in [[Lempel 1975](#)].

**Example 5.1.** Consider the identity matrix

$$S = I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

and note that it is a symmetric, parity indicative, full-rank matrix. Any frame with frame operator  $S$  is a Parseval frame. By [Theorem 4.15](#), a minimal 4-element such Parseval frame must satisfy  $\|\alpha\|_0 = 4$ , where  $\alpha[i] = (f_i, f_i)$  for each  $i$ ; clearly, this follows from the Parseval frame necessarily being an orthonormal basis. [Corollary 4.16](#) guarantees Parseval frames in  $\mathbb{Z}_2^4$  of length  $K = 5$  with either two or four odd vectors corresponding to any  $\alpha \in \mathbb{Z}_2^5$  with  $\|\alpha\|_0 = 2, 4$ . To begin the construction, factor  $S$  as  $I_4 I_4^*$ . Appending the zero-column to the left factor yields the matrix

$$\Theta_{\mathcal{F}} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix},$$

the columns of which constitute a frame with frame operator  $S$  and  $\alpha = (1, 1, 1, 1, 0)$ . To obtain any other  $\alpha \in \mathbb{Z}_2^5$  with  $\|\alpha\|_0 = 4$ , simply permute the columns.

We utilize [Lemma 4.13](#) to reduce the number of odd vectors by two. Let  $e$  be the zero-column and  $o_1, o_2, o_3$  be the first, second, and third columns of  $\Theta_{\mathcal{F}}$ , respectively. Replacing  $e, o_1, o_2, o_3$  with their counterparts constructed in [Lemma 4.13](#) results in

$$\Theta_{\mathcal{F}'} = \begin{bmatrix} 1 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}.$$

Taking the columns of  $\Theta_{\mathcal{F}'}$  as frame vectors builds the frame  $\mathcal{F}'$  satisfying  $\alpha = (0, 0, 0, 1, 1)$ . Again, the columns of  $\mathcal{F}'$  can be permuted to acquire any  $\alpha \in \mathbb{Z}_2^5$  with  $\|\alpha\|_0 = 2$ . Notice that a permutation of  $\mathcal{F}'$  appears in the proof of [Theorem 3.5](#).

**Example 5.2.** Suppose we wish to find a frame for  $\mathbb{Z}_2^3$  of length 7 with frame operator

$$S = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Since this rank-2, symmetric matrix is not parity indicative, we apply [Theorem 4.17](#). In doing so, we follow the proof of [Theorem 4.9](#) and factorize  $S$  as

$$S = P^* T \begin{bmatrix} L & 0 \\ 0 & 0 \end{bmatrix} T^* P,$$

where  $P$  is the  $3 \times 3$  identity matrix,

$$L = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \quad \text{and} \quad T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

Then

$$A = \begin{bmatrix} H & 0 \\ 0 & B \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix},$$

and we append three zero-columns to  $TA$  to build the frame  $\mathcal{F}$ :

$$\Theta_{\mathcal{F}} = [TA \ \vec{0} \ \vec{0} \ \vec{0}] = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 \end{bmatrix}.$$

Letting  $(f_i, f_i) = \alpha[i]$  for each  $i$ , we see that  $\mathcal{F}$  satisfies  $\alpha = (1, 0, 1, 1, 0, 0, 0)$ . We increase or decrease the number of odd vectors as desired, by applying [Lemma 4.13](#) first to  $\{f_1, f_3, f_4, f_5\}$  and then to  $\{f_4, f_5, f_6, f_7\}$ . We obtain frames  $\mathcal{F}_1$  and  $\mathcal{F}_2$  satisfying

$$\Theta_{\mathcal{F}_1} = \begin{bmatrix} 1 & 1 & 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}, \quad \alpha_1 = (0, 0, 0, 0, 1, 0, 0);$$

$$\Theta_{\mathcal{F}_2} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 \end{bmatrix}, \quad \alpha_2 = (1, 0, 1, 0, 1, 1, 1).$$

By [Theorem 4.17](#),  $\|\alpha\|_0 = 7$  is unattainable.

**Data.** An exhaustive search for frame operators  $\Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $\|\alpha\|_0$  associated with  $\mathcal{F} = \{f_j\}_{j=1}^K$  in  $\mathbb{Z}_2^d$  was performed, using Python 3.6, for various dimensions and frame lengths (i.e., various  $d$ 's and  $K$ 's). The tables contained in this subsection hold information about the number of symmetric matrices that are frame operators and the set of  $\|\alpha\|_0$  that accompany them. We include summaries for dimensions  $d = 2, \dots, 5$ . Because every frame in  $\mathbb{Z}_2^d$  must have at least  $d$  vectors, and because  $2d$  is the minimum number of vectors needed to ensure every symmetric matrix is a frame operator ([Theorems 4.5, 4.9](#)), the computations were performed for  $K = d, \dots, 2d$ .

For  $d = 2, \dots, 5$ , in the table containing information about the  $d$ -dimensional binary space, the entry in the row labeled  $\{\alpha_{\min}, \alpha_{\min} + 2, \dots, \alpha_{\min} + 2t\}$  and column labeled  $K = k_0$  shows the number of symmetric matrices  $S$  in  $d$ -dimensional space such that for each  $\alpha \in \{\alpha_{\min}, \alpha_{\min} + 2, \dots, \alpha_{\min} + 2t\}$  there exists a frame  $\mathcal{F} = \{f_j\}_{j=1}^{k_0}$  such that  $S = \Theta_{\mathcal{F}} \Theta_{\mathcal{F}}^*$  and  $\alpha[i] = (f_i, f_i)$  for  $1 \leq i \leq k_0$ .

In each table, the sum of the entries of the last column represents all possible symmetric  $d \times d$  binary matrices. There are  $2^{d(d+1)/2}$  such matrices, which becomes prohibitively large as the dimension  $d$  increases.

$\{\ \alpha\ _0\}$	$K$		
	2	3	4
{1}	2	2	0
{2}	1	3	2
{1, 3}	0	2	4
{2, 4}	0	0	2

**Table 1.** Number of attainable frame operators of frames for  $\mathbb{Z}_2^2$ .

$\{\ \alpha\ _0\}$	$K$			
	3	4	5	6
{1}	12	0	0	0
{2}	12	21	0	0
{3}	4	0	0	0
{4}	0	1	0	0
{1, 3}	0	28	24	0
{2, 4}	0	6	31	24
{1, 3, 5}	0	0	8	32
{2, 4, 6}	0	0	0	8

**Table 2.** Number of attainable frame operators of frames for  $\mathbb{Z}_2^3$ .

$\{\ \alpha\ _0\}$	$K$				
	4	5	6	7	8
{2}	168	0	0	0	0
{4}	28	0	0	0	0
{1, 3}	224	392	0	0	0
{2, 4}	0	420	441	0	0
{1, 3, 5}	0	56	504	448	0
{2, 4, 6}	0	0	63	511	448
{1, 3, 5, 7}	0	0	0	64	512
{2, 4, 6, 8}	0	0	0	0	64

**Table 3.** Number of attainable frame operators of frames for  $\mathbb{Z}_2^4$ .

$\{\ \alpha\ _0\}$	$K$					
	5	6	7	8	9	10
{5}	448	0	0	0	0	0
{6}	0	28	0	0	0	0
{1, 3}	6720	0	0	0	0	0
{2, 4}	6720	13020	0	0	0	0
{1, 3, 5}	0	13888	15120	0	0	0
{2, 4, 6}	0	840	15988	15345	0	0
{1, 3, 5, 7}	0	0	1008	16368	15360	0
{2, 4, 6, 8}	0	0	0	1023	16383	15360
{1, 3, 5, 7, 9}	0	0	0	0	1024	16384
{2, 4, 6, 8, 10}	0	0	0	0	0	1024

**Table 4.** Number of attainable frame operators of frames for  $\mathbb{Z}_2^5$ .

## Acknowledgement

We thank Erich McAlister for his very valuable feedback.

## References

- [Baker et al. 2018] Z. J. Baker, B. G. Bodmann, M. G. Bullock, S. N. Branum, and J. E. McLaney, “What is odd about binary Parseval frames?”, *Involve* **11**:2 (2018), 219–233.
- [Bodmann et al. 2009] B. G. Bodmann, M. Le, L. Reza, M. Tobin, and M. Tomforde, “Frame theory for binary vector spaces”, *Involve* **2**:5 (2009), 589–602. [MR](#) [Zbl](#)
- [Bodmann et al. 2014] B. G. Bodmann, B. Camp, and D. Mahoney, “Binary frames, graphs and erasures”, *Involve* **7**:2 (2014), 151–169. [MR](#) [Zbl](#)
- [Bownik and Jasper 2015] M. Bownik and J. Jasper, “Existence of frames with prescribed norms and frame operator”, pp. 103–117 in *Excursions in harmonic analysis, IV*, edited by R. Balan et al., Birkhäuser, Cham, 2015. [MR](#)
- [Cahill et al. 2013] J. Cahill, M. Fickus, D. G. Mixon, M. J. Poteet, and N. Strawn, “Constructing finite frames of a given spectrum and set of lengths”, *Appl. Comput. Harmon. Anal.* **35**:1 (2013), 52–73. [MR](#) [Zbl](#)
- [Casazza and Kutyniok 2013] P. G. Casazza and G. Kutyniok (editors), *Finite frames: theory and applications*, Birkhäuser, New York, 2013. [MR](#) [Zbl](#)
- [Casazza and Leon 2010] P. G. Casazza and M. T. Leon, “Existence and construction of finite frames with a given frame operator”, *Int. J. Pure Appl. Math.* **63**:2 (2010), 149–157. [MR](#) [Zbl](#)
- [Casazza et al. 2006] P. G. Casazza, M. Fickus, J. Kovačević, M. T. Leon, and J. C. Tremain, “A physical interpretation of tight frames”, pp. 51–76 in *Harmonic analysis and applications*, edited by C. Heil, Birkhäuser, Boston, 2006. [MR](#) [Zbl](#)
- [Christensen 2003] O. Christensen, *An introduction to frames and Riesz bases*, Birkhäuser, Boston, 2003. [MR](#) [Zbl](#)
- [Christensen et al. 2012] O. Christensen, A. M. Powell, and X. C. Xiao, “A note on finite dual frame pairs”, *Proc. Amer. Math. Soc.* **140**:11 (2012), 3921–3930. [MR](#) [Zbl](#)
- [Cohen et al. 1997] G. Cohen, I. Honkala, S. Litsyn, and A. Lobstein, *Covering codes*, North-Holland Mathematical Library **54**, North-Holland, Amsterdam, 1997. [MR](#) [Zbl](#)
- [Fickus et al. 2013] M. Fickus, D. G. Mixon, M. J. Poteet, and N. Strawn, “Constructing all self-adjoint matrices with prescribed spectrum and diagonal”, *Adv. Comput. Math.* **39**:3-4 (2013), 585–609. [MR](#) [Zbl](#)
- [Han et al. 2007] D. Han, K. Kornelson, D. Larson, and E. Weber, *Frames for undergraduates*, Student Mathematical Library **40**, American Mathematical Society, Providence, 2007. [MR](#) [Zbl](#)
- [Horn and Johnson 1985] R. A. Horn and C. R. Johnson, *Matrix analysis*, Cambridge Univ. Press, 1985. [MR](#) [Zbl](#)
- [Hotovy et al. 2015] R. Hotovy, D. R. Larson, and S. Scholze, “Binary frames”, *Houston J. Math.* **41**:3 (2015), 875–899. [MR](#) [Zbl](#)
- [Kovačević and Chebira 2007] J. Kovačević and A. Chebira, “Life beyond bases: the advent of frames, II”, *IEEE Signal Proc. Mag.* **24**:5 (2007), 115–125.
- [Lempel 1975] A. Lempel, “Matrix factorization over  $\text{GF}(2)$  and trace-orthogonal bases of  $\text{GF}(2^n)$ ”, *SIAM J. Comput.* **4**:2 (1975), 175–186. [MR](#) [Zbl](#)

Received: 2017-05-24

Accepted: 2017-07-17

[furst\\_v@fortlewis.edu](mailto:furst_v@fortlewis.edu)*Department of Mathematics, Fort Lewis College,  
Durango, CO, United States*[eric.powell.smith@gmail.com](mailto:eric.powell.smith@gmail.com)*Department of Mathematics, Fort Lewis College,  
Durango, CO, United States*

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2018 is US \$190/year for the electronic version, and \$250/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers



# involve

2018

vol. 11

no. 3

<a href="#">A mathematical model of treatment of cancer stem cells with immunotherapy</a>	361
ZACHARY J. ABERNATHY AND GABRIELLE EPELLE	
<a href="#">RNA, local moves on plane trees, and transpositions on tableaux</a>	383
LAURA DEL DUCA, JENNIFER TRIPP, JULIANNA TYMOCZKO AND JUDY WANG	
<a href="#">Six variations on a theme: almost planar graphs</a>	413
MAX LIPTON, EOIN MACKALL, THOMAS W. MATTMAN, MIKE PIERCE, SAMANTHA ROBINSON, JEREMY THOMAS AND ILAN WEINSCHELBAUM	
<a href="#">Nested Frobenius extensions of graded superrings</a>	449
EDWARD POON AND ALISTAIR SAVAGE	
<a href="#">On <math>G</math>-graphs of certain finite groups</a>	463
MOHAMMAD REZA DARAFSHEH AND SAFOORA MADADY MOGHADAM	
<a href="#">The tropical semiring in higher dimensions</a>	477
JOHN NORTON AND SANDRA SPIROFF	
<a href="#">A tale of two circles: geometry of a class of quartic polynomials</a>	489
CHRISTOPHER FRAYER AND LANDON GAUTHIER	
<a href="#">Zeros of polynomials with four-term recurrence</a>	501
KHANG TRAN AND ANDRES ZUMBA	
<a href="#">Binary frames with prescribed dot products and frame operator</a>	519
VERONIKA FURST AND ERIC P. SMITH	