

*Pacific
Journal of
Mathematics*

**RATIONAL CERTIFICATES OF POSITIVITY ON COMPACT
SEMIALGEBRAIC SETS**

VICTORIA POWERS

RATIONAL CERTIFICATES OF POSITIVITY ON COMPACT SEMIALGEBRAIC SETS

VICTORIA POWERS

Let $\mathbb{R}[X]$ denote the real polynomial ring $\mathbb{R}[X_1, \dots, X_n]$ and write $\sum \mathbb{R}[X]^2$ for the set of sums of squares in $\mathbb{R}[X]$. Given $g_1, \dots, g_s \in \mathbb{R}[X]$ such that the semialgebraic set $K := \{x \in \mathbb{R}^n \mid g_i(x) \geq 0 \text{ for all } i\}$ is compact, Schmüdgen's theorem says that if $f \in \mathbb{R}[X]$ such that $f > 0$ on K , then f is in the preordering in $\mathbb{R}[X]$ generated by the g_i 's, i.e., f can be written as a finite sum of elements $\sigma g_1^{e_1} \dots g_s^{e_s}$, where σ is a sum of squares in $\mathbb{R}[X]$ and each $e_i \in \{0, 1\}$. Putinar's theorem says that under a condition on the set of generators $\{g_1, \dots, g_s\}$ (which is a stronger condition than the compactness of K), any $f > 0$ on K can be written $f = \sigma_0 + \sigma_1 g_1 + \dots + \sigma_s g_s$, where $\sigma_i \in \sum \mathbb{R}[X]^2$. Both of these theorems can be viewed as statements about the existence of certificates of positivity on compact semialgebraic sets. In this note we show that if the defining polynomials g_1, \dots, g_s and polynomial f have coefficients in \mathbb{Q} , then in Schmüdgen's theorem we can find a representation in which the σ 's are sums of squares of polynomials over \mathbb{Q} . We prove a similar result for Putinar's theorem assuming that the set of generators contains $N - \sum X_i^2$ for some $N \in \mathbb{N}$.

1. Introduction

We write \mathbb{N} , \mathbb{R} , and \mathbb{Q} for the set of natural, real, and rational numbers. Let $n \in \mathbb{N}$ be fixed and let $\mathbb{R}[X]$ denote the polynomial ring $\mathbb{R}[X_1, \dots, X_n]$. We denote by $\sum \mathbb{R}[X]^2$ the set of sums of squares in $\mathbb{R}[X]$.

For $S = \{g_1, \dots, g_s\} \subseteq \mathbb{R}[X]$, the *basic closed semialgebraic set* generated by S , denoted K_S , is

$$\{x \in \mathbb{R}^n \mid g_1(x) \geq 0, \dots, g_s(x) \geq 0\}.$$

Associated to S are two algebraic objects: The *quadratic module generated by S* , denoted M_S , is the set of $f \in \mathbb{R}[X]$ which can be written

$$f = \sigma_0 + \sigma_1 g_1 + \dots + \sigma_s g_s,$$

MSC2010: primary 11E25, 12D15, 13J30, 14P10; secondary 14Q20.

Keywords: rational sums of squares, certificates of positivity, Schmüdgen's theorem, Putinar's theorem.

where each σ_i lies in $\sum \mathbb{R}[X]^2$, and the *preordering generated by S* , denoted T_S , is the quadratic module generated by all products of elements in S . In other words, T_S is the set of $f \in \mathbb{R}[X]$ which can be written as a finite sum of elements

$$\sigma g_1^{e_1} \dots g_s^{e_s}, \quad \text{for } \sigma \in \mathbb{R}[X] \text{ and each } e_i \in \{0, 1\}.$$

A polynomial $f \in \sum \mathbb{R}[X]^2$ is obviously globally nonnegative in \mathbb{R}^n and writing f explicitly as a sum of squares gives a “certificate of positivity” for the fact that f takes only nonnegative values in \mathbb{R}^n . (Note: To avoid having to write “nonnegativity or positivity” we use the term “positivity” to mean either.) More generally, for a basic closed semialgebraic set K_S , if $f \in T_S$ or $f \in M_S$, then f is nonnegative on K_S and an explicit representation of f in M_S or T_S gives a certificate of positivity for f on K_S .

Schmüdgen [1991] showed that if the semialgebraic set K_S is compact, then any $f \in \mathbb{R}[X]$ which is strictly positive on K_S is in the preordering T_S . A preordering or quadratic module is *archimedean* if it contains $N - \sum X_i^2$ for some $N \in \mathbb{N}$. We note that if M_S is archimedean, then it follows immediately that K_S is compact, however the converse is not true in general. Putinar [1993] showed that if M_S is archimedean then any $f \in \mathbb{R}[X]$ which is strictly positive on K_S is in M_S . In other words, these results say that under the given conditions a certificate of positivity for f on K_S exists.

Recently, techniques from semidefinite programming combined with Schmüdgen’s and Putinar’s theorems have been used to give numerical algorithms for applications such as optimization of polynomials on semialgebraic sets. However since these algorithms are numerical they might not produce exact certificates of positivity. With this in mind, Sturmfels asked whether any $f \in \mathbb{Q}[X]$ which is a sum of squares in $\mathbb{R}[X]$ is a sum of squares in $\mathbb{Q}[X]$. Hillar [2009] showed that the answer is yes in the case where f is known to be a sum of squares over a totally real field K . The general question remains unsolved.

It is natural to ask a similar question for Schmüdgen’s and Putinar’s theorems: If the polynomials defining the semialgebraic set and the positive polynomial f have rational coefficients, is there a certificate of positivity for f in which the sums of squares have rational coefficients? In this note, we show that in the case of Schmüdgen’s theorem the answer is yes. This follows from an algebraic proof of the theorem, originally due to T. Wörmann [1998]. In the case of Putinar’s theorem, we show that the answer is also yes as long as the generating set contains $N - \sum X_i^2$ for some $N \in \mathbb{N}$. This follows easily from an algorithmic proof of the theorem, due to Schweighofer [2005]. For Lasserre’s method [2001] for optimization of polynomials on compact semialgebraic sets, it is usual in concrete cases to add a polynomial of the type $N - \sum X_i^2$ to the generators in order to insure that Putinar’s theorem holds. Thus our assumption in this case is reasonable.

2. Rational certificates of for Schmüdgen’s theorem

Fix $S = \{g_1, \dots, g_s\} \subseteq \mathbb{R}[X]$ and define K_S and T_S as above.

Theorem 1 (Schmüdgen). *Suppose that K_S is compact. If $f \in \mathbb{R}[X]$ and $f > 0$ on K_S , then $f \in T_S$.*

In this section we show that if f and the generating polynomials g_1, \dots, g_s are in $\mathbb{Q}[X]$, then f has a representation in T_S in which all sums of squares σ_ϵ are in $\sum \mathbb{Q}[X]^2$. This follows from T. Wörmann’s algebraic proof of the theorem using the classical Abstract Positivstellensatz, and a generalization of Wörmann’s crucial lemma due to M. Schweighofer.

The abstract Positivstellensatz. We will need a version of the abstract Positivstellensatz, a result traditionally attributed to Kadison and Dubois, but now thought to have been proved earlier by Krivine or Stone. For details on its history, see [Prestel and Delzell 2001, Section 5.6]. The setting is preordered commutative rings, and we state the version we need as Theorem 2 below.

Let A be a commutative ring with $\mathbb{Q} \subseteq A$. A subset $T \subseteq A$ is a *preordering* if $T + T \subseteq T$, $T \cdot T \subseteq T$, and $-1 \notin T$. For $S = \{a_1, \dots, a_k\} \subseteq A$, we define the *preordering generated by S* , T_S , exactly as for $A = \mathbb{R}[X]$.

An *ordering* in A is a preordering P such that $P \cup -P = A$ and $P \cap -P$ is a prime ideal. Any $a \in A$ has a unique sign in $\{-1, 0, 1\}$ with respect to a fixed ordering P and we use the notation $a \geq_P 0$ if $a \in P$, $a >_P 0$ if $a \in P \setminus (P \cap -P)$, etc.

Fix a preordered ring (A, T) and denote by $\text{Sper } A$ the real spectrum of (A, T) , i.e., the set of orderings of A which contain T . Then define

$$H(A) = \{a \in A \mid \text{there exists } n \in \mathbb{N} \text{ with } n \pm a \geq_P 0 \text{ for all } P \in \text{Sper } A\},$$

the *ring of geometrically bounded elements in (A, T)* , and

$$H'(A) = \{a \in A \mid \text{there exists } n \in \mathbb{N} \text{ with } n \pm a \in T\},$$

the *ring of arithmetically bounded elements in (A, T)* . Clearly, $H'(A) \subseteq H(A)$. The preordering T is *archimedean* if $H'(A) = A$.

Theorem 2 [Schweighofer 2002, Theorem 1]. *Given the preordered ring (A, T) as above and suppose $A = H'(A)$. For any $a \in A$, if $a >_P 0$ for all $P \in \text{Sper } A$, then $a \in T$.*

Consider the case where $A = \mathbb{R}[X]$ and $T = T_S$ for $S = \{g_1, \dots, g_s\} \subseteq \mathbb{R}[X]$. Let $K = K_S$, then K embeds densely in $\text{Sper } A$ and hence $H(A) = \{f \in \mathbb{R}[X] \mid f \text{ is bounded on } S\}$. If S is compact, this implies $H(A) = A$ and Schmüdgen’s theorem follows from the following result:

Lemma 3 [Berr and Wörmann 2001, Lemma 1]. *With A , T , and S as above, if $H(A) = A$ then $H'(A) = A$.*

Our result follows from a generalization of this lemma:

Theorem 4 [Schweighofer 2002, Theorem 4.13]. *Let F be a subfield of \mathbb{R} and (A, T) a preordered F -algebra such that $F \subseteq H'(A)$ and A has finite transcendence degree over F . Then*

$$A = H(A) \implies A = H'(A).$$

We can now prove the existence of rational certificates of positivity in Schmüdgen's theorem. The argument is exactly that of the proof of the general theorem above.

Theorem 5. *Given $S = \{g_1, \dots, g_s\} \subseteq \mathbb{Q}[X]$ and suppose $K_S \subseteq \mathbb{R}^n$ is compact. Then for any and $f \in \mathbb{Q}[X]$ such that $f > 0$ on K_S , there is a representation of f in the preordering T_S ,*

$$f = \sum_{e \in \{0,1\}^s} \sigma_e g_1^{e_1} \cdots g_s^{e_s},$$

with all $\sigma_e \in \sum \mathbb{Q}[X]^2$.

Proof. Let T be the preordering in $\mathbb{Q}[X]$ generated by S . Since K_S is compact, every element of $\mathbb{Q}[X]$ is bounded on K_S . Then K_S dense in $\text{Sper } A$ implies that $H(\mathbb{Q}[X]) = \mathbb{Q}[X]$, hence by Theorem 4 we have $\mathbb{Q}[X] = H'(A)$. Note that the condition $F \subseteq H'(A)$ holds in this case since $\mathbb{Q}^+ = \sum \mathbb{Q}^2$. The result follows from Theorem 2. \square

3. Rational certificates for Putinar's theorem

Given $S = \{g_1, \dots, g_s\}$, recall that the quadratic module generated by S , M_S , is the set of elements in the preordering K_S with a "linear" representation, i.e.,

$$M_S = \left\{ \sigma_0 + \sigma_1 g_1 + \cdots + \sigma_s g_s \mid \sigma_i \in \sum \mathbb{R}[X]^2 \right\}.$$

In order to guarantee representations of positive polynomials in the quadratic module, we need a condition stronger than compactness of K_S , namely, we need M_S to be archimedean.

The quadratic module M_S is archimedean if all elements of $\mathbb{R}[X]$ are bounded by a positive integer with respect to M_S , i.e., if for every $f \in \mathbb{R}[X]$ there is some $N \in \mathbb{N}$ such that $N - f \in M_S$. It is not too hard to show that M_S is archimedean if there is some $N \in \mathbb{N}$ such that $N - \sum X_i^2 \in M_S$. Clearly, if M_S is archimedean, then K_S is compact; the polynomial $N - \sum X_i^2$ can be thought of as a "certificate of compactness". However, the converse is not true; see [Prestel and Delzell 2001, Example 6.3.1]. The key to the algebraic proof of Schmüdgen's theorem from the

previous section is showing that in the case of the preordering generated by a finite set of elements from $\mathbb{R}[X]$, the compactness of the semialgebraic set implies that the corresponding preordering is archimedean.

Putinar [1993] showed that if the quadratic module M_S is archimedean, we can replace the preordering T_S by the quadratic module M_S .

Theorem 6 (Putinar). *Suppose that the quadratic module M_S is archimedean. Then for every $f \in \mathbb{R}[X]$ with $f > 0$ on K_S , $f \in M_S$.*

Lasserre’s method [2001] for minimizing a polynomial on a compact semialgebraic set involves defining a sequence of semidefinite programs corresponding to representations of bounded degree in M_S whose solutions converge to the minimum. In this context, if M_S is archimedean then Putinar’s theorem implies the convergence of the semidefinite programs. In practice, it is not clear how to decide if M_S is archimedean for a given set of generators S , however in concrete cases a polynomial $N - \sum X_i^2$ can be added to the generators if an appropriate N is known or can be computed.

Using an algorithmic proof of Putinar’s theorem due to M. Schweighofer [2005] we can show that rational certificates exist for the theorem as long as we have a polynomial $N - \sum X_i^2$ as one of our generators:

Theorem 7. *Suppose $S = \{g_1, \dots, g_s\} \subseteq \mathbb{Q}[X]$ and $N - \sum X_i^2 \in M_S$ for some $N \in \mathbb{N}$. Then given any $f \in \mathbb{Q}[X]$ such that $f > 0$ on K_S , there exist $\sigma_0 \dots \sigma_s, \sigma \in \sum \mathbb{Q}[X]^2$ so that*

$$f = \sigma_0 + \sigma_1 g_1 + \dots + \sigma_s g_s + \sigma (N - \sum X_i^2).$$

Proof. The idea of Schweighofer’s proof is to reduce to Pólya’s theorem. We follow the proof, making sure that each step preserves rationality.

Let

$$\Delta = \left\{ y \in [0, \infty)^{2n} \mid y_1 + \dots + y_{2n} = 2n(N + \frac{1}{4}) \right\} \subseteq \mathbb{R}^{2n}$$

and let C be the compact subset of \mathbb{R}^n defined by $C = l(\Delta)$, where $l : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ is defined by

$$y \mapsto \left(\frac{y_1 - y_{n+1}}{2}, \dots, \frac{y_n - y_{2n}}{2} \right).$$

Scaling the g_i ’s by positive elements in \mathbb{Q} , we can assume that $g_i \leq 1$ on C for all i . The key to the proof is the observation that *there exists $\lambda \in \mathbb{R}^+$ such that $q := f - \lambda \sum (g_i - 1)^{2k} g_i > 0$ on C* [Schweighofer 2005, Lemma 2.3]. Since we can always replace λ by a smaller value, we can assume $\lambda \in \mathbb{Q}$, whence $q \in \mathbb{Q}[X]$.

Let $d = \deg q$ and let Q_i be the homogeneous part of q of degree i , so $q = \sum_{i=1}^d Q_i$. Let $Y = (Y_1, \dots, Y_{2n})$ and define in $\mathbb{Q}[Y]$

$$F(Y_1, \dots, Y_{2n}) := \sum_{i=1}^d Q_i \left(\frac{Y_1 - Y_{n+1}}{2}, \dots, \frac{Y_n - Y_{2n}}{2} \right) \left(\frac{Y_1 + \dots + Y_{2n}}{2n(N + \frac{1}{4})} \right)^{d-i}.$$

Then F is homogenous and $F > 0$ on $[0, \infty)^{2n} \setminus \{0\}$. By Pólya’s theorem, there is some $k \in \mathbb{N}$ so that

$$G := \left(\frac{Y_1 + \dots + Y_{2n}}{2n(N + \frac{1}{4})} \right)^k F$$

has nonnegative coefficients as a polynomial in $\mathbb{R}[Y]$. Furthermore, since $F \in \mathbb{Q}[Y_1, \dots, Y_{2n}]$, it is easy to see that $G \in \mathbb{Q}[Y]$.

Define $\phi : \mathbb{Q}[Y_1, \dots, Y_{2n}] \rightarrow \mathbb{Q}[X]$ by

$$\phi(Y_i) = N + \frac{1}{4} + X_i, \quad \phi(Y_{n+i}) = (N + \frac{1}{4}) - X_i \quad \text{for } i = 1, \dots, n$$

and note that $\phi(G) = q$ and

$$\begin{aligned} \phi(Y_i) &= (N + \frac{1}{4}) \pm X_i \\ &= \sum_{j \neq i} (X_j^2 + (X_i \pm \frac{1}{2})^2) + (N - \sum X_j^2) \in \sum \mathbb{Q}[X]^2 + (N - \sum X_j^2). \end{aligned}$$

Thus $\phi(G) = q$ implies there is a representation of q of the required type and then, since $f = q + \lambda \sum (g_i - 1)^{2k} g_i$ with $\lambda \in \mathbb{Q}$, we are done. □

Remark 8. In the preordering case (Schmüdgen’s theorem), as noted above if the semialgebraic set K_S is compact, then it follows that the preordering T_S in $\mathbb{Q}[X]$ is archimedean. However it is more subtle in the quadratic module case since it is not always clear how to decide if M_S is archimedean for a given set of generators S . Thus an open question is the following: Suppose $S \subseteq \mathbb{Q}[X]$ is a finite set of polynomials and M_S is archimedean as a quadratic module in $\mathbb{R}[X]$. Is it true that M_S is archimedean as a quadratic module in $\mathbb{Q}[X]$? To put it more concretely, suppose $S = \{g_1, \dots, g_s\} \subseteq \mathbb{Q}[X]$ and we know that there is some $N \in \mathbb{N}$ such that

$$N - \sum X_i^2 = \sigma_0 + \sigma_1 g_1 + \dots + \sigma_s g_s,$$

with $\sigma_i \in \sum \mathbb{R}[X]^2$. Does there exist a representation with $\sigma_i \in \sum \mathbb{Q}[X]^2$? Equivalently, does there exist $N \in \mathbb{N}$ such that for each $i = 1, \dots, n$ we can write

$$N \pm X_i = \sigma_0 + \sigma_1 g_1 + \dots + \sigma_s g_s,$$

with $\sigma_i \in \sum \mathbb{Q}[X]^2$?

References

- [Berr and Wörmann 2001] R. Berr and T. Wörmann, “Positive polynomials on compact sets”, *Manuscripta Math.* **104**:2 (2001), 135–143. MR 2002a:14061 Zbl 0992.14021
- [Hillar 2009] C. J. Hillar, “Sums of squares over totally real fields are rational sums of squares”, *Proc. Amer. Math. Soc.* **137**:3 (2009), 921–930. MR 2009h:12009 Zbl 1163.12005
- [Lasserre 2001] J. B. Lasserre, “Global optimization with polynomials and the problem of moments”, *SIAM J. Optim.* **11**:3 (2001), 796–817. MR 2002b:90054 Zbl 1010.90061
- [Prestel and Delzell 2001] A. Prestel and C. N. Delzell, *Positive polynomials: from Hilbert’s 17th problem to real algebra*, Springer, Berlin, 2001. MR 2002k:13044 Zbl 0987.13016
- [Putinar 1993] M. Putinar, “Positive polynomials on compact semi-algebraic sets”, *Indiana Univ. Math. J.* **42**:3 (1993), 969–984. MR 95h:47014 Zbl 0796.12002
- [Schmüdgen 1991] K. Schmüdgen, “The K -moment problem for compact semi-algebraic sets”, *Math. Ann.* **289**:2 (1991), 203–206. MR 92b:44011
- [Schweighofer 2002] M. Schweighofer, *Iterated rings of bounded elements and generalizations of Schmüdgen’s theorem*, Ph.D. thesis, Universität Konstanz, 2002, available at <http://tinyurl.com/SchweighoferPhD>.
- [Schweighofer 2005] M. Schweighofer, “Optimization of polynomials on compact semialgebraic sets”, *SIAM J. Optim.* **15**:3 (2005), 805–825. MR 2006d:90136 Zbl 1114.90098
- [Wörmann 1998] T. Wörmann, *Strikt positive polynome in der semialgebraischen geometrie*, Ph.D. thesis, Universität Dortmund, 1998, available at http://www.lulu.com/items/volume_63/2227000/2227267/2/print/diss.pdf.

Received January 2, 2011. Revised February 3, 2011.

VICTORIA POWERS
 DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
 EMORY UNIVERSITY
 ATLANTA, GA 30322
 UNITED STATES
vicki@mathcs.emory.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://www.pjmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or www.pjmath.org for submission instructions.

The subscription price for 2011 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2011 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 251 No. 2 June 2011

Two Kazdan–Warner-type identities for the renormalized volume coefficients and the Gauss–Bonnet curvatures of a Riemannian metric	257
BIN GUO, ZHENG-CHAO HAN and HAIZHONG LI	
Gonality of a general ACM curve in \mathbb{P}^3	269
ROBIN HARTSHORNE and ENRICO SCHLESINGER	
Universal inequalities for the eigenvalues of the biharmonic operator on submanifolds	315
SAÏD ILIAS and OLA MAKHOUL	
Multigraded Fujita approximation	331
SHIN-YAO JOW	
Some Dirichlet problems arising from conformal geometry	337
QI-RUI LI and WEIMIN SHENG	
Polycyclic quasiconformal mapping class subgroups	361
KATSUHIKO MATSUZAKI	
On zero-divisor graphs of Boolean rings	375
ALI MOHAMMADIAN	
Rational certificates of positivity on compact semialgebraic sets	385
VICTORIA POWERS	
Quiver grassmannians, quiver varieties and the preprojective algebra	393
ALISTAIR SAVAGE and PETER TINGLEY	
Nonautonomous second order Hamiltonian systems	431
MARTIN SCHECHTER	
Generic fundamental polygons for Fuchsian groups	453
AKIRA USHIJIMA	
Stability of the Kähler–Ricci flow in the space of Kähler metrics	469
KAI ZHENG	
The second variation of the Ricci expander entropy	499
MENG ZHU	