

*Pacific
Journal of
Mathematics*

**ULTRA-DISCRETIZATION OF THE $D_4^{(3)}$ -GEOMETRIC
CRYSTAL TO THE $G_2^{(1)}$ -PERFECT CRYSTALS**

MANA IGARASHI, KAILASH C. MISRA AND TOSHIKI NAKASHIMA

ULTRA-DISCRETIZATION OF THE $D_4^{(3)}$ -GEOMETRIC CRYSTAL TO THE $G_2^{(1)}$ -PERFECT CRYSTALS

MANA IGARASHI, KAILASH C. MISRA AND TOSHIKI NAKASHIMA

Let \mathfrak{g} be an affine Lie algebra and \mathfrak{g}^L its Langlands dual. It was conjectured by Kashiwara, Nakashima, and Okado that \mathfrak{g} has a positive geometric crystal whose ultra-discretization is isomorphic to the limit of certain coherent family of perfect crystals for \mathfrak{g}^L . We prove that the ultra-discretization of the positive geometric crystal for $\mathfrak{g} = D_4^{(3)}$ given by Igarashi and Nakashima is isomorphic to the limit of the coherent family of perfect crystals for $\mathfrak{g}^L = G_2^{(1)}$ constructed by Misra, Mohamad, and Okado.

1. Introduction

Let $A = (a_{ij})_{i,j \in I}$, where $I = \{0, 1, \dots, n\}$, be an affine Cartan matrix and let $(A, \{\alpha_i\}_{i \in I}, \{\alpha_i^\vee\}_{i \in I})$ be a given Cartan datum. Let $\mathfrak{g} = \mathfrak{g}(A)$ denote the associated affine Lie algebra [Kac 1990] and $U_q(\mathfrak{g})$ denote the corresponding quantum affine algebra. Let $P = \mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\Lambda_1 \oplus \dots \oplus \mathbb{Z}\Lambda_n \oplus \mathbb{Z}\delta$ denote the affine weight lattice and $P^\vee = \mathbb{Z}\alpha_0^\vee \oplus \mathbb{Z}\alpha_1^\vee \oplus \dots \oplus \mathbb{Z}\alpha_n^\vee \oplus \mathbb{Z}d$ the dual affine weight lattice. For a dominant weight $\lambda \in P^+ = \{\mu \in P \mid \mu(h_i) \geq 0 \text{ for all } i \in I\}$ of level $l = \lambda(\mathbf{c})$ (where \mathbf{c} is the canonical central element), Kashiwara [1990] defined the crystal base $(L(\lambda), B(\lambda))$ for the integrable highest weight $U_q(\mathfrak{g})$ -module $V(\lambda)$. The crystal $B(\lambda)$ is the $q = 0$ limit of the canonical basis [Lusztig 1990] or the global crystal basis [Kashiwara 1991]. It has many interesting combinatorial properties. To give an explicit realization of $B(\lambda)$, the notions of affine crystal and perfect crystal were introduced in [Kang et al. 1992a]. It is shown there that the affine crystal $B(\lambda)$ for the level $l \in \mathbb{Z}_{>0}$ integrable highest weight $U_q(\mathfrak{g})$ -module $V(\lambda)$ can be realized as the semi-infinite tensor product $\dots \otimes B_l \otimes B_l \otimes B_l$, where B_l is a perfect crystal of level l . This is known as the path realization.

Kang et al. [1994] remarked that one needs a coherent family of perfect crystals $\{B_l\}_{l \geq 1}$ in order to give a path realization of the Verma module $M(\lambda)$ (or $U_q^-(\mathfrak{g})$). In particular, the crystal $B(\infty)$ of $U_q^-(\mathfrak{g})$ can be realized as the semi-infinite tensor

Misra was supported in part by NSA Grant H98230-08-1-0080. Nakashima was supported in part by JSPS Grants in Aid for Scientific Research #22540031.

MSC2010: primary 17B37, 17B67; secondary 22E65, 14M15.

Keywords: geometric crystals, perfect crystals, ultra-discretization.

product $\cdots \otimes B_\infty \otimes B_\infty \otimes B_\infty$ where B_∞ , is the limit of the coherent family of perfect crystals $\{B_l\}_{l \geq 1}$.

At least one coherent family $\{B_l\}_{l \geq 1}$ of perfect crystals and its limit is known for $\mathfrak{g} = A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, A_{2n}^{(2)}, D_{n+1}^{(2)}, D_4^{(3)}, G_2^{(1)}$. (See [Kang et al. 1992b; 1994; Yamane 1998; Kashiwara et al. 2007; Misra et al. 2010].)

A perfect crystal is indeed a crystal for certain finite-dimensional modules of the quantum affine algebra $U_q(\mathfrak{g})$ named after Kirillov and Reshetikhin [1987], and known as KR-modules for short. KR-modules are parametrized by two integers, $i \in I \setminus \{0\}$ and $l > 0$. Let $\{\varpi_i\}_{i \in I \setminus \{0\}}$ be the set of level 0 fundamental weights [Kashiwara 2002]. Hatayama et al. [1999; 2002] conjectured that any KR-module $W(l\varpi_i)$ admits a crystal base $B^{i,l}$ in the sense of Kashiwara and that $B^{i,l}$ is perfect if l is a multiple of $c_i^\vee := \max(1, 2/(\alpha_i, \alpha_i))$. This conjecture has been proved for quantum affine algebras $U_q(\mathfrak{g})$ of classical types [Okado and Schilling 2008; Fourier et al. 2009; 2010]. When $\{B^{i,l}\}_{l \geq 1}$ is a coherent family of perfect crystals we denote its limit by $B_\infty(\varpi_i)$, or just B_∞ if there is no confusion.

The notion of geometric crystals is a geometric analog to Kashiwara's crystal [Kashiwara 1990]. It was defined in [Berenstein and Kazhdan 2000] for reductive algebraic groups and extended to general Kac–Moody groups in [Nakashima 2005a]. For a given Cartan datum $(A, \{\alpha_i\}_{i \in I}, \{\alpha_i^\vee\}_{i \in I})$, a geometric crystal is defined as a quadruple $\mathcal{V}(\mathfrak{g}) = (X, \{e_i\}_{i \in I}, \{\gamma_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$, where X is an algebraic variety, $e_i : \mathbb{C}^\times \times X \rightarrow X$ are rational \mathbb{C}^\times -actions and $\gamma_i, \varepsilon_i : X \rightarrow \mathbb{C}$ ($i \in I$) are rational functions satisfying certain conditions (see Definition 2.1). Geometric crystals have many properties similar to algebraic crystals. For instance, the product of two geometric crystals admits the structure of a geometric crystal if they are induced from unipotent crystals [Berenstein and Kazhdan 2000]. A geometric crystal is said to be a positive geometric crystal if it admits a positive structure (see Definition 2.5). A remarkable relation between positive geometric crystals and algebraic crystals is the ultra-discretization functor $\mathcal{U}\mathcal{D}$ between them (page 123). Applying this functor, positive rational functions are transferred to piecewise linear functions by the simple correspondence:

$$x \times y \mapsto x + y, \quad \frac{x}{y} \mapsto x - y, \quad x + y \mapsto \max(x, y).$$

Let G denote the affine Kac–Moody group associated with the affine Lie algebra \mathfrak{g} . Let B^\pm be fixed Borel subgroups and T the maximal torus of G such that $B^+ \cap B^- = T$. Set $y_i(c) := \exp(cf_i)$, and let $\alpha_i^\vee(c) \in T$ be the image of $c \in \mathbb{C}^\times$ under the group morphism $\mathbb{C}^\times \rightarrow T$ induced by the simple coroot α_i^\vee . We set $Y_i(c) := y_i(c^{-1}) \alpha_i^\vee(c) = \alpha_i^\vee(c) y_i(c)$. Let W and \bar{W} be the Weyl group and extended Weyl group associated with \mathfrak{g} . The Schubert cell

$$X_w := BwB/B,$$

where $w = s_{i_1} \cdots s_{i_k} \in W$, is birationally isomorphic to the variety

$$B_l^- := \{Y_{i_1}(x_1) \cdots Y_{i_k}(x_k) \mid x_1, \dots, x_k \in \mathbb{C}^\times\} \subset B^-,$$

and X_w has a natural geometric crystal structure, where $\iota = i_1, \dots, i_k$ is a reduced word for w . [Berenstein and Kazhdan 2000; Nakashima 2005a].

Let $W(\varpi_i)$ be the KR-module (also called the fundamental representation) of $U_q(\mathfrak{g})$ with ϖ_i as an extremal weight (see [Kashiwara 2002]). Denote its specialization at $q = 1$ by the same symbol, $W(\varpi_i)$. It is a finite-dimensional \mathfrak{g} -module (not necessarily irreducible). Let $\mathbb{P}(\varpi_i)$ be the projective space $(W(\varpi_i) \setminus \{0\})/\mathbb{C}^\times$. For any $i \in I$ the translation $t(c_i^\vee \varpi_i)$ belongs to \tilde{W} (see [Kashiwara et al. 2008]). For a subset J of I , let us denote by \mathfrak{g}_J the subalgebra of \mathfrak{g} generated by $\{e_i, f_i\}_{i \in J}$. For an integral weight μ , define $I(\mu) := \{j \in I \mid \langle \alpha_j^\vee, \mu \rangle \geq 0\}$.

Conjecture 1.1 [Kashiwara et al. 2008]. *For any $i \in I \setminus \{0\}$ there exist a unique variety X endowed with a positive \mathfrak{g} -geometric crystal structure and a rational mapping $\pi : X \rightarrow \mathbb{P}(\varpi_i)$ satisfying the following properties:*

- (i) *For an arbitrary extremal vector $u \in W(\varpi_i)_\mu$, writing the translation $t(c_i^\vee \mu)$ as $\tau w \in \tilde{W}$ with a Dynkin diagram automorphism τ and $w = s_{i_1} \cdots s_{i_k}$, there exists a birational mapping $\xi : B_{i_1, \dots, i_k}^- \rightarrow X$ such that ξ is a morphism of $\mathfrak{g}_{I(\mu)}$ -geometric crystals and that the composition $\pi \circ \xi : B_{i_1, \dots, i_k}^- \rightarrow \mathbb{P}(\varpi_i)$ coincides with $Y_{i_1}(x_1) \cdots Y_{i_k}(x_k) \mapsto Y_{i_1}(x_1) \cdots Y_{i_k}(x_k) \bar{u}$, where \bar{u} is the line including u .*
- (ii) *The ultra-discretization (Section 2) of X is isomorphic to the crystal $B_\infty = B_\infty(\varpi_i)$ of the Langlands dual \mathfrak{g}^L .*

In [Kashiwara et al. 2008], it was shown that this conjecture is true for $i = 1$ and $\mathfrak{g} = A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, A_{2n}^{(2)}, D_{n+1}^{(2)}$. In [Nakashima 2007], a positive geometric crystal for $\mathfrak{g} = G_2^{(1)}$ and $i = 1$ was constructed and it was shown in [Nakashima 2010] that the ultra-discretization of this positive geometric crystal is isomorphic to the limit of the coherent family of perfect crystals for $\mathfrak{g}^L = D_4^{(3)}$ given in [Kashiwara et al. 2007].

More recently, two of the authors have constructed a positive geometric crystal for $\mathfrak{g} = D_4^{(3)}$, $i = 1$ in [Igarashi and Nakashima 2010]. In this paper we describe the structure of the crystal obtained by the ultra-discretization of the geometric crystal $\mathcal{V}(\mathfrak{g})$ constructed in [Igarashi and Nakashima 2010] and then prove that it is isomorphic to the limit B_∞ of the coherent family of perfect crystals for its Langlands dual $\mathfrak{g}^L = G_2^{(1)}$ constructed in [Misra et al. 2010]. This proves Conjecture 4.5 in [Igarashi and Nakashima 2010].

This paper is organized as follows. In Section 2, we recall necessary definitions and facts about geometric crystals. In Section 3, we review needed facts about

affine crystals and perfect crystals. We recall from [Misra et al. 2010] the coherent family of perfect crystals for $\mathfrak{g} = G_2^{(1)}$ and its limit in Section 4. In Section 5, we review the positive geometric crystal $\mathcal{V}(\mathfrak{g})$ for $\mathfrak{g} = D_4^{(3)}$ constructed in [Igarashi and Nakashima 2010]. In Section 6, we state and prove our main result, Theorem 6.1.

2. Geometric crystals

In this section, we review Kac–Moody groups and geometric crystals following [Peterson and Kac 1983; Kumar 2002; Berenstein and Kazhdan 2000].

Kac–Moody algebras and Kac–Moody groups. Fix a symmetrizable generalized Cartan matrix $A = (a_{ij})_{i,j \in I}$ with a finite index set I . Let $(\mathfrak{t}, \{\alpha_i\}_{i \in I}, \{\alpha_i^\vee\}_{i \in I})$ be the associated root data, where \mathfrak{t} is a vector space over \mathbb{C} and $\{\alpha_i\}_{i \in I} \subset \mathfrak{t}^*$ and $\{\alpha_i^\vee\}_{i \in I} \subset \mathfrak{t}$ are linearly independent satisfying $\alpha_j(\alpha_i^\vee) = a_{ij}$.

The Kac–Moody Lie algebra $\mathfrak{g} = \mathfrak{g}(A)$ associated with A is the Lie algebra over \mathbb{C} generated by \mathfrak{t} , the Chevalley generators e_i and f_i ($i \in I$) with the usual defining relations [Kac and Peterson 1983; Peterson and Kac 1983]. There is the root space decomposition $\mathfrak{g} = \bigoplus_{\alpha \in \mathfrak{t}^*} \mathfrak{g}_\alpha$. Denote the set of roots by

$$\Delta := \{\alpha \in \mathfrak{t}^* \mid \alpha \neq 0, \mathfrak{g}_\alpha \neq (0)\}.$$

Set $Q = \sum_i \mathbb{Z}\alpha_i$, $Q_+ = \sum_i \mathbb{Z}_{\geq 0}\alpha_i$, $Q^\vee := \sum_i \mathbb{Z}\alpha_i^\vee$ and $\Delta_+ := \Delta \cap Q_+$. An element of Δ_+ is called a *positive root*. Let $P \subset \mathfrak{t}^*$ be a weight lattice such that $\mathbb{C} \otimes P = \mathfrak{t}^*$, whose element is called a weight.

Define simple reflections $s_i \in \text{Aut}(\mathfrak{t})$ ($i \in I$) by $s_i(h) := h - \alpha_i(h)\alpha_i^\vee$; they generate the Weyl group W , which acts on \mathfrak{t}^* by

$$s_i(\lambda) := \lambda - \lambda(\alpha_i^\vee)\alpha_i.$$

Set $\Delta^{\text{re}} := \{w(\alpha_i) \mid w \in W, i \in I\}$, whose elements are called real roots.

Let \mathfrak{g}' be the derived Lie algebra of \mathfrak{g} and G the Kac–Moody group associated with \mathfrak{g}' [Peterson and Kac 1983]. Let $U_\alpha := \exp \mathfrak{g}_\alpha$ ($\alpha \in \Delta^{\text{re}}$) be a one-parameter subgroup of G . The group G is generated by U_α ($\alpha \in \Delta^{\text{re}}$). Let U^\pm be the subgroup generated by $U_{\pm\alpha}$ ($\alpha \in \Delta_+^{\text{re}} = \Delta^{\text{re}} \cap Q_+$), i.e., $U^\pm := \langle U_{\pm\alpha} \mid \alpha \in \Delta_+^{\text{re}} \rangle$.

For any $i \in I$, there exists a unique homomorphism; $\phi_i : SL_2(\mathbb{C}) \rightarrow G$ such that

$$\phi_i\left(\begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix}\right) = c^{\alpha_i^\vee}, \quad \phi_i\left(\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}\right) = \exp(te_i), \quad \phi_i\left(\begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}\right) = \exp(tf_i),$$

where $c \in \mathbb{C}^\times$ and $t \in \mathbb{C}$. Set $\alpha_i^\vee(c) := c^{\alpha_i^\vee}$, $x_i(t) := \exp(te_i)$, $y_i(t) := \exp(tf_i)$, $G_i := \phi_i(SL_2(\mathbb{C}))$, $T_i := \phi_i(\{\text{diag}(c, c^{-1}) \mid c \in \mathbb{C}^\times\})$ and $N_i := N_{G_i}(T_i)$. Let T (resp. N) be the subgroup of G with the Lie algebra \mathfrak{t} (resp. generated by the N_i 's), which is called a *maximal torus* in G , and let $B^\pm = U^\pm T$ be the Borel subgroup of G . We have the isomorphism $\phi : W \xrightarrow{\sim} N/T$ defined by $\phi(s_i) = N_i T/T$. An element

$\bar{s}_i := x_i(-1)y_i(1)x_i(-1) = \phi_i\left(\begin{pmatrix} 0 & \pm 1 \\ \mp 1 & 0 \end{pmatrix}\right)$ is in $N_G(T)$, which is a representative of $s_i \in W = N_G(T)/T$.

Geometric crystals. Let X be an ind-variety, $\gamma_i : X \rightarrow \mathbb{C}$ and $\varepsilon_i : X \rightarrow \mathbb{C}$ ($i \in I$) rational functions on X , and $e_i : \mathbb{C}^\times \times X \rightarrow X$ ($(c, x) \mapsto e_i^c(x)$) a rational \mathbb{C}^\times -action.

Definition 2.1. A quadruple $(X, \{e_i\}_{i \in I}, \{\gamma_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$ is a G (or \mathfrak{g})-geometric crystal if it satisfies these conditions:

- (i) $\{1\} \times X \subset \text{dom}(e_i)$ for any $i \in I$.
- (ii) $\gamma_j(e_i^c(x)) = c^{a_{ij}} \gamma_j(x)$.
- (iii) The e_i satisfy

$$\begin{aligned} e_i^{c_1} e_j^{c_2} &= e_j^{c_2} e_i^{c_1} && \text{if } a_{ij} = a_{ji} = 0, \\ e_i^{c_1} e_j^{c_1 c_2} e_i^{c_2} &= e_j^{c_2} e_i^{c_1 c_2} e_j^{c_1} && \text{if } a_{ij} = a_{ji} = -1, \\ e_i^{c_1} e_j^{c_1^2 c_2} e_i^{c_1 c_2} e_j^{c_2} &= e_j^{c_2} e_i^{c_1 c_2} e_j^{c_1^2 c_2} e_i^{c_1} && \text{if } a_{ij} = -2, a_{ji} = -1, \\ e_i^{c_1} e_j^{c_1^3 c_2} e_i^{c_1^2 c_2} e_j^{c_1 c_2} e_i^{c_2} &= e_j^{c_2} e_i^{c_1 c_2} e_j^{c_1^3 c_2} e_i^{c_1^2 c_2} e_j^{c_1 c_2} e_i^{c_1} && \text{if } a_{ij} = -3, a_{ji} = -1. \end{aligned}$$

- (iv) $\varepsilon_i(e_i^c(x)) = c^{-1} \varepsilon_i(x)$ and $\varepsilon_i(e_j^c(x)) = \varepsilon_i(x)$ if $a_{i,j} = a_{j,i} = 0$.

Condition (iv) is slightly modified from the one in [Igarashi and Nakashima 2010; Nakashima 2007; 2010].

Let W be the Weyl group associated with \mathfrak{g} . Define $R(w)$ for $w \in W$ by

$$R(w) := \{(i_1, i_2, \dots, i_l) \in I^l \mid w = s_{i_1} s_{i_2} \cdots s_{i_l}\},$$

where l is the length of w . Then $R(w)$ is the set of reduced words of w . For a word $\mathbf{i} = (i_1, \dots, i_l) \in R(w)$ ($w \in W$), set $\alpha^{(j)} := s_{i_1} \cdots s_{i_{j+1}}(\alpha_{i_j})$ ($1 \leq j \leq l$) and

$$e_{\mathbf{i}} : T \times X \rightarrow X, \quad (t, x) \mapsto e_{\mathbf{i}}^t(x) := e_{i_1}^{\alpha^{(1)}(t)} e_{i_2}^{\alpha^{(2)}(t)} \cdots e_{i_l}^{\alpha^{(l)}(t)}(x).$$

Condition (iii) above amounts to saying that $e_{\mathbf{i}} = e_{\mathbf{i}'}$ for any $w \in W$ and $\mathbf{i}, \mathbf{i}' \in R(w)$.

Geometric crystal on Schubert cell. Let $w \in W$ be a Weyl group element and take a reduced expression $w = s_{i_1} \cdots s_{i_l}$. Let $X := G/B$ be the flag variety, which is an ind-variety and $X_w \subset X$ the Schubert cell associated with w , which has a natural geometric crystal structure [Berenstein and Kazhdan 2000; Nakashima 2005a]. For $\mathbf{i} := (i_1, \dots, i_k)$, set

$$(2-1) \quad B_{\mathbf{i}}^- := \{Y_{\mathbf{i}}(c_1, \dots, c_k) := Y_{i_1}(c_1) \cdots Y_{i_k}(c_k) \mid c_1 \cdots c_k \in \mathbb{C}^\times\} \subset B^-,$$

where $Y_i(c) := y_i(\frac{1}{c})\alpha_i^\vee(c)$. This has a geometric crystal structure [Nakashima 2005a] isomorphic to X_w . The explicit forms of the action e_i^c , the rational function ε_i and γ_i on $B_{\mathbf{i}}^-$ are given by

$$e_i^c(Y_{\mathbf{i}}(c_1, \dots, c_k)) = Y_{\mathbf{i}}(\mathcal{C}_1, \dots, \mathcal{C}_k),$$

where

$$(2-2) \quad \mathcal{C}_j := c_j \cdot \frac{\sum_{1 \leq m \leq j, i_m = i} \frac{c}{c_1^{a_{1,i}} \cdots c_{m-1}^{a_{m-1,i}} c_m} + \sum_{j < m \leq k, i_m = i} \frac{1}{c_1^{a_{1,i}} \cdots c_{m-1}^{a_{m-1,i}} c_m}}{\sum_{1 \leq m < j, i_m = i} \frac{c}{c_1^{a_{1,i}} \cdots c_{m-1}^{a_{m-1,i}} c_m} + \sum_{j \leq m \leq k, i_m = i} \frac{1}{c_1^{a_{1,i}} \cdots c_{m-1}^{a_{m-1,i}} c_m}},$$

$$(2-3) \quad \varepsilon_i(Y_{\mathbf{i}}(c_1, \dots, c_k)) = \sum_{1 \leq m \leq k, i_m = i} \frac{1}{c_1^{a_{1,i}} \cdots c_{m-1}^{a_{m-1,i}} c_m},$$

$$(2-4) \quad \gamma_i(Y_{\mathbf{i}}(c_1, \dots, c_k)) = c_1^{a_{1,i}} \cdots c_k^{a_{k,i}}.$$

Positive structure, ultra-discretizations and tropicalizations. The setting is the same as in [Kashiwara et al. 2008]. Let $T = (\mathbb{C}^\times)^l$ be an algebraic torus over \mathbb{C} , with character lattice $X^*(T) := \text{Hom}(T, \mathbb{C}^\times) \cong \mathbb{Z}^l$ and cocharacter lattice $X_*(T) := \text{Hom}(\mathbb{C}^\times, T) \cong \mathbb{Z}^l$. Set $R := \mathbb{C}(c)$ and define

$$v : R \setminus \{0\} \rightarrow \mathbb{Z}, \quad f(c) \mapsto \deg f(c),$$

where \deg is the degree of poles at $c = \infty$. Note that for $f_1, f_2 \in R \setminus \{0\}$, we have

$$(2-5) \quad v(f_1 f_2) = v(f_1) + v(f_2), \quad v\left(\frac{f_1}{f_2}\right) = v(f_1) - v(f_2).$$

A nonzero rational function on an algebraic torus T is called *positive* if it can be written as g/h where g and h are a positive linear combination of characters of T .

Definition 2.2. Let $f : T \rightarrow T'$ be a rational morphism between two algebraic tori T and T' . We say that f is *positive* if $\eta \circ f$ is positive for any character $\eta : T' \rightarrow \mathbb{C}$.

Denote by $\text{Mor}^+(T, T')$ the set of positive rational morphisms from T to T' .

Lemma 2.3 [Berenstein and Kazhdan 2000]. *For any $f \in \text{Mor}^+(T_1, T_2)$ and any $g \in \text{Mor}^+(T_2, T_3)$, the composition $g \circ f$ is well-defined and lies in $\text{Mor}^+(T_1, T_3)$.*

By Lemma 2.3, we can define a category \mathcal{T}_+ whose objects are algebraic tori over \mathbb{C} and arrows are positive rational morphisms.

Let $f : T \rightarrow T'$ be a positive rational morphism of algebraic tori T and T' . We define a map $\widehat{f} : X_*(T) \rightarrow X_*(T')$ by

$$\langle \eta, \widehat{f}(\xi) \rangle = v(\eta \circ f \circ \xi),$$

where $\eta \in X^*(T')$ and $\xi \in X_*(T)$.

Lemma 2.4 [Berenstein and Kazhdan 2000]. *For any algebraic tori T_1, T_2, T_3 , and positive rational morphisms $f \in \text{Mor}^+(T_1, T_2)$, $g \in \text{Mor}^+(T_2, T_3)$, we have $\widehat{g \circ f} = \widehat{g} \circ \widehat{f}$.*

Let \mathfrak{Set} denote the category of sets and set maps. By the lemma, we obtain a functor

$$\begin{aligned} \mathcal{U}\mathcal{D} : \quad \mathcal{T}_+ &\rightarrow \mathfrak{Set} \\ T &\mapsto X_*(T) \\ (f : T \rightarrow T') &\mapsto (\widehat{f} : X_*(T) \rightarrow X_*(T')). \end{aligned}$$

Definition 2.5 [Berenstein and Kazhdan 2000]. Let

$$\chi = (X, \{e_i\}_{i \in I}, \{\text{wt}_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$$

be a geometric crystal, T' an algebraic torus and $\theta : T' \rightarrow X$ a birational isomorphism. The isomorphism θ is called *positive structure* on χ if

- (i) for any $i \in I$ the rational functions $\gamma_i \circ \theta : T' \rightarrow \mathbb{C}$ and $\varepsilon_i \circ \theta : T' \rightarrow \mathbb{C}$ are positive, and
- (ii) for any $i \in I$, the rational morphism $e_{i,\theta} : \mathbb{C}^\times \times T' \rightarrow T'$ defined by $e_{i,\theta}(c, t) := \theta^{-1} \circ e_i^c \circ \theta(t)$ is positive.

Let $\theta : T \rightarrow X$ be a positive structure on a geometric crystal $\chi = (X, \{e_i\}_{i \in I}, \{\text{wt}_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$. Applying the functor $\mathcal{U}\mathcal{D}$ to positive rational morphisms $e_{i,\theta} : \mathbb{C}^\times \times T' \rightarrow T'$ and $\gamma_i, \varepsilon_i \circ \theta : T' \rightarrow \mathbb{C}$ (the notations are as above), we obtain

$$\begin{aligned} \tilde{e}_i &:= \mathcal{U}\mathcal{D}(e_{i,\theta}) : \mathbb{Z} \times X_*(T) \rightarrow X_*(T), \\ \text{wt}_i &:= \mathcal{U}\mathcal{D}(\gamma_i \circ \theta) : X_*(T') \rightarrow \mathbb{Z}, \\ \varepsilon_i &:= \mathcal{U}\mathcal{D}(\varepsilon_i \circ \theta) : X_*(T') \rightarrow \mathbb{Z}. \end{aligned}$$

Now, for given positive structure $\theta : T' \rightarrow X$ on a geometric crystal $\chi = (X, \{e_i\}_{i \in I}, \{\text{wt}_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$, we associate the quadruple $(X_*(T'), \{\tilde{e}_i\}_{i \in I}, \{\text{wt}_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$ with a free pre-crystal structure (see [Berenstein and Kazhdan 2000, 2.2]) and denote it by $\mathcal{U}\mathcal{D}_{\theta, T'}(\chi)$.

Theorem 2.6 [Berenstein and Kazhdan 2000; Nakashima 2005a]. *For any geometric crystal $\chi = (X, \{e_i\}_{i \in I}, \{\gamma_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$ and positive structure $\theta : T' \rightarrow X$, the associated pre-crystal $\mathcal{U}\mathcal{D}_{\theta, T'}(\chi) = (X_*(T'), \{\tilde{e}_i\}_{i \in I}, \{\text{wt}_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$ is a crystal (see [Berenstein and Kazhdan 2000, 2.2]).*

Now, let \mathcal{GC}^+ be the category whose objects are triplets (χ, T', θ) , where $\chi = (X, \{e_i\}, \{\gamma_i\}, \{\varepsilon_i\})$ is a geometric crystal and $\theta : T' \rightarrow X$ is a positive structure on χ , and whose morphisms $f : (\chi_1, T'_1, \theta_1) \rightarrow (\chi_2, T'_2, \theta_2)$ are given by morphisms $\varphi : X_1 \rightarrow X_2$ ($\chi_i = (X_i, \dots)$) such that

$$f := \theta_2^{-1} \circ \varphi \circ \theta_1 : T'_1 \rightarrow T'_2,$$

is a positive rational morphism. Let \mathcal{CR} be the category of crystals. **Theorem 2.6** yields:

Corollary 2.7. *The map $\mathcal{U}\mathcal{D} = \mathcal{U}\mathcal{D}_{\theta, T'}$ defined above is a functor*

$$\begin{aligned} \mathcal{U}\mathcal{D} : \quad & \mathcal{GC}^+ & \rightarrow & \mathcal{CR} \\ & (\chi, T', \theta) & \mapsto & X_*(T'), \\ & (f : (\chi_1, T'_1, \theta_1) \rightarrow (\chi_2, T'_2, \theta_2)) & \mapsto & (\widehat{f} : X_*(T'_1) \rightarrow X_*(T'_2)). \end{aligned}$$

We call the functor $\mathcal{U}\mathcal{D}$ “ultra-discretization” as in [Nakashima 2005a; 2005b] instead of “tropicalization” as in [Berenstein and Kazhdan 2000]. And for a crystal B , if there exists a geometric crystal χ and a positive structure $\theta : T' \rightarrow X$ on χ such that $\mathcal{U}\mathcal{D}(\chi, T', \theta) \cong B$ as crystals, we call an object (χ, T', θ) in \mathcal{GC}^+ a *tropicalization* of B , where it is not known that this correspondence is a functor.

3. Limit of perfect crystals

We review limit of perfect crystals following [Kang et al. 1994]. (See also [Kang et al. 1992a; 1992b].)

Crystals. First we review the theory of crystals, which is the notion obtained by abstracting the combinatorial properties of crystal bases. Let $(A, \{\alpha_i\}_{i \in I}, \{\alpha_i^\vee\}_{i \in I})$ be a Cartan data.

Definition 3.1. A *crystal* B is a set endowed with maps

$$\begin{aligned} \text{wt} : B &\rightarrow P, \\ \varepsilon_i : B &\rightarrow \mathbb{Z} \sqcup \{-\infty\}, \quad \varphi_i : B \rightarrow \mathbb{Z} \sqcup \{-\infty\} \quad \text{for } i \in I, \\ \tilde{e}_i : B \sqcup \{0\} &\rightarrow B \sqcup \{0\}, \quad \tilde{f}_i : B \sqcup \{0\} \rightarrow B \sqcup \{0\} \quad \text{for } i \in I, \\ \tilde{e}_i(0) = \tilde{f}_i(0) &= 0. \end{aligned}$$

satisfying the following axioms, for all $b, b_1, b_2 \in B$:

$$\begin{aligned} \varphi_i(b) &= \varepsilon_i(b) + \langle \alpha_i^\vee, \text{wt}(b) \rangle, \\ \text{wt}(\tilde{e}_i b) &= \text{wt}(b) + \alpha_i \quad \text{if } \tilde{e}_i b \in B, \\ \text{wt}(\tilde{f}_i b) &= \text{wt}(b) - \alpha_i \quad \text{if } \tilde{f}_i b \in B, \\ \tilde{e}_i b_2 = b_1 &\iff \tilde{f}_i b_1 = b_2, \\ \varepsilon_i(b) = -\infty &\implies \tilde{e}_i b = \tilde{f}_i b = 0. \end{aligned}$$

The following tensor product structure is one of the most crucial properties of crystals.

Theorem 3.2. *Let B_1 and B_2 be crystals, and set*

$$B_1 \otimes B_2 := \{b_1 \otimes b_2; b_j \in B_j (j = 1, 2)\}.$$

(i) $B_1 \otimes B_2$ is a crystal.

(ii) For $b_1 \in B_1$ and $b_2 \in B_2$, we have

$$\begin{aligned} \tilde{f}_i(b_1 \otimes b_2) &= \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \\ b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \leq \varepsilon_i(b_2), \end{cases} \\ \tilde{e}_i(b_1 \otimes b_2) &= \begin{cases} b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2), \\ \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \geq \varepsilon_i(b_2). \end{cases} \end{aligned}$$

Definition 3.3. Let B_1 and B_2 be crystals. A *strict morphism* of crystals

$$\psi : B_1 \rightarrow B_2$$

is a map $\psi : B_1 \sqcup \{0\} \rightarrow B_2 \sqcup \{0\}$ such that $\psi(0) = 0$, $\psi(B_1) \subset B_2$, ψ commutes with all \tilde{e}_i and \tilde{f}_i , and

$$\text{wt}(\psi(b)) = \text{wt}(b), \quad \varepsilon_i(\psi(b)) = \varepsilon_i(b), \quad \varphi_i(\psi(b)) = \varphi_i(b) \text{ for any } b \in B_1.$$

A bijective strict morphism is called an *isomorphism of crystals*.

Example 3.4. If (L, B) is a crystal base, then B is a crystal. Hence, for the crystal base $(L(\infty), B(\infty))$ of the nilpotent subalgebra $U_q^-(\mathfrak{g})$ of the quantum algebra $U_q(\mathfrak{g})$, $B(\infty)$ is a crystal.

Example 3.5. For $\lambda \in P$, set $T_\lambda := \{t_\lambda\}$. We define a crystal structure on T_λ by

$$\tilde{e}_i(t_\lambda) = \tilde{f}_i(t_\lambda) = 0, \quad \varepsilon_i(t_\lambda) = \varphi_i(t_\lambda) = -\infty, \quad \text{wt}(t_\lambda) = \lambda.$$

Definition 3.6. For a crystal B , a colored oriented graph structure is associated with B by

$$b_1 \xrightarrow{i} b_2 \iff \tilde{f}_i b_1 = b_2.$$

We call this graph the *crystal graph* of B .

Affine weights. Let \mathfrak{g} be an affine Lie algebra. The sets \mathfrak{t} , $\{\alpha_i\}_{i \in I}$ and $\{\alpha_i^\vee\}_{i \in I}$ be as in Section 2. We take $\dim \mathfrak{t} = \#I + 1$. Let $\delta \in Q_+$ be the unique element satisfying $\{\lambda \in Q \mid \langle \alpha_i^\vee, \lambda \rangle = 0 \text{ for any } i \in I\} = \mathbb{Z}\delta$ and $\mathbf{c} \in \mathfrak{g}$ be the canonical central element satisfying $\{h \in Q^\vee \mid \langle h, \alpha_i \rangle = 0 \text{ for any } i \in I\} = \mathbb{Z}\mathbf{c}$. We write, as in [Kac 1990, 6.1],

$$\mathbf{c} = \sum_i a_i^\vee \alpha_i^\vee, \quad \delta = \sum_i a_i \alpha_i.$$

Let $(\ , \)$ be the nondegenerate W -invariant symmetric bilinear form on \mathfrak{t}^* normalized by $(\delta, \lambda) = \langle \mathbf{c}, \lambda \rangle$ for $\lambda \in \mathfrak{t}^*$. Let us set $\mathfrak{t}_{\text{cl}}^* := \mathfrak{t}^*/\mathbb{C}\delta$ and let $\text{cl} : \mathfrak{t}^* \rightarrow \mathfrak{t}_{\text{cl}}^*$ be the canonical projection. Here we have $\mathfrak{t}_{\text{cl}}^* \cong \bigoplus_i (\mathbb{C}\alpha_i^\vee)^*$. Set $\mathfrak{t}_0^* := \{\lambda \in \mathfrak{t}^* \mid \langle \mathbf{c}, \lambda \rangle = 0\}$, $(\mathfrak{t}_{\text{cl}}^*)_0 := \text{cl}(\mathfrak{t}_0^*)$. Since $(\delta, \delta) = 0$, we have a positive definite symmetric form on $\mathfrak{t}_{\text{cl}}^*$ induced by the one on \mathfrak{t}^* . Let $\Lambda_i \in \mathfrak{t}_{\text{cl}}^*$ ($i \in I$) be a classical weight such that $\langle \alpha_i^\vee, \Lambda_j \rangle = \delta_{i,j}$, which is called a fundamental weight. We choose P so that $P_{\text{cl}} := \text{cl}(P)$ coincides with $\bigoplus_{i \in I} \mathbb{Z}\Lambda_i$ and we call P_{cl} a *classical weight lattice*.

Perfect crystals and their limits. Let \mathfrak{g} be an affine Lie algebra, let P_{cl} be a classical weight lattice as above and set $(P_{\text{cl}})_l^+ := \{\lambda \in P_{\text{cl}} \mid \langle \mathbf{c}, \lambda \rangle = l, \langle \alpha_i^\vee, \lambda \rangle \geq 0\}$ ($l \in \mathbb{Z}_{>0}$).

Definition 3.7. A crystal B is a *perfect crystal* of level l if the following conditions are satisfied:

- (i) $B \otimes B$ is connected as a crystal graph.
- (ii) There exists $\lambda_0 \in P_{\text{cl}}$ such that

$$\text{wt}(B) \subset \lambda_0 + \sum_{i \neq 0} \mathbb{Z}_{\leq 0} \text{cl}(\alpha_i), \quad \#B_{\lambda_0} = 1.$$

- (iii) There exists a finite-dimensional $U'_q(\mathfrak{g})$ -module V with a crystal pseudobase B_{ps} such that $B \cong B_{\text{ps}}/\pm 1$.
- (iv) For any $b \in B$, we have $\langle \mathbf{c}, \varepsilon(b) \rangle \geq l$.
- (v) The maps $\varepsilon, \varphi : B^{\min} := \{b \in B \mid \langle \mathbf{c}, \varepsilon(b) \rangle = l\} \rightarrow (P_{\text{cl}}^+)_l$ are bijective, where $\varepsilon(b) := \sum_i \varepsilon_i(b) \Lambda_i$ and $\varphi(b) := \sum_i \varphi_i(b) \Lambda_i$.

Let $\{B_l\}_{l \geq 1}$ be a family of perfect crystals of level l and set $J := \{(l, b) \mid l > 0, b \in B_l^{\min}\}$.

Definition 3.8. A crystal B_∞ with an element b_∞ is called a *limit of* $\{B_l\}_{l \geq 1}$ if

- (i) $\text{wt}(b_\infty) = \varepsilon(b_\infty) = \varphi(b_\infty) = 0$;
- (ii) for any $(l, b) \in J$, there exists an embedding of crystals

$$f_{(l,b)} : T_{\varepsilon(b)} \otimes B_l \otimes T_{-\varphi(b)} \hookrightarrow B_\infty, \quad t_{\varepsilon(b)} \otimes b \otimes t_{-\varphi(b)} \mapsto b_\infty;$$

- (iii) $B_\infty = \bigcup_{(l,b) \in J} \text{Im} f_{(l,b)}$.

As for the crystal T_λ , see [Example 3.5](#). If a limit exists for a family $\{B_l\}$, we say that $\{B_l\}$ is a *coherent family* of perfect crystals.

Here is one of the most important properties of limit of perfect crystals.

Proposition 3.9. *For the crystal $B(\infty)$ as in [Example 3.4](#), we have an isomorphism of crystals*

$$B(\infty) \otimes B_\infty \xrightarrow{\sim} B(\infty).$$

4. Perfect crystals of type $G_2^{(1)}$


In this section, we review the family of perfect crystals of type $G_2^{(1)}$ and its limit [[Misra et al. 2010](#)].

We fix the data for $G_2^{(1)}$. Let $\{\alpha_0, \alpha_1, \alpha_2\}$, $\{\alpha_0^\vee, \alpha_1^\vee, \alpha_2^\vee\}$ and $\{\Lambda_0, \Lambda_1, \Lambda_2\}$ be the set of simple roots, simple coroots and fundamental weights, respectively. The

Cartan matrix $A = (a_{ij})_{i,j=0,1,2}$ is given by

$$A = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -3 & 2 \end{pmatrix},$$

and its Dynkin diagram is as follows:



The standard null root δ and the canonical central element \mathbf{c} are given by

$$\delta = \alpha_0 + 2\alpha_1 + 3\alpha_2 \quad \text{and} \quad \mathbf{c} = \alpha_0^\vee + 2\alpha_1^\vee + \alpha_2^\vee,$$

where $\alpha_0 = 2\Lambda_0 - \Lambda_1 + \delta$, $\alpha_1 = -\Lambda_0 + 2\Lambda_1 - 3\Lambda_2$, and $\alpha_2 = -\Lambda_1 + 2\Lambda_2$.

For a positive integer l we introduce $G_2^{(1)}$ -crystals B_l and B_∞ as

$$B_l = \left\{ b = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1) \in (\mathbb{Z}_{\geq 0}/3)^6 \left| \begin{array}{l} 3b_3 \equiv 3\bar{b}_3 \pmod{2}, \\ \sum_{i=1,2} (b_i + \bar{b}_i) + \frac{1}{2}(b_3 + \bar{b}_3) \leq l \\ b_1, \bar{b}_1, b_2 - b_3, \bar{b}_3 - \bar{b}_2 \in \mathbb{Z} \end{array} \right. \right\},$$

$$B_\infty = \left\{ b = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1) \in (\mathbb{Z}/3)^6 \left| \begin{array}{l} 3b_3 \equiv 3\bar{b}_3 \pmod{2}, \\ b_1, \bar{b}_1, b_2 - b_3, \bar{b}_3 - \bar{b}_2 \in \mathbb{Z} \end{array} \right. \right\}.$$

Now we describe the explicit crystal structures of B_l and B_∞ . Indeed, most of them coincide with each other except for ε_0 and φ_0 . In the rest of this section, we use the following convention: $(x)_+ = \max(x, 0)$. For $b = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1)$ we define

$$(4-1) \quad s(b) = b_1 + b_2 + \frac{1}{2}(b_3 + \bar{b}_3) + \bar{b}_2 + \bar{b}_1,$$

and

$$(4-2) \quad z_1 = \bar{b}_1 - b_1, \quad z_2 = \bar{b}_2 - \bar{b}_3, \quad z_3 = b_3 - b_2, \quad z_4 = \frac{1}{2}(\bar{b}_3 - b_3).$$

Now we define conditions and (F_1) – (F_6) as follows:

$$(4-3) \quad \left\{ \begin{array}{l} (F_1) \quad z_1 + z_2 + z_3 + 3z_4 \leq 0, \quad z_1 + z_2 + 3z_4 \leq 0, \quad z_1 + z_2 \leq 0, \quad z_1 \leq 0, \\ (F_2) \quad z_1 + z_2 + z_3 + 3z_4 \leq 0, \quad z_2 + 3z_4 \leq 0, \quad z_2 \leq 0, \quad z_1 > 0, \\ (F_3) \quad z_1 + z_3 + 3z_4 \leq 0, \quad z_3 + 3z_4 \leq 0, \quad z_4 \leq 0, \quad z_2 > 0, \quad z_1 + z_2 > 0, \\ (F_4) \quad z_1 + z_2 + 3z_4 > 0, \quad z_2 + 3z_4 > 0, \quad z_4 > 0, \quad z_3 \leq 0, \quad z_1 + z_3 \leq 0, \\ (F_5) \quad z_1 + z_2 + z_3 + 3z_4 > 0, \quad z_3 + 3z_4 > 0, \quad z_3 > 0, \quad z_1 \leq 0, \\ (F_6) \quad z_1 + z_2 + z_3 + 3z_4 > 0, \quad z_1 + z_3 + 3z_4 > 0, \quad z_1 + z_3 > 0, \quad z_1 > 0. \end{array} \right.$$

Conditions (E_i) , for $1 \leq i \leq 6$, are defined from (F_i) by replacing $>$ with \geq and \leq with $<$.

We also define

$$(4-4) \quad A = (0, z_1, z_1 + z_2, z_1 + z_2 + 3z_4, z_1 + z_2 + z_3 + 3z_4, 2z_1 + z_2 + z_3 + 3z_4).$$

Then for $b = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1) \in B_l$ or B_∞ , the values of $\tilde{\varepsilon}_i b$, $\tilde{f}_i b$, $\varepsilon_i(b)$, and $\varphi_i(b)$, for $i = 0, 1, 2$, are as follows:

$$\tilde{\varepsilon}_0 b = \begin{cases} (b_1 - 1, \dots) & \text{if } (E_1), \\ (\dots, b_3 - 1, \bar{b}_3 - 1, \dots, \bar{b}_1 + 1) & \text{if } (E_2), \\ (\dots, b_2 - \frac{2}{3}, b_3 - \frac{2}{3}, \bar{b}_3 + \frac{4}{3}, \bar{b}_2 + \frac{1}{3}, \dots) & \text{if } (E_3) \text{ and } z_4 = -\frac{1}{3}, \\ (\dots, b_2 - \frac{1}{3}, b_3 - \frac{4}{3}, \bar{b}_3 + \frac{2}{3}, \bar{b}_2 + \frac{2}{3}, \dots) & \text{if } (E_3) \text{ and } z_4 = -\frac{2}{3}, \\ (\dots, b_3 - 2, \dots, \bar{b}_2 + 1, \dots) & \text{if } (E_3) \text{ and } z_4 \neq -\frac{1}{3}, -\frac{2}{3}, \\ (\dots, b_2 - 1, \dots, \bar{b}_3 + 2, \dots) & \text{if } (E_4), \\ (b_1 - 1, \dots, b_3 + 1, \bar{b}_3 + 1, \dots) & \text{if } (E_5), \\ (\dots, \bar{b}_1 + 1) & \text{if } (E_6), \end{cases}$$

$$\tilde{f}_0 b = \begin{cases} (b_1 + 1, \dots) & \text{if } (F_1), \\ (\dots, b_3 + 1, \bar{b}_3 + 1, \dots, \bar{b}_1 - 1) & \text{if } (F_2), \\ (\dots, b_3 + 2, \dots, \bar{b}_2 - 1, \dots) & \text{if } (F_3), \\ (\dots, b_2 + \frac{1}{3}, b_3 + \frac{4}{3}, \bar{b}_3 - \frac{2}{3}, \bar{b}_2 - \frac{2}{3}, \dots) & \text{if } (F_4) \text{ and } z_4 = \frac{1}{3}, \\ (\dots, b_2 + \frac{2}{3}, b_3 + \frac{2}{3}, \bar{b}_3 - \frac{4}{3}, \bar{b}_2 - \frac{1}{3}, \dots) & \text{if } (F_4) \text{ and } z_4 = \frac{2}{3}, \\ (\dots, b_2 + 1, \dots, \bar{b}_3 - 2, \dots) & \text{if } (F_4) \text{ and } z_4 \neq \frac{1}{3}, \frac{2}{3}, \\ (b_1 + 1, \dots, b_3 - 1, \bar{b}_3 - 1, \dots) & \text{if } (F_5), \\ (\dots, \bar{b}_1 - 1) & \text{if } (F_6), \end{cases}$$

$$\tilde{\varepsilon}_1 b = \begin{cases} (\dots, \bar{b}_2 + 1, \bar{b}_1 - 1) & \text{if } \bar{b}_2 - \bar{b}_3 \geq (b_2 - b_3)_+, \\ (\dots, b_3 + 1, \bar{b}_3 - 1, \dots) & \text{if } \bar{b}_2 - \bar{b}_3 < 0 \leq b_3 - b_2, \\ (b_1 + 1, b_2 - 1, \dots) & \text{if } (\bar{b}_2 - \bar{b}_3)_+ < b_2 - b_3, \end{cases}$$

$$\tilde{f}_1 b = \begin{cases} (b_1 - 1, b_2 + 1, \dots) & \text{if } (\bar{b}_2 - \bar{b}_3)_+ \leq b_2 - b_3, \\ (\dots, b_3 - 1, \bar{b}_3 + 1, \dots) & \text{if } \bar{b}_2 - \bar{b}_3 \leq 0 < b_3 - b_2, \\ (\dots, \bar{b}_2 - 1, \bar{b}_1 + 1) & \text{if } \bar{b}_2 - \bar{b}_3 > (b_2 - b_3)_+, \end{cases}$$

$$\tilde{\varepsilon}_2 b = \begin{cases} (\dots, \bar{b}_3 + \frac{2}{3}, \bar{b}_2 - \frac{1}{3}, \dots) & \text{if } \bar{b}_3 \geq b_3, \\ (\dots, b_2 + \frac{1}{3}, b_3 - \frac{2}{3}, \dots) & \text{if } \bar{b}_3 < b_3, \end{cases}$$

$$\tilde{f}_2 b = \begin{cases} (\dots, b_2 - \frac{1}{3}, b_3 + \frac{2}{3}, \dots) & \text{if } \bar{b}_3 \leq b_3, \\ (\dots, \bar{b}_3 - \frac{2}{3}, \bar{b}_2 + \frac{1}{3}, \dots) & \text{if } \bar{b}_3 > b_3, \end{cases}$$

$$\begin{aligned}\varepsilon_1(b) &= \bar{b}_1 + (\bar{b}_3 - \bar{b}_2 + (b_2 - b_3)_+)_+, & \varphi_1(b) &= b_1 + (b_3 - b_2 + (\bar{b}_2 - \bar{b}_3)_+)_+, \\ \varepsilon_2(b) &= 3\bar{b}_2 + \frac{3}{2}(b_3 - \bar{b}_3)_+, & \varphi_2(b) &= 3b_2 + \frac{3}{2}(\bar{b}_3 - b_3)_+, \\ \varepsilon_0(b) &= \begin{cases} l - s(b) + \max A - (2z_1 + z_2 + z_3 + 3z_4) & b \in B_l, \\ -s(b) + \max A - (2z_1 + z_2 + z_3 + 3z_4) & b \in B_\infty, \end{cases} \\ \varphi_0(b) &= \begin{cases} l - s(b) + \max A & b \in B_l, \\ -s(b) + \max A & b \in B_\infty. \end{cases}\end{aligned}$$

For $b \in B_l$ if $\tilde{e}_i b$ or $\tilde{f}_i b$ does not belong to B_l , namely, if b_j or \bar{b}_j for some j becomes negative or $s(b)$ exceeds l , we understand it to be 0.

Theorem 4.1 [Misra et al. 2010]. (i) *The $G_2^{(1)}$ -crystal B_l is a perfect crystal of level l .*

(ii) *The family of the perfect crystals $\{B_l\}_{l \geq 1}$ forms a coherent family and the crystal B_∞ is its limit with the vector $b_\infty = (0, 0, 0, 0, 0, 0)$.*

As was shown in [Misra et al. 2010], the minimal elements are given by

$$(B_l)_{\min} = \{(\alpha, \beta, \beta, \beta, \beta, \alpha) \mid \alpha \in \mathbb{Z}_{\geq 0}, \beta \in (\mathbb{Z}_{\geq 0})/3, 2\alpha + 3\beta \leq l\}.$$

Set $J = \{(l, b) \mid l \in \mathbb{Z}_{\geq 1}, b \in (B_l)_{\min}\}$ and let the maps $\varepsilon, \varphi : (B_l)_{\min} \rightarrow (P_{\text{cl}}^+)_l$ be as in Definition 3.7. Then we have $\text{wt } b_\infty = 0$ and

$$\varepsilon_i(b_\infty) = \varphi_i(b_\infty) = 0 \quad \text{for } i = 0, 1, 2.$$

For $(l, b_0) \in J$, since $\varepsilon(b_0) = \varphi(b_0)$, one can set $\lambda = \varepsilon(b_0) = \varphi(b_0)$. For

$$b = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1) \in B_l$$

we define a map

$$f_{(l, b_0)} : T_\lambda \otimes B_l \otimes B_{-\lambda} \rightarrow B_\infty$$

by

$$f_{(l, b_0)}(t_\lambda \otimes b \otimes t_{-\lambda}) = b' = (v_1, v_2, v_3, \bar{v}_3, \bar{v}_2, \bar{v}_1)$$

where $b_0 = (\alpha, \beta, \beta, \beta, \beta, \alpha)$, and

$$\begin{aligned}v_1 &= b_1 - \alpha, & \bar{v}_1 &= \bar{b}_1 - \alpha, \\ v_2 &= b_2 - \beta, & \bar{v}_2 &= \bar{b}_2 - \beta, \\ v_3 &= b_3 - \beta, & \bar{v}_3 &= \bar{b}_3 - \beta.\end{aligned}$$

Finally, we obtain

$$B_\infty = \bigcup_{(l, b) \in J} \text{Im } f_{(l, b)}.$$

5. Affine geometric crystal $\mathcal{V}_1(D_4^{(3)})$

Fundamental representation $W(\varpi_1)$ for $D_4^{(3)}$. Let $\mathbf{c} = \sum_i a_i^\vee \alpha_i^\vee$ be the canonical central element in an affine Lie algebra \mathfrak{g} (see [Kac 1990, 6.1]), $\{\Lambda_i \mid i \in I\}$ the set of fundamental weights as in the previous section and $\varpi_1 := \Lambda_1 - a_1^\vee \Lambda_0$ the fundamental weight (of level 0). Let $W(\varpi_1)$ be the fundamental representation of $U'_q(\mathfrak{g})$ associated with ϖ_1 [Kashiwara 2002].

By [Kashiwara 2002, Theorem 5.17], $W(\varpi_1)$ is a finite-dimensional irreducible integrable $U'_q(\mathfrak{g})$ -module and has a global basis with a simple crystal. Thus, we can consider the specialization $q = 1$ and obtain the finite-dimensional \mathfrak{g} -module $W(\varpi_1)$, which we call a fundamental representation of \mathfrak{g} and use the same notation as above.

We shall present the explicit form of $W(\varpi_1)$ for $\mathfrak{g} = D_4^{(3)}$.


$W(\varpi_1)$ for $D_4^{(3)}$. The Cartan matrix $A = (a_{i,j})_{i,j=0,1,2}$ of type $D_4^{(3)}$ is given by

$$A = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -3 \\ 0 & -1 & 2 \end{pmatrix}.$$

Then the simple roots are

$$\alpha_0 = 2\Lambda_0 - \Lambda_1 + \delta, \quad \alpha_1 = -\Lambda_0 + 2\Lambda_1 - \Lambda_2, \quad \alpha_2 = -3\Lambda_1 + 2\Lambda_2,$$

and the Dynkin diagram is this:



The $D_4^{(3)}$ -module $W(\varpi_1)$ is an 8-dimensional module with the basis

$$\{v_1, v_2, v_3, v_0, \emptyset, v_{\bar{3}}, v_{\bar{2}}, v_{\bar{1}}\}.$$

The explicit form of $W(\varpi_1)$ is given in [Kashiwara et al. 2007].

$$\begin{aligned} \text{wt}(v_1) &= \Lambda_1 - 2\Lambda_0, & \text{wt}(v_2) &= -\Lambda_0 - \Lambda_1 + \Lambda_2, & \text{wt}(v_3) &= -\Lambda_0 + 2\Lambda_1 - \Lambda_2, \\ \text{wt}(v_{\bar{i}}) &= -\text{wt}(v_i) \quad (i = 1, 2, 3), & \text{wt}(v_0) &= \text{wt}(\emptyset) = 0. \end{aligned}$$

The actions of e_i and f_i on these basis vectors are given as follows:

$$\begin{aligned} f_0(v_0, v_{\bar{3}}, v_{\bar{2}}, v_{\bar{1}}, \emptyset) &= (v_1, v_2, v_3, \emptyset + \frac{1}{2}v_0, \frac{3}{2}v_1), \\ f_1(v_1, v_3, v_0, v_{\bar{2}}) &= (v_2, v_0, 2v_{\bar{3}}, v_{\bar{1}}), & f_2(v_2, v_{\bar{3}}) &= (v_3, v_{\bar{2}}), \\ e_0(v_1, v_2, v_3, v_0, \emptyset) &= (\emptyset + \frac{1}{2}v_0, v_{\bar{3}}, v_{\bar{2}}, v_{\bar{1}}, \frac{3}{2}v_{\bar{1}}), \\ e_1(v_2, v_0, v_{\bar{3}}, v_{\bar{1}}) &= (v_1, 2v_3, v_0, v_{\bar{2}}), & e_2(v_3, v_{\bar{2}}) &= (v_2, v_{\bar{3}}), \end{aligned}$$

where we give nontrivial actions only.

Construction of the affine geometric crystal $\mathcal{V}_1(D_4^{(3)})$ in $W(\varpi_1)$. In this section, we follow [Igarashi and Nakashima 2010]. For $\xi \in (\mathfrak{t}_{\text{cl}}^*)_0$, let $t(\xi)$ be the translation as in [Kashiwara 2002, Section 4] and $\tilde{\varpi}_i$ as in [Kashiwara 2005]; indeed, $\tilde{\varpi}_i := \max(1, 2/(\alpha_i, \alpha_i))\varpi_i$. Then we have

$$\begin{aligned} t(\tilde{\varpi}_1) &= s_0 s_1 s_2 s_1 s_2 s_1 =: w_1, \\ t(\text{wt}(v_{\tilde{2}})) &= s_2 s_1 s_2 s_1 s_0 s_1 =: w_2. \end{aligned}$$

Associated with these Weyl group elements w_1 and w_2 , we define algebraic varieties $\mathcal{V}_1 = \mathcal{V}_1(D_4^{(3)})$ and $\mathcal{V}_2 = \mathcal{V}_2(D_4^{(3)}) \subset W(\varpi_1)$ respectively:

$$\begin{aligned} \mathcal{V}_1 &:= \{V_1(x) := Y_0(x_0)Y_1(x_1)Y_2(x_2)Y_1(x_3)Y_2(x_4)Y_1(x_5)v_1 \mid x_i \in \mathbb{C}^\times, 0 \leq i \leq 5\}, \\ \mathcal{V}_2 &:= \{V_2(y) := Y_2(y_2)Y_1(y_1)Y_2(y_4)Y_1(y_3)Y_0(y_0)Y_1(y_5)v_{\tilde{2}} \mid y_i \in \mathbb{C}^\times, 0 \leq i \leq 5\}. \end{aligned}$$

Owing to the explicit forms of f_i 's on $W(\varpi_1)$ as above, we have $f_0^3 = 0$, $f_1^3 = 0$ and $f_2^2 = 0$ and then

$$Y_i(c) = \left(1 + \frac{f_i}{c} + \frac{f_i^2}{2c^2}\right)\alpha_i^\vee(c) \quad (i = 0, 1), \quad Y_2(c) = \left(1 + \frac{f_2}{c}\right)\alpha_2^\vee(c).$$

We get explicit forms of $V_1(x) \in \mathcal{V}_1$ and $V_2(y) \in \mathcal{V}_2$ as in [Nakashima 2007]:

$$\begin{aligned} V_1(x) &= \sum_{1 \leq i \leq 3} (X_i v_i + X_{\bar{i}} v_{\bar{i}}) + X_0 v_0 + X_\emptyset \emptyset, \\ V_2(y) &= \sum_{1 \leq i \leq 3} (Y_i v_i + Y_{\bar{i}} v_{\bar{i}}) + Y_0 v_0 + Y_\emptyset \emptyset. \end{aligned}$$

where the rational functions X_i 's and Y_i 's are all positive in (x_0, \dots, x_5) and (y_0, \dots, y_5) respectively (as for their explicit forms, see [Igarashi and Nakashima 2010]) and for any x there exist a unique rational function $a(x)$ and y such that $V_2(y) = a(x)V_1(x)$. Using this result, we get the positive birational isomorphism $\bar{\sigma}: \mathcal{V}_1 \rightarrow \mathcal{V}_2$ ($V_1(x) \mapsto V_2(y)$) and we know that its inverse $\bar{\sigma}^{-1}$ is also positive. The actions of \bar{e}_0^c on $V_2(y)$ (respectively $\bar{\gamma}_0(V_2(y))$ and $\bar{\varepsilon}_0(V_2(y))$) are induced from the ones on $Y_2(y_2)Y_1(y_1)Y_2(y_4)Y_1(y_3)Y_0(y_0)Y_1(y_5)$ as an element of the geometric crystal \mathcal{V}_2 . We define the action e_0^c on $V_1(x)$ by

$$(5-1) \quad e_0^c(V_1(x)) := \bar{\sigma}^{-1} \circ \bar{e}_0^c \circ \bar{\sigma}(V_1(x)).$$

We also define $\gamma_0(V_1(x))$ and $\varepsilon_0(V_1(x))$ by

$$(5-2) \quad \gamma_0(V_1(x)) := \bar{\gamma}_0(\bar{\sigma}(V_1(x))), \quad \varepsilon_0(V_1(x)) := \bar{\varepsilon}_0(\bar{\sigma}(V_1(x))).$$

Theorem 5.1 [Igarashi and Nakashima 2010]. *Together with (5-1), (5-2) on \mathcal{V}_1 , we obtain a positive affine geometric crystal $\chi := (\mathcal{V}_1, \{e_i\}_{i \in I}, \{\gamma_i\}_{i \in I}, \{\varepsilon_i\}_{i \in I})$*

($I = \{0, 1, 2\}$), whose explicit form is as follows: first we have e_i^c , γ_i , and ε_i , for $i = 1, 2$, from (2-2), (2-3), and (2-4):

$$e_1^c(V_1(x)) = V_1(x_0, \mathcal{C}_1 x_1, x_2, \mathcal{C}_3 x_3, x_4, \mathcal{C}_5 x_5),$$

$$e_2^c(V_1(x)) = V_1(x_0, x_1, \mathcal{C}_2 x_2, x_3, \mathcal{C}_4 x_4, x_5),$$

where

$$\mathcal{C}_1 = \frac{\frac{c x_0}{x_1} + \frac{x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}}{\frac{x_0}{x_1} + \frac{x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}}, \quad \mathcal{C}_3 = \frac{\frac{c x_0}{x_1} + \frac{c x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}}{\frac{c x_0}{x_1} + \frac{x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}},$$

$$\mathcal{C}_5 = \frac{c \left(\frac{x_0}{x_1} + \frac{x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5} \right)}{\frac{c x_0}{x_1} + \frac{c x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}}, \quad \mathcal{C}_2 = \frac{\frac{c x_1^3}{x_2} + \frac{x_1^3 x_3^3}{x_2^2 x_4}}{\frac{x_1^3}{x_2} + \frac{x_1^3 x_3^3}{x_2^2 x_4}}, \quad \mathcal{C}_4 = \frac{c \left(\frac{x_1^3}{x_2} + \frac{x_1^3 x_3^3}{x_2^2 x_4} \right)}{\frac{c x_1^3}{x_2} + \frac{x_1^3 x_3^3}{x_2^2 x_4}},$$

$$\varepsilon_1(V_1(x)) = \frac{x_0}{x_1} + \frac{x_0 x_2}{x_1^2 x_3} + \frac{x_0 x_2 x_4}{x_1^2 x_3^2 x_5}, \quad \varepsilon_2(V_1(x)) = \frac{x_1^3}{x_2} + \frac{x_1^3 x_3^3}{x_2^2 x_4},$$

$$\gamma_1(V_1(x)) = \frac{x_1^2 x_3^2 x_5^2}{x_0 x_2 x_4}, \quad \gamma_2(V_1(x)) = \frac{x_2^2 x_4^2}{x_1^3 x_3^3 x_5^3}.$$

We also have e_0^c , ε_0 and γ_0 on $V_1(x)$:

$$e_0^c(V_1(x)) = V_1 \left(\frac{D}{c \cdot E} x_0, \frac{F}{c \cdot E} x_1, \frac{G}{c^3 \cdot E^3} x_2, \frac{D \cdot H}{c^2 \cdot E \cdot F} x_3, \frac{D^3}{c^3 \cdot G} x_4, \frac{D}{c \cdot H} x_5 \right),$$

$$\varepsilon_0(V_1(x)) = \frac{E}{x_0^3 x_2 x_3}, \quad \gamma_0(V_1(x)) = \frac{x_0^2}{x_1 x_3 x_5},$$

where

$$D = c^2 x_0^2 x_2 x_3 + x_1 x_2 x_3^2 x_5 + c x_0 (x_1 x_3^3 + x_2 (x_3^2 + x_1 x_4 + x_1 x_3 x_5)),$$

$$E = x_0^2 x_2 x_3 + x_1 x_2 x_3^2 x_5 + x_0 (x_1 x_3^3 + x_2 (x_3^2 + x_1 x_4 + x_1 x_3 x_5)),$$

$$F = x_2 x_3^2 (x_0 + x_1 x_5) + c x_0 (x_0 x_2 x_3 + x_1 (x_3^3 + x_2 x_4 + x_2 x_3 x_5)),$$

$$G = c^3 x_0^6 x_2^3 x_3^3 + 3c^2 x_0^5 x_2^3 x_3^4 + 3c^2 x_0^5 x_1 x_2^2 x_3^5 + 3c x_0^4 x_2^3 x_3^5$$

$$+ 6c x_0^4 x_1 x_2^2 x_3^6 + x_0^3 x_2^3 x_3^6 + 3c x_0^4 x_1^2 x_2 x_3^7 + 3x_0^3 x_1 x_2^2 x_3^7$$

$$+ 3x_0^3 x_1^2 x_2 x_3^8 + x_0^3 x_1^3 x_3^9 + 3c^3 x_0^5 x_1 x_2^3 x_3^2 x_4 + 6c^2 x_0^4 x_1 x_2^3 x_3^3 x_4$$

$$+ 3c x_0^4 x_1^2 x_2^2 x_3^4 x_4 + 3c^3 x_0^4 x_1^2 x_2^2 x_3^4 x_4 + 3c x_0^3 x_1 x_2^3 x_3^4 x_4$$

$$+ 3x_0^3 x_1^2 x_2^2 x_3^5 x_4 + 3c^2 x_0^3 x_1^2 x_2^2 x_3^5 x_4 + 2x_0^3 x_1^3 x_2 x_3^6 x_4$$

$$+ c^3 x_0^3 x_1^3 x_2 x_3^6 x_4 + 3c^3 x_0^4 x_1^2 x_2^3 x_3 x_4^2 + 3c^2 x_0^3 x_1^2 x_2^3 x_3^2 x_4^2$$

$$+ x_0^3 x_1^3 x_2^2 x_3^3 x_4^2 + 2c^3 x_0^3 x_1^3 x_2^2 x_3^3 x_4^2 + c^3 x_0^3 x_1^3 x_2^3 x_4^3$$

$$\begin{aligned}
 &+ 3c^3 x_0^5 x_1 x_2^3 x_3^3 x_5 + 9c^2 x_0^4 x_1 x_2^3 x_3^4 x_5 + 6c^2 x_0^4 x_1^2 x_2^2 x_3^5 x_5 \\
 &+ 9c x_0^3 x_1 x_2^3 x_3^5 x_5 + 12c x_0^3 x_1^2 x_2^2 x_3^6 x_5 + 3x_0^2 x_1 x_2^3 x_3^6 x_5 \\
 &+ 3c x_0^3 x_1^3 x_2 x_3^7 x_5 + 6x_0^2 x_1^2 x_2^2 x_3^7 x_5 + 3x_0^2 x_1^3 x_2 x_3^8 x_5 \\
 &+ 6c^3 x_0^4 x_1^2 x_2^3 x_3^2 x_4 x_5 + 12c^2 x_0^3 x_1^2 x_2^3 x_3^3 x_4 x_5 + 3c x_0^3 x_1^3 x_2^2 x_3^4 x_4 x_5 \\
 &+ 3c^3 x_0^3 x_1^3 x_2^2 x_3^4 x_4 x_5 + 6c x_0^2 x_1^2 x_2^3 x_3^4 x_4 x_5 + 3x_0^2 x_1^3 x_2^2 x_3^5 x_4 x_5 \\
 &+ 3c^2 x_0^2 x_1^3 x_2^2 x_3^5 x_4 x_5 + 3c^3 x_0^3 x_1^3 x_2^3 x_3 x_4^2 x_5 + 3c^2 x_0^2 x_1^3 x_2^3 x_3^2 x_4^2 x_5 \\
 &+ 3c^3 x_0^4 x_1^2 x_2^3 x_3^3 x_5^2 + 9c^2 x_0^3 x_1^2 x_2^3 x_3^4 x_5^2 + 3c^2 x_0^3 x_1^3 x_2^2 x_3^5 x_5^2 \\
 &+ 9c x_0^2 x_1^2 x_2^3 x_3^5 x_5^2 + 6c x_0^2 x_1^3 x_2^2 x_3^6 x_5^2 + 3x_0 x_1^2 x_2^3 x_3^6 x_5^2 \\
 &+ 3x_0 x_1^3 x_2^2 x_3^7 x_5^2 + 3c^3 x_0^3 x_1^3 x_2^3 x_3^2 x_4 x_5^2 + 6c^2 x_0^2 x_1^3 x_2^3 x_3^3 x_4 x_5^2 \\
 &+ 3c x_0 x_1^3 x_2^3 x_3^4 x_4 x_5^2 + c^3 x_0^3 x_1^3 x_2^3 x_3^3 x_5^3 + 3c^2 x_0^2 x_1^3 x_2^3 x_3^4 x_5^3 \\
 &+ 3c x_0 x_1^3 x_2^3 x_3^5 x_5^3 + x_1^3 x_2^3 x_3^6 x_5^3,
 \end{aligned}$$

$$H = c x_0^2 x_2 x_3 + x_0 x_2 x_3^2 + x_0 x_1 x_3^3 + x_0 x_1 x_2 x_4 + c x_0 x_1 x_2 x_3 x_5 + x_1 x_2 x_3^2 x_5.$$

6. Ultra-discretization

We denote the positive structure on χ as in the previous section by $\theta : T' := (\mathbb{C}^\times)^6 \rightarrow \mathcal{V}_1 (x \mapsto V_1(x))$. Then by [Corollary 2.7](#) we obtain the ultra-discretization $\mathcal{U}\mathcal{D}(\chi, T', \theta)$, which is a Kashiwara's crystal. Now we show that the conjecture in [[Igarashi and Nakashima 2010](#)] is correct.

Theorem 6.1. *The crystal $\mathcal{U}\mathcal{D}(\chi, T', \theta)$ as above is isomorphic to the crystal B_∞ of type $G_2^{(1)}$ as in [Section 4](#).*

To show this, we display the explicit crystal structure on $\mathcal{X} := \mathcal{U}\mathcal{D}(\chi, T', \theta)$. Note that $\mathcal{U}\mathcal{D}(\chi) = \mathbb{Z}^6$ as a set. Here as for variables in \mathcal{X} , we use the same notations c, x_0, x_1, \dots, x_5 as for χ .

For $x = (x_0, x_1, \dots, x_5) \in \mathcal{X}$, it follows from the results in the previous section that the functions wt_i and ε_i ($i = 0, 1, 2$) are given as

$$\begin{aligned}
 \text{wt}_0(x) &= 2x_0 - x_1 - x_3 - x_5, & \text{wt}_1(x) &= 2(x_1 + x_3 + x_5) - x_0 - x_2 - x_4, \\
 \text{wt}_2(x) &= 2(x_2 + x_4) - 3(x_1 - x_3 - x_5).
 \end{aligned}$$

Set

$$\begin{aligned}
 (6-1) \quad \alpha &:= 2x_0 + x_2 + x_3, & \beta &:= x_1 + x_2 + 2x_3 + x_5, & \gamma &:= x_0 + x_1 + 3x_3, \\
 \delta &:= x_0 + x_2 + 2x_3, & \epsilon &:= x_0 + x_1 + x_2 + x_4, & \phi &:= x_0 + x_1 + x_2 + x_3 + x_5.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 (6-2) \quad \varepsilon_0(x) &= \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi) - (3x_0 + x_2 + x_3), \\
 \varepsilon_1(x) &= \max(x_0 - x_1, x_0 + x_2 - 2x_1 - x_3, x_0 + x_2 + x_4 - 2x_1 - 2x_3 - x_5), \\
 \varepsilon_2(x) &= \max(3x_1 - x_2, 3x_1 + 3x_3 - 2x_2 - x_4).
 \end{aligned}$$

Indeed, from the explicit form of G in the previous section we have $\mathcal{U}\mathcal{D}(G)|_{c=-1} = \max(-3+3\alpha, -2+2\alpha+\delta, -2+2\alpha+\gamma, -1+\alpha+2\delta, -1+\alpha+\gamma+\delta, 3\delta, -1+\alpha+2\gamma, \gamma+2\delta, 2\gamma+\delta, 3\gamma, -3+2\alpha+\epsilon, -2+\alpha+\delta+\epsilon, -1+\alpha+\gamma+\epsilon, -1+2\delta+\epsilon, \gamma+\delta+\epsilon, 2\gamma+\epsilon, -3+\alpha+2\epsilon, -2+\delta+2\epsilon, \gamma+2\epsilon, -3+3\epsilon, -3+2\alpha+\phi, -2+\alpha+\delta+\phi, -2+\alpha+\gamma+\phi, -1+2\delta+\phi, -1+\gamma+\delta+\phi, \beta+2\delta, -1+2\gamma+\phi, \beta+\gamma+\delta, \beta+2\gamma, -3+\alpha+\epsilon+\phi, -2+\delta+\epsilon+\phi, -1+\gamma+\epsilon+\phi, -1+\beta+\delta+\epsilon, \beta+\gamma+\epsilon, -3+2\epsilon+\phi, -2+\beta+2\epsilon, -3+\alpha+2\phi, -2+\delta+2\phi, -2+\gamma+2\phi, -1+\alpha+2\beta, -1+\beta+\gamma+\phi, 2\beta+\delta, 2\beta+\gamma, -3+\epsilon+2\phi, -2+\beta+\epsilon+\phi, -1+2\beta+\epsilon, -3+3\phi, -2+\beta+2\phi, -1+2\beta+\phi, 3\beta)$.

We simplify this by using the following lemma:

Lemma 6.2. *For $m_1, \dots, m_k \in \mathbb{R}$ and $t_1, \dots, t_k \in \mathbb{R}_{\geq 0}$ such that $t_1 + \dots + t_k = 1$, we have*

$$\max\left(m_1, \dots, m_k, \sum_{i=1}^k t_i m_i\right) = \max(m_1, \dots, m_k).$$

Since we have

$$\begin{aligned} -2+2\alpha+\delta &= \frac{2(-3+3\alpha)+3\delta}{3}, & -2+2\alpha+\gamma &= \frac{2(-3+3\alpha)+3\gamma}{3}, \\ -1+\alpha+2\delta &= \frac{2 \cdot 3\delta + (-3+3\alpha)}{3}, & -1+\alpha+\gamma+\delta &= \frac{(-3+3\alpha)+3\gamma+3\delta}{3}, \\ -1+\alpha+2\gamma &= \frac{(-3+3\alpha)+2 \cdot 3\gamma}{3}, & \gamma+2\delta &= \frac{2 \cdot 3\delta + 3\gamma}{3}, \end{aligned}$$

and so on, we deduce using the lemma that

$$\mathcal{U}\mathcal{D}(G)|_{c=-1} = \max(-3+3\alpha, 3\beta, 3\gamma, 3\delta, -3+3\epsilon, -3+3\phi, -1+\alpha+\gamma+\epsilon, \gamma+\delta+\epsilon, \gamma+2\epsilon, 2\gamma+\epsilon, -1+\gamma+\epsilon+\phi, \beta+\gamma+\epsilon).$$

Next, we describe the actions of \tilde{f}_i ($i = 0, 1, 2$). Set $\mathfrak{E}_j := \mathcal{U}\mathcal{D}(\mathcal{C}_j)|_{c=-1}$, for $j = 1, \dots, 5$. Then we have

$$\begin{aligned} \mathfrak{E}_1 &= \max(-1+x_0-x_1, x_0+x_2-2x_1-x_3, x_0+x_2+x_4-2x_1-2x_3-x_5) \\ &\quad - \max(x_0-x_1, x_0+x_2-2x_1-x_3, x_0+x_2+x_4-2x_1-2x_3-x_5), \\ \mathfrak{E}_3 &= \max(-1+x_0-x_1, -1+x_0+x_2-2x_1-x_3, x_0+x_2+x_4-2x_1-2x_3-x_5) \\ &\quad - \max(-1+x_0-x_1, x_0+x_2-2x_1-x_3, x_0+x_2+x_4-2x_1-2x_3-x_5), \\ \mathfrak{E}_5 &= \max(-1+x_0-x_1, -1+x_0+x_2-2x_1-x_3, -1+x_0+x_2+x_4-2x_1-2x_3-x_5) \\ &\quad - \max(-1+x_0-x_1, -1+x_0+x_2-2x_1-x_3, x_0+x_2+x_4-2x_1-2x_3-x_5), \\ \mathfrak{E}_2 &= \max(-1+3x_1-x_2, 3x_1+3x_3-2x_2-x_4) \\ &\quad - \max(3x_1-x_2, 3x_1+3x_3-2x_2-x_4), \\ \mathfrak{E}_4 &= \max(-1+3x_1-x_2, -1+3x_1+3x_3-2x_2-x_4) \\ &\quad - \max(-1+3x_1-x_2, 3x_1+3x_3-2x_2-x_4). \end{aligned}$$

Therefore, for $x \in \mathcal{X}$ we have

$$\begin{aligned}\tilde{f}_1(x) &= (x_0, x_1 + \Xi_1, x_2, x_3 + \Xi_3, x_4, x_5 + \Xi_5), \\ \tilde{f}_2(x) &= (x_0, x_1, x_2 + \Xi_2, x_3, x_4 + \Xi_4, x_5).\end{aligned}$$

We obtain the action \tilde{e}_i ($i = 1, 2$) by setting $c = 1$ in ${}^{\mathcal{O}}\mathcal{U}\mathcal{D}(\mathcal{C}_i)$.

Finally, we describe the action of \tilde{f}_0 . Set

$$\begin{aligned}\Psi_0 &:= \max(-2 + \alpha, \beta, -1 + \gamma, -1 + \delta, -1 + \epsilon, -1 + \phi) - \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi) + 1, \\ \Psi_1 &:= \max(-1 + \alpha, \beta, -1 + \gamma, \delta, -1 + \epsilon, -1 + \phi) - \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi) + 1, \\ \Psi_2 &:= \max(-3 + 3\alpha, 3\beta, 3\gamma, 3\delta, -3 + 3\epsilon, -3 + 3\phi, -1 + \alpha + \gamma + \epsilon, \gamma + \delta + \epsilon, \\ &\quad \gamma + 2\epsilon, 2\gamma + \epsilon, -1 + \gamma + \epsilon + \phi, \beta + \gamma + \epsilon) - 3 \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi) + 3, \\ \Psi_3 &:= \max(-2 + \alpha, \beta, -1 + \gamma, -1 + \delta, -1 + \epsilon, -1 + \phi) \\ &\quad + \max(-1 + \alpha, \beta, \gamma, \delta, \epsilon, -1 + \phi) - \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi) \\ &\quad - \max(-1 + \alpha, \beta, -1 + \gamma, \delta, -1 + \epsilon, -1 + \phi) + 2, \\ \Psi_4 &:= 3 \max(-2 + \alpha, \beta, -1 + \gamma, -1 + \delta, -1 + \epsilon, -1 + \phi) \\ &\quad - \max(-3 + 3\alpha, 3\beta, 3\gamma, 3\delta, -3 + 3\epsilon, -3 + 3\phi, -1 + \alpha + \gamma + \epsilon, \gamma + \delta + \epsilon, \\ &\quad \gamma + 2\epsilon, 2\gamma + \epsilon, -1 + \gamma + \epsilon + \phi, \beta + \gamma + \epsilon) + 3, \\ \Psi_5 &:= \max(-2 + \alpha, \beta, -1 + \gamma, -1 + \delta, -1 + \epsilon, -1 + \phi) \\ &\quad - \max(1 + \alpha, \beta, \gamma, \delta, \epsilon, -1 + \phi) + 1,\end{aligned}$$

where $\alpha, \beta, \dots, \phi$ are as in (6-1). Therefore, by the explicit form of e_0^c as in the previous section, we have

$$(6-3) \quad \tilde{f}_0(x) = (x_0 + \Psi_0, x_1 + \Psi_1, x_2 + \Psi_2, x_3 + \Psi_3, x_4 + \Psi_4, x_5 + \Psi_5).$$

We have the explicit form of \tilde{e}_0 by setting $c = 1$ in ${}^{\mathcal{O}}\mathcal{U}\mathcal{D}(\mathcal{C}_i)$.

Proof of Theorem 6.1. Define the map

$$\begin{aligned}\Omega : \quad \mathcal{X} &\quad \rightarrow \quad B_\infty \\ (x_0, \dots, x_5) &\mapsto (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1),\end{aligned}$$

by

$$b_1 = x_5, \quad b_2 = \frac{1}{3}x_4 - x_5, \quad b_3 = x_3 - \frac{2}{3}x_4, \quad \bar{b}_3 = \frac{2}{3}x_2 - x_3, \quad \bar{b}_2 = x_1 - \frac{1}{3}x_2, \quad \bar{b}_1 = x_0 - x_1,$$

and Ω^{-1} is given by

$$\begin{aligned}x_0 &= b_1 + b_2 + \frac{1}{2}(b_3 + \bar{b}_3) + \bar{b}_2 + \bar{b}_1, & x_1 &= b_1 + b_2 + \frac{1}{2}(b_3 + \bar{b}_3) + \bar{b}_2, \\ x_2 &= 3b_1 + 3b_2 + \frac{3}{2}(b_3 + \bar{b}_3), & x_3 &= 2b_1 + 2b_2 + b_3, & x_4 &= 3b_1 + 3b_2, & x_5 &= b_1,\end{aligned}$$

which means that Ω is bijective. Note that $\frac{3}{2}(b_3 + \bar{b}_3) \in \mathbb{Z}$ by the definition of B_∞ as on page 127. We shall show that Ω is commutative with actions of \tilde{f}_i and

preserves the functions wt_i and ε_i , that is,

$$\tilde{f}_i(\Omega(x)) = \Omega(\tilde{f}_i x), \quad \text{wt}_i(\Omega(x)) = \text{wt}_i(x), \quad \varepsilon_i(\Omega(x)) = \varepsilon_i(x) \quad (i = 0, 1, 2),$$

Indeed, the commutativity $\tilde{e}_i(\Omega(x)) = \Omega(\tilde{e}_i x)$ is shown by a similar way. First, let us check wt_i :

Set $b = \Omega(x)$ and let (z_1, z_2, z_3, z_4) be as in (4-2). By the explicit forms of wt_i on \mathcal{X} and B_∞ , we have

$$\begin{aligned} \text{wt}_0(\Omega(x)) &= \varphi_0(\Omega(x)) - \varepsilon_0(\Omega(x)) = 2z_1 + z_2 + z_3 + 3z_4 \\ &= 2(\bar{b}_1 - b_1) + (\bar{b}_2 - \bar{b}_3) + (b_3 - b_2) + \frac{3}{2}(\bar{b}_3 - b_3) \\ &= 2(\bar{b}_1 - b_1) + \bar{b}_2 - b_2 + \frac{1}{2}(\bar{b}_3 - b_3) = 2x_0 - x_1 - x_3 - x_5 = \text{wt}_0(x), \end{aligned}$$

$$\begin{aligned} \text{wt}_1(\Omega(x)) &= \varphi_1(\Omega(x)) - \varepsilon_1(\Omega(x)) \\ &= b_1 + (b_3 - b_2 + (\bar{b}_2 - \bar{b}_3)_+)_+ - (\bar{b}_1 + (\bar{b}_3 - \bar{b}_2 + (b_2 - b_3)_+)_+) \\ &= b_1 - \bar{b}_1 - b_2 + \bar{b}_2 + b_3 - \bar{b}_3 = 2(x_1 + x_3 + x_5) - x_0 - x_2 - x_4 \\ &= \text{wt}_1(x), \end{aligned}$$

$$\begin{aligned} \text{wt}_2(\Omega(x)) &= \varphi_2(\Omega(x)) - \varepsilon_2(\Omega(x)) \\ &= 3b_2 + \frac{3}{2}(\bar{b}_3 - b_3)_+ - 3\bar{b}_2 - \frac{3}{2}(b_3 - \bar{b}_3)_+ \\ &= 3b_2 - 3\bar{b}_2 + \frac{3}{2}(\bar{b}_3 - b_3) = 2(x_2 + x_4) - 3(x_1 + x_3 + x_5) = \text{wt}_2(x). \end{aligned}$$

Next, we check ε_i :

$$\begin{aligned} \varepsilon_1(\Omega(x)) &= \bar{b}_1 + (\bar{b}_3 - \bar{b}_2 + (b_2 - b_3)_+)_+ \\ &= \max(\bar{b}_1, \bar{b}_1 + \bar{b}_3 - \bar{b}_2, \bar{b}_1 + \bar{b}_3 - \bar{b}_2 + b_2 - b_3) \\ &= \max(x_0 - x_1, x_0 - 2x_1 + x_2 - x_3, x_0 - 2x_1 + x_2 - 2x_3 + x_4 - x_5) = \varepsilon_1(x), \\ \varepsilon_2(\Omega(x)) &= 3\bar{b}_2 + \frac{3}{2}(b_3 - \bar{b}_3)_+ = \max(3\bar{b}_2, 3\bar{b}_2 + \frac{3}{2}(b_3 - \bar{b}_3)) \\ &= \max(3x_1 - x_2, 3x_1 - 2x_2 + 3x_3 - x_4) = \varepsilon_2(x). \end{aligned}$$

Now let us see ε_0 :

$$\begin{aligned} \varepsilon_0(\Omega(x)) &= -s(b) + \max A - (2z_1 + z_2 + z_3 + 3z_4) \\ &= -x_0 + \max(0, z_1, z_1 + z_2, z_1 + z_2 + 3z_4, \\ &\quad z_1 + z_2 + z_3 + 3z_4, 2z_1 + z_2 + z_3 + 3z_4) - (\alpha - \beta) \\ &= -x_0 + \max(-2x_0 + x_1 + x_3 + x_5, -x_0 + x_3, -x_0 + x_1 - x_2 + 2x_3, \\ &\quad -x_0 + x_1 - x_3 + x_4, -x_0 + x_1 + x_5, 0) \\ &= -(3x_0 + x_2 + x_3) + \max(x_1 + x_2 + 2x_3 + x_5, x_0 + x_2 + 2x_3, x_0 + x_1 + 3x_3, \\ &\quad x_0 + x_1 + x_2 + x_4, x_0 + x_1 + x_2 + x_3 + x_5, 2x_0 + x_2 + x_3) \\ &= -(3x_0 + x_2 + x_3) + \max(\beta, \delta, \gamma, \epsilon, \phi, \alpha). \end{aligned}$$

On the other hand, we have

$$\varepsilon_0(x) = -(3x_0 + x_2 + x_3) + \max(\alpha, \beta, \gamma, \delta, \epsilon, \phi).$$

which shows $\varepsilon_0(\Omega(x)) = \varepsilon_0(x)$.

Let us show $\tilde{f}_i(\Omega(x)) = \Omega(\tilde{f}_i(x))$ ($x \in \mathcal{X}$, $i = 0, 1, 2$). As for \tilde{f}_1 , set

$$A = x_0 - x_1, \quad B = x_0 + x_2 - 2x_1 - x_3, \quad C = x_0 + x_2 + x_4 - 2x_1 - 2x_3 - x_5.$$

Then we obtain

$$\Xi_1 = \max(A - 1, B, C) - \max(A, B, C),$$

$$\Xi_3 = \max(A - 1, B - 1, C) - \max(A - 1, B, C),$$

$$\Xi_5 = \max(A - 1, B - 1, C - 1) - \max(A - 1, B - 1, C).$$

Therefore, we have

$$\begin{aligned} \Xi_1 = -1, \quad \Xi_3 = 0, \quad \Xi_5 = 0, & \quad \text{if } A > B, C, \\ \Xi_1 = 0, \quad \Xi_3 = -1, \quad \Xi_5 = 0, & \quad \text{if } A \leq B > C, \\ \Xi_1 = 0, \quad \Xi_3 = 0, \quad \Xi_5 = -1, & \quad \text{if } A, B \leq C, \end{aligned}$$

which implies

$$\tilde{f}_1(x) = \begin{cases} (x_0, x_1 - 1, x_2, \dots, x_5) & \text{if } A > B, C, \\ (x_0, \dots, x_3 - 1, x_4, x_5) & \text{if } A \leq B > C, \\ (x_0, \dots, x_4, x_5 - 1) & \text{if } A, B \leq C. \end{cases}$$

Since $A = \bar{b}_1$, $B = \bar{b}_1 + \bar{b}_3 - \bar{b}_2$ and $C = \bar{b}_1 + \bar{b}_3 - \bar{b}_2 + b_2 - b_3$, we get ($b = \Omega(x)$)

$$\Omega(\tilde{f}_1(x)) = \begin{cases} (\dots, \bar{b}_2 - 1, \bar{b}_1 + 1) & \text{if } \bar{b}_2 - \bar{b}_3 > (b_2 - b_3)_+, \\ (\dots, b_3 - 1, \bar{b}_3 + 1, \dots) & \text{if } \bar{b}_2 - \bar{b}_3 \leq 0 < b_3 - b_2, \\ (b_1 - 1, b_2 + 1, \dots) & \text{if } (\bar{b}_2 - \bar{b}_3)_+ \leq b_2 - b_3, \end{cases}$$

which is the same as the action of \tilde{f}_1 on $b = \Omega(x)$ as on page 128. Hence, we have $\Omega(\tilde{f}_1(x)) = \tilde{f}_1(\Omega(x))$.

Let us see $\Omega(\tilde{f}_2(x)) = \tilde{f}_2(\Omega(x))$. Set

$$L = 3x_1 - x_2, \quad M = 3x_1 + 3x_3 - 2x_2 - x_4.$$

Then $\Xi_2 = \max(-1 + L, M) - \max(L, M)$ and $\Xi_4 = \max(-1 + L, -1 + M) - \max(-1 + L, M)$. Thus, one has

$$\begin{aligned} \Xi_2 = -1, \quad \Xi_4 = 0 & \quad \text{if } L > M, \\ \Xi_2 = 0, \quad \Xi_4 = -1 & \quad \text{if } L \leq M, \end{aligned}$$

which means

$$\tilde{f}_2(x) = \begin{cases} (x_0, x_1, x_2 - 1, x_3, x_4, x_5) & \text{if } L > M, \\ (x_0, x_1, x_2, x_3, x_4 - 1, x_5) & \text{if } L \leq M. \end{cases}$$

Since $L - M = x_2 - 3x_3 + x_4 = \frac{3}{2}(\bar{b}_3 - b_3)$, one gets

$$\Omega(\tilde{f}_2(x)) = \begin{cases} (\dots, \bar{b}_3 - \frac{2}{3}, \bar{b}_2 + \frac{1}{3}, \dots) & \text{if } \bar{b}_3 > b_3, \\ (\dots, b_2 - \frac{1}{3}, b_3 + \frac{2}{3}, \dots) & \text{if } \bar{b}_3 \leq b_3, \end{cases}$$

where $b = \Omega(x)$. This action coincides with the one of \tilde{f}_2 on $b \in B_\infty$ on page 128. Therefore, we get $\Omega(\tilde{f}_2(x)) = \tilde{f}_2(\Omega(x))$.

Finally, we shall check $\tilde{f}_0(\Omega(x)) = \Omega(\tilde{f}_0(x))$. For the purpose, we shall estimate the values Ψ_0, \dots, Ψ_5 explicitly.

First, the following cases are investigated:

$$(f_1) \quad \beta \geq \gamma, \delta, \epsilon, \phi, \quad \phi \geq \alpha, \delta \geq \alpha.$$

$$(f_2) \quad \beta < \delta \geq \alpha, \gamma, \epsilon, \quad \alpha > \phi, \beta \geq \phi.$$

$$(f_3) \quad \beta, \delta < \gamma \geq \alpha, \epsilon, \phi.$$

$$(f_4) \quad \beta, \delta < \epsilon \geq \alpha, \phi, \quad \epsilon = \gamma + 1.$$

$$(f'_4) \quad \beta, \delta < \epsilon \geq \alpha, \phi, \quad \epsilon = \gamma + 2.$$

$$(f''_4) \quad \beta, \delta < \epsilon \geq \alpha, \phi, \quad \epsilon > \gamma + 2.$$

$$(f_5) \quad \beta, \gamma, \epsilon < \phi \geq \alpha, \quad \alpha > \delta, \beta \geq \delta.$$

$$(f_6) \quad \alpha > \gamma, \delta, \epsilon, \phi, \quad \delta, \phi > \beta.$$

It is easy to see that each of these conditions are equivalent to the conditions (F_1) – (F_6) in (4-3); more precisely, we have $(f_i) \iff (F_i)$ ($i = 1, 2, 3, 5, 6$), $(f_4) \iff (F_4)$ and $z_4 = \frac{1}{3}$, $(f'_4) \iff (F_4)$ and $z_4 = \frac{2}{3}$ and $(f''_4) \iff (F_4)$ and $z_4 \neq \frac{1}{3}, \frac{2}{3}$, and that (f_1) – (f_6) cover all cases and they have no intersection.

Let us show $(f_1) \iff (F_1)$: the condition (f_1) means $\beta - \gamma = -(z_1 + z_2) \geq 0$, $\beta - \delta = -z_1 \geq 0$, $\beta - \epsilon = -(z_1 + z_2 + 3z_4) \geq 0$ and $\beta - \phi = -(z_1 + z_2 + z_3 + 3z_4) \geq 0$, which is equivalent to the condition $z_1 + z_2 \leq 0$, $z_1 \leq 0$, $z_1 + z_2 + 3z_4 \leq 0$ and $z_1 + z_2 + z_3 + 3z_4 \leq 0$. (Note that $\phi - \alpha = \beta - \delta$, $\delta - \alpha = \beta - \phi$.) This is just the condition (F_1) . Other cases $i = 2, 3, 5, 6$ are shown similarly. Next, let us see the cases (f_4) , (f'_4) and (f''_4) . Indeed,

$$\epsilon - \gamma = x_2 - 3x_3 + x_4 = \frac{3}{2}(\bar{b}_3 - b_3) = 3z_4.$$

Thus, we can easily get that $(f_4) \iff (F_4)$ and $z_4 = \frac{1}{3}$, $(f'_4) \iff (F_4)$ and $z_4 = \frac{2}{3}$ and $(f''_4) \iff (F_4)$ and $z_4 \neq \frac{1}{3}, \frac{2}{3}$.

Under the condition $(f_1) \iff (F_1)$, we have

$$\Psi_0 = \Psi_1 = \Psi_5 = 1, \Psi_2 = \Psi_4 = 3, \quad \Psi_3 = 2,$$

which means $\tilde{f}_0(x) = (x_0 + 1, x_1 + 1, x_2 + 3, x_3 + 2, x_4 + 3, x_5 + 1)$. Thus, we have

$$\Omega(\tilde{f}_0(x)) = (b_1 + 1, b_2, \dots, \bar{b}_1),$$

which coincides with the action of \tilde{f}_0 under (F_1) given on page 128. Similarly, we have

$$\begin{aligned} (f_2) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 1, 3, 1, 0, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1 + 1, x_2 + 3, x_3 + 1, x_4, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2, b_3 + 1, \bar{b}_3 + 1, \bar{b}_2, \bar{b}_1 - 1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_2) on the same page;

$$\begin{aligned} (f_3) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 0, 3, 2, 0, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1, x_2 + 3, x_3 + 2, x_4, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2, b_3 + 2, \bar{b}_3, \bar{b}_2 - 1, \bar{b}_1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_3) ;

$$\begin{aligned} (f_4) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 0, 2, 2, 1, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1, x_2 + 2, x_3 + 2, x_4 + 1, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2 + \frac{1}{3}, b_3 + \frac{4}{3}, \bar{b}_3 - \frac{2}{3}, \bar{b}_2 - \frac{2}{3}, \bar{b}_1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_4) and $z_4 = \frac{1}{3}$;

$$\begin{aligned} (f'_4) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 0, 1, 2, 2, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1, x_2 + 1, x_3 + 2, x_4 + 2, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2 + \frac{2}{3}, b_3 + \frac{2}{3}, \bar{b}_3 - \frac{4}{3}, \bar{b}_2 - \frac{1}{3}, \bar{b}_1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_4) and $z_4 = \frac{2}{3}$;

$$\begin{aligned} (f''_4) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 0, 0, 2, 3, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1, x_2, x_3 + 2, x_4 + 3, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2 + 1, b_3, \bar{b}_3 - 2, \bar{b}_2, \bar{b}_1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_4) and $z_4 \neq \frac{1}{3}, \frac{2}{3}$;

$$\begin{aligned} (f_5) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (0, 0, 0, 1, 3, 1) \\ &\Rightarrow \tilde{f}_0(x) = (x_0, x_1, x_2, x_3 + 1, x_4 + 3, x_5 + 1) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1 + 1, b_2, b_3 - 1, \bar{b}_3 - 1, \bar{b}_2, \bar{b}_1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_5) . Finally,

$$\begin{aligned} (f_6) &\Rightarrow (\Psi_0, \Psi_1, \Psi_2, \Psi_3, \Psi_4, \Psi_5) = (-1, 0, 0, 0, 0, 0) \\ &\Rightarrow \tilde{f}_0(x) = (x_0 - 1, x_1, x_2, x_3, x_4, x_5) \\ &\Rightarrow \Omega(\tilde{f}_0(x)) = (b_1, b_2, b_3, \bar{b}_3, \bar{b}_2, \bar{b}_1 - 1), \end{aligned}$$

which coincides with the action of \tilde{f}_0 under (F_6) , still on page 128. Now, we have $\Omega(\tilde{f}_0(x)) = \tilde{f}_0(\Omega(x))$. The proof of [Theorem 6.1](#) has been completed. \square

References

- [Berenstein and Kazhdan 2000] A. Berenstein and D. Kazhdan, “Geometric and unipotent crystals”, pp. 188–236 in *GAF A 2000* (Special volume of *Geom. Funct. Anal.*) (Tel Aviv, 1999), vol. I, edited by N. Alon et al., 2000. [MR 2003b:17013](#) [Zbl 1044.17006](#)
- [Fourier et al. 2009] G. Fourier, M. Okado, and A. Schilling, “Kirillov–Reshetikhin crystals for nonexceptional types”, *Adv. Math.* **222**:3 (2009), 1080–1116. [MR 2010j:17028](#) [Zbl 05609507](#)
- [Fourier et al. 2010] G. Fourier, M. Okado, and A. Schilling, “Perfectness of Kirillov–Reshetikhin crystals for nonexceptional types”, pp. 127–143 in *Quantum affine algebras, extended affine Lie algebras, and their applications*, edited by Y. Gao et al., *Contemp. Math.* **506**, Amer. Math. Soc., Providence, RI, 2010. [MR 2011b:17031](#) [Zbl 05901949](#)
- [Hatayama et al. 1999] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Y. Yamada, “Remarks on fermionic formula”, pp. 243–291 in *Recent developments in quantum affine algebras and related topics* (Raleigh, NC, 1998), edited by N. Jing and K. C. Misra, *Contemp. Math.* **248**, Amer. Math. Soc., Providence, RI, 1999. [MR 2001m:81129](#) [Zbl 1032.81015](#)
- [Hatayama et al. 2002] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, and Z. Tsuboi, “Paths, crystals and fermionic formulae”, pp. 205–272 in *MathPhys odyssey, 2001*, edited by M. Kashiwara and T. Miwa, *Prog. Math. Phys.* **23**, Birkhäuser, Boston, 2002. [MR 2003e:17020](#) [Zbl 1016.17011](#)
- [Igarashi and Nakashima 2010] M. Igarashi and T. Nakashima, “Affine geometric crystal of type $D_4^{(3)}$ ”, pp. 215–226 in *Quantum affine algebras, extended affine Lie algebras, and their applications*, edited by Y. Gao et al., *Contemp. Math.* **506**, Amer. Math. Soc., Providence, RI, 2010. [MR 2011h:17021](#) [Zbl 05901953](#)
- [Kac 1990] V. G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990. [MR 92k:17038](#) [Zbl 0716.17022](#)
- [Kac and Peterson 1983] V. G. Kac and D. H. Peterson, “Regular functions on certain infinite-dimensional groups”, pp. 141–166 in *Arithmetic and geometry*, vol. II, edited by M. Artin and J. Tate, *Progr. Math.* **36**, Birkhäuser, Boston, 1983. [MR 86b:17010](#) [Zbl 0578.17014](#)
- [Kang et al. 1992a] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, “Affine crystals and vertex models”, pp. 449–484 in *Infinite analysis* (Kyoto, 1991), edited by A. Tsuchiya et al., *Adv. Ser. Math. Phys.* **16**, World Sci. Publ., River Edge, NJ, 1992. [MR 94a:17008](#) [Zbl 0925.17005](#)
- [Kang et al. 1992b] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, “Perfect crystals of quantum affine Lie algebras”, *Duke Math. J.* **68**:3 (1992), 499–607. [MR 94j:17013](#) [Zbl 0774.17017](#)
- [Kang et al. 1994] S.-J. Kang, M. Kashiwara, and K. C. Misra, “Crystal bases of Verma modules for quantum affine Lie algebras”, *Compositio Math.* **92**:3 (1994), 299–325. [MR 95h:17016](#) [Zbl 0808.17007](#)

- [Kashiwara 1990] M. Kashiwara, “Crystalizing the q -analogue of universal enveloping algebras”, *Comm. Math. Phys.* **133**:2 (1990), 249–260. MR 92b:17018 Zbl 0724.17009
- [Kashiwara 1991] M. Kashiwara, “On crystal bases of the Q -analogue of universal enveloping algebras”, *Duke Math. J.* **63**:2 (1991), 465–516. MR 93b:17045 Zbl 0739.17005
- [Kashiwara 2002] M. Kashiwara, “On level-zero representations of quantized affine algebras”, *Duke Math. J.* **112**:1 (2002), 117–175. MR 2002m:17013
- [Kashiwara 2005] M. Kashiwara, “Level zero fundamental representations over quantized affine algebras and Demazure modules”, *Publ. Res. Inst. Math. Sci.* **41**:1 (2005), 223–250. MR 2005i:17021 Zbl 1147.17306
- [Kashiwara et al. 2007] M. Kashiwara, K. C. Misra, M. Okado, and D. Yamada, “Perfect crystals for $U_q(D_4^{(3)})$ ”, *J. Algebra* **317**:1 (2007), 392–423. MR 2009b:17035 Zbl 1140.17012
- [Kashiwara et al. 2008] M. Kashiwara, T. Nakashima, and M. Okado, “Affine geometric crystals and limit of perfect crystals”, *Trans. Amer. Math. Soc.* **360**:7 (2008), 3645–3686. MR 2009e:17020 Zbl 1219.17010
- [Kirillov and Reshetikhin 1987] A. N. Kirillov and N. Y. Reshetikhin, “Representations of Yangians and multiplicity of occurrence of the irreducible components of the tensor product of representations of simple Lie algebras”, pp. 211–221 in *Anal. Teor. Chisel i Teor. Funktsii.*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **8**, 1987. In Russian; translated in *J. Sov. Math.* **52** (1990), 3156–3164. MR 89b:17012
- [Kumar 2002] S. Kumar, *Kac–Moody groups, their flag varieties and representation theory*, Progress in Mathematics **204**, Birkhäuser, Boston, 2002. MR 2003k:22022 Zbl 1026.17030
- [Lusztig 1990] G. Lusztig, “Canonical bases arising from quantized enveloping algebras”, *J. Amer. Math. Soc.* **3**:2 (1990), 447–498. MR 90m:17023 Zbl 0703.17008
- [Misra et al. 2010] K. C. Misra, M. Mohamad, and M. Okado, “Zero action on perfect crystals for $U_q(G_2^{(1)})$ ”, *SIGMA* **6** (2010), Art. ID 022.
- [Nakashima 2005a] T. Nakashima, “Geometric crystals on Schubert varieties”, *J. Geom. Phys.* **53**:2 (2005), 197–225. MR 2006d:17016 Zbl 1156.17304
- [Nakashima 2005b] T. Nakashima, “Geometric crystals on unipotent groups and generalized Young tableaux”, *J. Algebra* **293**:1 (2005), 65–88. MR 2006j:20064 Zbl 1161.17319
- [Nakashima 2007] T. Nakashima, “Affine geometric crystal of type $G_2^{(1)}$ ”, pp. 179–192 in *Lie algebras, vertex operator algebras and their applications*, edited by Y.-Z. Huang and K. C. Misra, Contemp. Math. **442**, Amer. Math. Soc., Providence, RI, 2007. MR 2009e:17047 Zbl 1142.17010
- [Nakashima 2010] T. Nakashima, “Ultra-discretization of the $G_2^{(1)}$ -geometric crystals to the $D_4^{(3)}$ -perfect crystals”, pp. 273–296 in *Representation theory of algebraic groups and quantum groups*, edited by A. Gyoja et al., Progr. Math. **284**, Birkhäuser, New York, 2010. MR 2011m:17042 Zbl 05919687
- [Okado and Schilling 2008] M. Okado and A. Schilling, “Existence of Kirillov–Reshetikhin crystals for nonexceptional types”, *Represent. Theory* **12** (2008), 186–207. MR 2009c:17022 Zbl 05526467
- [Peterson and Kac 1983] D. H. Peterson and V. G. Kac, “Infinite flag varieties and conjugacy theorems”, *Proc. Nat. Acad. Sci. U.S.A.* **80**:6 i. (1983), 1778–1782. MR 84g:17017 Zbl 0512.17008
- [Yamane 1998] S. Yamane, “Perfect crystals of $U_q(G_2^{(1)})$ ”, *J. Algebra* **210**:2 (1998), 440–486. MR 2000f:17024 Zbl 0929.17013

MANA IGARASHI
DEPARTMENT OF MATHEMATICS
SOPHIA UNIVERSITY
KIOICHO 7-1, CHIYODA-KU
TOKYO 102-8554
JAPAN
mana-i@hoffman.cc.sophia.ac.jp

KAILASH C. MISRA
DEPARTMENT OF MATHEMATICS
NORTH CAROLINA STATE UNIVERSITY
2311 STINSON DRIVE
RALEIGH, NC 27695-8205
UNITED STATES
misra@math.ncsu.edu

TOSHIKI NAKASHIMA
DEPARTMENT OF MATHEMATICS
SOPHIA UNIVERSITY
KIOICHO 7-1, CHIYODA-KU
TOKYO 102-8554
JAPAN
toshiki@sophia.ac.jp

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 255 No. 1 January 2012

Averaging sequences	1
FERNANDO ALCALDE CUESTA and ANA RECHTMAN	
Affine group schemes over symmetric monoidal categories	25
ABHISHEK BANERJEE	
Eigenvalue estimates on domains in complete noncompact Riemannian manifolds	41
DAGUANG CHEN, TAO ZHENG and MIN LU	
Realizing the local Weil representation over a number field	55
GERALD CLIFF and DAVID MCNEILLY	
Lagrangian submanifolds in complex projective space with parallel second fundamental form	79
FRANKI DILLEN, HAIZHONG LI, LUC VRANCKEN and XIANFENG WANG	
Ultra-discretization of the $D_4^{(3)}$ -geometric crystal to the $G_2^{(1)}$ -perfect crystals	117
MANA IGARASHI, KAILASH C. MISRA and TOSHIKI NAKASHIMA	
Connectivity properties for actions on locally finite trees	143
KEITH JONES	
Remarks on the curvature behavior at the first singular time of the Ricci flow	155
NAM Q. LE and NATASA SESUM	
Stability of capillary surfaces with planar boundary in the absence of gravity	177
PETKO I. MARINOV	
Small hyperbolic polyhedra	191
SHAWN RAFALSKI	
Hurwitz spaces of coverings with two special fibers and monodromy group a Weyl group of type B_d	241
FRANCESCA VETRO	