

*Pacific
Journal of
Mathematics*

ON SLIM DOUBLE LIE GROUPOIDS

NICOLAS ANDRUSKIEWITSCH, JESUS OCHOA ARANGO
AND ALEJANDRO TIRABOSCHI

Volume 256 No. 1

March 2012

ON SLIM DOUBLE LIE GROUPOIDS

NICOLAS ANDRUSKIEWITSCH, JESUS OCHOA ARANGO
AND ALEJANDRO TIRABOSCHI

We prove that every slim double Lie groupoid with proper core action is completely determined by a factorization of a canonically defined “diagonal” Lie groupoid.

Introduction

A double groupoid is a set \mathcal{B} provided with two different but compatible groupoid structures. It is useful to represent the elements of \mathcal{B} as boxes that merge horizontally or vertically according to the groupoid multiplication into consideration. The vertical sides of a box belong to a groupoid \mathcal{V} , and the horizontal sides to a groupoid \mathcal{H} . A double groupoid is *slim* if any box is determined by its four sides. The notion of double groupoids was introduced by Ehresmann [1963], and later studied in [Brown 2004; Brown and Janelidze 2004; Brown and Mackenzie 1992; Brown and Spencer 1976] and references therein.

The notion of double Lie groupoid was defined and investigated by K. Mackenzie [1992; 2000]; see also [Pradines 1977; Mackenzie 1999; Lu and Weinstein 1989] for applications to differential and Poisson geometry. In particular the question of the classification of double Lie groupoids was raised in [Mackenzie 1992]; see also [Brown and Mackenzie 1992]. In the latter article, a complete answer was given in the restricted case of locally trivial double Lie groupoids. More recently, a description in two stages of discrete double groupoids was given in [Andruskiewitsch and Natale 2009]. To state them, let us recall that a diagram over a pair of groupoids \mathcal{V} and \mathcal{H} is a triple (\mathcal{D}, j, i) where \mathcal{D} is a groupoid and $i : \mathcal{H} \rightarrow \mathcal{D}$, $j : \mathcal{V} \rightarrow \mathcal{D}$ are morphisms of groupoids (over a fixed set of points). The stages are:

- (a) Any double groupoid is an extension of slim double groupoid (its *frame*) by an abelian group bundle.

This work was partially supported by Pontificia Universidad Javeriana, Bogotá, Colombia; CONICET, Fundación Antorchas, Agencia Córdoba Ciencia, ANPCyT and Secyt (UNC) (Argentina).
MSC2010: 18D05, 20L05, 22A22.

Keywords: Lie groupoid, core groupoid, proper action, double groupoid, core action, groupoid diagram.

- (b) The category of slim double groupoids, with fixed vertical and horizontal groupoids \mathcal{V} and \mathcal{H} , satisfying the filling condition, is equivalent to the category of diagrams over \mathcal{V} and \mathcal{H} .

In this paper, we extend stage (b) to the setting of double Lie groupoids. In this context the top-right map has to be a surjective submersion, a differentiable version of the filling condition [Mackenzie 1992]. As one may expect, there are some topological and geometrical ingredients in our main result, which is this:

Theorem 3.7. *The category of slim double Lie groupoids, with fixed vertical and horizontal Lie groupoids \mathcal{V} and \mathcal{H} , and proper core action, is equivalent to the category of diagrams of Lie groupoids (\mathcal{D}, j, i) such that the maps j and i are transversal at the identities.*

Our proof of this theorem relies on [Andruskiewitsch and Natale 2009, Theorem 2.8] and some topological and differentiable considerations. We should note that the properness of the core action really boils down to the closedness of the anchor map of the corresponding action groupoid, since the core action is actually free.

We have started to investigate the stage (a) of the above construction in the differentiable setting; we expect to present this analysis in a future paper.

1. Preliminaries on Lie groupoids and double Lie groupoids

We denote a groupoid in the form $\mathcal{G} \begin{smallmatrix} \xrightarrow{s} \\ \xrightarrow{e} \end{smallmatrix} \mathcal{P}$, where s and e stand for “source” and “end”; and the identity map is $\text{id} : \mathcal{P} \rightarrow \mathcal{G}$. Recall that a groupoid $\mathcal{G} \begin{smallmatrix} \xrightarrow{s} \\ \xrightarrow{e} \end{smallmatrix} \mathcal{P}$ is a *Lie groupoid* [Mackenzie 2005], if \mathcal{P} and \mathcal{G} are smooth manifolds, s and e are surjective submersions and the other structural maps are smooth. The *anchor* of \mathcal{G} is the map $\chi : \mathcal{G} \rightarrow \mathcal{P} \times \mathcal{P}$ given by $\chi(g) = (s(g), e(g))$.

Definition 1.1. Let $\mathcal{G} \begin{smallmatrix} \xrightarrow{s} \\ \xrightarrow{e} \end{smallmatrix} \mathcal{P}$ be a Lie groupoid and let $U \subseteq \mathcal{P}$ be an open subset. A *local bisection* of \mathcal{G} on U is a smooth map $\sigma : U \rightarrow \mathcal{G}$ which is a section of e and such that $V := (s \circ \sigma)(U)$ is an open subset of \mathcal{P} and $s \circ \sigma : U \rightarrow V$ is a diffeomorphism. Define $\mathcal{G}^U = s^{-1}(U)$ and $\mathcal{G}_U = e^{-1}(U)$; the *local left* and *right translations* induced by σ are the maps

$$(1-1) \quad \begin{aligned} L_\sigma : \mathcal{G}^U &\rightarrow \mathcal{G}^V, & g &\mapsto \sigma(s(g))g; \\ R_\sigma : \mathcal{G}_V &\rightarrow \mathcal{G}_U, & g &\mapsto g\sigma((s \circ \sigma)^{-1}(e(g))). \end{aligned}$$

For more on bisections see [Mackenzie 2005, Section 1.4].

Double Lie groupoids.

Definition 1.2 (Ehresmann). A *double groupoid* is a groupoid object internal to the category of groupoids. In other terms, a *double groupoid* consists of a set \mathcal{B} with two groupoid structures with *bases* \mathcal{H} and \mathcal{V} , which are themselves groupoids

over a common base \mathcal{P} , all subject to the compatibility condition that the structure maps of each structure are morphisms with respect to the other.

It is usual to represent a double groupoid $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ as a diagram of four related groupoids

$$\begin{array}{ccc}
 \mathcal{B} & \begin{array}{c} \xrightarrow{l} \\ \xrightarrow{r} \end{array} & \mathcal{V} \\
 \begin{array}{c} \Downarrow t \\ \Downarrow b \end{array} & & \begin{array}{c} \Downarrow t \\ \Downarrow b \end{array} \\
 \mathcal{H} & \begin{array}{c} \xrightarrow{l} \\ \xrightarrow{r} \end{array} & \mathcal{P}
 \end{array}$$

where t, b, l, r mean ‘‘top’’, ‘‘bottom’’, ‘‘left’’ and ‘‘right’’. We sketch the main axioms that these groupoids should satisfy and refer to, for instance, [Andruskiewitsch and Natale 2005, Section 2] and [Andruskiewitsch and Natale 2006, Section 1] for detailed exposition and other conventions.

The elements of \mathcal{B} are called boxes and will be denoted by

$$A = \begin{array}{c} t(A) \\ \boxed{A} \\ b(A) \end{array} \begin{array}{c} l(A) \\ r(A) \end{array} \in \mathcal{B}.$$

Here $t(A), b(A) \in \mathcal{H}$ and $l(A), r(A) \in \mathcal{V}$. The identity maps will be denoted $\mathbf{id} : \mathcal{V} \rightarrow \mathcal{B}$ and $\mathbf{id} : \mathcal{H} \rightarrow \mathcal{B}$. The product in the groupoid \mathcal{B} with base \mathcal{V} is called *horizontal* and denoted by AB or $\{AB\}$, for $A, B \in \mathcal{B}$ with $r(A) = l(B)$. The product in the groupoid \mathcal{B} with base \mathcal{H} is called *vertical* and we shall denote it by $\begin{smallmatrix} A \\ B \end{smallmatrix}$, for $A, B \in \mathcal{B}$ with $b(A) = t(B)$. This pictorial notation is useful for understanding products in the double structure. For instance, the compatibility axioms between the horizontal and vertical products are described by

$$\begin{array}{c} t \\ \boxed{A} \\ b \end{array} \begin{array}{c} i \\ r \end{array} \begin{array}{c} i' \\ \boxed{B} \\ b' \end{array} \begin{array}{c} r' \\ r' \end{array} = \begin{array}{c} ti' \\ \boxed{\{AB\}} \\ bb' \end{array} \begin{array}{c} r' \\ r' \end{array} \quad \text{and} \quad \begin{array}{c} t \\ \boxed{A} \\ b \end{array} \begin{array}{c} l \\ r \end{array} \begin{array}{c} b \\ b \end{array} \begin{array}{c} l' \\ \boxed{B} \\ b' \end{array} \begin{array}{c} r' \\ r' \end{array} = \begin{array}{c} t \\ \boxed{\begin{smallmatrix} A \\ B \end{smallmatrix}} \\ b' \end{array} \begin{array}{c} rr' \\ rr' \end{array}$$

We omit the letter inside the box if no confusion arises. We also write A^h and A^v to denote the inverse of $A \in \mathcal{B}$ with respect to the horizontal and vertical structures of groupoid over \mathcal{B} respectively. When one of the sides of a box is an identity, we draw this side as a double edge. For example, if $t(A) = \mathbf{id}_p$, we draw $\boxed{}$ and say that $t(A) \in \mathcal{P}$.

Definition 1.3 [Mackenzie 1992]. A double groupoid is a *double Lie groupoid* if all the four groupoids involved are Lie groupoids and the *top-right corner map*

$$(1-2) \quad \lrcorner : \mathcal{B} \rightarrow \mathcal{H} \times_t \mathcal{V}, \quad A \mapsto \lrcorner(A) = (t(A), r(A)),$$

is a surjective submersion.

We shall say that a double groupoid is *discrete* if no differentiable or topological structure is present. A discrete double groupoid satisfies the *filling condition* if the *top-right corner map* defined in (1-2) is surjective. We refer the reader to [Andruskiewitsch and Natale 2009] for more details on discrete double groupoids and the filling condition.

Definition 1.4 [Brown and Mackenzie 1992; Mackenzie 1992]. Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a double Lie groupoid. The core groupoid $\mathbf{E}(\mathcal{B})$ of \mathcal{B} is the set

$$\mathbf{E}(\mathcal{B}) = \{E \in \mathcal{B} : t(E), r(E) \in \mathcal{P}\} = (\lrcorner)^{-1}\{(\text{id}_p, \text{id}_p) : p \in \mathcal{P}\}$$

with source and target projections $s_{\mathbf{E}}, e_{\mathbf{E}} : \mathbf{E}(\mathcal{B}) \rightarrow \mathcal{P}$, given by $s_{\mathbf{E}}(E) = bl(E)$ and $e_{\mathbf{E}}(E) = tr(E)$ respectively; identity map given by $\text{id}_p = \Theta_p := \mathbf{id} \circ \text{id}(p)$; multiplication and inverse given by

$$(1-3) \quad E \circ F := \left\{ \begin{array}{cc} \mathbf{id}l(F) & F \\ E & \mathbf{id}(b(F)) \end{array} \right\}, \quad E^{(-1)} := (E \mathbf{id} b(E)^{-1})^v = \left\{ \begin{array}{cc} \mathbf{id}l(E)^{-1} \\ E^h \end{array} \right\},$$

for every compatible $E, F \in \mathbf{E}(\mathcal{B})$.

We observe that the elements of $\mathbf{E}(\mathcal{B})$ are of the form $E = \boxed{\quad}$; the source gives the bottom-left vertex and the target gives the top-right vertex of the box. Since the top-right corner map is a surjective submersion, the core groupoid $\mathbf{E}(\mathcal{B})$ is a closed embedded submanifold of \mathcal{B} and clearly $s_{\mathbf{E}}$ and $e_{\mathbf{E}}$ are surjective submersions. With these structural maps $\mathbf{E}(\mathcal{B})$ becomes a Lie groupoid with base \mathcal{P} (differentiability conditions being easily verified because $\mathbf{E}(\mathcal{B})$ is an embedded submanifold of \mathcal{B}).

Coarse double groupoid. Let \mathcal{P} be a set and \mathcal{V}, \mathcal{H} be groupoids over \mathcal{P} . Let $\square(\mathcal{V}, \mathcal{H})$ be the set

$$(\mathcal{V} \times_b \times_l \mathcal{H}) \times_{(t,r) \times (l,b)} (\mathcal{H} \times_t \times_r \mathcal{V});$$

that is, the set of quadruples $\begin{pmatrix} x & \\ f & g \\ & y \end{pmatrix}$ with $x, y \in \mathcal{H}$, $f, g \in \mathcal{V}$ such that

$$l(x) = t(f), \quad r(x) = t(g), \quad l(y) = b(f), \quad r(y) = b(g).$$

If no confusion arises, we shall denote a quadruple as above by a box as follows:

$$h \begin{array}{c} x \\ \square \\ y \end{array} g.$$

$\square(\mathcal{V}, \mathcal{H}) \rightrightarrows \mathcal{H}$
 The collection $\begin{array}{ccc} \square(\mathcal{V}, \mathcal{H}) & \rightrightarrows & \mathcal{H} \\ \Downarrow & & \Downarrow \\ \mathcal{V} & \rightrightarrows & \mathcal{P} \end{array}$ forms a double groupoid in the way, called the *coarse double groupoid* with sides in \mathcal{H} and \mathcal{V} .

Definition 1.5 [Andruskiewitsch and Natale 2009]. A double groupoid $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ is *slim* if the morphism of (discrete) double groupoids $\Pi : \mathcal{B} \rightarrow \square(\mathcal{V}, \mathcal{H})$ given by

$$\Pi \left(f \begin{array}{c} x \\ \square \\ A \\ y \end{array} g \right) = \left(f \begin{array}{c} x \\ \square \\ g \\ y \end{array} \right), \quad f \begin{array}{c} x \\ \square \\ A \\ y \end{array} g \in \mathcal{B},$$

is injective. That is, any box in \mathcal{B} is determined by its sides.

We now pass to the context of double Lie groupoids. Let M , N and P be smooth manifolds, let $f : M \rightarrow P$ and $g : N \rightarrow P$ be smooth maps, we remind that f and g are called *transversal* at $p \in \mathcal{P}$, if for every $m \in M$ and $n \in N$ such that $p = f(m) = g(n)$ the condition $(T_m f)(T_m M) + (T_n g)(T_n N) = T_p P$ holds. We said that f and g are *transversal* if they are transversal at every p as above.

Remark 1.6. Let \mathcal{V} and \mathcal{H} be Lie groupoids over the same manifold \mathcal{P} . If the anchors maps $\chi_{\mathcal{V}} : \mathcal{V} \rightarrow \mathcal{P} \times \mathcal{P}$ and $\chi_{\mathcal{H}} : \mathcal{H} \rightarrow \mathcal{P} \times \mathcal{P}$ are transversal then $\square(\mathcal{V}, \mathcal{H})$ is a double Lie groupoid [Brown and Mackenzie 1992].

2. Diagrams of groupoids

The following notions were introduced in [Andruskiewitsch and Natale 2009], in the context of discrete double groupoids. Let \mathcal{V} and \mathcal{H} be groupoids over the same base \mathcal{P} . A *diagram* over \mathcal{H} and \mathcal{V} is a triple (\mathcal{D}, j, i) where \mathcal{D} is a groupoid over \mathcal{P} and $i : \mathcal{H} \rightarrow \mathcal{D}$, $j : \mathcal{V} \rightarrow \mathcal{D}$ are morphisms of groupoids over \mathcal{P} . A diagram of groupoids (\mathcal{D}, j, i) over \mathcal{V} and \mathcal{H} is called a $(\mathcal{V}, \mathcal{H})$ -*factorization* of \mathcal{D} if $\mathcal{D} = j(\mathcal{V})i(\mathcal{H})$.

To each diagram of groupoids (\mathcal{D}, j, i) we associate a discrete double groupoid, denoted by $\square(\mathcal{D}, j, i)$ and defined as the double subgroupoid of $\square(\mathcal{V}, \mathcal{H})$ whose boxes are of the form

$$A = h \begin{array}{c} x \\ \square \\ g \\ y \end{array} \in \square(\mathcal{V}, \mathcal{H}),$$

with $x, y \in \mathcal{H}$, $g, h \in \mathcal{V}$, such that $i(x)j(g) = j(h)i(y)$ in \mathcal{D} .

Diagrams of Lie groupoids.

Definition 2.1. Let \mathcal{V} and \mathcal{H} be Lie groupoids over \mathcal{P} . A *diagram of Lie groupoids* over \mathcal{H} and \mathcal{V} is a diagram of groupoids (\mathcal{D}, j, i) such that \mathcal{D} is a Lie groupoid and the maps i and j are smooth. If this diagram is a factorization in the sense above, then we say that it is a $(\mathcal{V}, \mathcal{H})$ -*factorization of the Lie groupoid* \mathcal{D} .

Let (\mathcal{D}, j, i) be a diagram of Lie groupoids over \mathcal{H} and \mathcal{V} . Our aim in this Subsection is to determine when $\square(\mathcal{D}, j, i)$ is a double Lie groupoid. For this, we shall consider two maps; the first one is the composition

$$\mathcal{H}_{r \times_t} \mathcal{V} \xrightarrow{i \times j} \mathcal{D}_{e \times_s} \mathcal{D} \xrightarrow{m} \mathcal{D},$$

that is,

$$\Phi : \mathcal{H}_{r \times_t} \mathcal{V} \rightarrow \mathcal{D}, \quad (x, g) \mapsto i(x)j(g),$$

where s, e and m are the source, target and composition maps of \mathcal{D} . The second one is

$$\Psi : \mathcal{V}_{b \times_l} \mathcal{H} \rightarrow \mathcal{D}, \quad (f, y) \mapsto j(f)i(y).$$

Since t, b, l and r are surjective submersions, we have that the fiber products involved in the above maps, $\mathcal{V}_{b \times_l} \mathcal{H}$ and $\mathcal{H}_{r \times_t} \mathcal{V}$, are embedded submanifolds of $\mathcal{V} \times \mathcal{H}$ and $\mathcal{H} \times \mathcal{V}$ respectively, and since i and j are smooth then Φ and Ψ also are. With the above maps

$$\square(\mathcal{D}, j, i) = (\mathcal{V}_{b \times_l} \mathcal{H})_{\Psi \times \Phi} (\mathcal{H}_{r \times_t} \mathcal{V}),$$

and from general transversality theory [Lang 2002, Proposition 2.5], if Φ and Ψ are transversal then $\square(\mathcal{D}, j, i)$ is an embedded submanifold of $(\mathcal{V}_{b \times_l} \mathcal{H}) \times (\mathcal{H}_{r \times_t} \mathcal{V})$.

Lemma 2.2. *Let (\mathcal{D}, j, i) be a diagram of Lie groupoids. If i and j are transversal at the identities, then Ψ and Φ defined above are submersions.*

Proof. We take $((f, y), (x, g)) \in (\mathcal{V}_{b \times_r} \mathcal{H}) \times (\mathcal{H}_{r \times_t} \mathcal{V})$ such that $\Psi(f, y) = \Phi(x, g)$ i.e., $j(f)i(y) = i(x)j(g)$. Now, by [Lang 2002, Proposition 2.5], we have

$$T_{(x,g)}(\mathcal{H}_{r \times_t} \mathcal{V}) = \{(Y, X) \in (T_x \mathcal{H}) \times (T_g \mathcal{V}) / (T_x r)(Y) = (T_g t)(X)\}.$$

Let $W \in T_{(f,y)}(\mathcal{H}_{r \times_t} \mathcal{V})$. We need to prove that there is $(X_1, Y_1) \in T_{(f,y)}(\mathcal{V}_{b \times_l} \mathcal{H})$ such that $T_{(f,y)}\Psi(X_1, Y_1) = W$. Note that

$$\begin{aligned} T_{(f,y)}\Psi(X_1, Y_1) &= (T_{(j(f), i(y))}m)(T_{(f,y)}j \times i)(X_1, Y_1) \\ &= (T_{(j(f), i(y))}m)((T_f j)(X_1), (T_y i)(Y_1)). \end{aligned}$$

Now, we know that in $\mathcal{H} \rightrightarrows \mathcal{P}$ there is a local bisection $\tau : U \rightarrow \mathcal{H}$ with $r(y) \in U \subseteq \mathcal{P}$ open and $\tau(r(y)) = y$ [Mackenzie 2005, Proposition 1.4.9]. Since τ is a bisection, it induces local left and right translations as in (1-1). Concretely, set $V = (l \circ \tau)(U)$ open in \mathcal{P} , $\mathcal{H}^U = l^{-1}(U)$ and $\mathcal{H}_U = r^{-1}(U)$ and define

$$\begin{aligned} L_\tau : \mathcal{H}^U &\rightarrow \mathcal{H}^V, \quad z \mapsto \tau(l(z))z, \\ R_\tau : \mathcal{H}_V &\rightarrow \mathcal{H}_U, \quad z \mapsto z \tau((l \circ \tau)^{-1}r(z)). \end{aligned}$$

Define the map $\tau_{\mathcal{D}} : U \rightarrow \mathcal{D}$ by $i \circ \tau$; since i is a morphism of groupoids, $\tau_{\mathcal{D}}$ is a local bisection of \mathcal{D} . Also note that $\tau_{\mathcal{D}}(e(i(y))) = (i \circ \tau)(r(y)) = i(y)$.

Analogously, there is a local bisection $\sigma : U' \rightarrow \mathcal{V}$ such that $\sigma_{\mathcal{V}}(b(f)) = f$, with $U' \subseteq \mathcal{D}$ open and $b(f) \in U'$. Again this induces a local bisection $\sigma_{\mathcal{D}} : U' \rightarrow \mathcal{D}$ such that $\sigma_{\mathcal{D}}(e(j(f))) = j(f)$. Let $(X_1, Y_1) \in T_{(f,y)}(\mathcal{V}_b \times_l \mathcal{H})$, then by Xu's formula for product in the tangent groupoid [Mackenzie 2005, Theorem 1.4.14] we obtain:

$$\begin{aligned} T_{(f,y)}\Psi(X_1, Y_1) &= (T_{(j(f),i(y))}m)((T_f j)(X_1), (T_y i)(Y_1)) \\ &= (T_{i(y)}L_{\sigma_{\mathcal{D}}})(T_y i)(Y_1) + (T_{j(f)}R_{\tau_{\mathcal{D}}})(T_f j)(X_1) \\ &\quad - (T_{i(y)}L_{\sigma_{\mathcal{D}}})(T_{\text{Id}_{\mathcal{D}}l(y)}R_{\tau_{\mathcal{D}}})(T_{l(y)}\text{id}_{\mathcal{D}})(z), \end{aligned}$$

where we write $z = (T_{j(f)}e)(T_f j)(X_1) = (T_{i(y)}s)(T_y i)(Y_1)$.

Now $(T_{j(f)i(y)}L_{\sigma_{\mathcal{D}}}^{-1})(W) \in T_{i(y)}\mathcal{D}$ because we have

$$\begin{aligned} L_{\sigma_{\mathcal{D}}}^{-1}(j(f)i(y)) &= \sigma_{\mathcal{D}}^{-1}(s(j(f)i(y)))j(f)i(y) \\ &= \sigma_{\mathcal{D}}((s \circ \sigma_{\mathcal{D}})^{-1}(s(j(f)i(y))))^{-1}j(f)i(y) \\ &= \sigma_{\mathcal{D}}((s \circ \sigma_{\mathcal{D}})^{-1}s(j(f)))^{-1}j(f)i(y) \\ &= \sigma_{\mathcal{D}}(e(j(f)))^{-1}j(f)i(y) \\ &= j(f)^{-1}j(f)i(y) = i(y). \end{aligned}$$

In an analogous way, we have $(T_{i(y)}R_{\tau_{\mathcal{D}}}^{-1})(T_{j(f)i(y)}L_{\sigma_{\mathcal{D}}}^{-1})(W) \in T_{\text{Id}_{\mathcal{D}}l(y)}\mathcal{D}$ since

$$\begin{aligned} R_{\tau_{\mathcal{D}}}^{-1}(i(y)) &= i(y)\tau_{\mathcal{D}}^{-1}((s \circ \tau_{\mathcal{D}})^{-1}e(i(y))) \\ &= i(y)\tau_{\mathcal{D}}((s \circ \tau_{\mathcal{D}})^{-1}((s \circ \tau_{\mathcal{D}})^{-1}e(i(y))))^{-1} \\ &= i(y)\tau_{\mathcal{D}}((s \circ \tau_{\mathcal{D}})^{-1}((s \circ \tau_{\mathcal{D}})e(i(y))))^{-1} \\ &= i(y)\tau_{\mathcal{D}}(e(i(y)))^{-1} = i(y)^{-1}i(y) = \text{Id}_{\mathcal{D}}s(i(y)) = \text{Id}_{\mathcal{D}}(l(y)). \end{aligned}$$

Write $p = l(y)$, since i and j are transversal at $\text{Id}_{\mathcal{D}}(p)$ then

$$T_{\text{Id}_p}\mathcal{D} = (T_{\text{Id}_p}i)(T_{\text{Id}_p}\mathcal{H}) + (T_{\text{Id}_p}j)(T_{\text{Id}_p}\mathcal{V})$$

and in consequence we can find $X \in T_{\text{Id}_p}\mathcal{H}$ and $Y \in T_{\text{Id}_p}\mathcal{V}$ such that

$$(T_{i(y)}R_{\tau_{\mathcal{D}}}^{-1})(T_{j(f)i(y)}L_{\sigma_{\mathcal{D}}}^{-1})(W) = (T_{\text{Id}_p}i)(X) + (T_{\text{Id}_p}j)(Y).$$

Thus, if we take the vectors

$$X' = X + (T_p\text{Id}_{\mathcal{H}})(T_{\text{Id}_p}t)(Y), \quad Y' = Y + (T_p\text{Id}_{\mathcal{V}})(T_{\text{Id}_p}l)(X),$$

a direct calculation shows that

$$(2-1) \quad (T_{i(y)}R_{\tau_{\mathcal{D}}}^{-1})(T_{j(f)i(y)}L_{\sigma_{\mathcal{D}}}^{-1})(W) = (T_{\text{Id}_p}i)(X') + (T_{\text{Id}_p}j)(Y') - (T_p\text{Id}_{\mathcal{D}})(Z)$$

where $Z = (T_{\text{Id}_p}b)(Y) + (T_{\text{Id}_p}l)(X)$.

Since $(L_{\sigma_{\mathfrak{D}}} \circ R_{\tau_{\mathfrak{D}}} \circ i) = (L_{\sigma_{\mathfrak{D}}} \circ i \circ R_{\tau})$ and $(L_{\sigma_{\mathfrak{D}}} \circ R_{\tau_{\mathfrak{D}}} \circ j) = (R_{\tau_{\mathfrak{D}}} \circ j \circ L_{\sigma_{\mathfrak{V}}})$, we may apply $(T_{i(y)}L_{\sigma_D})(T_{\text{Id}_p}R_{\tau_{\mathfrak{D}}})$ to both sides of (2-1) and arrive at

$W =$

$$(T_{i(y)}L_{\sigma_{\mathfrak{D}}})(T_y i)(Y_1) + (T_{j(f)}R_{\tau_{\mathfrak{D}}})(T_f j)(X_1) - (T_{i(y)}L_{\sigma_{\mathfrak{D}}})(T_{\text{Id}_p}R_{\tau_{\mathfrak{D}}})(T_p \text{Id}_{\mathfrak{D}})(Z),$$

where $X_1 = (T_{\text{Id}_p}L_{\sigma_{\mathfrak{V}}})(Y')$ and $Y_1 = (T_{\text{Id}_p}R_{\tau})(X')$. It is clear that $(T_f b)(X_1) = Z = (T_y l)(Y_1)$, thus $T_{(f,y)}\Psi(X_1, Y_1) = W$.

In the same way we prove that Φ is a submersion. \square

From the above result we obtain the following immediate consequence.

Theorem 2.3. *Let (\mathfrak{D}, j, i) be a $(\mathcal{V}, \mathcal{H})$ -factorization of the Lie groupoid \mathfrak{D} . If i and j are transversal at the identities, then $\square(\mathfrak{D}, j, i)$ is a slim double Lie groupoid.*

Proof. By Lemma 2.2 we have that Φ and Ψ are transversal, thus $\square(\mathfrak{D}, j, i)$ is an embedded submanifold of $(\mathcal{V}_{b \times l} \times \mathcal{H}_{r \times t})$. Since Φ and Ψ are surjective submersions, both projections from $\square(\mathfrak{D}, j, i) = (\mathcal{V}_{b \times l} \times \mathcal{H}_{r \times t})_{\Psi \times \Phi}$ to $(\mathcal{V}_{b \times l} \times \mathcal{H}_{r \times t})$ and $\mathcal{H}_{r \times t} \times \mathcal{V}$ are surjective submersions and the same is true for the projections from the fiber products $\mathcal{V}_{b \times l} \times \mathcal{H}$ and $\mathcal{H}_{r \times t} \times \mathcal{V}$ to \mathcal{V} and \mathcal{H} ; then the top, bottom, left and right maps from $\square(\mathfrak{D}, j, i)$ are surjective submersions and the same for the top-right corner map. It is clear that the compositions, the identities maps and the inversions maps are smooth. \square

The diagonal groupoid. In this subsection, all groupoids are discrete. We review from [Andruskiewitsch and Natale 2009] the construction of the diagonal groupoid. Let \mathcal{B} be a double groupoid that satisfies the filling condition and let $\mathcal{V} \otimes \mathcal{H}$ be the free product (over \mathcal{P}) of the vertical and horizontal groupoids; see the same reference and [Higgins 2005]. We set

$$[A] := xgy^{-1}h^{-1} \in \mathcal{V} \otimes \mathcal{H} \quad \text{if } h \begin{array}{c} x \\ \boxed{A} \\ y \end{array} g \in \mathcal{B}.$$

We write $J_{\otimes}(\mathcal{B})$ for the subgroupoid of $\mathcal{V} \otimes \mathcal{H}$ generated by $\{[A] | A \in \mathcal{B}\}$. As $s_{\otimes}([A]) = e_{\otimes}([A]) = tl(A)$ we have that the groupoid $J_{\otimes}(\mathcal{B}) \rightrightarrows \mathcal{P}$ is in fact a group bundle. We know that the group bundle $J_{\otimes}(\mathcal{B})$ is a normal subgroupoid of $\mathcal{V} \otimes \mathcal{H}$ [Andruskiewitsch and Natale 2009, Lemma 3.5].

Assume that $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ is slim; then the associated *diagonal groupoid* is the quotient $\mathfrak{D}(\mathcal{B}) = \mathcal{V} \otimes \mathcal{H} / J_{\otimes}(\mathcal{B})$. If we compose the natural inclusions of \mathcal{V} and \mathcal{H} in $\mathcal{V} \otimes \mathcal{H}$ with the projections to $\mathfrak{D}(\mathcal{B})$ we get two groupoid morphisms

$$i : \mathcal{H} \rightarrow \mathfrak{D}(\mathcal{B}) \quad \text{and} \quad j : \mathcal{V} \rightarrow \mathfrak{D}(\mathcal{B}).$$

Thus we have a $(\mathcal{V}, \mathcal{H})$ -factorization $(\mathcal{D}(\mathcal{B}), i, j)$. In this subsection our aim is to give another presentation of the diagonal groupoid as a quotient of $\mathcal{V} \times_l \mathcal{H}$.

Proposition 2.4. *Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a slim double groupoid that satisfies the filling condition. We define on $\mathcal{V} \times_l \mathcal{H}$ the following relation $\sim_{\mathcal{B}}$:*

$$(v_1, h_1) \sim_{\mathcal{B}} (v_2, h_2) \iff r(h_1) = r(h_2), t(v_1) = t(v_2) \text{ and } v_1 h_1 h_2^{-1} v_2^{-1} \in J_{\otimes}(\mathcal{B}).$$

Then $\sim_{\mathcal{B}}$ is an equivalence relation and the map

$$\phi : \mathcal{V} \times_l \mathcal{H} / \sim_{\mathcal{B}} \rightarrow \mathcal{D}(\mathcal{B}), \quad [v, h] \mapsto j(v)i(h)$$

is a well defined isomorphism of quivers over \mathcal{P} .

Proof. Clearly $\sim_{\mathcal{B}}$ is an equivalence relation; we write $\mathcal{G} := (\mathcal{V} \times_l \mathcal{H}) / \sim_{\mathcal{B}}$ and denote by $[f, x]$ the equivalence class of $(f, x) \in \mathcal{V} \times_l \mathcal{H}$. Let $(f_1, x_1), (f_2, x_2)$ be elements of $\mathcal{V} \times_l \mathcal{H}$ such that $[f_1, x_1] = [f_2, x_2]$, then we have $f_1 x_1 x_2^{-1} f_2^{-1} \in J_{\otimes}(\mathcal{B})$ so that $\overline{f_1 x_1} = \overline{f_2 x_2}$ in $\mathcal{D}(\mathcal{B})$ (where \bar{w} denotes the image of w under i if w belongs to \mathcal{H} , or under j if w belongs to \mathcal{V}). This proves that ϕ is well defined.

Suppose that $g, g' \in \mathcal{V}$ and $x, x' \in \mathcal{H}$ is any collection satisfying $\bar{g} \bar{x} = \overline{g' x'}$. Then $g' x' x^{-1} g^{-1} \in J_{\otimes}(\mathcal{B})$, hence $[g', x'] = [g, x]$. Therefore ϕ is injective.

To prove that ϕ is surjective, let $d \in \mathcal{D}(\mathcal{B})$. Then $d = \overline{d_1 d_2 \cdots d_n}$ with d_i an element of \mathcal{V} or \mathcal{H} . Let $d_i \in \mathcal{H}$, $d_{i+1} \in \mathcal{V}$ with $r(d_i) = t(d_{i+1})$. Since \mathcal{B} satisfies the filling condition, the corner

$$(2-2) \quad \begin{array}{c} d_i \\ \square \\ d_{i+1} \end{array}$$

can be completed to a box in \mathcal{B} ; that is, there exists $B \in \mathcal{B}$ such that

$$B = f \begin{array}{c} d_i \\ \square \\ y \end{array} d_{i+1}.$$

Thus, $d_i d_{i+1} y^{-1} f^{-1} \in J_{\otimes}(\mathcal{B})$ and $\overline{d_i d_{i+1}} = \bar{f} \bar{y}$. So, we can commute the d_i 's in $d = \overline{d_1 d_2 \cdots d_n}$ in such a way that we can obtain $d = \bar{g} \bar{x}$ with $g \in \mathcal{V}$, $x \in \mathcal{H}$ and $b(g) = l(x)$. This proves that ϕ is surjective. \square

Remark 2.5. The map ϕ from the previous lemma induces a structure of groupoid on $\mathcal{G} = \mathcal{V} \times_l \mathcal{H} / \sim_{\mathcal{B}}$ with the following structural maps:

- The source and the target projections are

$$s : \mathcal{G} \rightarrow \mathcal{P}, \quad [v, h] \mapsto t(v); \quad e : \mathcal{G} \rightarrow \mathcal{P}, \quad [v, h] \mapsto r(h).$$

- The identity map is $\text{id} : \mathcal{P} \rightarrow \mathcal{G}$, $p \mapsto \text{id}_p = [\text{id}_p, \text{id}_p]$.

- Multiplication is given by $[v_1, h_1][v_2, h_2] = [v_1 f, z h_2]$, where $f \begin{array}{c} h_1 \\ \square \\ z \end{array} v_2 \in \mathcal{B}$, when it is defined;
- The inverse map is given by $[v, h]^{-1} = [f^{-1}, z^{-1}]$, for every $[v, h] \in \mathcal{G}$ and where $v \begin{array}{c} z \\ \square \\ h \end{array} f \in \mathcal{B}$.

Now we recall the following lemma which will be very useful for our purposes.

Lemma 2.6 [Andruskiewitsch and Natale 2009, Lemma 3.8]. *Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a slim double groupoid that satisfies the filling condition. Let $f \in \mathcal{V}$ and $x \in \mathcal{H}$ such that:*

- $l(x) = b(f)$ and $t(f) = r(x)$,
- $f x \in J_{\otimes}(\mathcal{B}) \subset \mathcal{V} \otimes \mathcal{H}$.

Then there exists $E \in \mathbf{E}(\mathcal{B})$ such that $E = f \begin{array}{c} \square \\ \square \\ x \end{array}$. □

We give an alternative description of $\sim_{\mathcal{B}}$ using this lemma. Indeed, $(f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2)$ if and only if $f_1 x_1 x_2^{-1} f_2^{-1} \in J_{\otimes}(\mathcal{B})$. Since $J_{\otimes}(\mathcal{B})$ is a normal subgroupoid, we have that $(f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2)$ if and only if $f_2^{-1} f_1 x_1 x_2^{-1} \in J_{\otimes}(\mathcal{B})$. Hence

$$(2-3) \quad (f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2) \iff \text{there exists } E = f_2^{-1} f_1 \begin{array}{c} \square \\ \square \\ x_1 x_2^{-1} \end{array} \in \mathcal{B}.$$

Thus, the graph of the relation $\sim_{\mathcal{B}}$ is

$$R = \left\{ (f_1, x_1, f_2, x_2) \in (\mathcal{V} \times_{b \times l} \mathcal{H}) \times_{\eta} (\mathcal{V} \times_{b \times l} \mathcal{H}) \mid \exists E \in \mathbf{E}(\mathcal{B}), E = f_2^{-1} f_1 \begin{array}{c} \square \\ \square \\ x_1 x_2^{-1} \end{array} \right\},$$

where

$$\eta : \mathcal{V} \times_{b \times l} \mathcal{H} \rightarrow \mathcal{P} \times \mathcal{P}, \quad (f, x) \mapsto (t(f), r(x));$$

from this we infer that the relation $\sim_{\mathcal{B}}$ is determined by the core groupoid of \mathcal{B} .

Lemma 2.7. *Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a slim double groupoid satisfying the filling condition. If $(f_1, x_1), (f_2, x_2) \in \mathcal{V} \times_{b \times l} \mathcal{H}$, then $(f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2)$ if and only if there exist $A, B \in \mathcal{B}$ such that*

$$A = f_1 \begin{array}{c} x \\ \square \\ x_1 \end{array} g, \quad \text{and} \quad B = f_2 \begin{array}{c} x \\ \square \\ x_2 \end{array} g.$$

Proof. In fact, if $A = f_1 \begin{array}{c} x \\ \square \\ x_1 \end{array} g$ and $B = f_2 \begin{array}{c} x \\ \square \\ x_2 \end{array} g$ belong to \mathcal{B} , then $x g x_1^{-1} f_1^{-1} \in$

$J_{\otimes}(\mathcal{B})$ and $x g x_2^{-1} f_2^{-1} \in J_{\otimes}(\mathcal{B})$; taking the inverse of the first one and composing with the second one we arrive at $f_1 x_1 x_2^{-1} f_2^{-1} \in J_{\otimes}(\mathcal{B})$, that is, $(f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2)$.

Conversely, if $(f_1, x_1) \sim_{\mathcal{B}} (f_2, x_2)$ then by (2-3) there is $E \in \mathbf{E}(\mathcal{B})$ such that

$$E = f_2^{-1} f_1 \begin{array}{|c|} \hline \square \\ \hline \end{array} \begin{array}{|c|} \hline x_1 x_2^{-1} \\ \hline \end{array}.$$

The filling condition guarantees that given $x_2 \in \mathcal{H}$ and $f_2 \in \mathcal{V}$ with $l(x_2) = b(f_2)$, there is a box $B' \in \mathcal{B}$ with $t(B') = x_2$ and $l(B') = f_2^{-1}$. Let

$$B' = f_2^{-1} \begin{array}{|c|} \hline x_2 \\ \hline \end{array} g^{-1} \quad \text{and} \quad A' := \left\{ \begin{array}{cc} E^v & \mathbf{id}(x_2) \\ \mathbf{id}(f_2) & B' \end{array} \right\} = f_1^{-1} \begin{array}{|c|} \hline x_1 \\ \hline \end{array} g^{-1}.$$

Let A, B be the vertical inverses of A' and B' . Thus $A = f_1 \begin{array}{|c|} \hline x \\ \hline \end{array} g$ and $B = f_2 \begin{array}{|c|} \hline x \\ \hline \end{array} g$ are both in \mathcal{B} and we get the result. \square

3. Slim double Lie groupoids

Actions of Lie groupoids.

Definition 3.1. A left action of a groupoid $\mathcal{G} \rightrightarrows \mathcal{P}$ along a map $\varepsilon : N \rightarrow \mathcal{P}$ is given by a map $\mathcal{G} \times_{e \times \varepsilon} N \rightarrow N$, denoted by $(g, n) \mapsto gn$, such that:

$$\varepsilon(hy) = s(h), \quad \mathbf{id}(\varepsilon(y)) y = y, \quad (gh)y = g(hy),$$

for all $g, h \in \mathcal{G}$ and $y \in N$ such that $e(g) = s(h)$ and $e(h) = \varepsilon(y)$. We shall simply say that the groupoid \mathcal{G} acts on N .

The *transformation* or *action groupoid* $\mathcal{G} \times N \rightrightarrows N$, associated with such an action, is a groupoid with set of arrows $\mathcal{G} \times_{e \times \varepsilon} N$. The source and target maps are

$$s', e' : \mathcal{G} \times N \rightarrow N, \quad \text{given by} \quad s'(g, n) = gn \quad \text{and} \quad e'(g, n) = n,$$

respectively, and composition $(g, n)(h, m) = (gh, m)$.

Let $\mathcal{G} \rightrightarrows \mathcal{P}$ be a Lie groupoid, N a smooth manifold and $\varepsilon : N \rightarrow \mathcal{P}$ a smooth map. A left action is *smooth* provided that the map $\mathcal{G} \times_{e \times \varepsilon} N \rightarrow N$ is smooth. The resulting action groupoid is again a Lie groupoid.

The *canonical left action* of a groupoid $\mathcal{G} \rightrightarrows \mathcal{P}$ along $\mathbf{id} : \mathcal{P} \rightarrow \mathcal{P}$ is given by $gp = s(g)$, where $g \in \mathcal{G}$, $p \in \mathcal{P}$ and $e(g) = p$. Recall that the orbit space \mathcal{P}/\mathcal{G} is the quotient space of \mathcal{P} where two points of \mathcal{P} are identified if there exists an arrow between them. Thus, if we have $\mathcal{G} \rightrightarrows \mathcal{P}$ an action along a map $\varepsilon : N \rightarrow \mathcal{P}$, the orbit space N/\mathcal{G} is equal to $N/\mathcal{G} \times N$, the orbit space of the action groupoid. Also, the stabilizer of $n \in N$ is defined in the usual way; it coincides with the isotropy group of n with respect to the action groupoid.

We recall that a continuous map $f : X \rightarrow Y$ between two topological spaces X and Y , is said to be *proper* if the inverse image of a compact subset of Y is compact.

Definition 3.2. A Lie groupoid $\mathcal{G} \rightrightarrows \mathcal{P}$ is *proper* if the anchor map is proper. An action of a Lie groupoid \mathcal{G} on a smooth manifold N is *proper* if the action groupoid $\mathcal{G} \ltimes N$ is proper.

The following proposition is useful to decide when an action is proper.

Proposition 3.3 [Tu 2004, 2.14]. *Let $\mathcal{G} \rightrightarrows \mathcal{P}$ be a Lie groupoid and N a smooth manifold endowed with a left action of \mathcal{G} . Then \mathcal{G} acts properly on N if and only if the anchor map $(s', e') : \mathcal{G} \ltimes N \rightarrow N \times N$ is closed and $\forall n \in N$ the stabilizer of n is compact.* \square

A groupoid is free when all the isotropy groups are trivial; thus, an action groupoid is free when all stabilizers are trivial.

Proposition 3.4 [Dragulete 2007, Theorem 3.3.1]. *If $\mathcal{G} \rightrightarrows \mathcal{P}$ is a free and proper groupoid, then the orbit space \mathcal{P}/\mathcal{G} is a smooth manifold and the orbit projection $\mathcal{P} \rightarrow \mathcal{P}/\mathcal{G}$ is a submersion.* \square

In particular, if \mathcal{G} is a Lie groupoid with a free and proper smooth action on N , then the orbit space N/\mathcal{G} is a smooth manifold and the orbit projection $N \rightarrow N/\mathcal{G}$ is a submersion.

The core action.

Proposition 3.5 [Andruskiewitsch and Natale 2009, Proposition 1.1]. *Suppose $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ is a slim double groupoid. Define $\gamma : \mathcal{B} \rightarrow \mathcal{P}$, $\gamma(A) = lb(A)$, the bottom-left vertex of A .*

(a) *There is an action of the core groupoid $\mathbf{E}(\mathcal{B})$ on $\gamma : \mathcal{B} \rightarrow \mathcal{P}$ given by*

$$(3-1) \quad E \rightarrow A := \left\{ \begin{array}{cc} \mathbf{id} l(A) & A \\ E & \mathbf{id} b(A) \end{array} \right\}, \quad A \in \mathcal{B}, E \in \mathbf{E}.$$

(b) *Let $B \in \mathcal{B}$. Then the stabilizer $\mathbf{E}(\mathcal{B})^B$ is trivial and the orbit of B is $\mathbb{O}_B = \{A \in \mathcal{B} : t(A) = t(B), r(A) = r(B)\}$.* \square

We introduce next the core action of the core groupoid on the fibred product $\mathcal{V}_{b \times l} \mathcal{H}$, that allows us to present the diagonal groupoid as a quotient of $\mathcal{V}_{b \times l} \mathcal{H}$.

Lemma 3.6. *Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a slim double groupoid and let $\eta : \mathcal{V}_{b \times l} \mathcal{H} \rightarrow \mathcal{P}$ be given by $\eta(f, x) = b(f) = l(x)$. Then $\mathbf{E}(\mathcal{B})$ acts on η by*

$$(3-2) \quad E \triangleright (f, x) = (f l(E), b(E) x), \quad \text{when } \eta(f, x) = e_{\mathbf{E}(\mathcal{B})}(E).$$

The orbit space $\mathcal{V}_{b \times l} \mathcal{H}/\mathbf{E}(\mathcal{B})$ coincides with $\mathcal{D}(\mathcal{B})$.

The map (3-2) is the *core action* of $\mathbf{E}(\mathcal{B})$ on $\mathcal{V}_{b \times l} \mathcal{H}$.

Proof. Since $b(f l(E)) = bl(E) = s_{\mathbf{E}(\mathcal{B})}(E)$ and $l(b(E) x) = lb(E) = s_{\mathbf{E}(\mathcal{B})}(E)$, the map $\triangleright : \mathbf{E}(\mathcal{B}) \times_{e_{\mathbf{E}(\mathcal{B})}} \times_{\eta} (\mathcal{V}_{b \times l} \mathcal{H}) \rightarrow \mathcal{V}_{b \times l} \mathcal{H}$ is well defined and \triangleright is an action:

$$\begin{aligned} (E \circ F) \triangleright (f, x) &= (f l(E \circ F), b(E \circ F) x) = (f l(F)l(E), b(E)b(F) x) \\ &= E \triangleright (f l(F), t(F) x) = E \triangleright (F \triangleright (f, x)). \end{aligned}$$

Also, $\eta(E \triangleright (f, x)) = \eta(f l(E), b(E) x) = l(b(E)x) = lb(E) = s_{\mathbf{E}(\mathcal{B})}(E)$.

For (b), if $(f, x) \sim_{\mathcal{B}} (g, y)$, there are $A, B \in \mathcal{B}$ such that $f \begin{array}{c} \boxed{A} \\ x \end{array} \begin{array}{c} z \\ h \end{array}$ and $g \begin{array}{c} \boxed{B} \\ y \end{array} \begin{array}{c} z \\ h \end{array}$;

see Lemma 2.7. Then, by Proposition 3.5, there is a box $E \in \mathbf{E}(\mathcal{B})$ such that $A = E \rightarrow B$, in consequence, $x = b(E)y$ and $f = g l(E)$.

Conversely, if $A, B \in \mathcal{B}$ and $E \in \mathbf{E}(\mathcal{B})$ satisfy $b(A) = b(E)b(B)$ and $l(A) = l(B)l(E)$, then the boxes $E \rightarrow B$ and B have the same top and right sides. By Lemma 2.7, we have

$$(l(E \rightarrow B), b(E \rightarrow B)) \sim_{\mathcal{B}} (l(B), b(B)),$$

that is $(l(A), b(A)) \sim_{\mathcal{B}} (l(B), b(B))$. From this we conclude that given (f, x) and (g, y) in $\mathcal{V}_{b \times l} \mathcal{H}$, we have $(f, x) \sim_{\mathcal{B}} (g, y)$ if and only if there exists $E \in \mathbf{E}(\mathcal{B})$ with $f = g l(E)$ and $x = b(E)y$. Thus the orbit space coincides with the diagonal groupoid. \square

Let $\pi : \mathcal{V}_{b \times l} \mathcal{H} \rightarrow \mathcal{V}_{b \times l} \mathcal{H}/\mathbf{E}(\mathcal{B})$ be the projection determined by (3-2). Let

$$\begin{aligned} \tilde{i} : \mathcal{H} &\rightarrow \mathcal{V}_{b \times l} \mathcal{H}, & x &\mapsto (\text{id}(l(x)), x) \\ \tilde{j} : \mathcal{V} &\rightarrow \mathcal{V}_{b \times l} \mathcal{H}, & f &\mapsto (f, \text{id}(b(f))), \end{aligned}$$

be the canonical inclusions of \mathcal{V} and \mathcal{H} , and consider the compositions $i = \pi \circ \tilde{i}$ and $j = \pi \circ \tilde{j}$.

The main theorem.

Theorem 3.7. *Let \mathcal{V} and \mathcal{H} be Lie groupoids. The assignments $\mathcal{B} \mapsto \mathcal{D}(\mathcal{B})$ and $(\mathcal{D}, j, i) \mapsto \square(\mathcal{D}, j, i)$ determine mutual category equivalences between*

- (a) *the category of slim double Lie groupoids $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ with proper core action, and*
- (b) *the category of $(\mathcal{V}, \mathcal{H})$ -factorizations of Lie groupoids (\mathcal{D}, j, i) such that the maps i and j are transversal at the identities.*

Proof. The equivalence of categories at the discrete level was obtained in [Andruskiewitsch and Natale 2009]. We see that this equivalence behaves well in the differentiable context. We start with the direction from (a) to (b).

Step 1. Let $(\mathcal{B}; \mathcal{V}, \mathcal{H}; \mathcal{P})$ be a slim double Lie groupoid. If the core action is proper, then $\mathcal{D}(\mathcal{B})$ is a Lie groupoid over \mathcal{P} . Furthermore, the maps i and j defined above are transversal at the identities.

Proof of the step. Since \mathcal{B} is slim, the action (3-2) is free. Hence, the orbit space $\mathcal{V}_{b \times l} \mathcal{H}/\mathbf{E}(\mathcal{B})$ has a unique manifold structure such that the canonical projection $\pi : \mathcal{V}_{b \times l} \mathcal{H} \rightarrow \mathcal{V}_{b \times l} \mathcal{H}/\mathbf{E}(\mathcal{B})$ is a surjective submersion, because the action is proper, see Proposition 3.4. We deduce using the identification in Lemma 3.6 that $\mathcal{D}(\mathcal{B})$ is a Lie groupoid over \mathcal{P} . In fact, the structure maps are described in Remark 2.5; using local sections of π , it is clear that the source and target maps are surjective submersions and that the other structural maps are smooth.

Let us now proceed with transversality. Let $p \in \mathcal{P}$ and we take a tangent vector $Z \in T_{[\text{id}_p, \text{id}_p]} \mathcal{D}(\mathcal{B})$, where $[\text{id}_p, \text{id}_p] = \pi(\text{id}_p, \text{id}_p)$. Since π is a surjective submersion, there is $(U, W) \in T_{(\text{id}_p, \text{id}_p)}(\mathcal{V}_{b \times l} \mathcal{H})$ such that $T_{(\text{id}_p, \text{id}_p)}\pi(U, W) = Z$. Choose

$$Y = U \in T_{\text{id}_p} \mathcal{V}, \quad X = W - (T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U) \in T_{\text{id}_p} \mathcal{H}.$$

It is clear that

$$(T_{\text{id}_p} \tilde{j})(Y) = (U, (T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U)) \text{ and } (T_{\text{id}_p} \tilde{i})(X) = ((T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} l)(X), X).$$

We compute

$$\begin{aligned} (T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} l)(X) &= (T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} l)(W) - (T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} l)(T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U) \\ &= (T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} l)(W) - (T_p \text{id}_{\mathcal{V}})(T_{\text{id}_p} b)(U) = 0; \end{aligned}$$

then $(T_{\text{id}_p} \tilde{i})(X) = (0, X)$. In consequence we have

$$\begin{aligned} (T_{\text{id}_p} \tilde{j})(Y) + (T_{\text{id}_p} \tilde{i})(X) &= (U, (T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U)) + (0, X) \\ &= (U, (T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U) + W - (T_p \text{id}_{\mathcal{H}})(T_{\text{id}_p} b)(U)) \\ &= (U, W). \end{aligned}$$

Then if we apply $T_{(\text{id}_p, \text{id}_p)}\pi$ to both sides of the above equation we arrive to

$$(T_{\text{id}_p} j)(Y) + (T_{\text{id}_p} i)(X) = Z,$$

that is, the maps i and j are transversal at the identities. \square

Step 2. Let (\mathcal{D}, j, i) be a $(\mathcal{V}, \mathcal{H})$ -factorization of Lie groupoids such that the maps i and j are transversal at the identities. Then the associated double groupoid $\square(\mathcal{D}, j, i)$ is a slim double Lie groupoid and its core action is proper.

Proof of the step. By Theorem 2.3, $\square(\mathcal{D}, j, i)$ is a slim double Lie groupoid. By [Brown and Mackenzie 1992], its core groupoid is a Lie groupoid with underlying

manifold

$$\mathcal{V}^{op} j \times_i \mathcal{H} = \{(h, y) \mid j(h^{-1}) = i(y)\}.$$

Up to this description, the core action on $\mathcal{V}_{b \times l} \mathcal{H}$ is given by

$$(3-3) \quad (h, y) \triangleright (f, x) = (fh, yx) \quad \text{when} \quad \eta(f, x) = t(h) = r(y).$$

It remains to prove that the core action (3-3) is proper. But the action (3-3) is free, that is the stabilizers are trivial. Thus, in order to prove that it is proper, we only need to check that the anchor map of the corresponding action groupoid

$$(s, t) : (\mathcal{V}^{op} j \times_i \mathcal{H}) \times (\mathcal{V}_{b \times l} \mathcal{H}) \rightarrow (\mathcal{V}_{b \times l} \mathcal{H}) \times (\mathcal{V}_{b \times l} \mathcal{H})$$

is closed; see Proposition 3.3. Let $A \subseteq (\mathcal{V} j \times_i \mathcal{H}) \times (\mathcal{V}_{b \times l} \mathcal{H})$ be a closed set and consider a sequence $\{(f_n, x_n, g_n, y_n)\}_{n \in \mathbb{N}}$ in A such that

$$\{(s, t)(f_n, x_n, g_n, y_n)\}_{n \in \mathbb{N}} = \{(g_n f_n, x_n y_n, g_n, y_n)\}_{n \in \mathbb{N}}$$

converges to $(a, b, g, y) \in (\mathcal{V}_{b \times l} \mathcal{H}) \times (\mathcal{V}_{b \times l} \mathcal{H})$. We need to see that $(a, b, g, y) \in (s, t)(A)$.

Clearly we have $g_n \rightarrow g$, $y_n \rightarrow y$, $g_n f_n \rightarrow a$, and $x_n y_n \rightarrow b$ as $n \rightarrow \infty$. Hence $(f_n, x_n, g_n, y_n) \rightarrow (g^{-1}a, by^{-1}, g, y)$. Since A is a closed set, we conclude that $((g^{-1}a, by^{-1}, g, y) \in A$. But $(a, b, g, y) = (s, t)(g^{-1}a, by^{-1}, g, y) \in (s, t)(A)$, hence $(s, t)(A)$ is closed. This finishes the proof of the step and of the theorem. \square

Example 3.8. Let G be a simply connected Poisson-Lie group, \mathfrak{g} its Lie algebra. It is well known that \mathfrak{g} is a Lie bialgebra; let \mathfrak{g}^* be the dual Lie algebra and let \mathfrak{d} be the corresponding Drinfeld double. Let G^* and D be simply connected Lie groups with Lie algebras \mathfrak{g}^* and \mathfrak{d} , respectively. Then the maps $G \rightarrow D$ taking u to \bar{u} and $G^* \rightarrow D$ taking $g \mapsto \bar{g}$ give rise to a symplectic double groupoid [Lu and Weinstein 1989, Theorem 3]:

$$(3-4) \quad \begin{array}{ccc} \Gamma & \begin{array}{c} \xrightarrow{l} \\ \xrightarrow{r} \end{array} & G^* \\ \begin{array}{c} \downarrow t \\ \downarrow b \end{array} & & \begin{array}{c} \downarrow t \\ \downarrow b \end{array} \\ G & \begin{array}{c} \xrightarrow{l} \\ \xrightarrow{r} \end{array} & \{*\} \end{array}$$

where $\Gamma = \{(g, u, v, h) \mid g, h \in G^*, u, v \in G, \text{ and } \bar{g}\bar{u} = \bar{v}\bar{h}\}$.

It is clear that $\mathbf{E}(\Gamma) = \{(g, u, e, e) \mid g \in G^*, u \in G, \text{ and } \bar{g}\bar{u} = e\}$, that the core action over $G \times G^*$ is given by $(g, v, e, e)(a, b) = (ag, vb)$, and that the anchor of the respective action groupoid is

$$\mathfrak{X} : \mathbf{E}(\Gamma) \times (G \times G^*) \rightarrow (G \times G^*) \times (G \times G^*), \quad ((g, v, e, e), (a, b)) = (ag, vb).$$

Arguing as in the in the proof of Step 2 of Theorem 3.7, we see that the core action is proper. Observe that $(\Gamma, G, G^*, *)$ does not satisfy, in general, the filling condition. If $\overline{G} \overline{G^*} = \overline{G^*} \overline{G}$, then Γ is a slim double Lie groupoid.

References

- [Andruskiewitsch and Natale 2005] N. Andruskiewitsch and S. Natale, “Double categories and quantum groupoids”, *Publ. Mat. Urug.* **10** (2005), 11–51. MR 2006b:16058 Zbl 1092.16021 arXiv math.QA/0308228
- [Andruskiewitsch and Natale 2006] N. Andruskiewitsch and S. Natale, “Tensor categories attached to double groupoids”, *Adv. Math.* **200**:2 (2006), 539–583. MR 2007f:18008 Zbl 1099.16016
- [Andruskiewitsch and Natale 2009] N. Andruskiewitsch and S. Natale, “The structure of double groupoids”, *J. Pure Appl. Algebra* **213**:6 (2009), 1031–1045. MR 2010e:20083 Zbl 1222.18004
- [Brown 2004] R. Brown, “Crossed complexes and homotopy groupoids as non commutative tools for higher dimensional local-to-global problems”, pp. 101–130 in *Galois theory, Hopf algebras, and semiabelian categories* (Toronto, ON, 2002), edited by G. Janelidze et al., Fields Institute Communications **43**, American Mathematical Society, Providence, RI, 2004. MR 2005f:18001 Zbl 1067.18004
- [Brown and Janelidze 2004] R. Brown and G. Janelidze, “Galois theory and a new homotopy double groupoid of a map of spaces”, *Appl. Categ. Structures* **12**:1 (2004), 63–80. MR 2005e:55016 Zbl 1050.18005
- [Brown and Mackenzie 1992] R. Brown and K. C. H. Mackenzie, “Determination of a double Lie groupoid by its core diagram”, *J. Pure Appl. Algebra* **80**:3 (1992), 237–272. MR 93g:55022 Zbl 0766.22001
- [Brown and Spencer 1976] R. Brown and C. B. Spencer, “Double groupoids and crossed modules”, *Cahiers Topologie Géom. Différentielle* **17**:4 (1976), 343–362. MR 55 #13427 Zbl 0344.18004
- [Dragulete 2007] O. M. Dragulete, *Some applications of symmetries in differential geometry and dynamical systems*, thesis, École Polytechnique Fédérale de Lausanne, 2007, Available at <http://library.epfl.ch/en/theses?nr=3937>.
- [Ehresmann 1963] C. Ehresmann, “Catégories doubles et catégories structurées”, *C. R. Acad. Sci. Paris* **256** (1963), 1198–1201. MR 27 #2539 Zbl 0132.25702
- [Higgins 2005] P. J. Higgins, “Categories and groupoids”, *Repr. Theory Appl. Categ.* **7** (2005), 1–195. MR 2005k:20137 Zbl 1087.20038
- [Lang 2002] S. Lang, *Introduction to differentiable manifolds*, 2nd ed., Springer, New York, 2002. MR 2003h:58002 Zbl 1008.57001
- [Lu and Weinstein 1989] J.-H. Lu and A. Weinstein, “Groupoïdes symplectiques doubles des groupes de Lie–Poisson”, *C. R. Acad. Sci. Paris Sér. I Math.* **309**:18 (1989), 951–954. MR 91i:58045 Zbl 0701.58025
- [Mackenzie 1992] K. C. H. Mackenzie, “Double Lie algebroids and second-order geometry, I”, *Adv. Math.* **94**:2 (1992), 180–239. MR 93f:58255 Zbl 0765.57025
- [Mackenzie 1999] K. C. H. Mackenzie, “On symplectic double groupoids and the duality of Poisson groupoids”, *Internat. J. Math.* **10**:4 (1999), 435–456. MR 2000g:58029 Zbl 0961.58008
- [Mackenzie 2000] K. C. H. Mackenzie, “Double Lie algebroids and second-order geometry, II”, *Adv. Math.* **154**:1 (2000), 46–75. MR 2002f:58025 Zbl 0971.58015

- [Mackenzie 2005] K. C. H. Mackenzie, *General theory of Lie groupoids and Lie algebroids*, London Mathematical Society Lecture Note Series **213**, Cambridge University Press, Cambridge, 2005. MR 2006k:58035 Zbl 1078.58011
- [Pradines 1977] J. Pradines, *Fibrés vectoriels doubles et calcul des jets non holonomes*, Esquisses Mathématiques **29**, Université d'Amiens U.E.R. de Mathématiques, Amiens, 1977. MR 83b:58010 Zbl 0396.53016
- [Tu 2004] J.-L. Tu, "Non-Hausdorff groupoids, proper actions and K -theory", *Doc. Math.* **9** (2004), 565–597. MR 2005h:22004 Zbl 1058.22005 arXiv math.OA/0403071

Received March 7, 2011. Revised September 13, 2011.

NICOLAS ANDRUSKIEWITSCH
FACULTAD DE MATEMÁTICAS ASTRONOMÍA Y FÍSICA
UNIVERSIDAD NACIONAL DE CÓRDOBA
FAMAF-CIEM (CONICET)
MEDINA ALLENDE S/N, CIUDAD UNIVERSITARIA
5000 CÓRDOBA
ARGENTINA
andrus@famaf.unc.edu.ar
<http://www.famaf.unc.edu.ar/~andrus/>

JESUS OCHOA ARANGO
DEPARTAMENTO DE MATEMÁTICAS
PONTIFICIA UNIVERSIDAD JAVERIANA
CRA. 7 No. 43-82 - EDIFICIO CARLOS ORTÍZ
BOGOTÁ
COLOMBIA
jesus.ochoa@javeriana.edu.co

ALEJANDRO TIRABOSCHI
FACULTAD DE MATEMÁTICAS ASTRONOMÍA Y FÍSICA
UNIVERSIDAD NACIONAL DE CÓRDOBA
FAMAF-CIEM (CONICET)
MEDINA ALLENDE S/N, CIUDAD UNIVERSITARIA
5000 CÓRDOBA
ARGENTINA
tirabo@famaf.unc.edu.ar

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 256 No. 1 March 2012

On slim double Lie groupoids	1
NICOLAS ANDRUSKIEWITSCH, JESUS OCHOA ARANGO and ALEJANDRO TIRABOSCHI	
Topological classification of quasitoric manifolds with second Betti number 2	19
SUYOUNG CHOI, SEONJEONG PARK and DONG YOUP SUH	
Refined Kato inequalities for harmonic fields on Kähler manifolds	51
DANIEL CIBOTARU and PENG ZHU	
Deformation retracts to the fat diagonal and applications to the existence of peak solutions of nonlinear elliptic equations	67
E. NORMAN DANCER, JONATHAN HILLMAN and ANGELA PISTOIA	
Descent for differential Galois theory of difference equations: confluence and q -dependence	79
LUCIA DI VIZIO and CHARLOTTE HARDOUIN	
Modulation and natural valued quiver of an algebra	105
FANG LI	
Willmore hypersurfaces with two distinct principal curvatures in \mathbb{R}^{n+1}	129
TONGZHU LI	
Variational inequality for conditional pressure on a Borel subset	151
YUAN LI, ERCAI CHEN and WEN-CHIAO CHENG	
New homotopy 4-spheres	165
DANIEL NASH	
Combinatorial constructions of three-dimensional small covers	177
YASUZO NISHIMURA	
On a theorem of Paul Yang on negatively pinched bisectonal curvature	201
AERYEONG SEO	
Orders of elements in finite quotients of Kleinian groups	211
PETER B. SHALEN	
A new algorithm for finding an l.c.r. set in certain two-sided cells	235
JIAN-YI SHI	
Addendum to the article Superconnections and parallel transport	253
FLORIN DUMITRESCU	



0030-8730(201203)256:1;1-A