

*Pacific
Journal of
Mathematics*

**ON A THEOREM OF PAUL YANG ON NEGATIVELY PINCHED
BISECTIONAL CURVATURE**

AERYEONG SEO

ON A THEOREM OF PAUL YANG ON NEGATIVELY PINCHED BISECTIONAL CURVATURE

AERYEONG SEO

We present examples of domains that do not admit any complete Kähler metric with bisectional curvature bounded between prescribed two negative constants by a modification of a method of P. Yang.

1. Introduction

Yang [1976] showed that the polydiscs of complex dimension at least two do not admit any complete Kähler metrics with their holomorphic bisectional curvature bounded between two negative constants. He also pointed out that the same argument applies to the bounded symmetric domains with rank higher than one; these domains therefore do not admit such metrics with the same curvature condition. On the other hand the Poincaré–Bergman metric of the unit ball is a complete Kähler metric with its bisectional curvature bounded between two negative constants. Thus the following question [Yau 1982] seems natural:

Which complex manifolds admit a complete Kähler metric with bisectional curvature bounded between two negative constants?

Yang's original interest was linked with a question on the holomorphic universal covering manifold of a compact Kähler manifold with negative curvature, it was conjectured that it should be biholomorphic to a bounded domain. Since such a universal cover was shown to be a Stein manifold by Greene and Wu [1971;1979] during that period, Yang's work was a natural one in the sense that he investigated the curvature of possible Kähler metrics on bounded symmetric domains. On the other hand, the question posed above seems, in its own right, to contain sufficient significance to deserve further study. Incidentally, the method developed in [Yang 1976], since it is the almost unique one known to this day, also seems worth deeper investigation.

The author was supported in part by the grant number 4.0006570 (2011, PI: Kang-Tae Kim) of The Basic Science Research Institute of Pohang University of Science and Technology, The Republic of Korea.

MSC2010: primary 32Q05; secondary 32Q15.

Keywords: bisectional curvature, negative curvature, complete Kähler metrics.

Indeed, [Seshadri and Zheng \[2008\]](#) investigated this line of thought and showed that Yang’s method works not only for domains but also for product manifolds. It occurred to us that it is meaningful to investigate whether there are manifolds which do not admit complete Kähler metrics with negative bisectional curvature that are neither biholomorphic to product nor homogeneous manifolds. In fact the main purpose of this paper is to present a modification of Yang’s method and applications, and the following theorems:

Theorem 1.1. *There exists a domain Ω in \mathbb{C}^2 satisfying the following properties:*

- (i) Ω is a bounded pseudoconvex domain with smooth boundary.
- (ii) $\text{Aut } \Omega$ does not act transitively on Ω .
- (iii) Ω is not biholomorphic to the product of complex manifolds.
- (iv) Ω does not admit complete Kähler metric with bisectional curvature bounded between two negative constants.

Theorem 1.2. *Let $\Omega = \{(z, w) \in \mathbb{C}^2 : |z| < 1, |w| < r(z)\}$, where $r : D \rightarrow \mathbb{R}$ is a smooth positive function on D such that $\frac{\partial^2}{\partial z \partial \bar{z}} \frac{1}{r(z)^2} < 0$. Then Ω cannot admit any complete Kähler metric with bisectional curvature bounded between two negative constants.*

The rest of the paper is organized as follows: [Section 2](#) introduces some basic facts and terminology. In [Section 3](#), our modification of Yang’s proof is presented with a proof of [Theorem 1.1](#). In [Section 4](#), we give yet another modification of Yang’s method and the proof of [Theorem 1.2](#).

2. Preliminary and fundamental facts

First, we introduce some facts and terminology. Let (M, J, h) be a Kähler manifold M of dimension n with a Kähler metric h and a complex structure J . The curvature tensor R on (M, J, h) is given by

$$R_{i\bar{j}k\bar{l}} = \frac{\partial^2 h_{i\bar{j}}}{\partial z_k \partial \bar{z}_l} - \sum_{\alpha, \beta=1}^n h^{\alpha\bar{\beta}} \frac{\partial h_{i\bar{\beta}}}{\partial z_k} \frac{\partial h_{\alpha\bar{j}}}{\partial \bar{z}_l}$$

in local coordinates (z_1, \dots, z_n) . The bisectional curvature $B(X, Y)$ for X, Y in $T_p M$ at $p \in M$ is given by

$$B(X, Y) = \frac{R(X, JX, Y, JY)}{h(X, X)h(Y, Y)}.$$

In terms of local coordinates,

$$B(X, Y) = -\frac{\sum_{i,j,k,l=1}^n R_{i\bar{j}k\bar{l}} X_i \bar{X}_j Y_k \bar{Y}_l}{\sum_{i,j=1}^n h_{i\bar{j}} X_i \bar{X}_j \sum_{i,j=1}^n h_{i\bar{j}} Y_i \bar{Y}_j}$$

where

$$X = \operatorname{Re} \sum_{j=1}^n X_j \frac{\partial}{\partial z_j} \quad \text{and} \quad Y = \operatorname{Re} \sum_{j=1}^n Y_j \frac{\partial}{\partial z_j}.$$

The following well known theorems will play important roles in our exposition:

Theorem 2.1 (generalization of Schwarz’s lemma [Yau 1978]). *Let (M, g) be a complete Kähler manifold with Ricci curvature bounded below by a constant $-k$ and let (N, h) be a Hermitian manifold with bisectional curvature bounded above by a negative constant $-K$. Then every holomorphic mapping $f : M \rightarrow N$ satisfies*

$$f^*h \leq \frac{k}{K}g.$$

Theorem 2.2 (generalized maximum principle [Omori 1967; Yau 1975]). *Let M be a complete Riemannian manifold with Ricci curvature bounded below. Then for every C^2 function $f : M \rightarrow \mathbb{R}$ that is bounded from above, there exists a sequence $\{p_k\}_{k=0}^\infty$ in M such that*

$$\lim_{k \rightarrow \infty} |\operatorname{grad} f(p_k)| = 0, \quad \limsup_{k \rightarrow \infty} \Delta f(p_k) \leq 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} f(p_k) = \sup_M f.$$

For $D_r = \{z \in \mathbb{C} : |z| < r\}$, denote by

$$g_r = \frac{r^2 dz d\bar{z}}{(r^2 - |z|^2)^2}$$

the Poincaré metric on D_r with curvature -4 . Let D be the unit disc and g the Poincaré metric on D . Denote by D^n the n -dimensional polydisc. Let grad_h and Δ_h be the gradient and the Laplacian with respect to the Riemannian metric h .

3. A modified proof of the theorem of Paul Yang

We begin this section by proving Yang’s theorem again, using a modification of his method; our modification lies in that we do not use integrals.

Theorem 3.1 [Yang 1976]. *The polydisc D^n ($n \geq 2$) and bounded symmetric domains with rank ≥ 2 do not admit any complete Kähler metrics with its bisectional curvature bounded between two negative constants.*

Proof. For simplicity, we will prove the result for D^2 . However, the same proof can be applied to higher-dimensional polydiscs and bounded symmetric domains with rank ≥ 2 . Suppose that there exists a complete Kähler metric h on D^2 with $-d \leq B(X, Y) \leq -c < 0$ for any X, Y , where c, d are positive constants. Fix $z \in D$ and define $i_z : D \rightarrow D^2$ by $i_z(w) = (z, w)$. By Theorem 2.1, we have $i_z^*h \leq \frac{4}{c}g$; that is,

$$h_{2\bar{2}}(z, w) \leq \frac{4}{c} \frac{1}{(1 - |w|^2)^2}.$$

Define $F : D \rightarrow \mathbb{R}$ by $F(z) = h_{2\bar{2}}(z, 0)$. Then F is a smooth, positive, bounded function. We will induce a contradiction by calculating the Laplacian of F :

$$\begin{aligned} \Delta_g F(z) &= (1 - |z|^2)^2 \frac{\partial^2 F}{\partial z \partial \bar{z}}(z) \\ &= (1 - |z|^2)^2 \left(R_{1\bar{1}2\bar{2}}(z, 0) + \sum_{\alpha, \beta=1}^2 h^{\alpha\bar{\beta}} \frac{\partial h_{2\bar{\beta}}}{\partial z} \frac{\partial h_{\alpha\bar{2}}}{\partial \bar{z}} \right) \\ &\geq c(1 - |z|^2)^2 h_{2\bar{2}}(z, 0) h_{1\bar{1}}(z, 0). \end{aligned}$$

From Schwarz’s lemma applied to $\pi : D^2 \rightarrow D$, $\pi(z, w) = z$, we get $\pi^*g \leq \frac{2d}{4}h$, that is, $(1 - |z|^2)^2 h_{1\bar{1}}(z, w) \geq 2/d$. So

$$\Delta_g F \geq \frac{2c}{d} F.$$

Since $\Delta_g \log F = \Delta_g F / |F| - |\text{grad}_g F|^2 / |F|^2$ and $\log F$ is a well defined bounded function, by the almost maximum principle ([Theorem 2.2](#)) there exists a sequence $\{p_k\}_{n=1}^\infty$ in D such that

$$\lim_{k \rightarrow \infty} F(p_k) = \sup_D F, \quad \lim_{k \rightarrow \infty} |\text{grad}_g \log F(p_k)| = 0 \quad \text{and} \quad \limsup_{k \rightarrow \infty} \Delta_g \log F(p_k) \leq 0.$$

But this contradicts $\limsup_{k \rightarrow \infty} \Delta_g \log F(p_k) \geq 2c/d$. □

Though clear, the crux of the above argument can be summarized as follows:

Proposition 3.2. *Let Ω be a domain in \mathbb{C}^n . Suppose that there exist an embedding*

$$\iota : D^2 \rightarrow \Omega, \quad \iota(z, w) = (z, w, 0, \dots, 0)$$

and a projection

$$\pi : \Omega \rightarrow D, \quad \pi(z_1, z_2, \dots, z_n) = z_1.$$

Then for any constants $c > d > 0$, there is no complete Kähler metric on Ω with

$$-c \leq \text{Bisec} \leq -d.$$

Example 3.3. Let $\Omega = \{(z, w, u) \in \mathbb{C}^2 : |z| < 1, |w| < \alpha, |z + w| < \beta\}$. If $\alpha \geq 1$ and $\beta \geq 2$, then Ω does not have any complete Kähler metrics with bisectonal curvature bounded between two negative constants. This Ω is a nonproduct and nonhomogeneous analytic polyhedron that is not biholomorphic to the bidisc: If Ω were a product manifold, Ω would be biholomorphic to D^2 by the Riemann mapping theorem. However, Ω cannot be biholomorphic to D^2 [[Fridman 1978](#)]. Likewise, Ω is not homogeneous; if it were, then it would be biholomorphic to D^2 [[Kim 1992](#)].

For a sequence of domains $\{\Omega_p\}_{p=1}^\infty$, it is possible that Ω_p does not admit any complete Kähler metrics with bisectional curvature bounded between two negative constants for all p , but that $\{\Omega_p\}$ converges to the ball as $p \rightarrow \infty$.

Corollary 3.4. *For any $0 < r < 1$ and positive constants $0 < c < d$,*

$$\Omega = \{(z, w) \in \mathbb{C} \times \mathbb{C}^n : |z|^2 + |w|^2 < 1, |z| < r\}$$

does not admit complete Kähler metric with

$$-d < \text{Bisec} < -c.$$

Proof. The mapping

$$(z, w) \mapsto \left(\frac{1}{r}z, \frac{1}{\sqrt{1-r^2}}w \right)$$

sends Ω to $\{(z, w) \in \mathbb{C} \times \mathbb{C}^n : r^2|z|^2 + (1-r^2)|w|^2 < 1, |z| < 1\}$. So an embedding and a projection like in [Proposition 3.2](#) exist. □

Remark 3.5. In contrast, the sequence of domains $\{\Omega_m\}_{m=1}^\infty$, where the domain $\Omega_m = \{(z, w) \in \mathbb{C}^n : |z|^2 + |w|^{2m} < 1\}$ has a complete Kähler metric with sectional curvature bounded between two negative constants for all m , converges to the bidisc. [Bland \[1986\]](#) proved that the Kähler–Einstein metric on Ω_m has sectional curvature bounded between two negative constants. However we could find another complete Kähler metric explicitly which has sectional curvature bounded between two negative constants.

On $\Omega_m = \{(z, w) \in \mathbb{C}^n : |z|^2 + |w|^{2m} < 1\}$, let

$$g_m(z, w) = -\partial\bar{\partial} \log \rho(z, w), \quad \text{where } \rho(z, w) = (1 - |z|^2)^{\frac{1}{m}} - |w|^2.$$

Then g_m is a complete Kähler metric on Ω_m . The metric $g_m(z, w)$ is given by

$$\frac{\frac{1}{m}(1 - |z|^2)^{\frac{1}{m}-2}}{((1 - |z|^2)^{\frac{1}{m}} - |w|^2)^2} \begin{pmatrix} (1 - |z|^2)^{\frac{1}{m}} - |w|^2 + \frac{1}{m}|z|^2|w|^2 & w\bar{z}(1 - |z|^2) \\ \bar{w}z(1 - |z|^2) & m(1 - |z|^2)^2 \end{pmatrix},$$

where

$$\det g_m = \frac{\frac{1}{m}(1 - |z|^2)^{\frac{2}{m}-2}}{((1 - |z|^2)^{\frac{1}{m}} - |w|^2)^3},$$

and the inverse $g_m^{-1}(z, w)$ is

$$\frac{(1 - |z|^2)^{\frac{1}{m}} - |w|^2}{(1 - |z|^2)^{\frac{1}{m}}} \begin{pmatrix} m(1 - |z|^2)^2 & -w\bar{z}(1 - |z|^2) \\ -\bar{w}z(1 - |z|^2) & (1 - |z|^2)^{\frac{1}{m}} - |w|^2 + \frac{1}{m}|z|^2|w|^2 \end{pmatrix}.$$

This metric g_m is invariant with respect to the automorphisms

$$f_\alpha(z, w) = \left(\frac{\alpha - z}{1 - \bar{\alpha}z}, \frac{(1 - \alpha^2)^{\frac{1}{2m}}}{(1 - \bar{\alpha}z)^{\frac{1}{m}}} w \right) \quad \text{for } \alpha \in D,$$

so it is enough to calculate the section curvature at $(0, w)$. By calculation,

$$\begin{aligned} 2mg_{1\bar{1}}^2 &> R_{1\bar{1}1\bar{1}} = 2g_{1\bar{1}}^2(|w|^2 + m(1 - |w|^2)) > 2g_{1\bar{1}}^2, \\ R_{2\bar{2}2\bar{2}} &= 2g_{2\bar{2}}^2, \\ R_{1\bar{1}2\bar{2}} &= R_{2\bar{2}1\bar{1}} = R_{1\bar{2}2\bar{1}} = R_{2\bar{1}1\bar{2}} = g_{1\bar{1}}g_{2\bar{2}}, \end{aligned}$$

and the other $R_{i\bar{j}k\bar{l}}$ are zero at $(0, w)$. This implies that

$$-2m < \text{sectional curvature of } \Omega_m < -\frac{1}{2}.$$

However the lower bound on the sectional curvature of this metric and of the Kähler–Einstein metric depends on m and goes to $-\infty$ as $m \rightarrow \infty$. So this question is still open:

If Ω_m has a complete Kähler metric with $-d < (bi)sect < -c$ for some constants $c, d > 0$ for all m and converges to Ω as $m \rightarrow \infty$, then does Ω have a complete Kähler metric with $-d < (bi)sect < -c$?

Proof of Theorem 1.1. For $\epsilon > 0, \alpha < 1$, let

$$(3-1) \quad \Omega_\epsilon = \left\{ (z, w) \in \mathbb{C}^2 : |z|^2 + \alpha e^{-1/(|w|^2 - \epsilon)} < 1, |w| > 1/\sqrt{\epsilon} \right\} \\ \cup \left\{ (z, w) \in \mathbb{C}^2 : |z| < 1, |w| \leq 1/\sqrt{\epsilon} \right\}$$

Then Ω_ϵ is a nonhomogeneous, nonproduct, bounded pseudoconvex domain with smooth boundary. The points $(z, w) \in \partial\Omega$ with $|w|^2 > \epsilon$ are strongly pseudoconvex. By dilating $(z, w) \mapsto (z, w/\sqrt{\epsilon})$, we see that Ω_ϵ is biholomorphic to

$$\left\{ (z, w) \in \mathbb{C}^2 : |z|^2 + \alpha^2 \exp(1 - 1/(\epsilon|w|^2 - \epsilon)) < 1, |z| < 1 \right\} \cup D^2.$$

Therefore we can apply [Proposition 3.2](#). □

4. Further modification

We now find more examples of domain that do not admit complete Kähler metric with bisectional curvature bounded between two negative constants. The domains we look for will be different from analytic polyhedra. We will achieve this by modifying Yang’s method one more time.

Proof of Theorem 1.2. Suppose that there exists a complete Kähler metric h on Ω with $-d \leq B(X, Y) \leq -c < 0$ for any X, Y for some positive constants c, d .

Define $i_z : D_{r(z)} \rightarrow \Omega$ by $i_z(w) = (z, w)$. By Schwarz's lemma, $i_z^*h \leq \frac{4}{c}g_{r(z)}$. So $h_{2\bar{2}}(z, w) \leq \frac{4}{c}r(z)^2/(r(z)^2 - |w|^2)^2$. Define $F : D \rightarrow \mathbb{R}$ by

$$F(z) = r(z)^2 h_{2\bar{2}}(z, 0).$$

Then F is a smooth, positive, bounded function. We have

$$\frac{\partial F}{\partial z}(z) = \frac{\partial r^2}{\partial z}(z) h_{2\bar{2}}(z, 0) + r(z)^2 \frac{\partial h_{2\bar{2}}}{\partial z}(z, 0)$$

and

$$\begin{aligned} \frac{\partial^2 F}{\partial z \partial \bar{z}} &= r^2 \frac{\partial^2 h_{2\bar{2}}}{\partial z \partial \bar{z}} + \frac{\partial^2 r^2}{\partial z \partial \bar{z}} h_{2\bar{2}} + 2 \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{\partial h_{2\bar{2}}}{\partial \bar{z}} \right) \\ &= r^2 \frac{\partial^2 h_{2\bar{2}}}{\partial z \partial \bar{z}} + \frac{\partial^2 r^2}{\partial z \partial \bar{z}} h_{2\bar{2}} + 2 \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{1}{r^2} \left(\frac{\partial F}{\partial \bar{z}} - \frac{\partial r^2}{\partial \bar{z}} h_{2\bar{2}} \right) \right) \\ &= r^2 \frac{\partial^2 h_{2\bar{2}}}{\partial z \partial \bar{z}} + \left(\frac{\partial^2 r^2}{\partial z \partial \bar{z}} - \frac{2}{r^2} \left| \frac{\partial r^2}{\partial z} \right|^2 \right) h_{2\bar{2}} + \frac{2}{r^2} \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{\partial F}{\partial \bar{z}} \right) \\ &= r^2 \frac{\partial^2 h_{2\bar{2}}}{\partial z \partial \bar{z}} - r^4 \frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} h_{2\bar{2}} + \frac{2}{r^2} \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{\partial F}{\partial \bar{z}} \right). \end{aligned}$$

Thus

$$\begin{aligned} \Delta_g F &\geq c(1 - |z|^2)^2 r^2 h_{2\bar{2}} h_{1\bar{1}} - (1 - |z|^2)^2 r^2 \frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} F + \frac{2(1 - |z|^2)^2}{r^2} \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{\partial F}{\partial \bar{z}} \right) \\ &\geq \frac{2c}{d} F - (1 - |z|^2)^2 r^2 \frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} F + \frac{2(1 - |z|^2)^2}{r^2} \operatorname{Re} \left(\frac{\partial r^2}{\partial z} \frac{\partial F}{\partial \bar{z}} \right). \end{aligned}$$

So if r satisfies $\frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} < 0$, then the desired result follows by the almost maximum principle as in the previous modification of Yang's proof. \square

Corollary 4.1. *Let Ω is the domain in the [Theorem 1.2](#). Suppose $1/r(z)^2 = \rho(|z|)$ for some positive function ρ on $[0, 1]$ and suppose*

$$t \frac{d\rho}{dt}(t)$$

is a decreasing function. Then Ω does not admit any complete Kähler metrics with bisectional curvature bounded between any two prescribed negative constants.

Proof. This follows by

$$\frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} = \frac{d^2 \rho}{dt^2} + \frac{1}{t} \frac{d\rho}{dt} = \frac{1}{t} \left(\frac{d}{dt} \left(t \frac{d\rho}{dt} \right) \right). \quad \square$$

Typical examples of ρ and Ω satisfying the conditions in [Corollary 4.1](#) are:

Example 4.2. Let $\rho(t) = \exp(-t)$ and

$$\Omega = \{(z, w) \in \mathbb{C}^2 : |z| < 1, |w|^2 < \exp(|z|)\}.$$

Example 4.3. Let $\rho(t) = \alpha - t^2$ and

$$\Omega = \{(z, w) \in \mathbb{C}^2 : |z| < 1, |w|^2(\alpha - |z|^2) < 1\}.$$

If we take $\alpha > 1$, then Ω is a bounded domain.

Remark 4.4. If

$$\frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} < 0 \quad \text{then} \quad \frac{\partial^2 r^2}{\partial z \partial \bar{z}} > \frac{2}{r^2} \left| \frac{\partial r^2}{\partial z} \right|^2.$$

Notice that the domains given in [Theorem 1.2](#) are in fact pseudoconcave.

The domains in [Theorem 1.2](#) converge to $\{(z, w) \in \mathbb{C}^2 : |z|^2 + \alpha \exp(\frac{-1}{|w|^2}) < 1\}$ as $\epsilon \rightarrow 0$, which does not admit a complete Kähler metric with $-d < \text{Bisec} < -c$ for some constants $c, d > 0$:

Corollary 4.5. *Let $\Omega = \{(z, w) \in \mathbb{C}^2 : |z|^2 + \alpha \exp(\frac{-1}{|w|^2}) < 1\}$, for some constant $\alpha > 1$. For any positive constants $c < d$, such that $c/d > (2 \log \alpha)^{-1}$, Ω cannot admit a complete Kähler metric with*

$$-d \leq \text{Bisec} \leq -c.$$

Proof. Notice that

$$\Omega = \{(z, w) \in \mathbb{C}^2 : |z| < 1, |w| < r(z)\}$$

where $r(z) = 1/(\log \alpha - \log(1 - |z|^2))^{1/2}$. Since $\frac{\partial^2 r^{-2}}{\partial z \partial \bar{z}} = \frac{1}{(1 - |z|^2)^2}$, we obtain for

$$F : D \rightarrow \mathbb{R}, \quad F(z) = r(z)^2 h_{2\bar{2}}(z, 0)$$

the inequality

$$\Delta_g F(z) \geq \frac{2c}{d} F(z) - \frac{F}{\log \alpha - \log(1 - |z|^2)} - \frac{2(1 - |z|^2)}{\log \alpha - \log(1 - |z|^2)} \text{Re} \left(\bar{z} \frac{\partial F}{\partial \bar{z}}(z) \right).$$

Since

$$0 < \frac{2(1 - |z|^2)}{\log \alpha - \log(1 - |z|^2)} \leq \frac{1}{\log \alpha}$$

and $\lim_{|z| \rightarrow 1} 1/(\log \alpha - \log(1 - |z|^2)) = 0$, for $c/d > (2 \log \alpha)^{-1}$, the desired result follows by the almost maximum principle. □

Acknowledgements

This work is part of the author’s Ph.D. dissertation at POSTECH. She would like to express her gratitude to her advisor Professor Kang-Tae Kim for his guidance.

References

- [Bland 1986] J. S. Bland, “The Einstein–Kähler metric on $\{|z|^2 + |w|^{2p} < 1\}$ ”, *Michigan Math. J.* **33**:2 (1986), 209–220. [MR 87i:32036](#) [Zbl 0602.32005](#)
- [Fridman 1978] B. L. Fridman, “A class of analytic polyhedra”, *Dokl. Akad. Nauk SSSR* **242**:5 (1978), 1020–1022. In Russian; translated in *Soviet Math. Dokl.* **19**:5 (1979), 1258–1261. [MR 80b:32014](#) [Zbl 0427.32013](#)
- [Greene and Wu 1971] R. E. Greene and H. Wu, “Curvature and complex analysis”, *Bull. Amer. Math. Soc.* **77** (1971), 1045–1049. [MR 44 #473](#) [Zbl 0225.32010](#)
- [Greene and Wu 1979] R. E. Greene and H. Wu, *Function theory on manifolds which possess a pole*, Lecture Notes in Mathematics **699**, Springer, Berlin, 1979. [MR 81a:53002](#) [Zbl 0414.53043](#)
- [Kim 1992] K.-T. Kim, “Domains in \mathbb{C}^n with a piecewise Levi flat boundary which possess a non-compact automorphism group”, *Math. Ann.* **292** (1992), 575–586. [MR 93h:32024](#) [Zbl 0735.32021](#)
- [Omori 1967] H. Omori, “Isometric immersions of Riemannian manifolds”, *J. Math. Soc. Japan* **19** (1967), 205–214. [MR 35 #6101](#) [Zbl 0154.21501](#)
- [Seshadri and Zheng 2008] H. Seshadri and F. Zheng, “Complex product manifolds cannot be negatively curved”, *Asian J. Math.* **12**:1 (2008), 145–149. [MR 2009d:53047](#) [Zbl 1147.53317](#)
- [Yang 1976] P. C. Yang, “On Kähler manifolds with negative holomorphic bisectional curvature”, *Duke Math. J.* **43**:4 (1976), 871–874. [MR 54 #7837](#) [Zbl 0358.32005](#)
- [Yau 1975] S. T. Yau, “Harmonic functions on complete Riemannian manifolds”, *Comm. Pure Appl. Math.* **28** (1975), 201–228. [MR 55 #4042](#) [Zbl 0291.31002](#)
- [Yau 1978] S. T. Yau, “A general Schwarz lemma for Kähler manifolds”, *Amer. J. Math.* **100**:1 (1978), 197–203. [MR 58 #6370](#) [Zbl 0424.53040](#)
- [Yau 1982] S. T. Yau (editor), *Seminar on Differential Geometry*, Annals of Mathematics Studies **102**, Princeton Univ. Press, 1982. [MR 83a:53002](#)

Received September 17, 2011. Revised December 5, 2011.

AERYEONG SEO
DEPARTMENT OF MATHEMATICS
POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH)
POHANG 790-784
SOUTH KOREA
inno827@postech.ac.kr

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by

E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Darren Long
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
long@math.ucsb.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Jonathan Rogawski
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
jonr@math.ucla.edu

PRODUCTION

pacific@math.berkeley.edu

Silvio Levy, Scientific Editor

Mathew Cargo, Senior Production Editor

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 256 No. 1 March 2012

On slim double Lie groupoids	1
NICOLAS ANDRUSKIEWITSCH, JESUS OCHOA ARANGO and ALEJANDRO TIRABOSCHI	
Topological classification of quasitoric manifolds with second Betti number 2	19
SUYOUNG CHOI, SEONJEONG PARK and DONG YOUP SUH	
Refined Kato inequalities for harmonic fields on Kähler manifolds	51
DANIEL CIBOTARU and PENG ZHU	
Deformation retracts to the fat diagonal and applications to the existence of peak solutions of nonlinear elliptic equations	67
E. NORMAN DANCER, JONATHAN HILLMAN and ANGELA PISTOIA	
Descent for differential Galois theory of difference equations: confluence and q-dependence	79
LUCIA DI VIZIO and CHARLOTTE HARDOUIN	
Modulation and natural valued quiver of an algebra	105
FANG LI	
Willmore hypersurfaces with two distinct principal curvatures in \mathbb{R}^{n+1}	129
TONGZHU LI	
Variational inequality for conditional pressure on a Borel subset	151
YUAN LI, ERCAI CHEN and WEN-CHIAO CHENG	
New homotopy 4-spheres	165
DANIEL NASH	
Combinatorial constructions of three-dimensional small covers	177
YASUZO NISHIMURA	
On a theorem of Paul Yang on negatively pinched bisectional curvature	201
AERYEONG SEO	
Orders of elements in finite quotients of Kleinian groups	211
PETER B. SHALEN	
A new algorithm for finding an l.c.r. set in certain two-sided cells	235
JIAN-YI SHI	
Addendum to the article Superconnections and parallel transport	253
FLORIN DUMITRESCU	