

*Pacific  
Journal of  
Mathematics*

**A NOTE ON INVERSE CURVATURE FLOWS IN  
ASYMPTOTICALLY ROBERTSON–WALKER SPACETIMES**

CLAUS GERHARDT

Volume 256    No. 2

April 2012

## A NOTE ON INVERSE CURVATURE FLOWS IN ASYMPTOTICALLY ROBERTSON–WALKER SPACETIMES

CLAUS GERHARDT

**We prove that the leaves of the rescaled curvature flow considered in earlier work converge to the graph of a constant function.**

### 1. Introduction

In [Gerhardt 2004] and [Gerhardt 2006a, Chapter 7] we considered the inverse mean curvature flow in a Lorentzian manifold  $N = N^{n+1}$  which we called an asymptotically Robertson–Walker space, and which is defined by the following conditions:

**Definition 1.1.** A cosmological spacetime  $N$ ,  $\dim N = n + 1$ , is said to be *asymptotically Robertson–Walker (ARW)* with respect to the future, if a future end of  $N$ ,  $N_+$ , can be written as a product  $N_+ = [a, b) \times \mathcal{S}_0$ , where  $\mathcal{S}_0$  is a compact Riemannian space, and there exists a future directed time function  $\tau = x^0$  such that the metric in  $N_+$  can be written as

$$(1-1) \quad d\tilde{s}^2 = e^{2\tilde{\psi}} \left\{ -(dx^0)^2 + \sigma_{ij}(x^0, x) dx^i dx^j \right\},$$

where  $\mathcal{S}_0$  corresponds to  $x^0 = a$ ,  $\tilde{\psi}$  is of the form

$$(1-2) \quad \tilde{\psi}(x^0, x) = f(x^0) + \psi(x^0, x),$$

and we assume that there exists a positive constant  $c_0$  and a smooth Riemannian metric  $\bar{\sigma}_{ij}$  on  $\mathcal{S}_0$  such that

$$(1-3) \quad \lim_{\tau \rightarrow b} e^{\psi} = c_0 \quad \text{and} \quad \lim_{\tau \rightarrow b} \sigma_{ij}(\tau, x) = \bar{\sigma}_{ij}(x),$$

and

$$(1-4) \quad \lim_{\tau \rightarrow b} f(\tau) = -\infty.$$

---

This work was supported by the DFG.

MSC2010: 35J60, 53C21, 53C44, 53C50, 58J05.

*Keywords:* Lorentzian manifold, mass, cosmological spacetime, general relativity, inverse curvature flow, ARW spacetimes.

Without loss of generality we shall assume  $c_0 = 1$ . Then  $N$  is ARW with respect to the future if the metric is close to the Robertson–Walker metric

$$(1-5) \quad d\bar{s}^2 = e^{2f} \{ -(dx^0)^2 + \bar{\sigma}_{ij}(x) dx^i dx^j \}$$

near the singularity  $\tau = b$ . By *close* we mean that the derivatives of arbitrary order with respect to space and time of the conformal metric  $e^{-2f} \check{g}_{\alpha\beta}$  in (1-1) should converge to the corresponding derivatives of the conformal limit metric in (1-5) when  $x^0$  tends to  $b$ . We emphasize that in our terminology Robertson–Walker metric does not imply that  $(\bar{\sigma}_{ij})$  is a metric of constant curvature, it is only the spatial metric of a warped product.

We assume, furthermore, that  $f$  satisfies the following five conditions:

$$(1-6) \quad -f' > 0.$$

There exists  $\omega \in \mathbb{R}$  such that

$$(1-7) \quad n + \omega - 2 > 0 \quad \text{and} \quad \lim_{\tau \rightarrow b} |f'|^2 e^{(n+\omega-2)f} = m > 0.$$

Set  $\tilde{\gamma} = \frac{1}{2}(n + \omega - 2)$ , then the limit

$$(1-8) \quad \lim_{\tau \rightarrow b} (f'' + \tilde{\gamma}|f'|^2)$$

exists and

$$(1-9) \quad |D_\tau^m (f'' + \tilde{\gamma}|f'|^2)| \leq c_m |f'|^m \quad \text{for all } m \geq 1,$$

as well as

$$(1-10) \quad |D_\tau^m f| \leq c_m |f'|^m \quad \text{for all } m \geq 1.$$

We call  $N$  a *normalized* ARW spacetime if

$$(1-11) \quad \int_{g_0} \sqrt{\det \bar{\sigma}_{ij}} = |S^n|.$$

**Remark 1.2.** (i) If these assumptions are satisfied, then we proved in [Gerhardt 2004] that the range of  $\tau$  is finite, hence, we shall assume without loss of generality that  $b = 0$ , that is,

$$(1-12) \quad a < \tau < 0.$$

(ii) Any ARW spacetime can be normalized as one easily checks. For normalized ARW spaces the constant  $m$  in (1-7) is defined uniquely and can be identified with the mass of  $N$ , see [Gerhardt 2006b].

(iii) In view of the assumptions on  $f$  the mean curvature of the coordinate slices  $M_\tau = \{x^0 = \tau\}$  tends to  $\infty$  if  $\tau$  goes to zero.

(iv) ARW spaces satisfy a strong volume decay condition, see [Gerhardt 2008, Definition 0.1].

(v) Similarly one can define  $N$  to be ARW with respect to the past. In this case the singularity would lie in the past, correspond to  $\tau = 0$ , and the mean curvature of the coordinate slices would tend to  $-\infty$ .

We assume that  $N$  satisfies the timelike convergence condition. Consider the future end  $N_+$  of  $N$  and let  $M_0 \subset N_+$  be a spacelike hypersurface with positive mean curvature  $\check{H}|_{M_0} > 0$  with respect to the past directed normal vector  $\check{\nu}$ —we shall explain in Section 2 why we use the symbols  $\check{H}$  and  $\check{\nu}$  and not the usual ones  $H$  and  $\nu$ . Then, as we have proved in [Gerhardt 2008], the inverse mean curvature flow

$$(1-13) \quad \dot{x} = -\check{H}^{-1} \check{\nu}$$

with initial hypersurface  $M_0$  exists for all time, is smooth, and runs straight into the future singularity.

If we express the flow hypersurfaces  $M(t)$  as graphs over  $\mathcal{S}_0$

$$(1-14) \quad M(t) = \text{graph } u(t, \cdot),$$

then one of the main results in our former paper was:

**Theorem 1.3.** (i) *Let  $N$  satisfy the above assumptions, then the range of the time function  $x^0$  is finite, that is, we may assume that  $b = 0$ . Set*

$$(1-15) \quad \tilde{u} = ue^{\gamma t},$$

where  $\gamma = \frac{1}{n}\check{\gamma}$ , then there are positive constants  $c_1, c_2$  such that

$$(1-16) \quad -c_2 \leq \tilde{u} \leq -c_1 < 0,$$

and  $\tilde{u}$  converges in  $C^\infty(\mathcal{S}_0)$  to a smooth function, if  $t$  goes to infinity. We shall also denote the limit function by  $\tilde{u}$ .

(ii) *Let  $\check{g}_{ij}$  be the induced metric of the leaves  $M(t)$ , then the rescaled metric*

$$(1-17) \quad e^{\frac{2}{n}t} \check{g}_{ij}$$

converges in  $C^\infty(\mathcal{S}_0)$  to

$$(1-18) \quad (\check{\gamma}^2 m)^{\frac{1}{\check{\nu}}} (-\tilde{u})^{\frac{2}{\check{\nu}}} \bar{\sigma}_{ij}.$$

(iii) *The leaves  $M(t)$  get more umbilical if  $t$  tends to infinity, namely,*

$$(1-19) \quad \check{H}^{-1} \left| \check{h}_i^j - \frac{1}{n} \check{H} \delta_i^j \right| \leq ce^{-2\gamma t}.$$

In case  $n + \omega - 4 > 0$ , we even get a better estimate

$$(1-20) \quad |\check{h}_i^j - \frac{1}{n} \check{H} \delta_i^j| \leq c e^{-\frac{1}{2n}(n+\omega-4)t}.$$

The results for the mean curvature flow have recently also been proved for other inverse curvature flows, where the mean curvature is replaced by a curvature function  $F$  of class  $(K^*)$  homogeneous of degree 1, which includes the  $n$ -th root of the Gaussian curvature, see Kröner [2011].

In this note we want to prove that the functions in (1-15) converge to a constant. This result will also be valid when, instead of the mean curvature, other curvature functions  $F$  homogeneous of degree one will be considered satisfying

$$(1-21) \quad F(1, \dots, 1) = n$$

provided the rescaled functions in (1-15) can be estimated as in (1-16) and converge in  $C^3(\mathcal{F}_0)$ . For simplicity we shall formulate the result only for the solution in Theorem 1.3, but it will be apparent from the proof that the result is also valid for different curvature functions.

**Theorem 1.4.** *The functions  $\tilde{u}$  in (1-15) converge to a constant.*

### 2. Proof of Theorem 1.4

When we proved the convergence results for the inverse mean curvature flow in [Gerhardt 2004], we considered the flow hypersurfaces to be embedded in  $N$  equipped with the conformal metric

$$(2-1) \quad d\tilde{s}^2 = -(dx^0)^2 + \sigma_{ij}(x^0, x) dx^i dx^j.$$

Though, formally, we have a different ambient space we still denote it by the same symbol  $N$  and distinguish only the metrics  $\check{g}_{\alpha\beta}$  and  $\bar{g}_{\alpha\beta}$

$$(2-2) \quad \check{g}_{\alpha\beta} = e^{2\check{\psi}} \bar{g}_{\alpha\beta}$$

and the corresponding geometric quantities of the hypersurfaces  $\check{h}_{ij}$ ,  $\check{g}_{ij}$ ,  $\check{\nu}$ , respectively  $h_{ij}$ ,  $g_{ij}$ ,  $\nu$ , and so on.

The second fundamental forms  $\check{h}_i^j$  and  $h_i^j$  are related by

$$(2-3) \quad e^{\check{\psi}} \check{h}_i^j = h_i^j + \check{\psi}_{\alpha} \nu^{\alpha} \delta_i^j$$

and, if we define  $F$  by

$$(2-4) \quad F = e^{\check{\psi}} \check{H},$$

then

$$(2-5) \quad F = H - n\tilde{\nu} f' + n\psi_{\alpha} \nu^{\alpha},$$

where

$$(2-6) \quad \tilde{v} = v^{-1},$$

and

$$(2-7) \quad v^2 = 1 - \sigma^{ij} u_i u_j \equiv 1 - |Du|^2.$$

The evolution equation can be written as

$$(2-8) \quad \dot{x} = -F^{-1}v,$$

since

$$(2-9) \quad \check{v} = e^{-\tilde{\psi}} v.$$

The flow (2-8) can also be considered to comprise more general curvature functions  $F$  by assuming that  $F = F(\check{h}_j^i)$ , where  $\check{h}_j^i$  is an abbreviation for the right-hand side of (2-3). Stipulating that indices of tensors will be raised or lowered with the help of the metric

$$(2-10) \quad g_{ij} = -u_i u_j + \sigma_{ij},$$

we may also consider  $F$  to depend on

$$(2-11) \quad \check{h}_{ij} = h_{ij} - \tilde{v} f' g_{ij} + \psi_\alpha v^\alpha g_{ij}$$

and we define accordingly

$$(2-12) \quad F^{ij} = \frac{\partial F}{\partial \check{h}_{ij}}.$$

Now, let us prove Theorem 1.4. We use the relation

$$(2-13) \quad \tilde{v}^2 = 1 + \|Du\|^2 = 1 + g^{ij} u_i u_j$$

and shall prove that

$$(2-14) \quad \lim_{t \rightarrow \infty} (\|Du\|^2)' e^{2\gamma t} = 2\gamma \Delta \tilde{u} \tilde{u},$$

where

$$(2-15) \quad \tilde{u} = \lim_{t \rightarrow \infty} u e^{\gamma t},$$

as well as

$$(2-16) \quad \lim_{t \rightarrow \infty} (\tilde{v}^2)' e^{2\gamma t} = -2\gamma \|D\tilde{u}\|^2$$

yielding

$$(2-17) \quad -\Delta \tilde{u} \tilde{u} = \|D\tilde{u}\|^2$$

on the compact limit hypersurface  $M$ . Since  $\tilde{u}$  is strictly negative we then conclude

$$(2-18) \quad \int_M \|D\tilde{u}\|^2 \tilde{u}^{-1} = 0,$$

hence  $\|D\tilde{u}\| = 0$ .

Let us first derive (2-14). Using

$$(2-19) \quad \dot{g}_{ij} = -2F^{-1}h_{ij},$$

see [Gerhardt 2006a, Lemma 2.3.1], where we write  $g_{ij} = g_{ij}(t, \xi)$ ,  $\xi = (\xi^i)$  are local coordinates for  $\mathcal{S}_0$ , and where

$$(2-20) \quad \dot{g}_{ij} = \frac{\partial g_{ij}}{\partial t} = \dot{u}_i u_j + u_i \dot{u}_j + \dot{\sigma}_{ij} \dot{u},$$

and  $\dot{\sigma}_{ij}$  is defined by

$$(2-21) \quad \dot{\sigma}_{ij} = \frac{\partial \sigma_{ij}}{\partial u},$$

we deduce

$$(2-22) \quad \begin{aligned} (\|Du\|^2)' &= (g^{ij}u_i u_j)' = 2g^{ij}\dot{u}_i u_j - \dot{g}_{ij}u^i u^j \\ &= 2F^{-1}H + g^{ij}\dot{\sigma}_{ij}\dot{u} - \dot{g}_{ij}u^i u^j \\ &= 2F^{-1}H + \tilde{v}F^{-1}g^{ij}\dot{\sigma}_{ij} + 2F^{-1}h_{ij}u^i u^j \\ &= 2F^{-1}H + \tilde{v}F^{-1}\sigma^{ij}\dot{\sigma}_{ij} + \tilde{v}^3 F^{-1}\dot{\sigma}_{ij}\check{u}^i \check{u}^j + 2F^{-1}h_{ij}u^i u^j, \end{aligned}$$

where we used the relation

$$(2-23) \quad g^{ij} = \sigma^{ij} + \tilde{v}^2 \check{u}^i \check{u}^j$$

and where  $\check{u}^i$  is defined by

$$(2-24) \quad \check{u}^i = \sigma^{ij}u_j.$$

The last two terms on the right-hand side of (2-22) are an  $o(e^{-2\gamma t})$ , thus we have

$$(2-25) \quad (\|Du\|^2)' = 2F^{-1}(H + \tilde{v}\frac{1}{2}\sigma^{ij}\dot{\sigma}_{ij}) + o(e^{-2\gamma t}).$$

On the other hand,

$$(2-26) \quad h_{ij}\tilde{v} = -u_{ij} + \bar{h}_{ij},$$

where  $\bar{h}_{ij}$  is the second fundamental form of the slices  $\{x^0 = \text{const}\}$

$$(2-27) \quad \bar{h}_{ij} = -\frac{1}{2}\dot{\sigma}_{ij}$$

and we infer

$$(2-28) \quad H\tilde{v} = -\Delta u + g^{ij}\bar{h}_{ij} = -\Delta u + \bar{H} + \tilde{v}^2\bar{h}_{ij}\tilde{u}^i\tilde{u}^j.$$

Combining (2-22), (2-27) and (2-28) we obtain

$$(2-29) \quad \begin{aligned} (\|Du\|^2)' &= 2F^{-1}(H - \tilde{v}\bar{H}) + o(e^{-2\gamma t}) \\ &= 2F^{-1}(H - \bar{H}) + o(e^{-2\gamma t}) \\ &= -2F^{-1}\Delta u + o(e^{-2\gamma t}). \end{aligned}$$

In view of [Gerhardt 2006a, Lemma 7.3.4], the estimates for  $h_{ij}$ ,  $u$ , and  $\psi$ , and the homogeneity of  $F$ , we have

$$(2-30) \quad \lim_{t \rightarrow \infty} F(-u) = n\tilde{\gamma}^{-1} = \gamma^{-1},$$

hence we deduce

$$(2-31) \quad \lim_{t \rightarrow \infty} (\|Du\|^2)' e^{2\gamma t} = 2\gamma \Delta \tilde{u} \tilde{u}.$$

Let us now differentiate  $\tilde{v}^2$ . From the relation

$$(2-32) \quad \tilde{v} = \eta_\alpha v^\alpha, \quad (\eta_\alpha) = (-1, 0, \dots, 0),$$

we infer

$$(2-33) \quad \dot{\tilde{v}} = \eta_{\alpha\beta} v^\alpha \dot{x}^\beta + \eta_\alpha \dot{v}^\alpha = -F^{-1}\eta_{\alpha\beta} v^\alpha v^\beta + (F^{-1})_k u^k,$$

where we used

$$(2-34) \quad \dot{v} = (-F^{-1})^k x_k,$$

see [Gerhardt 2006a, Lemma 2.3.2]. The first term on the right-hand side of (2-33) is an  $o(e^{-2\gamma t})$  in view of the asymptotic behavior of an ARW space, see the definition of *close* in Definition 1.1, while

$$(2-35) \quad \begin{aligned} (F^{-1})_k &= -F^{-2}F^{ij} \{ h_{ij;k} - \tilde{v}_k f' g_{ij} - \tilde{f}'' u_k g_{ij} + \psi_{\alpha\beta} v^\alpha x_k^\beta g_{ij} + \psi_\alpha x_l^\alpha h_k^l g_{ij} \}, \end{aligned}$$

where we applied the Weingarten equation to derive the last term on the right-hand side. Therefore, we infer

$$(2-36) \quad \lim_{t \rightarrow \infty} (F^{-1})_k u^k e^{2\gamma t} = \|D\tilde{u}\|^2 \frac{1}{n} \lim_{t \rightarrow \infty} \frac{f''}{|f'|^2} = -\frac{\tilde{\gamma}}{n} \|D\tilde{u}\|^2 = -\gamma \|D\tilde{u}\|^2,$$

in view of (1-8) and the definition of  $\gamma$  in Theorem 1.3, and we deduce further

$$(2-37) \quad \lim_{t \rightarrow \infty} (\tilde{v}^2)' e^{2\gamma t} = -2\gamma \|D\tilde{u}\|^2,$$



hence the limit function  $\tilde{u}$  satisfies

$$(2-38) \quad \|D\tilde{u}\|^2 = -\Delta\tilde{u}\tilde{u}$$

completing the proof of Theorem 1.4.

**Remark 2.1.** We believe that this method of proof will also work for other curvature flows driven by extrinsic curvatures, in Riemannian or Lorentzian manifolds, to prove that the leaves of the rescaled curvature flows converge to the graph of a constant function.

Indeed, applying this method we proved in [Gerhardt 2011, Lemma 6.12] that the rescaled curvature flow converges to a sphere.

### References

- [Gerhardt 2004] C. Gerhardt, “The inverse mean curvature flow in ARW spaces—transition from big crunch to big bang”, preprint, 2004. arXiv math/0403485
- [Gerhardt 2006a] C. Gerhardt, *Curvature problems*, Series in Geometry and Topology **39**, International Press, Somerville, MA, 2006. MR 2007j:53001 Zbl 1131.53001
- [Gerhardt 2006b] C. Gerhardt, “The mass of a Lorentzian manifold”, *Adv. Theor. Math. Phys.* **10**:1 (2006), 33–48. MR 2006m:53109 Zbl 1104.83019
- [Gerhardt 2008] C. Gerhardt, “The inverse mean curvature flow in cosmological spacetimes”, *Adv. Theor. Math. Phys.* **12**:6 (2008), 1183–1207. MR 2009i:53059 Zbl 1153.83016
- [Gerhardt 2011] C. Gerhardt, “Inverse curvature flows in hyperbolic space”, *J. Diff. Geom.* **89**:3 (2011), 487–527. arXiv 1101.2578
- [Kröner 2011] H. Kröner, “The inverse  $F$ -curvature flow in ARW spaces”, preprint, 2011. arXiv 1106.4703

Received July 5, 2011. Revised July 6, 2011.

CLAUS GERHARDT  
 INSTITUTE OF APPLIED MATHEMATICS  
 RUPRECHT-KARLS-UNIVERSITÄT HEIDELBERG  
 IM NEUENHEIMER FELD 294  
 D-69120 HEIDELBERG  
 GERMANY

gerhardt@math.uni-heidelberg.de  
<http://www.math.uni-heidelberg.de/studinfo/gerhardt/>

# PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by  
E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

## EDITORS

V. S. Varadarajan (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[pacific@math.ucla.edu](mailto:pacific@math.ucla.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Darren Long  
Department of Mathematics  
University of California  
Santa Barbara, CA 93106-3080  
[long@math.ucsb.edu](mailto:long@math.ucsb.edu)

Sorin Popa  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[popa@math.ucla.edu](mailto:popa@math.ucla.edu)

Robert Finn  
Department of Mathematics  
Stanford University  
Stanford, CA 94305-2125  
[finn@math.stanford.edu](mailto:finn@math.stanford.edu)

Jiang-Hua Lu  
Department of Mathematics  
The University of Hong Kong  
Pokfulam Rd., Hong Kong  
[jhlu@maths.hku.hk](mailto:jhlu@maths.hku.hk)

Jie Qing  
Department of Mathematics  
University of California  
Santa Cruz, CA 95064  
[qing@cats.ucsc.edu](mailto:qing@cats.ucsc.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Alexander Merkurjev  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[merkurev@math.ucla.edu](mailto:merkurev@math.ucla.edu)

Jonathan Rogawski  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[jonr@math.ucla.edu](mailto:jonr@math.ucla.edu)

## PRODUCTION

[pacific@math.berkeley.edu](mailto:pacific@math.berkeley.edu)

Silvio Levy, Scientific Editor

Matthew Cargo, Senior Production Editor

## SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI  
CALIFORNIA INST. OF TECHNOLOGY  
INST. DE MATEMÁTICA PURA E APLICADA  
KEIO UNIVERSITY  
MATH. SCIENCES RESEARCH INSTITUTE  
NEW MEXICO STATE UNIV.  
OREGON STATE UNIV.

STANFORD UNIVERSITY  
UNIV. OF BRITISH COLUMBIA  
UNIV. OF CALIFORNIA, BERKELEY  
UNIV. OF CALIFORNIA, DAVIS  
UNIV. OF CALIFORNIA, LOS ANGELES  
UNIV. OF CALIFORNIA, RIVERSIDE  
UNIV. OF CALIFORNIA, SAN DIEGO  
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ  
UNIV. OF MONTANA  
UNIV. OF OREGON  
UNIV. OF SOUTHERN CALIFORNIA  
UNIV. OF UTAH  
UNIV. OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

See inside back cover or [pacificmath.org](http://pacificmath.org) for submission instructions.

---

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

---

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS

at the University of California, Berkeley 94720-3840

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2012 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 256 No. 2 April 2012

---

$\mathbb{C}$ -operators on associative algebras and associative Yang–Baxter equations	257
CHENGMING BAI, LI GUO and XIANG NI	
Botany of irreducible automorphisms of free groups	291
THIERRY COULBOIS and ARNAUD HILION	
A note on inverse curvature flows in asymptotically Robertson–Walker spacetimes	309
CLAUS GERHARDT	
Total curvature of graphs after Milnor and Euler	317
ROBERT GULLIVER and SUMIO YAMADA	
Entire solutions of Donaldson’s equation	359
WEIYONG HE	
Energy identity and removable singularities of maps from a Riemann surface with tension field unbounded in $L^2$	365
YONG LUO	
Quotients by actions of the derived group of a maximal unipotent subgroup	381
DMITRI I. PANYUSHEV	
Invariants of totally real Lefschetz fibrations	407
NERMIN SALEPCI	
Stable trace formulas and discrete series multiplicities	435
STEVEN SPALLONE	
Small covers and the Halperin–Carlsson conjecture	489
LI YU	
Acknowledgement	509



0030-8730(201204)256:2;1-8