

*Pacific
Journal of
Mathematics*

**A RELATIVE TRACE FORMULA
FOR $\mathrm{PGL}(2)$ IN THE LOCAL SETTING**

BROOKE FEIGON

Volume 260 No. 2

December 2012

A RELATIVE TRACE FORMULA FOR $\mathrm{PGL}(2)$ IN THE LOCAL SETTING

BROOKE FEIGON

In memory of Jonathan Rogawski

We develop the local Kuznetsov trace formula on a unitary group in two variables for an unramified quadratic extension of local, non-Archimedean fields E/F and compare it to a local relative trace formula on $\mathrm{PGL}(2, E)$. To define the local distributions for the relative trace formula, we define a regularized local period integral and prove that it is a $\mathrm{PGL}(2, F)$ -invariant linear functional. By comparison of the two local trace formulas, we get an equality between a local $\mathrm{PGL}(2, F)$ -period and local Whittaker functionals.

| | |
|---|-----|
| 1. Introduction | 395 |
| 2. Notation | 403 |
| 3. The local Kuznetsov trace formula for $\mathrm{U}(2)$ | 405 |
| 4. The local relative trace formula and periods for $\mathrm{PGL}(2)$ | 413 |
| 5. Comparison of local trace formulas and applications | 427 |
| Acknowledgments | 430 |
| References | 431 |

1. Introduction

Base change is an important type of functoriality which is useful in the study of automorphic forms by relating automorphic representations on different groups. Hervé Jacquet shed light on a new technique for attacking certain cases of Robert Langlands' important functoriality conjectures by comparing the relative and Kuznetsov trace formulas in the global setting. Jacquet's comparison of trace formulas leads to global identities that characterize the image of the base change map associating automorphic representations of a unitary group for a quadratic extension of number fields E/F to automorphic representations of $\mathrm{GL}(2, \mathbb{A}_E)$ in terms of distinguished representations. While Jacquet's global identities factor, they do not give unique local identities.

The author was supported by NSF grant DMS-1201446.

MSC2010: 11F70, 22E50.

Keywords: relative trace formula.

This paper uses techniques of James Arthur to define and develop a local Kuznetsov trace formula on $U(2)$ and a local relative trace formula on $GL(2)$. Both local trace formulas are expanded geometrically in terms of orbital integrals and spectrally in terms of local Bessel distributions and local relative Bessel distributions. The latter involve regularized local period integrals. We then carry out Jacquet’s comparison in the local setting by relating these two local trace formulas for matching functions. This comparison yields identities between local Bessel distributions for automorphic representations on $U(2)$ and local relative Bessel distributions for automorphic representations on $GL(2)$.

Before we describe more precisely the local relative trace formula developed in this paper, let us recall the relative trace formula for $GL(2)$. Take E/F to be a quadratic extension of number fields and \mathbb{A}_F to be the adèles of F . Let ψ' be a character on $F \backslash \mathbb{A}_F \cong N(F) \backslash N(\mathbb{A}_F)$ where N is the upper triangular unipotent matrices of $GL(2)$. Let $\psi = \psi' \circ \text{tr}_{E/F}$.

A cuspidal automorphic representation π of $GL(2, \mathbb{A}_E)$ with central character trivial on $GL(2, \mathbb{A}_F)$ is *distinguished* by $GL(2, \mathbb{A}_F)$ if there exists a $\phi \in V_\pi$, the vector space associated to π , such that the *period integral*, $P(\phi)$, is nonzero:

$$P(\phi) := \int_{GL(2, F)Z(\mathbb{A}_F) \backslash GL(2, \mathbb{A}_F)} \phi(h) dh \neq 0.$$

Where π' is a cuspidal automorphic representation of the quasisplit unitary group $U(2, \mathbb{A}_F)$ and $\phi' \in V_{\pi'}$, let

$$W(\phi') = \int_{N(F) \backslash N(\mathbb{A}_F)} \phi'(n) \overline{\psi'(n)} dn \quad \text{and} \quad W(\phi) = \int_{N(E) \backslash N(\mathbb{A}_E)} \phi(n) \overline{\psi(n)} dn.$$

We define the *Bessel distribution* as

$$B_{\pi'}(f') := \sum_i W'(\pi'(f')\phi'_i) \overline{W'(\phi'_i)},$$

and the *relative Bessel distribution* as

$$B_\pi(f) := \sum_j P(\pi(f)\phi_j) \overline{W(\phi_j)},$$

where the summations are over an orthonormal basis of $V_{\pi'}$ and V_π respectively. Flicker [1991], following related work of Jacquet and Lai [1985] and Ye [1989], showed that for “matching functions” f' on $U(2, \mathbb{A}_F)$ and f on $GL(2, \mathbb{A}_E)$, if π' maps to π under the unstable base change, then

$$(1-1) \quad \sum_i W'(\pi'(f')\phi'_i) \overline{W'(\phi'_i)} = \sum_j P(\pi(f)\phi_j) \overline{W(\phi_j)}.$$

In particular, this equality characterizes the image of the unstable base change lift associating every automorphic representation of $U(2, \mathbb{A}_F)$ to an automorphic

representation of $GL(2, \mathbb{A}_E)$ in terms of $GL(2, \mathbb{A}_F)$ distinguished representations. The equality above is proved via the relative trace formula [Jacquet 2005], which tells us that for f and f' matching functions we have

$$\int_{(N(F)\backslash N(\mathbb{A}_F))^2} K_{f'}(n_1, n_2)\psi'(n_1^{-1}n_2) dn_1 dn_2 = \int_{GL(2, F)Z(\mathbb{A}_F)\backslash GL(2, \mathbb{A}_F)} \int_{N(E)\backslash N(\mathbb{A}_E)} K_f(h, n)\psi(n) dn dh$$

where

$$K_f(x, y) = \sum_{\delta \in Z(E)\backslash GL(2, E)} f(x^{-1}\delta y).$$

The distributions $B'_{\pi}(f')$ and $B_{\pi}(f)$ occur in the spectral expansions of the respective trace formulas.

In a different direction, Arthur [1989; 1991] developed a local version of the classical Arthur–Selberg trace formula. Let G be a connected reductive algebraic group over a local field F of characteristic zero. Diagonally embed $G(F)$ into $G(F) \times G(F)$. Then $L^2(G(F))$ is isomorphic to $L^2(G(F)\backslash G(F) \times G(F))$ by

$$\phi \mapsto ((y_1, y_2) \mapsto \phi(y_1^{-1}y_2)).$$

For $\phi \in L^2(G(F))$, let $(\rho(g_1, g_2)\phi)(x) = \phi(g_1^{-1}xg_2)$. The right regular representation of $G(F) \times G(F)$ on $L^2(G(F)\backslash G(F) \times G(F))$ is equivalent to ρ of $G(F) \times G(F)$ on $L^2(G(F))$. Thus to develop the local trace formula we look at $\rho(f)$ where $f = f_1 \otimes f_2 \in C_c^\infty(G(F) \times G(F))$. Then

$$(\rho(f)\phi)(x) = \int_{G(F)} \int_{G(F)} f_1(g)f_2(y)\phi(g^{-1}xy) dg dy$$

is an integral operator on $L^2(G(F))$ with kernel

$$K_f(x, y) = \int_{G(F)} f_1(g)f_2(x^{-1}gy) dg.$$

The local trace formula develops an explicit formula for the regularized trace of $\rho(f)$.

The main result of this paper is that, when evaluated with matching functions, the two local trace formulas described in Theorems 1.3 and 1.4 below, that is the local Kuznetsov trace formula and the local relative trace formula, are equal. Thus there is an equality between their local distributions on the spectral sides. This equality is stated in Theorem 1.1. This is the natural local counterpart to the global comparison from (1-1). In order to develop the local relative trace formula stated in Theorem 1.4, we have to define a local regularized period integral, prove it is a $GL(2, F) \times GL(2, F)$ -invariant linear functional and relate it to the truncated

period integral that initially appears in the relative trace formula. We state these properties about the local regularized period integral in Proposition 1.2.

To describe our results more precisely we need to introduce some further notation. Let E/F now denote an unramified extension of local non-Archimedean fields of characteristic 0. Let \mathbb{O}_F (respectively \mathbb{O}_E) denote the ring of integers in F (respectively E). Let $H = \text{GL}(2)/F$, $G = \text{Res}_{E/F} H$ and let

$$G' = \text{U}(2, F) = \left\{ g \in G : {}^t \bar{g} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} g = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}.$$

Let N' and N be the upper triangular unipotent matrices of G' and G , respectively, and let M' and M be the diagonal subgroup of G' and G , respectively. Let Z and Z' denote the center of G and G' , respectively. For any subgroup X of G let $\tilde{X} = Z \cap X \backslash X$ and let $X_H = X \cap H$. Let ψ' be an additive character on F with conductor \mathbb{O}_F and let $\psi(x) = \psi' \circ \text{tr}_{E/F}$. Let $f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$ and $f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$.

We define the local Kuznetsov trace formula as the equality between the geometric expansion (in terms of orbital integrals) and spectral expansion (in terms of representations) of

$$\lim_{t \rightarrow \infty} \int_{(N' \times N')(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2$$

and the local relative trace formula as the equality between the expansions of

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) dn dh.$$

In this local setting

$$K_f(x, y) = \int_{\tilde{G}(F)} f_1(g) f_2(x^{-1} g y) dg, \quad K_{f'}(x, y) = \int_{\tilde{G}'(F)} f'_1(g) f'_2(x^{-1} g y) dg$$

and $u(n, t)$ and $u(h, t)$ are truncation parameters that are needed due to convergence issues. They are defined analogously to Arthur’s truncation [1991, Section 3].

We use the following ideas in this paper to rewrite these local trace formulas in terms of orbital integrals and representations:

- methods of Arthur [1991] from the local trace formula,
- methods of Flicker [1991], Jacquet [2005] and Ye [1989] from the relative trace formula,
- Harish-Chandra’s Plancherel formula [Harish-Chandra 1984; Waldspurger 2003],
- Jacquet, Lapid and Rogawski’s methods for regularizing period integrals [Jacquet et al. 1999; Jacquet \geq 2012].

The power of the two trace formulas lies in the comparison. For “matching functions”, the geometric expansions of the two local relative trace formulas are equal. By comparing the spectral expansions in these two trace formulas, we get an analogue of (1-1), giving the following identity between local Bessel distributions for functions on $U(2)$ and local relative Bessel distributions for functions on $GL(2, E)$, and therefore local periods and local Whittaker functionals:

Theorem 1.1. *If σ is a supercuspidal representation on $\tilde{G}(F)$ that is the unstable base change lift of the supercuspidal representation σ' of $\tilde{G}'(F)$, and*

$$f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F)) \quad \text{and} \quad f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$$

are matching functions, then

$$(1-2) \quad d(\sigma') \sum_{S' \in \mathcal{B}(\sigma')} W'_{\sigma'}(\sigma'(f'_2)S'\sigma'(f'_1{}^\vee)) \overline{W'_{\sigma'}(S')} \\ = d(\sigma) \sum_{S \in \mathcal{B}(\sigma)} P_\sigma(\sigma(f_2)S\sigma(f_1{}^\vee)) \overline{W_\sigma(S)},$$

where $d(\sigma)$ is the formal degree of σ , $\mathcal{B}(\sigma)$ is an orthonormal basis of the Hilbert space of Hilbert–Schmidt operators on V_σ ,

$$W'_{\sigma'}(S') = \int_{N'(F)} \text{tr}(\sigma'(n)S')\psi'(n^{-1}) \, dn, \\ W_\sigma(S) = \int_{N(F)} \text{tr}(\sigma(n)S)\psi(n^{-1}) \, dn, \\ P_\sigma(S) = \int_{\tilde{H}(F)} \text{tr}(\sigma(h)S) \, dh.$$

The Bessel and relative Bessel distributions $B'_{\pi'}(f')$ and $B_\pi(f)$ factor into local (relative) Bessel distributions $B'_{\pi'_v}(f'_v)$ and $B_{\pi_v}(f_v)$, but it is not clear how to normalize the local distributions. The distributions on the left and right-hand side of (1-2) are each the product of two local distributions and (1-2) can be restated as

$$d(\sigma')B'_{\sigma'}(f'_2)B'_{\sigma'^*}(f'_1) = d(\sigma)B_\sigma(f_2)B_{\sigma^*}(f_1).$$

We note that the local period integral $P_\sigma(S)$ is not a convergent integral if σ is not a discrete series representation. To develop the local relative trace formula we have to define a local regularized period integral. Let $K = G(\mathbb{O}_F)$ and let $P = NM$. For $\lambda \in \mathbb{C}$ and $m = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix}$ let $e^{\lambda H_M(m)} = |\alpha/\beta|_E$ where $|\cdot|_E$ denotes the normalized valuation on E . For a principal series representation π of \tilde{G} and $u, v \in \pi$ we define the matrix coefficient $f_{u,v}(g) = \langle \pi(g)u, v \rangle$. Asymptotically on M , $f_{u,v}$ will equal a finite sum of functions of the form $e^{\lambda H_M(m)}$. We define the regularized period

integral as:

$$\int_{\tilde{H}(F)}^* f_{u,v}(h) dh := \int_{\tilde{H}(F)} f_{u,v}(h)u(h, t) dh + \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) f_{u,v}(k_1mk_2)(1 - u(m, t)) dm dk_1 dk_2$$

where

$$\int_{\tilde{M}_H^+(F)}^\sharp e^{\lambda H_M(m)}(1 - u(m, t)) dm$$

is the meromorphic continuation at $v = 0$ of

$$\int_{\tilde{M}_H^+(F)} e^{(v+\lambda)H_M(m)}(1 - u(m, t)) dm,$$

which is absolutely convergent for $\text{Re}(v) \ll 0$.

We prove that the regularized period integral is an $H(F) \times H(F)$ -invariant linear functional, and we relate it to the truncated period integral that initially appears in the local relative trace formula as follows. By abuse of notation we identify a character χ of $\tilde{M}(F)$ with a character χ of E^\times by letting $\chi \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} = \chi(a)\chi^{-1}(b)$. For $\lambda \in \mathbb{C}$ we let $\chi_\lambda(m) = \chi(m)e^{\lambda(H_M(m))}$. We let $I_P(\chi_\lambda)$ be the parabolically induced normalized representation acting on the Hilbert space $\mathcal{H}_P(\chi)$. Then for $S \in \mathcal{B}_P(\chi)$,

$$\text{tr}(I_P(\chi_\lambda, k_1gk_2)S) = E_P(g, \Psi_S, \lambda)_{k_1, k_2},$$

where $E_P(g, \Psi, \lambda)$ is the Eisenstein integral and

$$(C^P E_P)(m, \psi, \lambda) = (c_{P|P}(1, \lambda)\psi)(m)e^{\lambda H_M(m)} + (c_{P|P}(w, \lambda)\psi)(m)e^{-\lambda H_M(m)}.$$

We fix a uniformizer ϖ in F (and E) and $q^{-1} = |\varpi|_F$.

Proposition 1.2. *Fix a character χ of E^\times such that $\chi(\varpi) = 1$. Then for $t \gg 0$,*

$$\int_{\tilde{H}(F)} \text{tr}(I_P(\chi_\lambda, h)S)u(h, t) dh = \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) dh - \delta(\chi)(1 + q^{-1}) \left(\frac{q^{2\lambda(t+1)}}{1 - q^{2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda)\Psi_S(1)_{k_1, k_2} dk_1 dk_2 + \frac{q^{-2\lambda(t+1)}}{1 - q^{-2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda)\Psi_S(1)_{k_1, k_2} dk_1 dk_2 \right),$$

where $\delta(\chi) = 1$ if $\chi|_{\mathbb{O}_F^\times} = 1$ and $\delta(\chi) = 0$ if $\chi|_{\mathbb{O}_F^\times} \neq 1$.

Denote the action of the nontrivial element in $\text{Gal}(E/F)$ on $x \in E$ by \bar{x} . Denote by $N_{E/F}$ the norm map from E^\times to F^\times . Let $E^1 = \{x \in E^\times : N_{E/F}(x) = 1\}$. Let η denote an element in $G(F)$ such that $\bar{\eta}^{-1}\eta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

We define

$$D'_{\chi'_\lambda}(f') = \sum_{S' \in \mathcal{B}_P(\chi')} W'_{\chi'_\lambda}(S'_\lambda[f']) \overline{W'_{\chi'_\lambda}(S')}, \quad \text{and} \quad D_{\chi_\lambda}(f) = \sum_{S \in \mathcal{B}_P(\chi)} P_{\chi_\lambda}(S_\lambda[f]) \overline{W_{\chi_\lambda}(S)},$$

where

$$\begin{aligned} W'_{\chi'_\lambda}(S') &= \lim_{t \rightarrow \infty} \int_{N'(F)} \text{tr}(I_{P'}(\chi'_\lambda, n)S') \psi'(n^{-1})u(n, t) \, dn, \\ W_{\chi_\lambda}(S) &= \lim_{t \rightarrow \infty} \int_{N(F)} \text{tr}(I_P(\chi_\lambda, n)S) \psi(n^{-1})u(n, t) \, dn, \\ P_{\chi_\lambda}(S) &= \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) \, dh, \\ S_\lambda[f] &= I_P(\chi_\lambda, f_2) S I_P(\chi_\lambda, f_1^\vee). \end{aligned}$$

We let $\Pi_2(\tilde{G}'(F))$ be a set of equivalence classes of irreducible, tempered square integrable representations of $\tilde{G}'(F)$. We identify unitary characters on $\tilde{M}'(F)$ with characters on E^\times that are trivial on E^1 . We let $\{\Pi_2(\tilde{M}'(F))\}$ be a set of representatives of unitary characters χ' on $\tilde{M}'(F)$ such that $\chi'(\varpi) = 1$. We let $\mu(\chi'_\lambda)$ be Harish-Chandra's μ -function. We take the analogous definitions for $\tilde{G}(F)$.

Theorem 1.3 (local Kuznetsov trace formula). *For any*

$$f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F)),$$

we have

$$\begin{aligned} &\lim_{t \rightarrow \infty} \int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1}n_2) u(n_1, t) u(n_2, t) \, dn_1 \, dn_2 \\ &= \int_{a \in E^\times/E^1} O'(f_1, \psi', a) O'(f'_2, \bar{\psi}', a) |a|_E \, d^\times a \\ &= \sum_{\sigma' \in \Pi_2(\tilde{G}'(F))} d(\sigma') D_{\sigma'}(f') + \frac{1}{2} \sum_{\chi' \in \{\Pi_2(\tilde{M}'(F))\}} d(\chi') \int_0^{\pi i / \log q} \mu(\chi'_\lambda) D'_{\chi'_\lambda}(f') \, d\lambda, \end{aligned}$$

where

$$O'(f'_i, \psi', a) = \int_{N'(F)} \int_{N'(F)} f'_i(n_1^{-1} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} n_2) \overline{\psi'(n_1^{-1}n_2)} \, dn_1 \, dn_2.$$

Theorem 1.4 (local relative trace formula). *For any*

$$f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F)),$$

we have

$$\begin{aligned}
 \lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) \, dn \, dh \\
 = \int_{a \in E^\times / E^1} O(f_1, \psi, a) O(f_2, \bar{\psi}, a) |a|_E \, d^\times a \\
 = \sum_{\sigma \in \Pi_2(\tilde{G}(F))} d(\sigma) D_\sigma(f) + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi^2 \neq 1, \chi|_{F^\times} = 1}} \tilde{D}_\chi(f) \\
 + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi|_{E^1} = 1}} d(\chi) \int_0^{\pi i / \log q} \mu(\chi_\lambda) D_{\chi_\lambda}(f) \, d\lambda
 \end{aligned}$$

where

$$O(f_i, \psi, a) = \int_{\tilde{H}(F)} \int_{N(F)} f_i \left(h^{-1} \eta \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} n \right) \overline{\psi(n)} \, dn \, dh.$$

The representations that occur on the right-hand side of Theorem 1.4 are exactly the representations that are in the image of the unstable base change lift on $\tilde{G}'(F)$. The additional discrete term $\tilde{D}_\chi(f)$ corresponds to the representations that lift from discrete series on $\tilde{G}'(F)$ to principal series on $\tilde{G}(F)$.

In addition to the spectral comparison, these local trace formulas also have applications on the geometric side. If we define the inner product of two functions g_1, g_2 on E^\times / E^1 by

$$\langle g_1, g_2 \rangle = \int_{a \in E^\times / E^1} g_1(a) g_2(a) |a|_E \, d^\times a,$$

then:

Proposition 1.5 (orthogonality relations). *For f_1 and f_2 matrix coefficients of the supercuspidal representations σ_1 and σ_2 of $\tilde{G}(F)$ and f'_1 and f'_2 matrix coefficients of the supercuspidal representations σ'_1 and σ'_2 of $\tilde{G}'(F)$,*

$$\begin{aligned}
 \langle O'(f'_1, \psi', \cdot), O'(f'_2, \psi'^{-1}, \cdot) \rangle \neq 0 &\iff \sigma'_1 \sim \sigma'_2, \\
 \langle O(f_1, \psi, \cdot), O(f_2, \psi^{-1}, \cdot) \rangle \neq 0 &\iff \sigma_1 \sim \sigma_2.
 \end{aligned}$$

The rest of this paper is organized as follows. In Section 2 we define notation and give normalizations of measures. In Section 3 we develop the local Kuznetsov trace formula. For the geometric expansion we rewrite our trace formula in terms of orbital integrals corresponding to the $N' \backslash G' / N'$ double cosets. The orbital integrals for f'_1 and f'_2 initially depend on the truncation and are intertwined. It is only through the multiplication of the two orbital integrals, integration over the space of double cosets, and the nontriviality of the character ψ' , that we are able to untangle the orbital integral for f'_1 from the orbital integral for f'_2 . For the spectral

expansion we apply Harish-Chandra’s Plancherel formula to rewrite the local kernel in terms of representations. We are left with truncated integrals over the unipotent subgroup of matrix coefficients against the character ψ' . By the smoothness of the matrix coefficients and the appearance of the character, we show these distributions stabilize for t large.

In Section 4 we develop the local relative trace formula of $H \backslash G / N$. In the spectral expansion we have truncated integrals of matrix coefficients over H that do not converge without the truncation. We define the regularized period integral $P_{\chi_i}(S)$. We use the asymptotics of matrix coefficients of tempered representations to prove the truncated integral is a polynomial exponential function in the truncation parameter t . We define the regularized integral as the constant term of this polynomial, and prove that this is an $H \times H$ invariant linear functional and the relevant term in the local relative trace formula.

In Section 5 we compare our two local trace formulas. There is a bijection between the “admissible” $N' \backslash G' / N'$ cosets and the “admissible” $H \backslash G / N$ cosets and both of these sets can be parametrized by E^\times / E^1 . This bijection allows us to compare the geometric sides. By work of Ye and Flicker, we know that for any f' there is an f such that the orbital integrals are equal for corresponding cosets. Thus, by their geometric expansions, our local trace formulas are equal for matching functions. This gives an equality of the spectral expansions and of local distributions.

This paper would not have come into being had it not been for my teacher and advisor, Jonathan Rogawski. These thoughts originated as my PhD thesis under his direction, and his ideas, support, and guidance were critical to its completion. I am fortunate and will be forever grateful to have had him as a mentor. He could explain complicated math in a clear and simple way that aimed at the heart of the problem. He served, and continues to serve, as the role model of the inquisitive, patient, and approachable mathematician.

2. Notation

Let F be a non-Archimedean local field of characteristic 0 and odd residual characteristic q . Let E be an unramified quadratic extension of F . Let \mathbb{O}_F and \mathbb{O}_E denote the rings of integers in F and E , respectively. Let ϖ denote a uniformizer in the maximal ideal of \mathbb{O}_F . Thus ϖ is also a uniformizer in E . Let $v(\cdot)$ denote the valuation on F , extended to E . Let $|\cdot|_F$ and $|\cdot|_E$ denote the normalized valuations on F and E , respectively. Thus for $a \in F^\times$, $|a|_E = |a|_F^2$. Denote the action of the nontrivial element in $\mathrm{Gal}(E/F)$ on $x \in E$ by \bar{x} . Denote by $N_{E/F}$ the norm map from E^\times to F^\times . Let $E^1 = \{a \in E^\times : N_{E/F}(a) = 1\}$.

Let $H = \text{GL}(2)/F$ and let $G = \text{Res}_{E/F} H$, the restriction of scalars of $\text{GL}(2)$ from E to F . Thus $G(F) = \text{GL}(2, E)$. Let

$$G' = \text{U}(2, F) = \left\{ g \in G : {}^t \bar{g} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} g = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}.$$

We note that by defining the quasisplit unitary group in this way, $SL(2, F) \subset G'(F)$. Let N' and N be the upper triangular unipotent matrices of G' and G , respectively. Let M' and M be the diagonal subgroups of G' and G , respectively. That is,

$$M'(F) = \left\{ \begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} : a \in E^\times \right\} \quad \text{and} \quad M(F) = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} : a, b \in E^\times \right\}.$$

Occasionally by abuse of notation we let $n = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$ and $a = \begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix}$. Let $P = NM$ and $P' = N'M'$. Let $K = G(\mathbb{O}_F)$ and $K' = G'(\mathbb{O}_F)$. Let Z and Z' denote the centers of G and G' , respectively. For any subgroup X of G let $\tilde{X} = Z \cap X \backslash X$ and $X_H = X \cap H$. By abuse of notation we identify a character χ of $\tilde{M}(F)$ with a character χ of E^\times by letting $\chi \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} = \chi(a)\chi^{-1}(b)$.

Let ψ' be an additive character on F with conductor \mathbb{O}_F . Let ψ be the additive character on E defined by $\psi(x) = \psi'(x + \bar{x})$. By abuse of notation we will also denote by ψ and ψ' the corresponding characters on $N(F)$ and $N'(F)$, respectively. Let $f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$ and $f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$. For a function f on G , let $f^\vee(g) = f(g^{-1})$.

To define the local Kuznetsov trace formula and local relative trace formula we first multiply our function by the characteristic function of a large compact subset of $\tilde{G}(F)$ via Arthur's local truncation [1991, §3], and then take the limit of the integral of the truncated function. For $g \in G(F)$, $t \in \mathbb{Z}^+$, let

$$u(g, t) = \begin{cases} 1 & \text{if } g = zk_1 \begin{pmatrix} 1 & 0 \\ 0 & \alpha \end{pmatrix} k_2, \text{ for some } k_1, k_2 \in K, z \in Z(F), 0 \leq v(\alpha) \leq t, \\ 0 & \text{otherwise.} \end{cases}$$

We note that $u(\cdot, t)$ is well-defined on $\tilde{G}(F)$ and

$$u\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, t\right) = \begin{cases} 1 & \text{if } x \in \varpi^{\lceil -t/2 \rceil} \mathbb{O}_E, \\ 0 & \text{otherwise,} \end{cases}$$

where $[x]$ is the integral part of x .

If X is a closed subgroup of $\tilde{G}(F)$ with the subgroup topology, $\text{supp}(u(\cdot, t)) \cap X$ is a compact set.

We normalize the Haar measure dx on F so that $\text{vol}(\mathbb{O}_F) = 1$. We define the multiplicative measure $d^\times x$ on F^\times as

$$d^\times x = \frac{1}{1-q^{-1}} \frac{1}{|x|_F} dx.$$

Thus $\text{vol}(\mathbb{O}_F^\times) = 1$. We let $N(F)$ and $M(F)$ have the measures induced by dx and $d^\times x$. We normalize the Haar measure dk on K so that $\text{vol}(K) = 1$. We define the measure dg on $G(F)$ by

$$\int_{G(F)} f(g) dg = \int_{M(F)} \int_{N(F)} \int_K f(mnk) dk dn dm.$$

We define dg' on $G'(F)$ similarly. We normalize Haar measure on \tilde{K} by taking $\text{vol}(\tilde{K}) = 1$.

We let $d^\times a$ be the unique Haar measure on E^\times/E^1 such that

$$\text{vol}(\mathbb{O}_E^\times/E^1) = \frac{1}{1+q^{-1}}.$$

3. The local Kuznetsov trace formula for U(2)

In this section we develop a local Kuznetsov trace formula for the quasisplit unitary group in two variables. We expand this local Kuznetsov trace formula geometrically in terms of separate orbital integrals for f'_1 and f'_2 . Then we use Harish-Chandra's Plancherel formula to rewrite this expression spectrally in terms of representations.

We define the local Kuznetsov trace formula for

$$f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$$

as the equality between the geometric and spectral expansions of

$$\lim_{t \rightarrow \infty} \int_{(N' \times N')(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2$$

where

$$K_{f'}(n_1, n_2) = \int_{\tilde{G}'(F)} f'_1(g) f'_2(n_1^{-1} g n_2) dg.$$

We will show that for a fixed f' this limit stabilizes, that is, there exists a T such that for all $t' \geq T$,

$$\begin{aligned} & \int_{(N' \times N')(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t') u(n_2, t') dn_1 dn_2 \\ &= \lim_{t \rightarrow \infty} \int_{(N' \times N')(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2. \end{aligned}$$

3A. The geometric expansion. In this subsection we rewrite

$$\lim_{t \rightarrow \infty} \int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2$$

as an integral over admissible cosets of a product of an orbital integral for f'_1 and an orbital integral for f'_2 .

3A1. Integration formula. Let $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. For $a \in E^\times$, let $\beta_a = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$ and $\gamma_a = w \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$. By the Bruhat decomposition, $G' = P' \sqcup P'wP'$. Thus

$$\left\{ \beta_a : \begin{array}{l} a \text{ is in a set of} \\ \text{representatives for } E^\times/E^1 \end{array} \right\} \cup \left\{ \gamma_a : \begin{array}{l} a \text{ is in a set of} \\ \text{representatives for } E^\times/E^1 \end{array} \right\}$$

is a set of representatives for the double cosets of $N'(F) \backslash \tilde{G}'(F) / N'(F)$.

For $g \in G'(F)$ let

$$C_g(N'(F) \times N'(F)) = \{(n_1, n_2) \in N'(F) \times N'(F) : n_1^{-1}gn_2 = zg \text{ for some } z \in Z'(F)\}.$$

Definition 3.1. An element $g \in \tilde{G}'(F)$ and its corresponding orbit are called *admissible* if the map

$$C_g(N'(F) \times N'(F)) \rightarrow \mathbb{C} : (n_1, n_2) \mapsto \psi'(n_1^{-1}n_2)$$

is trivial.

By a simple calculation we see that

$$C_{\beta_a}(N'(F) \times N'(F)) = \left\{ \left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & \frac{x}{aa} \\ 0 & 1 \end{pmatrix} \right) : x \in F \right\},$$

$$C_{\gamma_a}(N'(F) \times N'(F)) = 1.$$

Thus the orbits represented by $\{\beta_1\} \cup \{\gamma_a : a \in E^\times/E^1\}$ are admissible.

We use the following integration formula to rewrite $K_{f'}(n_1, n_2)$ as an integral over the admissible cosets. Unlike in the global case the trivial admissible coset, β_1 , will not contribute to the trace formula.

For any $F \in C_c(\tilde{G}'(F))$,

$$(3-1) \quad \int_{\tilde{G}'(F)} F(g) dg = \int_{E^\times/E^1} \int_{(N' \times N')(F)} F(n_1^{-1}\gamma_a n_2) dn_1 dn_2 |a|_E d^\times a.$$

3A2. Separating the orbital integrals. Let

$$K^t(f') = \int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi(n_1^{-1}n_2) u(n_1, t) u(n_2, t) dn_1 dn_2.$$

Clearly $K^t(f')$ is absolutely convergent because f'_1 and $u(\cdot, t)$ have compact support on $\tilde{G}'(F)$ and $N'(F)$ respectively. By changing the order of integration and using (3-1), we see that $K^t(f')$ equals

$$\int_{E^\times/E^1} \int_{(N' \times N')(F)} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1}\gamma_a \hat{n}_2) f'_2(n_1^{-1}\hat{n}_1^{-1}\gamma_a \hat{n}_2 n_2) \\ \times \psi'(n_1^{-1}n_2) u(n_1, t) u(n_2, t) dn_1 dn_2 d\hat{n}_1 d\hat{n}_2 |a|_E d^\times a.$$

This integral is absolutely convergent because the map

$$N'(F) \times E^\times/E^1 \times N'(F) \rightarrow \tilde{G}'(F)$$

defined by

$$(n_1, a, n_2) \mapsto n_1^{-1} \gamma_a n_2$$

is injective and f'_1 has compact support. By a change of variables we have

$$K^t(f') = \int_{E^\times/E^1} K^t(\gamma_a, f') |a|_E d^\times a,$$

where

$$K^t(\gamma_a, f') = \int_{(N' \times N')(F)} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} \gamma_a \hat{n}_2) f'_2(n_1^{-1} \gamma_a n_2) \psi'(n_1^{-1} \hat{n}_1 \hat{n}_2^{-1} n_2) \\ \times u(\hat{n}_1^{-1} n_1, t) u(\hat{n}_2^{-1} n_2, t) dn_1 dn_2 d\hat{n}_1 d\hat{n}_2.$$

To complete the geometric expansion of the local Kuznetsov trace formula we rewrite $K^t(f')$ for $t \gg 0$ as an integral of two separate orbital integrals. We begin by examining the dependence of the integrand on the truncation.

Lemma 3.2. *Let $f'_1, f'_2 \in C_c(\tilde{G}'(F))$. For each $t_0 > 0$ there exists a $T > 0$ such that for all $t \geq T$,*

$$f'_1(x_1^{-1} \gamma y_1) f'_2(x_2^{-1} \gamma y_2) u(x_1^{-1} x_2, t) u(y_1^{-1} y_2, t_0) \\ = f'_1(x_1^{-1} \gamma y_1) f'_2(x_2^{-1} \gamma y_2) u(y_1^{-1} y_2, t_0)$$

for all $x_1, x_2, y_1, y_2, \gamma \in \tilde{G}'(F)$.

Proof. Let

$$\Omega_1 = \text{supp}(f'_1), \quad \Omega_2 = \text{supp}(f'_2), \quad \Omega_3 = \text{supp}(u(\cdot, t_0)) \cap \tilde{G}'(F).$$

These sets are all compact on $\tilde{G}'(F)$. If $f'_1(x_1^{-1} \gamma y_1) f'_2(x_2^{-1} \gamma y_2) u(y_1^{-1} y_2, t_0) \neq 0$, then the following conditions must hold:

- $x_1^{-1} \in \Omega_1 y_1^{-1} \gamma^{-1}$.
- $x_2 \in \gamma y_2 \Omega_2^{-1}$.
- $y_1^{-1} y_2 \in \Omega_3$.

Thus if $f'_1(x_1^{-1} \gamma y_1) f'_2(x_2^{-1} \gamma y_2) u(y_1^{-1} y_2, t_0) \neq 0$, then $x_1^{-1} x_2 \in \Omega_1 \Omega_3 \Omega_2^{-1}$. Because this is a compact set, there exists a $T > 0$ such that $\Omega_1 \Omega_3 \Omega_2^{-1} \subseteq \text{supp}(u(g, T))$. The lemma now follows. \square

Now we use this lemma, along with the character ψ' , to separate the two orbital integrals. By abuse of notation, in the proof of the following lemma we let

$$\varpi^n = \begin{pmatrix} \varpi^n & 0 \\ 0 & \varpi^{-n} \end{pmatrix} \quad \text{and} \quad a = \begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix}.$$

Lemma 3.3. *For $f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$, there exists a T such that for all $t \geq T$ and $n \in \mathbb{Z}$,*

$$(3-2) \quad \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} K^t(\gamma_a, f') d^\times a = \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} O'(f'_1, \psi', a) O'(f'_2, \bar{\psi}', a) d^\times a,$$

where

$$O'(f', \psi', a) = \int_{N'(F)} \int_{N'(F)} f'(n_1^{-1} \gamma_a n_2) \overline{\psi'(n_1^{-1} n_2)} dn_1 dn_2.$$

Proof. We show that there is a hidden truncation on the right-hand side of (3-2) that comes from the fact that the two orbital integrals are simultaneously evaluated at the same γ_a . Let K_1 be an open compact subgroup of $\tilde{G}'(F)$ such that f'_1 and f'_2 are bi- K_1 -invariant. There exists a positive constant c such that

$$\begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} \in K_1 \quad \text{for all } a \in (1 + \varpi^c \mathbb{O}_E) E^1.$$

By definition

$$\begin{aligned} & \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} K^t(\gamma_a, f') d^\times a \\ &= \int_{a \in \mathbb{O}_E^\times / E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} w \varpi^n a \hat{n}_2) \psi'(\hat{n}_1 \hat{n}_2^{-1}) \int_{(N' \times N')(F)} f'_2(n_1^{-1} w \varpi^n a n_2) \\ & \quad \times \psi'(n_1^{-1} n_2) u(\hat{n}_1^{-1} n_1, t) u(\hat{n}_2^{-1} n_2, t) dn_2 dn_1 d\hat{n}_2 d\hat{n}_1 d^\times a \\ &= \sum_{\eta \in \mathbb{O}_E^\times / (1 + \varpi^c \mathbb{O}_E) E^1} \int_{a \in (1 + \varpi^c \mathbb{O}_E) E^1 / E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} w \varpi^n \eta a \hat{n}_2) \psi'(\hat{n}_1 \hat{n}_2^{-1}) \\ & \quad \times \int_{(N' \times N')(F)} f'_2(n_1^{-1} w \varpi^n \eta a n_2) \psi'(n_1^{-1} n_2) u(\hat{n}_1^{-1} n_1, t) \\ & \quad \times u(\hat{n}_2^{-1} n_2, t) dn_2 dn_1 d\hat{n}_2 d\hat{n}_1 d^\times a. \end{aligned}$$

By a change of variables and the fact that f' is locally constant the right-hand side of this equation is equal to

$$\begin{aligned} & \sum_{\eta \in \mathbb{O}_E^\times / (1 + \varpi^c \mathbb{O}_E) E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} w \varpi^n \eta \hat{n}_2) \psi'(\hat{n}_1) \\ & \quad \times \int_{(N' \times N')(F)} f'_2(n_1^{-1} w \varpi^n \eta n_2) \psi'(n_1^{-1}) u(\hat{n}_1^{-1} n_1, t) dn_1 d\hat{n}_1 \\ & \quad \times \int_{a \in (1 + \varpi^c \mathbb{O}_E) E^1 / E^1} \psi'(a^{-1} \hat{n}_2^{-1} n_2 a) u(a^{-1} \hat{n}_2^{-1} n_2 a, t) d^\times a dn_2 d\hat{n}_2. \end{aligned}$$

We can rewrite the inner integral as

$$\begin{aligned} u(\hat{n}_2^{-1} n_2, t) & \int_{a \in (1 + \varpi^c \mathbb{O}_E) E^1 / E^1} \psi'((n_2 - \hat{n}_2)(a\bar{a})^{-1}) d^\times a \\ & = u(\hat{n}_2^{-1} n_2, t) \int_{b \in 1 + \varpi^c \mathbb{O}_F} \psi'(b(n_2 - \hat{n}_2)) d^\times b \\ & = u(\hat{n}_2^{-1} n_2, t) \frac{1}{1 - q^{-1}} \psi'(n_2 - \hat{n}_2) \int_{b \in \varpi^c \mathbb{O}_F} \psi'(b(n_2 - \hat{n}_2)) db \\ & = u(\hat{n}_2^{-1} n_2, t) u(\hat{n}_2^{-1} n_2, 2c) \frac{\mathrm{vol}(\varpi^c \mathbb{O}_F)}{1 - q^{-1}} \psi'(\hat{n}_2^{-1} n_2). \end{aligned}$$

Thus for $t \geq 2c$,

$$\begin{aligned} & \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} K^t(\gamma_a, f') d^\times a = \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} \gamma_a \hat{n}_2) \psi'(\hat{n}_1 \hat{n}_2^{-1}) \\ & \quad \times \int_{(N' \times N')(F)} f'_2(n_1^{-1} \gamma_a n_2) \psi'(n_1^{-1} n_2) u(\hat{n}_1^{-1} n_1, t) u(\hat{n}_2^{-1} n_2, 2c) dn_2 dn_1 d\hat{n}_2 d\hat{n}_1 d^\times a. \end{aligned}$$

By Lemma 3.2 there exists a $T > 0$ such that for all $t \geq \max\{T, 2c\}$,

$$\begin{aligned} & \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} K^t(\gamma_a, f') d^\times a \\ & = \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} \gamma_a \hat{n}_2) \psi'(\hat{n}_1 \hat{n}_2^{-1}) d\hat{n}_1 \\ & \quad \times \int_{(N' \times N')(F)} f'_2(n_1^{-1} \gamma_a n_2) \psi'(n_1^{-1} n_2) u(\hat{n}_2^{-1} n_2, 2c) dn_2 dn_1 d\hat{n}_2 d^\times a \\ & = \int_{a \in \varpi^n \mathbb{O}_E^\times / E^1} \int_{(N' \times N')(F)} f'_1(\hat{n}_1^{-1} \gamma_a \hat{n}_2) \psi'(\hat{n}_1 \hat{n}_2^{-1}) d\hat{n}_2 d\hat{n}_1 \\ & \quad \times \int_{(N' \times N')(F)} f'_2(n_1^{-1} \gamma_a n_2) \psi'(n_1^{-1} n_2) dn_2 dn_1 d^\times a. \quad \square \end{aligned}$$

We have shown that the truncated local Kuznetsov trace formula stabilizes.

Proposition 3.4. *For any $f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$ and $t \gg 0$,*

$$\int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1}n_2) u(n_1, t) u(n_2, t) dn_1 dn_2 = \int_{a \in E^\times/E^1} O'(f'_1, \psi', a) O'(f'_2, \bar{\psi}', a) |a|_E d^\times a.$$

3B. The spectral expansion. Now we derive a spectral expansion for the local Kuznetsov trace formula,

$$\lim_{t \rightarrow \infty} \int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1}n_2) u(n_1, t) u(n_2, t) dn_1 dn_2.$$

Our main tool is the Plancherel formula for p -adic groups, which was first stated, with an outlined proof, by Harish-Chandra [1984]. Silberger [1996] later filled in an important proof of one of the steps in the theorem. More recently Waldspurger [2003] provided a complete proof.

As in [Arthur 1991, §2], we begin by rewriting $K_{f'}(x, y)$ using the Plancherel formula. First we introduce some additional notation. For an irreducible representation (σ, V_σ) of $G'(F)$ let $\mathcal{B}(\sigma)$ be the Hilbert space of Hilbert–Schmidt operators on V_σ . The inner product on $\mathcal{B}(\sigma)$ is defined as

$$\langle S, S' \rangle := \text{tr}(SS'^*)$$

for $S, S' \in \mathcal{B}(\sigma)$, where $\text{tr}(SS'^*) = \sum_{\text{o.n.b. } V_\sigma} \langle SS'^* u_i, u_i \rangle$ and this sum converges absolutely and does not depend on the basis. For a discrete series representation σ of a group G let $d(\sigma)$ be the formal degree of σ .

Let $\Pi_2(\tilde{G}'(F))$ be a set of representatives for the equivalence classes of irreducible, tempered square integrable representations of $\tilde{G}'(F)$ and let $\{\Pi_2(\tilde{M}'(F))\}$ be a set of representatives of unitary characters χ on $\tilde{M}'(F)$ such that $\chi(\varpi) = 1$. For a character χ of $M'(F)$ and $\lambda \in \mathbb{C}$, let $\chi_\lambda(m) = \chi(m) e^{\lambda(H_{P'}(m))}$. For $\chi \in \{\Pi_2(\tilde{M}'(F))\}$, $I_{P'}^{G'}(\chi_\lambda) = I_{P'}(\chi_\lambda)$ is the normalized induced representation of $\tilde{G}'(F)$ acting on a Hilbert space $\mathcal{H}_{P'}(\chi)$ of vector-valued functions on K' . Let $\mathcal{B}_{P'}(\chi)$ be a fixed K' -finite orthonormal basis of the Hilbert space of Hilbert–Schmidt operators on $\mathcal{H}_{P'}(\chi)$.

Let $m(\sigma)$ be the Plancherel density. We normalize our measures following [Arthur 1991, §1]. The Plancherel density satisfies $m(\chi_\lambda) = d(\chi) \mu(\chi_\lambda)$, where $\mu(\chi_\lambda)$ is Harish-Chandra’s μ -function.

For a fixed $x \in G'(F)$, let

$$h(v) = \int_{\tilde{G}'(F)} f'_1(xu) f'_2(uxv) du.$$

Then $h \in C_c^\infty(\tilde{G}'(F))$ and $K_{f'}(x, y) = h(yx^{-1})$, so by the Plancherel formula,

$$K_{f'}(x, y) = \sum_{\sigma \in \Pi_2(\tilde{G}'(F))} d(\sigma) \operatorname{tr}(\sigma(R(yx^{-1})h)) + \frac{1}{2} \sum_{\chi \in \{\Pi_2(\tilde{M}'(F))\}} \int_0^{\frac{\pi i}{\log q}} \operatorname{tr}(I_{P'}(\chi, R(yx^{-1})h)) m(\chi_\lambda) d\lambda.$$

Because $I_{P'}(\chi_\lambda, R(yx^{-1})h) = I_{P'}(\chi_\lambda, f_1^{\vee}) I_{P'}(\chi_\lambda, x) I_{P'}(\chi_\lambda, f_2') (I_{P'}(\chi_\lambda, y))^*$, we have

$$\begin{aligned} \operatorname{tr}(I_{P'}(\chi_\lambda, R(yx^{-1})h)) &= \sum_{S \in \mathcal{B}_{P'}(\chi)} (I_{P'}(\chi_\lambda, f_1^{\vee}) I_{P'}(\chi_\lambda, x) I_{P'}(\chi_\lambda, f_2'), S^*) \overline{(I_{P'}(\chi_\lambda, y), S^*)} \\ &= \sum_{S \in \mathcal{B}_{P'}(\chi)} \operatorname{tr}(I_{P'}(\chi_\lambda, f_1^{\vee}) I_{P'}(\chi_\lambda, x) I_{P'}(\chi_\lambda, f_2') S) \overline{\operatorname{tr}(I_{P'}(\chi_\lambda, y) S)} \\ &= \sum_{S \in \mathcal{B}_{P'}(\chi)} \operatorname{tr}(I_{P'}(\chi_\lambda, x) S_\lambda[f']) \overline{\operatorname{tr}(I_{P'}(\chi_\lambda, y) S)}, \end{aligned}$$

where $S_\lambda[f'] = I_{P'}(\chi_\lambda, f_2') S I_{P'}(\chi_\lambda, f_1^{\vee})$.

For $f' \in C_c^\infty(\tilde{G}'(F))$, π an admissible representation, $\pi(f')$ has finite rank. Thus the sum over S is a finite sum of an orthonormal basis of operators on $\mathcal{H}_P(\chi)^{K_0}$ for some open compact K_0 .

Putting everything together we have

$$\begin{aligned} &\int_{N'(F)} \int_{N'(F)} K_{f_1' \otimes f_2'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2 \\ &= \sum_{\sigma \in \Pi_2(\tilde{G}'(F))} d(\sigma) \sum_{S \in \mathcal{B}(\sigma)} \left(\int_{N'(F)} \operatorname{tr}(\sigma(n)(\sigma(f_2') S \sigma(f_1^{\vee}))) \psi'(n^{-1}) u(n, t) dn \right. \\ &\quad \left. \times \overline{\int_{N'(F)} \operatorname{tr}(\sigma(n) S) \psi'(n^{-1}) u(n, t) dn} \right) \\ &+ \frac{1}{2} \sum_{\chi \in \{\Pi_2(\tilde{M}'(F))\}} d(\chi) \times \int_0^{\frac{\pi i}{\log q}} \left(\sum_{S \in \mathcal{B}_{P'}(\chi)} \int_{N'(F)} \operatorname{tr}(I_{P'}(\chi_\lambda, n) S_\lambda[f']) \psi'(n^{-1}) u(n, t) dn \right. \\ &\quad \left. \times \overline{\int_{N'(F)} \operatorname{tr}(I_{P'}(\chi_\lambda, n) S) \psi'(n^{-1}) u(n, t) dn} \right) \mu(\chi_\lambda) d\lambda. \end{aligned}$$

To finish the spectral expansion we show that the above unipotent integrals stabilize. We first note that in the discrete series case the above integrals are absolutely convergent without any truncation for reasons similar to those in Section 4B1.

Lemma 3.5 (spectral stabilization). *For any complex-valued function ϕ on $\tilde{G}'(F)$ that is biinvariant under a fixed open compact subgroup, there exists a positive*

integer c such that for all $t \geq c$,

$$\int_{N'(F)} \phi(n)\psi'(n)u(n, t) \, dn = \int_{N'(F)} \phi(n)\psi'(n)u(n, c) \, dn.$$

This c only depends on the open compact subgroup under which ϕ is biinvariant.

Proof. Let K_1 be an open compact subgroup of $\tilde{G}'(F)$ under which ϕ is biinvariant. K_1 must contain a neighborhood of the identity, so there exists a positive integer c' such that

$$\begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} \in K_1 \quad \text{for all } a \in (1 + \varpi^{c'} \mathbb{O}_E)E^1.$$

We show that for $m > c'$,

$$\int_{\varpi^{-m} \mathbb{O}_F^\times} \phi\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) \psi'(x) \, dx = 0.$$

We note that

$$\begin{pmatrix} a & 0 \\ 0 & \bar{a}^{-1} \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a^{-1} & 0 \\ 0 & \bar{a} \end{pmatrix} = \begin{pmatrix} 1 & a\bar{a}x \\ 0 & 1 \end{pmatrix}.$$

Thus for $x' \in 1 + \varpi^{c'} \mathbb{O}_F$,

$$\phi\left(\begin{pmatrix} 1 & x'x \\ 0 & 1 \end{pmatrix}\right) = \phi\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right).$$

Hence

$$\begin{aligned} & \int_{\varpi^{-m} \mathbb{O}_F^\times} \phi\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) \psi'(x) \, dx \\ &= \sum_{\alpha \in \mathbb{O}_F^\times / (1 + \varpi^{c'} \mathbb{O}_F)} \int_{\varpi^{-m}(1 + \varpi^{c'} \mathbb{O}_F)} \phi\left(\begin{pmatrix} 1 & \alpha x \\ 0 & 1 \end{pmatrix}\right) \psi'(\alpha x) \, dx \\ &= \sum_{\alpha \in \mathbb{O}_F^\times / (1 + \varpi^{c'} \mathbb{O}_F)} \phi\left(\begin{pmatrix} 1 & \varpi^{-m} \alpha \\ 0 & 1 \end{pmatrix}\right) \psi'(\varpi^{-m} \alpha) \int_{\varpi^{c'-m} \mathbb{O}_F} \psi'(x) \, dx. \end{aligned}$$

The last line equals 0 for $m > c'$. Thus for $t > 2c'$,

$$\int_{N'(F)} \phi(n)\psi'(n)u(n, t) \, dn = \int_{N'(F)} \phi(n)\psi'(n)u(n, 2c') \, dn. \quad \square$$

We have now proved the following.

Proposition 3.6. For any $f' = f'_1 \otimes f'_2 \in C_c^\infty(\tilde{G}'(F) \times \tilde{G}'(F))$,

$$\begin{aligned} & \lim_{t \rightarrow \infty} \int_{N'(F)} \int_{N'(F)} K_{f'}(n_1, n_2) \psi'(n_1^{-1} n_2) u(n_1, t) u(n_2, t) dn_1 dn_2 \\ &= \sum_{\sigma' \in \Pi_2(\tilde{G}'(F))} d(\sigma') D'_{\sigma'}(f') + \frac{1}{2} \sum_{\chi' \in \{\Pi_2(\tilde{M}'(F))\}} d(\chi') \int_0^{\frac{\pi i}{\log q}} D'_{\chi'}(f') \mu(\chi'_\lambda) d\lambda, \end{aligned}$$

where

$$\begin{aligned} D'_{\sigma'}(f') &= \sum_{S \in \mathcal{B}(\sigma')} W'_{\sigma'}(\sigma'(f'_2) S \sigma'(f'_1{}^\vee)) \overline{W'_{\sigma'}(S)}, \\ W'_{\sigma'}(S) &= \int_{N'(F)} \text{tr}(\sigma'(n) S) \psi'(n^{-1}) dn, \\ D'_{\chi'_\lambda}(f') &= \sum_{S \in \mathcal{B}_{P'}(\chi'_\lambda)} W'_{\chi'_\lambda}(I_{P'}(\chi'_\lambda, f'_2) S I_{P'}(\chi'_\lambda, f'_1{}^\vee)) \overline{W'_{\chi'_\lambda}(S)}, \\ W'_{\chi'_\lambda}(S) &= \lim_{t \rightarrow \infty} \int_{N'(F)} \text{tr}(I_{P'}(\chi'_\lambda, n) S) \psi'(n^{-1}) u(n, t) dn. \end{aligned}$$

We note that Theorem 1.3 now follows from the results of Propositions 3.4 and 3.6.

4. The local relative trace formula and periods for PGL(2)

In this section we define a local relative trace formula for PGL(2). We expand this local relative trace formula geometrically in terms of separate orbital integrals of f_1 and f_2 . Then we use Harish-Chandra’s Plancherel formula to rewrite this expression spectrally in terms of representations. We define a regularized period integral, show that it is an $H \times H$ -invariant linear functional and that it is the term that appears in the spectral expansion of the local relative trace formula.

We define the local relative trace formula for $f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$ as the equality between the geometric and spectral expansions of

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) dn dh$$

where

$$K_f(h, n) = \int_{\tilde{G}(F)} f_1(g) f_2(h^{-1} gn) dg.$$

As we did with the local Kuznetsov trace formula, we show that for a fixed f this limit stabilizes.

4A. The geometric expansion. We will rewrite

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) dn dh$$

as an integral over admissible cosets of a product of an orbital integral for f_1 and an orbital integral f_2 .

4A1. Integration formula. As pointed out in [Jacquet et al. 1999, §VI.13], by [Springer 1985], $G(F) = H(F)P(F) \sqcup H(F)\eta P(F)$, where η is any element in $G(F)$ such that $\bar{\eta}^{-1}\eta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Let $\eta_a = \eta \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$ and $\gamma_\alpha = \begin{pmatrix} 1 & 0 \\ 0 & \alpha + \sqrt{\tau} \end{pmatrix}$, where $E = F(\sqrt{\tau})$. Then

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\} \cup \{ \gamma_\alpha : \alpha \in F \} \cup \{ \eta_a : a \text{ is in a set of representatives for } E^\times/E^1 \}$$

is a set of representatives for the double cosets of $\tilde{H}(F) \backslash \tilde{G}(F) / N(F)$.

For $g \in G(F)$, let

$$C_g(\tilde{H}(F) \times N(F)) = \{ (h, n) \in \tilde{H}(F) \times N(F) : h^{-1}gn = zg \text{ for some } z \in Z(F) \}.$$

Definition 4.1. An element $g \in \tilde{G}(F)$ and its corresponding orbit is called *admissible* if the map $C_g(\tilde{H}(F) \times N(F)) \rightarrow \mathbb{C} : (h, n) \mapsto \psi(n)$ is trivial.

By a short calculation we see that

$$C_{\gamma_\alpha}(\tilde{H}(F) \times N(F)) = \left\{ \left(\begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & y(\alpha + \sqrt{\tau}) \\ 0 & 1 \end{pmatrix} \right) : y \in F \right\},$$

$$C_{\eta_a}(\tilde{H}(F) \times N(F)) = 1.$$

Thus the orbits represented by $\{ \eta_a : a \in E^\times/E^1 \} \cup \{ \gamma_0 \}$ are admissible.

We have the following integration formula. For any $F \in C_c(\tilde{G}(F))$,

$$(4-1) \quad \int_{\tilde{G}(F)} F(g) dg = \int_{E^\times/E^1} \int_{\tilde{H}(F) \times N(F)} F(h^{-1}\eta_a n) dn dh |a|_E d^\times a.$$

4A2. Separating the orbital integrals. Let

$$R^t(f) = \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) dn dh.$$

$R^t(f)$ is absolutely convergent because $f_1(g)$, $u(h, t)$ and $u(n, t)$ have compact support on $\tilde{G}(F)$, $\tilde{H}(F)$ and $N(F)$ respectively. By changing the order of integration and applying (4-1) we see that $R^t(f)$ equals

$$\int_{E^\times/E^1} \int_{\tilde{H}(F) \times N(F)} \int_{\tilde{H}(F) \times N(F)} f_1(h_1^{-1}\eta_a n_1) f_2(h_2^{-1}h_1^{-1}\eta_a n_1 n_2) \\ \times \psi(n_2) u(h_2, t) u(n_2, t) dn_2 dh_2 dn_1 dh_1 |a|_E d^\times a.$$

By a change of variables we have

$$R^t(f) = \int_{E^\times/E^1} R^t(\eta_a, f) |a|_E d^\times a,$$

where

$$R^t(\eta_a, f) = \int_{\tilde{H}(F) \times N(F)} \int_{\tilde{H}(F) \times N(F)} f_1(h_1^{-1} \eta_a n_1) f_2(h_2^{-1} \eta_a n_2) \psi(n_1^{-1} n_2) \times u(h_1^{-1} h_2, t) u(n_1^{-1} n_2, t) dn_2 dh_2 dn_1 dh_1.$$

To complete the geometric expansion of the local relative trace formula we rewrite $R^t(f)$ for $t \gg 0$ as an integral of a product of two separate orbital integrals that are not truncated. We omit the proof the lemma below as it is very similar to the proof of Lemma 3.3.

Lemma 4.2. *For $f \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$, there exists a $T > 0$ such that for all $t \geq T$ and $n \in \mathbb{Z}$,*

$$\int_{\mathfrak{o}^n \mathbb{O}_E^\times / E^1} R^t(\eta_a, f) d^\times a = \int_{\mathfrak{o}^n \mathbb{O}_E^\times / E^1} O(f_1, \psi, a) O(f_2, \bar{\psi}, a) d^\times a$$

where

$$O(f, \psi, a) = \int_{\tilde{H}(F)} \int_{N(F)} f(h^{-1} \eta_a n) \overline{\psi(n)} dn dh.$$

We have proved the following proposition.

Proposition 4.3. *For any $f = f_1 \otimes f_2 \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$,*

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K(h, n) \psi(n) u(h, t) u(n, t) dn dh \\ = \int_{a \in E^\times / E^1} O(f_1, \psi, a) O(f_2, \bar{\psi}, a) |a|_E d^\times a. \end{aligned}$$

Here, as in the local Kuznetsov trace formula, we have actually shown that the limit of the truncated local relative trace formula stabilizes.

4B. The spectral expansion and period integrals. We want to develop a spectral expansion for the local relative trace formula,

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n) \psi(n) u(h, t) u(n, t) dn dh.$$

As in the previous section, we expand the kernel via the Plancherel formula:

$$\begin{aligned} (4-2) \quad & \int_{\tilde{H}(F)} \int_{N(F)} K(h, n) \psi(n) u(h, t) u(n, t) dn dh \\ & = \sum_{\sigma \in \Pi_2(\tilde{G}(F))} d(\sigma) D_\sigma^t(f) + \frac{1}{2} \sum_{\chi \in \{\Pi_2(\tilde{M}(F))\}} d(\chi) \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda \end{aligned}$$

where

$$\begin{aligned}
 D_\sigma^t(f) &= \sum_{S \in \mathcal{B}(\sigma)} P_\sigma^t(\sigma(f_2)S\sigma(f_1^\vee)) \overline{W_\sigma^t(S)}, \\
 D_{\chi_\lambda}^t(f) &= \sum_{S \in \mathcal{B}_P(\chi)} P_{I_P(\chi_\lambda)}^t(I_P(\chi_\lambda, f_2)SI_P(\chi_\lambda, f_1^\vee)) \overline{W_{I_P(\chi_\lambda)}^t(S)}, \\
 W_\pi^t(S) &= \int_{N(F)} \text{tr}(\pi(n)S)\psi(n^{-1})u(n, t) \, dn, \\
 P_\pi^t(S) &= \int_{\tilde{H}(F)} \text{tr}(\pi(h)S)u(h, t) \, dh.
 \end{aligned}$$

By Lemma 3.5, there exists a positive integer c , such that for $t > c$,

$$W_\pi^t(S) = \int_{N(F)} \text{tr}(\pi(n)S)\psi(n^{-1})u(n, c) \, dn.$$

Thus as in the previous section, we define

$$\begin{aligned}
 W_\sigma(S) &= \int_{N(F)} \text{tr}(\sigma(n)S)\psi(n^{-1}) \, dn, \\
 W_{\chi_\lambda}(S) &= \lim_{t \rightarrow \infty} \int_{N(F)} \text{tr}(I_P(\chi_\lambda, n)S)\psi(n^{-1})u(n, t) \, dn.
 \end{aligned}$$

To finish the spectral expansion of the local relative trace formula we need to define the regularized integral

$$\int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) \, dh$$

because $\text{tr}(I_P(\chi_\lambda, -)S)$ is not integrable over $\tilde{H}(F)$.

Many of the techniques in this section are inspired by the work of Jacquet, Lapid and Rogawski in [Jacquet et al. 1999]. In that paper they define a regularized period integral for an automorphic form ϕ on $G(\mathbb{A})$ integrated over H where G is a reductive group over a number field F and H is the fixed point set of an involution of G . They focus on the case $G = \text{Res}_{E/F} H$ where E/F is a quadratic extension and they obtain explicit results for $G = \text{GL}(n, E)$, $H = \text{GL}(n, F)$.

For $\lambda \in \mathbb{C}$ and $m = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \in M(F)$, let $e^{\lambda H_M(m)} = |\alpha/\beta|_E^\lambda$. If $g = m(g)n(g)k(g)$, $m(g) \in M(F)$, $n(g) \in N(F)$, $k(g) \in K$, we let $e^{\lambda H_P(g)} = e^{\lambda H_M(m(g))}$. Let $\delta_P(m) = e^{H_M(m)}$. We give analogous definitions for $e^{\lambda H_{M_H}}$ and δ_{P_H} so that for $m \in M_H(F)$, $e^{\lambda H_M(m)} = e^{2\lambda H_{M_H}(m)}$.

We recall the Cartan decomposition $H(F) = K_H M_H^+(F) K_H$, where

$$M_H^+(F) = \left\{ \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \in M(F) : v\left(\frac{\alpha}{\beta}\right) \leq 0 \right\}.$$

Then for any absolutely integrable function f

$$\int_{\tilde{H}(F)} f(h) dh = \int_{\tilde{K}_H} \int_{\tilde{K}_H} \int_{\tilde{M}_H^+(F)} D_{P_H}(m) f(k_1 m k_2) dm dk_2 dk_1,$$

where

$$D_{P_H} \left(\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \right) = \begin{cases} |\alpha/\beta|_F (1 + |\varpi|_F) & v(\alpha/\beta) \leq 0, \\ 0 & v(\alpha/\beta) > 0. \end{cases}$$

To define the regularized integral, we begin by defining a regularized integral on $M_H^+(F)$. We note that

$$1 - u \left(\begin{pmatrix} 1 & \\ & \alpha \end{pmatrix}, t \right) = \begin{cases} 0 & 0 \leq v(\alpha) \leq t, \\ 1 & v(\alpha) > t. \end{cases}$$

For $\text{Re } \nu < -\text{Re } \lambda$,

$$(4-3) \quad \int_{\tilde{M}_H^+(F)} e^{(\nu+\lambda)H_M(m)} (1 - u(m, t)) dm = \sum_{n=t+1}^{\infty} q^{2n(\nu+\lambda)} = \frac{q^{(t+1)2(\nu+\lambda)}}{1 - q^{2(\nu+\lambda)}}.$$

We write

$$\int_{\tilde{M}_H^+(F)}^{\sharp} e^{\lambda H_M(m)} (1 - u(m, t)) dm$$

to denote the meromorphic continuation at $\nu = 0$ of (4-3). This is well-defined so long as $\lambda \neq 0$. Let

$$(4-4) \quad \begin{aligned} \phi(k_1 m k_2) &= \sum_{i=1}^r \phi_i(k_1, k_2) f_i(m) e^{\lambda_i H_M(m)}, \quad k_1, k_2 \in K_H, \\ m &= \begin{pmatrix} 1 & \\ & \varpi^n \end{pmatrix}, \quad n \geq 0, \quad f_i \in C_c(\tilde{M}(F)) \end{aligned}$$

with $\lambda_i \neq -\frac{1}{2}$. We define for $t \gg 0$,

$$\begin{aligned} &\int_{\tilde{H}(F)}^{\sharp} \phi(h) (1 - u(h, t)) dh \\ &= \sum_{i=1}^r \int_{\tilde{K}_H \times \tilde{K}_H} \phi_i(k_1, k_2) \int_{\tilde{M}_H^+(F)}^{\sharp} D_{P_H}(m) e^{\lambda_i H_M(m)} (1 - u(m, t)) dm \\ &= (1 + q^{-1}) \sum_{i=1}^r \int_{\tilde{K}_H \times \tilde{K}_H} \phi_i(k_1, k_2) \int_{\tilde{M}_H^+(F)}^{\sharp} e^{(\lambda_i+1/2)H_M(m)} (1 - u(m, t)) dm. \end{aligned}$$

If ϕ is a matrix coefficient of $I_P(\chi_\lambda)$ where $\chi(\varpi) = 1$ then by smoothness and the asymptotics of matrix coefficients there exists a function $C^P \phi$ of the form in

(4-4) with $\lambda_i \in \{\lambda - \frac{1}{2}, -\lambda - \frac{1}{2}\}$ and for $n \gg 0$,

$$C^P \phi \left(k_1 \begin{pmatrix} 1 & \\ & \varpi^n \end{pmatrix} k_2 \right) = \phi \left(k_1 \begin{pmatrix} 1 & \\ & \varpi^n \end{pmatrix} k_2 \right).$$

Note that the condition for the regularized integral to exist is now that $\lambda \neq 0$.

Definition 4.4. For any matrix coefficient ϕ of $I_P(\chi_\lambda)$ such that $\chi(\varpi) = 1$ and $\lambda \neq 0$,

$$\int_{\tilde{H}(F)}^* \phi(h) dh := \int_{\tilde{H}(F)} \phi(h)u(h, t) dh + \int_{\tilde{H}(F)}^\sharp \phi(h)(1 - u(h, t)) dh$$

for $t \gg 0$.

One can check that this definition of the regularized integral is independent of t and agrees with the usual integral if we start with something that is integrable. Now we will prove that it is H -invariant and then we will explicitly relate the regularized period to the truncated period that occurs in the local trace formula.

Let $\phi^{h_0}(x) = \phi(xh_0)$ for $h_0 \in \tilde{H}$. Note that if ϕ is a matrix coefficient of π then ϕ^{h_0} is as well.

Lemma 4.5. Fix $h_0 \in H$, $\lambda \neq 0$ and a character χ of E^\times with $\chi(\varpi) = 1$. Then for any matrix coefficient ϕ of $I_P(\chi_\lambda)$ and $t \gg 0$,

$$\begin{aligned} \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H(F)}^\sharp D_{P_H}(m) \phi^{h_0}(k_1 m k_2) (1 - u(k_1 m k_2 h_0, t)) dm dk_1 dk_2 \\ = \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H(F)}^\sharp D_{P_H}(m) \phi(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2. \end{aligned}$$

Proof. For $g \in G(F)$ let $\mathcal{M}(g) \in M^+(F)$ be such that $g = k_1 \mathcal{M}(g) k_2$, $k_1, k_2 \in K$. For $\text{Re } \nu \ll 0$ and $t \gg 0$,

$$\begin{aligned} \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H(F)} D_{P_H}(m) \phi^{h_0}(k_1 m k_2) \\ \times e^{\nu(H_M(\mathcal{M}(k_1 m k_2 h_0)))} (1 - u(k_1 m k_2 h_0, t)) dm dk_1 dk_2 \\ = \int_{\tilde{H}(F)} \phi(hh_0) e^{\nu(H_M(\mathcal{M}(hh_0)))} (1 - u(hh_0, t)) dh \\ = \int_{\tilde{H}(F)} \phi(h) e^{\nu(H_M(\mathcal{M}(h)))} (1 - u(h, t)) dh \\ = \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H(F)} D_{P_H}(m) \phi(k_1 m k_2) e^{\nu(H_M(m))} (1 - u(m, t)) dm dk_1 dk_2 \end{aligned}$$

by the invariance of Haar measure, since both sides are absolutely convergent. For $t \gg 0$, if $h \in \text{supp}(1 - u(\cdot, t))$, then $\mathcal{M}(hh_0) = \mathcal{M}(h)\mathcal{M}(k_2 h_0)$. Thus both sides of the equation above have a meromorphic continuation whose value at $\nu = 0$ gives the statement of the lemma. \square

Proposition 4.6 (*H*-invariance). *Let ϕ be a matrix coefficient of $I_P(\chi_\lambda)$, where $\chi(\varpi) = 1$ and $\lambda \neq 0$, and let $h_0 \in H(F)$. Then*

$$\int_{\tilde{H}(F)}^* \phi^{h_0}(h) dh = \int_{\tilde{H}(F)}^* \phi(h) dh.$$

Proof. By the definition of the regularized integrals, the statement of the proposition will follow once we prove the following equality:

$$\begin{aligned} & \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi^{h_0}(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2 \\ & \quad - \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2 \\ & \quad = \int_{\tilde{H}(F)} \phi(h) u(h, t) dh - \int_{\tilde{H}(F)} \phi^{h_0}(h) u(h, t) dh. \end{aligned}$$

First we note that by Lemma 4.5

$$\begin{aligned} & \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi^{h_0}(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2 \\ & \quad - \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2 \\ & \quad = \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi(k_1 m k_2) (1 - u(m k_2 h_0^{-1}, t)) dm dk_1 dk_2 \\ & \quad \quad - \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)}^\sharp D_{P_H}(m) \phi(k_1 m k_2) (1 - u(m, t)) dm dk_1 dk_2. \end{aligned}$$

For fixed h_0 and t sufficiently large, $u(\cdot h_0^{-1}, t) - u(\cdot, t)$ has support contained in an annulus. From this fact one can easily check that the previous line is equal to the convergent integral

$$\begin{aligned} & \int_{\tilde{K}_H \times \tilde{K}_H} \int_{\tilde{M}_H^+(F)} D_{P_H}(m) \phi(k_1 m k_2) [u(m, t) - u(m k_2 h_0^{-1}, t)] dm dk_1 dk_2 \\ & \quad = \int_{\tilde{H}(F)} \phi(h) [u(h, t) - u(h h_0^{-1}, t)] dh \\ & \quad = \int_{\tilde{H}(F)} \phi(h) u(h, t) dh - \int_{\tilde{H}(F)} \phi^{h_0}(h) u(h, t) dh. \quad \square \end{aligned}$$

We note that Proposition 4.6 also holds if we replace ϕ^{h_0} with $\phi(h_0 -)$ so our regularized integral is $H \times H$ invariant.

Now we derive an explicit formula relating regularized periods to truncated periods for the matrix coefficients that appear in the trace formula. We begin by recalling

some definitions of Harish-Chandra's. For σ an admissible, tempered representation of G , $\mathcal{A}_\sigma(G)$ is the space of functions on G spanned by K -finite matrix coefficients of σ , $\mathcal{A}_{\text{temp}}(G)$ is the sum of $\mathcal{A}_\sigma(G)$ over all admissible tempered representations of G and $\mathcal{A}_2(G)$ is the sum of $\mathcal{A}_\sigma(G)$ over all unitary, square integrable representations. For τ a finite dimensional, unitary, two-sided representation of K ,

$$\mathcal{A}_\sigma(G, \tau) = \{f \in \mathcal{A}_\sigma(G) \otimes V_\tau : f(k_1 g k_2) = \tau(k_1) f(g) \tau(k_2), g \in G, k_1, k_2 \in K\}.$$

Then $\mathcal{A}_{\text{temp}}(G, \tau)$ and $\mathcal{A}_2(G, \tau)$ are defined similarly.

Let $\tau_M = \tau|_{K \cap M}$. By [Harish-Chandra 1984, §3] for $f \in \mathcal{A}_\sigma(G, \tau)$ there exists a unique function $C^P f \in \mathcal{A}(M, \tau_M)$ such that

$$\lim_{|\frac{g}{\beta}|_E \rightarrow \infty} \left| \delta_P \left(\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \right)^{\frac{1}{2}} f \left(\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \right) - (C^P f) \left(\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \right) \right| = 0.$$

We call $C^P f$ the weak constant term of f .

For two parabolics P_1, P_2 with Levi component M , let

$$V_{P_1 P_2} = \{v \in V : \tau(n_1) v \tau(n_2) = v, n_1 \in N_{P_1} \cap K, n_2 \in N_{P_2} \cap K\}$$

and let $\tau_{P_1 P_2}$ be the subrepresentation of τ_M on $V_{P_1 P_2}$. For $\Psi \in \mathcal{A}_2(M, \tau_{P_1 P_2})$ and $\lambda \in [0, \pi i / \log q]$, the Eisenstein integral $E_P(g, \Psi, \lambda) \in \mathcal{A}_{\text{temp}}(G, \tau)$ is defined as

$$E_P(g, \Psi, \lambda) = \int_K \tau(k)^{-1} \Psi_P(kg) e^{(\lambda+1/2)(H_M(kg))} dk$$

where Ψ_P extends Ψ to G by $\Psi_P(nmk) = \Psi(m) \tau(k)$ for $n \in N, m \in M, k \in K$.

The weak constant term of the Eisenstein integral uniquely defines Harish-Chandra's c -functions [1984, §6]. For each element w in the Weyl group W of \tilde{G} , the c -function $c_{P|P}(w, \lambda)$ is a linear map from $\mathcal{A}_2(M, \tau_{P|P})$ to $\mathcal{A}_2(M, \tau_{P|\bar{P}})$ such that

$$(C^P E_P)(m, \Psi, \lambda) = (c_{P|P}(1, \lambda) \Psi)(m) e^{\lambda H_M(m)} + (c_{P|P}(w, \lambda) \Psi)(m) e^{-\lambda H_M(m)}$$

where w is a representative for the nontrivial element in the Weyl group of \tilde{G} . Let $c_{P|P}(s, \lambda)_\chi$ denote the restriction of $c_{P|P}(s, \lambda)$ to $\mathcal{A}_\chi(M, (\tau_\Gamma)_{P|P})$. We have

$$\mu(\chi_\lambda)^{-1} = c_{P|P}(s, \lambda)_\chi^* c_{P|P}(s, \lambda)_\chi.$$

For the rest of this section we let $c(1, \lambda) = c_{P|P}(1, \lambda)_\chi$ and $c(w, \lambda) = c_{P|P}(w, \lambda)_\chi$.

We note that the S we consider are actually in $\mathcal{H}_P(\chi)^{K_0}$ for some open compact K_0 . Harish-Chandra [1976, §7] gives an isomorphism $S \rightarrow \Psi_S$ from $\text{End}(\mathcal{H}_P(\chi)^K)$ onto $\mathcal{A}_\chi(M, (\tau)_{P|P})$ where V_τ is a particular subspace of $L^2(K \times K)$ such that

$$\text{tr}(I_P(\chi_\lambda, k_1 g k_2) S) = E_P(g, \Psi_S, \lambda)_{k_1, k_2}.$$

We can now relate the regularized integral to what appears in the local relative trace formula.

Proposition 4.7. For $\chi = (\chi, \chi^{-1}) \in \{\Pi_2(\tilde{M}(F))\}$, $\chi(\varpi) = 1$, $\lambda \neq 0$, $t \gg 0$,

$$\begin{aligned} & \int_{\tilde{H}(F)}^* \mathrm{tr}(I_P(\chi_\lambda, h)S) dh \\ &= \int_{\tilde{H}(F)} \mathrm{tr}(I_P(\chi_\lambda, h)S)u(h, t) dh \\ & \quad + \delta(\chi)(1+q^{-1}) \left(\frac{q^{2\lambda(t+1)}}{1-q^{2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda)\Psi_S(1)_{k_1, k_2} dk_1 dk_2 \right. \\ & \quad \left. + \frac{q^{-2\lambda(t+1)}}{1-q^{-2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda)\Psi_S(1)_{k_1, k_2} dk_1 dk_2 \right), \end{aligned}$$

where $\delta(\chi) = 1$ if $\chi|_{\mathbb{O}_F^\times} = 1$ and $\delta(\chi) = 0$ if $\chi|_{\mathbb{O}_F^\times} \neq 1$.

Proof. For $S \in \mathcal{B}_P(\chi)$, $\Psi_S \in \mathcal{A}_\chi(M, (\tau_\Gamma)_{P|P})$ and

$$c(1, \lambda)\Psi_S, c(w, \lambda)\Psi_S \in \mathcal{A}_\chi(M, (\tau_\Gamma)_{P|\bar{P}}).$$

Therefore $\Psi = \Psi_S$ can be written as a sum of matrix coefficients of χ . Thus

$$\begin{aligned} C^P E_P(m, \Psi, \lambda)_{k_1, k_2} &= c(1, \lambda)\Psi(m)_{k_1, k_2} e^{\lambda H_M(m)} + c(w, \lambda)\Psi(m)_{k_1, k_2} e^{-\lambda H_M(m)} \\ &= \chi(m) \left[(c(1, \lambda)\Psi)(1)_{k_1, k_2} e^{\lambda H_M(m)} + (c(w, \lambda)\Psi)(1)_{k_1, k_2} e^{-\lambda H_M(m)} \right] \end{aligned}$$

where $\chi(m) \in \mathbb{C}^\times$. Hence for $t \gg 0$,

$$\begin{aligned} & \int_{\tilde{M}_H^+(F)} D_{P_H}(m) \mathrm{tr}(I_P(\chi_\lambda, k_1 m k_2)S) e^{\nu(H_M(m))} (1-u(m, t)) dm \\ &= \int_{\tilde{M}_H^+(F)} D_{P_H}(m) \delta_P^{-\frac{1}{2}}(m) (c(1, \lambda)\Psi)(1)_{k_1, k_2} e^{(\lambda+\nu)(H_M(m))} \chi(m) (1-u(m, t)) dm \\ & \quad + \int_{\tilde{M}_H^+(F)} D_{P_H}(m) \delta_P^{-\frac{1}{2}}(m) (c(w, \lambda)\Psi)(1)_{k_1, k_2} e^{(-\lambda+\nu)(H_M(m))} \chi(m) (1-u(m, t)) dm \\ &= (1+q^{-1}) (c(1, \lambda)\Psi)(1)_{k_1, k_2} \int_{\tilde{M}_H^+(F)} e^{(\lambda+\nu)(H_M(m))} \chi(m) (1-u(m, t)) dm \\ & \quad + (1+q^{-1}) (c(w, \lambda)\Psi)(1)_{k_1, k_2} \int_{\tilde{M}_H^+(F)} e^{(-\lambda+\nu)(H_M(m))} \chi(m) (1-u(m, t)) dm \\ &= (1+q^{-1}) \int_{\mathbb{O}_F^\times} \chi(\alpha) d^\times \alpha \sum_{n=t+1}^{\infty} \left[(c(1, \lambda)\Psi)(1)_{k_1, k_2} q^{2(\lambda+\nu)n} \right. \\ & \quad \left. + (c(w, \lambda)\Psi)(1)_{k_1, k_2} q^{2(-\lambda+\nu)n} \right]. \end{aligned}$$

Clearly $\int_{\mathbb{O}_F^\times} \chi(\alpha) d^\times\alpha = 0$ unless $\chi|_{\mathbb{O}_F^\times} = 1$. If $\chi|_{\mathbb{O}_F^\times} = 1$, the previous line equals

$$(1 + q^{-1}) \left(\frac{q^{2(\lambda+\nu)(t+1)}}{1 - q^{2(\lambda+\nu)}} c(1, \lambda) \Psi(1)_{k_1, k_2} + \frac{q^{2(-\lambda+\nu)(t+1)}}{1 - q^{2(-\lambda+\nu)}} c(w, \lambda) \Psi(1)_{k_1, k_2} \right).$$

Therefore for $t \gg 0$

$$\begin{aligned} & \int_{\tilde{M}_H^+(F)}^\# D_{P_H}(m) \operatorname{tr}(I_P(\chi_\lambda, k_1 m k_2) S) (1 - u(m, t)) dm \\ &= \delta(\chi) (1 + q^{-1}) \left(\frac{q^{2\lambda(t+1)}}{1 - q^{2\lambda}} c(1, \lambda) \Psi_S(1)_{k_1, k_2} + \frac{q^{-2\lambda(t+1)}}{1 - q^{-2\lambda}} c(w, \lambda) \Psi_S(1)_{k_1, k_2} \right) \end{aligned}$$

and the proposition now follows. \square

Lemma 4.8. *Let $\chi = (\chi, \chi^{-1})$ where χ is a character of E^\times such that $\chi(\varpi) = 1$. Then*

(1) *If $\chi|_{F^\times} \neq 1$ and $\chi|_{E^1} \neq 1$, then*

$$\int_{\tilde{H}(F)}^* \operatorname{tr}(I_P(\chi_\lambda, h) S) dh = \int_{\tilde{H}(F)} \operatorname{tr}(I_P(\chi_\lambda, h) u(h, t)) dh = 0.$$

(2) *If $\chi|_{F^\times} \neq 1$ and $\chi|_{E^1} = 1$, then for $t \gg 0$,*

$$\int_{\tilde{H}(F)}^* \operatorname{tr}(I_P(\chi_\lambda, h) S) dh = \int_{\tilde{H}(F)} \operatorname{tr}(I_P(\chi_\lambda, h) S) u(h, t) dh.$$

(3) *If $\chi|_{F^\times} = 1$ and $\chi|_{E^1} \neq 1$, then $\int_{\tilde{H}}^* \operatorname{tr}(I_P(\chi_\lambda, h) S) dh$ is 0 whenever defined and*

$$\int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2 = \int_{\tilde{K}_H \times \tilde{K}_H} c(s, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2$$

at $\lambda = 0$.

(4) *If $\chi|_{F^\times} = 1$ and $\chi|_{E^1} = 1$, then $\chi^2 = 1$. In this case $c(1, \lambda)$ and $c(s, \lambda)$ have a simple pole at $\lambda = 0$ and so $\mu(\chi_\lambda)$ has a zero of order two at $\lambda = 0$ and $\mu(\chi_\lambda)c(1, \lambda) = \mu(\chi_\lambda)c(s, \lambda) = 0$ at $\lambda = 0$.*

In all cases,

$$\mu(\chi_\lambda) \int_{\tilde{H}(F)}^* \operatorname{tr}(I_P(\chi_\lambda, h) S) dh$$

is holomorphic for all $\lambda \in i\mathbb{R}$, $S \in \mathcal{B}_P(\chi)$.

Proof. In this proof we follow the techniques of [Jacquet \geq 2012]. Case 2 is obvious from the above work. Case 1 is obvious from the above work and the H -invariance of $\int_{\tilde{H}}^* \operatorname{tr}(I_P(\chi_\lambda, h) S) dh$ [Jacquet et al. 1999, Proposition 22].

The vanishing of the regularized period for $\lambda \neq 0$ in case 3 also follows from H -invariance. Then by the previous proposition we know that for $\lambda \neq 0$,

$$\int_{\tilde{H}(F)} \text{tr}(I_P(\chi_\lambda, h)S)u(h, t) dh = -(1 + q^{-1}) \left(\frac{q^{2\lambda(t+1)}}{1 - q^{2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2 + \frac{q^{-2\lambda(t+1)}}{1 - q^{-2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2 \right).$$

Both sides are holomorphic and the left-hand side is also defined and holomorphic for $\lambda = 0$. As

$$\text{Res}_{\lambda=0} \frac{q^{2\lambda(t+1)}}{1 - q^{2\lambda}} = \frac{-1}{2 \log q} \quad \text{and} \quad \text{Res}_{\lambda=0} \frac{q^{-2\lambda(t+1)}}{1 - q^{-2\lambda}} = \frac{1}{2 \log q},$$

we must have that

$$\int_{\tilde{K}_H \times \tilde{K}_H} c(1, 0) \Psi(1)_{k_1, k_2} dk_1 dk_2 = \int_{\tilde{K}_H \times \tilde{K}_H} c(w, 0) \Psi(1)_{k_1, k_2} dk_1 dk_2.$$

In case 4 the poles and zeros are well-known and can also be seen by explicit computations of the intertwining operators. We have that

$$\begin{aligned} &\mu(\chi_\lambda) \int_{\tilde{H}(F)} \text{tr}(I_P(\chi_\lambda, h)S)u(h, t) dh \\ &= \mu(\chi_\lambda) \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) dh \\ &\quad - (1 + q^{-1}) \mu(\chi_\lambda) \left(\frac{q^{2\lambda(t+1)}}{1 - q^{2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2 + \frac{q^{-2\lambda(t+1)}}{1 - q^{-2\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_S(1)_{k_1, k_2} dk_1 dk_2 \right). \end{aligned}$$

The left-hand side is 0 at $\lambda = 0$ and the last two terms are holomorphic at $\lambda = 0$ so the first term must be holomorphic at $\lambda = 0$. □

Let

$$\begin{aligned}
 D_{\chi_\lambda}(f) &= \sum_{S \in \mathcal{B}_P(\chi)} P_{\chi_\lambda}(S_\lambda[f]) \overline{W_{\chi_\lambda}(S)}, \\
 P_{\chi_\lambda}(S) &= \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) dh, \\
 S_\lambda[f] &= I_P(\chi_\lambda, f_2) S I_P(\chi_\lambda, f_1^\vee), \\
 \tilde{D}_\chi(f) &= (1 + q^{-1})\mu(\chi_0) \sum_{S \in \mathcal{B}_P(\chi)} \overline{W_{\chi_0}(S)} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, 0) \psi_{S_0[f]}(1)_{k_1, k_2} dk_1 dk_2.
 \end{aligned}$$

We now relate the distributions above to the truncated distributions from (4-2).

Lemma 4.9. *Let $\chi = (\chi, \chi^{-1})$ where χ is a character of E^\times such that $\chi(\varpi) = 1$.*

(1) *If $\chi|_{F^\times} \neq 1$ and $\chi|_{E^1} \neq 1$, then*

$$\lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda = 0.$$

(2) *If $\chi|_{F^\times} \neq 1$ and $\chi|_{E^1} = 1$, then*

$$\lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda = \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}(f) d\lambda.$$

(3) *If $\chi|_{F^\times} = 1$ and $\chi|_{E^1} \neq 1$, then*

$$\lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda = \tilde{D}_\chi(f).$$

(4) *If $\chi|_{F^\times} = 1$ and $\chi|_{E^1} = 1$, then*

$$\lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda = \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}(f) d\lambda.$$

Proof. First we note that

$$\begin{aligned}
 \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}^t(f) d\lambda &= \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) \sum_{S \in \mathcal{B}_P(\chi)} P_{I_P(\chi_\lambda)}^t(S_\lambda[f]) \overline{W_{\chi_\lambda}(S)} d\lambda \\
 &= \sum_{S \in \mathcal{B}_P(\chi)} \int_0^{\frac{\pi i}{\log q}} \left(\overline{\int_{N(F)} \text{tr}(I_P(\chi_\lambda, n)S) \psi(n^{-1})u(n, t) dn} \right) \\
 &\quad \times \mu(\chi_\lambda) \int_{\tilde{H}(F)} \text{tr}(I_P(\chi_\lambda, h)S_\lambda[f])u(h, t) dh d\lambda.
 \end{aligned}$$

Cases 1 and 2 now follow directly from Lemma 4.8. For the remaining cases we note that by Proposition 4.7 for $t \gg 0$,

$$\begin{aligned} & \int_{\tilde{H}(F)} \text{tr}(I_P(\chi_\lambda, h) S_\lambda[f]) u(h, t) dh \\ &= \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h) S_\lambda[f]) dh \\ & \quad + \delta(\chi) \frac{1+q^{-1}}{q^\lambda - q^{-\lambda}} \left(q^{2\lambda(t+\frac{1}{2})} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right. \\ & \quad \left. - q^{-2\lambda(t+\frac{1}{2})} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right). \end{aligned}$$

In case 3, by Lemma 4.8 the regularized period vanishes and we are left computing

$$\begin{aligned} (4-5) \quad & (1+q^{-1}) \lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) \overline{W_{\chi_\lambda}(S)} \\ & \left(\frac{q^{2\lambda(t+1/2)}}{q^\lambda - q^{-\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right. \\ & \quad \left. - \frac{q^{-2\lambda(t+1/2)}}{q^\lambda - q^{-\lambda}} \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right) d\lambda. \end{aligned}$$

Let

$$\begin{aligned} f_1(\lambda) &= \frac{1+q^{-1}}{2} \mu(\chi_\lambda) \overline{W_{\chi_\lambda}(S)} \left(\int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right. \\ & \quad \left. - \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right), \\ f_2(\lambda) &= \frac{1+q^{-1}}{2} \mu(\chi_\lambda) \overline{W_{\chi_\lambda}(S)} \left(\int_{\tilde{K}_H \times \tilde{K}_H} c(1, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right. \\ & \quad \left. + \int_{\tilde{K}_H \times \tilde{K}_H} c(w, \lambda) \Psi_{S_\lambda[f]}(1)_{k_1, k_2} dk_1 dk_2 \right). \end{aligned}$$

Then (4-5) equals

$$\begin{aligned} & \lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \frac{f_1(\lambda)(q^{2\lambda(t+\frac{1}{2})} + q^{-2\lambda(t+\frac{1}{2})})}{q^\lambda - q^{-\lambda}} \\ & \quad + \lim_{t \rightarrow \infty} \int_0^{\frac{\pi i}{\log q}} \frac{f_2(\lambda)(q^{2\lambda(t+\frac{1}{2})} - q^{-2\lambda(t+\frac{1}{2})})}{q^\lambda - q^{-\lambda}} d\lambda. \end{aligned}$$

By Lemma 4.8, $f_1(0) = 0$. Hence by Fourier analysis the first integral will vanish. The limit of the second integral will be $f_2(0)$, which, by the identity in case 3 of

Lemma 4.8, equals

$$(1 + q^{-1})\mu(\chi_0)\overline{W_{\chi_0}(S)} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, 0)\Psi_{S_0[f]}(1)_{k_1, k_2} dk_1 dk_2.$$

For case 4 by Lemma 4.8 when multiplied by $\mu(\chi_\lambda)\overline{W_{\chi_\lambda}(S)}$, $f_1(\lambda)$ and $f_2(\lambda)$ are holomorphic functions of λ and vanish at $\lambda = 0$, thus by similar analysis as above the last two terms vanish in the limit and we are left with the statement of the lemma. \square

4B1. Discrete series representations. Because the matrix coefficient of a supercuspidal representation σ has compact support it is obvious that

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \text{tr}(\sigma(h)S)u(h, t) dh = \int_{\tilde{H}(F)} \text{tr}(\sigma(h)S) dh.$$

Now we will prove that this is also true for Steinberg representations.

Lemma 4.10. *For $\sigma = St(\chi)$, $\chi^2 = 1$, the matrix coefficients are absolutely convergent over $\tilde{H}(F)$. Thus the limit*

$$\lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \text{tr}(\sigma(h)S)u(h, t) dh$$

exists and equals

$$\int_{\tilde{H}(F)} \text{tr}(\sigma(h)S) dh.$$

Proof. By [Borel and Wallach 1980, XI.4.3; Casselman 1995, 4.2.3], a matrix coefficient for σ evaluated at $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ is equal to a matrix coefficient for the Jacquet functor σ_N , evaluated at the same value, for $|\frac{a}{b}|_E$ sufficiently small. The Jacquet functor of σ is δ_P . Thus outside some compact set, our original matrix coefficient will behave like δ_P on $M_H^-(F)$. When we integrate over $\tilde{H}(F)$, using the $K_H M_H^-(F) K_H$ decomposition, we get a measure factor of $\delta_P^{-1/2}$. Thus outside a set of compact support our integral will look like $\int_{|a| < c} |a|_F d^\times a$ for some $c > 0$. \square

Putting everything together we have proved the following.

Proposition 4.11. *For any $f \in C_c^\infty(\tilde{G}(F) \times \tilde{G}(F))$,*

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_f(h, n)\psi(n)u(h, t)u(n, t) dn dh \\ = \sum_{\sigma \in \Pi_2(\tilde{G}(F))} d(\sigma)D_\sigma(f) + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi^2 \neq 1, \chi|_{F^\times} = 1}} \tilde{D}_\chi(f) \\ + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi|_{E^1} = 1}} d(\chi) \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda)D_{\chi_\lambda}(f) d\lambda, \end{aligned}$$

where

$$\begin{aligned}
 D_{\chi_\lambda}(f) &= \sum_{S \in \mathcal{B}_P(\chi)} P_{\chi_\lambda}(S_\lambda[f]) \overline{W_{\chi_\lambda}(S)}, \\
 P_{\chi_\lambda}(S) &= \int_{\tilde{H}(F)}^* \text{tr}(I_P(\chi_\lambda, h)S) dh, \\
 \tilde{D}_\chi(f) &= (1 + q^{-1})\mu(\chi_0) \sum_{S \in \mathcal{B}_P(\chi)} \overline{W_{\chi_0}(S)} \int_{\tilde{K}_H \times \tilde{K}_H} c(1, 0)\psi_{S_0[f]}(1)_{k_1, k_2} dk_1 dk_2, \\
 D_\sigma(f) &= \sum_{S \in \mathcal{B}(\sigma)} P_\sigma(\sigma(f_2)S\sigma(f_1^\vee)) \overline{W_\sigma(S)}, \\
 P_\sigma(S) &= \int_{\tilde{H}(F)} \text{tr}(\sigma(h)S) dh.
 \end{aligned}$$

This proposition combined with Proposition 4.3 proves Theorem 1.4.

5. Comparison of local trace formulas and applications

We now combine the results of the previous two sections to compare the two trace formulas. Let $\omega_{E/F}$ be the quadratic character of F^\times associated to E/F and let ω denote its trivial extension to E^\times .

Definition 5.1. We say that $f' \in C_c^\infty(\tilde{G}'(F))$ and $f \in C_c^\infty(\tilde{G}(F))$ are *matching functions* if $O'(f', \psi', a) = \omega(a)O(f, \psi, a)$ for all $a \in E^\times$.

By work of Ye [1989] and Flicker [1991, Proposition 3], we know that for any $f' \in C_c^\infty(\tilde{G}'(F))$ there exists a matching $f \in C_c^\infty(\tilde{G}(F))$ and vice versa. In fact, by the Fundamental Lemma, for f' spherical, we know that f is the corresponding function from the base change map between their Hecke algebras. Thus by the geometric expansion of the trace formulas in Propositions 3.4 and 4.3 we have the following statement.

Proposition 5.2. For $f'_i \in C_c^\infty(\tilde{G}'(F))$ and $f_i \in C_c^\infty(\tilde{G}(F))$ matching functions for $i = 1, 2$,

$$\begin{aligned}
 \lim_{t \rightarrow \infty} \int_{N'(F)} \int_{N'(F)} K_{f'_1 \otimes f'_2}(n_1, n_2)\psi'(n_1^{-1}n_2)u(n_1, t)u(n_2, t) dn_1 dn_2 \\
 = \lim_{t \rightarrow \infty} \int_{\tilde{H}(F)} \int_{N(F)} K_{f_1 \otimes f_2}(h, n)\psi(n)u(h, t)u(n, t) dn dh.
 \end{aligned}$$

Now we use the equality of the trace formulas to compare the spectral expansions. By Propositions 3.6, 4.11 and 5.2 we have the following result.

Theorem 5.3. For f_i and f'_i matching functions for $i = 1, 2$,

$$\begin{aligned} & \sum_{\sigma' \in \Pi_2(\tilde{G}'(F))} d(\sigma') D'_{\sigma'}(f'_1 \otimes f'_2) + \frac{1}{2} \sum_{\chi' \in \{\Pi_2(\tilde{M}'(F))\}} d(\chi') \int_0^{\frac{\pi i}{\log q}} \mu(\chi'_\lambda) D'_{\chi'_\lambda}(f'_1 \otimes f'_2) d\lambda \\ &= \sum_{\sigma \in \Pi_2(\tilde{G}(F))} d(\sigma) D_\sigma(f_1 \otimes f_2) + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi^2 \neq 1, \chi|_{F^\times} = 1}} \tilde{D}_\chi(f_1 \otimes f_2) \\ & \quad + \frac{1}{2} \sum_{\substack{\chi \in \{\Pi_2(\tilde{M}(F))\} \\ \chi|_{E^1} = 1}} d(\chi) \int_0^{\frac{\pi i}{\log q}} \mu(\chi_\lambda) D_{\chi_\lambda}(f_1 \otimes f_2) d\lambda. \end{aligned}$$

The unstable base change map associated to ω lifts principal series representations of \tilde{G}' to principal series representations $I_P(\chi)$ of \tilde{G} such that $\chi|_{E^1} = 1$. It also lifts certain square integrable representations of \tilde{G}' to the principal series representations of \tilde{G} defined by $I_P(\chi\omega)$ such that $\chi^2 \neq 1, \chi|_{F^\times} = 1$. It lifts the remaining square integrable representations of \tilde{G}' to square integral representations of \tilde{G} [Rogawski 1990; Flicker 1982]. Thus we could rephrase the right-hand side of Theorem 5.3 in terms of summing over the representations of \tilde{G} that are the unstable base change lifts of representations of \tilde{G}' . The extra discrete term $\tilde{W}_\chi(f)$ corresponds exactly to the representations that lift from the discrete series of \tilde{G}' to the principal series of \tilde{G} .

We also note that the only representations that appear on the right-hand side of Theorem 5.3 are those σ or $I_P(\chi_\lambda)$ for which there is a matrix coefficient such that the regularized integral over H is nonzero. This gives us a more explicit description of the nonvanishing H invariant linear functional that characterizes the image of the unstable base change map.

We would like to relate our distributions to the local factors in the Bessel and relative Bessel distributions. Recall from the introduction that Jacquet’s global relative trace formula tells us that for f' on $U(2, \mathbb{A}_F)$ and f on $GL(2, \mathbb{A}_E)$ matching functions, if a cuspidal representation π' of $U(2, \mathbb{A}_F)$ maps to π of $GL(2, \mathbb{A}_E)$ under unstable base change, then

$$B'_{\pi'}(f') = B_\pi(f)$$

where

$$\begin{aligned}
 B'_{\pi'}(f') &= \sum_{\phi' \in \text{o.n.b.}(V_{\pi'})} W'(\pi'(f')\phi') \overline{W'(\phi')}, \\
 B_{\pi}(f) &= \sum_{\phi \in \text{o.n.b.}(V_{\pi})} P(\pi(f)\phi) \overline{W(\phi)}, \\
 W'(\phi') &= \int_{N'(F)\backslash N'(\mathbb{A}_F)} \phi'(n) \overline{\psi'(n)} \, dn, \\
 W(\phi) &= \int_{N(E)\backslash N(\mathbb{A}_E)} \phi(n) \overline{\psi(n)} \, dn, \\
 P(\phi) &= \int_{\text{GL}(2,F)\text{Z}(\mathbb{A}_F)\backslash \text{GL}(2,\mathbb{A}_F)} \phi(h) \, dh \neq 0.
 \end{aligned}$$

While $B'_{\pi'}(f')$ and $B_{\pi}(f)$ factor into local Bessel distributions $B'_{\pi'_v}(f'_v)$ and $B_{\pi_v}(f_v)$, it is not clear how to normalize the local Bessel distributions. We can rewrite our local distributions as a product of two local Bessel (or local relative Bessel) distributions:

Lemma 5.4. (1) *For σ' an irreducible supercuspidal representation of $\tilde{G}'(F)$, there exists a local Bessel distribution $B'_{\sigma'}$, unique up to a constant of absolute value 1, such that*

$$D'_{\sigma'}(f'_1 \otimes f'_2) = B'_{\sigma'}(f'_2) B'_{\sigma'^*}(f'_1).$$

(2) *For σ an irreducible supercuspidal representation of $\tilde{G}(F)$, there exists a local relative Bessel distribution B_{σ} , unique up to a constant of absolute value 1, such that*

$$D_{\sigma}(f_1 \otimes f_2) = B_{\sigma}(f_2) B_{\sigma^*}(f_1).$$

Proof. We recall that

$$D'_{\sigma'}(f') = \sum_{S' \in \mathcal{B}(\sigma')} \int_{N'(F)} \text{tr}(\sigma'(n_1)\sigma'(f'_2)S'\sigma'^*(f'_1))\psi'(n_1)^{-1} \, dn_1 \overline{\int_{N'(F)} \text{tr}(\sigma'(n_2)S')\psi'(n_2)^{-1} \, dn_2}.$$

Let $V = V_{\sigma'}$. As S' is an endomorphism on V there exist $v \in V, v^* \in V^*$ such that $S' = v \otimes v^*$. Then the linear functional on $V \otimes V^*$ that acts by

$$v \otimes v^* \mapsto \int_{N'(F)} \text{tr}(\sigma'(n)v \otimes v^*)\psi'(n)^{-1} \, dn$$

transforms under n on v and v^* by ψ' . Thus it is a Whittaker functional on $V \otimes V^*$. By the uniqueness of Whittaker models,

$$\int_{N'(F)} \text{tr}(\sigma'(n)S')\psi'(n)^{-1} dn = W'(v)W'(v^*).$$

Thus

$$\begin{aligned} D'_{\sigma'}(f') &= \sum_{v \otimes v^*} W'(\sigma'(f'_2)v)\overline{W'(v)}W'(\sigma'^*(f'_1)v^*)\overline{W'(v^*)} \\ &= B'_{\sigma'}(f'_2)B'_{\sigma'^*}(f'_1). \end{aligned}$$

We note that if we change $B'_{\sigma'}$ by a constant c , then $B'_{\sigma'^*}$ will change by \bar{c} .

The proof for the local relative Bessel distributions is similar, using the uniqueness of the H -invariant linear functional [Hakim 1991; Flicker 1991, Proposition 11]. \square

We can also describe matching functions by an equality of all the Bessel distributions.

Lemma 5.5 (density). (1) *If $f'_1 \in C_c^\infty(\tilde{G}'(F))$ is such that $D'_{\sigma'}(f'_1 \otimes f'_2) = 0$ for all irreducible tempered representations σ' of $\tilde{G}'(F)$ and all f'_2 , then $O'(f'_1, \psi'^{-1}, a) = 0$ for all $a \in E^\times$.*

(2) *If $f_1 \in C_c^\infty(\tilde{G}(F))$ is such that $D_\sigma(f_1 \otimes f_2) = 0$ and $\tilde{D}_\sigma(f_1 \otimes f_2) = 0$ for all irreducible tempered representations σ of $\tilde{G}(F)$ and f_2 , then $O(f_1, \psi^{-1}, a) = 0$ for all $a \in E^\times$.*

Proof. If $D'_{\sigma'}(f'_1 \otimes f'_2) = 0$ for all σ' , then by Theorem 1.3,

$$\int_{a \in E^\times/E^1} |a|_E O'(f'_1, \psi'^{-1}, a) O'(f'_2, \psi', a) d^\times a = 0$$

for all $f'_2 \in C_c^\infty(\tilde{G}'(F))$. As $O'(f'_1, \psi'^{-1}, a)$ is a locally constant function of a there exists some open compact U such that $O'(f'_1, \psi'^{-1}, a)$ is biinvariant under it. Then by choosing f'_2 such that $O'(f'_2, \psi', a)$ has support contained in U we see that $O'(f'_1, \psi'^{-1}, a) = 0$. The second case follows from the first one. \square

Combining Theorem 5.3 with Lemma 5.4 and the global relative trace formula, we have the following result:

Corollary 5.6. *If σ is the supercuspidal representation of $\tilde{G}(F)$ that is the unstable base change lift of the supercuspidal representation σ' on $\tilde{G}'(F)$, and f'_i and f_i are matching functions for $i = 1, 2$, then*

$$d(\sigma')D_{\sigma'}(f'_1 \otimes f'_2) = d(\sigma)D_\sigma(f_1 \otimes f_2).$$

Proof. From the global comparison of relative trace formulas [Flicker 1991; Lapid 2006; Ye 1989] and a standard globalization argument we know there exists a constant c_σ such that $B'_{\sigma'}(f'_i) = c_\sigma B_\sigma(f_i)$ for all matching f_i, f'_i . Take f'_1 and f_2 to

be matrix coefficients of σ' and σ such that $B'_{\sigma'}(f'_1) \neq 0$ and $B_\sigma(f_2) \neq 0$. Take f'_2 a matching function to f_2 and f_1 a matching function to f'_1 . Then by Theorem 5.3,

$$d(\sigma')D'_{\sigma'}(f') = d(\sigma)D_\sigma(f). \quad \square$$

In addition to the spectral comparison, these local trace formulas also have applications on the geometric side. If we define the inner product of two functions g_1, g_2 on E^\times/E^1 by

$$\langle g_1, g_2 \rangle = \int_{a \in E^\times/E^1} g_1(a)g_2(a)|a|_E d^\times a,$$

then:

Corollary 5.7 (orthogonality relations). *For f_1 and f_2 matrix coefficients of the supercuspidal representations σ_1 and σ_2 of $\tilde{G}(F)$,*

$$\langle O(f_1, \psi, \cdot), O(f_2, \psi^{-1}, \cdot) \rangle \neq 0 \iff \sigma_1 \sim \sigma_2.$$

For f'_1 and f'_2 matrix coefficients of the supercuspidal representations σ'_1 and σ'_2 of $\tilde{G}'(F)$,

$$\langle O'(f'_1, \psi', \cdot), O'(f'_2, \psi'^{-1}, \cdot) \rangle \neq 0 \iff \sigma'_1 \sim \sigma'_2.$$

Proof. This follows directly from the local Kuznetsov and local relative trace formulas. □

Acknowledgments

The author thanks Erez Lapid for his helpful suggestions in an early stage of this project and Omer Offen for his useful comments. Finally, the author thanks the referee for helpful comments.

References

- [Arthur 1989] J. Arthur, “Towards a local trace formula”, pp. 1–23 in *Algebraic analysis, geometry, and number theory* (Baltimore, MD, 1988), edited by J.-I. Igusa, Johns Hopkins Univ. Press, Baltimore, MD, 1989. MR 98h:22020 Zbl 0765.22010
- [Arthur 1991] J. Arthur, “A local trace formula”, *Inst. Hautes Études Sci. Publ. Math.* 73 (1991), 5–96. MR 92f:22029 Zbl 0741.22013
- [Borel and Wallach 1980] A. Borel and N. R. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, Annals of Mathematics Studies 94, Princeton University Press, 1980. MR 83c:22018 Zbl 0443.22010
- [Casselman 1995] B. Casselman, “Introduction to the theory of admissible representations of p -adic reductive groups”, unpublished notes, 1995, Available at www.math.ubc.ca/~cass/research/pdf/p-adic-book.pdf.
- [Flicker 1982] Y. Z. Flicker, “Stable and labile base change for $U(2)$ ”, *Duke Math. J.* 49:3 (1982), 691–729. MR 84i:22016 Zbl 0502.12013

- [Flicker 1991] Y. Z. Flicker, “On distinguished representations”, *J. Reine Angew. Math.* **418** (1991), 139–172. MR 92i:22019 Zbl 0725.11026
- [Hakim 1991] J. Hakim, “Distinguished p -adic representations”, *Duke Math. J.* **62**:1 (1991), 1–22. MR 92c:22037 Zbl 0724.22016
- [Harish-Chandra 1976] Harish-Chandra, “Harmonic analysis on real reductive groups, III: The Maass–Selberg relations and the Plancherel formula”, *Ann. of Math. (2)* **104**:1 (1976), 117–201. MR 55 #12875 Zbl 0331.22007
- [Harish-Chandra 1984] Harish-Chandra, “The plancherel formula for reductive p -adic groups”, pp. 353–367 in *Collected papers Vol. IV*, edited by V. S. Varadarajan, Springer, New York, 1984. MR 85e:01061d
- [Jacquet 2005] H. Jacquet, “A guide to the relative trace formula”, pp. 257–272 in *Automorphic representations, L-functions and applications: Progress and prospects*, edited by J. W. Cogdell et al., Ohio State Univ. Math. Res. Inst. Publ. **11**, de Gruyter, Berlin, 2005. MR 2006g:11100 Zbl 1106.11016
- [Jacquet \geq 2012] H. Jacquet, “Truncation for the Fourier trace formula: the $GL(2)$ case”, unpublished notes.
- [Jacquet and Lai 1985] H. Jacquet and K. F. Lai, “A relative trace formula”, *Compositio Math.* **54**:2 (1985), 243–310. MR 86j:11059 Zbl 0587.12006
- [Jacquet et al. 1999] H. Jacquet, E. Lapid, and J. Rogawski, “Periods of automorphic forms”, *J. Amer. Math. Soc.* **12**:1 (1999), 173–240. MR 99c:11056 Zbl 1012.11044
- [Lapid 2006] E. M. Lapid, “On the fine spectral expansion of Jacquet’s relative trace formula”, *J. Inst. Math. Jussieu* **5**:2 (2006), 263–308. MR 2007d:11059 Zbl 1195.11071
- [Rogawski 1990] J. D. Rogawski, *Automorphic representations of unitary groups in three variables*, Annals of Mathematics Studies **123**, Princeton University Press, 1990. MR 91k:22037 Zbl 0724.11031
- [Silberger 1996] A. J. Silberger, “Harish-Chandra’s Plancherel theorem for p -adic groups”, *Trans. Amer. Math. Soc.* **348**:11 (1996), 4673–4686. MR 99c:22026 Zbl 0869.22007
- [Springer 1985] T. A. Springer, “Some results on algebraic groups with involutions”, pp. 525–543 in *Algebraic groups and related topics* (Kyoto/Nagoya, 1983), edited by R. Hotta, Adv. Stud. Pure Math. **6**, North-Holland, Amsterdam, 1985. MR 86m:20050 Zbl 0628.20036
- [Waldspurger 2003] J.-L. Waldspurger, “La formule de Plancherel pour les groupes p -adiques (d’après Harish-Chandra)”, *J. Inst. Math. Jussieu* **2**:2 (2003), 235–333. MR 2004d:22009 Zbl 1029.22016
- [Ye 1989] Y. Ye, “Kloosterman integrals and base change for $GL(2)$ ”, *J. Reine Angew. Math.* **400** (1989), 57–121. MR 90i:11134 Zbl 0665.10020

Received July 19, 2012. Revised October 9, 2012.

BROOKE FEIGON
DEPARTMENT OF MATHEMATICS
THE CITY COLLEGE OF NEW YORK, CUNY
NAC 8/133
NEW YORK, NY 10031
UNITED STATES
bfeigon@ccny.cuny.edu

PACIFIC JOURNAL OF MATHEMATICS

<http://pacificmath.org>

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Alexander Merkurjev
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
merkurev@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

PRODUCTION

Silvio Levy, Scientific Editor, pacific@math.berkeley.edu

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or pacificmath.org for submission instructions.

The subscription price for 2012 is US \$420/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Periodicals Service Company, 11 Main Street, Germantown, NY 12526-5635. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and the Science Citation Index.

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 969 Evans Hall, Berkeley, CA 94720-3840, is published monthly except July and August. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright © 2012 Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 260 No. 2 December 2012

Special issue devoted to the memory of Jonathan Rogawski

| | |
|--|-----|
| In memoriam: Jonathan Rogawski | 257 |
| DON BLASIUS, DINAKAR RAMAKRISHNAN and V. S. VARADARAJAN | |
| p -adic Rankin L -series and rational points on CM elliptic curves | 261 |
| MASSIMO BERTOLINI, HENRI DARMON and KARTIK PRASANNA | |
| The syntomic regulator for K_4 of curves | 305 |
| AMNON BESSER and ROB DE JEU | |
| Unique functionals and representations of Hecke algebras | 381 |
| BENJAMIN BRUBAKER, DANIEL BUMP and SOLOMON FRIEDBERG | |
| A relative trace formula for $\mathrm{PGL}(2)$ in the local setting | 395 |
| BROOKE FEIGON | |
| On the degrees of matrix coefficients of intertwining operators | 433 |
| TOBIAS FINIS, EREZ LAPID and WERNER MÜLLER | |
| Comparison of compact induction with parabolic induction | 457 |
| GUY HENNIART and MARIE-FRANCE VIGNERAS | |
| The functional equation and beyond endoscopy | 497 |
| P. EDWARD HERMAN | |
| A correction to <i>Conducteur des Représentations du groupe linéaire</i> | 515 |
| HERVÉ JACQUET | |
| Modular L -values of cubic level | 527 |
| ANDREW KNIGHTLY and CHARLES LI | |
| On occult period maps | 565 |
| STEPHEN KUDLA and MICHAEL RAPOPORT | |
| A prologue to “Functoriality and reciprocity”, part I | 583 |
| ROBERT LANGLANDS | |
| Truncation of Eisenstein series | 665 |
| EREZ LAPID and KEITH OUELLETTE | |
| Some comments on Weyl’s complete reducibility theorem | 687 |
| JONATHAN ROGAWSKI and V. S. VARADARAJAN | |
| On equality of arithmetic and analytic factors through local Langlands correspondence | 695 |
| FREYDOON SHAHIDI | |



0030-8730(201212)260:2;1-A