

*Pacific
Journal of
Mathematics*

**NOTES ON THE STRUCTURE CONSTANTS OF
HECKE ALGEBRAS OF INDUCED REPRESENTATIONS OF
FINITE CHEVALLEY GROUPS**

CHARLES W. CURTIS

NOTES ON THE STRUCTURE CONSTANTS OF HECKE ALGEBRAS OF INDUCED REPRESENTATIONS OF FINITE CHEVALLEY GROUPS

CHARLES W. CURTIS

This paper is dedicated to the memory of Robert Steinberg.

This paper contains an algorithm for the structure constants of the Hecke algebra of a Gelfand–Graev representation of a finite Chevalley group.

1. Introduction

Let G be a Chevalley group over a finite field $k = F_q$ of characteristic p (as in [Chevalley 1955] or [Steinberg 1968]). Let B be a Borel subgroup of G with $U = O_p(B)$ (the unipotent radical of B), and let T be a maximal torus such that $B = UT$. Let W be the Weyl group of G . Then W is a finite Coxeter group with distinguished generators $S = \{s_1, \dots, s_n\}$.

Let Φ be the root system associated with W , with $\{\alpha_1, \dots, \alpha_n\}$ the set of simple roots corresponding to the generators $s_i \in S$, and Φ_{\pm} the set of positive roots (respectively, negative roots) associated with them. For each root α , let U_{α} be the root subgroup of G corresponding to it. The subgroup U is generated by the root subgroups U_{α} , $\alpha > 0$.

From [Steinberg 1968, §3], the Chevalley group G has a B, N -pair, with Borel subgroup B , N the subgroup generated by all elements $w_{\alpha}(t)$, and $B \cap N$ equal to T , the subgroup generated by all elements $h_{\alpha}(t)$ (see the definitions of $w_{\alpha}(t)$ and $h_{\alpha}(t)$ in Section 2). Then $N/T \cong W$. (If the field k contains more than three elements, then N is the normalizer $N = N_G(T)$; see [Steinberg 1968, p. 36]).

By the Bruhat decomposition, the (U, U) -double cosets are parametrized by the elements of N , while the (B, B) -double cosets are parametrized by the elements of W .

We consider induced representations γ of the form ψ^G , for a linear representation ψ of U . Let

$$e = |U|^{-1} \sum_{u \in U} \psi(u^{-1})u$$

MSC2010: 20G40.

Keywords: representation theory, finite Chevalley groups.

be the primitive idempotent affording ψ in the group algebra $\mathbb{C}U$ of U over the field of complex numbers. Then $\gamma = \psi^G$ is afforded by the left $\mathbb{C}G$ -module $\mathbb{C}Ge$. The *Hecke algebra* of γ is the subalgebra $H = e\mathbb{C}Ge$ of $\mathbb{C}G$, and is isomorphic to $(\text{End}_{\mathbb{C}G} \mathbb{C}Ge)^\circ$. These representations and their Hecke algebras were first investigated by Gelfand and Graev [1962a; 1962b]. In particular, they introduced the important class of *Gelfand–Graev representations* of G , which are the induced representation ψ^G , for a linear representation ψ of U in *general position*, that is, $\psi|_{U_{\alpha_i}} \neq 1$ for each simple root subgroup U_{α_i} , $1 \leq i \leq n$, and $\psi|_{U_\alpha} = 1$ for each positive and not simple root α .

It is known (see [Gelfand and Graev 1962b] for the case of $G = \text{SL}_n(k)$ for a finite field k , and [Steinberg 1968, Theorem 49] for the general case) that the Hecke algebra H of a Gelfand–Graev representation is a commutative algebra, so that a Gelfand–Graev representation is multiplicity-free.

A basis for the Hecke algebra H of a Gelfand–Graev representation ψ^G is given by the nonzero elements of the form ene with $n \in N$. The *standard basis elements* are the nonzero elements of the form $c_n = \text{ind}(n) ene$, where $\text{ind}(n) = |U : nUn^{-1} \cap U|$. The structure constants for the standard basis elements, defined by the formulas

$$c_\ell c_m = \sum_n [c_\ell c_m : c_n] c_n,$$

with $\ell, m, n \in N^*$, are algebraic integers (here N^* is the set of elements $n \in N$ such that $ene \neq 0$).

The structure constants of H are given by the formula

$$[c_\ell c_m : c_n] = \sum_{ulu_1 = nvm^{-1} \in U\ell U \cap nU_{m-1}m^{-1}} \psi((uu_1)^{-1}v),$$

by [Curtis and Reiner 1981, Proposition 11.30], and the fact that $U\ell U \cap nU_{m-1}m^{-1}$ is a set of representatives of the left U -cosets in $U\ell U \cap nU_{m-1}m^{-1}U$. As in [Curtis 1988; 2009], $U_n = U \cap nU_{-n}^{-1}$ for $n \in N$. The structure constants are exponential sums involving the linear character ψ of U and combinatorial information about multiplication and intersections patterns of (U, U) -double cosets. The latter information is also given at least partially for the algebraic group $G(\bar{k})$ over the algebraic closure \bar{k} of k corresponding to G , with some questions about the geometry not completely settled at this time. A main result in the paper is an algorithm given in Section 4 for the solutions (u, u_1, v) of the equation $ulu_1 = nvm^{-1}$ in the formula above, so that in some sense the structure constants are computable. The approach taken here is based on the theory of *cells* U_τ developed in [Curtis 1988; 2009]. The algorithm for the solutions of the equations is a refined version of an algorithm for them given in [Curtis 2009, Theorem 2.1]. At the end of Section 4, some problems for further research are mentioned.

In case ψ^G is a Gelfand–Graev representation, the values of the irreducible representations of the commutative semisimple algebra H on standard basis elements are obtained as eigenvalues of matrices giving the regular representation of H and whose entries are the structure constants $[c_\ell c_m : c_n]$; see [Curtis 2009, Proposition 1.1].

Formulas for the structure constants based on different algorithms and a different set of representatives of the cosets of U were obtained by Simion [2015].

The irreducible representations of H were obtained in [Curtis 1993] using the results of Deligne and Lusztig [1976] on representations of G defined on the ℓ -adic cohomology of locally closed subsets of the algebraic group $G(\bar{k})$ with Frobenius endomorphism F on which the finite group G acts. The formulas for the irreducible representations of H in [Curtis 1993] involve a homomorphism of algebras $f_T : H \rightarrow \mathbb{C}T$ for each F -stable maximal torus T of G , proved using the character formula of Deligne and Lusztig [1976] for the virtual representations $R_{T,\theta}$. The homomorphisms f_T provide an approach to the representations of H , and are of independent interest (see [Bonnafé and Kessar 2008]).

A combinatorial approach to the representations of H based on the structure constants of the Hecke algebra H and the internal structure of the finite Chevalley group G is a main objective of this paper.

Two final sections contain examples in which a combinatorial construction of the homomorphisms f_T is obtained. These include the *Bessel functions over finite fields* of Gelfand and Graev [1962a], for the groups $SL_2(k)$ and k a finite field of odd characteristic, and a construction of the homomorphisms $f_T : H \rightarrow \mathbb{C}T$ for the split torus T in a general Chevalley group.

2. Background and preliminary results

For each root α , there is a homomorphism (see [Steinberg 1968, page 46]) $\varphi = \varphi_\alpha : SL_2(k) \rightarrow G$ such that φ takes

$$\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \rightarrow x_\alpha(t), \quad \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix} \rightarrow x_{-\alpha}(t), \quad \begin{pmatrix} 0 & t \\ -t^{-1} & 0 \end{pmatrix} \rightarrow w_\alpha(t) \in N, \quad \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \rightarrow h_\alpha(t) \in T$$

for all $t \in k$. The elements $w_\alpha(t)$ and $h_\alpha(t)$ are given by

$$w_\alpha(t) = x_\alpha(t)x_{-\alpha}(-t^{-1})x_\alpha(t), \quad h_\alpha(t) = w_\alpha(t)w_\alpha(1)^{-1},$$

by [Steinberg 1968, p. 30]. If $w = s_k \cdots s_1$ is a reduced expression of an element $w \in W$ then $\dot{w} = \dot{s}_k \cdots \dot{s}_1$, with $\dot{s}_i = w_{\alpha_i}(t_i)$ for some fixed choice of $t_i \in k^* = k - \{0\}$, is a representative in N of w which is independent of the choice of the reduced expression chosen, by [Steinberg 1968, Lemma 83, p. 242]. In what follows we assume that representatives $\dot{x} \in N$ of all elements $x \in W$ have been chosen in this way, for a fixed choice of representatives \dot{s}_i of the generators $s_i \in S$.

We may assume that

$$\dot{s}_k = \varphi_{\alpha_k} \left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$$

for a simple root α_k , $1 \leq k \leq n$.

Using the homomorphisms φ_α , we obtain the so-called SL_2 -IDENTITY:

$$\dot{s}_k^{-1} x_{\alpha_k}(t) \dot{s}_k = x_{\alpha_k}(-t^{-1}) \dot{s}_k h_{\alpha_k}(t) x_{\alpha_k}(-t^{-1}),$$

for a simple root α_k and $t \in k^*$ (cf. [Curtis 2009, Lemma 2.1]).

As in [Deodhar 1985], a *subexpression* τ of a fixed reduced expression $w = s_k \cdots s_1$ is a sequence $\tau = (\tau_k, \dots, \tau_1, \tau_0)$ of elements of W such that $\tau_i \tau_{i-1}^{-1} \in \{1, s_i\}$ for $i = 1, \dots, k$ and $\tau_0 = 1$. Then the set of terminal elements τ_k of subexpressions of $w = s_k \cdots s_1$ coincides with the set of elements $x \in W$ such that $x \leq w$ in the Chevalley–Bruhat order. In what follows, the *length* of an element $w \in W$ in terms of the generators $s_i \in S$ is denoted by $\ell(w)$. A subexpression $\tau = (\tau_k, \dots, \tau_1, \tau_0)$ is called a *K-sequence* relative to the triple $w = s_k \cdots s_1, x, y$ of elements of W if it satisfies conditions (2.10)(a-c) of [Kawanaka 1975]. It is understood that a *K-sequence* for the triple (w, x, y) is always given with reference to a fixed reduced expression $w = s_k \cdots s_1$. Let $J_\tau = \{j : \tau_j \tau_{j-1}^{-1} = s_j\} \cup \{0\}$. Then the defining conditions for a *K-sequence* state that $\tau_k x = y$ and

$$\ell(s_p \tau_j x) < \ell(\tau_j x)$$

for each $j \in J_\tau$ and p in the interval between j and the next element in J_τ (or simply all $p > j$ if j is the maximal element of J_τ). For each *K-sequence* τ , set

$$J_\tau^- = \{j \in J_\tau : \ell(s_j \tau_{j'} x) < \ell(\tau_{j'} x)\}$$

where $j' \in J_\tau$ is the predecessor of j , and define a pair of nonnegative integers by

$$a(\tau) = |J_\tau^-|, b(\tau) = k - |J_\tau| + 1 = \text{card}\{j > 0 : \tau_j \tau_{j-1}^{-1} = 1\}.$$

For each element $w \in W$, let $U_w = U \cap^w U_-$ where $U_- = {}^{w_0} U$ and w_0 is the element of maximal length in W . Then $U = U_w U_{ww_0}$ and $BwB = U_w \dot{w} B$, in both cases with uniqueness of expression. Let $w = s_k \cdots s_1$ be a reduced expression of $w \in W$. Then $U_w = U_{\alpha_k} \dot{s}_k U_{s_{k-1} \cdots s_1} \dot{s}_k^{-1}$ with uniqueness of expression. An element of U_w expressed in this way, for a fixed reduced expression of w , is said to be in *standard form* (see [Deodhar 1985, Lemma 2.2]), and can be assigned *coordinates* in the field k .

Let w, x, y be elements of W , and $\dot{w}, \dot{x}, \dot{y}$ corresponding elements of N . Let

$$U(w, x, y) = \{u \in U_w : u \dot{w} B \cap \dot{y} U_{x^{-1} \dot{x}^{-1}} \neq \emptyset\}.$$

Then $U(w, x, y)$ is independent of the choice of representatives $\dot{w}, \dot{x}, \dot{y}$ of w, x, y in N . Moreover, $U_w \dot{w} B \cap \dot{y} U_{x^{-1} \dot{x}^{-1}}$ is a set of representatives of the left B -cosets

in $BwB \cap y(BxB)^{-1}$, and its cardinality is the structure constant $[e_w e_x : e_y]$ of the standard basis elements e_w, e_x, e_y , for $w, x, y \in W$, in the Iwahori Hecke algebra.

The K -sequences were first applied by Kawanaka to prove the following result [Kawanaka 1975, Lemma 2.14b]. For a finite Chevalley group G over $k = F_q$ the nonzero structure constants of the Iwahori Hecke algebra are given by the formula

$$[e_w e_x : e_y] = |B\dot{w}B \cap \dot{y}U_{x^{-1}}\dot{x}^{-1}| = |U(w, x, y)| = \sum_{\tau} q^{a(\tau)}(q-1)^{b(\tau)}$$

where the sum is taken over all K -sequences τ for w, x, y , and $a(\tau)$ and $b(\tau)$ are the nonnegative integers defined above.

As a consequence, it follows that $U(w, x, y) \neq \emptyset$ if and only if there exist K -sequences for w, x, y (see also [Borel and Tits 1972, Remark 3.19], where the conditions are stated in a different way).

In [Curtis 1988] a geometric version of Kawanaka’s formula was proved. It states that $U(w, x, y)$, viewed as a subset of the algebraic group $G(\bar{k})$, is a disjoint union of subsets U_{τ} , which we shall call (in this paper) *cells*. The cells U_{τ} are subsets of $G(\bar{k})$ parametrized by K -sequences τ for w, x, y relative to a fixed reduced expression of the element w , with corresponding subsets U_{τ} , also called cells (defined in [Curtis 1988]), in the finite Chevalley group $G = G(k)$ (see Lemma 3.3 below for a review of the definition of cells). The result extends Deodhar’s decomposition ([Deodhar 1985], and [Curtis 2009, §4]) of the intersection $ByB \cap B_{-}xB$, viewed as subsets of the flag variety G/B in the algebraic group $G(\bar{k})$, with B_{-} the Borel subgroup opposite to B . Each cell U_{τ} is isomorphic (in bijective correspondence as a set, or isomorphic as a variety in $G(\bar{k})$) to a product,

$$U_{\tau} \cong \prod_{\alpha} U_{\alpha} \times \prod_{\beta} U_{\beta}^*$$

for certain subsets $\{\alpha\}$ and $\{\beta\}$ of cardinalities $a(\tau)$ and $b(\tau)$ of the positive root subgroups determined by τ and where U_{β}^* is the set of nonidentity elements in U_{β} . From the decomposition of $U(w, x, y)$ as a union of cells U_{τ} , it follows that $U_w \dot{w}B \cap \dot{y}U_{x^{-1}}\dot{x}^{-1}$ can be identified with the set of triples (u, b, v) with $u \in U_{\tau}$ for some τ , $b \in B$, and $v \in U_{x^{-1}}$ satisfying the equation $u\dot{w}b = \dot{y}v\dot{x}^{-1}$ with b and v uniquely determined by u by [Curtis 2009, Lemma 2.4].

3. Relations between cells

Let ℓ, m, n in N^* correspond to elements w, x, y in W . Then ℓ, m, n are multiples by elements of T of representatives $\dot{w}, \dot{x}, \dot{y}$ in N determined as above. The set $U_{\ell}\ell U \cap nU_{m^{-1}}m^{-1}$ will be obtained by an algorithm based on a fixed reduced expression $w = s_k \cdots s_1$ of the element $w \in W$ in terms of the generators $s_i \in S$, and the theory of cells U_{τ} associated with K -sequences τ for w, x, y .

As the cells U_τ are contained in the set $U(w, x, y)$ each element $u \in U_\tau$ satisfies a structure equation

$$u\dot{w}b = \dot{y}v\dot{x}^{-1}$$

with $b \in B$ and $v \in U_{x^{-1}}$. The subgroup B is a semidirect product $B = UT$, so one has $b = u_1s$ with $u_1 \in U, s \in T$, and it will be important to keep track of these factors in the discussion to follow.

In this section, it will be shown how elements $u \in U_\tau \subseteq U(w, x, y)$, with $\tau = (\tau_k, \dots, \tau_1, \tau_0)$ a K -sequence for w, x, y , are related to elements u' in cells $U_{\tau'}$ with $\tau' = (\tau_{k-1}, \dots, \tau_1, \tau_0)$ a K -sequence for $s_{k-1} \cdots s_1, x', y'$, and how the structure equations for u and u' are related. We keep in mind that $U(w, x, y) \neq \emptyset$ if and only if there exist K -sequences for w, x, y .

Lemma 3.1. *Let $\tau = (\tau_k, \dots, \tau_1, \tau_0)$ be a K -sequence for w, x, y for $k \geq 1$, and consider $\tau' = (\tau_{k-1}, \dots, \tau_0)$.*

- (i) τ' is a K -sequence for $s_k^{-1}w, x, s_k^{-1}y$ if $\tau_k\tau_{k-1}^{-1} = s_k$ and $\ell(s_k y) < \ell(y)$.
- (ii) τ' is a K -sequence for $s_k^{-1}w, x, y$ if $\tau_k\tau_{k-1}^{-1} = 1$ and $\ell(s_k y) < \ell(y)$.
- (iii) τ' is a K -sequence for $s_k^{-1}w, x, s_k^{-1}y$ if $\ell(s_k y) > \ell(y)$ and $\tau_k\tau_{k-1}^{-1} = s_k$.

It is understood that $\tau_0 = 1$ is a K -sequence for $(1, x, x)$ and that $a(\tau_0) = b(\tau_0) = 0$. These sets of conditions are the only possibilities for τ' to be a K -sequence, and one of them must occur.

We first note that either $\ell(s_k y) < \ell(y)$ or $\ell(s_k y) > \ell(y)$, since either $y^{-1}(\alpha_k) \in \Phi_+$ or $y^{-1}(\alpha_k) \in \Phi_-$. The proof then follows immediately from the definition of K -sequence (see the proof of Lemma 2.14 of [Kawanaka 1975]). For example, we verify that the condition $\ell(s_k y) > \ell(y)$ implies $\tau_k \neq \tau_{k-1}$, and hence $\tau_k\tau_{k-1}^{-1} = s_k$. Otherwise $\tau_k = \tau_{k-1}$, $\tau_{k-1}x = y$, and $k \notin J_\tau$. This implies that $\ell(s_k\tau_{k-1}x) < \ell(\tau_{k-1}x)$ by a defining property of K -sequences, and hence $\ell(s_k y) < \ell(y)$, contrary to assumption.

The next result is background for the relation between cells U_τ and $U_{\tau'}$, with τ and τ' as in the preceding lemma. It is a version of Lemma 2.3 of [Curtis 2009]. (Parts (i) and (ii) were misstated in that article and are corrected here. We also take the opportunity to correct the statement on page 220 of [Curtis 2009] that the cells U_τ are invariant under conjugation by elements of T ; this was not shown there.)

Lemma 3.2. *Let $w = s_k \cdots s_1$ be a reduced expression with $k \geq 1$ and let $x, y \in W$. Then $U(w, x, y)$ is either empty or is related to sets $U(s_k^{-1}w, x', y')$, with x' and y' depending on the K -sequence τ associated with w, x, y as follows.*

- (i) *Let $\ell(s_k y) < \ell(y)$ and assume $\tau_k\tau_{k-1}^{-1} = s_k$. Then $\dot{s}_k U_{s_k^{-1}w} \dot{s}_k^{-1} \cap U(w, x, y)$ is either empty or*

$$\dot{s}_k U_{s_k^{-1}w} \dot{s}_k^{-1} \cap U(w, x, y) = \dot{s}_k U(s_k^{-1}w, x, s_k^{-1}y) \dot{s}_k^{-1}.$$

- (ii) Let $\ell(s_k y) < \ell(y)$ and assume $\tau_k \tau_{k-1}^{-1} = 1$. Then the part $U(w, x, y)^b$ of $U(w, x, y)$ not in $\dot{s}_k U_{s_k^{-1}w} \dot{s}_k^{-1}$ consists of the elements $u = x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1}$, with $x_{\alpha_k}(t) \in U_{\alpha_k}^*$ and $\tilde{u} \in U_{s_k^{-1}w}$ such that $\pi(x_{\alpha_k}(-t^{-1})\tilde{u}) \in U(s_k^{-1}w, x, y)$, and $t \in k^*$; here π is the projection $\pi : U \rightarrow U_{s_k^{-1}w}$ accompanying the decomposition $U = U_{s_k^{-1}w} U_{s_k^{-1}w w_0}$. The map

$$u = x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \rightarrow \pi(x_{\alpha_k}(-t^{-1})\tilde{u})$$

from $U(w, x, y)^b$ to $U(s_k^{-1}w, x, y)$ is surjective. There is a bijection of sets $U(w, x, y)^b \cong U_{\alpha_k}^* \times U(s_k^{-1}w, x, y)$.

- (iii) Let $\ell(s_k y) > \ell(y)$ and $\tau_k \tau_{k-1}^{-1} = s_k$. Then

$$U(w, x, y) = U_{\alpha_k} \dot{s}_k U(s_k^{-1}w, x, s_k^{-1}y) \dot{s}_k^{-1}$$

and there is a bijection of sets $U(w, x, y) \cong U_{\alpha_k} \times U(s_k^{-1}w, x, s_k^{-1}y)$.

The proof is included in the proof of Lemma 2.3 of [Curtis 2009].

Lemma 3.3. Let w, x, y be elements of W and let $w = s_k \cdots s_1$ be a reduced expression for w . Let $\tau = (\tau_k, \dots, \tau_1, \tau_0)$ be a K -sequence for w, x, y with $\tau_0 = 1$, and let U_τ be the corresponding cell, viewed as a subset of $U(w, x, y) \subseteq U_w$. Let $\tau' = (\tau_{k-1}, \dots, \tau_1, \tau_0)$ be a K -sequence for $s_{k-1} \cdots s_1, x', y'$ as in one of the cases in Lemma 3.1, and let $U_{\tau'}$ be the corresponding cell in $U(s_{k-1} \cdots s_1, x', y')$. The construction of the cell U_τ from $U_{\tau'}$, reviewed below, defines a surjective map of sets $\lambda : U_\tau \rightarrow U_{\tau'}$. Let $U_\tau(\bar{k})$ and $U_{\tau'}(\bar{k})$ be the corresponding cells in the algebraic group $G(\bar{k})$ over the algebraic closure \bar{k} of k . Then the map $\lambda : U_\tau(\bar{k}) \rightarrow U_{\tau'}(\bar{k})$, defined as in part (i), is a surjective morphism of algebraic sets, defined over k .

The construction of $U_\tau(\bar{k})$ from $U_{\tau'}(\bar{k})$ was given in the three cases of Lemma 3.1 in the proof of Theorem 1.6 of [Curtis 1988] and in [Curtis 2009, page 220], and will be reviewed here in the case of the algebraic group $G(\bar{k})$. We abbreviate $U_\tau(\bar{k})$ to U_τ , etc.

(i) $\tau_k \tau_{k-1}^{-1} = s_k$ and $\ell(s_k y) < \ell(y)$. In this case, we have $U_\tau \subseteq \dot{s}_k U_{s_k^{-1}w} \dot{s}_k^{-1}$ and $U_{\tau'} \subseteq U(s_k^{-1}w, x, s_k^{-1}y)$, and one has $U_\tau = \dot{s}_k^{-1} U_{\tau'} \dot{s}_k$. The map $\lambda : u \rightarrow \dot{s}_k^{-1} u \dot{s}_k$ is clearly a surjective morphism from U_τ to $U_{\tau'}$ and is defined over k because \dot{s}_k belongs to the finite Chevalley group $G(k)$.

(ii) $\tau_k \tau_{k-1}^{-1} = 1$ and $\ell(s_k y) < \ell(y)$. This time U_τ is in the part of $U(w, x, y)$ which is not contained in $\dot{s}_k U_{s_k^{-1}w} \dot{s}_k^{-1}$ and consists of the elements $x_{\alpha_k}(t_k) \dot{s}_k \tilde{u} \dot{s}_k^{-1}$ such that $t_k \neq 0$, $\tilde{u} \in U_{s_k^{-1}w}$ and $\pi(x_{\alpha_k}(-t_k^{-1})\tilde{u}) \in U_{\tau'}$, where π is the projection $U \rightarrow U_{s_k^{-1}w}$ associated with the factorization $U = U_{s_k^{-1}w} U_{s_k^{-1}w w_0}$. The map

$$\lambda : x_{\alpha_k}(t_k) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \rightarrow \pi(x_{\alpha_k}(-t_k^{-1})\tilde{u})$$

is a surjective morphism defined over k from U_τ to $U_{\tau'}$.

(iii) $\tau_k \tau_{k-1}^{-1} = s_k$ and $\ell(s_k y) > \ell(y)$. In this situation, we have $U_\tau = U_{\alpha_k} \dot{s}_k U_{\tau'} \dot{s}_k^{-1}$, in terms of the factorization: $U_w = U_{\alpha_k} \dot{s}_k U_{s_{k-1} \cdots s_1} \dot{s}_k^{-1}$, and $U_{\tau'} \subseteq U(s_k^{-1} w, x, s_k y)$. Then the map

$$\lambda : x_{\alpha_k}(t_k) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \rightarrow \tilde{u}$$

is a surjective morphism defined over k from U_τ to $U_{\tau'}$ (as the projection from U_w to $\dot{s}_k U_{s_{k-1} \cdots s_1} \dot{s}_k^{-1}$ in the factorization given above, followed by the inner automorphism by an element of $G(k)$). This completes our discussion of the proof of the lemma.

We now have a reduction process for cells, $U_\tau \rightarrow U_{\tau'}$, as in the preceding lemma. Let $u \in U_\tau$ correspond to $u' \in U_{\tau'}$ as in the lemma. Then the structure equation $u \dot{w} u_1 s = \dot{y} v \dot{x}^{-1}$ satisfied by u corresponds to the structure equation $u' \dot{s}_k^{-1} \dot{w} u_1' s' = \dot{y}' v' \dot{x}'^{-1}$ satisfied by u' , with uniquely determined factors $\{u, u_1, s, v\}$ and $\{u', u_1', s', v'\}$. The next lemma shows how the elements u', u_1', v' in U and $s' \in T$ are related to u, u_1, v in U and $s \in T$, using the standard form and facts about the multiplicative structure of the Chevalley group such as the decomposition $U = U_x U_{xw_0}$ for elements $x \in W$. It is also shown that the process is reversible, assuming $u \in U_\tau$ is known.

Lemma 3.4. *Suppose that the cell $U_\tau \subseteq U(w, x, y)$ maps onto the cell $U_{\tau'} \subseteq U(s_{k-1} \cdots s_1, x', y')$ as in cases (i)–(iii) of Lemma 3.1, and let the structure equation satisfied by $u \in U_\tau$ be $u \dot{w} u_1 s = \dot{y} v \dot{x}^{-1}$ with factors $u \in U_\tau, u_1 \in U, s \in T$, and $v \in U_{x^{-1}}$ uniquely determined by $u \in U_\tau$. Let $u \rightarrow u' = \lambda(u)$ with $u' \in U_{\tau'}$ as in Lemma 3.3, and consider the structure equation satisfied by u' with factors $u' \in U, s' \in T, u_1' \in U$, and $v' \in U_{x^{-1}}$ in each of the cases. Then the factors u', u_1', s' and v' are given as in the proof of the lemma. Conversely, assuming $u \in U_\tau$ is known, u_1, s and v are obtained from u', s', u_1' , and v' , as shown in the proof of the lemma.*

In case (i), we have $\tau_k \tau_{k-1}^{-1} = s_k$, $\ell(s_k y) < \ell(y)$ and $U_{\tau'} = \dot{s}_k^{-1} U_\tau \dot{s}_k$. Then the equation satisfied by $u \in U_\tau$ is $u \dot{w} u_1 s = \dot{y} v \dot{x}^{-1}$ with $u = \dot{s}_k u' \dot{s}_k^{-1}$ and $u' \in U_{\tau'}$. It becomes the equation for $u' \in U_{\tau'}$ after multiplication by \dot{s}_k^{-1} , and the lemma is proved in this case.

For the proof in case (ii) recall that $\tau_k \tau_{k-1}^{-1} = 1$ and $\ell(s_k y) < \ell(y)$. Then, using the standard form for u , the structure equation satisfied by $u = x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \in U_\tau$ is

$$x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \dot{w} u_1 s = \dot{y} v \dot{x}^{-1},$$

with $x_{\alpha_k}(t) \in U_{\alpha_k}^*$, $\tilde{u} \in U_{s_k^{-1} w}$, $s \in T, u_1 \in U, v \in U_{x^{-1}}$. We want to derive an equation of the form

$$u' \dot{s}_k^{-1} \dot{w} u_1' s' = \dot{y}' v' \dot{x}'^{-1}$$

with $u' = \pi(x_{\alpha_k}(-t^{-1}) \tilde{u}) \in U_{\tau'}, s' \in T, u_1' \in U$ and $v' \in U_{x^{-1}}$, where π is the

projection $U \rightarrow U_{s_k^{-1}w}$ as in the proof of Lemma 3.3. As

$$\dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} = \dot{s}_k^{-1} x_{\alpha_k}(-t^{-1}) \dot{s}_k,$$

we can multiply the equation for $x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1} \in U_\tau$ by $\dot{s}_k \dot{s}_k^{-1}$ and apply the SL_2 -IDENTITY from Section 2 to obtain

$$\dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k h_{\alpha_k}(t) x_{\alpha_k}(-t^{-1}) \tilde{u} \dot{s}_k^{-1} \dot{w} u_1 s = \dot{y} v \dot{x}^{-1}.$$

One has $x_{\alpha_k}(-t^{-1}) \tilde{u} = \pi(x_{\alpha_k}(-t^{-1}) \tilde{u}) u^*$ for $u^* \in U_{s_k^{-1}w w_0}$ so the equation becomes

$$\pi(x_{\alpha_k}(-t^{-1}) \tilde{u}) \dot{s}_k^{-1} \dot{w} (\dot{s}_k^{-1} \dot{w})^{-1} u^* \dot{s}_k^{-1} \dot{w} u_1 s = \dot{y} \dot{y}^{-1} (\dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} \dot{s}_k^2 h_{\alpha_k}(t))^{-1} \dot{y} v \dot{x}^{-1}$$

where $(\dot{s}_k^{-1} \dot{w})^{-1} u^* \dot{s}_k^{-1} \dot{w} \in U$ because $u^* \in U_{s_k^{-1}w w_0}$. Note also that $\dot{s}_k^2 h_{\alpha_k}(t) = h_{\alpha_k}(-t)$, and that the right side of the equation is

$$\dot{y} \dot{y}^{-1} (h_{\alpha_k}(-t))^{-1} \dot{y} \dot{y}^{-1} \dot{s}_k (x_{\alpha_k}(-t^{-1})^{-1}) \dot{s}_k^{-1} \dot{y} v \dot{x}^{-1}.$$

Because $\ell(s_k y) < \ell(y)$, one has $\dot{y}^{-1} \dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} \dot{y} \in U$, and we consider first the case where $\dot{y}^{-1} \dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} \dot{y} \in U_{x^{-1}}$. Then the equation above becomes the structure equation for $u' \in U_{\tau'}$ with $u' = \pi(x_{\alpha_k}(-t^{-1}) \tilde{u})$, $u'_1 = (\dot{s}_k^{-1} \dot{w})^{-1} u^* \dot{s}_k^{-1} \dot{w} u_1$,

$$s' = s (\dot{x} \dot{y}^{-1} (h_{\alpha_k}(-t))^{-1} \dot{y} \dot{x}^{-1})^{-1},$$

and v' is

$$\dot{y}^{-1} \dot{s}_k (x_{\alpha_k}(-t^{-1}))^{-1} \dot{s}_k^{-1} \dot{y} v \in U_{x^{-1}}$$

conjugated by $\dot{y}^{-1} h_{\alpha_k}(-t)^{-1} \dot{y}$. Note that $\dot{y}^{-1} h_{\alpha_k}(-t)^{-1} \dot{y} \in T$, and that we have conjugated this element past $\dot{y} v \dot{x}^{-1}$ and brought the result to the left-hand side as a factor of s' . We have also used the fact that $U_{x^{-1}}$ is invariant under conjugation by elements of T .

For the reversibility, consider u' , u'_1 , s' , v' and $u = x_{\alpha_k}(t) \dot{s}_k \tilde{u} \dot{s}_k^{-1}$ in U_τ . Then $x_{\alpha_k}(-t^{-1}) \tilde{u} = u' u^*$, so $u^* = (u')^{-1} (x_{\alpha_k}(-t^{-1}) \tilde{u})$. Then $s = s' (\dot{x} \dot{y}^{-1} h_{\alpha_k}(-t) \dot{y} \dot{x}^{-1})$, $u_1 = ((\dot{s}_k^{-1} \dot{w})^{-1} u^* \dot{s}_k^{-1} \dot{w})^{-1} u'_1$ and

$$v = (\dot{y}^{-1} \dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} \dot{y})^{-1} \dot{y}^{-1} h_{\alpha_k}(-t)^{-1} \dot{y} v' \dot{y}^{-1} h_{\alpha_k}(-t) \dot{y} \in U_{x^{-1}},$$

completing the proof of reversibility in this case, using the fact again that $U_{x^{-1}}$ is invariant under conjugation by elements of T .

Now we have to discuss the case $\dot{y}^{-1} \dot{s}_k x_{\alpha_k}(-t^{-1}) \dot{s}_k^{-1} \dot{y} \notin U_{x^{-1}}$. Then we obtain a new formula for v' as follows. We have

$$\dot{y}^{-1} \dot{s}_k x_{\alpha_k}(-t^{-1})^{-1} \dot{s}_k^{-1} \dot{y} v = v' v'',$$

with uniquely determined factors $v' \in U_{x^{-1}}$ and $v'' \in U_{x^{-1}w_0}$. Then $v'' \dot{x}^{-1} = \dot{x}^{-1} \dot{x} v'' \dot{x}^{-1}$ with $\dot{x} v'' \dot{x}^{-1} \in U$. Then the structure equation in this case for $u' =$

$\pi(x_{\alpha_k}(-t^{-1})\tilde{u})$ has as factors $u'_1 = (\dot{s}_k^{-1}\dot{w})^{-1}u^*\dot{s}_k^{-1}\dot{w}u_1s'(\dot{x}v''\dot{x}^{-1})^{-1}(s')^{-1}$, s' as in the first case, and v' as defined at the beginning of this paragraph, conjugated by $\dot{y}^{-1}h_{\alpha_k}(-t)^{-1}\dot{y}$.

For the reversibility in this case, suppose we have $u = x_{\alpha_k}(t)\dot{s}_k\tilde{u}\dot{s}_k^{-1} \in U_\tau$, $u' = \pi(x_{\alpha_k}(-t^{-1})\tilde{u})$, and u'_1, s' , and v' as above. We have to solve for u_1, s , and v . Then, after reversing the conjugation by $\dot{y}^{-1}h_{\alpha_k}(-t)^{-1}\dot{y}$, we obtain

$$(\dot{y}^{-1}\dot{s}_kx_{\alpha_k}(-t^{-1})\dot{s}_k^{-1}\dot{y})^{-1}v' = v(v'')^{-1}.$$

Then v is the projection of the left-hand side in $U_{x^{-1}}$. Then from

$$u'_1 = (\dot{s}_k^{-1}\dot{w})^{-1}u^*\dot{s}_k^{-1}\dot{w}u_1s'(\dot{x}v''\dot{x}^{-1})^{-1}(s')^{-1},$$

we can express u^* as the projection of $x_{\alpha_k}(-t^{-1})\tilde{u}$ in $U_{s_k^{-1}ww_0}$ and $(\dot{x}v''\dot{x}^{-1})^{-1}$ from $x_{\alpha_k}(-t^{-1})$ and v' , so we recover u_1 . Finally the element s is computed as in the first case. Therefore we have obtained u_1, s , and v , completing the proof of reversibility in this case.

In case (iii), the structure equation satisfied by $u = x_{\alpha_k}(t)\dot{s}_k\tilde{u}\dot{s}_k^{-1} \in U_\tau$ is

$$x_{\alpha_k}(t)\dot{s}_k\tilde{u}\dot{s}_k^{-1}\dot{w}u_1s = \dot{y}v\dot{x}^{-1}$$

as in case (ii). This time $\ell(s_k y) > \ell(y)$, so $\dot{y}^{-1}x_{\alpha_k}(t)\dot{y} \in U$ and the structure equation for $\tilde{u} \in U_{\tau'}$ becomes

$$\tilde{u}\dot{s}_k^{-1}\dot{w}u_1s = \dot{s}_k^{-1}\dot{y}\dot{y}^{-1}x_{\alpha_k}(t)^{-1}\dot{y}v\dot{x}^{-1},$$

and the rest of the proof is handled as in case (ii), depending on whether $\dot{y}^{-1}x_{\alpha_k}(t)\dot{y} \in U_{x^{-1}}$ or not. This completes the proof of the lemma.

4. Solution of the structure equation

Let $\ell, m, n \in N^*$ be representatives of w, x, y in W , and let $w = s_k \cdots s_1$ be a fixed reduced expression of w . From the Introduction, the structure constants of standard basis elements c_ℓ, c_m, c_n of the Hecke algebra H of an induced representation $\gamma = \psi^G$ of G are given by the formula

$$[c_\ell c_m : c_n] = \sum_{u\ell u_1 = nvm^{-1} \in U\ell U \cap nU_{m-1}m^{-1}} \psi((uu_1)^{-1}v),$$

where $U\ell U \cap nU_{m-1}m^{-1}$ is the set of elements $u \in U, u_1 \in U, v \in U_{m-1}$ satisfying the equation

$$u\ell u_1 = nvm^{-1}.$$

The elements ℓ, m, n are multiples of $\dot{w}, \dot{x}, \dot{y}$ by elements of T , $\ell = \dot{w}s, m = \dot{x}s',$ etc. for elements s, s', s'' in T . The equations $u\ell u_1 = nvm^{-1}$ as above can be

rewritten in the form $u\dot{w}\hat{u}_1\hat{s} = \dot{y}\hat{v}\dot{x}^{-1}$ for $u \in U$, $\hat{s} = s(\dot{x}s''(s')^{-1}\dot{x}^{-1})^{-1} \in T$, $\hat{u}_1 = su_1s^{-1} \in U$, $\hat{v} = s''v(s'')^{-1} \in U_{x^{-1}}$, using the fact that the subsets U_x , for $x \in W$, are invariant under conjugation by elements of T . Each element $u \in U$ satisfying the equation above belongs to the set $U(w, x, y)$, and consequently $u \in U_\tau$ for a cell U_τ defined by a K -sequence τ for the elements $\{w, x, y\}$ in W . An algorithm for the solutions u, \hat{u}_1, \hat{v} of these equations is the main result of this section.

As $u \in U_\tau$ is known in terms of the root subgroups from the main result of [Curtis 1988], the problem is to calculate \hat{u}_1 and \hat{v} in terms of a given expression of u .

The following remarks may throw some light on these problems. Let τ be a K -sequence for w, x, y in W , and let U_τ be the corresponding cell in $U(w, x, y)$. Each element $u \in U_\tau$ satisfies a structure equation

$$u\dot{w}b = \dot{y}v\dot{x}^{-1}$$

with $b \in B$ and $v \in U_{x^{-1}}$. It is known that the elements b and v in the structure equation are uniquely determined by u [Curtis 2009, Lemma 2.4]. A main theorem in [Curtis 2009] was an inductive construction of the solutions of the structure equation. Theorem 4.1 below gives more information, and in a sense, calculates b and v from an expression of u in standard form using a fixed reduced expression of w . In particular, this result determines, for each element $u \in U_\tau$, the solutions (u_1, v) , of the equations $u\ell u_1 = nv m^{-1}$ with $u \in U_\tau$, $u_1 \in U$, and $v \in U_{x^{-1}}$, needed for the structure constants of H .

Theorem 4.1. *Let w, x, y be elements of W , and let U_τ be a cell associated with a K -sequence τ for w, x, y , and a fixed reduced expression $w = s_k \cdots s_1$ for w . The algorithm given below determines the set of elements $s \in T$, $u_1 \in U$ and $v \in U_{x^{-1}}$ satisfying the equation $u\dot{w}u_1s = \dot{y}v\dot{x}^{-1}$, for a given element $u \in U_\tau$. The possibility that the set of solutions is empty is not excluded.*

Let w, x, y , the cell U_τ , and $w = s_k \cdots s_1$ be as in the hypothesis of the theorem. Let u be a fixed element of U_τ . With these as a starting point, the algorithm gives the elements $s \in T$, $u_1 \in U$ and $v \in U_{x^{-1}}$ satisfying the equation stated in the theorem. It is proved by induction on $\ell(w)$.

We begin with the case $\ell(w) = 1$, so $\dot{w} = \dot{s}_1$ and let $\tau = (\tau_1, \tau_0)$ be a K -sequence for (s_1, x, y) corresponding to one of the three cases in Lemma 3.1.

Case (i). $\tau_1 = s_1$, $\ell(s_1y) < \ell(y)$. Then $U_{\tau_0} = 1$, $U_\tau = \dot{s}_1U_{\tau_0}\dot{s}_1^{-1} = 1$, $s_1x = y$ and it is easily proved using Lemma 83 of [Steinberg 1968] that $\dot{s}_1\dot{x} = \dot{y}$. Then there is a unique solution, namely $(1, 1, 1)$, of the structure equation $u\dot{s}_1b = \dot{s}_1\dot{x}v(\dot{x})^{-1}$.

Case (ii). $\tau_1 = \tau_0$, and $\ell(s_1y) < \ell(y)$. In this case the definition of K -sequence implies $x = y$. We also have (by part (ii) of the proof of Lemma 3.3) $U_\tau = U_{\alpha_1}^*$. First assume $\ell(x) = 1$. Then the assumptions imply that $x = s_1$, $\dot{x} = \dot{y} = \dot{s}_1$, and for each element $u \in U_{\alpha_1}^*$ there is a unique solution of the structure equation $u\dot{s}_1b = \dot{s}_1v\dot{s}_1^{-1}$

by the SL_2 -IDENTITY, so that quadruples (u, s, u_1, v) with $u \in U_\tau$, $b = u_1s \in B$, and $v \in U_{x^{-1}}$ satisfying the equation exist, and are known. Now let $\ell(x) > 1$; then $\ell(s_1x) < \ell(x)$ implies $\dot{x} = \dot{s}_1\dot{x}_1$ with $\ell(s_1x_1) > \ell(x_1)$. For each $v \in U_{\alpha_1}^*$ one has $\dot{x}_1^{-1}v\dot{x}_1 \in U$ because $\ell(s_1x_1) > \ell(x_1)$. Moreover

$$\dot{s}_1\dot{x}_1(\dot{x}_1^{-1}v\dot{x}_1)\dot{x}_1^{-1}\dot{s}_1^{-1} = \dot{s}_1v\dot{s}_1^{-1} \in U_-$$

so $\dot{x}_1^{-1}v\dot{x}_1 \in U_{x^{-1}}$. The unique solution of the structure equation $u\dot{s}_1b = \dot{s}_1v\dot{s}_1^{-1}$ with $u \in U_{\alpha_1}^*$ from the case $\ell(x) = 1$ now yields the unique solution $(u, b, \dot{x}_1^{-1}v\dot{x}_1)$ of the structure equation for (s_1, x, y) , namely

$$u\dot{s}_1b = \dot{y}\dot{x}_1^{-1}v\dot{x}_1\dot{x}^{-1}$$

as $\dot{y}\dot{x}_1^{-1} = \dot{s}_1$ and $\dot{x}_1\dot{x}^{-1} = \dot{s}_1^{-1}$. Note that in case (ii) there is no solution of the structure equation $u\dot{s}_1b = \dot{y}v\dot{x}^{-1}$ in case $u = 1$ and $x = y$ as this would contradict the fact that $B\dot{y}^{-1}\dot{s}_1B \neq B\dot{x}^{-1}B$ by the uniqueness part of the Bruhat decomposition.

Case (iii). $\tau_1 = s_1$, and $\ell(s_1y) > \ell(y)$. In this case $U_\tau = U_{\alpha_1}$ and the unique solution of the structure equation $u\dot{s}_1b = \dot{y}v\dot{x}^{-1}$ for $u \in U_{\alpha_1}$ is

$$(u, h_{\alpha_1}(-1), \dot{y}^{-1}u\dot{y}),$$

for each $u \in U_{\alpha_1}$, noting that $\dot{y}^{-1}u\dot{y} \in U_{x^{-1}}$, and $\dot{y}\dot{x}^{-1} = \dot{s}_1^{-1} = \dot{s}_1h_{\alpha_1}(-1)$ by [Steinberg 1968, Lemma 83] again. This completes the discussion of the solutions of the structure equation for the case $\ell(w) = 1$.

We now proceed to the general case, with $\ell(w) > 1$. Let τ be a K -sequence for w, x, y and let $u \in U_\tau$. Let u correspond to $u' = \lambda(u) \in U_{\tau'}$ (as in Lemma 3.3) for the K -sequence τ' for $s_{k-1} \dots s_1, x', y'$ in one of the three cases of Lemma 3.1, and let

$$u's_{k-1} \dots s_1u'_1s' = \dot{y}'v'\dot{x}'$$

be the structure equation satisfied by u' . By the induction hypothesis, the factors u'_1, s' , and v' of the structure equation for u' are determined by u' . As $u' = \lambda(u)$, the elements u_1, s , and v satisfying the equation $u\dot{w}b = u\dot{w}u_1s = \dot{y}v\dot{x}^{-1}$ are determined by u , using Lemma 3.4. This completes the proof of the theorem.

At the beginning of the section, it was explained how the solutions $u \in U, u_1 \in U$, and $v \in U_{x^{-1}}$ of the equations $ulu_1 = nvm^{-1}$, for $\ell, m, n \in N^*$, required for the formulas for the structure constants are obtained from the solutions $u \in U_\tau, u_1 \in U, s \in T, v \in U_{x^{-1}}$ of the equations $u\dot{w}u_1s = \dot{y}v\dot{x}^{-1}$ solved by the algorithm in Theorem 4.1. For this step, it is necessary to determine the elements t, t', t'' in T such that $\ell = \dot{w}t, m = \dot{x}t', n = \dot{y}t''$, in order to transform the first set of equations to the second by the algorithms for multiplication the Chevalley group. This information can be obtained from Steinberg's proof of Theorem 49 in [Steinberg 1968, §14], in case ψ^G is a Gelfand–Graev representation. The theorem states that

the Hecke algebra H of a Gelfand–Graev representation is a commutative algebra, and the proof is obtained by constructing a certain antiautomorphism f of the Chevalley group G whose extension to the group algebra is at the same time an antiautomorphism of the group algebra and whose restriction to the Hecke algebra H is the identity. As shown in [Steinberg 1968], the representatives $\ell \in N^*$ of the basis elements of H have the form $t\dot{w}$ for elements $w \in W$ such that $w = w_0w_\pi$, where w_0 is the element of maximal length in W and w_π is the element of maximal length in the subgroup of the Weyl group generated by the reflections taken from a subset π of the set of simple roots, and t is an element of T such that $t\dot{w}$ is fixed by the antiautomorphism f . From the discussion on page 262 of [Steinberg 1968], it follows directly that $\dot{w}t$, with $w = w_0w_\pi$ as above, represents a basis element of H , fixed by the antiautomorphism f , whenever t commutes with w_π .

We recall the connection between the solutions of the equation $u\ell u_1 = nvm^{-1}$ and the solutions of the equation $u\dot{w}\hat{u}_1\hat{s} = \dot{y}\hat{v}\hat{x}^{-1}$ with $\ell = \dot{w}s$, $m = \dot{x}s'$, $n = \dot{y}s''$, $s, s', s'' \in T$ and $\hat{u}_1 = su_1s^{-1} \in U$, $\hat{v} = s''v(s'')^{-1} \in U_{x^{-1}}$, for $u \in U$. For a solution $u \in U$, we have $u \in U(w, x, y)$ so $u \in U_\tau$ for a K -sequence τ for w, x, y . We can now state a formula for the structure constants $[c_\ell c_m : c_n]$ based on Theorem 4.1.

Corollary 4.2. *The structure constants are given by the formula*

$$[c_\ell c_m : c_n] = \sum_{\tau} \sum_{u \in U_\tau} \psi((uu_1)^{-1}v)$$

where for each K -sequence τ for w, x, y , the sum is taken over solutions of the equation $u\dot{w}\hat{u}_1\hat{s} = \dot{y}\hat{v}\hat{x}^{-1}$ obtained by Theorem 4.1, with $u \in U_\tau$ and $\hat{u}_1, \hat{v}, \hat{s}$ satisfying the conditions $\hat{u}_1 = su_1s^{-1} \in U$, $\hat{v} = s''v(s'')^{-1} \in U_{x^{-1}}$ and $\hat{s} = s(\dot{x}s''(s')^{-1}\dot{x}^{-1})^{-1} \in T$. If there are no solutions satisfying these conditions, then the structure constant is zero.

We end this section with two problems for further research.

1. The first problem is to apply the algorithm obtained in Theorem 4.1 and Corollary 4.2 to obtain formulas for the structure constants $[c_\ell c_m : c_n]$ which can be used to give a combinatorial proof of the existence of the homomorphisms f_T mentioned in the Introduction.

2. The second problem is to develop a theory of cells for Chevalley groups over a p -adic field K , using the Bruhat decomposition for these groups obtained by Iwahori and Matsumoto [1965].

5. Example: application to $SL_2(k)$

Let G be the Chevalley group $SL_2(k)$ for a finite field k of odd characteristic, and let H be the Hecke algebra of a Gelfand–Graev representation of G . Gelfand and

Graev [1962a] stated formulas for the structure constants of the standard basis of H , and calculated the irreducible representations of H .

As an application of the ideas in Section 4, we shall calculate the structure constants of H relative to the standard basis of the Hecke algebra, and apply them to give a self-contained proof, different from the one obtained by Gelfand and Graev, of formulas for the irreducible representations of H (for another approach, using the Deligne-Lusztig character formula, see [Curtis 1993, §5]).

We begin with a Gelfand–Graev character ψ^G of $G = \mathrm{SL}_2(k)$, for a linear character ψ of U in general position. Then we may assume that

$$\psi \begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix} = \chi(\alpha), \quad \begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix} \in U, \quad \alpha \in k$$

for a nontrivial additive character χ on k . The standard basis elements of the Hecke algebra H of ψ^G are the elements

$$c_\lambda = q e_\psi n_\lambda e_\psi, \quad n_\lambda = \begin{pmatrix} 0 & \lambda \\ -\lambda^{-1} & 0 \end{pmatrix}, \quad \lambda \neq 0, \quad q = |k|$$

together with the identity element e_ψ and one other basis element $e_{-1} = e_\psi \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} e_\psi$, where

$$e_\psi = |U|^{-1} \sum_{u \in U} \psi(u^{-1})u$$

as in the Introduction.

Lemma 5.1. *The algebra H is commutative with identity element e_ψ . One has $e_{-1}^2 = e_\psi$, and $e_{-1}c_\lambda = c_{-\lambda}$ for each $\lambda \neq 0$. The other nonzero structure constants of H for the standard basis elements are as follows. For c_λ, c_μ, c_ν as above one has*

$$[c_\lambda c_\mu : c_\nu] = \chi(\lambda\mu\nu^{-1} + \lambda\mu^{-1}\nu + \lambda^{-1}\mu\nu),$$

and

$$[c_\lambda c_\lambda : e_{-1}] = q, \quad [c_\lambda c_{-\lambda} : e_\psi] = q,$$

for $\lambda, \mu, \nu \neq 0$ in k , and $q = |k|$.

The structure constants are computed using the formula at the beginning of §4 and the solutions of the structure equation

$$u\ell u_1 = nv(m)^{-1}$$

with $\ell, m, n \in N$ (see [Curtis 2009, §3] for more details).

The group $G = \mathrm{SL}_2(k)$ can be viewed as the group of fixed points by the usual Frobenius endomorphism F of the semisimple algebraic group $\mathrm{SL}_2(\bar{k})$, over the algebraic closure \bar{k} of k . There are two conjugacy classes of F -stable maximal tori in $\mathrm{SL}_2(\bar{k})$ with representatives in the finite group G given by the *split torus*

T_0 consisting of the matrices $\begin{pmatrix} \mu & 0 \\ 0 & \mu^{-1} \end{pmatrix}$ with $\mu \neq 0$ in k , and the Coxeter torus T_1 , isomorphic to the set C of elements ξ of norm 1 in the quadratic extension of k , that is, $\xi^{q+1} = 1$. The main theorem on the representations of the Hecke algebra H of a Gelfand–Graev representation of G states that the irreducible representations of H factor through the group algebra of one of the maximal tori of G . More precisely, one has:

Theorem 5.2. *Each irreducible representation f of the Hecke algebra H of a Gelfand–Graev representation of G can be factored as*

$$f = \theta \circ f_T,$$

where f_T is a homomorphism, independent of θ , of H into the group algebra of a maximal torus T of G , and θ is an irreducible representation of the group algebra of the maximal torus. The homomorphisms from H into the group algebras of the two types of maximal tori are given as follows. For the split torus T_0 , consisting of diagonal matrices with entries in k^* , the homomorphism $f_{T_0} : H \rightarrow \mathbb{C}T_0$ is given by

$$f_{T_0}(c_\lambda) = \sum_t \chi(\lambda(t + t^{-1})) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}, \quad t \in k^*, \quad \text{and} \quad f_{T_0}(e_{-1}) = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix},$$

where c_λ is a standard basis element of H as above. For the Coxeter torus, the homomorphism $f_{T_1} : H \rightarrow \mathbb{C}C$ is given by

$$f_{T_1}(c_\lambda)(\xi) = -\chi(\lambda(\xi + \xi^{-1})), \quad \xi \in C, \quad \text{and} \quad f_{T_1}(e_{-1}) = \xi_{-1},$$

where ξ_{-1} is the unique element in C of order two.

Lemma 5.3. *Let $a, b \in k$. Then:*

- (i)
$$\sum_{t \in k^*} \chi(at) = -1 + q\delta_{a,0}.$$
- (ii)
$$\sum_{t \in k^*} \chi(at + bt^{-1}) = \sum_{t \in k^*} \chi(t + abt^{-1}) + q\delta_{a,0}\delta_{b,0}.$$
- (iii) *For the Coxeter torus C , we first note that $\xi + \xi^{-1} \in k$ because $\xi^{q+1} = 1$ for $\xi \in C$ implies that $\xi + \xi^{-1} = \xi + \xi^q \in k$. Let $\xi \in C, \eta \in F_{q^2}$. Then*

$$\sum_{\xi \in C} \chi(\xi\eta + \xi^q\eta^q) = - \sum_{t \in k^*} \chi(t + \eta\eta^q t^{-1}) + \delta_{\eta,0}q.$$

We refer to [Chang 1976, Lemma 1.2]. Part (iii) is proved using an analysis of quadratic equations over k . The result, and extensions of it to F -stable maximal tori in general finite reductive groups, are suggested by the Davenport–Hasse Theorem on Gauss sums.

For the proof that f_{T_0} is a homomorphism, let c_λ, c_μ, c_ν be standard basis elements of H as above. Then

$$c_\lambda c_\mu = \sum_\nu [c_\lambda c_\mu : c_\nu] c_\nu + \delta_{\lambda, \mu} q e_{-1} + \delta_{\lambda, -\mu} q e_\psi,$$

with the structure constants as in Lemma 5.1. We have

$$\begin{aligned} f_{T_0}(c_\lambda) f_{T_0}(c_\mu) &= \sum_{t \in k^*} \sum_{s \in k^*} \chi(\lambda(ts + t^{-1}s^{-1})) \chi(\mu(s + s^{-1})) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \\ &= \sum_t \sum_s \chi((\lambda t + \mu)s + (\lambda t^{-1} + \mu)s^{-1}) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \\ &= \sum_t \sum_{s'} \chi(s' + (\lambda^2 + \mu^2 + \lambda\mu(t + t^{-1})(s')^{-1})) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \end{aligned}$$

by Lemma 5.3(ii), and have to show this is equal to

$$\begin{aligned} \sum_t \sum_\nu [c_\lambda c_\mu : c_\nu] f_{T_0}(c_\nu)(t) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} + \delta_{\lambda, \mu} q \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} + \delta_{\lambda, -\mu} q \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ = \sum_t \sum_\nu \chi(\lambda\mu\nu^{-1} + \lambda\mu^{-1}\nu + \lambda^{-1}\mu\nu + \nu(t + t^{-1})) \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}, \end{aligned}$$

etc. The result is clear, by another application of Lemma 5.3(ii), in case $\lambda \neq \pm\mu$.

Now let $\lambda = \mu$. The expressions to be checked agree except possibly at $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$. At $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, the first expression becomes

$$\sum_s \chi(\lambda(t+1)s + \lambda(t^{-1}+1)s^{-1})$$

with $t = -1$, which is $q - 1$ by Lemma 5.3. The second expression at $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ becomes

$$\sum_\nu \chi(\nu + (2\lambda^2 + (-2\lambda^2))\nu^{-1}) + q = -1 + q$$

by Lemma 5.3 again, completing the proof in this case. The proof in case $\lambda = -\mu$ is similar and will be omitted.

For the homomorphism from H into the group algebra of C , we first have

$$\begin{aligned} f_{T_1}(c_\lambda) f_{T_1}(c_\mu) &= \sum_{\xi \in C} \sum_{\eta \in C} \chi(\lambda(\xi\eta + (\xi\eta)^{-1}) + \mu(\eta + \eta^{-1})) \xi \\ &= \sum_{\xi} \sum_{\eta} \chi((\lambda\xi + \mu)\eta + (\lambda\xi^{-1} + \mu)\eta^{-1}) \xi \\ &= - \sum_{\xi} \sum_{t \in k^*} \chi(t + (\lambda\xi + \mu)(\lambda\xi^q + \mu)t^{-1}) \xi \end{aligned}$$

by Lemma 5.3(ii). We have to show that this is equal to

$$\begin{aligned} & \sum_{\xi \in C} \sum_{v \in k^*} \chi(\lambda \mu v^{-1} + \lambda \mu^{-1} v + \lambda^{-1} \mu v) f_{T_1}(c_v) \xi + \delta_{\lambda, \mu} q f_{T_1}(e_{-1}) + \delta_{\lambda, -\mu} q f_{T_1}(e_{\psi}) \\ &= - \sum_{\xi \in C} \sum_{v \in k^*} \chi(v(\xi + \xi^{-1} + \lambda \mu^{-1} + \lambda^{-1} \mu) + v^{-1} \lambda \mu) \xi + \delta_{\lambda, \mu} q f_{T_1}(e_{-1}) + \delta_{\lambda, -\mu} q f_{T_1}(e_{\psi}) \\ &= - \sum_{\xi} \sum_{v'} \chi(v' + v'^{-1}(\xi + \xi^q) \lambda \mu + \lambda^2 + \mu^2) \xi + \delta_{\lambda, \mu} q f_{T_1}(e_{-1}) + \delta_{\lambda, -\mu} q f_{T_1}(e_{\psi}), \end{aligned}$$

where we have used Lemma 5.3(ii) and Lemma 5.1 for the structure constant formula. Together, these formulas prove the multiplication formula in case $\lambda \neq \pm \mu$.

In case $\lambda = \mu$, it is only necessary to check the expressions at $\xi = \xi_{-1}$, where ξ_{-1} is the unique element of C such that $\xi_{-1}^2 = 1$, $\xi_{-1} \neq 1$, so $\xi_{-1} = -1$ in F_{q^2} . The contribution from the first expression is

$$\sum_{\eta} \chi(\lambda(\xi_{-1} + 1)\eta + \lambda(\xi_{-1}^{-1} + 1)\eta^{-1}) = - \sum_{t \in k^*} \chi(t + (2\lambda^2 + \lambda^2(\xi_{-1} + \xi_{-1}^q))t^{-1}) + q$$

by Lemma 5.3(iii). As $\xi_{-1} + \xi_{-1}^q = -2$ in F_{q^2} , this expression is equal to $1 + q$ by Lemma 5.3. For the second expression at ξ_{-1} we obtain

$$- \sum_{v'} \chi(v' + (v')^{-1}((\xi_{-1} + \xi_{-1}^q)\lambda^2 + 2\lambda^2)) + q = q + 1,$$

completing the proof in this case.

For the remaining case $\lambda = -\mu$ it is only necessary to check both expressions at $\xi = 1$ and this is immediate.

Corollary 5.4. *The formulas for the irreducible representations of the Hecke algebra H are*

$$f(c_{\lambda}) = \theta \circ f_T(c_{\lambda}) = \sum_{t \in T} \chi(\lambda(t + t^{-1}))\theta(t), \quad f(e_{-1}) = \theta(-1)$$

for the split torus T , and an irreducible representation θ of T , and

$$f(c_{\lambda}) = \pi \circ f_{T_1}(c_{\lambda}) = - \sum_{\xi \in C} \chi(\lambda(\xi + \xi^{-1}))\pi(\xi), \quad f(e_{-1}) = \pi(-1)$$

for the Coxeter torus T_1 represented by C , and an irreducible representation π of C .

The fact that all the irreducible representations of H are obtained in this way follows by a counting argument.

Gelfand and Graev [1962a] obtained these formulas, and pointed out that they are similar to the integral formulas for Bessel functions over \mathbb{C} (see [Whittaker and Watson 1927, Chapter XVII]). They mentioned that the formulas in Corollary 5.4 can be called *Bessel functions over finite fields*.

The group $G = \mathrm{SL}_2(k)$, for $q = |k|$ odd, has two classes of Gelfand–Graev representations. For a determination of the irreducible characters of G , and how they appear in the Gelfand–Graev representations, including for example the subtle cases of those of degree $\frac{1}{2}(q+1)$ and $\frac{1}{2}(q-1)$, see [Gelfand and Graev 1962a, §4] and [Digne and Michel 1991, §15.9].

6. Example: the homomorphisms f_T associated with principal series representations of finite Chevalley groups

We return to the set-up described in the Introduction, with G a Chevalley group over a finite field k , with a Borel subgroup $B = UT$ containing the torus T , and Weyl group W . Let ψ^G be a Gelfand–Graev representation of G , and $H = e\mathbb{C}Ge$ the Hecke algebra associated with it, with

$$e = |U|^{-1} \sum_{u \in U} \psi(u^{-1})u$$

(remember that H is a commutative algebra!). In this section, we give a character theoretic construction of a homomorphism $f_T : H \rightarrow \mathbb{C}T$ and the resulting irreducible representations of H . An open problem is to find a combinatorial construction of homomorphisms f_T for twisted tori T . Such a result would define a family of functions associated with the Hecke algebra H and maximal tori in G , starting from the Bessel functions over k in the case of $\mathrm{SL}_2(k)$ and the Coxeter torus C . A proof would require information about the structure constants, and extensions of Lemma 5.3(iii), which would be of independent interest. Homomorphisms $f_T : H \rightarrow \mathbb{C}T$ from a Gelfand–Graev Hecke algebra H to the group algebra of a maximal torus are known to exist, for a connected reductive algebraic group G defined over a finite field, with Frobenius endomorphism F ([Curtis 1993] and [Bonnafe and Kessar 2008]), and are derived using the trace formula in ℓ -adic cohomology.

We are concerned with the *principal series* representations of G . These are the induced representations λ^G , where λ is a linear character of the Borel subgroup B with U in its kernel, and the irreducible representations of G which occur as constituents of λ^G for some choice of lambda. We require the following result of Kilmoyer [1978, Proposition 6.1].

Lemma 6.1. *Let ψ^G be a fixed Gelfand–Graev character of G . Each induced character λ^G , as above, contains a unique irreducible constituent ξ_λ which appears with multiplicity one in both λ^G and the Gelfand–Graev character ψ^G .*

As in [Curtis 1993], we introduce the notation \mathbf{a} for the element of the group algebra $\mathbb{C}G$ given by

$$\mathbf{a} = \sum_{g \in G} \alpha(g^{-1})g$$

for a complex valued function α on G . If α is an irreducible character of G , then α is a multiple of the central primitive idempotent in the group algebra associated with α .

Theorem 6.2. *Let l_λ and x_λ be the elements of the group algebra corresponding to the induced character λ^G and the irreducible character ξ_λ as in Lemma 6.1. Then*

$$el_\lambda = ex_\lambda \neq 0$$

and affords an irreducible representation $f_{\xi_\lambda} : H \rightarrow \mathbb{C}^$ of H of degree one, such that*

$$h(ex_\lambda) = f_{\xi_\lambda}(h)ex_\lambda, \quad h \in H.$$

The representation f_{ξ_λ} is the restriction to H of the unique irreducible character of the group algebra $\mathbb{C}G$ obtained from the character ξ_λ of G as in Lemma 6.1.

By Lemma 6.1, the class function $l_\lambda = x_\lambda + y$ where y is a linear combination of central primitive idempotents corresponding to irreducible characters of G which do not appear in the Gelfand–Graev representation ψ^G . Then $ey = 0$, so $el_\lambda = ex_\lambda \neq 0$ and ex_λ is a nonzero multiple of the primitive central idempotent in H affording the irreducible representation f_{ξ_λ} of H of degree one, as in the statement of the theorem, by [Curtis and Reiner 1981, Corollary 11.26 and Theorem 11.25]. The last statement of the theorem also follows from [Curtis and Reiner 1981, Theorem 11.25].

Theorem 6.3. *Let λ be an irreducible character of B with U in its kernel. Let ξ_λ be the irreducible character of G which appears with multiplicity one in λ^G and in the Gelfand–Graev character ψ^G , and let $f_{\xi_\lambda} : H \rightarrow \mathbb{C}$ be the irreducible representation of the Hecke algebra H of the Gelfand–Graev representation ψ^G defined in Theorem 6.2. There exists a unique homomorphism of algebras $f_T : H \rightarrow \mathbb{C}T$, independent of λ , such that, for each linear character λ of T , one has*

$$f_{\xi_\lambda}(h) = \tilde{\lambda} \circ f_T(h), \quad h \in H,$$

where $\tilde{\lambda}$ is the extension of $\lambda : T \rightarrow \mathbb{C}$ to the group algebra $\mathbb{C}T$. The homomorphism f_T is given by the formula $f_T(c_n) = \sum_{t \in T} f_T(c_n)(t)t$, where

$$f_T(c_n)(t) = \text{ind } n|B|^{-1}|U|^{-1} \sum_{g \in G, u \in U, gung^{-1} = tu'} \psi(u^{-1}),$$

for a standard basis element c_n of H and $gung^{-1} = tu'$, $t \in T$, $u' \in U$, is an element of B which projects onto the element $t \in T$ by the homomorphism $B \rightarrow T$. If there are no solutions to the equation $gung^{-1} = tu'$, then $f_T(c_n)(t) = 0$.

By the proof of Theorem 6.2, the representation f_{ξ_λ} of the Hecke algebra H is the restriction to H of the unique irreducible character (see Lemma 6.1) ξ_λ extended

to the group algebra $\mathbb{C}G$. Moreover, the proof of Theorem 6.2 shows that

$$f_{\xi_\lambda}(h) = \lambda^G(h), \quad h \in H,$$

with λ^G extended to the group algebra, because $eI_\lambda = ex_\lambda$ and ex_λ affords the representation f_{ξ_λ} of $H = e\mathbb{C}Ge \subseteq \mathbb{C}G$. In more detail, h lies in H , and is viewed as an element of the group algebra $\mathbb{C}G$. Then $\lambda^G(h)$ is the trace of the action of h on a module M affording the induced character λ^G . As $he = h$, the trace is computed on the module eM , which is one dimensional, and affords the representation f_{ξ_λ} of H , by Theorem 6.2.

For a standard basis element c_n of H , we have

$$c_n = \text{ind } n \text{ en}e = |U|^{-1} \sum_{u_1 n u_2 \in UnU} \psi(u_1^{-1} u_2^{-1}) u_1 n u_2,$$

by [Curtis and Reiner 1981, Proposition 11.30(i)]. Then, with λ^G extended to the group algebra, we obtain

$$\lambda^G(c_n) = |U|^{-1} \sum_{u_1 n u_2 \in UnU} \psi(u_1^{-1} u_2^{-1}) \lambda^G(u_1 n u_2) = \text{ind } n |U|^{-1} \sum_{u \in U} \psi(u^{-1}) \lambda^G(un).$$

We have used the fact that the double coset UnU contains $\text{ind } n$ one sided cosets. For the induced character we have, by [Curtis and Reiner 1981, 10.3],

$$\lambda^G(un) = |B|^{-1} \sum_{g \in G} \dot{\lambda}(g^{-1}ung),$$

where $\dot{\lambda}(x) = 0$ if $x \notin B$. Then $\dot{\lambda}(g^{-1}ung) \neq 0$ only if $g^{-1}ung = u't$ with $u' \in U$ and $t \in T$, and in that case, $\dot{\lambda}(g^{-1}ung) = \lambda(t)$. Therefore

$$\lambda^G(c_n) = \text{ind } n |B|^{-1} |U|^{-1} \sum_{t \in T} \sum_{g^{-1}ung=u't} \psi(u^{-1}) \lambda(t).$$

Then, for $t \in T$,

$$f_T(c_n)(t) = \text{ind } n |B|^{-1} |U|^{-1} \sum_{g^{-1}ung=u't} \psi(u^{-1})$$

is independent of λ , and we have

$$f_{\xi_\lambda}(h) = \tilde{\lambda} \circ f_T(h), \quad h \in H.$$

The facts that $f_T : H \rightarrow \mathbb{C}T$ is a homomorphism of algebras and is a uniquely determined linear map with the factorization property stated in the theorem both follow from the orthogonality relations for the linear characters λ of T . This completes the proof of the theorem.

It is a nice exercise to derive the formula for the homomorphism $f_{T_0} : H \rightarrow \mathbb{C}T_0$ given in Theorem 5.2 from the statement of the preceding theorem.

Theorem 6.3, for principal series representations of finite Chevalley groups, is a special case of Theorem 4.2 in [Curtis 1993] for representations $R_{T,\theta}$ of connected reductive algebraic groups defined over finite fields. The point of including it here is that in the special case of principal series representations, it is possible to give a combinatorial proof of the existence of the homomorphisms f_T .

References

- [Bonnafé and Kessar 2008] C. Bonnafé and R. Kessar, “On the endomorphism algebras of modular Gelfand–Graev representations”, *J. Algebra* **320**:7 (2008), 2847–2870. MR 2010f:20046 Zbl 1197.20037
- [Borel and Tits 1972] A. Borel and J. Tits, “Compléments à l’article: “Groupes réductifs””, *Inst. Hautes Études Sci. Publ. Math.* 41 (1972), 253–276. MR 47 #3556 Zbl 0254.14018
- [Chang 1976] B. Chang, “Decomposition of Gelfand–Graev characters of $LGL_3(q)$ ”, *Comm. Algebra* **4**:4 (1976), 375–401. MR 53 #5766 Zbl 0356.20051
- [Chevalley 1955] C. Chevalley, “Sur certains groupes simples”, *Tôhoku Math. J. (2)* **7** (1955), 14–66. MR 17,457c Zbl 0066.01503
- [Curtis 1988] C. W. Curtis, “A further refinement of the Bruhat decomposition”, *Proc. Amer. Math. Soc.* **102**:1 (1988), 37–42. MR 89d:20038 Zbl 0656.20049
- [Curtis 1993] C. W. Curtis, “On the Gelfand–Graev representations of a reductive group over a finite field”, *J. Algebra* **157**:2 (1993), 517–533. MR 94i:20075 Zbl 0787.20026
- [Curtis 2009] C. W. Curtis, “On problems concerning the Bruhat decomposition and structure constants of Hecke algebras of finite Chevalley groups”, *Michigan Math. J.* **58**:1 (2009), 213–230. MR 2011a:20007 Zbl 1241.20004
- [Curtis and Reiner 1981] C. W. Curtis and I. Reiner, *Methods of representation theory with applications to finite groups and orders, vol I.*, John Wiley & Sons, New York, 1981. MR 82i:20001 Zbl 0469.20001
- [Deligne and Lusztig 1976] P. Deligne and G. Lusztig, “Representations of reductive groups over finite fields”, *Ann. of Math. (2)* **103**:1 (1976), 103–161. MR 52 #14076 Zbl 0336.20029
- [Deodhar 1985] V. V. Deodhar, “On some geometric aspects of Bruhat orderings, I: a finer decomposition of Bruhat cells”, *Invent. Math.* **79**:3 (1985), 499–511. MR 86f:20045 Zbl 0563.14023
- [Digne and Michel 1991] F. Digne and J. Michel, *Representations of finite groups of Lie type*, London Mathematical Society Student Texts **21**, Cambridge University Press, Cambridge, 1991. MR 92g:20063 Zbl 0815.20014
- [Gelfand and Graev 1962a] I. M. Gelfand and M. I. Graev, “Categories of group representations and the classification problem of irreducible representations”, *Dokl. Akad. Nauk SSSR* **146** (1962), 757–760. In Russian; translated in *Sov. Math. Dokl.* **3** (1962), 1378–1381. MR 26 #237
- [Gelfand and Graev 1962b] I. M. Gelfand and M. I. Graev, “Construction of irreducible representations of simple algebraic groups over a finite field”, *Dokl. Akad. Nauk SSSR* **147** (1962), 529–532. In Russian; translated in *Sov. Math. Dokl.* **3** (1962), 1646–1649. MR 26 #6271 Zbl 0119.26902
- [Iwahori and Matsumoto 1965] N. Iwahori and H. Matsumoto, “On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups”, *Inst. Hautes Études Sci. Publ. Math.* 25 (1965), 5–48. MR 32 #2486 Zbl 0228.20015
- [Kawanaka 1975] N. Kawanaka, “Unipotent elements and characters of finite Chevalley groups”, *Osaka J. Math.* **12**:2 (1975), 523–554. MR 52 #5784 Zbl 0314.20031

- [Kilmoyer 1978] R. W. Kilmoyer, “Principal series representations of finite Chevalley groups”, *J. Algebra* **51**:1 (1978), 300–319. MR 81e:20047 Zbl 0389.20008
- [Simion 2015] I. I. Simion, “On refinements of the Bruhat decomposition”, *J. Algebraic Combin.* (online publication August 2015).
- [Steinberg 1968] R. Steinberg, “Lectures on Chevalley groups”, mimeographed notes, Yale University, New Haven, 1968. MR 57 #6215 Zbl 1196.22001
- [Whittaker and Watson 1927] E. T. Whittaker and G. N. Watson, *A course of modern analysis: an introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions*, 4th ed., Cambridge University Press, Cambridge, 1927. MR 97k:01072 Zbl 53.0180.04

Received March 4, 2015. Revised July 24, 2015.

CHARLES W. CURTIS
DEPARTMENT OF MATHEMATICS AND INSTITUTE OF THEORETICAL SCIENCE
UNIVERSITY OF OREGON
EUGENE, OR 97403
UNITED STATES
cwc@uoregon.edu

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2015 is US \$420/year for the electronic version, and \$570/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 279 No. 1-2 December 2015

In memoriam: Robert Steinberg

Robert Steinberg (1922–2014): In memoriam V. S. VARADARAJAN	1
Cellularity of certain quantum endomorphism algebras HENNING H. ANDERSEN, GUSTAV I. LEHRER and RUIBIN ZHANG	11
Lower bounds for essential dimensions in characteristic 2 via orthogonal representations ANTONIO BABIC and VLADIMIR CHERNOUSOV	37
Cocharacter-closure and spherical buildings MICHAEL BATE, SEBASTIAN HERPEL, BENJAMIN MARTIN and GERHARD RÖHRLE	65
Embedding functor for classical groups and Brauer–Manin obstruction EVA BAYER-FLUCKIGER, TING-YU LEE and RAMAN PARIMALA	87
On maximal tori of algebraic groups of type G_2 CONSTANTIN BELI, PHILIPPE GILLE and TING-YU LEE	101
On extensions of algebraic groups with finite quotient MICHEL BRION	135
Essential dimension and error-correcting codes SHANE CERNELE and ZINOVY REICHSTEIN	155
Notes on the structure constants of Hecke algebras of induced representations of finite Chevalley groups CHARLES W. CURTIS	181
Complements on disconnected reductive groups FRANÇOIS DIGNE and JEAN MICHEL	203
Extending Hecke endomorphism algebras JIE DU, BRIAN J. PARSHALL and LEONARD L. SCOTT	229
Products of partial normal subgroups ELLEN HENKE	255
Lusztig induction and ℓ -blocks of finite reductive groups RADHA KESSAR and GUNTER MALLE	269
Free resolutions of some Schubert singularities MANOJ KUMMINI, VENKATRAMANI LAKSHMIBAI, PRAMATHANATH SASTRY and C. S. SESHADRI	299
Free resolutions of some Schubert singularities in the Lagrangian Grassmannian VENKATRAMANI LAKSHMIBAI and REUVEN HODGES	329
Distinguished unipotent elements and multiplicity-free subgroups of simple algebraic groups MARTIN W. LIEBECK, GARY M. SEITZ and DONNA M. TESTERMAN	357
Action of longest element on a Hecke algebra cell module GEORGE LUSZTIG	383
Generic stabilisers for actions of reductive groups BENJAMIN MARTIN	397
On the equations defining affine algebraic groups VLADIMIR L. POPOV	423
Smooth representations and Hecke modules in characteristic p PETER SCHNEIDER	447
On CRDAHA and finite general linear and unitary groups BHAMA SRINIVASAN	465
Weil representations of finite general linear groups and finite special linear groups PHAM HUU TIEP	481
The pro- p Iwahori Hecke algebra of a reductive p -adic group, V (parabolic induction) MARIE-FRANCE VIGNÉRAS	499
Acknowledgement	531



0030-8730(2015)279:1;1-1