

*Pacific
Journal of
Mathematics*

**ON WEAK CONVERGENCE
OF QUASI-INFINITELY DIVISIBLE LAWS**

ALEXEY KHARTOV

Volume 322 No. 2

February 2023

ON WEAK CONVERGENCE OF QUASI-INFINITELY DIVISIBLE LAWS

ALEXEY KHARTOV

We study a new class of so-called quasi-ininitely divisible laws, which is a wide natural extension of the well-known class of infinitely divisible laws through the Lévy–Khinchin representations. We are interested in criteria of weak convergence within this class. Under rather natural assumptions, we state assertions, which connect a weak convergence of quasi-ininitely divisible distribution functions with one special type of convergence of their Lévy–Khinchin spectral functions. The latter convergence is not equivalent to the weak convergence. So we complement known results by Lindner, Pan, and Sato (2018) in this field.

1. Introduction

This paper is devoted to the questions concerning weak convergence within a new class of so-called quasi-ininitely divisible probability laws.

Let F be a distribution function of a probability law on the real line \mathbb{R} with the characteristic function

$$f(t) := \int_{\mathbb{R}} e^{itx} dF(x), \quad t \in \mathbb{R}.$$

Recall that F (and the corresponding law) is called *infinitely divisible* if for every positive integer n there exists a distribution function $F_{1/n}$ such that $F = (F_{1/n})^{*n}$, where $*$ is the convolution, i.e., F is n -fold convolution power of $F_{1/n}$. It is well known that F is infinitely divisible if and only if the characteristic function f is represented by the *Lévy–Khinchin formula*:

$$(1) \quad f(t) = \exp \left\{ it\gamma + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG(x) \right\}, \quad t \in \mathbb{R},$$

with some $\tau > 0$, *shift parameter* $\gamma \in \mathbb{R}$, and with a bounded nondecreasing *spectral function* $G : \mathbb{R} \rightarrow \mathbb{R}$ that is assumed to be right-continuous with condition $G(-\infty) = 0$ (all over the paper, $G(\pm\infty)$ denote the limits at $\pm\infty$ correspondingly).

MSC2020: 60E05, 60E07, 60E10, 60F05.

Keywords: quasi-ininitely divisible laws, characteristic functions, the Lévy–Khinchin formula, weak convergence.

We use $u \mapsto \frac{1}{\tau} \sin(\tau u)$ as the *centering function* in the integral in (1) following Zolotarev [33; 34]. If (1) holds for some $\tau = \tau_0 > 0$, then it holds for any $\tau > 0$, where γ will depend on τ , but G will not. It is well known that the *spectral pair* (γ, G) is uniquely determined by f and hence by F . The Lévy–Khinchin formula plays a fundamental role in probability theory; it also has a lot of applications in related fields (see [3; 30; 31]).

It turns out that there exists a rather wide class of probability laws that are very similar to infinitely divisible laws. This class of so-called *quasi-infinitely divisible* laws was introduced by Lindner and Sato [22]. Following them, a distribution function F (and the corresponding law) is called *quasi-infinitely divisible* if its characteristic function f admits the representation (1) with some *shift parameter* $\gamma \in \mathbb{R}$, *spectral function* $G : \mathbb{R} \rightarrow \mathbb{R}$ of bounded variation on \mathbb{R} (not necessarily monotone), and for some (any) $\tau > 0$. Here G is assumed to be right-continuous with condition $G(-\infty) = 0$ as before and so f (and F) uniquely determines the *spectral pair* (γ, G) (see [13, p. 80]). Observe that, due to the Jordan decomposition, we can represent

$$G(x) = G^+(x) - G^-(x), \quad x \in \mathbb{R},$$

with some bounded nondecreasing functions G^+ and G^- on \mathbb{R} . Also we can always write $\gamma = \gamma^+ - \gamma^-$ with some numbers γ^+ and γ^- from \mathbb{R} . Then it is clear that

$$f(t) = f^+(t)/f^-(t), \quad t \in \mathbb{R},$$

where f^+ and f^- are characteristic functions of some two infinitely divisible distribution functions F^+ and F^- with the spectral pairs (γ^+, G^+) and (γ^-, G^-) correspondingly, and so $F * F^- = F^+$. Starting from this point of view, it is rather natural to call distribution function F (and the corresponding law) *rationally infinitely divisible*. So every infinitely divisible law is quasi-infinitely divisible, but the converse is not true. There are a lot of interesting examples of quasi-infinitely divisible laws, which are not infinitely divisible (see [13, pp. 82–83; 24, p. 165; 25, pp. 123–124]). Moreover, it seems that the class of quasi-infinitely divisible laws is essentially wider than the class of infinitely divisible ones. In particular, it is clearly seen within discrete probability laws (see [2; 18; 19; 23]).

Various forms of definition and the first detailed analysis of the class of quasi-infinitely divisible laws on \mathbb{R} was performed in [23], the multivariate case is considered in the recent papers [6; 7; 21]. There are some results for discrete probability laws in this field (see [1; 2; 17; 18; 19]) and for mixed laws (see [4; 5]). It should be noted that quasi-infinitely divisible laws now have interesting applications in theory of stochastic processes (see [22; 28]), number theory (see [26]), physics (see [11; 12]), and insurance mathematics (see [32]).

We now focus on a weak convergence of quasi-infinitely divisible laws. Recall that, by definition, the sequence $(F_n)_{n \in \mathbb{N}}$ (where \mathbb{N} is the set of positive integers) of

distribution functions *weakly converges* to a distribution function F (we will write $F_n \xrightarrow{w} F, n \rightarrow \infty$) if

$$(2) \quad \int_{\mathbb{R}} h(x) dF_n(x) \rightarrow \int_{\mathbb{R}} h(x) dF(x), \quad n \rightarrow \infty$$

for any bounded continuous function $h : \mathbb{R} \rightarrow \mathbb{R}$. It is a well known fact that this is equivalent to the following convergence:

$$(3) \quad F_n(x) \rightarrow F(x), \quad n \rightarrow \infty \quad \text{for any } x \in S,$$

where S is an arbitrary dense subset of \mathbb{R} and, in particular, it can be chosen as the set of all continuity points of F . The latter convergence is usually called *weak* too (see [25]).

The weak convergence is also introduced for the class of real-valued functions of bounded variation on the real line (or for corresponding signed measures). Following Bogachev [10], it is analogously defined by (2), but instead of $F_n, n \in \mathbb{N}$, and F we write some functions of bounded variation $G_n, n \in \mathbb{N}$, and G correspondingly. We will save the notation $G_n \xrightarrow{w} G, n \rightarrow \infty$, in this case. It should be noted that weak convergence here is not equivalent to the analog of convergence (3) with functions of bounded variation (see [10, Section 1.4]).

There are rather general results by Lindner, Pan, and Sato in [23] concerning the weak convergence of quasi-infinately divisible distribution functions. The authors state the conditions under which the weak convergence of distribution functions implies the weak convergence of the corresponding spectral functions together with the convergence of the shift parameters and vice versa. Namely, let $(F_n)_{n \in \mathbb{N}}$ be a sequence of quasi-infinately divisible distribution functions and let (γ_n, G_n) be the spectral pair of F_n for every $n \in \mathbb{N}$. Let F be a quasi-infinately divisible distribution function with the spectral pair (γ, G) . Then the results from [23] are in fact the following: (1) If $\gamma_n \rightarrow \gamma$ and $G_n \xrightarrow{w} G, n \rightarrow \infty$, then $F_n \xrightarrow{w} F, n \rightarrow \infty$. (2) If we suppose $F_n \xrightarrow{w} F, n \rightarrow \infty$, then, under some assumptions on tightness and uniform boundedness for $(G_n)_{n \in \mathbb{N}}$, we have $\gamma_n \rightarrow \gamma$ and $G_n \xrightarrow{w} G, n \rightarrow \infty$. Here we omitted some details, the full formulation will be given in Section 3.

In this work we complement the results by Lindner, Pan, and Sato [23]. We connect the weak convergence of quasi-infinately divisible distribution functions with one type of convergence of their spectral functions. The latter convergence is a special modification of the convergence (3) (see the next section for details), and we think that it is more natural and explicit than the weak convergence for the class of functions of bounded variation. A very similar convergence has appeared in [10, Theorem 1.4.7], but we are not aware of the existence of a definition for such a convergence. So we introduce the necessary definition in Section 2. We also show that the introduced convergence for functions of bounded variation follows from

the pointwise convergence of their Fourier–Stieltjes transforms under some natural assumptions. This and other close propositions are key tools for our main results devoted to the weak convergence of quasi-infinitely divisible distribution functions. The main results are presented in Section 3. All proofs are provided in Section 4.

2. Preliminaries and tools

Let us consider the class of all functions $G : \mathbb{R} \rightarrow \mathbb{R}$ of bounded variations on \mathbb{R} . Since we will be interested in the functions G , which generate the Lebesgue–Stieltjes signed measures μ_G , we will focus only on right-continuous functions G . So for the measures we will have $\mu_G((a, b]) = G(b) - G(a)$ for all $a, b \in \mathbb{R}$ and $a \leq b$. Recall that all intervals $(a, b]$ constitute the generating semiring for μ_G . So if there are two right-continuous functions G_1 and G_2 of bounded variations such that $G_2(x) = G_1(x) + C$, $x \in \mathbb{R}$, where $C \in \mathbb{R}$ is a constant, then the corresponding measures are the same. Therefore we will consider only functions G that satisfy $G(-\infty) = 0$.

Let \mathbf{V} denote the class of all functions $G : \mathbb{R} \rightarrow \mathbb{R}$ of bounded variation on \mathbb{R} , which are right-continuous at every point $x \in \mathbb{R}$ and satisfy $G(-\infty) = 0$. For every $G \in \mathbf{V}$ its total variation on \mathbb{R} will be denoted by $\|G\|$ and the total variation on $(-\infty, x]$ by $|G|(x)$, $x \in \mathbb{R}$. So we have

$$(4) \quad |G(x)| \leq |G|(x) \leq \|G\| \quad \text{for any } x \in \mathbb{R}$$

and $|G|(+\infty) = \|G\|$.

We now introduce a special type of convergence on the class \mathbf{V} . Suppose that a whole sequence $(G_n)_{n \in \mathbb{N}}$ and a function G are from \mathbf{V} . We say that $(G_n)_{n \in \mathbb{N}}$ *converges basically* to G , and write $G_n \Rightarrow G$, $n \rightarrow \infty$, if each of its subsequences contains a further subsequence $(G_{n_k})_{k \in \mathbb{N}}$ such that

$$G_{n_k}(x_2) - G_{n_k}(x_1) \rightarrow G(x_2) - G(x_1), \quad k \rightarrow \infty$$

for any $x_1, x_2 \in \mathbb{R}$ except at most a countable set, which in general depends on the choice of the subsequences.

Let us show that the basic convergence is equivalent to the weak convergence for distribution functions. Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of distribution functions and let F be a distribution function. Suppose that $F_n \xrightarrow{w} F$, $n \rightarrow \infty$. Then we have (3), where S is the set of all continuity points of F . Hence

$$F_n(x_2) - F_n(x_1) \rightarrow F(x_2) - F(x_1), \quad n \rightarrow \infty \quad \text{for all } x_1, x_2 \in S.$$

Since $\mathbb{R} \setminus S$ is at most countable set, we conclude that $F_n \Rightarrow F$, $n \rightarrow \infty$, by definition. We now suppose that $F_n \Rightarrow F$, $n \rightarrow \infty$. Let $(F_{n_k})_{k \in \mathbb{N}}$ be an arbitrary subsequence

of $(F_n)_{n \in \mathbb{N}}$ such that

$$(5) \quad F_{n_k}(x_2) - F_{n_k}(x_1) \rightarrow F(x_2) - F(x_1), \quad k \rightarrow \infty$$

for any $x_1, x_2 \in \mathbb{R}$ except at most a countable set D . Let us fix $\varepsilon > 0$ and choose $r_\varepsilon > 0$ such that $\pm r_\varepsilon \in \mathbb{R} \setminus D$ and $1 - F(r_\varepsilon) + F(-r_\varepsilon) < \varepsilon$. We define

$$T_k(r) := 1 - F_{n_k}(r) + F_{n_k}(-r), \quad k \in \mathbb{N}, r > 0.$$

Due to (5), there exists $k_\varepsilon \in \mathbb{N}$ such that $T_k(r_\varepsilon) < \varepsilon$ for all $k \geq k_\varepsilon$. Taking r_ε greater to provide $T_k(r_\varepsilon) < \varepsilon$ for all $k < k_\varepsilon$, we obtain $\sup_{k \in \mathbb{N}} T_k(r_\varepsilon) < \varepsilon$ because, due to monotonicity of every F_{n_k} , the inequality $T_k(r_\varepsilon) < \varepsilon$ still holds for all $k \geq k_\varepsilon$. Thus $\sup_{k \in \mathbb{N}} T_k(r) \rightarrow 0, r \rightarrow \infty$, and in particular, $\sup_{k \in \mathbb{N}} F_{n_k}(-r) \rightarrow 0, r \rightarrow \infty$. Due to the latter, it is easy to check that (5) yields the convergence $F_{n_k}(x) \rightarrow F(x)$ for any $x \in \mathbb{R}$ except at most countable set D . Since $\mathbb{R} \setminus D$ is a dense subset of \mathbb{R} , we have $F_{n_k} \xrightarrow{w} F, k \rightarrow \infty$. Thus, according to definition of basic convergence, every subsequence of $(F_n)_{n \in \mathbb{N}}$ contains a further subsequence $(F_{n_k})_{k \in \mathbb{N}}$ that satisfies (5) and hence weakly converges to F . By the well-known fact in [8, p. 337], it means that the whole sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F .

The proved assertion can be generalized for bounded nondecreasing functions $F \in \mathbf{V}$ and $F_n \in \mathbf{V}, n \in \mathbb{N}$, but here the basic convergence $F_n \Rightarrow F, n \rightarrow \infty$, must be taken together with an additional condition that $F_n(+\infty) \rightarrow F(+\infty), n \rightarrow \infty$ (see [13, p. 39]). It should be noted that basic and weak convergences are not equivalent in a general case for functions from \mathbf{V} . Indeed, the weak convergence implies the basic one that will follow from Theorem 5 below, and also it is seen from Theorem 1.4.7 in [10]. However, the converse is not true. The latter assertion is concluded from the following simple examples. Below $\mathbb{1}_a$ with fixed $a \in \mathbb{R}$ denotes the following functions: $\mathbb{1}_a(x) = 1$ for $x \geq a$ and $\mathbb{1}_a(x) = 0$ for $x < a$.

Example 1. Let us define $G_n(x) := \mathbb{1}_n(x) - \mathbb{1}_{n+1}(x), x \in \mathbb{R}, n \in \mathbb{N}$. It is easily seen that $G_n \in \mathbf{V}, n \in \mathbb{N}$, and $G_n(x) \rightarrow 0, n \rightarrow \infty$ for all $x \in \mathbb{R}$. Setting $G(x) := 0, x \in \mathbb{R}$, we have the basic convergence $G_n \Rightarrow G, n \rightarrow \infty$. Here $G_n(+\infty) = G(+\infty) = 0$ and $\|G_n\| = 2, n \in \mathbb{N}$. However, for the continuous and bounded function $x \mapsto \cos(\pi x), x \in \mathbb{R}$, we conclude that

$$\int_{\mathbb{R}} \cos(\pi x) dG_n(x) \not\rightarrow \int_{\mathbb{R}} \cos(\pi x) dG(x) = 0, \quad n \rightarrow \infty.$$

Indeed,

$$\begin{aligned} \int_{\mathbb{R}} \cos(\pi x) dG_n(x) &= \cos(\pi n) - \cos(\pi(n+1)) \\ &= (-1)^n - (-1)^{n+1} = 2 \cdot (-1)^n \not\rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

Thus $(G_n)_{n \in \mathbb{N}}$ doesn't weakly converge to G . \square

Example 2. Let $G_n(x) := n\mathbb{1}_0(x) - n\mathbb{1}_{1/n^2}(x)$, $x \in \mathbb{R}$, $n \in \mathbb{N}$. So $G_n \in \mathbf{V}$, $n \in \mathbb{N}$, and $G_n(x) \rightarrow 0$, $n \rightarrow \infty$ for all $x \neq 0$. We set $G(x) := 0$, $x \in \mathbb{R}$, and we obtain that $G_n \Rightarrow G$, $n \rightarrow \infty$. Observe that $\|G_n\| = 2n \rightarrow \infty$, $n \rightarrow \infty$. Hence $(G_n)_{n \in \mathbb{N}}$ cannot be weak convergent sequence, because, under the weak convergence, total variations must be uniformly bounded (see Proposition 1.4.4. in [10, p. 22]). Moreover, it even fails to hold that

$$(6) \quad \int_{\mathbb{R}} h(x) dG_n(x) \rightarrow \int_{\mathbb{R}} h(x) dG(x), \quad n \rightarrow \infty$$

for any continuous function h with compact support. Indeed, let $h(x) := \sqrt{x}$ for $x \in [0, 1]$, $h(x) := \sqrt{2-x}$ for $x \in [1, 2]$, and $h(x) := 0$ for $x \notin [0, 2]$. Obviously, the function h satisfies the required properties. So we have

$$\int_{\mathbb{R}} h(x) dG_n(x) = h(0) \cdot n - h(1/n^2) \cdot n = 0 - \sqrt{1/n^2} \cdot n = -1 \quad \text{for every } n \in \mathbb{N},$$

but $\int_{\mathbb{R}} h(x) dG(x) = 0$. Thus (6) does not hold. However, it is interesting to note that there is a convergence of Fourier–Stieltjes transforms. Indeed, for any $t \in \mathbb{R}$

$$\int_{\mathbb{R}} e^{itx} dG_n(x) = (1 - e^{it/n^2}) \cdot n = -\frac{it}{n} (1 + o(1)) \rightarrow 0 = \int_{\mathbb{R}} e^{itx} dG(x), \quad n \rightarrow \infty. \quad \square$$

The next example shows that the use of the subsequences in the definition of basic convergence is essential.

Example 3. For any $n \in \mathbb{N}$ we set $k_n \in \mathbb{N} \cup \{0\}$ satisfying $2^{k_n} \leq n < 2^{k_n+1}$. We define

$$G_n(x) := \mathbb{1}_{a_n}(x) - \mathbb{1}_{b_n}(x), \quad x \in \mathbb{R},$$

where

$$a_n := \frac{n - 2^{k_n}}{2^{k_n}} \quad \text{and} \quad b_n := \frac{n + 1 - 2^{k_n}}{2^{k_n}}, \quad n \in \mathbb{N}.$$

It is seen that the interval $[a_n, b_n]$ is vanishing ($b_n - a_n = 2^{-k_n} \rightarrow 0$) and shifting over $[0, 1]$ as $n \rightarrow \infty$. Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a bounded continuous function. Due to the uniform continuity of h on $[0, 1]$, we have that

$$\int_{\mathbb{R}} h(x) dG_n(x) = h(a_n) - h(b_n) \rightarrow 0, \quad n \rightarrow \infty.$$

So $(G_n)_{n \in \mathbb{N}}$ weakly converges to $G(x) := 0$ for all $x \in \mathbb{R}$. Then, by the comments above, $(G_n)_{n \in \mathbb{N}}$ basically converges to G that can be also checked directly by definition. However, for any $x_0, x_1, x_2 \in [0, 1)$ there is no limit either for $G_n(x_0)$ or $G_n(x_2) - G_n(x_1)$ as $n \rightarrow \infty$, because $G_n(x)$ takes an infinite number of times each of the values 1 or 0, when $x \in [a_n, b_n]$ or not correspondingly. Note that, due to the weak convergence of $(G_n)_{n \in \mathbb{N}}$, there is a convergence of Fourier–Stieltjes transforms:

$$\int_{\mathbb{R}} e^{itx} dG_n(x) \rightarrow \int_{\mathbb{R}} e^{itx} dG(x) = 0 \quad \text{as } n \rightarrow \infty \quad \text{for every } t \in \mathbb{R}. \quad \square$$

The following example shows that it is important to use the differences of values of the functions at points x_1 and x_2 in the definition of the basic convergence in order to stay within \mathbf{V} .

Example 4. For every $n \in \mathbb{N}$ we define $G_n(x) := 1 + \frac{x}{n}$ for $x \in [-n, n]$, $G_n(x) = 0$ for $x < -n$, and $G_n(x) = 2$ for $x > n$. So G_n are nondecreasing continuous functions and $G_n(+\infty) = \|G_n\| = 2$, $n \in \mathbb{N}$. We see that $G_n(x) \rightarrow 1$ as $n \rightarrow \infty$ for any $x \in \mathbb{R}$. However, an identical 1 doesn't belong to \mathbf{V} (it must be 0 at $-\infty$). At the same time for any real x_1 and x_2 we have $G_n(x_2) - G_n(x_1) \rightarrow 0$, $n \rightarrow \infty$, and we conclude that $(G_n)_{n \in \mathbb{N}}$ basically converges to the function $G(x) := 0$ for all $x \in \mathbb{R}$, which is from \mathbf{V} . Of course, $(G_n)_{n \in \mathbb{N}}$ doesn't weakly converge to G here, because

$$\int_{\mathbb{R}} dG_n(x) = 2 \not\rightarrow \int_{\mathbb{R}} dG(x) = 0, \quad n \rightarrow \infty.$$

Note that for $t \neq 0$ we have

$$\int_{\mathbb{R}} e^{itx} dG_n(x) = \frac{1}{n} \int_{-n}^n e^{itx} dx = \frac{e^{itn} - e^{-itn}}{itn} \rightarrow 0, \quad n \rightarrow \infty$$

and

$$\int_{\mathbb{R}} e^{itx} dG_n(x)|_{t=0} = \int_{\mathbb{R}} dG_n(x) = G_n(+\infty) = 2, \quad n \in \mathbb{N}.$$

Thus the Fourier–Stieltjes transforms of G_n , $n \in \mathbb{N}$, pointwisely converge to the Fourier–Stieltjes transform of G (i.e., to identical 0) for almost all $t \in \mathbb{R}$. \square

We now consider a general question about the relationship between the basic convergence of functions from \mathbf{V} and the convergence of their Fourier–Stieltjes transforms. We do not pretend to study this question in full here; instead, we present only those assertions that will be used in the main results of the article.

Let $(G_n)_{n \in \mathbb{N}}$ be a sequence of functions from \mathbf{V} . Let us define the corresponding sequence of Fourier–Stieltjes integrals:

$$g_n(t) = \int_{\mathbb{R}} e^{itx} dG_n(x), \quad t \in \mathbb{R}, n \in \mathbb{N}.$$

The results below in fact show that, under the rather weak and natural assumptions, the pointwise convergence of g_n implies the basic convergence of G_n as $n \rightarrow \infty$.

We will use the following assumption:

$$(7) \quad \overline{\lim}_{n \rightarrow \infty} \|G_n\| = B < \infty.$$

Theorem 5. *Let $(G_n)_{n \in \mathbb{N}}$ satisfy (7). Suppose that $g_n(t) \rightarrow g(t)$, $n \rightarrow \infty$, for almost all $t \in \mathbb{R}$ with some function $g : \mathbb{R} \rightarrow \mathbb{C}$. Then there exists a function $G \in \mathbf{V}$ such that $\|G\| \leq B$ and the equality*

$$(8) \quad g(t) = \int_{\mathbb{R}} e^{itx} dG(x)$$

holds for almost all $t \in \mathbb{R}$ including all continuity points of the function g . The function G is uniquely determined in the class \mathbf{V} , and $G_n \Rightarrow G$, $n \rightarrow \infty$. If also $g_n(0) \rightarrow g(0)$, $n \rightarrow \infty$, and g is continuous at $t = 0$, then additionally $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.

We are not aware of any results with such assertion. There are some close remarks in [9] and [29]. It is seen that this theorem complements and partially generalizes the well-known Levy’s continuity theorem, which was stated for sequences of probability distribution functions.

Suppose that the sequence $(G_n)_{n \in \mathbb{N}}$ weakly converges to a function G from \mathbf{V} with Fourier–Stieltjes transform g . Then (7) is satisfied (see Proposition 1.4.4. in [10, p. 22]) and $g_n(t) \rightarrow g(t)$, $n \rightarrow \infty$ for every $t \in \mathbb{R}$. According to the theorem, we have the basic convergence $G_n \Rightarrow G$ and also $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$. Thus we showed that the weak convergence implies the basic convergence.

We next formulate the analog of Theorem 5 using the decompositions

$$(9) \quad G_n(x) = G_n^+(x) - G_n^-(x), \quad x \in \mathbb{R}, n \in \mathbb{N},$$

where G_n^+ and G_n^- are nondecreasing functions from \mathbf{V} . Note that (9) are not necessarily corresponding to the Hahn–Jordan decomposition (see (13) below). Here we assume

$$(10) \quad \overline{\lim}_{n \rightarrow \infty} G_n^-(+\infty) = M < \infty,$$

which can sometimes be more convenient for checking than (7).

Proposition 6. *Let $(G_n)_{n \in \mathbb{N}}$ satisfy (10) for some decompositions (9). Suppose that $g_n(0) \rightarrow c \in \mathbb{R}$, $n \rightarrow \infty$. Then (7) holds with some $B \leq c + 2M$.*

Thus if there is a convergence of g_n , $n \in \mathbb{N}$, at $t = 0$, then for some (9) assumptions (7) and (10) are equivalent. So we come to the following assertion.

Theorem 7. *Let $(G_n)_{n \in \mathbb{N}}$ satisfy (10) for some decompositions (9). Suppose that $g_n(t) \rightarrow g(t)$, $n \rightarrow \infty$, for almost all $t \in \mathbb{R}$ including $t = 0$ with some function $g : \mathbb{R} \rightarrow \mathbb{C}$. Then $g(0) \in \mathbb{R}$, the condition (7) is satisfied with $B \leq g(0) + 2M$, and the assertions of Theorem 5 hold. If also g is continuous at $t = 0$, then additionally $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.*

3. Main results

Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of quasi-infinitely divisible distribution function with corresponding sequence of characteristic function $(f_n)_{n \in \mathbb{N}}$. Let every f_n admit the representation

$$(11) \quad f_n(t) = \exp \left\{ it\gamma_n + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \right\}, \quad t \in \mathbb{R}, n \in \mathbb{N},$$

where $\gamma_n \in \mathbb{R}$, $G_n \in \mathbf{V}$, $n \in \mathbb{N}$, and $\tau > 0$ is a fixed number. We are interested in criteria of the weak convergence of $(F_n)_{n \in \mathbb{N}}$ in terms of the spectral pairs (γ_n, G_n) , $n \in \mathbb{N}$.

Assertions of the following Theorems 8 and 9 were obtained by Lindner, Pan, and Sato in [23] (where the results were presented in another form).

Theorem 8. *If $\gamma_n \rightarrow \gamma$ and $G_n \xrightarrow{w} G$, $n \rightarrow \infty$, with some $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$, then (γ, G) is the spectral pair for some quasi-infininitely divisible distribution function F , and $(F_n)_{n \in \mathbb{N}}$ weakly converges to F .*

We next use the decompositions

$$(12) \quad G_n(x) = G_n^+(x) - G_n^-(x), \quad x \in \mathbb{R}, n \in \mathbb{N},$$

where G_n^+ and G_n^- are nondecreasing functions from \mathbf{V} . There exists an important way of choosing G_n^+ and G_n^- . Let μ_{G_n} be the signed measure that is generated by G_n for every $n \in \mathbb{N}$, i.e., such that $\mu_{G_n}((a, b]) = G_n(b) - G_n(a)$ for all $a, b \in \mathbb{R}$, $a \leq b$, $n \in \mathbb{N}$. Every measure μ_{G_n} is uniquely represented by the Hahn–Jordan decomposition $\mu_{G_n} = \mu_{G_n}^+ - \mu_{G_n}^-$, where $\mu_{G_n}^+$ and $\mu_{G_n}^-$ are nonnegative finite measures concentrated on some disjoint sets (see [10, p. 3]). So we can choose

$$(13) \quad G_n^+(x) = \mu_{G_n}^+((-\infty, x]) \quad \text{and} \quad G_n^-(x) = \mu_{G_n}^-((-\infty, x]), \quad x \in \mathbb{R}, n \in \mathbb{N}.$$

In this case we will have (12) and additionally that $|G_n|(x) = G_n^+(x) + G_n^-(x)$, $x \in \mathbb{R}$, $n \in \mathbb{N}$.

Theorem 9. *Let F be a distribution function and $(F_n)_{n \in \mathbb{N}}$ weakly converge to F . Suppose that G_n^+ and G_n^- from (12) are defined according to the Hahn–Jordan decomposition by (13) for every $n \in \mathbb{N}$. Suppose that the sequence $(G_n^-)_{n \in \mathbb{N}}$ satisfies the assumptions*

$$\sup_{n \in \mathbb{N}} \|G_n^-\| < \infty \quad \text{and} \quad \lim_{r \rightarrow \infty} \sup_{n \in \mathbb{N}} (1 - |G_n^-|(r) + |G_n^-|(-r)) = 0$$

(uniform boundedness in variation and tightness, correspondingly). Then F is quasi-infininitely divisible with some spectral pair (γ, G) . Moreover, $\gamma_n \rightarrow \gamma$ and $G_n \xrightarrow{w} G$, $n \rightarrow \infty$.

Theorems 8 and 9 connect the weak convergence of quasi-infininitely divisible distribution functions with the weak convergence of their spectral functions. We are interested in analogs of these theorems but with the basic convergence of the spectral functions.

We will use the following assumption:

$$(14) \quad \overline{\lim}_{n \rightarrow \infty} \|G_n\| = B < \infty.$$

Theorem 10. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (14) with some $B \geq 0$. Let $(F_n)_{n \in \mathbb{N}}$ weakly converge to a distribution function F . Then F is quasi-infinitely divisible with some spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$ with $\|G\| \leq B$. Moreover, $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$ and $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.*

The next theorem is an analog of this one, but with the assumption

$$(15) \quad \overline{\lim}_{n \rightarrow \infty} G_n^-(+\infty) = M < \infty$$

on decompositions (12) for G_n , $n \in \mathbb{N}$. If we choose G_n^+ and G_n^- according to the Hahn–Jordan decomposition by (13) for every $n \in \mathbb{N}$, then (15) is weaker than (14). Also observe that (15) is satisfied, when we deal with nondecreasing functions G_n , $n \in \mathbb{N}$. It should be noted, however, that it is not required in the theorems and corollaries below that G_n^+ and G_n^- in (12) must be chosen according to the Hahn–Jordan decomposition.

Theorem 11. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (15) with some $M \geq 0$ and for some decompositions (12). Let $(F_n)_{n \in \mathbb{N}}$ weakly converge to a distribution function F . Then (14) holds for some $B \geq 0$ and all assertions of Theorem 10 are true. Also we have that $B \leq G(+\infty) + 2M$.*

Theorems 10 and 11 yield necessary conditions for the weak convergence within the class of quasi-infinitely divisible distribution functions under the assumption (14) or (15).

Corollary 12. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (14) or (15) for some decompositions (12). Let F be a quasi-infinitely divisible distribution function F with spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. If the sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F , then $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$ and $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.*

Also Theorems 10 and 11 state sufficient conditions for membership of the class of quasi-infinitely divisible distribution functions.

Corollary 13. *A distribution function F is quasi-infinitely divisible if it is a weak limit of a sequence $(F_n)_{n \in \mathbb{N}}$ of quasi-infinitely divisible distribution functions (with characteristic functions (11)), which satisfies (14) or (15) for some decompositions (12).*

Note that this corollary is a stronger version of the same assertion in Theorem 9, because we don't assume the tightness for $(G_n^-)_{n \in \mathbb{N}}$ and we don't require the use of the Hahn–Jordan decomposition.

It is known (see [23, p. 17]) that a weak limit of quasi-infinitely divisible distribution functions is not necessarily quasi-infinitely divisible. Hence assumptions (14) or (15) cannot be simply omitted in Corollary 13. However, it seems that they can be done weaker (see [23, Example 4.4]).

We will use a notion of relative compactness for $(F_n)_{n \in \mathbb{N}}$ in the next theorem. Recall that $(F_n)_{n \in \mathbb{N}}$ is said to be *relatively compact* if every its subsequence contains a further subsequence that weakly converges to a distribution function. It is clear that a weakly convergent sequence of distribution functions is relatively compact. In general, the property of relative compactness is not difficult for checking due to Prokhorov's theorem and various probability inequalities. Also some criteria of relative compactness are known for particular important sequences of distribution functions (for example, see [15; 16; 17]).

Theorem 14. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (14). If $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$, $n \rightarrow \infty$, with some $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$, then (γ, G) is the spectral pair for a quasi-infinitely divisible distribution function F and the sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F .*

This theorem yields sufficient conditions for the weak convergence within the class of quasi-infinitely divisible distribution functions under the assumption (14).

Corollary 15. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (14). Let F be a quasi-infinitely divisible distribution function F with spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. If $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$, $n \rightarrow \infty$, then $(F_n)_{n \in \mathbb{N}}$ weakly converges to F .*

Corollaries 12 and 15 directly yield the following criterion.

Theorem 16. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (14). Let F be a quasi-infinitely divisible distribution function F with spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. The sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F if and only if $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$, $n \rightarrow \infty$. Moreover, the convergence $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$, can be added to the necessary conditions.*

We now formulate the analogs of Theorems 14 and 16, and of Corollary 15 under the assumption (15). They are directly stated due to the following simple note.

Suppose that a sequence $(G_n)_{n \in \mathbb{N}}$ from \mathbf{V} satisfies (15) for some decompositions (12). If $\overline{\lim}_{n \rightarrow \infty} G_n(+\infty)$ is finite, then (14) holds. Indeed, according to (12), it follows from the inequalities

$$\|G_n\| \leq \|G_n^+\| + \|G_n^-\| = G_n^+(+\infty) + G_n^-(+\infty) = G_n(+\infty) + 2G_n^-(+\infty), \quad n \in \mathbb{N}.$$

So we obtain the following results.

Theorem 17. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (15) for some decompositions (12). If $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$, and $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$, with some $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$, then all assertions of Theorem 14 hold.*

So Theorems 14 and 17 complement Theorem 8: we use weaker convergence for the spectral functions $(G_n)_{n \in \mathbb{N}}$, but we additionally assume the relative compactness of $(F_n)_{n \in \mathbb{N}}$.

Corollary 18. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (15) for some decompositions (12). Let F be a quasi-infinitely divisible distribution function F with the spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. If $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$ and $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$, then $(F_n)_{n \in \mathbb{N}}$ weakly converges to F .*

Corollaries 12 and 18 directly yield the following criterion.

Theorem 19. *Suppose that $(F_n)_{n \in \mathbb{N}}$ satisfies (15) for some decompositions (12). Let F be a quasi-infinitely divisible distribution function F with spectral pair (γ, G) , where $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. The sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F if and only if $(F_n)_{n \in \mathbb{N}}$ is relatively compact and $\gamma_n \rightarrow \gamma$, $G_n \Rightarrow G$, and $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.*

On account of comments before Example 1 in Section 2, Theorems 16 and 19 complement similar well-known results for the weak convergence of infinitely divisible distribution functions (see [13, p. 87]).

4. Proofs

Proof of Theorem 5. First, observe that the function g is measurable, because it is an almost everywhere limit of continuous (hence measurable) functions g_n , $n \in \mathbb{N}$. So we have

$$(16) \quad \int_{\mathbb{R}} g_n(t) \rho(t) dt \rightarrow \int_{\mathbb{R}} g(t) \rho(t) dt, \quad n \rightarrow \infty$$

for any function $\rho \in L_1(\mathbb{R})$. Indeed, due to (7), there exists a constant $B_0 > 0$ such that $|g_n(t)| \leq \|G_n\| \leq B_0$ for all $n \in \mathbb{N}$, and convergence (16) holds by the Lebesgue dominated convergence theorem.

Let us define the function

$$\varphi(x) := \int_{\mathbb{R}} e^{itx} \rho(t) dt, \quad x \in \mathbb{R}.$$

Observe that for every $n \in \mathbb{N}$ we have

$$(17) \quad \begin{aligned} \int_{\mathbb{R}} g_n(t) \rho(t) dt &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{itx} dG_n(x) \right) \rho(t) dt \\ &= \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{itx} \rho(t) dt \right) dG_n(x) = \int_{\mathbb{R}} \varphi(x) dG_n(x). \end{aligned}$$

Let us consider the last integral. Due to (4) and (7), by Helly's first theorem (see [27, pp. 222 and 240]), there exists a subsequence $(G_{n_k})_{k \in \mathbb{N}}$ in $(G_n)_{n \in \mathbb{N}}$ and a function of bounded variation $G_* : \mathbb{R} \rightarrow \mathbb{R}$ such that $G_{n_k}(x) \rightarrow G_*(x)$ as $k \rightarrow \infty$ for all $x \in \mathbb{R}$. Note that, in general, G_* may not be right-continuous (see Example 3). But φ is bounded and continuous on \mathbb{R} and hence there exists the Riemann–Stieltjes integral $\int_{\mathbb{R}} \varphi(x) dG_*(x)$. Also the (Lebesgue–Stieltjes) integrals $\int_{\mathbb{R}} \varphi(x) dG_n(x)$ coincide with the corresponding Riemann–Stieltjes integrals. Next, due to the well-known

fact that $\varphi(x) \rightarrow 0$ as $x \rightarrow \pm\infty$, by Helly's second theorem (see [27, p. 240]), we have the following convergence for the Riemann–Stieltjes integrals:

$$\int_{\mathbb{R}} \varphi(x) dG_{n_k}(x) \rightarrow \int_{\mathbb{R}} \varphi(x) dG_*(x), \quad k \rightarrow \infty.$$

Let us define $G(x) := G_*(x+) - G_*(-\infty)$, $x \in \mathbb{R}$ (note that $G_*(-\infty) \neq 0$ in general, see Example 4). So G is right-continuous on \mathbb{R} and $G(-\infty) = 0$, i.e., $G \in V$. Since $G(x)$ equals $G_*(x) - G_*(-\infty)$ for all $x \in \mathbb{R}$ except at most countable set, due to the continuity of φ , we have

$$\int_{\mathbb{R}} \varphi(x) dG_*(x) = \int_{\mathbb{R}} \varphi(x) dG(x),$$

where the integral on the right-hand side can be considered as Lebesgue–Stieltjes integral. Thus we have the following convergence with the Lebesgue–Stieltjes integrals:

$$\int_{\mathbb{R}} \varphi(x) dG_{n_k}(x) \rightarrow \int_{\mathbb{R}} \varphi(x) dG(x), \quad k \rightarrow \infty.$$

The integral on the right-hand side admits the following representation analogously to (17):

$$\int_{\mathbb{R}} \varphi(x) dG(x) = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{itx} dG(x) \right) \rho(t) dt.$$

Due to (16) and (17), we also have

$$\int_{\mathbb{R}} \varphi(x) dG_{n_k}(x) \rightarrow \int_{\mathbb{R}} g(t) \rho(t) dt, \quad k \rightarrow \infty.$$

Thus we obtain

$$(18) \quad \int_{\mathbb{R}} g(t) \rho(t) dt = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{itx} dG(x) \right) \rho(t) dt$$

for any function $\rho \in L_1(\mathbb{R})$. This implies that

$$(19) \quad g(t) = \int_{\mathbb{R}} e^{itx} dG(x) \quad \text{for almost every } t \in \mathbb{R}.$$

Indeed, conversely, suppose that there exists a bounded set E of nonzero Lebesgue measure such that $\Delta(t) := g(t) - \int_{\mathbb{R}} e^{itx} dG(x) \neq 0$, $t \in E$. Let us introduce the sets

$$E_1 := \{t \in E : \operatorname{Re} \Delta(t) > 0\}, \quad E_2 := \{t \in E : \operatorname{Re} \Delta(t) < 0\}, \\ E_3 := \{t \in E : \operatorname{Im} \Delta(t) > 0\}, \quad E_4 := \{t \in E : \operatorname{Im} \Delta(t) < 0\}.$$

It is easily seen that $E = E_1 \cup E_2 \cup E_3 \cup E_4$. Hence at least one E_j has nonzero Lebesgue measure. We denote any such set by E_* . Next, according to the property of strict positivity of integral, we obtain

$$\left| \int_{E_*} \Delta(t) dt \right| \geq \left| \int_{E_*} \operatorname{Re} \Delta(t) dt \right| = \int_{E_*} |\operatorname{Re} \Delta(t)| dt > 0 \quad \text{for } E_* = E_1 \text{ or } E_* = E_2,$$

and

$$\left| \int_{E_*} \Delta(t) dt \right| \geq \left| \int_{E_*} \operatorname{Im} \Delta(t) dt \right| = \int_{E_*} |\operatorname{Im} \Delta(t)| dt > 0 \quad \text{for } E_* = E_3 \text{ or } E_* = E_4.$$

Thus we have

$$\left| \int_{E_*} g(t) dt - \int_{E_*} \left(\int_{\mathbb{R}} e^{itx} dG(x) \right) dt \right| = \left| \int_{E_*} \Delta(t) dt \right| > 0.$$

This contradicts (18) when we choose ρ as follows: $\rho(t) = 1$, $t \in E_*$ and $\rho(t) = 0$, $t \notin E_*$. It is valid since $\rho \in L_1(\mathbb{R})$ due to the boundedness $E_* \subset E$. Thus (19) is true.

Let us show that (19) holds for every continuity point of the function g . Let \mathcal{T} be the set of all $t \in \mathbb{R}$ for which (19) holds. Hence the Lebesgue measure of $\mathbb{R} \setminus \mathcal{T}$ equals zero. Let g be continuous at the fixed point t_0 . So we can choose $t_m \in \mathcal{T}$, $m \in \mathbb{N}$ such that $t_m \rightarrow t_0$, $m \rightarrow \infty$. Then $g(t_m) \rightarrow g(t_0)$, $m \rightarrow \infty$, and at the same time

$$g(t_m) = \int_{\mathbb{R}} e^{it_m x} dG(x) \rightarrow \int_{\mathbb{R}} e^{it_0 x} dG(x), \quad m \rightarrow \infty$$

due to continuity of the function $t \mapsto \int_{\mathbb{R}} e^{itx} dG(x)$ on \mathbb{R} . Thus we have

$$g(t_0) = \int_{\mathbb{R}} e^{it_0 x} dG(x).$$

According to (19), the function g almost everywhere coincides with the continuous function $t \mapsto \int_{\mathbb{R}} e^{itx} dG(x)$, $t \in \mathbb{R}$. So the latter function is uniquely determined by g within the class of all continuous complex-valued functions on \mathbb{R} . Next, it is well known that $t \mapsto \int_{\mathbb{R}} e^{itx} dG(x)$, $t \in \mathbb{R}$, uniquely determines G within the class \mathbf{V} . Therefore g uniquely determines G in the class \mathbf{V} .

Let's return to the sequence $(G_{n_k})_{k \in \mathbb{N}}$. From the above we know that $G_{n_k}(x) \rightarrow G_*(x)$ for all $x \in \mathbb{R}$ and $G(x) = G_*(x) - G_*(-\infty)$ for all $x \in \mathbb{R}$ except at most a countable set D where G_* is not right-continuous. Then for all $x_1, x_2 \in \mathbb{R} \setminus D$ we have

$$(20) \quad G_{n_k}(x_2) - G_{n_k}(x_1) \rightarrow (G(x_2) + G_*(-\infty)) - (G(x_1) + G_*(-\infty)) = G(x_2) - G(x_1), \quad k \rightarrow \infty.$$

Let $(G_{m_l})_{l \in \mathbb{N}}$ be an arbitrary subsequence of $(G_n)_{n \in \mathbb{N}}$. Analogously to the above, there exists a further subsequence $(G_{m'_k})_{k \in \mathbb{N}}$ in $(G_{m_k})_{k \in \mathbb{N}}$, which pointwise converges to some function of bounded variation $H_*: \mathbb{R} \rightarrow \mathbb{R}$, i.e., $G_{m'_k}(x) \rightarrow H_*(x)$, $k \rightarrow \infty$ for all $x \in \mathbb{R}$. Defining $H(x) := H_*(x+) - H_*(-\infty)$, $x \in \mathbb{R}$, we as before will obtain $g(t) = \int_{\mathbb{R}} e^{itx} dH(x)$ for almost all $t \in \mathbb{R}$, with $H \in \mathbf{V}$. Since G is a unique function within \mathbf{V} , which represents g by (8), we have $H(x) = G(x)$, $x \in \mathbb{R}$. We also have

$$G_{m'_k}(x_2) - G_{m'_k}(x_1) \rightarrow G(x_2) - G(x_1), \quad k \rightarrow \infty$$

for all $x_1, x_2 \in \mathbb{R}$ except at most countable set D' where H_* is not right-continuous (in general $D' \neq D$). So we proved that $G_n \Rightarrow G$, $n \rightarrow \infty$.

Let us consider the numbers $g_n(0) = \int_{\mathbb{R}} dG_n(x) = G_n(+\infty)$, $n \in \mathbb{N}$. If we suppose that g is continuous at $t = 0$, then, by the above remarks, we will have $g(0) = \int_{\mathbb{R}} dG(x) = G(+\infty)$. Therefore, assuming to hold $g_n(0) \rightarrow g(0)$, $n \rightarrow \infty$, we will obtain $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.

It remains to prove that $\|G\| \leq B$. On the contrary, suppose that this is false. Then we can find $y_0, y_1, \dots, y_N \in \mathbb{R}$ such that

$$(21) \quad B < \sum_{j=1}^N |G(y_j) - G(y_{j-1})| \leq \|G\|.$$

Let us take our sequence $(G_{n_k})_{k \in \mathbb{N}}$ and the set D , which is at most countable. Since G is right-continuous and the set $\mathbb{R} \setminus D$ is dense, we can assume that y_0, y_1, \dots, y_N are chosen from $\mathbb{R} \setminus D$. Next, due to the convergence (20) and assumption (7), we have

$$\sum_{j=1}^N |G(y_j) - G(y_{j-1})| = \lim_{k \rightarrow \infty} \sum_{j=1}^N |G_{n_k}(y_j) - G_{n_k}(y_{j-1})| \leq \overline{\lim}_{n \rightarrow \infty} \|G_n\| \leq B,$$

which contradicts (21). □

Proof of Proposition 6. By the assumption $g_n(0) \rightarrow c \in \mathbb{R}$, $n \rightarrow \infty$. Since $g_n(0) = \int_{\mathbb{R}} dG_n(x) = G_n(+\infty)$, $n \in \mathbb{N}$, we have the convergence $G_n(+\infty) \rightarrow c$, $n \rightarrow \infty$. Let us consider decompositions (9). We have $G_n(+\infty) = G_n^+(+\infty) - G_n^- (+\infty)$, $n \in \mathbb{N}$. Also observe that

$$\|G_n\| \leq \|G_n^+\| + \|G_n^-\| = G_n^+(+\infty) + G_n^- (+\infty) = G_n(+\infty) + 2G_n^- (+\infty), \quad n \in \mathbb{N}.$$

Therefore

$$B = \overline{\lim}_{n \rightarrow \infty} \|G_n\| \leq \lim_{n \rightarrow \infty} G_n(+\infty) + 2 \overline{\lim}_{n \rightarrow \infty} G_n^- (+\infty) = c + 2M.$$

Thus we have (7) with $B \leq g(0) + 2M$. □

Proof of Theorem 7. By the assumption $g_n(0) \rightarrow g(0)$, $n \rightarrow \infty$. Since $g_n(0) = \int_{\mathbb{R}} dG_n(x) = G_n(+\infty)$, $n \in \mathbb{N}$, we have the convergence $G_n(+\infty) \rightarrow g(0)$, $n \rightarrow \infty$. So the sequence $G_n(+\infty) \in \mathbb{R}$, $n \in \mathbb{N}$, has a finite limit $g(0)$ that must be real. According to Proposition 6, condition (7) holds with some $B \leq g(0) + 2M$. Using Theorem 5, we get all its assertions. So $g(t) = \int_{\mathbb{R}} e^{itx} dG(x)$ holds for some $G \in \mathcal{V}$ and for all $t \in \mathbb{R}$ that are continuity points of the function g . Under the assumption, g is continuous at $t = 0$, and we have $g(0) = \int_{\mathbb{R}} dG(x) = G(+\infty)$. Since $G_n(+\infty) \rightarrow g(0)$, $n \rightarrow \infty$, we obtain that

$$G_n(+\infty) \rightarrow G(+\infty), \quad n \rightarrow \infty. \quad \square$$

We need the following lemma for proving Theorem 10.

Lemma 20. *For any $t \in \mathbb{R}$ and $\tau > 0$ the following representations hold:*

$$(22) \quad e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) = \int_{A_{t,\tau}} e^{isx} dU_{t,\tau}(s),$$

$$\left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1}{x^2} = \int_{A_{t,\tau}} e^{isx} dV_{t,\tau}(s),$$

$$(23) \quad \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} = \int_{A_{t,\tau}} e^{isx} dW_{t,\tau}(s), \quad x \in \mathbb{R},$$

where $A_{t,\tau} := \{s \in \mathbb{R} : |s| \leq \max\{|t|, \tau\}\}$, and

$$(24) \quad U_{t,\tau}(s) := \mathbb{1}_t(s) - \mathbb{1}_0(s) - \frac{t}{2\tau}(\mathbb{1}_\tau(s) - \mathbb{1}_{-\tau}(s)), \quad s \in \mathbb{R},$$

$$(25) \quad V_{t,\tau}(s) := \int_{-\infty}^s \rho_{t,\tau}(y) dy,$$

$$(26) \quad \begin{aligned} \rho_{t,\tau}(s) &:= -\frac{1}{2}(|s-t| - |s| - \frac{t}{2\tau}(|s-\tau| - |s+\tau|)), \quad s \in \mathbb{R}, \\ W_{t,\tau}(s) &:= U_{t,\tau}(s) + V_{t,\tau}(s), \quad s \in \mathbb{R}. \end{aligned}$$

For any $t \in \mathbb{R}$ and $\tau > 0$ it is true that $U_{t,\tau}(s) = 0$ and $\rho_{t,\tau}(s) = 0$ for all $s \notin A_{t,\tau}$, $\rho_{t,\tau}$ is a continuous function on \mathbb{R} with a broken-line graph, and, in particular, $\rho_{t,\tau} \in L_1(\mathbb{R})$, the functions $U_{t,\tau}$, $V_{t,\tau}$, and $W_{t,\tau}$ belong to the class \mathbf{V} .

Proof of Lemma 20. Let us fix $t \in \mathbb{R}$, $\tau > 0$, and define $A_{t,\tau}$ as in the formulation. We write

$$(27) \quad e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) = e^{itx} - 1 - \frac{t}{2\tau}(e^{i\tau x} - e^{-i\tau x}) = \int_{\mathbb{R}} e^{isx} dU_{t,\tau}(s), \quad x \in \mathbb{R},$$

where $U_{t,\tau}$ is defined by (24). Using the definition of the function $\mathbb{1}_a(\cdot)$, $a \in \mathbb{R}$, it is easily seen that $U_{t,\tau}$ is an right-continuous function on \mathbb{R} , $U_{t,\tau}(s) = 0$ for all $s \notin A_{t,\tau}$, and in particular, $U_{t,\tau} \in \mathbf{V}$. Therefore the set \mathbb{R} can be changed by $A_{t,\tau}$ in the integral (27).

Let us consider the function

$$\varphi_{t,\tau}(x) := \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1}{x^2}, \quad x \in \mathbb{R}.$$

Observe that $\varphi_{t,\tau} \in L_1(\mathbb{R})$. So we define

$$(28) \quad \rho_{t,\tau}(s) := \frac{1}{2\pi} \int_{\mathbb{R}} e^{-isx} \varphi_{t,\tau}(x) dx, \quad s \in \mathbb{R}.$$

Let us find an explicit formula for $\rho_{t,\tau}(s)$ for every $s \in \mathbb{R}$. Observe that $x \mapsto \operatorname{Re} \varphi_{t,\tau}(x)$, $x \in \mathbb{R}$, is an even function and $x \mapsto \operatorname{Im} \varphi_{t,\tau}(x)$, $x \in \mathbb{R}$, is an odd

function. Therefore

$$\begin{aligned} \rho_{t,\tau}(s) &= \frac{1}{2\pi} \int_{\mathbb{R}} (\operatorname{Re} \varphi_{t,\tau}(x) \cos(sx) + \operatorname{Im} \varphi_{t,\tau}(x) \sin(sx)) dx \\ &= \frac{1}{\pi} \int_0^\infty (\operatorname{Re} \varphi_{t,\tau}(x) \cos(sx) + \operatorname{Im} \varphi_{t,\tau}(x) \sin(sx)) dx \\ &= \frac{1}{\pi} \int_0^\infty \left(\frac{\cos(tx) - 1}{x^2} \cos(sx) + \frac{\sin(tx) - \frac{t}{\tau} \sin(\tau x)}{x^2} \sin(sx) \right) dx, \quad s \in \mathbb{R}. \end{aligned}$$

Next, using the known trigonometric formulas, we write

$$\begin{aligned} \rho_{t,\tau}(s) &= \frac{1}{\pi} \int_0^\infty \left(\frac{\cos(tx) \cos(sx) + \sin(tx) \sin(sx) - \cos(sx)}{x^2} - \frac{\frac{t}{\tau} \sin(\tau x) \sin(sx)}{x^2} \right) dx \\ &= \frac{1}{\pi} \int_0^\infty \left(\frac{\cos((s-t)x) - \cos(sx)}{x^2} - \frac{t}{\tau} \frac{\cos((s-\tau)x) - \cos((s+\tau)x)}{2x^2} \right) dx \\ &= \frac{1}{\pi} \int_0^\infty \frac{\cos(|s-t|x) - \cos(|s|x)}{x^2} dx \\ &\quad - \frac{t}{2\tau} \cdot \frac{1}{\pi} \int_0^\infty \frac{\cos(|s-\tau|x) - \cos(|s+\tau|x)}{x^2} dx, \quad s \in \mathbb{R}. \end{aligned}$$

It is known (see [14, p. 447, formula 3.782(2)]) that

$$\int_0^\infty \frac{1 - \cos(ax)}{x^2} dx = \frac{a\pi}{2}, \quad a \geq 0.$$

Hence

$$\rho_{t,\tau}(s) = -\frac{1}{2} (|s-t| - |s| - \frac{t}{2\tau} (|s-\tau| - |s+\tau|)), \quad s \in \mathbb{R},$$

as in (25). We see that $\rho_{t,\tau}$ is a continuous function with a broken-line graph. Also observe that $\rho_{t,\tau}(s) = 0$ for all $s \notin A_{t,\tau}$. Indeed, if $s > \max\{|t|, \tau\}$, then

$$\rho_{t,\tau}(s) = -\frac{1}{2} (s-t - s - \frac{t}{2\tau} (s-\tau - (s+\tau))) = -\frac{1}{2} (-t - \frac{t}{2\tau} \cdot (-2\tau)) = -\frac{1}{2} (-t+t) = 0,$$

and if $s < -\max\{|t|, \tau\}$, then

$$\rho_{t,\tau}(s) = -\frac{1}{2} (-(s-t) + s - \frac{t}{2\tau} (-(s-\tau) + s+\tau)) = -\frac{1}{2} (t - \frac{t}{2\tau} \cdot 2\tau) = -\frac{1}{2} (t-t) = 0.$$

Thus $\rho_{t,\tau} \in L_1(\mathbb{R})$. By the way, observe that $V_{t,\tau}$, which is defined by (25), is a continuous function on \mathbb{R} and it vanishes at $-\infty$, i.e., $V_{t,\tau} \in \mathcal{V}$. Then, according to these remarks and (28), we have

$$\varphi_{t,\tau}(x) = \int_{\mathbb{R}} e^{isx} \rho_{t,\tau}(s) ds = \int_{A_{t,\tau}} e^{isx} \rho_{t,\tau}(s) ds = \int_{A_{t,\tau}} e^{isx} dV_{t,\tau}(s), \quad x \in \mathbb{R}.$$

Next, summing the proved equalities in (22), we get (23) with $W_{t,\tau}$ defined by (26). Since $U_{t,\tau}$ and $V_{t,\tau}$ belong to \mathcal{V} , we conclude that $W_{t,\tau} \in \mathcal{V}$. \square

Proof of Theorem 10. Let f be a characteristic function of the limit distribution function F . By the continuity theorem, we have

$$(29) \quad f_n(t) \rightarrow f(t), \quad n \rightarrow \infty \quad \text{for every } t \in \mathbb{R}.$$

Moreover, it is well known (see [25]) that

$$(30) \quad \sup_{t \in [-T, T]} |f_n(t) - f(t)| \rightarrow 0, \quad n \rightarrow \infty \quad \text{for any } T > 0.$$

First let us recall that characteristic functions of quasi-infinitely divisible distributions have no zeroes on the real line (see [23] or (32) below). So, in particular, $f_n(t) \neq 0, t \in \mathbb{R}, n \in \mathbb{N}$. We now show that $f(t) \neq 0$ for all $t \in \mathbb{R}$. For any fixed $n \in \mathbb{N}$ and $t \in \mathbb{R}$ we consider

$$\begin{aligned} |f_n(t)| &= \exp \left\{ \int_{\mathbb{R}} (\cos(tx) - 1) \frac{1+x^2}{x^2} dG_n(x) \right\} \\ &\geq \exp \left\{ - \left| \int_{\mathbb{R}} (\cos(tx) - 1) \frac{1+x^2}{x^2} dG_n(x) \right| \right\} \\ &\geq \exp \left\{ - \int_{\mathbb{R}} (1 - \cos(tx)) \frac{1+x^2}{x^2} d|G_n|(x) \right\}. \end{aligned}$$

Let us estimate the inner function $x \mapsto (1 - \cos(tx)) \frac{1+x^2}{x^2}, x \in \mathbb{R}$, which is equal to $\frac{t^2}{2}$ at $x = 0$ for the continuity by the well known convention. Due to the inequality $1 - \cos y \leq \frac{y^2}{2}, y \in \mathbb{R}$, for the case $|tx| \leq 2$ we have

$$(1 - \cos(tx)) \frac{1+x^2}{x^2} \leq \frac{t^2 x^2}{2} \cdot \frac{1+x^2}{x^2} = \frac{t^2 + t^2 x^2}{2} \leq \frac{t^2}{2} + 2.$$

Using the simple inequality $1 - \cos y \leq 2, y \in \mathbb{R}$, for the case $|tx| > 2$ we obtain

$$(1 - \cos(tx)) \frac{1+x^2}{x^2} \leq 2 \cdot \frac{1+x^2}{x^2} = 2 \cdot \left(\frac{1}{x^2} + 1 \right) \leq 2 \cdot \left(\frac{t^2}{4} + 1 \right) = \frac{t^2}{2} + 2.$$

Thus

$$(31) \quad (1 - \cos(tx)) \frac{1+x^2}{x^2} \leq \frac{t^2}{2} + 2 \quad \text{for any } x \in \mathbb{R}, t \in \mathbb{R}.$$

Thus for any $n \in \mathbb{N}$ and $t \in \mathbb{R}$ we obtain

$$(32) \quad |f_n(t)| \geq \exp \left\{ - \int_{\mathbb{R}} \left(\frac{t^2}{2} + 2 \right) d|G_n|(x) \right\} = \exp \left\{ - \left(\frac{t^2}{2} + 2 \right) \|G_n\| \right\} > 0.$$

Hence, due to (14) and (29), we have

$$|f(t)| = \lim_{n \rightarrow \infty} |f_n(t)| \geq \exp \left\{ - \left(\frac{t^2}{2} + 2 \right) \overline{\lim}_{n \rightarrow \infty} \|G_n\| \right\} = \exp \left\{ - \left(\frac{t^2}{2} + 2 \right) B \right\} > 0, \quad t \in \mathbb{R},$$

i.e., $f(t) \neq 0$ for any $t \in \mathbb{R}$.

Due to the above remarks, the distinguished logarithms $t \mapsto \text{Ln } f(t)$ and $t \mapsto \text{Ln } f_n(t)$, $n \in \mathbb{N}$, are defined for all $t \in \mathbb{R}$. According to (11), we have

$$(33) \quad \text{Ln } f_n(t) = it\gamma_n + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x), \quad t \in \mathbb{R}, n \in \mathbb{N}.$$

Due to the convergence (29), we have that

$$(34) \quad \text{Ln } f_n(t) \rightarrow \text{Ln } f(t), \quad n \rightarrow \infty \quad \text{for every } t \in \mathbb{R}.$$

Hence, in particular,

$$\gamma_n = \frac{\text{Im}(\text{Ln } f_n(\tau))}{\tau} \rightarrow \frac{\text{Im}(\text{Ln } f(\tau))}{\tau} \in \mathbb{R}, \quad n \rightarrow \infty.$$

We denote this limit by γ . So we have

$$(35) \quad \gamma_n \rightarrow \gamma, \quad n \rightarrow \infty.$$

We next introduce the following functions

$$\psi(t, s) := \text{Ln } f(t) - \frac{1}{2}(\text{Ln } f(t-s) + \text{Ln } f(t+s)), \quad t \in \mathbb{R}, s \geq 0,$$

and analogously

$$(36) \quad \psi_n(t, s) := \text{Ln } f_n(t) - \frac{1}{2}(\text{Ln } f_n(t-s) + \text{Ln } f_n(t+s)), \quad t \in \mathbb{R}, s \geq 0, n \in \mathbb{N}.$$

From (34) we conclude that

$$(37) \quad \psi_n(t, s) \rightarrow \psi(t, s), \quad n \rightarrow \infty \quad \text{for any } t \in \mathbb{R}, s \geq 0.$$

Moreover, since (30) implies the convergence (see [20, p. 15], or [31, p. 34])

$$\sup_{t \in [-T, T]} |\text{Ln } f_n(t) - \text{Ln } f(t)| \rightarrow 0, \quad n \rightarrow \infty \quad \text{for any } T > 0,$$

it is clear that

$$(38) \quad \sup_{t, s \in [-T, T]} |\psi_n(t, s) - \psi(t, s)| \rightarrow 0, \quad n \rightarrow \infty \quad \text{for any } T > 0.$$

We next show that ψ_n , $n \in \mathbb{N}$, are uniformly bounded over $t \in \mathbb{R}$ and $n \in \mathbb{N}$ for any fixed $s \geq 0$. Using (33) in (36), we have

$$(39) \quad \begin{aligned} \psi_n(t, s) &= it\gamma_n + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \\ &\quad - \frac{1}{2} \left(i2t\gamma_n + \int_{\mathbb{R}} \left(e^{itx} (e^{-isx} + e^{isx}) - 2 - \frac{i2t}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \right) \\ &= \int_{\mathbb{R}} e^{itx} (1 - \cos(sx)) \frac{1+x^2}{x^2} dG_n(x), \quad t \in \mathbb{R}, s \geq 0, n \in \mathbb{N}. \end{aligned}$$

The estimate (31) yields

$$\begin{aligned} \sup_{t \in \mathbb{R}} |\psi_n(t, s)| &\leq \sup_{t \in \mathbb{R}} \int_{\mathbb{R}} \left| e^{itx} (1 - \cos(sx)) \frac{1+x^2}{x^2} \right| d|G_n|(x) \\ &= \int_{\mathbb{R}} (1 - \cos(sx)) \frac{1+x^2}{x^2} d|G_n|(x) \leq \left(\frac{s^2}{2} + 1 \right) \|G_n\|, \quad s \geq 0, \quad n \in \mathbb{N}. \end{aligned}$$

According to (14), there exists a constant $B_0 \geq 0$ such that $\|G_n\| \leq B_0$ for all $n \in \mathbb{N}$. Then we conclude

$$(40) \quad \sup_{n \in \mathbb{N}} \sup_{t \in \mathbb{R}} |\psi_n(t, s)| \leq B_0 \cdot \left(\frac{s^2}{2} + 1 \right), \quad s \geq 0.$$

Additionally, in view of (37), we obtain

$$(41) \quad \sup_{t \in \mathbb{R}} |\psi(t, s)| \leq B_0 \cdot \left(\frac{s^2}{2} + 1 \right), \quad s \geq 0.$$

Next, since $\int_0^\infty (s^2 + 1) e^{-s} ds < \infty$, we can define the functions

$$g_n(t) := \int_0^\infty \psi_n(t, s) e^{-s} ds, \quad n \in \mathbb{N} \quad \text{and} \quad g(t) := \int_0^\infty \psi(t, s) e^{-s} ds, \quad t \in \mathbb{R},$$

and, due to (37), conclude at once by the Lebesgue dominated convergence theorem that

$$g_n(t) \rightarrow g(t), \quad n \rightarrow \infty \quad \text{for every } t \in \mathbb{R}.$$

Let us prove that

$$(42) \quad \sup_{t \in [-T, T]} |g_n(t) - g(t)| \rightarrow 0, \quad n \rightarrow \infty \quad \text{for any } T > 0.$$

We fix any $T > 0$ and $\varepsilon > 0$. It is clear that for every $n \in \mathbb{N}$

$$(43) \quad \sup_{t \in [-T, T]} |g_n(t) - g(t)| \leq \int_0^\infty \sup_{t \in [-T, T]} |\psi_n(t, s) - \psi(t, s)| e^{-s} ds.$$

We denote by $J_n(T)$ the last integral for every $n \in \mathbb{N}$. Let us choose a constant $h_\varepsilon > 0$ such that

$$(44) \quad B_0 \int_{h_\varepsilon}^\infty (s^2 + 2) e^{-s} ds < \varepsilon.$$

Then we write $J_n(T) = J_{n,1}(T) + J_{n,2}(T)$, $n \in \mathbb{N}$, where

$$\begin{aligned} J_{n,1}(T) &:= \int_0^{h_\varepsilon} \sup_{t \in [-T, T]} |\psi_n(t, s) - \psi(t, s)| e^{-s} ds, \\ J_{n,2}(T) &:= \int_{h_\varepsilon}^\infty \sup_{t \in [-T, T]} |\psi_n(t, s) - \psi(t, s)| e^{-s} ds. \end{aligned}$$

All the integrals $J_n(T)$, $J_{n,1}(T)$, and $J_{n,2}(T)$ are nonnegative. Observe that

$$\begin{aligned} J_{n,1}(T) &\leq \sup_{\substack{t \in [-T, T] \\ s \in [0, h_\varepsilon]}} |\psi_n(t, s) - \psi(t, s)| \int_0^{h_\varepsilon} e^{-s} ds \\ &\leq \sup_{t, s \in [-T_\varepsilon, T_\varepsilon]} |\psi_n(t, s) - \psi(t, s)|, \quad n \in \mathbb{N}, \end{aligned}$$

where $T_\varepsilon := \max\{T, h_\varepsilon\}$. Due to (38), the last supremum vanishes as $n \rightarrow \infty$. So there exists $n_\varepsilon \in \mathbb{N}$ such that $J_{n,1}(T) < \varepsilon$ for any $n \geq n_\varepsilon$. Let us turn to $J_{n,2}(T)$. According to (40), (41), and (44), we have

$$J_{n,2}(T) \leq \int_{h_\varepsilon}^\infty \left(\sup_{t \in \mathbb{R}} |\psi_n(t, s)| + \sup_{t \in \mathbb{R}} |\psi(t, s)| \right) e^{-s} ds \leq \int_{h_\varepsilon}^\infty B_0(s^2 + 2) e^{-s} ds < \varepsilon.$$

Then $J_n(T) = J_{n,1}(T) + J_{n,2}(T) < 2\varepsilon$ for any $n \geq n_\varepsilon$. Since $\varepsilon > 0$ was chosen arbitrarily, $J_n(T) \rightarrow 0$ as $n \rightarrow \infty$. Thus, according to (43), we obtain (42). Since g is a uniform limit of continuous functions g_n on any segment $[-T, T]$ as $n \rightarrow \infty$, the function g is continuous on \mathbb{R} .

Let us consider the functions g_n , $n \in \mathbb{N}$. Using (39), we write

$$\begin{aligned} g_n(t) &= \int_0^\infty \psi_n(t, s) e^{-s} ds \\ &= \int_0^\infty \left(\int_{\mathbb{R}} e^{itx} (1 - \cos(sx)) \frac{1+x^2}{x^2} dG_n(x) \right) e^{-s} ds \\ &= \int_{\mathbb{R}} \left(\int_0^\infty (1 - \cos(sx)) e^{-s} ds \right) e^{itx} \frac{1+x^2}{x^2} dG_n(x), \quad t \in \mathbb{R}, n \in \mathbb{N}. \end{aligned}$$

The inner integral is calculated (see [14, p. 486, formula 3.893(2)]):

$$\int_0^\infty (1 - \cos(sx)) e^{-s} ds = 1 - \int_0^\infty \cos(sx) e^{-s} ds = 1 - \frac{1}{1+x^2} = \frac{x^2}{1+x^2}, \quad x \in \mathbb{R}.$$

Therefore we have

$$g_n(t) = \int_{\mathbb{R}} e^{itx} dG_n(x), \quad t \in \mathbb{R}, n \in \mathbb{N}.$$

We now use Theorem 5. So there exists a unique function $G \in V$ such that $\|G\| \leq B$ and the equality

$$g(t) = \int_{\mathbb{R}} e^{itx} dG(x)$$

holds for all $t \in \mathbb{R}$, because g is continuous on \mathbb{R} . Moreover, due to the theorem, we have $G_n \Rightarrow G$ and also $G_n(+\infty) \rightarrow G(+\infty)$, $n \rightarrow \infty$.

We now prove that for any $t \in \mathbb{R}$ and $\tau > 0$

$$(45) \quad \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \\ \rightarrow \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG(x), \quad n \rightarrow \infty.$$

Let us fix $t \in \mathbb{R}$ and $\tau > 0$. From Lemma 20 we know that

$$\left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} = \int_{A_{t,\tau}} e^{isx} dW_{t,\tau}(s), \quad x \in \mathbb{R},$$

where $A_{t,\tau} = \{s \in \mathbb{R} : |s| \leq \max\{|t|, \tau\}\}$ and $W_{t,\tau} \in V$. Hence for every $n \in \mathbb{N}$

$$\int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \\ = \int_{\mathbb{R}} \left(\int_{A_{t,\tau}} e^{isx} dW_{t,\tau}(s) \right) dG_n(x) \\ = \int_{A_{t,\tau}} \left(\int_{\mathbb{R}} e^{isx} dG_n(x) \right) dW_{t,\tau}(s) = \int_{A_{t,\tau}} g_n(s) dW_{t,\tau}(s).$$

Also we have analogously that

$$\int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG(x) = \int_{A_{t,\tau}} g(s) dW_{t,\tau}(s).$$

Thus (45) takes the form

$$\int_{A_{t,\tau}} g_n(s) dW_{t,\tau}(s) \rightarrow \int_{A_{t,\tau}} g(s) dW_{t,\tau}(s), \quad n \rightarrow \infty.$$

This convergence holds. Indeed, for every $n \in \mathbb{N}$

$$\left| \int_{A_{t,\tau}} g_n(s) dW_{t,\tau}(s) - \int_{A_{t,\tau}} g(s) dW_{t,\tau}(s) \right| \leq \int_{A_{t,\tau}} |g_n(s) - g(s)| d|W_{t,\tau}|(s) \\ \leq \sup_{s \in A_{t,\tau}} |g_n(s) - g(s)| \cdot \|W_{t,\tau}\|,$$

where, due to (42), the supremum vanishes as $n \rightarrow \infty$. Thus we proved (45).

From (33), (35), and (45), for any $t \in \mathbb{R}$ we have

$$\text{Ln } f_n(t) = it\gamma_n + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG_n(x) \\ \rightarrow it\gamma + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG(x), \quad n \rightarrow \infty.$$

According to (34), we conclude that

$$\text{Ln } f(t) = it\gamma + \int_{\mathbb{R}} \left(e^{itx} - 1 - \frac{it}{\tau} \sin(\tau x) \right) \frac{1+x^2}{x^2} dG(x), \quad t \in \mathbb{R},$$

where, as we have already proved, $\gamma \in \mathbb{R}$ and $G \in \mathbf{V}$. Thus f has the Lévy–Khinchin type representation with (γ, G) , i.e., the distribution function F corresponding to f is quasi-infininitely divisible. \square

Proof of Theorem 11. Let f be a characteristic function of the limit distribution function F . So we have (29) and also (30) (see comments in the proof of Theorem 10).

Recall that $f_n(t) \neq 0$, $t \in \mathbb{R}$, $n \in \mathbb{N}$. Let us choose $\delta > 0$ such that $f(t) \neq 0$, $|t| \leq \delta$ (it is possible because f is continuous on \mathbb{R} and $f(0) = 1$). Let us consider values of the Khinchin functional $\chi_\delta(\cdot)$ (see [24, p. 79]) with parameter δ on f and f_n , $n \in \mathbb{N}$:

$$\chi_\delta(f) = -\frac{1}{\delta} \int_0^\delta \ln |f(s)| ds, \quad \chi_\delta(f_n) = -\frac{1}{\delta} \int_0^\delta \ln |f_n(s)| ds, \quad n \in \mathbb{N}.$$

These quantities are finite and nonnegative. Due to (30), we have

$$(46) \quad \chi_\delta(f_n) \rightarrow \chi_\delta(f), \quad n \rightarrow \infty.$$

Observe that

$$\begin{aligned} (47) \quad \chi_\delta(f_n) &= -\frac{1}{\delta} \int_0^\delta \left(\int_{\mathbb{R}} (\cos(sx) - 1) \frac{1+x^2}{x^2} dG_n(x) \right) ds \\ &= \int_{\mathbb{R}} \left(\frac{1}{\delta} \int_0^\delta (1 - \cos(sx)) ds \right) \frac{1+x^2}{x^2} dG_n(x) \\ &= \int_{\mathbb{R}} \left(1 - \frac{\sin(\delta x)}{\delta x} \right) \frac{1+x^2}{x^2} dG_n(x), \quad n \in \mathbb{N}. \end{aligned}$$

where we set

$$(48) \quad (\cos(sx) - 1) \frac{1+x^2}{x^2} \Big|_{x=0} = -\frac{s^2}{2}, \quad \left(1 - \frac{\sin(\delta x)}{\delta x} \right) \frac{1+x^2}{x^2} \Big|_{x=0} = \frac{\delta^2}{3!},$$

according to known expansions $\cos y = 1 - \frac{y^2}{2} + o(y^2)$ and $\sin y = y - \frac{y^3}{3!} + o(y^3)$, $y \rightarrow 0$. Let us consider the inner function of the integral in (47):

$$(49) \quad x \mapsto \left(1 - \frac{\sin(\delta x)}{\delta x} \right) \frac{1+x^2}{x^2}, \quad x \in \mathbb{R}.$$

By convention (48), it is continuous at the point $x = 0$. We see that this function is continuous and strictly positive on \mathbb{R} . Also observe that it tends to 1 as $x \rightarrow \pm\infty$.

Hence it is clear that there exist positive constants c_δ and C_δ such that

$$(50) \quad 0 < c_\delta \leq \left(1 - \frac{\sin(\delta x)}{\delta x}\right) \frac{1+x^2}{x^2} \leq C_\delta < \infty, \quad x \in \mathbb{R}.$$

Let us take some decompositions (12) for G_n , $n \in \mathbb{N}$. According to (47):

$$\begin{aligned} & \chi_\delta(f_n) \\ &= \int_{\mathbb{R}} \left(1 - \frac{\sin(\delta x)}{\delta x}\right) \frac{1+x^2}{x^2} dG_n^+(x) - \int_{\mathbb{R}} \left(1 - \frac{\sin(\delta x)}{\delta x}\right) \frac{1+x^2}{x^2} dG_n^-(x), \quad n \in \mathbb{N}. \end{aligned}$$

Due to (50), we obtain

$$\chi_\delta(f_n) \geq c_\delta \int_{\mathbb{R}} dG_n^+(x) - C_\delta \int_{\mathbb{R}} dG_n^-(x) = c_\delta G_n^+(\infty) - C_\delta G_n^-(\infty), \quad n \in \mathbb{N}.$$

From this we have

$$G_n^+(\infty) \leq \frac{\chi_\delta(f_n) + C_\delta G_n^-(\infty)}{c_\delta}, \quad n \in \mathbb{N}.$$

Hence, due to (15) and (46), we get

$$(51) \quad \begin{aligned} \overline{\lim}_{n \rightarrow \infty} G_n^+(\infty) &\leq \frac{1}{c_\delta} \left(\overline{\lim}_{n \rightarrow \infty} \chi_\delta(f_n) + C_\delta \overline{\lim}_{n \rightarrow \infty} G_n^-(\infty) \right) \\ &= \frac{1}{c_\delta} (\chi_\delta(f) + C_\delta M) < \infty. \end{aligned}$$

According to (12) and conventions there, it is true that

$$(52) \quad \|G_n\| \leq \|G_n^+\| + \|G_n^-\| = G_n^+(\infty) + G_n^-(\infty), \quad n \in \mathbb{N}.$$

So we conclude from (15) and (51) that (14) holds for some $B < \infty$.

Thus all assertions of Theorem 10 hold. In particular, $G_n \Rightarrow G$ and $G_n(\infty) \rightarrow G(\infty)$, $n \rightarrow \infty$, where G is some function from \mathcal{V} . It remains to prove that $B \leq G(\infty) + 2M$. Using inequality (52), we write

$$\begin{aligned} B &= \overline{\lim}_{n \rightarrow \infty} \|G_n\| \leq \overline{\lim}_{n \rightarrow \infty} (G_n^+(\infty) + G_n^-(\infty)) \\ &\leq \overline{\lim}_{n \rightarrow \infty} (G_n^+(\infty) - G_n^-(\infty)) + 2 \overline{\lim}_{n \rightarrow \infty} G_n^-(\infty), \end{aligned}$$

but $G_n(\infty) = G_n^+(\infty) - G_n^-(\infty)$, $n \in \mathbb{N}$, and we obtain

$$B \leq \overline{\lim}_{n \rightarrow \infty} G_n(\infty) + 2 \overline{\lim}_{n \rightarrow \infty} G_n^-(\infty) = G(\infty) + 2M,$$

as required. □

Proof of Theorem 14. Let $(F_{n_k})_{k \in \mathbb{N}}$ be an arbitrary subsequence of $(F_n)_{n \in \mathbb{N}}$, which weakly converges to some distribution function F_* . Due to the assumption of relative compactness of $(F_n)_{n \in \mathbb{N}}$, such subsequence exists. By Theorem 10, F_* is quasi-infinitely divisible with some spectral pair (γ_*, G_*) , where $\gamma_* \in \mathbb{R}$ and $G_* \in \mathcal{V}$.

Moreover, $\gamma_{n_k} \rightarrow \gamma_*$ and $G_{n_k} \Rightarrow G_*$, $k \rightarrow \infty$. According to the assumption that $\gamma_n \rightarrow \gamma$, $n \rightarrow \infty$, we conclude that $\gamma_* = \gamma$. Let us show that $G_* = G$. By definition, the convergence $G_{n_k} \Rightarrow G_*$, $k \rightarrow \infty$, implies the existence of a subsequence $(G_{n'_l})_{l \in \mathbb{N}}$ in $(G_{n_k})_{k \in \mathbb{N}}$ such that

$$G_{n'_l}(x_2) - G_{n'_l}(x_1) \rightarrow G_*(x_2) - G_*(x_1), \quad l \rightarrow \infty$$

for all $x_1, x_2 \in \mathbb{R}$ except at most countable set D' . Due to the assumption that $G_n \Rightarrow G$, $n \rightarrow \infty$, we can choose a further subsequence $(G_{n''_l})_{l \in \mathbb{N}}$ in $(G_{n'_l})_{l \in \mathbb{N}}$ such that

$$G_{n''_l}(x_2) - G_{n''_l}(x_1) \rightarrow G(x_2) - G(x_1), \quad l \rightarrow \infty$$

for all $x_1, x_2 \in \mathbb{R}$ except at most countable set D'' (and let $D' \neq D''$ in general). Therefore

$$G_*(x_2) - G_*(x_1) = G(x_2) - G(x_1)$$

for all $x_1, x_2 \in \mathbb{R}$ except at most countable set $D' \cup D''$. Letting $x_1 \rightarrow -\infty$ over $x_1 \in \mathbb{R} \setminus (D' \cup D'')$ we have $G_*(x_2) = G(x_2)$ for every $x_2 \in \mathbb{R} \setminus (D' \cup D'')$ and, consequently for all $x_2 \in \mathbb{R}$, because G_* , $G \in \mathcal{V}$, i.e., they are right-continuous and

$$G_*(-\infty) = G(-\infty) = 0.$$

Thus we proved that $\gamma_* = \gamma$ and $G_* = G$.

The previous remark means that (γ, G) is the spectral pair for some quasi-infininitely divisible distribution function F . We also saw that every subsequence $(F_{n_k})_{k \in \mathbb{N}}$, which weakly converges to some distribution function, converges exactly to F , because a spectral pair uniquely determines a distribution function. Therefore, since $(F_n)_{n \in \mathbb{N}}$ is relatively compact, we conclude that whole sequence $(F_n)_{n \in \mathbb{N}}$ weakly converges to F (this is a known fact, see [8, p. 337]). \square

5. Acknowledgment

This research was supported by the Ministry of Science and Higher Education of the Russian Federation, agreement 075-15-2019-1620 date 08/11/2019 and 075-15-2022-289 date 06/04/2022.

References

- [1] I. A. Alexeev and A. A. Khartov, "On convergence and compactness in variation with a shift of discrete probability laws", *Vestn. St.-Peterbg. Univ. Mat. Mekh. Astron.* **8**:3 (2021), 385–393. In Russian; translated in *Vestnik St. Petersburg Univ. Math.* **54**:3 (2021), 221–226. MR Zbl
- [2] I. Alexeev and A. Khartov, "Spectral representations of characteristic functions of discrete probability laws", *Bernoulli* **29**:2 (2023), 1392–1409. MR Zbl
- [3] D. Applebaum, *Lévy processes and stochastic calculus*, Cambridge Stud. Adv. Math. **93**, Cambridge Univ. Press, 2004. MR Zbl

- [4] D. Berger, “On quasi-infinitely divisible distributions with a point mass”, *Math. Nachr.* **292**:8 (2019), 1674–1684. MR Zbl
- [5] D. Berger and M. Kutlu, “Quasi-infinite divisibility of a class of distributions with discrete part”, *Proc. Amer. Math. Soc.* **151**:5 (2023), 2211–2224. MR Zbl
- [6] D. Berger and A. Lindner, “A Cramér–Wold device for infinite divisibility of \mathbb{Z}^d -valued distributions”, *Bernoulli* **28**:2 (2022), 1276–1283. MR Zbl
- [7] D. Berger, M. Kutlu, and A. Lindner, “On multivariate quasi-infinitely divisible distributions”, pp. 87–120 in *A lifetime of excursions through random walks and Lévy processes*, edited by L. Chaumont and A. E. Kyprianou, Progr. Probab. **78**, Birkhäuser, Cham, 2021. MR Zbl
- [8] P. Billingsley, *Probability and measure*, 3rd ed., Wiley, New York, 1995. MR Zbl
- [9] S. Bochner, “A theorem on Fourier–Stieltjes integrals”, *Bull. Amer. Math. Soc.* **40**:4 (1934), 271–276. MR Zbl
- [10] V. I. Bogachev, *Weak convergence of measures*, Math. Surv. Monogr. **234**, Amer. Math. Soc., Providence, RI, 2018. MR Zbl
- [11] H. Chhaiba, N. Demni, and Z. Mouayn, “Analysis of generalized negative binomial distributions attached to hyperbolic Landau levels”, *J. Math. Phys.* **57**:7 (2016), art. id. 072103. MR Zbl
- [12] N. Demni and Z. Mouayn, “Analysis of generalized Poisson distributions associated with higher Landau levels”, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **18**:4 (2015), art. id. 1550028. MR Zbl
- [13] B. V. Gnedenko and A. N. Kolmogorov, *Limit distributions for sums of independent random variables*, Addison-Wesley, Cambridge, MA, 1954. MR Zbl
- [14] I. S. Gradshteyn and I. M. Ryzhik, “Definite integrals of elementary functions”, pp. 249–519 in *Table of integrals, series, and products*, 7th ed., Elsevier, Amsterdam, 2007. MR Zbl
- [15] A. A. Khartov, “Characteristic functions and compactness of distributions of sums of independent random variables”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **454**:24 (2016), 292–308. In Russian; translated in *J. Math. Sci. (N.Y.)* **229**:6 (2018), 792–802. MR Zbl
- [16] A. A. Khartov, “Criteria of relative and stochastic compactness for distributions of sums of independent random variables”, *Teor. Veroyatn. Primen.* **63**:1 (2018), 70–88. In Russian; translated in *Theory Probab. Appl.* **63**:1 (2018), 57–71. MR Zbl
- [17] A. A. Khartov, “Compactness criteria for quasi-infinitely divisible distributions on the integers”, *Statist. Probab. Lett.* **153** (2019), 1–6. MR Zbl
- [18] A. A. Khartov, “A criterion of quasi-infinite divisibility for discrete laws”, *Statist. Probab. Lett.* **185** (2022), art. id. 109436. MR Zbl
- [19] A. A. Khartov and I. A. Alexeev, “Quasi-infinite divisibility and three-point probability laws”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **495**:29 (2020), 305–316. In Russian; translated in *J. Math. Sci. (N.Y.)* **268**:5 (2022), 731–738. MR Zbl
- [20] A. Y. Khinchin, *Предельные законы для сумм независимых случайных величин*, ONTI, Moscow, 1938.
- [21] M. Kutlu, “On a denseness result for quasi-infinitely divisible distributions”, *Statist. Probab. Lett.* **176** (2021), art. id. 109139. MR Zbl
- [22] A. Lindner and K.-i. Sato, “Properties of stationary distributions of a sequence of generalized Ornstein–Uhlenbeck processes”, *Math. Nachr.* **284**:17–18 (2011), 2225–2248. MR Zbl
- [23] A. Lindner, L. Pan, and K.-i. Sato, “On quasi-infinitely divisible distributions”, *Trans. Amer. Math. Soc.* **370**:12 (2018), 8483–8520. MR Zbl

- [24] J. V. Linnik and I. V. Ostrovskii, *Decomposition of random variables and vectors*, Transl. Math. Monogr. **48**, Amer. Math. Soc., Providence, RI, 1977. MR Zbl
- [25] E. Lukacs, *Characteristic functions*, 2nd ed., Hafner, New York, 1970. MR Zbl
- [26] T. Nakamura, “A complete Riemann zeta distribution and the Riemann hypothesis”, *Bernoulli* **21**:1 (2015), 604–617. MR Zbl
- [27] I. P. Natanson, *Theory of functions of a real variable*, revised ed., Ungar, New York, 1983.
- [28] R. Passeggeri, “Spectral representations of quasi-infinately divisible processes”, *Stochastic Process. Appl.* **130**:3 (2020), 1735–1791. MR Zbl
- [29] R. S. Phillips, “On Fourier–Stieltjes integrals”, *Trans. Amer. Math. Soc.* **69** (1950), 312–323. MR Zbl
- [30] A. Rocha-Arteaga and K.-i. Sato, *Topics in infinitely divisible distributions and Lévy processes*, Aportaciones Mat. Investigación **17**, Soc. Mat. Mexicana, Mexico City, 2003. MR Zbl
- [31] K.-i. Sato, *Lévy processes and infinitely divisible distributions*, Cambridge Stud. Adv. Math. **68**, Cambridge Univ. Press, 1999. MR Zbl
- [32] H. Zhang, Y. Liu, and B. Li, “Notes on discrete compound Poisson model with applications to risk theory”, *Insurance Math. Econom.* **59** (2014), 325–336. MR Zbl
- [33] V. M. Zolotarev, *One-dimensional stable distributions*, Transl. Math. Monogr. **65**, Amer. Math. Soc., Providence, RI, 1986. MR Zbl
- [34] V. M. Zolotarev, *Modern theory of summation of random variables*, de Gruyter, Berlin, 1997. MR Zbl

Received June 15, 2022.

ALEXEY KHARTOV
LABORATORY FOR APPROXIMATION PROBLEMS OF PROBABILITY
SMOLENSK STATE UNIVERSITY
SMOLENSK
RUSSIA
alexeykharov@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 322 No. 2 February 2023

Elements of higher homotopy groups undetectable by polyhedral approximation	221
JOHN K. ACETI and JEREMY BRAZAS	
Regularity for free multiplicative convolution on the unit circle	243
SERBAN T. BELINSCHI, HARI BERCOVICI and CHING-WEI HO	
Invariant theory for the free left-regular band and a q -analogue	251
SARAH BRAUNER, PATRICIA COMMINS and VICTOR REINER	
Irredundant bases for finite groups of Lie type	281
NICK GILL and MARTIN W. LIEBECK	
Local exterior square and Asai L -functions for $GL(n)$ in odd characteristic	301
YEONGSEONG JO	
On weak convergence of quasi-infinitely divisible laws	341
ALEXEY KHARTOV	
C^* -irreducibility of commensurated subgroups	369
KANG LI and EDUARDO SCARPARO	
Local Maass forms and Eichler–Selberg relations for negative-weight vector-valued mock modular forms	381
JOSHUA MALES and ANDREAS MONO	
Representations of orientifold Khovanov–Lauda–Rouquier algebras and the Enomoto–Kashiwara algebra	407
TOMASZ PRZEŹDZIECKI	